

THE CONFORMAL FLOW OF METRICS AND THE GENERAL PENROSE INEQUALITY

QING HAN AND MARCUS KHURI

ABSTRACT. The conformal flow of metrics [2] has been used to successfully establish a special case of the Penrose inequality, which yields a lower bound for the total mass of a spacetime in terms of horizon area. Here we show how to adapt the conformal flow of metrics, so that it may be applied to the Penrose inequality for general initial data sets of the Einstein equations. The Penrose conjecture without the assumption of time symmetry is then reduced to solving a system of PDE with desirable properties.

1. INTRODUCTION

Let (M, g, k) be an initial data set for the Einstein equations with a single asymptotically flat end. This triple consists of a 3-manifold M , on which a Riemannian metric g and symmetric 2-tensor k are defined and satisfy the constraint equations

$$(1.1) \quad 16\pi\mu = R + (Trk)^2 - |k|^2, \quad 8\pi J = \operatorname{div}(k + (Trk)g).$$

The quantities μ and J represent the energy and momentum densities of the matter fields, respectively, whereas R denotes the scalar curvature of g . The dominant energy condition will be assumed $\mu \geq |J|$. This asserts that all measured energy densities are nonnegative, and implies that matter cannot travel faster than the speed of light.

Null expansions measure the strength of the gravitational field around a hypersurface $S \subset M$ and are given by

$$(1.2) \quad \theta_{\pm} := H_S \pm Tr_S k,$$

where H_S denotes the mean curvature with respect to the unit normal pointing towards spatial infinity. More precisely, $\theta_{\pm}$ may be interpreted as the rate at which the area of a shell of light is changing as it moves away from the surface in the outward future/past direction $(+/-)$. A future or past trapped surface is defined by the inequality $\theta_{+} < 0$ or $\theta_{-} < 0$, respectively, and according to the above interpretation such a surface lies in a strong gravitation field. If $\theta_{+} = 0$ or $\theta_{-} = 0$, then S is referred to as a future or past apparent horizon. These surfaces naturally arise as boundaries of future or past trapped regions. An apparent horizon will be referred to as outermost if it is not enclosed by any other apparent horizon. According to [9, 10] each component of an outermost apparent horizon must have spherical topology. These surfaces replace the role of event horizons in the formulation of the Penrose inequality. In particular, under the above hypotheses on the initial data, the standard version of the Penrose conjecture [19] asserts that

$$(1.3) \quad M_{ADM} \geq \sqrt{\frac{A_{min}^{\pm}}{16\pi}}$$

The first author acknowledges the support of NSF Grant DMS-1105321. The second author acknowledges the support of NSF Grant DMS-1708798.

where M_{ADM} is the ADM mass and A_{min}^+ (A_{min}^-) is the minimum area required to enclose the outermost future (past) apparent horizon. The Riemannian version of this inequality in the time symmetric case, when $k = 0$, along with the corresponding rigidity statement in the case of equality, have been established. For a single component horizon the proof was given by Huisken and Ilmanen in [12] using inverse mean curvature flow, and for a multiple component horizon the proof was given by Bray in [2] using the conformal flow of metrics. However for general k , the inequality (1.3) remains an important open problem.

In this paper we aim to study a slightly different inequality. Suppose that the initial data have boundary ∂M , consisting of an outermost apparent horizon. Each component of the outermost apparent horizon is either a future or past horizon, however it will be assumed that these two types of components do not intersect. Then a version of the Penrose conjecture asserts that

$$(1.4) \quad M_{ADM} \geq \sqrt{\frac{A_{min}}{16\pi}},$$

where A_{min} is the minimum area required to enclose ∂M . Moreover the rigidity statement is as follows. Equality holds in (1.4) if and only if (M, g, k) arises from a spacelike slice of the Schwarzschild spacetime, with outerminimizing boundary.

In [3, 4] Bray and the second author proposed a generalized version of the Jang equation [20], designed specifically for application to the Penrose conjecture. This consists of searching for a graph $\Sigma = \{t = f(x)\}$ within the warped product manifold $(M \times \mathbb{R}, g + \phi^2 dt^2)$. The function $\phi \geq 0$ on M is to be chosen naturally depending on the strategy of the proof, and should reduce to the Schwarzschild warping factor in the case of equality. In [3, 4] the strategy was based on inverse mean curvature flow (IMCF), and although the resulting ϕ had some desirable properties it also has the potential to vanish on the interior of M , rendering the coupling of the generalized Jang equation to IMCF degenerate. The purpose of the current paper is to utilize an alternate strategy, namely the conformal flow of metrics [2], and find a new choice for ϕ (see (3.6) below) which is strictly positive away from ∂M and hence leads to a nondegenerate coupling with the generalized Jang equation.

The generalized Jang equation is motivated by the need for the hypersurface Σ to have weakly nonnegative scalar curvature. As in the classical case [20] this may be achieved by solving

$$(1.5) \quad H_\Sigma - Tr_\Sigma K = 0,$$

where H_Σ denotes mean curvature and K is a symmetric 2-tensor on $M \times \mathbb{R}$ which is an extension of the initial data k . The precise definition of K is given in [3, 4] along with the following formula for the scalar curvature of Σ

$$(1.6) \quad \overline{R} = 16\pi(\mu - J(w)) + |h - K|_\Sigma|^2 + 2|q|^2 - 2\phi^{-1}\overline{\text{div}}(\phi q),$$

where h is the second fundamental form of Σ , $\overline{\text{div}}$ is the divergence operation with respect to the induced metric $\overline{g} = g + \phi^2 df^2$, and q and w are the 1-forms

$$(1.7) \quad w_i = \frac{\phi f_i}{\sqrt{1 + \phi^2 |\nabla f|^2}}, \quad q_i = \frac{\phi f^j}{\sqrt{1 + \phi^2 |\nabla f|^2}} (h_{ij} - (K|_\Sigma)_{ij}).$$

Observe that the dominant energy condition implies that the right-hand side of (1.6) is manifestly nonnegative except for the divergence term. In this way, we may view $\overline{R}$ as being ‘weakly’ nonnegative in that multiplying by ϕ and integrating by parts produces a nonnegative quantity modulo boundary integrals.

The expression for the generalized Jang equation (1.5) in local coordinates is given by

$$(1.8) \quad \left(g^{ij} - \frac{\phi^2 f^i f^j}{1 + \phi^2 |\nabla f|^2} \right) \left(\frac{\phi \nabla_{ij} f + \phi_i f_j + \phi_j f_i}{\sqrt{1 + \phi^2 |\nabla f|^2}} - k_{ij} \right) = 0.$$

Note that this quasilinear elliptic equation is degenerate where f blows-up or where $\phi = 0$. Moreover, this equation reduces to the classical Jang equation utilized by Schoen and Yau in their proof of the positive mass theorem [20], when $\phi = 1$.

In [11] blow-up solutions of the generalized Jang equation have been studied in detail, and the main result may be described as follows. Let $\tau(x) = \text{dist}(x, \partial M)$ and denote the corresponding level sets by S_τ . According to the assumptions above, the boundary may be decomposed into future (+) and past (−) apparent horizon components $\partial M = \partial_+ M \cup \partial_- M$. We then specify that in a neighborhood of $\partial_\pm M$ there are constants $l \geq 1$ and $c > 0$ such that

$$(1.9) \quad |\theta_\pm(S_\tau)| \leq c\tau^l.$$

Suppose also that ϕ is positive on the interior of M , and near ∂M the following structure condition holds

$$(1.10) \quad \phi(x) = \tau^b(x) \tilde{\phi}(x),$$

where the constant b is nonnegative and $\tilde{\phi} > 0$ is a smooth function up to the boundary. If $\frac{1}{2} \leq b < \frac{l+1}{2}$, then there exists a smooth solution f of the generalized Jang equation (1.8) with the property that $f(x) \rightarrow \pm\infty$ as $x \rightarrow \partial_\pm M$. Furthermore, the asymptotics for this blow-up are given by

$$(1.11) \quad \begin{aligned} \alpha^{-1} \tau^{1-2b} + \beta^{-1} &\leq \pm f \leq \alpha \tau^{1-2b} + \beta & \text{if} & \quad \frac{1}{2} < b < \frac{l+1}{2}, \\ -\alpha^{-1} \log \tau + \beta^{-1} &\leq \pm f \leq -\alpha \log \tau + \beta & \text{if} & \quad b = \frac{1}{2}, \end{aligned}$$

for some positive constants α, β . If in addition the warping factor satisfies the following asymptotics at spatial infinity

$$(1.12) \quad \phi(x) = 1 + \frac{C}{|x|} + O\left(\frac{1}{|x|^2}\right) \quad \text{as} \quad |x| \rightarrow \infty$$

where C is a constant, then

$$(1.13) \quad |\nabla^m f|(x) = O(|x|^{-\frac{1}{2}-m}) \quad \text{as} \quad |x| \rightarrow \infty, \quad m = 0, 1, 2.$$

This fall-off for the solution guarantees that the ADM energy of the Jang surface and that of the initial data are the same. Furthermore, the blow-up (1.11) implies that the Jang surface Σ is a manifold with boundary, when $\frac{1}{2} \leq b < 1$. This is in contrast to the blow-up solutions of the classical Jang equation which approximate an infinitely long cylinder over the horizon. Moreover, simple computations shows that $\partial\Sigma$ is a minimal surface, and that the area of the boundaries of the Jang surface and initial data agree $|\partial\Sigma| = |\partial M|$.

It is not always the case that blow-up is the correct boundary behavior for application to the Penrose inequality. Moreover instead of working with M , it will become apparent that we should focus on M_0 , where $M_0 \subset M$ is the region outside of the outermost minimal area enclosure of ∂M . Assuming that the warping factor ϕ vanishes at ∂M_0 and is positive on the interior of M_0 , the correct boundary condition for the generalized Jang equation is

$$(1.14) \quad \overline{H}_{\partial\Sigma_0} = 0$$

where Σ_0 represents the Jang surface over M_0 , and $\overline{H}_{\partial\Sigma_0}$ is the mean curvature of $\partial\Sigma_0$. It is conjectured [4] that the following Neumann type condition at ∂M_0 implies that (1.14) holds:

$$(1.15) \quad \frac{\phi \partial_\nu f(x)}{\sqrt{1 + \phi^2 |\nabla f|^2}} = \begin{cases} +1 & \text{if } \theta_+(x) = 0 \text{ and } \theta_-(x) \neq 0, \\ -1 & \text{if } \theta_+(x) \neq 0 \text{ and } \theta_-(x) = 0, \\ 0 & \text{if } \theta_+(x) = 0 \text{ and } \theta_-(x) = 0, \end{cases}$$

where ν denotes the unit outer normal to ∂M_0 (pointing away from spatial infinity). Notice that since $\phi|_{\partial M_0} = 0$ we still have $|\partial\Sigma_0| = |\partial M_0|$, even if blow-up does not occur.

These results show that the Jang surface $(\Sigma, \bar{g})$ may be interpreted as a deformation of the initial data (M, g, k) , which preserves the geometric quantities involved in the Penrose inequality, and yields weakly nonnegative scalar curvature. All of this suggests that in order to establish (1.4), one should try to apply the techniques used to prove the Riemannian Penrose inequality, inside the Jang surface. In fact such a method was outlined in [3] and [4], for the inverse mean curvature flow. This leads to a coupling of the inverse mean curvature flow with the generalized Jang equation, through a specific choice of warping factor ϕ . Unfortunately, this choice of warping factor lacks regularity and vanishes identically when the weak inverse mean curvature flow jumps, which in turn leads to a very degenerate generalized Jang equation. Thus, it is of significant interest to find a more appropriate choice for ϕ that leads to a better system of equations. In [3, 4] it was conjectured that such a ϕ exists, and may be found by coupling the generalized Jang equation to the conformal flow of metrics. It is the purpose of this paper to confirm this, and to give an explicit and simple expression for ϕ (see (3.6)). This choice of warping factor is positive on the interior of M , and has better regularity properties than that arising from the inverse mean curvature flow. It will also be shown that (3.6) reduces to the correct expression, that is, the warping factor for the Schwarzschild spacetime, in the case of equality in (1.4).

Theorem 1.1. *The Penrose conjecture (1.4), along with its corresponding rigidity statement, reduce to the problem of solving a system of PDE arising from a natural coupling of the (nondegenerate) generalized Jang equation and the conformal flow of metrics.*

Beyond the Penrose inequality, the generalized Jang equation has been applied to other related geometric inequalities in [8, 7, 13, 14, 15]. Moreover in [5, 6], a further modification of the Jang equation has been adapted to treat inequalities for which the model spacetime is stationary. Lastly, a charged version of the conformal flow has been developed in [16, 17, 18], and was used to establish the Penrose inequality with charge for multiple black holes, in the time-symmetric case. It is likely that the methods of the current note may be extended to this setting, yielding a coupling of the charged conformal flow with the generalized Jang equation, which may be applied to the Penrose inequality with charge in the non-time-symmetric case.

2. THE CONFORMAL FLOW OF METRICS AND A REFINED VERSION OF THE RIEMANNIAN PENROSE INEQUALITY

The purpose of this section is to review the basic properties of the conformal flow of metrics [2], and to obtain a strengthened version of the Riemannian inequality. We will also slightly modify a portion of Bray's proof of the Riemannian Penrose inequality, so as to make the conformal flow applicable to the setting described in the introduction.

Let (M, g) be a 3-dimensional Riemannian manifold, which is asymptotically flat (having one end), and has an outerminimizing minimal surface boundary consisting of a finite number of components.

Unlike [2], however, we do *not* assume that the scalar curvature R_g is nonnegative. The conformal flow of metrics is given by $g_t = u_t^4 g$ where

$$(2.1) \quad \frac{d}{dt} u_t = v_t u_t,$$

and

$$(2.2) \quad \Delta_{g_t} v_t = 0, \quad \text{on } M_t,$$

$$(2.3) \quad v_t|_{\partial M_t} = 0, \quad v_t(x) \rightarrow -1 \quad \text{as } |x| \rightarrow \infty.$$

Here M_t denotes the region outside of the outermost minimal surface (denoted ∂M_t) in (M, g_t) . Note that $u_t(x) = \exp\left(\int_0^t v_s(x) ds\right)$. Moreover if L_g denotes the conformal Laplacian, then by a standard formula

$$(2.4) \quad u_t^5 R_{g_t} = -8 L_g u_t := -8 \left(\Delta_g u_t - \frac{1}{8} R_g u_t \right),$$

so that

$$(2.5) \quad \begin{aligned} \frac{d}{dt} (u_t^5 R_{g_t}) &= -8 L_g \frac{d}{dt} u_t \\ &= -8 L_g (v_t u_t) \\ &= -8 u_t^5 L_{g_t} v_t \\ &= -8 u_t^5 \left(\Delta_{g_t} v_t - \frac{1}{8} R_{g_t} v_t \right) \\ &= (u_t^5 R_{g_t}) v_t. \end{aligned}$$

We then have

$$(2.6) \quad u_t^5 R_{g_t} = \exp\left(\int_0^t v_s ds\right) R_g = u_t R_g.$$

Let $\widetilde{M}_t = M_t^- \cup M_t^+$ denote the doubled manifold, reflected across the minimal surface ∂M_t . Here M_t^- and M_t^+ each represent a copy of M_t , and we endow each of these copies with the metric $g_t^\pm = (w_t^\pm)^4 g_t$ where $w_t^\pm = (1 \pm v_t)/2$. The metric on $\widetilde{M}_t$ will be denoted by $\widetilde{g}_t = g_t^- \cup g_t^+$. Let $c : Cl(T\widetilde{M}_t) \rightarrow \text{End}(\mathfrak{S})$ be the usual representation of the Clifford algebra on the bundle of spinors $\mathfrak{S}$, so that

$$(2.7) \quad c(X)c(Y) + c(Y)c(X) = -2g(X, Y).$$

Choose orthonormal frame fields e_i , $i = 1, 2, 3$ for g , and observe that $e_i^\pm = (w_t^\pm u_t)^{-2} e_i$, $i = 1, 2, 3$ are the corresponding orthonormal frames for $g_t^\pm$. There exists a positive definite inner product on $\mathfrak{S}$, denoted by $\langle \cdot, \cdot \rangle$, with respect to which $c(e_i)$ is anti-hermitian

$$(2.8) \quad \langle c(e_i) \psi, \eta \rangle = -\langle \psi, c(e_i) \eta \rangle.$$

Define the respective spin connections by

$$(2.9) \quad \nabla_{e_i} = e_i + \frac{1}{4} \sum_{j,l=1}^3 \Gamma_{ij}^l c(e_j) c(e_l), \quad \nabla_{e_i^\pm} = e_i^\pm + \frac{1}{4} \sum_{j,l=1}^3 \Gamma_{ij}^{\pm l} c(e_j^\pm) c(e_l^\pm),$$

where Γ_{ij}^l and $\Gamma_{ij}^{\pm l}$ are Levi-Civita connection coefficients for g and $g_t^\pm$. The corresponding Dirac operators are given by

$$(2.10) \quad \mathcal{D} = \sum_{i=1}^3 c(e_i) \nabla_{e_i}, \quad \mathcal{D}^\pm = \sum_{i=1}^3 c(e_i^\pm) \nabla_{e_i^\pm}.$$

Let $\tilde{\psi}_t$ be a harmonic spinor on $\widetilde{M}_t$ which converges to a constant spinor of unit norm at spatial infinity; for the present discussion it is assumed that such a spinor exists. We will denote the restriction of $\tilde{\psi}_t$ to $M_t^\pm$ by $\psi_t^\pm$, then $\psi_t^\pm$ is a harmonic spinor on $(M_t^\pm, g_t^\pm)$. We may now apply the Lichnerowicz formula [1] on each part of the doubled manifold to find that

$$(2.11) \quad 4\pi \tilde{E}_{ADM}(t) - \int_{\partial M_t^-} \langle \psi_t^-, c(e_3^-) \sum_{i=1}^2 c(e_i^-) \nabla_{e_i^-} \psi_t^- \rangle = \int_{M_t^-} \left(|\nabla^- \psi_t^-|^2 + \frac{1}{4} R_{g_t^-} |\psi_t^-|^2 \right) d\omega_{g_t^-},$$

and

$$(2.12) \quad - \int_{\partial M_t^+} \langle \psi_t^+, c(e_3^+) \sum_{i=1}^2 c(e_i^+) \nabla_{e_i^+} \psi_t^+ \rangle = \int_{M_t^+} \left(|\nabla^+ \psi_t^+|^2 + \frac{1}{4} R_{g_t^+} |\psi_t^+|^2 \right) d\omega_{g_t^+},$$

where e_i , $i = 1, 2$ are tangent and e_3 is normal (pointing towards spatial infinity) to the boundary ∂M_t , and $\tilde{E}_{ADM}(t)$ is the ADM energy of $(\widetilde{M}_t, \tilde{g}_t)$. A standard calculation shows that

$$(2.13) \quad c(e_3^\pm) \sum_{i=1}^2 c(e_i^\pm) \nabla_{e_i^\pm} \psi_t^\pm = \mathcal{D}_{\partial M_t^\pm} \psi_t^\pm - \frac{1}{2} H_t^\pm \psi_t^\pm,$$

where $H_t^\pm$ are the mean curvatures of $\partial M_t^\pm$ with respect to the metrics $g_t^\pm$, and the boundary Dirac operator is given by

$$(2.14) \quad \mathcal{D}_{\partial M_t^\pm} = c(e_3^\pm) \sum_{i=1}^2 c(e_i^\pm) \tilde{\nabla}_{e_i^\pm}^\pm$$

with

$$(2.15) \quad \tilde{\nabla}_{e_i^\pm}^\pm = e_i^\pm + \frac{1}{4} \sum_{j,l=1}^2 \Gamma_{ij}^{\pm l} c(e_j^\pm) c(e_l^\pm), \quad i = 1, 2.$$

We claim that the sum of the boundary terms in (2.11) and (2.12) all cancel. To see this note that $\psi_t^-|_{\partial M_t^-} = \psi_t^+|_{\partial M_t^+}$, and since $v_t|_{\partial M_t} = 0$ we have $g_t^-|_{\partial M_t^-} = g_t^+|_{\partial M_t^+}$. Moreover, as ∂M_t^- and ∂M_t^+ represent the same surface in $\widetilde{M}_t$ we have that $e_3^- = -e_3^+$, and thus

$$(2.16) \quad \mathcal{D}_{\partial M_t^-} = -\mathcal{D}_{\partial M_t^+}.$$

Now consider the mean curvature terms. According to a standard formula for the change of mean curvature under conformal deformation

$$(2.17) \quad H_t^\pm = (w_t^\pm)^{-2} [H_t + 4u_t^{-2} e_3(\log w_t^\pm)] = \pm 4u_t^{-2} e_3(v_t),$$

where $H_t = 0$ is the mean curvature of ∂M_t with respect to g_t . It follows that

$$(2.18) \quad H_t^- = -H_t^+.$$

We may now add (2.11) and (2.12), and apply (2.16) and (2.18), to obtain

$$(2.19) \quad \begin{aligned} 4\pi \tilde{E}_{ADM}(t) &= \int_{M_t^-} \left(|\nabla^- \psi_t^-|^2 + \frac{1}{4} R_{g_t^-} |\psi_t^-|^2 \right) d\omega_{g_t^-} \\ &\quad + \int_{M_t^+} \left(|\nabla^+ \psi_t^+|^2 + \frac{1}{4} R_{g_t^+} |\psi_t^+|^2 \right) d\omega_{g_t^+}. \end{aligned}$$

Moreover, since v_t is harmonic we find

$$(2.20) \quad R_{g_t^\pm} = -8(w_t^\pm)^{-5} \left(\Delta_{g_t} \left(\frac{1 \pm v_t}{2} \right) - \frac{1}{8} R_{g_t} \left(\frac{1 \pm v_t}{2} \right) \right) = (w_t^\pm u_t)^{-4} R_g.$$

It follows that

$$(2.21) \quad \begin{aligned} 4\pi \tilde{E}_{ADM}(t) &= \int_{M_t} [(w_t^- u_t)^6 |\nabla^- \psi_t^-|^2 + (w_t^+ u_t)^6 |\nabla^+ \psi_t^+|^2] d\omega_g \\ &\quad + \int_{M_t} \frac{1}{4} ((w_t^-)^2 |\psi_t^-|^2 + (w_t^+)^2 |\psi_t^+|^2) u_t^2 R_g d\omega_g, \end{aligned}$$

It should be noted that a formula similar to (2.21) may be obtained without the use of spinors. To see this, solve the zero scalar curvature equation

$$(2.22) \quad \Delta_{\tilde{g}_t} z_t - \frac{1}{8} \tilde{R}_t z_t = 0 \quad \text{on} \quad \tilde{M}_t, \quad z_t \rightarrow 1 \quad \text{as} \quad |x| \rightarrow \infty,$$

where $\tilde{R}_t$ is the scalar curvature of $\tilde{g}_t$; for the present discussion it is assumed that such a solution exists. Then the conformally related metric $z_t^4 \tilde{g}_t$ has zero scalar curvature. It follows that the ADM energy is nonnegative $E_{ADM}(z_t^4 \tilde{g}_t) \geq 0$. Moreover, the solution has an asymptotic expansion of the form

$$(2.23) \quad z_t = 1 + \frac{C_t}{|x|} + O\left(\frac{1}{|x|^2}\right) \quad \text{as} \quad |x| \rightarrow \infty.$$

A small calculation then shows that

$$(2.24) \quad 0 \leq E_{ADM}(z_t^4 \tilde{g}_t) = \tilde{E}_{ADM}(t) + 2C_t.$$

Hence, integrating by parts produces

$$(2.25) \quad \begin{aligned} 2\pi \tilde{E}_{ADM}(t) &\geq -4\pi C_t \\ &= \int_{S_\infty} z_t \partial_\nu z_t \\ &= \int_{\tilde{M}_t} \left(|\nabla_{\tilde{g}_t} z_t|^2 + \frac{1}{8} \tilde{R}_t z_t^2 \right) d\omega_{\tilde{g}_t} \\ &= \int_{M_t} \left((w_t^- u_t)^6 |\nabla_{\tilde{g}_t} z_t^-|^2 + (w_t^+ u_t)^6 |\nabla_{\tilde{g}_t} z_t^+|^2 \right) d\omega_g \\ &\quad + \int_{M_t} \frac{1}{8} [(w_t^- z_t^-)^2 + (w_t^+ z_t^+)^2] u_t^2 R_g d\omega_g, \end{aligned}$$

where $z_t^\pm$ denotes z_t restricted to $M_t^\pm$. This formula is analogous to (2.21), and was obtained without the use of spinors. Note that part of the integrand on the right-hand side of (2.25) satisfies an elliptic equation with respect to the metric g , even though the solution depends on t :

$$(2.26) \quad 0 = (w_t^\pm u_t)^5 L_{\tilde{g}_t^\pm} z_t^\pm = L_g(w_t^\pm z_t^\pm u_t)$$

where $\tilde{g}_t^\pm = (z_t^\pm)^4 g_t^\pm$.

We mention that an expression similar to (2.21) and (2.25) may also be obtained using the inverse mean curvature flow starting from a point in $\widetilde{M}_t$.

Equations (2.21) and (2.25) imply that $\tilde{E}_{ADM}(t) \geq 0$ under the hypothesis of nonnegative scalar curvature $R_g \geq 0$. Actually, this fact may be considered the crux of the argument for Bray's proof of the Riemannian Penrose inequality. Although the explicit expression above for the energy $\tilde{E}_{ADM}(t)$ is not necessary for Bray's proof, it will however be needed in the next section where we outline an argument for the general Penrose inequality. The purpose of (2.21) and (2.25) is to show that the ADM energy of (M_t, g_t) , which will be denoted by m_t , is decreasing when $R_g \geq 0$. For the Riemannian Penrose inequality m_t is decreasing instantaneously, that is the derivative $\frac{dm_t}{dt} \leq 0$ for all t . However for the general Penrose inequality, where one does not necessarily have nonnegative scalar curvature, m_t will only be shown to be decreasing on the interval $[0, \infty)$, that is $m_0 \geq \lim_{t \rightarrow \infty} m_t$. In order to obtain such a result, let us first conclude this review of the proof of the Riemannian Penrose inequality.

As in [2], we may assume without loss of generality, that the initial data metric g is harmonically flat at infinity. This means that outside a large compact set, g is scalar flat and conformal to the Euclidean metric, with a conformal factor that approaches a positive constant at infinity. In this outer region we may then write

$$(2.27) \quad g = \left(1 + \frac{m}{r} + O\left(\frac{1}{r^2}\right)\right)^4 \delta.$$

Recall that according to a standard formula, if $w = 1 + \frac{c}{r} + \dots$ at spatial infinity, then $E_{ADM}(w^4 g) = E_{ADM}(g) + 2c$. Therefore the parameter m in (2.27) represents half the mass of g , or rather $m_0 = 2m$. Since v_t is a harmonic function, both u_t and v_t have similar asymptotic expansions at spatial infinity

$$(2.28) \quad u_t = \alpha_t + \frac{\beta_t}{r} + O\left(\frac{1}{r^2}\right), \quad v_t = -1 + \frac{\gamma_t}{r} + O\left(\frac{1}{r^2}\right),$$

where $\alpha_0 = 1$ and $\beta_0 = 0$. It follows that

$$(2.29) \quad g_t = u_t^4 g = \left(\alpha_t + \frac{\beta_t + m\alpha_t}{r} + O\left(\frac{1}{r^2}\right)\right)^4 \delta,$$

and hence

$$(2.30) \quad m_t = 2\alpha_t(\beta_t + m\alpha_t).$$

In order to calculate the derivative $m'_t = \frac{dm_t}{dt}$, observe that (2.1) and (2.28) imply

$$(2.31) \quad \alpha'_t = -\alpha_t, \quad \beta'_t = \alpha_t \gamma_t - \beta_t \quad \Rightarrow \quad \alpha_t = e^{-t}, \quad \beta_t = e^{-t} \int_0^t \gamma_s ds.$$

From this we obtain

$$(2.32) \quad m'_t = 2\alpha'_t(\beta_t + m\alpha_t) + 2\alpha_t(\beta'_t + m\alpha'_t) = 2e^{-2t}\gamma_t - 4e^{-t}(\beta_t + e^{-t}m) = -2(m_t - e^{-2t}\gamma_t).$$

However if we set $\tilde{g}_t = \left(\frac{1-v_t}{2}\right)^4 g_t$ then

$$(2.33) \quad \tilde{g}_t = \left(\alpha_t + \frac{\beta_t + m\alpha_t - \alpha_t \gamma_t / 2}{r} + O\left(\frac{1}{r^2}\right)\right)^4 \delta,$$

and thus the energy of $\tilde{g}_t$ is given by

$$(2.34) \quad \tilde{m}_t = 2\alpha_t \left(\beta_t + m\alpha_t - \frac{\alpha_t \gamma_t}{2}\right) = m_t - e^{-2t}\gamma_t,$$

which yields

$$(2.35) \quad m'_t = -2\tilde{m}_t.$$

Therefore upon integrating and rewriting in terms of the previous notation, we find that

$$(2.36) \quad E_{ADM}(0) - E_{ADM}(\infty) = 2 \int_0^\infty \tilde{E}_{ADM}(t) dt$$

where $E_{ADM}(t)$ denotes the total energy of (M_t, g_t) .

Together, (2.21) (or (2.25)) and (2.36) show that the function $E_{ADM}(t)$ is decreasing over all time. There are two other crucial properties of the conformal flow which yield the Penrose inequality. First, due to the boundary condition (2.3) and the fact that ∂M_t is a minimal surface, it follows that the area function $A(t)$ remains constant in time, where $A(t)$ denotes the area of ∂M_t . Second, it is shown in [2] that after rescaling (M_t, g_t) converges to the exterior region of a constant time slice of the Schwarzschild spacetime, so that $E_{ADM}(\infty) = \sqrt{\frac{A(\infty)}{16\pi}}$. By combining all these results together we obtain the following refined version of the Penrose inequality

$$(2.37) \quad E_{ADM} - \sqrt{\frac{|\partial M|}{16\pi}} = 2 \int_0^\infty \tilde{E}_{ADM}(t) dt.$$

In the case of equality, when $E_{ADM} = \sqrt{\frac{|\partial M|}{16\pi}}$, standard arguments combined with (2.21) (or (2.25)) and (2.36) show that (M, g) is conformally equivalent to $(\mathbb{R}^3 - B_1(0), \delta)$ with zero scalar curvature. Since the boundary ∂M is a minimal surface, it follows that (M, g) is isometric to a constant time slice of the Schwarzschild spacetime.

3. COUPLING THE CONFORMAL FLOW TO THE GENERALIZED JANG EQUATION

Let Σ_0 be a solution of the generalized Jang equation (1.8) satisfying boundary condition (1.14), as described in Section 1. Then the boundary $\partial\Sigma_0$ is a minimal surface, and is in fact the outermost minimal surface. To see this, suppose that a surface $\bar{S}$ encloses but is not equal to $\partial\Sigma_0$ and satisfies $|\bar{S}| \leq |\partial\Sigma_0|$. Since $\bar{g} \geq g$, we have $|S| \leq |\bar{S}|$, where $S \subset M$ is the projection of $\bar{S}$. Thus $|S| \leq |\partial\Sigma_0| = |\partial M_0|$, which is impossible since ∂M_0 is the outermost minimal area enclosure of ∂M .

We will now perform the conformal flow of metrics on the Jang surface Σ_0 , and show how this leads naturally to a choice for the warping factor ϕ . Following the notation from the previous section let $(g_t = u_t^4 \bar{g}, \Sigma_t)$ be a conformal flow, where Σ_t denotes the region outside of the outermost minimal surface $\partial\Sigma_t$. Then from (2.6) the scalar curvature evolves by

$$(3.1) \quad R_{g_t} = u_t^{-4} \bar{R}.$$

Moreover, since the ADM energies of (M, g, k) and $(\Sigma, \bar{g})$ agree, we may apply (2.21) or (2.25) and (2.36) to obtain

$$(3.2) \quad E_{ADM} - \sqrt{\frac{|\partial\Sigma_0|}{16\pi}} \geq 2 \int_0^\infty \tilde{E}_{ADM}(t) dt,$$

where $\tilde{E}_{ADM}(t)$ denotes the total energy of $(\tilde{\Sigma}_t, g_t^- \cup g_t^+)$ and is given by

$$(3.3) \quad \begin{aligned} 4\pi \tilde{E}_{ADM}(t) = & \int_{\Sigma_t} [(w_t^- u_t)^6 |\nabla^- \psi_t^-|^2 + (w_t^+ u_t)^6 |\nabla^+ \psi_t^+|^2] d\omega_{\bar{g}} \\ & + \int_{\Sigma_t} \frac{1}{4} ((w_t^-)^2 |\psi_t^-|^2 + (w_t^+)^2 |\psi_t^+|^2) u_t^2 \bar{R} d\omega_{\bar{g}}, \end{aligned}$$

or

$$(3.4) \quad \begin{aligned} 4\pi \tilde{E}_{ADM}(t) &= \int_{\Sigma_t} 2 \left((w_t^- u_t)^6 |\nabla_{\tilde{g}_t} z_t^-|^2 + (w_t^+ u_t)^6 |\nabla_{\tilde{g}_t} z_t^+|^2 \right) d\omega_{\tilde{g}} \\ &\quad + \int_{\Sigma_t} \frac{1}{4} \left[(w_t^- z_t^-)^2 + (w_t^+ z_t^+)^2 \right] u_t^2 \bar{R} d\omega_{\tilde{g}}. \end{aligned}$$

Notice that we cannot conclude as in Section 2 that $\tilde{E}_{ADM}(t) \geq 0$, since the scalar curvature $\bar{R}$ of the Jang surface is not necessarily nonnegative. According to (1.6), it is a divergence term which is the possible obstruction to nonnegativity. However by choosing ϕ appropriately, this divergence term may be integrated away so that the right-hand side of (2.37) is nonnegative. To accomplish this let

$$(3.5) \quad \chi_t(x) = \begin{cases} 1 & \text{if } x \in \Sigma_t, \\ 0 & \text{if } x \in \Sigma_0 - \Sigma_t, \end{cases}$$

and set

$$(3.6) \quad \phi = 2 \int_0^\infty \chi_t \left((w_t^-)^2 |\psi_t^-|^2 + (w_t^+)^2 |\psi_t^+|^2 \right) u_t^2 dt,$$

or

$$(3.7) \quad \phi = 2 \int_0^\infty \chi_t \left[(w_t^- z_t^-)^2 + (w_t^+ z_t^+)^2 \right] u_t^2 dt.$$

Notice that ϕ should be strictly positive away from $\partial\Sigma_0$,

$$(3.8) \quad \phi(x) \rightarrow 1 \quad \text{as } |x| \rightarrow \infty, \quad \text{and} \quad \phi|_{\partial\Sigma_0} = 0,$$

where we have used (2.3) and the fact that $u_t \rightarrow e^{-t}$ as $|x| \rightarrow \infty$. By applying Fubini's Theorem, using the boundary condition (3.8), as well as the scalar curvature formula for the Jang surface (1.6), we may calculate

$$(3.9) \quad \begin{aligned} \int_0^\infty \int_{\Sigma_t} \left((w_t^-)^2 |\psi_t^-|^2 + (w_t^+)^2 |\psi_t^+|^2 \right) u_t^2 \bar{R} d\omega_{\tilde{g}} dt &= \frac{1}{2} \int_{\Sigma_0} \phi \bar{R} d\omega_{\tilde{g}} \\ &\geq \frac{1}{2} \int_{\Sigma_0} \phi \left(2|q|^2 - \frac{2}{\phi} \overline{\text{div}}(\phi q) \right) d\omega_{\tilde{g}} \\ &= \int_{\Sigma_0} \phi |q|^2 d\omega_{\tilde{g}} + \int_{\partial\Sigma_0} \phi q(\bar{N}) d\sigma_{\tilde{g}} \\ &= \int_{\Sigma_0} \phi |q|^2 d\omega_{\tilde{g}}, \end{aligned}$$

where $\bar{N}$ is the unit normal to $\partial\Sigma_0$ pointing towards spatial infinity. We could also have used (3.7). The above calculation also uses the reasonable assumption that $|q|$ remains uniformly bounded and decays very fast at spatial infinity; this behavior is consistent with that of solutions to the classical Jang equation [20]. This may be combined with (1.6), (3.2), (3.3), and (3.9) to yield

$$(3.10) \quad \begin{aligned} &E_{ADM} - \sqrt{\frac{|\partial\Sigma_0|}{16\pi}} \\ &\geq \frac{1}{2\pi} \int_0^\infty \int_{\Sigma_t} \left[(w_t^- u_t)^6 |\nabla^- \psi_t^-|^2 + (w_t^+ u_t)^6 |\nabla^+ \psi_t^+|^2 \right] d\omega_{\tilde{g}} dt \\ &\quad + \frac{1}{8\pi} \int_0^\infty \int_{\Sigma_t} \left((w_t^-)^2 |\psi_t^-|^2 + (w_t^+)^2 |\psi_t^+|^2 \right) u_t^2 \left(16\pi(\mu - J(w)) + |h - K|_\Sigma|^2 + 2|q|^2 \right) d\omega_{\tilde{g}}. \end{aligned}$$

The Penrose inequality (1.4) (with ADM energy replacing ADM mass) now follows. This is due to the fact that the Jang metric $\bar{g}$ measures areas to be at least as large as those measure by g , and hence $|\partial\Sigma_0| \geq A_{min}$. We remark that because of the asymptotics (1.13), only the ADM energy is encoded in the solution of the generalized Jang equation. At this time it is unknown how to encode the ADM linear momentum into a solution of Jang's equation.

Now consider the case of equality in (1.4). We will use (3.6), although a similar argument holds for (3.7). First note that in this case $|\partial\Sigma_0| = A_{min}$. Next, according to (3.10), both spinors $\psi_t^\pm$ are parallel. This implies that $|\psi_t^\pm| \equiv 1$, in light of the boundary condition at spatial infinity. It follows that

$$(3.11) \quad \mu = |J| \equiv 0, \quad h = K|_\Sigma, \quad q \equiv 0,$$

where we have used the dominant energy condition and the fact that $|w| < 1$ away from ∂M . Hence $\bar{R} \equiv 0$. Moreover the existence of a basis of parallel spinors shows that $(\tilde{\Sigma}_t, g_t^- \cup g_t^+)$ is isometric to $(\mathbb{R}^3, \delta)$. The Jang surface Σ_0 is then conformally flat with zero scalar curvature, and since $\partial\Sigma_0$ is minimal, we find that $(\Sigma_0, \bar{g})$ is isometric to the exterior region of the $t = 0$ slice of the Schwarzschild spacetime

$$(3.12) \quad \mathbb{SC}^4 = (\mathbb{R} \times (\mathbb{R}^3 - B_{\frac{m_0}{2}}(0)), -\phi_{SC}^2 dt^2 + g_{SC}),$$

where $r = |x|$ and

$$(3.13) \quad \phi_{SC}(x) = \frac{1 - \frac{m_0}{2r}}{1 + \frac{m_0}{2r}}, \quad g_{SC} = \left(1 + \frac{m_0}{2r}\right)^4 \delta.$$

We will show that $\phi = \phi_{SC}$. Since $\bar{g}$ is isometric to the spatial Schwarzschild metric, the conformal flow of metrics may be given explicitly, namely

$$(3.14) \quad v_t(x) = \begin{cases} \frac{-e^{-t} + \frac{m_0}{2r} e^t}{e^{-t} + \frac{m_0}{2r} e^t} & \text{if } r \geq \frac{m_0}{2} e^{2t} \\ 0 & \text{if } r < \frac{m_0}{2} e^{2t} \end{cases}, \quad u_t(x) = \begin{cases} \frac{e^{-t} + \frac{m_0}{2r} e^t}{1 + \frac{m_0}{2r}} & \text{if } r \geq \frac{m_0}{2} e^{2t} \\ \frac{\sqrt{\frac{2m_0}{r}}}{1 + \frac{m_0}{2r}} & \text{if } r < \frac{m_0}{2} e^{2t} \end{cases}.$$

It follows that on Σ_t

$$(3.15) \quad w_t^- = \frac{1 - v_t}{2} = \frac{e^{-t}}{e^{-t} + \frac{m_0}{2r} e^t}, \quad w_t^+ = \frac{1 + v_t}{2} = \frac{\frac{m_0}{2r} e^t}{e^{-t} + \frac{m_0}{2r} e^t},$$

and therefore

$$(3.16) \quad \begin{aligned} \phi(x) &= 2 \int_0^{\log \sqrt{\frac{2r}{m_0}}} \left[\left(\frac{e^{-t}}{e^{-t} + \frac{m_0}{2r} e^t} \right)^2 + \left(\frac{\frac{m_0}{2r} e^t}{e^{-t} + \frac{m_0}{2r} e^t} \right)^2 \right] \left(\frac{e^{-t} + \frac{m_0}{2r} e^t}{1 + \frac{m_0}{2r}} \right)^2 dt \\ &= 2 \int_0^{\log \sqrt{\frac{2r}{m_0}}} \frac{e^{-2t} + \left(\frac{m_0}{2r}\right)^2 e^{2t}}{\left(1 + \frac{m_0}{2r}\right)^2} dt \\ &= \left(\frac{1 - \frac{m_0}{2r}}{1 + \frac{m_0}{2r}} \right) = \phi_{SC}. \end{aligned}$$

Consider now the graph map $G : M \rightarrow \mathbb{SC}^4$ given by $x \mapsto (x, f(x))$. Since $\phi \equiv \phi_{SC}$, the map G yields an isometric embedding of the initial data (M_0, g, k) into the Schwarzschild spacetime, as

$$(3.17) \quad g = \bar{g} - \phi^2 df^2 = g_{SC} - \phi_{SC}^2 df^2.$$

Moreover since $h = K|_\Sigma$, a calculation [3], [4] guarantees that the second fundamental form of this embedding agrees with k . Lastly observe that since ϕ vanishes at ∂M_0 , we must have that ∂M_0 is

an apparent horizon. However ∂M is an outermost apparent horizon, so in fact $\partial M_0 = \partial M$, and hence $M_0 = M$.

Remark 3.1. Without further modification, the arguments of this paper can only yield the inequality

$$(3.18) \quad E_{ADM} \geq \sqrt{\frac{A_{min}}{16\pi}},$$

where the ADM mass is replaced by the ADM energy in (1.4). The rigidity statement for (3.18) is as follows, equality holds only if (M, g, k) arises from a spacelike slice of the Schwarzschild spacetime, with outermimizing boundary. Note that the ‘if’ direction is not included.

A proof of the ‘if’ part in the rigidity statement for the inequality (1.4) is as follows. Suppose that (M, g, k) is a spacelike slice of the Schwarzschild spacetime with outermimizing boundary. Since $\phi_{SC}|_{\partial M} = 0$, we have that $|\partial M| = |\partial \widehat{M}|$, where $(\widehat{M}, g_{SC}, 0)$ is the $t = 0$ slice of the Schwarzschild spacetime. Furthermore

$$(3.19) \quad M_{ADM} = M_{ADM}(\widehat{M}) = E_{ADM}(\widehat{M}),$$

so that

$$(3.20) \quad M_{ADM} = E_{ADM}(\widehat{M}) = \sqrt{\frac{|\partial \widehat{M}|}{16\pi}} = \sqrt{\frac{|\partial M|}{16\pi}} \geq \sqrt{\frac{A_{min}}{16\pi}}.$$

It follows that equality holds in (1.4) since ∂M is outermimizing, that is $|\partial M| = A_{min}$. These same arguments also show why the boundary must be outermimizing in the case of equality. Namely, in light of (3.20), the only way that equality can hold in (1.4) is if $|\partial M| = A_{min}$.

REFERENCES

- [1] R. Bartnik, P. Chrusciel, *Boundary value problems for Dirac-type equations*, J. Reine Angew. Math., **579** (2005), 13-73. arXiv:math/0307278
- [2] H. Bray, *Proof of the Riemannian Penrose inequality using the positive mass theorem*, J. Differential Geom., **59** (2001), no. 2, 177-267.
- [3] H. Bray, M. Khuri, *A Jang equation approach to the Penrose Inequality*, Discrete Contin. Dyn. Syst., **27** (2010), 741-766. arXiv:0910.4785
- [4] H. Bray, M. Khuri, *P.D.E's which imply the Penrose conjecture*, Asian J. Math., **15** (2011), no. 4, 557-610. arXiv:0905.2622
- [5] Y.-S. Cha, M. Khuri, *Deformations of axially symmetric initial data and the mass-angular momentum inequality*, Ann. Henri Poincaré, **16** (2015), no. 3, 841-896. arXiv:1401.3384
- [6] Y.-S. Cha, M. Khuri, *Deformations of charged axially symmetric initial data and the mass-angular momentum-charge inequality*, Ann. Henri Poincaré, **16** (2015), no. 12, 2881-2918. arXiv:1407.3621
- [7] Y.-S. Cha, M. Khuri, and Anna Sakovich, *Reduction arguments for geometric inequalities associated with asymptotically hyperboloidal slices*, Class. Quantum Grav., **33** (2016), 035009. arXiv:1509.06255
- [8] M. Disconzi, and M. Khuri, *On the Penrose inequality for charged black holes*, Class. Quantum Grav., **29** (2012), 245019, arXiv:1207.5484.
- [9] G. Galloway, *Rigidity of marginally trapped surfaces and the topology of black holes*, Commun. Anal. Geom., **16** (2008), 217-229. arXiv:gr-qc/0608118
- [10] G. Galloway, R. Schoen, *A generalization of Hawking's black hole topology theorem to higher dimensions*, Commun. Math. Phys., **266** (2006), no. 2, 571-576. arXiv:gr-qc/0509107
- [11] Q. Han, M. Khuri, *Existence and blow-up behavior for solutions of the generalized Jang equation*, Comm. Partial Differential Equations, **38** (2013), 2199-2237. arXiv:1206.0079.
- [12] G. Huisken, T. Ilmanen, *The inverse mean curvature flow and the Riemannian Penrose inequality*, J. Differential Geom., **59** (2001), 353-437.
- [13] M. Khuri, *A Penrose-like inequality for general initial data sets*, Commun. Math. Phys., **290** (2009), 779-788; *Erratum*, **317** (2013), 267. arXiv:0910.5039

- [14] M. Khuri, *A Penrose-Like inequality with charge*, Gen. Relativity Gravitation, **45** (2013), 2341-2361. arXiv:1308.3591
- [15] M. Khuri, and G. Weinstein, *Rigidity in the positive mass theorem with charge*, J. Math. Phys., **54** (2013), 092501. arXiv:1307.5499
- [16] M. Khuri, G. Weinstein, and S. Yamada, *Extensions of the charged Riemannian Penrose inequality*, Class. Quantum Grav., **32** (2015), 035019. arXiv:1410.5027
- [17] M. Khuri, G. Weinstein, and S. Yamada, *The Riemannian Penrose inequality with charge for multiple black holes*, Proceedings of the Complex Analysis & Dynamical Systems VI Conference (Nahariya, Israel, May 2013), Contemporary Mathematics, **653** (2015), 219-226. arXiv:1308.3771
- [18] M. Khuri, G. Weinstein, and S. Yamada, *Proof of the Riemannian Penrose inequality with charge for multiple black holes*, J. Differential Geom., **106** (2017), 451-498. arXiv:1409.3271
- [19] M. Mars, *Present status of the Penrose inequality*, Classical Quantum Gravity, **26** (2009), no. 19, 193001. arXiv:0906.5566
- [20] R. Schoen, S.-T. Yau, *Proof of the positive mass theorem II*, Commun. Math. Phys., **79** (1981), no. 2, 231-260.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
E-mail address: qhan@nd.edu

BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, BEIJING, 100871, CHINA
E-mail address: qhan@math.pku.edu.cn

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794
E-mail address: khuri@math.sunysb.edu