

Research



Cite this article: Pang Y, Sezgin E. 2020 On the consistency of a class of R -symmetry gauged $6D, \mathcal{N} = (1,0)$ supergravities. *Proc. R. Soc. A* **476**: 20200115.
<http://dx.doi.org/10.1098/rspa.2020.0115>

Received: 21 February 2020

Accepted: 3 July 2020

Subject Areas:

high energy physics, string theory

Keywords:

supergravity, anomalies, string theory

Author for correspondence:

Ergin Sezgin

e-mail: sezgin@tamu.edu

This paper is dedicated to Michael Duff on the occasion of his 70th birthday.

[†]Present address: Center for Joint Quantum Studies and Department of Physics, School of Science, Tianjin University, Tianjin 300350, People's Republic of China.

On the consistency of a class of R -symmetry gauged $6D, \mathcal{N} = (1,0)$ supergravities

Yi Pang^{1,†} and Ergin Sezgin²

¹Mathematical Institute, University of Oxford, Woodstock Road, Oxford OX2 6GG, UK

²George and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA

ES, 0000-0002-1415-8023

R -symmetry gauged $6D, (1,0)$ supergravities free from all local anomalies, with gauge groups $G \times G_R$ where G_R is the R -symmetry group and G is semisimple with rank greater than one, and which have no hypermultiplet singlets, are extremely rare. There are three such models known in which the gauge symmetry group is $G_1 \times G_2 \times U(1)_R$, where the first two factors are $(E_6/\mathbb{Z}_3) \times E_7$, $G_2 \times E_7$ and $F_4 \times Sp(9)$. These are models with single tensor multiplet, and hyperfermions in the $(1,912)$, $(14,56)$ and $(52,18)$ dimensional representations of $G_1 \times G_2$, respectively. So far, it is not known if these models follow from string theory. We highlight key properties of these theories, and examine constraints which arise from the consistency of the quantization of anomaly coefficients formulated in their strongest form by Monnier and Moore. Assuming that the gauged models accommodate dyonic string excitations, we find that these constraints are satisfied only by the model with the $F_4 \times Sp(9) \times U(1)_R$ symmetry. We also discuss aspects of dyonic strings and potential caveats they may pose in applying the stated consistency conditions to the R -symmetry gauged models.

1. Introduction

The work described herein is on a subject in which Michael has made magnificent contributions. Let us also remember that his advocacy of supermembranes and eleven dimensions (11D) prior to their wide acceptance, is in the annals of physics. He has related amusing anecdotes about the era prior to this

acceptance. One of us (ES) has this one to add: In an Aspen Conference in 1987, in a conversation on supermembranes and 11D, a very distinguished colleague said that ‘in Cambridge meeting there were 5 h of talk on the subject, this must be a Thatcherian plot to destroy the British physics!’ Asked about 11D supergravity, he replied, ‘it is a curiosity!’

In the spirit of exploring some other ‘curiosities’, here we aim at drawing attention to a class of supergravities in six dimensions (6D) that are free from all local anomalies in a rather remarkable fashion. To begin with, let us recall that the requirement of anomaly freedom has considerably constraining consequences for supergravity theories. The gauge groups and matter content are restricted, and on-shell supersymmetry, in the presence of the Green–Schwarz anomaly counterterm, *requires* the introduction of an infinite tower of higher derivative couplings. While this is not expected to fix uniquely an effective theory of quantum supergravity [1–4], it may nonetheless provide a framework for a ‘ α' -deformation’ program in which extra consistency requirements, including those arising from global anomalies, may restrict further the effective theory. If consistent such theories exist, in principle, they may offer viable spots in a region outside the string lamp post in a search for UV completeness. In fact, even in 10D it would be instructive to determine if and how the α' deformation approach runs into problems unless it is uniquely determined by string theory. If all roads lead to string theory that too would be a valuable outcome in this program, providing more evidence for what is referred to as the ‘string lamp post principle’ [5,6].

In this note, as mentioned above, we draw attention to a class of 6D supergravities which are remarkably anomaly free, and yet it is not known if they can be embedded into string theory. These are $U(1)_R$ gauged supergravities with specific gauge groups, and from the string theory perspective unusual hypermatter content. The qualification ‘remarkable’ is due to the fact that R -symmetry gauged 6D (1,0) supergravities free from all local anomalies, and with gauge groups $G \times G_R$, where G_R is the R -symmetry group and G is semisimple with rank greater than one, and without any hypermultiplet singlets, are extremely rare; see for example [7]. By contrast, there is a huge number of anomaly free ungauged 6D (1,0) supergravities¹ one can construct directly, and a very large class that can be embedded, indeed directly be obtained from, string theory; see for example [8]. One can also find several R -symmetry gauged models in which there are several hypermultiplet singlets, and gauge group G that involves a number of $U(1)$ factors [7,9].

If we insist on semisimple gauge groups and exclude hyperfermion singlets, then there are only three anomaly free gauged 6D (1,0) supergravities known so far. They have the gauge symmetry $G_1 \times G_2 \times U(1)_R$ where the last factor is the gauged R -symmetry group, and the first two factors are $(E_6/\mathbb{Z}_3) \times E_7$ [10], $G_2 \times E_7$ [11] and $F_4 \times Sp(9)$ [7]. These are models with a single tensor multiplet, and hyperfermions in the (1,912), (14,56) and (52,18) dimensional representations of $G_1 \times G_2$, respectively. The embedding of these theories in string theory is not known.² In particular, the $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$ model contains a representation of E_7 beyond the fundamental and adjoint what normally one encounters in string theory constructions. Neither R -symmetry gauging, nor such representations seem to arise in string/F theory constructions.³ Another landmark of these models is that they come with a positive definite potential proportional to the exponential function of the dilaton, even in the absence of the hypermultiplets. This has significant consequences. For example, these models do not admit maximally symmetric 6D space–time vacua, but rather Minkowski₄ $\times S^2$ with a monopole configuration on S^2 [12].

¹There is a crucial difference between gauging of R -symmetry compared to gauging of non- R -symmetries. Here, we shall reserve the word ‘gauged’ to mean ‘ R -symmetry gauged’.

²If one considers strictly the $U(1)_R$ gauged theory with $n_T = 1$ and no other gauge sector and hypermatter [12], it has been shown [13] to follow from pure Type I supergravity in 10D, on a non-compact hyperboloid $\mathcal{H}_{2,2}$ times S^1 and a consistent chiral truncation. However, the inclusion and origin the Yang–Mills hypermatter remains an open problem.

³It has been suggested in [14] that higher Kac–Moody level string worldsheet algebras may lead to such representations but this matter is far from being settled.

In its simplest form, such gauged supergravities seem to have attractive features for cosmology (see, for example, [15–19]). The presence of the potential also has interesting consequences for the important question of whether dyonic string excitations are supported by the theory. While some encouraging results have been obtained in that direction [20–22], much remains to be investigated.

The GS mechanism ensures the cancellation of local anomalies. Demanding the absence of possible global anomalies, on the other hand, can impose additional constraints. Such anomalies can arise from different aspects of the data furnished by the local anomaly free theory, and they can be rather difficult to compute. The best understood global anomaly has to do with large gauge transformations not connected to the identity. The models in question are free from these anomalies. Other global considerations, motivated in part by lessons learned from the F-theory construction of anomaly free 6D theories [23], have led to additional constraints.

To begin with, Seiberg & Taylor [24], employed the properties of the dyonic charge lattice and the Dirac quantization conditions they must satisfy, to deduce the consequences for the coefficients of the anomaly polynomial. They observed that these coefficients form a sublattice of dyonic string charge lattice, and that the consistency requires that this can be extended to a unimodular (self-dual) lattice. A stronger condition was put forward relatively recently by Monnier *et al.* [25] who assumed that a consistent supergravity theory may be put on an arbitrary spin manifold and that any smooth gauge field configurations should be allowed in the supergravity ‘path integral’, referring to this assumption as the ‘generalized completeness hypothesis’. They find a constraint which states that the anomaly coefficients for the gauge group G should be an element of $2H^4(BG; \mathbb{Z}) \otimes \Lambda_5$ where BG is the classifying space of the gauge group G , and Λ_5 is the unimodular string charge lattice.

Monnier & Moore [26,27] have further argued that these constraints are necessary but not sufficient for the cancellation of all anomalies, local and global, and proposed the necessary and sufficient conditions. They do so by requiring that the Green–Schwarz anomaly counterterm is globally well defined, and show that this leads to the requirement that for any given Green–Schwarz counterterm, there must exist a certain 7D spin topological field theory that is trivial. These authors arrive at a proposition in [26] which states the conditions that need to be satisfied for the 6D theory to be free from all anomalies, in the case of connected gauge groups. Assuming that the theories under consideration here support dyonic string excitations, we will take this proposition as a basis for testing the consistency of these theories. We will see that the model based on $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$ satisfies the weaker set of constraints mentioned above, but not all the conditions of the stronger criterion of Monnier and Moore, and that the model based on $F_4 \times Sp(9) \times U(1)_R$ remarkably satisfies them all.

The paper is organized as follows. In the next section, we shall describe the anomaly freedom aspects of the class of models being considered here. In §3, we recall the structure of the bosonic field equations, discuss the issue of higher derivative corrections, and survey the key properties of the few known dyonic string solutions. In §4, we summarize the Monnier–Park constraints, and in §5, we study these constraints for the models at hand. In the conclusions, we discuss the possible caveats in the interpretation of our results, and the appendix contain useful formula for the anomalies of the models under consideration, and in particular, we provide more details for the ones based on $F_4 \times Sp(9) \times U(1)_R$ gauge group.

2. $U(1)_R$ gauged anomaly free models

We shall focus on $U(1)_R$ gauged $6D, \mathcal{N}=(1,0)$ supergravities coupled to one tensor multiplet, vector multiples associated with group $G = G_1 \times G_2 \times U(1)_R$, and (half)hypermultiplets transforming in $(R_1, R_2)_0$ representation of $G_1 \times G_2$, with the subscript denoting the $U(1)_R$ charge of hyperfermions. The gravitino, dilatino and gauginos have unit $U(1)_R$ charge. The three models

we shall consider have the gauge groups and hyperfermion contents [7,10,11]

$$(A) \quad (E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R \quad (1, 912)_0 \quad (2.1)$$

$$(B) \quad G_2 \times E_7 \times U(1)_R \quad (14, 56)_0 \quad (2.2)$$

and $(C) \quad F_4 \times Sp(9) \times U(1)_R \quad (52, 18)_0. \quad (2.3)$

The perturbative gravitational, gauge and mixed anomalies are encoded in an 8-form anomaly polynomial I_8 , and they are cancelled by Green–Schwarz mechanism that exploits its factorization as

$$I_8 = \frac{1}{2} \Omega_{\alpha\beta} Y^\alpha Y^\beta \quad \text{and} \quad \Omega_{\alpha\beta} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (2.4)$$

where

$$Y^\alpha = \frac{1}{16\pi^2} \left(\frac{1}{2} a^\alpha \text{tr} R^2 + b_r^\alpha \left(\frac{2}{\lambda_r} \text{tr} F_r^2 \right) + 2 c^\alpha F^2 \right) \quad (2.5)$$

$r = 1, 2$ labels the group $G_1 \times G_2$. Here, F_r is the field strength of the r th component of the gauge group, F denotes the $U(1)_R$ field strength, tr is the trace in the fundamental representation, and summation over r is understood,⁴ and λ_r is normalization factor which is fixed by demanding that the smallest topological charge of an embedded $SU(2)$ instanton is 1 [31]. These factors, which are equal to the Dynkin indices of the fundamental representations, are listed below for the groups needed here.

G	E_7	E_6	F_4	G_2	$Sp(9)$	
λ	12	6	6	2	1	(2.6)

The vectors (a, b_i, c) in the space $\mathbb{R}^{1,1}$ should belong to an integral lattice, referred to as the *anomaly lattice*. For the three models, we are considering, these vectors are [7,10,11]

$$(A) \quad a = (2, -2), \quad b_6 = (1, 3), \quad b_7 = (3, -9), \quad c = (2, 18), \quad (2.7)$$

$$(B) \quad a = (2, 2), \quad b_2 = (3, 15), \quad b_7 = (3, 1), \quad c = \left(2, -\frac{38}{3} \right) \quad (2.8)$$

and $(C) \quad a = (2, -2), \quad b_4 = (2, -10), \quad b_9 = \left(1, \frac{1}{2} \right), \quad c = (2, 19). \quad (2.9)$

Note that in models (A) and (C), the anomaly polynomial starts with $-\text{tr}(R^2)^2$, while in the model (B) it starts with $+\text{tr}(R^2)^2$. The anomaly polynomials for models (A) [10,21], (B) [11] and (C) [7] are provided in the appendix.

For groups with a non-trivial sixth homotopy group, there may be global anomalies associated with large gauge transformations not connected to the identity. Among the gauge groups of the three models above, only G_2 has a non-trivial such homotopy group given by $\pi_6(G_2) = \mathbb{Z}_3$. In that case, the vanishing of the global anomaly requires that [32]

$$G_2: \quad 1 - 4 \sum_R n_R d_R = 0 \bmod 3, \quad (2.10)$$

where n_R is the number of (half)hypermultiplets transforming in the representation R of G_2 , and d_R is defined by $\text{tr}_R F^4 = d_R (\text{tr} F^2)^2$. Since $n_{14} = \frac{1}{2} \times 56$ and $d_{14} = \frac{5}{2}$ for model (B), the global anomaly is absent [11].

3. Supersymmetry, bosonic field equations and dyonic strings

In order to highlight the issues that arise in the context of finding dyonic string solutions of the $U(1)_R$ gauged theory, here we shall review the bosonic field equations, with the assumptions that

⁴There exists the identity $(\text{tr} F^2)/\lambda = (\text{Tr} F^2)/(2h^\vee)$, where Tr is the trace in the adjoint representation and $h^\vee$ is the dual Coxeter number. λ is in fact the index of the fundamental representation of group G .

the hyperscalar fields are set to zero in these equations. We start by introducing a metric $G_{\alpha\beta}$, and $SO(1, 1)$ invariant tensor $\Omega_{\alpha\beta}$ in terms of Ω -orthogonal vectors e_α and j_α as follows:

$$\begin{aligned} G_{\alpha\beta} &= e_\alpha e_\beta + j_\alpha j_\beta, & \Omega_{\alpha\beta} &= -e_\alpha e_\beta + j_\alpha j_\beta, \\ e_\alpha &= \frac{1}{\sqrt{2}}(e^{-\varphi}, -e^\varphi), & j_\alpha &= \frac{1}{\sqrt{2}}(e^{-\varphi}, e^\varphi). \end{aligned} \quad (3.1)$$

They satisfy $e \cdot e = -1, j \cdot j = 1$ and $e \cdot j = 0$ where the inner product is with respect to $\Omega^{\alpha\beta} = (\Omega_{\alpha\beta})^{-1}$. It is also convenient to introduce the notation

$$v_L^\alpha := \frac{1}{2}a^\alpha \quad \text{and} \quad v_i^\alpha := \left(\frac{2b_1^\alpha}{\lambda_1}, \frac{2b_2^\alpha}{\lambda_2}, 2c^\alpha \right), \quad (3.2)$$

where $i = 1, 2, 3$ labels the group $G_1 \times G_2 \times U(1)_R$. Then, (2.5) can be written as

$$Y^\alpha = \frac{1}{16\pi^2} \left(v_L^\alpha \text{tr} R^2 + v_i^\alpha \text{tr} F_i^2 \right), \quad (3.3)$$

where $v_3^\alpha \text{tr} F^2 \equiv v^\alpha F^2$. The constant vectors v_i^α can be directly read off from (2.7), (2.8) and (2.9). Since $dY^\alpha = 0$, one can locally define the associated Chern–Simons form through $Y^\alpha = d\Gamma^\alpha$. The 3-form field strength is then defined as⁵

$$H^\alpha = dB^\alpha + \alpha' \Gamma^\alpha, \quad \text{with } d\Gamma^\alpha = 16\pi^2 Y^\alpha, \quad (3.4)$$

where α' is the ‘inverse string tension’.

If we set $v_L^\alpha = 0$, and either $v_i^1 = 0$ or $v_i^2 = 0$, then a classically locally supersymmetric and gauge-invariant action exists for arbitrary v_i^1 or v_i^2 [33]. If we switch on both v_i^α simultaneously, local supersymmetric field equations of motion have also been constructed, but anomalies in gauge transformation and local supersymmetry arise [34,35]. This is to be expected, since Green–Schwarz counterterms required for the cancellation of one-loop anomalies are present, and therefore the classical and one-loop quantum effects are mixed. The Green–Schwarz counterterm also requires that the parameters v_L^1 and v_L^2 are turned on, and fixes v_i^1 , and v_i^2 in terms of a single dimensionful parameter α' . Furthermore, supersymmetry is now broken already at order α' , since R and F have the same dimension, and arise in the field equations already at order α' .

For $v_L^\alpha = 0$, the bosonic field equations with hypermultiplet scalars set to zero, and in Einstein frame, take the form [34,36,37]

$$\star \left(G_{\alpha\beta} H^\beta \right) = \Omega_{\alpha\beta} H^\beta, \quad (3.5)$$

$$\alpha' D \left(\star e \cdot v_i F_i \right) = 2\sqrt{2} \alpha' v_i^\alpha G_{\alpha\beta} \star H^\beta \wedge F_i, \quad (3.6)$$

$$R_{\mu\nu} = \partial_\mu \varphi \partial_\nu \varphi + G_{\alpha\beta} (H^\alpha \cdot H^\beta)_{\mu\nu} + 2\sqrt{2} \alpha' e \cdot v_i \text{tr} \left((F_i^2)_{\mu\nu} - \frac{1}{8} g_{\mu\nu} F_i^2 \right) + \frac{1}{8\sqrt{2}} \frac{1}{\alpha'} (e \cdot v)^{-1} g_{\mu\nu} \quad (3.7)$$

$$\text{and } \nabla_\mu \partial^\mu \varphi = -\frac{1}{\sqrt{2}} \alpha' j \cdot v_i \text{tr} F_i^2 - \frac{2}{3} e_\alpha j_\beta H^\alpha \cdot H^\beta + \frac{1}{4\sqrt{2}} \frac{1}{\alpha'} \frac{j \cdot v}{(e \cdot v)^2}, \quad (3.8)$$

with self explanatory meaning of the notations $H^\alpha \cdot H^\beta, (F^2)_{\mu\nu}$ and F^2 . It follows from (3.5) and (3.4) that

$$d \star \left(G_{\alpha\beta} H^\beta \right) = 16\pi^2 \alpha' \Omega_{\alpha\beta} Y^\beta \quad \text{and} \quad dH^\alpha = 16\pi^2 \alpha' Y^\alpha. \quad (3.9)$$

Thus $\Omega_{\alpha\beta} Y^\beta$ and Y^α are the electric and magnetic sources, respectively. Note also that $\star e \cdot H = -e \cdot H$ belongs to the supergravity multiplet, and $\star j \cdot H = j \cdot H$ is in the single tensor multiplet. We also see from (3.6) that there are terms proportional to α'^2 that break the gauge

⁵We use the notation and conventions of [25] to large extent. One particular exception is that we take $e \cdot H$ to belong to the supergravity multiplet, rather than the tensor multiplet. This is in accordance with the conventions of [34]. We also redefine α' by a factor of $16\pi^2$ for convenience in notation. We thank Hao-Yuan Chang for pointing out certain errors in such factors in an earlier version of this paper.

invariance. Therefore, these equations should be treated as order α' equations, and thus letting $H \rightarrow dB$ in (3.6), and $H \cdot H \rightarrow dB \cdot (dB + 2\alpha' \Gamma)$ in (3.7) and (3.8).

Turning on v_L^α breaks supersymmetry even at order α' . This phenomenon has been well studied in particular in 10D [1] and it is known that restoring supersymmetry at order α' requires the addition of a Riemann curvature-squared term into the action roughly by letting, schematically, $\alpha' F^2 \rightarrow \alpha' (R^2 + F^2)$. In the ungauged 6D theory, similar phenomenon occurs, and such terms have been considered in [38] in the context of heterotic–heterotic string duality, and in [39], in the context of constructing Killing spinors.

Considering the gauged supergravities, while a Noether procedure has not been carried out completely as yet for the full system at order α' , taking into account [40], we expect the following result in the absence of hypermultiplet⁶

$$S = \int \left\{ \frac{1}{4} R(\omega) \star \mathbb{1} - \frac{1}{4} \star d\varphi \wedge d\varphi - \frac{1}{2} G_{\alpha\beta} \star (dB^\alpha) \wedge (dB^\beta + 2\alpha' \Gamma^\beta) + 16\pi^2 \alpha' \Omega_{\alpha\beta} B^\alpha \wedge Y^\beta \right. \\ \left. - \frac{1}{\sqrt{2}} \alpha' e \cdot (v_L \text{tr} \star R(\omega) \wedge R(\omega) + v_i \text{tr} \star F_i \wedge F_i) - \frac{1}{8\sqrt{2}\alpha'} (e \cdot v)^{-1} \star \mathbb{1} + \dots \right\}, \quad (3.10)$$

where the ellipses are yet to be determined $H = dB$ and dilaton-dependent terms.⁷ A similar action for the ungauged theory in string frame, albeit in a non-manifestly $SO(1,1)$ invariant form, was given in [38]. In obtaining the field equations from this action, the duality equation (3.5) is to be imposed *after* the variation of the action. With this in mind, it can be checked that this action gives the equations of motion (3.6)–(3.9), if v_L^α is set to zero. The inclusion of v_L^α effects will clearly introduce higher derivative terms in the Einstein's and dilaton field equations, though the consequences for the other field equations remain to be investigated, since the Noether procedure for the full system at order α' has not been established as yet. As for the term $\int \Omega_{\alpha\beta} B^\alpha \wedge Y^\beta$ in the above action, naturally it plays a crucial role in the discussion of Dirac quantization of dyonic string charges, as we shall see later.

Turning to the action (3.10), the requirement that the gauge kinetic terms are ghost-free imposes the constraints $e \cdot v_i > 0$ [37,41,42]. These kinetic terms for models A, B and C are given by

$$A: -\frac{\alpha'}{6} (e^{-\varphi} - 3e^\varphi) \text{tr} \star F_6 \wedge F_6 \\ - \frac{\alpha'}{4} (e^{-\varphi} + 3e^\varphi) \text{tr} \star F_7 \wedge F_7 - 2\alpha' (e^{-\varphi} - 9e^\varphi) \star F_1 \wedge F_1, \\ B: -\frac{3\alpha'}{2} (e^{-\varphi} - 5e^\varphi) \text{tr} \star F_2 \wedge F_2 \\ - \frac{\alpha'}{12} (3e^{-\varphi} - e^\varphi) \text{tr} \star F_7 \wedge F_7 - \frac{2\alpha'}{3} (3e^{-\varphi} + 19e^\varphi) \star F_1 \wedge F_1 \\ \text{and} \\ C: -\frac{\alpha'}{3} (e^{-\varphi} + 5e^\varphi) \text{tr} \star F_4 \wedge F_4 \\ - \frac{\alpha'}{2} (2e^{-\varphi} - e^\varphi) \text{tr} \star F_9 \wedge F_9 - \alpha' (2e^{-\varphi} - 19e^\varphi) \star F_1 \wedge F_1. \quad (3.11)$$

It is easy to check that the positivity condition for these kinetic terms are satisfied for

$$(A): \quad e^{-\varphi} > 3, \quad (B): \quad e^{-2\varphi} > 5 \quad \text{and} \quad (C): \quad e^{-2\varphi} > \frac{19}{2}. \quad (3.12)$$

The perturbative results are reliable for sufficiently negative values of φ , while the lower bounds on $e^{-\varphi}$ stated above correspond to the strong Yang–Mills coupling regime. It is also clear that there are a number of values for φ where some of the Yang–Mills couplings vanish. As discussed in detail in [38], these are points where phase transitions are expected to occur.

⁶In path integral considerations, the convenient normalization of the topological term is $\frac{1}{2}\alpha'\Omega_{\alpha\beta}B^\alpha \wedge Y^\beta$, which can be achieved by suitable rescalings of fields and parameters [25].

⁷In the absence of $U(1)_R$ gauging and if $v_2^\alpha = 0$, then such terms would be accounted for by shifting the spin connection occurring in the Riem^2 term by torsion as $\omega \rightarrow \omega - \frac{1}{2}H$.

The last terms in Einstein and dilaton field equations above involve a potential function, and arise as a consequence of the $U(1)_R$ gauging, and that they are absent in the ungauged 6D models, even if the gauge groups include ‘external’ $U(1)$ factors. These terms clearly have significant impact on the structure of the vacuum as well as the non-perturbative exact solutions. For example, it is easy to check that these terms forbid Minkowski₆ and $(A)dS$ vacuum solutions.

As for dyonic string solutions of $U(1)_R$ gauge theory, to our best knowledge, few solutions exist to equations in which only the classically exactly supersymmetric supergravity equations are solved. The action with $v_L^\alpha = 0$, $v_i^2 = 0$ [33] has been used to obtain the dyonic solutions mentioned above. Here, $B = B^1$ which represents the combination of the 2-forms residing in supergravity and single tensor multiplet, and therefore it is free from (anti)self duality condition. The action which can be read off from (3.10) by taking $v_L^\alpha = 0$, $v_i^2 = 0$, $B^2 = 0$ and setting $B^1 \equiv B$, $v^1 \equiv v$, takes the form⁸

$$S = \int \left(\frac{1}{4} R \star \mathbb{1} - \frac{1}{4} \star d\varphi \wedge d\varphi - \frac{1}{2} e^{-2\varphi} \star G \wedge G - \frac{1}{2} \alpha' e^{-\varphi} v_i \text{tr} \star F_i \wedge F_i - \frac{1}{8\alpha' v} e^\varphi \star \mathbb{1} \right), \quad (3.13)$$

where $G = dB^1 + \alpha' \Gamma^1|_{v_L=0}$. The solution found for the resulting equations has only the following non-vanishing fields [20], and it takes the form

$$ds^2 = c^2 dx^\mu dx_\mu + a^2 dr^2 + b^2 \left(\sigma_1^2 + \sigma_2^2 + \frac{4gP}{k} \sigma_3^2 \right),$$

$$G = P\sigma_1 \wedge \sigma_2 \wedge \sigma_3 - u(r) d^2x \wedge dr$$

and

$$F = k\sigma_1 \wedge \sigma_2, \quad e^{2\varphi} = \left(Q_0 + \frac{Q}{r^2} \right) \left(P_0 + \frac{P}{r^2} \right)^{-1}, \quad (3.14)$$

where a, b, c, u are functions of r which can be found in [20], k, P_0, Q_0, P, Q are constants, g is the $U(1)_R$ coupling constant, σ_i are left-invariant one-forms on the 3-sphere satisfying $d\sigma_i = -\frac{1}{2}\epsilon_{ijk}\sigma_j \wedge \sigma_k$. The solution also requires that⁹

$$4gP = k(1 - 2kg), \quad (3.15)$$

which is a condition not arising in the ungauged 6D theory, and it has 1/4 of the 6D supersymmetry. It is asymptotic to a cone over $(\text{Minkowski})_2 \times \text{squashed } S^3$, as opposed to the expected maximally symmetric known vacuum solution given by $\text{Minkowski}_4 \times S^2$ [12], and the dilaton blows up asymptotically [20]. The near horizon limit of the gauged dyonic string is given by $AdS_3 \times \text{squashed } S^3$ with fraction of supersymmetry increased from 1/4 to 1/2. A dyonic string solution of the $U(1)_R$ gauged theory in which an additional $U(1)$ gauge field residing in E_7 is activated was found in [21], under the assumptions that are similar to those of [20] outlined above. In particular, 1/4 supersymmetry also arises and again the dilaton blows up asymptotically.

4. Constraints on anomaly coefficients

The factorization of the anomaly polynomial has been shown to imply that [23]

$$a \cdot a, \quad a \cdot b_r \quad \text{and} \quad b_r \cdot b_s \in \mathbb{Z} \quad \text{for all } r, s, \quad (4.1)$$

where the products are in $\mathbb{R}^{1,1}$ with metric $\Omega_{\alpha\beta}$. The condition above can be checked explicitly for all three models studied here. The fact that the anomaly coefficients belong to an integral lattice is not sufficient for the consistency of the theory. To elaborate further on this point, it is convenient

⁸We have let $\varphi \rightarrow -\varphi/\sqrt{2}$ in the results of [33].

⁹Note that upon letting $A_\mu \rightarrow A_\mu/g$, this condition becomes $4P = k(1 - k(\alpha')^{-1})$.

to first re-express the form Y^α appearing in the Bianchi identity $dH^\alpha =$ in terms of characteristic forms, applied to the models considered here taking the form [25,27]

$$Y^\alpha = \frac{1}{4} a^\alpha p_1 - b_r^\alpha c_2^r + \frac{1}{2} c^\alpha (c_1)^2, \quad (4.2)$$

where p_1 , c_2 and c_1 are the Chern–Weyl representatives of the indicated cohomology classes defined as

$$p_1 = \frac{1}{8\pi^2} \text{tr } R^2, \quad c_2^r = -\frac{1}{8\pi^2} \left(\frac{1}{\lambda_r} \text{tr}_r F^2 \right) \quad \text{and} \quad c_1 = \frac{F_1}{2\pi}. \quad (4.3)$$

It is then argued in [25] that the string charge defined by the integral $\int_{\Sigma_4} Y$, where Σ_4 is any integral 4-cycle, must be cancelled by background self-dual strings. Consequently, it is argued that this charge must yield an element of the *unimodular string charge lattice* Λ_S , and this ‘string quantization condition’ is explicitly stated as¹⁰

$$\int_{\Sigma_4} Y \in \Lambda_S. \quad (4.4)$$

The fact that Λ_S is a unimodular, equivalently self-dual, lattice can be seen from basic arguments that can be found, for example, in [24,25].

The completeness hypothesis was taken a step further by Monnier *et al.* [25] who assumed that a consistent supergravity theory may be put on an arbitrary spin manifold and that any smooth gauge field configurations should be allowed in the supergravity ‘path integral’. The strategy employed in [25] is then to assume the generalized completeness hypothesis and obtain strong constraints by evaluating (4.4) on suitable chosen space–times M and gauge bundles. In particular, taking $M = CP^3$, and evaluating (4.4) along a suitable 4-cycle, they derive the condition (applied to the groups considered here) [25]

$$a, b_r, \frac{1}{2}c \in \Lambda_S, \quad \Lambda_S \text{ unimodular}. \quad (4.5)$$

A special case of this condition was derived earlier by Seiberg & Taylor [24] in the form

$$b_r \in \Lambda_S, \quad \Lambda_S \text{ unimodular}. \quad (4.6)$$

by demanding consistency of the theory by means of Dirac quantization of charges, once it is compactified on various spaces, such as T^2 , T^4 and CP^2 . It was also argued that the presence or absence of the Abelian factors in the gauge group does not effect their results, which depends only on the non-Abelian part of the gauge group.

In [25], it has also been shown that the constraint (4.5) is equivalent to the statement [25]

$$a \in \Lambda_S, \quad \frac{1}{2}\mathfrak{b} \in H^4(B\tilde{G}; \mathbb{Z}) \times \Lambda_S, \quad \Lambda_S \text{ unimodular}, \quad (4.7)$$

where $B\tilde{G}$ is the classifying space of the universal cover of the semisimple part of the gauge group G . The bilinear form $\mathfrak{b}$ in our case, where $G = \bigotimes_r G_r \times U(1)_R$ with $r = 1, 2$, can be written as

$$\mathfrak{b} = \bigoplus_r b_r K_r \oplus c. \quad (4.8)$$

Here, K_r is the canonically normalized Killing form,¹¹ with respect to which, the length squared of the longest simple root is 2 (for $U(1)$, the root length squared is 1). It has also been shown that this is equivalent to the statement that $\mathfrak{b}$ is an even Λ_S -valued bilinear form when restricted to the

¹⁰The anomaly coefficients are measured in units of α' which we set equal to one.

¹¹Here, we use standard math convention in which K_r is unit matrix of dimension spanning the rank of the underlying Lie algebra, upon its restriction to the Cartan subalgebra.

coroot lattice [25]. Specifically, (4.7) implies for any x, y inside the coroot lattice,

$$\frac{1}{2}\mathfrak{b}(x, x) \in \Lambda_S, \quad \text{and} \quad \mathfrak{b}(x, y) \in \Lambda_S \quad \text{for } x \neq y. \quad (4.9)$$

Taking into account the global structure of the gauge group, the condition (4.7) has been strengthened to [25]

$$a \in \Lambda_S, \quad \frac{1}{2}\mathfrak{b} \in H^4(BG; \mathbb{Z}) \times \Lambda_S, \quad \Lambda_S \text{ unimodular.} \quad (4.10)$$

which leads to conditions similar to (4.9) with x, y now belonging to the cocharacter lattice. For a detailed description of various lattices of G , see [43]. We only emphasize the following key aspects here. There is a general relation among the coroot lattice, cocharacter lattice and the coweight lattice for a given semisimple Lie algebra $\mathfrak{g}$ [43,44]

$$\Lambda^{\text{coroot}} \subseteq \Lambda^{\text{cocharacter}} \subseteq \Lambda^{\text{coweight}}. \quad (4.11)$$

These inclusions are determined by the global structure of the group G . Specifically, [43,44]

$$\frac{\Lambda^{\text{cocharacter}}}{\Lambda^{\text{coroot}}} = \pi_1(G) \quad \text{and} \quad \frac{\Lambda^{\text{coweight}}}{\Lambda^{\text{cocharacter}}} = Z(G), \quad (4.12)$$

where $\pi_1(G)$ is the first homotopy group of G and $Z(G)$ denotes the centre of G . For connected Lie groups, $H^4(BG; \mathbb{Z})$ is torsion free. For disconnected groups, there could potentially be a torsion class whose coefficient should be quantized in terms of the string charge lattice [26].

As mentioned in the introduction, Monnier and Moore extended the above considerations and arrived at a stronger criterion by seeking the conditions under which the Green–Schwarz counterterm is well defined. This leads to the requirement for the existence of a topologically trivial field theory in 7D, referred to as Wu–Chern–Simons theory, and a set of conditions for the 6D theory to be free from all anomalies. Applied to the cases under consideration, where the gauge groups are connected, the proposition states that given string charge lattice Λ_S , and the anomaly polynomial A_8 , and 4-form Y as defined in (4.2), assume that [26]

$$1. \quad A_8 = \frac{1}{2}Y \wedge Y; \quad (4.13)$$

$$2. \quad \Lambda_S \text{ is unimodular}; \quad (4.14)$$

$$3. \quad \mathfrak{b} \in 2H^4(BG; \Lambda_S); \quad (4.15)$$

$$4. \quad a \in \Lambda_S \text{ is a characteristic element}; \quad (4.16)$$

$$5. \quad \Omega_7^{\text{Spin}}(BG) = 0, \quad (4.17)$$

where $\Omega_7^{\text{Spin}}(BG) = 0$ is the spin cobordism group associated with Lie group G . Then all anomalies of the 6D theory, local and global, cancel. The ways in which this proposition extends (4.7) are as follows. Firstly, the derivation of the third condition does not rely on the generalized completeness hypothesis. Furthermore, the fourth condition states not only that $a \in \Lambda_S$ but it is also a characteristic element. Finally, the fifth condition clearly goes beyond what is required in (4.7).

5. Application of the consistency conditions

Monnier and Moore also tacitly assumes that string defects are included whenever they are necessary to satisfy the tadpole condition, and that their worldsheet anomalies cancel the boundary contributions to the anomaly of the supergravity theory through the anomaly inflow mechanism, as has been stated in [26]. Very recently, [30] proposed that using the gravitational and gauge anomaly inflow on the probe string, one can compute the worldsheet gravitational central charge and the gauge group’s current algebra level depending on the string charge and the bulk anomaly coefficients. (For earlier work in this context, see [28,29].) The requirement that the left-moving central charge should be large enough to allow the unitary representations of the current algebra for a given level imposes a constraint on the allowed gauge group content. However, the

fact that the near horizon limit of the gauged dyonic string is given by $1/2$ BPS $\text{AdS}_3 \times$ squashed S^3 suggests that the IR CFT of the probe string coupled to the gauged supergravity should be a two-dimensional $\mathcal{N} = 2$ CFT, in contrast to [30] where the worldsheet IR CFT is described by a $(0,4)$ CFT. Thus one cannot directly apply the result of [30] here before a careful study on the low energy dynamics of the probe string is carried out. Altogether, whether the tacit assumptions made as prelude to the Monnier–Moore proposition are satisfied by the $U(1)_R$ gauged 6D $(1,0)$ supergravities is not entirely clear, and remain to be investigated. Nonetheless, we shall at least assume that suitably behaved dyonic string solutions exist and proceed below with the analysis of the consequences of the above proposition for these models.

To begin with, condition 1 is obviously satisfied by models A , B and C . Next, we look at condition 5. To this end, we note that¹²

$$\begin{aligned} \Omega_7^{\text{Spin}}(BG_2) &= 0, & \Omega_7^{\text{Spin}}(BF_4) &= 0, & \Omega_7^{\text{Spin}}(BE_7) &= 0, & \Omega_7^{\text{Spin}}(BSp(9)) &= 0, \\ \Omega_7^{\text{Spin}}\left(\frac{BE_6}{\mathbb{Z}_3}\right) &= D_3, \end{aligned} \quad (5.1)$$

where D_3 is yet to be determined group of exponent 6. Since it is not known yet whether D_3 is trivial or not, we shall examine the other conditions required by the proposition in the case of model A which has the symmetry $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$. As for models B and C , given the results (5.1), they pass the fifth condition of the proposition.

For the convenience of further discussion, we introduce the notation

$$\mathcal{M}(x, y) = \begin{pmatrix} x \cdot x & x \cdot y \\ y \cdot x & y \cdot y \end{pmatrix}, \quad (5.2)$$

where x, y are $\mathbb{R}^{1,1}$ vectors and the product is defined with respect to $\Omega_{\alpha\beta}$. The fact that string charge lattice Λ_S is unimodular implies that $-\det \mathcal{M}(x, y)$ must be a square of a positive integer for any $x, y \in \Lambda_S$.

In using the relations (4.11), it is also useful to note that as far as the non-Abelian groups appearing in models A , B and C are concerned, $\pi_1(G) = \mathbb{1}$ and $Z(G) = \mathbb{1}$ for all, except that

$$\pi_1(E_6/\mathbb{Z}_3) = \mathbb{Z}_3, \quad Z(E_6) = \mathbb{Z}_3, \quad Z(E_7) = \mathbb{Z}_2 \quad \text{and} \quad Z(Sp(9)) = \mathbb{Z}_2. \quad (5.3)$$

In the following, we will test constraints stated in the proposition for models A , B and C , though, we will also see if only the weaker constraints (4.5) and/or (4.10) are satisfied in some cases.

(a) The $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$ invariant model

We first compute $-\det \mathcal{M}(x, y)$ for x, y being any two distinct $\mathbb{R}^{1,1}$ vectors among $a, b_6, b_7, \frac{1}{2}c$. The result is given by

$$\begin{aligned} -\det \mathcal{M}(a, b_6) &= 8^2, & -\det \mathcal{M}(a, b_7) &= 12^2, & -\det \mathcal{M}(a, \tfrac{1}{2}c) &= 20^2, \\ -\det \mathcal{M}(b_6, b_7) &= 18^2, & -\det \mathcal{M}(b_6, \tfrac{1}{2}c) &= 6^2, & -\det \mathcal{M}(b_7, \tfrac{1}{2}c) &= 36^2. \end{aligned} \quad (5.4)$$

therefore the anomaly coefficients in this model are compatible with the second condition of the proposition, namely with (4.14), that the anomaly coefficients lie on a unimodular string charge lattice. To verify that the lattice is indeed unimodular, we proceed by choosing as a basis of a unimodular charge lattice

$$e_1 = (1, 0) \quad \text{and} \quad e_2 = (0, 1), \quad (5.5)$$

and observe that the anomaly coefficients can be recast as linear combination of e_1, e_2 with integer coefficients. Note that this lattice is *even*.

Next, we inspect the anomaly coefficients against the stronger constraint (4.10). In order to do so, we need to evaluate the bilinear form b on the cocharacter lattice of $E_6 \times E_7 \times U(1)_R$.

¹²We are very grateful to I. García-Etxebarria and M. Montero, for explaining to us their results for $\Omega_7^{\text{Spin}}(BG)$ for G_2 (unpublished), F_4, E_6 and E_7 , and $E_6/\mathbb{Z}_3$ (unpublished).

In the model with gauged $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$ symmetry, E_6 appears only in the adjoint representation. Therefore, a vector v on the E_6 cocharacter lattice should satisfy

$$e^{2\pi i v_i h_i} = \mathbb{1}_{78 \times 78}, \quad (5.6)$$

where h_i , $i = 1, \dots, 6$ are generators of Cartan subalgebra (in the Cartan–Weyl basis) of E_6 in the adjoint representation. Clearly, such v lies in the coweight lattice of E_6 spanned by $\check{w}_m$ that obey $\sum_i (\check{w}_m)_i (r_n)_i = \delta_{mn}$, for simple roots labelled by r_n . Using the definition of coweights $\check{w}_m$, we can evaluate the bilinear form K_6 on the coweight lattice and obtain

$$K_6(\check{w}_r, \check{w}_s) = \frac{1}{3} \begin{pmatrix} 4 & 5 & 6 & 4 & 2 & 3 \\ 5 & 10 & 12 & 8 & 4 & 6 \\ 6 & 12 & 18 & 12 & 6 & 9 \\ 4 & 8 & 12 & 10 & 5 & 6 \\ 2 & 4 & 6 & 5 & 4 & 3 \\ 3 & 6 & 9 & 6 & 3 & 6 \end{pmatrix}, \quad (5.7)$$

which is equal to the inverse of the E_6 Cartan matrix. This happens to be so because Lie algebra of E_6 is simply laced and thus the length squared of every simple root equals 2, implying the coweight vector coincides with the fundamental weight vector. From the expression above, we single out a particular element $K_6(\check{w}_1, \check{w}_1)$ whose product with b_6 leads to the following vector on $\mathbb{R}^{1,1}$

$$\tilde{b}_6 = \frac{1}{2} b_6 K_6(\check{w}_1, \check{w}_1) = \frac{2}{3} b_6. \quad (5.8)$$

This gives $-\det \mathcal{M}(a, \tilde{b}_6) = (\frac{16}{3})^2$, which means that $\tilde{b}_6$ and a cannot belong to the same unimodular lattice. Thus, the third condition of the proposition, namely (4.15), is not satisfied.

(b) The $E_7 \times G_2 \times U(1)_R$ invariant model

Similar to the previous case, we first investigate whether the anomaly coefficients can be embedded in a unimodular lattice, by computing $-\det \mathcal{M}(x, y)$ for x, y being any two distinct $\mathbb{R}^{1,1}$ vectors among $a, b_2, b_7, \frac{1}{2}c$. It turns out that

$$\begin{aligned} -\det \mathcal{M}(a, b_2) &= 24^2, & -\det \mathcal{M}(a, b_7) &= 4^2, & -\det \mathcal{M}(a, \frac{1}{2}c) &= (\frac{44}{3})^2, \\ -\det \mathcal{M}(b_2, b_7) &= 42^2, & -\det \mathcal{M}(b_2, \frac{1}{2}c) &= 34^2, & -\det \mathcal{M}(b_7, \frac{1}{2}c) &= 20^2. \end{aligned} \quad (5.9)$$

As $-\det \mathcal{M}(a, \frac{1}{2}c)$ is not given by a positive integer squared, the anomaly coefficients $a, b_2, b_7, \frac{1}{2}c$ cannot all belong to a unimodular lattice, thus violating the second condition of the proposition, namely (4.14).

(c) The $F_4 \times Sp(9) \times U(1)_R$ invariant model

For this model, we obtain $-\det \mathcal{M}(x, y)$ for x, y being any two distinct $\mathbb{R}^{1,1}$ vectors among $a, b_4, b_9, \frac{1}{2}c$ as

$$\begin{aligned} -\det \mathcal{M}(a, b_4) &= 16^2, & -\det \mathcal{M}(a, b_9) &= 3^2, & -\det \mathcal{M}(a, \frac{1}{2}c) &= 21^2, \\ -\det \mathcal{M}(b_4, b_9) &= 11^2, & -\det \mathcal{M}(b_4, \frac{1}{2}c) &= 29^2, & -\det \mathcal{M}(b_9, \frac{1}{2}c) &= 9^2, \end{aligned} \quad (5.10)$$

which shows that the necessary condition for the lattice Λ_S being unimodular is satisfied. To establish that it is indeed unimodular, we proceed as follows. We choose the following basis for a unimodular charge lattice

$$e_1 = (2, 0) \quad \text{and} \quad e_2 = (1, \frac{1}{2}), \quad (5.11)$$

Next, we observe that the anomaly coefficients in this model can be expressed as linear combinations of e_1, e_2 with integer coefficients. This shows that condition (4.5) is indeed satisfied. Note also that the lattice here is *odd*, since $e_2 \cdot e_2 = 1$. Furthermore, in this model, the group F_4 appears only in the adjoint representation, whereas the hypermultiplet carries also the

fundamental representation of $Sp(9)$. One should also note that since the hyperfermions are singlet under $U(1)_R$, it is not possible to form an identity by combining a centre element of $Sp(9)$ with an element of $U(1)_R$.¹³ Since $Z(F_4)$ and $\pi_1(F_4)$ are all trivial, the coroot, cocharacter and coweight lattices are equivalent (4.12), the third condition of the proposition (4.15) reduces to the condition (4.5), which we have shown above to be satisfied.

We now move on to discuss the stronger constraint imposed on $Sp(9)$. We recall that $Z(Sp(9)) = \mathbb{Z}_2$ and $\pi_1(Sp(9)) = \mathbb{1}$. Thus the cocharacter lattice is different from coweight lattice but coincides with the coroot lattice. Indeed the transformation matrix from the standard coroot basis to the standard cocharacter basis is given by the unimodular matrix

$$\mathcal{T}_9 = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 & 2 & 2 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 & 3 & 3 & 3 & 3 & 3 \\ 1 & 2 & 3 & 4 & 4 & 4 & 4 & 4 & 4 \\ 1 & 2 & 3 & 4 & 5 & 5 & 5 & 5 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 & 6 & 6 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 7 & 7 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 8 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \end{pmatrix}. \quad (5.12)$$

Thus again, the third condition of the proposition (4.15) becomes equivalent to the condition (4.5) already shown to be satisfied by the explicit construction of the string charge lattice basis given in (5.11). Using (5.11) as the basis, an element on the charge lattice can be parameterized as

$$x = (2n + m, \frac{1}{2}m), \quad m, n \in \mathbb{Z}. \quad (5.13)$$

Thus one can easily show that

$$a \cdot x = x \cdot x \pmod{2}. \quad (5.14)$$

Given also that $\Omega_7^{\text{spin}}(BG) = 0$ for $G = F_4 \times Sp(9) \times U(1)_R$,¹⁴ we see that all the conditions of the proposition, namely (4.13)–(4.17) are satisfied, and therefore this model is free from all anomalies.

6. Conclusion

We have highlighted the significance of R -symmetry gauging in $6D, \mathcal{N} = (1, 0)$ supergravity, and focused on three such models that stand out in their accommodation of Green–Schwarz mechanism for the cancellation of all local anomalies in a non-trivial way. We have examined constraints imposed on the anomaly coefficients that are associated with the factorized anomaly polynomials in these models, as proposed in their strongest form by Monnier & Moore [26]. Adopting the assumptions made by these authors, we have found that only model C, based on the gauge group $F_4 \times Sp(9) \times U(1)_R$, satisfies all the conditions required for freedom from all anomalies, local and global. We have also seen that model A based on the gauge group $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$ does have a unimodular lattice, thus satisfying the weaker version of the consistency conditions on the anomaly coefficients [24], but it fails the stronger conditions of [25,26].

A word of caution is appropriate in applying the Monnier–Moore criteria to the R -symmetry gauged 6D supergravities for the following reason. It is assumed that dyonic strings with proper behaviour that give well defined string charge lattice exist. On the other hand, the existence of dyonic string excitations in these models are yet to be firmly established. The task is primarily complicated by the fact that the $U(1)_R$ gauging gives rise to a potential function which effects in a significant way the solution space, and in particular the asymptotic behaviour. The potential comes with an inverse power of α' , and certain dyonic string solutions in the presence of a

¹³This is different from the $U(2)$ example studied in [25], where element of the cocharacter lattice is formed by combining a centre element of $SU(2)$ with an element of the remaining $U(1)$.

¹⁴See [45] for $\Omega_7^{\text{spin}}(BF_4)$, where co-bordism groups have also been used in constraining varieties of models of physical interest. In the latter context, see also [46,47].

potential, and in which the anomaly coefficients v_α^2 (arising in the source term in the 2-form field equation) are set to zero, [20,21] require a relation among the parameters not seen in the usual dyonic string solutions of the ungauged 6D supergravities. Search and in depth study of the dyonic strings solutions of R-symmetry gauged 6D (1,0) supergravities is needed before a robust conclusion can be reached with regard to their global anomalies. In particular, the consequences for the existence of a worldsheet theory, and the attendant inflow anomalies require scrutiny, as they may impose yet further constraints on the consistency of the anomaly coefficients, as has been found to be the case for certain *ungauged* 6D supergravities with minimal supersymmetry [30].

Notwithstanding the caveat mentioned, we conclude by noting that it is still remarkable that the R-symmetry gauged model with $F_4 \times Sp(9) \times U(1)_R$ satisfies all the constraints of the Monnier–Moore proposition, which are most stringent ones known as yet. As such, it certainly deserves a closer look, to address further questions such as their place in the arena of swampland conjectures, even though, being conjectures, they are not as firm as the requirement of anomaly freedom so far. It would also be interesting to explore the dyonic string solutions and the charges they are allowed to carry, which can serve as a consistency check to the proposed charge lattice (5.11) implying that the minimal charge carried by a purely electric string (labelled by e_1 (5.11)) is twice as big as that of a purely magnetic string (labelled by $2e_2 - e_1$ (5.11)). A study of α' corrections due to supersymmetry, likely combined with other considerations such as unitarity and causality, may shed some light on the UV completion of the theory, if such a completion exists at all. Finally, it would be interesting to explore the application of the model to cosmology, as it may yield significantly different results compared to those of standard string cosmology, in view of the positive potential afforded by the R-symmetry gauging.

Data accessibility. This article does not contain any additional data.

Authors' contributions. Y.P. and E.S. carried out the research, findings of which are reported in this paper. Furthermore, both authors revised the paper and approve the final version and agree to be held accountable for all aspects of the work.

Competing interests. We declare we have no competing interest.

Funding. The work of Y.P. is supported by a Newton International Fellowship NF170385 of the Royal Society, and the work of E.S. is supported in part by NSF grant no. PHY-1803875.

Acknowledgements. We thank Eric Bergshoeff, Stanley Deser, André Henriques, Alex Kehagias and Yoshiaki Tanii for useful discussions. We are very grateful to Greg Moore for explaining his results with S. Monnier and D.S. Park on anomaly constraints, and Inaki Garcia-Etxebarria and Miguel Montero for explaining their results on spin bordism groups. We also thank Istanbul Technical University for hospitality during our visit.

Appendix A. The anomaly polynomials

The fields that contribute to gravitational, gauge and mixed anomalies in $n_T = 1$, $\mathcal{N} = (1,0)$ supergravity with gauge group $G = G_1 \times G_2 \times U(1)_R$ in 6D are as follows:

$$\psi_{\mu^A+}, \quad \chi_-^A, \quad \lambda_+^{IA} \quad \text{and} \quad \psi_-^{aa'}, \quad (\text{A } 1)$$

with chiralities denoted by $\pm$. The fermions are symplectic Majorana Weyl, the index $A = 1, 2$ labels the $SO(2)_R \subset Sp(1)_R$ fundamental, I labels the adjoint representation of the group G , and (aa') label the representation content of the hyperfermions under $G_1 \times G_2$.

(a) The $(E_6/\mathbb{Z}_3) \times E_7 \times U(1)_R$ model

From [10], we have

$$\left. \begin{aligned} Y^1 &= \frac{1}{16\pi^2} \left(\text{tr } R^2 + \frac{1}{3} \text{tr } F_6^2 + \frac{1}{2} \text{tr } F_7^2 + 4F_1^2 \right) \\ \text{and} \quad Y^2 &= -\frac{1}{16\pi^2} \left(\text{tr } R^2 - \text{tr } F_6^2 + \frac{3}{2} \text{tr } F_7^2 - 36F_1^2 \right). \end{aligned} \right\} \quad (\text{A } 2)$$

The computation of the anomaly polynomial can be found in [10] where the details of the computation are spelled out.¹⁵ The generators of the gauge group are taken to be Hermitian, and the strength of the $U(1)_R$ coupling constant to be unity, i.e. $D_\mu = \partial_\mu - iA_\mu$. It should be noted that the normalizations in Y^α are taken differently in various papers. However, following [25], we take them to be $1/(16\pi^2)$, motivated by the fact that this is the appropriate normalization in the integrals $\int_{\Sigma_4} Y$ discussed in §4, in which these integrals are related to Chern–Weyl classes. The freedom to do so stems from the fact that the anomaly coefficients are fixed in terms of α' which we can normalize appropriately, and set equal to one, after having done so.

(b) The $G_2 \times E_7 \times U(1)_R$ model

From [11], we have

$$\left. \begin{aligned} Y^1 &= \frac{1}{16\pi^2} \left(\text{tr} R^2 + 3\text{tr} F_2^2 + \frac{1}{2}\text{tr} F_7^2 + 4F_1^2 \right) \\ \text{and} \quad Y^2 &= \frac{1}{16\pi^2} \left(\text{tr} R^2 + 15\text{tr} F_2^2 + \frac{1}{6}\text{tr} F_7^2 - \frac{76}{3}F_1^2 \right) \end{aligned} \right\} \quad (\text{A } 3)$$

The details of the computations for this anomaly polynomial are provided in [11], where the generators of the gauge group are taken to be anti-Hermitian while here we are employing Hermitian generators. The $U(1)_R$ covariant derivative $D_\mu = \partial_\mu - iA_\mu$ is assumed.

(c) The $F_4 \times Sp(9) \times U(1)_R$ model

From the data provided in [7], we find

$$\left. \begin{aligned} Y^1 &= \frac{1}{16\pi^2} \left(\text{tr} R^2 + \frac{2}{3}\text{tr} F_4^2 + 2\text{tr} F_9^2 + 4F_1^2 \right) \\ \text{and} \quad Y^2 &= -\frac{1}{16\pi^2} \left(\text{tr} R^2 + \frac{10}{3}\text{tr} F_4^2 - \text{tr} F_9^2 - 38F_1^2 \right) \end{aligned} \right\} \quad (\text{A } 4)$$

As hyperinos transform as $(52, 18)_0$ under $F_4 \times Sp(9)$, and are neutral under $U(1)_R$, the contributions to the gravitation, gauge and mixed anomalies to the anomaly polynomial are

$$P(\psi_\mu) = \frac{245}{360}\text{tr} R^4 - \frac{43}{288}(\text{tr} R^2)^2 - \frac{19}{6}F_1^2\text{tr} R^2 + \frac{10}{3}F_1^4, \quad (\text{A } 5)$$

$$P(\chi) = -\left(\frac{1}{360}\text{tr} R^4 + \frac{1}{288}(\text{tr} R^2)^2 \right) - \frac{1}{6}F_1^2\text{tr} R^2 - \frac{2}{3}F_1^4, \quad (\text{A } 6)$$

$$\begin{aligned} P(\psi^{aa'}) &= -\frac{1}{2}(52 \times 18) \left(\frac{1}{360}\text{tr} R^4 + \frac{1}{288}(\text{tr} R^2)^2 \right) - \frac{1}{2} \times \frac{1}{6} (18\text{Tr} F_4^2 + 52\text{tr} F_9^2) \text{tr} R^2 \\ &\quad - \frac{1}{2} \times \frac{2}{3} (18\text{Tr} F_4^4 + 52\text{tr} F_9^4) - \frac{1}{2} \times 4\text{Tr} F_4^2\text{tr} F_9^2 \end{aligned} \quad (\text{A } 7)$$

$$\begin{aligned} \text{and} \quad P(\lambda) &= (52 + 171 + 1) \left(\frac{1}{360}\text{tr} R^4 + \frac{1}{288}(\text{tr} R^2)^2 \right) + \frac{1}{6}F_1^2\text{tr} R^2 + \frac{2}{3}F_1^4 \\ &\quad + \frac{1}{6}\text{Tr} F_4^4\text{tr} R^2 + \frac{1}{6}\text{Tr} F_9^2\text{tr} R^2 + \frac{2}{3}(\text{Tr} F_4^4 + \text{Tr} F_9^4) + 4F_1^2\text{Tr} F_4^2 + 4F_1^2\text{Tr} F_9^2, \end{aligned} \quad (\text{A } 8)$$

where Tr and tr denote the traces in the adjoint and fundamental representations, respectively. Here, the group generators are taken to be Hermitian, and for $U(1)$ we have $D_\mu = \partial_\mu - iA_\mu$, and

¹⁵See also [21], where few typos were corrected in the expressions for the individual contributions to the anomaly polynomial, without any effect on the total and, of course, its factorization.

F_4, F_9, F_1 are associated with $F_4 \times Sp(9) \times U(1)_R$. Using the relations

$$\text{Tr } F_4^2 = 3 \text{tr } F_4^2, \quad \text{Tr } F_4^4 = \frac{5}{12} \left(\text{tr } F_4^2 \right)^2 \quad (\text{A } 9)$$

and

$$\text{Tr } F_9^2 = 20 \text{tr } F_9^2, \quad \text{Tr } F_9^4 = 26 \text{tr } F_9^4 + 3 \left(\text{tr } F_9^2 \right)^2, \quad (\text{A } 10)$$

the sum I_8 becomes

$$\begin{aligned} I_8 = & - \left(\text{tr } R^2 \right)^2 + 34 F_1^2 \text{tr } R^2 + 152 F_1^4 - 4 \text{tr } F_4^2 \text{tr } R^2 - \text{tr } F_9^2 \text{tr } R^2 \\ & - \frac{20}{9} \left(\text{tr } F_4^2 \right)^2 + 2 \left(\text{tr } F_9^2 \right)^2 - 6 \text{tr } F_4^2 \text{tr } F_9^2 + 12 F_1^2 \text{tr } F_4^2 + 80 F_1^2 \text{tr } F_9^2. \end{aligned} \quad (\text{A } 11)$$

Arranging this data into a 4×4 matrix, it has rank 2, and it factorizes as in (A 4).

References

1. Bergshoeff EA, de Roo M. 1989 The quartic effective action of the heterotic string and supersymmetry. *Nucl. Phys. B* **328**, 439–468. (doi:10.1016/0550-3213(89)90336-2)
2. de Roo M, Suelmann H, Wiedemann A. 1993 The supersymmetric effective action of the heterotic string in ten-dimensions. *Nucl. Phys. B* **405**, 326–366. (doi:10.1016/0550-3213(93)90550-9)
3. de Roo M, Suelmann H, Wiedemann A. 1992 Supersymmetric R^{*4} actions in ten-dimensions. *Phys. Lett. B* **280**, 39–46. (doi:10.1016/0370-2693(92)90769-Z)
4. Tseytlin AA. 1996 Heterotic type I superstring duality and low-energy effective actions. *Nucl. Phys. B* **467**, 383–395. (<http://arxiv.org/abs/hep-th/9512081>) (doi:10.1016/0550-3213(96)00080-6)
5. Kim HC, Tarazi HC, Vafa C. 2019 Four dimensional $\mathcal{N} = 4$ SYM and the Swampland. (<http://arxiv.org/abs/1912.06144>)
6. Palti E. 2019 The swampland: introduction and review. *Fortsch. Phys.* **67**, 1900037. (<http://arxiv.org/abs/1903.06239>) (doi:10.1002/prop.201900037)
7. Avramis SD, Kehagias A. 2005 A Systematic search for anomaly-free supergravities in six dimensions. *J. High Energy Phys.* **0510**, 052. (<http://arxiv.org/abs/hep-th/0508172>) (doi:10.1088/1126-6708/2005/10/052)
8. Green MB, Schwarz JH, West PC. 1985 Anomaly free chiral theories in six-dimensions. *Nucl. Phys. B* **254**, 327–348. (doi:10.1016/0550-3213(85)90222-6)
9. Suzuki R, Tachikawa Y. 2006 More anomaly-free models of six-dimensional gauged supergravity. *J. Math. Phys.* **47**, 062302. (<http://arxiv.org/abs/hep-th/0512019>) (doi:10.1063/1.2209767)
10. Randjbar-Daemi S, Salam A, Sezgin E, Strathdee JA. 1985 An anomaly free model in six-dimensions. *Phys. Lett.* **151B**, 351–356. (doi:10.1016/0370-2693(85)91653-3)
11. Avramis SD, Kehagias A, Randjbar-Daemi S. 2005 A new anomaly-free gauged supergravity in six dimensions. *J. High Energy Phys.* **0505**, 057. (<http://arxiv.org/abs/hep-th/0504033>) (doi:10.1088/1126-6708/2005/05/057)
12. Salam A, Sezgin E. 1984 Chiral compactification on $Minkowski \times S^2$ of $N = 2$ Einstein-Maxwell supergravity in six-dimensions. *Phys. Lett.* **147B**, 47–51. (doi:10.1016/0370-2693(84)90589-6)
13. Cvetič M, Gibbons GW, Pope CN. 2004 A string and M theory origin for the Salam-Sezgin model. *Nucl. Phys. B* **677**, 164–180. (<http://arxiv.org/abs/hep-th/0308026>) (doi:10.1016/j.nuclphysb.2003.10.016)
14. Distler J, Sharpe E. 2010 Heterotic compactifications with principal bundles for general groups and general levels. *Adv. Theor. Math. Phys.* **14**, 335–397. (<http://arxiv.org/abs/hep-th/0701244>) (doi:10.4310/ATMP.2010.v14.n2.a1)
15. Maeda K i., Nishino H. 1985 Attractor universe in six-dimensional $N = 2$ supergravity Kaluza-Klein theory. *Phys. Lett.* **158B**, 381–387. (doi:10.1016/0370-2693(85)90437-X)
16. Halliwell JJ. 1987 Classical and quantum cosmology of the Salam-Sezgin model. *Nucl. Phys. B* **286**, 729–750. (doi:10.1016/0550-3213(87)90461-5)

17. Gibbons GW, Townsend PK. 1987 Cosmological evolution of degenerate vacua. *Nucl. Phys. B* **282**, 610–625. (doi:10.1016/0550-3213(87)90700-0)
18. Gibbons GW, Maeda K-i. 1988 Black holes and membranes in higher dimensional theories with dilaton fields. *Nucl. Phys. B* **298**, 741–775. (doi:10.1016/0550-3213(88)90006-5)
19. Anchordoqui LA, Antoniadis I, Lst D, Soriano JF, Taylor TR. H_0 tension and the string swampland. (<http://arxiv.org/abs/1912.00242>)
20. Gueven R, Liu JT, Pope CN, Sezgin E. 2004 Fine tuning and six-dimensional gauged $N = (1, 0)$ supergravity vacua. *Class. Quant. Grav.* **21**, 1001–1014. (<http://arxiv.org/abs/hep-th/0306201>) (doi:10.1088/0264-9381/21/4/019)
21. Randjbar-Daemi S, Sezgin E. 2004 Scalar potential and dyonic strings in 6-D gauged supergravity. *Nucl. Phys. B* **692**, 346–362. (<http://arxiv.org/abs/hep-th/0402217>) (doi:10.1016/j.nuclphysb.2004.05.023)
22. Jong DC, Kaya A, Sezgin E. 2006 6D Dyonic string with active hyperscalars. *J. High Energy Phys.* **0611**, 047. (<http://arxiv.org/abs/hep-th/0608034>) (doi:10.1088/1126-6708/2006/11/047)
23. Kumar V, Morrison DR, Taylor W. 2010 Global aspects of the space of 6D $N = 1$ supergravities. *J. High Energy Phys.* **1011**, 118. (<http://arxiv.org/abs/arXiv:1008.1062>) (doi:10.1007/JHEP11(2010)118)
24. Seiberg N, Taylor W. 2011 Charge lattices and consistency of 6D supergravity. *J. High Energy Phys.* **1106**, 001. (<http://arxiv.org/abs/arXiv:1103.0019>) (doi:10.1007/JHEP06(2011)001)
25. Monnier S, Moore GW, Park DS. 2018 Quantization of anomaly coefficients in 6D $\mathcal{N} = (1, 0)$ supergravity. *J. High Energy Phys.* **1802**, 020. (<http://arxiv.org/abs/1711.04777>) (doi:10.1007/JHEP02(2018)020)
26. Monnier S, Moore GW. 2019 Remarks on the Green-Schwarz terms of six-dimensional supergravity theories. *Commun. Math. Phys.* **372**, 963–1025. (<http://arxiv.org/abs/1808.01334>) (doi:10.1007/s00220-019-03341-7)
27. Monnier S, Moore GW. A brief summary of global anomaly cancellation in six-dimensional supergravity. (<http://arxiv.org/abs/1808.01335>)
28. Kim HC, Kim S, Park J. 2016 6d strings from new chiral gauge theories. (<http://arxiv.org/abs/1608.03919>)
29. Shimizu H, Tachikawa Y. 2016 Anomaly of strings of 6d $\mathcal{N} = (1, 0)$ theories. *J. High Energy Phys.* **1611**, 165. (<http://arxiv.org/abs/arXiv:1608.05894>) (doi:10.1007/JHEP11(2016)165)
30. Kim HC, Shiu G, Vafa C. 2019 Branes and the Swampland. *Phys. Rev. D* **100**, 066006. (<http://arxiv.org/abs/1905.08261> [hep-th]) (doi:10.1103/PhysRevD.100.066006)
31. Bernard CW, Christ NH, Guth AH, Weinberg EJ. 1977 Instanton parameters for arbitrary gauge groups. *Phys. Rev. D* **16**, 2967–2977. (doi:10.1103/PhysRevD.16.2967)
32. Bershadsky M, Vafa C. 2017 Global anomalies and geometric engineering of critical theories in six-dimensions. (<http://arxiv.org/abs/hep-th/9703167>)
33. Nishino H, Sezgin E. 1986 The complete $N = 2, d = 6$ supergravity with matter and Yang-Mills couplings. *Nucl. Phys. B* **278**, 353–379. (doi:10.1016/0550-3213(86)90218-X)
34. Nishino H, Sezgin E. 1997 New couplings of six-dimensional supergravity. *Nucl. Phys. B* **505**, 497–516. (<http://arxiv.org/abs/hep-th/9703075>) (doi:10.1016/S0550-3213(97)00357-X)
35. Riccioni F, Sagnotti A. 1998 Consistent and covariant anomalies in six-dimensional supergravity. *Phys. Lett. B* **436**, 298–305. (<http://arxiv.org/abs/hep-th/9806129>) (doi:10.1016/S0370-2693(98)00846-6)
36. Riccioni F. 2001 All couplings of minimal six-dimensional supergravity. *Nucl. Phys. B* **605**, 245–265. (<http://arxiv.org/abs/hep-th/0101074>) (doi:10.1016/S0550-3213(01)00199-7)
37. Sagnotti A. 1992 A note on the Green-Schwarz mechanism in open string theories. *Phys. Lett. B* **294**, 196–203. (<http://arxiv.org/abs/hep-th/9210127>) (doi:10.1016/0370-2693(92)90682-T)
38. Duff MJ, Minasian R, Witten E. 1996 Evidence for heterotic/heterotic duality. *Nucl. Phys. B* **465**, 413–438. (<http://arxiv.org/abs/hep-th/9601036>) (doi:10.1016/0550-3213(96)00059-4)
39. Fontanella A, Ortín T. 2019 On the supersymmetric solutions of the Heterotic Superstring effective action. (<http://arxiv.org/abs/1910.08496>)
40. Han SK, Kim JK, Koh IG, Tanii Y. 1986 Supersymmetrization of six-dimensional anomaly free $E_6 \times E_7 \times U(1)$ theory with Lorentz Chern-Simons term. *Phys. Lett. B* **177**, 167–170. (doi:10.1016/0370-2693(86)91048-8)

41. Seiberg N, Witten E. 1996 Comments on string dynamics in six-dimensions. *Nucl. Phys. B* **471**, 121–134. (<http://arxiv.org/abs/hep-th/9603003>) (doi:10.1016/0550-3213(96)00189-7)
42. Duff MJ, Lu H, Pope CN. 1996 Heterotic phase transitions and singularities of the gauge dyonic string. *Phys. Lett. B* **378**, 101–106. (<http://arxiv.org/abs/hep-th/9603037>) (doi:10.1016/0370-2693(96)00420-0)
43. Gukov S, Witten E. Gauge theory, ramification, and the geometric langlands program. (<http://arxiv.org/abs/hep-th/0612073>)
44. Simon B. 1996 *Representations of finite and compact groups*. Graduate Studies in Mathematics, vol. 10. Providence, RI: American Mathematical Society.
45. García-Etxebarria I, Montero M. 2019 Dai-Freed anomalies in particle physics. *J. High Energy Phys.* **1908**, 003. (<http://arxiv.org/abs/arXiv:1808.00009>) (doi:10.1007/JHEP08(2019)003)
46. Davighi J, Gripaos B, Lohitsiri N. 2019 Global anomalies in the standard model(s) and beyond. (<http://arxiv.org/abs/arXiv:1910.11277>)
47. Wan Z, Wang J. 2020 Beyond standard models and grand unifications: anomalies, topological terms, and dynamical constraints via cobordisms. (<http://arxiv.org/abs/arXiv:1910.14668>)