

Chern classes of automorphic vector bundles, II

Hélène Esnault and Michael Harris

Abstract. We prove that the ℓ -adic Chern classes of canonical extensions of automorphic vector bundles, over toroidal compactifications of Shimura varieties of Hodge type over $\bar{\mathbb{Q}}_p$, descend to classes in the ℓ -adic cohomology of the minimal compactifications. These are invariant under the Galois group of the p -adic field above which the variety and the bundle are defined.

Keywords. Automorphic bundles, Chern classes, Shimura varieties, perfectoids

2010 Mathematics Subject Classification. 11G18; 14G35

[Français]

Titre. Classes de Chern des fibrés vectoriels automorphes, II

Résumé. Nous démontrons que, sur les compactifications toroïdales des variétés de Shimura de type Hodge sur $\bar{\mathbb{Q}}_p$, les classes de Chern ℓ -adiques des extensions canoniques des fibrés vectoriels automorphes descendent en des classes de cohomologie ℓ -adique sur les compactifications minimales. Elles sont invariantes sous l'action du groupe de Galois du corps p -adique sur lequel la variété et le fibré sont définis.

Received by the Editors on July 25, 2018, and in final form on July 25, 2019.
Accepted on September 2, 2019.

Hélène Esnault

Freie Universität Berlin, Mathematik, Arnimallee 3, 14195 Berlin, Germany

e-mail: esnault@math.fu-berlin.de

Michael Harris

Department of Mathematics, Columbia University, 2990 Broadway New York, New York 10027, USA

e-mail: harris@math.columbia.edu

The research leading to these results has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) / ERC Advanced Grants 22224001 and 290766 (AAMOT), the Einstein Foundation, the NSF grants DMS-1404769 and DMS-1701651.

Contents

Introduction	2
1. Generalities on automorphic bundles	4
2. Chern classes for compactified Shimura varieties	5
3. Construction of perfectoid covers	13
4. Construction of ℓ-adic Chern classes	16
5. Comparison of torsors	22
Appendix A. Geometry	24
References	26

Introduction

Let S be a Shimura variety. It is defined over a number field [Del71, Corollaire 5.5], called the reflex field E , and carries a family of automorphic vector bundles $\mathcal{E}$, defined (collectively) over the same number field ([Har85, Theorem 4.8]; the rationality conventions of [EH17, §1] are recalled in §1 below). The Shimura variety has a minimal compactification $S \hookrightarrow S^{\min}$ which in dimension ≥ 2 is singular. This is minimal in the sense that each of the toroidal compactifications S_{Σ}^{tor} , subject to the extra choice of a fan Σ , admits a birational proper morphism $S_{\Sigma}^{\text{tor}} \rightarrow S^{\min}$, which is an isomorphism on S and (under additional hypotheses on the choice of fan and level subgroup, that we will make systematically) is a desingularization of $S^{\min}$. The automorphic bundle $\mathcal{E}$ admits a canonical extension $\mathcal{E}^{\text{can}}$ on S_{Σ}^{tor} constructed by Mumford [Mum77, Theorem 3.1] and more generally in [Har89]; $\mathcal{E}^{\text{can}}$ is locally free and defined over a finite extension of the reflex field. On $S^{\min}$, on the other hand, there is generally no locally free extension of $\mathcal{E}$, except in dimension 1 when $S_{\Sigma}^{\text{tor}} \rightarrow S^{\min}$ is an isomorphism. In [EH17] we studied the continuous ℓ -adic Chern classes of $\mathcal{E}$ in case S is proper. In particular, we proved that the higher Chern classes in continuous ℓ -adic cohomology die if $\mathcal{E}$ is flat [EH17, Theorem 0.2]. A key ingredient in the proof is the study of the action of Hecke algebra on continuous ℓ -adic cohomology, and the fact that Chern classes lie in the eigenspace for the volume character [EH17, Corollary 1.18]. When S is not proper, the Hecke algebra does not act on any cohomology of S_{Σ}^{tor} . Thus we cannot apply the methods of *loc.cit.* to prove the analogous theorem for the canonical extensions of flat automorphic bundles on S^{tor} .

On the other hand, the Hecke algebra does act on the ℓ -adic cohomology of $S^{\min}$. In addition, over the field of complex numbers $\mathbb{C}$, Goresky-Pardon [GP02, Theorem] used explicit estimates on differential forms to construct classes $c_n(\mathcal{E})^{GP} \in H^{2n}(S^{\min}, \mathbb{C})$ which descend the Chern classes of $\mathcal{E}^{\text{can}}$ in Betti cohomology $H^{2n}(S_{\Sigma}^{\text{tor}}, \mathbb{Q}(n))$. They do not come from classes in $H^{2n}(S^{\min}, \mathbb{Q}(n))$; precisely this fact enabled Looijenga [Loo17, Theorem 5.1] to construct mixed Tate extensions in the Hodge category on the Siegel modular variety (see also the result announced by Nair in [Nai14, §0.4]).

Our main theorem gives an ℓ -adic version of the Goresky-Pardon construction.

Theorem 0.1 (Theorem 4.9, Theorem 4.13). *Let S be a Shimura variety of Hodge type. Let ℓ be a prime number, E_w the completion of E at a place w dividing the prime p different from ℓ , and $E_w \hookrightarrow \bar{\mathbb{Q}}_p$ be an algebraic closure. Then the Chern classes of $\mathcal{E}^{\text{can}}$ in $H^{2n}(S_{\Sigma, \bar{\mathbb{Q}}_p}^{\text{tor}}, \mathbb{Q}_{\ell}(n))$ descend to well defined classes $c_n(\mathcal{E})$ in $H^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(n))$. The $c_n(\mathcal{E})$ are contravariant for the change of level and of Shimura data, verify the Whitney*

product formula, lie in the eigenspace for the volume character of the Hecke algebra, and are invariant under the action of the Galois group of E_w on $H^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n))$. Their pull-back under

$$H^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n)) \rightarrow H^{2n}(S_{\bar{\mathbb{Q}}_p}^{\text{tor}}, \mathbb{Q}_\ell(n)) = H^{2n}(S_{\Sigma, \bar{\mathbb{Q}}}^{\text{tor}}, \mathbb{Q}_\ell(n))$$

are the Chern classes of $\mathcal{E}^{\text{can}}$ which are invariant under the Galois group of E .

Here, as in [EH17, Lemma 1.15], the *volume character* is the character by which the Hecke algebra acts on the constant functions. The theorem can easily be extended to Shimura varieties of abelian type (see Remarks 4.19) but we leave the precise statement to the reader.

As the classes of $\mathcal{E}^{\text{can}}$ are even defined in continuous ℓ -adic cohomology of $S_{\bar{\mathbb{Q}}}^{\text{tor}}$, where $E \hookrightarrow \bar{\mathbb{Q}}$ is an algebraic closure, they are in particular invariant under the Galois group of E . The classes we construct rely on p -adic geometry and have no reason a priori not to depend on the chosen p , let alone to lie in $H^{2n}(S_{\bar{\mathbb{Q}}}^{\min}, \mathbb{Q}_\ell(n))^{G_E}$ via the isomorphism $H^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n)) = H^{2n}(S_{\bar{\mathbb{Q}}}^{\min}, \mathbb{Q}_\ell(n))$.

We now describe the method of proof. The main tool used is the existence of a perfectoid space $P^{\min}$ above¹ $S^{\min}$ which has been constructed by Peter Scholze in [Sch15, Theorem 4.1.1] for Shimura varieties of Hodge type, together with a Hodge-Tate period map $\pi_{HT} : P^{\min} \rightarrow \hat{\mathcal{X}}$ (see [CS17, Theorem 2.3] for the final form) with values in the adic space of the compact dual variety defined over the completion of $\bar{\mathbb{Q}}_p$. This allows us to define *vector bundles* $\pi_{HT}^*(\mathcal{E})$ on $P^{\min}$ where $\mathcal{E}$ are equivariant vector bundles on $\hat{\mathcal{X}}$ defining the automorphic bundles. So while the bundles $\mathcal{E}$ do not extend to $S^{\min}$, they do on Scholze's limit space $P^{\min}$.

On the other hand, suppose S is the Siegel modular variety. Then Pilloni and Stroth [PS16, Corollaire 1.6] have constructed a perfectoid space P^{tor} above S^{tor} . On P^{tor} one has the pull-back of the bundles $\pi_{HT}^*(\mathcal{E})$ and the pull-back via the tower defining the perfectoid space of the canonical extensions $\mathcal{E}^{\text{can}}$. Theorem 2.22 asserts that both pull-backs on P^{tor} are the same. This argument generalizes to Shimura varieties satisfying a certain technical Hypothesis 2.18. We expect Hypothesis 2.18 to hold in general; however, for general Shimura varieties of Hodge type we provide a more indirect route in §3 to the same comparison (Theorem 3.12), following a suggestion of Bhargav Bhatt.

Descent for cohomology with torsion coefficients enables one to construct the classes as claimed in Theorem 0.1, see Appendix A.

The classes $c_n(\mathcal{E}) \in H^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n))$ map in particular to well-defined classes in ℓ -adic intersection cohomology $IH^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n))$. On intersection cohomology of $S^{\min}$ over $\bar{\mathbb{Q}}$, we identify the eigenspace under the volume character with the cohomology of the compact dual $\hat{\mathcal{X}}$ of S (Proposition 4.16). If the classes in $IH^{2n}(S_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n))$, identified with $IH^{2n}(S_{\bar{\mathbb{Q}}}^{\min}, \mathbb{Q}_\ell(n))$, descended to continuous ℓ -adic intersection cohomology, we could try to apply the method developed in [EH17] to show that the classes of $\mathcal{E}^{\text{can}}$ in continuous ℓ -adic cohomology on S^{tor} over E die when $\mathcal{E}$ is flat. This would be in accordance with [EV02, Theorem 1.1] where it is shown that the higher Chern classes of $\mathcal{E}^{\text{can}}$ in the rational Chow groups on the Siegel modular variety vanish when $\mathcal{E}$ is flat. However, we are not yet able to work with continuous ℓ -adic intersection cohomology. We prove in Lemma 4.18 the vanishing $c_n(\mathcal{E}) = 0$ for all $n > 0$ when $\mathcal{E}$ is flat.

While we were writing the present note, it was brought to our attention that Nair in the unpublished manuscript [Nai14] independently mentioned the possibility of using Scholze's Hodge-Tate morphism to construct Chern classes in the cohomology of minimal compactifications, see *loc. cit.* §0.4.

Acknowledgements. This note is based on Peter Scholze's construction of a perfectoid space above the minimal compactification of Shimura varieties of Hodge type together with the Hodge-Tate period map.

¹ ↑ We are using a simplified notation here for the space denoted $K^p\mathcal{S}(G, X)^{\min}$ in the body of the text. Automorphic vector bundles are also denoted $[\mathcal{E}]$ rather than $\mathcal{E}$, and the superscript $^{\text{tor}}$ is absent in most cases.

We thank him profoundly for the numerous discussions on his theory, for his precise and generous answers to our questions. We thank Vincent Pilloni for explaining to us the content of [PS16, Corollaire 1.6] which plays an important role in our note, and for pointing out some oversights in an earlier version of this note. We thank David Hansen for providing the keys to navigating the rapidly expanding literature on perfectoid geometry; we thank Hansen again, as well as Laurent Fargues, for their patient and detailed explanations that allowed us to correct misunderstandings that one of the referees pointed out in an earlier version of this paper. We thank Mark Goresky for helpful discussions, Simon Pépin Lehalleur who brought the reference [Nai14, §0.4] to our attention, and Ana Caraiani, who helped us to understand the issues that forced us to take an unexpected detour in the proof of Proposition 3.11. We are deeply indebted to the two referees who helped us improving our article with their precise and helpful comments. We apologize for the work we unnecessarily gave them through our numerous typos. Finally, we are grateful to Bhargav Bhatt for a number of crucial suggestions, notably for the construction of Z_s in (3.9).

1. Generalities on automorphic bundles

If G is a reductive algebraic group over $\mathbb{Q}$, by an *admissible irreducible representation* of $G(\mathbf{A})$ we mean an irreducible admissible $(\mathfrak{g}, K) \times G(\mathbf{A}_f)$ -module, where $\mathfrak{g}$ is the complexified Lie algebra of G , $K \subset G(\mathbb{R})$ is a connected subgroup generated by the center of $G(\mathbb{R})$ and a maximal compact connected subgroup, $G(\mathbf{A})$ ($G(\mathbf{A}_f)$) is the group of (finite) adèles of G . If π is such a representation then we write

$$\pi \simeq \pi_\infty \otimes \pi_f$$

where π_∞ is an irreducible admissible $(\mathfrak{g}, K)$ -module and π_f is an irreducible admissible representation of $G(\mathbf{A}_f)$.

Let (G, X) be a Shimura datum: G is a reductive algebraic group over $\mathbb{Q}$, and X is a (finite union of) hermitian symmetric spaces, endowed with a transitive action of $G(\mathbb{R})$ satisfying a familiar list of axioms. The compact dual $\hat{X}$ of X is a (projective) flag variety for G . Thus one can speak of the category $\text{Vect}_G(\hat{X})$ of G -equivariant vector bundles on $\hat{X}$; the choice of a base point $h \in X \subset \hat{X}$ determines an equivalence of categories $\text{Rep}(P_h) \simeq \text{Vect}_G(\hat{X})$, where $P_h \subset G$ is the stabilizer of h , a maximal parabolic subgroup of G , and for any algebraic group H , $\text{Rep}(H)$ denotes the tensor category of its representations. Let K_h be the Levi quotient of P_h ; the group $K_{h,\infty} := K_h(\mathbb{C}) \cap G(\mathbb{R})$ can be identified with a maximal Zariski connected subgroup of $G(\mathbb{R})$ that is compact modulo the center of G . If $K \subset G(\mathbf{A}_f)$ is an open compact subgroup, we let ${}_K S(G, X)$ denote the Shimura variety attached to (G, X) at level K ; it has a canonical model over the *reflex field* $E(G, X)$ which is a number field (see [Del71, Corollaire 5.5]). We always assume that K is *neat*; then ${}_K S(G, X)$ is a smooth quasi-projective variety.

Furthermore, we use the notation and conventions of [EH17, §1.1–1.2], specifically $\mathcal{E} \in \text{Vect}(\hat{X})$ the notation for the vector bundle on the compact dual, $[\mathcal{E}]_K \in \text{Vect}({}_K S(G, X))$ for the automorphic vector bundle associated to the underlying representation of the compact (mod center) group $K_{h,\infty} \subset G(\mathbb{R})$, the stabilizer of a chosen point $h \in X$. As in [EH17], we always assume that K_h is defined over a CM field and that every irreducible representation of K_h has a model rational over the CM field E'_h ; in particular, P_h is also defined over E'_h . For the purposes of constructing Chern classes, we need only consider semisimple representations of P_h , which necessarily factor through representations of K_h .

The action of the Hecke algebra $\mathcal{H}_K = \{T(g), g \in G(\mathbf{A}_f)\}$ is recalled in Section 4. The Chern character

$$ch_{\hat{X}} : \text{Vect}_G(\hat{X}) \rightarrow CH(\hat{X})_{\mathbb{Q}} \rightarrow \oplus_i H^{2i}(\hat{X}, \mathbb{Q}(i))$$

induces isomorphisms

$$K_0(\text{Rep}(K_h)) \otimes_{K_0(\text{Rep}(G))} \mathbb{Q} \xrightarrow{\sim} CH(\hat{X})_{\mathbb{Q}} \xrightarrow{\sim} \oplus H^{2i}(\hat{X}, \mathbb{Q}(i)) \quad (1.1)$$

and analogously in ℓ -adic cohomology. On the other hand, the construction of automorphic vector bundles gives rise to a homomorphism

$$ch_K : \begin{cases} \text{Vect}_G(\hat{X}) & \rightarrow CH({}_K S(G, X))_{\mathbb{Q}, v} \\ \mathcal{E} & \mapsto ch([\mathcal{E}]_K) \end{cases} \quad (1.2)$$

where v indicates the invariants under the volume character. If $\mathcal{E} \in \text{Vect}_G(\hat{X})$ is in the image of $\text{Rep}(G)$ via the factorization

$$\text{Rep}(G) \rightarrow \text{Rep}(K_h) \rightarrow \text{Vect}_G(\hat{X}),$$

then $[\mathcal{E}]$ is endowed with a natural flat connection, so its higher Chern classes in $\oplus_i H^{2i}({}_K S(G, X), \mathbb{Q}(i))$ are equal to 0. We obtain a morphism

$$K_0(\text{Rep}(K_h)) \otimes_{K_0(\text{Rep}(G))} \mathbb{Q} \rightarrow \oplus_i H^{2i}({}_K S(G, X), \mathbb{Q}(i))_v \subset \oplus_i H^{2i}({}_K S(G, X), \mathbb{Q}(i)), \quad (1.3)$$

analogous to (1.1), where the subscript v denotes a certain eigenspace for the action of the unramified Hecke algebra.

2. Chern classes for compactified Shimura varieties

2.A. Toroidal and minimal compactifications, and canonical extensions of automorphic vector bundles

Henceforward we assume the Shimura variety ${}_K S(G, X)$ is *not projective*; equivalently, the derived subgroup G^{der} of G is isotropic over $\mathbb{Q}$. In this case the automorphic theory naturally gives information about Chern classes of canonical extensions on toroidal compactifications on the one hand; on the other hand, the v -eigenspace most naturally appears in intersection cohomology of the minimal compactification. This has been worked out in detail in the $\mathcal{C}^\infty$ and the L_2 theory by Goresky and Pardon in [GP02]. In what follows, we let $j_K : {}_K S(G, X) \hookrightarrow {}_K S(G, X)^{\min}$ denote the minimal (Baily-Borel-Satake) compactification. The minimal compactification is functorial, thus if $K' \subset K$, there is a unique morphism ${}_K S(G, X)^{\min} \rightarrow {}_{K'} S(G, X)^{\min}$ extending the natural map of open Shimura varieties, and if $g \in G(\mathbf{A}_f)$, then the Hecke correspondence $T(g)$ extends canonically to a correspondence on ${}_K S(G, X)^{\min} \times {}_K S(G, X)^{\min}$ where the product is taken over the reflex field. In particular, for any cohomology theory H as above, we have Hecke operators

$$T(g) \in \text{End}(H^*({}_K S(G, X)^{\min})).$$

The minimal compactification is always singular, except when G^{ad} is a product of copies of $\text{PGL}(2)$, and the automorphic vector bundles do not in general extend as bundles to ${}_K S(G, X)^{\min}$. Thus the classical theory does not automatically attach Chern classes to automorphic vector bundles on non-proper Shimura varieties in some cohomology theory $H^*({}_K S(G, X)^{\min}, *)$.

On the other hand, there is a large collection of *toroidal compactifications* ${}_K S(G, X) \hookrightarrow {}_K S(G, X)_\Sigma$ indexed by combinatorial data Σ (see [AMRT75, III, §6, Main Theorem] for details; the adelic construction is in [Har89, Pin90]). The set of Σ is adapted to the level subgroup K . It is partially ordered by *refinement*: if Σ' is a refinement of Σ , then there is a natural proper morphism

$$p_{\Sigma', \Sigma} : {}_K S(G, X)_{\Sigma'} \rightarrow {}_K S(G, X)_\Sigma$$

extending the identity map on the open Shimura variety. Any two Σ and Σ' can be simultaneously refined by some Σ'' . Further, the open embedding ${}_K S(G, X) \hookrightarrow {}_K S(G, X)_\Sigma$, is completed to a diagram

$$\begin{array}{ccc} & & {}_K S(G, X)_\Sigma \\ & \nearrow j_{K, \Sigma} & \downarrow \varphi_\Sigma \\ {}_K S(G, X) & \xrightarrow{j_K} & {}_K S(G, X)^{\min} \end{array} \quad (2.1)$$

where φ_Σ is a proper morphism, which is an isomorphism on ${}_K S(G, X)$. Now assume that K is a *neat* open compact subgroup of $G(\mathbf{A}_f)$, in the sense of [Har89] or [Pin90, 0.6]. In that case, we can choose Σ so that ${}_K S(G, X)_\Sigma$ is smooth and projective, and the boundary divisor has normal crossings. *We do so unless we specify otherwise.* With these conventions, φ_Σ is always a desingularization of the minimal compactification, which is constructed as in [AMRT75], *loc.cit.*; moreover, for any refinement Σ' of Σ , ${}_K S(G, X)_{\Sigma'}$ is again smooth and projective.

Mumford proved in [Mum77, Theorem 3.1] that, if $\mathcal{E} \in \text{Vect}_G^{\text{ss}}(\hat{X})$ (recall from [EH17] that the upper index ss stands for semi-simple) the automorphic vector bundle $[\mathcal{E}]_K$ on ${}_K S(G, X)$ admits a *canonical extension* $[\mathcal{E}]_K^{\text{can}}$ to ${}_K S(G, X)_\Sigma$; we write $[\mathcal{E}]_K^\Sigma$ if we want to emphasize the toroidal data. The adelic construction is carried out in Section 4 of [Har89], where Mumford's result was generalized to arbitrary $\mathcal{E} \in \text{Vect}_G(\hat{X})$. In particular it was shown there that if $\mathcal{E} = (G \times W)/P_h$, where W is the restriction to P_h of a representation of G , the action of P_h is diagonal, and we have identified G/P_h with $\hat{X}$, then $[\mathcal{E}]_K$ is a vector bundle on ${}_K S(G, X)$ with a flat connection, and its canonical extension $[\mathcal{E}]_K^{\text{can}}$ on ${}_K S(G, X)_\Sigma$ is exactly Deligne's canonical extension. In particular, the connection has logarithmic poles along ${}_K S(G, X)_\Sigma \setminus {}_K S(G, X)$. Moreover, if Σ' is a refinement of Σ , then

$$p_{\Sigma', \Sigma}^* [\mathcal{E}]_K^\Sigma = [\mathcal{E}]_K^{\Sigma'}$$

(see [Har90, Lemma 4.2.4]), which in particular implies

$$p_{\Sigma', \Sigma, *} [\mathcal{E}]_K^{\Sigma'} = R p_{\Sigma', \Sigma, *} [\mathcal{E}]_K^{\Sigma'} = [\mathcal{E}]_K^\Sigma.$$

Finally, for any fixed Σ ,

$$\mathcal{E} \rightarrow [\mathcal{E}]_K^{\text{can}}$$

is a monoidal functor from $\text{Vect}_G(\hat{X})$ to the category of vector bundles on ${}_K S(G, X)_\Sigma$.

Unfortunately, the $g \in G(\mathbf{A}_f)$ generally permute the set of Σ and thus the Hecke correspondences $T(g)$ in general *do not* extend to correspondences on a given ${}_K S(G, X)_\Sigma$. Thus the arguments of [EH17], which are based on the study of the v -eigenspace for the Hecke operators in the cohomology of ${}_K S(G, X)$, cannot be applied directly to prove vanishing of higher Chern classes of $[\mathcal{E}]_K^{\text{can}}$ in continuous ℓ -adic cohomology of ${}_K S(G, X)_\Sigma$ over its reflex field, when $[\mathcal{E}]_K$ is a flat automorphic vector bundle.

One can obtain information about the classes of $[\mathcal{E}]_K$ on the open Shimura variety, but these lose information. From the standpoint of automorphic forms, the natural target of the topological Chern classes of automorphic vector bundles should be the *intersection cohomology* of the minimal compactification. We fix cohomological notation: $H^*(Z, \mathbb{Q})$, resp. $IH^*(Z, \mathbb{Q})$ denotes Betti resp. intersection cohomology of a complex variety Z , while $H^*(Z, \mathbb{Q}_\ell)$, resp. $IH^*(Z, \mathbb{Q}_\ell)$ denote $\mathbb{Q}_\ell$ -étale cohomology.

Remark 2.2. The action of Hecke correspondences on Betti intersection cohomology $IH({}_K S(G, X), \mathbb{Q})$ are defined analytically by reference to Zucker's conjecture. For a purely geometric construction of the action of Hecke correspondences on $IH({}_K S(G, X), \mathbb{Q})$ and thus on $IH({}_K S(G, X), \mathbb{Q}_\ell)$ see [GM03, (13.3)] (the argument applies more generally to *weighted cohomology* as defined there).

The following statement generalizes Proposition 1.20 of [EH17] and is proved in the same way.

Proposition 2.3. *There is a canonical isomorphism of algebras*

$$\begin{aligned} H^*(\hat{X}, \mathbb{Q}) &\xrightarrow{\sim} IH^*({}_K S(G, X)^{\min}, \mathbb{Q})_v \\ H^*(\hat{X}, \mathbb{Q}_\ell) &\xrightarrow{\sim} IH^*({}_K S(G, X)^{\min}, \mathbb{Q}_\ell)_v. \end{aligned}$$

Proof. The second statement is deduced directly from the first one by the comparison isomorphism [BBD82, Section 6]. As in the proof of the analogous fact in [EH17], it suffices to prove the corresponding statement over $\mathbb{C}$. By Zucker's Conjecture [Loo88, SS90], $IH^*({}_K S(G, X)^{\min}, \mathbb{C})$ is computed using *square-integrable*

automorphic forms and Matsushima's formula [Bor83, 3.6, formula (I)]. Say the space $\mathcal{A}_{(2)}(G)$ of square-integrable automorphic forms on $G(\mathbb{Q}) \backslash G(\mathbb{A})$ decomposes as the direct sum

$$\mathcal{A}_{(2)}(G) = \oplus_{\pi} m(\pi) \pi$$

where π runs over irreducible admissible representations of $G(\mathbb{A})$ and $m(\pi)$ is a non-negative integer, which is positive for a countable set of π ². Then

$$IH^i({}_K S(G, X)^{\min}, \mathbb{C}) \xrightarrow{\sim} \oplus_{\pi} m(\pi) (H^i(\mathfrak{g}, K_h; \pi_{\infty}) \otimes \pi_f^K).$$

This implies

$$IH^i({}_K S(G, X), \mathbb{C})_v \xrightarrow{\sim} \oplus_{\pi} m(\pi) (H^i(\mathfrak{g}, K_h; \pi_{\infty}) \otimes (\pi_f^K)_v), \quad (2.4)$$

where $(\pi_f^K)_v$ is the eigenspace in π_f^K for the volume character of $\mathcal{H}_K$. Write $\pi_f = \otimes'_q \pi_q$, where q runs over rational primes. Now if q is unramified for K then $\pi_f^K = 0$ unless π_q is spherical. Now the trivial representation of $G(\mathbb{Q}_q)$ has the property that it equals its spherical subspace and the corresponding representation of the local Hecke algebra is the (local) volume character. By the Satake parametrization the trivial representation is the only spherical representation of $G(\mathbb{Q}_q)$ with this property. Thus $(\pi_f^K)_v \neq 0$ implies that π_q is the trivial representation for all q that are unramified for K . It then follows from weak approximation that π is in fact the trivial representation. Thus for all i ,

$$IH^i({}_K S(G, X), \mathbb{C})_v \xrightarrow{\sim} H^i(\mathfrak{g}, K_h; \mathbb{C}). \quad (2.5)$$

But this is equal to $H^i(\hat{X}, \mathbb{C}) = IH^i(\hat{X}, \mathbb{C})$ by a standard calculation; see [GP02, Remark 16.6]. The rest of the proof follows as in the proof of [EH17, Proposition 1.20]. $\square$

The point of the preceding proposition is that the Chern classes of automorphic vector bundles on the *non-proper* Shimura variety ${}_K S(G, X)$ are represented by square-integrable automorphic forms, more precisely, by the differential forms on $G(\mathbb{Q}) \backslash G(\mathbb{A})$ that are invariant under the action of $G(\mathbb{A})$. In other words, the only π that contributes is the space of constant functions on $G(\mathbb{Q}) \backslash G(\mathbb{A})$, which are square integrable modulo the center of $G(\mathbb{R})$. Thus $IH^{2*}({}_K S(G, X)^{\min}, \mathbb{Q}(*))_v$ can be viewed as L_2 -Chern classes of automorphic vector bundles. In addition $IH^{2*+1}({}_K S(G, X)^{\min}, \mathbb{Q}(*))_v = 0$. Although most automorphic vector bundles do not extend as bundles to the minimal compactification ${}_K S(G, X)^{\min}$, it was proved by Goresky and Pardon in [GP02, Main Theorem] that under the natural homomorphism

$$H^*({}_K S(G, X)^{\min}, \mathbb{C})_v \rightarrow IH^*({}_K S(G, X)^{\min}, \mathbb{C}). \quad (2.6)$$

the classes in $IH^*({}_K S(G, X)^{\min}, \mathbb{C})_v$ lift canonically, as differential forms, to ordinary singular cohomology $H^*({}_K S(G, X)^{\min}, \mathbb{C})_v$.

We use the notation of the diagram (2.1). Let K be a neat open compact subgroup of $G(\mathbb{A}_f)$, as above. Let $[\mathcal{E}]_K$ be the automorphic vector bundle on ${}_K S(G, X)$ attached to the homogeneous vector bundle $\mathcal{E} \in \text{Vect}_G(\hat{X})$. Let

$$c_n([\mathcal{E}]_K^{\text{can}}) \in H^{2n}({}_K S(G, X)_{\Sigma}, \mathbb{Q}(n)), \quad c_n([\mathcal{E}]_K) \in H^{2n}({}_K S(G, X), \mathbb{Q}(n))$$

be the Chern classes in Betti cohomology. The following theorem summarizes the main results of the article [GP02].

Theorem 2.7. (i) *There are canonical classes $c_n([\mathcal{E}]_K)^{GP} \in H^{2n}({}_K S(G, X)^{\min}, \mathbb{C})$ such that*

$$\varphi_{\Sigma}^*(c_n([\mathcal{E}]_K)^{GP}) = c_n([\mathcal{E}]_K^{\text{can}}) \in H^{2n}({}_K S(G, X)_{\Sigma}, \mathbb{Q}(n)).$$

² $\uparrow$ Automorphic forms are understood to be C^{∞} and K_h -finite, and to satisfy the remaining conditions introduced by Harish-Chandra, hence no completion is needed.

(ii) *In particular*

$$j_K^*(c_n([\mathcal{E}]_K)^{GP}) = c_n([\mathcal{E}]_K) \in H^{2n}({}_K S(G, X), \mathbb{Q}(n)).$$

(iii) *The classes $c_n([\mathcal{E}])^{GP}$ (we drop the subscript K) are represented by square integrable automorphic forms. More precisely, the image of $c_n([\mathcal{E}])^{GP}$ under the morphism of (2.6) is contained in the subspace of intersection cohomology that corresponds, under the isomorphism (2.4), to the relative Lie algebra cohomology of $\oplus \pi$ with trivial archimedean component π_∞ .*

(iv) *Suppose $G^{\text{der}}(\mathbb{R}) = \prod_i G_i$ is a product of simple Lie groups, all of which are of the form $\text{Sp}(2n, \mathbb{R})$, $SU(p, q)$, $SO^*(2n)$, or $SO(p, 2)$ with p odd³. The $\mathbb{Q}$ -subalgebra*

$$H_{\text{Chern}}^{2*}({}_K S(G, X)^{\min}) \subset H^{2*}({}_K S(G, X)^{\min}, \mathbb{C})$$

generated by the $c_n([\mathcal{E}])^{GP}$, as $\mathcal{E}$ varies over $\text{Vect}_G(\hat{X})$ is endowed with a naturally defined surjective homomorphism

$$H_{\text{Chern}}^{2*}({}_K S(G, X)^{\min}) \xrightarrow{h} H^{2*}(\hat{X}, \mathbb{Q}(*))$$

such that for any n the diagram

$$\begin{array}{ccc} H_{\text{Chern}}^{2n}({}_K S(G, X)^{\min}) & \xrightarrow[\text{natural map}]{\quad} & HH^{2n}({}_K S(G, X)^{\min}, \mathbb{Q}(n))_v \\ \downarrow h & & \uparrow \sim \text{Prop. 2.3} \\ H^{2n}(\hat{X}, \mathbb{Q}(n)) & \xrightarrow{\quad} & H^{2n}(\hat{X}, \mathbb{Q}(n)) \end{array}$$

commutes. In other words, the isomorphism of Proposition 2.3 tensor $\mathbb{C}$ factors through

$$H^{2n}({}_K S(G, X)^{\min}, \mathbb{C}) \xrightarrow{\text{natural map}} HH^{2n}({}_K S(G, X)^{\min}, \mathbb{C}).$$

In the next sections, we use Peter Scholze's perfectoid geometry and his Hodge-Tate morphism to prove an analogue of the Goresky-Pardon theorem for ℓ -adic cohomology of Shimura varieties of abelian type.

2.A.a. Even-dimensional quadrics

Theorem 2.7 excludes the case where $G^{\text{der}}(\mathbb{R})$ contains a factor isomorphic to $SO(2k-2, 2)$ with $k > 2$. Assuming G^{der} is $\mathbb{Q}$ -simple, there is then a totally real field F , with $[F : \mathbb{Q}] = d$, say, such that $\hat{X}_{\mathbb{C}}$ is isomorphic to a product Q_n^d of d smooth projective complex quadrics Q_n , each of dimension $n = 2k-2$. The reason for this exclusion is explained in §16.6 of [GP02]. Following §16.5 of [BH58], we consider the cohomology algebra $A := H^*(Q_n, \mathbb{C})$ of a complex quadric $Q_n = \text{SO}(n+2)/\text{SO}(n) \times \text{SO}(2)$ of dimension n , with $C = \mathbb{C}$ or $\mathbb{Q}_{\ell}$. Then A contains a subalgebra A^+ isomorphic to $C[c_1]/(c_1^{n+1})$, with $c_1 \in H^2(Q_n, C)$ given by the Chern class of the line bundle corresponding to the standard representation of $\text{SO}(2)$. (There is a misprint in [GP02]; the total dimension of A^+ as C -vector space is $n+1$, not n .) Moreover there is an isomorphism

$$A = A^+[e]/(e^2 - c_1^n) \tag{2.8}$$

with e the Euler class of the the vector bundle arising from the standard representation of $\text{SO}(n)$. In [BH58], the class c_1 is denoted x_1 and the class e is denoted $\prod_{i=2}^n x_i$; the equation $e^2 = c_1^n$ then follows immediately for formula (6) of [BH58].

More generally, if $\hat{X}_{\mathbb{C}}$ is isomorphic to Q_n^d as above, we denote by $c_{1,r} \in H^2(\hat{X}_{\mathbb{C}}, C(1))$ the class of the line bundle defined above corresponding to the r -th factor of Q_n^d , $r = 1, \dots, d$, and let e_r correspond to

³ ↑ Some of the G_i can also be of the form $SO(2, 2)$, as in the statement of [GP02].

the Euler class in the r -th factor. The isomorphism of Proposition 2.3 is valid in all cases, and Goresky and Pardon showed that the image in $IH^{2j}({}_K S(G, X)^{\min}, \mathbb{Q}(j))_v$ of the classes $c_{1,r}^j \in H^{2j}(\hat{X}_{\mathbb{C}}, C(j))$ lift canonically to the cohomology of the minimal compactification (the twist j here is unnecessary for $C = \mathbb{C}$, we write it for the case $C = \mathbb{Q}_{\ell}$). However, when $n = 2k - 2 > 2$, they were unable to show that the e_r lift. We can extend the statement of Theorem 2.7 with the following definition.

Definition 2.9. (i) Suppose G^{der} is $\mathbb{Q}$ -simple and $G^{\text{der}}(\mathbb{R}) \simeq \text{SO}(2k - 2, 2)^d$ for some integer d , with $k > 2$. Suppose $C = \mathbb{C}, \mathbb{Q}_{\ell}$, or $\mathbb{Q}$. Define

$$\begin{aligned} H^*(\hat{X}, C)^+ &\subset H^*(\hat{X}, C); \\ IH^*({}_K S(G, X)^{\min}, \mathbb{Q}(n))_v^+ &\subset IH^*({}_K S(G, X)^{\min}, \mathbb{Q}(n))_v \end{aligned} \quad (2.10)$$

to be the subalgebras generated respectively by the classes $c_{1,r}$, $r = 1, \dots, d$, and their images under the isomorphism of Proposition 2.3. We similarly define

$$[K_0(\text{Rep}(K_h)) \otimes_{K_0(\text{Rep}(G))} C]^+ \subset K_0(\text{Rep}(K_h)) \otimes_{K_0(\text{Rep}(G))} C \quad (2.11)$$

to be the inverse image of $H^*(\hat{X}, C)^+$ under the isomorphism (1.1).

(ii) If G satisfies condition (iv) of Theorem 2.7, we let

$$\begin{aligned} H^*(\hat{X}, C)^+ &= H^*(\hat{X}, C); \\ IH^*({}_K S(G, X)^{\min}, \mathbb{Q}(n))_v^+ &= IH^*({}_K S(G, X)^{\min}, \mathbb{Q}(n))_v; \\ [K_0(\text{Rep}(K_h)) \otimes_{K_0(\text{Rep}(G))} C]^+ &= K_0(\text{Rep}(K_h)) \otimes_{K_0(\text{Rep}(G))} C. \end{aligned}$$

(iii) In either case, we define

$$\text{Vect}_G(\hat{X})^+ = ch_{\hat{X}}^{-1}(H^*(\hat{X}, C)^+) \subset K_0(\text{Vect}_G(\hat{X})).$$

We continue to use the notation $\mathcal{E}$ to denote (virtual) bundles in $\text{Vect}_G(\hat{X})^+$.

Theorem 2.12. *Suppose G satisfies either (i) or (ii) of Definition 2.9. Then the conclusions of Theorem 2.7 hold with $H^*(\hat{X}, C)$, $\text{Vect}_G(\hat{X})$, and $IH^*({}_K S(G, X)^{\min}, \mathbb{Q}(n))_v$ replaced by the versions with superscript $^+$, and for $\mathcal{E} \in \text{Vect}_G(\hat{X})^+$.*

The theorem, and its application in Proposition 4.16 below, naturally extend to groups G such that G^{der} is a product of groups of type (i) and (ii) in Definition 2.9. We omit the details.

2.B. Perfectoid Shimura varieties and the Hodge-Tate morphism

The results of the present section are entirely due to Scholze [Sch15], then to Caraiani-Scholze [CS17] and Pilloni-Stroh [PS16].

Fix a level subgroup $K \subset G(\mathbf{A}_f)$. Let $G(\mathbf{A}_f) \rightarrow G(\mathbb{Q}_p)$, $k \mapsto k_p$ be the projection. Denote by K_p its projection to $G(\mathbb{Q}_p)$. For $r \geq 0$ we let $K_{p,r} \subset K_p$ be a decreasing family of subgroups of finite index, with $K_{p,r} \supset K_{p,r+1}$ for all r , and such that $\bigcap_r K_{p,r} = \{1\}$. Let $K_r = \{k \in K \mid k_p \in K_{p,r}\}$, and let $K^p = \bigcap_r K_r$. We identify K^p with its projection to the prime-to- p adèles $G(\mathbf{A}_f^p)$; then K^p is an open compact subgroup of $G(\mathbf{A}_f^p)$ called a “tame level subgroup”.

We assume that the Shimura datum (G, X) is of *Hodge type*. Thus, up to replacing K by a subgroup of finite index, ${}_K S(G, X)$ admits an embedding of Shimura varieties in a Siegel modular variety of some level attached to the Shimura datum $(\text{GSp}(2g), X_{2g})$ for some g , where X_{2g} is the union of the Siegel upper and lower half-spaces. We let $\hat{X}_{2g}$ denote the compact dual flag variety of X_{2g} . Let C denote the completion of

an algebraic closure $\overline{\mathbb{Q}_p}$ of $\mathbb{Q}_p$. Denote by $_{K^p}\mathcal{S}(G, X)$, resp. $_{K^p}\mathcal{S}(G, X)^{\min}$ the adic space over C attached to $_{K^p}\mathcal{S}(G, X)$, resp. $_{K^p}\mathcal{S}(G, X)^{\min}$. We assume that K^p is contained in the principal congruence subgroup of level N for some $N \geq 3$, in the sense explained in §4 of [Sch15]. We restate one of the main theorems of Scholze's article [Sch15]. In §4.1 of *loc. cit.* it is shown that there is a level K'_r for $(\mathrm{GSp}(2g), X_{2g})$ such that $K_r = K'_r \cap G(\mathbf{A}_f)$, and such that the scheme theoretic image of $_{K^p}\mathcal{S}(G, X)^{\min}$ in $_{K'_r}\mathcal{S}(\mathrm{GSp}(2g), X_{2g})^{\min}$ does not depend on the choice of K'_r . We denote it by $\overline{_{K^p}\mathcal{S}(G, X)^{\min}}$, and by $\overline{_{K^p}\mathcal{S}(G, X)^{\min}}$ its associated C -adic space.

Theorem 2.13 (Theorem 4.1.1 in [Sch15]). *There is a commutative diagram of morphisms of adic spaces*⁴

$$\begin{array}{ccc} _{K^p}\mathcal{S}(G, X) & \sim & \varprojlim_{\neq} _{K_r}\mathcal{S}(G, X) \\ \downarrow j & & \downarrow \varprojlim_r j_r \\ \overline{_{K^p}\mathcal{S}(G, X)^{\min}} & \sim & \varprojlim_{\neq} \overline{_{K_r}\mathcal{S}(G, X)^{\min}} \end{array} \quad (2.14)$$

Here, on the left side, j is an open embedding of perfectoid Shimura varieties over C , on the right side, $\varprojlim_{\neq} j_r$ denotes the formal inverse system of open embeddings of adic spaces over C , and the notation $\sim$ is defined precisely in [SW13, Definition 2.4.1]. Moreover, there is a $G(\mathbb{Q}_p)$ -equivariant Hodge-Tate morphism

$$\pi_{HT} : \overline{_{K^p}\mathcal{S}(G, X)^{\min}} \rightarrow \hat{\mathcal{X}}_{2g}$$

which is compatible with change of tame level subgroup K^p .

Discussion 2.15. Here we use the notation $\hat{\mathcal{X}}_{2g}$ for the adic space over C attached to the flag variety X_{2g} . In *loc. cit.* the notation $\mathcal{X}_{K^p}^*$ is used for the perfectoid Shimura variety $_{K^p}\mathcal{S}(G, X)^{\min}$. The notation $\sim$ indicates that the right hand side of (2.14) is not to be viewed as a projective limit in the category of adic spaces, which in general does not make sense. Rather, what is meant is that, for each r , there is a commutative diagram

$$\begin{array}{ccc} _{K^p}\mathcal{S}(G, X) & \longrightarrow & _{K_r}\mathcal{S}(G, X) \\ \downarrow j & & \downarrow j_r \\ \overline{_{K^p}\mathcal{S}(G, X)^{\min}} & \longrightarrow & \overline{_{K_r}\mathcal{S}(G, X)^{\min}} \end{array}; \quad (2.16)$$

that these are compatible with the natural maps from level K_{r+1} to level K_r ; and that the objects on the left have the properties indicated in [SW13, Definition 2.4.1].

The target of the Hodge-Tate morphism was clarified in [CS17]. Let $\hat{\mathcal{X}} \subset \hat{\mathcal{X}}_{2g}$ be the closed embedding of C -adic spaces corresponding to the closed embedding $\hat{X} \subset \hat{X}_{2g}$ of the compact duals of X and X_{2g} .

Theorem 2.17 (Theorem 2.1.3 in [CS17]). *The Hodge-Tate morphism π_{HT} in Theorem 2.13 factors through the inclusion $\hat{\mathcal{X}} \subset \hat{\mathcal{X}}_{2g}$, yielding a $G(\mathbb{Q}_p)$ -equivariant Hodge-Tate morphism*

$$\pi_{HT} : \overline{_{K^p}\mathcal{S}(G, X)^{\min}} \rightarrow \hat{\mathcal{X}}.$$

Proof. The existence of π_{HT} is stated for the *open* perfectoid Shimura variety $_{K^p}\mathcal{S}(G, X)$ with values in $\hat{\mathcal{X}}$. Since $\hat{\mathcal{X}}$ is closed in $\hat{\mathcal{X}}_{2g}$, the extension to the boundary then follows from Theorem 2.13 by continuity, as in [C+6, Theorem 3.3.4]. $\square$

⁴ $\uparrow$ The projective limits on the right-hand side are not literally adic spaces; see Discussion 2.15 for the precise meaning of this statement.

Assume for the moment that $(G, X) = (\mathrm{GSp}(2g), X_{2g})$. Fix $K_g \subset \mathrm{GSp}(2g, \mathbf{A}_f)$ a neat compact open subgroup, with $K_p = \mathrm{GSp}(2g, \mathbb{Z}_p)$ and write ${}_{K_g}\mathcal{A}_g$ instead of ${}_{K_g}S(\mathrm{GSp}(2g), X_{2g})$ for the Siegel modular variety of genus g and level K_g , viewed as an adic space over C . Let $K_{g,r} \subset K_g$ be the principal congruence subgroup of K_g of level p^r . Let ${}_{K_g}\mathcal{A}_g^{\mathrm{tor}} = {}_{K_g}\mathcal{A}_{g, \Sigma_g}$ be a smooth projective toroidal compactification of ${}_{K_g}\mathcal{A}_g$ for some combinatorial datum Σ_g as above. Following Pilloni-Stroh in [PS16], §1.3, but with a change of notation, we let ${}_{K_{g,r}}\mathcal{A}_g^{\mathrm{tor}}$ denote the corresponding toroidal compactification of ${}_{K_{g,r}}\mathcal{A}_g$ for each r , with the same Σ_g . The ${}_{K_{g,r}}\mathcal{A}_g^{\mathrm{tor}}$ form a projective system of adic spaces over C . The authors construct a projective system of normal models ${}_{K_{g,r}}\mathcal{A}_{g, \mathcal{O}_C}^{\mathrm{tor}}$ over $\mathrm{Spec}(\mathcal{O}_C)$ for each r , and define a perfectoid space ${}_{K_g^p}\mathcal{A}_g^{\mathrm{tor}}$ which is the generic fiber of the projective limit of the ${}_{K_{g,r}}\mathcal{A}_{g, \mathcal{O}_C}^{\mathrm{tor}}$, in the sense of [SW13, Section 2.2]. See [PS16, Section A.12 and Corollaire A.19] for the statement that the projective limit, which they denote $\mathcal{X}(p^\infty)^{\mathrm{tor-mod}}$, is indeed perfectoid. There is a natural map

$$\eta_g : {}_{K_g^p}\mathcal{A}_g^{\mathrm{tor}} \rightarrow {}_{K_g^p}\mathcal{A}_g^{\mathrm{min}}.$$

Let (G, X) be any Shimura datum of Hodge type, with $\iota : (G, X) \hookrightarrow (\mathrm{GSp}(2g), X_{2g})$ a fixed symplectic embedding. In the remainder of this section we make the following hypothesis, which (as we have seen) is true for the Shimura datum $(\mathrm{GSp}(2g), X_{2g})$:

Hypothesis 2.18. *The projective system of ${}_{K_r}S(G, X)$ has a compatible projective system of toroidal compactifications ${}_{K_r}S(G, X)_\Sigma$ such that, if ${}_{K_r}S(G, X)_\Sigma$ is the associated adic space over C , then there is a perfectoid space ${}_{K^p}S(G, X)_\Sigma$ such that*

$${}_{K^p}S(G, X)_\Sigma \sim \varprojlim_{K_r} {}_{K_r}S(G, X)_\Sigma$$

in the sense of [SW13, Section 2.2].

It is likely the Hypothesis 2.18 is valid, but its proof is probably as elaborate as the construction of integral toroidal compactifications in [Lan13]⁵. Thus we will provide an alternative proof of our main theorems, that does not depend on this hypothesis, in section §3. However, the proof assuming Hypothesis 2.18 is considerably simpler, and it is valid, by [PS16], in the Siegel modular case.

Under Hypothesis 2.18, there is automatically a map

$$q : {}_{K^p}S(G, X)_\Sigma \rightarrow {}_{K^p}S(G, X)^{\mathrm{min}}$$

as well as the maps

$$q_r : {}_{K_r}S(G, X)_\Sigma \rightarrow {}_{K_r}S(G, X)^{\mathrm{min}}$$

at finite level (the existence of which does not depend on the hypothesis).

By [Har85, Section 3.4], there is a correspondence

$$\begin{array}{ccc} {}_K I(G, X) & \xrightarrow{a'} & \hat{X} \\ b' \downarrow & & \\ {}_K S(G, X) & & \end{array} \quad (2.19)$$

where b' is a family of G -torsors, functorial with respect to inclusions $K' \subset K$ and translation by elements $g \in G(\mathbf{A}_f)$, and a' is a G -equivariant morphism. For any G -equivariant vector bundle $\mathcal{E}$ over $\hat{X}$, the pullback $a'^*(\mathcal{E})$ is G -equivariant, and thus descends to the automorphic vector bundle $[\mathcal{E}]$ over ${}_K S(G, X)$. The G -torsor ${}_K I(G, X)$ is constructed as the moduli space $\mathrm{Isom}^\otimes([V_\rho], V)$ of trivializations of a flat automorphic vector bundle $[V_\rho]$ attached to a faithful representation $\rho : G \rightarrow \mathrm{GL}(V)$; the superscript $\otimes$ indicates that the isomorphisms respect absolute Hodge cycles. Thus the construction requires *a priori* the

⁵ ↑ Ana Caraiani has informed us that a proof of Hypothesis 2.18, for the case of a Shimura variety attached to a quasi-split unitary similitude group of even dimension, will appear in her forthcoming joint work with Scholze.

existence of the automorphic vector bundle $[V_\rho]$. When ${}_K S(G, X)$ is of Hodge type, one can take ρ to be a symplectic embedding $\rho : G \rightarrow \mathrm{GSp}(V)$, and define $[V_\rho]$ to be the pullback to ${}_K S(G, X)$ of the (dual of the) relative de Rham H^1 of the universal abelian scheme over the Siegel modular variety attached to $\mathrm{GSp}(V)$. Then the morphism a is defined by transferring the Hodge filtration on $[V_\rho]$ to the constant vector bundle defined by V . That this construction is canonically independent of the choice of symplectic embedding ρ is explained in [Har85, Remark 4.9.1]; see also [CS17, Lemma 2.3.4].

Choose a base point $h \in X \subset \hat{X}$ as in Section 1; we may as well assume h to be a CM point. Recall that P_h denotes the stabilizer of h and K_h its Levi quotient (in [CS17] this group is denoted M_μ). Any faithful representation $\alpha : K_h \rightarrow \mathrm{GL}(W_h)$ defines by pullback to P_h a G -equivariant vector bundle $\mathcal{W}_h$ over $\hat{X}$, and thus an automorphic vector bundle $[\mathcal{W}_h]$ over ${}_K S(G, X)$ that varies functorially in K . We can then define a family (depending on K) of K_h -torsors $b : T = {}_K T_h(G, X) \rightarrow {}_K S(G, X)$ with a K_h -equivariant morphism $a : {}_K T_h(G, X) \rightarrow \hat{X}$ as the moduli space of trivializations of $[\mathcal{W}_h]$ as above. More precisely, letting $R_u P_h$ denote the unipotent radical of P_h , the natural morphism $\hat{X}_h := G/R_u P_h \rightarrow \hat{X}$ is canonically a G -equivariant K_h -torsor, whose pullback $a'^{-1} \hat{X}_h$ descends to a K_h -torsor over ${}_K S(G, X)$, over the reflex field of the CM point h , that is naturally identified with T defined above. Moreover, the construction is canonically independent of the choice of base point. This is also constructed in Section 2.3, especially Lemma 2.3.5, of [CS17].

The pull-back of T via the ringed space morphism $\hat{\mathcal{X}} \rightarrow \hat{X}$ is denoted by $\mathcal{T}$. The K_h -torsor b has a Mumford extension $T^{\mathrm{can}} \rightarrow {}_K S(G, X)_\Sigma$ as a G -torsor. See [HZ94], specifically Lemma 4.4.2 and pages 320–321, where the Mumford extension of the G -torsor b' in (2.19) is constructed; the K_h -torsor is constructed in the same way. One denotes by $\mathcal{M}^{\mathrm{can}} = {}_K \mathcal{M}^{\mathrm{can}}$ the pull-back of T^{can} via the ringed space morphism ${}_K S(G, X)_\Sigma \rightarrow {}_K S(G, X)$.

We define the K_h -torsor

$$\mathcal{M}_p := \pi_{HT}^* \mathcal{T}$$

on ${}_K S(G, X)^{\mathrm{min}}$, which coincides by definition with the K_h -torsor defined in [CS17, Lemma 2.3.8]. We define the K_h -torsor

$$\mathcal{M}_{dR, \Sigma, K} := \pi_{K, K^p}^* ({}_K \mathcal{M}^{\mathrm{can}}) \quad (2.20)$$

on ${}_K S(G, X)_\Sigma$.

Proposition 2.21 (Proposition 2.3.9 in [CS17]). *For any neat level subgroup K , there is a canonical isomorphism*

$$j^* \mathcal{M}_p \xrightarrow{\sim} j_\Sigma^* \mathcal{M}_{dR, \Sigma, K}$$

of K_h -torsors over ${}_K S(G, X)$.

Although the article [CS17] is written for compact Shimura varieties, the argument developed there for this point is valid for any Shimura variety of Hodge type. Strictly speaking, as explained in [CS17], the torsors $\mathcal{M}_p$ and $\mathcal{M}_{dR, \Sigma, K}$ have natural extensions to G -torsors by pullback to torsors for opposite parabolics, followed by pushforward to G , so the comparison only applies to semisimple automorphic vector bundles.

The following theorem is essentially due to Pilloni and Stroth.

Theorem 2.22. *Assume Hypothesis 2.18. Then, for any neat level subgroup K , the isomorphism of Proposition 2.21 extends to a canonical isomorphism*

$$q^* \mathcal{M}_p \xrightarrow{\sim} \mathcal{M}_{dR, \Sigma, K}$$

of K_h -torsors over ${}_K S(G, X)_\Sigma$. As K^p varies, these isomorphisms are equivariant under the action of $G(\mathbf{A}_f^p)$.

In [PS16] this is proved for the Siegel modular variety, although it is not stated in this form. In Section 5 we explain their result and show how to obtain Theorem 2.22 for general Shimura varieties of Hodge type, under Hypothesis 2.18. In the following section, we show how to dispense with Hypothesis 2.18 and prove Theorem 3.12 as an alternative to Theorem 2.22. In Section 4 we construct ℓ -adic Chern classes using either Theorem 2.22 (when it is available) or Theorem 3.12 for general Shimura varieties of Hodge type.

3. Construction of perfectoid covers

Let (G, X) be any Shimura datum of Hodge type, with $\iota : (G, X) \hookrightarrow (\mathrm{GSp}(2g), X_{2g})$ a fixed symplectic embedding. Choose a combinatorial datum Σ for ${}_K S(G, X)$ that is compatible with the fixed Σ_g chosen above, so that there is a morphism

$$[\iota] : {}_K S(G, X)_\Sigma \rightarrow {}_{K_g} \mathcal{A}_{g, \Sigma_g}$$

which factors through

$$\overline{{}_K S(G, X)_\Sigma} = {}_{K_g} \mathcal{A}_{g, \Sigma_g} \times_{{}_{K_g} \mathcal{A}_g^{\min}} \overline{{}_K S(G, X)}^{\min}. \quad (3.1)$$

We denote the corresponding C -adic spaces by ${}_K S(G, X)_\Sigma$ and $\overline{{}_K S(G, X)_\Sigma}$.

If we admitted Hypothesis 2.18 then we would be able to replace $\overline{{}_K S(G, X)_\Sigma}$ by ${}_K S(G, X)_\Sigma$ in what follows; however, we will not assume Hypothesis 2.18 in the remainder of this section. By [Sch12, Prop. 6.18] the fibre product

$$\overline{{}_K S(G, X)_\Sigma} := {}_{K_g^p} \mathcal{A}_g^{\mathrm{tor}} \times_{{}_{K_g^p} \mathcal{A}_g^{\min}} \overline{{}_K S(G, X)}^{\min} \quad (3.2)$$

exists in the category of perfectoid spaces. We denote by

$$\bar{q} : \overline{{}_K S(G, X)_\Sigma} \rightarrow \overline{{}_K S(G, X)}^{\min} \quad (3.3)$$

the projection on the second factor, and by

$$\bar{q}_r : \overline{{}_K S(G, X)_\Sigma} \rightarrow \overline{{}_K S(G, X)}^{\min}_r$$

the composition with the canonical map of $\overline{{}_K S(G, X)}^{\min}$ to $\overline{{}_K S(G, X)}^{\min}_r$. By definition, in the sense of [SW13, Definition 2.4.1], one has

$${}_{K_g^p} \mathcal{A}_g^{\mathrm{tor}} \sim \varprojlim_r {}_{K_{g,r}} \mathcal{A}_g^{\mathrm{tor}}, \quad {}_{K_g^p} \mathcal{A}_g^{\min} \sim \varprojlim_r {}_{K_{g,r}} \mathcal{A}_g^{\min}, \quad \overline{{}_K S(G, X)}^{\min} \sim \varprojlim_r \overline{{}_K S(G, X)}^{\min}_r.$$

One deduces

$$\overline{{}_K S(G, X)_\Sigma} \sim \varprojlim_r \overline{{}_K S(G, X)_\Sigma}_r. \quad (3.4)$$

In particular, by [Sch12, Corollary 7.1.8], one has the relation

$$H^*(\overline{{}_K S(G, X)_\Sigma}, \mathbb{Z}/\ell^n) = \varprojlim_r H^*(\overline{{}_K S(G, X)_\Sigma}_r, \mathbb{Z}/\ell^n). \quad (3.5)$$

On the other hand, Scholze's Hodge-Tate map yields the composite map

$$\overline{{}_K S(G, X)_\Sigma} \rightarrow {}_{K_g^p} \mathcal{A}_g^{\min} \rightarrow \hat{\mathcal{X}}_{2g}$$

which by definition is the same as the composite map

$$\overline{{}_{K^p}\mathcal{S}(G, X)_\Sigma} \rightarrow \overline{{}_{K^p}\mathcal{S}(G, X)}^{\min} \xrightarrow{\pi_{HT}} \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}_{2g}.$$

This defines a map

$$\overline{{}_{K^p}\mathcal{S}(G, X)_\Sigma} \rightarrow \hat{\mathcal{X}} \quad (3.6)$$

in the category of C -adic spaces. It is $G(\mathbb{Q}_p)$ -equivariant.

In what follows the idea to use the space denoted $\mathbb{P}^{N, \text{perf}}$ to define a perfectoid space above ${}_{K_s}\mathcal{S}(G, X)_\Sigma$ is due to Bhargav Bhatt. We fix a natural number s . Associated to it we have the normalization morphism $\nu_s : {}_{K_s}\mathcal{S}(G, X)_\Sigma \rightarrow \overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma}$. It is a finite morphism, thus it factors as

$$\nu_s : {}_{K_s}\mathcal{S}(G, X)_\Sigma \xrightarrow{\eta_s} \overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^N \xrightarrow{\text{projection}} \overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma}$$

for some natural number N , where η_s is a closed embedding. We set

$$Y_s = \text{Image}(\eta_s).$$

By definition η_s induces an isomorphism between ${}_{K_s}\mathcal{S}(G, X)_\Sigma$ and Y_s . For any $\rho \geq 0$ we also define

$$Y_{s+\rho} = Y_s \times_{\overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma}} \overline{{}_{K_{s+\rho}}\mathcal{S}(G, X)_\Sigma}. \quad (3.7)$$

We fix coordinates $(x_0 : \dots : x_N)$ of $\mathbb{P}^N$ and define for $\rho \in \mathbb{N}$ the finite flat morphisms $\phi_\rho : \mathbb{P}^N \rightarrow \mathbb{P}^N$ by $\phi_\rho(x_0 : \dots : x_N) = (x_0^{p^\rho} : \dots : x_N^{p^\rho})$. Over C , a perfectoid space $\mathbb{P}^{N, \text{perf}}$ is associated to $(\mathbb{P}^N, \phi_\rho)$ [Sch14, Section 17], such that

$$\mathbb{P}^{N, \text{perf}} \sim \varprojlim_{\rho} \phi_\rho.$$

Then the fiber product

$${}_{K^p}\mathcal{S}(G, X)_{\Sigma, N} := \overline{{}_{K^p}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^{N, \text{perf}}$$

exists as a perfectoid space, by [Sch12, Proposition 6.18], and we have

$${}_{K^p}\mathcal{S}(G, X)_{\Sigma, N} \sim \varprojlim_{\rho} \overline{{}_{K_{s+\rho}}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^N, \quad (3.8)$$

where the map

$$\overline{{}_{K_{s+\rho}}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^N \rightarrow \overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^N$$

is given by the natural projection in the first factor and ϕ_ρ in the second factor.

Proposition 3.9. *There is a closed perfectoid subspace*

$$Z_s \subset {}_{K^p}\mathcal{S}(G, X)_{\Sigma, N}$$

such that

$$Z_s \sim \varprojlim_{\rho \geq 0} Y_{s+\rho}.$$

Proof. We define $Z_s \subset {}_{K^p}\mathcal{S}(G, X)_{\Sigma, N}$ by the pull-back of the ideal of Y_s in $\overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^N$ via (3.8)

$${}_{K^p}\mathcal{S}(G, X)_{\Sigma, N} \rightarrow \overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma} \times_C \mathbb{P}^N.$$

It follows from [Bha17, Proposition 9.4.1] that Z_s is perfectoid. $\square$

By construction, Z_s maps to $\overline{{}_{K^p}\mathcal{S}(G, X)_\Sigma}$ which itself maps to $\overline{{}_{K^p}\mathcal{S}(G, X)}^{\min}$, and to ${}_{K_s}\mathcal{S}(G, X)_\Sigma$, but not necessarily to ${}_{K_r}\mathcal{S}(G, X)_\Sigma$ for $r > s$. On the other hand, ${}_{K_s}\mathcal{S}(G, X)_\Sigma$ naturally maps to ${}_{K_s}\mathcal{S}(G, X)^{\min}$, and thus we have a composite diagram

$$Z_s \xrightarrow{\phi} {}_{K_s}\mathcal{S}(G, X)_\Sigma \xrightarrow{q_s} {}_{K_s}\mathcal{S}(G, X)^{\min} \xrightarrow{u_s} \overline{{}_{K_s}\mathcal{S}(G, X)}^{\min} \quad (3.10)$$

We let Q denote the composite of the three maps, and let $\underline{q}_s = u_s \circ q_s$ denote the composite of the last two maps.

Proposition 3.11. *The perfectoid space Z_s has the following properties.*

(i) *There are morphisms*

$$\begin{aligned} Q : Z_s &\xrightarrow{\psi} \overline{{}_{K^p}\mathcal{S}(G, X)_\Sigma} \xrightarrow{\bar{q}} \overline{{}_{K^p}\mathcal{S}(G, X)}^{\min} \\ \phi : Z_s &\rightarrow {}_{K_s}\mathcal{S}(G, X)_\Sigma \end{aligned}$$

such that the following diagram commutes

$$\begin{array}{ccc} Z_s & \xrightarrow{\psi} & \overline{{}_{K^p}\mathcal{S}(G, X)_\Sigma} \\ \phi \downarrow & \searrow Q_s & \downarrow \bar{q}_s \\ Y_s = {}_{K_s}\mathcal{S}(G, X)_\Sigma & \xrightarrow{q_s} & \overline{{}_{K_s}\mathcal{S}(G, X)}^{\min} \end{array}$$

(ii) *The homomorphism on cohomology*

$$\phi^* : H^\bullet(Y_s, \mathbb{F}_\ell) \rightarrow H^\bullet(Z_s, \mathbb{F}_\ell),$$

induced from (i), is injective, where ℓ is a prime different from p . The injectivity holds also true with $\mathbb{Q}_\ell$ -coefficients when $s = 0$.

Proof. In what follows we assume G^{ad} contains no $\mathbb{Q}$ -simple factor isomorphic to $\mathrm{PGL}(2)$ – in other words, that $S(G, X)$ contains no factor isomorphic to the elliptic modular curve – since in that case the minimal and toroidal compactifications coincide and there is nothing to prove. The point (i) is just a restatement of the above construction. We prove (ii). By Proposition 3.9 together with [Sch12, Corollary 7.18], we have

$$H^\bullet(Z_s, \mathbb{F}_\ell) = \varinjlim_{\rho \geq 0} H^\bullet(Y_{s+\rho}, \mathbb{F}_\ell).$$

On the other hand, the proper maps $Y_{s+\rho} \rightarrow Y_{s+\rho-1}$ for $\rho \geq 1$ are finite, of degree a p -power for $s \geq 1$. Indeed, the finiteness of this map is reduced by Definition (3.7) to the corresponding assertion for the maps

$$\overline{{}_{K_{s+\rho}}\mathcal{S}(G, X)_\Sigma} \rightarrow \overline{{}_{K_s}\mathcal{S}(G, X)_\Sigma}.$$

This in turn is reduced by (3.1) to the well known fact that the corresponding morphisms for change of level in the toroidal compactification ${}_{K_g}\mathcal{A}_{g, \Sigma_g}$ and the compactifications $\overline{{}_{K}\mathcal{S}(G, X)}^{\min}$ are finite. Indeed, let ω_g denote the determinant of the relative cotangent bundle of the universal abelian scheme over ${}_{K_g}\mathcal{A}_g$; in other words, ω_g is the line bundle whose global sections are Siegel modular forms of weight 1 and level K_g . Then it is well known (from the theory of the minimal compactification, and because we have excluded the case of modular curves) that ω_g extends to an ample line bundle over ${}_{K_g}\mathcal{A}_g^{\min}$, and that the pullback of this extension to ${}_{K}\mathcal{S}(G, X)^{\min}$ is ample as well. Thus – although we have been told there is a problem with the

injectivity claim in Corollary 10.2.3 of [SW17] – the map from ${}_K\mathcal{S}(G, X)^{\min}$ to its image in ${}_{K_s}\mathcal{A}_s^{\min}$ is finite, and the morphisms from ${}_K\mathcal{S}(G, X)^{\min}$ to $\overline{{}_K\mathcal{S}(G, X)}^{\min}$ are finite, and from the commutative square

$$\begin{array}{ccc} {}_K\mathcal{S}(G, X)_s^{\min} & \xrightarrow{\text{finite}} & \overline{{}_K\mathcal{S}(G, X)}_s^{\min} \\ \text{finite} \downarrow & & \downarrow \\ {}_K\mathcal{S}(G, X)_r^{\min} & \xrightarrow{\text{finite}} & \overline{{}_K\mathcal{S}(G, X)}_r^{\min} \end{array}$$

for $s \geq r$ we conclude that the level-changing maps $\overline{{}_K\mathcal{S}(G, X)}_s^{\min} \rightarrow \overline{{}_K\mathcal{S}(G, X)}_r^{\min}$ are finite.

This implies (ii) for $s \geq 1$ – and also for $s = 0$, after replacing $\mathbb{F}_\ell$ by $\mathbb{Q}_\ell$ in case the order of the group K_0/K_1 is divisible by ℓ . $\square$

The analogue of Theorem 2.22 is

Theorem 3.12. *For any neat level subgroup K and any $s \geq 0$, the isomorphism of Proposition 2.21 extends to a canonical isomorphism*

$$Q^*\mathcal{M}_p \xrightarrow{\sim} \phi^*\mathcal{M}_{dR, \Sigma, K_s}$$

of K_h -torsors over Z_s . As K^p varies, these isomorphisms are equivariant under the action of $G(\mathbf{A}_f^p)$.

The proof is given in Section 5.

4. Construction of ℓ -adic Chern classes

As in the previous sections, all level subgroups K are neat and all toroidal compactifications are smooth and projective. The constructions in the present section depend in an essential way on Theorem 3.12 – or on Theorem 2.22, if we assume Hypothesis 2.18. Proofs are given without assuming Hypothesis 2.18; to pass from the situation with Hypothesis 2.18, we just formally set $Q = q$.

4.A. Construction using π_{HT}

As in [CS17, Theorem 2.1.3 (2)], the existence of the K_h -torsor $\mathcal{M}_p$ over $\overline{{}_K\mathcal{S}(G, X)}^{\min}$ implies that there is a functor

$$\begin{array}{ccc} \text{Vect}_G^{\text{ss}}(\hat{X})[\overset{r}{\simeq} \text{Rep}_{K_h}] & \rightarrow & \{\text{Vector bundles over } \overline{{}_K\mathcal{S}(G, X)}^{\min}\} \\ \mathcal{E} & \mapsto & [\mathcal{E}]_p. \end{array} \quad (4.1)$$

Similarly, the existence of the K_h torsor $\mathcal{M}_{dR, \Sigma, K}$ over ${}_K\mathcal{S}(G, X)_\Sigma$ gives rise to a functor

$$\begin{array}{ccc} \text{Vect}_G^{\text{ss}}(\hat{X})[\overset{r}{\simeq} \text{Rep}_{K_h}] & \rightarrow & \{\text{Vector bundles over } {}_K\mathcal{S}(G, X)_\Sigma\} \\ \mathcal{E} & \mapsto & [\mathcal{E}^{\text{can}}]. \end{array} \quad (4.2)$$

With the notation of Theorem 3.12, we define

$$[\mathcal{E}]_{p, \Sigma, K_s} = Q^*[\mathcal{E}]_p, \quad [\mathcal{E}]_{dR, \Sigma, K_s} = \phi^*[\mathcal{E}^{\text{can}}]. \quad (4.3)$$

Let $Z_s^o = {}_K\mathcal{S}(G, X) \times_{\overline{{}_K\mathcal{S}(G, X)}^{\min}} Z_s$, $j_s : Z_s^o \hookrightarrow Z_s$ the canonical map. The isomorphism of torsors in Proposition 2.21 gives rise to a canonical isomorphism of tensor functors over Z_s^o

$$(\mathcal{E} \mapsto j_s^*([\mathcal{E}]_{p, \Sigma, K_s})) \xrightarrow{\sim} (\mathcal{E} \mapsto j_s^*([\mathcal{E}]_{dR, \Sigma, K_s})). \quad (4.4)$$

From Theorem 3.12 one immediately obtains the following corollary.

Corollary 4.5. *There is a canonical isomorphism of tensor functors*

$$(\mathcal{E} \mapsto [\mathcal{E}]_{p,\Sigma,K_s}) \xrightarrow{\sim} (\mathcal{E} \mapsto [\mathcal{E}]_{dR,\Sigma,K_s})$$

over Z_s , lifting the isomorphism (4.4) over Z_s° .

We return to this theme in the next paragraph. Recall briefly how the Hecke algebra acts on automorphic bundles on ${}_K S(G, X)$. See [EH17, Section 1.2]. Fix $K \subset G(\mathbf{A}_f)$. Let $h \in G(\mathbf{A}_f)$ and consider $K_h = K \cap hKh^{-1} \subset K$. Right multiplication by h defines an isomorphism $r_h : {}_{hKh^{-1}} S(G, X) \xrightarrow{\sim} {}_K S(G, X)$. One defines

$$T(h)[\mathcal{E}]_K = r_{h*} \circ \pi_{hKh^{-1}, K_h^*} \circ \pi_{K, K_h}^* [\mathcal{E}]_K. \quad (4.6)$$

Projection formula implies

$$T(h)[\mathcal{E}]_K = [\mathcal{E}]_K \otimes \pi_{K, K_h^*} \mathcal{O}_{K_h S(G, X)}.$$

As one has pull-backs and push-downs for $K_{0, \mathbb{Q}}$ and for ℓ -adic cohomology, one has on them an action of the Hecke algebra $\mathcal{H}_K$, spanned by the $T(h)$ for $h \in G(\mathbf{A}_f)$, with h trivial at p and at ramified places for K . In particular, formula (4.6) implies the formula

$$\text{class of } T(h)[\mathcal{E}]_K = \text{class of } [\mathcal{E}]_K^{[K:K_h]} \text{ in } K_0({}_K S(G, X))_{\mathbb{Q}}. \quad (4.7)$$

Recall that the volume character of the Hecke algebra spanned by the $T(h)$ with values in a ring R , is the character sending $T(h)$ to $[K : K_h]$ viewed as an element in R [EH17, Lemma 1.15]. Thus (4.7) is saying that classes of automorphic bundles in $K_0({}_K S(G, X))_{\mathbb{Q}}$ are eigenvectors for the volume character under the action of $\mathcal{H}_K$.

Let $\ell \neq p$ be a prime number.

Lemma 4.8. *The isomorphism r_h induces an isomorphism*

$$r_h^{\min} : {}_K S(G, X)^{\min} \xrightarrow{\sim} {}_{hKh^{-1}} S(G, X)^{\min}.$$

The finite morphism $\pi_{K, K_h} : {}_{K_h} S(G, X) \rightarrow {}_K S(G, X)$ extends to a finite morphism

$$\pi_{K, K_h}^{\min} : {}_{K_h} S(G, X)^{\min} \rightarrow {}_K S(G, X)^{\min}$$

and the action of $\mathcal{H}_K$ on $H^{2n}({}_K S(G, X), \mathbb{Z}_\ell(n))$ extends to an action on $H^{2n}({}_K S(G, X)^{\min}, \mathbb{Z}_\ell(n))$.

Proof. When $G = \text{GL}(2)_{\mathbb{Q}}$ the Shimura varieties are finite unions of modular curves, and the result in that case is standard. We may thus assume G^{ad} contains no factor isomorphic to $\text{PGL}(2)_{\mathbb{Q}}$. Then we know by Theorem 10.14 of [BB66] that $H^0({}_K S(G, X), \omega^n)$ is finite dimensional for any $n \geq 0$, where ω is the dualizing sheaf. The isomorphism r_h induces an algebra isomorphism

$$\bigoplus_{n \in \mathbb{N}} H^0({}_K S(G, X), \omega^n) \xrightarrow{\sim} \bigoplus_{n \in \mathbb{N}} H^0({}_{hKh^{-1}} S(G, X), \omega^n).$$

Similarly the finite cover π_{K, K_h} induces an injective algebra morphism

$$\bigoplus_{n \in \mathbb{N}} H^0({}_K S(G, X), \omega^n) \hookrightarrow \bigoplus_{n \in \mathbb{N}} H^0({}_{K_h} S(G, X), \omega^n).$$

Since ${}_K S(G, X)^{\min}$ (resp. ${}_{K_h} S(G, X)$) is Proj of the left-hand side (resp. right-hand side) of the last diagram [BB66, Theorem 10.11], the induced map on the Proj defines the extensions $r_h^{\min}$ and $\pi_{K, K_h}^{\min}$. On the other hand, pull-back on cohomology is defined, while the trace map $\text{Tr} : \pi_{K, K_h^*} \mathbb{Z}_\ell \rightarrow \mathbb{Z}_\ell$ extends to $j_{K*} \pi_{K, K_h^*} \mathbb{Z}_\ell = \pi_{K, K_h^*}^{\min} j_{K_h^*} \mathbb{Z}_\ell = \pi_{K, K_h^*}^{\min} \mathbb{Z}_\ell \rightarrow j_{K*} \mathbb{Z}_\ell = \mathbb{Z}_\ell$. This proves the second part. $\square$

Let $E(G, X) =: E$ be the reflex field. Fix an embedding $E \hookrightarrow \bar{\mathbb{Q}}$. We choose a place w of E dividing the prime p which we assume to be different from ℓ . We extend $E \hookrightarrow \bar{\mathbb{Q}}$ to the completion $E_w \hookrightarrow \bar{\mathbb{Q}}_p$. We denote by G_{E_w}, G_E the Galois groups of E_w, E . We add a subscript on the lower right to indicate the field over which we regard the varieties considered. We denote by $H^i({}_K S(G, X)_{\bar{\mathbb{Q}}}^{\min}, \mathbb{Q}_\ell)_v = H^i({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell)_v$ the eigenspace under the volume character of the action of $\mathcal{H}_K$.

Through the end of this section, we set $K_0 = K$ for the neat subgroup $K \subset G(\mathbf{A}_f)$ and use the notation $H^i(\Lambda(j))$ for étale cohomology with coefficients Λ , which are either $\mathbb{Z}_\ell$ or $\mathbb{Q}_\ell$.

Theorem 4.9. *Let (G, X) be a Shimura datum of Hodge type. For any prime $\ell \neq p$, any $\mathcal{E} \in \text{Vect}_G^{\text{ss}}(\hat{X}_E)$, we define ℓ -adic Chern classes*

$$c_{n,\ell}([\mathcal{E}]_p)_{K_r} \in H^{2n}({}_{K_r} S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \Lambda(n))_v^{G_{E_w}}, \quad n \in \mathbb{N}$$

where $\Lambda = \mathbb{Q}_\ell$ for $r = 0$ and $\mathbb{Z}_\ell$ for $r \geq 1$, with the following properties:

- (1) $j_{K_r}^*(c_{n,\ell}([\mathcal{E}]_p)_{K_r}) = c_{n,\ell}([\mathcal{E}]_{K_r}) \in H^{2n}({}_{K_r} S(G, X)_{\bar{\mathbb{Q}}}^{\min}, \Lambda(n))$, where the right hand side is the ℓ -adic Chern class of $[\mathcal{E}]$ on ${}_K S(G, X)$.
- (2) If $K' \subset K$ then $\pi_{K_r, K'_r}^{\min*}(c_{n,\ell}([\mathcal{E}]_p)_{K_r}) = c_{n,\ell}([\mathcal{E}]_p)_{K'_r}$.
- (3) $r_h^{\min*}(c_{n,\ell}([\mathcal{E}]_p)_{K_r}) = c_{n,\ell}([\mathcal{E}]_p)_{hK_r h^{-1}}$ for $h \in G(\mathbf{A}_f)$ with trivial component at p and at places ramified for K .
- (4) Whitney product formula: If $0 \rightarrow \mathcal{E}_1 \rightarrow \mathcal{E} \rightarrow \mathcal{E}_2 \rightarrow 0$ is an exact sequence in $\text{Vect}_G^{\text{ss}}(\hat{X}_E)$, then

$$c_{n,\ell}([\mathcal{E}]_p)_{K_r} = \oplus_{a+b=n} c_{a,\ell}([\mathcal{E}_1]_p)_{K_r} \cdot c_{b,\ell}([\mathcal{E}_2]_p)_{K_r}.$$

- (5) If $(G', X') \rightarrow (G, X)$ is a morphism of Shimura data of Hodge type, $K' \rightarrow K$ is a compatible level, with K_r and K'_r compatibly chosen, and

$$f : {}_{K'_r} S(G', X')^{\min} \rightarrow {}_{K_r} S(G, X)^{\min}$$

the corresponding morphism of Shimura varieties, then

$$f^*(c_{n,\ell}([\mathcal{E}]_p)_{K_r}) = (c_{n,\ell}([f^* \mathcal{E}]_p)_{K'_r}) \in H^{2n}({}_{K'_r} S(G', X')_{\bar{\mathbb{Q}}_p}^{\min}, \Lambda(n))_v^{G_{E_w}}.$$

Proof. We first construct $c_{n,\ell}([\mathcal{E}]_p)_{K_r}$ without the v invariance. The vector bundle $[\mathcal{E}]_p$ from (4.1) has Chern classes

$$\bar{c}_{n,\ell}([\mathcal{E}]_p) \in \varprojlim_m H^{2n}(\overline{{}_{K^p} S(G, X)}^{\min}, \mathbb{Z}/\ell^m(n)), \quad n \in \mathbb{N}$$

by Proposition A.1. On the other hand, $[\mathcal{E}]_p$ is defined as $\pi_{HT}^* \mathcal{E}$, where $\mathcal{E}$ is viewed as a vector bundle on the C -adic space $\hat{\mathcal{X}}$. But $\mathcal{E}$ is already defined on $\hat{X}_E$. In particular, with $K_{p,r}$ defined as in Section 2.B, the classes $\bar{c}_{n,\ell}([\mathcal{E}]_p)$ are invariant under the action of $K_{p,r} \subset G(\mathbb{Q}_p)$ for all r and by G_{E_w} . By Proposition A.4 it follows that the classes $\bar{c}_{n,\ell}([\mathcal{E}]_p)$ uniquely define classes

$$c_{n,\ell}([\mathcal{E}]_p) \in H^{2n}(\overline{{}_{K_r} S(G, X)}_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Z}_\ell(n))^{G_{E_w}}$$

for $r \geq 1$. Since K/K_1 is finite, we can descend to $\mathbb{Q}_\ell$ -cohomology at level K . We thus have descended classes

$$\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_r} \in H^{2n}(\overline{{}_K S(G, X)}_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(n))^{G_{E_w}}.$$

Letting $u_r : {}_{K_r}S(G, X)^{\min} \rightarrow \overline{{}_{K_r}S(G, X)^{\min}}$ denote the natural map, as in (3.10), the classes

$$c_{n,\ell}([\mathcal{E}]_p)_{K_r} := u_r^*(\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_r})$$

are the desired ones.

Property (1) is then a direct consequence of Proposition 2.21. Property (2) follows directly from the construction using descent via Proposition A.4. As for Property (3), using descent again, it is enough to see that $r_h^{\min}$ extends to an isomorphism

$$r_h^{\text{perf},\min} : {}_{K_r}S(G, X)^{\min} \rightarrow {}_{hK_r h^{-1}}S(G, X)^{\min}$$

of perfectoid spaces, which respects $[\mathcal{E}]_p$. (See [Sch15, Theorem 4.1.1(iv)] for the Siegel modular case; the general case is identical.) Our notation $[\mathcal{E}]_p$ does not refer to K_p . One should replace the notation $[\mathcal{E}]_p$ with $[\mathcal{E}]_p^{K_p}$. Then the compatibility means $r_h^{\text{perf},\min*}[\mathcal{E}]_p^{hK_p h^{-1}} = [\mathcal{E}]_p^{K_p}$, which follows from Theorem 2.13 together with the addendum Theorem 2.17. Property (4) follows directly from the Whitney product formula for the Chern classes of $[\mathcal{E}]_p$ and Property (5) of the functoriality of the construction of ${}_{K^p}S(G, X)^{\min}$ ([Sch15, *loc.cit.*, comes from the construction, even if this is not explicitly mentioned). It remains to see that $c_{n,\ell}([\mathcal{E}]_p)_{K_r} \in H^{2n}({}_{K_r}S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \Lambda(n))_v$. By construction, this follows from

$$\bar{c}_{n,\ell}([\mathcal{E}]_p) \in \varprojlim_m H^{2n}(\overline{{}_{K^p}S(G, X)^{\min}}, \mathbb{Z}/\ell^m(n))_v, \quad n \in \mathbb{N},$$

which again follows by the projection formula from the trivial relation

$$T(g)[\mathcal{E}]_p = [\mathcal{E}]_p \otimes_{\mathcal{O}_{{}_{K^p}S(G, X)^{\min}}} \mathcal{O}_{{}_{K_h^p}S(G, X)^{\min}}.$$

□

Notation 4.10. The functor

$$\begin{cases} \text{Rep}_{\bar{\mathbb{Q}}}(K_h) & \rightarrow \{\text{Vector Bundles on } {}_{K^p}S(G, X)^{\min}\} \\ \mathcal{E} & \mapsto [\mathcal{E}]_p \end{cases},$$

is a tensor functor, which sends $\text{Rep}_{\bar{\mathbb{Q}}}(G)$ to trivial bundles. It induces the Chern character

$$K_0(\text{Rep}_{\bar{\mathbb{Q}}}(K_h) \otimes_{\text{Rep}_{\bar{\mathbb{Q}}}(G)} \mathbb{Q}) \xrightarrow{ch_\ell} H^{2*}({}_{K^p}S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(*))_v^{G(E_w)} \subset \left(\varprojlim_m H^{2*}({}_{K^p}S(G, X)^{\min}, \mathbb{Z}/\ell^m(*)) \right) \otimes \mathbb{Q}.$$

4.B. The image of $H^*({}_{K^p}S(G, X)^{\min})$ in $H^*({}_{K^p}S(G, X)_\Sigma)$

We now return to the topological setting. In this section, ${}_{K^p}S(G, X)^{\min}$ and ${}_{K^p}S(G, X)_\Sigma$ are identified with the analytic spaces underlying their $\mathbb{C}$ -valued points. We fix Σ , use the notation (2.1) and let $\varphi := \varphi_\Sigma$ denote the desingularization map $\varphi_\Sigma : {}_{K^p}S(G, X)_\Sigma \rightarrow {}_{K^p}S(G, X)^{\min}$. It is a projective morphism. Recall from the general theory [dCM05, Theorem 2.8.1] that the choice of a polarization $\mathcal{L}$ for φ induces a factorization

$$\begin{array}{ccc} H^i({}_{K^p}S(G, X)^{\min}, \mathbb{Q}) & \xrightarrow{\text{natural map}} & IH^i({}_{K^p}S(G, X)^{\min}, \mathbb{Q}) \\ & \searrow \varphi^* & \downarrow \varphi_{\mathcal{L}}^* \\ & & H^i({}_{K^p}S(G, X)_\Sigma, \mathbb{Q}) \end{array} \quad (4.11)$$

in which all morphisms are compatible with the polarized Hodge structure, and $\varphi_{\mathcal{L}}^*$ is injective (see e.g. [dCM05, Corollary 2.8.2]). For a prime $\ell \neq p$ and any $s \geq 0$, the diagram (4.11) has an ℓ -adic version

$$\begin{array}{ccc} H^i(K_s S(G, X)^{\min}, \mathbb{Q}_{\ell}(j)) & \xrightarrow{\text{natural map}} & IH^i(K_s S(G, X)^{\min}, \mathbb{Q}_{\ell}(j)) \\ & \searrow \varphi^* & \downarrow \varphi_{\mathcal{L}}^* \\ & & H^i(K_s S(G, X)_{\Sigma}, \mathbb{Q}_{\ell}(j)) \end{array} \quad (4.12)$$

Thus by the comparison isomorphism [BBD82, Section 6], we conclude that $\varphi_{\mathcal{L}}^*$ in the diagram (4.12) is injective as well. The image of $\varphi_{\mathcal{L}}^*$ might depend on the choice of $\mathcal{L}$, see [dCM05, Example 2.9]. Furthermore, $IH^i(K S(G, X)^{\min}, \mathbb{Q})$, thus

$$\varphi_{\mathcal{L}}^*(IH^i(K S(G, X)^{\min}, \mathbb{Q})) \subset H^i(K S(G, X)_{\Sigma}, \mathbb{Q}),$$

is pure of weight i , and $\varphi^*(H^i(K S(G, X)^{\min}, \mathbb{Q})) \subset H^i(K S(G, X)_{\Sigma}, \mathbb{Q})$ is the maximal pure weight i quotient of $H^i(K S(G, X)^{\min}, \mathbb{Q})$ by [Del74, Proposition 8.2.5].

Theorem 4.13. *For any $\mathcal{E} \in \text{Vect}_G^{\text{ss}}(\hat{X})$, any neat level subgroup $K \subset G(\mathbf{A}_f)$ and all $s \geq 0$, we have*

$$\varphi^*((c_{n,\ell}([\mathcal{E}]_p)_{K_s}) = c_{n,\ell}([\mathcal{E}]^{\text{can}})_{K_s}) \in H^{2n}(K_s S(G, X)_{\Sigma, \bar{\mathbb{Q}}}, \Lambda(n))^{G_E}.$$

Here $\Lambda = \mathbb{Q}_{\ell}$ for $s = 0$ and $\mathbb{Z}_{\ell}$ for $s \geq 1$, as in Theorem 4.9.

Proof. We place ourselves in the situation of Theorem 3.12 which we use; in particular we fix the natural number s . The proof assuming Hypothesis 2.18 is the same without fixing s . It suffices to check the claims with $c_{n,\ell}([\mathcal{E}]_p)_{K_s}$ replaced by $\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_s}$.

Notation is as in Section 3. For $s \geq 0$ we have the diagram with commutative squares

$$\begin{array}{ccccc} & & Q & & \\ & \nearrow & & \searrow & \\ Z_s & \xrightarrow{\psi} & \overline{K^p S(G, X)_{\Sigma}} & \xrightarrow{\bar{q}} & \overline{K^p S(G, X)^{\min}} \\ \downarrow \phi & & \downarrow \pi_{K_s, K^p} & & \downarrow \pi_{K_s, K^p}^{\min} \\ K_s S(G, X)_{\Sigma} & \xrightarrow{\gamma} & \overline{K_s S(G, X)_{\Sigma}} & \xrightarrow{\delta} & \overline{K_s S(G, X)^{\min}} \end{array} \quad (4.14)$$

Here $\delta \circ \gamma = \varphi$ and $\delta, \bar{q}$ come from the definitions (3.1) and (3.2).

By Theorem 3.12 one has

$$\begin{aligned} (\pi_{K_s, K^p}^{\min} \circ Q)^*(\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_s}) &= \phi^*(\delta \circ \gamma)^*(\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_s}) \\ &= Q^*(\bar{c}_{n,\ell}([\mathcal{E}]_p)) = \phi^*(\bar{c}_{n,\ell}([\mathcal{E}]^{\text{can}})) \\ &\in Q^*H^{2n}(K^p S(G, X)^{\min}, \Lambda(n)) \subset H^{2n}(Z_s, \Lambda(n)). \end{aligned}$$

Since (4.14) commutes, and since ϕ^* is injective by Proposition 3.11, we thus have

$$c_{n,\ell}([\mathcal{E}]^{\text{can}}) = (\delta \circ \gamma)^*(\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_s}) = \varphi^*((\bar{c}_{n,\ell}([\mathcal{E}]_p)_{K_s})), \quad (4.15)$$

as classes in $H^{2n}(K_s S(G, X)_{\Sigma, \bar{\mathbb{Q}}}, \Lambda(n))$. On the other hand,

$$H^{2n}(K_s S(G, X)_{\Sigma, \bar{\mathbb{Q}}}, \Lambda(n)) = H^{2n}(K_s S(G, X)_{\Sigma, \bar{\mathbb{Q}}}, \Lambda(n))$$

by the comparison isomorphism [Hub98, Theorem 4.2]. In addition, the automorphic bundles $[\mathcal{E}]^{\text{can}}$ are defined over E . Thus the classes in (4.15) lie in $H^{2n}(K_s S(G, X)_{\Sigma, \bar{\mathbb{Q}}}, \Lambda(n))^{G_E}$. This finishes the proof. $\square$

Let

$$H_{\text{Chern}}^{2*} \left({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(*) \right) \subset H^{2*} \left({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(*) \right)$$

denote the $\mathbb{Q}_\ell$ -subalgebra generated by the images of $c_{*,\ell}([\mathcal{E}]_p)_K$, and

$$H_{\text{Chern}}^{2*} ({}_K S(G, X)_{\Sigma, \bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*)) \subset H^{2*} ({}_K S(G, X)_{\Sigma, \bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))$$

denote the $\mathbb{Q}_\ell$ -subalgebra generated by the images of $c_{*,\ell}([\mathcal{E}]_K^{\text{can}})_K$. Here, when G satisfies (i) of Definition 2.9, we take $\mathcal{E}$ to belong to $\text{Vect}_G(\hat{X})^+$.

Proposition 4.16. *The commutative diagram (4.12) restricts to the commutative diagram of $\mathbb{Q}_\ell$ -algebras*

$$\begin{array}{ccc}
 & & \left[K_0(\text{Rep}_{\bar{\mathbb{Q}}}(\bar{K}_h)) \otimes_{K_0(\text{Rep}_{\bar{\mathbb{Q}}}(G))} \mathbb{Q}_\ell \right]^+ \\
 & \swarrow \text{ch}_\ell \sim & \downarrow \sim (\text{ch } \circ (1.1)) \\
 & & H^{2*}(\hat{X}_{\bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))^+ \\
 & \swarrow & \downarrow \sim (\text{Prop. 2.3}) \\
 H_{\text{Chern}}^{2*} \left({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(*) \right) & \xrightarrow{\text{natural map } \sim} & IH^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(*))_v^+ \\
 & \searrow \varphi^* \sim & \downarrow \varphi_{\mathcal{L}}^* \sim \\
 & & H_{\text{Chern}}^{2*} ({}_K S(G, X)_{\Sigma, \bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))
 \end{array} \tag{4.17}$$

in which $\varphi_{\mathcal{L}}^*$, ch_ℓ , the natural map and φ^* are isomorphisms of $\mathbb{Q}_\ell$ -algebras. In particular

$$\varphi_{\mathcal{L}}^* \left(IH^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_\ell(*))_v^+ \right) \subset H^{2*} ({}_K S(G, X)_{\Sigma, \bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))$$

is a $\mathbb{Q}_\ell$ -sub-vectorspace which does not depend on the choice of the relative polarization $\mathcal{L}$.

Proof. By Theorem 2.12, the image of the Goresky-Pardon Chern classes $c_n([\mathcal{E}]_K)^{GP}$ in the algebra $IH^{2*}({}_K S(G, X)^{\min}, \mathbb{Q}(*))$ lies in $IH^{2*}({}_K S(G, X)^{\min}, \mathbb{Q}(*))_v^+$ and spans this subalgebra over $\mathbb{Q}$. In particular, the image of

$$H^{2*}({}_K S(G, X)^{\min}, \mathbb{Q}(*))_v \xrightarrow{\text{natural map}} IH^{2*}({}_K S(G, X)^{\min}, \mathbb{Q}(*))_v$$

contains $IH^{2*}({}_K S(G, X), \mathbb{Q}(*))_v^+$, and the map

$$\varphi_{\mathcal{L}}^* : IH^{2*}({}_K S(G, X), \mathbb{Q}(*))_v^+ \rightarrow H_{\text{Chern}}^{2*} ({}_K S(G, X)_{\Sigma}, \mathbb{Q}(*))^+$$

is surjective. By comparison with ℓ -adic intersection cohomology, the image of

$$H^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))_v \xrightarrow{\text{natural map}} IH^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))_v$$

contains $IH^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))_v^+$, and

$$\varphi_{\mathcal{L}}^* : IH^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))_v^+ \rightarrow H_{\text{Chern}}^{2*} ({}_K S(G, X)_{\Sigma, \bar{\mathbb{Q}}_p}, \mathbb{Q}_\ell(*))$$

is surjective as well.

We have no direct way to compare the Goresky-Pardon classes to our classes from Theorem 4.9 on ${}_K S(G, X)_{\bar{\mathbb{Q}}_p}$. However, applying Proposition 4.13 for $s = 0$, i.e. for $K_0 = K$, one has

$$\varphi_{\mathcal{L}}^* \left(H_{\text{Chern}}^{2*} \left({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(\ast) \right) \right) = H_{\text{Chern}}^{2*} \left({}_K S(G, X)_{\Sigma, \bar{\mathbb{Q}}_p}, \mathbb{Q}_{\ell}(\ast) \right).$$

As $\varphi_{\mathcal{L}}^*$ is injective, we deduce that the natural map sends the $\mathbb{Q}_{\ell}$ -subalgebra $H_{\text{Chern}}^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(\ast))$ onto $IH^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(\ast))_v^+$. This proves that the natural map in the proposition and φ^* are surjective. It remains to prove that they are injective as well. The Chern character has by definition values in $H_{\text{Chern}}^{2*}({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(\ast))$ and the whole diagram commutes. This proves that φ^* is injective as well. It follows that the homomorphisms ch_{ℓ} , φ^* and the natural map are isomorphisms. Finally, the image of φ^* being equal to the image of $\varphi_{\mathcal{L}}^*$, the latter does not depend on $\mathcal{L}$. This finishes the proof. $\square$

Lemma 4.18. *The classes*

$$c_{n,\ell}([\mathcal{E}]_p)_{K_r} \in H^{2n} \left({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(n) \right)_v^{G_{E_w}}$$

constructed in Theorem 4.9 vanish for all $n > 0$ when $\mathcal{E}$ is flat, that is when $\mathcal{E}$ comes from a G -representation.

Proof. Under the assumption of the lemma, $[\mathcal{E}]_p$ is trivial (see Notation 4.10). The construction of Theorem 4.9 thus implies the lemma. $\square$

Remarks 4.19. (i) If we ignore the action of the Galois group, the Chern classes constructed above depend only on the connected components of ${}_K S(G, X)^{\min}$. Thus the results above extend to general Shimura varieties of abelian type. We leave the precise formulation to the reader.

(ii) We naturally expect that Proposition 4.16 remains true without the superscripts $^+$, which in any case change nothing when G satisfies condition (iv) of Theorem 2.7. In the excluded case, it is easy to see that the classes denoted e_r in §2.A.a have non-trivial images in $IH^n({}_K S(G, X)_{\bar{\mathbb{Q}}_p}^{\min}, \mathbb{Q}_{\ell}(\ast))$; indeed, this follows from the results of [GP02] and the equality $e_r^2 = c_{1,r}^n$ for each r . However, we also have

$$(c_{1,r}^{k-1})^2 = c_{1,r}^n,$$

so it is possible that $c_{1,r}$ and $\pm e_r$ have the same image in IH^n . We have not attempted to prove that this is not the case.

5. Comparison of torsors

Let $(G, X) = (\text{GSp}(2g), X_{2g})$, write $\hat{X}_{2g}$ for the compact dual of X_{2g} , fix a base point $h \in X_{2g} \subset \hat{X}_{2g}$, and let Ω_h denote the fiber at h of the cotangent bundle to $\hat{X}_{2g}$. Let V_g denote the vector group $\mathbb{G}_a^g$. Let $K_{h,g} \subset G$ denote the stabilizer of h . Then $K_{h,g}$ can be identified with $\text{GL}(g) \times \mathbb{G}_m = \text{Aut}(V_g) \times \mathbb{G}_m$ in such a way that the restriction to $\text{GL}(g)$ of the isotropy action of $K_{h,g}$ on Ω_h is equivalent to $\text{Sym}^2(V_g)$. We take St to be the standard representation $K_{h,g} \mapsto \text{Aut}(V_g)$, and let $\mathcal{E}_{\text{St}}$ be the corresponding equivariant vector bundle on $\hat{X}_{2g}$. Fix $K_g = K_{g,p} \cdot K_g^p$ a neat open compact subgroup of $\text{GSp}(2g)(\mathbf{A}_f)$, and fix a toroidal datum Σ for K . As in §2, we write

$$\mathfrak{q}_g : {}_{K_g^p} \mathcal{A}_g^{\text{tor}} \rightarrow {}_{K_g^p} \mathcal{A}_g^{\min}$$

for the morphism denoted q above.

Proposition 5.1 ([PS16]). *There is a canonical isomorphism*

$$\theta_g : \mathfrak{q}_g^*[\mathcal{E}_{\text{St}}]_p \xrightarrow{\sim} [\mathcal{E}_{\text{St}}]_{dR, \Sigma}$$

of vector bundles over ${}_{K_g^p} \mathcal{A}_g^{\text{tor}}$.

Proof. This is essentially equivalent to Corollaire 1.16 of [PS16]. More precisely, the generic fiber of the bundle denoted ω_A^{mod} in [PS16] is exactly $[\mathcal{E}_{\text{St}}]_{dR,\Sigma}$. Indeed, with our notation, the restriction of $[\mathcal{E}_{\text{St}}]^{\text{can}}$ to the open Siegel modular variety in (neat) finite level K is the sheaf of relative sections of the relative 1-forms of the universal polarized abelian scheme, and its canonical extension to the toroidal compactification is isomorphic to the sheaf of relative sections of the relative invariant 1-forms on the corresponding semi-abelian scheme; see [Lan12, Proposition 6.9] or [Lan17, (1.3.3.15)]; the comparison between the algebraic and analytic constructions is Theorem 5.2.12 of [Lan12a]. $\square$

We now restrict the information to Shimura varieties of Hodge type. Let (G, X) be a Shimura datum of Hodge type, with compact dual $\hat{X}$. We fix a symplectic embedding

$$\iota : (G, X) \rightarrow (\text{GSp}(2g), X_{2g}).$$

Let $h \in X$ be a base point, let $K_h \subset G$ be its stabilizer, let $K_{h,g} \subset \text{GSp}(2g)$ be the stabilizer of $\iota(h)$, and denote by ι_h the inclusion of K_h in $K_{h,g}$. Let $\text{St}_h = \text{St} \circ \iota_h : K_h \rightarrow \text{Aut}(V_g)$, with St the standard faithful representation of $K_{h,g}$. Finally, let $\mathcal{E}_{\text{St}_h}$ be the equivariant vector bundle on $\hat{X}$ with isotropy representation St_h . Let $K = K_p \cdot K^p$ be a neat open compact subgroup of $G(\mathbf{A}_f)$.

Corollary 5.2. *We fix a natural number s . The morphism θ_g induces by pullback for any s a canonical isomorphism*

$$\theta(s) : Q^*[\mathcal{E}_{\text{St}_h}]_p \xrightarrow{\sim} [\mathcal{E}_{\text{St}_h}]_{dR,\Sigma,K_s}$$

of vector bundles over Z_s .

Proof of Theorem 3.12. We fix s as in Section 3. As St_h is faithful, any irreducible representation V of K_h defined over $\bar{\mathbb{Q}}$ is a direct factor of the representation $\text{St}_h^{\otimes m} \otimes \text{St}_h^{\vee \otimes n}$ for some pair of natural numbers (m, n) [DMOS82, I, Proposition 3.1(a), II, Proposition 2.23]. Write

$$V \xrightarrow{\sigma} \text{St}_h^{\otimes m} \otimes \text{St}_h^{\vee \otimes n} \xrightarrow{\tau} V$$

for the splitting. The isomorphism of vector bundles $\theta(s)$ induces an isomorphism of vector bundles

$$\theta(s)^{\otimes m} \otimes \theta(s)^{\vee \otimes n} : \mathfrak{q}_g^*([\mathcal{E}_{\text{St}_h}]_p^{\otimes m} \otimes [\mathcal{E}_{\text{St}_h}]_p^{\vee \otimes n}) \xrightarrow{\sim} [\mathcal{E}_{\text{St}_h}]_{dR,\Sigma,K_s}^{\otimes m} \otimes [\mathcal{E}_{\text{St}_h}]_{dR,\Sigma,K_s}^{\vee \otimes n}$$

while σ and τ induce the splittings of vector bundles over $K^p\mathcal{S}(G, X)^{\min}$ and over Z_s

$$\begin{aligned} [\mathcal{V}]_p &\xrightarrow{\sigma_p} [\mathcal{E}_{\text{St}_h}]_p^{\otimes m} \otimes [\mathcal{E}_{\text{St}_h}]_p^{\vee \otimes n} \xrightarrow{\tau_p} [\mathcal{V}]_p \\ [\mathcal{V}]_{dR,\Sigma,K_s} &\xrightarrow{\sigma_{dR,\Sigma,K_s}} [\mathcal{E}_{\text{St}_h}]_{dR,\Sigma,K_s}^{\otimes m} \otimes [\mathcal{E}_{\text{St}_h}]_{dR,\Sigma,K_s}^{\vee \otimes n} \xrightarrow{\tau_{dR,\Sigma,K_s}} [\mathcal{V}]_{dR,\Sigma,K_s}. \end{aligned}$$

This yields the following morphisms of coherent sheaves over Z_s

$$\begin{aligned} \alpha &:= \tau_{dR,\Sigma,K_s} \circ \theta(s)^{\otimes m} \otimes \theta(s)^{\vee \otimes n} \circ Q^* \sigma_p : Q^*[\mathcal{V}]_p \rightarrow [\mathcal{V}]_{dR,\Sigma,K_s} \\ \beta &:= Q^* \tau_p \circ (\theta(s)^{\otimes m} \otimes \theta(s)^{\vee \otimes n})^{-1} \circ \sigma_{dR,\Sigma,K_s} : [\mathcal{V}]_{dR,\Sigma,K_s} \rightarrow Q^*[\mathcal{V}]_p \end{aligned}$$

which pull back to inverse isomorphisms on Z_s^o under j_s (see (4.4)). Thus

$$\beta \circ \alpha - \text{Id}_{Q^*[\mathcal{V}]_p} : Q^*[\mathcal{V}]_p \rightarrow Q^*[\mathcal{V}]_p, \quad \alpha \circ \beta - \text{Id}_{[\mathcal{V}]_{dR,\Sigma,K_s}} : [\mathcal{V}]_{dR,\Sigma,K_s} \rightarrow [\mathcal{V}]_{dR,\Sigma,K_s}$$

are homomorphisms of coherent sheaves which pull back to zero on Z_s^o . As $Q^*[\mathcal{V}]_p$ and $[\mathcal{V}]_{dR,\Sigma,K_s}$ are vector bundles, thus are both torsion free, we conclude that both maps are 0.

This finishes the proof of Theorem 3.12. As for Theorem 2.22, the proof is identical, except that we do not have to fix s in the beginning and we replace Q by q everywhere. $\square$

Appendix A. Geometry

We collect here the statements of some standard results in the theory of schemes, in versions adapted to perfectoid spaces. The proofs are due to Peter Scholze.

Proposition A.1 (Scholze). *Let $\mathcal{W}$ be a vector bundle of rank r over a perfectoid space $\mathcal{X}$ over C . Let $\mathbb{P}(\mathcal{W})$ denote the corresponding projective bundle over $\mathcal{X}$, viewed as an adic space over C (see Remark A.2 below). Then $\mathcal{O}_{\mathbb{P}(\mathcal{W})}(1)$ has a first Chern class*

$$(z_m) \in \varprojlim_m H^2(\mathbb{P}(\mathcal{W}), \mathbb{Z}/\ell^m(1))$$

such that the homomorphism

$$\varprojlim_m H^b(\mathbb{P}(\mathcal{W}), \mathbb{Z}/\ell^m(a)) \leftarrow \oplus_{i=0}^{r-1} \varprojlim_m H^{b-2i}(\mathcal{X}, \mathbb{Z}/\ell^m(a-i)) \cdot (z_n)^i$$

is an isomorphism.

Remark A.2. One of the referees asked how $\mathbb{P}(\mathcal{W})$ can be defined as an adic space. We forwarded the question to Laurent Fargues and David Hansen, who sent essentially identical replies. Here is an (almost literal) paraphrase of Hansen's argument. First, without loss of generality, we may localize on $\mathcal{X}$ and assume that $\mathcal{X}$ is affinoid perfectoid, say $\mathrm{Spa}(R, R^+)$, and that $\mathcal{W}$ is the trivial bundle. Thus we need to define $\mathrm{Spa}(R, R^+) \times \mathbb{P}^{r-1}$ as an adic space. Next, by the usual covering of $\mathbb{P}^{r-1}$ by r affinoid balls, we are reduced to showing that the ring $R\langle T_1, \dots, T_{r-1} \rangle$ (power series with coefficients in R that tend to 0) is *sheafy* [SW17, Definition 3.1.11]. In fact, $R\langle T_1, \dots, T_{r-1} \rangle$ is *sousperfectoid*, in the terminology of Hansen and Kedlaya, and in particular is sheafy; see §6.3 of [SW17] for the proofs. Moreover, $R\langle T_1, \dots, T_{r-1} \rangle$ is *analytic* [SW17, Definition 4.4.2].

We are warned, however, that there is as yet no reference for the étale cohomology in the generality of sousperfectoid spaces. We have seen that $\mathbb{P}(\mathcal{W})$ is an analytic adic space; thus, in order to define the étale cohomology spaces used in Proposition A.1, one can pass to the associate locally spatial diamond (see [Sch17, §15]), and apply the cohomological formalism developed in [Sch17].

We define the Chern class $c_i(\mathcal{W})$ by Grothendieck's standard equation

$$\varprojlim_m H^{2r}(\mathbb{P}(\mathcal{W}), \mathbb{Z}/\ell^m(r)) \ni \sum_{i=0}^r (-1)^i c_i(\mathcal{W}) \cdot (z_m)^{r-i} = 0, \quad (c_i(\mathcal{W})_m) \in \varprojlim_m H^{2i}(\mathbb{P}(\mathcal{W}), \mathbb{Z}/\ell^m(i)).$$

Proof. One defines z as usual, as the projective limit over m of the images $z_m \in H^2(\mathbb{P}(\mathcal{W}), \mathbb{Z}/\ell^m(1))$ of the class of $\mathcal{O}_{\mathbb{P}(\mathcal{W})}(1)$ in $H^1(\mathbb{P}(\mathcal{W}), \mathbb{G}_m)$ via the connecting homomorphism of the étale Kummer exact sequence $1 \rightarrow \mu_{\ell^m} \rightarrow \mathbb{G}_m \xrightarrow{\ell^m} \mathbb{G}_m \rightarrow 1$. To prove the statement, it is enough to prove that the map

$$H^b(\mathbb{P}(\mathcal{W}), \mathbb{Z}/\ell^m(a)) \leftarrow \oplus_{i=0}^{r-1} H^{b-2i}(\mathcal{X}, \mathbb{Z}/\ell^m(a-i)) \cdot z_m^i$$

is an isomorphism. This is a local property on $\mathcal{X}$, reduced by [CS17, Lemma 4.4.1] to the computation of the étale cohomology of $\mathbb{P}(\mathcal{W})$ over a geometric point $\bar{x} = \mathrm{Spa}(C(\bar{x}), C(\bar{x})^+)$, which then is the standard computation. $\square$

Remark A.3. We avoid here the delicate question whether the surjection

$$H^j(\mathcal{X}, \mathbb{Z}_{\ell}(i)) \rightarrow \varprojlim_m H^j(\mathcal{X}, \mathbb{Z}/\ell^m(i))$$

is an isomorphism as it is irrelevant for our purpose.

Recall that we have level subgroups $K_{p,r} \subset G(\mathbb{Q}_p)$. We denote by Σ^H the invariants under a group H acting on the set Σ ; we use this notation for cohomology invariants.

Proposition A.4 (Scholze). *The morphisms of ringed spaces*

$$\overline{K^p \mathcal{S}(G, X)}^{\min} \rightarrow_{K^p \cdot K_{p,r}} \overline{\mathcal{S}(G, X)}^{\min}$$

induce for all pairs of integers (i, j) and all $r \geq 1$ homomorphisms

$$H^j \left(\overline{K^p \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right)^{K_{p,r}} \rightarrow H^j \left(\overline{K^p \cdot K_{p,r} \mathcal{S}(G, X)}^{\min}_{\mathbb{Q}_p}, \mathbb{Z}/\ell^m(i) \right)$$

and

$$\left[\lim_{\leftarrow m} H^j \left(\overline{K^p \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right) \right]^{K_{p,r}} = \lim_{\leftarrow m} \left[H^j \left(\overline{K^p \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right) \right]^{K_{p,r}} \rightarrow H^j \left(\overline{K^p \cdot K_{p,r} \mathcal{S}(G, X)}^{\min}_{\mathbb{Q}_p}, \mathbb{Z}/\ell(i) \right).$$

Proof. As before, we write $K_r = K^p \cdot K_{p,r}$. By Theorem 2.13 and [Sch12, Corollary 7.18], one has

$$H^j \left(\overline{K^p \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right) = \lim_{\leftarrow r} H^j \left(\overline{K_r \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right) = \lim_{\leftarrow r} H^j \left(\overline{K_r \mathcal{S}(G, X)}^{\min}_{\mathbb{Q}_p}, \mathbb{Z}/\ell^m(i) \right).$$

The second equality comes from the comparison isomorphism [Hub98, Theorem 4.2].

On the other hand, for $r' > r \geq 1$ one has $K_r/K_{r'} = K_{p,r}/K_{p,r'}$, which is a p -group. It follows that the transition maps in the inductive system are injective as $\overline{K_{r+1} \mathcal{S}(G, X)}^{\min} \rightarrow \overline{K_r \mathcal{S}(G, X)}^{\min}$ is finite surjective of degree prime to ℓ . This implies

$$\begin{aligned} H^j \left(\overline{K^p \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right)^{K_{p,r}} &= \lim_{\leftarrow r'} H^j \left(\overline{K_{r'} \mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right)^{K_{p,r}/K_{p,r'}} \\ &= \lim_{\leftarrow r'} H^j \left([\overline{K_{r'} \mathcal{S}(G, X)}^{\min}] / [K_{p,r}/K_{p,r'}], \mathbb{Z}/\ell^m(i) \right) \end{aligned} \quad (\text{A.5})$$

if $r \geq 1$, because $K_{p,r}/K_{p,r'}$ is a p -group. Here $[\overline{K_{r'} \mathcal{S}(G, X)}^{\min}] / [K_{p,r}/K_{p,r'}]$ denotes the quotient of the (projective) scheme $\overline{K_{r'} \mathcal{S}(G, X)}^{\min}$ by the finite p -group $K_{p,r}/K_{p,r'}$.

If $r = 0$, the last two groups are equal up to finite error that is bounded independently of r' and m ; we return to this case below.

For any $r' \geq r$, the quotient $\overline{K_{r'} \mathcal{S}(G, X)}^{\min} / [K_{p,r}/K_{p,r'}]$ lies in a sequence of finite morphisms

$$\overline{K_r \mathcal{S}(G, X)}^{\min} \rightarrow \overline{K_{r'} \mathcal{S}(G, X)}^{\min} / [K_{p,r}/K_{p,r'}] \rightarrow \overline{K_r \mathcal{S}(G, X)}^{\min}.$$

Indeed, the minimal compactification is normal, which explains the left arrow, and the right one is just the image of the middle term in the minimal compactification of the Siegel space at level K_r . Define

$$\overline{K_r \mathcal{S}(G, X)}^{\min, \#} = \lim_{\leftarrow r'} \overline{K_{r'} \mathcal{S}(G, X)}^{\min} / [K_{p,r}/K_{p,r'}].$$

This is a scheme of finite type as it is dominated by $\overline{K_r \mathcal{S}(G, X)}^{\min}$. Again we have a sequence of finite morphisms

$$\overline{K_r \mathcal{S}(G, X)}^{\min} \rightarrow \overline{K_r \mathcal{S}(G, X)}^{\min, \#} \rightarrow \overline{K_r \mathcal{S}(G, X)}^{\min}.$$

It follows from the above considerations that, for any m, j , and i , we have

$$H^j \left(\overline{{}_{K^p}\mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right)^{K_{p,r}} = \varprojlim_r H^j \left([{}_{K_r}\overline{\mathcal{S}(G, X)}^{\min, \#}], \mathbb{Z}/\ell^m(i) \right)$$

if $r \geq 1$, and with finite error independent of m if $r = 0$.

So composing with the natural map

$$H^* \left({}_{K_r}\mathcal{S}(G, X)^{\min, \#}, \mathbb{Z}/\ell^m(i) \right) \rightarrow H^* \left({}_{K_r}\mathcal{S}(G, X)^{\min}, \mathbb{Z}/\ell^m(i) \right)$$

one obtains the homomorphism

$$H^j \left(\overline{{}_{K^p}\mathcal{S}(G, X)}^{\min}, \mathbb{Z}/\ell^m(i) \right)^{K_{p,r}} \rightarrow H^j \left({}_{K_r}\mathcal{S}(G, X)^{\min}, \mathbb{Z}/\ell^m(i) \right) = H^j \left({}_{K_r}\mathcal{S}(G, X)_{\overline{\mathbb{Q}_p}}^{\min}, \mathbb{Z}/\ell^m(i) \right) \quad (\text{A.6})$$

(see again [Hub98], *loc.cit.*, for the comparison isomorphism), if $r \geq 1$.

Finally, by definition, one has

$$\varprojlim_m H^j \left({}_{K^p}\mathcal{S}(G, X)^{\min}, \mathbb{Z}/\ell^m(i) \right)^{K_{p,r}} = \left[\varprojlim_m H^j \left({}_{K^p}\mathcal{S}(G, X)^{\min}, \mathbb{Z}/\ell^m(i) \right) \right]^{K_{p,r}},$$

which thus maps to

$$\varprojlim_m H^j \left({}_{K_r}\mathcal{S}(G, X)_{\overline{\mathbb{Q}_p}}^{\min}, \mathbb{Z}/\ell^m(i) \right) = H^j \left({}_{K_r}\mathcal{S}(G, X)_{\overline{\mathbb{Q}_p}}^{\min}, \mathbb{Z}_\ell(i) \right).$$

□

If $r = 0$ this remains true with $\mathbb{Z}/\ell^m(i)$ replaced by $\mathbb{Q}_\ell(i)$: the maps in (A.5) are not necessarily isomorphisms but the kernels and cokernels are of order bounded independently of m and i .

References

- [AMRT75] A. Ash, D. Mumford, M. Rapoport, and Y. Tai, *Smooth compactification of locally symmetric varieties*. In: Lie Groups: History, Frontiers and Applications, vol. IV, Math. Sci. Press, Brookline, Mass., 1975. [MR-0457437](#)
- [BB66] W. L. Baily, Jr. and A. Borel, *Compactification of arithmetic quotients of bounded symmetric domains*, Ann. of Math. (2) **84** (1966), 442–528. [MR-0216035](#)
- [BBD82] A. A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers*. In: Analysis and topology on singular spaces, I (Luminy, 1981), pp. 5–171, Astérisque, vol. 100, Soc. Math. France, Paris, 1982. [MR-0751966](#)
- [Bha17] B. Bhatt, *Lecture notes for a class on perfectoid spaces*, 2017. <http://www-personal.umich.edu/~bhattb/teaching/mat679w17/lectures.pdf>
- [BH58] A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces. I*. Amer. J. Math. **80** (1958), 458–538. [MR-0102800](#)
- [Bor83] A. Borel, *Regularization theorems in Lie algebra cohomology. Applications*. Duke Math. J. **50** (1983), no. 3, 605–623. [MR-0714820](#)
- [CS17] A. Caraiani and P. Scholze, *On the generic part of the cohomology of compact unitary Shimura varieties*, Ann. of Math. (2) **186** (2017), no. 3, 649–766. [MR-3702677](#)

- [C+6] A. Caraiani, D. R. Gulotta, C.-Y. Hsu, C. Johansson, L. Mocz, E. Reinecke, and S.-C. Shih, *Shimura varieties at level $\Gamma_1(p^\infty)$ and Galois representations*, preprint 2018. [arXiv:1804.00136](#)
- [dCM05] M. A. A. de Cataldo and L. Migliorini, *Hodge theoretic aspects of the Decomposition Theorem*. In: Algebraic Geometry – Seattle 2005, part 2, pp. 489–504, Proc. Sympos. Pure Math., vol. 80, part 2, Amer. Math. Soc., Providence, RI, 2009. [MR-2483945](#)
- [Del71] P. Deligne, *Travaux de Shimura*. In: Séminaire Bourbaki, 23ème année (1970/71), Exp. No. 389, pp. 123–165, Lectures Notes in Math., vol. 244, Springer, Berlin, 1971. [MR-0498581](#)
- [Del74] P. Deligne, *Théorie de Hodge. III*. Inst. Hautes Études Sci. Publ. Math. **44** (1974), 5–77. [MR-0498552](#)
- [DMOS82] P. Deligne, J. S. Milne, A. Ogus, and K. Shih, *Hodge cycles, motives and Shimura varieties*, Lecture Notes in Mathematics, vol. 900, Springer-Verlag, Berlin-New York, 1982. [MR-0654325](#)
- [EV02] H. Esnault and E. Viehweg, *Chern classes of Gauss-Manin bundles of weight 1 vanish*, K-Theory **26** (2002), no. 3, 287–305. [MR-1936356](#)
- [EH17] H. Esnault and M. Harris, *Chern classes of automorphic vector bundles*, Pure Appl. Math. Q. **13** (2017), no. 2, 193–213. [MR-3858008](#)
- [GP02] M. Goresky and W. Pardon, *Chern classes of automorphic vector bundles*, Invent. math. **147** (2002), no. 3, 561–612. [MR-1893006](#)
- [GM03] M. Goresky and R. MacPherson, *The topological trace formula*, J. Reine Angew. Math. **560** (2003), 77–150. [MR-1992803](#)
- [Har85] M. Harris, *Arithmetic vector bundles and automorphic forms on Shimura varieties. I*. Invent. Math. **82** (1985), no. 1, 151–189. [MR-0808114](#)
- [Har89] M. Harris, *Functorial properties of toroidal compactifications of locally symmetric varieties*, Proc. London Math. Soc. (3) **59** (1989), no. 1, 1–22. [MR-0997249](#)
- [Har90] M. Harris, *Automorphic forms of $\bar{\partial}$ -cohomology type as coherent cohomology classes*, J. Differential Geom. **32** (1990), no. 1, 1–63. [MR-1064864](#)
- [HZ94] M. Harris and S. Zucker, *Boundary cohomology of Shimura varieties. I. Coherent cohomology on toroidal compactifications*. Ann. Sci. École Norm. Sup. (4) **27** (1994), no. 3, 249–344. [MR-1272293](#)
- [Hub98] R. Huber, *A comparison theorem for ℓ -adic cohomology*, Compositio Math. **112** (1998), no. 2, 217–235. [MR-1626021](#)
- [Lan12] K.-W. Lan, *Toroidal compactifications of PEL-type Kuga families*, Algebra Number Theory **6** (2012), no. 5, 885–966. [MR-2968629](#)
- [Lan12a] K.-W. Lan, *Comparison between analytic and algebraic constructions of toroidal compactifications of PEL-type Shimura varieties*, J. Reine Angew. Math. **664** (2012), 163–228. [MR-2980135](#)
- [Lan13] K.-W. Lan, *Arithmetic compactifications of PEL-type Shimura varieties*, London Mathematical Society Monographs, vol. 36, Princeton University Press, Princeton, NJ, 2013. [MR-3186092](#)
- [Lan17] K.-W. Lan, *Compactifications of PEL-type Shimura varieties and Kuga families with ordinary loci*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2018. [MR-3729423](#)
- [Loo88] E. Looijenga, *L^2 -cohomology of locally symmetric varieties*, Compositio Math. **67** (1988), no. 1, 3–20. [MR-0949269](#)

- [Loo17] E. Looijenga, *Goresky-Pardon lifts of Chern classes and some Tate extension*, *Compositio Math.* **153** (2017), no. 7, 1349–1371. [MR-3705260](#)
- [Mum77] D. Mumford, *Hirzebruch's proportionality theorem in the noncompact case*, *Invent. math.* **42** (1977), 239–272. [MR-0471627](#)
- [Nai14] A. N. Nair, *Chern classes of automorphic bundles and the reductive Borel-Satake compactification*, preprint 2014. http://www.math.tifr.res.in/~arvind/preprints/chernrbs_v2.pdf
- [PS16] V. Pilloni and B. Stroh, *Cohomologie cohérente et représentations Galoisiennes*, *Ann. Math. Qué.* **40** (2016), no. 1, 167–202. [MR-3512528](#)
- [Pin90] R. Pink, *Arithmetical compactification of mixed Shimura varieties*, *Bonner Mathematische Schriften*, vol. 209, Universität Bonn, Mathematisches Institut, Bonn, 1990. [MR-1128753](#)
- [SS90] L. Saper and M. Stern, *L_2 -cohomology of arithmetic varieties*, *Ann. of Math. (2)* **132** (1990), no. 1, 1–69. [MR-1059935](#)
- [Sch12] P. Scholze, *Perfectoid spaces*, *Publ. Math. Inst. Hautes Études Sci.* **116** (2012), 245–313. [MR-3090258](#)
- [SW13] P. Scholze and J. Weinstein, *Moduli of p -divisible groups*, *Camb. J. Math.* **1** (2013), no. 2, 145–237. [MR-3272049](#)
- [Sch14] P. Scholze, *Perfectoid spaces and their applications*. In: *Proceedings of the International Congress of Mathematicians – Seoul 2014*, vol. II, pp. 461–486, Kyung Moon Sa, Seoul, 2014. [MR-3728623](#)
- [Sch15] P. Scholze, *On torsion in the cohomology of locally symmetric varieties*, *Ann. of Math. (2)* **182** (2015), no. 3, 945–1066. [MR-3418533](#)
- [SW17] P. Scholze and J. Weinstein, *Berkeley lectures on p -adic geometry*, 2014. <http://www.math.uni-bonn.de/people/scholze/Berkeley.pdf>
- [Sch17] P. Scholze, *Étale cohomology of diamonds*, preprint 2017. [arXiv:1709.07343](#)