

Quantifying CDS sortability of permutations by strategic pile size

Marisa Gaetz

*Department of Mathematics, Massachusetts Institute of Technology
77 Massachusetts Ave, Cambridge, MA 02139, USA mgaetz@mit.edu*

Bethany Flanagan

*Department of Mathematics, Purdue University
610 Purdue Mall, West Lafayette, IN 47907, USA bmolokac@purdue.edu*

Marion Scheepers

*Department of Mathematics, Boise State University
1910 West University Dr, Boise, ID 83725, USA mscheepe@boisestate.edu*

Meghan Shanks

*Department of Computer Science
University of Illinois at Urbana-Champaign
201 N Goodwin Ave, Urbana, IL 61801, USA megshanks2@illinois.edu*

Received 28 December 2018

Revised 6 September 2019

Accepted 13 November 2019 Published

10 January 2020

The special purpose sorting operation, *context directed swap* (CDS), is an example of the block interchange sorting operation studied in prior work on permutation sorting. CDS has been postulated to model certain molecular sorting events that occur in the genome maintenance program of some species of ciliates. We investigate the mathematical structure of permutations not sortable by the CDS sorting operation. In particular, we present substantial progress towards quantifying permutations with a given *strategic pile* size, which can be understood as a measure of CDS non-sortability. Our main results include formulas for the number of permutations in S_n with maximum size strategic pile. More generally, we derive a formula for the number of permutations in S_n with strategic pile size k , in addition to an algorithm for computing certain coefficients of this formula, which we call *merge numbers*.

Keywords: Permutation sorting; context directed swap; strategic pile; factorization into cycles.

1. Introduction

Sorting is a fundamental step in numerous natural, industrial, commercial, and scientific computing processes. Correspondingly, the mathematical analysis of sorting operations has a long history. The typical concerns with a sorting process include the efficiency of the sorting operation, a characterization of the situations in which the sorting operation achieves the sorting objective, and a characterization of the situations in which the sorting operation does not achieve the sorting objective. In this paper, we focus on the third of these concerns. In particular, we seek to quantify for a specific sorting operation the prevalence of what can be seen as the worst case obstruction to sortability.

The specific sorting operation we consider aims to sort a permuted list of the numbers $1, 2, \dots, n$ to the canonical ordered list $(1, 2, \dots, n)$. This sorting operation appears in two prior works. It appears in the 2003 template model for the construction of a new macronucleus from its scrambled precursor micronucleus in certain ciliate species [9]. In this model the sorting operation is named *dlad*. For more on this fascinating biological background the reader may consult the review [8] and the textbook [5]. It turns out, by hindsight, that this sorting operation also includes special cases of the block interchange sorting operation examined in [4] by Christie. The *minimal block interchanges* identified by Christie are special cases of the *dlad* operation.

In yet another investigation into genome rearrangement combinatorics, the double cut and join operation, denoted DCJ, is introduced by Yancopoulos *et al.* [11] to establish a mathematical measure of distance between two genomes. In the DCJ theory, generic block interchanges studied by Christie [4] are modeled by a very specific sequence of DCJ events, visualized in [11, Fig. 6]. Modeling *dlad* as a DCJ operation requires specifying additional DCJ constraints. To emphasize the specific mathematical nature of the sorting operation we consider here, the operation will be called *context directed swap*, denoted CDS; CDS is an example of a block interchange sorting algorithm. We base our treatment on the mathematical counterpart of the essential features identified in the paper [9].

A permuted list of numbers is said to be *CDS-sortable* if there is a sequence of applications of the CDS sorting operation (to be defined in Sec. 2) that results in the numbers listed in increasing order. Not every permutation is sortable by CDS. CDSsortability criteria have been given previously (for instance, see [6]). Also, from prior work one can deduce that when CDS can sort a permutation, it is the most efficient block interchange sorting algorithm [4]. Mathematically interesting phenomena arise from the study of permutations not sortable by applications of CDS. The essential structural obstacle to a permutation's CDS-sortability was identified in [1], giving rise to the notion of the *strategic pile* of a permutation.

The notions of CDS-sortability, the strategic pile of a permutation, and appropriate notation and terminology will be introduced in Sec. 2. In this section, we explicitly describe the problem being treated in this paper, and we report our findings in Secs. 3–5.

In Sec. 3, we determine the number of elements in S_n that have the maximum size strategic pile among all elements of S_n . This counting problem reduces to a variation of the cycle factoring problem for S_n , studied previously, and depends on the cycle factoring results of [2, 3]. In Sec. 4, we investigate how prevalent it is for permutations in S_n to have strategic piles of cardinality k . As a result of this work we develop formulas in closed form that produce the terms of the integer sequences A267323, A267324, and A267391 in [10]. We also contribute the integer sequence A281259 to [10], as well as its formula. In Sec. 5, we highlight a more challenging component of our formula from Sec. 4.

2. Preliminaries

For a positive integer n , the symbol S_n denotes the set of one-to-one functions from the set $\{1, 2, \dots, n\}$ to itself, also known as permutations of $\{1, 2, \dots, n\}$. The notation

$$[a_1 \ a_2 \ \cdots \ a_{n-1} \ a_n] \quad (2.1)$$

denotes the permutation π for which $\pi(i) = a_i$ for $1 \leq i \leq n$. In current literature, the notation in (2.1) is called *one-line notation*. This one-line notation should be distinguished from

$$(c_1 \ c_2 \ \cdots \ c_{k-1} \ c_k), \quad (2.2)$$

which is the so-called *cycle notation* that denotes the permutation π where $\pi(c_1) = c_2, \pi(c_2) = c_3, \dots, \pi(c_{k-1}) = c_k, \pi(c_k) = c_1$, and where $\pi(i) = i$ for $i \notin \{c_1, \dots, c_k\}$

(note that this notation is very similar to notation we will later use to describe ordered lists. The two can be distinguished by noting that we do not use commas to describe a cycle permutation, but will use them to describe ordered lists).

To define the CDS sorting operation, associate with each entry of the permutation $\pi \in S_n$ left and right pointers as follows: For an entry $k \in \{1, 2, \dots, n\}$ of π , the *left pointer* of k is $\langle k-1, k \rangle$, while the *right pointer* of k is $\langle k, k+1 \rangle$. By convention, the smallest entry, 1, does not have a left pointer, and the largest entry, n , does not have a right pointer.

Example 2.1. Equation (2.3) shows the permutation $\pi = [2 \ 4 \ 3 \ 1 \ 5]$ with all pointers marked:

$$\pi = [{}_{(1,2)}2{}_{(2,3)} \ {}_{(3,4)}4{}_{(4,5)} \ {}_{(2,3)}3{}_{(3,4)} \ {}_1{}_{(1,2)} \ {}_{(4,5)}5] \quad (2.3)$$

Observe that each pointer in a permutation occurs twice. Given two pointers, p and q , in the permutation π , the sorting operation CDS at these pointers acts as follows on π : If the pointers do not appear in the order $\cdots p \cdots q \cdots p \cdots q \cdots$ in π , then CDS does not apply and we say that the pointer context is invalid. Otherwise, the two segments of π that are flanked by the pointer context $p \cdots q$ are interchanged. The pointers $p = \langle 3, 4 \rangle$ and $q = \langle 4, 5 \rangle$ appear in $\cdots p \cdots q \cdots p \cdots q \cdots$ context in the permutation $\pi = [2 \ 4 \ 3 \ 1 \ 5]$. CDS applied to π for this pointer context produces the permutation $[2 \ 1 \ 3 \ 4 \ 5]$. On the other hand, as the pointers $r = \langle 1, 2 \rangle$ and $s = \langle 3, 4 \rangle$ appear in $\cdots r \cdots s \cdots s \cdots r \cdots$ context in π , CDS cannot be applied.

When there are no pointers p and q that appear in context $\cdots p \cdots q \cdots p \cdots q \cdots$ in π , the permutation π is said to be a CDS *fixed point*. For each positive integer n , there are exactly n CDS fixed points in S_n , namely the permutations $[k+1 \ \cdots \ n \ 1 \ 2 \ \cdots \ k]$ for $1 \leq k < n$, and the identity permutation $[1 \ 2 \ \cdots \ n-1 \ n]$.

By [1], we know that for each permutation π in S_n that is not a CDS fixed point, some sequence of applications of CDS to π terminate in a CDS fixed point. If a sequence of applications of CDS to the permutation π terminates in the identity permutation $[1 \ 2 \ \cdots \ n]$, we say that π is CDS-*sortable*. The CDS-sortability of permutations has been characterized in prior works such as [1, 6]. In [1], the obstacle to CDS-sortability of a permutation $\pi \in S_n$ is identified as follows. Suppose $\pi =$

$[a_1 \ a_2 \ \cdots \ a_n]$. Define the cycle permutations X_n and Y_π by

$$X_n := (0 \quad 1 \quad 2 \quad \cdots \quad n) \quad \text{and} \quad (2.4)$$

$$Y_\pi := (0 \quad a_n \quad a_{n-1} \quad \cdots \quad a_1). \quad (2.5)$$

Then define

$$C_\pi := Y_\pi \circ X_n. \quad (2.6)$$

In Eq. (2.6), the symbol “ $\circ$ ” denotes functional composition, and we use the standard convention that $f \circ g(x)$ denotes the value $f(g(x))$.

When the entries 0 and n occur in the same cycle in the disjoint cycle decomposition of C_π , we shall write this cycle in the form

$$(0 \quad u_1 \quad u_2 \quad \cdots \quad u_j \quad n \quad b_1 \quad b_2 \quad \cdots \quad b_k). \quad (2.7)$$

The set $SP(\pi) = \{b_1, b_2, \dots, b_k\}$ is said to be the *strategic pile* of π . If 0 and n do not appear in the same cycle, we define $SP(\pi)$ to be the empty set. The ordered list $SP^*(\pi) = (b_1, b_2, \dots, b_k)$ is called the *ordered strategic pile* of π , and its ordering is determined by the order of appearance in (2.7). In [1], it was proven that a permutation π is CDS-

sortable if and only if its strategic pile is the empty set (i.e., if and only if 0 and n do not appear in the same cycle).

Example 2.2. For the permutation $\pi = [2\ 5\ 1\ 4\ 3]$ we have $C_\pi = Y_\pi \circ X_5 = (0\ 3\ 4\ 1\ 5\ 2)(0\ 1\ 2\ 3\ 4\ 5) = (0\ 5\ 3\ 1)(2\ 4)$, written in disjoint cycle form. Thus, the strategic pile of π is the set $SP(\pi) = \{1, 3\}$, while $SP^*(\pi) = (3, 1)$.

The strategic pile of a permutation π is intimately related to the set of achievable CDS fixed points.

Theorem 2.3 ([1, Theorem 2.22]). *If a permutation $\pi \in S_n$ is not CDS-sortable, then the following are equivalent for $1 \leq k < n$:*

- (1) *There is a sequence of applications of CDS to π that terminates in the CDS fixed point $[k + 1\ k + 2\ \dots\ n\ 1\ 2\ \dots\ k]$.* (2) *k is a member of the strategic pile of π .*

3. Maximum Size Strategic Piles

We now investigate the number of permutations in S_n with maximum size strategic piles; these permutations can be considered to have maximal CDS nonsortability. Since there are n CDS fixed points (including the identity permutation), Theorem 2.3 implies that a strategic pile of a permutation in S_n can have at most $n - 1$ elements.

Lemma 3.1. *If there is a permutation in S_n which has a strategic pile of size $n-1$, then n is even.*

Proof. By (2.7), if the strategic pile of permutation π has size $n - 1$, then

$$C_\pi = (0\ n\ b_1\ b_2\ \dots\ b_{n-1}). \quad (3.1)$$

But C_π is the composition of two $(n + 1)$ -cycles, and thus an even permutation. Therefore n is even. $\square$

As we shall see later, the converse of Lemma 3.1 also holds. As a consequence of Lemma 3.1 we get the following.

Corollary 3.2. *If n is odd, then the strategic pile of an element of S_n has at most $n - 2$ elements.*

We shall also later see that there are permutations in S_n with strategic pile of size $n - 2$ for every odd integer $n \geq 1$. In Secs. 3.1 and 3.2 we count for each n the number

of permutations in S_n with strategic pile of maximal size for n . Section 3.1 is dedicated to the case when n is even, and Sec. 3.2 is dedicated to the case when n is odd.

3.1. Maximum size strategic piles for even values of n

Theorem 3.3. *For each even number n , the number of permutations in S_n with strategic pile of size $n - 1$ is*

$$\frac{2(n-1)!}{n}.$$

As noted in the proof of Lemma 3.1, an element of S_n having a strategic pile of size $n - 1$ is related to the possibility of factoring certain $(n + 1)$ -cycles into two

$(n+1)$ -cycles. As a result, to prove Theorem 3.3, we first introduce some additional notation, which we will use to define injective maps between sets of factorizations.

Notation 3.4.

- Let A denote the set of all factorizations of X_{n-2} into two $(n-1)$ -cycles.
- Let B denote the set of all factorizations of X_n into two $(n+1)$ -cycles where the rightmost factor is of the form $(0 \ n \ \cdots)$.
- Let B_i denote the subset of B whose elements have rightmost factors of form $(0 \ n \ i \ \cdots)$.
- Define $\lambda_n = (0 \ n \ 1)$ and $c_n = (1 \ 2 \ \cdots \ n-1)$.

We begin by constructing a bijection between the sets A and B_1 in Lemmas 3.6 and 3.8. We then show in Lemma 3.9 that $B_1 \rightarrow B_2, B_2 \rightarrow B_3, \dots, B_{n-2} \rightarrow B_{n-1}$, and $B_{n-1} \rightarrow B_1$, where $\rightarrow$ indicates an injective map. Since the injective maps between the B_i sets form a cycle, it follows that $|A| = |B_1| = \cdots = |B_{n-1}|$. Finally, we determine $|A|$ using the following prior result that counts the number of factorizations of an arbitrary $(n-1)$ -cycle into two cycles of length $n-1$.

Lemma 3.5 ([3, Theorem 3]). *Let $\sigma \in S_{n-1}$ be an even $(n-1)$ -cycle. Then the number of factorizations of σ into two $(n-1)$ -cycles is*

$$\frac{2(n-2)!}{n}.$$

We now establish the previously described injections.

Lemma 3.6. *There is an injective map from A to B_1 .*

Proof. Let $\gamma \circ \delta$ be a factorization in A . Namely, suppose γ and δ are $(n-1)$ -cycles satisfying $\gamma \circ \delta = X_{n-2}$. Define γ_1 and δ_1 as follows:

$$\gamma_1 := \lambda_n \circ c_n \circ \gamma \circ (c_n)^{-1},$$

$$\delta_1 := c_n \circ \delta \circ (c_n)^{-1} \circ \lambda_n.$$

Note that if $\gamma_1 = \gamma'_1$, then $\gamma = \gamma'$, which shows that the map is injective. Therefore, to complete the proof it suffices to show that γ_1 and δ_1 are $(n+1)$ -cycles, that δ_1 is of the form $(0 \ n \ 1 \ \cdots)$, and that $\gamma_1 \circ \delta_1 = X_n$.

Since conjugation preserves cycle structure, the factors $c_n \circ \gamma \circ (c_n)^{-1}$ of γ_1 form an $(n - 1)$ -cycle with elements $\{0, 2, 3, \dots, n - 1\}$. Composing λ_n with this $(n - 1)$ -cycle creates an $(n + 1)$ -cycle with elements $\{0, 1, 2, \dots, n\}$.

Similarly, the factors $c_n \circ \delta \circ (c_n)^{-1}$ of δ_1 form an $(n - 1)$ -cycle with elements $\{0, 2, 3, \dots, n - 1\}$. Composing this $(n - 1)$ -cycle with λ_n adds the elements n and 1 to form an $(n + 1)$ -cycle of the form $(0 \ n \ 1 \ \dots)$.

Finally,

$$\begin{aligned} \gamma_1 \circ \delta_1 &= (\lambda_n \circ c_n \circ \gamma \circ (c_n)^{-1}) \circ (c_n \circ \delta \circ (c_n)^{-1} \circ \lambda_n) \\ &= \lambda_n \circ c_n \circ X_{n-2} \circ (c_n)^{-1} \circ \lambda_n \\ &= \lambda_n \circ (0 \ 2 \ 3 \ \dots \ n - 1) \circ \lambda_n = X_n. \end{aligned} \quad \square$$

Example 3.7. Let $n = 6$. Then $X_{n-2} = X_4 = (0 \ 1 \ 2 \ 3 \ 4)$. Consider the factorization

$$X_4 = (0 \ 1 \ 2 \ 3 \ 4) = (0 \ \underline{2 \ 4} \ \underline{1 \ 3}) (0 \ \underline{4 \ 3} \ \underline{2 \ 1})$$

$\gamma \qquad \qquad \delta$

Using the maps defined in Lemma 3.6, we get $\gamma_1 =$

$$\begin{aligned} &\lambda_6 \circ c_6 \circ \gamma \circ (c_6)^{-1} \\ &= (0 \ 6 \ 1)(1 \ 2 \ 3 \ 4 \ 5)(0 \ 2 \ 4 \ 1 \ 3)(5 \ 4 \ 3 \ 2 \ 1) \\ &= (0 \ 3 \ 5 \ 2 \ 4 \ 6 \ 1) \end{aligned}$$

and $\delta_1 = c_6 \circ \delta \circ (c_6)^{-1} \circ \lambda_6$

$$\begin{aligned} &= (1 \ 2 \ 3 \ 4 \ 5)(0 \ 4 \ 3 \ 2 \ 1)(5 \ 4 \ 3 \ 2 \ 1)(0 \ 6 \ 1) = (0 \\ &\quad 6 \ 1 \ 5 \ 4 \ 3 \ 2). \end{aligned}$$

Note that these are $(n + 1)$ -cycles, that δ_1 is of the form $(0 \ n \ 1 \ \dots)$, and that

$$\begin{aligned} \gamma_1 \circ \delta_1 &= (0 \ 3 \ 5 \ 2 \ 4 \ 6 \ 1)(0 \ 6 \ 1 \ 5 \ 4 \ 3 \ 2) \\ &= (0 \ 1 \ 2 \ 3 \ 4 \ 5 \ 6) = X_n, \end{aligned}$$

as desired.

Lemma 3.8. *There is an injective map from B_1 to A .*

Proof. Let $\gamma_1 \circ \delta_1 = (0 \ t_1 \ t_2 \ \dots \ t_n)(0 \ n \ 1 \ v_1 \ \dots \ v_{n-2})$ be an arbitrary factorization in B_1 . It suffices to show that we can associate to $\gamma_1 \circ \delta_1$ a unique factorization $\gamma \circ \delta$ of X_{n-2} in A . To this end, let $\delta := (c_n)^{-1} \circ \delta_1 \circ (\lambda_n)^{-1} \circ c_n$. Then,

$$\begin{aligned} \delta &= (c_n)^{-1} \circ (0 \quad n \quad 1 \quad v_1 \quad \dots \quad v_{n-2}) \circ (\lambda_n)^{-1} \circ c_n \\ &= (c_n)^{-1} \circ (0 \quad v_1 \quad \dots \quad v_{n-2})(1)(n) \circ c_n \\ &= (0 \quad v_1 - 1 \quad \dots \quad v_{n-2} - 1). \end{aligned}$$

It follows that δ is an $(n-1)$ -cycle. Moreover, changing δ_1 also changes δ , implying that the map is injective.

Since $\gamma_1 = (0 \ t_1 \ t_2 \ \dots \ t_n)$ and $\gamma_1 \circ \delta_1 = X_n$, we have that $t_n = 1$ and $t_{n-1} = n$.

Let $\gamma = (c_n)^{-1} \circ (\lambda_n)^{-1} \circ \gamma_1 \circ c_n$. Then,

$$\begin{aligned} \gamma &= (c_n)^{-1} \circ (\lambda_n)^{-1} \circ (0 \quad t_1 \quad \dots \quad t_{n-2} \quad n \\ &\quad 1) \circ c_n = (c_n)^{-1} \circ (0 \quad t_1 \quad \dots \\ &\quad t_{n-2})(n)(1) \circ c_n. \end{aligned}$$

Since conjugation preserves cycle structure, γ is an $(n-1)$ -cycle.

Finally,

$$\begin{aligned} \gamma \circ \delta &= (c_{n-1} \circ \lambda_{n-1} \circ \gamma_1 \circ c_n) \circ (c_{n-1} \circ \delta_1 \circ \lambda_{n-1} \circ c_n) \\ &= c_{n-1} \circ \lambda_{n-1} \circ X_n \circ \lambda_{n-1} \circ c_n = X_{n-2}. \end{aligned}$$

We have shown that for each factorization $\gamma_1 \circ \delta_1$ in B_1 , there are unique corresponding $(n-1)$ -cycles γ and δ such that $\gamma \circ \delta = X_{n-2}$. It follows that $\gamma \circ \delta \in A$, and this completes the proof. $\square$

Since the injective maps defined in the proof of Lemma 3.8 are merely inverses of those defined in the proof of Lemma 3.6, these maps in fact serve as bijective maps between the sets A and B_1 . It follows that $|A| = |B_1|$. The next lemma will function to show that $|B_1| = \dots = |B_{n-1}|$.

Lemma 3.9.

(1) For every $1 \leq i \leq n - 2$, there is an injection from B_i to B_{i+1} .

(2) There is an injection from B_{n-1} to B_1 .

In other words, $B_1 \rightarrow B_2 \rightarrow \cdots \rightarrow B_{n-1} \rightarrow B_1$, where each $\rightarrow$ indicates an injective map.

Proof. We prove the two statements separately.

(1). Let i satisfy $1 \leq i \leq n - 2$. Let γ_i and δ_i be $(n + 1)$ -cycles, where δ_i is of the form $(0 \ n \ i \ \cdots)$, and where $\gamma_i \circ \delta_i = X_n$. Let r_n denote the cycle $(2 \ 1 \ n)$. Define

$$\gamma_{i+1} = r_n \circ c_n \circ \gamma_i \circ (c_n)^{-1}, \delta_{i+1} = c_n \circ$$

$$\delta_i \circ (c_n)^{-1}.$$

It suffices to show that γ_{i+1} and δ_{i+1} are $(n + 1)$ -cycles, that δ_{i+1} is of the form $(0 \ n \ i + 1 \ \cdots)$, and that $\gamma_{i+1} \circ \delta_{i+1} = X_n$.

Since conjugation preserves the cycle structure of a permutation, both $c_n \circ \gamma_i \circ (c_n)^{-1}$ and $\delta_{i+1} = c_n \circ \delta_i \circ (c_n)^{-1}$ are $(n + 1)$ -cycles. One can also check that composition with r_n does not affect the cycle structure of $c_n \circ \gamma_i \circ (c_n)^{-1}$, meaning γ_{i+1} is also an $(n + 1)$ -cycle.

Next, observe that

$$\delta_{i+1}(0) = c_n(\delta_i(0)) = c_n(n) = n$$

and

$$\delta_{i+1}(n) = c_n(\delta_i(n)) = c_n(i) = i + 1.$$

Therefore, δ_{i+1} is of the form $(0 \ n \ i + 1 \ \cdots)$.

Finally,

$$\begin{aligned} \gamma_{i+1} \circ \delta_{i+1} &= (r_n \circ c_n \circ \gamma_i \circ (c_n)^{-1}) \circ (c_n \circ \delta_i \circ (c_n)^{-1}) \\ &= r_n \circ c_n \circ X_n \circ (c_n)^{-1} \\ &= r_n \circ (0 \quad 2 \quad 3 \quad \cdots \quad n-1 \quad 1 \quad n) \\ &= X_n. \end{aligned}$$

(2). Statement (2) follows from the observations that $\delta_1 \neq \delta_i \neq \delta_n$ for all $1 < i < n$, and that $\delta_n = \delta_1$. The latter observation follows directly from the fact that the order of c_n in the group of permutations is $n - 1$. $\square$

Example 3.7 (Continued). One can check that under the $\delta_i \rightarrow \delta_{i+1}$ map defined in the proof of Lemma 3.9, we get

$$\begin{aligned} \delta_1 = (0 \quad 6 \quad 1 \quad 5 \quad 4 \quad 3 \quad 2) &\rightarrow (0 \quad 6 \quad 2 \quad 1 \quad 5 \quad 4 \quad 3) \rightarrow (0 \quad 6 \quad 3 \quad 2 \quad 1 \quad 5 \quad 4) \\ &\rightarrow (0 \quad 6 \quad 4 \quad 3 \quad 2 \quad 1 \quad 5) \rightarrow \cdots \rightarrow (0 \quad 6 \quad 5 \quad 4 \quad 3 \quad 2 \quad 1) \\ &\rightarrow (0 \quad 6 \quad 1 \quad 5 \quad 4 \quad 3 \quad 2) = \delta_n. \end{aligned}$$

We now prove the main result of this section, Theorem 3.3.

Proof of Theorem 3.3. Since $X_n = Y_{\pi}^{-1} \circ C_{\pi}$, we count the factorizations of X_n into two $(n+1)$ -cycles where the second factor has the form $(0 \ n \ \cdots)$. This is the sum $\sum_{1 \leq i \leq n-1} |B_i|$. By Lemma 3.5, there are $\frac{2(n-2)!}{n}$ factorizations of X_{n-2} into two $(n-1)$ -cycles. In other words, $|A| = \frac{2(n-2)!}{n}$. By Lemmas 3.6, 3.8, and 3.9, we have that $|A| = |B_1| = \cdots = |B_{n-1}|$. It follows that for each even n , the number of permutations in S_n with strategic pile size $n-1$ is

$$\sum_{1 \leq i \leq n-1} |B_i| = (n-1)|A| = \frac{2(n-1)!}{n},$$

as desired. □

3.2. Maximum size strategic piles for odd values of n

We now prove an analog of Theorem 3.3 to address the case that n is odd.

Theorem 3.10. *For each odd number n , the number of permutations in S_n with strategic pile size $n-2$ is $2(n-2)!$.*

A permutation $\pi = [a_1 a_2 \cdots a_n]$ is said to have an *adjacency* if there is an index $i < n$ such that $a_{i+1} = a_i + 1$.

Lemma 3.11. *Let $n > 1$ be an odd number and let π be an element of S_n . If π has a strategic pile of cardinality $n-2$, then π has a single adjacency.*

Proof. Let $\pi = [a_1 a_2 \cdots a_n]$. We have that

$$C_{\pi} = (0 \quad a_n \quad \cdots \quad a_1) \circ (0 \quad 1 \quad \cdots n),$$

which is an even permutation. It follows from Lemma 3.1 that C_π is not a single cycle. Since π has a strategic pile of cardinality $n - 2$, we have that C_π is of the form

$$C_\pi = (0 \quad n \quad c_1 \quad \cdots \quad c_{n-2}) \circ (x).$$

The singleton cycle (x) comes about on account of the following configuration in the computation of C_π :

$$(\cdots \quad x+1 \quad x \quad \cdots) \circ (0 \quad 1 \quad \cdots \quad x \quad x+1 \quad \cdots \quad n).$$

Thus, in π we have that for some i , $a_i = x$ and $a_{i+1} = x + 1$. □

When a permutation π in S_n has a single adjacency, it can be projected to a unique corresponding permutation $P(\pi)$ in S_{n-1} which has no adjacencies, as follows: Let $\pi = [a_1 \ a_2 \ \cdots \ a_i \ a_{i+1} \ \cdots \ a_n] \in S_n$ have the single adjacency $a_{i+1} = a_i + 1$. We define $P(\pi)$ by removing the second element of the adjacency and reducing all larger elements by one. More precisely, $P(\pi) = [a'_1 \ a'_2 \ \cdots \ a'_{n-1}]$, where

$$\text{and } a_j \leq a_i, \quad a'_j = \begin{cases} a_j - 1 & \text{if } j \leq i \\ a_{j+1} - 1 & \text{if } n > j > i \text{ and } a_{j+1} \geq a_{i+1} \\ a_{j+1} & \text{if } j \leq i \text{ and } a_j > a_i, \\ & n > j > i \text{ and } a_{j+1} < a_{i+1} \end{cases},$$

Example 3.12. The permutation $\pi = [2 \ 3 \ 6 \ 1 \ 5 \ 4]$ has one adjacency. $P(\pi) = [2 \ 5 \ 1 \ 4 \ 3]$ has no adjacencies. Observe that there are five different elements of S_6 , each with a single adjacency, that give rise in this way to $[2 \ 5 \ 1 \ 4 \ 3]$, namely: $[2 \ \underline{3} \ 6 \ 1 \ 5 \ 4]$, $[2 \ 5 \ \underline{6} \ 1 \ 4 \ 3]$, $[3 \ 6 \ 1 \ \underline{2} \ 5 \ 4]$, $[2 \ 6 \ 1 \ 4 \ \underline{5} \ 3]$, and $[2 \ 6 \ 1 \ 5 \ 3 \ \underline{4}]$.

Remark 3.13. If $n > 1$ is an odd number and $\pi \in S_n$ is a permutation with a strategic pile of cardinality $n - 2$, then $P(\pi) \in S_{n-1}$ is a permutation with a strategic pile of cardinality $n - 2$.

Conversely, if we are given a permutation $\mu \in S_{n-1}$ which has no adjacencies, say $[\]$, and any position i , we can construct a unique permutation $(\mu, i) = [\]_1 \cdots [\]_n$ in S_n which has a single adjacency, and for which $P(E(\mu, i)) = \mu$: Namely, define a_{i+1} to be $a'_i + 1$; for $j < i$ define $a_j = a'_j + 1$ if $a_i < a_j$, and $a_j = a'_j$ otherwise; for $j > i$ define $a_{j+1} = a'_j$ if $a'_j < a'_i$ and $a_{j+1} = a'_j + 1$ otherwise.

Remark 3.14. If $n > 1$ is an odd number and $\mu \in S_{n-1}$ is a permutation with a strategic pile of cardinality $n-2$, and if $i \leq n-1$, then $E(\mu, i) \in S_n$ is a permutation with a strategic pile of cardinality $n-2$.

With these facts at our disposal we now prove Theorem 3.10.

Proof of Theorem 3.10. By Remark 3.14 each permutation $\mu \in S_{n-1}$ with full strategic pile produces $n-1$ permutations $\pi_i = E(\mu, i)$ for $i \leq n-1$ in S_n with strategic pile of size $n-2$. Thus by Theorem 3.3 there are at least $(n-1) \cdot \frac{2(n-2)!}{n-1} = 2(n-2)!$ elements of S_n with strategic pile of size $n-2$. Conversely, by Lemma 3.11 each element of S_n that has a strategic pile of size $n-2$ arises in this way. $\square$

4. Strategic Piles of Size k

Having quantified the number of permutations with maximum size strategic piles, we next produce an analogous quantification for permutations with strategic piles of arbitrary size. Before stating the main result of this section, we first establish terminology and structural properties of permutations with strategic piles of size k .

4.1. Structure of permutations with strategic pile of size k

Proposition 4.1. For a permutation π in S_n , $SP^*(\pi) = (b_1, b_2, \dots, b_k)$ if and only if the following are true:

- (1) $\pi(1) = b_k + 1$.
- (2) $\pi(n) = b_1$.
- (3) For all $j \in \{2, 3, \dots, k-1\}$, the element b_j appears to the immediate left of $b_{j-1} + 1$ in π (when written in one-line notation).

Proof. First note that $C_\pi(b_k) = 0$ if and only if $Y_\pi(b_k + 1) = 0$, since $C_\pi(b_k) = Y_\pi(X(b_k)) = Y_\pi(b_k + 1)$. Also, by definition, $Y_\pi(b_k + 1) = 0$ if and only if $\pi(1) = b_k + 1$. Therefore, $C_\pi(b_k) = 0$ if and only if $\pi(1) = b_k + 1$.

Second, $C_\pi(n) = b_1$ if and only if $Y_\pi(0) = b_1$, since $C_\pi(n) = Y_\pi(X(n)) = Y_\pi(0)$. Also, by definition, $Y_\pi(0) = b_1$ if and only if $\pi(n) = b_1$. Therefore, $C_\pi(n) = b_1$ if and only if $\pi(n) = b_1$.

Finally, for $j \in \{2, 3, \dots, k-1\}$, $C_\pi(b_{j-1}) = b_j$ if and only if $Y_\pi(b_{j-1} + 1) = b_j$, since $C_\pi(b_{j-1}) = Y_\pi(X(b_{j-1})) = Y_\pi(b_{j-1} + 1)$. Also, by definition, $Y_\pi(b_{j-1} + 1) = b_j$ if and only if b_j immediately

precedes $b_{j-1} + 1$ in π . Therefore, $C_\pi(b_{j-1}) = b_j$ if and only if b_j appears immediately to the left of $b_{j-1} + 1$ in π .

Since $SP^*(\pi) = (b_1, b_2, \dots, b_k)$ if and only if $C_\pi(b_k) = 0$, $C_\pi(n) = b_1$, and $C_\pi(b_{j-1}) = b_j$ for all $j \in \{2, 3, \dots, k-1\}$, our proposition holds. $\square$

With b_j denoting the j th element of the ordered strategic pile of a permutation π , adjacent entries of the form $b_j b_{j-1} + 1$ in π are called a *pair*. Viewing subscripts modulo k , we also consider $b_1 b_k + 1$ a pair. In general, a permutation π with $SP^*(\pi) = (b_1, b_2, \dots, b_k)$ has the following form in terms of its pairs:

$$[b_k + 1 \cdots b_{x_1} b_{x_1-1} + 1 \cdots b_{x_2} b_{x_2-1} + 1 \cdots b_{x_{k-1}} b_{x_{k-1}-1} + 1 \cdots b_1]. \quad (4.1)$$

Definition 4.2. The ordered list

$$\sigma_\pi = (b_{x_1}, b_{x_2}, \dots, b_{x_{k-1}}, b_1),$$

consisting of the first member of each pair, in the order of occurrence in π , is said to be the *ordered pair list* of π .

Since b_1 is the final entry of a permutation π with a nonempty strategic pile, b_1 is always the terminating member of the ordered pair list σ_π .

Example 4.3. The permutation $\pi = [6 \ 4 \ 5 \ 8 \ 7 \ 2 \ 3 \ 1]$ has strategic pile $SP(\pi) = \{1, 5, 7\}$, and $SP^*(\pi) = (1, 7, 5) = (b_1, b_2, b_3)$. Therefore, $\pi =$

$[6 \ 4 \ b_3 \ 8 \ b_2 \ 2 \ 3 \ b_1] = [b_3 + 1 \ 4 \ b_3 \ b_2 + 1 \ b_2 \ b_1 + 1 \ 3 \ b_1]$, as suggested by Proposition 4.1. This gives that $\sigma_\pi = (b_3, b_2, b_1)$.

In Definition 4.2 we defined the ordered pair list with respect to a specified permutation π . Note, however, that we can instead define an ordered pair list independently of a specific permutation. Using this interpretation, any permutation where the x_i th strategic pile element leads the i th pair for all $1 \leq i \leq k-1$ will be said to have the ordered pair list $\sigma = (b_{x_1}, b_{x_2}, \dots, b_{x_{k-1}}, b_1)$.

Example 4.4. Consider the ordered pair list $\sigma = (b_2, b_3, b_1)$, defined independently of a specific permutation. Any permutation with $SP^* = (b_1, b_2, b_3)$ that is of the form

$$[b_3 + 1 \quad \cdots \quad b_2 \quad b_1 + 1 \quad \cdots \quad b_3 \quad b_2 + 1 \quad \cdots \quad b_1]$$

will have ordered pair list σ . In particular, the permutation $\pi = [2 \ 3 \ 6 \ 1 \ 4 \ 5]$ has $SP^* = (5, 3, 1)$ and thus $\sigma_\pi = (3, 1, 5) = (b_2, b_3, b_1) = \sigma$. Similarly, the permutation $\nu = [4 \ 1 \ 6 \ 3 \ 2 \ 5]$ has $SP^* = (5, 1, 3)$ and thus $\sigma_\nu = (1, 3, 5) = (b_2, b_3, b_1) = \sigma$.

As Example 4.6 will illustrate, for subsequent pairs $b_{x_i} b_{x_i-1} + 1$ and $b_{x_{i+1}} b_{x_{i+1}-1} + 1$ of a permutation π it may happen that $b_{x_i-1} + 1 = b_{x_{i+1}}$, in which case b_{x_i} and $b_{x_{i+1}}$ are consecutive entries of π . As these adjacencies will be of central importance in the proof of Theorem 4.10, we formalize their definition as follows.

Definition 4.5. An adjacency of strategic pile members b_{x_i} and $b_{x_{i+1}}$ in π is said to be a *merge between b_{x_i} and $b_{x_{i+1}}$* in π . Such a merge will be denoted $b_{x_i} b_{x_{i+1}}$.

Example 4.6. The permutation $\pi = [5 \ 4 \ 6 \ 3 \ 2 \ 1]$ has strategic pile $\text{SP}(\pi) = \{1, 3, 4, 5\}$, and $\text{SP}^*(\pi) = (1, 3, 5, 4) = (b_1, b_2, b_3, b_4)$. Moreover, $\sigma_\pi =$

(b_3, b_4, b_2, b_1) since π has the form $[b_3 \ b_4 \ 6 \ b_2 \ 2 \ b_1]$. The strategic pile members b_3 and b_4 are adjacent in π , and thus there is a merge in π . Since we are also considering $b_1 b_4 + 1$ a pair in π , $b_1 b_3$ is also ruled a merge in π .

When considering strategic piles of size k , we refer to an arrangement of the strategic pile variables b_i and $b_i + 1$ for $1 \leq i \leq k$ as a *strategic pile variable arrangement* if the arrangement satisfies the properties described in Proposition 4.1. All possible strategic pile variable arrangements can be obtained by shifting and merging pairs within the possible frameworks of the form (4.1).

Example 4.7. The following are five of the possible strategic pile variable arrangements for permutations in S_7 with $\text{SP}^* = (b_1, b_2, b_3)$ and ordered pair list $\sigma = (b_2, b_3, b_1)$, where the ___'s can be filled in by any remaining permutation elements:

- (1) $[b_3 + 1 \ ___ \ b_2 \ b_1 + 1 \ b_3 \ b_2 + 1 \ b_1]$ (no merges),
- (2) $[b_3 + 1 \ b_2 \ b_1 + 1 \ ___ \ b_3 \ b_2 + 1 \ b_1]$ (no merges),
- (3) $[b_2 \ b_1 + 1 \ ___ \ b_3 \ b_2 + 1 \ b_1]$ (merge $b_1 b_2$),
- (4) $[b_2 \ b_1 + 1 \ b_3 \ b_2 + 1 \ ___ \ ___ \ b_1]$ (merge $b_1 b_2$),
- (5) $[b_2 \ b_1 + 1 \ ___ \ ___ \ ___ \ b_3 \ b_1]$ (merges $b_3 b_1$ and $b_1 b_2$).

The above definitions and structural properties regarding permutations with strategic piles of size k suggest an approach for quantifying such permutations. Since a permutation has strategic pile size k if and only if it takes the form described in Proposition 4.1, we start by counting the number of strategic pile variable arrangements. To this end, we define *merge numbers*.

Definition 4.8. Consider the set of permutations $\pi \in S_n$ with $\text{SP}^*(\pi) = (b_1, \dots, b_k)$. Given $\ell \geq 0$, the symbol c_k denotes the number of ways to choose an ordered pair list σ_π along with merges. The number c_k is said to be a *merge number*.

Example 4.9. For permutations with $SP^* = (b_1, b_2, b_3)$, the only possible ordered pair lists are (b_2, b_3, b_1) and (b_3, b_2, b_1) , which correspond to the following permutation structures:

$$(1) [b_3 + 1 \cdots b_2 b_1 + 1 \cdots b_3 b_2 + 1 \cdots b_1],$$

$$(2) [b_3 + 1 \cdots b_3 b_2 + 1 \cdots b_2 b_1 + 1 \cdots b_1].$$

In the first form, each of the merges $b_2 b_3$, $b_3 b_1$, and $b_1 b_2$ are possible, so there are three ways to create a single merge with this ordered pair list. In the second form, a merge cannot occur at all, since it would require that $b_i + 1 = b_i$, which is impossible. Therefore, $c_{3,1} = 1 \cdot 3 + 1 \cdot 0 = 3$.

We are now ready to state the main result of this section.

4.2. Main result

Theorem 4.10. For $1 \leq k \leq n - 1$ and even n , or $1 \leq k \leq n - 2$ and odd n , the number of permutations in S_n with strategic pile of size k is

$$(n - k)! \sum_{i=0}^{\infty} c_{k,i} \binom{n - (k + 1)}{k - (i + 1)}.$$

As there is a limit on the number of merges that can occur in a permutation, each merge number $c_{k,i}$ will be zero for all i above a certain value. We leave determining this maximum number of merges, as well as the general method for computing merge numbers, to Sec. 5. To prove Theorem 4.10, we will

- use merge numbers to determine the number of strategic pile variable arrangements (see Lemma 4.12 and Corollary 4.13) and
- determine the number of ways to assign numerical values to the resulting variable arrangements (see Lemma 4.14).

Assuming we can compute each merge number $c_{k,i}$, we can suppose we are given a framework comprised of an ordered pair list and a set of merges. To quantify the possible strategic pile variable arrangements, we are left to account for how this framework can shift within n positions. To this end, we develop terminology to refer to the components of this framework.

Example 4.11. Consider an ordered pair list $\sigma = (b_{x_1}, b_{x_2}, \dots, b_{x_{k-1}}, b_1)$, which by Lemma

4.1 yields a permutation of the form $[b_k + 1 \quad \cdots \quad b_{x_1} \quad b_{x_1-1} + 1 \cdots$
 $b_{x_2} \quad b_{x_2-1} + 1 \quad \underline{\hspace{1cm}} \quad \underline{\hspace{1cm}}]$

$$\cdots \quad b_{x_{k-1}} \quad \underline{b_{x_{k-1}-1} + 1} \quad \cdots \quad b_1].$$

After a merge, say between b_{x_1} and b_{x_2} , we get

$$[b_k + 1 \quad \cdots \quad b_{x_1} \quad \underline{b_{x_2}} \quad \underline{b_{x_2}-1} + 1 \quad \cdots \quad b_{x_{k-1}} \quad \underline{b_{x_{k-1}-1} + 1} \quad \cdots \quad b_1]$$

In Example 4.11, observe that it may not be intuitive to call $b_{x_1} b_{x_2} b_{x_2-1} + 1$ a pair; we use the term *grouping* to refer to pairs as well as any set of pairs joined by merges.

Recall that $b_k + 1$ and b_1 are always in the first and last positions of a permutation, respectively. Moreover, observe that the position of each underlined element in Example 4.11 is determined by the placement of the leftmost element in its grouping. We call both of these types of elements *determined*.

In a permutation with strategic pile of size k with no merges, there are $k+1$ determined elements (i.e., b_1 , $b_k + 1$, and $b_{j-1} + 1$ for $2 \leq j \leq k$). Furthermore, observe that “merging” groupings do not affect the total number of determined elements, since a merge has the effect of equating a determined element with an undetermined element. In Example 4.11, the merge between b_{x_1} and b_{x_2} equates $b_{x_1-1} + 1$ (a determined element) with b_{x_2} (an undetermined element), making a grouping with two determined elements, the same total number that the pairs $b_{x_1} b_{x_1-1} + 1$ and $b_{x_2} b_{x_2-1} + 1$ had to begin with. Therefore, any grouping arrangement, despite the number of merges, will have $k + 1$ determined elements.

Lemma 4.12. *Given an ordered pair list $\sigma = (b_{x_1}, \dots, b_{x_k})$ and a set of i merges, there are*

$$\binom{n - (k + 1)}{(k - 1) - i}$$

ways to place the resulting groupings within a permutation of length n .

Proof. Recall that in a permutation with strategic pile size k , there are always $k + 1$ determined elements. For each determined element, we set aside one space in the permutation. This leaves $n - (k + 1)$ unoccupied spaces in which to place the groupings. Since the leftmost variable of each grouping is the only undetermined variable in the grouping, we must only place these $(k-1)-i$ undetermined variables, and the placement of all other variables follows. Because there are $n-(k+1)$ spaces in which to place these undetermined variables, $\binom{n-(k+1)}{(k-1)-i}$ represents the number of ways to place the groupings. $\square$

Corollary 4.13. *The number of strategic pile variable arrangements in the set of permutations in S_n with strategic piles of size k and i merges is*

$$c_{k,i} \cdot \binom{n - (k + 1)}{k - i - 1}.$$

Proof. By definition of the merge number $c_{k,i}$, there are $c_{k,i}$ ways to choose an ordered pair list $\sigma = (b_{x_1}, \dots, b_{x_k})$ along with i merges. Each fixed ordered pair list and set of merges defines a framework of groupings, which by Lemma 4.12 can be placed $\binom{n-(k+1)}{k-i-1}$ different ways among n positions. Thus, there are a total of $c_{k,i} \binom{n-(k+1)}{k-i-1}$ different strategic pile variable arrangements. $\square$

We now determine the number of ways to assign a numerical value to each variable and to the remaining elements of the permutation.

Lemma 4.14. *Given a fixed strategic pile variable arrangement (i.e., an ordered pair list σ of strategic pile elements $\{b_1, \dots, b_k\}$, a set of merges, and a fixed set of grouping positions), there are $(n-k)!$ permutations in S_n with that arrangement.*

Proof. Given a strategic pile variable arrangement, the only thing left to do is assign values to the variables comprising the permutation. For each $b_j \in \text{SP}$, there is some variable $b_j + 1$, whose value follows immediately from a value assignment of b_j . Therefore, only $n-k$ values need to be assigned, and there are $(n-k)!$ possible assignments. $\square$

We are now ready to prove our main result of this section.

Proof of Theorem 4.10. By Corollary 4.13, there are $c_{k,i} \binom{n-(k+1)}{k-i-1}$ strategic pile variable arrangements in the set of permutations in S_n with strategic piles of size k and i merges. Summing over the number of merges i , we get that the total number of strategic pile variable arrangements for permutations in S_n with strategic piles of size k is

$$\sum_{i=0}^{\infty} c_{k,i} \binom{n-(k+1)}{k-i-1}.$$

We have by Lemma 4.14 that there are $(n-k)!$ permutations corresponding to each strategic pile variable arrangement. It follows that there are

$$(n-k)! \sum_{i=0}^{\infty} c_{k,i} \binom{n-(k+1)}{k-i-1}$$

permutations in S_n with strategic pile size k . $\square$

5. Determining the Values of Merge Numbers

As mentioned in Sec. 4, there is a limit to the number of merges that can occur in a permutation with strategic pile of size k . Thus, there exists an i such that $c_{k,i'} = 0$

Table 1. Merge numbers found using *ad hoc* methods.

k	1	2	3	4	5	6	..	k					..	OEIS
							.						.	
ck	0	1	2	3	4	5!								A0001
$,0$	!	!	!	!	!	57								42
ck			3	1	9	6								A1307
$,1$			3	6	0	111	0!	1!	2!	3!	4!	5!	44	
ck				1	13	6							$k(k -$	
$,2$				6	0		?							
ck					8	108							?	
$,3$					0	0								
ck					9	54							?	
$,4$					0	0								
.														
.														
.														
$ck,$														
i														

Table 2. Known formulas for the number of permutations with strategic piles of size k .

k	Number of elements of S_n with strategic piles of size k	OEIS
1	$(n-1)! \left[\binom{n-2}{0} 0! \right]$	A000142
2	$(n-2)! \left[\binom{n-3}{1} 1! \right]$	A062119
3	$(n-3)! \left[\binom{n-4}{2} 2! + \binom{n-4}{0} 3 + \binom{n-4}{0} 3 \right]$	A267323
4	$(n-4)! \left[\binom{n-5}{3} 3! + \binom{n-5}{2} 16 + \binom{n-5}{1} 16 \right]$	A267324
5	$(n-5)! \left[\binom{n-6}{4} 4! + \binom{n-6}{3} 90 + \binom{n-6}{2} 130 + \binom{n-6}{1} 80 + \binom{n-6}{0} 90 \right]$	A267391
6	$(n-6)! \left[\binom{n-7}{5} 5! + \binom{n-7}{4} 576 + \binom{n-7}{3} 1116 + \binom{n-7}{2} 1080 + \binom{n-7}{1} 540 \right]$	A281259
$\cdot$		
$\cdot$		
$\cdot$		
k	$(n-k)! \left[\binom{n-(k+1)}{k-1} (k-1)! + \binom{n-(k+1)}{k-2} c_{k,1} + \dots + \binom{n-(k+1)}{1} c_{k,k-2} \right] (k \text{ odd})$	

for all $i' \geq i$. In this section, we determine this i and derive an algorithm for computing merge numbers. As described later this section, determining the efficiency of this algorithm is dependent on the solutions of certain open problems. However, we can explicitly compute merge numbers in some limited cases (see Table 1). Using these merge numbers, Theorem 4.10 gives the formulas given in Table 2.

We now discuss how to compute merge numbers in general. First, however, we define two graph theoretic tools, which will be useful for accomplishing both of the aforementioned goals.

5.1. Merge graphs and τ -graphs

As in Example 4.9, the ordered pair list determines the set of possible merges for permutations with that ordered pair list. As a result, it will often be of use to classify permutations based on ordered pair list.

Definition 5.1. Let σ be an ordered pair list. Define $T_\sigma := \{\pi \in S_n \mid \sigma_\pi = \sigma\}$.

Moreover, to count the number of ways to choose an ordered pair list along with merges (i.e., to compute merge numbers), it is crucial that we are first able to determine the set of allowable merges corresponding to an ordered pair list. To this end, we define the following.

Definition 5.2. Consider the set of permutations in S_n with ordered strategic pile $SP^*(\pi) = (b_1, b_2, \dots, b_k)$ and ordered pair list $\sigma = (b_{x_1}, b_{x_2}, \dots, b_{x_{k-1}}, b_1)$. Define

- $\psi := (b_1 b_2 \dots b_k)$,
- $\sigma^* := (b_{x_1} \ b_{x_2} \ \dots \ b_{x_{k-1}} \ b_1$
- $\tau_\sigma := \sigma^* \circ \psi$.

), and

Observe that there exists a permutation $\pi \in T_\sigma$ with $b_i + 1 = b_j$ if and only if $\tau_\sigma(b_i) = b_j$. In other words, τ_σ describes exactly the allowable merges for the set of permutations T_σ . It will often be useful for us to encode this information graphically.

Definition 5.3. The τ -graph corresponding to an ordered pair list σ is an at most in-degree one, out-degree one directed graph $T_\sigma = (V, E)$, where $V = \{b_1, b_2, \dots, b_k\}$ and $(b_i, b_j) \in E$ if and only if $b_i + 1 = b_j$ in some permutation in T_σ .

Note that an edge (b_i, b_j) in a τ -graph T_σ corresponds to the equality $b_i + 1 = b_j$, and not to the merge $b_i b_j$. Rather, the edge (b_i, b_j) represents the existence of a merge $b_{i+1} b_j$ in some permutation $\pi \in T_\sigma$. Furthermore, it may be useful to note that τ -graphs are comprised completely of cycles and isolated vertices, and that each cycle in the τ -graph T_σ corresponds to a cyclic factor of the cycle permutation τ_σ .

Example 5.4. Consider the set of permutations T_σ corresponding to the ordered pair list $\sigma = (b_1, b_6, b_7, b_5, b_2, b_4, b_3)$. Then

$$\begin{aligned} \tau_\sigma &= \sigma^* \circ \psi = (b_1 \ b_6 \ b_7 \ b_5 \ b_2 \ b_4 \ b_3) \circ (b_1 \ b_2 \ b_3 \ b_4 \ b_5 \ b_6 \ b_7) \\ &= (b_1 \ b_4 \ b_2)(b_5 \ b_6 \ b_7)(b_3), \end{aligned}$$

which indicates that there are two cycles in T_σ formed by the edges $\{(b_1, b_4), (b_4, b_2), (b_2, b_1)\}$ and $\{(b_5, b_7), (b_7, b_6), (b_6, b_5)\}$. Figure 1 shows this graph.

We now define a second graph theoretic tool, the *merge graph*, which will be similar to the τ -graph, but will correspond to a specific permutation rather than to an ordered pair list.




Fig. 1. The τ -graph T_σ corresponding to $\sigma = (b_5, b_6, b_4, b_2, b_3, b_7, b_1)$.

Definition 5.5. The *merge graph* of π is an in-degree at most one, out-degree at most one directed graph $M_\pi = (V, E)$, where $V = \{b_1, b_2, \dots, b_k\}$ and $(b_i, b_j) \in E$ if and only if $b_i + 1 = b_j$ in the permutation π .

Observe that if π is a permutation with ordered pair list σ , then the merge graph M_π , is an edge subgraph of the τ -graph T_σ . Moreover, Lemma 5.6 will show that M_π is a proper subgraph of T_σ , and that any acyclic edge subgraph of T_σ is a merge graph.

Lemma 5.6. *A graph is a merge graph if and only if it is an acyclic edge subgraph of a τ -graph.*

Proof. Suppose on the contrary that π is a permutation such that its merge graph M_π contains an ℓ -cycle between consecutive vertices $v_1, v_2, \dots, v_\ell$ for some $\ell > 0$, and let b_{x_i} be the strategic pile element corresponding to the vertex v_i for $1 \leq i \leq \ell$. Then, by definition of M_π , $b_{x_1} + 1 = b_{x_2} = b_{x_3} - 1$, so $b_{x_1} = b_{x_3} - 2$. Continuing in this manner, we see that $b_{x_1} = b_{x_\ell} - (\ell - x_1 - x_\ell + 1)$. However, as the directed edges among these vertices form a cycle, we also have that $b_{x_1} = b_{x_\ell} + 1$. Since $\ell > 0$, this is impossible.

Conversely, given an acyclic edge subgraph of a τ -graph, one can construct an ordered pair list with the corresponding merges. This can be done because our only constraint on merges that can occur is cyclic relationship between strategic pile variables. $\square$

Example 5.7. For $\pi = [5\ 4\ 6\ 3\ 2\ 1]$ we have $\text{SP}^*(\pi) = (1, 3, 5, 4) = (b_1, b_2, b_3, b_4)$ and ordered pair list $\sigma = (5, 4, 3, 1) = (b_3, b_4, b_2, b_1)$. Thus, $\psi = (1\ 3\ 5\ 4) = (b_1\ b_2\ b_3\ b_4)$ and $\sigma^* = (5\ 4\ 3\ 1) = (b_3\ b_4\ b_2\ b_1)$. Note that $\tau_\sigma = \sigma^* \circ \psi = (b_3\ b_4\ b_2\ b_1) \circ (b_1\ b_2\ b_3\ b_4) = (b_1) \circ (b_2\ b_4\ b_3)$.

The cycle $(b_2\ b_4\ b_3)$ in the cycle decomposition of the permutation τ_σ indicates that the τ -graph T_σ will have edges (b_2, b_4) , (b_4, b_3) , and (b_3, b_2) . Lemma 5.6 implies that any

proper subset of these three edges can occur in M_π . Indeed, $b_2 + 1 = b_4$ and $b_4 + 1 = b_3$ in π , while $b_3 + 1 \neq b_2$ in π . The merge graph M_π and τ -graph T_σ are depicted in Fig. 2.

5.2. Maximum number of merges

We will now determine the maximum number of merges that can occur in a permutation in S_n with strategic pile size k . This will function to show that $c_{k,i} = 0$ for all i greater than a certain value. We start with the following observation.

Remark 5.8. Let $\sigma = (b_{x1}, b_{x2}, \dots, b_{xk-1}, b_1)$. Since $\tau_\sigma \in S_k$ is the composition of two cycles of the same length k , τ_σ is an even permutation. It follows that if k is even, then τ_σ cannot be a k -cycle.




Fig. 2. Merge graph of $\pi = [5\ 4\ 6\ 3\ 2\ 1]$ (solid) as a subgraph of the τ -graph corresponding to $\sigma\pi$ (solid and dashed).

This observation, in conjunction with our graph theoretic tools, will give us our desired result.

Lemma 5.9. Consider the set T of permutations with strategic piles of size k .

- (1) If k is odd, then the number of merges for any permutation in T is at most $k - 1$.
- (2) If k is even, then the number of merges for any permutation in T is at most $k - 2$.

Proof. Case 1: Let k be odd. Suppose $\pi \in S_n$ has ordered strategic pile $SP^*(\pi) = (b_1, \dots, b_k)$. Consider the merge graph M_π , which will have k vertices. By definition, M_π is at most in-degree one and out-degree one. Since Lemma 5.6 gives that M_π is acyclic, it follows that the number of edges in M_π does not exceed $k - 1$. Thus, there are at most $k - 1$ merges in π .

Case 2: Let k be even. By Remark 5.8, the cycle decomposition of τ does not contain a k -cycle, and thus the largest possible cycle in T is a $(k-1)$ -cycle. It follows that there are at most $k-2$ merges in any permutation $\alpha \in T$. $\square$

Lemma 5.9 tells us that $c_{k,i} = 0$ for large enough i , or in other words, that the number of permutations of n elements with strategic pile of size k can be written

$$(n-k)! \sum_{i=0}^t c_{k,i} \binom{n-(k+1)}{k-i-1},$$

where $t = k-1$ if k is odd and $t = k-2$ if k is even.

5.3. Merge number algorithm

In Sec. 5.2, we established that $c_{k,i} = 0$ for $i > k-1$ when k is odd and for $i > k-2$ when k is even. We now discuss how to compute the merge numbers corresponding to smaller i . We present Algorithm 5.10 for computing such merge numbers, prove its correctness, and discuss its complexity and what work still needs to be done in order to make this algorithm more efficient.

Algorithm 5.10. Merge Number Computation

Input: Integers k and ℓ where $\ell \leq k-1$ if k is odd and $\ell \leq k-2$ if k is even.

Output: The merge number $c_{k,\ell}$.

- (1) Consider all possible cycle structures for a τ -graph with k vertices.
- (2) For each of these cycle structures:
 - (a) Determine the number of ordered pair lists $\sigma = (b_{x_1}, b_{x_2}, \dots, b_{x_{k-1}}, b_1)$ which yield the given cycle structure.
 - (b) Multiply by the number of ways merges can be chosen from the given cycle structure.
- (3) Sum the results of the calculation for each cycle structure.

5.3.1. Correctness and complexity of Step 1

Step 1 of Algorithm 5.10 requires considering all possible cycle structures for a τ -graph with k vertices. Recall that a τ -graph consists only of cycles and isolated vertices. As a result, we can use the following notation to refer to the cycle structure of a τ -graph.

Notation 5.11. Let $[a_1, a_2, \dots, a_m]$ denote the cycle structure of a τ -graph T_σ , where $a_1 \geq a_2 \geq \dots \geq a_m > 0$, and where each a_i corresponds to the number of edges of a cycle in T_σ . We do not include isolated vertices in our cycle structure representation.

Since any τ -graph is in-degree at most one and out-degree at most one, a τ -graph on k vertices has at most k edges. As a result, the cycle structure $[a_1, a_2, \dots, a_m]$ corresponding to T_σ satisfies $\sum_{1 \leq i \leq m} a_i \leq k$. Therefore, to consider all possible cycle structures for a τ -graph with k vertices, it would suffice to consider all integer partitions of at most k . However, to speed up Step 1, we would like to be able to consider a smaller set of partitions.

To this end, note that a τ -graph T_σ on k vertices contains no self-loops, since this would imply that $b_i + 1 = b_i$ for some strategic pile element b_i . Moreover, note that the cycle structure of T_σ must have an even number of even parts, since τ_σ is an even permutation. To summarize:

Remark 5.12. Every τ -graph on k vertices has a cycle structure in the form of an integer partition $[a_1, a_2, \dots, a_m]$, with no parts of size one, with an even number of even parts, and with $\sum_{1 \leq i \leq m} a_i \leq k$.

Example 5.13. Consider the graph in Fig. 3, which has cycle structure $[3, 3]$. Notice that this cycle structure satisfies the conditions described in Remark 5.12, which are necessary conditions for a cycle structure to correspond to a τ -graph.




Fig.3.A τ -graph with cycle structure $[3, 3]$.

Indeed, this graph can be derived from the ordered pair list $\sigma = (b_5, b_3, b_4, b_2, b_6, b_1)$, and is thus a τ -graph.

We will not prove that every integer partition satisfying the conditions of Remark 5.12 corresponds to the cycle structure of a τ -graph on k vertices (i.e., the converse of Remark 5.12), since it will not affect the correctness of our algorithm. If it happens that we consider in Step 1 a partition that does not correspond to the cycle structure of a τ -graph on k vertices, Step 2(a) will yield a zero, so we will not be over-counting.

Unfortunately, the best way currently known to determine the set of integer partitions satisfying the properties of Remark 5.12 is the brute force method of checking every partition of every integer from 1 to k . Since the number of integer partitions of an integer n grows exponentially with n [7], Step 1 is inefficient for large k .

5.3.2. Correctness and complexity of Step 2(a)

Step 2(a) requires determining the number of ordered pair lists $\sigma = (b_{x_1}, b_{x_2}, \dots, b_{x_{k-1}}, b_1)$ which yield a given τ -graph cycle structure. As in Step 1, this can be done through brute force; namely, one can generate all $O(k!)$ possible ordered pair lists, and for each ordered pair list σ , can compute $\tau_\sigma = \sigma^* \circ \psi$ to determine whether τ_σ has the given cycle structure. Since each computation of τ_σ requires $O(k)$ time, Step 2(a) can be completed in $O(k \cdot k!)$ with this brute force method.

It is possible, however, that this step could be accomplished in polynomial time using a recursive formula. We will now derive such a formula, though a method for efficiently computing the base cases for this formula is currently unknown. Our derivation will involve understanding the relationship between τ -graphs on k vertices with cycle structure $[a_1, a_2, \dots, a_m]$ and τ -graphs on $k - 1$ vertices with the




Fig. 4. τ -graphs for Example 5.14. (a) Original τ -graph corresponding to the ordered pair list $(b_2, b_5, b_4, b_3, b_1)$. (b) Rotated τ -graph with b_5 as an isolated vertex. (c) τ -graph after removing b_5 .

same cycle structure. To build intuition for this relationship, let us consider an example.

Example 5.14. The aforementioned relationship will be established by rotating and removing vertices from τ -graphs. For example, consider the τ -graph corresponding to the ordered pair list $(b_2, b_5, b_4, b_3, b_1)$ (see Fig. 4(a)). In Lemma 5.15, we will prove that any rotation of a τ -graph is also a τ -graph. In particular, any rotation of the τ -graph in Fig. 4(a) is a τ -graph; Fig. 4(b) shows the rotation that is the τ -graph corresponding to the ordered pair list $(b_5, b_3, b_4, b_2, b_1)$. Note that this τ -graph has b_5 as an isolated vertex.

In a graph with b_k (in this case b_5) as an isolated vertex, removing b_k will give us another τ -graph; this is because when b_k is an isolated vertex, removing the vertex b_k corresponds to removing $b_k + 1 \dots b_k$ from the beginning of the pair ordering. This will leave $b_{k-1} + 1$ at the beginning of the pair ordering, which will yield a valid ordered pair list on $k-1$ strategic pile elements. Figure 4(c) shows the τ -graph corresponding to the ordered pair list (b_3, b_4, b_2, b_1) that occurs when b_5 is removed from our example τ -graph.

We will now formalize this idea. Let $X_{k,[a_1, a_2, \dots, a_m]}$ be the set of τ -graphs with k vertices and with cycle structure $[a_1, a_2, \dots, a_m]$. We are interested in finding a relationship between $|X_{k,[a_1, a_2, \dots, a_m]}|$ and $|X_{k-1,[a_1, a_2, \dots, a_m]}|$. We begin by showing that if a given τ -graph is in $X_{k,[a_1, a_2, \dots, a_m]}$, then so are all rotations of that graph.

Thus we get a group action on $X_{k,[a_1, a_2, \dots, a_m]}$.

Lemma 5.15. *Let $\sigma = (b_{x_1}, \dots, b_{x_{k-1}}, b_1)$ be an ordered pair list, and define ϕ :*

$$\mathbb{Z}_k \times X_{k,[a_1, \dots, a_m]} \rightarrow X_{k,[a_1, \dots, a_m]} \text{ as } \phi(i, \tau_\sigma) = \beta^i \circ \tau_\sigma \circ \beta^{-i},$$

where $\beta = (1 \ 2 \ 3 \ \dots \ k)$. Then ϕ is a group action.

Proof. Let $\sigma = (b_{x_1}, \dots, b_{x_{k-1}}, b_1)$ be an ordered pair list and suppose τ_σ has cycle structure $[a_1, \dots, a_m]$.

We first show that $\phi(i, \tau_\sigma) \in X_{k,[a_1, \dots, a_m]}$. Since conjugation preserves cycle structure, it is clear that $\phi(i, \tau_\sigma)$ will be a graph on k vertices with cycle structure $[a_1, \dots, a_m]$. We have left to show that $\phi(i, \tau_\sigma)$ corresponds to an ordered pair list σ (i.e., that $\varphi(i, \tau_\sigma) = \tau_{\sigma'}$).

Recall that $\tau_\sigma = \sigma^* \circ \psi$. Therefore, $\phi(i, \tau_\sigma) = \beta^i \circ \tau_\sigma \circ \beta^{-i} = \beta^i \circ \sigma^* \circ \psi \circ \beta^{-i} = \beta^i \circ \sigma^*$

$$\circ \beta^{-i} \circ \beta^i \circ \psi \circ \beta^{-i}.$$

Observe that $\beta^i \circ \psi \circ \beta^{-i} = \psi$. Furthermore, $\beta^i \circ \sigma^* \circ \beta^{-i} = \sigma'$ is a k -cycle containing the elements $\{b_1, \dots, b_k\}$, and thus represents an ordered pair list. As a result,

$\varphi(i, \tau_\sigma) = \psi \circ \sigma' = \tau_{\sigma'}$. It follows that $\phi(i, \tau_\sigma) \in X_{k,[a_1, \dots, a_m]}$, as desired. Finally, we check that ϕ satisfies the axioms of group actions. Clearly, $\beta^0 = \beta^k$ is the identity permutation,

and therefore $\phi(0, \tau_\sigma) = \tau_\sigma$. In addition, $\phi(i + j, \tau_\sigma) = \beta_{i+j} \circ \tau_\sigma \circ \beta_{-(i+j)} = \beta_i \beta_j \circ \tau_\sigma \circ \beta_{-j} \beta_{-i} = \phi(i, \phi(j, \tau_\sigma))$. $\square$

We can use Lemma 5.15 to give a process for deriving $X_{k,[a_1, a_2, \dots, a_m]}$ from $X_{k-1,[a_1, a_2, \dots, a_m]}$.

Lemma 5.16. *Let $G = (V, E)$ with $V = \{b_1, \dots, b_k\}$, and let $a_1, \dots, a_m \in \mathbb{Z}$ be such that $a_1 + a_2 + \dots + a_m < k$. Then $G \in X_{k,[a_1, a_2, \dots, a_m]}$ if and only if there exists some $G_r = (V_r, E_r) \in \text{orb}_{\mathbb{Z}_k}(G)$ such that b_k is an isolated vertex in G_r and $G_s := (V_r \setminus \{b_k\}, E_r) \in X_{k-1,[a_1, a_2, \dots, a_m]}$.*

Proof. Assume $G \in X_{k,[a_1, a_2, \dots, a_m]}$. Since $a_1 + a_2 + \dots + a_m < k$, there exists at least one isolated vertex in G . Therefore, some rotation of G has b_k as an isolated vertex. Let this rotation be G_r and let G_s be defined as in the lemma statement. We have left to show that G_s is a τ -graph. Since G_r has b_k as an isolated vertex, any permutation with ordered pair list corresponding to G_r must be of the form

$$[b_k + 1 \quad \dots \quad b_k \quad b_{k-1} + 1 \quad \dots \quad b_{\ell_1} \quad b_{\ell_2-1} + 1 \quad \dots \quad b_{\ell_{k-1}} \\ b_{\ell_{k-2}-1} + 1 \quad \dots \quad b_1].$$

Removing the $b_k b_{k-1} + 1$ pair, we are left with an ordered pair list $\sigma =$

$(b_{\ell_1}, b_{\ell_2}, \dots, b_{\ell_{k-2}}, b_1)$ of $k-1$ strategic pile elements. This ordered pair list clearly corresponds to the graph G_s , meaning G_s is a τ -

$-1, [a_1, a_2, \dots, a_m]$.

Conversely, let $G_s = (V_s, E_s)$ graph; it follows that $G_s \in X_k$

) be an arbitrary graph in $X_{k-1,[a_1, a_2, \dots, a_m]}$, and

let $G_r := (V_s \cup \{b_k\}, E_s)$. Then G_r is also a τ -graph, since the ordered pair list associated with G_r is the ordered pair list associated with G_s with the addition of b_k as the first element. Note that $\text{orb}_{\mathbb{Z}_k}(G_r)$ is a subset of $X_{k,[a_1, a_2, \dots, a_m]}$ since $\mathbb{Z}_k$ acts on $X_{k,[a_1, a_2, \dots, a_m]}$. It follows that $G \in X_{k,[a_1, a_2, \dots, a_m]}$ for any $G \in \text{orb}_{\mathbb{Z}_k}(G_r)$. $\square$

Using Lemma 5.16, we can determine the number of elements in $X_{k,[a_1, a_2, \dots, a_m]}$ by adding a vertex to each graph in $X_{k-1,[a_1, a_2, \dots, a_m]}$ and then considering all rotations of each of those graphs. However, the graphs formed through this method are not necessarily distinct. One of the reasons this is true is due to the fact that two graphs in $X_{k-1,[a_1, a_2, \dots, a_m]}$

may be in the same orbit when the vertex b_k is added. The following lemma addresses this issue.

Lemma 5.17. *For all $\tau \in X_{k,[a_1,a_2,\dots,a_m]}$, let Z_τ be the set of all τ -graphs in the orbit of τ under Z_k which do not have b_k as an isolated vertex. Then*

$$|Z_\tau| = \frac{a_1 + a_2 + \dots + a_m}{|\text{stab}(\tau)|}.$$

Proof. Define $\ell := a_1 + a_2 + \dots + a_m$, and note that this is the number of edges in any member of $X_{k,[a_1,a_2,\dots,a_m]}$. Let $\tau \in X_{k,[a_1,a_2,\dots,a_m]}$ and label the edges of τ as $e_1, e_2, \dots, e_\ell$. Let z_i be the rotation of τ such that e_i is a directed edge terminating at b_k . Then $Z_\tau = \{z_1, z_2, \dots, z_\ell\}$, since b_k is not an isolated vertex if and only if some edge points to b_k . However these z_i are not necessarily distinct.

Observe that for a given i , the number of times z_i appears in Z_τ is given by $|\text{stab}(z_i)|$. Moreover, since $z_i \in \text{orb}_{Z_k}(\tau)$, we have that $|\text{stab}(z_i)| = |\text{stab}(\tau)|$. Therefore,

$$|Z_\tau| = \frac{a_1 + a_2 + \dots + a_m}{|\text{stab}(\tau)|} = \frac{1}{|\text{stab}(\tau)|} \sum_{i=1}^{\ell} |\text{stab}(z_i)| = \frac{\ell}{|\text{stab}(\tau)|}. \quad \square$$

We now have what we need to prove the main relationship between $|X_{k,[a_1,a_2,\dots,a_m]}|$ and $|X_{k-1,[a_1,a_2,\dots,a_m]}|$.

Theorem 5.18. *For any $a_1, a_2, \dots, a_m \in \mathbb{Z}$ such that $a_1 + a_2 + \dots + a_m < k$*

$$|X_{k,[a_1,a_2,\dots,a_m]}| = \frac{k |X_{k-1,[a_1,a_2,\dots,a_m]}|}{k - (a_1 + a_2 + \dots + a_m)}.$$

Proof. By Lemma 5.16, in order to count $|X_{k,[a_1,a_2,\dots,a_m]}|$, we can add a vertex to every graph in $X_{k-1,[a_1,a_2,\dots,a_m]}$ and consider all rotations of these new graphs. Each of these graphs has k possible rotations. However, this does not produce distinct elements of $X_{k,[a_1,a_2,\dots,a_m]}$. In fact, for each $\tau \in X_{k,[a_1,a_2,\dots,a_m]}$, we have counted it $|\text{stab}_{Z_k}(\tau)| \cdot |\text{orb}_{Z_k}(\tau)| \cdot |Z_\tau|$ times. Since the addition of the vertex b_k can cause nonisomorphic graphs in $X_{k-1,[a_1,a_2,\dots,a_m]}$ to be in the same orbit under Z_k (see Fig. 5), we have over-counted each orbit in $X_{k,[a_1,a_2,\dots,a_m]}$ by a factor of $|\text{orb}_{Z_k}(\tau)| \cdot |Z_\tau|$. Due to rotational symmetry, we over-count $\tau \in \text{orb}_{Z_k}(\tau)$ by a factor of $|\text{stab}_{Z_k}(\tau)|$.

By Lemma 5.17, for all $\tau \in X_{k,[a_1,a_2,\dots,a_m]}$, we have that $|Z_\tau| = \frac{a_1+a_2+\dots+a_m}{|\text{stab}(\tau)|}$. Recall that $Z_\tau \subseteq \text{orb}_{Z_k}(\tau)$. Therefore,

$$\begin{aligned} |\text{stab}_{Z_k}(\tau)| \cdot |\text{orb}_{Z_k}(\tau) \setminus Z_\tau| &= |\text{stab}_{Z_k}(\tau)| (|\text{orb}_{Z_k}(\tau)| - |Z_\tau|) \\ &= |\text{stab}_{Z_k}(\tau)| \cdot |\text{orb}_{Z_k}(\tau)| - |\text{stab}_{Z_k}(\tau)| \cdot |Z_\tau| \\ &= |\text{stab}_{Z_k}(\tau)| \cdot |\text{orb}_{Z_k}(\tau)| - (a_1 + a_2 + \dots + a_m). \end{aligned}$$




Fig. 5. Two non-isomorphic graphs that will be in the same Z_k -orbit after the addition of the vertex .

Then, by the orbit stabilizer theorem, $|\text{stab}_{Z_k}(\tau)| \cdot |\text{orb}_{Z_k}(\tau)| = |Z_k| = k$, so we have counted each rotation $k - (a_1 + a_2 + \dots + a_m)$ times. Therefore,

$$|X_{k,[a_1,a_2,\dots,a_m]}| = \frac{k|X_{k-1,[a_1,a_2,\dots,a_m]}|}{k - (a_1 + a_2 + \dots + a_m)}.$$

□

This recursive relationship could be useful for addressing Step 2(a) of the merge number algorithm. However it is only useful when the base cases, $|X_{\ell,[a_1,a_2,\dots,a_m]}|$ (where $\ell = a_1 + a_2 + \dots + a_m$), are already known. Unfortunately, there is no known efficient way to compute these base cases. Using brute force in the same way as we can for Step 2(a) (see the beginning of Sec. 5.3.2), these base cases could be computed in $O(\ell \cdot \ell!)$ time. Since $\ell = O(k)$ in the worst case, this is not a significant improvement over the original brute force algorithm for Step 2(a).

5.3.3. Step 2(b)

Step 2(b) of the algorithm requires determining the number of ways merges can be picked with the given cycle structure. From Lemma 5.6, we can know that this is equivalent to choosing edges so that no cycle is formed.

Given a graph with cycle structure $[a_1, a_2, \dots, a_m]$, let e be the number of edges in the graph. This means that $e = a_1 + a_2 + \dots + a_m$. The total number of ways to choose edges

is $\binom{e}{\ell}$. The total number of ways to choose edges that include at least one cycle can be found using the inclusion–exclusion principle as follows:

$$\sum_{i=1}^m (-1)^{i+1} \sum \left\{ \binom{e - a_{k_1} - a_{k_2} - \cdots - a_{k_i}}{\ell - a_{k_1} - a_{k_2} - \cdots - a_{k_i}} : 1 \leq k_1, \dots, k_i \leq m, \text{ all distinct} \right\}$$

Subtracting this from $\binom{e}{\ell}$ yields the total number of ways to choose edges without picking a cycle, which is what we wanted.

6. Future Work

According to Theorem 3.10, for an odd natural number n , the number of elements of S_n that have a maximum size strategic pile is $2 \cdot (n - 2)!$ This number is related to the number of factorizations given in the following result from Bertram and Wei.

Theorem 6.1 ([2, Theorem 1]). *For $n \geq 3$, each odd permutation in S_n has exactly $2(n - 2)!$ factorizations of the form $\alpha \circ \beta$ where α is an n -cycle and β is an $(n - 1)$ -cycle.*

Viewing Theorem 6.1 in our context, let $n \geq 3$ be an odd integer, and let π be an element of S_n . With X_n , Y_π , and C_π as defined in Eqs. (2.4)–(2.6), we are considering factorizations of X_n of the form

$$X_n = Y_{\pi^{-1}} \circ C_\pi,$$

where C_π is a single cycle of length n , while X_n and Y_π are cycles of length $n + 1$. Applying Theorem 6.1, we see that according that theorem there are $2(n - 1)!$ factorizations of X_n of the form $\mu \circ \nu$ where μ is an $(n+1)$ -cycle and ν is an n -cycle. In each of these cases, we can write μ as a $Y_{\pi^{-1}}$ for some $\pi \in S_n$, and for $2(n - 2)!$ of these π the corresponding ν is a C_π of the form $(0 \ n \ i \ \cdots)$.

Example 6.2. Consider $n = 5$. The following table indicates that X_5 has factorizations into a 6-cycle and a 5-cycle for which the corresponding permutations π have various strategic pile sizes.

π	C_π	Strategic pile	Strategic pile size
[2 4 1 3 5]	(0 4 3 2 1)	$\emptyset$	0
[5 2 3 1 4]	(0 3 1 5 4)	$\{4\}$	1
[2 1 5 3 4]	(0 2 5 4 1)	$\{1, 4\}$	2
[3 5 1 2 4]	(0 5 4 3 2)	$2, 3, 4$	3

{ }

Thus, it can happen that the cycle C_π of length n in the factorization of X_n represents a strategic pile of size less than the maximal possible size for n . It would be interesting to determine, for odd integers n and for each strategic pile size $0 \leq k \leq n - 2$ how many of the permutations in S_n for which C_π is a cycle of length n have strategic pile size k . We have also not addressed the analogous question for the case when n is an even integer.

In addition to the problem just described, we would like to either (1) improve the merge number algorithm described in Sec. 5.3 or (2) construct an alternative algorithm for computing merge numbers.

Accomplishing (1) would require improving the following aspects of our algorithm. Let k indicate strategic pile size. Recall that Step 1 of this algorithm requires determining the set of integer partitions of k with no parts of size one, and with an even number of even parts. As previously mentioned, the number of integer partitions of k grows exponentially in k [7], meaning Step 1 is inefficient for large k . To make this step of the algorithm less costly, we would like a better method for computing the number of partitions with the aforementioned properties. Recall also that Step 2(a) of this algorithm can be done through brute force in $O(k \cdot k!)$ time. We offer a recursive method for completing Step 2(a) with runtime polynomial in k . However, this recursive method is only useful when the base cases, $|X_{\ell, [a_1, a_2, \dots, a_m]}|$ (where $\ell = a_1 + a_2 + \dots + a_m$), are already known. Unfortunately, the best known method for computing the base cases of this algorithm requires $O(\ell \cdot \ell!)$ time. Since this is no better than the brute force method for Step 2(a), we would like an efficient method for computing base cases so that our recursive method can be used to make Step 2(a) more efficient.

Alternatively, it would be ideal to (2) construct an algorithm for computing merge numbers that completely circumvents the dependency on exponential time computations. However, due to the nature of merge numbers described above, it seems that these dependencies might be unavoidable. Consequently, it is not clear how realistic it would be to accomplish (2).

Acknowledgment

This work was supported by NSF grant DMS-1359425 as a part of the 2016 Boise State University Complexity Across Disciplines Research Experience for Undergraduates (REU).

References

- [1] K. L. M. Adamyk, E. Holmes, G. R. Mayfield, D. J. Moritz, M. Scheepers, B. E. Tenner and H. C. Wauck, Sorting permutations: Games, genomes, and cycles, *Discrete Math. Algorithms Appl.* **9** (2017) 5.
- [2] E. A. Bertram and V. K. Wei, Decomposing a permutation into two large cycles: An enumeration, *SIAM J. Algebr. Discrete Methods* **1**(4) (1980) 450–461.
- [3] L. Cangelmi, Factorization of an n -cycle into two n -cycles, *European J. Combin.* **24**(7) (2003) 849–853.
- [4] D. A. Christie. Sorting permutations by block-interchanges, *Inform. Process. Lett.* **60**(4) (1996) 165–169.
- [5] A. Ehrenfeucht, T. Harju, I. Petre, D. M. Prescott and G. Rozenberg, *Computation in Living Cells: Gene Assembly in Ciliates*, Natural Computing Series (Springer, 2003).
- [6] A. Ehrenfeucht, T. Harju, I. Petre and G. Rozenberg. Characterizing the micronuclear gene patterns in ciliates, *Theory Comput. Syst.* **35**(5) (2002) 501–519.
- [7] G. H. Hardy and S. Ramanujan, Asymptotic formulae in combinatory analysis, *Proc. London Math. Soc.* **2**(1) (1918) 75–115.
- [8] D. M. Prescott, Genome gymnastics: Unique modes of DNA evolution and processing in ciliates, *Nat. Rev. Genet.* **1**(3) (2000) 191–198.
- [9] D. M. Prescott, A. Ehrenfeucht and G. Rozenberg, Template-guided recombination for IES elimination and unscrambling of genes in stichotrichous ciliates, *J. Theor. Biol.* **222**(3) (2003) 323–330.
- [10] N. J. A. Sloane, The on-line encyclopedia of integer sequences, <http://www.oeis.org/>.
- [11] S. Yancopoulos, O. Attie and R. Friedberg. Efficient sorting of genomic permutations by translocation, inversion and block interchanges, *Bioinformatics* **21**(16) (2005) 3340–3346.