

Algebra & Number Theory

Volume 14
2020
No. 2

**Characteristic cycles and Gevrey series solutions
of A -hypergeometric systems**

Christine Berkesch and María-Cruz Fernández-Fernández

Characteristic cycles and Gevrey series solutions of A -hypergeometric systems

Christine Berkesch and María-Cruz Fernández-Fernández

We compute the L -characteristic cycle of an A -hypergeometric system and higher Euler–Koszul homology modules of the toric ring. We also prove upper semicontinuity results about the multiplicities in these cycles and apply our results to analyze the behavior of Gevrey solution spaces of the system.

Introduction

Let D denote the Weyl algebra on $X = \mathbb{C}^n$ with coordinates $x = x_1, \dots, x_n$. Let ∂_i denote the variable that acts on $\mathbb{C}[x]$ as $\partial/\partial x_i$ and write $\partial = \partial_1, \dots, \partial_n$. A *weight vector* on D is $L = (L_x, L_\partial) \in \mathbb{Q}^n \times \mathbb{Q}^n$ such that $L_x + L_\partial \geq 0$. Such a vector induces an exhaustive increasing filtration L on D by, for $k \in \mathbb{Q}$,

$$L^k D := \mathbb{C} \cdot \{x^u \partial^v \mid L \cdot (u, v) \leq k\}.$$

Write $L^{<k} D := \bigcup_{\ell < k} L^\ell D$. For any P in $L^k D \setminus L^{<k} D$, set

$$\mathrm{in}_L(P) := P + L^{<k} D \in \mathrm{gr}^{L,k} D := L^k D / L^{<k} D \subseteq \mathrm{gr}^L D \quad \text{and} \quad \deg^L(P) := k.$$

For a left D -ideal I and the D -module $M = D/I$, set

$$\mathrm{gr}^L(I) := \langle \mathrm{in}_L(P) \mid P \in I \rangle \subseteq \mathrm{gr}^L(D) \quad \text{and} \quad \mathrm{gr}^L(M) := \mathrm{gr}^L(D) / \mathrm{gr}^L(I).$$

If $L_x + L_\partial = 0$, the associated graded ring $\mathrm{gr}^L D$ is isomorphic to D and $\mathrm{gr}^L(I)$ can be identified with a left D -ideal, which is also called a *Gröbner deformation* of I in [Saito et al. 2000]. It is suggestive to call $\mathrm{gr}^L(M)$ the Gröbner deformation of M with respect to L . On the other hand, if $L_x + L_\partial > 0$, the associated graded ring $\mathrm{gr}^L D$ is isomorphic to the coordinate ring of $T^*X \cong \mathbb{C}^{2n}$, which is a polynomial ring in $2n$ variables. In this latter case, the L -characteristic variety of M is

$$\mathrm{Char}^L(M) := \mathrm{Var}(\mathrm{gr}^L(I)) \subseteq T^*X \cong \mathbb{C}^{2n}. \quad (0-1)$$

The L -characteristic cycle of M is the finite formal sum

$$\mathrm{CC}^L(M) := \sum_C \mu^{L,C}(M) \cdot C,$$

CB was partially supported by NSF Grants DMS 1661962, DMS 1440537, OISE 0964985. MCFF was partially supported by MTM2016-75024-P, PP2014-2397, P12-FQM-2696 and FEDER.

MSC2010: primary 13N10; secondary 14M25, 32C38, 33C70.

Keywords: A -hypergeometric system, toric ring, D -module, characteristic cycle, irregularity sheaf, Gevrey series.

where C runs over the irreducible components of $\text{Char}^L(M)$, and

$$\mu^{L,C}(M) := \ell((\text{gr}^L(M))_{P_C})$$

is the multiplicity of $\text{gr}^L(M)$ along C , where P_C is the defining ideal of C in $\text{gr}^L(D)$ and ℓ denotes the length of a $\text{gr}^L(D)_{P_C}$ -module.

The weight vector $F = (\mathbf{0}_n, \mathbf{1}_n) := (0, \dots, 0, 1, \dots, 1) \in \mathbb{Q}^n \times \mathbb{Q}^n$ induces the *order filtration* on D . We notice that $\text{Char}^F(M)$ and $\text{CC}^F(M)$ are called, respectively, the *characteristic variety* and the *characteristic cycle* of M . If M is *holonomic*, that is, the dimension of its characteristic variety is n , then the *rank* of M , defined $\text{rank}(M) := \dim_{\mathbb{C}(x)} \mathbb{C}(x) \otimes_{\mathbb{C}[x]} M$, coincides with the dimension of the space of germs of its holomorphic solutions at any nonsingular point by a result of Kashiwara (see e.g., [Saito et al. 2000, Theorem 1.4.19]). Notice that $\text{rank}(M) = \mu^{F,C}(M)$ for $C = T_X^*X$.

One motivation for the study of L -characteristic cycles comes from the theory of irregularity of holonomic D -modules. For a flavor of this deep and involved theory that fits the goals of this paper, a projective weight vector of the form

$$L = F + (s - 1)V$$

where $s \in \mathbb{Q}$ and $V = (-w, w)$, with $w = (0, \dots, 0, 1)$, induces the Kashiwara–Malgrange filtration along the coordinate hyperplane $Y = \{x_n = 0\} \subseteq X = \mathbb{C}^n$. In this case, the L -characteristic variety $\text{Char}^{F+(s-1)V}(M)$ is locally constant with respect to $s \in \mathbb{Q}$, except for at a finite set of values called *algebraic slopes* of M along Y . This is a global version of the algebraic slopes defined and studied by Laurent [1987]. On the other hand, the *analytic slopes* of M along Y were defined as jumps in the *Gevrey* filtration of the *irregularity sheaf* of M along Y by Mebkhout [1990]. The comparison theorem for slopes states that the algebraic and analytic slopes for M along Y coincide, and, even more, the Euler–Poincaré characteristic of the irregularity sheaf can be computed in terms of the L -characteristic cycles of M [Laurent and Mebkhout 1999]. In particular, certain multiplicities in the L -characteristic cycles are closely related to the dimension of the space of Gevrey solutions of M along Y .

Another motivating idea of this article is that the F -characteristic cycle of a Gröbner deformation of a holonomic D -module M is equal to the L -characteristic cycle of M for an appropriate L (see Lemma 3.1 for the precise statement). In particular, the holonomic rank of such a Gröbner deformation is the multiplicity of the component T_X^*X in $\text{CC}^L(M)$.

Our main interest is A -hypergeometric D -modules, also known as GKZ-systems after their introduction and study by Gelfand, Graev, Kapranov, and Zelevinsky [Gelfand et al. 1987; 1989; 1990]. Let

$$A = [a_{ij}] = [a_1 \cdots a_n] \in \mathbb{Z}^{d \times n}$$

be an integral matrix such that the group generated by the columns of A , $\mathbb{Z}A$, is equal to $\mathbb{Z}^d$, and the positive real cone $\mathbb{R}_{\geq 0}A$ over the columns is pointed. Let

$$I_A := \langle \partial^u - \partial^v \mid Au = Av \rangle \subseteq \mathbb{C}[\partial]$$

denote the *toric ideal* of A . For $\beta \in \mathbb{C}^d$, write $E - \beta$ for the sequence of *Euler operators* given by

$$E_i - \beta_i := \sum_{j=1}^n a_{ij} x_j \partial_j - \beta_i$$

for $i = 1, \dots, d$. The A -hypergeometric system of A at $\beta \in \mathbb{C}^d$ is

$$H_A(\beta) := D \cdot (I_A + \langle E - \beta \rangle) \quad \text{with associated module } M_A(\beta) := D/H_A(\beta).$$

A weight vector $L = (L_x, L_\partial) \in \mathbb{Q}^n \times \mathbb{Q}^n$ as above is called *projective* if $L_x + L_\partial = c \cdot \mathbf{1}_n := c \cdot (1, \dots, 1)$ for some constant $c > 0$. Notice that any Euler operator E_i is homogeneous with respect to such a filtration. In [Schulze and Walther 2008], the irreducible components of $\text{CC}^L(M_A(\beta))$ were enumerated, and when β is generic (or not rank-jumping), $\text{CC}^L(M_A(\beta))$ was computed. In this article, we compute $\text{CC}^L(M_A(\beta))$ for any β , along with the characteristic cycles of higher Euler–Koszul homology modules (see Section 1) of the toric ring $\mathbb{C}[\partial]/I_A$. We also provide upper semicontinuity results for some of these multiplicities and apply our results to the Gevrey solution spaces of $M_A(\beta)$.

Outline. In Sections 1–2, we provide background and preliminary results on Euler–Koszul homology and L -characteristic cycles of A -hypergeometric systems. We compute the multiplicities in the characteristic cycles of the Euler–Koszul homology of the toric ring in Section 4, with consequences in Section 5. We provide upper semicontinuity results in Section 6 and study Gevrey solutions of $H_A(\beta)$ in Section 7.

1. Euler–Koszul homology

In this section, we present background related to Euler–Koszul homology, as found in [Matusevich et al. 2005; Schulze and Walther 2009], with some additions needed in the sequel. We use the convention that $0 \in \mathbb{N}$. Recall that a_i denotes the i -th column of the matrix A . Given a subset $\tau \subseteq A$ of the column set of A , the semigroup generated by τ ,

$$\mathbb{N}\tau := \left\{ \sum_{a_i \in \tau} j_i a_i \mid j_i \in \mathbb{N} \text{ for all } a_i \in \tau \right\},$$

generates the semigroup ring $S_\tau := \mathbb{C}[\mathbb{N}\tau]$. With $\pi_\tau: \mathbb{C}[\partial_\tau] := \mathbb{C}[\partial_j \mid a_j \in \tau] \rightarrow S_\tau$ denoting the map induced by τ , we have the isomorphism of rings $S_\tau \cong \mathbb{C}[\partial_\tau]/\ker \pi_\tau$. When convenient, we will abuse notation and also view τ as a matrix.

A subset G of the columns of the matrix A is a *face* of A , denoted $G \preceq A$, if $\mathbb{R}_{\geq 0}G$ is a face of the cone $\mathbb{R}_{\geq 0}A$ and $G = A \cap \mathbb{R}G$. The codimension of a nonempty face G is $\text{codim}(G) := d - \dim(\mathbb{R}G)$, with $\text{codim}(\emptyset) = d$ by convention. Let G^c denote the complement of G in A .

Define a $\mathbb{Z}^d$ -grading on D via $\deg(x_i) := -a_i$ and $\deg(\partial_{x_i}) := a_i$. A $\mathbb{Z}^d$ -graded $\mathbb{C}[\partial]$ -module N is *toric* if it has a filtration

$$0 = N^{(0)} \subseteq N^{(1)} \subseteq \dots \subseteq N^{(\ell-1)} \subseteq N^{(\ell)} = N$$

such that $N^{(i)}/N^{(i-1)}$, for each i , is a $\mathbb{Z}^d$ -graded translate of S_{G_i} for some face $G_i \preceq A$. The *degree set* of a finitely generated $\mathbb{Z}^d$ -graded $\mathbb{C}[\partial]$ -module N is $\deg(N) := \{\alpha \in \mathbb{Z}^d \mid N_\alpha \neq 0\}$. The *quasidegree set* of N , denoted $\text{qdeg}(N)$, is the Zariski closure of $\deg(N)$ under the natural embedding $\mathbb{Z}^d \hookrightarrow \mathbb{C}^d$. A $\mathbb{Z}^d$ -graded $\mathbb{C}[\partial]$ -module N is *weakly toric* if there is a filtered partially ordered set $(\mathfrak{S}, \leq)$ and a $\mathbb{Z}^d$ -graded direct limit

$$\phi_s: N^{(s)} \rightarrow \lim_{\longrightarrow s \in \mathfrak{S}} N^{(s)} = N,$$

where $N^{(s)}$ is a toric $\mathbb{C}[\partial]$ -module for each $s \in \mathfrak{S}$. The *quasidegrees* of N are

$$\text{qdeg}(N) := \bigcup_{s \in \mathfrak{S}} \text{qdeg}(\phi_s(N^{(s)})),$$

where each $\text{qdeg}(\phi_s(N^{(s)}))$ is already defined since $\phi_s(N^{(s)})$ is toric for each s .

Let N be a weakly toric module. Given a homogeneous $y \in D \otimes_{\mathbb{C}[\partial]} N$, define an action of the Euler operators for $1 \leq i \leq d$ by

$$(E_i - \beta_i) \circ y = (E_i - \beta_i + \deg_i(y))y,$$

and extend this action $\mathbb{C}$ -linearly to $D \otimes N$. With this sequence of commuting endomorphisms on $D \otimes N$, let $\mathcal{K}_\bullet^A(N, \beta)$ denote the Koszul complex on the left D -module $D \otimes_{\mathbb{C}[\partial]} N$, which we call the *Euler–Koszul complex* of N at β . Its homology is denoted $\mathcal{H}_i^A(N, \beta) := H_i(\mathcal{K}_\bullet^A(N, \beta))$ or simply $\mathcal{H}_i(N, \beta)$ when A is clear from the context. Euler–Koszul homology was first introduced in [Matusevich et al. 2005] for toric modules and extended to weakly toric modules in [Schulze and Walther 2009].

If $b \in \mathbb{Z}^d$, we denote by $N(b)$ a $\mathbb{Z}^d$ -graded translated copy of N such that $N(b)_v = N_{v-b}$ for all $v \in \mathbb{Z}^d$. Thus, $\deg(N(b)) = b + \deg(N)$. For example, if $N = S_A = \mathbb{C}[\mathbb{N}A]$ then $N(b) = \mathbb{C}[\mathbb{N}A]t^b$. Euler–Koszul homology is compatible with these graded shifts. Namely, we have

$$\mathcal{H}_q(N(b), \beta) \cong \mathcal{H}_q(N, \beta - b)(b). \quad (1-1)$$

Theorem 1.1 [Schulze and Walther 2009, Theorem 5.4]. *For a weakly toric module N , the following are equivalent:*

- (1) $\mathcal{H}_i(N, \beta) = 0$ for all $i \geq 0$.
- (2) $\mathcal{H}_0(N, \beta) = 0$.
- (3) $\beta \notin \text{qdeg}(N)$. □

Theorem 1.2 [Matusevich et al. 2005, Theorem 6.6; Schulze and Walther 2009]. *Let N be a weakly toric module. Then $\mathcal{H}_i(N, \beta) = 0$ for all $i > 0$ and for all $\beta \in \mathbb{C}^d$ if and only if N is a maximal Cohen–Macaulay S_A -module.* □

For a subset $\tau \subseteq A$, given an $\mathbb{N}\tau$ -module $\mathbb{S}$, define the S_τ -module $\mathbb{C}\{\mathbb{S}\} := \bigoplus_{s \in \mathbb{S}} \mathbb{C} \cdot t^s$ as a $\mathbb{C}$ -vector space with S_τ -action given by $\partial_i \cdot t^s = t^{s+a_i}$. Then $\mathbb{C}\{\mathbb{S}\}$ has a multiplicative structure given by $t^s \cdot t^{s'} = t^{s+s'}$, and $S_\tau \cong \mathbb{C}\{\mathbb{N}\tau\}$ as rings. The *saturation* of τ in $\mathbb{Z}\tau$ is the semigroup $\widetilde{\mathbb{N}\tau} = \mathbb{R}_{\geq 0}\tau \cap \mathbb{Z}\tau$. The *saturation*

of S_τ is the semigroup ring of the saturation of τ in $\mathbb{Z}\tau$, which is given by $\tilde{S}_\tau = \mathbb{C}\{\widetilde{\mathbb{N}\tau}\}$ as a $\mathbb{Z}^d$ -graded S_τ -module. By [Hochster 1972], $\tilde{S}_\tau$ is a Cohen–Macaulay S_τ -module.

2. Characteristic cycles of A -hypergeometric systems

Let $L = (L_x, L_\partial) \in \mathbb{Q}^{2n}$ be a projective weight vector on D . In this section, we recall from [Schulze and Walther 2008] the description of the L -characteristic variety of an A -hypergeometric system, which includes the computation of the L -characteristic cycle of $H_A(\beta)$ when β is not rank-jumping for A .

Let $h = (h_1, \dots, h_d) \in \mathbb{Q}^d$ be such that $h \cdot a_i > 0$ for $i = 1, \dots, n$. Choose $\varepsilon > 0$ such that $h \cdot a_i + \varepsilon L_{\partial_i} > 0$ for $i = 1, \dots, n$, and denote by H_ε the hyperplane in $\mathbb{P}_{\mathbb{Q}}^d$ given by

$$\{[y_0 : y_1 : \dots : y_d] \in \mathbb{P}_{\mathbb{Q}}^d \mid \varepsilon y_0 + h_1 y_1 + \dots + h_d y_d = 0\}.$$

The L -polyhedron of A is the convex hull of $\{[1 : \mathbf{0}_d], [L_{\partial_1} : a_1], \dots, [L_{\partial_n} : a_n]\}$ in the affine space $\mathbb{P}_{\mathbb{Q}}^d \setminus H_\varepsilon$. The (A, L) -umbrella, denoted Φ_A^L , is the set of faces of the L -polyhedron of A that do not contain $[1 : \mathbf{0}_d]$.

We denote by $\Phi_A^{L,k} \subset \Phi_A^L$ the subset of faces τ of dimension k (equivalently, $\dim(\mathbb{C}\tau) = k + 1$). A face τ of Φ_A^L will be identified with $\{j \in \{1, \dots, n\} \mid [L_{\partial_j} : a_j] \in \tau\}$ or with the submatrix of A indexed by this set, when necessary. With this identification, Φ_A^L is an abstract polyhedral complex. For any face $G \preceq A$, set $\Phi_G^L := \{\tau \in \Phi_A^L \mid \tau \subseteq G\}$.

Let (x, ξ) denote the coordinates on $T^*X = T^*\mathbb{C}^n$. For any $\tau \subseteq \{1, \dots, n\}$, let

$$C_A^\tau := \left\{ (x, \xi) \in T^*X \mid \xi_i = 0 \text{ for } i \notin \tau, \sum_{i \in \tau} a_i x_i \xi_i = 0 \text{ and } \exists t \in (\mathbb{C}^*)^d, \xi_j = t^{a_j}, \forall j \in \tau \right\},$$

and let $\overline{C_A^\tau}$ denote the Zariski closure of C_A^τ in T^*X , with defining ideal $P_\tau \subseteq \mathbb{C}[x, \xi]$. In particular, $\overline{C_A^\emptyset} = T_X^*X$ and $\overline{C_A^{\{j\}}} = T_{(x_j=0)}^*X$.

If N is a $\mathbb{Z}^d$ -graded $\mathbb{C}[\partial]$ -module and $C = \overline{C_A^\tau}$ for some $\tau \in \Phi_A^L$, we write

$$\mu_{A,i}^{L,\tau}(N, \beta) := \mu_{A,i}^{L,C}(\mathcal{H}_i(N, \beta)) = \ell((\text{gr}^L(\mathcal{H}_i(N, \beta)))_{P_\tau}). \quad (2-1)$$

We will also denote $\mu_{A,i}^{L,\tau}(\beta) := \mu_{A,i}^{L,\tau}(S_A, \beta)$. By [Schulze and Walther 2008, Corollary 4.13],

$$\mu_A^{L,\tau} := \sum_{j=0}^d (-1)^j \mu_{A,j}^{L,\tau}(\beta) = \mu_{A,0}^{L,\tau}(\tilde{S}_A, \beta) = \mu_{A,0}^{L,\tau}(S_A[\partial_A^{-1}], \beta) \quad (2-2)$$

is independent of $\beta \in \mathbb{C}^d$.

Note that $\text{rank}(M_A(\beta))$ is equal to $\mu_{A,0}^{F,\emptyset}(\beta)$. Since $M_A(\beta)$ is always holonomic [Gelfand et al. 1987; Adolphson 1994], its rank is always finite. Further, the rank of $M_A(\beta)$ is upper semicontinuous as a function of the parameter β , with a generic value equal to $\text{vol}_{\mathbb{Z}^d}(A)$, the normalized volume in $\mathbb{Z}A = \mathbb{Z}^d$ of the convex hull of the columns of A and the origin [Matusevich et al. 2005; Adolphson 1994; Gelfand et al. 1990]. We recall that the normalized volume function in a lattice Ω , denoted by vol_Ω , is defined so

that the volume of the unit simplex in Ω (that is, the convex hull of the origin and a lattice basis of Ω) is one.

A parameter β is said to be *rank-jumping* when $\text{rank}(M_A(\beta)) > \text{vol}_{\mathbb{Z}^d}(A)$. The set of rank-jumping parameters is described in [Matusevich et al. 2005]; namely, with $\varepsilon_A := \sum_{i=1}^n a_i$,

$$\mathcal{E}_A := \{\beta \in \mathbb{C}^d \mid \text{rank}(M_A(\beta)) > \text{vol}_{\mathbb{Z}^d}(A)\} = -\text{qdeg}\left(\bigoplus_{i=0}^{d-1} \text{Ext}_{\mathbb{C}[\partial]}^{n-i}(S_A, \mathbb{C}[\partial])(-\varepsilon_A)\right).$$

Schulze and Walther provided a description of $\text{CC}^L(H_A(\beta))$ when β is not rank-jumping, as summarized through the following two results.

Theorem 2.1 [Schulze and Walther 2008, Theorem 4.21]. *For all $G \preceq A$, if $\tau \in \Phi_G^L$, then*

$$\mu_G^{L,\tau} = \sum_{\tau' \subseteq \tau \in \Phi_G^{L,d'-1}} [\mathbb{Z}G : \mathbb{Z}\tau'] \cdot [(\mathbb{Z}\tau' \cap \mathbb{Q}\tau) : \mathbb{Z}\tau] \cdot \text{vol}_{\pi(\mathbb{Z}\tau')}(P_{\tau,\tau'} \setminus Q_{\tau,\tau'}),$$

where $d' = \dim(\mathbb{C}G)$, $\pi : \mathbb{Z}\tau' \rightarrow \mathbb{Z}\tau' / (\mathbb{Z}\tau' \cap \mathbb{Q}\tau)$ is the natural projection and $P_{\tau,\tau'}$ and $Q_{\tau,\tau'}$ denote the convex hull of $\pi(\tau' \cup \{0\})$ and $\pi(\tau' \setminus \tau)$ respectively.

In [Schulze and Walther 2008], Theorem 4.21 is only stated for $G = A$. Theorem 2.1 is a straightforward adaptation that will be useful in the sequel. Note that here we are using (2-1) and (2-2) with A replaced by G , but we still write L for the filtration induced on the Weyl Algebra D_G in the variables $\{x_j \mid j \in G\}$ by the projective weight vector given by the G -coordinates of L_x and L_∂ .

Theorem 2.2 [Schulze and Walther 2008, Corollary 4.12]. *The L -characteristic variety of $M_A(\beta)$ is independent of $\beta \in \mathbb{C}^d$ and given by*

$$\text{Char}^L(M_A(\beta)) = \bigcup_{\tau \in \Phi_A^L} \overline{C}_A^\tau,$$

where each component $\overline{C}_A^\tau$ is irreducible. Moreover, $\mu_{A,0}^{L,\tau}(\beta) \geq \mu_A^{L,\tau}$, and equality holds if β is not rank-jumping.

Theorem 2.2 implies that when β is not rank-jumping,

$$\text{CC}^L(M_A(\beta)) = \sum_{\tau \in \Phi_A^L} \mu_A^{L,\tau} \cdot \overline{C}_A^\tau,$$

and for each $\tau \in \Phi_A^L$, the multiplicity $\mu_A^{L,\tau}$ is computed in Theorem 2.1.

A subset $\tau \subseteq A$ is called *F-homogeneous* if the set of columns of A indexed by τ lie in a common affine hyperplane off the origin. For a subset $\tau \subseteq A$, let $\Delta_\tau = \text{conv}(\tau \cup \{0\}) \subseteq \mathbb{R}^d$ denote the convex hull of the origin and all the columns of τ .

By [Schulze and Walther 2008, Corollary 4.22 and Remark 4.23],

$$\mu_A^{L,\emptyset} = \text{vol}_{\mathbb{Z}^d}\left(\bigcup_{\tau' \in \Phi_A^{L,d-1}} \Delta_{\tau'} \setminus \text{conv}(\tau')\right). \quad (2-3)$$

Hence if all the facets of the (A, L) -umbrella are F -homogeneous, then

$$\mu_A^{L, \emptyset} = \text{vol}_{\mathbb{Z}^d} \left(\bigcup_{\tau' \in \Phi_A^{L, d-1}} \Delta_{\tau'} \right). \quad (2-4)$$

3. F -characteristic cycles of initial ideals are L -characteristic cycles

Given any real vector $w \in \mathbb{R}^n$ and any left ideal $J \subseteq D$, we can consider the initial ideal $\text{in}_{(-w, w)}(J)$ as defined in [Saito et al. 2000]. We recall that by [loc. cit., Theorem 2.2.1], if $M = D/J$ is a holonomic D -module, then so is $\text{gr}^{(-w, w)}(M) := D/\text{in}_{(-w, w)}(J)$ and, moreover,

$$\text{rank}(\text{gr}^{(-w, w)}(M)) \leq \text{rank}(M). \quad (3-1)$$

On the other hand, by [loc. cit., Lemma 2.1.6], for any weight vector $(u, v) \in \mathbb{R}^{2n}$ and $L = (-w, w) + \epsilon(u, v)$ with $\epsilon > 0$ small enough,

$$\text{gr}^{(u, v)}(\text{gr}^{(-w, w)}(M)) = \text{gr}^L(M). \quad (3-2)$$

Lemma 3.1. *If $M = D/J$ is a holonomic D -module, then for L chosen as in (3-2) with $(u, v) = F$,*

$$\text{CC}^F(\text{gr}^{(-w, w)}(M)) = \text{CC}^L(M).$$

The holonomic rank of $\text{in}_{(-w, w)}(M_A(\beta))$, a central object of study in [loc. cit.], equals the multiplicity $\mu_{A,0}^{L, \emptyset}(\beta)$ for $L = (-w, w) + \epsilon F$ and $\epsilon > 0$ small enough. Notice that, by the form of L , all the facets of Φ_A^L are F -homogeneous. We will see in Section 4 that for any projective weight vector L , the multiplicity $\mu_{A,0}^{L, \emptyset}(\beta)$ equals the rank of a Gröbner deformation of $M_A(\beta)$ (see Corollaries 4.3 and 4.5).

4. Computing multiplicities in L -characteristic cycles

In this section, we use the approach of [Berkesch 2011] to compute the multiplicities in the L -characteristic cycles of Euler–Koszul homology modules of the toric ring S_A . We first recall some definitions from [Berkesch 2011; Berkesh et al. 2018].

For a face $G \preceq A$, consider the union of the lattice translates

$$\mathbb{E}_G^\beta := [\mathbb{Z}^d \cap (\beta + \mathbb{C}G)] \setminus (\mathbb{N}A + \mathbb{Z}G) = \bigsqcup_{b \in B_G^\beta} (b + \mathbb{Z}G), \quad (4-1)$$

where B_G^β is a set of lattice translate representatives. As such, $|B_G^\beta|$ is the number of translates of $\mathbb{Z}G$ appearing in $\mathbb{E}_G^\beta$, which is by definition equal to the difference between $[\mathbb{Z}^d \cap \mathbb{Q}G : \mathbb{Z}G]$ and the number of translates of $\mathbb{Z}G$ along $\beta + \mathbb{C}G$ that are contained in $\mathbb{N}A + \mathbb{Z}G$.

For a face $\tau \in \Phi_A^L$ of the (A, L) -umbrella, let $\mathbb{E}_\tau^\beta$ denote the union of the ranking lattices $\mathbb{E}_G^\beta$, where $G \preceq A$ contains τ .

Theorem 4.1. *Let L be a projective weight vector and $\tau \in \Phi_A^L$ be a face of the (A, L) -umbrella. For each i and β , the multiplicity $\mu_{A,i}^{L, \tau}(\beta)$, which is the coefficient of $\bar{C}_A^\tau$ in the characteristic cycle $\text{CC}^L(\mathcal{H}_i(S_A, \beta))$*

(see (2-1)), can be computed from the combinatorics of the ranking lattices at β and the (A, L) -umbrella Φ_A^L . More precisely, there is a spectral sequence involving the faces of Φ_A^L that contain τ and the ranking lattices in $\mathbb{E}_\tau^\beta$, from which $\mu_{A,i}^{L,\tau}(\beta)$ can be computed.

Before proving [Theorem 4.1](#), we state some consequences.

Corollary 4.2. *For all $\beta \in \mathbb{C}^d$ and all projective weight vectors L, L' ,*

$$\mathrm{CC}^L(M_A(\beta)) = \mathrm{CC}^{L'}(M_A(\beta)) \quad \text{if and only if} \quad \Phi_A^L = \Phi_A^{L'}.$$

Proof. While the only if direction follows from [Theorem 2.2](#), the if direction uses [Theorems 2.1, 2.2](#), and [4.1](#). □

Corollary 4.3. *For any projective weight vector $L = (u, v)$ on D such that all the facets of Φ_A^L are F -homogeneous,*

$$\mathrm{CC}^F(\mathrm{gr}^{(-v,v)}(M_A(\beta))) = \mathrm{CC}^L(M_A(\beta)).$$

In particular, $\mathrm{rank}(\mathrm{gr}^{(-v,v)}(M_A(\beta))) = \mu_{A,0}^{L,\emptyset}(\beta)$.

Proof. Let $\epsilon > 0$ be as small as necessary in the sequel. Notice first that $\mathrm{CC}^F(\mathrm{gr}^{(-v,v)}(M_A(\beta))) = \mathrm{CC}^{L_\epsilon}(M_A(\beta))$ for $L_\epsilon := (-v, v) + \epsilon F$ by [Lemma 3.1](#). Moreover, by the assumption on the (A, L) -umbrella, we have $\Phi_A^L = \Phi_A^{L+\epsilon F}$. On the other hand, the last n coordinates of $L + \epsilon F$ and L_ϵ are equal to $v + \epsilon \cdot \mathbf{1}_n$, and hence $\Phi_A^L = \Phi_A^{L_\epsilon}$. Thus, the result follows from [Corollary 4.2](#). □

As a particular case of [Corollary 4.3](#), the characteristic cycles, and hence the ranks, of the modules $\mathrm{gr}^{(-\mathbf{1}_n, \mathbf{1}_n)}(M_A(\beta))$ and $M_A(\beta)$ are equal. We next show that [[Saito et al. 2000](#), Corollary 3.2.14] holds with weakened hypotheses.

Corollary 4.4. *For any $\beta \in \mathbb{C}^d$ and any (not necessarily homogeneous) A , the small Gröbner fan of the hypergeometric ideal $H_A(\beta)$ refines the secondary fan of A .*

Proof. It suffices to see that each open cone of the small Gröbner fan of $H_A(\beta)$ is contained in an open cone of the secondary fan of A . Since such an open cone corresponds to a Gröbner deformation with respect to a generic weight vector $w \in \mathbb{R}^n$, it follows that $L = (-w + c \cdot \mathbf{1}_n, w)$ is a projective weight vector for any $c > 0$ and Φ_A^L , which only depends on w , has only F -homogeneous facets. Thus, beginning with generic vectors w, w' with

$$\mathrm{gr}^{(-w,w)}(M_A(\beta)) = \mathrm{gr}^{(-w',w')}(M_A(\beta)),$$

[Corollaries 4.2](#) and [4.3](#) imply that $\Phi_A^L = \Phi_A^{L'}$ where the last coordinates of L and L' are w and w' , respectively. This means that w and w' belong to the same cone of the secondary fan of A . □

Corollary 4.5. *Any projective weight vector $L = (u, v)$ on D has a perturbation L' such that all the facets of the (A, L') -umbrella $\Phi_A^{L'}$ are F -homogeneous and $\mu_{A,0}^{L,\emptyset}(\beta) = \mu_{A,0}^{L',\emptyset}(\beta)$.*

Proof. If $L'(\epsilon) := L + \epsilon(\mathbf{1}_n, -\mathbf{1}_n)$ for $\epsilon > 0$, then there is an $\epsilon_0 > 0$ such that the $L'(\epsilon)$ -umbrella is constant for $\epsilon \in (0, \epsilon_0]$. Thus, if we fix $L' = L'(\epsilon_0)$, then all the facets of $\Phi_A^{L'}$ are F -homogeneous. Moreover, by the choice of L' , any F -homogeneous facet of Φ_A^L is a facet of $\Phi_A^{L'}$, while each non- F -homogeneous facet τ of Φ_A^L is replaced in $\Phi_A^{L'}$ by the set of facets of $\Phi_\tau^{L''}$, where $L'' := (c\mathbf{1}_n, -\mathbf{1}_n)$ is a projective weight vector for any $c > 1$. This latter set is the set of facets of $\text{conv}(\tau)$ that are not facets of Δ_τ . This proves that $\mu_A^{L,\emptyset} = \mu_A^{L',\emptyset}$ by using (2-3) to compute $\mu_A^{L,\emptyset}$ and (2-4) to compute $\mu_A^{L',\emptyset}$. Analogously, $\mu_G^{L,\emptyset} = \mu_G^{L',\emptyset}$ for any face $G \leq A$. Finally, the result follows from previous equality and Theorem 4.1. $\square$

Corollary 4.6. *Given any projective weight vector L and $\beta \in \mathbb{C}^d$,*

$$\mu_{A,0}^{L,\emptyset}(\beta) \leq \text{rank}(M_A(\beta)) \leq 4^{(d+1)} \text{vol}(A).$$

Proof. The first inequality is a consequence of (3-1) and Corollaries 4.5 and 4.3. The second is [Berkesch et al. 2018, Corollary 6.2]. $\square$

To prove Theorem 4.1, we will follow the approach used to compute the rank of an A -hypergeometric system from [Berkesch 2011] (see also [Berkesch et al. 2018]). We will use the set

$$\mathcal{C}_A(\beta) := \mathbb{Z}^d \cap (\text{Re } \beta + \mathbb{R}_{\geq 0}A).$$

Note that $\mathcal{C}_A(\beta)$ here is defined differently than in [Berkesch 2011]. However, the quotient between the subsequent modules with the same names, defined using $\mathcal{C}_A(\beta)$ here or as in [loc. cit.], all have quasidegree sets that do not contain β . Hence, by Theorem 1.1, the Euler–Koszul homology modules for modules with the same names here and in [loc. cit.] are isomorphic.

Given a subset

$$J \subseteq \mathcal{J}(\beta) := \{(G, b) \mid G \leq A, b \in B_G^\beta, \mathbb{E}_G^\beta \neq \emptyset\}, \quad (4-2)$$

define

$$\mathbb{E}_J^\beta := \bigcup_{(G,b) \in J} (b + \mathbb{Z}G) \quad \text{and} \quad \mathbb{P}_J^\beta := \mathcal{C}_A(\beta) \cap \mathbb{E}_J^\beta.$$

Now define the respective sets and S_A -modules

$$\mathbb{T}^\beta := \mathbb{N}A \cup \left[\bigcup_{b \in \mathbb{P}_{\mathcal{J}(\beta)}^\beta} (b + \widetilde{\mathbb{N}A}) \right], \quad \mathbb{T}^\beta := \mathbb{C}\{\mathbb{T}^\beta\}, \quad \mathbb{S}_J^\beta := \mathbb{T}^\beta \setminus \mathbb{P}_J^\beta, \quad \mathbb{S}_J^\beta := \mathbb{C}\{\mathbb{S}_J^\beta\}, \quad \text{and} \quad P_J^\beta := \frac{\mathbb{T}^\beta}{\mathbb{S}_J^\beta}.$$

The degree set of P_J^β is $\deg(P_J^\beta) = \mathbb{P}_J^\beta$. If a toric module N is isomorphic to P_J^β for some $J \subseteq \mathcal{J}(\beta)$ and β , then we say that N is a *ranking toric module* determined by J . A *simple ranking toric module* is a module isomorphic to $P_{G,J}^\beta := P_{J(G)}^\beta$, where $G \leq A$ is a fixed face of A such that $\mathbb{E}_G^\beta \neq \emptyset$ and

$$J(G) := \{(G, b) \in J \mid b \in B_G^\beta\}.$$

When $J = \mathcal{J}(\beta)$, we suppress it from the notation and write P^β and P_G^β in place of P_J^β and $P_{G,J}^\beta$, respectively. If $(G, b) \in J$ and there is not any other pair $(G', b') \in J$ such that $b + \mathbb{Z}G \subsetneq b' + \mathbb{Z}G'$ we say that (G, b) is a maximal pair in J . We denote by $\max(J)$ the set of all maximal pairs in J .

Lemma 4.7. *If $G \preceq A$ and $\tau \in \Phi_A^L$, then the multiplicity $\mu_{A,q}^{L,\tau}(P_G^\beta, \beta)$ of the simple ranking toric module P_G^β is*

$$\mu_{A,q}^{L,\tau}(P_G^\beta, \beta) = |B_G^\beta| \cdot \mu_{A,q}^{L,\tau}(\tilde{S}_G(b), \beta) = \begin{cases} |B_G^\beta| \cdot \binom{\text{codim}(G)}{q} \cdot \mu_G^{L,\tau} & \text{if } \tau \subseteq G, \\ 0 & \text{otherwise.} \end{cases}$$

for any $b \in B_G^\beta$.

Proof. For all $j \notin G$ we have that $\partial_j \cdot \tilde{S}_G = 0$, hence that $\xi_j \cdot \text{gr}^L(\mathcal{H}_0(\tilde{S}_G(b), \beta)) = 0$ where $\xi_j = \text{in}_L(\partial_j) \in \mathbb{C}[x, \xi] \cong \text{gr}^L(D)$. On the other hand, by the definition of P_τ , it is clear that $\xi_j \in P_\tau$ if and only if $j \notin \tau$. Thus, we have that $(\text{gr}^L(\mathcal{H}_0(\tilde{S}_G(b), \beta)))_{P_\tau} = 0$ if $\tau \not\subseteq G$. Now, with $\mu^{L, \overline{C}_A}$ in place of rank, the arguments in the proof of [Berkesch 2011, Theorem 6.1] yield this result. $\square$

Proof of Theorem 4.1. The argument proving [Berkesch 2011, Theorem 6.6] can be used to obtain this result, when J is chosen to be the right hand side of (4-2) and $\mu^{L, \overline{C}_A}$ in place of rank. We make note of the necessary modifications below.

To begin, it follows from Theorem 1.2 and (2-2) that

$$\mu_{A,i}^{L,\tau}(\beta) = \begin{cases} \mu_{A,i}^{L,\tau} + \mu_{A,1}^{L,\tau}(Q_A, \beta) - \mu_{A,0}^{L,\tau}(Q_A, \beta) & \text{if } i = 0, \\ \mu_{A,i+1}^{L,\tau}(Q_A, \beta) & \text{if } i > 0, \end{cases} \quad (4-3)$$

where Q_A sits in the short exact sequence $0 \rightarrow S_A \rightarrow S_A[\partial^{-1}] \rightarrow Q_A \rightarrow 0$. Then [Berkesch 2011, Proposition 5.10] implies that

$$\mu_{A,i}^{L,\tau}(Q_A, \beta) = \mu_{A,i}^{L,\tau}(P_J^\beta, \beta), \quad (4-4)$$

where J is equal to the right hand side of (4-2). Now [loc. cit., Lemmas 6.9, 6.10, 6.11, and 6.14] can be applied verbatim, while Lemma 4.7 replaces the need for [loc. cit., Lemma 6.13]. Finally, as [loc. cit., Lemmas 6.12 and 6.15] hold when rank is replaced with $\mu_{A,i}^{L, \overline{C}_A}$, which is possible since localization at P_τ and $\text{gr}^L(-)$ are exact functors and length is additive, the arguments of the proof of [loc. cit., Theorem 6.6] yield the desired result. In particular, the spectral sequence involved begins with the cellular resolution of P_J^β as constructed in [loc. cit., (6.3)]:

$$0 \rightarrow P_J \rightarrow I_J^0 \rightarrow I_J^1 \rightarrow \cdots \rightarrow I_J^r \rightarrow 0, \quad (4-5)$$

where $I_J^\bullet$ is constructed as follows. Set

$$\begin{aligned} \Delta_J^0 &= \{F \preceq A \mid \exists (F, b) \in \max(J)\}, \\ \Delta_J^p &= \{s \subseteq \Delta_J^0 \mid |s| = p+1\}, \text{ and} \\ F_s &= \bigcap_{G \in s} G \text{ for } s \in \Delta_J^p. \end{aligned}$$

With $r+1 = |\Delta_J^0|$, let $\Delta = \Delta_J^\beta$ be the standard r -simplex with vertices corresponding to the elements of Δ_J^0 . To the p -face of Δ spanned by the vertices corresponding to the elements in $s \in \Delta_J^p$, assign

the ranking toric module $P_{F_s, J}^\beta$. Choosing the natural maps $P_{F_s, J}^\beta \rightarrow P_{F_t, J}^\beta$ for $s \subseteq t$ induces a cellular complex supported on Δ ,

$$I_J^\bullet: I_J^0 \rightarrow I_J^1 \rightarrow \cdots \rightarrow I_J^r \rightarrow 0 \quad \text{with } I_J^p = \bigoplus_{s \in \Delta_J^p} P_{F_s, J}^\beta. \quad (4-6)$$

Applying Euler–Koszul homology to (4-6) yields a double complex. The desired spectral sequence arises from this double complex after localizing at P_τ and applying $\text{gr}^L(-)$. $\square$

Remark 4.8. If $\beta \in \mathbb{C}^d$ is such that $\max(\mathcal{J}(\beta))$ involves two faces, F_1, F_2 , then the proof of [Theorem 4.1](#) shows that

$$\mu_{A,0}^{L,\tau}(\beta) - \mu_A^{L,\tau} = \sum_{i=1}^2 (|B_{F_i}^\beta| \cdot [\text{codim}(F_i) - 1] \cdot \mu_{F_i}^{L,\tau}) + |B_G^\beta| \cdot C^\beta \cdot \mu_G^{L,\tau}, \quad (4-7)$$

where $G = F_1 \cap F_2$ and the constant C^β is given by

$$C^\beta = \binom{\text{codim}(G)}{2} - \text{codim}(G) + 1 - \binom{\text{codim}(F_1)}{2} - \binom{\text{codim}(F_2)}{2} + \binom{\text{codim}(\mathbb{C}F_1 + \mathbb{C}F_2)}{2}.$$

Example 4.9. The values of the $\mu_{A,0}^{L,\tau}(\beta)$ for a fixed β are dependent upon the choice of face $\tau \in \Phi_A^L$. For example, consider the matrix

$$A = \begin{bmatrix} 2 & 3 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 2 & 3 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \end{bmatrix},$$

and the parameter $\beta = (0, 0, -1)^t$, which lies outside the cone $\mathbb{R}_{\geq 0}A$. It turns out that

$$\widetilde{\mathbb{N}}A \setminus \mathbb{N}A = (\beta + \mathbb{N}G_1) \cap \mathbb{R}_{\geq 0}A = (\beta + \mathbb{N}G_1) \setminus \{\beta\},$$

where $G_1 = \{a_3, a_6\}$ and $G_2 = \{a_1, a_2, a_4, a_5, a_7\}$ are facets of A . In particular, $\mathcal{E}_A = \{\beta\}$ and the ranking lattices at β are

$$\mathbb{E}^\beta = (\beta + \mathbb{Z}G_1) \cup (\beta + \mathbb{Z}G_2).$$

By [Remark 4.8](#), $\mu_{A,0}^{L,\emptyset}(\beta) - \mu_A^{L,\emptyset} = 1$ for any projective weight vector L . On the other hand, $\mu_{A,0}^{L,\tau}(\beta) = \mu_A^{L,\tau}$ if $\tau \neq \emptyset$.

Example 4.10. The choice of projective weight vector impacts the resulting stratification via multiplicities of $\mathcal{E}_A$. For example, consider the matrix

$$A = \begin{bmatrix} 2 & 3 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 & 1 \end{bmatrix},$$

which has

$$\widetilde{\mathbb{N}}A \setminus \mathbb{N}A = (\beta + \mathbb{N}G_1) \cup (\beta + \mathbb{N}G_2),$$

where $G_1 = \{a_3, a_4\}$, $G_2 = \{a_5, a_6\} \preceq A$ and $\beta = (1, 0, 0)^t$. Moreover, we also have

$$\mathbb{E}^\beta = (\beta + \mathbb{Z}G_1) \cup (\beta + \mathbb{Z}G_2) \quad \text{and} \quad \mathcal{E}_A = (\beta + \mathbb{C}G_1) \cup (\beta + \mathbb{C}G_2).$$

If $L_\partial = (1, 4, 1, 4, 1, 3, 1)$ and $L_x = 5 \cdot \mathbf{1}_7 - L_\partial$, then $\mu_{A,0}^{L,\emptyset}(\beta') - \mu_A^{L,\emptyset} = 1$ for any $\beta' \in \mathcal{E}_A$. On the other hand, the stratification of $\mathcal{E}_A$ by the rank jump is different:

$$\mu_{A,0}^{F,\emptyset}(\beta') - \mu_A^{F,\emptyset} = \begin{cases} 2 & \text{if } \beta' \in (\beta + \mathbb{C}G_2) \setminus \{\beta\}, \\ 3 & \text{if } \beta' \in (\beta + \mathbb{C}G_1) \setminus \{\beta\}, \\ 4 & \text{if } \beta' = \beta. \end{cases}$$

5. More consequences of the multiplicity computation

For $\tau \in \Phi_A^L$, let

$$j_A^{L,\tau}(\beta) := \mu_{A,0}^{L,\tau}(\beta) - \mu_A^{L,\tau}$$

be the (L, τ) -multiplicity jump at β , and let

$$\mathcal{E}_A^{L,\tau} := \{\beta \in \mathbb{C}^d \mid j_A^{L,\tau}(\beta) > 0\}$$

be the (L, τ) -exceptional set of A . In this section, we record consequences of [Theorem 4.1](#) and its implications for $\mathcal{E}_A^{L,\tau}$. We also propose a description of $\mathcal{E}_A^{L,\tau}$ and prove it holds in a special case.

Corollary 5.1. *If $\tau \in \Phi_A^L$ is a face of the (A, L) -umbrella such that τ is not contained in any face of A of codimension 2, then $\mathcal{E}_A^{L,\tau} = \emptyset$.*

Proof. Fix $\beta \in \mathbb{C}^d$. By hypothesis, τ is contained in at most one facet of A . Recall that the cellular resolution of P_J^β is made of ranking toric modules P_G^β for faces $G \preceq A$ such that $\mathbb{E}_G^\beta \neq \emptyset$.

If τ is not contained in any proper face of A or it is contained in a unique facet $F \preceq A$ with $\mathbb{E}_F^\beta = \emptyset$, then [Lemma 4.7](#) guarantees that $\mu_{A,q}^{L,\tau}(P_G^\beta, \beta) = 0$ for all $q \geq 0$ for any proper face $G \preceq A$ with $\mathbb{E}_G^\beta \neq \emptyset$. Thus, the formula from [Theorem 4.1](#) computes that $\mu_{A,i}^{L,\tau}(P^\beta, \beta) = 0$ for all $i \geq 0$.

For the remaining case when τ is contained in a unique facet $F \preceq A$ and $\mathbb{E}_F^\beta \neq \emptyset$,

$$\mu_{A,i}^{L,\tau}(P^\beta, \beta) = \mu_{A,i}^{L,\tau}(P_F^\beta, \beta) = |B_F^\beta| \cdot \mu_{A,i}^{L,\tau}(\tilde{S}_F, \beta) = |B_F^\beta| \cdot \binom{1}{i} \cdot \mu_F^{L,\tau}$$

for all $i \geq 0$. Therefore, as in the proof of [Theorem 4.1](#),

$$j_A^{L,\tau}(\beta) = \mu_{A,1}^{L,\tau}(P^\beta, \beta) - \mu_{A,0}^{L,\tau}(P^\beta, \beta) = 0. \quad \square$$

Remark 5.2. As an immediate consequence of [Corollary 5.1](#), if $\dim(\mathbb{C}\tau) \geq d - 1$, then $\mu_{A,0}^{L,\tau}(\beta)$ is independent of β . Notice that this fact was known when $\dim(\mathbb{C}\tau) = d$ (see [\[Schulze and Walther 2008, Theorem 3.10\]](#)). $\square$

Corollary 5.3. *If $\tau \in \Phi_A^L$ is a face of the (A, L) -umbrella such that τ is contained in a unique face $G \preceq A$ of codimension 2, then*

$$j_A^{L,\tau}(\beta) = \begin{cases} |B_G^\beta| \cdot \mu_G^{L,\tau} & \text{if } (G, b) \in \max(\mathcal{J}(\beta)) \text{ for } b \in B_G^\beta, \text{ see (4-2),} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By the proof of [Theorem 4.1](#) and [Lemma 4.7](#),

$$j_A^{L,\tau}(\beta) = \mu_{A,1}^{L,\tau}(P^\beta, \beta) - \mu_{A,0}^{L,\tau}(P^\beta, \beta) = \mu_{A,1}^{L,\tau}(P_{J'}^\beta, \beta) - \mu_{A,0}^{L,\tau}(P_{J'}^\beta, \beta),$$

where $J' = \{(F, b) \in J \mid \tau \subseteq F\}$ for $J = \mathcal{J}(\beta)$. If $(G, b) \in \max(J)$ and $b \in B_G^\beta$, then $J' = J(G)$ and $P_{J'}^\beta = P_G^\beta$. Thus, it is enough to consider the case when $(G, b) \notin \max(J)$ for any b but there exists at least one facet F such that $\tau \subseteq G \preceq F$ and $(F, b) \in \max(J)$. In this case, either $\max(J') = J(F)$ or $J' = J(F) \cup J(F')$ for some other facet F' such that $F \cap F' = G$. Either way, it follows that $\mu_{A,1}^{L,\tau}(P_{J'}^\beta, \beta) - \mu_{A,0}^{L,\tau}(P_{J'}^\beta, \beta) = 0$. $\square$

By [Corollaries 5.1](#) and [5.3](#), if $\dim(\mathbb{C}\tau) = d - 2$, then $j_A^{L,\tau}(\beta) > 0$ only when there is a (unique) codimension 2 face G of A containing τ and $(G, b) \in \max(\mathcal{J}(\beta))$ for some $b \in B_G^\beta$.

Notation 5.4. For any $\tau \in \Phi_A^L$, let us denote $S_A^\tau := \mathbb{C}[(\mathbb{N}A + \mathbb{Z}\tau) \cap \mathbb{R}_{\geq 0}A]$.

Conjecture 5.5. *There is an equality*

$$\mathcal{E}_A^{L,\tau} = -\text{qdeg} \left(\bigoplus_{q=0}^{d-1} \text{Ext}_{\mathbb{C}[\partial]}^{n-q}(S_A^\tau, \mathbb{C}[\partial])(-\varepsilon_A) \right),$$

where $\varepsilon_A := \sum_{i=1}^n a_i$. In particular, $\mathcal{E}_A^{L,\tau} = \emptyset$ if and only if S_A^τ is Cohen–Macaulay.

As evidence of the truth of [Conjecture 5.5](#), we exhibit a containment between the two sets involved. We then prove the second part of conjecture in the case that $\mathbb{R}_{\geq 0}A$ is a simplicial cone.

Proposition 5.6. *There is a containment*

$$\mathcal{E}_A^{L,\tau} \subseteq -\text{qdeg} \left(\bigoplus_{q=0}^{d-1} \text{Ext}_{\mathbb{C}[\partial]}^{n-q}(S_A^\tau, \mathbb{C}[\partial])(-\varepsilon_A) \right).$$

Proof. By the definition of S_A^τ , it is clear that $S_A^\tau[\partial_\tau^{-1}] = S_A[\partial_\tau^{-1}]$ and thus,

$$\mu_{A,0}^{L,\tau}(S_A, \beta) = \mu_{A,0}^{L,\tau}(S_A[\partial_\tau^{-1}], \beta) = \mu_{A,0}^{L,\tau}(S_A^\tau[\partial_\tau^{-1}], \beta) = \mu_{A,0}^{L,\tau}(S_A^\tau, \beta),$$

where the first and third equalities follows from the definition of $\mu_{A,0}^{L,\tau}$ (see [\(2-1\)](#)) and the fact that $\xi_j = \text{in}_L(\partial_j) \notin P_\tau$ if and only if $j \in \tau$.

If $\beta \notin -\text{qdeg}(\text{Ext}_{\mathbb{C}[\partial]}^{n-q}(S_A^\tau, \mathbb{C}[\partial])(-\varepsilon_A))$ for any $q = 0, \dots, d - 1$, then $\mathcal{H}_i(S_A^\tau, \beta) = 0$ for all $i > 0$ by [\[Matusevich et al. 2005, Theorem 6.6\]](#). Thus, $\mu_{A,0}^{L,\tau}(S_A^\tau, \beta) = \sum_{j=0}^d (-1)^j \mu_{A,j}^{L,\tau}(S_A^\tau, \beta)$, which is independent of β by [\[Schulze and Walther 2008, Theorem 4.11\]](#) and hence equal to the generic value $\mu_A^{L,\tau}$. In particular, $\beta \notin \mathcal{E}_A^{L,\tau}$. $\square$

Proposition 5.7. Fix $\beta \in \mathbb{C}^d$ and let J be as in (4-2). If J involves only facets of A satisfying that the intersection of r of them is a face of codimension at most r , then $\mathcal{H}_q(P_J^\beta, \beta) = 0$ for all $q \geq 2$.

Proof. Consider the cellular resolution of P_J^β as constructed in [Berkesch 2011, (6.3)]:

$$0 \rightarrow P_J \rightarrow I_J^0 \rightarrow I_J^1 \rightarrow \cdots \rightarrow I_J^r \rightarrow 0,$$

where $r + 1$ is the cardinality of J . On the other hand, if $K_p := \ker(I_J^p \rightarrow I_J^{p+1})$ for $0 \leq p \leq r - 1$ and $K_r = I_J^r$, then there are short exact sequences

$$0 \rightarrow P_J \rightarrow I_J^0 \rightarrow K_1 \rightarrow 0 \quad \text{and} \quad 0 \rightarrow K_p \rightarrow I_J^p \rightarrow K_{p+1} \rightarrow 0 \quad \text{for } 1 \leq p \leq r - 1.$$

By the assumption on J , I_J^p is a direct sum of simple ranking toric modules P_G for faces G of codimension at most $p + 1$, so by [Berkesch 2011, Proposition 3.2], $\mathcal{H}_q(I_J^p, \beta) = 0$ for all $q \geq p + 2$ and $p = 0, \dots, r$. Therefore

$$\mathcal{H}_q(P_J, \beta) \cong \mathcal{H}_{q+1}(K_1, \beta) \cong \cdots \cong \mathcal{H}_{q+r-1}(K_{r-1}, \beta) \cong \mathcal{H}_{q+r}(I_J^r, \beta) = 0$$

for all $q \geq 2$, as desired. $\square$

Note that if $\mathbb{R}_{\geq 0}A$ is simplicial then any set of facets of A satisfies the property required in Proposition 5.7. To the contrary, Example 4.9 does not satisfy this property.

Theorem 5.8. Let $\tau \in \Phi_A^L$ and assume that $\mathbb{R}_{\geq 0}A$ is a simplicial cone. Then $\mathcal{E}_A^{L, \tau} = \emptyset$ if and only if S_A^τ is Cohen–Macaulay.

Proof. The if direction is proven in Proposition 5.6. By the definition of S_A^τ we have that

$$\text{rank}(\mathcal{H}_0(S_A^\tau, \beta)) = \text{vol}(A) + \mu_{A,1}^{F, \emptyset}(P_{J'}) - \mu_{A,0}^{F, \emptyset}(P_{J'}),$$

where $J' := \{(G, b) \in \mathcal{J}(\beta) \mid \tau \subseteq G\}$. If S_A^τ is not Cohen–Macaulay, then by Theorem 1.2, there exists $\beta \in \mathbb{C}^d$ such that $\text{rank}(\mathcal{H}_0(S_A^\tau, \beta)) > \text{vol}(A)$. Since $\mathbb{R}_{\geq 0}A$ is simplicial, by Proposition 5.7 there must be a face G of codimension at least 2 such that $(G, b) \in \max(J')$. Thus, for generic $\beta' \in b + \mathbb{C}G$, we have that $\max(\mathcal{J}(\beta')) = \{(G, b_1), \dots, (G, b_r)\}$ with $r = |B_G^{\beta'}|$. Now, using Lemma 4.7, we have that

$$\mu_A^{L, \tau}(S_A, \beta') = \mu_A^{L, \tau}(S_A^\tau, \beta') = \mu_A^{L, \tau} + r(\text{codim}(G) - 1) \cdot \mu_G^{L, \tau} > \mu_A^{L, \tau}$$

and thus $\beta' \in \mathcal{E}_A^{L, \tau} \neq \emptyset$. $\square$

6. Upper-semicontinuity and convex filtrations

It was conjectured in [Schulze and Walther 2008] that the multiplicities $\mu_A^{L, \tau}(\beta)$ are upper semicontinuous in $\beta \in \mathbb{C}^d$ for any projective L and $\tau \in \Phi_A^L$. We prove this conjecture when L and τ satisfy certain conditions with respect to A (see Theorem 6.1 and Corollary 6.6). We also prove Conjecture 5.5 in this setting when $\tau = \emptyset$ (see Corollary 6.5).

Given a submatrix $\sigma \subseteq A$ with rank d , denote by E_i^σ the Euler operator associated with the i -th row of the matrix σ . Let D_σ denote the Weyl algebra associated to the variables $x_\sigma = \{x_i \mid a_i \in \sigma\}$. We have that $\mathbb{Z}A = \mathbb{Z}^d = \bigoplus_{j=1}^r \Lambda_j$, where $r = [\mathbb{Z}^d : \mathbb{Z}\sigma]$ and $\Lambda_j = b_j + \mathbb{Z}\sigma$ for some $b_j \in \mathbb{Z}^d$ with $j = 1, \dots, r$.

If N is a $\mathbb{Z}^d$ -graded S_A -module, then $N_j := \bigoplus_{\alpha \in \Lambda_j} N_\alpha$ is an S_σ -module. Let $\mathcal{K}_\bullet^\sigma(N, \beta)$ denote the direct sum over j of the Euler–Koszul complexes on $D_\sigma \otimes_{\mathbb{C}[\partial_\sigma]} N_j(-b_j)$ given by the operators $\{E_i^\sigma - \beta_i + (b_j)_i\}_{i=1}^d$, where each such Euler–Koszul complex is placed in degree b_j . That is,

$$\mathcal{K}^\sigma(N, \beta) := \bigoplus_{j=1}^r \mathcal{K}^\sigma(N_j(-b_j), \beta - b_j)(b_j),$$

where the right-hand side Euler–Koszul complexes were defined before since $N_j(-b_j)$ is a $\mathbb{Z}\sigma$ -graded S_σ -module. This definition is independent of the chosen elements $b_1, \dots, b_r \in \mathbb{Z}^d$ by (1-1). With this setup, $D_\sigma \otimes N \cong \bigoplus_{j=1}^r (D_\sigma \otimes N_j)$, and $\mathcal{K}_\bullet^\sigma(N, \beta)$ is a $\mathbb{Z}^d$ -graded complex of left D_σ -modules. Set

$$\mathcal{H}_i^\sigma(N, \beta) := H_i(\mathcal{K}_\bullet^\sigma(N, \beta)),$$

and note that these definitions make (1-1) and Theorem 1.1 also valid for the homology modules $\mathcal{H}_i^\sigma(N, \beta)$.

Let L be a projective weight vector, which induces a filtration on D as considered in the introduction. We denote by A^L the submatrix of A whose columns belong to facets of Φ_A^L . We say that L is a *convex* filtration with respect to A if all facets of Φ_A^L are F -homogeneous and

$$\bigcup_{\tau' \in \Phi_A^{L, d-1}} \Delta_{\tau'} \quad (6-1)$$

is a convex polytope, and thus equal to Δ_{A^L} . Notice that, by the inclusion $S_{A^L} \subseteq S_A$, the ring S_A is an S_{A^L} -module.

Theorem 6.1. *If L is a convex filtration with respect to A , then $\mu_{A,0}^{L,\varnothing}(\beta) = \text{rank}(\mathcal{H}_0^{A^L}(S_A, \beta))$. In particular, $\mu_{A,0}^{L,\varnothing}(\beta)$ is upper-semicontinuous in β .*

Before proving Theorem 6.1, we first consider the simple case.

Proposition 6.2. *Let L be a convex filtration of D with respect to A and $G \preceq A$. Then for all $(G, b) \in \mathcal{J}(\beta)$,*

$$\mu_G^{L,\varnothing} = \text{vol}_{\mathbb{Z}G}(G^L) = \text{rank}(\mathcal{H}_0^{G^L}(P_{(G,b)}^\beta, \beta)),$$

where G^L denotes the submatrix of A whose columns belong to facets of Φ_G^L .

Proof. The first equality follows from Theorem 2.1 and (2-4) since L is convex. For the second equality, by definition of the (G, L) -umbrella Φ_G^L , the submatrix G^L of G is such that $\mathbb{R}_{\geq 0}G = \mathbb{R}_{\geq 0}G^L$ and $\text{rank}(G) = \text{rank}(G^L)$. This implies that S_G is a toric S_{G^L} -module. Further,

$$P_{(G,b)}^\beta \subseteq S_G[\partial_G^{-1}](b) \cong \bigoplus_{\alpha \in \Lambda} S_{G^L}[\partial_{G^L}^{-1}](\alpha),$$

where Λ is a finite subset of $b + \mathbb{Z}G$ of cardinality $[\mathbb{Z}G : \mathbb{Z}G^L]$. Since

$$\deg(S_G[\partial_G^{-1}](b)/P_{(G,b)}^\beta) = (b + \mathbb{Z}G) \setminus \mathbb{P}_{(G,b)}^\beta,$$

it follows from the definition of $C_A(\beta)$ that the parameter β does not belong to the quasidegrees set of the weakly toric module $S_G[\partial_G^{-1}](b)/P_{(G,b)}^\beta$. Thus, since $S_{G^L}[\partial_{G^L}^{-1}]$ is a Cohen–Macaulay S_{G^L} -module, by [Theorem 1.1](#) and [Theorem 1.2](#), $\mathcal{H}_i^{G^L}(P_{(G,b)}^\beta, \beta) = 0$ for all $i \geq 1$ and

$$\text{rank}(\mathcal{H}_0^{G^L}(P_{(G,b)}^\beta, \beta)) = [\mathbb{Z}G : \mathbb{Z}G^L] \cdot \text{vol}_{\mathbb{Z}G^L}(G^L) = \text{vol}_{\mathbb{Z}G}(G^L). \quad \square$$

Remark 6.3. Notice that any weakly toric S_A -module $M \subseteq S_A[\partial_A^{-1}]$ can be viewed as a weakly toric S_{A^L} -module. Indeed, since A^L and A have the same rank, then $\mathbb{Z}A = \bigoplus_{j=1}^r (b_j + \mathbb{Z}A^L)$ for some $b_j \in \mathbb{Z}A$ with $j = 1, \dots, r$. Thus $S_A[\partial_A^{-1}] = \bigoplus_{j=1}^r S_{A^L}[\partial_{A^L}^{-1}](b_j)$ as S_{A^L} -modules. Setting $M_j := M \cap S_{A^L}[\partial_{A^L}^{-1}](b_j)$, then M is the direct sum of the weakly toric S_{A^L} -modules M_j . Moreover, for any face $G \preceq A$,

$$\mathbb{E}_G^\beta = \bigsqcup_{b \in B_G^\beta} (b + \mathbb{Z}G) = \bigsqcup_{c \in B_{G^L}^\beta} (c + \mathbb{Z}G^L), \quad (6-2)$$

where B_G^β and $B_{G^L}^\beta$ is a set of lattice representatives (see [\(4-1\)](#)).

Lemma 6.4. *The module P^β is a direct sum of toric S_{A^L} -modules, and for any face $G \preceq A$ and $q \geq 0$,*

$$\mu_{A,q}^{L,\emptyset}(P_G^\beta, \beta) = \mu_{A^L,q}^{F,\emptyset}(P_G^\beta, \beta).$$

Proof. The decomposition of $M = S_A$ as a direct sum of weakly toric S_A^L -modules M_j given in [Remark 6.3](#) induces a decomposition of $S_A[\partial_A^{-1}]/M$ as a direct sum of the weakly toric $S_{A^L}^L$ -modules $S_{A^L}[\partial_{A^L}^{-1}](b_j)/M_j$. Then, by the two short exact sequences in the proof of [\[Berkesch 2011, Proposition 5.10\]](#), P^β is a direct sum of weakly toric S_{A^L} -modules. Moreover, since $\mathbb{P}^\beta = \mathbb{E}^\beta \cap C_A(\beta)$ and $C_A(\beta) = C_{A^L}(\beta)$, it follows that P^β is a direct sum of toric S_{A^L} -modules.

On the other hand, if G is a face of A , then by [\(6-2\)](#), $|B_G^\beta|[\mathbb{Z}G : \mathbb{Z}G^L] = |B_{G^L}^\beta|$. Thus, using [Lemma 4.7](#) and [Proposition 6.2](#),

$$\mu_{A,q}^{L,\emptyset}(P_G^\beta, \beta) = |B_G^\beta| \cdot \binom{\text{codim}(G)}{q} \cdot \text{vol}_{\mathbb{Z}G}(G^L) = |B_{G^L}^\beta| \cdot \binom{\text{codim}(G)}{q} \cdot \text{vol}_{\mathbb{Z}G^L}(G^L) = \mu_{A^L,q}^{F,\emptyset}(P_G^\beta, \beta). \quad \square$$

The proof of [Theorem 6.1](#) makes use of the notion of a *holonomic family* from [\[Matusevich et al. 2005, Definition 2.1\]](#), which we now recall. While defined over any algebraic variety B with structure sheaf $\mathcal{O}_B$, we will need only the case when $B = \mathbb{A}_{\mathbb{C}}^d$, affine d -space over $\mathbb{C}$.

If $\beta \in B$, denote by p_β the prime ideal (sheaf) of β and set $\kappa_\beta = \mathcal{O}_{B,\beta}/p_\beta \mathcal{O}_{B,\beta}$, the residue field of the stalk $\mathcal{O}_{B,\beta}$. A *coherent sheaf* of $(D \otimes_{\mathbb{C}} \mathcal{O}_B)$ -modules is a quasicohherent sheaf of $\mathcal{O}_B$ -modules on B whose sections over each open affine subset $U \subset B$ are finitely generated over the ring of global sections $H^0(B, D \otimes_{\mathbb{C}} \mathcal{O}_U)$. Let $\mathcal{O}_B(x)$ denote the localization at $\langle 0 \rangle \in \text{Spec}(\mathbb{C}[x])$ of $\mathcal{O}_B[x] := \mathbb{C}[x] \otimes_{\mathbb{C}} \mathcal{O}_B \subset D \otimes_{\mathbb{C}} \mathcal{O}_B$. The sheaf-spectrum of $\mathcal{O}_B(x)$ is the base-extended scheme $B(x) := \text{Spec } \mathbb{C}(x) \times_{\text{Spec } \mathbb{C}} B$.

A *holonomic family* over B is a coherent sheaf $\widetilde{\mathcal{M}}$ of left $(D \otimes_{\mathbb{C}} \mathcal{O}_B)$ -modules such that

- (1) the fibers $\mathcal{M}_\beta = \widetilde{\mathcal{M}} \otimes_{\mathcal{O}_B} \kappa_\beta$ are holonomic D -modules for all $\beta \in B$, and
 (2) $\mathcal{O}_B(x) \otimes_{\mathcal{O}_B} \widetilde{\mathcal{M}}$ is coherent on $B(x)$.

Proof of Theorem 6.1. Since $S_A[\partial^{-1}]$ is a maximal Cohen–Macaulay weakly toric S_{A^L} -module,

$$\mathcal{H}_i^{A^L}(S_A[\partial^{-1}], \beta) = 0$$

for all $i > 0$ by Theorem 1.2. Thus, applying Euler–Koszul homology with respect to A^L to the short exact sequence

$$0 \rightarrow S_A \rightarrow S_A[\partial^{-1}] \rightarrow Q \rightarrow 0$$

and using that $\mathcal{H}_q^{A^L}(P^\beta, \beta) \simeq \mathcal{H}_q^{A^L}(Q, \beta)$ (see the proof of [Berkesch 2011, Proposition 5.10], which can be adapted to this case), it follows that

$$\text{rank}(\mathcal{H}_0^{A^L}(S_A, \beta)) = \text{rank}(\mathcal{H}_0^{A^L}(S_A[\partial^{-1}], \beta)) + \mu_{A^L,1}^{F,\varnothing}(P^\beta, \beta) - \mu_{A^L,0}^{F,\varnothing}(P^\beta, \beta).$$

The proofs of [loc. cit., Theorem 6.6] and Theorem 4.1 and the induction argument in the proof of [loc. cit., Proposition 6.18] reduces the computation of

$$\mu_{A^L,1}^{F,\varnothing}(P^\beta, \beta) - \mu_{A^L,0}^{F,\varnothing}(P^\beta, \beta) \quad (\text{and respectively } \mu_{A,1}^{L,\varnothing}(P^\beta, \beta) - \mu_{A,0}^{L,\varnothing}(P^\beta, \beta))$$

to that of $\mu_{A^L,q}^{F,\varnothing}(N, \beta)$ (and respectively $\mu_{A,q}^{L,\varnothing}(N, \beta)$) for $q \geq 0$ and simple toric modules $N = P_G^\beta$ with $\mathbb{E}_G^\beta \neq \varnothing$. Thus, by Lemma 6.4,

$$\mu_{A^L,1}^{F,\varnothing}(P^\beta, \beta) - \mu_{A^L,0}^{F,\varnothing}(P^\beta, \beta) = \mu_{A,1}^{L,\varnothing}(P^\beta, \beta) - \mu_{A,0}^{L,\varnothing}(P^\beta, \beta) = \mu_{A,0}^{L,\varnothing}(\beta) - \mu_A^{L,\varnothing}$$

which yields the desired equality.

Finally, since S_A is a toric S_{A^L} -module, $\mathcal{H}_0^{A^L}(S_A, b)$ is a holonomic family by [Matusевич et al. 2005, Theorem 7.5]. Hence [loc. cit., Theorem 2.6] guarantees that

$$\beta \mapsto \text{rank}(\mathcal{H}_0^{A^L}(S_A, \beta))$$

is an upper semicontinuous function. □

Theorem 6.1 provides a way to prove Conjecture 5.5 when L is a convex filtration of D with respect to A and $\tau = \varnothing$.

Corollary 6.5. *If L is a convex filtration of D with respect to A , then*

$$\mathcal{E}_A^{L,\varnothing} = -\text{qdeg} \left(\bigoplus_{q=0}^{d-1} \text{Ext}_{\mathbb{C}[\partial]}^{n-q}(S_A, \mathbb{C}[\partial])(-\varepsilon_A) \right).$$

Proof. By the proof of Theorem 6.1, $\mathcal{H}_0^{A^L}(S_A, b)$ is a holonomic family and $\mathcal{E}_A^{L,\varnothing} = \mathcal{E}_A^{F,\varnothing}$, and thus by [Matusевич et al. 2005, Theorem 9.1],

$$\mathcal{E}_A^{L,\varnothing} = \deg \left(2 \bigoplus_{i=0}^{d-1} H_{\mathfrak{m}_L}^i(S_A) \right)^{\text{Zariski}},$$

where $\mathfrak{m}_L$ denotes the maximal homogeneous ideal in S_{A^L} . However, since $\mathbb{R}_{\geq 0}A = \mathbb{R}_{\geq 0}A^L$, the radical of the extended ideal $\mathfrak{m}_L S_A$ in S_A equals $\mathfrak{m}$. Therefore, by applying graded Matlis duality, we obtain the desired result. $\square$

Let L be a filtration on D induced by a projective weight vector. For $\tau \in \Phi_A^L$, we denote by $A^{L,\tau}$ the submatrix of A whose columns belong to facets $\tau' \in \Phi_A^{L,d-1}$ such that $\tau \subseteq \tau'$. We say that L is τ -convex if all facets of Φ_A^L containing τ are F -homogeneous and the polytope

$$\bigcup_{\tau \subseteq \tau' \in \Phi_A^{L,d-1}} \Delta_{\tau'} \quad (6-3)$$

is convex, and thus equal to $\Delta_{A^{L,\tau}}$.

We recall that a subset $\eta' \subseteq A$ is said to be a *pyramid* over $\eta \subseteq \eta'$ if

$$\text{rank}_{\mathbb{Z}}(\mathbb{Z}\eta) + |\eta' \setminus \eta| = d,$$

where we denote by $|\lambda|$ the cardinality of a set λ .

[Theorem 6.1](#) can now be generalized as follows.

Corollary 6.6. *If L induces a τ -convex filtration for some $\tau \in \Phi_A^L$ and any $\tau' \in \Phi_A^{L,d-1}$ such that $\tau \subseteq \tau'$ is a pyramid over $\tau' \setminus \tau$, then*

$$\mu_{A,0}^{L,\tau}(\beta) = \text{rank}(\mathcal{H}_0^{A^{L,\tau}}(S_A, \beta)).$$

In particular, $\mu_{A,0}^{L,\tau}(\beta)$ is upper-semicontinuous in β .

Proof. Recall the formula in [Theorem 2.1](#). For any $\tau' \in \Phi_A^L$ containing τ , since τ' is a pyramid over $\tau' \setminus \tau$, it follows that

$$\mathbb{Z}\tau' \cap \mathbb{Q}\tau = \mathbb{Z}\tau, \quad \pi_{\tau,\tau'}(\mathbb{Z}\tau') = \mathbb{Z}(\tau' \setminus \tau), \quad P_{\tau,\tau'} = \Delta_{\tau' \setminus \tau},$$

$\mathcal{Q}_{\tau,\tau'}$ is the convex hull of $\tau' \setminus \tau$ (whose volume is zero because τ' is F -homogeneous), and $\text{vol}_{\mathbb{Z}\tau'}(\tau') = \text{vol}_{\mathbb{Z}(\tau' \setminus \tau)}(\tau' \setminus \tau)$. Thus, for any face $G \preceq A$ that contains τ ,

$$\mu_G^{L,\tau} = \sum_{\tau \subseteq \tau' \in \Phi_G^{L,d-1}} [\mathbb{Z}G : \mathbb{Z}\tau'] \cdot \text{vol}_{\mathbb{Z}\tau'}(\Delta_{\tau'}) = \text{vol}_{\mathbb{Z}G} \left(\bigcup_{\tau \subseteq \tau' \in \Phi_G^{L,d-1}} \Delta_{\tau'} \right) = \text{vol}_{\mathbb{Z}G}(\Delta_{G^{L,\tau}}).$$

When $(G, b) \in \mathcal{J}(\beta)$, to obtain the equality

$$\text{rank}(\mathcal{H}_0^{G^{L,\tau}}(P_{(G,b)}^\beta, \beta)) = \text{vol}_{\mathbb{Z}G}(\Delta_{G^{L,\tau}}) \quad (6-4)$$

we can proceed as in the proof of [Proposition 6.2](#), but now $\mathbb{R}_{\geq 0}G$ is not equal to $\mathbb{R}_{\geq 0}G^{L,\tau}$, so S_A is only a direct sum of weakly toric $S_{A^{L,\tau}}$ -modules (by [Remark 6.3](#)) instead of a toric $S_{A^{L,\tau}}$ -module. On the other hand, in the proof of [Theorem 6.1](#) we can use P_J^β with $J = \{(G, b) \in \mathcal{J}(\beta) \mid \tau \subseteq G\}$ instead of P^β and consider each P_G^β as a direct sum of weakly toric Cohen–Macaulay $S_{G^{L,\tau}}$ -modules.

Finally, by [Schulze and Walther 2009, Remark 5.5.(5)], in the analytic topology, $\mathcal{H}_0^{A^{L,\tau}}(S_A, \beta)$ is locally a holonomic family on $\mathbb{A}^d$. This fact along with [Matusevich et al. 2005, Theorem 2.6] and Theorem 4.1 imply that the function $\beta \mapsto \text{rank}(\mathcal{H}_0^{A^{L,\tau}}(S_A, \beta))$ is upper-semicontinuous. $\square$

7. Gevrey series solutions associated to slopes

Let $\mathcal{D}$ be the sheaf of linear partial differential operators with coefficients in the sheaf $\mathcal{O}_X^{\text{an}}$ of holomorphic functions on $X = \mathbb{C}^n$. The irregularity sheaf of order $s > 1$ of a holonomic $\mathcal{D}$ -module $\mathcal{M}$ along a hypersurface Y was introduced and proved to be a perverse sheaf on Y by Mebkhout [1990]. In particular, higher cohomology of the irregularity sheaf vanishes at generic points of Y .

In this section, for a coordinate hyperplane $Y \subset X$, we compute the dimension of the stalk at a generic point $p \in Y$ of the irregularity sheaf of order s of $\mathcal{M}_A(\beta) := \mathcal{D} \otimes_D M_A(\beta)$ along Y for any parameter $\beta \in \mathbb{C}^d$, generalizing results from [Fernández-Fernández 2010]. As a consequence, we provide some formulas for the dimension of the Gevrey solution spaces of $\mathcal{M}_A(\beta)$ in particular cases, and we show that the dimension of the generic stalk of the irregularity sheaf of $\mathcal{M}_A(\beta)$ along Y is upper-semicontinuous in β .

We assume for simplicity that $Y = \text{Var}(x_n)$ and write s instead of $L(s)$ for the filtration given by $L(s) := F + (s-1)V_n$ with $s \geq 1$, where $F = (\mathbf{0}_n, \mathbf{1}_n)$ is the filtration by the order of the differential operators and V_n is the Kashiwara–Malgrange filtration along Y . Recall that this filtration is induced by the weight vector $V_n := (0, \dots, 0, -1, 0, \dots, 0, 1)$, where -1 is the weight for the variable x_n . More precisely, the filtration $L(s)$ is determined by

$$\deg_s \partial_i = \begin{cases} 1 & \text{if } 1 \leq i \leq n-1, \\ s & \text{if } i = n, \end{cases} \quad \text{and} \quad \deg_s(x_i) = 1 - \deg_s(\partial_i).$$

In this section, we call the $(A, L(s))$ -umbrella the (A, s) -umbrella, and we denote $\Phi_A^s := \Phi_A^{L(s)}$ for $s \geq 1$.

A global version of Laurent’s slope theory [1987] proceeds as follows. Let M be a holonomic D -module. A number $s > 1$ is said to be a *slope* of M along $Y = \text{Var}(x_n)$ if and only if the s -characteristic variety $\text{Char}^s(M)$ of M along Y is not homogeneous with respect to the weight vector $F = (\mathbf{0}_n, \mathbf{1}_n)$.

Remark 7.1. Denote by A' the submatrix of A defined by the first $n-1$ columns and by Δ' the convex hull of the columns of A' and the origin. Note that a_n/s belongs to a hyperplane off the origin that contains a facet of $\Delta_{A'}$ if and only if there exists a facet of the (A, s) -umbrella, in other words an element of $\Phi_A^{s,d-1}$, that is not F -homogeneous. Moreover, by [Schulze and Walther 2008, Corollary 4.18], this condition holds if and only if $s > 1$ is a slope of $M_A(\beta)$ along $\text{Var}(x_n)$.

Let $\widehat{\mathcal{O}_{X|Y}}$ denote the formal completion of $\mathcal{O}_X$ along Y . A *germ* $f \in \widehat{\mathcal{O}_{X|Y}, p}$ with $p \in Y$ is a formal series

$$f = \sum_{m=0}^{\infty} f_m(x_1, \dots, x_{n-1}) x_n^m$$

such that there exists some open subset $U \subseteq \mathbb{C}^{n-1}$ so that f_m is a holomorphic function in U for all $m \geq 0$. The formal series $f \in \mathcal{O}_{\widehat{X|Y},p}$ is said to be a *Gevrey series* of order $s \in \mathbb{R}$ along Y at $p \in Y$ if the series

$$\rho_s(f) := \sum_{m=0}^{\infty} \frac{f_m(x_1, \dots, x_{n-1})}{(m!)^{s-1}} x_n^m$$

is convergent at p . Moreover, if $\rho_{s'}(f)$ is not convergent at p for any $s' < s$, then s is said to be the *Gevrey index* of f along Y at p . Denote by $\mathcal{O}_{X|Y}(s)$ the subsheaf of $\mathcal{O}_{\widehat{X|Y}}$ whose germs are Gevrey series of order s along Y .

The *irregularity sheaf* of a $\mathcal{D}$ -module $\mathcal{M}$ along Y of order $s > 1$ is

$$\mathrm{Irr}_Y^{(s)}(\mathcal{M}) := \mathbb{R}\mathrm{Hom}_{\mathcal{D}}(\mathcal{M}, \mathcal{O}_{\widehat{X|Y}}(s)/\mathcal{O}_{X|Y}).$$

For $s = \infty$, the sheaf $\mathrm{Irr}_Y^{\infty}(\mathcal{M})$ is simply called the *irregularity sheaf* of $\mathcal{M}$ along Y . If M is a D -module, we define $\mathrm{Irr}_Y^{(s)}(M) := \mathrm{Irr}_Y^{(s)}(\mathcal{M})$, where $\mathcal{M} := \mathcal{D} \otimes_D M$.

Set $d_s(A, \beta) := \dim H^0(\mathrm{Irr}_Y^{(s)}(M_A(\beta)))_p$ for a generic point $p \in Y = \mathrm{Var}(x_n)$. Applying Théorème 2.3.1 and (2.3.1) in [Laurent and Mebkhout 1999] to this setting yields the equality

$$d_s(A, \beta) = \mu_{A,0}^{s+\epsilon, \emptyset}(\beta) - \mu_{A,0}^{1+\epsilon, \emptyset}(\beta) + \mu_{A,0}^{1+\epsilon, \{n\}}(\beta) - \mu_{A,0}^{s+\epsilon, \{n\}}(\beta) \quad (7-1)$$

for $\epsilon > 0$ small enough. In particular, if β is not rank-jumping for A , then by Theorem 2.1 and [Fernández-Fernández 2010, Theorem 7.5], $d_s(A, \beta)$ is equal to

$$d_s(A) := \mu_A^{s+\epsilon, \emptyset} - \mu_A^{1+\epsilon, \emptyset} + \mu_A^{1+\epsilon, \{n\}} - \mu_A^{s+\epsilon, \{n\}} = \sum_{n \notin \tau \in \Phi_A^{s+\epsilon, d-1} \setminus \Phi_A^{1+\epsilon, d-1}} \mathrm{vol}_{\mathbb{Z}^d}(A_{\tau}). \quad (7-2)$$

Remark 7.2. Notice that (7-2) also holds for any face $G \preceq A$ in place of A when $a_n \in G$. Moreover, $d_s(G) = \dim H^0(\mathrm{Irr}_Y^{(s)}(M_G(\beta'))_p$ for a generic point $p \in Y' = \mathrm{Var}(x_n) \subseteq \mathbb{C}^G$ and $\beta' \in \mathbb{C}^G$ that is not rank-jumping for G . The genericity condition on p requires that it avoids any other irreducible component of the singular locus of $M_G(\beta')$ (which is independent of β' as a consequence of Theorem 2.2). On the other hand, if $a_n \notin G$, then the coordinates indexed by G of the projective weight vectors $L(s)$ and F are the same. Hence the two induced filtrations over (any cyclic module over) the Weyl algebra in the variables indexed by G are also the same. Thus, $\mu_G^{1+\epsilon, \tau} = \mu_G^{s+\epsilon, \tau}$ for $\tau = \{n\}$ and $\tau = \emptyset$ in this case, so $d_s(G) = 0$.

Proposition 7.3. For any $\beta \in \mathbb{C}^d$, there is a lower bound $d_s(A, \beta) \geq d_s(A)$.

Proof. For a $\mathbb{Z}^d$ -graded $\mathbb{C}[\partial]$ -module N , define $d_s^{(j)}(N, \beta) := \dim H^0(\mathrm{Irr}_Y^{(s)}(\mathcal{H}_j(N, \beta)))_p$ for a generic point $p \in Y = \mathrm{Var}(x_n)$. Then by the same argument as in (7-1),

$$d_s^{(j)}(N, \beta) = \mu_{A,j}^{s+\epsilon, \emptyset}(N, \beta) - \mu_{A,j}^{1+\epsilon, \emptyset}(N, \beta) + \mu_{A,j}^{1+\epsilon, \{n\}}(N, \beta) - \mu_{A,j}^{s+\epsilon, \{n\}}(N, \beta) \quad (7-3)$$

for $\epsilon > 0$ small enough. Notice that $d_s(A, \beta) = d_s^{(0)}(S_A, \beta)$. By [Schulze and Walther 2008, Corollary 4.13] and (7-3), $d_s^{(0)}(\tilde{S}_A, \beta) = d_s(A)$. Moreover, if $j \geq 1$, then $d_s^{(j)}(\tilde{S}_A, \beta) = 0$ because $\mathcal{H}_j(\tilde{S}_A, \beta) = 0$ by Theorem 1.2.

On the other hand, if N is a toric module with dimension lower than d , it follows that

$$d_s^{(0)}(N, \beta) \leq d_s^{(1)}(N, \beta)$$

by the same argument as in the proof of [Schulze and Walther 2008, Lemma 4.29], with the replacement, for each D -module M that appears in that proof, of the role of $\mathrm{CC}^L(M)$ by $\dim H^0(\mathrm{Irr}_Y^{(s)}(M)_p)$ for a generic point $p \in Y = \mathrm{Var}(x_n)$. This is allowable because $H^1(\mathrm{Irr}_Y^{(s)}(M)_p) = 0$ for generic points $p \in Y$ when M is holonomic (see [Mebkhout 1990]). Thus, with the previous ingredients, the proof of [Schulze and Walther 2008, Theorem 4.28] gives the result with $d_s^{(0)}$ in place of $\mu_{A,0}^{L,\tau}$. $\square$

Corollary 7.4. *For $s > 1$, the dimension $d_s(A, \beta)$ of the stalk of $\mathrm{Irr}_Y^{(s)}(M_A(\beta))$ at a generic point p of Y can be computed from the combinatorics of $\Phi_A^{s+\epsilon} \setminus \Phi_A^{1+\epsilon}$ for $\epsilon > 0$ small enough and the ranking lattices $\mathbb{E}_G^\beta$ at β such that $a_n \in G \preceq A$.*

Proof. It follows from (7-1) and Theorem 4.1 that $d_s(A, \beta)$ can be computed from the combinatorics of the (A, s') -umbrellas for $s' \in \{1 + \epsilon, s + \epsilon\}$ and the ranking lattices $\mathbb{E}^\beta$. Thus, by Remark 7.2, it is enough to consider the ranking lattices $\mathbb{E}_G^\beta$ at β corresponding to the faces $G \preceq A$ containing a_n . $\square$

We now state further consequences for $d_s(A, \beta)$.

Corollary 7.5. *If $P^\beta = P_G^\beta$ for some $G \preceq A$, then*

$$d_s(A, \beta) = d_s(A) + |B_G^\beta| \cdot (\mathrm{codim}(G) - 1) \cdot d_s(G).$$

In particular, if $a_n \notin G$ or $\mathrm{codim}(G) = 1$, then $d_s(A, \beta) = d_s(A)$.

Proof. It is a direct consequence of (4-3), (4-4), Lemma 4.7, (7-1), (7-2), and Remark 7.2. $\square$

Corollary 7.6. *If $d = 2$, then $d_s(A, \beta) = d_s(A)$ for any $\beta \in \mathbb{C}^d$.*

Proof. Since $d = 2$, the matrix A has only two proper faces $G_1, G_2 \preceq A$, which both have codimension 1. Moreover, a_n belongs to at most one of these two facets. Thus, by Corollaries 7.4 and 7.5, it is enough to consider the case when $a_n \in G_1$ and $\max(\mathcal{J}(\beta))$ involves G_1 . In this case, $d_s(A, \beta)$ can be computed as in the simple case, so the formula in Corollary 7.5 can be applied, giving $d_s(A, \beta) = d_s(A)$ since $\mathrm{codim}(G_1) = 1$. $\square$

Notice that Corollary 7.6 also follows from [Schulze and Walther 2008, Proposition 4.25] and (7-1).

Corollary 7.7. *If $d = 3$, then $d_s(A, \beta) > d_s(A)$ if and only if $\max(\mathcal{J}(\beta))$ involves a face G with $a_n \in G$ and $\dim G = 1$. If this is the case, $d_s(A, \beta) = d_s(A) + |B_G^\beta| \cdot d_s(G)$.*

Proof. Again by Corollary 7.4, we only need to consider the ranking lattices $\mathbb{E}_G^\beta$ such that $a_n \in G$. Thus, by the reduction given in [Berkesch 2011, Section 5.3], it is enough to prove the result in the following two cases.

The first case is that a_n belongs to a unique face G among those involved in $\max(\mathcal{J}(\beta))$. In this case, the computation follows as in the simple case, and we obtain the same formula as in Corollary 7.5.

In the second case, we may assume that there are exactly two faces G_1 and G_2 involved in $\max(\mathcal{J}(\beta))$ that contain a_n . Since the face $G_1 \cap G_2$ contains a_n and $d = 3$, it follows that G_1 and G_2 are two facets intersecting in a face of codimension 2. In this case, [Remark 4.8](#) shows that $\mu_{A,0}^{L,\tau}(\beta) = \mu_A^{L,\tau}$ for any filtration L and any $\tau \in \Phi_A^L$, so $d_s(A, \beta) = d_s(A)$. $\square$

Lemma 7.8. *Let $s > 1$ be such that the (A, s) -umbrella Φ_A^s has a unique facet τ that is not F -homogeneous, p is a generic point of $Y = \text{Var}(x_n)$, and $\epsilon > 0$ small enough. Then the function*

$$\beta \mapsto d(A, \beta, s) := \dim \mathcal{H}om_{\mathcal{D}}(\mathcal{M}_A(\beta), \mathcal{O}_{\widehat{X|Y}}(s + \epsilon) / \mathcal{O}_{\widehat{X|Y}}(s - \epsilon))_p$$

is upper-semicontinuous.

Proof. Notice first that by the assumption and [Remark 7.1](#), s is a slope of $M_A(\beta)$ along Y and $a_n \in \tau$. Indeed, the assumption implies that $\tau' := \tau \setminus \{n\}$ is the unique facet of $\Phi_A^{s+\epsilon}$ that does not contain a_n and is also not a facet of $\Phi_A^{s-\epsilon}$. On the other hand,

$$d(A, \beta, s) = d_{s+\epsilon}(A, \beta) - d_{s-\epsilon}(A, \beta) = \mu_{A,0}^{s+\epsilon,\emptyset}(\beta) - \mu_{A,0}^{s-\epsilon,\emptyset}(\beta) + \mu_{A,0}^{s-\epsilon,\{n\}}(\beta) - \mu_{A,0}^{s+\epsilon,\{n\}}(\beta).$$

Thus, setting $d(A, s) := \mu_{A,0}^{s+\epsilon,\emptyset} - \mu_{A,0}^{s-\epsilon,\emptyset} + \mu_{A,0}^{s-\epsilon,\{n\}} - \mu_{A,0}^{s+\epsilon,\{n\}}$ yields

$$d(A, s) = \text{vol}_{\mathbb{Z}A}(\Delta_{\tau'}) = \text{rank}(\mathcal{H}_0^{\tau'}(\tilde{S}_A, \beta)),$$

where the first equality follows by the assumption, (2-4), and [\[Fernández-Fernández 2010, Lemma 7.4\]](#). The second equality follows as in the proof of (6-4), since $A_{\tau'}$ is a rank d submatrix of A . Similarly, for faces G of A such that $a_n \in G$ and $\tau'' := \tau' \cap G$ is a facet of $\Phi_G^{s+\epsilon}$, we also have that $d(G, s) = \text{rank}(\mathcal{H}_0^{\tau''}(\tilde{S}_G, \beta))$. Thus, arguments similar to those in [Corollary 6.6](#) show that $d(A, \beta, s) = \text{rank}(\mathcal{H}_0^{\tau'}(S_A, \beta))$ and that the function $\beta \mapsto d(A, \beta, s) = \text{rank}(\mathcal{H}_0^{\tau'}(S_A, \beta))$ is upper-semicontinuous in β . $\square$

Theorem 7.9. *Assume that for all $s > 1$, a_n/s is in at most one of the hyperplanes off the origin supported in a facet of Δ' (see [Remark 7.1](#)). Then the function $\beta \mapsto d_s(A, \beta)$ is upper-semicontinuous for all $s > 1$.*

Proof. Let $1 < s_1 < \dots < s_r$ be the set of slopes of $M_A(\beta)$ along Y that are lower or equal to s . Then $d_s(A, \beta) = \sum_{j=1}^r d(A, \beta, s_j)$, and the result follows by [Lemma 7.8](#). $\square$

In view of the preceding results we state the following conjecture.

Conjecture 7.10. *The map $\beta \mapsto d_s(A, \beta)$ is upper-semicontinuous. Moreover, there is an equality*

$$\mathcal{E}_A^n(s) := \{\beta \in \mathbb{C}^d \mid d_s(A, \beta) > d_s(A)\} = -\text{qdeg} \left(2 \bigoplus_{q=0}^{d-1} \text{Ext}_{\mathbb{C}[\partial]}^{n-q}(S_A^{[n]}, \mathbb{C}[\partial])(-\varepsilon_A) \right),$$

where $\varepsilon_A := \sum_{i=1}^n a_i$. In particular, $\mathcal{E}_A^n(s) = \emptyset$ if and only if $S_A^{[n]}$ is Cohen–Macaulay.

The values $d_s(A, \beta)$ and $d_s(A)$ defined in this section depend on the variety Y along which we are considering the irregularity sheaf of $M_A(\beta)$. Although we assumed $Y = \text{Var}(x_n)$ for simplicity, we can consider any $Y_j := \text{Var}(x_j) \subseteq \mathbb{C}^n$ since reordering the variables is equivalent to reordering the columns

of A . Let $d_s(A, \beta, j)$ and $d_s(A, j)$ denote the values of $d_s(A, \beta)$ and $d_s(A)$ respectively for Y_j in place of Y . In the following example, we compute the difference $d_s(A, \beta, j) - d_s(A, j)$ for different j by using [Corollary 7.7](#).

Example 7.11. Let us consider the matrix A in [Example 4.10](#). The hyperplanes contained in the singular locus of $M_A(\beta)$ are exactly Y_j for $j \in \{2, 4, 6, 7\}$ and there is exactly one slope $s_j \geq 1$ of $M_A(\beta)$ along each Y_j . More precisely, by [Remark 7.1](#), $s_2 = \frac{3}{2}$, $s_4 = 3$, $s_6 = 2$, and $s_7 = \frac{7}{6}$. It is clear that $d_s(A, \beta', j) = 0$ if $1 \leq s < s_j$ for any $\beta' \in \mathbb{C}^d$, so let us assume that $s \geq s_j$ in each case. We have that $d_s(A, \beta', j) = d_s(A, j)$ for all β' and $j \in \{2, 7\}$. On the other hand, $d_s(A, \beta', 4) - d_s(A, 4)$ is 1 if $\beta' \in \beta + \mathbb{C}G_1$ and 0 otherwise. Finally, $d_s(A, \beta', 6) - d_s(A, 6)$ is 1 if $\beta' \in \beta + \mathbb{C}G_2$ and zero otherwise.

One natural problem after the computation of $d_s(A, \beta) = m$ is to construct an explicit set of Gevrey series $\varphi_1, \dots, \varphi_m$ along Y at a nonsingular point $p \in Y$ so that their classes in the space $(\mathcal{O}_{\widehat{X|Y}}(s)/\mathcal{O}_{X|Y})_p$ form a basis of $H^0(\text{Irr}_Y^{(s)}(M_A(\beta)))_p$. This was done in [\[Fernández-Fernández 2010\]](#) when β is generic enough. At any parameter β , this problem is much more involved in general. However, it is easy to compute some examples by using a slightly modified version of a method used in [\[Fernández-Fernández 2013\]](#). In order to do so, recall that the direct sum of two matrices $A_1 \in \mathbb{Z}^{d_1 \times n_1}$, $A_2 \in \mathbb{Z}^{d_2 \times n_2}$ is the following $(d_1 + d_2) \times (n_1 + n_2)$ matrix:

$$A_1 \oplus A_2 = \begin{pmatrix} A_1 & \mathbf{0}_{d_1 \times n_2} \\ \mathbf{0}_{d_2 \times n_1} & A_2 \end{pmatrix},$$

where $\mathbf{0}_{d \times n}$ denotes the $d \times n$ zero matrix. Let $\beta = (\beta^{(1)}, \beta^{(2)})$ denote a complex vector in $\mathbb{C}^{d_1+d_2} \cong \mathbb{C}^{d_1} \times \mathbb{C}^{d_2}$. It is easy to show using [\[Fernández-Fernández 2013, Lemma 2.2\]](#) that

$$d_s(A, \beta, n_1) = d_s(A_1, \beta^{(1)}, n_1) \cdot \text{rank}(M_{A_2}(\beta^{(2)})).$$

Now, let us take $(A_1, \beta^{(1)})$ such that $M_{A_1}(\beta^{(1)})$ has slopes along $\{x_{n_1} = 0\}$, and let consider the subset of Gevrey series $\{g_1, \dots, g_{r(1)}\} \subseteq \mathcal{O}_{\widehat{X|Y}}(s)$ whose classes form a basis of

$$H^0(\text{Irr}_{\{x_{n_2}=0\}}^{(s)}(M_{A_1}(\beta^{(1)})))_p.$$

Let us take also a pair $(A_2, \beta^{(2)})$ for which a basis $\{f_1, \dots, f_{r(2)}\}$ of convergent series solutions of $M_{A_2}(\beta^{(2)})$ at a nonsingular point p' is known for a rank-jumping parameter $\beta^{(2)} \in \mathbb{C}^{d_2}$. Then $\{g_i f_j \mid 1 \leq i \leq r(1), 1 \leq j \leq r(2)\}$ is a basis of

$$H^0(\text{Irr}_{\{x_{n_2}=0\}}^{(s)}(M_{A_1 \oplus A_2}(\beta)))_{(p, p')},$$

where $\beta = (\beta^{(1)}, \beta^{(2)})$. Note that

$$\dim H^0(\text{Irr}_{\{x_{n_2}=0\}}^{(s)}(M_{A_1 \oplus A_2}(\beta)))_{(p, p')} = r(1) \cdot r(2) > d_s(A, n_1) = d_s(A_1, n_1) \cdot \mu_{A_2, 0}^F.$$

In particular, the smallest example of this family is the one obtained by taking $A = A_1 \oplus A_2$ for $A_1 = (1 \ 2)$ and $A_2 = (\tilde{0}, \tilde{1}, \tilde{3}, \tilde{4})$, where $\tilde{a} = (1, a)^t$ and $\beta = (b, 1, 2)^t$ for any $b \in \mathbb{C} \setminus \mathbb{Z}$. We notice that $M_{A_2}((1, 2)^t)$ was the first example known of an A -hypergeometric system for which the rank is greater than the

normalized volume [Sturmfels and Takayama 1998]. Indeed, a basis of $H^0(\mathrm{Irr}_{\{x_{n_2}=0\}}^{(s)}(M_{A_1}(b))_p)$ is $\{\bar{\phi}_v\} \subset (\mathcal{O}_{\widehat{X|Y}}(s)/\mathcal{O}_{X|Y})_p$, where ϕ_v is the Γ -series associated to $v = (b, 0)$ (see [Fernández-Fernández 2010]) and $\mathrm{rank}(M_{A_2}(\beta^{(2)})) = \mathrm{vol}_{\mathbb{Z}^2}(A_2) + 1 = 5$ (see [Sturmfels and Takayama 1998], where a basis of solutions is also described). Thus, in this case, $M_A(\beta)$ has the slope $s = 2$ along $x_2 = 0$ and for $s \geq 2$, $d_s(A, \beta, 2) = d_s(A, 2) + 1 = 5$.

Acknowledgements

We are grateful to Francisco Jesús Castro Jiménez, Laura Felicia Matusevich, and Uli Walther for helpful conversations related to this work. Fernández-Fernández would like to thank the School of Mathematics of the University of Minnesota for the hospitality during her visit to work on this paper with Berkesch.

References

- [Adolphson 1994] A. Adolphson, “Hypergeometric functions and rings generated by monomials”, *Duke Math. J.* **73**:2 (1994), 269–290. [MR](#) [Zbl](#)
- [Berkesch 2011] C. Berkesch, “The rank of a hypergeometric system”, *Compos. Math.* **147**:1 (2011), 284–318. [MR](#) [Zbl](#)
- [Berkesch et al. 2018] C. Berkesch, J. Forsgård, and L. F. Matusevich, “On the parametric behavior of A -hypergeometric series”, *Trans. Amer. Math. Soc.* **370**:6 (2018), 4089–4109. [MR](#) [Zbl](#)
- [Fernández-Fernández 2010] M.-C. Fernández-Fernández, “Irregular hypergeometric $\mathcal{D}$ -modules”, *Adv. Math.* **224**:5 (2010), 1735–1764. [MR](#) [Zbl](#)
- [Fernández-Fernández 2013] M.-C. Fernández-Fernández, “Exponential growth of rank jumps for A -hypergeometric systems”, *Rev. Mat. Iberoam.* **29**:4 (2013), 1397–1404. [MR](#) [Zbl](#)
- [Gelfand et al. 1987] I. M. Gelfand, M. I. Graev, and A. V. Zelevinsky, “Holonomic systems of equations and series of hypergeometric type”, *Dokl. Akad. Nauk SSSR* **295**:1 (1987), 14–19. In Russian; translated in *Soviet Math. Dokl.* **36**:1 (1988), 5–10. [MR](#)
- [Gelfand et al. 1989] I. M. Gelfand, A. V. Zelevinsky, and M. M. Kapranov, “Hypergeometric functions and toric varieties”, *Funkt. Anal. i Prilozhen.* **23**:2 (1989), 12–26. In Russian; translated in *Funct. Anal. Appl.* **23**:2 (1989), 94–106. Correction in *Funkt. Anal. i Prolozhen* **27**:4 (1993), 91; correction translated in *Funct. Anal. Appl.* **27**:4 (1993), 295. [MR](#)
- [Gelfand et al. 1990] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, “Generalized Euler integrals and A -hypergeometric functions”, *Adv. Math.* **84**:2 (1990), 255–271. [MR](#) [Zbl](#)
- [Hochster 1972] M. Hochster, “Rings of invariants of tori, Cohen–Macaulay rings generated by monomials, and polytopes”, *Ann. of Math. (2)* **96** (1972), 318–337. [MR](#) [Zbl](#)
- [Laurent 1987] Y. Laurent, “Polygône de Newton et b -fonctions pour les modules microdifférentiels”, *Ann. Sci. École Norm. Sup. (4)* **20**:3 (1987), 391–441. [MR](#) [Zbl](#)
- [Laurent and Mebkhout 1999] Y. Laurent and Z. Mebkhout, “Pentes algébriques et pentes analytiques d’un $\mathcal{D}$ -module”, *Ann. Sci. École Norm. Sup. (4)* **32**:1 (1999), 39–69. [MR](#) [Zbl](#)
- [Matusevich et al. 2005] L. F. Matusevich, E. Miller, and U. Walther, “Homological methods for hypergeometric families”, *J. Amer. Math. Soc.* **18**:4 (2005), 919–941. [MR](#) [Zbl](#)
- [Mebkhout 1990] Z. Mebkhout, “Le théorème de positivité de l’irrégularité pour les $\mathcal{D}_X$ -modules”, pp. 83–132 in *The Grothendieck Festschrift, III*, edited by P. Cartier et al., Progr. Math. **88**, Birkhäuser, Boston, 1990. [MR](#) [Zbl](#)
- [Saito et al. 2000] M. Saito, B. Sturmfels, and N. Takayama, *Gröbner deformations of hypergeometric differential equations*, Algor. Comput. Math. **6**, Springer, 2000. [MR](#) [Zbl](#)
- [Schulze and Walther 2008] M. Schulze and U. Walther, “Irregularity of hypergeometric systems via slopes along coordinate subspaces”, *Duke Math. J.* **142**:3 (2008), 465–509. [MR](#) [Zbl](#)

- [Schulze and Walther 2009] M. Schulze and U. Walther, “Hypergeometric D-modules and twisted Gauß–Manin systems”, *J. Algebra* **322**:9 (2009), 3392–3409. [MR](#) [Zbl](#)
- [Sturmfels and Takayama 1998] B. Sturmfels and N. Takayama, “Gröbner bases and hypergeometric functions”, pp. 246–258 in *Gröbner bases and applications* (Linz, Austria, 1998), edited by B. Buchberger and F. Winkler, Lond. Math. Soc. Lecture Note Ser. **251**, Cambridge Univ. Press, 1998. [MR](#) [Zbl](#)

Communicated by Gavril Farkas

Received 2019-02-12 Revised 2019-09-25 Accepted 2019-11-13

cberkesc@umn.edu

*School of Mathematics, University of Minnesota, Minneapolis, MN,
United States*

mcferfer@algebra.us.es

*Departamento de Álgebra e Instituto de Matemáticas-IMUS,
Universidad de Sevilla, Spain*

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Anand Pillay	University of Notre Dame, USA
Samit Dasgupta	Duke University, USA	Michael Rapoport	Universität Bonn, Germany
Hélène Esnault	Freie Universität Berlin, Germany	Victor Reiner	University of Minnesota, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Sergey Fomin	University of Michigan, USA	Michael Singer	North Carolina State University, USA
Edward Frenkel	University of California, Berkeley, USA	Christopher Skinner	Princeton University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Ben J. Green	University of Oxford, UK	Shunsuke Takagi	University of Tokyo, Japan
Joseph Gubeladze	San Francisco State University, USA	Pham Huu Tiep	University of Arizona, USA
Christopher Hacon	University of Utah, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Marie-France Vignéras	Université Paris VII, France
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 14 No. 2 2020

On the definition of quantum Heisenberg category	275
JONATHAN BRUNDAN, ALISTAIR SAVAGE and BEN WEBSTER	
Characteristic cycles and Gevrey series solutions of A -hypergeometric systems	323
CHRISTINE BERKESCH and MARÍA-CRUZ FERNÁNDEZ-FERNÁNDEZ	
Singularity categories of deformations of Kleinian singularities	349
SIMON CRAWFORD	
Iwasawa main conjecture for Rankin–Selberg p -adic L -functions	383
XIN WAN	
Positivity results for spaces of rational curves	485
ROYA BEHESHTI and ERIC RIEDL	
Generalized Schur algebras	501
ALEXANDER KLESHCHEV and ROBERT MUTH	