

Observation of Road Salt Aerosol Driving Inland Wintertime Atmospheric Chlorine Chemistry

Stephen M. McNamara, Katheryn R. Kolesar, Siyuan Wang, Rachel M. Kirpes, Nathaniel W. May, Matthew J. Gunsch, Ryan D. Cook, Jose D. Fuentes, Rebecca S. Hornbrook, Eric C. Apel, Swarup China, Alexander Laskin, and Kerri A. Pratt*



Cite This: *ACS Cent. Sci.* 2020, 6, 684–694



Read Online

ACCESS |

Metrics & More

Article Recommendations


Supporting Information

ABSTRACT: Inland sources of particulate chloride for atmospheric nitryl chloride (ClNO_2) formation remain unknown and unquantified, hindering air quality assessments. Globally each winter, tens of millions of tons of road salt are spread on roadways for deicing. Here, we identify road salt aerosol as the primary chloride aerosol source, accounting for 80–100% of ClNO_2 formation, at an inland urban area in the wintertime. This study provides experimental evidence of the connection between road salt and air quality through the production of this important reservoir for nitrogen oxides and chlorine radicals, which significantly impact atmospheric composition and pollutant fates. A numerical model was employed to quantify the contributions of chloride sources to ClNO_2 production. The traditional method for simulating ClNO_2 considers chloride to be homogeneously distributed across the atmospheric particle population; yet, we show that only a fraction of the particulate surface area contains chloride. Our new single-particle parametrization considers this heterogeneity, dramatically lowering overestimations of ClNO_2 levels that have been routinely reported using the prevailing methods. The identification of road salt as a ClNO_2 source links this common deicing practice to atmospheric composition and air quality in the urban wintertime environment.



1. INTRODUCTION

In coastal urban areas, dinitrogen pentoxide (N_2O_5), formed by NO_2 reaction with the photolabile nitrate radical (NO_3 , reaction R1), reacts on the surface of chloride (Cl^-)-containing sea spray aerosol particles to produce nitryl chloride (ClNO_2 , R2).^{1–3} Upon sunrise, ClNO_2 photolysis releases chlorine atoms (Cl) and nitrogen dioxide (NO_2) (R3), altering pollutant fate, ozone levels, and particulate nitrate formation.^{1,4}



While this chlorine chemistry was originally thought to only occur in marine and coastal locations, Thornton et al.⁵ first showed observations of ClNO_2 far inland (near Boulder, Colorado, 1400 km inland). Model predictions, constrained to measurements of particulate chloride across the United States, indicate that the majority of ClNO_2 formation occurs over land, particularly during winter.^{5,6} However, the source(s) of inland particulate chloride are highly uncertain.^{2,7}

The wintertime application of road salt for deicing purposes is ubiquitous,^{8,9} with tens of millions of tons of salt applied globally each year.^{8–12} In the United States, over 20 million tons of salt were applied to roadways in 2014 alone.⁸ This road

salt contributes to chloride-containing atmospheric particles,¹¹ saline snowpacks,¹³ and urban grime.^{14,15} Road salt is mechanically aerosolized by vehicular traffic,¹⁶ with deposition observed hundreds of meters away from roadways.¹⁷ Observed increases in wintertime $\text{PM}_{2.5}$ (particulate matter $<2.5 \mu\text{m}$) chloride concentrations are often correlated with snowfall, due to the application of deicing salts before, during, and after snowfall events.¹¹ These wintertime inland $\text{PM}_{2.5}$ chloride concentrations can even rival those of coastal areas influenced by sea spray.¹¹ For flights downwind from New York City, NY, after a snow storm, Haskins et al.¹⁸ reported observations of gas phase Cl_y ($\equiv \text{HCl} + \text{ClNO}_2 + \text{HOCl} + 2\text{Cl}_2$) that could not be explained by sea salt displacement, suggesting road salt influence. In Calgary, Alberta, where road salts are used, increasing ClNO_2 was observed following the first snowfall in November 2010,¹⁹ suggesting that road salt could be an important chloride source for ClNO_2 production. Further, the slower thermal dissociation of N_2O_5 (R1, reverse) under the lower winter temperatures,²⁰ when road salts are used, leads to

Received: September 30, 2019

Published: May 13, 2020



greater importance of this chemistry in wintertime environments. However, the contribution of road salt chloride to ClNO_2 formation is unknown,² despite being hypothesized for several decades.¹

To model ClNO_2 production in the atmosphere, laboratory-based measurements of N_2O_5 uptake ($\gamma_{\text{N}_2\text{O}_5}$) and ClNO_2 yields (φ_{ClNO_2}) have informed the current parametrization based on aerosol mass concentrations of chloride, nitrate, and liquid water content.²¹ This “bulk” parametrization²¹ assumes that all aerosol chemical components are equally distributed across the entire aerosol size distribution and that all particles contain chloride.²² Yet, even in the polluted marine atmosphere, chloride is not homogeneously mixed across the entire aerosol population.^{23,24} The $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values are often overestimated when compared to field-derived values,^{5,25–29} including recent comprehensive assessments by McDuffie et al.^{30,31} Further, observed variability and trends in $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} are often not reproduced, pointing to an inaccurate parametrization and reducing the accuracy of associated air quality simulations.^{26,27,30,31}

2. RESULTS AND DISCUSSION

2.1. Winter Measurements of ClNO_2 and Identification of Road Salt Aerosol. From February 1 to March 10, 2016 in Ann Arbor, Michigan, ClNO_2 production was investigated using a comprehensive suite of atmospheric trace gas and particle measurements. Using chemical ionization mass spectrometry (CIMS),³² N_2O_5 and ClNO_2 were continuously measured at 12 m above ground level, with average nighttime levels of 141 ± 7 ppt (pmol mol^{-1} , 95% confidence interval) and 23 ± 1 ppt, respectively (Figure S1 and Section S2.1 of the Supporting Information). Nocturnal ClNO_2 reached a maximum of 220 ppt. Lower temperatures in the winter favor N_2O_5 (reducing thermal dissociation),²⁰ and the application of road salt for deicing provides a unique chloride source. Previously, in Calgary, Alberta, Mielke et al.^{19,33} observed elevated ClNO_2 in March and April (up to 338 and 250 ppt, respectively), compared to September (max. 30 ppt), prior to road salt application. Similarly, in Ann Arbor, N_2O_5 and ClNO_2 levels were lower in October 2016, with average nighttime levels of 71 ± 5 ppt and 3.5 ± 0.5 ppt, respectively, and maximum ClNO_2 of 29 ppt (see Section S2.2).

To test the hypothesis that road salt aerosol chloride contributes to ClNO_2 production, we present comprehensive observations of individual particle chemical composition with source attribution, complemented by modeling of two representative case periods (February 17–18 and March 7–8, 2016). These case periods encompass a range of observed ClNO_2 levels, and no precipitation or rapid changes in wind speed and direction occurred, making them well-suited for numerical modeling. Although surface-level observations can vary from those made aloft,^{34,35} ClNO_2 and N_2O_5 mole ratios during the case periods were not impacted by N_2O_5 depletion from NO_3^- titration,³⁶ or morning ClNO_2 enhancement due to entrainment from the residual layer³⁷ (Figure S1). The model results, using our new parametrization, show that the ClNO_2 observed in Ann Arbor was predominantly produced from N_2O_5 reaction with particulate chloride from widespread road salt application, thereby quantifying the link between wintertime deicing practices and chlorine activation (Figure 1).



Figure 1. The role of road salt aerosol in wintertime ClNO_2 production in inland areas. Vehicular traffic causes lofting of road salt, providing a chloride-containing particle surface for N_2O_5 uptake, producing ClNO_2 at night. Upon sunrise, ClNO_2 photolysis leads to chlorine radical formation and NO_x recycling, linking the common deicing practice to wintertime inland air quality.

Individual particle measurements of chemical composition and morphology were conducted using computer-controlled scanning electron microscopy with energy dispersive X-ray spectroscopy (CCSEM-EDX)³⁸ and aerosol time-of-flight mass spectrometry (ATOFMS),³⁹ with 8,052 and 83,597 individual particles analyzed by each method, respectively. These methods identified and quantified the chloride-containing particles. This analysis complemented bulk $\text{PM}_{2.5}$ inorganic ion measurements using ambient ion monitor-ion chromatography (AIM-IC).⁴⁰ Concurrent measurements of atmospheric $\text{PM}_{2.5}$ chloride ranged from 0.01 to $0.35 \mu\text{g m}^{-3}$ (Section S2.1), similar to concentrations during elevated ClNO_2 levels in Boulder, CO.⁵

Five individual particle types—road salt, aged road salt, biomass burning, soot, and road dust (Figures 2 and S2, and discussed in Section 4.3)—were observed by ATOFMS and CCSEM-EDX. Only road salt particles contained significant chloride. Road salt particles were primarily composed of sodium and chloride (average Cl/Na atomic ratio of 0.87 ± 0.03 , 95% confidence interval, determined by CCSEM-EDX), consistent with road salt used by the city of Ann Arbor and the University of Michigan for winter maintenance (Section S2.3). Sulfate, known to suppress aerosol ClNO_2 production,⁴¹ was not detected in the individual nascent road salt particles. Aged road salt particles were depleted in chloride (average Cl/Na atomic ratio of 0.10 ± 0.01) and enriched in nitrate and sulfate, compared to the nascent road salt particles. Biomass burning particles have been suggested as a possible ClNO_2 source, depending on their fuel chloride content.⁴² However, these particles were identified as a mixture of mainly organic carbon, nitrate, sulfate, ammonium, and potassium (Figure S2),⁴³ and contained less than 2% chlorine (atomic percentage), representing a minor chloride source. Therefore, the observed ammonium was primarily in the form of ammonium nitrate and sulfate, rather than ammonium chloride (see Section S2.4). Less than 3% of road dust particles, by number, and no soot particles contained chloride (Section 4.5.2). The frequent nightly observations of elevated N_2O_5 and ClNO_2 , concurrent with nascent and aged road salt particles, suggest that ClNO_2 is mainly formed from the multiphase reaction of N_2O_5 on road salt aerosols, as confirmed by the model simulations described below.

2.2. Considering Aerosol Heterogeneity in Model Simulations of ClNO_2 Production. A one-dimensional




Figure 2. Identification and quantitation of nascent and aged road salt aerosol. Representative EDX spectra and SEM images and average ATOFMS mass spectra of individual (a, b) nascent and (c, d) aged road salt particles. *Al and Si peaks in the EDX spectra are from substrate and detector backgrounds. Aged road salt is characterized by chloride depletion and nitrate and/or sulfate enrichment. (e) Average aerosol surface area fractions (0.015–20 μm) attributed to the five particle types identified by CCSEM-EDX and ATOFMS for the February 17–18 and March 7–8 cases.

atmospheric numerical model simulated ClNO_2 production constrained by measurements of N_2O_5 , HCl , and aerosol surface area and composition for the February 17–18 and March 7–8 cases. The influence of aerosol chemical composition on the uptake of N_2O_5 ($\gamma_{\text{N}_2\text{O}_5}$) and yield of ClNO_2 (φ_{ClNO_2}) was accounted for through two different model scenarios: (1) using the “traditional” parametrization by Bertram & Thornton,²¹ which is based on bulk particulate inorganic ion mass concentrations, here constrained by hourly $\text{PM}_{2.5}$ measurements using AIM-IC, and (2) a new parametrization developed based on measured single-particle chemical composition. Our single-particle measurements showed that chloride-containing particles, dominated by nascent and aged road salt, only contributed 20–26% of the aerosol surface area concentration (Figure 2), suggesting that the bulk parametrization should overestimate ClNO_2 by assuming that all particles contain chloride. Indeed, for southeast Michigan, previously simulated mean wintertime ClNO_2 mole ratios⁶ (using the bulk parametrization) were up to an order of magnitude higher than our observations.

With knowledge of the distribution of chloride among individual particles, the new single particle parametrization, described in Section 4.5, considers that N_2O_5 reacts with a heterogeneous aerosol population, producing ClNO_2 only from particles that contain chloride. We assigned chemically specific $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values to the five observed particle types using the most relevant laboratory model systems^{20,21,44–46} (Table S1). Since N_2O_5 reacts on the surface of particles,³ these $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values were then weighted by the fraction of the measured temporally varying aerosol surface area concentration corresponding to each particle type. This method results in significantly lower estimates of $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} compared to the bulk parametrization, by directly

accounting for additional chemical components that alter $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} , such as sulfate, carboxylate, and nitrate,^{21,41} within the heterogeneous individual particle population. A primary advantage of this single-particle approach is that only chloride-containing particles can produce ClNO_2 , and that aerosol components that suppress production are accounted for within the individual particle types. The main uncertainty in the single-particle method comes from the choice of laboratory proxy $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values for the observed particle types, and this uncertainty can be reduced with further laboratory studies of varying aerosol composition.

The average $\gamma_{\text{N}_2\text{O}_5}$ values calculated for the single-particle method ($\gamma_{\text{N}_2\text{O}_5} = 0.018$ and 0.013 for February 17–18 and March 7–8) were approximately half of those predicted by the bulk method ($\gamma_{\text{N}_2\text{O}_5,\text{bulk}} = 0.030$ and 0.031 , respectively). Likewise, the average φ_{ClNO_2} values for the single-particle method ($\varphi_{\text{ClNO}_2} = 0.28$ and 0.20 for the two cases, respectively) were about one-third of the bulk chemical composition method ($\varphi_{\text{ClNO}_2,\text{bulk}} = 0.8 \pm 0.1$ and 0.7 ± 0.2 , respectively). Notably, the single-particle-weighted $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} are both closer to previous wintertime field-based estimates ($\gamma_{\text{N}_2\text{O}_5} = 0.02$; $\varphi_{\text{ClNO}_2} = 0.005–0.15$) in Calgary, Alberta,^{19,33} where road salting is also commonplace.

By constraining to measurements of N_2O_5 , two model simulations of ClNO_2 production were conducted: using the (1) traditional bulk parametrization and (2) new single-particle parametrization, which employs the chemically dependent $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values. The predicted ClNO_2 mole ratios using the single-particle parametrization generally agree within the measurement uncertainty of the observations, while the bulk parametrization overpredicts ClNO_2 by an average of 120




Figure 3. Single-particle parametrization of N_2O_5 uptake ($\gamma_{\text{N}_2\text{O}_5}$) and ClNO_2 yield (φ_{ClNO_2}) improves model agreement with measured ClNO_2 in Ann Arbor, Michigan. (a) Schematic comparing the bulk and single-particle methods to parametrize N_2O_5 uptake and ClNO_2 yield. For both (b) February 17–18 and (c) March 7–8 cases, the modeled ClNO_2 , using $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} from the single-particle chemical composition and surface area parametrization (solid green lines), agrees well with the magnitude and shape of the ClNO_2 measurements (black lines), while the traditional bulk aerosol composition parametrization (gray lines) significantly overpredicts ClNO_2 . The dashed blue lines represent modeled ClNO_2 from road salt aerosol (nascent + aged) only, demonstrating that road salt aerosol is the dominant ClNO_2 source (80–100%).

± 20 ppt (340%) and 12 ± 3 ppt (150%), respectively, for the February and March cases (Figure 3). The results of the single-particle method identify aerosolized road salt as the dominant ClNO_2 source, accounting for 80–100%, on average, of the total simulated ClNO_2 (Figure 3). This study demonstrates road salt as the dominant source of photolabile chlorine in the wintertime inland environment, despite only representing a fraction (20–26%) of the aerosol surface area available for N_2O_5 uptake and ClNO_2 production (Figure 2). Application of this new parametrization to modeling in both marine and inland environments will improve predictions of ClNO_2 production because chloride is typically not distributed equally among all particles, even in the coastal marine atmosphere.²³

3. ATMOSPHERIC IMPLICATIONS

For decades, road salt use for wintertime deicing has been increasing across North America^{8,9} and has led to increased salinity of surface and ground waters with severe ecosystem

impacts.⁴⁷ Here, this common deicing practice is linked to wintertime air quality due to the role of road salt chloride as a large and potent source of photolabile chlorine. Single-particle measurements identified aerosolized road salt as the dominant source of particulate chloride. This road salt aerosol was calculated to be responsible for 80–100% of atmospheric ClNO_2 in this inland wintertime environment. The key to the improved modeling of ClNO_2 is for only the chloride-containing particle surface area to produce ClNO_2 , as was achieved with our single-particle parametrization. Laboratory studies of additional model aerosols, as well as authentic road salt aerosol, are needed to measure $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} to improve the accuracy of the application of the single-particle parametrization. Although the chloride-containing particles were primarily fresh road salt in this study, environments influenced by other particle types may need to consider additional single-particle properties. For example, different classes of organic compounds can result in particle phase separation,⁴⁸ impact

N_2O_5 uptake,^{49,50} and/or limit chloride availability,⁵¹ thereby altering ClNO_2 production. If additional properties are known or can be modeled, then appropriate laboratory $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} can be applied through the single-particle, surface area-weighted approach, which can also be applied to other multiphase reactions. While additional measurements of single-particle chemical composition are needed for other environments influenced by chlorine chemistry, modeling approaches that describe the distribution of chemical components across the aerosol population, from zero-dimensional to regional/global model levels, are now available²² and should be used to further evaluate the single-particle, surface area-weighted approach presented here.

The chlorine radicals produced from ClNO_2 photolysis, which contributed 44–99% of the daytime Cl production rate (Figure S3), impact the fate of atmospheric hydrocarbons, particularly in the winter when hydroxyl radicals are less abundant due to reduced sunlight and water vapor.⁵² In addition, ClNO_2 photolysis also significantly alters the levels of NO_x , O_3 , and particulate matter,^{53,54} impacting downwind areas struggling with wintertime air quality.⁵⁵ Other road salt contaminated surfaces, including buildings, road surfaces, and snowpacks, which were not explored in this study, likely also support ClNO_2 production¹⁵ and should be examined in future work. Future efforts are also needed to quantify road salt emissions and generate inventories for models, as this substantial chloride source is currently not included in emissions inventories for atmospheric modeling of nitryl chloride,^{6,56} thereby limiting the quantitation of the broad implications of this chlorine chemistry.

4. METHODS

Atmospheric sampling of trace gases and particles was conducted at 12 m above ground level (agl) in Ann Arbor, MI, at the University of Michigan (UM, 42.2786°N, 83.7369°W) from February 1 to March 10, 2016. ClNO_2 , N_2O_5 , HNO_3 , and Cl_2 were monitored using CIMS (Section 4.1 and Section S1.1), and soluble inorganic trace gases, including HCl and SO_2 , and $\text{PM}_{2.5}$ Cl^- and NO_3^- were measured using the AIM-IC system (Section S1.2). Measurements of individual particle chemical composition were conducted online using ATOFMS (Section 4.2.1) and offline using CCSEM-EDX (Section 4.2.2). Sections S1.3–S1.6 of the Supporting Information contain additional methods details, including measurements of NO , volatile organic compounds, and aerosol size distributions. Single-particle identifications are described in Section 4.3. Measurements and calculations pertaining to the single-particle parametrization for ClNO_2 formation are described in Sections 4.4 and 4.5. Two 24 h case study periods (February 17–18 and March 7–8, 2016), each starting and ending at 12:00 EST, were investigated using a one-dimensional atmospheric model described in Section 4.6.

Roadway and sidewalk deicing maintenance using road salt and brine was regularly conducted adjacent to the sampling site by the city of Ann Arbor and UM. Meteorological data (temperature, relative humidity, wind speed, and wind direction) were obtained from a weather station (42.2769°N 83.7655°W) located 2.3 km west of the measurement site. Wind speed and direction were measured at ~12 m agl. Temperature and relative humidity were measured at ~5 m agl. No unexpected or unusually high safety hazards were encountered.

4.1. Chemical Ionization Mass Spectrometry. A chemical ionization mass spectrometer (CIMS, THS Instruments)³² quantified ambient ClNO_2 , N_2O_5 , and HNO_3 from February 1 to March 10, 2016, and again from October 23–28, 2016 for comparison to autumn conditions prior to road salting. The CIMS used $\text{I}(\text{H}_2\text{O})_n$ reagent ions to react with analytes, forming iodide ion adducts that were subsequently detected by a quadrupole mass analyzer. Ambient air was pulled through a heated (25–30 °C), 0.95 cm I.D., 2.5 m long FEP tube at 7.1 L min^{-1} , at atmospheric pressure, into a custom three-way valve used for calibration and background measurements, as described by Liao et al.³² The CIMS ion molecule reaction region (IMR, maintained at 19 Torr) sampled 0.9 L min^{-1} , and an ozone monitor (model 205, 2B Technologies) sampled 1.7 L min^{-1} , with the remaining flow sent to exhaust. The $\text{I}^-(\text{H}_2\text{O})_n$ reagent ions were formed by passing 5 L min^{-1} of CH_3I (5 ppm in N_2 , Scott-Marrin, Inc.) through a ^{210}Po source (20 mCi), forming I^- , and combining with humidified N_2 (from a room temperature 1 L bubbler) in the CIMS IMR. Water vapor was added, as in previous work,⁵⁷ to prevent fluctuations in CIMS sensitivity due to ambient humidity. The following ions were monitored (with dwell times noted): ClNO_2 as $\text{I}^{35}\text{ClNO}_2^-$ at m/z 208 (1 s) and $\text{I}^{37}\text{ClNO}_2^-$ at m/z 210 (0.5 s), N_2O_5 as IN_2O_5^- at m/z 235 (0.5 s), HNO_3 as IHNO_3^- at m/z 190 (0.2 s), and Cl_2 as $\text{I}^-(^{35}\text{Cl}^{35}\text{Cl})$ at m/z 197 (1 s) and $\text{I}^-(^{37}\text{Cl}^{35}\text{Cl})$ at m/z 199 (0.5 s). The measured isotopic ratio for $^{37}\text{ClNO}_2$ to $^{35}\text{ClNO}_2$ (0.31) confirmed the identity of the observed ClNO_2 (theoretical ratio = 0.32). Ambient Cl_2 was not observed above the detection limit during the study and was therefore not quantified. Reagent ions were monitored at m/z 147 ($\text{IH}_2^{18}\text{O}^-$). Considering all monitored masses, the total measurement time for the CIMS was 14.6 s. Background measurements were conducted for 2 min every 15 min by passing the airflow through a glass wool and stainless steel wool scrubber heated to 120 °C, removing ClNO_2 and N_2O_5 with >95% efficiency, and HNO_3 with 88% efficiency.

Online Cl_2 calibration was completed every 2 h by adding 200 mL min^{-1} of $190 \pm 10 \text{ ppb Cl}_2$ (in N_2) from a permeation source (VICI Metronics) to the ambient airflow. The Cl_2 permeation rate (110 ng min^{-1}) was confirmed every 1–2 days using the optical absorption method described by Liao et al.³² The Cl_2 sensitivity at m/z 197 was $3.0 \pm 0.4 \text{ Hz ppt}^{-1}$. Calibrations of HNO_3 , N_2O_5 , and ClNO_2 were completed offline using calibration factors relative to Cl_2 (Section S1.1). Limits of detection (LODs, 3σ), corresponding to 2 min background periods, were 0.9, 4, 34, and 2 ppt for ClNO_2 , N_2O_5 , HNO_3 , and Cl_2 , respectively. Taking into account counting statistics,³² we report 10 min averaged LODs of 0.4, 2, 15, and 1 ppt for ClNO_2 , N_2O_5 , HNO_3 , and Cl_2 , respectively. Measurement uncertainties, which include the propagated calibration uncertainties and fluctuations in CIMS background signals, for 10 min averaged ClNO_2 , N_2O_5 , and HNO_3 were 17% + 0.4 ppt, 17% + 2 ppt, and 27% + 15 ppt, respectively. Following the study, the CIMS sampling line was characterized for wall losses of ClNO_2 , N_2O_5 , and HNO_3 by flowing each trace gas through the sampling line and comparing to its direct injection into the CIMS inlet. Measured line losses for each species ($\text{ClNO}_2 = 20 \pm 7\%$ (\pm standard deviation), $\text{N}_2\text{O}_5 = 14 \pm 15\%$, and $\text{HNO}_3 = 3 \pm 9\%$) were used to adjust the CIMS measured values to ambient mole ratios. The CIMS sampling line was tested postcampaign for potential inlet artifacts, and little to no ClNO_2 generation was observed following the

addition of N_2 , N_2O_5 , and Cl_2 into the sampling line, as discussed in [Section S1.1](#).

4.2. Measurements of Individual Particle Chemical Composition and Morphology. 4.2.1. *Aerosol Time-of-Flight Mass Spectrometry (ATOFMS).* An aerosol time-of-flight mass spectrometer (ATOFMS), based on the design of Pratt et al.,³⁹ measured individual particle size and chemical composition for particles 0.1–1.5 μm (vacuum aerodynamic diameter) from February 3 to March 9, 2016. Air was sampled from a manifold (shared with additional aerosol instruments, [Section S1.4](#)) through a 0.17 cm ID copper sampling line at 0.1 L min^{-1} . The operation of the ATOFMS is described in detail elsewhere.³⁹ Positive and negative ion mass spectra were collected for 83,597 individual particles during the study. Mass spectral peak lists were generated using custom LabVIEW software and imported into Matlab for analysis with YAADA, a custom toolkit. Individual particle mass spectra were clustered using an ART-2a algorithm with a vigilance factor of 0.8 and a learning rate of 0.05 for 20 iterations.⁵⁸ Particle types were identified based on the most likely m/z from previous field and laboratory studies ([Section 4.3](#)).

4.2.2. *Computer-Controlled Scanning Electron Microscopy with Energy Dispersive X-ray Spectroscopy (CCSEM-EDX).* Atmospheric particles were impacted onto transmission electron microscopy (TEM) grids (Formvar Type-B Copper grids, Ted Pella, Inc.) and aluminum foil substrates (MSP Corp.) using a rotating 10-stage micro-orifice uniform deposit impactor (MOUDI, model 110R, MSP Corp.). Ambient air flowed at 20 L min^{-1} through a 3.7 m long, 1.1 cm ID unheated, insulated copper sampling inlet with 10 L min^{-1} of particle-free dilution air sampled through a HEPA capsule (Pall Laboratory), resulting in a 30 L min^{-1} flow into the MOUDI. Particles analyzed herein were collected on stages with 50% efficiency upper-limit size cuts of 3.2, 1.0, 0.32, and 0.10 μm (aerodynamic diameter, d_a). Individual particles were analyzed by computer-controlled scanning electron microscopy with energy dispersive X-ray spectroscopy (CCSEM-EDX) using FEI Quanta SEM and FEI Helios SEM instruments. Both instruments were equipped with a field emission gun operating at 20 keV and high angle annular dark field scanning TEM detectors and Everhart-Thornley secondary electron detectors, for analysis of morphology, including projected area diameter, of particles on TEM grids and aluminum foil substrates, respectively. An EDX detector (EDAX, Inc.) collected X-ray spectra for individual particles to determine relative atomic abundances of C, N, O, Na, Mg, P, S, Cl, K, Ca, Ti, Fe, and Ni. Samples from the two case days were analyzed by CCSEM-EDX: February 17–18, 2016 19:00–07:45 EST (2,915 particles analyzed) and March 7–8, 2016 19:30–07:15 EST (5,137 particles analyzed). K-means cluster analysis of the individual particle EDX spectra³⁸ resulted in 50 clusters, which were grouped into particle classes based on elemental composition. Particle classes were identified based on similarity to individual particle EDX spectral signatures in published literature.³⁸ The CCSEM-EDX relative atomic percentages for Cl and Na were used to calculate individual road salt particle Cl/Na atomic ratios.

4.3. Identification of Atmospheric Particle Types from Single-Particle Measurements. The clustering analyses of ATOFMS and CCSEM-EDX individual particle spectra ([Section 4.2](#)) resulted in the identification of five major particle types: road salt, aged road salt, biomass burning, soot, and road dust ([Figure 2](#) and [Figure S2](#)). Road salt was

characterized by intense Na and Cl peaks in the EDX spectra;^{38,59} by ATOFMS, road salt was identified by an intense m/z 23 (Na^+) peak, as well as less intense ion peaks at m/z 39 (K^+), 40 (Ca^+), 81 (Na_2Cl^+), –35/–37 (Cl^-), and –93/95 (NaCl_2^-) ([Figure 2](#)). The road salt mass spectra were consistent with previous ATOFMS ambient observations,⁶⁰ as well as our laboratory measurements of local road salt and brine samples, as discussed in [Section S2.3](#). Aged road salt also contained N and/or S, in addition to Na, and was depleted in Cl, as determined by EDX. Through ATOFMS analysis, aged road salt was also characterized by positive ions at m/z 23 (Na^+), 39 (K^+), and 40 (Ca^+), with negative ion peaks at m/z –46 (NO_2^-), –62 (NO_3^-), and –97 (HSO_4^-), indicative of atmospheric processing.⁶¹ Biomass burning particles, attributed to residential wood burning,⁴³ were identified by ATOFMS by an intense peak at m/z 37 (C_3H^+) and 39 (K^+), with less intense organic carbon peaks at m/z 27 (C_2H_3^+), 43 ($\text{C}_2\text{H}_3\text{O}^+$), and 50 (C_4H_2^+), as well as ammonium (m/z 18 (NH_4^+)) ([Figure S2](#)). The EDX spectra of these particles were characterized by intense C and O peaks, in addition to S, N, and/or K.^{62,63} Road dust particles were identified by intense Fe and Al and/or Si peaks, in addition to O, in the EDX spectra;³⁸ corresponding ATOFMS spectra were characterized by an intense peak at m/z 56 (Fe^+), with less intense peaks at m/z 72 (FeO^+) and 88 (FeO_2^+), with iron likely from vehicular brake wear consistent with local road dust ([Figure S2](#)).⁶⁴ Using CCSEM-EDX, soot particles were primarily composed of carbon and characterized by chain-like agglomerate morphology ([Figure S2](#)).^{62,65} Corresponding soot particle mass spectra contained elemental carbon peaks at m/z 12, 24, 36, 48, and 60 (C^+ , C_2^+ , C_3^+ , C_4^+ , C_5^+); these particles are consistent with local vehicle combustion.^{66,67}

4.4. Calculation of Chemically Resolved Surface Area Concentrations. Combined aerosol size distributions were obtained, using the method of Khlystov et al.,⁶⁸ assuming a shape factor of 1 and a density of 1.5 g cm^{-3} , using measurements from a scanning mobility particle sizer spectrometer (SMPS) for 15–600 nm particles (mobility diameter, d_m) and an aerodynamic particle sizer (APS) for 0.60–20 μm (d_a) particles ([Section S1.4](#)). This results in continuous particle size distributions from 0.015 to 20 μm (d_a). Total particle surface area concentrations were then calculated based on these size-resolved number concentrations for the February 17–18 and March 7–8 cases. These surface area concentrations were then scaled by the size-resolved number fractions of the five particle types (identified in [Section 4.3](#)) to calculate time-resolved fractions of the aerosol surface area corresponding to each particle type using the following method.

First, particle diameters measured by CCSEM-EDX were converted from projected area diameter to d_a using the method of Wagner & Leith⁶⁹ ([Section S1.5](#)). Then, the number fractions of the five particle types, determined by CCSEM-EDX for 0.13–2.29 μm , were applied to each size bin of the full aerosol size distribution, measured by SMPS and APS for 0.015–20 μm , to calculate size-resolved number fractions for each particle type. Since the measurements of aerosol size distributions extended down to 0.015 μm , the number fractions of each particle type below 0.13 μm were assumed to be the same as for the smallest CCSEM-EDX size bin (0.13–0.17 μm), which consisted of 20% road salt, 25% biomass burning, 2% aged road salt, 29% soot, and 24% road dust. Likewise, the aerosol size distribution measurements

extended up to 20 μm ; therefore, particles larger than 2.29 μm were assumed to be the same as for the number fractions of particle types from the largest CCSEM-EDX size bin (1.84–2.29 μm), which consisted of 70% road salt, 3% biomass burning, 10% aged road salt, 0% soot, and 18% road dust. Particles smaller than 0.13 μm composed 46 \pm 11% and 36 \pm 3%, on average, of the total particle surface area concentrations during the February 17–18 and March 7–8 case days, respectively. The percentage of total surface area corresponding to 2.29–20 μm particles was an average of 3 \pm 2% during February 17–18 and 2 \pm 1% during March 7–8.

For each case period, the size-resolved particle type number fractions were scaled by the time-varying particle number distributions (0.015–20 μm , d_a , [Section S1.4](#)) using the method described by Reinard et al.⁷⁰ This generated number concentrations for each of the five particle types as a function of time. These chemically resolved, time-dependent number fractions were then used to determine the surface area concentrations for each particle type, with particles assumed to be spherical. The resulting chemically resolved, time-dependent surface area concentrations were used to calculate the single-particle weighted N_2O_5 uptake and ClNO_2 yield ([Section 4.5](#)).

4.5. Determination of the Single-Particle Weighted $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} . Reactions between particles and gas-phase N_2O_5 ($\gamma_{\text{N}_2\text{O}_5}$), and the yield of ClNO_2 (φ_{ClNO_2}), are dependent on particle chemical composition. To generate time-resolved $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values used for the new single-particle parametrization, previously determined laboratory values of $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} for laboratory proxies were weighted by the time-resolved fractions of aerosol surface area for each ambient particle type ([Section 4.4](#)). These laboratory-based $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} are provided in [Table S1](#), and the selection criteria are discussed below.

4.5.1. Road Salt Particles (Nascent and Aged). There are no reported studies of N_2O_5 uptake onto road salt. Given the observed RH range for the case days ([Section S2.5](#)) and the mechanism of road salt aerosolization (spray generated from vehicular traffic on wet roads),¹⁶ the road salt particles were likely deliquesced at the relative humidities observed for February 17–18 and March 7–8: 54–87% and 58–78%, with minimum temperatures of –12 and 10 °C, respectively ([Section S2.5](#)). For reference, the deliquescence and efflorescence RH points for NaCl particles were previously determined to be 75–80% and 40%, respectively, over a wide range of temperatures (253–298 K).⁷¹ Since the road salt aerosol was primarily NaCl (Cl/Na atomic ratio of 0.87 \pm 0.03; [Figure 2](#)), the corresponding $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values for road salt aerosol were assumed to be that of NaCl.^{20,21,44} However, the aged road salt particles were largely depleted in chloride and enriched in nitrate, as observed by ATOFMS; CCSEM-EDX showed that 64% of these particles contained little chlorine (2–5% atomic percentage), with an overall average Cl/Na atomic ratio of 0.10 \pm 0.01 ([Figure 2](#)). The particulate nitrate suppresses N_2O_5 uptake, and the lack of available chloride reduces the ClNO_2 yield.²¹ Therefore, the $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values for aged road salt were chosen based on studies of mixed nitrate and NaCl particles.^{21,44}

4.5.2. Biomass Burning, Road Dust, and Soot Particles. Recently, Ahern et al.⁴² confirmed the production of ClNO_2

from reaction of N_2O_5 on biomass burning particles, showing that it is a function of the chloride mass fraction, but $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values were not quantified in that study. However, only 8–18%, by number, of the observed biomass burning particles contained chloride (determined by ATOFMS), with measured atomic percentages of less than 2% chlorine (determined by CCSEM-EDX). It is likely that these particles corresponded to both fresh and aged residential wood burning emissions.⁷² Therefore, the uptake of N_2O_5 onto malonic acid (RH = 50–70%)²⁰ was used as a proxy for the primarily organic biomass burning particles. The φ_{ClNO_2} of the biomass burning particles was assumed to be the same as the aged road salt particles, approximately five times lower than that for nearly pure NaCl (nascent road salt), likely representing an upper limit of ClNO_2 production. For the road dust particles, $\gamma_{\text{N}_2\text{O}_5}$ values are based on literature review recommendations,⁴⁵ and since <3% of these particles, by number, contained Cl, φ_{ClNO_2} values were set to zero. For the soot particles, $\gamma_{\text{N}_2\text{O}_5}$ values are based on chamber studies by Saathoff et al.,⁴⁶ and φ_{ClNO_2} values were set to zero, since no chloride was observed by CCSEM-EDX or ATOFMS. Surface area fractions of the single-particle types ([Figure 2](#)) were then used to calculate the single-particle surface area-weighted $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} values.

4.6. One-Dimensional (1-D) Atmospheric Model Description.

A one-dimensional (1-D) numerical model simulated ClNO_2 mole ratios for the February 17–18 and March 7–8 case study periods. The 1-D model incorporates turbulent transport, dry deposition, and multiphase aerosol chemistry in the urban atmospheric boundary layer, building upon the 0-D model described by Wang & Pratt.⁷³ The model consists of 22 vertical atmospheric layers (heights: 0.02, 0.3, 1, 2, 3, 4, 6, 8, 12, 20, 30, 60, 100, 200, 300, 400, 500, 600, 700, 800, 900, and 1,000 m agl). The differential equations describing the temporal evolution of chemical species are solved using IGOR Pro software (<https://www.wavemetrics.com/>).

Temporal evolution of each chemical species is described by

$$\frac{dC}{dt} = \frac{d}{dz} \left(K \frac{\partial C}{\partial z} \right) + P - L - D + \sum_p k_{t,p} \left(C \cdot LWC_p - \frac{C_p}{HRT} \right) \quad (E1)$$

The term $\frac{d}{dz} \left(K \frac{\partial C}{\partial z} \right)$ describes the vertical transport, where z is the layer height (m agl), and K is the turbulent eddy diffusivity ($\text{m}^2 \text{s}^{-1}$), described in [Section S1.6](#). P , L , and D are chemical production, loss, and deposition terms, respectively (all in $\text{molecules cm}^{-3} \text{s}^{-1}$). Chemical reactions are based on those in the 0-D model described by Wang & Pratt⁷³ and from the Master Chemical Mechanism (C1–C4 hydrocarbon precursors from the MCM version 3.2, <http://mcm.leeds.ac.uk/MCM/>), except that bromine chemistry is not included. Time-dependent photolysis frequencies were calculated using the Tropospheric Ultraviolet and Visible Radiation Model (<https://www2.acm.ucar.edu/modeling/tropospheric-ultraviolet-and-visible-tuv-radiation-model>). Trace gas dry deposition was calculated using resistance-analog methods, with the exception of N_2O_5 and NO_2 , for which deposition velocities previously determined for snow-covered ground

were applied.^{74,75} The term $\sum_p k_{t,p} \left(C \cdot LWC_p - \frac{C_p}{HRT} \right)$ describes the mass transport between the gas and aqueous particle phases. C is the concentration of a given species (molecules cm^{-3}) in the gas-phase. C_p is the corresponding particle-phase concentration of liquid water, and $k_{t,p}$ (s^{-1}) is the phase-transfer coefficient for particles.

A number of measurements were used as model inputs: (1) Daily sounding data and ground-based meteorological data (Section S2.5) were used to calculate altitude- and time-varying eddy diffusivities. (2) Ambient measurements of N_2O_5 , O_3 , HCl , and $\text{PM}_{2.5}$ chloride and nitrate were constrained in the corresponding model layer (12 m agl). Measurements of NO (Figure S4) and select volatile organic compounds (VOCs, Table S2) were also used as constraints. (3) Measured aerosol total surface area (Figure S5) was constrained and assumed to be the same throughout all model atmospheric layers. (4) $\text{PM}_{2.5}$ inorganic composition was used as inputs for the thermodynamic model (E-AIM Model IV, <http://www.aim.env.uea.ac.uk/aim/model4/model4a.php>) to calculate aerosol liquid water content. When ambient temperatures were below 265 K, the input temperature was set to 265 K due to the specified limits of the E-AIM model. The traditional parametrization of $\gamma_{\text{N}_2\text{O}_5}$ and φ_{ClNO_2} , based on Bertram and Thornton²¹ and constrained by hourly measurements of $\text{PM}_{2.5}$ Cl^- and NO_3^- , and our new parametrization, constrained to measured single-particle chemical composition (Section 4.5), were employed in the model simulations.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscentsci.9b00994>.

Supplementary tables and figures, and additional methods details, including the $\text{PM}_{2.5}$ sampling information and other measurements used to constrain the 1-D model. Additional ambient observations, analysis of the road salt and brine standards, and meteorology information (PDF)

AUTHOR INFORMATION

Corresponding Author

Kerri A. Pratt — Department of Chemistry and Department of Earth and Environmental Sciences, University of Michigan, Ann Arbor, Michigan, United States;  orcid.org/0000-0003-4707-2290; Phone: (734) 763-2871; Email: prattka@umich.edu

Authors

Stephen M. McNamara — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States

Katherlyn R. Kolesar — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States

Siyuan Wang — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States

Rachel M. Kirpes — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States;  orcid.org/0000-0002-2998-0108

Nathaniel W. May — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States

Matthew J. Gunsch — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States

Ryan D. Cook — Department of Chemistry, University of Michigan, Ann Arbor, Michigan, United States

Jose D. Fuentes — Department of Meteorology and Atmospheric Science, The Pennsylvania State University, University Park, Pennsylvania, United States

Rebecca S. Hornbrook — Atmospheric Chemistry Observations & Modeling Laboratory, National Center for Atmospheric Research, Boulder, Colorado, United States

Eric C. Apel — Atmospheric Chemistry Observations & Modeling Laboratory, National Center for Atmospheric Research, Boulder, Colorado, United States

Swarup China — Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, Washington, United States;  orcid.org/0000-0001-7670-335X

Alexander Laskin — Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, Washington, United States;  orcid.org/0000-0002-7836-8417

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acscentsci.9b00994>

Notes

The authors declare no competing financial interest.

The CIMS dataset is archived through PANGAEA: McNamara, S. M., Pratt, K. A. 10-min averages of ambient ClNO_2 , N_2O_5 , HNO_3 , and O_3 measured at 12 m above ground level in Ann Arbor, Michigan, USA, from February 1 to March 10, 2016. PANGAEA, 2020, <https://doi.org/10.1594/PANGAEA.912746>.

ACKNOWLEDGMENTS

Funding was provided by the University of Michigan (UM), an Alfred P. Sloan Foundation Research Fellowship in Chemistry (FG-2017-9431), and the National Science Foundation (NSF) Atmospheric Chemistry program (AGS-1738588). K.R.K. was partially funded by a Dow Postdoctoral Fellowship in Sustainability from UM. S.M.M. and N.W.M. were partially funded by U.S. Department of Education Graduate Assistance in Areas of National Need fellowships. CCSEM-EDX analyses were performed at the Environmental Molecular Sciences Laboratory, a national scientific user facility located at the Pacific Northwest National Laboratory and sponsored by the Office of Biological and Environmental Research of the U.S. Department of Energy. Additional CCSEM-EDX analyses were conducted at the Michigan Center for Materials Characterization, which is thanked for use of the instruments and staff assistance. The National Center for Atmospheric Research is sponsored by the NSF; any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the NSF. The authors thank D. J. Tanner and L. G. Huey (Georgia Institute of Technology) for providing the NO analyzer, T. B. Ryerson (National Oceanic and Atmospheric Administration) for providing the photolytic NO_2 converter, C. Mattson, A. Barget, A. Kevelin, M. Morales, M. Parks, G. Verwer, and A. Leemon (UM) for experimental assistance, C. R. Thompson (NOAA) for assistance with NO_x measurements and helpful discussions, and A. P. Ault, P. K. Peterson (UM), and H. Osthoff (University of Calgary) for helpful discussions.

■ REFERENCES

(1) Finlayson-Pitts, B. J.; Ezell, M. J.; Pitts, J. N. Formation of Chemically Active Chlorine Compounds by Reactions of Atmospheric NaCl Particles with Gaseous N₂O₅ and ClONO₂. *Nature* **1989**, *337* (6204), 241–244.

(2) Simpson, W. R.; Brown, S. S.; Saiz-Lopez, A.; et al. Tropospheric Halogen Chemistry: Sources, Cycling, and Impacts. *Chem. Rev.* **2015**, *115*, 4035–4062.

(3) Gaston, C. J.; Thornton, J. A. Reacto-Diffusive Length of N₂O₅ in Aqueous Sulfate- and Chloride-Containing Aerosol Particles. *J. Phys. Chem. A* **2016**, *120*, 1039.

(4) Osthoff, H. D.; Roberts, J. M.; Ravishankara, A. R.; et al. High Levels of Nitryl Chloride in the Polluted Subtropical Marine Boundary Layer. *Nat. Geosci.* **2008**, *1* (5), 324–328.

(5) Thornton, J. A.; Kercher, J. P.; Riedel, T. P.; et al. A Large Atomic Chlorine Source Inferred from Mid-Continental Reactive Nitrogen Chemistry. *Nature* **2010**, *464* (7286), 271–274.

(6) Sarwar, G.; Simon, H.; Bhade, P.; et al. Examining the Impact of Heterogeneous Nitryl Chloride Production on Air Quality across the United States. *Atmos. Chem. Phys.* **2012**, *12* (14), 6455–6473.

(7) Sherwen, T.; Schmidt, J. A.; Evans, M. J.; Carpenter, L. J.; Großmann, K.; Eastham, S. D.; Jacob, D. J.; Dix, B.; Koenig, T. K.; Sinreich, R.; Ortega, I.; Volkamer, R.; Saiz-Lopez, A.; Prados-Roman, C.; Mahajan, A. S.; Ordonez, C. Global Impacts of Tropospheric Halogens (Cl, Br, I) on Oxidants and Composition in GEOS-Chem. *Atmos. Chem. Phys.* **2016**, *16*, 12239–12271.

(8) Lilek, J. Roadway deicing in the United States, Factsheet. <https://www.americangeosciences.org/critical-issues/factsheet/roadway-deicing-united-states>.

(9) Ramakrishna, D. M.; Viraraghavan, T. Environmental Impact of Chemical Deicers - A Review. *Water, Air, Soil Pollut.* **2005**, *166* (1–4), 49–63.

(10) Breining, G. We're pouring millions of tons of salt on roads each winter. Here's why that's a problem. <https://ensia.com/features/road-salt/> (accessed September 18, 2019).

(11) Kolesar, K. R.; Mattson, C. N.; Peterson, P. K.; et al. Increases in Wintertime PM_{2.5} Sodium and Chloride Linked to Snowfall and Road Salt Application. *Atmos. Environ.* **2018**, *177*, 195–202.

(12) Denby, B. R.; Ketzel, M.; Ellermann, T.; et al. Road Salt Emissions: A Comparison of Measurements and Modelling Using the NORTRIP Road Dust Emission Model. *Atmos. Environ.* **2016**, *141*, 508–522.

(13) Sansalone, J. J.; Glenn, D. W. Accretion and Partitioning of Heavy Metals Associated with Snow Exposed to Urban Traffic and Winter Storm Maintenance Activities. I. *J. Environ. Eng.* **2002**, *128* (2), 151–166.

(14) Baergen, A. M.; Styler, S. A.; Van Pinxteren, D.; et al. Chemistry of Urban Grime: Inorganic Ion Composition of Grime vs Particles in Leipzig, Germany. *Environ. Sci. Technol.* **2015**, *49* (21), 12688–12696.

(15) Baergen, A. M.; Donaldson, D. J. Seasonality of the Water-Soluble Inorganic Ion Composition and Water Uptake Behavior of Urban Grime. *Environ. Sci. Technol.* **2019**, *53* (10), 5671–5677.

(16) Denby, B. R.; Kupiainen, K. J.; Gustafsson, M. Review of Road Dust Emissions. In *Non-Exhaust Emissions: An Urban Air Quality Problem for Public Health; Impact and Mitigation Measures*; Elsevier Inc., 2018; pp 183–203. DOI: [10.1016/b978-0-12-811770-5.00009-1](https://doi.org/10.1016/b978-0-12-811770-5.00009-1).

(17) Lazarcik, J.; Dibb, J. E. Evidence of Road Salt in New Hampshire's Snowpack Hundreds of Meters from Roadways. *Geosciences* **2017**, *7* (3), 54.

(18) Haskins, J. D.; Jaeglé, L.; Shah, V.; et al. Wintertime Gas-Particle Partitioning and Speciation of Inorganic Chlorine in the Lower Troposphere over the Northeast United States and Coastal Ocean. *J. Geophys. Res.: Atmos.* **2018**, *123*, 1–20.

(19) Mielke, L. H.; Furgeson, A.; Odame-Ankrah, C. A.; Osthoff, H. D.; et al. Ubiquity of ClONO₂ in the Urban Boundary Layer of Calgary, Alberta, Canada. *Can. J. Chem.* **2016**, *94*, 414.

(20) Chang, W. L.; Bhade, P. V.; Brown, S. S.; et al. Heterogeneous Atmospheric Chemistry, Ambient Measurements, and Model Calculations of N₂O₅: A Review. *Aerosol Sci. Technol.* **2011**, *45* (6), 665–695.

(21) Bertram, T. H.; Thornton, J. A. Toward a General Parameterization of N₂O₅ Reactivity on Aqueous Particles: The Competing Effects of Particle Liquid Water, Nitrate and Chloride. *Atmos. Chem. Phys.* **2009**, *9*, 8351–8363.

(22) Riemer, N.; Ault, A. P.; West, M.; Craig, R. L.; Curtis, J. H. Aerosol Mixing State: Measurements, Modeling, and Impacts. *Rev. Geophys.* **2019**, *57*, 187–6.

(23) Gaston, C. J.; Quinn, P. K.; Bates, T. S.; et al. The Impact of Shipping, Agricultural, and Urban Emissions on Single Particle Chemistry Observed Aboard the R/V Atlantis during CalNex. *J. Geophys. Res. Atmos.* **2013**, *118* (10), 5003–5017.

(24) Healy, R. M.; Hellebust, S.; Kourtchev, I.; et al. Source Apportionment of PM_{2.5} in Cork Harbour, Ireland Using a Combination of Single Particle Mass Spectrometry and Quantitative Semi-Continuous Measurements. *Atmos. Chem. Phys.* **2010**, *10* (19), 9593–9613.

(25) Tham, Y. J.; Wang, Z.; Li, Q.; Wang, W.; Wang, X.; Lu, K.; Ma, N.; Yan, C.; Kecorius, S.; Wiedensohler, A.; Zhang, Y.; Wang, T. Heterogeneous N₂O₅ Uptake Coefficient and Production Yield of ClONO₂ in Polluted Northern China: Roles of Aerosol Water Content and Chemical Composition. *Atmos. Chem. Phys.* **2018**, *18*, 13155–13171.

(26) Phillips, G. J.; Thieser, J.; Tang, M. J.; et al. Estimating N₂O₅ Uptake Coefficients Using Ambient Measurements of NO₃, N₂O₅, ClONO₂ and Particle-Phase Nitrate. *Atmos. Chem. Phys.* **2016**, *3* (August), 1–34.

(27) Mielke, L. H.; Stutz, J.; Tsai, C.; et al. Heterogeneous Formation of Nitryl Chloride and Its Role as a Nocturnal NO_x Reservoir Species during CalNex-LA 2010. *J. Geophys. Res. Atmos.* **2013**, *118* (18), 10638–10652.

(28) Riedel, T. P.; Bertram, T. H.; Ryder, O. S.; et al. Direct N₂O₅ Reactivity Measurements at a Polluted Coastal Site. *Atmos. Chem. Phys.* **2012**, *12* (6), 2959–2968.

(29) Bertram, T. H.; Thornton, J. A.; Riedel, T. P.; Middlebrook, A. M.; Bahreini, R.; Bates, T. S.; Quinn, P. K.; Coffman, D. J. Direct Observations of N₂O₅ Reactivity on Ambient Aerosol Particles. *Geophys. Res. Lett.* **2009**, *36* (19), 1–5.

(30) McDuffie, E. E.; Fibiger, D. L.; Dubé, W. P.; et al. ClONO₂ Yields from Aircraft Measurements during the 2015 WINTER Campaign and Critical Evaluation of the Current Parameterization. *J. Geophys. Res.: Atmos.* **2018**, *123*, 1–22.

(31) McDuffie, E. E.; Fibiger, D. L.; Dubé, W. P.; et al. Heterogeneous N₂O₅ Uptake During Winter: Aircraft Measurements During the 2015 WINTER Campaign and Critical Evaluation of Current Parameterizations. *J. Geophys. Res. Atmos.* **2018**, *123* (8), 4345–4372.

(32) Liao, J.; Sihler, H.; Huey, L. G.; et al. A Comparison of Arctic BrO Measurements by Chemical Ionization Mass Spectrometry and Long Path-Differential Optical Absorption Spectroscopy. *J. Geophys. Res.* **2011**, *116*, D00R02.

(33) Mielke, L. H.; Furgeson, A.; Osthoff, H. D. Observation of ClONO₂ in a Mid-Continental Urban Environment. *Environ. Sci. Technol.* **2011**, *45* (20), 8889–8896.

(34) Riedel, T. P.; Wagner, N. L.; Dubé, W. P.; et al. Chlorine Activation within Urban or Power Plant Plumes: Vertically Resolved ClONO₂ and Cl₂ Measurements from a Tall Tower in a Polluted Continental Setting. *J. Geophys. Res. Atmos.* **2013**, *118* (15), 8702–8715.

(35) Wagner, N. L.; Riedel, T. P.; Young, C. J.; et al. N₂O₅ Uptake Coefficients and Nocturnal NO₂ Removal Rates Determined from Ambient Wintertime Measurements. *J. Geophys. Res. Atmos.* **2013**, *118* (16), 9331–9350.

(36) Stutz, J.; Aliche, B.; Ackermann, R.; et al. Vertical Profiles of NO₃, N₂O₅, O₃, and NO_x in the Nocturnal Boundary Layer: 1. Observations during the Texas Air Quality Study 2000. *J. Geophys. Res.* **2004**, *109* (12), 1–15.

(37) Tham, Y. J.; Wang, Z.; Li, Q.; et al. Significant Concentrations of Nitryl Chloride Sustained in the Morning: Investigations of the Causes and Impacts on Ozone Production in a Polluted Region of Northern China. *Atmos. Chem. Phys.* **2016**, *16*, 14959–14977.

(38) Ault, A. P.; Peters, T. M.; Sawvel, E. J.; et al. Single-Particle SEM-EDX Analysis of Iron-Containing Coarse Particulate Matter in an Urban Environment: Sources and Distribution of Iron within Cleveland, Ohio. *Environ. Sci. Technol.* **2012**, *46* (8), 4331–4339.

(39) Pratt, K. A.; Mayer, J. E.; Holecek, J. C.; et al. Development and Characterization of an Aircraft Aerosol Time-of-Flight Mass Spectrometer. *Anal. Chem.* **2009**, *81* (5), 1792–1800.

(40) Markovic, M. Z.; VandenBoer, T. C.; Murphy, J. G. Characterization and Optimization of an Online System for the Simultaneous Measurement of Atmospheric Water-Soluble Constituents in the Gas and Particle Phases. *J. Environ. Monit.* **2012**, *14* (7), 1872.

(41) Staudt, S.; Gord, J. R.; Karimova, N. V.; et al. Sulfate and Carboxylate Suppress the Formation of ClNO_2 at Atmospheric Interfaces. *ACS Earth Sp. Chem.* **2019**, *3* (9), 1987–1997.

(42) Ahern, A. T.; Goldberger, L.; Jahl, L.; et al. Production of N_2O_5 and ClNO_2 through Nocturnal Processing of Biomass-Burning Aerosol. *Environ. Sci. Technol.* **2018**, *52* (2), 550–559.

(43) Pratt, K. A.; Murphy, S. M.; Subramanian, R.; et al. Flight-Based Chemical Characterization of Biomass Burning Aerosols within Two Prescribed Burn Smoke Plumes. *Atmos. Chem. Phys.* **2011**, *11* (24), 12549–12565.

(44) Behnke, W.; George, C.; Scheer, V.; et al. Production and Decay of ClNO_2 from the Reaction of Gaseous N_2O_5 with NaCl Solution: Bulk and Aerosol Experiments. *J. Geophys. Res.* **1997**, *102* (D3), 3795–3804.

(45) Crowley, J. N.; Ammann, M.; Cox, R. A.; et al. Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry: Volume v -Heterogeneous Reactions on Solid Substrates. *Atmos. Chem. Phys.* **2010**, *10* (18), 9059–9223.

(46) Saathoff, H.; Naumann, K.-H.; Riemer, N.; et al. The Loss of NO_2 , HNO_3 , $\text{NO}_3/\text{N}_2\text{O}_5$, and $\text{HO}_2/\text{HOONO}_2$ on Soot Aerosol: A Chamber and Modeling Study. *Geophys. Res. Lett.* **2001**, *28* (10), 1957–1960.

(47) Dugan, H. A.; Bartlett, S. L.; Burke, S. M.; et al. Salting Our Freshwater Lakes. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114* (17), 4453–4458.

(48) Gaston, C. J.; Thornton, J. A.; Ng, N. L. Reactive Uptake of N_2O_5 to Internally Mixed Inorganic and Organic Particles: The Role of Organic Carbon Oxidation State and Inferred Organic Phase Separations. *Atmos. Chem. Phys.* **2014**, *14* (11), 5693–5707.

(49) Anttila, T.; Kiendler-Scharr, A.; Tillmann, R.; et al. On the Reactive Uptake of Gaseous Compounds by Organic-Coated Aqueous Aerosols: Theoretical Analysis and Application to the Heterogeneous Hydrolysis of N_2O_5 . *J. Phys. Chem. A* **2006**, *110* (35), 10435–10443.

(50) Ryder, O. S.; Campbell, N. R.; Morris, H.; et al. Role of Organic Coatings in Regulating N_2O_5 Reactive Uptake to Sea Spray Aerosol. *J. Phys. Chem. A* **2015**, *119* (48), 11683–11692.

(51) Ryder, O. S.; Ault, A. P.; Cahill, J. F.; et al. On the Role of Particle Inorganic Mixing State in the Reactive Uptake of N_2O_5 to Ambient Aerosol Particles. *Environ. Sci. Technol.* **2014**, *48* (3), 1618–1627.

(52) Lelieveld, J.; Gromov, S.; Pozzer, A.; et al. Global Tropospheric Hydroxyl Distribution, Budget and Reactivity. *Atmos. Chem. Phys.* **2016**, *16* (19), 12477–12493.

(53) Wang, T.; Tham, Y. J.; Xue, L.; Li, Q.; Zha, Q.; Wang, Z.; Poon, S. C. N.; Dube, W. P.; Blake, D. R.; Louie, P. K. K.; Luk, C. W. Y.; Tsui, W.; Brown, S. S. Observations of Nitryl Chloride and Modeling Its Source and Effect on Ozone in the Planetary Boundary Layer of Southern China. *J. Geophys. Res. Atmos.* **2016**, *121*, 2476–2489.

(54) Sarwar, G.; Simon, H.; Xing, J.; et al. Importance of Tropospheric ClNO_2 Chemistry across the Northern Hemisphere. *Geophys. Res. Lett.* **2014**, *41* (11), 4050–4058.

(55) Stanier, C.; Singh, A.; Adamski, W.; et al. Overview of the LADCO Winter Nitrate Study: Hourly Ammonia, Nitric Acid and PM2.5 Composition at an Urban and Rural Site Pair during PM2.5 Episodes in the US Great Lakes Region. *Atmos. Chem. Phys.* **2012**, *12* (22), 11037–11056.

(56) Wang, X.; Jacob, D. J.; Eastham, S. D.; Sulprizio, M. P.; Zhu, L.; Chen, Q.; Alexander, B.; Sherwen, T.; Evans, M. J.; Lee, B. H.; Haskins, J. D.; Lopez-Hilfiker, F. D.; Thornton, J. A.; Huey, G. L.; Liao, H. The Role of Chlorine in Tropospheric Chemistry. *Atmos. Chem. Phys.* **2019**, *19*, 3981–4003.

(57) McNamara, S. M.; Raso, A. R. W.; Wang, S.; et al. Springtime Nitrogen Oxide-Influenced Chlorine Chemistry in the Coastal Arctic. *Environ. Sci. Technol.* **2019**, *53* (14), 8057–8067.

(58) Song, X. H.; Hopke, P. K.; Fergenson, D. P.; et al. Classification of Single Particles Analyzed by ATOFMS Using an Artificial Neural Network, ART-2A. *Anal. Chem.* **1999**, *71* (4), 860–865.

(59) Kumar, P.; Hopke, P. K.; Raja, S.; et al. Characterization and Heterogeneity of Coarse Particles across an Urban Area. *Atmos. Environ.* **2012**, *46*, 449–459.

(60) Corbin, J. C.; Rehbein, P. J. G.; Evans, G. J.; et al. Combustion Particles as Ice Nuclei in an Urban Environment: Evidence from Single-Particle Mass Spectrometry. *Atmos. Environ.* **2012**, *51*, 286–292.

(61) Sullivan, R. C.; Guazzotti, S. A.; Sodeman, D. A.; et al. Direct Observations of the Atmospheric Processing of Asian Mineral Dust. *Atmos. Chem. Phys.* **2007**, *7* (5), 1213–1236.

(62) Pósfai, M.; Simonics, R.; Li, J.; Hobbs, P. V.; Buseck, P. R.; Individual Aerosol Particles from Biomass Burning in Southern Africa: 1. Compositions and Size Distributions of Carbonaceous Particles. *J. Geophys. Res. Atmos.* **2003**, *108* (D13). DOI: 10.1029/2002JD002291.

(63) Moffet, R. C.; Henn, T. R.; Tivanski, A. V.; et al. Microscopic Characterization of Carbonaceous Aerosol Particle Aging in the Outflow from Mexico City. *Atmos. Chem. Phys.* **2010**, *10* (3), 961–976.

(64) Dall’Osto, M.; Beddows, D. C. S.; Gietl, J. K.; et al. Characteristics of Tyre Dust in Polluted Air: Studies by Single Particle Mass Spectrometry (ATOFMS). *Atmos. Environ.* **2014**, *94*, 224–230.

(65) Németh, Z.; Pósfai, M.; Nyiro-Kósa, I.; et al. Images and Properties of Individual Nucleated Particles. *Atmos. Environ.* **2015**, *123*, 166–170.

(66) Toner, S. M.; Sodeman, D. A.; Prather, K. A. Single Particle Characterization of Ultrafine and Accumulation Mode Particles from Heavy Duty Diesel Vehicles Using Aerosol Time-of-Flight Mass Spectrometry. *Environ. Sci. Technol.* **2006**, *40* (12), 3912–3921.

(67) Toner, S. M.; Shields, L. G.; Sodeman, D. A.; et al. Using Mass Spectral Source Signatures to Apportion Exhaust Particles from Gasoline and Diesel Powered Vehicles in a Freeway Study Using UF-ATOFMS. *Atmos. Environ.* **2008**, *42* (3), 568–581.

(68) Khlystov, A.; Stanier, C.; Pandis, S. N. An Algorithm for Combining Electrical Mobility and Aerodynamic Size Distributions Data When Measuring Ambient Aerosol Special Issue of Aerosol Science and Technology on Findings from the Fine Particulate Matter Supersites Program. *Aerosol Sci. Technol.* **2004**, *38* (sup1), 229–238.

(69) Wagner, J.; Leith, D. Passive Aerosol Sampler. Part I: Principle of Operation. *Aerosol Sci. Technol.* **2001**, *34* (2), 186–192.

(70) Reinard, M. S.; Adou, K.; Martini, J. M.; et al. Source Characterization and Identification by Real-Time Single Particle Mass Spectrometry. *Atmos. Environ.* **2007**, *41* (40), 9397–9409.

(71) Cziczo, D. J.; Abbatt, J. P. D. Infrared Observations of the Response of NaCl , MgCl_2 , NH_4HSO_4 , and NH_4NO_3 Aerosols to Changes in Relative Humidity from 298 to 238 K. *J. Phys. Chem. A* **2000**, *104* (10), 2038–2047.

(72) Tan, P. V.; Evans, G. J.; Tsai, J.; et al. On-Line Analysis of Urban Particulate Matter Focusing on Elevated Wintertime Aerosol Concentrations. *Environ. Sci. Technol.* **2002**, *36* (16), 3512–3518.

(73) Wang, S.; Pratt, K. A. Molecular Halogens Above the Arctic Snowpack: Emissions, Diurnal Variations, and Recycling Mechanisms. *J. Geophys. Res. Atmos.* **2017**, *122* (21), 11991–12007.

(74) Valdez, M. P.; Bales, R. C.; Stanley, D. A.; et al. Gaseous Deposition to Snow: 1. Experimental Study of SO_2 and NO_2 Deposition. *J. Geophys. Res.* **1987**, *92* (D8), 9779.

(75) Huff, D. M.; Joyce, P. L.; Fochesatto, G. J.; et al. Deposition of Dinitrogen Pentoxide, N_2O_5 , to the Snowpack at High Latitudes. *Atmos. Chem. Phys.* **2011**, *11* (10), 4929–4938.