

Early-life exposure to 17 β -estradiol and 4-nonylphenol impacts the growth hormone/insulin-like growth-factor system and estrogen receptors in Mozambique tilapia, *Oreochromis mossambicus*

Fritzie T. Celino-Brady^a, Cody K. Petro-Sakuma^a, Jason P. Breves^b, Darren T. Lerner^c, Andre P. Seale^{a,*}

^a Department of Human Nutrition, Food and Animal Sciences, University of Hawai'i at Mānoa, 1955 East-West Road, Honolulu, HI 96822, USA

^b Department of Biology, Skidmore College, 815 N. Broadway, Saratoga Springs, NY 12866, USA

^c University of Hawai'i Sea Grant College Program, University of Hawai'i at Mānoa, 2525 Correa Road, Honolulu, HI 96822, USA

ARTICLE INFO

Keywords:

Endocrine disruption
Growth
Insulin-like growth-factor binding proteins
Liver
Mozambique tilapia
Pituitary

ABSTRACT

It is widely recognized that endocrine disrupting chemicals (EDCs) released into the environment through anthropogenic activities can have short-term impacts on physiological and behavioral processes and/or sustained or delayed long-term developmental effects on aquatic organisms. While numerous studies have characterized the effects of EDCs on temperate fishes, less is known on the effects of EDCs on the growth and reproductive physiology of tropical species. To determine the long-term effects of early-life exposure to common estrogenic chemicals, we exposed Mozambique tilapia (*Oreochromis mossambicus*) yolk-sac fry to 17 β -estradiol (E2) and nonylphenol (NP) and subsequently characterized the expression of genes involved in growth and reproduction in adults. Fry were exposed to waterborne E2 (0.1 and 1 μ g/L) and NP (10 and 100 μ g/L) for 21 days. After the exposure period, juveniles were reared for an additional 112 days until males were sampled. Gonadosomatic index was elevated in fish exposed to E2 (0.1 μ g/L) while hepatosomatic index was decreased by exposure to NP (100 μ g/L). Exposure to E2 (0.1 μ g/L) induced hepatic *growth hormone receptor (ghr)* mRNA expression. The high concentration of E2 (1 μ g/L), and both concentrations of NP, increased hepatic *insulin-like growth-factor 1 (igf1)* expression; E2 and NP did not affect hepatic *igf2* and pituitary *growth hormone (gh)* levels. Both E2 (1 μ g/L) and NP (10 μ g/L) induced hepatic *igf binding protein 1b (igfbp1b)* levels while only NP (100 μ g/L) induced hepatic *igfbp2b* levels. By contrast, hepatic *igfbp6b* was reduced in fish exposed to E2 (1 μ g/L). There were no effects of E2 or NP on hepatic *igfbp4* and *igfbp5a* expression. Although the expression of three *vitellogenin* transcripts was not affected, E2 and NP stimulated hepatic *estrogen receptor (era and erb)* mRNA expression. We conclude that tilapia exposed to E2 and NP as yolk-sac fry exhibit subsequent changes in the endocrine systems that control growth and reproduction during later life stages.

1. Introduction

Particular compounds released into the environment through anthropogenic activities impact the endocrine systems of vertebrates, including fishes (Colborn et al., 1996). These compounds, known as endocrine disrupting chemicals (EDCs), include hormones, pharmaceuticals, pesticides, plasticizers, and naturally occurring compounds. Fish are employed as indicator species for environmental pollution in aquatic systems because they are among the first animals exposed to waterborne chemicals. The adverse activities of EDCs in fish are known to include impacts on fertility, sexual maturation, somatic

growth, and circulating hormone levels. Moreover, EDCs can activate stress responses and induce cellular damage, effects that may increase the incidence of disease and mortality (Ankley et al., 2009; Bernanke and Kohler, 2009; Bhandari et al., 2015; Breves et al., 2018; Celino et al., 2009; Jones et al., 2000; Lerner et al., 2007a, b).

Many EDCs act as agonists or antagonists of estrogen receptors (Er) (cf. Ankley et al., 2009). Among the most pervasive EDCs in the aquatic environment are 17 β -estradiol (E2) and nonylphenol (NP) (Aris et al., 2014; Giger et al., 1984; Xu et al., 2014). E2 is one of the most common feminizing compounds found in sewage effluent discharged into rivers (Desbrow et al., 1998). Nonylphenol ethoxylates (NPEs) are widely

* Corresponding author.

E-mail addresses: fbrady@hawaii.edu (F.T. Celino-Brady), cps3@hawaii.edu (C.K. Petro-Sakuma), jbrevs@skidmore.edu (J.P. Breves), lerner@hawaii.edu (D.T. Lerner), seale@hawaii.edu (A.P. Seale).

<https://doi.org/10.1016/j.aquatox.2019.105336>

Received 4 July 2019; Received in revised form 28 September 2019; Accepted 14 October 2019

Available online 24 October 2019

0166-445X/ © 2019 Elsevier B.V. All rights reserved.

used as surfactants in industrial processes and products, including cleaners, detergents, and plastics. As in the case of E2, NPEs are also discharged through domestic and industrial wastewater (Mao et al., 2012; Servos et al., 2003). NPEs are degraded into NP, which persists in the environment (Ahel and Giger, 1993). Free NP is presumed to be widely distributed in surface waters (Ekelund et al., 1990, 1993) with concentrations ranging from approximately 30 to 30,000 ng/L in Guangzhou riverine waters in China, the Seine estuary in France, and the European river basin in Spain (Brix et al., 2010; Cailleaud et al., 2007; Peng et al., 2008). NP accumulates in various aquatic organisms at concentrations ranging from 0.68 to 160 ng/g tissue weight (Vethaak et al., 2005; Zhou et al., 2019). NP exerts feminizing effects in mice (Hernandez et al., 2006), reduces fecundity and fertility in Japanese medaka (*Oryzias latipes*) (Ishibashi et al., 2006; Kang et al., 2003), reduces semen volume in rainbow trout (*Oncorhynchus mykiss*) (Lahnsteiner et al., 2005), and diminishes plasma testosterone in male carp (*Cyprinus carpio*) (Amaninejad et al., 2018). Moreover, the presence of NP and NPEs in the environment was linked to a low male:female sex ratio in wild Nile tilapia (*Oreochromis niloticus*) (Chen et al., 2014). Most studies reporting on the effects of E2 and NP on growth and reproduction in fishes have been conducted with temperate species (Harries et al., 2000; Filby et al., 2006; Goetz et al., 2009; Duffy et al., 2014; Breves et al., 2018).

Given its importance to worldwide aquaculture (FAO, 2005), the Mozambique tilapia (*Oreochromis mossambicus*) is one of the most thoroughly studied tropical fishes with respect to how environmental conditions impact growth and reproductive endocrinology (Davis et al., 2009a, b; Gaigher and Krause, 1983; Kiilerich et al., 2011; Moorman et al., 2016; Kajimura et al., 2005). Tilapia are widely distributed in tropical areas where they are cultured for human consumption. They inhabit regions where agricultural, municipal, and industrial waters are discharged and are therefore exposed to persistent environmental EDCs based on the contaminants detected in their tissues (Authman, 2008; Babu and Ozbay, 2013; Hemmatinezhad et al., 2017; Osman, 2012).

The endocrine system of fishes mediates the effects of environmental stimuli, including contaminants, on growth and reproduction. The growth hormone (Gh)/insulin-like growth-factor (Igf) system plays a major role in regulating the growth and development of vertebrates, including teleosts (Duan et al., 2010; Reindl and Sheridan, 2012). Upon binding to the Gh receptor (Ghr), Gh stimulates the release of Igf1 which has growth-promoting actions in target tissues (Butler and LeRoith, 2001; Duan, 1998; Fan et al., 2009; Le Roith et al., 2001; Le Roith and Roberts, 2003). Igfs interact with a family of binding proteins, known as Igf binding proteins (Igfbps), which influence their availability and activities (Duan and Xu, 2005; Duan et al., 2010; Rajaram et al., 1997) and teleost fishes possess an expanded suite of Igfbps (Allard and Duan, 2018). Steroid hormone receptors mediate target-tissue responsiveness to the actions of steroid hormones, in addition to compounds that mimic hormone actions (Park et al., 2007; Gross and Yee, 2002). The production of vitellogenin (Vtg), a precursor of egg yolk protein produced by the liver of female oviparous animals (Denslow, 1999; Hiramatsu et al., 2005), is stimulated by activation of *ers* and *Ers* (Bowman et al., 2002; Flouriot et al., 1996; Jalabert, 2005; Nelson and Habibi, 2010). These same *Ers* are the pathway in which estrogenic EDCs interfere with normal estrogen signaling (Shanle and Xu, 2011). Hence, Vtg/vtg and *Ers/ers* are often used as indicators of estrogenic EDC exposure (Jones et al., 2000; Matozzo et al., 2008; Leet et al., 2011; Park et al., 2007). Plasma Vtg has been detected in male white sucker (*Catostomus commersoni*) and rainbow trout inhabiting waters contaminated by sewage effluent (Purdum et al., 1994; Vajda et al., 2008). Estrogenic EDCs such as E2, 17 α -ethinylestradiol (EE2), diethylstilbestrol, and NP induce vtg expression and plasma Vtg levels in male fish (Davis et al., 2007, 2009b; Folmar et al., 2000; Hemmer et al., 2001). In male tilapia, injection of E2 induces Vtg production while concurrently suppressing the Gh/Igf system (Davis et al., 2007, 2008). Moreover, in juvenile Atlantic salmon, vtg transcripts were induced,

while hepatic *er*, *ghr*, *igfs*, and several *igfbps* were reduced following waterborne exposure to EE2 and NP (Breves et al., 2018). Filby et al. (2006) previously showed that male fathead minnows (*Pimephales promelas*) exposed to E2 exhibited reduced hepatic *igf1* expression levels. Here, we examined the long-term effects of waterborne exposure to NP and E2 (as positive control) in Mozambique tilapia. We reared tilapia fry in water containing E2 (0.1 and 1 μ g/L) and NP (10 and 100 μ g/L) for 21 days, and then measured the expression of pituitary *gh*, and hepatic *ghr*, *igfs*, *igfbps*, *vtgs*, and *ers* after an additional 112 days of growth and development.

2. Materials and methods

2.1. Animals

Mozambique tilapia yolk-sac fry were obtained from broodstock tanks maintained in dechlorinated city water at the Hawai'i Institute of Marine Biology (Kāne'ohe, HI). Fry were initially reared in 7-L conical tanks supplied with filtered dechlorinated city water for 2–3 days. Fry were then distributed to 5-L aerated flow-through tanks (33 fry/tank) supplied by 19-L header tanks and reared for an additional 5–10 days until yolk absorption was ~90% complete. Header and fish-holding tanks were lined with modified polytetrafluoroethylene (MPTFE) (Welch Fluorocarbon, Inc., Dover, NH). Two replicate tanks were used for each treatment and subsequent rearing. Water temperature was maintained at ~26–28 °C under a 12L:12D photoperiod. After yolk absorption, fry were fed crushed trout chow pellets (Skrretting, Tooele, UT) twice daily for the remainder of the experiment. Fecal material and uneaten food were siphoned out from the tanks before 60–70 % of the water volume was changed daily. All housing and experimental procedures were conducted in accordance with the principles and procedures approved by the Institutional Animal Care and Use Committee of the University of Hawai'i.

2.2. Chemical exposures and rearing

Fry (0.029 $\pm$ 0.001 g body weight) were exposed to E2 (0.1 and 1 μ g/L) and NP (10 and 100 μ g/L) in fresh water for 21 days. The duration of the exposures was chosen following previous studies aimed at identifying responses to chronic EDC exposures (Woltering, 1984; Lerner et al., 2007a, b). The range of nominal concentrations of E2 used in this study were above those typically found in the environment inasmuch as they were included as a positive control. The concentrations of E2 and NP were based on those employed in previous studies where Atlantic salmon yolk-sac larvae, fry, juvenile, and smolts were subjected to the aqueous exposure of these chemicals; the concentrations of the chemicals in the water have been previously validated (Breves et al., 2018; Duffy et al., 2014; Lerner et al., 2007a, b). E2 and NP were purchased from Sigma-Aldrich (St. Louis, MO) and Acros Organics (Fair Lawn, NJ). All chemicals were solubilized in ethanol and then added to fresh water at a final concentration of 0.0001% ethanol to minimize solvent toxicity. Control treatments received solvent only. Header tanks were covered and refilled every two days with filtered and dechlorinated city water with or without the experimental chemicals. The flow-rate from the header tanks to the rearing tanks averaged 0.2 L/h. The fry were maintained in 5-L MPTFE-lined tanks until the end of the exposures (see Fig. 1 for the experimental setup) at which time the juveniles were transferred to 19-L aerated flow-through MPTFE-lined containers and reared for an additional 112 days until sampling. MPTFE was used to prevent the leaching of chemicals from the plastic containers used as tanks and header buckets. This approach has been previously used in a similar study employing coho salmon (Harding et al., 2016). Bodyweight (BW) and total length (TL) were measured every two weeks throughout the duration of the experiment.

At the end of the experiment, male tilapia were netted and anesthetized with 2-phenoxyethanol (0.3 mL/L; Sigma-Aldrich) and BW




Fig. 1. Schematic illustration of header and experimental tanks. Nineteen-liter capacity header tanks supplied water either containing solvent only (control), 0.1 µg/L or 1 µg/L 17β -estradiol (E2), or 10 µg/L or 100 µg/L nonylphenol (NP) to fish tanks. Both header and fish tanks were lined with modified polytetrafluoroethylene. Black squares = aeration, white rectangles = pipes.

and TL were measured. Anesthetized fish were euthanized by rapid decapitation. Testes and liver were removed and weighed for calculation of gonadosomatic index (GSI; (gonad weight/BW)*100) and hepatosomatic index (HSI; (liver weight/BW)*100). Condition factor (CF) was calculated as $CF = (BW/TL^3) \times 100$. Liver and pituitary were collected, immediately snap-frozen in liquid nitrogen, and stored at -80°C until RNA extraction.

2.3. Quantitative real-time PCR (qRT-PCR)

Total RNA was extracted from liver and pituitary using TRI Reagent (MRC, Cincinnati, OH) according to the manufacturer's protocols. The concentration and purity of extracted RNA were assessed using a NanoDrop (NanoDrop One, Thermo Scientific). Total RNA (100–500 ng) was reverse-transcribed using a High Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific, Waltham, MA). The mRNA levels of reference and target genes were determined by the relative quantification method using a StepOnePlus real-time PCR system (Thermo Fisher Scientific). The qRT-PCR reaction mix (15 µL) contained Power SYBR Green PCR Master Mix (Thermo Fisher Scientific), 200 nM of forward and reverse primers, and 1 µL cDNA. Dilution of experimental cDNA from liver ranged from 10- to 175-fold. PCR cycling parameters were: 2 min at 50°C , 10 min at 95°C followed by 40 cycles at 95°C for 15 s and 60°C for 1 min. All qRT-PCR primers have been previously described; PCR efficiencies are reported in Table 1. Since hepatic *elongation factor 1a* (*ef1a*), *β-actin*, and *18s* levels varied across treatments, the geometric mean ($\sqrt[3]{x_1 \times x_2 \times x_3}$; where x = quantity of each reference gene) of these reference genes was used to normalize target genes in the liver. *ef1a* levels were used to normalize pituitary *gh* expression after verification that *ef1a* expression did not

vary across treatments. Data are expressed as a fold-change relative to control.

2.4. Statistical analysis

Group comparisons were performed by one-way ANOVA followed by Fisher's protected LSD test. In order to meet assumptions of normality (assessed by Kolmogorov-Smirnov), individual values were log-transformed when necessary prior to statistical analysis. Pearson correlation coefficients were used to describe the relationships between *gh*, *ghr*, *igf1*, and *igf2* mRNA levels. Statistical calculations were performed using Prism 8.0 (GraphPad, La Jolla, CA). Significance for all tests was set at $P < 0.05$.

3. Results

3.1. Effects of E2 and NP on physical characteristics

BW was significantly higher in fish exposed to 10 µg/L NP compared with controls (Fig. 2A) whereas TL and CF were unaffected by E2 and NP (Fig. 2B, C). The high concentration of NP (100 µg/L) reduced HSI relative to controls (Fig. 2D). GSI was elevated in fish exposed to E2 at 0.1 µg/L (Fig. 2E).

3.2. Effects of E2 and NP on *ghr*, *igf1*, *igf2*, and *gh* gene expression

Hepatic *ghr* levels were > 2-fold higher in fish exposed to E2 (0.1 µg/L) compared with controls; E2 at 1 µg/L and NP at both concentrations did not impact *ghr* levels (Fig. 3A). By contrast, hepatic *igf1* was elevated > 2-fold in fish exposed to the high concentration (1 µg/L)

Table 1
List of primers used in qPCR assays.

Gene	Forward primer Sequence (5'-3')	Reverse primer Sequence (5'-3')	R ²	% efficiency	Accession Number	Reference
<i>18s</i>	GCTACCACATCCAAGGAAGGC	TTCGTCACACTCCCGAGT	0.999	89.1	AF497908	Magdeldin et al., 2007
<i>ef1a</i>	AGCAAGTACTACGTGACCATCAATTG	AGTCAGCCTGGGAGGTACCA	0.999	95.1	AB075952	Breves et al., 2010
<i>β-actin</i>	CTCTTCAGCCTTCCTTCCT	ACAGGTCCTTACGGATGTGCG	0.998	96.5	FN673689	Tipsmark et al., 2011
<i>ghr</i>	CACACCTCGATCTGGACATATTACA	CGGTTGGACAATGTCAATTAACAA	0.997	96.8	EF452496	Pierce et al., 2007
<i>igf1</i>	CTGCTTCCAAAGCTGTGAGCT	GATCGAGAAATCTTGGGAGTCTTG	0.999	92.3	AF033796	Kajimura et al., 2004
<i>igf2</i>	GCITTTATTTTCAGTAGGCCAACCA	CACAGCTACAGAAAAGACACTCCTCTA	0.997	90.1	AH006117	Davis et al., 2008
<i>igfbp1b</i>	CCITTCCTTTGATCACCAAG	GTGTGACATGGACCTGTGTG	0.997	90.8	XM_003438121	Breves et al., 2014
<i>igfbp2b</i>	CCGACTTCCTTTACAGCAG	TCAGTCCCATGCACCTCATA	0.998	92.2	XM_0054504847	Breves et al., 2014
<i>igfbp4</i>	ATCCCATACCCAACCTGTGA	TGATCCACACACAGCAGATTT	0.999	85.6	XM_003454633	Breves et al., 2014
<i>igfbp5a</i>	AACTGGACGGGATCAITTCAG	GCACTGTTTTCGTTTGAAGA	0.999	106.7	XM_003443250.2	Breves et al., 2014
<i>igfbp6b</i>	TCCTACCTGCAGAGGAAAGC	CGCAGCTCAGAGTGTAGACG	0.975	96.5	XM_003441337	Breves et al., 2014
<i>gh</i>	TTACATCATCAGCCGATCG	AGATCGACAGCAGCTTCAGGA	0.999	94.3	AF033806	Magdeldin et al., 2007
<i>vtga</i>	GAATGTGAATGGGCTGGAAATAC	TTTGTGTTGATCTGGATCTCAGCTT	0.999	90.3	EF408235	Davis et al., 2007
<i>vtgb</i>	AAGTTGCAGACTGGATGAAAGGA	GCGGTACTCGTCTCCGACAT	1.000	97.7	EF408236	Davis et al., 2007
<i>vtgc</i>	GGACCTTGCAGAACCCAAAG	CATCGTTTCTTGGCAGTTCCA	0.998	94.9	EF408237	Davis et al., 2007
<i>era</i>	GGCTCAGCAGCAGTCAAGAA	TGCCTTGAGGTCTGAACTG	0.990	87.0	AM284390	Park et al., 2007
<i>erβ</i>	ACCTTCCGGCAGCAGTACAC	TCCAACATCTCCAGCAACAG	0.994	108.0	AM 284391	Park et al., 2007




Fig. 2. Body weight (BW) (A), total length (TL) (B), condition factor (CF) (C), hepatosomatic index (HSI) (D), and gonadosomatic index (GSI) (E) of Mozambique tilapia adults 112 days after 21-day exposure as fry to water containing 0 (control), 0.1 µg/L or 1 µg/L 17β-estradiol (E2) or 10 µg/L or 100 µg/L nonylphenol (NP). Values are means ± SEM ($n = 5$ –23; BW, TL, and CF; $n = 5$ –12, HSI and GSI). Asterisk indicates significant difference between treatment and control group (One-way ANOVA; Fisher's protected LSD; $P < 0.05$).

of E2 and > 1-fold to 2-fold to both concentrations of NP (Fig. 3B). Hepatic *igf2* levels were highly variable and were not impacted by E2 and NP (Fig. 3C). Pituitary *gh* levels in the E2 and NP treatments were not different from control levels (Fig. 3D). Hepatic *ghr* was significantly correlated with *igf1* ($r^2 = 0.33$) and *igf2* ($r^2 = 0.22$).

3.3. Effects of E2 and NP on *igfbp* gene expression

Exposures to E2 and NP at 1 and 10 µg/L, respectively, induced hepatic *igfbp1b* levels by > 2-fold from controls (Fig. 4A). Fish exposed to NP (100 µg/L) exhibited elevated *igfbp2b* levels compared with controls (Fig. 4B). *igfbp4* and *igfbp5a* levels were unaffected by E2 and NP exposures (Fig. 4C, D) whereas *igfbp6b* was the only *igfbp* transcript suppressed by E2 (Fig. 4E).

3.4. Effects of E2 and NP on *vtg* and *er* gene expression

Although all three *vtg* transcripts had a tendency to be elevated in fish exposed to E2, no significant effects of E2 or NP were detected (Fig. 5A–C). On the other hand, *era* levels were increased following exposures to all tested concentrations of E2 and NP (Fig. 5D); *erβ* levels were stimulated by E2 and NP at 0.1 and 100 µg/L, respectively (Fig. 5E).

4. Discussion

The objective of this study was to determine whether Mozambique tilapia exposed to estrogenic chemicals as fry exhibit long-term responses that impact physiological systems underlying growth and reproduction as adults. Mature tilapia exhibit concentration-dependent responses to estrogenic compounds commonly found in sewage effluents, such as E2, o,p'-DDE (dichlorodiphenyl dichloroethene), heptachlor, and NP (Davis et al., 2007, 2008, 2009b). Little is known,

however, on whether exposure to estrogenic compounds during early-life stages may impart long-term physiological effects on adult tilapia. To our knowledge, this is the first study demonstrating that early-life exposure to estrogenic EDCs affects the Gh/Igf system and *er* expression in adult tilapia. At very low levels, E2 and NP still elicited physiological responses in exposed individuals.

In previous studies, BW was significantly reduced in fry and adult male fish after exposure to estrogenic compounds such as EE2 and NP (Meredith et al., 1999; Breves et al., 2018). In the current study, however, the BW of adults exposed to NP (10 µg/L) as fry was actually greater than controls. This may be attributed to a capacity for the somatotropic axis to compensate for poor growth during early life-stages (Bertram et al., 1993; Chambers et al., 1988; Gagliano and McCormick, 2007; Segers et al., 2012), in this case when the EDCs were present. Compensatory growth following periods of suppressed growth, such as food restriction, occurs in several teleosts, including striped bass (*Morone saxatilis*), Atlantic halibut (*Hippoglossus hippoglossus*), channel catfish (*Ictalurus punctatus*), rainbow trout, and hybrid tilapia (*O. mosambicus* x *O. niloticus*) (Gaylord and Gatlin, 2000; Heide et al., 2006; Montserrat et al., 2007; Picha et al., 2008). In the current study, we observed that HSI was lower in fish exposed to NP (100 µg/L) as fry. Since the liver is a major site for metabolism, detoxification, and vitellogenesis, there are a variety of factors that likely contributed to this response (Roberts, 2012; Asem-Hiablíe et al., 2013). HSI is naturally elevated during reproductive periods as a result of increased protein synthesis and Vtg production (Jia et al., 2019). In salmon fry and smolts, E2, EE2, and NP elevated HSI (Lerner et al., 2012; Duffy et al., 2014). While elevations in HSI have been correlated with the occurrence of xenobiotics in polluted zones (Karels et al., 1998; Billiard and Khan, 2003), in some cases, such as in Mozambique tilapia, African catfish (*Clarias gariepinus*), spotted pim (*Pimelodus maculatus*), and Japanese medaka, lower HSIs were observed in fish exposed to sewage effluents and agricultural runoffs (Ma et al., 2005; Asem-Hiablíe et al.,




Fig. 3. Hepatic *ghr* (A), *igf1* (B), *igf2* (C), and pituitary *gh* (D) mRNA levels in Mozambique tilapia adults 112 days after 21-day exposure as fry to water containing 0 (control), 0.1 µg/L or 1 µg/L 17β-estradiol (E2) or 10 µg/L or 100 µg/L nonylphenol (NP). mRNA levels are presented as fold-change relative to the control group. Values are means ± SEM (n = 5–12). Asterisk indicates significant difference between treatment and control group (One-way ANOVA; Fisher's protected LSD; $P < 0.05$).

2013; Sadekarpawar and Parikh, 2013; Araújo et al., 2018). Further investigation is needed to determine the mechanisms that underlie reductions in HSI following EDCs exposures.

GSI has been extensively used as an indicator of sexual maturation as well as a biomarker for exposure of aquatic organisms to estrogenic EDCs. Several laboratory studies have reported that exposure to estrogenic chemicals inhibits testicular development (Gimeno et al., 1997; Komen et al., 1989; Christiansen et al., 1998). Field studies have also documented a correlation between estrogenic compounds and lower GSI in exposed male fish (Andersson et al., 1988; Harries et al., 1997; Kukkonen et al., 1999; Hassanin et al., 2002). In the current study, we observed that fry exposed to E2 (0.1 µg/L) exhibited elevated GSI as juvenile males. Jobling et al. (1996) found that the inhibitory effects of estrogenic compounds on sexually maturing rainbow trout were not evident in mature or regressing fish. In juvenile male salmon, a relatively low concentration of E2 (2 µg/L) increased GSI (Lerner et al., 2007a). Furthermore, E2 also plays a role in male gonad development. Although high concentrations of E2 could inhibit testicular development in some fishes such as Japanese eel (*Anguilla japonica*) and threespot wrasse (*Halichoeres trimaculatus*), low concentrations of E2 were found to induce spermatogonial stem cell renewal and spermatogonial proliferation, suggesting a modulatory role of E2 in normal testicular development (Miura et al., 1999, 2003; Kobayashi et al., 2011). These previous studies, therefore, may explain the increase in GSI in fish exposed to the low concentration of E2. The other concentrations of E2 and NP employed in this study were seemingly insufficient to affect GSI,

especially given that fish were exposed to these concentrations as fry. Taken together with previous findings, our results suggest that the impact of estrogenic chemicals on GSI is dependent on both concentration and timing of exposure.

Our results indicate that the somatotrophic axis of adult tilapia was impacted by early-life exposure to the tested EDCs. Both stimulatory and inhibitory effects of estrogenic chemicals on *gh* transcript levels and Gh synthesis were previously reported in teleosts (Elango et al., 2006; Holloway and Leatherland, 1997; Shved et al., 2007, 2008; Zou et al., 1997). In juvenile Atlantic salmon, E2 and NP injection had no effect on pituitary transcript levels of *gh* (Yadete and Male, 2002), a pattern that is consistent with our current observations. Diminished hepatic *ghr* gene expression was associated with reductions in Gh binding capacity, circulating Igf1 levels, and *igf1* expression following exposure to estrogenic compounds in salmonids (Breves et al., 2018; Hanson et al., 2017; Lerner et al., 2012; Norbeck and Sheridan, 2011). By contrast, we observed elevations in hepatic *ghr* and *igf1* gene expression in addition to positive correlations between *ghr* and both *igf1* and *igf2*, following exposure to estrogenic compounds. As discussed above, these patterns may be associated with a compensatory growth response. In this instance, the Gh/Igf system is seemingly 'activated' following the withdrawal of estrogenic chemicals. During restricted feeding, for example, the catabolic state preceding compensatory growth is characterized by depressed levels of hepatic *ghr*, *igf1*, and plasma Igf1 (Gray et al., 1992; Duan et al., 1995; Pierce et al., 2005; Norbeck et al., 2007; Picha et al., 2008). Upon re-feeding, a rapid increase in specific growth rate and




Fig. 4. Hepatic *igfbp1b* (A), *igfbp2b* (B), *igfbp4* (C), *igfbp5a* (D), and *igfbp6b* (E) mRNA levels in Mozambique tilapia adults 112 days after 21-day exposure as fry to water containing 0 (control), 0.1 µg/L or 1 µg/L 17β-estradiol (E2) or 10 µg/L or 100 µg/L nonylphenol (NP). mRNA levels are presented as fold-change relative to the control group. Values are means ± SEM ($n = 5-12$). Asterisk indicates significant difference between treatment and control group (One-way ANOVA; Fisher's protected LSD; $P < 0.05$).

hepatic *ghr*, *igf1*, and *igf2* expression occurs (Picha et al., 2008). Although there was no clear inhibition of pituitary *gh* expression, we found a negative correlation between pituitary *gh* and hepatic *igf1*, a possible indication of feedback regulation of Gh by Igf1 (Reinecke, 2010). No correlation was observed, however, between hepatic *igf2* and pituitary *gh*. Moreover, the differing responses by hepatic *igf1* and *igf2* to estrogenic EDCs observed in this study were similar to patterns in male Mozambique tilapia injected with E2 (Davis et al., 2008), in which hepatic *igf2* was not affected. In mammals, Igf2 is mainly associated with fetal growth and development (Constancia et al., 2002; Daughaday and Rotwein, 1989). In teleosts, however, some studies suggest that Igf2 is also an important factor in adult growth (Pierce et al., 2011; Reindl and Sheridan, 2012). The varying responses of *igf2* to estrogenic compound may be due to differences among species and tissue sensitivity.

Igfbps are key modulators of Igf activity (Duan and Xu, 2005). Only a few studies in mammals and fishes have described how steroid hormones regulate Igfbps (Duan et al., 2010; Garcia de la Serrana et al., 2017; Rajaram et al., 1997; Reindl and Sheridan, 2012), and fewer yet have examined the long-term effects of EDC exposure on *igfbps* during early developmental stages (Breves et al., 2018). In teleosts, it is generally accepted that the liver is a major site for Igfbp synthesis and secretion (Shimizu and Dickhoff, 2017; Zhou et al., 2008). In Mozambique tilapia, *igfbp1b*, -2b, -5a, -4, and -6b are expressed in the liver (Breves et al., 2014). In vertebrates, Igfbp1 plays a highly conserved role as a negative regulator of somatic growth by restricting Igf1 from binding to its receptor (Kajimura et al., 2005; Kamei et al., 2008). Knockdown of Igfbp1 in zebrafish, for instance, alleviates hypoxia-induced retardation of growth, while its overexpression causes growth and developmental retardation (Kajimura et al., 2005). In chinook (*O. tshawytscha*) and Atlantic salmon, Igfbp1b paralogs are important

modulators of Igf signaling in response to nutrient availability (Hevrøy et al., 2011; Shimizu et al., 2005, 2006, 2009). Therefore, the increase in *igfbp1b* transcript levels after E2 and NP exposures may provide a mechanism for EDCs to impact growth. In other words, by enhancing *igfbp1b*, a negative regulator of growth, estrogenic EDCs inhibit somatic growth. Alternatively, the increase in *igf1* mRNA levels (and plasma Igf1) and other factors within the Gh/Igf system following EDC exposures may counterbalance the *igfbp1b* response. Previous studies in mature male Mozambique tilapia (Riley et al., 2004) and striped bass (*Morone saxatilis*) (Fukazawa et al., 1995) reported that E2 stimulated the release of putative Igfbp1s from hepatocytes. In Atlantic salmon fry and smolts, however, estrogenic compounds inhibited *igfbp1b* (Breves et al., 2018). Diets supplemented with E2 also inhibited hepatic *igfbp1b1* expression (along with hepatic *igf1* and *igf2*) in rainbow trout (Cleveland and Weber, 2016). These findings support the notion that responses to estrogenic compounds are both species and life stage-dependent.

In salmonids, Igfbp2 paralogs are major carriers of plasma Igf1 (Shimizu and Dickhoff, 2017). In mammals and teleosts, varying patterns exist on the regulation of *igfbp2* gene expression. For example, overexpression of Igfbp2 in mouse embryos reduces growth rates, which was proposed to be related to a reduction in Igf availability (Hoeftlich et al., 1999). On the other hand, in salmonids, circulating Igfbp2 increase in response to Gh (García de la Serrana and Macqueen, 2018; Shimizu et al., 1999, 2003). In tilapia, *igfbp2b* expression increased with an increased plasma Igf1 induced by Gh injection (Breves et al., 2014). Moreover, *igfbp2b* expression was increased by treatment of NP but decreased by treatment of E2 in salmon smolts (Breves et al., 2018). By contrast, in rats, hepatic *igfbp2* expression is induced by E2 (Hoeftlich et al., 2014; Ricciarelli et al., 1991). In rainbow trout, *igfbp2*




Fig. 5. Hepatic *vtga* (A), *vtgb* (B), *vtgc* (C), *era* (D), and *erβ* (E) mRNA levels in Mozambique tilapia adults 112 days after 21-day exposure as fry to water containing 0 (control), 0.1 µg/L or 1 µg/L 17β-estradiol (E2) or 10 µg/L or 100 µg/L nonylphenol (NP). mRNA levels are presented as fold-change relative to the control group. Values are means ± SEM ($n = 5-12$). Asterisk indicates significant difference between treatment and control group (One-way ANOVA; Fisher's protected LSD; $P < 0.05$).

expression in ovarian follicles is also increased by E2 treatment (Kamangar et al., 2006). Hence, the observed increase in *igfbp2b* levels following exposure to NP, and its tendency to increase after E2 exposure may be either associated with the increase in *igf1* levels or modulated by E2 and E2 analogues. Further studies, however, are needed to assess whether there are direct actions of *Igf1* and E2 on hepatic *igfbp2b* expression. In Atlantic salmon, *igfbp4*, *-5a*, and *-6b* regulate the binding of *Igfs* to its receptor in the tissues where they are produced (Breves et al., 2017; Cleveland and Weber, 2015; Macqueen et al., 2013). Unlike patterns observed in Atlantic salmon (Breves et al., 2018), we found no significant effect of estrogenic EDCs on *igfbp4* and *-5a*. *igfbp6*, on the other hand, was significantly decreased following E2 exposure. In teleosts, *Igfbp6* inhibits *Igf*-signaling that supports growth and development (Wang et al., 2009). While the decrease in *igfbp6b* may be a residual effect from the earlier exposure to E2, additional work should address whether E2 and NP act directly on the liver to regulate *igfbp6*.

To assess the effects of the tested EDCs on estrogenic biomarkers in males, we measured hepatic *vtg* and *er* transcripts. While a trend in all *vtg* transcripts was observed, no significant effects of E2 or NP were detected. Alternatively, both *era* and *erβ* were stimulated by E2 and NP. In previous studies, *vtg* and *er* (α and β) were induced in liver and testis after injection of E2 and other estrogenic compounds in mature male Mozambique tilapia (Davis et al., 2009b). A concurrent increase in *vtg* and *era* expression was also observed in Atlantic salmon embryos, yolk-sac fry, feeding fry, and smolts in response to E2, EE2, and NP (Duffy et al., 2014; Breves et al., 2018). The lack of effects on *vtg* in the current study may be linked to the time of EDC exposures and life stage. Indeed, it is noteworthy that even after 112 days since E2 and NP exposures, both *ers* were still elevated. This elevation suggests that males may be more sensitive to E2 and similar chemicals after a previous EDC

exposure. Increased sensitivity to estrogenic compounds through enhanced expression of *ers* may render males more susceptible to further detrimental effects on their reproductive development.

5. Conclusion

Our current findings indicate that early aqueous exposure to estrogenic EDCs exerts long lasting effects on the somatotrophic axis of tilapia, a central mediator of adaptive patterns of growth and development throughout the life cycle in vertebrates. Thus, an improved understanding of how EDCs impact the endocrine systems controlling growth and reproduction attest to the importance of fish as sentinels for assessing the health of the aquatic ecosystem. Moreover, studies such as this one shall be instrumental in optimizing culture practices for tropical fishes in environments where EDCs are pervasive. Nonetheless, future work that include female tilapia is needed to characterize the long-term effects of estrogenic EDCs in both sexes. Moreover, additional analyses, such as histological examination of testicular tissue would further shed light on the long-term effects of early exposure to estrogenic EDCs on testicular development. Future investigations should also seek to determine the effects of these estrogenic chemicals on the indices of reproductive capacity such as spawning efficiency, fertilization success, and viability of embryos.

Contributions

F.T.C.B. conducted experiments, conceived and designed experiments, collected and analyzed data, and wrote the manuscript. C.K.P.S. collected data and conducted experiments. J.P.B. designed experiments and revised the manuscript. D.T.L. coordinated the study and revised the manuscript. A.P.S. conceived and designed experiments,

coordinated the study, and wrote the manuscript. All authors approved the final article.

Funding

This work was supported in part by grants from the National Oceanic and Atmospheric Administration (NA18OAR4170347 to D.T.L. and A.P.S. and NA14OAR4170071, which is sponsored by the University of Hawai'i Sea Grant College Program project R/SB-18 to A.P.S.), National Science Foundation (IOS-1119693), National Institute of Food and Agriculture Hatch (HAW02051-H) and National Institutes of Diabetes and Digestive and Kidney Diseases (1R21DK111775-01) to A.P.S. The views expressed herein are those of the authors and do not necessarily reflect the views of the aforementioned granting agencies. University of Hawai'i Sea Grant publication number UNIH-SEAGRANT-JC-16-27.

Declaration of Competing Interest

The authors declare no conflicts of interest.

Acknowledgements

The authors would like to thank the laboratory assistance of Austin Macpherson and Daniel Woo during the course of the study.

References

- Ahel, M., Giger, W., 1993. Partitioning of alkylphenols and polyethoxylates between water and organic solvents. *Chemosphere* 26, 1471–1478.
- Allard, J.B.I., Duan, C.I., 2018. IGF-binding proteins: why do they exist and why are there so many? *Front. Endocrinol. (Lausanne)* 9, 117. <https://doi.org/10.3389/fendo.2018.00117>.
- Amaninejad, P., Sahafi, H.H., Soltani, M., Shekarabi, S.P.H., 2018. Endocrine disrupting effects of 4-nonylphenol on plasma vitellogenin, reproductive system and histology in koi carp (*Cyprinus carpio*). *Int. Aquat. Res.* 10, 263–274.
- Andersson, T., Friln, L., Hrdig, J., Larsson, K., 1988. Physiological disturbances in fish living in coastal water polluted with bleached kraft pulp mill effluents. *Can. J. Fish. Aquat. Sci.* 45, 1525–1536.
- Ankley, G.T., Bencic, D.C., Breen, M.S., Collette, T.W., Conolly, R.B., Denslow, N.D., Edwards, S.W., Ekman, D.R., Garcia-Reyero, N., Jensen, K.M., Lazorchak, J.M., Martinovic, D., Miller, D.H., Perkins, E.J., Orlando, E.F., Villeneuve, D.L., Wang, R.L., Watanabe, K.H., 2009. Endocrine disrupting chemicals in fish: developing exposure indicators and predictive models of effects based on mechanism of action. *Aquat. Toxicol.* 92, 168–178.
- Araújo, F.G., Moradoa, C.N., Parenteb, T.T.E., Paumgartenb, F.J.R., Gomes, I.D., 2018. Biomarkers and bioindicators of the environmental condition using a fish species (*Pimelodus maculatus* Lacepède, 1803) in a tropical reservoir in Southeastern Brazil. *Braz. J. Biol.* 78, 351–359.
- Aris, A.Z., Shamsuddin, A.S., Praveena, S.M., 2014. Occurrence of 17alpha-ethynylestradiol (EE2) in the environment and effect on exposed biota: a review. *Environ. Int.* 69, 104–119.
- Asem-Hiablie, S., Church, C.D., Elliott, H.A., Shappell, N.W., Schoenfuss, H.L., Drechsel, P., Williams, C.F., Knopf, A.L., Dabie, M.Y., 2013. Serum estrogenicity and biological responses in African catfish raised in wastewater ponds in Ghana. *Sci. Total Environ.* 463–464, 1182–1191.
- Authman, M.M.N., 2008. *Oreochromis niloticus* as a biomonitor of heavy metal pollution with emphasis on potential risk and relation to some biological aspects. *Glob. Vet.* 2, 104–109.
- Babu, B., Ozbay, G., 2013. Screening of imported tilapia filets for heavy metals and veterinary drug residues in the Mid-Atlantic Region, USA. *J. Food Process. Technol.* 4, 266. <https://doi.org/10.4172/2157-7110.1000266>.
- Bernanke, J., Kohler, H.R., 2009. The impact of environmental chemicals on wildlife vertebrates. *Rev. Environ. Contam. Toxicol.* 198, 1–47.
- Bertram, D.F., Chambers, R.C., Leggett, W.C., 1993. Negative correlations between larval and juvenile growth rates in winter flounder: implications of compensatory growth for variation in size-at-age. *Mar. Ecol. Prog. Ser.* 96, 209–215.
- Bhandari, R.K., Deem, S.L., Holliday, D.K., Jandegian, C.M., Kassotis, C.D., Nagel, S.C., Tillitt, D.E., Vom Saal, F.S., Rosenfeld, C.S., 2015. Effects of the environmental estrogenic contaminants bisphenol A and 17alpha-ethinyl estradiol on sexual development and adult behaviors in aquatic wildlife species. *Gen. Comp. Endocrinol.* 214, 195–219.
- Billiard, S.M., Khan, R.A., 2003. Chronic stress in cunner, *Tautoglabrus adspersus*, exposed to municipal and industrial effluents. *Ecotoxicol. Environ. Saf.* 55, 9–18.
- Bowman, C.J., Kroll, K.J., Gross, T.G., Denslow, N.D., 2002. Estradiol-induced gene expression in largemouth bass (*Micropterus salmoides*). *Mol. Cell. Endocrinol.* 196, 67–77.
- Breves, J.P., Duffy, T.A., Einarsdottir, I.E., Björnsson, B.T., McCormick, S.D., 2018. In vivo effects of 17alpha-ethinylestradiol, 17beta-estradiol and 4-nonylphenol on insulin-like growth-factor binding proteins (igfbps) in Atlantic salmon. *Aquat. Toxicol.* 203, 28–39.
- Breves, J.P., Fujimoto, C.K., Phipps-Costin, S.K., Einarsdottir, I.E., Björnsson, B.T., McCormick, S.D., 2017. Variation in branchial expression among insulin-like growth factor binding proteins (igfbps) during Atlantic salmon smoltification and seawater exposure. *BMC Physiol.* 17, 2.
- Breves, J.P., Hirano, T., Grau, E.G., 2010. Ionoregulatory and endocrine responses to disturbed salt and water balance in Mozambique tilapia exposed to confinement and handling stress. *Comp. Biochem. Physiol.* 155, 294–300.
- Breves, J.P., Tipsmark, C.K., Stough, B.A., Seale, A.P., Flack, B.R., Moorman, B.P., Lerner, D.T., Grau, E.G., 2014. Nutritional status and growth hormone regulate insulin-like growth factor binding protein (igfbp) transcripts in Mozambique tilapia. *Gen. Comp. Endocrinol.* 207, 66–73.
- Brix, R., Postigo, C., Gonzalez, S., Villagrana, M., Navarro, A., Kuster, M., de Alda, M.J., Barcelo, D., 2010. Analysis and occurrence of alkylphenolic compounds and estrogens in a European river basin and an evaluation of their importance as priority pollutants. *Anal. Bioanal. Chem.* 396, 1301–1309.
- Butler, A.A., LeRoith, D., 2001. Minireview: tissue-specific versus generalized gene targeting of the igf1 and igf1r genes and their roles in insulin-like growth factor physiology. *Endocrinology* 142, 1685–1688.
- Cailleaud, K., Forget-Leray, J., Souissi, S., Lardy, S., Augagneur, S., Budzinski, H., 2007. Seasonal variation of hydrophobic organic contaminant concentrations in the water-column of the Seine Estuary and their transfer to a planktonic species *Eurytemora affinis* (Calanoid, copepod). Part 2: Alkylphenol-polyethoxylates. *Chemosphere* 70, 281–287.
- Celino, F.T., Yamaguchi, S., Miura, C., Miura, T., 2009. Arsenic inhibits in vitro spermatogenesis and induces germ cell apoptosis in Japanese eel (*Anguilla japonica*). *Reproduction* 138, 279–287.
- Chambers, R.C., Leggett, W.C., Brown, J.A., 1988. Variation in and among early life-history traits of laboratory-reared winter flounder *Pseudopleuronectes americanus*. *Mar. Ecol. Prog. Ser.* 47, 1–15.
- Chen, W.L., Gwo, J.C., Wang, G.S., Chen, C.Y., 2014. Distribution of feminizing compounds in the aquatic environment and bioaccumulation in wild tilapia tissues. *Environ. Sci. Pollut. Res. Int.* 21, 11349–11360.
- Christiansen, T., Korsgaard, B., Jespersen, A., 1998. Induction of vitellogenin synthesis by nonylphenol and 17β-Estradiol and effects on the testicular structure in the eelpout *Zoarces viviparus*. *Mar. Environ. Res.* 46, 141–144.
- Cleveland, B.M., Weber, G.M., 2015. Effects of sex steroids on expression of genes regulating growth-related mechanisms in rainbow trout (*Oncorhynchus mykiss*). *Gen. Comp. Endocrinol.* 216, 103–115.
- Cleveland, B.M., Weber, G.M., 2016. Effects of steroid treatment on growth, nutrient partitioning, and expression of genes related to growth and nutrient metabolism in adult triploid rainbow trout (*Oncorhynchus mykiss*). *Domest. Anim. Endocrinol.* 56, 1–12.
- Colborn, T., Dumanoski, D., Myers, J.P., 1996. Our Stolen Future: Are We Threatening Our Fertility, Intelligence, and Survival? A Scientific Detective Story. Dutton, New York.
- Constancia, M., Hemberger, M., Hughes, J., Dean, W., Ferguson-Smith, A., Fundele, R., Stewart, F., Kelsey, G., Fowden, A., Sibley, C., Reik, W., 2002. Placental-specific IGF-II is a major modulator of placental and fetal growth. *Nature* 417, 945–948.
- Daughaday, W.H., Rotwein, P., 1989. Insulin-like growth factors I and II. Peptide, messenger ribonucleic acid and gene structures, serum, and tissue concentrations. *Endocr. Rev.* 10, 68–91.
- Davis, L.K., Hiramatsu, N., Hiramatsu, K., Reading, B.J., Matsubara, T., Hara, A., Sullivan, C.V., Pierce, A.L., Hirano, T., Grau, E.G., 2007. Induction of three vitellogenins by 17beta-estradiol with concurrent inhibition of the growth hormone-insulin-like growth factor 1 axis in a euryhaline teleost, the tilapia (*Oreochromis mossambicus*). *Biol. Reprod.* 77, 614–625.
- Davis, L.K., Pierce, A.L., Hiramatsu, N., Sullivan, C.V., Hirano, T., Grau, E.G., 2008. Gender-specific expression of multiple estrogen receptors, growth hormone receptors, insulin-like growth factors and vitellogenins, and effects of 17 beta-estradiol in the male tilapia (*Oreochromis mossambicus*). *Gen. Comp. Endocrinol.* 156, 544–551.
- Davis, L.K., Fox, B.K., Lim, C., Hiramatsu, N., Sullivan, C.V., Hirano, T., Grau, E.G., 2009a. Induction of vitellogenin production in male tilapia (*Oreochromis mossambicus*) by commercial fish diets. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* 154, 249–254.
- Davis, L.K., Visitacion, N., Riley, L.G., Hiramatsu, N., Sullivan, C.V., Hirano, T., Grau, E.G., 2009b. Effects of o,p'-DDE, heptachlor, and 17beta-estradiol on vitellogenin gene expression and the growth hormone/insulin-like growth factor-I axis in the tilapia, *Oreochromis mossambicus*. *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* 149, 507–514.
- Denslow, N.D., 1999. Vitellogenin as a biomarker of exposure for estrogen or estrogen mimics. *Exotoxicology* 8, 385–398.
- Desbrow, C., Routledge, E.J., Brighty, G.C., Sumpter, J.P., Waldock, M., 1998. Identification of estrogenic chemicals in STW effluent. 1. Chemical fractionation and in vitro biological screening. *Environ. Sci. Technol.* 32, 1549–1558.
- Duan, C., Plisetskaya, E.M., Dickhoff, W.W., 1995. Expression of insulin-like growth factor I in normally and abnormally developing coho salmon (*Oncorhynchus kisutch*). *Endocrinology* 136, 446–452.
- Duan, C., 1998. Nutritional and developmental regulation of insulin-like growth factors in fish. *J. Nutr.* 128, 306S–314S.
- Duan, C., Ren, H., Gao, S., 2010. Insulin-like growth factors (IGFs), IGF receptors, and IGF-binding proteins: roles in skeletal muscle growth and differentiation. *Gen. Comp. Endocrinol.* 167, 344–351.

- Duan, C., Xu, Q.J., 2005. Roles of insulin-like growth factor (IGF) binding proteins in regulating IGF actions. *Gen. Comp. Endocrinol.* 142, 44–52.
- Duffy, T.A., Iwanowicz, L.R., McCormick, S.D., 2014. Comparative responses to endocrine disrupting compounds in early life stages of Atlantic salmon, *Salmo salar*. *Aquat. Toxicol.* 152, 1–10.
- Ekelund, R., Bergman, A., Granmo, A., Berggren, M., 1990. Bioaccumulation of 4-nonylphenol in marine animals—a re-evaluation. *Environ. Pollut.* 64, 107–120.
- Ekelund, R., Granmo, A., Magnusson, K., Berggren, M., Bergman, A., 1993. Biodegradation of 4-nonylphenol in seawater and sediment. *Environ. Pollut.* 79, 59–61.
- Elango, A., Shepherd, B., Chen, T.T., 2006. Effects of endocrine disruptors on the expression of growth hormone and prolactin mRNA in the rainbow trout pituitary. *Gen. Comp. Endocrinol.* 145, 116–127.
- Fan, Y., Menon, R.K., Cohen, P., Hwang, D., Clemens, T., DiGirolamo, D.J., Kopchick, J.J., Le Roith, D., Trucco, M., Sperling, M.A., 2009. Liver-specific deletion of the growth hormone receptor reveals essential role of growth hormone signaling in hepatic lipid metabolism. *J. Biol. Chem.* 284, 19937–19944.
- FAO, 2005. 2005–2019. National Aquaculture Sector Overview. Mozambique. In: Omar, I. (Ed.), National Aquaculture Sector Overview Fact Sheets. FAO Fisheries and Aquaculture Department [online], Rome Updated 10 October.
- Filby, A.L., Thorpe, K.L., Tyler, C.R., 2006. Multiple molecular effect pathways of an environmental oestrogen in fish. *J. Mol. Endocrinol.* 37, 121–134.
- Flouriot, G., Pakdel, F., Valotaire, Y., 1996. Transcriptional and post-transcriptional regulation of rainbow trout estrogen receptor and vitellogenin gene expression. *Mol. Cell. Endocrinol.* 124, 173–183.
- Folmar, L.C., Hemmer, M., Hemmer, R., Bowman, C., Kroll, K., Denslow, N.D., 2000. Comparative estrogenicity of estradiol, ethynyl estradiol and diethylstilbestrol in an in vivo, male sheephead minnow (*Cyprinodon variegatus*), vitellogenin bioassay. *Aquat. Toxicol.* 49, 77–88.
- Fukazawa, Y., Siharath, K., Iguchi, T., Bern, H.A., 1995. In vitro secretion of insulin-like growth factor-binding proteins from liver of striped bass, *Morone saxatilis*. *Gen. Comp. Endocrinol.* 99, 239–247.
- Gagliano, M., McCormick, M.I., 2007. Compensating in the wild: is flexible growth the key to early juvenile survival? *Oikos* 116, 111–120.
- Gaigher, I.G., Krause, J.B., 1983. Growth rates of Mozambique tilapia (*Oreochromis mossambicus*) and silver carp (*Hypophthalmichthys molitrix*) without artificial feeding in floating cages in plankton-rich waste water. *Aquaculture* 31, 361–367.
- García de la Serrana, D., Fuentes, E.N., Martín, S.A.M., Johnston, I.A., Macqueen, D.J., 2017. Divergent regulation of insulin-like growth factor binding protein genes in cultured Atlantic salmon myotubes under different models of catabolism and anabolism. *Gen. Comp. Endocrinol.* 247, 53–65.
- García de la Serrana, D., Macqueen, D.J., 2018. Insulin-like growth factor-binding proteins of teleost fishes. *Front. Endocrinol.* 9, 80. <https://doi.org/10.3389/fendo.2018.00080>.
- Gaylord, T.G., Gatlin III, D.M., 2000. Assessment of compensatory growth in Channel catfish, *Ictalurus punctatus*, R. and associated changes in body condition indices. *J. World Aquac. Soc.* 31, 326–336. <https://doi.org/10.1111/j.1749-7345.2000.tb00884.x>.
- Giger, W., Brunner, P.H., Schaffner, C., 1984. 4-Nonylphenol in sewage sludge: accumulation of toxic metabolites from nonionic surfactants. *Science* 225, 623–625.
- Gimeno, S., Komen, H., Venderbosch, P.W.M., Bowmer, T., 1997. Disruption of sexual differentiation in genetic male common carp (*Cyprinus carpio*) exposed to an alkylphenol during different life stages. *Environ. Sci. Technol.* 31, 2884–2890.
- Goetz, F.W., Rise, M.L., Rise, M., Goetz, G.W., Binkowski, F., Shepherd, B.S., 2009. Stimulation of growth and changes in the hepatic transcriptome by 17 β -estradiol in the yellow perch (*Perca flavescens*). *Physiol. Genomics* 8, 261–280.
- Gray, E.S., Kelley, K.M., Law, S., Tsai, R., Young, G., Bern, H.A., 1992. Regulation of hepatic growth hormone receptors in coho salmon (*Oncorhynchus kisutch*). *Gen. Comp. Endocrinol.* 88, 243–252.
- Gross, J.M., Yee, D., 2002. How does the estrogen receptor work? *Breast Cancer Res.* 4, 62–64.
- Hanson, A.M., Ickstadt, A.T., Marquart, D.J., Kittilson, J.D., Sheridan, M.A., 2017. Environmental estrogens inhibit mRNA and functional expression of growth hormone receptors as well as growth hormone signaling pathways in vitro in rainbow trout (*Oncorhynchus mykiss*). *Gen. Comp. Endocrinol.* 246, 120–128.
- Harding, L.B., Schultz, I.R., da Silva, D.A.M., Ylitalo, G.M., Ragsdale, D., Harris, S.I., Bailey, S., Pepich, B.V., Swanson, P., 2016. Wastewater treatment plant effluent alters pituitary glandgonadotropin mRNA levels in juvenile coho salmon (*Oncorhynchus kisutch*). *Aquat. Toxicol.* 178, 118–131.
- Harries, J.E., Runnalls, T., Hill, E., Harris, C.A., Maddix, S., Sumpter, J.P., Tyler, C.R., 2000. Development of a reproductive performance for endocrine disrupting chemicals using pair-breeding fathead minnows (*Pimephales promelas*). *Environ. Sci. Technol.* 34, 3003–3011.
- Harries, J.E., Sheahan, D.A., Jobling, S., Matthiessen, P., Neall, P., Sumpter, J.P., Tyler, C.R., Zaman, N., 1997. Estrogenic activity in five United Kingdom rivers detected by measurement of vitellogenesis in caged male trout. *Environ. Toxicol. Chem.* 16, 534–542.
- Hassanin, A.I., Kuwahara, S., Nurhidayat, Tsukamoto, Y., Ogawa, K., Hiramatsu, K., Sasaki, F., 2002. Gonadosomatic index and testis morphology of common carp (*Cyprinus carpio*) in rivers contaminated with estrogenic chemicals. *J. Vet. Med. Sci.* 64, 921–926.
- Heide, A., Foss, A., Stefansson, S.O., Mayer, I., Norberg, B., Roth, B., Jenssen, M.D., Nortvedt, R., Imsland, A.K., 2006. Compensatory growth and fillet crude composition in juvenile Atlantic halibut: effects of short term starvation periods and subsequent feeding. *Aquaculture* 261, 109–117.
- Hemmatinezhad, B., Sarmeidani, M.M., Yoosefi, A.H., Fadaeifard, F., 2017. Assessment of four heavy metals mercury, lead, copper and cadmium levels in muscles of imported Tilapia to Iran. *J. Chem. Health Risk* 7, 133–138.
- Hemmer, M.J., Hemmer, B.L., Bowman, C.J., Kroll, K.J., Folmar, L.C., Marcovich, D., Hoglund, M.D., Denslow, N.D., 2001. Effects of p-nonylphenol, methoxychlor, and endosulfan on vitellogenin induction and expression in sheephead minnow (*Cyprinodon variegatus*). *Environ. Toxicol. Chem.* 20, 336–343.
- Hernandez, J.P., Chapman, L.M., Kretschmer, X.C., Baldwin, W.S., 2006. Gender-specific induction of cytochrome P450s in nonylphenol-treated FVB/NJ mice. *Toxicol. Appl. Pharmacol.* 216, 186–196.
- Hevroy, E.M., Azpeleta, C., Shimizu, M., Lanzén, A., Kaiya, H., Espe, M., Olsvik, P.A., 2011. Effects of short-term starvation on ghrelin, GH-IGF system, and IGF-binding proteins in Atlantic salmon. *Fish Physiol. Biochem.* 37, 217–232.
- Hoeflich, A., Wirthgen, E., David, R., Classen, C.F., Spitschak, M., Brenmoehl, J., 2014. Control of IGFBP-2 expression by steroids and peptide hormones in vertebrates. *Front. Endocrinol.* 5, 43. <https://doi.org/10.3389/fendo.2014.00043>.
- Hoeflich, A., Wu, M., Mohan, S., Foll, J., Wanke, R., Froehlich, T., Arnold, G.J., Lahm, H., Kolb, H.J., Wolf, E., 1999. Overexpression of insulin-like growth factor-binding protein-2 in transgenic mice reduces postnatal body weight gain. *Endocrinology* 140, 5488–5496.
- Holloway, A.C., Leatherland, J.F., 1997. Effect of gonadal steroid hormones on plasma growth hormone concentrations in sexually mature immature rainbow trout, *Oncorhynchus mykiss*. *Gen. Comp. Endocrinol.* 105, 246–254.
- Ishibashi, H., Hirano, M., Matsumura, N., Watanabe, N., Takao, Y., Arizono, K., 2006. Reproductive effects and bioconcentration of 4-nonylphenol in medaka fish (*Oryzias latipes*). *Chemosphere* 65, 1019–1026.
- Jalabert, B., 2005. Particularities of reproduction and oogenesis in teleost fish compared to mammals. *Reprod. Nutr. Dev. EDP Sci.* 45, 261–279.
- Jia, Y., Jing, Q., Gao, Y., Huang, B., 2019. Involvement and expression of growth hormone/insulin-like growth factor member mRNAs in the ovarian development of turbot (*Scophthalmus maximus*). *Fish Physiol. Biochem.* 45, 955–964. <https://doi.org/10.1007/s10695-018-0604-z>.
- Jobling, S., Sheahan, D., Osborne, J.A., Matthiessen, P., Sumpter, J.P., 1996. Inhibition of testicular growth in rainbow trout (*Oncorhynchus mykiss*) exposed to estrogenic alkylphenolic chemicals. *Environ. Toxicol. Chem.* 15, 194–202.
- Jones, P.D., Tremblay, L.A., De Coen, W.M., Giesy, J.P., 2000. Vitellogenin as a biomarker for environmental estrogens. *Australasian J. Ecotoxicol.* 6, 45–58.
- Kajimura, S., Kawaguchi, N., Kaneko, T., Kawazoe, I., Hirano, T., Visitacion, N., Grau, E.G., Aida, K., 2004. Identification of the growth hormone receptor in an advanced teleost, the tilapia (*Oreochromis mossambicus*) with special reference to its distinct expression pattern in the ovary. *J. Endocrinol.* 181, 65–76.
- Kajimura, S., Seale, A.P., Hirano, T., Cooke, I.M., Grau, E.G., 2005. Physiological concentrations of ouabain rapidly inhibit prolactin release from the tilapia pituitary. *Gen. Comp. Endocrinol.* 143, 240–250.
- Kamangar, B.B., Gabillard, J.C., Bobe, J., 2006. Insulin-like growth factor-binding protein (IGFBP)-1, -2, -3, -4, -5, and -6 and IGFBP-related protein 1 during rainbow trout postvitellogenesis and oocyte maturation: molecular characterization, expression profiles, and hormonal regulation. *Endocrinology* 147, 2399–2410.
- Kamei, H., Lu, L., Jiao, S., Li, Y., Gyrop, C., Laursen, L.S., Oxvig, C., Zhou, J., Duan, C., 2008. Duplication and diversification of the hypoxia-inducible IGFBP-1 gene in zebrafish. *PLoS One* 3, e3091.
- Kang, I.J., Yokota, H., Oshima, Y., Tsuruda, Y., Hano, T., Maeda, M., Imada, N., Tadokoro, H., Honjo, T., 2003. Effects of 4-nonylphenol on reproduction of Japanese medaka, *Oryzias latipes*. *Environ. Toxicol. Chem.* 22, 2438–2445.
- Karels, A.E., Soimasuo, M., Lappivaara, J., Leppänen, H., Aaltonen, T., Mellanen, P., Oikari, A.O.J., 1998. Effects of ECF-bleached kraft mill effluent on reproductive steroids and liver MFO activity in populations of perch and roach. *Ecotoxicology* 7, 123–132.
- Kiilerich, P., Tipsmark, C.K., Borski, R.J., Madsen, S.S., 2011. Differential effects of cortisol and 11-deoxycorticosterone on ion transport protein mRNA levels in gills of two euryhaline teleosts, Mozambique tilapia (*Oreochromis mossambicus*) and striped bass (*Morone saxatilis*). *J. Endocrinol.* 209, 115–126. <https://doi.org/10.1530/JOE-10-0326>.
- Kobayashi, Y., Nozu, R., Nakamura, M., 2011. Role of estrogen in spermatogenesis in initial phase males of the three-spot wrasse (*Halichoeres trimaculatus*): effect of aromatase inhibitor on the testis. *Dev. Dyn.* 240, 116–121.
- Komen, J., Lodder, P.A.J., Huskensi, F., Richter, C.J.J., Huisman, E.A., 1989. Effects of oral administration of 17 α -methyltestosterone and 17 β -estradiol on gonadal development in common carp, *Cyprinus carpio* L. *Aquaculture* 78, 349–363.
- Kukkonen, J.V.K., Punta, E., Koponen, P., Paranko, J., Leppänen, H., Holopainen, I.J., Hyvärinen, H., 1999. Biomarker responses by Crucian carp (*Carassius Carassius*) living in a pond of secondary treated pulp mill effluent. *Water Sci. Technol.* 40, 123–130.
- Lahnsteiner, F., Berger, B., Grubinger, F., Weismann, T., 2005. The effect of 4-nonylphenol on semen quality, viability of gametes, fertilization success, and embryo and larvae survival in rainbow trout (*Oncorhynchus mykiss*). *Aquat. Toxicol.* 71, 297–306.
- Leet, J.K., Gall, H.E., Sepúlveda, M.S., 2011. A review of studies on androgen and estrogen exposure in fish early life stages: effects on gene and hormonal control of sexual differentiation. *J. Aquat. Toxicol.* 31, 379–398.
- Lerner, D.T., Björnsson, B.T., McCormick, S.D., 2007a. Aqueous exposure to 4-nonylphenol and 17 β -estradiol increases stress sensitivity and disrupts ion regulatory ability of juvenile Atlantic salmon. *Environ. Toxicol. Chem.* 26, 1433–1440.
- Lerner, D.T., Björnsson, B.T., McCormick, S.D., 2007b. Larval exposure to 4-nonylphenol and 17 β -estradiol affects physiological and behavioral development of seawater adaptation in Atlantic salmon smolts. *Environ. Sci. Technol.* 41, 4479–4485.
- Lerner, D.T., Sheridan, M.A., McCormick, S.D., 2012. Estrogenic compounds decrease growth hormone receptor abundance and alter osmoregulation in Atlantic salmon.

- Gen. Comp. Endocrinol. 179, 196–204.
- Le Roith, D., Roberts, C.T.J., 2003. The insulin-like growth factor system and cancer. *Cancer Lett.* 195, 127–137.
- Le Roith, D., Scavo, L., Butler, A., 2001. What is the role of circulating IGF-I? *Trends Endocrinol. Metab.* 12, 48–52.
- Ma, T., Wan, X., Huang, Q., Wang, Z., Liu, J., 2005. Biomarker responses and reproductive toxicity of the effluent from a Chinese large sewage treatment plant in Japanese medaka (*Oryzias latipes*). *Chemosphere* 59, 281–288.
- Macqueen, D.J., Garcia de la Serrana, Daniel, Johnston, I.A., 2013. Evolution of ancient functions in the vertebrate insulin-like growth factor system uncovered by study of duplicated salmonid fish genomes. *Mol. Biol. Evol.* 30, 1060–1076.
- Magdeldin, S., Uchida, K., Hirano, T., Grau, E.G., Abdelfattah, A., Nozaki, M., 2007. Effects of environmental salinity on somatic growth and growth hormone/insulin-like growth factor-I axis in juvenile tilapia, *Oreochromis mossambicus*. *Fish. Sci.* 73, 1023–1032.
- Mao, Z., Zheng, X.F., Zhang, Y.Q., Tao, X.X., Li, Y., Wang, W., 2012. Occurrence and biodegradation of nonylphenol in the environment. *Int. J. Mol. Sci.* 13, 491–505.
- Matozzo, V., Gagne, F., Marin, M.G., Ricciardi, F., Blaise, C., 2008. Vitellogenin as a biomarker of exposure to estrogenic compounds in aquatic invertebrates: a review. *Environ. Int.* 34, 531–545.
- Meredith, H.O., Richman III, N.H., Collier, J.T., Seale, A.P., Riley, L.G., Ball, C.H., Shimoda, S.K., Stetson, M.H., Grau, E.G., 1999. Pesticide effects on prolactin release from the rostral pars distalis in vitro and their effects on growth in vivo in the tilapia (*Oreochromis mossambicus*). In: Hanshel, D.S., Black, M.C., Harrasse, M.C. (Eds.), *Environmental Toxicology and Risk Assessment*. American Society for Testing and Materials, West Conshohocken, PA, pp. 239–253.
- Miura, T., Miura, C., Ohta, T., Nader, M.R., Todo, T., Yamauchi, K., 1999. Estradiol-17 β stimulates the renewal of spermatogonial stem cells in males. *Biochem. Biophys. Res. Commun.* 264, 230–234.
- Miura, T., Ohta, T., Miura, C., Yamauchi, K., 2003. Complementary deoxyribonucleic acid cloning of spermatogonial stem cell renewal factor. *Endocrinology* 144, 5504–5510.
- Montserrat, N., Gabillard, J.C., Capilla, E., Navarro, M.I., Gutierrez, J., 2007. Role of insulin, insulin-like growth factors, and muscle regulatory factors in the compensatory growth of the trout (*Oncorhynchus mykiss*). *Gen. Comp. Endocrinol.* 150, 462–472.
- Moorman, B.P.I., Yamaguchi, Y., Lerner, D.T., Grau, E.G., Seale, A.P., 2016. Rearing Mozambique tilapia in tidally-changing salinities: Effects on growth and the growth hormone/insulin-like growth factor I axis. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* 198, 8–14. <https://doi.org/10.1016/j.cbpa.2016.03.014>.
- Nelson, E.R., Habibi, H.R., 2010. Functional significance of nuclear estrogen receptor subtypes in the liver of goldfish. *Endocrinology* 151, 1668–1676.
- Norbeck, L.A., Kittilson, J.D., Sheridan, M.A., 2007. Resolving the growth-promoting and metabolic effects of growth hormone: differential regulation of GH-IGF-I system components. *Gen. Comp. Endocrinol.* 151, 332–341.
- Norbeck, L.A., Sheridan, M.A., 2011. An in vitro model for evaluating peripheral regulation of growth in fish: effects of 17 β -estradiol and testosterone on the expression of growth hormone receptors, insulin-like growth factors, and insulin-like growth factor type 1 receptors in rainbow trout (*Oncorhynchus mykiss*). *Gen. Comp. Endocrinol.* 173, 270–280.
- Osman, A.G.M., 2012. Biomarkers in Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758) to assess the impacts of river Nile pollution: bioaccumulation, biochemical and tissues biomarkers. *J. Environ. Prot.* 3, 966–997.
- Park, C.B., Takemura, A., Aluru, N., Park, Y.J., Kim, B.H., Lee, C.H., Lee, Y.D., Moon, T.W., Vijayan, M.M., 2007. Tissue-specific suppression of estrogen, androgen and glucocorticoid receptor gene expression in feral vitellogenic male Mozambique tilapia. *Chemosphere* 69, 32–40.
- Picha, M.E., Turano, M.J., Tipsmark, C.K., Borski, R.J., 2008. Regulation of endocrine and paracrine sources of Igfs and Gh receptor during compensatory growth in hybrid striped bass (*Morone chrysops* X *Morone saxatilis*). *J. Endocrinol.* 199, 81–94. <https://doi.org/10.1677/JOE-07-0649>.
- Pierce, A.L., Breves, J.P., Moriyama, S., Hirano, T., Grau, E.G., 2011. Differential regulation of Igf1 and Igf2 mRNA levels in tilapia hepatocytes: effects of insulin and cortisol on GH sensitivity. *J. Endocrinol.* 211, 201–210.
- Pierce, A.L., Fox, B.K., Davis, L.K., Visitation, N., Kitahashi, T., Hirano, T., Grau, E.G., 2007. Prolactin receptor, growth hormone receptor, and putative somatolactin receptor in Mozambique tilapia: tissue specific expression and differential regulation by salinity and fasting. *Gen. Comp. Endocrinol.* 154, 31–40.
- Pierce, A.L., Shimizu, M., Beckman, B.R., Baker, D.M., Dickhoff, W.W., 2005. Time course of the GH/IGF axis response to fasting and increased ration in Chinook salmon (*Oncorhynchus tshawytscha*). *Gen. Comp. Endocrinol.* 140, 192–202.
- Peng, X., Yu, Y., Tang, C., Tan, J., Huang, Q., Wang, Z., 2008. Occurrence of steroid estrogens, endocrine-disrupting phenols, and acid pharmaceutical residues in urban riverine water of the Pearl River Delta, South China. *Sci. Total Environ.* 397, 158–166.
- Purdom, C.E., Hardiman, P.A., Bye, V.V.J., Eno, N.C.C., Tyler, R., Sumpter, J.P., 1994. Estrogenic effects of effluents from sewage treatment works. *Chem. Ecol.* 8, 275–285. <https://doi.org/10.1080/02757549408038554>.
- Rajaram, S., Baylink, D.J., Mohan, S., 1997. Insulin-like growth factor-binding proteins in serum and other biological fluids: regulation and functions. *Endocr. Rev.* 18, 801–831.
- Reindl, K.M., Sheridan, M.A., 2012. Peripheral regulation of the growth hormone-insulin-like growth factor system in fish and other vertebrates. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* 163, 231–245.
- Reinecke, M., 2010. Influences of the environment on the endocrine and paracrine fish growth hormone–insulin-like growth factor-I system. *J. Fish Biol.* 76, 1233–1254.
- Ricciarelli, E., Hernandez, E.R., Hurwitz, A., Kokia, E., Rosenfeld, R.G., Schwander, J., Adashi, E., 1991. The ovarian expression of the antigonadotropic insulin-like growth factor binding protein-2 is theca-interstitial cell-selective: evidence for hormonal regulation. *Endocrinology* 129, 2266–2268.
- Riley, L.G., Hirano, T., Grau, E.G., 2004. Estradiol-17 β and dihydrotestosterone differentially regulate vitellogenin and insulin-like growth factor-I production in primary hepatocytes of the tilapia *Oreochromis mossambicus*. *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* 138, 177–186.
- Roberts, R.J., 2012. *Fish Pathology*. Wiley-Blackwell, West Sussex, UK.
- Sadekarpawar, S., Parikh, P., 2013. Gonadosomatic and Hepatosomatic Indices of Freshwater Fish *Oreochromis mossambicus* in Response to a Plant Nutrient. *World J. Zool.* 8, 110–118.
- Segers, F.H., Berishvili, G., Taborsky, B., 2012. Egg size-dependent expression of growth hormone receptor accompanies compensatory growth in fish. *Proc. Biol. Sci. R. Soc.* 279, 592–600.
- Servos, M.R., Maguire, R.J., Bennie, D.T., Lee, H.B., Cureton, P.M., Davidson, N., Sutcliffe, R., Rawn, D.F.K., 2003. An ecological risk assessment of nonylphenol and its ethoxylates in the aquatic environment. *Hum. Ecol. Risk Assess.* 9, 569–587.
- Shanley, E.K., Xu, W., 2011. Endocrine disrupting chemicals targeting estrogen receptor signaling: identification and mechanisms of action. *Chem. Res. Toxicol.* 24, 6–19.
- Shimizu, M., Beckman, B.R., Hara, A., Dickhoff, W.W., 2006. Measurement of circulating salmon IGF binding protein-1: assay development, response to feeding ration and temperature, and relation to growth parameters. *J. Endocrinol.* 188, 101–110.
- Shimizu, M., Cooper, K.A., Dickhoff, W.W., Beckman, B.R., 2009. Postprandial changes in plasma growth hormone, insulin, insulin-like growth factor (IGF)-I, and IGF-binding proteins in coho salmon fasted for varying periods. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* 297, 352.
- Shimizu, M., Dickey, J.T., Fukada, H., Dickhoff, W.W., 2005. Salmon serum 22 kDa insulin-like growth factor-binding protein (IGFBP) is IGFBP-1. *J. Endocrinol.* 184, 267–276.
- Shimizu, M., Dickhoff, W.W., 2017. Circulating insulin-like growth factor binding proteins in fish: their identities and physiological regulation. *Gen. Comp. Endocrinol.* 252, 150–161.
- Shimizu, M., Swanson, P., Dickhoff, W.W., 1999. Free and protein-bound insulin-like growth factor-I (IGF-I) and IGF-binding proteins of coho salmon, *Oncorhynchus kisutch*. *Gen. Comp. Endocrinol.* 115, 398–405.
- Shimizu, M., Swanson, P., Hara, A., Dickhoff, W.W., 2003. Purification of a 41-kDa insulin-like growth factor binding protein from serum of chinook salmon, *Oncorhynchus tshawytscha*. *Gen. Comp. Endocrinol.* 132, 103–111.
- Shved, N., Berishvili, G., D'Cotta, H., Baroiller, J., Segner, H., Eppler, E., Reinecke, M., 2007. Ethinylestradiol differentially interferes with IGF-I in liver and extrahepatic sites during development of male and female bony fish. *J. Endocrinol.* 195, 513–523.
- Shved, N., Berishvili, G., Baroiller, J., Segner, H., Reinecke, M., 2008. Environmentally relevant concentrations of 17 α -ethinylestradiol (EE2) interfere with the growth hormone (GH)/insulin-like growth factor (IGF)-I system in developing bony fish. *Toxicol. Sci.* 106, 93–102.
- Tipmark, C.K., Breves, J.P., Seale, A.P., Lerner, D.T., Hirano, T., Grau, E.G., 2011. Switching of Na⁺ + K⁺-ATPase isoforms by salinity and prolactin in the gill of a cichlid fish. *J. Endocrinol.* 209, 237–244.
- Vajda, A.M., Barber, L.B., Gray, J.L., Lopez, E.M., Woodling, J.D., Norris, D.O., 2008. Reproductive disruption in fish downstream from an estrogenic wastewater effluent. *Environ. Sci. Technol.* 42, 3407–3414.
- Vethaak, A.D., Lahr, J., Schrap, S.M., Belfroid, A.C., Rijs, G.B., Gerritsen, A., de Boer, J., Bulder, A.S., Grinwis, G.C., Kuiper, R.V., Legler, J., Murk, T.A., Peijnenburg, W., Verhaar, H.J., de Voogt, P., 2005. An integrated assessment of estrogenic contamination and biological effects in the aquatic environment of The Netherlands. *Chemosphere* 59, 511–524.
- Wang, X., Lu, L., Li, Y., Li, M., Chen, C., Feng, Q., Zhang, C., Duan, C., 2009. Molecular and functional characterization of two distinct IGF binding protein-6 genes in zebrafish. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* 296, 1348–1357.
- Woltering, D.M., 1984. The growth response in fish chronic and early life stage toxicity tests: A critical review. *Aquat. Toxicol.* 5, 1–21.
- Xu, W., Yan, W., Huang, W., Miao, L., Zhong, L., 2014. Endocrine-disrupting chemicals in the Pearl River Delta and coastal environment: sources, transfer, and implications. *Environ. Geochem. Health* 36, 1095–1104.
- Yadette, F., Male, R., 2002. Effects of 4-nonylphenol on gene expression of pituitary hormones in juvenile Atlantic salmon (*Salmo salar*). *Aquat. Toxicol.* 58, 113–129.
- Zhou, X., Yang, Z., Luo, Z., Li, H., Chen, G., 2019. Endocrine disrupting chemicals in wild freshwater fishes: species, tissues, sizes and human health risks. *Environ. Pollut.* 244, 462–468.
- Zou, J.J., Trudeau, V.L., Cui, Z., Brechin, J., Mackenzie, K., Zhu, Z., Houlihan, D.F., Peter, R.E., 1997. Estradiol stimulates growth hormone production in female goldfish. *Gen. Comp. Endocrinol.* 106, 102–112.