

Abraham Model Correlations for Describing Solute Transfer Processes into Diethyl Carbonate

Jingyi Dai^a, Shrika Eddula^a, Carina Jiang^a, Alex Zhang^a, Kelly Liu^a, Siqi Zhu^a, Shang Wang^a, Avi Gupta^a, Brittani Churchill^a, Estefania Garcia^a, William E. Acree, Jr.^a and Michael H. Abraham^b

^a Department of Chemistry, University of North Texas, 1155 Union Circle Drive #305070, Denton, TX 76203 (USA)

^b Department of Chemistry, University College London, 20 Gordon Street, London WC1H 0AJ (UK)

Abstract

Experimental solubilities have been determined for 1-chloroanthraquinone, anthracene, biphenyl, pyrene, benzil, benzoic acid, acetylsalicylic acid, 4-*tert*-butylbenzoic acid, 3,4-dichlorobenzoic acid, 2-chloro-5-nitrobenzoic acid, 4-chloro-3-nitrobenzoic acid, 2-methoxybenzoic acid, 4-methoxybenzoic acid, 3,4-dimethoxybenzoic acid, 3,4,5-trimethoxybenzoic acid, 2-methylbenzoic acid, 3-methylbenzoic acid, 4-methyl-3-nitrobenzoic acid, 4-nitrobenzoic acid, benzoin, salicylamine, thioxanthen-9-one, 1,4-dichloro-2-nitrobenzene and phenothiazine dissolved in diethyl carbonate at 298.15 K. Results of our experimental measurements, combined with published solubility and activity coefficient data, have been used to derive Abraham model correlations for describing solute transfer into diethyl carbonate. The derived mathematical correlations back-calculate the observed experimental data to within an overall standard deviation of 0.13 log units.

Key Words and Phrases:

Diethyl carbonate solvent; Solubilities of crystalline organic compounds; Abraham model correlations; Solute transfer processes

*To whom correspondence should be addressed. (E-mail: acree@unt.edu)

1. Introduction

Organic solvents play an important role in many chemical manufacturing processes. They provide a liquid medium for the synthesis of new products, for the development of new analytical methodologies based on spectroscopic measurements and volumetric titrations, and for the extraction and isolation of phytochemicals from plant materials. Each year the chemical manufacturing sector consumes and disposes of several million tons of organic solvents that are harmful to the environment. The environmental impact is profound, and in many instances is irreversible. If not checked the environmental damage is likely to increase as the annual consumption of organic solvents is projected to grow in future years to accommodate the expected population growth. Increased environmental awareness, combined public concern regarding workplace safety, have resulted in new governmental regulations regarding the utilization and subsequent disposal of toxic chemicals in industrial manufacturing processes. To address these concerns the manufacturing sector is continuously searching for safer solvent alternatives and for “solvent free” synthetic methods. Replacement of harmful organic solvents is not an easy task as very little information is available regarding the solubilizing properties of many of the potential solvent candidates.

In recent years [1-15] there has been renewed interest in performing solubility measurements for crystalline organic compounds dissolved both in organic mono-solvents and in binary solvent mixtures. The primary objective of these studies has been to identify suitable solvents for purifying medicinal and cosmetic compounds, pesticides and herbicides, and other synthesized chemical products through recrystallization. For the most part the fore-mentioned studies have been limited to only a handful of different organic mono-solvents and solvent

mixtures. Very little attention was given to extending the measured data to predict the solubility of the studied compounds in additional organic solvents.

Our solubilization studies [16-24] differ from those performed by other research groups in that we have used our experimental data to develop Abraham model correlations [25-27]:

$$\log P \text{ and } \log\left(\frac{c_{\text{solute,org solv}}^{\text{sat}}}{c_{\text{solute,water}}^{\text{sat}}}\right) = c_p + e_p \cdot \mathbf{E} + s_p \cdot \mathbf{S} + a_p \cdot \mathbf{A} + b_p \cdot \mathbf{B} + v_p \cdot \mathbf{V} \quad (1)$$

$$\log K \text{ and } \log\left(\frac{c_{\text{solute,org solv}}^{\text{sat}}}{c_{\text{solute,gas phase}}^{\text{sat}}}\right) = c_k + e_k \cdot \mathbf{E} + s_k \cdot \mathbf{S} + a_k \cdot \mathbf{A} + b_k \cdot \mathbf{B} + l_k \cdot \mathbf{L} \quad (2)$$

that can be used to predict molar solute solubilities and infinite dilution activity coefficients in additional organic solvents. The Abraham solvation parameter model is based on a firm, fundamental understanding of how molecules interact in fluid solution. Each term on the right-hand side of Eqns. 1 and 2 represent a different type of molecular interaction as quantified by the product of a solute property times the complementary solvent property. Solute properties (called solute descriptors) are defined as follows: **E** corresponds to the molar refraction of the given solute in excess of that of a linear alkane having a comparable molecular size; **S** is a combination of the electrostatic polarity and polarizability of the dissolved solute; **A** and **B** quantify the respective hydrogen-bond donating and hydrogen-bond accepting abilities of the solute; **V** refers to the solute's McGowan molecular volume (calculated from atomic sizes and chemical bond numbers [28]); and **L** denotes the logarithm of the solute's measured gas-to-hexadecane partition coefficient at 298.15 K.

The complementary solvent properties ($c_p, e_p, s_p, a_p, b_p, v_p, c_k, e_k, s_k, a_k, b_k$, and l_k) in Eqns. 1 and 2 are obtained by curve-fitting logarithms of experimental water-to-organic solvent partition/transfer coefficients, $\log P$, logarithms of experimental gas-to-organic solvent

partition/transfer coefficients, $\log K$, and logarithms of molar solubility ratios, $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,water}^{sat}}\right)$

and $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,gas\,phase}^{sat}}\right)$, in accordance to the Abraham model. The subscripts “org solv”, “water” and “gas phase” associated with the molar solubilities indicate the phase to which the molar solubility pertains. For example $C_{solute,org\,solv}^{sat}$ and $C_{solute,water}^{sat}$ denote the molar solubility of the solute in the organic solvent and in water, respectively. The quantity $C_{solute,gas}^{gas}$ is the molar concentration of the solute in the gas phase whose numerical value is often determined as part of the solute descriptor calculations. The Abraham model is described in greater detail in several earlier publications [25, 26, 29-31]. The advantage that the Abraham model offers over solution models used by other researchers is that once an Abraham model equation coefficients have been determined for a given solvent, one can then predict the solubility and/or infinite dilution activity coefficients of more than 8,000 different compounds in the given solvent. Abraham model solute descriptors are readily available for over 8,000 organic, organometallic and inorganic compounds [32].

To date we have determined Abraham model process coefficients for several different classes of organic compounds including alkanes, alkylbenzene, chloroalkanes, perfluoroalkanes, halogenated benzenes, alkanols and alkanediols, alkyl alkanoates, dialkyl ethers, alkoxyalcohols, alkanenitriles, alkylformamides, alkylacetamides and many other miscellaneous organic solvents (dimethyl sulfoxides, aniline, acetophenone, benzonitrile, nitrobenzene, carbon disulfide, acetic acid, tributyl phosphate, propylene carbonate) [16-24, 27, 33-35]. Process equation coefficients have also been determined for binary aqueous-methanol [36] and binary aqueous-ethanol [37, 38] solvent mixtures. The aforementioned correlations provide the chemical manufacturer sector with predictive solubility and solute transfer expressions for a wide range of chemically diverse solvent

media. One solvent class that is not very well represented by our earlier studies is alkyl carbonates. We have studied only a single alkyl carbonate, namely propylene carbonate [39].

In the present communication we extend our solubilization studies to include diethyl carbonate, which is an important organic co-solvent used in lithium ion batteries [40, 41] and as an extraction solvent for the removal of carboxylic acids from aqueous solutions [42 - 44]. Organic carbonates have also been studied as possible “green” solvents in polymer membrane preparation [45]. As noted by Rasool and coworkers [45] “Organic carbonates in general are environmentally friendly, have low eco-toxicity, and show good biodegradability.” Experimental solubilities have been determined for 1-chloroanthraquinone, anthracene, biphenyl, pyrene, benzil, benzoic acid, acetylsalicylic acid, 4-*tert*-butylbenzoic acid, 3,4-dichlorobenzoic acid, 2-chloro-5-nitrobenzoic acid, 4-chloro-3-nitrobenzoic acid, 2-methoxybenzoic acid, 4-methoxybenzoic acid, 3,4-dimethoxybenzoic acid, 3,4,5-trimethoxybenzoic acid, 2-methylbenzoic acid, 3-methylbenzoic acid, 4-methyl-3-nitrobenzoic acid, 4-nitrobenzoic acid, benzoin, salicylamine, thioxanthen-9-one, 1,4-dichloro-2-nitrobenzene and phenothiazine dissolved in diethyl carbonate at 298.15 K. Results of our measurements, combined with published solubility data for helium [46], neon [46], argon [46], krypton [46], xenon [46], hydrogen [46], nitrogen [46], oxygen [47], sulfur hexafluoride [46], carbon dioxide [48], tetrafluorocarbon [46], hydrogen sulfide [48], methane [46], ethane [46], ethene [46], 1,1,1,2-tetrafluoroethane [49] and diphenyl carbonate [50] and published infinite dilution activity coefficient data for hexane [51], octane [52], dodecane [53], cyclohexane [54], benzene [55], carbon tetrachloride [56], methyl *tert*-butyl ether [57] and dimethyl carbonate [58, 59], have been used to derive Abraham model correlations for solute transfer into diethyl carbonate from both water (Eqn. 1) and from the gas phase (Eqn. 2).

2. Chemical Materials and Experimental Methodology

In assembling the crystalline organic compounds for the solubility determinations we selected compounds that exhibit a wide range of chemical diversity and that span a wide range of solute descriptors values. All chemicals used in the solubility measurements were purchased from commercial sources. The sample of diethyl carbonate was dried over activated molecular sieves and distilled shortly before use. Samples of 1-chloroanthraquinone, benzil, salicylamide, thioxanthen-9-one, pyrene, 1,4-dichloro-2-nitrobenzene, and biphenyl were recrystallized three times from anhydrous methanol and dried for two days at 333 K prior to use. Anthracene was recrystallized three times from anhydrous acetone and dried for two days at 333 K. The carboxylic acid solutes were dried for two days at 333 K prior to use. Gas chromatographic analysis (flame ionization detector) indicated that the purities of purified samples of diethyl carbonate, 1-chloroanthraquinone, benzil, anthracene, biphenyl, pyrene, 1,4-dichloro-2-nitrobenzene, thioxanthen-9-one and salicylamide were 0.997 mass fraction. The purities of the fourteen carboxylic acid samples were determined by a non-aqueous titrimetric method based on the published method of Fritz and Lisicki [60]. The method was modified for health reasons in that benzene was replaced by toluene in preparing the titration solvent. Acid-base titrations with freshly standardized sodium methoxide (thymol blue indicator) indicated that the purities of acetylsalicylic acid, benzoic acid, 4-*tert*-butylbenzoic acid, 3,4-dichlorobenzoic acid, 2-chloro-5-nitrobenzoic acid, 4-chloro-3-nitrobenzoic acid, 2-methoxybenzoic acid, 4-methoxybenzoic acid, 3,4-dimethoxybenzoic acid, 3,4,5-trimethoxybenzoic acid, 2-methylbenzoic acid, 3-methylbenzoic acid, 4-methyl-3-nitrobenzoic acid, and 4-nitrobenzoic acid were 0.998 (± 0.003) mass fraction. The commercial suppliers and chemical purities of the chemicals used in the solubility measurements are summarized in Table 1.

Table 1. Chemical Sources and Mass Fraction Purities of Chemicals Used in the Solubility Studies

Chemical	Supplier	Purification Method	Purity (Mass Fraction)
Diethyl carbonate	Acros Organics, Morris Plains, New Jersey, USA	Stored over molecular and distilled	0.997
1-Chloroanthraquinone	Aldrich Chemical Company, Milwaukee, Wisconsin, USA	Recrystallization from anhydrous methanol	0.997
Phenothiazine	Acros Organics	Used as received	0.99
Biphenyl	Aldrich Chemical Company	Recrystallization from anhydrous methanol	0.997
Benzil	Aldrich Chemical Company	Recrystallization from anhydrous methanol	0.997
Anthracene	Aldrich Chemical Company	Recrystallization from anhydrous acetone	0.997
Pyrene	Aldrich Chemical Company	Recrystallization from anhydrous methanol	0.997
Acetylsalicylic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
Thioxanthen-9-one	Aldrich Chemical Company	Recrystallization from anhydrous methanol	0.997
Salicylamide	Aldrich Chemical Company	Recrystallization from anhydrous methanol	0.997
Benzoin	Aldrich Chemical Company	Recrystallization from anhydrous methanol	0.997
1,4-Dichloro-2-nitrobenzene	TCI America Chemical Company, Portland, Oregon, USA	Recrystallization from anhydrous methanol	0.997
Benzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
4- <i>tert</i> -Butylbenzoic acid	TCI America Chemical Company	Dried for two days at 333 K	0.998
2-Chloro-5-nitrobenzoic acid	Acros Organics	Dried for two days at 333 K	0.998
4-Chloro-3-nitrobenzoic acid	Acros Organics	Dried for two days at 333 K	0.998
3,4-Dichlorobenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
2-Methoxybenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
4-Methoxybenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998

3,4-Dimethoxybenzoic acid	Across Organics	Dried for two days at 333 K	0.998
3,4,5-Trimethoxybenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
2-Methylbenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
3-Methylbenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
4-Methyl-3-nitrobenzoic acid	Aldrich Chemical Company	Dried for two days at 333 K	0.998
4-Nitrobenzoic acid	Acros Organics	Dried for two days at 333 K	0.998
Toluene	Aldrich Chemical Company	None	0.998, anhydrous
Sodium methoxide, 25 mass % solution in methanol	Aldrich Chemical Company	None	
2-Propanol	Aldrich Chemical Company	None	0.99

Solubilities of the 24 crystalline organic compounds were measured using a static equilibration method followed by a spectroscopic determination of the concentration of the dissolved solute in the saturated solution based on the Beer-Lambert law. The experimental methodology that we use has been given in many of our earlier publications [61-65], and to conserve journal space will only be briefly described in this communication. Aliquots of the saturated solutions were transferred by syringe into weighed volumetric flasks after the samples had equilibrated in sealed amber glass bottles in a constant temperature water bath at 298.15 ± 0.05 K for at least three days. The samples were periodically agitated to facilitate dissolution and mixing. The volumetric flasks containing the transferred aliquots were weighed on an electronic analytical balance. The transferred solutions were diluted quantitatively with 2-propanol. Absorbances of the diluted solutions and of the nine standard solutions of known solute concentrations were recorded on a Milton Roy Spectronic 1000 Plus spectrophotometer (Milton Roy, Rochester, NY, USA). The concentration of each diluted solution was computed from a

Beer-Lambert law plot of absorbance versus concentration curve obtained from the measured absorbances of nine standard solutions of known solute molarity. The analysis wavelengths and concentration ranges used for each solute have been given in our earlier publications [61-81]. We checked to make sure that diethyl carbonate did not absorb light at the analysis wavelengths and at the diethyl carbonate concentrations in the diluted solutions.

Molar concentrations of the diluted samples deduced from the Beer-Law law graph were converted into mole fraction solubilities using the volume of the volumetric flasks that the solute was transferred into, the mass of the sample analyzed, molar masses of the solvent and the respective solutes, and any dilutions that may have been needed in order for the sample's measured absorbance to fall on the Beer-Lambert law curve. We checked each equilibrated solid phase for possible solvate formation by determining the melting point temperature of the equilibrated solid phases recovered from the saturated solutions after the solubility measurements of each solute were completed. The melting point temperature of each equilibrated solid phase was within ± 0.5 K of the melting point temperature of the commercial sample or recrystallized compound prior to being in contact with the diethyl carbonate solvent.

The experimental mole fraction solubilities, x_{solute}^{sat} , of the crystalline organic solutes in diethyl carbonate are tabulated in the second column of Table 2. The numerical values represent the average of 4 to 10 independent experimental determinations, which were reproducible to within ± 2 % (relative error). To the best of our knowledge there are no published solubility data for these organic solutes in diethyl carbonate that we can compare our experimental values against. The only published solubility data that we were able to find for solid organic compounds dissolved in diethyl carbonate were the value previously published by Li and coworkers [50] for diphenyl carbonate.

Table 2. Mole Fraction Solubilities, x_{solute}^{sat} , of 24 Crystalline Organic Compounds Dissolved in Diethyl Carbonate at a Temperature of 298.15 K

Crystalline Organic Compound	x_{solute}^{sat}
1-Chloroanthraquinone	0.00775
Phenothiazine	0.0376
Biphenyl	0.234
Benzil	0.133
Anthracene	0.00547
Pyrene	0.0451
Acetylsalicylic acid	0.0226
Thioxanthen-9-one	0.00323
Salicyl amide	0.0364
Benzoin	0.109
1,4-Dichloro-2-nitrobenzene	0.438
Benzoic acid	0.112
4- <i>tert</i> -Butylbenzoic acid	0.0332
2-Chloro-5-nitrobenzoic acid	0.0309
4-Chloro-3-nitrobenzoic acid	0.0133
3,4-Dichlorobenzoic acid	0.00809
2-Methoxybenzoic acid	0.0366
4-Methoxybenzoic acid	0.00600
3,4-Dimethoxybenzoic acid	0.00401
3,4,5-Trimethoxybenzoic acid	0.00698
2-Methylbenzoic acid	0.0996
3-Methylbenzoic acid	0.0951
4-Methyl-3-nitrobenzoic acid	0.0123
4-Nitrobenzoic acid	0.00614

3. Development of Abraham Model Correlations for Solute Transfer into Diethyl Carbonate

Development of Abraham model correlations to describe solute transfer into diethyl carbonate requires constructing an Eqn 1 and Eqn. 2 for each of the 50 individual organic and inorganic solutes considered in the current study. The solute descriptors needed for the right-hand

sides of the two equations are tabulated in Table 3. The transfer properties needed on the left-hand side of Eqns. 1 and 2 are calculated from the experimental mole fraction solubility data given in Table 2 or from the experimental solubility data and activity coefficients retrieved from the published chemical and engineering. Mole fraction based solubilities are converted into molar solubilities by dividing x_{solute}^{sat} by the ideal molar volume of the saturated solution:

$$C_{solute,org\ solv}^{sat} = \frac{x_{solute}^{sat}}{(1-x_{solute}^{sat})V_{solvent} + x_{solute}^{sat}V_{solute}} \quad (3)$$

The numerical values used for the molar volumes of the hypothetical subcooled liquid solutes are given in our earlier publications [63-86], along with the aqueous molar solubilities, $C_{solute,water}^{sat}$, and solute molar gas concentrations, $C_{solute,gas}^{gas}$, needed in calculating the two molar solubility ratios. The mole fraction solubility of diphenyl carbonate in diethyl carbonate that was retrieved from the published literature [50] was similarly converted to a molar solubility. The molar solubility ratios are tabulated in the eighth and ninth columns of Table 3.

Table 3. Logarithms of Experimental Partition Coefficients, $\log P$ and $\log K$, Logarithms of Molar Solubility Ratios, $\log(\frac{C_{solute,org\ solv}^{sat}}{C_{solute,water}^{sat}})$ and $\log(\frac{C_{solute,org\ solv}^{sat}}{C_{solute,gas\ phase}^{sat}})$, and Solute Descriptors for Organic Compounds and Inorganic Gases Dissolved in Diethyl Carbonate at 298.15 K

Solute	E	S	A	B	L	V	Log K^a	Log P^b	Ref.
Helium	0.000	0.000	0.000	0.000	-1.741	0.0680	-1.504	0.516	[46]
Neon	0.000	0.000	0.000	0.000	-1.575	0.0850	-1.378	0.582	[46]
Argon	0.000	0.000	0.000	0.000	-0.688	0.1900	-0.571	0.899	[46]
Krypton	0.000	0.000	0.000	0.000	-0.211	0.2460	-0.132	1.078	[46]
Xenon	0.000	0.000	0.000	0.000	0.378	0.3290	0.420	1.390	[46]
Hydrogen	0.000	0.000	0.000	0.000	-1.200	0.1086	-1.107	0.613	[46]
Nitrogen	0.000	0.000	0.000	0.000	-0.978	0.2222	-0.813	0.987	[46]
Oxygen	0.000	0.000	0.000	0.000	-0.723	0.1830	-0.713	0.787	[47]

Sulfur hexafluoride	-0.600	-0.200	0.000	0.000	-0.120	0.4643	0.018	2.238	[46]
Tetrafluoromethane	-0.580	-0.260	0.000	0.000	-0.817	0.3203	-0.667	1.623	[46]
Carbon dioxide	0.000	0.280	0.050	0.100	0.058	0.2809	0.713	0.793	[48]
Hydrogen sulfide	0.350	0.310	0.100	0.070	0.723	0.2721	1.056	0.656	[48]
Methane	0.000	0.000	0.000	0.000	-0.323	0.2495	-0.261	1.199	[46]
Ethane	0.000	0.000	0.000	0.000	0.492	0.3904	0.495	1.835	[46]
Ethene	0.107	0.100	0.000	0.070	0.289	0.3474	0.499	1.439	[46]
Hexane	0.000	0.000	0.000	0.000	2.668	0.9540	2.887	4.707	[51]
Octane	0.000	0.000	0.000	0.000	3.677	1.2358	3.437	5.547	[52]
Dodecane	0.000	0.000	0.000	0.000	5.696	1.7994	5.441	7.971	[53]
Cyclohexane	0.305	0.100	0.000	0.000	2.964	0.8454	2.610	3.510	[54]
1,1,1,2-Tetrafluoroethane	-0.410	0.410	0.060	0.020	0.530	0.4612	1.562	1.972	[49]
Tetrachloromethane	0.460	0.380	0.000	0.000	2.823	0.7391	3.134	3.194	[56]
Benzene	0.610	0.520	0.000	0.140	2.786	0.7164	3.232	2.602	[55]
Methyl <i>tert</i> -butyl ether	0.024	0.220	0.000	0.590	2.380	0.8718	2.674	1.084	[57]
Dimethyl carbonate	0.142	0.540	0.000	0.570	2.328	0.6644	3.308		[58,59]
Diethyl carbonate	0.060	0.580	0.000	0.530	3.412	0.9462	4.099	1.659	Unity
Diphenyl carbonate	1.280	1.530	0.000	0.590	7.843	1.5982	9.374	4.242	[50]
1-Chloroanthraquinone	1.900	1.790	0.000	0.570	9.171	1.6512	10.376	4.342	This work
Anthracene	2.290	1.340	0.000	0.280	7.568	1.4544	8.111	5.081	This work
Biphenyl	1.360	0.990	0.000	0.260	6.014	1.3242	6.540	4.590	This work
Pyrene	2.808	1.710	0.000	0.280	8.833	1.5850	9.211	5.711	This work
Benzil	1.445	1.590	0.000	0.620	7.611	1.6374	8.930	4.060	This work
Benzoic acid	0.730	0.900	0.590	0.400	4.657	0.9317	6.660	1.520	This work
Acetylsalicylic acid	0.781	1.690	0.710	0.670	6.279	1.2879	9.446	0.946	This work
4- <i>tert</i> -Butylbenzoic acid	0.730	1.111	0.551	0.443	6.547	1.4953	8.553	3.329	This work
3,4-Dichlorobenzoic acid	0.950	0.920	0.670	0.260	5.623	1.1766	7.542	2.802	This work
2-Chloro-5-nitrobenzoic acid	1.250	1.400	0.670	0.460	6.513	1.2283	8.941	1.991	This work
4-Chloro-3-nitrobenzoic acid	1.250	1.470	0.700	0.440	6.685	1.2283	9.249	2.039	This work
2-Methoxybenzoic acid	0.899	1.410	0.450	0.620	5.636	1.1313	7.831	1.031	This work
4-Methoxybenzoic acid	0.899	1.250	0.620	0.520	5.741	1.1313	8.192	1.492	This work
3,4-Dimethoxybenzoic acid	0.950	1.646	0.570	0.755	6.746	1.3309	9.459	1.012	This work
3,4,5-Trimethoxybenzoic acid	1.001	1.760	0.603	0.850	7.711	1.5309	10.562	1.307	This work
2-Methylbenzoic acid	0.730	0.840	0.420	0.440	4.677	1.0726	6.272	1.972	This work
3-Methylbenzoic acid	0.730	0.890	0.600	0.400	4.819	1.0726	7.012	2.032	This work
4-Methyl-3-nitrobenzoic acid	1.040	1.461	0.659	0.521	6.434	1.2468	8.972	1.687	This work
4-Nitrobenzoic acid	0.990	1.520	0.680	0.400	5.770	1.1059	8.582	1.682	This work
Thioxanthen-9-one	1.940	1.441	0.000	0.557	8.436	1.5357	9.031	3.963	This work
Benzoin	1.585	2.115	0.196	0.841	9.159	1.6804	11.351	2.620	This work
Salicylamide	1.160	1.650	0.630	0.480	5.910	1.0315	8.912	1.227	This work
1,4-Dichloro-2-nitrobenzene	1.120	1.289	0.000	0.199	5.783	1.1354	6.747	3.846	This work

Phenothiazine	1.890	1.560	0.310	0.300	8.389	1.4789	9.987	4.584	This work
---------------	-------	-------	-------	-------	-------	--------	-------	-------	-----------

^a For crystalline solutes the tabulated value is $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,gas\,phase}^{sat}}\right)$.

^b For crystalline solutes the tabulated value is $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,water}^{sat}}\right)$.

The experimental data for the dissolved liquid solutes was given in the published literature as the infinite dilution activity coefficient, γ_{solute}^{∞} . Infinite dilution activity coefficients are converted to the logarithm of the gas-to-organic solvent partition coefficients through standard thermodynamic relationships:

$$\log K = \log\left(\frac{RT}{\gamma_{solute}^{\infty} p_{solute}^o V_{solvent}}\right) \quad (4)$$

where R is the universal gas constant and p_{solute}^o is the saturated vapor pressure of a pure solute at the solution temperature, which in this study is $T = 298.15$ K. As an informational note infinite dilution activity coefficient of diethyl carbonate dissolved in itself is unity.

The published solubility data for the gaseous solutes was given either in terms of gas-to-diethyl carbonate partition coefficient, K , or in terms of the Henrys law constant, K_{Henry} . Henrys law constants are converted to $\log K$ values through Eqn. 5.

$$\log K = \log\left(\frac{RT}{K_{\text{Henry}} V_{solvent}}\right) \quad (5)$$

The $\log K$ values for the liquid and gaseous solutes are converted to $\log P$ values by:

$$\log P = \log K - \log K_{\text{water}} \quad (6)$$

where K_{water} is the solute's gas-to-water partition coefficient, which is available for the different solutes considered in the current study [82-86]. We have also tabulated the numerical values of $\log K$ and $\log P$ values for the liquid and gases in the eighth and ninth columns of Table 3. As the

column headings indicate the tabulated numerical values for the crystalline organic solutes pertain

to the two respective solubility ratios, $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,gas\,phase}^{sat}}\right)$ and $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,water}^{sat}}\right)$.

Our experimental measurements, combined with our search of the published literature, have experimental partition coefficients and solubility ratios for 50 different organic and inorganic compounds. The solutes are chemically diverse, they cover a wide range of solute polarities and hydrogen-bonding capability, and include several fairly strong H-bond donors (acetylsalicylic acid, **A** = 0.710; 4-chloro-3-nitrobenzoic acid, **A** = 0.700) and several fairly strong H-bond acceptors (3,4,5-trimethoxybenzoic acid, **B** = 0.850; benzoin, **B** = 0.841). Regression analysis of the experimental values yielded the following two Abraham model expressions:

$$\log P \text{ and } \log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,water}^{sat}}\right) = 0.133(0.044) + 0.135(0.075) \mathbf{E} - 0.309(0.121) \mathbf{S} - 1.532(0.103) \mathbf{A} - 4.816(0.183) \mathbf{B} + 4.398(0.069) \mathbf{V} \quad (7)$$

(with N = 49, SD = 0.134, R^2 = 0.994, F = 1333)

$$\log K \text{ and } \log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,gas\,phase}^{sat}}\right) = 0.092(0.027) - 0.598(0.057) \mathbf{E} + 1.527(0.078) \mathbf{S} + 1.942(0.088) \mathbf{A} + 0.948(0.014) \mathbf{L} \quad (8)$$

(with N = 50, SD = 0.117, R^2 = 0.999, F = 13898)

The correlations were obtained using the IBM SPSS Statistical 22 commercial software. The statistical information pertaining to each correlation is given below the respective correlation, where N refers to the number of experimental data points used in determining the equation coefficients, SD is the standard deviation, R^2 corresponds to the squared correlation coefficient, and F represents the Fisher F-statistic. Standard errors in the equation coefficients (at 95 % level) are given in parenthesis after the coefficient itself. As an informational note the $b_k \cdot \mathbf{B}$ was

removed from Eqn. 8 because diethyl carbonate lacks an acidic hydrogen and therefore cannot act as a hydrogen-bond donor. Removal of the $b_k \cdot \mathbf{B}$ led to an insignificant increase in the standard deviation, from $SD = 0.115$ to $SD = 0.117$. The $b_p \cdot \mathbf{B}$ term was retained in Eqn. 7, however, as the expression pertains to solute transfer into diethyl carbonate from water. Here the b_p equation coefficient represents the difference in the acidity of diethyl carbonate and water, which does possess two acidic hydrogens capable of engagement in hydrogen-bond formation.

As noted in the introduction a major advantage that the Abraham model offers over the solution models that other research groups are using to mathematically describe experimental solubility data is that the derived correlations can be used to predict the solubility and/or infinite dilution activity coefficient of many additional organic, organometallic and inorganic solutes dissolved in the given solvent media. An indication of Eqns. 7 and 8 ability to estimate partition coefficients and solubility ratios can be obtained by examining each correlation's ability to describe the 50 experimental values given in Table 3. The small standard deviations, $SD = 0.134$ log units and $SD = 0.117$ log units, and near unity squared correlation coefficients, $R^2 = 0.994$ and $R^2 = 0.999$, indicate that Eqs. 7 and 8 provide reasonably accurate mathematical descriptions of the observed experimental data used in both regression analyses. Figures 1 and 2 provide a graphical comparison of the experimental data versus back-calculated values based on Eqs. 7 and

8, respectively. As an informational note the experimental $\log K$ and $\log(\frac{c_{solute,org\,solv}^{sat}}{c_{solute,gas\,phase}^{sat}})$ data span

a range of just under 12 log units, from $\log K = -1.504$ for helium to $\log(\frac{c_{solute,org\,solv}^{sat}}{c_{solute,gas\,phase}^{sat}}) = 11.351$

for benzoin. A slightly smaller range of approximately 7.5 log units is covered by the experimental

$\log P$ and $\log(\frac{c_{solute,org\,solv}^{sat}}{c_{solute,water}^{sat}})$ data.

Diethyl carbonate is the second carbonate solvent that we have studied. Previously we determined Abraham model correlations for propylene carbonate [39]:

$$\log P \text{ and } \log\left(\frac{C_{\text{solute,org solv}}^{\text{sat}}}{C_{\text{solute,water}}^{\text{sat}}}\right) = 0.004 + 0.168 \mathbf{E} + 0.504 \mathbf{S} - 1.283 \mathbf{A} - 4.407 \mathbf{B} + 3.421 \mathbf{V} \quad (9)$$

$$\log K \text{ and } \log\left(\frac{C_{\text{solute,org solv}}^{\text{sat}}}{C_{\text{solute,gas phase}}^{\text{sat}}}\right) = -0.356 - 0.413 \mathbf{E} + 2.587 \mathbf{S} + 2.207 \mathbf{A} + 0.455 \mathbf{B} + 0.717 \mathbf{L} \quad (10)$$

based on experimental solubility and activity coefficient data for 69 compounds. The equation coefficients of the two carbonate solvents are somewhat different. Propylene carbonate exhibits greater dipolarity/polarizability, and is a better H-bond acceptor and H-bond donor as evidenced by the larger s_k , b_k and a_k equation coefficients.

4. Conclusion

Mathematical expressions based on the Abraham solvation parameter model have been shown to provide reasonably accurate correlations of the solute transfer properties of a wide range of organic and inorganic compounds into diethyl carbonate from both water and from the gas phase. Experimental properties that were described by the derived Abraham model expressions include the logarithm of the water-to-diethyl carbonate transfer coefficient, the logarithm of the gas-to-diethyl carbonate transfer coefficient, and the logarithms of two molar solubility ratios given by $\log\left(\frac{C_{\text{solute,org solv}}^{\text{sat}}}{C_{\text{solute,water}}^{\text{sat}}}\right)$, $\log\left(\frac{C_{\text{solute,org solv}}^{\text{sat}}}{C_{\text{solute,gas phase}}^{\text{sat}}}\right)$. The derived correlations described the fore-mentioned experimental solute transfer properties to within 0.13 log units (or less). Based on our more than 20 years of experience in using the Abraham model we believe that our derived Eqns. 7 and 8 will provide reasonably accurate predictions of the $\log P$, $\log K$, $\log\left(\frac{C_{\text{solute,org solv}}^{\text{sat}}}{C_{\text{solute,water}}^{\text{sat}}}\right)$ and

$\log \left(\frac{C_{\text{solute,org solv}}^{\text{sat}}}{C_{\text{solute,gas phase}}^{\text{sat}}} \right)$ values for additional organic and inorganic compounds dissolved in diethyl carbonate at 298.15 K, provided of course that the solute descriptors of the compounds fall within the range of numerical values used in obtaining these correlations. Partition coefficients predicted by the Abraham model can be transformed into $\gamma_{\text{solute}}^{\infty}$ values through Eqs. 4 and 6. In closing we note that predicted $\log P$ and $\log K$ values at 298.15 K can be extrapolated to slightly lower and higher temperatures using our published enthalpy of solvation correlations, ΔH_{solv} , [87] based on experimental ΔH_{solv} data for 80 solutes dissolved in diethyl carbonate. Our published Abraham model correlations provide the manufacturing sector with important tools to aid in the solvent selection process.

Acknowledgements

The authors acknowledge the National Science Foundation's Research Experiences for Undergraduates (REU) program (CHE-1757946) for supporting Estefania Garcia's participation in this research. Kelly Liu, Jingyi Dai, Shrika Eddula and Carina Jiang thank the University of North Texas Texas Academy of Mathematics and Science (TAMS) program for providing a summer research scholarship to each student.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- [1] Fan F, Yuan H, Wu S, Zhai J, Liu F, Zhang L, Liu X, Lu J. Solubility measurement and molecular simulation of 1,1'-binaphthyl-2,2'-diyl hydrogenphosphate racemate and enantiomers in organic solvents with and without reverse micelles. *J Chem Thermodyn.* 2020; 140: 105905/1-105905/16.
- [2] Li Z, Wang Z, Li M, Gao Z, Wang B. Measurement and correlation of solubility of methyl gallate in nine pure and ethanol + n-propanol mixed solvents at temperatures within 293.15-333.15 K. *J Mol Liq.* 2019; 293: 111531/1-111531/9.
- [3] Wu K, Li Y. Solubility and solution thermodynamics of p-toluenesulfonamide in 16 solvents from $T = 273.15$ to 324.75 K. *J Mol Liq.* 2019; 293: 111577/1-111577/10.
- [4] Yang Z, Shao D, Zhou G. Analysis of solubility parameters of fenbendazole in pure and mixed solvents and evaluation of thermodynamic model. *J Chem Thermodyn.* 2020; 140: 105876/1-105876/7.
- [5] He Z, Zhang J, Gao X, Tang T, Yin X, Zhao J, Li R, Han D. Solubility determination, correlation, and solute-solvent molecular interactions of 5-aminotetrazole in various pure solvents. *J Chem Eng Data.* 2019; Ahead of Print. DOI: 10.1021/acs.jced.9b00385.
- [6] Zou Z, Sun X, Yu Y, Fang X, Xu L. Determination and correlation of solubility and solution thermodynamics of musk ketone in twelve pure solvents. *J Mol Liq.* 2019; 293: 111437/1-111437/10.
- [7] Zhang P, Wan Y, Zhang C, Zhao R, Sha J, Li Y, Li T, Ren B. Solubility and mixing thermodynamic properties of levamisole hydrochloride in twelve pure solvents at various temperatures. *J Chem Thermodyn.* 2019; 139: 105882/1-105882/10.

[8] Huang C, Yun F, Yongmei X, Jun W, Ye Y, Hang L, Yixin L. Correlation of solubility and solution thermodynamics of DL-malic acid in organic solvents at different temperatures. Russ J Phys Chem A. 2018; 92: 2710-2717.

[9] Gong T, Han D, Chen Y, Wang S, Tang W. Solubility and data correlation of isoniazid in different pure and binary mixed solvent systems from 283.15 K to 323.15 K. J Chem Eng Data. 2018; 63: 4767-4778.

[10] Li S, Liu Y, Yin F, Ye X. Solubility measurement and thermodynamic properties of levetiracetam in pure and mixed solvents. J Chem Eng Data. 2018; 63: 4669-4681.

[11] Shen J, Liang X, Lei H. Measurements, thermodynamic modeling, and a hydrogen bonding study on the solubilities of metoprolol succinate in organic solvents. Molecules. 2018; 23: 1-11.

[12] Fan B, Li Q, Lu M, Guo F, Cheng C, Gao Ying. Measurement and correlation of solubility of 3-hydroxy-2-naphthoic acid in ten pure and binary mixed organic solvents from $T = (293.15 \text{ to } 333.15 \text{ K})$. J Chem Eng Data. 2018; 63: 4459-4467.

[13] Baluja S, Lava D, Hirpara A, Bhesaniya K. Thermodynamics of alloxan solubility in various solvents at different temperatures. Eur Chem Bull. 2017; 6: 206-210.

[14] Ahmad A, Raish M, Alkhafry KM, Alsarra IA, Khan A, Ahad A, Jan BL, Shakeel F. Solubility, solubility parameters and solution thermodynamics of thymoquinone in different mono solvents. J Mol Liq. 2018; 272: 912-918.

[15] Asadzadeh B, Zhong J, Yan W. Solid-liquid equilibrium of rebaudioside A in pure and binary mixed solvents at $T = (288.15 \text{ to } 328.15) \text{ K}$. J Chem Eng Data. 2018; 63: 4269-4276.

[16] Qian E, Gupta A, Neal R, Lee G, Che M, Wang L, Yue D, Wang S, Liu K, Zhang A, Acree WE Jr, Abraham MH. Abraham model correlations for describing solute transfer into 4-methyl-2-pentanol from both water and the gas phase. *J Mol Liq.* 2019; 278: 335-341.

[17] Hart E, Jodray M, Rodriguez K, Barrera M, Qian E, Zha O, Tong XY, Woods D, Liu Y, Lee G, Acree WE Jr, Abraham MH. Abraham model correlations for describing the solubilizing character of 3-methoxy-1-butanol and 1-tert-butoxy-2-propanol solvents. *Phys Chem Liq.* 2019; 57: 163-173.

[18] Hart E, Klein A, Barrera M, Jodray M, Rodriguez K, Acree WE Jr, Abraham MH. Development of Abraham model correlations for describing the transfer of molecular solutes into propanenitrile and butanenitrile from water and from the gas phase. *Phys Chem Liq.* 2018; 56: 821-833.

[19] Tong X, Woods D, Acree WE Jr, Abraham MH. Updated Abraham model correlations for correlating solute transfer into dry butanone and dry cyclohexanone solvents. *Phys Chem Liq.* 2018; 56: 571-583.

[20] Liu K, Wang S, Hart E, Klein A, Stephens TW, Bates B, Calzada J, Garcia M, Abban-Manesah J, Acree WE Jr, Abraham MH. Development of Abraham model correlations for solute transfer into 2-ethyl-1-hexanol from both water and the gas phase based on measured solubility ratios. *Phys Chem Liq.* 2019; Ahead of Print. DOI: 10.1080/00319104.2018.1564306.

[21] Sedov IA, Salikov TM, Wadawadigi A, Zha O, Qian E, Acree WE Jr, Abraham, M H. Abraham model correlations for describing the thermodynamic properties of solute transfer into pentyl acetate based on headspace chromatographic and solubility measurements. *J Chem Thermodyn.* 2018; 124: 133-140.

[22] Sedov IA, Magsumov TI, Hart E, Higgins E, Grover D, Zettl H, Zad M, Acree WE Jr, Abraham MH. Abraham model expressions for describing water-to-diethylene glycol and gas-to-diethylene glycol solute transfer processes at 298.15 K. *J Solut Chem.* 2017; 46: 331-351.

[23] Qian E, Gupta A, Neal R, Lee G, Che M, Wang L, Yue D, Fischer R, Jodray M, Connolly E, Wang S, Liu K, Zhang AW, Dai J, Zhu S, Acree WE Jr, Abraham MH. Development of Abraham model correlations for describing solute transfer into 2-methyl-1-butanol from both water and the gas phase from experimental solubility data for crystalline organic compounds. *Phys Chem Liq.* 2019; Ahead of Print. doi: 10.1080/00319104.2019.1625050.

[24] Fischer R, Jodray M, Qian E, Wang L, Lee G, Yue D, Che M, Liu Y, Acree WE Jr, Abraham MH. Abraham model correlations for solute transfer into benzyl alcohol from both water and the gas phase. *Phys Chem Liq.* 2018; Ahead of Print. doi: 10.1080/00319104.2018.1550778.

[25] Abraham MH. Scales of solute hydrogen-bonding: their construction and application to physicochemical and biochemical processes. *Chem Soc Reviews* 1993; 22: 73-83.

[26] Abraham MH, Ibrahim A, Zissimos AM. Determination of sets of solute descriptors from chromatographic measurements. *J Chromatog A.* 2004; 1037: 29-47.

[27] Abraham MH, Smith RE, Luchefeld R, Boorem AJ, Luo R, Acree WE Jr. Prediction of solubility of drugs and other compounds in organic solvents. *J Pharm Sci.* 2010; 99: 1500-1515.

[28] Abraham MH, McGowan JC. The use of characteristic volumes to measure cavity terms in reversed phase liquid chromatography. *Chromatographia.* 1987; 23: 243-246.

[29] Endo S, Goss K-U. Applications of polyparameter linear free energy relationships in environmental chemistry. *Environ Sci Technol*. 2014; 48: 12477-12491.

[30] Poole CF, Ariyasena TC, Lenca N. Estimation of the environmental properties of compounds from chromatographic measurements and the solvation parameter model. *J Chromatogr A*. 2013; 1317: 85-104.

[31] Jalan A, Ashcraft RW, West RH, Green WH. Predicting solvation energies for kinetic modeling. *Ann Rep Prog Chem, Sect C: Phys Chem*. 2010; 106: 211-258.

[32] Ulrich N, Endo S, Brown TN, Watanabe N, Bronner G, Abraham MH, Goss K-U. UFZ-LSER database v 3.2.1 [Internet], Leipzig, Germany, Helmholtz Centre for Environmental Research-UFZ. 2017 [accessed on 27.08.2019]. Available from <http://www.ufz.de/lserd>.

[33] Abraham MH, Acree WE Jr. Descriptors for ferrocene and some substituted ferrocenes. *J Mol Liq*. 2017; 232: 325-331.

[34] Sedov IA, Salikov TM, Qian E, Wadawadigi A, Zha O, Acree WE Jr, Abraham MH. Abraham model correlations for solute transfer into 2-methyl-2-butanol based on measured activity coefficient and solubility data at 298.15 K. *J Mol Liq*. 2019; 293: 111454/1-111454/10.

[35] Abraham MH, Acree WE Jr. Gas-solvent and water-solvent partition of trans-stilbene at 298 K. *J Mol Liq*. 2017; 238: 58-61.

[36] Abraham MH, Acree WE Jr. Equations for the partition of neutral molecules, ions and ionic species from water to water-methanol mixtures. *J Solut Chem*. 2016; 45: 861-874.

[37] Abraham MH, Acree WE Jr. Equations for the partition of neutral molecules, ions and ionic species from water to water-ethanol mixtures. *J Solut Chem*. 2012; 41: 730-740.

[38] Abraham MH, Acree WE Jr. Partition coefficients and solubilities of compounds in the water-ethanol solvent system. *J Solut Chem.* 2011; 40: 1279-1290.

[39] Abraham MH, Acree WE Jr. The transfer of neutral molecules, ions and ionic species from water to ethylene glycol and to propylene carbonate; descriptors for pyridinium cations. *New J Chem.* 2010; 34: 2298-2305.

[40] Logan ER, Tonita EM, Gering KL, Li J, Ma X, Beaulieu LY, Dahn JR. A study of the physical properties of li-ion battery electrolytes containing esters. *J Electrochem Soc.* 2018; 165: A21-A30.

[41] Zhang Y, Hu X-G, Zhang C-F. Effects of organic solvents on performance of Li-ion battery. *Dianchi.* 2004; 34: 30-32.

[42] Bilgin M, Birman I. Separation of propionic acid by diethyl carbonate or diethyl malonate or diethyl fumarate and the synergistic effect of phosphorus compounds and amines. *Fluid Phase Equilib.* 2010; 292: 13-19.

[43] Uslu H, Kirbaslar SI. Extraction of aqueous of malic acid by trioctylamine extractant in various diluents. *Fluid Phase Equilib.* 2010; 287: 134-140.

[44] Uslu H, Kirbaslar SI. Purification of L-malic acid from aqueous solution by a method of reactive extraction. *J Chem Eng Data.* 2009; 54: 2819-2826.

[45] Rasool MA, Pescarmona PP, Vankelecom IFJ. Applicability of organic carbonates as green solvents for membrane preparation. *ACS Sustainable Chem Eng.* 2019; 7: 13774-13785.

[46] de la Iglesia O, Mainar AM, Pardo JI, Urieta JS. Solubilities of nonpolar gases in triethylene glycol dimethyl ether, tetraethylene glycol dimethyl ether, dimethyl carbonate, and diethyl carbonate at 298.15 K and 101.33 kPa partial pressure of gas. *J Chem Eng Data.* 2003; 48: 657-661.

[47] Read J, Mutolo K, Ervin M, Behl W, Wolfenstine J, Driedger A, Foster D. Oxygen transport properties of organic electrolytes and performance of lithium/oxygen battery. *J Electrochem Soc.* 2003; 150: A1351-A1356.

[48] Zhao Z, Xing X, Tang Z, Zhao Y, Fei W, Liang X, He Z, Zhang S, Guo D. Solubility of CO₂ and H₂S in carbonates solvent: Experiment and quantum chemistry calculation. *Int J Greenhouse Gas Control.* 2017; 59: 123-135.

[49] Lopez ER, Mainar AM, Urieta JS, Fernandez J. Solubility of HFC134a (1,1,1,2-tetrafluoroethane) in two dialkyl carbonates. *J Chem Eng Data.* 2009; 54: 2609-2615.

[50] Li, Wenchao; Xuan, Aiguo; Wu, Yuanxin; Ai, Jun; Liang, Lei. Measurement and correlation of solid-liquid equilibrium of diphenyl carbonate in ethanol-diethyl carbonate. *Huagong Xuebao.* 2009; 60: 1357-1361.

[51] Cocero MJ, Gonzalez JA, Garcia I, Cobos JC, Mato F, Kehiaian HV. Excess Gibbs energy. Diethyl carbonate-hexane system. *Int DATA Ser, Sel Data Mix, Ser A.* 1991; 19: 132.

[52] Cocero, M. J.; Gonzalez, J. A.; Garcia, I.; Cobos, J. C.; Mato, F.; Kehiaian, H. V. Excess Gibbs energy. Diethyl carbonate-octane system. *Int DATA Ser, Sel Data Mix, Ser A.* 1991; 19: 135.

[53] Cocero MJ, Gonzalez JA, Garcia I, Cobos JC, Mato F, Kehiaian HV. Excess Gibbs energy. Diethyl carbonate-dodecane system. *Int DATA Ser, Sel Data Mix, Ser A.* 1991; 19: 138.

[54] Cocero MJ, Gonzalez JA, Garcia I, Cobos JC, Mato F, Kehiaian HV. Excess Gibbs energy. Diethyl carbonate-cyclohexane system. *Int DATA Ser, Sel Data Mix, Ser A.* 1991; 19: 123.

[55] Cocero MJ, Gonzalez JA, Garcia I, Cobos JC, Mato F, Kehiaian HV. Excess Gibbs energy. Diethyl carbonate-benzene system. *Int DATA Ser, Sel Data Mix, Ser A.* 1991; 19: 126.

[56] Cocero MJ, Gonzalez JA, Garcia I, Cobos JC, Mato F, Kehiaian HV. Excess Gibbs energy. Tetrachloromethane-diethyl carbonate system. Int DATA Ser, Sel Data Mix, Ser A. 1991; 19: 129.

[57] Francesconi R. Excess Gibbs energy. Diethyl carbonate - tert-butyl methyl ether system. Int DATA Ser, Sel Data Mix, Ser A. 1998; 26: 136.

[58] Francesconi R, Ottani S, Comelli F. Excess Gibbs energy. Dimethyl carbonate-diethyl carbonate system. Int DATA Ser, Sel Data Mix, Ser A. 1997; 25: 211.

[59] Comelli F, Francesconi R, Ottani S. Isothermal vapor-liquid equilibria of dimethyl carbonate + diethyl carbonate in the range (313.15 to 353.15) K. J Chem Eng Data. 1996; 41: 534-536.

[60] Fritz JS, Lisicki NM. Titration of acids in nonaqueous solvents. Anal Chem. 1951; 23: 589-591.

[61] Dabadge N, Tian A, Willis B, Wilson A, Acree WE Jr. Solubility of anthracene in binary ethylbenzene + alcohol solvent mixtures at 298.15 K. Phys Chem Liq. 2013; 51: 715-720.

[62] Monarrez CI, Woo JH, Taylor PG, Tran AM, Acree WE Jr. Solubility in binary solvent mixtures: pyrene dissolved in alcohol + acetonitrile mixtures at 299.2 K. J Chem Eng Data. 2003; 48: 736-738.

[63] Stovall DM, Givens C, Keown S, Hoover KR, Barnes R, Harris C, Lozano J, Nguyen M, Rodriguez E, Acree WE Jr, Abraham MH. Solubility of crystalline nonelectrolyte solutes in organic solvents: mathematical correlation of 4-chloro-3-nitrobenzoic acid and 2-chloro-5-nitrobenzoic acid solubilities with the Abraham solvation parameter model. Phys Chem Liq. 2005; 43: 351-360.

[64] Hoover KR, Stovall DM, Pustejovsky E, Coaxum R, Pop K, Acree WE Jr, Abraham MH. Solubility of crystalline nonelectrolyte solutes in organic solvents - mathematical correlation of 2-methoxybenzoic acid and 4-methoxybenzoic acid solubilities with the Abraham solvation parameter model. *Can J Chem*. 2004; 82: 1353-1360.

[65] Wilson A, Tian A, Chou V, Quay AN, Acree WE Jr, Abraham MH. Experimental and predicted solubilities of 3,4-dichlorobenzoic acid in select organic solvents and in binary aqueous-ethanol mixtures. *Phys Chem Liq*. 2012; 50: 324-335.

[66] Charlton AK, Daniels CR, Acree WE Jr, Abraham MH. Solubility of crystalline nonelectrolyte solutes in organic solvents: mathematical correlation of acetylsalicylic acid solubilities with the Abraham general solvation model. *J Solut Chem*. 2003; 32: 1087-1102.

[67] Flanagan KB, Hoover KR, Garza O, Hizon A, Soto T, Villegas N, Acree WE Jr, Abraham MH. Mathematical correlation of 1-chloroanthraquinone solubilities in organic solvents with the Abraham solvation parameter model. *Phys Chem Liq*. 2006; 44: 377-386.

[68] Daniels CR, Charlton AK, Wold RM, Acree WE Jr, Abraham MH. Thermochemical behavior of dissolved carboxylic acid solutes: Solubilities of 3-methylbenzoic acid and 4-chlorobenzoic acid in organic solvents. *Can J Chem* 2003; 81: 1492-1501.

[69] Hoover KR, Coaxum R, Pustejovsky E, Stovall DM, Acree WE Jr, Abraham MH. Thermochemical behavior of dissolved carboxylic acid solutes: part 4 - mathematical correlation of 4-nitrobenzoic acid solubilities with the Abraham solvation parameter model. *Phys Chem Liq*. 2004; 42: 339-347.

[70] Coaxum R, Hoover KR, Pustejovsky E, Stovall DM, Acree WE Jr, Abraham MH. Thermochemical behavior of dissolved carboxylic acid solutes: Part 3 - Mathematical

correlation of 2-methylbenzoic acid solubilities with the Abraham solvation parameter model. *Phys Chem Liq.* 2004; 42: 313-322.

[71] Hoover KR, Acree WE Jr, Abraham MH. Mathematical correlation of phenothiazine solubilities in organic solvents with the Abraham solvation parameter model. *Phys Chem Liq.* 2006; 44: 367-376.

[72] Acree WE Jr, Abraham MH. Solubility of crystalline nonelectrolyte solutes in organic solvents: mathematical correlation of benzil solubilities with the Abraham general solvation model. *J Solution Chem.* 2002; 31: 293-303.

[73] Fletcher KA, Pandey S, McHale MER, Acree WE Jr. Solubility of benzil in organic nonelectrolyte solvents. Comparison of observed versus predicted values based upon mobile theory. *Phys Chem Liq.* 1996; 33: 181-190.

[74] Stephens TW, Loera M, Calderas M, Diaz R, Montney N, Acree WE Jr, Abraham MH. Determination of Abraham model solute descriptors for benzoin based on measured solubility ratios. *Phys Chem Liq.* 2012; 50: 254-265.

[75] Blake-Taylor BH, Deleon VH, Acree WE Jr, Abraham MH. Mathematical correlation of salicylamide solubilities in organic solvents with the Abraham solvation parameter model. *Phys Chem Liq.* 2007 45: 389-398.

[76] Wang S, Liu K, Zhang A, Dai J, Gupta A, Zhu S, Eddula S, Jiang C, Acree WE Jr, Abraham MH. Solubility of 4-methyl-3-nitrobenzoic acid in organic mono-solvents: calculation of Abraham model solute descriptors. *Phys Chem Liq.* 2019; in press, <https://doi.org/10.1080/00319104.2019.1660982>.

[77] Hart E, Lee G, Qian E, Jodray M, Barrera M, Fischer R, Che M, Liu Y, Zha O, Woods D, Acree WE Jr, Abraham MH. Determination of Abraham model solute descriptors for 4-

tert-butylbenzoic acid from experimental solubility data in organic mono-solvents. *Phys Chem Liq.* 2019; 57: 445-452.

[78] Brumfield M, Wadawadigi A, Kuprasertkul N, Mehta S, Stephens TW, Barrera M, De La Rosa J, Kennemer L, Meza J, Acree WE Jr, Abraham MH. Determination of Abraham model solute descriptors for three dichloronitrobenzenes from measured solubilities in organic solvents. *Phys Chem Liq.* 2015; 53: 163-173.

[79] Hart E, Klein A, Zha O, Wadawadigi A, Qian E, Dunn S, Herron J, Kankolongo K, Ryan S, Acree WE Jr.; Abraham MH, Determination of Abraham model solute descriptors for monomeric 3,4,5-trimethoxybenzoic acid from experimental solubility data in organic solvents measured at 298.2 K. *Phys Chem Liq.* 2018; 56: 381-390.

[80] Bowen KR, Stephens TW, Lu H, Satish K, Shan D, Acree WE Jr, Abraham MH. Experimental and predicted solubilities of 3,4-dimethoxybenzoic acid in select organic solvents of varying polarity and hydrogen-bonding character. *Eur Chem Bull* 2013; 2: 577-583.

[81] De Fina KM, Sharp TL, Acree WE Jr. Solubility of biphenyl in organic nonelectrolyte solvents. Comparison of observed versus predicted values based upon mobile order theory. *Can J Chem.* 1999; 77: 1589-1593.

[82] Abraham MH, Zissimos AM, Acree WE Jr. Partition of solutes from the gas phase and from water to wet and dry di-n-butyl ether: a linear free energy relationship analysis. *Phys Chem Chem Phys.* 2001; 3: 3732-3736.

[83] Sprunger LM, Proctor A, Acree WE Jr, Abraham MH, Benjelloun-Dakhama N. Correlation and prediction of partition coefficient between the gas phase and water, and

the solvents dry methyl acetate, dry and wet ethyl acetate, and dry and wet butyl acetate. Fluid Phase Equilib. 2008; 270: 30-44.

- [84] Abraham MH, Acree WE Jr, Leo AJ, Hoekman D. The partition of compounds from water and from air into wet and dry ketones. New J Chem. 2009; 33: 568-573.
- [85] Abraham MH, Acree WE Jr, Leo AJ, Hoekman D. Partition of compounds from water and from air into the wet and dry monohalobenzenes. New J Chem. 2009; 33: 1685-1692.
- [86] Abraham MH, Acree WE Jr, Cometto-Muniz JE. Partition of compounds from water and from air into amides. New J Chem. 2009; 33: 2034-2043.
- [87] Hart E, Grover D, Zettl H, Acree WE Jr, Abraham MH. Abraham model enthalpy of solvation correlations for solutes dissolved in dimethyl carbonate and diethyl carbonate. Phys Chem Liq. 2015; 53: 732-747.

Figure 1




Figure 1. Comparison between the experimental $\log P$ and $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,water}^{sat}}\right)$ data and back-calculated values based on Eq. 7.

Figure 2




Figure 2. Comparison between the experimental $\log K$ and $\log\left(\frac{C_{solute,org\,solv}^{sat}}{C_{solute,gas\,phase}^{sat}}\right)$ data and back-calculated values based on Eq. 8.