

Note on Lagrangian-Eulerian Methods for Uniqueness in Hydrodynamic Systems

Peter Constantin ^{*1} and Joonhyun La ^{†1}

¹Department of mathematics, Princeton University

November 30, 2018

Abstract

We discuss the Lagrangian-Eulerian framework for hydrodynamic models and provide a proof of Lipschitz dependence of solutions on initial data in path space. The paper presents a corrected version of the result in [1].

MSC Classification: 35Q30, 35Q35.

1 Introduction

Many hydrodynamical systems consist of evolution equations for fluid velocities forced by external stresses, coupled to evolution equations for the external stresses. In the simplest cases, the Eulerian velocity u can be recovered from the stresses σ via a linear operator

$$u = \mathbb{U}(\sigma) \tag{1}$$

and the stress matrix σ obeys a transport and stretching equation of the form

$$\partial_t \sigma + u \cdot \nabla \sigma = F(\nabla u, \sigma),$$

where F is a nonlinear coupling depending on the model. The Eulerian velocity gradient is obtained in terms of the operator

$$\nabla_x u = \mathbb{G}(\sigma), \tag{2}$$

and, in many cases, \mathbb{G} is bounded in Hölder spaces of low regularity. Then, passing to Lagrangian variables,

$$\tau = \sigma \circ X$$

where X is the particle path transformation $X(\cdot, t) : \mathbb{R}^d \rightarrow \mathbb{R}^d$, a volume preserving diffeomorphism, the system becomes

$$\begin{cases} \partial_t X = \mathcal{U}(X, \tau), \\ \partial_t \tau = \mathcal{T}(X, \tau). \end{cases} \tag{3}$$

with

$$\begin{aligned} \mathcal{U}(X, \tau) &= \mathbb{U}(\tau \circ X^{-1}) \circ X, \\ \mathcal{T}(X, \tau) &= F(\mathbb{G}(\tau \circ X^{-1}) \circ X, \tau). \end{aligned} \tag{4}$$

In particular, τ solves an ODE

$$\frac{d}{dt} \tau = F(g, \tau) \tag{5}$$

where $g = \nabla_x u \circ X$ is of the same order of magnitude as τ in appropriate spaces, and so the size of τ is readily estimated from the information provided by the ODE model, analysis of \mathbb{G} and of the

^{*}const@math.princeton.edu

[†]joonhyun@math.princeton.edu

operation of composition with X . The main additional observation that leads to Lipschitz dependence in path space is that derivatives with respect to parameters of expressions of the type encountered in the Lagrangian evolution (4),

$$\mathbb{U}(\tau \circ X^{-1}) \circ X, \quad \mathbb{G}(\tau \circ X^{-1}) \circ X,$$

introduce commutators, and these are well behaved in spaces of relatively low regularity. The Lagrangian-Eulerian method of [2] formalized these considerations leading to uniqueness and Lipschitz dependence on initial data in path space, with application to several examples including incompressible 2D and 3D Euler equations, the surface quasi-geostrophic equation (SQG), the incompressible porous medium equation, the incompressible Boussinesq system, and the Oldroyd-B system coupled with the steady Stokes system. In all these examples the operators \mathbb{U} and \mathbb{G} are time-independent.

The paper [1] considered time-dependent cases. When the operators \mathbb{U} and \mathbb{G} are time-dependent, in contrast to the time-independent cases studied in [2], \mathbb{G} is not necessarily bounded in $L^\infty(0, T; C^\alpha)$. This was addressed in [1] by using a Hölder continuity $\sigma \in C^\beta(0, T; C^\alpha)$. While this treated the Eulerian issue, it was tacitly used but never explicitly stated in [1] that this kind of Hölder continuity is transferred to σ from τ by composition with a smooth time-depending diffeomorphism close to the identity. This is false. In fact, we can easily give examples of C^α functions τ which are time-independent (hence analytic in time with values in C^α) and diffeomorphisms $X(t)(a) = a + vt$ with constant v , such that $\sigma = \tau \circ X^{-1}$ is not continuous in C^α as a function of time. In this paper we present a correct version of the results in [1]. Instead of relying on the time regularity of τ alone, we also use the fact that \mathbb{G} is composed from a time-independent bounded operator and an operator whose kernel is smooth and rapidly decaying in space. Then the time singularity is resolved by using the Lipschitz dependence in L^1 of Schwartz functions composed with smoothly varying diffeomorphisms near the identity.

A typical example of the systems we can treat is the Oldroyd-B system coupled with Navier-Stokes equations:

$$\left\{ \begin{array}{l} \partial_t u - \nu \Delta u = \mathbb{H}(\operatorname{div}(\sigma - u \otimes u)), \\ \nabla \cdot u = 0, \\ \partial_t \sigma + u \cdot \nabla \sigma = (\nabla u)\sigma + \sigma(\nabla u)^T - 2k\sigma + 2\rho K((\nabla u) + (\nabla u)^T), \\ u(x, 0) = u_0(x), \sigma(x, 0) = \sigma_0(x). \end{array} \right. \quad (6)$$

Here $(x, t) \in \mathbb{R}^d \times [0, T]$. The Leray-Hodge projector $\mathbb{H} = \mathbb{I} + R \otimes R$ is given in terms of the Riesz transforms $R = (R_1, \dots, R_d)$, and $\nu, \rho K, k$ are fixed positive constants. This system is viscoelastic, and the behavior of the solution depends on the history of its deformation.

The non-resistive MHD system

$$\left\{ \begin{array}{l} \partial_t u - \nu \Delta u = \mathbb{H}(\operatorname{div}(b \otimes b - u \otimes u)), \\ \nabla \cdot u = 0, \\ \nabla \cdot b = 0, \\ \partial_t b + u \cdot \nabla b = (\nabla u)b, \\ u(x, 0) = u_0(x), b(x, 0) = b_0(x). \end{array} \right. \quad (7)$$

can also be treated by this method. The systems (6) and (7) have been studied extensively, and a review of the literature is beyond the scope of this paper.

2 The Lagrangian-Eulerian formulation

We show calculations for (6) in order to be explicit, and because the calculations for (7) are entirely similar. The solution map for $u(x, t)$ of (6) is

$$u(x, t) = \mathbb{L}_\nu(u_0)(x, t) + \int_0^t g_{\nu(t-s)} * (\mathbb{H}(\operatorname{div}(\sigma - u \otimes u)))(x, s) ds. \quad (8)$$

where

$$\mathbb{L}_\nu(u_0)(x, t) = g_{\nu t} * u_0(x) = \int_{\mathbb{R}^d} \frac{1}{(4\pi\nu t)^{\frac{d}{2}}} e^{-\frac{|x-y|^2}{4\nu t}} u_0(y) dy. \quad (9)$$

Throughout the paper we use

$$g_{\nu t}(x) = \frac{1}{(4\pi\nu t)^{\frac{d}{2}}} e^{-\frac{|x|^2}{4\nu t}}.$$

The velocity gradient satisfies

$$(\nabla u)(x, t) = \mathbb{L}_\nu(\nabla u_0)(x, t) + \int_0^t (g_{\nu(t-s)} * (\mathbb{H}\nabla \operatorname{div} (\sigma - u \otimes u)))(x, s) ds. \quad (10)$$

We denote the Eulerian velocity and gradient operators

$$\begin{cases} \mathbb{U}(f)(x, t) = \int_0^t (g_{\nu(t-s)} * \mathbb{H}\operatorname{div} f)(x, s) ds, \\ \mathbb{G}(f)(x, t) = \int_0^t (g_{\nu(t-s)} * \mathbb{H}\nabla \operatorname{div} f)(x, s) ds. \end{cases} \quad (11)$$

Note that for a second order tensor f , $\mathbb{G}(f) = \nabla_x \mathbb{U}(f) = R \otimes R(\mathbb{U}(\nabla_x f))$. Let X be the Lagrangian path diffeomorphism, v the Lagrangian velocity, and τ the Lagrangian added stress,

$$\begin{aligned} v &= \frac{\partial X}{\partial t} = u \circ X, \\ \tau &= \sigma \circ X. \end{aligned} \quad (12)$$

We also set

$$\begin{aligned} g(a, t) &= (\nabla u)(X(a, t), t) = \mathbb{L}_\nu(\nabla u_0) \circ X(a, t) \\ &+ \mathbb{G}(\tau \circ X^{-1}) \circ X(a, t) - \mathbb{U}(\nabla_x((v \otimes v) \circ X^{-1})) \circ X(a, t). \end{aligned} \quad (13)$$

In Lagrangian variables the system is

$$\begin{cases} X(a, t) = a + \int_0^t \mathcal{V}(X, \tau, a, s) ds, \\ \tau(a, t) = \sigma_0(a) + \int_0^t \mathcal{T}(X, \tau, a, s) ds, \\ v(a, t) = \mathcal{V}(X, \tau, t) \end{cases} \quad (14)$$

where the Lagrangian nonlinearities \mathcal{V}, \mathcal{T} are

$$\begin{cases} \mathcal{V}(X, \tau, a, s) = \mathbb{L}_\nu(u_0) \circ X(a, s) + (\mathbb{U}((\tau - v \otimes v) \circ X^{-1})) \circ X(a, s), \\ \mathcal{T}(X, \tau, a, s) = (g\tau + \tau g^T - 2k\tau + 2\rho K(g + g^T))(a, s), \end{cases} \quad (15)$$

and g is defined above in (13). The main result of the paper is

Theorem 1. *Let $0 < \alpha < 1$ and $1 < p < \infty$, be given. Let also $v_1(0) = u_1(0) \in C^{1+\alpha, p}$ and $v_2(0) = u_2(0) \in C^{1+\alpha, p}$ be given divergence-free initial velocities, and $\sigma_1(0), \sigma_2(0) \in C^{\alpha, p}$ be given initial stresses. Then there exists $T_0 > 0$ and $C > 0$ depending on the norms of the initial data such that $(X_1, \tau_1, v_1), (X_2, \tau_2, v_2)$, with initial data $(Id, \sigma_1(0), u_1(0)), (Id, \sigma_2(0), u_2(0))$, are bounded in $Id + \operatorname{Lip}(0, T_0; C^{1+\alpha, p}) \times \operatorname{Lip}(0, T_0; C^{\alpha, p}) \times L^\infty(0, T_0; C^{1+\alpha, p})$ and solve the Lagrangian form (14) of (6). Moreover,*

$$\begin{aligned} \|X_2 - X_1\|_{\operatorname{Lip}(0, T_0; C^{1+\alpha, p})} + \|\tau_2 - \tau_1\|_{\operatorname{Lip}(0, T_0; C^{\alpha, p})} + \|v_2 - v_1\|_{L^\infty(0, T_0, C^{1+\alpha, p})} \\ \leq C(\|u_2(0) - u_1(0)\|_{1+\alpha, p} + \|\tau_2(0) - \tau_1(0)\|_{\alpha, p}) \end{aligned} \quad (16)$$

Remark 1. *The solutions' Lagrangian stresses τ are Lipschitz in time with values in C^α . Their Lagrangian counterparts $\sigma = \tau \circ X^{-1}$ are bounded in time with values in C^α and space-time Hölder continuous with exponent α . The Eulerian version of the equations (6) is satisfied in the sense of distributions, and solutions are unique in this class.*

The spaces $C^{\alpha,p}$ are defined in the next section. The proof of the theorem occupies the rest of the paper. We start by considering variations of Lagrangian variables. We take a family $(X_\epsilon, \tau_\epsilon)$ of flow maps depending smoothly on a parameter $\epsilon \in [1, 2]$, with initial data $u_{\epsilon,0}$ and $\sigma_{\epsilon,0}$. Note that $v_\epsilon = \partial_t X_\epsilon$. We use the following notations

$$\left\{ \begin{array}{l} u_\epsilon = \partial_t X_\epsilon \circ X_\epsilon^{-1}, g'_\epsilon = \frac{d}{d\epsilon} g_\epsilon, \\ X'_\epsilon = \frac{d}{d\epsilon} X_\epsilon, \eta_\epsilon = X'_\epsilon \circ X_\epsilon^{-1}, \\ v'_\epsilon = \frac{d}{d\epsilon} v_\epsilon, \\ \sigma_\epsilon = \tau_\epsilon \circ X_\epsilon^{-1}, \\ \tau'_\epsilon = \frac{d}{d\epsilon} \tau_\epsilon, \delta_\epsilon = \tau'_\epsilon \circ X_\epsilon^{-1}, \end{array} \right. \quad (17)$$

and

$$u'_{\epsilon,0} = \frac{d}{d\epsilon} u_\epsilon(0), \sigma'_{\epsilon,0} = \frac{d}{d\epsilon} \sigma_\epsilon(0). \quad (18)$$

We represent

$$\left\{ \begin{array}{l} X_2(a, t) - X_1(a, t) = \int_1^2 \mathcal{X}'_\epsilon d\epsilon, \\ \tau_2(a, t) - \tau_1(a, t) = \int_1^2 \pi_\epsilon d\epsilon, \\ v_2(a, t) - v_1(a, t) = \int_1^2 \frac{d}{d\epsilon} \mathcal{V}_\epsilon d\epsilon, \end{array} \right. \quad (19)$$

where

$$\begin{aligned} \mathcal{X}'_\epsilon &= \int_0^t \frac{d}{d\epsilon} \mathcal{V}_\epsilon ds, \quad \pi_\epsilon = \int_0^t \frac{d}{d\epsilon} \mathcal{T}_\epsilon ds + \sigma'_{\epsilon,0}, \\ \mathcal{V}_\epsilon &= \mathcal{V}(X_\epsilon, \tau_\epsilon), \quad \mathcal{T}_\epsilon = \mathcal{T}(X_\epsilon, \tau_\epsilon). \end{aligned} \quad (20)$$

We have the following commutator expressions arising by differentiating in ϵ ([1], [2]):

$$\left(\frac{d}{d\epsilon} (\mathbb{U}(\tau_\epsilon \circ X_\epsilon^{-1}) \circ X_\epsilon) \right) \circ X_\epsilon^{-1} = [\eta_\epsilon \cdot \nabla_x, \mathbb{U}](\sigma_\epsilon) + \mathbb{U}(\delta_\epsilon), \quad (21)$$

where

$$[\eta_\epsilon \cdot \nabla_x, \mathbb{U}](\sigma_\epsilon) = \eta_\epsilon \cdot \nabla_x (\mathbb{U}(\sigma_\epsilon)) - \mathbb{U}(\eta_\epsilon \cdot \nabla_x \sigma_\epsilon) \quad (22)$$

and

$$\begin{aligned} &\left(\frac{d}{d\epsilon} \mathbb{U}(v_\epsilon \otimes v_\epsilon \circ X_\epsilon^{-1}) \circ X_\epsilon \right) \circ X_\epsilon^{-1} \\ &= [\eta_\epsilon \cdot \nabla_x, \mathbb{U}](u_\epsilon \otimes u_\epsilon) + \mathbb{U}((v'_\epsilon \otimes v_\epsilon + v_\epsilon \otimes v'_\epsilon) \circ X_\epsilon^{-1}). \end{aligned} \quad (23)$$

We note, by the chain rule,

$$\nabla_a \mathcal{V} = (\nabla_a X) g. \quad (24)$$

Consequently, differentiating \mathcal{V}_ϵ , g_ϵ and the relation (24) we have

$$\left\{ \begin{array}{l} \left(\frac{d}{d\epsilon} \mathcal{V}_\epsilon \right) \circ X_\epsilon^{-1} = \eta_\epsilon \cdot (\mathbb{L}_\nu(\nabla_x u_{\epsilon,0})) + \mathbb{L}_\nu(u'_{\epsilon,0}) \\ + [\eta_\epsilon \cdot \nabla_x, \mathbb{U}](\sigma_\epsilon - u_\epsilon \otimes u_\epsilon) + \mathbb{U}(\delta_\epsilon - (v'_\epsilon \otimes v_\epsilon + v_\epsilon \otimes v'_\epsilon) \circ X_\epsilon^{-1}), \\ g_\epsilon = \mathbb{L}(\nabla_x u_{\epsilon,0}) \circ X_\epsilon + \mathbb{G}(\sigma_\epsilon) \circ X_\epsilon - \mathbb{U}(\nabla_x(u_\epsilon \otimes u_\epsilon)) \circ X_\epsilon, \\ g'_\epsilon \circ X_\epsilon^{-1} = \eta_\epsilon \cdot \mathbb{L}_\nu(\nabla_x \nabla_x u_{\epsilon,0}) + \mathbb{L}_\nu(\nabla_x u'_{\epsilon,0}) + [\eta_\epsilon \cdot \nabla_x, \mathbb{G}](\sigma_\epsilon) + \mathbb{G}(\delta_\epsilon) \\ - [\eta_\epsilon \cdot \nabla_x, \mathbb{U}](\nabla_x(u_\epsilon \otimes u_\epsilon)) - \mathbb{U}(\nabla_x((v'_\epsilon \otimes v_\epsilon + v_\epsilon \otimes v'_\epsilon) \circ X_\epsilon^{-1})), \\ \frac{d}{d\epsilon} (\nabla_a \mathcal{V}_\epsilon) = (\nabla_a X'_\epsilon) g_\epsilon + (\nabla_a X_\epsilon) g'_\epsilon, \\ \frac{d}{d\epsilon} \mathcal{T}_\epsilon = g'_\epsilon \tau_\epsilon + g_\epsilon \tau'_\epsilon + \tau'_\epsilon g_\epsilon^T + \tau_\epsilon(g'_\epsilon)^T - 2k\tau'_\epsilon + 2\rho K(g'_\epsilon + (g'_\epsilon)^T). \end{array} \right. \quad (25)$$

3 Functions, operators, commutators

We consider function spaces

$$C^{\alpha,p} = C^\alpha(\mathbb{R}^d) \cap L^p(\mathbb{R}^d) \quad (26)$$

with norm

$$\|f\|_{\alpha,p} = \|f\|_{C^\alpha(\mathbb{R}^d)} + \|f\|_{L^p(\mathbb{R}^d)} \quad (27)$$

for $\alpha \in (0, 1), p \in (1, \infty)$, $C^{1+\alpha}(\mathbb{R}^d)$ with norm

$$\|f\|_{C^{1+\alpha}(\mathbb{R}^d)} = \|f\|_{L^\infty(\mathbb{R}^d)} + \|\nabla f\|_{C^\alpha(\mathbb{R}^d)}, \quad (28)$$

and

$$C^{1+\alpha,p} = C^{1+\alpha}(\mathbb{R}^d) \cap W^{1,p}(\mathbb{R}^d) \quad (29)$$

with norm

$$\|f\|_{1+\alpha,p} = \|f\|_{C^{1+\alpha}(\mathbb{R}^d)} + \|f\|_{W^{1,p}(\mathbb{R}^d)}. \quad (30)$$

We also use spaces of paths, $L^\infty(0, T; Y)$ with the usual norm,

$$\|f\|_{L^\infty(0, T; Y)} = \sup_{t \in [0, T]} \|f(t)\|_Y, \quad (31)$$

spaces $Lip(0, T; Y)$ with norm

$$\|f\|_{Lip(0, T; Y)} = \sup_{t \neq s, t, s \in [0, T]} \frac{\|f(t) - f(s)\|_Y}{|t - s|} + \|f\|_{L^\infty(0, T; Y)} \quad (32)$$

where Y is $C^{\alpha,p}$ or $C^{1+\alpha,p}$ in the following. We use the following lemmas.

Lemma 1 ([2]). *Let $0 < \alpha < 1, 1 < p < \infty$. Let $\eta \in C^{1+\alpha}(\mathbb{R}^d)$ and let*

$$(\mathbb{K}\sigma)(x) = P.V. \int_{\mathbb{R}^d} k(x - y)\sigma(y)dy \quad (33)$$

be a classical Calderon-Zygmund operator with kernel k which is smooth away from the origin, homogeneous of degree $-d$ and with mean zero on spheres about the origin. Then the commutator $[\eta \cdot \nabla, \mathbb{K}]$ can be defined as a bounded linear operator in $C^{\alpha,p}$ and

$$\|[\eta \cdot \nabla, \mathbb{K}]\sigma\|_{C^{\alpha,p}} \leq C \|\eta\|_{C^{1+\alpha}(\mathbb{R}^d)} \|\sigma\|_{C^{\alpha,p}}. \quad (34)$$

Lemma 2 (Generalized Young's inequality). *Let $1 \leq q \leq \infty$ and $C > 0$. Suppose K is a measurable function on $\mathbb{R}^d \times \mathbb{R}^d$ such that*

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} |K(x, y)|dy \leq C, \quad \sup_{y \in \mathbb{R}^d} \int_{\mathbb{R}^d} |K(x, y)|dx \leq C. \quad (35)$$

If $f \in L^q(\mathbb{R}^d)$, the function Tf defined by

$$Tf(x) = \int_{\mathbb{R}^d} K(x, y)f(y)dy \quad (36)$$

is well defined almost everywhere and is in L^q , and $\|Tf\|_{L^q} \leq C \|f\|_{L^q}$.

The proof of this lemma for $1 < q < \infty$ is done using duality, a straightforward application of Young's inequality and changing order of integration. The extreme cases $q = 1$ and $q = \infty$ are proved directly by inspection.

For simplicity of notation, let us denote

$$M_X = 1 + \|X - \text{Id}\|_{L^\infty(0, T; C^{1+\alpha})}. \quad (37)$$

Theorem 2. Let $0 < \alpha < 1, 1 < p < \infty$ and let $T > 0$. Also let X be a volume preserving diffeomorphism such that $X - \text{Id} \in \text{Lip}(0, T; C^{1+\alpha})$. Then

$$\|\tau \circ X^{-1}\|_{L^\infty(0, T; C^{\alpha, p})} \leq \|\tau\|_{L^\infty(0, T; C^{\alpha, p})} M_X^\alpha. \quad (38)$$

If $X' \in \text{Lip}(0, T; C^{1+\alpha})$, then

$$\|X' \circ X^{-1}\|_{L^\infty(0, T; C^{1+\alpha})} \leq \|X'\|_{L^\infty(0, T; C^{1+\alpha})} M_X^{1+2\alpha}. \quad (39)$$

If $v \in \text{Lip}(0, T; W^{1,p})$, then

$$\|v \circ X^{-1}\|_{L^\infty(0, T; W^{1,p})} \leq \|v\|_{L^\infty(0, T; W^{1,p})} M_X. \quad (40)$$

If in addition $\partial_t X', \partial_t X$ exist in $L^\infty(0, T; C^{1+\alpha})$, then

$$\|X' \circ X^{-1}\|_{\text{Lip}(0, T; C^\alpha)} \leq \|X'\|_{\text{Lip}(0, T; C^{1+\alpha})} \|X - \text{Id}\|_{\text{Lip}(0, T; C^{1+\alpha})} M_X^{1+3\alpha}. \quad (41)$$

Proof.

$$\|\tau \circ X^{-1}\|_{L^p \cap L^\infty} = \|\tau\|_{L^p \cap L^\infty}, \quad (42)$$

and, denoting the seminorm

$$[\tau]_\alpha = \sup_{a \neq b, a, b \in \mathbb{R}^2} \frac{|\tau(a) - \tau(b)|}{|a - b|^\alpha}$$

we have

$$[\tau \circ X^{-1}(t)]_\alpha \leq [\tau(t)]_\alpha \|\nabla_x X^{-1}(t)\|_{L^\infty}^\alpha \leq [\tau(t)]_\alpha (1 + \|X - \text{Id}\|_{L^\infty(0, T; C^{1+\alpha})})^\alpha. \quad (43)$$

Note that this shows that the same bound holds when we replace X^{-1} by X . For the second and third part, it suffices to remark that

$$\nabla_x(X' \circ X^{-1}) = ((\nabla_a X) \circ X^{-1})^{-1} ((\nabla_a X') \circ X^{-1}) \quad (44)$$

and the previous part gives the bound in terms of Lagrangian variables. For the last part, we note that

$$\begin{aligned} & \frac{1}{t-s} (X'(X^{-1}(x, t), t) - X'(X^{-1}(x, s), s)) \\ &= \int_0^1 ((\partial_t X') (X^{-1}(x, \beta_\tau), \beta_\tau) + (\partial_t X^{-1})(x, \beta_\tau) (\nabla_a X') (X^{-1}(x, \beta_\tau), \beta_\tau)) d\tau, \end{aligned} \quad (45)$$

where

$$\beta_\tau = \tau t + (1 - \tau)s. \quad (46)$$

Now noting that

$$\partial_t X^{-1} = -((\partial_t X) \circ X^{-1}) ((\nabla_a X)^{-1} \circ X^{-1}) \quad (47)$$

we have

$$\begin{aligned} & \frac{1}{t-s} \|X' \circ X^{-1}(t) - X' \circ X^{-1}(s)\|_{C^\alpha} \\ & \leq \left(\|\partial_t X'\|_{L^\infty(0, T; C^\alpha)} + \|\partial_t X\|_{L^\infty(0, T; C^\alpha)} \|X'\|_{L^\infty(0, T; C^{1+\alpha})} \right) \left(1 + \|X - \text{Id}\|_{L^\infty(0, T; C^{1+\alpha})} \right)^{1+3\alpha} \end{aligned} \quad (48)$$

so that

$$\|X' \circ X^{-1}\|_{\text{Lip}(0, T; C^\alpha)} \leq \|X'\|_{\text{Lip}(0, T; C^{1+\alpha})} \|X - \text{Id}\|_{\text{Lip}(0, T; C^{1+\alpha})} \left(1 + \|X - \text{Id}\|_{L^\infty(0, T; C^{1+\alpha})} \right)^{1+3\alpha}. \quad (49)$$

□

Theorem 3. Let $0 < \alpha < 1, 1 < p < \infty$ and let $T > 0$. There exists a constant C independent of T and ν such that for any $0 < t < T$,

$$\begin{aligned} \|\mathbb{L}_\nu(u_0)\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \|u_0\|_{\alpha,p}, \\ \|\mathbb{L}_\nu(u_0)\|_{L^\infty(0,T;C^{1+\alpha,p})} &\leq C \|u_0\|_{1+\alpha,p}, \\ \|\mathbb{L}_\nu(\nabla u_0)(t)\|_{\alpha,p} &\leq \frac{C}{(\nu t)^{\frac{1}{2}}} \|u_0\|_{\alpha,p}, \\ \|\mathbb{L}_\nu(\nabla u_0)\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \|u_0\|_{1+\alpha,p} \end{aligned} \quad (50)$$

hold.

Proof.

$$\begin{aligned} \|\mathbb{L}_\nu(u_0)(t)\|_{\alpha,p} &\leq \|g_{\nu t}\|_{L^1} \|u_0\|_{\alpha,p} = \|u_0\|_{\alpha,p}, \\ \|\mathbb{L}_\nu(u_0)(t)\|_{1+\alpha,p} &\leq \|g_{\nu t}\|_{L^1} \|u_0\|_{1+\alpha,p} = \|u_0\|_{1+\alpha,p}, \\ \|\mathbb{L}_\nu(\nabla u_0)(t)\|_{\alpha,p} &\leq \|\nabla g_{\nu t}\|_{L^1} \|u_0\|_{1+\alpha,p} = \frac{C}{(\nu t)^{\frac{1}{2}}} \|u_0\|_{\alpha,p}, \\ \|\mathbb{L}_\nu(\nabla u_0)(t)\|_{\alpha,p} &\leq \|g_{\nu t}\|_{L^1} \|\nabla u_0\|_{\alpha,p} \leq \|u_0\|_{1+\alpha,p}. \end{aligned} \quad (51)$$

□

Theorem 4. Let $0 < \alpha < 1, 1 < p < \infty$ and let $T > 0$. There exists a constant C such that

$$\|\mathbb{U}(\sigma)\|_{L^\infty(0,T;C^{\alpha,p})} \leq C \left(\frac{T}{\nu} \right)^{\frac{1}{2}} \|\sigma\|_{L^\infty(0,T;C^{\alpha,p})}. \quad (52)$$

Proof.

$$\begin{aligned} \|\mathbb{U}(\sigma)(t)\|_{C^{\alpha,p}} &\leq C \int_0^t \|\nabla g_{\nu(t-s)}\|_{L^1} \|\sigma(s)\|_{\alpha,p} ds \\ &\leq \frac{C}{\nu^{\frac{1}{2}}} \int_0^t \frac{1}{(t-s)^{\frac{1}{2}}} ds \|\sigma\|_{L^\infty(0,T;C^{\alpha,p})} \leq \frac{C}{\nu^{\frac{1}{2}}} \sqrt{T} \|\sigma\|_{L^\infty(0,T;C^{\alpha,p})}. \end{aligned} \quad (53)$$

□

Theorem 5. Let $0 < \alpha < 1, 1 < p < \infty$ and let $T > 0$. There exist constants C_1, C_2 depending only on α and ν , and $C_3(T, X), C_4(T, X)$ such that

$$\begin{aligned} \|\mathbb{G}(\tau \circ X^{-1})\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C_1 \|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})}^\alpha \|\tau(0)\|_{\alpha,p} (1 + C_3(T, X)) \\ &\quad + C_2 \|\tau\|_{Lip(0,T;C^{\alpha,p})} C_4(T, X) \end{aligned} \quad (54)$$

where $C_3(T, X)$ and $C_4(T, X)$ are of the form $CT^{\frac{1}{2}} \left(\|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})}^\alpha + \|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})}^4 \right)$.

Proof. Since $\mathbb{G} = (R \otimes R)\mathbb{H}\Gamma$ where

$$\Gamma(\tau \circ X^{-1}) = \int_0^t \Delta g_{\nu(t-s)} * (\tau \circ X^{-1}(s)) ds, \quad (55)$$

we can replace \mathbb{G} by Γ . Then $\Gamma(\tau \circ X^{-1})$ can be written as

$$\begin{aligned} \Gamma(\tau \circ X^{-1})(t) &= \int_0^t \Delta g_{\nu(t-s)} * \left((\tau \circ X^{-1})(s) - (\tau \circ X^{-1})(t) \right) ds \\ &\quad + \int_0^t \Delta g_{\nu(t-s)} * (\tau \circ X^{-1})(t) ds. \end{aligned} \quad (56)$$

But

$$\int_0^t \Delta g_{\nu(t-s)} * (\tau \circ X^{-1})(t) ds = \tau \circ X^{-1}(t) - g_{\nu t} * (\tau \circ X^{-1})(t) \quad (57)$$

so the second term is bounded by $2 \|\tau\|_{L^\infty(0,T;C^{\alpha,p})} M_X^\alpha$ by Theorem 2. Now we let

$$\tau \circ X^{-1}(x, s) - \tau \circ X^{-1}(x, t) = \Delta_1 \tau(x, s, t) + \Delta_2 \tau(x, s, t), \quad (58)$$

where

$$\begin{aligned} \Delta_1 \tau(x, s, t) &= \tau(X^{-1}(x, s), s) - \tau(X^{-1}(x, s), t), \\ \Delta_2 \tau(x, s, t) &= \tau(X^{-1}(x, s), t) - \tau(X^{-1}(x, t), t). \end{aligned} \quad (59)$$

But since

$$\|\Delta_1 \tau(s, t)\|_{C^{\alpha,p}} \leq |t - s| M_X^\alpha \|\tau\|_{Lip(0,T;C^{\alpha,p})}, \quad (60)$$

by the proof of Theorem 2 we get

$$\left\| \int_0^t \Delta g_{\nu(t-s)} * \Delta_1 \tau(s, t) ds \right\|_{\alpha,p} \leq \frac{Ct}{\nu} \|\tau\|_{Lip(0,T;C^{\alpha,p})} M_X^\alpha, \quad (61)$$

On the other hand,

$$\int_0^t \Delta g_{\nu(t-s)} * \Delta_2 \tau(s, t) ds = \int_0^t \int_{\mathbb{R}^d} K(x, z, t, s) \tau(z, t) dz ds, \quad (62)$$

where

$$K(x, z, t, s) = \Delta g_{\nu(t-s)}(x - X(z, s)) - \Delta g_{\nu(t-s)}(x - X(z, t)). \quad (63)$$

We use the following lemma.

Lemma 3. $K(x, z, t, s)$ is L^1 in both the x variable and the z variable, and

$$\sup_z \|K(\cdot, z, t, s)\|_{L^1}, \sup_x \|K(x, \cdot, t, s)\|_{L^1} \leq \frac{C \|X - \text{Id}\|_{Lip(0,T;L^\infty)}}{|t - s|^{\frac{1}{2}} \nu^{\frac{3}{2}}}. \quad (64)$$

Proof. We define

$$S(x) = 4\pi e^{-|x|^2} \left(|x|^2 - \frac{d}{2} \right) \quad (65)$$

so that

$$(\Delta g_{\nu(t-s)}) = (4\pi\nu(t-s))^{-(\frac{d}{2}+1)} S\left(\frac{x}{(4(t-s))^{\frac{1}{2}}}\right). \quad (66)$$

Then

$$\begin{aligned} \int |K(x, z, t, s)| dz &= \int (4\pi\nu(t-s))^{-(\frac{d}{2}+1)} \left| S\left(\frac{x - X(z, s)}{(4\nu(t-s))^{\frac{1}{2}}}\right) - S\left(\frac{x - X(z, t)}{(4\nu(t-s))^{\frac{1}{2}}}\right) \right| dz \\ &= \int (4\pi\nu(t-s))^{-(\frac{d}{2}+1)} \left| S\left(\frac{x - y}{(4\nu(t-s))^{\frac{1}{2}}}\right) - S\left(\frac{x - X(y, t-s)}{(4\nu(t-s))^{\frac{1}{2}}}\right) \right| dy \\ &= (4\pi\nu(t-s))^{-1} \pi^{-(\frac{d}{2}+1)} \int \left| S(u) - S\left(u - \frac{(X - \text{Id})(x - (4(t-s))^{\frac{1}{2}}u, t-s)}{(4\nu(t-s))^{\frac{1}{2}}}\right) \right| du. \end{aligned} \quad (67)$$

However, for each u

$$\begin{aligned} \left| S(u) - S\left(u - \frac{(X - \text{Id})(x - (4\nu(t-s))^{\frac{1}{2}}u, t-s)}{(4\nu(t-s))^{\frac{1}{2}}}\right) \right| &\leq \left| \frac{(X - \text{Id})(x - (4\nu(t-s))^{\frac{1}{2}}u, t-s)}{(4\nu(t-s))^{\frac{1}{2}}} \right| \\ &\times \sup \left\{ |\nabla S(u - z)| : |z| \leq \left| \frac{(X - \text{Id})(x - (4\nu(t-s))^{\frac{1}{2}}u, t-s)}{(4\nu(t-s))^{\frac{1}{2}}} \right| \right\} \end{aligned} \quad (68)$$

and we have

$$\left| \frac{(X - \text{Id})(x - (4\nu(t-s))^{\frac{1}{2}}u, t-s)}{(4\nu(t-s))^{\frac{1}{2}}} \right| \leq \|(X - \text{Id})\|_{Lip(0,T;L^\infty)} \frac{|t - s|^{\frac{1}{2}}}{\nu^{\frac{1}{2}}} \leq CT^{\frac{1}{2}} \quad (69)$$

and obviously

$$\tilde{S}(u) = \sup_{z \leq CT^{\frac{1}{2}}} |(\nabla S)(u - z)| \quad (70)$$

is integrable in \mathbb{R}^d ; because ∇S is Schwartz,

$$|(\nabla S)(x)| \leq \frac{C_d}{(1 + 2C^2T + |x|^2)^d} \quad (71)$$

for some constant C_d , but if $|z| \leq CT^{\frac{1}{2}}$, then $|u - z|^2 \geq |u|^2 - C^2T$ and

$$|(\nabla S)(u - z)| \leq \frac{C_d}{(1 + C^2T + |u|^2)^d} \quad (72)$$

and the right side of above is clearly integrable with bound depending only on d and T . Therefore, we have

$$\int |K(x, z, t, s)| dz \leq |t - s|^{-\frac{1}{2}} \nu^{-\frac{3}{2}} \|(X - \text{Id})\|_{Lip(0, T; L^\infty)} C(d, T). \quad (73)$$

Similarly,

$$\begin{aligned} \int |K(x, z, t, s)| dx &= \int (4\pi\nu(t-s))^{-(\frac{d}{2}+1)} \left| S\left(\frac{x - X(z, s)}{(4\nu(t-s))^{\frac{1}{2}}}\right) - S\left(\frac{x - X(z, t)}{(4\nu(t-s))^{\frac{1}{2}}}\right) \right| dx \\ &= \int (4\pi\nu(t-s))^{-1} \pi^{-(\frac{d}{2}+1)} \left| S(y) - S\left(y + \frac{X(z, s) - X(z, t)}{(4\nu(t-s))^{\frac{1}{2}}}\right) \right| dy \end{aligned} \quad (74)$$

and again we have

$$\left| \frac{X(z, s) - X(z, t)}{(4\nu(t-s))^{\frac{1}{2}}} \right| \leq \|(X - \text{Id})\|_{Lip(0, T; L^\infty)} |t - s|^{\frac{1}{2}} \nu^{-\frac{1}{2}} \leq CT^{\frac{1}{2}}. \quad (75)$$

Therefore, we have the bound

$$\int |K(x, z)| dx \leq |t - s|^{-\frac{1}{2}} \nu^{-\frac{3}{2}} \|(X - \text{Id})\|_{Lip(0, T; L^\infty)} C(d, T). \quad (76)$$

□

From Lemma 3 and generalized Young's inequality, we have

$$\left\| \int_0^t \Delta g_{\nu(t-s)} * \Delta_2 \tau(s, t) ds \right\|_{L^p \cap L^\infty} \leq \frac{C}{\nu} \left(\left(\frac{t}{\nu} \right)^{\frac{1}{2}} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} \right) \|\tau\|_{L^\infty(0, T; L^p \cap L^\infty)}. \quad (77)$$

For the Hölder seminorm, we measure the finite difference. Let us denote $\delta_h f(x, t) = f(x+h, t) - f(x, t)$. If $|h| < t$, then

$$\delta_h \left(\int_0^t \Delta g_{\nu(t-s)} * \Delta_2 \tau(s, t) ds \right) = \int_0^t \delta_h(\Delta g_{\nu(t-s)}) * \Delta_2 \tau(s, t) ds. \quad (78)$$

If $0 < t - s < |h|$, then $\|\delta_h \Delta g_{\nu(t-s)}\|_{L^1} \leq 2 \|\Delta g_{\nu(t-s)}\|_{L^1} \leq \frac{C}{\nu(t-s)}$ and since

$$\|\Delta_2 \tau(s, t)\|_{L^\infty} \leq |t - s|^\alpha \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau\|_{L^\infty(0, T; C^{\alpha, p})} \quad (79)$$

we have

$$\left\| \int_{t-|h|}^t \delta_h(\Delta g_{\nu(t-s)}) * \Delta_2 \tau(s, t) ds \right\|_{L^\infty} \leq \frac{C}{\nu \alpha} |h|^\alpha \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau\|_{L^\infty(0, T; C^{\alpha, p})}. \quad (80)$$

If $|h| < t - s < t$, then following lines of Lemma 3 $\delta_h(\Delta g_{\nu(t-s)})$ is a L^1 function with

$$\|\delta_h(\Delta g_{\nu(t-s)})\|_{L^1} \leq \frac{C|h|}{(\nu(t-s))^{\frac{3}{2}}} \quad (81)$$

and we have

$$\begin{aligned} & \left\| \int_0^{t-|h|} \delta_h(\Delta g_{\nu(t-s)}) * \Delta_2 \tau(s, t) ds \right\|_{L^\infty} \\ & \leq \begin{cases} \frac{C}{\nu^{\frac{3}{2}}} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau\|_{L^\infty(0, T; C^{\alpha, p})} |h|^{\frac{1}{2} \frac{t^\alpha}{\alpha}} & \alpha \leq \frac{1}{2}, \\ \frac{C}{\nu^{\frac{3}{2}}} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau\|_{L^\infty(0, T; C^{\alpha, p})} |h|^{\frac{t^{\alpha-\frac{1}{2}}}{\alpha-\frac{1}{2}}} & \alpha > \frac{1}{2}. \end{cases} \end{aligned} \quad (82)$$

If $|h| \geq t$, then we only have the first term. Therefore, we have

$$\frac{1}{|h|^\alpha} \left\| \delta_h \left(\int_0^t \Delta g_{\nu(t-s)} * \Delta_2 \tau(s, t) ds \right) \right\|_{L^\infty} \leq \frac{C(\alpha)}{\nu} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau\|_{L^\infty(0, T; C^{\alpha, p})}. \quad (83)$$

We note that

$$\|\tau(t)\|_{\alpha, p} \leq \|\tau(0)\|_{\alpha, p} + t \|\tau\|_{Lip(0, T; C^{\alpha, p})}. \quad (84)$$

To summarize, we have

$$\begin{aligned} & \|\Gamma(\tau \circ X^{-1})\|_{L^\infty(0, T; C^{\alpha, p})} \\ & \leq C(\alpha) \left(1 + \frac{1}{\nu} \right) \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau(0)\|_{\alpha, p} + C(\alpha) \left(1 + \frac{1}{\nu} \right) \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha T \|\tau\|_{Lip(0, T; C^{\alpha, p})} \\ & \quad + \frac{C(\alpha)}{\nu} \left(\frac{T}{\nu} \right)^{\frac{1}{2}} \max\{\|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha, \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^4\} (\|\tau(0)\|_{\alpha, p} + T \|\tau\|_{Lip(0, T; C^{\alpha, p})}), \end{aligned} \quad (85)$$

and this completes the proof. \square

Theorem 6. Let $0 < \alpha < 1, 1 < p < \infty$ and let $T > 0$. Let $X' \in Lip(0, T; C^{1+\alpha})$ with $\partial_t X' \in L^\infty(0, T; C^{1+\alpha})$. There exists a constant C such that

$$\begin{aligned} & \| [X' \circ X^{-1} \cdot \nabla, \mathbb{U}] (\sigma) \|_{L^\infty(0, T; C^{\alpha, p})} \\ & \leq C \left(\left(\frac{T}{\nu} \right)^{\frac{1}{2}} + \frac{T}{\nu} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} \right) M_X^{1+3\alpha} \|X'\|_{Lip(0, T; C^{1+\alpha})} \|\sigma\|_{L^\infty(0, T; C^{\alpha, p})} \end{aligned} \quad (86)$$

Proof. First, we denote

$$\eta = X' \circ X^{-1}. \quad (87)$$

Then we have

$$\begin{aligned} & [\eta \cdot \nabla, \mathbb{U}] (\sigma)(t) \\ & = \eta(t) \cdot \nabla \int_0^t g_{\nu(t-s)} * \mathbb{H} \text{div} \sigma(s) ds - \int_0^t g_{\nu(t-s)} * \mathbb{H} \text{div} (\eta(s) \cdot \nabla \sigma(s)) ds \\ & = [\eta(t) \cdot \nabla, \mathbb{H}] \int_0^t g_{\nu(t-s)} * \text{div} \sigma(s) ds + \mathbb{H} \int_0^t (\nabla g_{\nu(t-s)}) * (\nabla \cdot \eta(s) \sigma(s)) ds \\ & \quad - \mathbb{H} \int_0^t (\nabla \nabla g_{\nu(t-s)}) * (\eta(s) - \eta(t)) \sigma(s) ds \\ & \quad + \mathbb{H} \int_0^t (\eta(t) \cdot (\nabla \nabla g_{\nu(t-s)}) * \sigma(s) - (\nabla \nabla g_{\nu(t-s)}) * (\eta(t) \sigma(s))) ds, \end{aligned} \quad (88)$$

where $(\nabla \nabla g_{\nu(t-s)}) * (\eta(s) - \eta(t)) \sigma(s)$, $\eta(t) \cdot (\nabla \nabla g_{\nu(t-s)}) * \sigma(s)$, and $(\nabla \nabla g_{\nu(t-s)}) * (\eta(s) \sigma(s))$ represent

$$\begin{aligned} & \sum_{i,j} (\partial_i \partial_j g_{\nu(t-s)} *) (\eta_i(s) - \eta_i(t)) \sigma_{jk}(s), \\ & \sum_{i,j} \eta_i(t) (\partial_i \partial_j g_{\nu(t-s)}) * \sigma_{jk}(s), \text{ and respectively } \sum_{i,j} (\partial_i \partial_j g_{\nu(t-s)}) * (\eta_i(s) \sigma_{jk}(s)). \end{aligned} \quad (89)$$

The first term is bounded by Lemma 1 and the second term is estimated directly

$$\begin{aligned} \left\| [\eta(t) \cdot \nabla, \mathbb{H}] \int_0^t g_{\nu(t-s)} * \operatorname{div} \sigma(s) ds \right\|_{\alpha, p} &\leq C \|\eta(t)\|_{C^{1+\alpha}} \left(\frac{t}{\nu} \right)^{\frac{1}{2}} \|\sigma\|_{L^\infty(0, T; C^{\alpha, p})}, \\ \left\| \mathbb{H} \int_0^t (\nabla g_{\nu(t-s)}) * (\nabla \cdot \eta(s) \sigma(s)) ds \right\|_{\alpha, p} &\leq C \left(\frac{t}{\nu} \right)^{\frac{1}{2}} \|\eta\|_{L^\infty(0, T; C^{1+\alpha})} \|\sigma\|_{L^\infty(0, T; C^{\alpha, p})}. \end{aligned} \quad (90)$$

The third term is bounded by

$$\frac{Ct}{\nu} \|\eta\|_{Lip(0, T; C^\alpha)} \|\sigma\|_{L^\infty(0, T; C^{\alpha, p})} \quad (91)$$

by the virtue of Theorem 2. For the last term, note that

$$\begin{aligned} &(\eta(t) \cdot (\nabla \nabla g_{\nu(t-s)}) * \sigma(s) - (\nabla \nabla g_{\nu(t-s)}) * (\eta(t) \sigma(s))) (x) \\ &= \int_{\mathbb{R}^d} \nabla \nabla g_{\nu(t-s)}(z) z \cdot \left(\int_0^1 \nabla \eta(x - (1-\lambda)z, t) d\lambda \right) \sigma(x - z, s) dz \end{aligned} \quad (92)$$

and note that $\nabla \nabla g_{\nu(t-s)}(z) z$ is a L^1 function with

$$\|\nabla \nabla g_{\nu(t-s)}(z) z\|_{L^1} \leq \frac{C}{(\nu(t-s))^{\frac{1}{2}}}. \quad (93)$$

Therefore,

$$\begin{aligned} &\|(\eta(t) \cdot (\nabla \nabla g_{\nu(t-s)}) * \sigma(s) - (\nabla \nabla g_{\nu(t-s)}) * (\eta(t) \sigma(s)))\|_{\alpha, p} \\ &\leq \frac{C}{(\nu(t-s))^{\frac{1}{2}}} \|\eta(t)\|_{C^{1+\alpha}} \|\sigma(s)\|_{\alpha, p} \end{aligned} \quad (94)$$

so that the last term is bounded by

$$C \left(\frac{t}{\nu} \right)^{\frac{1}{2}} \|\eta(t)\|_{C^{1+\alpha}} \|\sigma\|_{L^\infty(0, T; C^{\alpha, p})}. \quad (95)$$

We finish the proof by replacing η by X' using Theorem 2. \square

Theorem 7. *Let $0 < \alpha < 1$, $1 < p < \infty$ and let $T > 0$. Let $X' \in Lip(0, T; C^{1+\alpha})$ with $\partial_t X' \in L^\infty(0, T; C^{1+\alpha})$. There exists a constant $C(\alpha)$ depending only on α such that*

$$\begin{aligned} &\left\| [X' \circ X^{-1} \cdot \nabla, \mathbb{G}] (\tau \circ X^{-1}) \right\|_{L^\infty(0, T; C^{\alpha, p})} \\ &\leq (\|X'\|_{L^\infty(0, T; C^{1+\alpha})} + \|X'\|_{Lip(0, T; C^{1+\alpha})} T^{\frac{1}{2}}) R \end{aligned} \quad (96)$$

where R is a polynomial function on $\|\tau\|_{Lip(0, T; C^{\alpha, p})}$, $\|X - \operatorname{Id}\|_{Lip(0, T; C^{1+\alpha})}$, whose coefficients depend on α , ν , and T , and in particular it grows polynomially in T and bounded below.

Proof. Again we denote $\eta = X' \circ X^{-1}$. Also it suffices to bound

$$[\eta \cdot \nabla, \Gamma] (\tau \circ X^{-1}) = \eta(t) \cdot \nabla \Gamma (\tau \circ X^{-1}) - \Gamma (\eta \cdot \nabla (\tau \circ X^{-1})) \quad (97)$$

where Γ is as defined in (55), since

$$[\eta \cdot \nabla, \mathbb{G}] = (R \otimes R) \mathbb{H} [\eta \cdot \nabla, \Gamma] + [\eta(t) \cdot \nabla, (R \otimes R) \mathbb{H}] \Gamma \quad (98)$$

and the second term is bounded by Lemma 1. For the first term, we have

$$[\eta \cdot \nabla, \Gamma] (\tau \circ X^{-1})(t) = I_1 + I_2 + I_3 + I_4 + I_5 + I_6, \quad (99)$$

where

$$\begin{aligned}
I_1 &= \int_0^t \eta(t) \cdot (\nabla \Delta g_{\nu(t-s)} * (\tau \circ X^{-1}(t))) - \nabla \Delta g_{\nu(t-s)} * (\eta(t) \tau \circ X^{-1}(t)) \, ds, \\
I_2 &= \int_0^t \eta(t) \cdot (\nabla \Delta g_{\nu(t-s)} * (\tau \circ X^{-1}(s) - \tau \circ X^{-1}(t))) \\
&\quad - \nabla \Delta g_{\nu(t-s)} * (\eta(t) (\tau \circ X^{-1}(s) - \tau \circ X^{-1}(t))) \, ds, \\
I_3 &= - \int_0^t \nabla \Delta g_{\nu(t-s)} * ((\eta(s) - \eta(t)) (\tau \circ X^{-1}(s))) \, ds, \\
I_4 &= \int_0^t \Delta g_{\nu(t-s)} * (\nabla \cdot (\eta(s) - \eta(t)) \tau \circ X^{-1}(s)) \, ds, \\
I_5 &= \int_0^t \Delta g_{\nu(t-s)} * (\nabla \cdot \eta(t) (\tau \circ X^{-1}(s) - \tau \circ X^{-1}(t))) \, ds, \\
I_6 &= - \frac{1}{\nu} (\nabla \cdot \eta(t) \tau \circ X^{-1}(t) - g_{\nu t} * (\nabla \cdot \eta(t) \tau \circ X^{-1}(t))).
\end{aligned} \tag{100}$$

First, $I_1 + I_6$ can be bounded:

$$\begin{aligned}
I_1 + I_6 &= \frac{1}{\nu} (\eta(t) \cdot \nabla (g_{\nu t} * (\tau \circ X^{-1}(t))) - \nabla (g_{\nu t} * (\eta(t) \tau \circ X^{-1}(t)))) \\
&\quad - \frac{1}{\nu} g_{\nu t} * (\nabla \cdot \eta(t) (\tau \circ X^{-1}(t)))
\end{aligned} \tag{101}$$

and the first term is treated in the same way as (92). Since the first term is

$$\frac{1}{\nu} \left(\int_{\mathbb{R}^d} \nabla g_{\nu t}(y) y \cdot \int_0^1 \nabla \eta(x - (1 - \lambda)y, t) d\lambda (\tau \circ X^{-1})(x - y, t) dy \right) \tag{102}$$

and

$$\|\nabla g_{\nu t}(y) y\|_{L^1} \leq C, \tag{103}$$

the $C^{\alpha,p}$ -norm of the first term is bounded by

$$\frac{C}{\nu} \|\eta(t)\|_{C^{1+\alpha}} \|\tau \circ X^{-1}(t)\|_{\alpha,p}. \tag{104}$$

The $C^{\alpha,p}$ -norm of the second term is also bounded by the same bound. Therefore,

$$\|I_1 + I_6\|_{L^\infty(0,T;C^{\alpha,p})} \leq \frac{C}{\nu} M_X^{1+3\alpha} \|X'\|_{L^\infty(0,T;C^{1+\alpha})} \|\tau\|_{L^\infty(0,T;C^{\alpha,p})}. \tag{105}$$

The term I_3 is bounded due to Theorem 2. Since $\eta \in Lip(0, T; C^\alpha)$ we have

$$\begin{aligned}
\|I_3\|_{L^\infty(0,T;C^{\alpha,p})} &\leq \frac{C}{\nu} \left(\frac{T}{\nu} \right)^{\frac{1}{2}} M_X^{1+4\alpha} \|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})} \\
&\quad \|X'\|_{Lip(0,T;C^{1+\alpha})} \|\tau\|_{L^\infty(0,T;C^{\alpha,p})}.
\end{aligned} \tag{106}$$

The terms I_4 , and I_5 are treated in the spirit of Theorem 5. We treat $L^p \cap L^\infty$ norm and Hölder seminorm separately. For the term I_5 , we have

$$I_5 = \int_0^t \Delta g_{\nu(t-s)} * (\nabla \cdot \eta(t) (\Delta_1 \tau(s, t) + \Delta_2 \tau(s, t))) \, ds \tag{107}$$

where $\Delta_1 \tau$ and $\Delta_2 \tau$ are the same as (59). From the same arguments from the above,

$$\begin{aligned}
&\left\| \int_0^t \Delta g_{\nu(t-s)} * (\nabla \cdot \eta(t) \Delta_1 \tau(s, t)) \, ds \right\|_{\alpha,p} \\
&\leq \frac{Ct}{\nu} \|\eta\|_{L^\infty(0,T;C^{1+\alpha})} \|\tau\|_{Lip(0,T;C^{\alpha,p})} M_X^\alpha.
\end{aligned} \tag{108}$$

On the other hand,

$$\begin{aligned} \Delta g_{\nu(t-s)} * (\nabla \cdot \eta(t) \Delta_2 \tau(s, t))(x) &= \int_{\mathbb{R}^d} (K(x, z, t, s) (\nabla \cdot \eta)(X(z, t), t) \\ &\quad + \Delta g_{\nu(t-s)}(x - X(z, t)) ((\nabla \cdot \eta)(X(z, s), t) - (\nabla \cdot \eta)(X(z, t), t))) dz, \end{aligned} \quad (109)$$

where K is as in (63). Then as in the proof of Lemma 3, by the generalized Young's inequality we have

$$\begin{aligned} \left\| \int_0^t \Delta g_{\nu(t-s)} * (\nabla \cdot \eta(t) \Delta_2 \tau(s, t)) ds \right\|_{L^p \cap L^\infty} &\leq C \|\tau(t)\|_{L^p \cap L^\infty} \|\eta\|_{L^\infty(0, T; C^{1+\alpha})} \\ \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} \left(\frac{t^\alpha}{\nu^\alpha} + \left(\frac{t}{\nu^3} \right)^{\frac{1}{2}} + \frac{t^2}{\nu^3} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^3 \right). \end{aligned} \quad (110)$$

For the Hölder seminorm, we repeat the same argument in the proof of Theorem 5, using the bound (81). Then we obtain

$$\begin{aligned} &\frac{1}{|h|^\alpha} \left\| \delta_h \left(\int_0^t \Delta g_{\nu(t-s)} * \Delta_2 \tau(s, t) ds \right) \right\|_{L^\infty} \\ &\leq \frac{C(\alpha)}{\nu} \left(1 + \left(\frac{t}{\nu} \right)^{\frac{1}{2}} + \left(\frac{t}{\nu} \right)^2 \right) \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\tau\|_{L^\infty(0, T; C^{\alpha, p})} \|\eta\|_{L^\infty(0, T; C^{1+\alpha})}. \end{aligned} \quad (111)$$

Therefore,

$$\begin{aligned} \|I_5\|_{L^\infty(0, T; C^{\alpha, p})} &\leq \frac{C(\alpha)}{\nu} \left(1 + t + \left(\frac{t}{\nu} \right)^2 \right) \left(1 + \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} \right)^3 M_X^{1+2\alpha} \\ &\quad \|X'\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{Lip(0, T; C^{\alpha, p})}. \end{aligned} \quad (112)$$

The term $I_4(t)$ is treated in the exactly same way, by noting that

$$\begin{aligned} \nabla \cdot (\eta(s) - \eta(t)) &= \nabla_x X^{-1}(s) : (\Delta_1 \nabla_a X'(s, t)) + \nabla_x X^{-1}(s) : (\Delta_2 \nabla_a X'(s, t)) \\ &\quad + (\nabla_x X^{-1}(s) - \nabla_x X^{-1}(t)) : (\nabla_a X' \circ X^{-1})(t), \end{aligned} \quad (113)$$

where as in (59)

$$\begin{aligned} \Delta_1 \nabla_a X'(x, s, t) &= \nabla_a X'(X^{-1}(x, s), s) - \nabla_a X'(X^{-1}(x, s), t), \\ \Delta_2 \nabla_a X'(x, s, t) &= \nabla_a X'(X^{-1}(x, s), t) - \nabla_a X'(X^{-1}(x, t), t), \end{aligned} \quad (114)$$

and

$$\nabla_x (X^{-1}(x, s) - X^{-1}(x, t)) = (\nabla_a X \circ X^{-1})(x, t) (\nabla_a (X - \text{Id}))(X^{-1}(x, t), t - s) \quad (115)$$

so that

$$\|\nabla_x X^{-1}(s) - \nabla_x X^{-1}(t)\|_{C^\alpha} \leq |t - s| \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} M_X^{1+2\alpha}. \quad (116)$$

Also note that

$$\|\Delta_2 \nabla_a X'(s, t)\|_{L^\infty} \leq \|\nabla_a X'(t)\|_{C^\alpha} \|X - \text{Id}\|_{Lip(0, T; L^\infty)}^\alpha |t - s|^\alpha \quad (117)$$

so that

$$\begin{aligned} &\left\| \int_0^t \Delta g_{\nu(t-s)} * (\nabla_x X^{-1}(s) : (\Delta_2 \nabla_a X'(s, t)) \tau \circ X^{-1}(s)) ds \right\|_{C^{\alpha, p}} \\ &\leq \frac{C(\alpha)}{\nu} \left(1 + t^\alpha + \left(\frac{t}{\nu} \right)^2 \right) M_X^{1+2\alpha} \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \\ &\quad \|X'\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{L^\infty(0, T; C^{\alpha, p})}. \end{aligned} \quad (118)$$

The final result is

$$\begin{aligned} \|I_4(t)\|_{\alpha, p} &\leq \frac{C(\alpha)}{\nu} \left(1 + t + \left(\frac{t}{\nu} \right)^2 \right) M_X^{2+4\alpha} \|X'\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{L^\infty(0, T; C^{\alpha, p})} \\ &\quad + C \frac{t}{\nu} M_X^{1+3\alpha} \|X'\|_{Lip(0, T; C^{1+\alpha})} \|\tau\|_{L^\infty(0, T; C^{\alpha, p})}. \end{aligned} \quad (119)$$

Finally, I_2 can be bounded using the combination of the technique in Theorem 5 and Theorem 6. First, we have

$$\begin{aligned} I_2(x, t) = & \int_0^t \int_{\mathbb{R}^d} \nabla \Delta g_{\nu(t-s)}(y) \cdot y \cdot \left(\int_0^1 \nabla \eta(x - (1-\lambda)y, t) d\lambda (\Delta_1 \tau(x - y, s, t)) \right) dy ds \\ & + \int_0^t \int_{\mathbb{R}^d} \nabla \Delta g_{\nu(t-s)}(x - z) \cdot (x - z) \cdot \left(\int_0^1 \nabla \eta(\lambda x + (1-\lambda)z, t) d\lambda (\Delta_2 \tau(z, s, t)) \right) dz ds. \end{aligned} \quad (120)$$

Then applying the argument of the proof of Theorem 6, the first term is bounded by

$$\frac{C}{\nu} t M_X^\alpha \|\eta\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{Lip(0, T; C^{\alpha, p})}. \quad (121)$$

The second term is treated using the method used in Theorem 5. By changing variables to form a kernel similar to (63), and applying generalized Young's inequality, the $L^p \cap L^\infty$ norm of the second term is bounded by

$$\frac{C(\alpha)}{\nu} \left(t^\alpha + \left(\frac{t}{\nu} \right)^{\frac{1}{2}} + \left(\frac{t}{\nu} \right)^2 \right) \left(1 + \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} \right)^4 \|\eta\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{L^\infty(0, T; L^p \cap L^\infty)}. \quad (122)$$

Finally, the Hölder seminorm of the second term is bounded by the same method as Theorem 5. The only additional point is the finite difference of $\nabla \eta$ term, but this term is bounded by a straightforward estimate. The bound for the Hölder seminorm of the second term is

$$\frac{C(\alpha)}{\nu} \left(1 + t^\alpha + \left(\frac{t}{\nu} \right)^{\frac{1}{2}} + \left(\frac{t}{\nu} \right)^2 \right) \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}^\alpha \|\eta\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{L^\infty(0, T; C^{\alpha, p})}. \quad (123)$$

To sum up, we have

$$\begin{aligned} \|I_2(t)\|_{\alpha, p} \leq & \frac{C(\alpha)}{\nu} \left(1 + t + \left(\frac{t}{\nu} \right)^2 \right) \left(1 + \|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})} \right)^4 M_X^{1+3\alpha} \\ & \|X'\|_{L^\infty(0, T; C^{1+\alpha})} \|\tau\|_{Lip(0, T; C^{\alpha, p})}. \end{aligned} \quad (124)$$

If we put this together,

$$\begin{aligned} & \| [X' \circ X^{-1} \cdot \nabla, \mathbb{G}] (\tau \circ X^{-1}) \|_{L^\infty(0, T; C^{\alpha, p})} \\ & \leq C \|X'\|_{L^\infty(0, T; C^{1+\alpha})} M_X^{1+2\alpha} \|\Gamma(\tau \circ X^{-1})\|_{L^\infty(0, T; C^{\alpha, p})} \\ & + (\|X'\|_{L^\infty(0, T; C^{1+\alpha})} + \|X'\|_{Lip(0, T; C^{1+\alpha})} T^{\frac{1}{2}}) F_1(\nu, \alpha, X, \|\tau\|_{Lip(0, T; C^{\alpha, p})}, T) \end{aligned} \quad (125)$$

where F_1 depends on the written variables and grows like polynomial in T , $\|\tau\|_{Lip(0, T; C^{\alpha, p})}$, and $\|X - \text{Id}\|_{Lip(0, T; C^{1+\alpha})}$. The bound on $\Gamma(\tau \circ X^{-1})$ is given by Theorem 5. \square

4 Bounds on variations and variables

Using the results from the previous section we find bounds for variations and variables. For simplicity, we adopt the notation

$$M_\epsilon = 1 + \|X_\epsilon - \text{Id}\|_{L^\infty(0, T; C^{1+\alpha})}. \quad (126)$$

First, we bound $\frac{d}{d\epsilon} \mathcal{V}_\epsilon$. Note that $X_\epsilon(0) = \text{Id}$, so $X'_\epsilon(0) = 0$ and by Theorem 2 and since $X'_\epsilon \in Lip(0, T; C^{1+\alpha, p})$ we have

$$\begin{aligned} \|X'_\epsilon\|_{L^\infty(0, T; C^{1+\alpha})} & \leq T \|X'_\epsilon\|_{Lip(0, T; C^{1+\alpha, p})}, \\ \|\eta_\epsilon(t)\|_{C^\alpha} & \leq t \|X'\|_{Lip(0, T; C^{1+\alpha, p})} M_\epsilon^\alpha. \end{aligned} \quad (127)$$

Then by the Theorem 3, we have

$$\begin{aligned} \|\eta_\epsilon \cdot \mathbb{L}_\nu(\nabla_x u_{\epsilon,0})\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \left(\frac{T}{\nu}\right)^{\frac{1}{2}} M_\epsilon^\alpha \|X'_\epsilon\|_{Lip(0,T;C^{1+\alpha,p})} \|u_{\epsilon,0}\|_{1+\alpha,p}, \\ \|\mathbb{L}_\nu(u'_{\epsilon,0})\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \|u'_{\epsilon,0}\|_{\alpha,p}. \end{aligned} \quad (128)$$

By Theorem 6, we have

$$\begin{aligned} \|\eta_\epsilon \cdot \nabla_x \mathbb{U} (\sigma_\epsilon - u_\epsilon \otimes u_\epsilon)\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \left(\left(\frac{T}{\nu}\right)^{\frac{1}{2}} + \left(\frac{T}{\nu}\right)\right) M_\epsilon^{2+4\alpha} \\ \|X'_\epsilon\|_{Lip(0,T;C^{1+\alpha})} \|\tau_\epsilon - v_\epsilon \otimes v_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})}, \end{aligned} \quad (129)$$

and by Theorem 4, we have

$$\begin{aligned} \|\mathbb{U}(\delta_\epsilon - (v'_\epsilon \otimes v_\epsilon + v_\epsilon \otimes v'_\epsilon) \circ X_\epsilon^{-1})\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \left(\frac{T}{\nu}\right)^{\frac{1}{2}} M_\epsilon^\alpha \\ \|\tau'_\epsilon - (v'_\epsilon \otimes v_\epsilon + v_\epsilon \otimes v'_\epsilon)\|_{L^\infty(0,T;C^{\alpha,p})}. \end{aligned} \quad (130)$$

Therefore,

$$\begin{aligned} \left\| \frac{d}{d\epsilon} \mathcal{V}_\epsilon \right\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C \|u'_{\epsilon,0}\|_{\alpha,p} \\ + S_1(T) (\|X'_\epsilon\|_{Lip(0,T;C^{1+\alpha,p})} + \|v'_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} + \|\sigma'_{\epsilon,0}\|_{\alpha,p} + \|\tau'_\epsilon\|_{Lip(0,T;C^{\alpha,p})}) Q_1 \end{aligned} \quad (131)$$

where $S_1(T)$ vanishes as $T^{\frac{1}{2}}$ as $T \rightarrow 0$ and Q_1 is a polynomial in $\|u_{\epsilon,0}\|_{1+\alpha,p}$, $\|X_\epsilon - \text{Id}\|_{Lip(0,T;C^{1+\alpha,p})}$, $\|\tau_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})}$, and $\|v_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})}$, whose coefficients depend on ν . Similarly,

$$\|g_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} \leq M_X^\alpha \|u_0\|_{1+\alpha,p} + C_1 \|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})}^\alpha \|\sigma_{\epsilon,0}\|_{\alpha,p} + S_2(T) Q_2, \quad (132)$$

where $S_2(T)$ vanishes as $T^{\frac{1}{2}}$ as $T \rightarrow 0$ and Q_2 is polynomial in $\|\tau\|_{Lip(0,T;C^{\alpha,p})}$ and $\|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})}$, whose coefficients depend on α and ν . Also

$$\begin{aligned} \|g'_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} &\leq C (\|u'_{\epsilon,0}\|_{1+\alpha,p} + \|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha})}^\alpha \|\tau'_{\epsilon,0}\|_{\alpha,p}) \\ + S_3(T) (\|X'_\epsilon\|_{Lip(0,T;C^{1+\alpha,p})} + \|\sigma'_{\epsilon,0}\|_{\alpha,p} + \|\tau'_\epsilon\|_{Lip(0,T;C^{\alpha,p})} + \|v'_\epsilon\|_{L^\infty(0,T;C^{1+\alpha,p})}) Q_3, \end{aligned} \quad (133)$$

where $S_3(T)$ vanishes as $T^{\frac{1}{2}}$ as $T \rightarrow 0$ and Q_3 is polynomial in $\|u_{\epsilon,0}\|_{1+\alpha,p}$, $\|X - \text{Id}\|_{Lip(0,T;C^{1+\alpha,p})}$, $\|\tau\|_{Lip(0,T;C^{\alpha,p})}$, and $\|v_\epsilon\|_{L^\infty(0,T;C^{1+\alpha,p})}$, whose coefficients depend on ν and α . Then we have

$$\left\| \nabla_a \frac{d}{d\epsilon} \mathcal{V}_\epsilon \right\|_{L^\infty(0,T;C^{\alpha,p})} \leq T \|X'_\epsilon\|_{Lip(0,T;C^{1+\alpha})} \|g_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} + M_\epsilon \|g'_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} \quad (134)$$

and

$$\begin{aligned} \left\| \frac{d}{d\epsilon} \mathcal{T}_\epsilon \right\|_{L^\infty(0,T;C^{\alpha,p})} &\leq 2 \|g'_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} \left(\|\tau_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} + 2\rho K \right) \\ + \|\tau'_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} \left(\|g_\epsilon\|_{L^\infty(0,T;C^{\alpha,p})} + 2k \right). \end{aligned} \quad (135)$$

5 Local existence

We define the function space \mathcal{P}_1 and the set \mathcal{I} ,

$$\begin{aligned} \mathcal{P}_1 &= Lip(0,T;C^{1+\alpha,p}) \times Lip(0,T;C^{\alpha,p}) \times L^\infty(0,T;C^{1+\alpha,p}) \\ \mathcal{I} &= \{(X, \tau, v) : \|(X - \text{Id}, \tau, v)\|_{\mathcal{P}_1} \leq \Gamma, v = \frac{dx}{dt}\}, \end{aligned} \quad (136)$$

where $\Gamma > 0$ and $T > 0$ are to be determined. Now, for given $u_0 \in C^{1+\alpha,p}$ divergence free and $\sigma_0 \in C^{\alpha,p}$ we define the map

$$(X, \tau, v) \rightarrow \mathcal{S}(X, \tau, v) = (X^{new}, \tau^{new}, v^{new}) \quad (137)$$

where

$$\begin{cases} X^{new}(t) = \text{Id} + \int_0^t \mathcal{V}(X(s), \tau(s), v(s)) ds, \\ \tau^{new}(t) = \sigma_0 + \int_0^t \mathcal{T}(X(s), \tau(s), v(s)) ds, \\ v^{new}(t) = \mathcal{V}(X, \tau, v). \end{cases} \quad (138)$$

If $(X - \text{Id}, \tau, v) \in \mathcal{P}_1$, then $(X^{new} - \text{Id}, \tau^{new}, v^{new}) \in \mathcal{P}_1$ for any choice of $T > 0$. Moreover, we have the following:

Theorem 8. *For given $u_0 \in C^{1+\alpha,p}$ divergence free and $\sigma_0 \in C^{\alpha,p}$, there is a $\Gamma > 0$ and $T > 0$ such that the map \mathcal{S} of (138) maps \mathcal{I} to itself.*

Proof. It is obvious that $\frac{d}{dt} X^{new} = v^{new}$. For the size of $\mathcal{S}(X, \tau, v)$, first note that if $(X - \text{Id}, \tau, v)_{\mathcal{P}_1} \leq \Gamma$, then

$$M_X = 1 + \|X - \text{Id}\|_{L^\infty(0,T;C^{1+\alpha})} \leq 1 + TT. \quad (139)$$

Applying Theorem 3 and Theorem 4, we know that

$$\|\mathcal{V}\|_{L^\infty(0,T;C^{\alpha,p})} \leq \|u_0\|_{\alpha,p} + A_1(T)B_1(\Gamma, \|u_0\|_{\alpha,p}, \|\sigma_0\|_{\alpha,p}), \quad (140)$$

where $A_1(T)$ vanishes like $T^{\frac{1}{2}}$ for small $T > 0$ and B_1 is a polynomial in its arguments, and some coefficients depend on ν . We estimate

$$\|g\|_{L^\infty(0,T;C^{\alpha,p})} \leq \|u_0\|_{1+\alpha,p} + C_1\Gamma^\alpha \|\sigma_0\|_{\alpha,p} + A_2(T)B_2(\Gamma, \|u_0\|_{1+\alpha,p}, \|\sigma_0\|_{\alpha,p}), \quad (141)$$

where C_1 is as in Theorem 5, depending only on α and ν , $A_2(T)$ vanishes in the same order as $A_1(T)$ as $T \rightarrow 0$, and B_2 is a polynomial in its arguments, and some coefficients depend on ν and α . From (24) we conclude

$$\|\mathcal{V}\|_{L^\infty(0,T;C^{1+\alpha,p})} \leq K_1(\|u_0\|_{1+\alpha,p} + \Gamma^\alpha \|\sigma_0\|_{\alpha,p}) + A_3(T)B_3(\Gamma, \|u_0\|_{1+\alpha,p}, \|\sigma_0\|_{\alpha,p}), \quad (142)$$

where K_1 is a constant depending only on ν and α , and A_3 and B_3 have the same properties as previous A_i s and B_i s. Now we measure \mathcal{T} . From (84) and the previous estimate on g we have

$$\begin{aligned} \|\mathcal{T}\|_{L^\infty(0,T;C^{\alpha,p})} &\leq K_2(\|u_0\|_{1+\alpha,p}(\rho K + \|\sigma_0\|_{\alpha,p}) + \|\sigma_0\|_{\alpha,p}(\Gamma^\alpha \|\sigma_0\|_{\alpha,p} + \rho K \Gamma^\alpha + k)) \\ &\quad + A_4 B_4, \end{aligned} \quad (143)$$

where K_2 is a constant depending on ν and α , and A_4 and B_4 are as before. Since $\alpha < 1$, we can appropriately choose large $\Gamma > \|\sigma_0\|_{\alpha,p} + \|u_0\|_{1+\alpha,p}$ and correspondingly small $\frac{1}{6} > T > 0$ so that the right side of (142) and (143) are bounded by $\frac{\Gamma}{6}$. Then $\|(X^{new} - \text{Id}, \tau^{new}, v^{new})\|_{\mathcal{P}_1} \leq \Gamma$. \square

We show now that \mathcal{S} is a contraction mapping on \mathcal{I} for a short time.

Theorem 9. *For given $u_0 \in C^{1+\alpha,p}$ divergence free and $\sigma_0 \in C^{\alpha,p}$, there is a Γ and $T > 0$, depending only on $\|u_0\|_{1+\alpha,p}$ and $\|\sigma_0\|_{\alpha,p}$, such that the map \mathcal{S} is a contraction mapping on $\mathcal{I} = \mathcal{I}(\Gamma, T)$, that is*

$$\|\mathcal{S}(X_2, \tau_2, v_2) - \mathcal{S}(X_1, \tau_1, v_1)\|_{\mathcal{P}_1} \leq \frac{1}{2} \|(X_2 - X_1, \tau_2 - \tau_1, v_2 - v_1)\|_{\mathcal{P}_1}. \quad (144)$$

Proof. First from Theorem 8 we can find a Γ and $T_0 > 0$, depending only on the size of initial data, say

$$N = \max\{\|u_0\|_{1+\alpha,p}, \|\sigma_0\|_{\alpha,p}\}, \quad (145)$$

which guarantees that \mathcal{S} maps \mathcal{I} to itself. This property still holds if we replace T_0 by any smaller $T > 0$. In view of the fact that \mathcal{I} is convex, we put

$$\begin{aligned} X_\epsilon &= (2 - \epsilon)X_1 + (\epsilon - 1)X_2, \\ \tau_\epsilon &= (2 - \epsilon)\tau_1 + (\epsilon - 1)\tau_2, 1 \leq \epsilon \leq 2. \end{aligned} \quad (146)$$

Then $(X_\epsilon, \tau_\epsilon, v_\epsilon) \in \mathcal{I}$, $v_\epsilon = (2 - \epsilon)v_1 + (\epsilon - 1)v_2$, $u_{\epsilon,0} = u_0$, and $\sigma_{\epsilon,0} = \sigma_0$. This means that

$$X'_\epsilon = X_2 - X_1, v'_\epsilon = v_2 - v_1, u'_{\epsilon,0} = 0, \sigma'_{\epsilon,0} = 0. \quad (147)$$

Then from the results of Section 4, we see that

$$\begin{aligned} \left\| \frac{d}{d\epsilon} \mathcal{V}_\epsilon \right\|_{L^\infty(0,T;C^{1+\alpha,p})} &\leq (\|X_2 - X_1\|_{Lip(0,T;C^{1+\alpha,p})} + \|v_2 - v_1\|_{L^\infty(0,T;C^{\alpha,p})} \\ &\quad + \|\tau_2 - \tau_1\|_{Lip(0,T;C^{\alpha,p})}) S'_1(T) Q'_1(\Gamma), \\ \|\mathcal{X}'_\epsilon\|_{Lip(0,T;C^{1+\alpha,p})} &\leq (\|X_2 - X_1\|_{Lip(0,T;C^{1+\alpha,p})} + \|v_2 - v_1\|_{L^\infty(0,T;C^{\alpha,p})} \\ &\quad + \|\tau_2 - \tau_1\|_{Lip(0,T;C^{\alpha,p})}) S'_2(T) Q'_2(\Gamma), \end{aligned} \quad (148)$$

$$\begin{aligned} \|\pi_\epsilon\|_{Lip(0,T;C^{\alpha,p})} &\leq (\|X_2 - X_1\|_{Lip(0,T;C^{1+\alpha,p})} + \|v_2 - v_1\|_{L^\infty(0,T;C^{\alpha,p})} \\ &\quad + \|\tau_2 - \tau_1\|_{Lip(0,T;C^{\alpha,p})}) S'_3(T) Q'_3(\Gamma), \end{aligned}$$

where \mathcal{X}'_ϵ and π_ϵ are defined in (20), $S'_1(T), S'_2(T), S'_3(T)$ vanish at the rate of $T^{\frac{1}{2}}$ as $T \rightarrow 0$, and $Q'_1(\Gamma), Q'_2(\Gamma), Q'_3(\Gamma)$ are polynomials in Γ , whose coefficients depend only on ν and α . By choosing $0 < T < T_0$ small enough, depending on the size of $Q'_i(\Gamma)$ s, we conclude the proof. \square

We have obtained a solution to the system (6) in the path space \mathcal{P}_1 for a short time, that is, we have (X, τ, v) satisfying $v = \frac{dX}{dt}$ and satisfying (14). We also have Lipschitz dependence on initial data, Theorem 1.

Proof. We repeat the calculation of the Theorem 9, but this time $u'_{\epsilon,0} = u_1(0) - u_2(0)$ and $\sigma'_{\epsilon,0} = \sigma_1(0) - \sigma_2(0)$. Then we choose T_0 small enough that $S'_i(T_0)Q'_1(\Gamma) < \frac{1}{2}$. \square

Acknowledgements The research of P.C. was partially supported by NSF grant DMS-1713985. The research of J.L. was partially supported by a Samsung scholarship.

References

- [1] P. Constantin. Lagrangian-Eulerian methods for uniqueness in hydrodynamic systems *Adv. Math.*, 278:67-102, 2015.
- [2] P. Constantin. *Analysis of hydrodynamic models*, volume 90 of *CBMS-NSF Regional Conference Series in Applied Mathematics*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2017.