

Mitochondrial DNA Part B

Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete mitogenome of the invasive Japanese mud snail *Batillaria attramentaria* (Gastropoda: Batillariidae) from Elkhorn Slough, California, USA

Hartnell College Genomics Group, Paulina Andrade, Lisbeth Arreola, Melissa Belnas, Estefania Bland, Araceli Castillo, Omar Cisneros, Valentin Contreras, Celeste Diaz, Kevin T. Do, Carlos Donate, Estevan Espinoza, Nathan Frater, Garry G. Gabriel, Eric A. Gomez, Gino F. Gonzalez, Myrka Gonzalez, Paola Guido, Dylan Guidotti, Mishell Guzman Espinoza, Ivan Haro, Javier Hernandez Lopez, Caden E. Hernandez, Karina Hernandez, Jazmin A. Hernandez-Salazar, Jeffery R. Hughey, Héctor Jácome-Sáenz, Luis A. Jimenez, Eli R. Kallison, Mylisa S. King, Luis J. Lazaro, Feifei Zhai Lorenzo, Isaac Madrigal, Savannah Madruga, Adrian J. Maldonado, Alexander M. Medina, Marcela Mendez-Molina, Ali Mendez, David Murillo Martinez, David Orozco, Juan Orozco, Ulises Ortiz, Jennifer M. Pantoja, Alejandra N. Ponce, Angel R. Ramirez, Israel Rangel, Eliza Rojas, Adriana Roque, Beatriz Rosas, Colt Rubbo, Justin A. Saldana, Eilan Sanchez, Alicia Steinhardt, Maria O. Taveras Dina, Judith Torres, Silvestre Valdez-Mata, Valeria Vargas, Paola Vazquez, Michelle M. Vazquez, Irene Vidales, Frances L. Wong, Christian S. Zagal, Santiago Zamora & Jesus Zepeda Amador

To cite this article: Hartnell College Genomics Group, Paulina Andrade, Lisbeth Arreola, Melissa Belnas, Estefania Bland, Araceli Castillo, Omar Cisneros, Valentin Contreras, Celeste Diaz, Kevin T. Do, Carlos Donate, Estevan Espinoza, Nathan Frater, Garry G. Gabriel, Eric A. Gomez, Gino F. Gonzalez, Myrka Gonzalez, Paola Guido, Dylan Guidotti, Mishell Guzman Espinoza, Ivan Haro, Javier Hernandez Lopez, Caden E. Hernandez, Karina Hernandez, Jazmin A. Hernandez-Salazar, Jeffery R. Hughey, Héctor Jácome-Sáenz, Luis A. Jimenez, Eli R. Kallison, Mylisa S. King, Luis J. Lazaro, Feifei Zhai Lorenzo, Isaac Madrigal, Savannah Madruga, Adrian J. Maldonado, Alexander M. Medina, Marcela Mendez-Molina, Ali Mendez, David Murillo Martinez, David Orozco, Juan Orozco, Ulises Ortiz, Jennifer M. Pantoja, Alejandra N. Ponce, Angel R. Ramirez, Israel Rangel, Eliza Rojas, Adriana Roque, Beatriz Rosas, Colt Rubbo, Justin A. Saldana, Eilan Sanchez, Alicia Steinhardt, Maria O. Taveras Dina, Judith Torres, Silvestre Valdez-Mata, Valeria Vargas, Paola Vazquez, Michelle M. Vazquez, Irene Vidales, Frances L. Wong, Christian S. Zagal, Santiago Zamora & Jesus Zepeda Amador (2019) The complete mitogenome of the invasive Japanese mud snail *Batillaria attramentaria* (Gastropoda: Batillariidae) from Elkhorn Slough, California, USA, Mitochondrial DNA Part B, 4:2, 4031-4032, DOI: [10.1080/23802359.2019.1688719](https://doi.org/10.1080/23802359.2019.1688719)

To link to this article: <https://doi.org/10.1080/23802359.2019.1688719>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 13 Nov 2019.



Submit your article to this journal



Article views: 528



View related articles



CrossMark View Crossmark data

The complete mitogenome of the invasive Japanese mud snail *Batillaria attramentaria* (Gastropoda: Batillariidae) from Elkhorn Slough, California, USA

Hartnell College Genomics Group*, Paulina Andrade, Lisbeth Arreola, Melissa Belnas, Estefania Bland, Araceli Castillo, Omar Cisneros, Valentin Contreras, Celeste Diaz, Kevin T. Do, Carlos Donate, Estevan Espinoza, Nathan Frater, Garry G. Gabriel, Eric A. Gomez, Gino F. Gonzalez, Myrka Gonzalez, Paola Guido, Dylan Guidotti, Mishell Guzman Espinoza, Ivan Haro, Javier Hernandez Lopez, Caden E. Hernandez, Karina Hernandez, Jazmin A. Hernandez-Salazar, Jeffery R. Hughey , Héctor Jácome-Sáenz, Luis A. Jimenez, Eli R. Kallison, Mylisa S. King, Luis J. Lazaro, Feifei Zhai Lorenzo, Isaac Madrigal, Savannah Madruga, Adrian J. Maldonado, Alexander M. Medina, Marcela Mendez-Molina, Ali Mendez, David Murillo Martinez, David Orozco, Juan Orozco, Ulises Ortiz, Jennifer M. Pantoja, Alejandra N. Ponce, Angel R. Ramirez, Israel Rangel, Eliza Rojas, Adriana Roque, Beatriz Rosas, Colt Rubbo, Justin A. Saldana, Erian Sanchez, Alicia Steinhardt, Maria O. Taveras Dina, Judith Torres, Silvestre Valdez-Mata, Valeria Vargas, Paola Vazquez, Michelle M. Vazquez, Irene Vidales, Frances L. Wong, Christian S. Zagal, Santiago Zamora and Jesus Zepeda Amador

Division of Mathematics, Science, and Engineering, Hartnell College, Salinas, CA, USA

ABSTRACT

Genomic analysis of the invasive marine snail *Batillaria attramentaria* from Elkhorn Slough, Moss Landing, California, USA using 150 bp paired-end Illumina sequences resulted in the assembly of its complete mitogenome. The mitogenome is 16,095 bp in length and contains 2 rRNA, 13 protein-coding, and 22 tRNA genes (GenBank Accession MN557850). Gene content and organization of *B. attramentaria* are identical to the Turritellidae and Pachychilidae. The phylogenetic analysis of *B. attramentaria* resolves it in a fully supported clade with these same two families in the superfamily Cerithioidea. Nucleotide BLAST searches of the Elkhorn Slough *cox1* gene of *B. attramentaria* yielded identical sequences from invasive populations from California and British Columbia, and native populations from northeastern and central Japan. These data show that mitogenome sequencing is a useful tool for studying the classification and phylogenetic history Cerithioidea.

ARTICLE HISTORY

Received 22 October 2019
Accepted 27 October 2019

KEYWORDS

Batillaria attramentaria; Cerithioidea; Elkhorn Slough; invasive species; mitogenome

The Batillariidae are aquatic snails that occur along coastal intertidal and shallow estuarine lagoons. The family contains four genera *sensu stricto* and 14 species (Ozawa et al. 2009). One of these genera, *Batillaria*, includes a highly invasive marine snail *B. attramentaria* (Byers 1999, 2000; Miura et al. 2006). *Batillaria* was first introduced in March of 1930 into Elkhorn Slough, California, USA in a shipment of 150 boxes of oysters that arrived from Japan (Bonnot 1935). This shipment and perhaps others that followed inadvertently introduced *B. attramentaria* and more than 50 other invertebrates into the slough (Wasson et al. 2001). The population of *B. attramentaria* in Elkhorn Slough was estimated at 1 billion, and in California, it was responsible for displacing the native snail *Cerithidea californica* (Byers 1999). Here, we performed high-throughput sequencing on a specimen of *B. attramentaria* from Elkhorn Slough to determine its mitogenome structure and phylogenomic relationship to other gastropods in the superfamily Cerithioidea.

DNA was extracted from *B. attramentaria* (Voucher Specimen-Hartnell College #263) following the protocol of Lindstrom et al. (2011). The 150 bp PE Illumina library construction and sequencing was performed by myGenomics, LLC (Alpharetta, Georgia, USA). The mitogenome was assembled using 22,355,422 reads with the default de novo settings in MEGAHIT (Li et al. 2015). The genes were annotated with NCBI ORFfinder and MITOS (Bernt et al. 2013). Alignment of the *B. attramentaria* mitogenome to other Gastropoda was completed with MAFFT (Katoh and Standley 2013). The RAxML analysis was run using T-REX (Boc et al. 2012) with the GTR+gamma model and 1000 bootstraps. The tree was visualized with TreeDyn 198.3 at Phylogeny.fr (Dereeper et al. 2008).

The mitogenome of *B. attramentaria* (GenBank MN557850) is 16,095 bp in length and has a base composition of 30.14% A, 35.24% T, 17.01% G, and 17.61% C. It contains 2 rRNA (*rnl*, *rns*), 13 electron transport and oxidative phosphorylation

CONTACT Jeffery R. Hughey   jhughey@hartnell.edu  Division of Mathematics, Science, and Engineering, Hartnell College, 411 Central Ave., Salinas, CA 93901, USA

*All authors contributed equally to the analysis and writing of this paper.

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Maximum-likelihood phylogram of *Batillaria attramentaria* and related gastropod mitogenomes. Superfamilies are listed to the right. Numbers along branches are bootstrap supports based on 1000 nreps. The legend below represents the scale for nucleotide substitutions.

genes, and 22 tRNAs (*tRNA-Leu* and *tRNA-Ser* are duplicated). All 13 protein-coding genes initiate with the ATG codon. Most of the genes terminate with the TAA codon, however, *cox2*, *nad1*, and *nad2* terminate with TAG. The mitogenome of *B. attramentaria* is identical in gene content and organization to *Turritella* (Family Turritellidae) and *Tylomelania* (Pachychilidae) (Hilgers et al. 2016; Zeng et al. 2016). These three families have tandemly arranged *tRNA-Arg* and *tRNA-Gln* genes. In other published Cerithioidea mitogenomes, *tRNA-Gln* is situated on the chromosome with *tRNA-Ser*, and *tRNA-Arg* is positioned in between *tRNA-Cys* and *tRNA-Ala*. Phylogenetic analysis of *B. attramentaria* resolves it in a fully supported clade with *Turritella* and *Tylomelania* in the superfamily Cerithioidea (Figure 1). Strong et al. (2011) previously reported a similar phylogenetic relationship between the Batillariidae, Turritellidae, and Pachychilidae based on 16S and 28S rRNA DNA sequences. Analysis of the *B. attramentaria* *cox1* sequence found identical sequences from invasive specimens from Elkhorn Slough, California and British Columbia, Canada, and native specimens from northeastern and central Japan (Kojima et al. 2004; Miura et al. 2006; Castelin et al. 2016).

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

Funding

This research was supported by NSF award number [1832446] to Hartnell College.

ORCID

Jeffery R. Hughey <http://orcid.org/0000-0003-4053-9150>

References

- Boc A, Diallo AB, Makarenkov V. 2012. T-REX: a web server for inferring, validating and visualizing phylogenetic trees and networks. Nucleic Acids Res. 40(W1):W573–W579.
- Bonnot P. 1935. A recent introduction of exotic species of molluscs into California waters from Japan. Nautilus. 49:1–2.
- Byers JE. 1999. The distribution of an introduced mollusc and its role in the long-term demise of a native confamilial species. Biol Invasions. 1(4):339–352.
- Byers JE. 2000. Competition between two estuarine snails: implications for invasions of exotic species. Ecology. 81(5):1225–1239.
- Castelin M, Van Steenkiste N, Pante E, Harbo R, Lowe G, Gilmore SR, Therriault TW, Abbott CL. 2016. A new integrative framework for large-scale assessments of biodiversity and community dynamics, using littoral gastropods and crabs of British Columbia, Canada. Mol Ecol Resour. 16(6):1322–1339.
- Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S, Lefort V, Lescot M, et al. 2008. Phylogeny.fr: robust phylogenetic analysis for the non-specialist. Nucleic Acids Res. 36(Web Server):W465–W469.
- Hilgers L, Grau JH, Pfäender J, von Rintelen T. 2016. The complete mitochondrial genome of the viviparous freshwater snail *Tylomelania sasinorum* (Caenogastropoda: Cerithioidea). Mitochondrial DNA B Resour. 1(1):330–331.
- Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: improvements in performance and usability. Mol Biol Evol. 30(4):772–780.
- Kojima S, Hayashi I, Kim D, Iijima A, Eurota T. 2004. Phylogeography of an intertidal direct-developing gastropod *Batillaria cumingi* around the Japanese Islands. Mar Ecol Prog Ser. 276:161–172.
- Li D, Liu CM, Luo R, Sadakane K, Lam TW. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 31(10):1674–1676.
- Lindstrom SC, Hughey JR, Martone PT. 2011. New, resurrected and redefined species of *Mastocarpus* (Phyllophoraceae, Rhodophyta) from the northeast Pacific. Phycologia. 50(6):661–683.
- Miura O, Torchin ME, Kuris AM, Hechinger RF, Chiba S. 2006. Introduced cryptic species of parasites exhibit different invasion pathways. Proc Natl Acad Sci USA. 103(52):19818–19823.
- Ozawa T, Köhler F, Reid DG, Glaubrecht M. 2009. Tethyan relicts on continental coastlines of the northwestern Pacific Ocean and Australasia: molecular phylogeny and fossil record of batillariid gastropods (Caenogastropoda, Cerithioidea). Zool Scr. 38(5):503–525.
- Strong EE, Colgan DJ, Healy JM, Lydeard C, Ponder WF, Glaubrecht M. 2011. Phylogeny of the gastropod superfamily Cerithioidea using morphology and molecules. Zool J Linn Soc. 162(1):43–89.
- Wasson K, Zabin CJ, Bedinger L, Diaz MC, Pearse JS. 2001. Biological invasions of estuaries without international shipping: the importance of intraregional transport. Biol Conserv. 102(2):143–153.
- Zeng L, Wang Y, Zhang J, Wu C. 2016. Complete mitochondrial genome of *Turritella terebra bacillum*. Mitochondrial DNA B Resour. 1(1):350–351.