THE HABITABLE-ZONE PLANET FINDER REVEALS A HIGH MASS AND A LOW OBLIQUITY FOR THE YOUNG NEPTUNE K2-25B

Gudmundur Stefansson^{1,2,3,4,5}, Suvrath Mahadevan^{1,2}, Marissa Maney¹, Joe P. Ninan^{1,2}, Paul Robertson⁶, Jayadev Rajagopal⁷, Flynn Haase⁷, Lori Allen⁷, Eric B. Ford^{1,2,8}, Joshua Winn⁴, Angie Wolfgang^{1,2}, Rebekah I. Dawson^{1,2,9}, John Wisniewski¹⁰, Chad F. Bender¹¹, Caleb Cañas^{1,2,3}, William Cochran¹², Scott A. Diddams^{13,14}, Connor Fredrick^{13,14}, Samuel Halverson¹⁵, Fred Hearty^{1,2}, Leslie Hebb¹⁶, Shubham Kanodia^{1,2}, Eric Levi¹, Andrew J. Metcalf^{17,13,14}, Andrew Monson^{1,2}, Lawrence Ramsey^{1,2}, Arpita Roy^{18,19}, Christian Schwab²⁰, Ryan Terrien²¹, and Jason T. Wright^{1,2}

ABSTRACT

Using radial-velocity data from the Habitable-zone Planet Finder, we have measured the mass of the Neptune-sized planet K2-25b, as well as the obliquity of its M4.5-dwarf host star in the 600-800MYr Hyades cluster. This is one of the youngest planetary systems for which both of these quantities have been measured, and one of the very few M dwarfs with a measured obliquity. Based on a joint analysis of the radial velocity data, time-series photometry from the K2 mission, and new transit light curves obtained with diffuser-assisted photometry, the planet's radius and mass are $3.44 \pm 0.12R_{\oplus}$ and $24.5^{+5.7}_{-5.2}M_{\oplus}$. These properties are compatible with a rocky core enshrouded by a thin hydrogen-helium atmosphere (5% by mass). We measure an orbital eccentricity of $e = 0.43 \pm 0.05$. The sky-projected stellar obliquity is $\lambda = 3 \pm 16^{\circ}$, compatible with spin-orbit alignment, in contrast to other "hot Neptunes" that have been studied around older stars.

Keywords: Exoplanets, Transits, Rossiter-McLaughlin Effect, Radial Velocities

1. INTRODUCTION

The observed orbital properties of planetary systems are influenced both by the formation process as well as subsequent dynamical interactions that can take place after planets are formed. Dynamical interactions over billions of years can

modify or even rearrange planetary orbits, making it difficult to learn about the initial conditions. Young systems have had less time to evolve, and their orbital properties may provide a clearer view of the early stages of planet formation. As such, young systems are valuable benchmarks for testing models of planet formation and evolution.

The *K2* mission (Howell et al. 2014) enabled the detection of many planets in young associations and clusters, including a number of planets discovered by The Zodical Ex-

¹Department of Astronomy & Astrophysics, The Pennsylvania State University, 525 Davey Lab, University Park, PA 16802, USA

²Center for Exoplanets & Habitable Worlds, University Park, PA 16802, USA

³NASA Earth and Space Science Fellow

⁴Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA

⁵Henry Norris Russell Fellow

⁶Department of Physics and Astronomy, University of California-Irvine, Irvine, California 92697, USA

⁷NSF's OIR Lab, 950 N. Cherry Ave. Tucson, AZ 85719, USA

⁸Institute for Computational & Data Sciences, The Pennsylvania State University, 203 Computer Building, University Park, PA 16802, USA

⁹Alfred P. Sloan Foundation Fellow

¹⁰Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, 440 W. Brooks Street, Norman, OK 73019, USA

¹¹Steward Observatory, University of Arizona, Tucson, Arizona 85721, USA

¹²McDonald Observatory and Center for Planetary Systems Habitability, The University of Texas at Austin, Austin TX 78712, USA

¹³Time and Frequency Division, National Institute of Standards and Technology, 325 Broadway, Boulder, CO 80305, USA

¹⁴Department of Physics, University of Colorado, 2000 Colorado Avenue, Boulder, CO 80309, USA

¹⁵ Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

¹⁶Department of Physics, Hobart and William Smith Colleges, 300 Pulteney Street, Geneva, NY 14456, USA

¹⁷Space Vehicles Directorate, Air Force Research Laboratory, 3550 Aberdeen Ave SE, Kirtland AFB, NM 87117, USA

¹⁸Millikan Prize Postdoctoral Fellow

¹⁹California Institute of Technology, 1200 E California Blvd, Pasadena, California 91125, USA

²⁰Department of Physics and Astronomy, Macquarie University, Balaclava Road, North Ryde, NSW 2109, Australia

²¹Department of Physics and Astronomy, Carleton College, Northfield, Minnesota 55057, USA

oplanets In Time (ZEIT) project (e.g., Mann et al. 2016a,b, 2017), and the four newborn transiting planets around V1298 Tau (David et al. 2019). The ongoing Transiting Exoplanet Survey Satellite (TESS) mission (Ricker et al. 2014) is also observing young stars, and has led to the discovery of the 45 Myr old Neptune-sized planet DS Tuc Ab (Newton et al. 2019; Benatti et al. 2019), and the 22 MYr old Neptune orbiting the pre-main-sequence star AU Microscopii (AU Mic) (Playchan et al. 2020). Follow-up spectroscopic observations of both DS Tuc Ab (Zhou et al. 2019; Montet et al. 2019), and AU Mic b (Addison et al. 2020; Hirano et al. 2010; Palle et al. 2020) have revealed that both stars have low obliquities—the angle between its rotation axis and the planet's orbital axis. This is interesting because close-orbiting Neptunes around older stars seem to have a broad range of obliquities (e.g., Winn et al. 2010b; Bourrier et al. 2018b), although the number of measurements is still quite limited.

Both the stellar obliquity and orbital eccentricity are clues about the formation and subsequent dynamical history of planetary systems. For example, based on direct imaging data, Bowler et al. (2019) reported a difference in the eccentricity distributions of planets and brown dwarfs, evidence that these objects form in different ways. Planets are expected to form on circular and coplanar orbits within protoplanetary disks, although can develop non-zero eccentricities via planet/planet interactions (Rasio & Ford 1996), secular von Zeipel-Lidov-Kozai cycles (Naoz 2016; Ito & Ohtsuka 2019), planet-disk interactions (Goldreich & Sari 2003), or other dynamical processes. The same dynamical processes can also alter orbital inclinations of planetary orbits, changing the obliquity of the star (Winn & Fabrycky 2015). In particular, von Zeipel-Kozai-Lidov cycles combined with tidal friction (Fabrycky & Tremaine 2007) can leave a planet stranded on a polar orbit (e.g., GJ 436b; Bourrier et al. 2018b) or even a retrograde orbit (e.g., HAT-P-7b; Winn et al. 2009). Obliquities can be measured by exploiting the Rossiter-McLaughlin (RM) effect, the alteration of the star's absorption line profiles during a planetary transit, which is often manifested as a radial-velocity anomaly (Rossiter 1924; McLaughlin 1924).

Although hundreds of obliquities have been measured with the RM effect, the current list includes only three M dwarfs: GJ 436 (Bourrier et al. 2018a), TRAPPIST-1 (Hirano et al. 2020a), and AU Mic (Addison et al. 2020; Hirano et al. 2020b; Palle et al. 2020). The obliquity of GJ 436b was found to be $\lambda = 72^{+33}_{-24}$, suggesting a strong misalignment, and AU Mic was found to be well-aligned with the equator of its host star. For TRAPPIST-1, the current data are compatible with a low obliquity but with large uncertainties. RM measurements of M dwarfs have been limited as they tend to be optically faint, hindering the detection of the RM effect. If we could expand the sample of M-dwarfs with measured obliquities, we might be able to gain clues about the dynamical histories of late-type stars with close-in Neptune and Jupiter mass planets.

This paper reports on a suite of follow-up observations of K2-25, a young M4.5 dwarf in the Hyades with a close-orbiting and transiting Neptune-sized planet. The new data

allow us to measure the planet's mass and the star's obliquity. Section 2 introduces the K2-25 system. Section 3 presents the new photometric and spectroscopic observations. Section 4 describes the data reduction. Section 5 presents an updated determination of the stellar parameters, along with new estimates of the projected rotation velocity, rotation period, and stellar inclination with respect to the line of sight. Section 6 presents a joint analysis of the photometric and spectroscopic data and provides the results for the planet's mass and other system parameters. Section 7 focuses on the RM effect. All of the results are discussed in Section 8, along with the feasibility of future observations of the planet's transmission spectrum. We conclude with a summary of our key findings in Section 9.

2. THE K2-25 SYSTEM

K2-25b was originally discovered by Mann et al. (2016a) and David et al. (2016) in data from the Kepler spacecraft as part of the K2 mission. K2-25b is a Neptune-sized planet $(R \sim 3.5R_{\oplus})$ in a P = 3.5 day orbit around its M4.5 dwarf host star in the Hyades. With its large transit depth of 1.1% and its brightness at near-infrared (NIR) wavelengths (J = 11.3), K2-25b has been discussed (see e.g., Mann et al. 2016a; David et al. 2016) as a prime candidate for atmospheric characterization in the future with JWST, ARIEL (Tinetti et al. 2016), and large ground-based observatories. In addition, due to its large transit depth and the rapid stellar rotation $(P = 1.878 \,\mathrm{days})$, K2-25b has an estimated RM effect amplitude of $\sim 60-70\,\mathrm{m/s}$, making the Rossiter-McLaughlin effect detectable with high precision radial velocities in the NIR. Further, planets with well-constrained ages are scarce, making the determination of K2-25b's 3D orbit valuable for constraining theories of planet formation and migration mechanisms that aim to explain planetary and orbital parameters as a function of age.

Recently, two groups discussed additional transit followup observations of K2-25b. Thao et al. (2020) study the transit depth of K2-25b as a function of wavelength using photometric observations at different wavelengths from K2, the MEarth Observatories (Irwin et al. 2015), and the Las Cumbres Global Telescope Network (LCOGT) in the optical, and from the Spitzer Telescope in the NIR, and find no significant evidence of changes in transit depth as a function of wavelength. To explain the flat broadband transmission spectrum, they favor a scenario where K2-25b has a cloudy atmosphere using a predicted mass from exoplanet mass-radius relations. Although K2-25 could have a cloudy atmosphere, we show in this paper that the apparent flat broad-band transmission spectrum of K2-25b could also be partially explained by the larger as-observed mass of K2-25b than the mass assumed in Thao et al. (2020).

Kain et al. (2019) performed long-term photometric monitoring of K2-25 using the MEarth Observatories (Irwin et al. 2015), to monitor the activity of the star, and to look for evidence of transiting timing variations (TTVs) in the transits of K2-25b, which could be suggestive of an additional planet orbiting in the system. They show that the photometric variability of the star was significantly smaller between

2016-2018 than what *K2* observed in its 71 day observing window in 2015. Kain et al. (2019) do not identify any definite starspot-crossing events in their transit data, although a few of their transits could contain tentative evidence for such events. They further searched for TTVs in the system from the additional transits, and find no evidence of significant TTVs, placing constraints on planetary companions orbiting close to mean-motion resonances of K2-25b's orbit. This agrees with the transit observations presented in this work.

3. OBSERVATIONS

3.1. *ARCTIC*

We obtained five transits of K2-25b using the Astrophysical Research Consortium Telescope Imaging Camera (ARC-TIC) imager (Huehnerhoff et al. 2016) on the 3.5m Astrophysical Research Consortium (ARC) 3.5m Telescope at Apache Point Observatory (APO) on the nights of UT 20170917, 20190104, 20190118, 20190125, and 20190201. All of the transit observations were performed with the Engineered Diffuser available on ARCTIC, which we designed specifically to enable very high-precision photometric observations (see e.g., Stefansson et al. 2017, 2018a,b, 2020). We used the SDSS i' filter except on the night of 20190118, when we used the SDSS z' filter without the diffuser to minimize background Moon contamination that night. Only data from the egress from this night were usable due to the moon contamination. The observations were performed in the quadreadout and 4×4 binning mode, resulting in a readout time of 2.7 s. The first transit was observed with an exposure time of 20s, and the rest of the transits with an exposure time of 30 s. In this binning mode ARCTIC has a gain of 2.0e/ADU, and a plate scale of 0.44"/pixel. Table 1 further summarizes the observations.

3.2. Half Degree Imager

We observed four transits of K2-25b using the Half-Degree Imager (HDI; Deliyannis 2013) at the WIYN 0.9m Telescope at Kitt Peak National Observatory on the nights of UT 20180207, 20180221, 20181214, and 20181221. HDI has a 4096 × 4096 pixel back-illuminated CCD from e2v, covering a $29.2' \times 29.2'$ Field-of-View (FOV) at a plate scale of 0.425''/pixel, with a gain of 1.3 e/ADU in the 1×1 binning mode. The observations were performed as part of commissioning observations of the newly installed Engineered Diffuser for the telescope, which is now available for high precision photometric observations. The Engineered Diffuser on HDI uses the same custom-optimized top-hat Engineered Diffuser pattern as we developed for the ARC 3.5m Telescope with a diffuser opening angle of $\theta = 0.34^{\circ}$ (Stefansson et al. 2017). The diffuser is placed in a filter wheel holder in the dual filter wheel 45 mm away from the focal plane, resulting in a stabilized Point Spread Function (PSF) with a Full-Width-at-Half-Maximum (FWHM) of 7.6". The size of the diffuser is $50.8 \times 50.8 \, \text{mm}$ and vignettes the field to an effective FOV of $\sim 20' \times 20'$, still allowing for a number of available reference stars in the field.

Table 1 summarizes the observations, and lists the num-

ber of reference stars and exposure time used. All of the observations were performed in the SDSS z' filter using the 1×1 binning mode. The first three observations were performed with the Engineered Diffuser with an exposure time of 120s. The diffuser was not used on the night of 20181221 due to the high degree of Moon contamination (Moon was \sim 97% full and at a separation of \sim 6° from the target). Further, during this night, we experienced issues with the camera shutter causing the shutter to be stuck for a periods of time which led to gaps in the datastream seen in Figure 2g. As the observations on that night were performed close to infocus, the exposure time used was scaled down to 30s to minimize risk of saturation. Two different readout modes were used for the observations: a quad readout mode, and a singleamplifier readout mode, with readout times of 11s and 37s, respectively. The observations on the nights of 20180207 and 20181221 used the single amplifier readout mode, with the quad readout mode used for the other two.

3.3. Habitable-zone Planet Finder

We obtained precision NIR RVs of K2-25 with the Habitable-zone Planet Finder (HPF) (Mahadevan et al. 2012, 2014) with the twofold goal to constrain the mass of K2-25b and to constrain the obliquity of the host star. HPF is a high resolution ($R \sim 55,000$) NIR fiber-fed spectrograph on the 10m Hobby-Eberly Telescope (HET) at McDonald Observatory in Texas. HPF is actively temperature stabilized to the milli-Kelvin level to enable precision radial velocities in the NIR (Hearty et al. 2014; Stefansson et al. 2016). The HET is a fully queue-scheduled telescope with all observations executed in a queue by the HET resident astronomers (Shetrone et al. 2007). HPF has a NIR laser-frequency comb (LFC) calibrator which has been shown to enable $\sim 20\,\mathrm{cm\,s^{-1}}$ calibration precision and $1.5\,\mathrm{m\,s^{-1}}$ RV precision on-sky on the bright and stable M-dwarf Barnard's Star (Metcalf et al. 2019).

In total, we obtained 105 spectra in 34 different visits with HPF. We removed 16 spectra after performing a S/N > 25 quality cut (S/N per pixel); these spectra were adversely affected by weather and deemed to be too low S/N. The median S/N of the 89 remaining spectra was 45, which were obtained in 32 different HET visits/tracks¹.

For the out-of-transit observations, we obtained 69 spectra of K2-25b, in 32 different tracks with HPF with a median S/N of 52 and median RV errorbar of 42 m/s. For most of the spectra (58 in total) the exposure time was 969 – 1160s, with two spectra obtained per HET visit. In addition to these spectra, we also use 11 spectra which were obtained during the out-of-transit baseline for the three transit observations described below. These spectra had an exposure time of 309s and a median S/N of 32. To simplify the analysis and obtain the highest precision errorbar per RV visit, we performed a weighted average of the RVs of the out-of-transit RVs following the methodology described in Metcalf et al. (2019); Stefansson et al. (2020). The final RVs used for the out-of-

 $^{^1}$ The unique design of the HET only enables acquisition of K2-25 during a discrete East or West track with a $\sim\!1$ hour maximum track duration.

transit modeling are shown in Table C3 in the Appendix.

To constrain the obliquity of the K2-25b system, we obtained spectra of K2-25 during three transits of K2-25b on the nights of 20181221 UT, 20181228 UT, and 20190104 UT. We used an exposure time of 309s, corresponding to 29 up-the-ramp reads on the HPF Hawaii-2RG NIR detector, to obtain the required time resolution to resolve the RM waveform. Overall, we obtained 11, 11, and 9 spectra in the three different visits, respectively. The three sets of spectra had a median S/N of 33, 36, and 35 (per extracted 1D pixel), and median RV errors of $85 \, \text{m/s}$, $72 \, \text{m/s}$, and $74 \, \text{m/s}$, respectively. Out of these 31 spectra, 20 spectra were in transit and 11 out-of-transit. The S/N of the first night (20181221 UT), was the lowest as the Moon was $\sim 97\%$ full and only $\sim 6^\circ$ away from the target during the observations. Making the S/N > 25 quality cut described above removed one out-of-

transit baseline spectrum that was observed at the edge of the available HET track during the transit on 20190104 UT. Table 1 further summarizes the in-transit observations. The final RVs used for the in-transit modeling are shown in Table C4 in the Appendix.

Following our observational setup described in Stefansson et al. (2020), due to the faintness of the target (J = 11.3), we elected to not have the HPF LFC on for any of the observations described, to minimize the risk that the bright LFC lines would contaminate the stellar spectrum. This does not significantly impact the drift correction for this target, as the drift of HPF is linear during a night (amplitude of $\sim 10 \, \text{m/s}$), and is well traced and calibrated to $< 1 \, \text{m/s}$ RV precision by daily HPF calibrations and a linear drift model as has been detailed in Stefansson et al. (2020). For all of the observations above, the HPF sky fiber was used to subtract any Moon and/or other sources of background light contamination.

Table 1. Summary of transit observations analyzed in this work. The aperture setting column lists the aperture setting in pixels used to extract the photometry in AstroImageJ (Collins et al. 2017), showing the radius of the photometric aperture, and the radius of the inner and outer background annulli, respectively. The Airmass Range column shows the airmass range of the observations from high to low airmass. The observations on UT 20181221 and UT 20190118 were not performed with the diffuser due to the large Moon contamination on those nights.

Date (UT)	Instrument	Filter/Bandpass	Exposure Time (s)	Diffuser	Airmass Range	# Ref. Stars	Aperture setting (pixels)
Photometric T	ransit Observ	rations:					
20180207	HDI	SDSS z'	120	Yes	2.10, 1.16	9	21, 40, 60
20180221	HDI	SDSS z'	120	Yes	1.84, 1.09	12	21, 40, 60
20181214	HDI	SDSS z'	120	Yes	1.70, 1.04	7	22, 39, 59
20181221	HDI	SDSS z'	30	No	1.37, 1.08	6	9, 16, 24
20170917	ARCTIC	SDSS i'	20	Yes	1.68, 1.09	8	14, 35, 50
20190104	ARCTIC	SDSS i'	30	Yes	1.41, 1.06	3	16, 28, 42
20190118	ARCTIC	SDSS z'	30	No	1.23, 1.10	6	16, 50, 70
20190125	ARCTIC	SDSS i'	30	Yes	1.34, 1.05	13	19, 30, 45
20190201	ARCTIC	SDSS i'	30	Yes	1.11, 1.04	8	16, 28, 42
Spectroscopic	Transit Obse	rvations:					
20181221	HPF	820-1280nm	300	-	1.37, 1.17	-	-
20181228	HPF	820-1280nm	300	-	1.36, 1.17	-	-
20190104	HPF	820-1280nm	300	-	1.34, 1.18	-	-

4. DATA REDUCTION

4.1. Photometric Observations

We reduced the photometry using AstroImageJ (Collins et al. 2017), following a similar methodology as in Stefansson et al. (2017) and Stefansson et al. (2018a). In short, we experimented using a number of different aperture settings, varying the radii of the software aperture, and inner, and outer background annulli, selecting the aperture setting that resulted in the minimum RMS scatter in the resulting photometry. Table 1 summarizes the aperture setting used for each observation in pixels that led to the lowest pho-

tometric noise in the light curve. We add the scintillation error bar estimates to the photometric error bars estimated by AstroImageJ following the methodology in Stefansson et al. (2017). The individual exposures were calibrated using standard median bias, dark, and flat field procedures in AstroImageJ following Stefansson et al. (2017). The median bias, dark, and flat-field exposures were taken at the beginning or end of each night of observation. We converted the timestamps of the ground-based observations to Barycentric Julian Date time (BJD_{TDB}) using the Python package barycorrpy (Kanodia & Wright 2018), which uses the barycentric correction algorithm from Wright & Eastman

(2014).

K2-25b was observed by the *K2* mission as part of Campaign 4 in long-cadence (30min exposures) mode, from February 8, 2015 to April 20th 2015, resulting in 71 days of continuous photometric observations. We use the light curve from the Everest pipeline (Luger et al. 2016), which is capable of correcting the periodic correlated errors in the *K2* data due to imperfect pointing of the spacecraft. The corrected Everest light curve improved the 6hour CDPP standard deviation of the raw *K2* long-cadence data of K2-25 from 432 ppm to 295 ppm.

4.2. Spectroscopic Observations

The HPF 1D spectra were reduced and extracted with the custom HPF data-extraction pipeline following the procedures outlined in Ninan et al. (2018), Kaplan et al. (2018), and Metcalf et al. (2019). For the RV extractions, we used the Spectrum Radial Velocity Analyzer (SERVAL) which we have adapted for use for the HPF spectra, following the methodology described in Metcalf et al. (2019) and Stefansson et al. (2020). In short, SERVAL uses the templatematching method to measure the radial velocities (see e.g., Anglada-Escudé & Butler 2012; Zechmeister et al. 2018). We extracted the HPF RVs using the 8 orders with the least telluric contamination in the HPF bandpass (orders covering the wavelength regions from 8540-8890Å, and 9940-10760Å). Although we plan to include more orders in the RV extraction to potentially enable even high RV precision, we elected using only these 8 orders that we have extensively tested to produce reliable RVs with HPF. We subtracted the estimated sky-background from the stellar spectrum using the dedicated HPF sky fiber. Following the methodology described in Metcalf et al. (2019), we explicitly masked out telluric lines and sky-emission lines to minimize their impact on the RV determination. To minimize the impact of the asymmetric spectral-line variations seen during the RM effect, we generated the master RV template to extract the RV using only out-of-transit spectra. As described in Subsection 3.3, this resulted in 40 spectra after a quality cut of S/N > 25used to generate the master RV template.

We calculated the RVs in two steps. First, to generate a first-pass master template, we ignored any possible planetary induced RVs and coadded the out-of-transit spectra (after barycentric correction) using the SERVAL template creation functionality, which we then used to derive a first set of RVs. We then used this first set of RVs to further align the out-of-transit spectra to create a second-pass RV template, and re-extracted RVs using this more refined template. The out-of-transit RVs used for the mass determination of K2-25b are shown in Table C3 in the Appendix, and the in-transit RVs are shown in Table C4.

5. STELLAR PARAMETERS

The stellar parameters used in this work are summarized in Table 2, and are broadly adopted from the values presented in Mann et al. (2016a) and Thao et al. (2020). Thao et al. (2020) estimate their stellar parameters (including the stellar radius and mass) using the empirically calibrated relations

of Mann et al. (2015) and Mann et al. (2019). As an additional test of these parameters, we performed an independent SED and isochrone fit using the EXOFASTv2 package (Eastman 2017; Eastman et al. 2019) placing informative Gaussian priors on the known metallicity ([Fe/H] = 0.15 ± 0.03) and age Age = $730 \pm 50 \,\mathrm{Myr}$ of the star derived from its Hyades cluster membership (Mann et al. 2016a), along with its known distance from Gaia. The values we obtained are in good agreement with the values from Thao et al. (2020) in particular obtaining a fully consistent stellar density value from the stellar mass and radius (we obtain a density of $\rho = 14.7 \pm 1.2 \,\mathrm{g \, cm^{-3}}$, and they obtain $\rho = 14.7 \pm 1.5 \,\mathrm{g \, cm^{-3}}$), which carries important information on the eccentricity of the planet via the photoeccentric effect (see e.g., Dawson & Johnson 2012). We elected to incorporate the values from Thao et al. (2020) for our analysis, as they are derived from precise empirically calibrated relations, rather than through model-dependent SED and isochrone fitting. We note that with an effective temperature of $T_{\rm eff} = 3207 \, \rm K$ and radius of $R = 0.29R_{\oplus}$, K2-25 lies at the higher radius end of the radius discontinuity for low-mass M-dwarfs in Rabus et al. (2019), which they observe for M-dwarfs with effective temperatures between 3200K and 3300K. This radius discontinuity has been interpreted as the boundary between partially convective and fully-convective M-dwarfs, and as K2-25 is observed to be just above the radius discontinuity, suggests that K2-25 is potentially partially convective and just at the onset of being a fully-convective star. Our estimate of the stellar rotation period, projected rotational velocity, and stellar inclination are discussed in the next two subsections.

Table 2. Summary of stellar parameters used in this work.

Parameter	Description	Value	Reference
Main identifier	s:		
EPIC	-	210490365	Huber
Stellar magnitu	des:		
B	APASS Johnson B mag	17.449 ± 0.144	APASS
V	APASS Johnson V mag	15.891 ± 0.180	APASS
g'	APASS Sloan g' mag	16.567 ± 0.018	APASS
r'	APASS Sloan r' mag	15.300 ± 0.183	APASS
i'	APASS Sloan i' mag	13.698 ± 0.206	APASS
Kepler-mag	Kepler magnitude	14.528	Huber
J	2MASS J mag	11.303 ± 0.021	2MASS
H	2MASS H mag	10.732 ± 0.020	2MASS
K_S	2MASS K _S mag	10.444 ± 0.019	2MASS
WISE1	WISE1 mag	10.275 ± 0.024	WISE
WISE2	WISE2 mag	10.086 ± 0.020	WISE
WISE3	WISE3 mag	9.936 ± 0.057	WISE
Stellar Paramet	ers:		
M_*	Mass in M_{\odot}	0.2634 ± 0.0077	Thao et al. (2020)
R_*	Radius in R_{\odot}	0.2932 ± 0.0093	Thao et al. (2020)
$ ho_*$	Density in g cm ⁻³	14.7 ± 1.5	Thao et al. (2020)
log(g)	Surface gravity in cgs units	4.944 ± 0.031	Derived from M & R
$T_{ m eff}$	Effective temperature in K	3207 ± 58	Thao et al. (2020)

Table 2 (continued)

Parameter	Description	Value	Reference
[Fe/H]	Metallicity in dex	0.15 ± 0.03	Mann et al. (2016a)
Age	Age in Gyrs	$0.730^{+0.050}_{-0.052}$	Mann et al. (2016a)
L_*	Luminosity in L_{\odot}	0.00816 ± 0.00029	Thao et al. (2020)
d	Distance in pc	$45.01_{-0.17}^{+0.16}$	Gaia
π	Parallax in mas	$22.218^{+0.081}_{-0.083}$	Gaia
$P_{\rm rot}$	Rotational period in days	1.878 ± 0.005	This work
$v \sin i_*$	Stellar rotational velocity in km s ⁻¹	8.8 ± 0.6	This work
i_*	Stellar inclination	$90\pm12^{\circ}$	This work

References are: Huber (Huber et al. 2016), Lepine (Lépine & Shara 2005), Reid (Reid et al. 2004), Gaia (Gaia Collaboration 2018), APASS (Henden et al. 2015), UCAC2 (Zacharias et al. 2004), 2MASS (Cutri et al. 2003), WISE (Cutri & et al. 2014).

5.1. Rotation Period

The K2 data of K2-25b is modulated at the 1% level, suggestive of starspots and/or active regions that rotate in and out-of-view with the rotation period of the star. We independently measure the rotation period of K2-25b following the methodology in Stefansson et al. (2020). In short, we model the stellar active regions using a quasi-periodic Gaussian Process (GP) as quasi-periodic GP kernels have been shown to yield reliable stellar rotation rates (Angus et al. 2018). For computational efficiency, we adopt the quasi-periodic kernel from the celerite package (Foreman-Mackey et al. 2017) as implemented in juliet. The form of this kernel is further discussed in Section 6.1. To estimate the rotation period, we removed points in a 2x transit window around the expected transit times, and fit the resulting photometry using the celerite quasi-periodic GP kernel (see Foreman-Mackey et al. 2017, and Equation 3 in Subsection 6.1). We placed non-informative priors on the GP period, amplitude and GP decay timescale parameters.

Figure 1 shows the resulting posteriors of the GP period hyperparameter along with the phase-folded photometry using our best-fit rotation period of $P_{\rm rot} = 1.878 \pm 0.005\,{\rm days}$. In the phase-folded photometry, we see that the photometric modulation remains relatively stable throughout the 71 days of K2 dataset, with a slight evolution observed. From Figure 1 we also see evidence of flares. Our period estimate agrees well with the rotation period reported in Mann et al. (2016a) of $P_{\rm rot} = 1.881 \pm 0.021\,{\rm days}$ estimated using an Autocorrelation-Function method, and Dmitrienko & Savanov (2017) of $P_{\rm rot} = 1.878 \pm 0.030\,{\rm days}$ from power-spectra analysis. The top panel in Figure 2 in Section 6 shows the modulation in the K2 photometry as a function of time for the full 71 day baseline.

5.2. Projected Rotational Velocity and Stellar Inclination

We measured the projected rotation velocity using the empirical spectral matching algorithm described in Stefansson et al. (2020), which closely follows the SpecMatchEmp algorithm described in Yee et al. (2017). In short, the algorithm




Figure 1. a) Phase-folded *K2* photometry of K2-25 using our best estimate of the rotation period (see b), showing a clear periodic photometric modulation. The photometric modulation remains stable over the 71 day *K2* photometric baseline, with a slight evolution seen. b) The posteriors of the period of the quasi-periodic GP kernel, which we interpret as the stellar rotation period.

compares the as-observed target star spectrum to a library of as-observed slowly rotating stellar spectra using a χ^2 metric, and refer the reader to Stefansson et al. (2020) for a more detailed discussion.

We used the algorithm to measure independent $v \sin i_*$ values for the 8 HPF orders cleanest of tellurics. Table 3 shows the resulting values, showing that the independent orders agree well on the resulting value, with a mean value of $v \sin i_* = 8.8 \text{km/s}$ and a scatter of 0.3 km/s. The values in Table 3 show a small formal scatter at the 0.3 km/s level. However, given our experience calculating $v \sin i_*$ values of other stars from high resolution spectra with HPF a more realistic error estimate is a factor of 2 larger. We adopt an error estimate of 0.6km/s to account for possible systematics in our method as $v \sin i_*$ measurements are generally dominated by systematics (see e.g., Reiners et al. 2012). As such, we adopt a $v \sin i_* = 8.8 \pm 0.6$ km/s. We note that value is somewhat larger than the value presented in Mann et al. (2016a) of $v \sin i_* = 7.8 \pm 0.5$ km/s determined from their as-observed spectra obtained with the IGRINS spectrograph. The method used in Mann et al. (2016a) used an overall similar χ^2 method as presented here, but used artificially rotationally broadened theoretical BT-SETTL spectra for the χ^2 comparison instead of as-observed spectra. Although this value is formally slightly higher than the maximum equatorial velocity of the star of 7.9 ± 0.25 km/s (assuming a stellar radius of $R = 0.2932 \pm 0.0093 \,\mathrm{R}_{\odot}$ and stellar rotation period of $P = 1.878 \pm 0.005$ days) the two values overlap within the

Table 3. Resulting $v \sin i_*$ values for the 8 different HPF orders cleanest of tellurics. The resulting median value is $v \sin i_* = 8.8 \,\text{km/s}$ with a scatter of $0.3 \,\text{km/s}$. As is mentioned in the text, we adopt a value of $v \sin i_* = 8.8 \pm 0.6 \,\text{km/s}$.

Order	Wavelength Region [Å]	$v \sin i_* \text{ [km/s]}$
4	[8540, 8640]	8.3
5	[8670, 8750]	8.9
6	[8790, 8885]	8.6
14	[9940, 10055]	8.8
15	[10105, 10220]	8.8
16	[10280, 10395]	9.0
17	[10460, 10570]	9.3
18	[10640, 10760]	8.9

 2σ uncertainties. The high $v \sin i_*$ suggests that the stellar inclination is close to 90°.

To estimate accurate posteriors for the stellar inclination i_* from the stellar rotational velocity v estimated from R and P_{rot} and its sky-projection $v \sin i_*$ measured from the HPF spectra, we use the formalism in Masuda & Winn (2020), which accurately accounts for the correlated dependence between $v \sin i_*$ and v. The $v \sin i_*$ measurement does not distinguish between solutions between i and $180^\circ - i$ and we thus calcu-

late two independent solutions between 0° and 90° , and 90° and 180° , respectively. Using the values listed in Table 2 for $P_{\rm rot}$, $v\sin i_*$, and R, we obtain two mirrored posteriors with a highest-likelihood inclination at 90° . Taken together, the two solutions result in an inclination constraint of $90 \pm 12^{\circ}$, consistent with viewing K2-25's stellar equator edge on. This agrees well with the stellar inclination constraint provided by Mann et al. (2016a) of $i_* > 79^{\circ}$ at 1σ (68.4% confidence; they considered inclinations between 0 and 90°).

Table 4. Summary of priors used for the three joint transit and RV fits performed. $\mathcal{N}(m,\sigma)$ denotes a normal prior with mean m, and standard deviation σ ; $\mathcal{U}(a,b)$ denotes a uniform prior with a start value a and end value b, $\mathcal{J}(a,b)$ denotes a Jeffreys prior with a start value a and end value b. A Gaussian prior on the stellar density was placed for all fits. The dilution parameters in juliet were fixed to 1 for all transit observations. For the three models considered we sampled 70, 72, and 69 parameters.

Parameter	Description	Model RV1 $(e = 0, GP \text{ for RVs})$	Model RV2 $(e \neq 0, GP \text{ for RVs})$	Model RV3 $(e \neq 0, \text{No GP for RVs})$
Orbital Param	eters:			
P (days)	Orbital period	$\mathcal{N}(3.484548, 0.000042)$	$\mathcal{N}(3.484548, 0.000042)$	N(3.484548, 0.000042)
T_C	Transit Midpoint - 2400000 (BJD _{TDB})	U(58515.63, 58515.66)	U(58515.63, 58515.66)	U(58515.63, 58515.66)
$r_1{}^a$	Radius ratio / Impact parameter	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$
r_2^a	Radius ratio / Impact parameter	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$
a/R_*	Scaled semi-major axis	U(1,50)	U(1,50)	U(1,50)
$m_{ m flux}$	Transit baseline parameter	$\mathcal{U}(0,0.1)$	$\mathcal{U}(0,0.1)$	U(0, 0.1)
$\sigma_{ m K2}$	Photometric errorbar (ppm)	$\mathcal{J}(1, 1000)$	$\mathcal{J}(1,1000)$	$\mathcal{J}(1, 1000)$
e	Eccentricity	0 (fixed)	U(0, 0.95)	U(0, 0.95)
ω	Argument of Periastron	90 (fixed)	U(0, 360)	U(0, 360)
K	RV semi-amplitude (m/s)	U(0, 200)	U(0, 200)	U(0, 200)
Other constrai	ints:			
$ ho_*$	Stellar density (g cm ⁻³)	$\mathcal{N}(14.7, 1.5)$	$\mathcal{N}(14.7, 1.5)$	$\mathcal{N}(14.7, 1.5)$
litter and othe	r instrumental terms:			
$q_1{}^b$	Limb-darkening parameter	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$
$q_2^{\ b}$	Limb-darkening parameter	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$	$\mathcal{U}(0,1)$
$\sigma_{ m phot}{}^c$	Photometric jitter (ppm)	J(1,1000)	J(1,1000)	J(1,1000)
$\mu_{ ext{phot}}{}^c$	Photometric baseline	$\mathcal{N}(0,0.1)$	$\mathcal{N}(0, 0.1)$	$\mathcal{N}(0, 0.1)$
$\sigma_{ m HPF}$	HPF RV jitter (m/s)	$\mathcal{J}(0.1, 300)$	$\mathcal{J}(0.1, 300)$	$\mathcal{J}(0.1, 300)$
$\mu_{ ext{HPF}}$	HPF RV offset (m/s)	$\mathcal{U}(-200, 200)$	$\mathcal{U}(-200, 200)$	$\mathcal{U}(-200, 200)$
Shared Photor	netric and RV Quasi-Periodic GP Paramete	ers:		
$P_{ m GP}$	GP Period (days)	$\mathcal{N}(1.8784, 0.005)$	$\mathcal{N}(1.8784, 0.005)$	$\mathcal{N}(1.8784, 0.005)$
K2 Quasi-Peri	odic GP Parameters:			
B	Photometric GP Amplitude (ppm ²)	$\mathcal{J}(10^{-6}, 10^5)$	$\mathcal{J}(10^{-6}, 10^5)$	$\mathcal{J}(10^{-6}, 10^5)$
C	GP Additive Factor	$\mathcal{J}(10^{-6}, 10^5)$	$\mathcal{J}(10^{-6}, 10^5)$	$\mathcal{J}(10^{-6}, 10^5)$
L	GP Length Scale (days)	$\mathcal{J}(10^3,10^6)$	$\mathcal{J}(10^3,10^6)$	$\mathcal{J}(10^3, 10^6)$
Ground-based	Approximate Matern GP Parameters ^d :			
$\sigma_{ ext{GP}}$	Photometric GP Amplitude (ppm)	$\mathcal{J}(10^{-1}, 10^4)$	$\mathcal{J}(10^{-1}, 10^4)$	$\mathcal{J}(10^{-1}, 10^4)$
L	Timescale of exponential kernel (days)	$\mathcal{J}(10^{-2}, 10^5)$	$\mathcal{J}(10^{-2}, 10^5)$	$\mathcal{J}(10^{-2}, 10^5)$
ho	Timescale of Matern kernel (days)	$\mathcal{J}(10^{-2}, 10^5)$	$\mathcal{J}(10^{-2}, 10^5)$	$\mathcal{J}(10^{-2}, 10^5)$
RV GP Param	eters:			
$\sigma_{ ext{GP}}$	RV GP Amplitude (m/s)	$\mathcal{J}(10^1,10^5)$	$\mathcal{U}(10^1,10^5)$	-
Γ	Harmonic structure / scaling parameter	$\mathcal{N}(8.0, 1.9)$	$\mathcal{N}(8.0, 1.9)$	-
α	Inverse length scale (days ⁻²)	$\mathcal{J}(10^{-12}, 10^{-3})$	$\mathcal{J}(10^{-12}, 10^{-3})$	-

Table 4 (continued)

Parameter	Description	Model RV1	Model RV2	Model RV3
		(e = 0, GP for RVs)	$(e \neq 0, GP \text{ for RVs})$	$(e \neq 0, \text{No GP for RVs})$

^a Using the efficient sampling of the r_1 and r_2 parameterization for the impact parameter b and radius ratio $p = R_p/R_*$ as described in Espinoza (2018).

6. PLANET PARAMETERS FROM TRANSIT PHOTOMETRY AND RVS

6.1. Transit, RV and Gaussian Process model

We jointly model the *K2* and ground-based transits and the HPF out-of-transit RVs using the juliet Python package (Espinoza et al. 2019), which uses the batman Python package (Kreidberg 2015) for the transit model, and the radvel package (Fulton et al. 2018) for the RV model. We used the dynesty sampler (Speagle 2019) available in juliet to perform dynamic nested sampling to obtain both posterior and evidence estimates, where we used the default weight and stopping functions, and stopping criteria in dynesty². The total log likelihood in juliet is the sum of the individual log-likelihoods of each dataset considered³.

Following the implementation in juliet, we parametrize the radius ratio $p = R_p/R_*$ and the impact parameter b using the efficient r_1 and r_2 parametrization described in Espinoza (2018). We sampled the limb darkening parameters using the quadratic q_1 and q_2 limb-darkening parametrization from Kipping (2013). Following the suggestion in Kipping (2010), we resampled and rebinned our transit model to the effective 30 minute exposure time of the long-cadence K2 data to account for the smoothing of the transit shape. We assumed that there was no source of dilution, as no obvious close-by companion is seen in the adaptive optics imaging presented in Mann et al. (2016a).

We used a Gaussian Process (GP) model with 3 different kernels to account for the characteristic correlated noise behavior in the *K*2 photometry, ground-based photometry, and the radial velocities. GPs have been used by a number of groups to jointly model correlated noise observed in photometric and radial velocity data due to stellar active regions on the surface of the star (e.g., spots, faculae, plages; see e.g., Haywood et al. 2014; Grunblatt et al. 2015; López-Morales et al. 2016; Dai et al. 2017; Angus et al. 2018; Haywood et al. 2018). For our GP modeling, we use the GP kernels available in juliet, which are based on the GP imple-

mentations from the george (Ambikasaran et al. 2015) and celerite (Foreman-Mackey et al. 2017) Python packages. In juliet, the elements of the covariance matrix C_i for instrument i are assumed to be of the form,

$$C_{i,l,m} = k_i(x_l, x_m) + (\sigma_{i,w}^2 + \sigma_{t,i}^2) \delta_{l,m}, \tag{1}$$

where $\delta_{l,m}$ is the Kronecker delta function, $k_i(x_l, x_m)$ is the kernel of the GP for instrument i, and $\sigma_{t_l,i}$ is the error estimated at time t_l , and $\sigma_{i,w}$ is an additional white-noise jitter parameter. For all of the GP fits considered, we fit a kernel function as well as a separate white noise jitter term for each instrument as is shown in Equation 1.

We choose three different GP kernels as a balance between computational speed and to adequately account for the characteristic correlated noise properties of the different datasets. First, to model the correlated noise in the radial velocity observations, we elect to use the quasi-periodic rotational kernel, which has been shown to be effective at modeling rotational variations in RV datasets and has well-studied hyperparameters (Haywood et al. 2014; Grunblatt et al. 2015; Haywood et al. 2018). In juliet, the quasi-periodic kernel is given with the following form,

$$k(x_l, x_m) = \sigma_{\rm GP}^2 \exp\left(-\alpha \tau^2 - \Gamma \sin^2\left[\frac{\pi \tau}{P_{\rm GP}}\right]\right),\tag{2}$$

where $\tau = |x_l - x_m|$, with hyperparameters σ_{GP} (RV amplitude in m/s), α (inverse square timescale in units of days⁻²), Γ (a unitless scaling parameter), and P_{GP} (the periodicity of the GP in days, which we interpret as the rotation period)⁴. The Γ parameter changes the amplitude of the \sin^2 term, and controls the harmonic structure of the resulting GP model (see e.g., discussion in Angus et al. 2018). We follow Haywood et al. (2018) and place an informative prior on the Γ parameter; Haywood et al. (2018) place an informative Gaussian prior of $\eta_4 = 0.5 \pm 0.05$, which corresponds to $\Gamma = 8 \pm 1.9$ (in their notation $\Gamma = 2/\eta_4^2$). In doing so enforces the RV curve to have up to two or three maxima and two or three minima per rotation, as is typical of stellar light curves and RV curves (Haywood et al. 2018).

b Overall we modeled 4 pairs of limb darkening parameters q₁ and q₂ (parametrization from Kipping (2013)): a) one pair for K2, b) one pair for all of the transits observed with ARCTIC in the SDSS i' filter and c) one pair for the ARCTIC SDSS z' filter observations, and d) one pair for the HDI transits in the SDSS z' filter.

^C We placed a separate photometric jitter term and baseline offset term for each of the photometric observations.

d We place one set of three parameters (σ_{GP}, L, ρ) for each of the ground-based transits.

² For a description of the weight and stopping functions in the dynamic nested sampler in dynesty, see Section 3 in Speagle (2019).

 $^{^3}$ See Equations and discussion surrounding Equations 6 and 7 in Espinoza et al. (2019) to see the explicit likelihood used in juliet.

⁴ As mentioned by Espinoza et al. (2019), the hyperparameters in Equation 2 correspond to the following parameters in the notation of Haywood et al. (2018): $\sigma_{GP} = \eta_1$, $\alpha = 1/2\eta_2^2$, $P_{rot} = \eta_3$, and $\Gamma = 2/\eta_4^2$.

Second, for a computationally efficient analysis of the *K2* data, we elect to use the quasi-periodic kernel available in celerite (Foreman-Mackey et al. 2017), where the kernel function is given with the following form,

$$k(x_l, x_m) = \frac{B}{2+C} e^{-\tau/L} \left[\cos \left(\frac{2\pi\tau}{P_{\text{GP}}} \right) + (1+C) \right], \quad (3)$$

where $\tau = |x_l - x_m|$, and where B, C, L, and P_{rot} are the hyperparameters of the kernel. B and C tune the weight of the exponential decay component of the kernel with a decay constant of L (in days), and P_{GP} corresponds to the periodicity of the quasi-periodic oscillations which we interpret as the stellar rotation period. Although not exactly of the same form as the kernel in Equation 2, we selected this kernel as it shares similar quasi-periodic properties as the kernel in Equation 2, but is orders of magnitude faster to evaluate on the large number of K2 datapoints (see discussion in Foreman-Mackey et al. 2017). Given the high precision and the clear photometric modulation seen in the K2 data, we share the P_{GP} parameter between the quasi-periodic K2 GP kernel and our quasiperiodic RV kernel, to allow the high precision K2 photometry to accurately constrain the P_{GP} parameter. We did not share any other GP parameters between the K2 and the RV kernels, given the different forms of the kernels used. Table 4 further lists all of the priors used and which parameters are linked between different GP kernels.

Lastly, the ground-based photometric transit observations are not long enough to be be measurably impacted by the starspot modulation seen in the *K*2 light curve. Instead, the characteristic timescales of the observed correlated noise is much shorter or < 1 hour, originating from observational and/or instrumental systematics. Therefore, for the ground-based observations, we adopt the Approximate Matern-3/2 kernel multiplied by an exponential kernel which has covariance properties that are better matched to these timescales (see e.g., Pepper et al. 2017; Espinoza et al. 2019). As implemented in juliet, this kernel has the following form (see also Foreman-Mackey et al. (2017)),

$$k(x_l, x_m) = \sigma_{\text{GP}}^2 e^{-\tau/L} \left[(1 + 1/\epsilon) e^{-1(1-\epsilon)s} + (1 - 1/\epsilon) e^{-1(1+\epsilon)s} \right],$$
(4)

where $s = \sqrt{3}\tau/\rho$, and $\tau = |x_l - x_m|$, with hyperparameters $\sigma_{\rm GP}$ (photometric amplitude in ppm), L (length scale of the exponential component in days), and ρ (length scale of the Matern-3/2 kernel in days), and with $\epsilon = 0.01$, where we note that as ϵ approaches 0 the factor inside the brackets converges to a Matern 3/2 kernel (Foreman-Mackey et al. 2017; Espinoza et al. 2019). To allow for sufficient flexibility in modeling out different systematics seen in the different observing setups in the ground-based light curves, we assigned each ground-based transit an independent Approximate Matern 3-2 kernel with independent hyperparameters (σ_{GP} , L, and ρ). As an additional test, we also experimented using a pure exponential kernel to model the correlations seen in the groundbased data (with a timescale L and σ amplitude parameters) as implemented in juliet. Although both kernels yielded consistent planet parameters within the 68.3% credible intervals of the posteriors ($\sim 1\sigma$ uncertainties for a Gaussian distribution) in our experiments, the pure exponential kernel tended to favor small decay values that visually over-fit the noise structures in the data. The additional capability of the composite kernel in Equation 4 to account for both lower and higher frequency correlated noise in the ground-based light curves led to less over-fitting of the noise (see e.g., additional discussions in Pepper et al. 2017; Espinoza et al. 2019), and we thus favor this kernel in our analysis.

To investigate the evidence for eccentricity in the system, and to study the impact that our GP RV model has on the derived orbital parameters, we ran three different models. First, we performed a fit assuming a circular orbit (e=0) (Model RV1 in Table 4) with a simultaneous GP fit for the transits and radial velocity data. Second, we performed a fit letting the eccentricity e and argument of periastron ω float (Model RV2 in Table 4), while also performing a simultaneous GP fit like in Model RV1. Third, we let e and ω float, but removed the GP model on the radial velocity data (keeping the GP model for the photometry; Model RV3 in Table 4). The priors for these three models are summarized in Table 4. In total, we sampled 70, 72 and 69 parameters for the three fits, where we obtained 64077, 54258, and 52522 posterior samples for the three models, respectively.

In all three models we place a Gaussian prior on the stellar density of $\rho = 14.7 \pm 1.5 \,\mathrm{g \, cm^{-3}}$ estimated from the mass and radius of the star in Table 2. In doing so, allows us to place an important constraint on the eccentricity and argument of periastron derived from the transit from the photo-eccentric effect (Dawson & Johnson 2012). Mann et al. (2016a) and Thao et al. (2020) also report evidence of non-zero eccentricity in the system from their analyses of $e = 0.27^{+0.16}_{-0.21}$, and $e = 0.27^{+0.16}_{-0.06}$, respectively. Our data also favors an eccentric solution. In performing a circular fit placing no constraint on the stellar density, the resulting best-fit fit yielded a stellar density of $\rho = 40 \pm 4 \,\mathrm{g \, cm^{-3}}$, significantly larger than the expected stellar density of $\rho = 14.7 \pm 1.5 \,\mathrm{g \, cm^{-3}}$. Therefore, to ensure an a/R_* value that results in a stellar density consistent with our estimated stellar density derived from the stars mass and radius, we placed an informative Gaussian prior on the stellar density in all fits reported in Table 5.

To estimate the expected photometric amplitude from ellipsoidal variations caused by tidal interactions between the planet and the host star, we used Equation 7 in Shporer (2017), which gives the expected ellipsoidal variation amplitude in parts-per-million as a function of the stellar mass, planetary mass, the orbital distance of the planet, and the limb darkening and gravity darkening parameters of the host star. Using our parameter constraints, nominal vales for the limb-darkening and gravity darkening parameters, and assuming a circular orbit with an $a/R_* = 21$, we obtain an expected ellipsoidal variation amplitude of 0.03ppm. Even if we assume a circular orbit at K2-25b's periastron distance of $a/R_* \sim 11.5$, we obtain a small expected ellipsoidal variation amplitude of 0.2ppm. Both values are substantially below the photometric precision of our datasets. Given the low amplitude of the signal, we did not attempt to fit any ellipsoidal variations as part of our transit model.

Table 5. Median values and 68% credible intervals for the three joint-fit models considered. We adopt the values for Model RV2 as it is statistically favored over the other two models.

Parameter	Description	Model RV1 $(e = 0, \text{ with GP for RVs})$	Model RV2 (adopted) ($e \neq 0$, with GP for RVs)	Model RV3 $(e \neq 0, \text{ no GP for RVs})$
Model Evidence:	:			
$\ln Z$	Model log evidence	25828.7 ± 9.9	25875.4 ± 4.1	25869.9 ± 4.2
Planet Parameter	s:			
T_C (BJD _{TDB})	Transit Midpoint	$2458515.64215^{+0.00008}_{-0.00008}$	$2458515.64206^{+0.00010}_{-0.00009}$	2458515.642134 ^{+0.00008} _{-0.00008}
P	Orbital period (days)	$3.48456408^{+0.0000005}_{-0.0000005}$	$3.48456408^{+0.0000006}_{-0.0000005}$	$3.48456407^{+0.0000005}_{-0.0000006}$
(R_p/R_*)	Radius ratio	$0.1146^{+0.0010}_{-0.0011}$	$0.1075^{+0.0018}_{-0.0018}$	$0.108^{+0.0018}_{-0.0019}$
R_p	Planet radius (R_{\oplus})	$3.66_{-0.12}^{+0.12}$	$3.44^{+0.12}_{-0.12}$	$3.45^{+0.12}_{-0.12}$
R_p	Planet radius (R_J)	$0.327^{+0.011}_{-0.011}$	$0.306^{+0.011}_{-0.011}$	$0.308^{+0.011}_{-0.011}$
δ_p	Transit depth	$0.01314^{+0.00024}_{-0.00025}$	$0.01155^{+0.00038}_{-0.00039}$	$0.01167^{+0.00040}_{-0.00041}$
a/R_*	Scaled semi-major axis	$24.47^{+0.33}_{-0.33}$	$21.09^{+0.57}_{-0.59}$	$21.29^{+0.55}_{-0.64}$
a	Semi-major axis (from a/R_* and R_*)	$0.0334^{+0.0012}_{-0.0011}$	$0.0287^{+0.0012}_{-0.0012}$	$0.029^{+0.0012}_{-0.0012}$
i	Transit inclination (°)	$88.068^{+0.046}_{-0.046}$	$87.16^{+0.18}_{-0.21}$	$87.24^{+0.18}_{-0.23}$
b	Impact parameter	$0.8252^{+0.0092}_{-0.0096}$	$0.628^{+0.032}_{-0.037}$	$0.619^{+0.037}_{-0.043}$
e	Eccentricity	$0.0^{+0.0}_{-0.0}$	$0.428^{+0.050}_{-0.049}$	$0.409^{+0.041}_{-0.039}$
ω	Argument of periastron (°)	$90.0^{+0.0}_{-0.0}$	$120.0^{+12.0}_{-14.0}$	$106.0^{+13.0}_{-16.0}$
$T_{ m eq}$	Equilibrium temp. (assuming $a = 0$)	458.3 ^{+8.8} _{-8.7}	$494.0^{+11.0}_{-11.0}$	$492.0^{+11.0}_{-11.0}$
$T_{ m eq}$	Equilibrium temp. (assuming $a = 0.3$)	$320.8^{+6.2}_{-6.1}$	$345.7^{+8.0}_{-7.8}$	$344.3^{+8.0}_{-7.8}$
S	Insolation Flux (S_{\oplus})	$7.35^{+0.58}_{-0.55}$	$9.91^{+0.95}_{-0.86}$	$9.75^{+0.94}_{-0.85}$
T_{14}	Transit duration (days)	$0.03398^{+0.00027}_{-0.00024}$	$0.03182^{+0.00036}_{-0.00037}$	$0.03162^{+0.00037}_{-0.00035}$
T_{23}	Transit duration (days)	$0.01454^{+0.00091}_{-0.00090}$	$0.02212^{+0.00063}_{-0.00065}$	$0.02208^{+0.00068}_{-0.00071}$
au	Ingress/egress duration (days)	$0.00972^{+0.00042}_{-0.00042}$	$0.00485^{+0.00040}_{-0.00039}$	$0.00477^{+0.00046}_{-0.00042}$
K	RV semi-amplitude (m/s)	$24.7^{+7.2}_{-6.8}$	$27.9^{+6.5}_{-6.0}$	$32.2^{+9.7}_{-9.4}$
m_p	Planet mass (M_{\oplus})	$24.0^{+7.1}_{-6.6}$	$24.5^{+5.7}_{-5.2}$	$28.5^{+8.5}_{-8.3}$
$\sigma_{ ext{HPF}}$	HPF RV jitter (m/s)	$13.9^{+6.9}_{-5.3}$	$1.8^{+5.1}_{-1.4}$	$42.9^{+7.2}_{-6.5}$
γ	HPF RV offset (m/s)	$-5.0^{+13.0}_{-12.0}$	$4.0^{+16.0}_{-16.0}$	$-4.0^{+7.5}_{-7.5}$
Derived Stellar P	arameters:			
ρ	Stellar density (g cm ⁻³)	$22.83^{+0.92}_{-0.91}$	$14.6^{+1.2}_{-1.2}$	$15.0^{+1.2}_{-1.3}$
GP Hyperparame	eters for RVs:			
$\sigma_{ ext{GP}}$	GP RV amplitude (m/s)	$34.8^{+8.6}_{-6.6}$	$41.0^{+11.0}_{-7.9}$	-
α	GP inverse timescale (day ⁻²)	$13^{+90}_{-10} \times 10^{-6}$	$15^{+70}_{-1} \times 10^{-8}$	-
Γ	GP Frequency Structure parameter	$8.0^{+1.1}_{-1.1}$	$6.75^{+0.93}_{-1.2}$	-
P_{GP}	GP Kernel Periodicity parameter (days)	1.88219 ^{+0.00034} _{-0.00032}	1.88203 ^{+0.00032} _{-0.00031}	_

6.2. Derived Parameters

Figure 2 shows the *K*2 transits along with the 9 ground-based transits observed and analyzed in this work. The transits on nights of December 21, 2018 and January 4, 2019 were performed simultaneously with the RM effect observations further discussed in Section 7. The shorter cadence of the ground-based observations compared to the 30 minute cadence *K*2 observations allows us to resolve the transit shape better, resulting in more precise planet parameters. Table 5 compares the best-fit parameters from the three models considered in this work.

Figure 3 compares the resulting phased RV plots for the three different models considered. The derived best-fit RV semi-amplitudes for the three models are $K_{\rm RV1} = 24.7^{+7.2}_{-6.8} \, {\rm m/s}, \ K_{\rm RV2} = 27.9^{+6.5}_{-6.0} \, {\rm m/s}, \ {\rm and} \ K_{\rm RV3} = 32.2^{+9.7}_{-9.4} \, {\rm m/s},$

respectively. This results in mass estimates of $M_{\rm RV1} = 24.0^{+7.1}_{-6.6}\,\rm M_{\oplus}$, $M_{\rm RV2} = 24.5^{+5.7}_{-5.2}\,\rm M_{\oplus}$, and $M_{\rm RV3} = 28.5^{+8.5}_{-8.3}\,\rm M_{\oplus}$ for the three models, respectively. We see that all three models result in consistent mass estimates. We note that the GP RV amplitudes for Model RV1 and RV2 are similar with $\sigma_{\rm GP,RV1} = 35^{+9}_{-7}\,\rm m/s$, and $\sigma_{\rm GP,RV2} = 41^{+11}_{-8}\,\rm m/s$. As expected, we see that for Model RV3—which does not employ a GP—the white noise error term significantly increases to compensate for the additional correlated-noise, with $\sigma_{\rm w,RV1} = 13.9^{+6.9}_{-5.3}\,\rm m/s$, $\sigma_{\rm w,RV2} = 1.8^{+5.1}_{-1.4}\,\rm m/s$, and $\sigma_{\rm w,RV3} = 42.9^{+7.2}_{-6.5}\,\rm m/s$.

To check which model between models RV1, RV2, and RV3, is statistically favored, we use the log-evidence values calculated by the dynesty dynamic nested sampler. As the estimated errors of the log-evidences can be underestimated by nested sampling algorithms (see e.g., Nelson et al. 2020; Espinoza et al. 2019), to get an accurate estimate of the distri-




Figure 2. a) K2 light curve of K2-25 corrected for instrument systematics using the Everest pipeline, showing clear starspot induced rotational modulations. b) Detrended and flattened K2 light curve shown in black points using the best-fit Gaussian Process model from Model RV2 in Table 5, along with the best-fit transit model from Juliet shown in red. c) Phased K2 light curve (black) along with the best-fit transit model (red). d)-g) Ground-based transits as observed with HDI showing the best-fit transit model from Juliet in red. We note that the HDI transit on 20181221 (g) was observed without the diffuser during sub-optimal observing conditions (Moon illumination 97%). Although the light curve was modelled unbinned, it is shown here binned to a cadence of \sim 6 minutes for clarity. h-l) Ground-based transits observed with the 3.5m ARC Telescope at Apache Point. All of the ARCTIC observations were observed using a diffuser, except the transit on the night of 20190118 UT (j).

bution in the log-evidence values for each model, we ran each joint-fit model RV1, RV2, RV3, six separate times. In doing so, we obtain log evidence values of $\ln Z = 25828.7 \pm 9.9$, $\ln Z = 25875.4 \pm 4.1$, and $\ln Z = 25869.9 \pm 4.2$, respectively, where the value reported is the mean of the 6 runs for each model and the uncertainty estimate is the standard deviation of the 6 runs. We note that the scatter in the log-evidence values (\sim 10 for Model RV1, and \sim 4 for Models RV2 and RV3) is substantially larger than the internal uncertainty estimate of ~ 0.5 reported from the dynesty sampler for each individual run. Even so, within each group of 6 runs for Models RV1, RV2, and RV3, the resulting output posteriors are fully consistent. Between the three models, we see that the two eccentric Models RV2 and RV3 are favored over the circular Model RV1. Between Models RV2 and RV3, we see a statistical preference ($\Delta \ln Z \sim 5$) in favor of Model RV2. From Table 5, we see that the derived planet parameters, including the semi-amplitude, mass, and eccentricity for Models RV2 and RV3 are fully consistent. Given the statistical preference, and the fact we know that K2-25 is a young and active star with a well characterized stellar rotation period that is explicitly modelled in model RV2, we elect to adopt the values from Model RV2. We compared the distribution of the bestfit residuals of the HPF RVs from model RV2 to a normal

distribution using the Kolmogorov-Smirnov test, from which we obtain a p-value of 6.5×10^{-8} . This suggests that the distribution of the RV residuals is indistinguishable from a normal distribution, and that our model (1 planet Keplerian along with a quasi-periodic GP) can accurately model the observed RVs.

6.3. Stellar Activity from HPF Spectra

To further study the activity of the star, we measured a number of stellar activity indicators from the HPF spectra. Figure 4 shows the Generalized Lomb-Scargle (LS) periodograms of the out-of-transit RVs we used for our mass measurement, along with an array of different activity indicators measured from the HPF spectra, including: the Differential Line Width (dLW), the Chromatic Index (CRX), and line indices of the three Calcium II Infrared Triplet (Ca II IRT) lines. To calculate the LS periodograms, we used the periodogram functions in the astropy.timeseries package, and the false alarm probabilities⁵ were calculated using

⁵ Although the False Alarm Probability is a commonly used in periodogram analysis in radial velocity data it has known limitations (see e.g., discussion in Fischer et al. 2016).




Figure 3. Out-of-transit RVs of K2-25 from HPF comparing the resulting RV models from a) Model RV1, b) Model RV2, and c) Model RV3, with associated residuals shown in the corresponding lower panels. The 50th percentile best-fit models are shown in red. The grey shaded regions show the 1σ (grey) and 3σ (lighter grey) estimated model confidence bands. The log-evidence for the three models are $\ln Z = 2582.7 \pm 9.9$, $\ln Z = 25875.4 \pm 4.1$, and $\ln Z = 25869.9 \pm 4.2$, respectively, showing a significant preference for the eccentric models (Models RV2 and RV3). We adopt Model RV2 due to the statistical preference over Model RV3, with $\Delta \ln Z \sim 5$ over Model RV3. d-e) The non-phased RVs as a function of time for two observing Seasons (approximately a year apart). The red curve shows the best-fit GP and Keplerian model. The grey shaded regions show the 1σ (grey) and 3σ (lighter grey) estimated model confidence bands.

the bootstrap method implemented in this same package. Additionally, in Figure 4, we show the Window Function (WF) of our RV observations. All of the periodograms in Figure 4 are normalized using the formalism in Zechmeister & Kürster (2009), except the window function is normalized such that the highest peak has a power of 1. Table C3 in the appendix lists the values of the RVs and the activity indicators used in this work.

To measure these indicators, we follow the definition and procedures in the SERVAL pipeline (Zechmeister et al. 2018). To measure the Ca II IRT indicators, we measure the mean flux in a 30km/s wide region centered on the three Ca II IRT line centers and we use two offset bands (to the right and to the left of the line center, 100km/s wide) as reference regions to measure line indices as defined in Equation 27 in (Zechmeister et al. 2018),

$$I = \frac{\langle f_0 \rangle}{0.5(\langle f_1 \rangle + \langle f_2 \rangle)},\tag{5}$$

where $\langle f_0 \rangle$ is the mean flux around the line center, and $\langle f_1 \rangle$ and $\langle f_2 \rangle$ are the mean fluxes around the reference regions. The exact locations of the line center regions and the offset regions we used are given in Table A1 in the Appendix.

From Figure 4a, we see a clear peak in the RV periodogram at the known rotation period, indicative of rotationally modulated RV variations e.g., due to starspots. This interpretation is further strengthened by the fact that we see a clear peak in the periodogram of the dLW indicator at the known rotation period (Figure 4c). In Figure 4b, we show the periodogram of the RVs after subtracting the best-fit GP activity model, demonstrating that after removing the GP model, the peak at the stellar rotation is significantly suppressed and the peak at the planet period increases in significance. The CRX indicator does not show any clear evidence of periodic variations in the chromaticity of the RVs at either the stellar rotation period or the planet period. Interestingly, in Figures 4e, f, and g, we do not see clear peaks in the Ca II IRT activity indicators at the rotation period, but rather, we see a clear peak in all three indices at 2.46 days. We speculate that this could indicate that the active chromospheric regions the Ca II IRT lines trace, could have a different characteristic evolution timescale than the rotation period of the star.

6.4. RV Injection and Recovery Tests

To test that our RV precision and the RV sampling is sufficient to accurately constrain the Keplerian parameters of K2-25b, we performed two series of injection and recovery tests, broadly following the methodology in Klein & Donati (2020). For both series of tests, we injected a signal with known Keplerian parameters (P, T_C, K, ω, e) , and Gaussian process hyper-parameters $(P_{GP}, \sigma_{GP}, \alpha_{GP}, \Gamma_{GP})$, along with a white noise parameter (σ_w) . The values of the parameters for the two tests, along with the priors used for the recovery, are given in Table B2 in the Appendix. We placed informative Gaussian priors on the orbital period (P), transit center (T_C) , and the GP period (P_{GP}) , and the GP frequency structure parameter (Γ_{GP}) , and we placed broad uninformative priors on the other parameters (same priors as




Figure 4. Lomb-Scargle periodograms of the HPF RVs along with different activity indicators. The stellar rotation period ($P = 1.88 \, \text{days}$) and the planet period ($P = 3.48 \, \text{days}$) are highlighted with the dashed blue and red lines, respectively. False alarm probabilities of 1% and 0.1% calculated using a bootstrap method are denoted with the grey solid and grey dashed lines, respectively. a) the HPF out-of-transit RVs used for the mass measurement of K2-25b. b) same as a) but after removing the best-fit GP model from Model RV2. This shows that the known planet peak becomes more significant. c) Differential line width (dLW) activity indicator, showing a clear peak at the known rotation period. d) Chromatic Index activity indicator. e-g) Ca II IRT indices for the three Ca II IRT lines. h) The window function of the HPF RVs, showing a clear sampling peak at 1 day. The power in a-g) is normalized using the formalism in Zechmeister & Kürster (2009), and h) is normalized so that the highest peak is unity.

used for fit RV2). For the first series of tests (Test I), we set the GP amplitude $\sigma_{\rm GP}=42\,{\rm m/s}$, and other parameters to values similar to the nominal 50th quantile values from our adopted RV2 fit in Table 5 (see exact values injected in Table B2 in the Appendix). For the second series of tests (Test II), we increased the injected GP amplitude to its 95th percentile value from our adopted RV2 model in Table 5, corresponding to $\sigma_{\rm GP}=64.6\,{\rm m/s}$, to check if the injected Keplerian parameters—in particular K and e—could be reliably recovered if the level of the correlated noise is higher. For both tests, we repeated the injection and recovery 200 times, and we then inspect the resulting posteriors calculated using juliet from each individual run.

Figure B1 in the Appendix shows the distribution of mean values from Test I for a few select parameters of interest: K, ω , e, σ_{GP} , α_{GP} , and σ_{w} . We see that for all parameters, the

distribution of the recovered values is fully consistent with the known injected value. Further, Figure B2 in the Appendix compares the distribution of all 200 posteriors for σ_{GP} , K, and e, for both series of injection and recovery tests. To compare the distribution of the synthetic residuals to a normal distribution, we used the Kolmogorov-Smirnov test, from which we see that all datasets have a p-value $< 10^{-5}$ (with most datasets showing a p value $< 10^{-7}$), suggesting that the best-fit residuals from the synthetic tests are indistinguishable from a normal distribution. Although σ_{GP} shows broad posteriors and is relatively poorly constrained, in both series of tests, the true values of K and e are consistently recovered. For Test I, which assumes $\sigma_{GP} = 42 \,\mathrm{m/s}$, the true value of K is within the 68% and 95% credible intervals in 80% and 99.7% of the cases, respectively. For Test II, even with the higher injected value of $\sigma_{GP} = 64.6 \,\mathrm{m/s}$, the true value of K is reliably recovered within the 68% and 95% credible intervals in 75% and 97% of the cases, respectively. We conclude that the number and the sampling of the HPF RVs is sufficient to accurately constrain the Keplerian planet parameters.

7. THE RM EFFECT OF K2-25

7.1. RM model

We model the Rossiter-McLaughlin effect using the prescription given in Hirano et al. (2010) and Hirano et al. (2011). Specifically, we use Equation 26 from Hirano et al. (2011), which gives the RM velocity anomaly as,

$$\Delta v = -\left(\frac{2(\beta + \sigma_{\rm RM})^2}{2\beta^2 + \sigma_{\rm RM}^2}\right)^{3/2} f v_p \left(1 - \frac{v_p^2}{(2\beta^2 + \sigma_{\rm RM}^2)} + \frac{v_p^4}{2(2\beta^2 + \sigma_{\rm RM}^2)^2}\right), \tag{6}$$

where β indicates the best-fit Gaussian dispersion (in km/s) of the intrinsic line broadening in the absence of stellar rotation, and σ_{RM} indicates the Gaussian line width component arising from stellar rotation (see Hirano et al. 2010 for details). Here we set β to the width of the HPF resolution element, i.e., $\beta = 5.45 \pm 0.5$ km/s, where the errorbar is to account for any effects of macroturbulence and/or other nonstellar rotation processes that could broaden the line profile. The f parameter denotes the fraction of the star being blocked by the planet during the transit as a function of time (i.e., f = 1 - F, where F is the photometric transit model), and v_p denotes the sub-planet velocity (in km/s) i.e., the velocity of the star being blocked by the planet as a function of time during the transit. As discussed in Hirano et al. (2011), $\sigma_{\rm RM}$ describes the dispersion of a Gaussian approximating the stellar rotational kernel, and here we follow Hirano et al. (2010) and assume $\sigma_{\rm RM} = v \sin i_* / 1.31$. We neglect any differential rotation, as Dmitrienko & Savanov (2017) showed that the differential rotation of K2-25b is small, or $\Delta\Omega = 0.0071 \pm 0.002 \,\mathrm{rad/day}$, and thus negligible during the

To model the RM effect, we model all three transits jointly. For the fit, we included the bulk RV motion imposing informative priors on the Keplerian orbital parameters from our best-fit orbital values from Model RV2 in Table 5. In addition, we placed an informative Gaussian prior on the $v \sin i_*$ using our $v \sin i_*$ value measured from the HPF spectra. To

account for possible systematics between the transits (e.g., due to stellar activity and/or instrumental systematics), we added an independent RV offset parameter γ for each individual transit. We sampled the limb-darkening parameters using the q_1 and q_2 parametrization described in Kipping (2013), and fully sample them across the whole valid range from 0 to 1 to minimize any biases on the RM effect anomaly due to limb-darkening effects. To account for the smoothening of the finite exposure times of our RV observations, we supersampled the model 7 times and resampled the model according to the exposure time of \sim 300s. We calculated the bulk RV model using the radvel python package (Fulton et al. 2018). To calculate the transit fraction f, we used the batman package (Kreidberg 2015), calculating the transit flux and setting f = 1 - F, where F is the transit model from batman. Before we started the MCMC sampling, we found the global most probable solution using the PyDE differential evolution optimizer (Parviainen 2016). We then initialized 100 MCMC walkers in the vicinity of the global most probable solution using the emcee MCMC affine-invariant MCMC sampling package (Foreman-Mackey et al. 2013). We ran the 100 walkers for 25,000 steps, and after removing the first 2,000 steps as burn-in and thinning the chains by a factor of 100, the Gelman-Rubin statistic of the resulting chains was within < 1% of unity, which we consider wellmixed.

To test the impact of our assumptions about stellar activity on λ , we performed four additional fits:

- A joint fit with a single RV offset parameter γ .
- A joint fit with a single RV offset parameter γ after subtracting our best-fit GP activity model (from Model RV2 in Table 5) from the RVs.
- A joint fit with three independent RV offset parameters γ and removing the prior on the semi-amplitude.
- A joint fit with three independent RV offset parameters γ and removing the prior on the $v \sin i_*$.

Table 6. Median values and associated 16th and 84th percentile values from our RM fit.

Parameter	Description	Value
Model Likeli	hood Parameters:	
$\chi^2_ u$	Reduced χ^2	1.04
DOF	Degrees of freedom	25
MCMC Para	meters:	
λ	Sky-projected obliquity (°)	3 ± 16
$v \sin i_*$	Rotational velocity (km/s)	8.9 ± 0.6
γ_1	RV offset (m/s)	-10^{+28}_{-28}
γ_2	RV offset (m/s)	58^{+27}_{-26}
γ_3	RV offset (m/s)	-95^{+29}_{-28}
q_1	Linear Limb darkening parameter	$0.47^{+0.35}_{-0.33}$
q_2	Quadratic Limb darkening parameter	$0.44^{+0.36}_{-0.31}$
β	Intrinsic stellar line width	$5.41^{+0.49}_{-0.50}$
Derived Para	meters:	
ψ	3D obliquity (°)	17^{+11}_{-8}




Figure 5. Upper panels: Three RM effect transits as observed with HPF (black points), along with the best-fit joint RM effect model shown in red. The RM model includes the bulk RV shift and an independent RV offset parameter γ is included for each transit to account for stellar activity and/or instrumental effects. The grey shaded regions show the 1σ , 2σ , and 3σ confidence regions. Lower panels: Two simultaneous photometric transits from HDI (Dec 21, 2018), and ARCTIC (Jan 4, 2019) shown in black, along with the best-fit transit model (model RV2 in Table 4 and 5). The HDI observations were performed without a diffuser due to the Moon contamination that night. No starspot crossing events are seen in the simultaneous transit observations, which otherwise would complicate the RM-effect analysis. Figure 6 shows the three RM effect observations phased together and binned to 5 minute bins for further visual inspection.

All fits resulted in fully consistent constraints on λ suggesting a well-aligned system, although we note that the fit with a single RV offset parameter resulted in slightly higher uncertainty estimate on λ , or a constraint of $\lambda = 0 \pm 24^{\circ}$. From the GP activity model constrained from our out-oftransit RVs, we note that the expected RV variation during the three transits observed is slowly varying and significantly smaller ($< 5 \,\mathrm{m/s}$) than the observed RM amplitude of \sim 65 m/s, and thus is effectively modeled out with independent offsets between the transits. In comparing the Baysian Information Criterion (BIC) and Akaike Information Criterion (AIC; Akaike 1974), the models show a statistical preference ($\triangle AIC \sim 9$) for models that allowed for independent offsets between the transits. We note that using the BIC and AIC in this case is heuristic, as more datapoints would be formally needed for these criteria to be in the asymptotic regime where they are accurate and justified for model comparison. To allow for flexibility to take out potential systematic offsets between the three transits, we report the resulting posterior constraint from the fit assuming three independent RV offset γ parameters in Table 6. We further note that the fit where we allowed $v \sin i_*$ to vary freely resulted in a fully consistent λ value and yielded a $v \sin i_* = 11.6 \pm 3 \,\mathrm{km/s}$, which is consistent with the as-measured $v \sin i_* = 8.8 \pm 0.6$ km/s from the HPF-spectra at the 1σ level.

7.2. Results

Figure 5 shows the three RM effect observations using HPF along with our best-fit RM model (RM amplitude

 \sim 65 m/s), along with 1σ , 2σ , and 3σ shaded regions. Table 6 summarizes the resulting best-fit posterior values, showing that we obtain a sky-projected obliquity constraint of $\lambda = 3 \pm 16^{\circ}$. To further visualize and compare the observed RVs to the best-fit RM model, in Figure 6 we show all of the three transits in Figure 5 phased to the transit ephemeris and binned to a 5 minute cadence using a weighted average. The resulting median RV error is 55 m/s in the 5 minute bins. The RM model is shown with the bulk RV model and RV offset parameters removed for clarity.

In addition to the sky-projected obliquity listed in Table 6, we calculate the true obliquity angle ψ , using the following equation,

$$\cos \psi = \sin i_* \cos \lambda \sin i + \cos i_* \cos i, \tag{7}$$

where i_* is the stellar inclination, i is the transit inclination, and λ is the sky-projected obliquity angle. Using Equation 7 above and our $\lambda = 3 \pm 16^{\circ}$ constraint from Table 6, our $i = 87.13 \pm 0.17^{\circ}$ from Table 5 (Model RV2), and our $i_* = 90 \pm 12^{\circ}$ constraint from Table 2, we obtain the following constraint on $\psi = 17^{+11}_{-8}$ °. Integrating the resulting posterior, we can say that $\psi < 30^{\circ}$ with 89% confidence, which is compatible with a well aligned system.

8. DISCUSSION

With its known age and characterized 3D orbital architecture, K2-25b is an interesting laboratory to test different formation scenarios. Below, in Section 8.1, we compare our improved ephemerides to the ephemerides of other recent lit-




Figure 6. The RM effect of K2-25b after combining the three jointly fitted RM effect observations shown in the upper panels in Figure 5. The best-fit RM model (red curve) has the bulk RV motion and RV offset parameters removed. The black points show the data binned to a 5min cadence. The grey shaded regions show the 1σ , 2σ and 3σ confidence bands.




Figure 7. Updated transit ephemeris for K2-25b into the JWST era. Our ephemeris derived from the K2 photometry only (blue shaded region) agrees well with the ephemeris provided in Mann et al. (2016a) (purple). Our improved ephemeris (Model RV2 in Table 5) results in an error of \sim 20sec at the beginning of the JWST era in nominally 2021, and is in excellent agreement with the ephemeris provided in Thao et al. (2020) derived from data from K2, Spitzer, and additional ground-based observations. The shaded regions show the 1σ error estimates.

erature on this system. In Section 8.2, we discuss the composition of K2-25b showing that at its observed mass and radius it is consistent with a water-rich world, or a rocky core with a small H/He envelope. In Section 8.3, using our mass constraint we discuss the possibility for atmospheric characterization through transmission spectroscopy with JWST in the future. We discuss our obliquity constraint in Section 8.4, which is suggestive of a well-aligned orbit, and place it in context to other measurements in the literature. Finally, given our detailed characterization of the different orbital parameters of the planet allows us to place informative constraints on different formation scenarios, which we discuss in Section 8.6.1.

8.1. Improved Ephemeris

Figure 7 shows the improved ephemeris derived by jointly modeling the K2 and ground-based transits (Model RV2 in Table 5). In blue we show our ephemeris derived from the K2-data only $(T_{0.K2} = 2457062.5790 \pm 0.0005)$ and $P_{K2} =$ 3.484547 ± 0.000040), which agrees well with the ephemeris derived in Mann et al. (2016a) also from the K2 data. Our K2only ephemeris results in a transit timing uncertainty of \sim 35-40minutes at the start of the JWST era nominally in 2021. Further, we see that our joint-fit ephemeris (shown in black in Figure 7) is fully consistent within the 1σ errorbars of our K2-only ephemeris. Our joint-fit ephemeris results in a factor of \sim 150 improvement in the transit timing precision from the K2 data only, yielding a timing uncertainty of $\sim 20 \,\mathrm{sec}$ at the start of the JWST era nominally in 2021, which will be important for scheduling follow-up observations in the future. Additionally, Figure 7 shows that our joint-fit is in excellent agreement with the transit ephemeris presented in Thao et al. (2020) derived from photometry from K2, Spitzer and ground-based observations.

8.2. Composition

To compare the possible composition of K2-25b to other planets, in Figure 8 we plot K2-25b along with other planets in the exoplanet mass-radius plane⁶. In Figure 8, we only show planets with fractional errors on mass and radius that are better than 25%, as otherwise, their mass and radius values are consistent with a wide array of planet composition models. The shaded grey region indicates planets with iron content exceeding the maximum value predicted from models of collisional stripping (Marcus et al. 2010). The solid lines are theoretical mass-radius curves assuming a constant density from Zeng et al. (2019). From Figure 8, we see that at a radius of $R = 3.44R_{\oplus}$, K2-25b is similar in size to two other well-studied M-dwarf planets: GJ 436b (4.2 R_{\oplus} ; Maciejewski et al. 2014), and GJ 3470b (4.57 R_{\oplus} ; Awiphan et al. 2016), which we highlight in orange in Figure 8. With a mass of $M = 24^{+5.7}_{-5.2} M_{\oplus}$, K2-25b is similar in mass to GJ 436b $(M = 22.1 \,\mathrm{M}_{\oplus}; \,\mathrm{Maciejewski} \,\mathrm{et \,al.} \,\,2014), \,\mathrm{but \,more \,massive}$ than GJ 3470b ($M = 13.9 \,\mathrm{M}_{\oplus}$; Awiphan et al. 2016). Interestingly, both GJ 436b and GJ 3470b are known to be experiencing substantial atmospheric outflows, resulting in significant atmospheric mass-loss throughout their lifetime (see e.g., Ehrenreich et al. 2015, and Bourrier et al. 2018a; Ninan et al. 2019, respectively). The possibility that K2-25b is experiencing outflows is further discussed in Subsection 8.3.

There are degeneracies in the composition of planets with radii between $2-4R_{\oplus}$: these planets could either have rocky cores with H/He envelopes, or they could be water-rich worlds that contain a significant amount of multi-component water dominated ices/fluids in addition to rock and gas (Zeng et al. 2019). As such, from knowing the masses and radii of a planet alone, we can not discern between the different solutions (Adams et al. 2008).

⁶ Data retrieved from the NASA Exoplanet Archive Akeson et al. (2013) in November 2019




Figure 8. K2-25b compared to other similar-sized planets in the exoplanet mass radius plane for M-dwarf planets. M-dwarf planets ($T_{\rm eff}$ < 4000 K) are denoted with the filled black circles, and planets orbiting hotter stars are shown with the faint grey circles. Blue squares shows solar system planets. The red curve shows our best-fit model from interpolating the two-layer composition model of Lopez & Fortney (2014) assuming a rocky core capped by a H/He envelope, resulting in a H/He mass fraction of 5%. The red shaded region shows the associated 68% credible interval. The other solid lines show the composition models of Zeng et al. (2019). The shaded grey region indicates planets with iron content exceeding the maximum value predicted from models of collisional stripping (Marcus et al. 2010). K2-25b is similar in size to the well-studied M-dwarf planets GJ 436b and GJ 3470b.

Given this known degeneracy, to explore the range of possible compositions for K2-25b, we overplot a number of different compositional growth models in the exoplanet massradius plane from Zeng et al. (2019). To place a quantitative estimate on the possible H/He fraction of K2-25b, we modeled the composition of K2-25b assuming a two-layer thermal model consisting of a rocky core and a H/He atmosphere using the models presented in Lopez & Fortney (2014). In this model, the H/He envelope is the dominant driver of the size of the planet. Assuming this two component model, we linearly interpolated the tables presented in Lopez & Fortney (2014) and together with the posteriors in the observed mass and radius of K2-25b, we estimate that K2-25b has an envelope mass fraction of $5.3^{+1.2}_{-0.9}\%$. The 50th percentile model that best fits the observed mass and radius constraints of K2-25b is shown in the red curve in Figure 8, with the 1σ error intervals denoted by the red shaded region.

Despite these degeneracies, with more mass measurements of planets in young clusters (see e.g., Barragán et al. 2019), we can start to gain further insights onto the exoplanet Mass-Radius distribution as a function of age, which can help place further constraints on planet formation mechanisms and how time-dependent processes such as photo-evaporation sculpt exoplanet Mass-Radius plane.

8.3. Prospects for Transmission Spectroscopy

With its large transit depth, and brightness at NIR wavelengths (J = 11.3), K2-25b has been mentioned as a prime

candidate for transmission spectroscopy (e.g., Mann et al. 2016a; Thao et al. 2020). Thao et al. (2020) measured and studied the transit depths of K2-25b in different broad-band filters via precision ground-based photometry, and spacebased photometry from Kepler, and Spitzer. In Figure 9, we plot the transit depth measurements from Thao et al. (2020) along with our transit depth measurements in the SDSS i' and SDSS z' bands. As our transit depth measurement presented in Table 5 assumed a single transit depth for all bands, to measure these transit depths, we conducted two separate transit fits using the Juliet program: first, we jointly modeled our five SDSS z' band transits, and second, we jointly modeled our four SDSS i' band transits. From Figure 9, we see that our transit depths in these bands agree well with the optical transit depth measurements in Thao et al. (2020). However, as is seen in Figure 9, the infrared *Spitzer* transits from Thao et al. (2020) are statistically smaller than the optical transits. As is detailed in Thao et al. (2020), this could potentially be due to starspots causing the transit depths to be less deep. However, like they argue, the impact of starspots on the transit depth in the NIR is likely lower than in the optical. Further, no clear starspot crossing events are observed in their data, the data studied in Kain et al. (2019) (although Kain et al. (2019) mention a few candidate starspot crossing events) or in the transits presented here. As concluded by Thao et al. (2020), while spots can have an impact on the overall transmission spectrum, it is unlikely that starspots alone could cause the difference in the NIR transit




Figure 9. Comparison of transit depth as a function of wavelength for K2-25b, from Thao et al. (2020) (black points) and from this work (blue points) in the SDSS i' and z' bands. We further overlay two expected cloud-free transmission spectra as calculated with the pandexo tool for our measured mass of $25M_{\oplus}$ and compare that with the case using the predicted mass of $11.5M_{\oplus}$ estimated using the mass-radius relation in the Forecaster package. As expected, our higher mass increases the surface gravity of the planet, which further mutes the expected transmission spectroscopic features.

depths. One possible explanation, as mentioned in Thao et al. (2020) is that K2-25b has a predominantly cloudy or hazy atmosphere, causing a flat transmission curve as a function of wavelength.

Thao et al. (2020) use the parametric mass-radius relation from Wolfgang et al. (2016) to estimate the most likely mass for K2-25b of $M = 13M_{\oplus}$, and consider different transmission models assuming planet surface gravities of 6 m s⁻², 9 m s⁻², and 12 m s⁻², corresponding to planet masses of $8M_{\oplus}$, $11.5M_{\oplus}$, and $15M_{\oplus}$, respectively. All of these mass estimates are lower than our measured mass of $M = 24.5^{+5.7}_{-5.2} M_{\oplus}$, which corresponds to a surface gravity of $g = 20.3 \pm 4.7 \,\mathrm{m \, s^{-2}}$. As an independent test, we also predicted a mass of $M = 11.5^{+9.1}_{-4.8} M_{\oplus}$ using the mass-radius relation in the Forecaster package from Chen & Kipping (2017). In Figure 9, we compare the expected transmission spectrum as calculated with pandexo package (Batalha et al. 2017) for both the predicted mass of $11.5M_{\oplus}$, and our measured mass of $25M_{\oplus}$. As expected, we see that the transmission features of K2-25b are muted in the more massive case, as the increased mass increases the surface gravity of the planet. This can also be seen if we calculate the Transmission Spectroscopy Metric (TSM) defined in Kempton et al. (2018): using the predicted mass of $11.5 M_{\oplus}$ we obtain a TSM=143, while if we use our mass estimate, we estimate a lower of TSM = 66^{+21}_{-14} . Our larger mass measurement which mutes the expected transmission features of K2-25b could further help—at least partially—explain the flat transmission features observed by Thao et al. (2020).

Despite our larger mass estimate causing the expected transmission features to be muted, the cloud-rich or hazy atmosphere scenario suggested in Thao et al. (2020) is still a likely possibility. In fact, this would conform with the rising statistical trend that colder planets preferentially show muted and/or flat features in their transmission spectra Crossfield & Kreidberg (2017). However, to confidently quantify the exact amplitude of the transmission features—which strongly depend on the mass of the planet, which is now known—we argue that additional observations with JWST will be valu-

able to further confidently rule out or confirm a flat transmission spectrum (particularly the information-rich $1-2 \mu m$ region). As an example, in Figure 9, we overplot the expected S/N of JWST/NIRISS in Single Object Slitless Spectroscopy (SOSS) mode using the gr700xd grism after 5 transit observations binned to a resolving power of R = 50, showing that even for our large mass of $24.5M_{\oplus}$, JWST/NIRISS should have the sensitivity to confidently discern between a clear or cloudy atmosphere. Gaining further insights into K2-25b's atmosphere will be particularly valuable, as this will allow us to put constraints on the atmospheric constituents of this relatively young planet, giving key insights into the atmospheric composition of adolescent planets and planetary atmospheres as a function of time. We note that Wang & Dai (2019) suggest that young low-density 'super-puff' planets planets with mean density $\rho < 10^{-1} \,\mathrm{g/cm^3}$ —could be susceptible to extreme hydrodynamic mass loss which can carry large numbers of small dust particles to high altitudes, which in turn can create featureless transmission spectra. However, with K2-25b having a bulk density of $\rho = 3.3 \pm 0.8 \,\mathrm{g/cm^3}$, which is substantially larger than the cutoff for 'super-puff' planets, this scenario is unlikely to be the case.

8.4. Obliquity and Orbital Alignment

Figure 10 compares the sky-projected obliquity of K2-25b to currently available sky-projected obliquity measurements of other planetary systems from the TEPCAT⁷ database (Southworth 2011), where M-dwarfs are shaded in red and the onset of fully-convective stars is highlighted with the red solid line. Our obliquity constraint of K2-25b marks the fourth obliquity constraint of an M-dwarf planet system via the RM effect, the other being GJ 436b as measured by Bourrier et al. (2018b), AU Mic b as measured by Addison et al. (2020), Hirano et al. (2020a), and Palle et al. (2020a), and TRAPPIST-1 as measured by Hirano et al. (2020a). At

⁷ TEPCAT database: https://www.astro.keele.ac.uk/jkt/ tepcat/obliquity.html




Figure 10. Currently available sky-projected obliquity constraints λ obtained from the TEPCAT database (Southworth 2011) as a function of stellar effective temperature. Small planets ($R < 6R_{\oplus}$) are highlighted in black, and larger planets in blue. The grey area shows stellar hosts past the Kraft-break (Kraft 1967), where stars loose their outer convective layers and become fully-radiative. M-dwarf systems are shaded in red, with the onset of fully-convective stars shown with the red solid line. Currently only four M-dwarf planetary systems have their obliquity measured via the RM effect: GJ 436b (Bourrier et al. 2018b), TRAPPIST-1 (Hirano et al. 2018), AU Mic b (Addison et al. 2020; Hirano et al. 2020b; Palle et al. 2020), and K2-25b (this work).

an age of 600-800 MYr, K2-25b has an intermediate age among these systems: TRAPPIST-1 has an estimated age of 7.6 ± 2.2 GYr (Burgasser & Mamajek 2017), GJ 436 an age between 4-8 GYr (Bourrier et al. 2018b), and AU Mic is the youngest with an age of 22 MYr. Interestingly, both K2-25b and GJ 436b have eccentric orbits ($e = 0.43 \pm 0.05$ and $e = 0.1616 \pm 0.004$, respectively) and are similar in size, but they show different orbital architectures: GJ 436b observed to have a misaligned orbit while K2-25b is observed to have an aligned orbit. This could be suggestive of a different formation and subsequent dynamical history. Bourrier et al. (2018b) suggest that von Zeipel-Lidov-Kozai migration induced by a candidate perturber could explain both GJ 436b's eccentricity and misaligned orbit. However, as we do not see that K2-25b is heavily misaligned this disfavors von Zeipel-Lidov-Kozai cycles acting on the planet. We further discuss potential formation pathways for K2-25b in Subsection 8.6.

With only a few obliquity measurements of M-dwarf planetary systems published via the RM effect, the orbital architectures of individual M-dwarf planetary systems remain relatively unexplored. However, we note that statistical studies suggest that planets orbiting cooler planet hosts have orbits that are on average better aligned to their stellar equators than planets orbiting hotter stars. In comparing the rotation distribution of Kepler Objects of Interests (KOIs) hosting transiting planet candidates to a control sample of single stars without transiting planet candidates, Mazeh et al. (2015) showed that hotter stars ($T_{\rm eff} > 6000\,\rm K$) show on average lower amplitude photometric modulations suggesting a broad distribution of obliquities, while cooler KOIs $(3500 \,\mathrm{K} < T_{\mathrm{eff}} < 6000 \,\mathrm{K})$ showed on average higher amplitude modulations, suggesting well-aligned systems. This result is in broad alignment with the findings of Winn et al. (2010a) Schlaufman (2010), and Albrecht et al. (2012), that hotter stars ($T_{\rm eff} > 6250\,{\rm K}$) hosting close-in gas giants show a broad distribution of obliquities and misalignments, while cooler stars ($T_{\rm eff} < 6250\,{\rm K}$) tend to host well-aligned systems. We note that neither of these studies encompassed mid-to-late M-dwarfs. In general, across the M-dwarf spectral type—and especially for mid-to-late M-dwarfs—the occurrence of gas giants is lower than seen around FGK stars (Johnson et al. 2010; Dressing & Charbonneau 2015; Hardegree-Ullman et al. 2019). By further studying the orbital architectures of M-dwarf systems, which have a lower occurrence of massive planets, we can gain further insights into the role massive planets play in sculpting the orbital architectures across different exoplanet host stars. The advent of precise NIR spectrographs, such as HPF (Mahadevan et al. 2012, 2014), the Infrared Doppler Instrument (IRD; Kotani et al. 2018), CARMENES (Quirrenbach et al. 2018), SPIROU (Artigau et al. 2014), NIRPS (Wildi et al. 2017), GIANO-B (Claudi et al. 2018), and red-optical spectrographs on large telescopes such as MAROON-X (Seifahrt et al. 2016), ESPRESSO (Pepe 2018), and KPF (Gibson et al. 2016), is opening the doors to the ensemble study of obliquities of M-dwarfs.

8.5. Search for Additional non-transiting Planets in the HPF RVs

To place an upper limit on a potentially non-transiting second planet in the system, we performed an additional two-planet RV fit of the HPF RVs using <code>juliet</code>. As there is no obvious signs of another transiting planet in the system in the K2 data, we only considered the HPF RVs for this fit. We assumed a two planet model along with a quasi-periodic Gaussian Process to account for correlated noise due to stellar activity at the known rotation period of the star. We placed Gaussian priors on the orbital parameters of K2-25b derived from the joint fit of the RVs and the photometry, and used the same priors we placed on the Gaussian Process hyperparameters as listed in Table 4 for fits RV1 and RV2. For the hypothetical planet c, we placed broad priors on the period (Jeffreys prior from 0.5days to 50days), time of conjunction (modeled as a transit midpoint uniform from 50days before

the first RV point to 50days after the last RV point), eccentricity (uniform from 0 to 0.95), argument of periastron (uniform prior from 0° to 360°), and RV semi-amplitude (uniform from 0 to 500m/s). To check for evidence of a long-term slope, we also added a radial velocity slope with broad uniform priors.

Figure 11 shows the constraint on the orbital period and mass $(m \sin i)$ of the hypothetical planet c. From this, we see that no obvious preferred solution is found. From the posteriors, we place an upper limit on the mass of a possible secondary planet of $m \sin i < 82 M_{\oplus}$ at 99.7% confidence (3σ) for periods between 0.5days to 50days. The red curve in Figure 11 shows a running 99.7% upper limit on the mass as a function of period for smaller period bins. We further note that our constraint on an additional radial velocity slope is consistent with zero slope within the 1σ uncertainties. We additionally compared the log-evidence values we obtained from our juliet fit for the 2-planet model (2 planets, a GP to account for stellar activity, and an RV slope), to a null model assuming only K2-25b in the system (1 planet and GP to account for stellar activity). In doing so, the two planet model had a log-evidence value of $ln(Z) = -183.7 \pm 0.5$, while the null one planet model had a log-evidence value of $ln(Z) = -170.6 \pm 0.1$, where we have reported the mean and standard deviation of 6 independent runs of each model to get an accurate estimate of the spread in log-evidence values. We see that the one planet model is statistically favored with a higher evidence of $\Delta \ln(Z) = 13.1$. With the current RV data in hand, we rule out massive companions with masses $m\sin i > 82M_{\oplus}$ at 99.7% confidence with periods between 0.5 and 50 days, and conclude that we do not have sufficient evidence to claim another small short period planet in the system.

8.6. Formation and Subsequent Evolution

From our detailed constraints of K2-25b's planetary properties, including constraints on its mass, eccentricity, volatile content, and obliquity of the host star, allows us to put informative constraints on potential formation scenarios, which we discuss below.

8.6.1. Potential Formation Scenarios

In the core accretion model of planet formation, planetesimals collide to form protoplanetary cores, which then attract a gaseous envelope (e.g., Bodenheimer & Pollack 1986; Pollack et al. 1996). If the planet core becomes sufficiently massive—after reaching a critical mass M_{crit} —before the gaseous protoplanetary disk dissipates, the protoplanetary core can enter a phase of runaway gas accretion where the planet attracts a massive gaseous envelope. Although the critical core mass is typically quoted as 10 Earth masses, it can vary by orders of magnitude depending on the disk conditions and planetesimal accretion rate (see e.g., Rafikov 2006). The gaseous envelope is accreted onto the planetary core from the gaseous component of the protoplanetary disk, which only remains present for a few million years around young stars (Williams & Cieza 2011; Ribas et al. 2015), suggesting that core formation must happen within this timeframe. However, traditional core accretion models suffer




Figure 11. Posterior constraints on a hypothetical non-transiting planet c in the HPF RVs in mass $(m_c \sin i)$ and orbital period (P_c) space. We see no obvious evidence for another massive planet in the HPF RVs given the broad posterior constraints. The red curve shows a running 99.7% percentile upper limit (3σ) on the mass as a function of period. Across the full period range considered, we use the posteriors to place an upper mass limit of $82M_{\oplus}$ at 99.7% confidence for periods between 0.5 and 50days.

from predicting formation timescales for gaseous planets that are much larger than the disk dissipation timescale (Dodson-Robinson et al. 2009; Rafikov 2011).

In the pebble accretion model of planet formation, small millimeter to centimeter sized pebbles accrete onto a planetary core (Lambrechts & Johansen 2012; Lambrechts et al. 2014; Lambrechts & Lega 2017). These pebbles are marginally coupled to the nebular gas on orbital timescales, creating sufficient gas drag to enable efficient core formation possible within the disk-dissipation timescale even at large orbital separations (Lambrechts & Johansen 2012). The pebble accretion model predicts that pebble accretion terminates when the planet reaches the 'pebble isolation mass', M_{iso} , the mass when the gravity of the core is strong enough to open a gap in the disk that hinders further accretion of pebbles (Lambrechts et al. 2014). If the pebble isolation mass is greater than the critical core mass and the critical core mass is reached before the disk dissipates of gas, the massive core can rapidly accrete gas to form a gas giant. However, if the pebble isolation mass is not reached and/or if it is reached after most of the disk dissipated and/or it is below the critical core mass, the planet core consists primarily of accreted icy pebbles with a minimal H/He atmosphere. Although the exact value of the pebble isolation mass depends strongly on a number of uncertain disk properties including the aspect ratio, viscosity, local disk structure, and the pebble size, the pebble isolation mass for solar-type stars at 5 AU has been estimated to be $10-20\,\mathrm{M}_{\oplus}$ (Lambrechts et al. 2014), although

larger values are also possible depending on the assumed disk properties (see e.g., Bitsch et al. 2018). The pebble isolation mass increases with the orbital semi-major axis proportional to $a^{3/4}$ (Lambrechts et al. 2014), where a is the semi-major axis of the planetary core. This suggests that planets with massive dense cores can not form too close to the host star where the pebble isolation mass is smaller. Further, the pebble isolation mass is expected to decrease for less massive later-type stars according to the following formula from Liu et al. (2019),

$$M_{\rm iso} = 25 \left(\frac{M_*}{1 \,\mathrm{M}_\odot}\right)^{4/3} M_{\oplus},\tag{8}$$

where M_* is the mass of the host star, and assuming that the planet reaches the isolation mass at the ice-line distance. Assuming K2-25b formed at the ice-line and subsequently migrated further in, we calculate an expected isolation mass of $\sim 5M_{\oplus}$.

At its currently observed mass of 24.5 M_⊕ with a thin 5% H/He envelope, K2-25b is nominally at odds with the predictions of the core-accretion and pebble-accretion models if it reached its final mass during the full gas disk stage beyond the ice line under certain disk conditions and assumed planetesimal accretion rates. From the core-accretion model, at its currently observed mass we would have predicted K2-25b to have experienced runaway gas accretion and to have formed a gas giant, while we instead infer today only a thin H/He envelope of \sim 5%. From the viewpoint of the pebbleaccretion model, we would expect that K2-25b would have reached a nominal maximum core mass mass that is closer to the isolation mass of $\sim 5M_{\oplus}$ that we estimated from Equation 8 if it formed beyond the ice line. We again acknowledge that that the exact value of the pebble isolation mass is uncertain as it depends strongly on a number of gas disk properties for K2-25b which are not well known.

To arrive at the presently observed mass, and to explain K2-25b's low inferred H/He content and moderate eccentricity, a possible scenario is that K2-25b grew its mass through the merging of planetary cores. The merging of icy cores has been postulated to explain the observed bimodality in the mass distribution of Neptunes between 2-4 R_{\oplus} (Zeng et al. 2019). Although collisional growth could happen at K2-25b's current orbital location after e.g., in-situ formation or disk-driven migration, exciting the planet's current high eccentricity could be more easily excited at larger orbital distances (we discuss its high eccentricity and migration pathways further below). If the seed cores were composed primarily of water ices, Marcus et al. (2010) show that the collisions of two icy cores tend to stick yielding a final core of doubled mass. During collisions, it is possible that substantial fractions of H/He envelopes get stripped away (see e.g., Inamdar & Schlichting 2015, applicable for masses lower than Neptune), which could mean its atmospheric fraction was somewhat higher in the past (though still not a gas giant).

Lastly, a possible scenario is that K2-25b formed in-situ close to its current observed orbit close to the host-star (e.g., Batygin et al. 2016), followed by dynamical interactions to

explain its current eccentricity. In-situ formation has been shown to be able to describe the observed compositional and orbital diversity of super-Earths and mini-Neptunes through inherent variations in the initial formation conditions in the disk (MacDonald et al. 2020). To avoid runaway accretion in an in-situ formation scenario, as Lee et al. (2014); MacDonald et al. (2020) and others have argued, the accretion could happen at a later stage when the disk is partially depleted, then grow through giant impacts, and accrete a gaseous envelope in a depleted nebula. However, a challenge for this formation scenario is that in-situ formation generally predicts eccentricities that are lower than K2-25b's currently observed eccentricity (see e.g., MacDonald et al. 2020).

8.6.2. Potential migration histories and eccentricity

From the viewpoint of core and/or pebble-accretion, it is possible that K2-25b formed at large orbital distances where these models predict the creation of more massive planetary cores which later collided together to assemble a more massive planet. If K2-25b formed further out, then K2-25b must have needed to migrate closer to the star to arrive at its current short $P=3.48\,\mathrm{day}$ orbit with its moderate eccentricity of e=0.428. As such, K2-25b could still be in the process of migrating to an even shorter orbit through tidal migration and could represent a precursor of hot Neptunes seen around older stars. There are two major possibilities for migration of K2-25b: disk-driven migration, and high-eccentricity migration

Disk-driven migration (see e.g., Kley & Nelson 2012, and references therein) relies on the exchange of angular momentum between the disk and the planet through mutual gravitational interactions within the plane of the disk. These interactions tend to migrate planets from long period orbits to shorter period orbits while damping eccentricities and thus resulting in circular orbits within the timescale of disk-dissipation (Kley & Nelson 2012). Therefore, to explain K2-25b's current moderate eccentricity, a round of dynamical interactions would be needed. However, K2-25b's eccentricity is much larger than we expect from eccentricity excitation following migration (Petrovich et al. 2014).

Another possible scenario is that K2-25b arrived at its present orbit via high-eccentricity migration, a process which appears to be the dominant process for generating shortperiod giant planets (Dawson & Johnson 2018), and has been used to explain the migration process for most observed warm giants with e > 0.4 (Petrovich & Tremaine 2016). High eccentricity migration can often be approximated as a two step process: reducing the planet's orbital angular momentum, and then reducing its energy (Dawson & Johnson 2018). During the first step, a perturber extract orbital angular momentum from the planet by perturbing it into a highly elliptical orbit. In the second step, the planet tidally dissipates its orbital energy through interactions with the central star. If so, what originally excited K2-25b's eccentricity? Several theories have been proposed to explain the original excitation of eccentricities, including planet-planet scattering and/or merging events (Rasio & Ford 1996; Chatterjee et al. 2008; Jurić & Tremaine 2008), secular chaos (Wu & Lithwick 2011), and stellar flybys (e.g., Kaib et al. 2013). Addi-




Figure 12. Eccentricity as a function of orbital period for known exoplanet systems: grey points show known non-transiting systems, and the blue points show known transiting planets. The blue region shows the region where scattering tends to leads to eccentricity excitation rather than collisions (calculated assuming K2-25b's current mass of $25M_{\oplus}$). Seed planetary cores could have formed at these orbital distances, and then scattered through gravitational interactions to a high eccentricity orbit. Subsequently, K2-25b could have migrated closer to its host star via tidal interactions, arriving at its current orbit. The red curve shows the track of constant angular momentum for K2-25b, showing a possible migration pathway. Only systems with eccentricity errors less than < 0.1 are shown.

tionally, secular interactions—e.g., via the von Zeipel-Lidov-Kozai mechanism from a widely separated perturber (Naoz 2016; Ito & Ohtsuka 2019), or coplanar secular interactions (Petrovich 2015)—can trigger high eccentricity migration. However, significant spin-orbit misalignment and retrograde motion can result from the von Zeipel-Lidov-Kozai mechanism (e.g., Storch et al. 2017; Bourrier et al. 2018b). Therefore, to explain the observed low obliquity of K2-25b, we argue that this mechanism is less likely the cause for K2-25b's moderate eccentricity. Instead, to explain K2-25's low inferred obliquity, planet-planet scattering events and/or coplanar secular interactions are more likely mechanisms as they can produce systems with high eccentricities but low inclinations (see e.g., Chatterjee et al. 2008; Petrovich et al. 2014; Petrovich 2015).

Figure 12 compares the orbital eccentricities of exoplanets as a function of orbital period⁸, showing that K2-25b is among select few planets with moderate eccentricities (e > 0.4) at short orbital periods (< 10 days). The region to the right of the grey curve in Figure 12 is the region where planet-planet scattering can take place to excite eccentricities, and not preferentially cause collisions⁹. As discussed above, a possible formation scenario for K2-25b is that initial planet seed cores at the isolation mass formed via pebble accretion at long orbital periods, which then scattered via dynamical interactions to high eccentricities. These highly eccentric orbits then potentially led to orbit crossings and subsequent collisions resulting in the dense planet we see today, which could then be currently migrating towards a shorter circular orbit through tidal interactions with the star. Figure

Using tidal dissipation theory, we can gain further insight into the plausibility of this formation scenario. Tidal dissipation mechanisms vary strongly with the internal structure of the planet (Goldreich & Soter 1966; Guenel et al. 2014), with higher tidal quality factors O seen for the gas giant planets than the denser rocky planets in the solar system (Goldreich & Soter 1966)¹⁰. Using the equations in Jackson et al. (2009), Kain et al. (2019) estimate a circularization timescale of 410 MYr for K2-25b, assuming a modified tidal quality factor¹¹ of $Q'_* = 10^5$ for the host star, and $Q'_p = 5 \times 10^4$ for K2-25b which they selected based on Neptune's most likely Q_p' value from Zhang & Hamilton (2008). As further noted by Kain et al. (2019), the exact value of the tidal circularization timescale for K2-25b scales directly with Q'_n . Assuming the same tidal quality factors for the planet and the star as Kain et al. (2019), but using the planetary mass, radius, and a/R_* values derived in this work, we estimate a circularization timescale of 306 MYr, which is slightly lower but broadly consistent with their value. We formally require a tidal quality factor of $Q'_p > 1 \times 10^5$ to achieve a tidal circularization timescale consistent with or longer than the age of the K2-25 system (650-800MYr). If the tidal quality factor of K2-25b is indeed $Q_p' = 1 \times 10^5$ or larger, this could explain the moderate eccentricity of K2-25b we observe today, suggesting that at its current age of 650-800 MYr, K2-25b has not had sufficient time to circularize its orbit. Although

¹² highlights a nominal migration track assuming a track of constant angular momentum. Extrapolating this track suggests a fully circularized orbit with a period of \sim 2.5 days (see Figure 12).

⁸ Data obtained from the NASA Exoplanet Archive (Akeson et al. 2013).

 $^{^9}$ Calculated assuming K2-25b's current mass of $\sim 25 M_{\oplus}$ using Equation 10 in (Dawson & Johnson 2018) which compares the escape velocity of the planet's surface to the escape velocity from the star at the planet's semi-major axis.

 $^{^{10}}$ The tidal quality factor Q is a factor quantifying the degree of tidal dissipation in gravitational systems and is inversely proportional to the degree of dissipation

¹¹ The modified tidal quality factor Q' is defined as $3Q/2k_2$, where k_2 is the Love number of degree 2 (Goldreich & Soter 1966) for the system.

larger than Neptune's quality factor, we note that it has been hypothesized that the interior structures of close-in Neptune-sized planets may differ from those of the more distant ice giants in our solar system; in particular, Morley et al. (2017) reported a high dissipation factor for GJ 436b of $Q_p' = 10^5 - 10^6$, which has been theorized to help explain the moderate eccentricity e = 0.16 observed for GJ 436b.

If however, we assume a lower tidal quality of $Q'_p = 5 \times$ 10³—a representative tidal quality factor between the tidal quality factors of the rocky planets and the gas giants in the Solar system (Goldreich & Soter 1966)—we obtain a circularization timescale of 30 MYr, which is substantially shorter than the age of the system (650-800 MYr). If the tidal quality factor is indeed this low, K2-25b would likely require additional ongoing eccentricity excitation to account for its currently observed moderate eccentricity. Although ongoing eccentricity excitation via Kozai-Lidov cycles is less likely given the low measured obliquity, coplanar secular excitation—as has been explored by Batygin et al. (2009) to account for the moderate eccentricity of GJ 436 b—is a possibility. However, as is discussed in Section 8.5, we do not see any clear evidence for a second massive planetary companion in the HPF RVs, although more precision RVs are needed to confidently rule out the presence of smaller planets that could be present. Another more fine-tuned possibility that we can not rule out from the available data, is that K2-25b was only recently (around 30 MYr ago) excited to an eccentric orbit far from the star, and could then still be in the process of circularizing its orbit.

With a single transiting planet seen in the K2 data, K2-25b is consistent with the trend observed in Van Eylen et al. (2019), that single transiting planets tend to have higher eccentricities than planets in systems with multiple transiting planets, and the trend that Dong et al. (2018) observed that hot-Neptunes—planets with $2-6R_{\oplus}$ and an orbital period $P < 10\,\mathrm{days}$ —are most commonly found in single transiting systems. Further, Petigura et al. (2017) note that more massive sub-Saturns tend to have moderately eccentric orbits and orbit stars without other detected planets. By detecting more young Neptune-sized systems, we can compare their observed properties (e.g., eccentricities and obliquities), and compare them to the predictions from different formation and migration mechanisms to start to establish a clearer picture on how short period Neptune-sized planets originate.

8.7. Independent Analysis of K2-25b by IRD

During the preparation of this manuscript, we became aware that Gaidos et al. 2020 (submitted) performed a complimentary analysis of the K2-25b system to constrain its obliquity using the Infrared Doppler Instrument (IRD) on the 8.2m Subaru Telescope. Although the submissions of these complementary studies were coordinated between the groups, the data analyses and interpretations were performed independently.

9. SUMMARY

We present the first mass and obliquity constraint for the young Neptune-sized planet K2-25b orbiting its M4.5-dwarf

host star in the Hyades cluster. Given its known age and well characterized orbital parameters, K2-25b is a benchmark system to study M-dwarf planet formation and subsequent dynamics, giving us further insights into the formation and migration mechanisms that produce other hot Neptune exoplanets

To characterize the planet properties, we jointly fit the available K2 photometry along with precision diffuserassisted ground-based photometry obtained with the Engineered Diffuser on the ARCTIC imager on the 3.5m telescope at Apache Point Observatory and the newly installed Engineered Diffuser on the Half-Degree Imager (HDI) on the 0.9m WIYN Telescope at Kitt Peak Observatory, along with precision out-of-transit NIR RVs from the Habitablezone Planet Finder (HPF) spectrograph at the 10m Hobby-Eberly Telescope (HET) at McDonald Observatory. We see clear evidence for starspot activity in both the K2 data, and the HPF RVs and associated activity indicators. Jointly fitting the available photometry and RVs suggests a best-fit radius of $R = 3.44 \pm 0.12 R_{\oplus}$, an eccentric orbit of $e = 0.41 \pm 0.05$, and a mass of $M = 24.5^{+5.7}_{-5.2} M_{\oplus}$. We tested the robustness of our HPF mass measurement by conducting injection-andrecovery tests in synthetic RV streams. Using our radius and mass constraints, and assuming a two-component composition model of a rocky core enshrouded by a thin H/He envelope, we obtain a H/He envelope mass fraction of 5%. No obvious long-period massive companion is detected in the HPF RV data and continued precise RV monitoring is required to confidently detect or exclude such a companion.

To constrain the obliquity of the system, we present three Rossiter-McLaughlin effect observations of K2-25b obtained with HPF, yielding a sky-projected obliquity constraint of $\lambda=3\pm16^\circ$. Using our constraint for the stellar inclination derived from the stellar radius and our $v\sin i_*=8.8\pm0.6$ km/s constraint from the high resolution HPF spectra, we obtain a true 3D obliquity of $\psi=17^{+11}_{-8}^{\circ}$. Our obliquity and eccentricity constraints paints a picture of a well-aligned, but eccentric system.

With precisely determined age and orbital parameters, we discuss a few possible formation scenarios for K2-25b. If K2-25b reached its current mass during the gas disk phase beyond the ice line, K2-25b would be at odds with the predictions of core and pebble-accretion models—with certain assumptions about the disk properties and planetesimal accretion luminosity—as those models predict that K2-25b should have experienced runaway gas accretion resulting in a gas giant planet. To explain its currently observed mass, we surmise that K2-25b could be the product of planet merging events of smaller planetary cores to produce a more massive planet. Such a dynamical environment could have excited K2-25b into an eccentric orbit, and K2-25b could be in the process of migrating to a shorter period orbit through tidal interactions with the host star. To explain K2-25b's current moderate eccentricity from tidal circularization theory, we place a lower limit on the tidal quality factor of $Q'_p \sim 10^5$, corresponding to a circularization timescale consistent with the age of the system. This tidal quality factor is higher than the tidal quality factor for Neptune, which suggest that K225b's internal structure could be different than that of the small gas-giants (Uranus, Neptune) in the Solar System.

We thank the anonymous referee for a thoughtful reading of the manuscript, and for useful suggestions and comments which made for a clearer and stronger manuscript. This work was partially supported by funding from the Center for Exoplanets and Habitable Worlds. The Center for Exoplanets and Habitable Worlds is supported by the Pennsylvania State University, the Eberly College of Science, and the Pennsylvania Space Grant Consortium. This work was supported by NASA Headquarters under the NASA Earth and Space Science Fellowship Program through grants NNX16AO28H and 80NSSC18K1114. We acknowledge support from NSF grants AST-1006676, AST-1126413, AST-1310885, AST-1517592, AST-1310875, AST-1907622, the NASA Astrobiology Institute (NAI; NNA09DA76A), and PSARC in our pursuit of precision radial velocities in the NIR. We acknowledge support from the Heising-Simons Foundation via grant 2017-0494 and 2019-1177. We acknowledge support from NSF grant AST-1909506 and the Research Corporation for precision photometric observations with diffuser-assisted photometry. Computations for this research were performed on the Pennsylvania State University's Institute for Computational & Data Sciences (ICDS). A portion of this work was enabled by support from the Mt Cuba Astronomical Foundation. RID is supported by NASA XRP 80NSSC18K0355 and the Alfred P. Sloan Foundation's Sloan Research Fellowship. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (NASA). GKS wishes to thank Kento Masuda for informative discussions on determining stellar inclinations from projected rotational velocities, stellar radii, and rotation periods.

These results are based on observations obtained with the Habitable-zone Planet Finder Spectrograph on the Hobby-Eberly Telescope. We thank the Resident astronomers and Telescope Operators at the HET for the skillful execution of our observations of our observations with HPF. The Hobby-Eberly Telescope is a joint project of the University of Texas at Austin, the Pennsylvania State University, Ludwig-Maximilians-Universität München, and Georg-August Universität Gottingen. The HET is named in honor of its principal benefactors, William P. Hobby and Robert E. Eberly. The HET collaboration acknowledges the support and resources from the Texas Advanced Computing Center. This is University of Texas Center for Planetary Systems Habitability Contribution 0001.

These results are based on observations obtained with the Apache Point Observatory 3.5-meter telescope which is owned and operated by the Astrophysical Research Consortium. We wish to thank the APO 3.5m telescope operators in their assistance in obtaining these data.

Based in part on observations at the Kitt Peak National

Observatory, NSF's NOIRLab (Prop. ID 0925-2018B; PI: G. Stefansson), managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. The WIYN 0.9m telescope is operated by WIYN Inc. on behalf of a Consortium of ten partner Universities and Organizations. WIYN is a joint facility of the University of Wisconsin–Madison, Indiana University, NSF's NOIRLab, the Pennsylvania State University, Purdue University, University of California, Irvine, and the University of Missouri. The authors are honored to be permitted to conduct astronomical research on Iolkam Du'ag (Kitt Peak), a mountain with particular significance to the Tohono O'odham. We wish to dearly thank James Winsky at NSF's NOIRLab for his help to conduct some of the HDI observations.

This paper includes data collected by the *Kepler* telescope. The *Kepler* and *K2* data presented in this paper were obtained from the Mikulski Archive for Space Telescopes (MAST). Space Telescope Science Institute is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NNX09AF08G and by other grants and contracts. Funding for the K2 Mission is provided by the NASA Science Mission directorate. This research made use of the NASA Exoplanet Archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program. This work has made use of data from the European Space Agency (ESA) mission Gaia (https://www.cosmos.esa.int/gaia), processed by the Gaia Data Processing and Analysis Consortium (DPAC, https://www.cosmos.esa.int/web/ gaia/dpac/consortium). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement.

Facilities: K2, Gaia, HDI/WIYN 0.9m, ARCTIC/ARC 3.5m, HPF/HET 10m.

Software: AstroImageJ (Collins et al. 2017), astroplan (Morris et al. 2018), astropy (Astropy Collaboration et al. 2013), astroquery (Ginsburg et al. 2018), barycorrpy (Kanodia & Wright 2018), batman (Kreidberg 2015), corner.py (Foreman-Mackey 2016), celerite (Foreman-Mackey et al. 2017), dynesty (Speagle 2019), emcee (Foreman-Mackey et al. 2013), everest (Luger et al. 2017), EXOFASTv2 (Eastman 2017), HxRGproc (Ninan et al. 2018), iDiffuse (Stefansson et al. 2018b), Jupyter (Kluyver et al. 2016), juliet (Espinoza et al. 2019), matplotlib (Hunter 2007), numpy (Van Der Walt et al. 2011), MRExo (Kanodia et al. 2019), pandas (McKinney 2010), PyAstronomy (Czesla et al. 2019), pyde (Parviainen 2016), radvel (Fulton et al. 2018), SERVAL (Zechmeister et al. 2018).

```
Akaike, H. 1974, IEEE Transactions on Automatic Control, 19, 716
```

Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, PASP, 125, 989

Albrecht, S., Winn, J. N., Johnson, J. A., et al. 2012, ApJ, 757, 18

Ambikasaran, S., Foreman-Mackey, D., Greengard, L., Hogg, D. W., & O'Neil, M. 2015, IEEE Transactions on Pattern Analysis and Machine Intelligence, 38, 252

Anglada-Escudé, G. & Butler, R. P. 2012, ApJS, 200, 15

Angus, R., Morton, T., Aigrain, S., Foreman-Mackey, D., & Rajpaul, V. 2018, MNRAS, 474, 2094

 Artigau, É., Kouach, D., Donati, J.-F., et al. 2014, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9147,
 SPIRou: the near-infrared spectropolarimeter/high-precision velocimeter for the Canada-France-Hawaii telescope, 914715

Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33

Awiphan, S., Kerins, E., Pichadee, S., et al. 2016, Monthly Notices of the Royal Astronomical Society, 463, 2574

Barragán, O., Aigrain, S., Kubyshkina, D., et al. 2019, MNRAS, 490, 698 Batalha, N. E., Mandell, A., Pontoppidan, K., et al. 2017, PASP, 129, 064501

Batygin, K., Bodenheimer, P. H., & Laughlin, G. P. 2016, ApJ, 829, 114

Batygin, K., Laughlin, G., Meschiari, S., et al. 2009, ApJ, 699, 23

Benatti, S., Nardiello, D., Malavolta, L., et al. 2019, A&A, 630, A81

Bitsch, B., Morbidelli, A., Johansen, A., et al. 2018, A&A, 612, A30

Bodenheimer, P. & Pollack, J. B. 1986, Icarus, 67, 391

Bourrier, V., Lecavelier des Etangs, A., Ehrenreich, D., et al. 2018a, Astronomy and Astrophysics, 620, A147

Bourrier, V., Lovis, C., Beust, H., et al. 2018b, Nature, 553, 477

Bowler, B. P., Blunt, S. C., & Nielsen, E. L. 2019, arXiv e-prints, arXiv:1911.10569

Burgasser, A. J. & Mamajek, E. E. 2017, ApJ, 845, 110

Chatterjee, S., Ford, E. B., Matsumura, S., & Rasio, F. A. 2008, ApJ, 686, 580

Chen, J. & Kipping, D. 2017, ApJ, 834, 17

Claudi, R., Benatti, S., Carleo, I., et al. 2018, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 10702, Proc. SPIE, 107020Z

Collins, K. A., Kielkopf, J. F., Stassun, K. G., & Hessman, F. V. 2017, ArXiv e-prints, arXiv:1701.04817 [astro-ph.IM]

Crossfield, I. J. M. & Kreidberg, L. 2017, AJ, 154, 261

Cutri, R. M. & et al. 2014, VizieR Online Data Catalog, 2328

Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, VizieR Online Data Catalog, 2246

Czesla, S., Schröter, S., Schneider, C. P., et al. 2019, PyA: Python astronomy-related packages

Dai, F., Winn, J. N., Gandolfi, D., et al. 2017, AJ, 154, 226

David, T. J., Petigura, E. A., Luger, R., et al. 2019, ApJL, 885, L12

David, T. J., Conroy, K. E., Hillenbrand, L. A., et al. 2016, AJ, 151, 112

Dawson, R. I. & Johnson, J. A. 2012, ApJ, 756, 122

—. 2018, ARA&A, 56, 175

Deliyannis, C. P. 2013, in American Astronomical Society Meeting Abstracts, Vol. 222, American Astronomical Society Meeting Abstracts, 111.06

Dmitrienko, E. S. & Savanov, I. S. 2017, Astronomy Reports, 61, 871Dodson-Robinson, S. E., Veras, D., Ford, E. B., & Beichman, C. A. 2009, ApJ, 707, 79

Dong, S., Xie, J.-W., Zhou, J.-L., Zheng, Z., & Luo, A. 2018, Proceedings of the National Academy of Sciences, 115, 266

Dressing, C. D. & Charbonneau, D. 2015, ApJ, 807, 45

Eastman, J. 2017, EXOFASTv2: Generalized publication-quality exoplanet modeling code, Astrophysics Source Code Library, ascl:1710.003

Eastman, J. D., Rodriguez, J. E., Agol, E., et al. 2019, arXiv e-prints, arXiv:1907.09480 [astro-ph.EP]

Ehrenreich, D., Bourrier, V., Wheatley, P. J., et al. 2015, Nature, 522, 459 Espinoza, N. 2018, Research Notes of the American Astronomical Society, 2, 209

Espinoza, N., Kossakowski, D., & Brahm, R. 2019, MNRAS, 490, 2262

Fabrycky, D. & Tremaine, S. 2007, ApJ, 669, 1298

Fischer, D. A., Anglada-Escude, G., Arriagada, P., et al. 2016, PASP, 128, 066001

Foreman-Mackey, D. 2016, The Journal of Open Source Software, 24

Foreman-Mackey, D., Agol, E., Ambikasaran, S., & Angus, R. 2017, AJ, 154, 220

Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306

Fulton, B. J., Petigura, E. A., Blunt, S., & Sinukoff, E. 2018, PASP, 130, 044504

Gaia Collaboration. 2018, VizieR Online Data Catalog, I/345

Gibson, S. R., Howard, A. W., Marcy, G. W., et al. 2016, in Ground-based and Airborne Instrumentation for Astronomy VI, ed. C. J. Evans,
 L. Simard, & H. Takami, Vol. 9908, International Society for Optics and Photonics (SPIE), 2093

Ginsburg, A., Sipocz, B., Parikh, M., et al. 2018, astropy/astroquery: v0.3.7 release

Goldreich, P. & Sari, R. 2003, ApJ, 585, 1024

Goldreich, P. & Soter, S. 1966, Icarus, 5, 375

Grunblatt, S. K., Howard, A. W., & Haywood, R. D. 2015, ApJ, 808, 127

Guenel, M., Mathis, S., & Remus, F. 2014, A&A, 566, L9

Hardegree-Ullman, K. K., Cushing, M. C., Muirhead, P. S., & Christiansen, J. L. 2019, AJ, 158, 75

Haywood, R. D., Collier Cameron, A., Queloz, D., et al. 2014, MNRAS, 443, 2517

Haywood, R. D., Vanderburg, A., Mortier, A., et al. 2018, AJ, 155, 203

Hearty, F., Levi, E., Nelson, M., et al. 2014, in Proc. SPIE, Vol. 9147,

Ground-based and Airborne Instrumentation for Astronomy V, 914752 Henden, A. A., Levine, S., Terrell, D., & Welch, D. L. 2015, in American

Astronomical Society Meeting Abstracts, Vol. 225, American Astronomical Society Meeting Abstracts, 336.16

Hirano, T., Suto, Y., Taruya, A., et al. 2010, ApJ, 709, 458

Hirano, T., Suto, Y., Winn, J. N., et al. 2011, ApJ, 742, 69

Hirano, T., Dai, F., Livingston, J. H., et al. 2018, ArXiv e-prints, arXiv:1801.06957 [astro-ph.EP]

Hirano, T., Gaidos, E., Winn, J. N., et al. 2020a, arXiv e-prints, arXiv:2002.05892

Hirano, T., Krishnamurthy, V., Gaidos, E., et al. 2020b, arXiv e-prints, arXiv:2006.13243

Howell, S. B., Sobeck, C., Haas, M., et al. 2014, PASP, 126, 398

Huber, D., Bryson, S. T., Haas, M. R., et al. 2016, ApJS, 224, 2

Huehnerhoff, J., Ketzeback, W., Bradley, A., et al. 2016, in Proc. SPIE, Vol. 9908, Ground-based and Airborne Instrumentation for Astronomy VI, 99085H

Hunter, J. D. 2007, Computing in Science and Engineering, 9, 90

Inamdar, N. K. & Schlichting, H. E. 2015, MNRAS, 448, 1751

Irwin, J. M., Berta-Thompson, Z. K., Charbonneau, D., et al. 2015, in Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, Vol. 18, 18th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, 767

Ito, T. & Ohtsuka, K. 2019, Monographs on Environment, Earth and Planets, 7, 1

Jackson, B., Barnes, R., & Greenberg, R. 2009, ApJ, 698, 1357

Johnson, J. A., Aller, K. M., Howard, A. W., & Crepp, J. R. 2010, PASP, 122, 905

Jurić, M. & Tremaine, S. 2008, ApJ, 686, 603

Kaib, N. A., Raymond, S. N., & Duncan, M. 2013, Nature, 493, 381

Kain, I. J., Newton, E. R., Dittmann, J. A., et al. 2019, arXiv e-prints, arXiv:1912.05552

Kanodia, S., Wolfgang, A., Stefansson, G. K., Ning, B., & Mahadevan, S. 2019, arXiv e-prints, arXiv:1903.00042

Kanodia, S. & Wright, J. 2018, Research Notes of the American Astronomical Society, 2, 4

Kaplan, K. F., Bender, C. F., Terrien, R., et al. 2018, in The 28th International Astronomical Data Analysis Software & Systems

Kempton, E. M.-R., Bean, J. L., Louie, D. R., et al. 2018, ArXiv e-prints, arXiv:1805.03671 [astro-ph.EP]

```
Kipping, D. M. 2010, MNRAS, 408, 1758
—. 2013, MNRAS, 435, 2152
Klein, B. & Donati, J. F. 2020, MNRAS, L9
Kley, W. & Nelson, R. P. 2012, ARA&A, 50, 211
Kluyver, T., Ragan-Kelley, B., Pérez, F., et al. 2016, in Positioning and
  Power in Academic Publishing: Players, Agents and Agendas, ed.
  F. Loizides & B. Scmidt (IOS Press), 87
Kotani, T., Tamura, M., Nishikawa, J., et al. 2018, in Society of
  Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol.
  10702, Proc. SPIE, 1070211
Kraft, R. P. 1967, ApJ, 150, 551
Kreidberg, L. 2015, PASP, 127, 1161
Lambrechts, M. & Johansen, A. 2012, A&A, 544, A32
Lambrechts, M., Johansen, A., & Morbidelli, A. 2014, A&A, 572, A35
Lambrechts, M. & Lega, E. 2017, A&A, 606, A146
Lee, E. J., Chiang, E., & Ormel, C. W. 2014, ApJ, 797, 95
Lépine, S. & Shara, M. M. 2005, AJ, 129, 1483
Liu, B., Lambrechts, M., Johansen, A., & Liu, F. 2019, arXiv e-prints,
  arXiv:1909.00759
Lopez, E. D. & Fortney, J. J. 2014, ApJ, 792, 1
López-Morales, M., Haywood, R. D., Coughlin, J. L., et al. 2016, The
  Astronomical Journal, 152, 204
Luger, R., Agol, E., Kruse, E., et al. 2016, AJ, 152, 100
Luger, R., Kruse, E., Foreman-Mackey, D., Agol, E., & Saunders, N. 2017,
  ArXiv e-prints, arXiv:1702.05488 [astro-ph.IM]
MacDonald, M. G., Dawson, R. I., Morrison, S. J., Lee, E. J., &
  Khandelwal, A. 2020, ApJ, 891, 20
Maciejewski, G., Niedzielski, A., Nowak, G., et al. 2014, Acta
  Astronomica, 64, 323
Mahadevan, S., Ramsey, L., Bender, C., et al. 2012, in Proc. SPIE, Vol.
  8446, Ground-based and Airborne Instrumentation for Astronomy IV,
  84461S
Mahadevan, S., Ramsey, L. W., Terrien, R., et al. 2014, in Society of
  Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol.
  9147, Ground-based and Airborne Instrumentation for Astronomy V,
  91471G
Mann, A. W., Feiden, G. A., Gaidos, E., Boyajian, T., & von Braun, K.
  2015, ApJ, 804, 64
Mann, A. W., Gaidos, E., Mace, G. N., et al. 2016a, ApJ, 818, 46
Mann, A. W., Newton, E. R., Rizzuto, A. C., et al. 2016b, AJ, 152, 61
Mann, A. W., Gaidos, E., Vanderburg, A., et al. 2017, AJ, 153, 64
Mann, A. W., Dupuy, T., Kraus, A. L., et al. 2019, ApJ, 871, 63
Marcus, R. A., Sasselov, D., Hernquist, L., & Stewart, S. T. 2010, ApJL,
Masuda, K. & Winn, J. N. 2020, arXiv e-prints, arXiv:2001.04973
Mazeh, T., Perets, H. B., McQuillan, A., & Goldstein, E. S. 2015, ApJ, 801,
McKinney, W. 2010, in Proceedings of the 9th Python in Science
  Conference, ed. S. van der Walt & J. Millman, 51
McLaughlin, D. B. 1924, ApJ, 60
Metcalf, A. J., Anderson, T., Bender, C. F., et al. 2019, Optica, 6, 233
Montet, B. T., Feinstein, A. D., Luger, R., et al. 2019, arXiv e-prints,
  arXiv:1912.03794
Morley, C. V., Knutson, H., Line, M., et al. 2017, AJ, 153, 86
Morris, B. M., Tollerud, E., Sipőcz, B., et al. 2018, AJ, 155, 128
Naoz, S. 2016, ARA&A, 54, 441
Nelson, B. E., Ford, E. B., Buchner, J., et al. 2020, AJ, 159, 73
Newton, E. R., Mann, A. W., Tofflemire, B. M., et al. 2019, ApJL, 880, L17
Ninan, J. P., Bender, C. F., Mahadevan, S., et al. 2018, in Society of
  Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol.
  10709, High Energy, Optical, and Infrared Detectors for Astronomy
  VIII, 107092U
Ninan, J. P., Mahadevan, S., Stefansson, G., et al. 2019, arXiv e-prints,
  arXiv:1903.06614 [astro-ph.IM]
Palle, E., Oshagh, M., Casasayas-Barris, N., et al. 2020, arXiv e-prints,
```

arXiv:2006.13609

Parviainen, H. 2016, PyDE: v1.5

```
Pepe, F. 2018, in Ground-based and Airborne Instrumentation for
  Astronomy VII, ed. C. J. Evans, L. Simard, & H. Takami, Vol. 10702,
  International Society for Optics and Photonics (SPIE)
Pepper, J., Gillen, E., Parviainen, H., et al. 2017, AJ, 153, 177
Petigura, E. A., Sinukoff, E., Lopez, E. D., et al. 2017, AJ, 153, 142
Petrovich, C. 2015, ApJ, 805, 75
Petrovich, C. & Tremaine, S. 2016, ApJ, 829, 132
Petrovich, C., Tremaine, S., & Rafikov, R. 2014, ApJ, 786, 101
Playchan, P., Barclay, T., Gagné, J., et al. 2020, Nature, 582, 497
Pollack, J. B., Hubickyj, O., Bodenheimer, P., et al. 1996, Icarus, 124, 62
Quirrenbach, A., Amado, P. J., Ribas, I., et al. 2018, in Society of
  Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol.
  10702, Proc. SPIE, 107020W
Rabus, M., Lachaume, R., Jordán, A., et al. 2019, MNRAS, 484, 2674
Rafikov, R. R. 2006, ApJ, 648, 666
 -. 2011, ApJ, 727, 86
Rasio, F. A. & Ford, E. B. 1996, Science, 274, 954
Reid, I. N., Cruz, K. L., Allen, P., et al. 2004, AJ, 128, 463
Reiners, A., Joshi, N., & Goldman, B. 2012, AJ, 143, 93
Ribas, Á., Bouy, H., & Merín, B. 2015, A&A, 576, A52
Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2014,
  arXiv:arXiv:1406.0151v3
Rossiter, R. A. 1924, ApJ, 60
Schlaufman, K. C. 2010, ApJ, 719, 602
Seifahrt, A., Bean, J. L., Stürmer, J., et al. 2016, in Society of
  Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol.
  9908, Proc. SPIE, 990818
Shetrone, M., Cornell, M. E., Fowler, J. R., et al. 2007, PASP, 119, 556
Shporer, A. 2017, PASP, 129, 072001
Southworth, J. 2011, MNRAS, 417, 2166
Speagle, J. S. 2019, arXiv e-prints, arXiv:1904.02180
Stefansson, G., Li, Y., Mahadevan, S., et al. 2018a, AJ, 156, 266
Stefansson, G., Hearty, F., Robertson, P., et al. 2016, ApJ, 833, 175
Stefansson, G., Mahadevan, S., Hebb, L., et al. 2017, ApJ, 848, 9
Stefansson, G., Mahadevan, S., Wisniewski, J., et al. 2018b, in Society of
  Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol.
  10702, Ground-based and Airborne Instrumentation for Astronomy VII,
  1070250
Stefansson, G., Cañas, C., Wisniewski, J., et al. 2020, AJ, 159, 100
Storch, N. I., Lai, D., & Anderson, K. R. 2017, MNRAS, 465, 3927
Thao, P. C., Mann, A. W., Johnson, M. C., et al. 2020, AJ, 159, 32
Tinetti, G., Drossart, P., Eccleston, P., et al. 2016, in Proc. SPIE, Vol. 9904,
  Space Telescopes and Instrumentation 2016: Optical, Infrared, and
  Millimeter Wave, 99041X
Van Der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, ArXiv e-prints,
  arXiv:1102.1523 [cs.MS]
Van Eylen, V., Albrecht, S., Huang, X., et al. 2019, AJ, 157, 61
Wang, L. & Dai, F. 2019, ApJL, 873, L1
Wildi, F., Blind, N., Reshetov, V., et al. 2017, in Society of Photo-Optical
  Instrumentation Engineers (SPIE) Conference Series, Vol. 10400,
  Proc. SPIE, 1040018
Williams, J. P. & Cieza, L. A. 2011, ARA&A, 49, 67
Winn, J. N., Fabrycky, D., Albrecht, S., & Johnson, J. A. 2010a, ApJL, 718,
Winn, J. N. & Fabrycky, D. C. 2015, ARA&A, 53, 409
Winn, J. N., Johnson, J. A., Albrecht, S., et al. 2009, ApJL, 703, L99
Winn, J. N., Johnson, J. A., Howard, A. W., et al. 2010b, ApJL, 723, L223
Wolfgang, A., Rogers, L. A., & Ford, E. B. 2016, ApJ, 825, 19
Wright, J. T. & Eastman, J. D. 2014, Publications of the Astronomical
  Society of the Pacific, 126, 838
Wu, Y. & Lithwick, Y. 2011, ApJ, 735, 109
Yee, S. W., Petigura, E. A., & von Braun, K. 2017, ApJ, 836, 77
Zacharias, N., Urban, S. E., Zacharias, M. I., et al. 2004, AJ, 127, 3043
Zechmeister, M. & Kürster, M. 2009, A&A, 496, 577
Zechmeister, M., Reiners, A., Amado, P. J., et al. 2018, A&A, 609, A12
Zeng, L., Jacobsen, S. B., Sasselov, D. D., et al. 2019, Proceedings of the
  National Academy of Science, 116, 9723
```

Zhang, K. & Hamilton, D. P. 2008, Icarus, 193, 267Zhou, G., Winn, J. N., Newton, E. R., et al. 2019, arXiv e-prints, arXiv:1912.04095

APPENDIX

A. CALCIUM INFRARED TRIPLET LINE POSITIONS

Table A1 lists the line index positions used to measure the Ca II IRT indices in the HPF spectra.

Table A1. Line index positions used for Ca II IRT activity indicators. Wavelengths are given in vacuum wavelengths.

Line #	Description	Start & End Wavelengths [Å]
1	Line Center	[8499.930, 8500.780]
1	Left Reference Region	[8493.200, 8495.467]
1	Right Reference Region	[8505.202, 8507.472]
2	Line Center	[8544.009, 8544.864]
2	Left Reference Region	[8535.887, 8538.737]
2	Right Reference Region	[8551.562, 8554.412]
3	Line Center	[8664.086, 8664.953]
3	Left Reference Region	[8657.294, 8660.184]
3	Right Reference Region	[8670.300, 8673.190]

B. INJECTION AND RECOVERY TESTS

Table B2, and Figures B1, and B2 summarize the results from our two series (Tests I and II) of synthetic injection and recovery tests. The methodology is further described in Section 6.4. Figures B1, and B2, show that the injected Keplerian parameters are consistently accurately recovered.

Table B2. Summary of injected parameters for both series of tests considered. The results from the injection and recovery test are summarized in Figures B1 and B2. $\mathcal{N}(m,\sigma)$ denotes a normal prior with mean m, and standard deviation σ ; $\mathcal{U}(a,b)$ denotes a uniform prior with a start value a and end value b, $\mathcal{J}(a,b)$ denotes a Jeffreys prior with a start value a and end value b.

Parameter	Description	Prior	Value: Test 1	Value: Test 2
Keplerian Pa	arameters:			
P (days)	Orbital period	$\mathcal{N}(3.484548, 0.000042)$	3.484	56424
T_C	Transit Midpoint - 2400000 (BJD _{TDB})	$\mathcal{U}(58515.63, 58515.66)$	58515.	642096
e	Eccentricity	U(0, 0.95)	0.43	
ω	Argument of Periastron	U(0, 360)	195	
K	RV semi-amplitude (m/s)	$\mathcal{U}(0,200)$	34	
GP Hyperpa	rameters:			
$P_{ m GP}$	GP Period (days)	$\mathcal{N}(1.8784, 0.005)$	1.88	3178
$\sigma_{ ext{GP}}$	RV GP Amplitude (m/s)	$\mathcal{U}(10^1, 10^5)$	42	64.6
Γ_{GP}	Harmonic structure / scaling parameter	$\mathcal{N}(8.0, 1.9)$	7	.4
$lpha_{ ext{GP}}$	Inverse length scale (days ⁻²)	$\mathcal{J}(10^{-12}, 10^{-3})$	3.8×10^{-9}	
Other HPF I	Parameters:			
$\sigma_{ m w}$	HPF white noise RV jitter (m/s)	$\mathcal{J}(0.1, 300)$	1	.5
$\mu_{ ext{HPF}}$	HPF RV offset (m/s)	U(-200, 200)		0




Figure B1. Injection and Recovery Test I: results of 200 injection and recovery tests in synthetic HPF RV streams. The panels show the distribution of the means of individual posteriors. The known injected values are highlighted with the blue lines. The distribution of mean values agree well with the known values. Plot generated using corner.py (Foreman-Mackey 2016).




Figure B2. Posteriors (black curves, 200 each) of the GP amplitude σ_{GP} , the radial velocity semi-amplitude K, and eccentricity e, after two series of 200 injection and recovery tests in the HPF out-of-transit RVs. True values for σ_{GP} , K, and e, are indicated by the blue lines. The first test (Test I; top panels) set $\sigma_{GP} = 42 \,\mathrm{m/s}$, $K = 34 \,\mathrm{m/s}$ and $e = 0.43 \,\mathrm{similar}$ to the nominal median expected values from fit RV2. The true value of the semi-amplitude is within the 68% and 95% credible intervals in 80% and 99.7% of the cases, respectively. The second test (Test II; bottom panels) assumed that $\sigma_{GP} = 64.6 \,\mathrm{m/s}$, or at the 95th quantile value from fit RV2, while keeping the other parameters the same. Even with the higher assumed correlated noise, the true value of the semi-amplitude is still reliably recovered at a high coverage probability: the true value of the semi-amplitude is within the 68% and 95% credible intervals in 75% and 97% of the cases, respectively.

C. HPF RADIAL VELOCITIES

C.1. Out of transit

Table C3 lists the RVs from HPF used for the mass measurement of K2-25b, along with associated stellar activity indicators.

 $\textbf{Table C3}. \ \ \text{Out-of-transit RVs from HPF along with the differential line width (dLW), chromatic index (CRX), and the line indices for the three Ca II IRT triplet lines (Ca II IRT 1, 2 and 3), along with associated errors.$

BJD	RV [m s ⁻¹]	dLW [m ² s ⁻²]	CRX [m s ⁻¹ Np ⁻¹]	Ca II IRT 1	Ca II IRT 2	Ca II IRT 3
2458424.95976	29.8 ± 15.6	288.7 ± 74.2	88.1 ± 233.7	0.878 ± 0.005	0.790 ± 0.006	0.662 ± 0.005
2458425.74905	-90.6 ± 35.6	312.2 ± 170.0	-1649.2 ± 478.6	1.097 ± 0.014	0.931 ± 0.017	0.828 ± 0.013
2458429.93716	39.5 ± 16.8	64.9 ± 80.0	-445.4 ± 209.2	0.930 ± 0.006	0.826 ± 0.007	0.703 ± 0.005
2458433.92567	44.7 ± 38.0	311.3 ± 180.5	1049.5 ± 630.4	0.854 ± 0.012	0.764 ± 0.016	0.685 ± 0.012
2458436.91691	-56.8 ± 23.9	-266.8 ± 114.7	-145.4 ± 332.6	0.975 ± 0.008	0.871 ± 0.010	0.708 ± 0.008
2458437.90840	18.0 ± 17.3	88.2 ± 82.9	317.4 ± 224.3	1.045 ± 0.006	0.926 ± 0.008	0.776 ± 0.006
2458439.91091	-5.9 ± 21.2	301.2 ± 101.1	174.7 ± 394.9	0.908 ± 0.007	0.801 ± 0.009	0.686 ± 0.007
2458441.89899	62.8 ± 22.3	176.8 ± 106.5	-132.8 ± 183.3	0.893 ± 0.007	0.784 ± 0.009	0.668 ± 0.007
2458442.89348	-28.7 ± 29.1	123.3 ± 139.4	437.9 ± 522.9	0.988 ± 0.010	0.882 ± 0.012	0.760 ± 0.009
2458443.71423	38.9 ± 24.4	296.1 ± 116.1	19.8 ± 352.8	0.908 ± 0.007	0.810 ± 0.009	0.712 ± 0.007
2458444.71135	39.9 ± 19.6	-315.5 ± 94.6	34.1 ± 181.4	0.946 ± 0.006	0.844 ± 0.008	0.720 ± 0.006
2458449.88697	-80.4 ± 22.3	169.7 ± 107.3	-236.0 ± 321.6	0.970 ± 0.007	0.876 ± 0.009	0.761 ± 0.007
2458451.68266	-19.1 ± 28.1	319.9 ± 134.9	387.9 ± 453.2	0.900 ± 0.009	0.859 ± 0.012	0.713 ± 0.009
2458473.80425	-4.4 ± 42.5	1752.3 ± 196.0	178.1 ± 669.2	0.893 ± 0.015	0.762 ± 0.019	0.644 ± 0.014
2458480.80404	21.5 ± 52.9	43.1 ± 252.9	127.4 ± 632.2	0.860 ± 0.020	0.744 ± 0.027	0.635 ± 0.021
2458487.78924	-153.6 ± 44.8	-523.5 ± 218.1	137.8 ± 710.1	0.891 ± 0.017	0.718 ± 0.024	0.640 ± 0.017
2458546.61407	1.0 ± 33.4	-390.2 ± 162.4	434.9 ± 462.0	0.895 ± 0.010	0.829 ± 0.013	0.677 ± 0.010
2458549.60481	15.6 ± 28.0	473.1 ± 134.3	-178.2 ± 444.0	0.885 ± 0.008	0.825 ± 0.011	0.684 ± 0.008
2458741.89127	-36.4 ± 37.6	-179.1 ± 180.4	-55.6 ± 587.0	0.907 ± 0.012	0.831 ± 0.017	0.708 ± 0.012
2458744.87566	11.1 ± 31.4	-48.3 ± 149.9	-1098.4 ± 405.1	1.040 ± 0.011	0.960 ± 0.014	0.823 ± 0.010
2458752.85672	-43.7 ± 31.5	393.2 ± 149.1	-267.1 ± 441.1	1.030 ± 0.011	0.907 ± 0.015	0.803 ± 0.011
2458804.90940	-107.8 ± 41.2	486.7 ± 196.0	1583.8 ± 570.3	0.902 ± 0.014	0.801 ± 0.018	0.667 ± 0.014
2458805.71647	-117.9 ± 28.5	433.8 ± 135.7	-1673.7 ± 264.1	0.874 ± 0.010	0.753 ± 0.013	0.674 ± 0.010
2458808.70580	7.6 ± 23.4	-300.2 ± 112.8	64.8 ± 464.9	0.946 ± 0.008	0.889 ± 0.011	0.726 ± 0.008
2458808.90127	31.0 ± 22.0	-61.7 ± 105.8	-96.6 ± 260.6	1.024 ± 0.008	0.921 ± 0.010	0.775 ± 0.007
2458811.88979	28.6 ± 30.4	-132.5 ± 146.7	-412.6 ± 421.3	0.918 ± 0.010	0.859 ± 0.013	0.708 ± 0.010
2458824.66447	-97.2 ± 44.0	722.8 ± 209.6	247.7 ± 747.0	0.907 ± 0.015	0.797 ± 0.020	0.677 ± 0.015
2458825.66903	32.9 ± 55.7	-289.9 ± 271.4	365.9 ± 792.5	0.939 ± 0.018	0.878 ± 0.024	0.757 ± 0.018
2458832.64612	58.1 ± 35.4	-230.3 ± 172.1	738.2 ± 645.5	0.899 ± 0.011	0.792 ± 0.015	0.692 ± 0.012
2458849.59344	39.7 ± 54.1	254.3 ± 256.9	616.0 ± 873.0	0.907 ± 0.017	0.774 ± 0.021	0.714 ± 0.017
2458849.77940	41.6 ± 32.7	-606.0 ± 158.7	661.1 ± 496.7	0.924 ± 0.010	0.817 ± 0.013	0.683 ± 0.010
2458852.77670	-61.7 ± 27.8	271.2 ± 132.7	-400.7 ± 457.1	0.901 ± 0.008	0.821 ± 0.011	0.675 ± 0.008

 Table C4. In-transit RVs from HPF.

Time (BJD _{TDB})	RV (m/s)	RV Error (m/s)	S/N	Transit #
2458473.798377	21.502	92.473	31.5	1
2458473.802296	-99.316	85.383	32.9	1
2458473.806148	32.915	87.976	33.0	1
2458473.810188	25.879	76.650	36.0	1
2458473.814183	58.845	68.489	38.9	1
2458473.818066	22.483	81.288	35.4	1
2458473.822004	-29.955	70.497	38.1	1
2458473.825933	-107.874	82.864	33.7	1
2458473.829840	10.968	101.494	28.8	1
2458473.833884	5.058	112.039	26.2	1
2458473.837847	-5.540	111.578	26.1	1
2458480.779096	93.172	81.592	32.5	2
2458480.783031	84.061	74.764	35.2	2
2458480.786978	289.359	70.453	36.9	2
2458480.790926	118.139	59.718	42.5	2
2458480.794850	147.978	61.526	41.0	2
2458480.798729	-91.458	63.679	40.3	2
2458480.802696	-89.542	72.040	36.7	2
2458480.806659	-88.708	71.505	36.4	2
2458480.810610	82.751	78.649	34.5	2
2458480.814557	-30.039	93.241	29.1	2
2458480.818470	-30.570	103.872	27.2	2
2458487.763970	-32.264	92.523	29.1	3
2458487.767653	-108.714	72.947	35.5	3
2458487.771528	-163.954	69.931	37.3	3
2458487.775525	-131.220	67.386	38.7	3
2458487.779335	-91.287	66.037	38.8	3
2458487.783313	-202.220	74.173	35.5	3
2458487.787296	-203.813	88.859	30.1	3
2458487.791204	-20.690	97.731	28.4	3
2458487.795138	-138.737	108.448	25.3	3