

Preparation of Enantioenriched Alkylcarbastannatranes via Nucleophilic Inversion of Alkyl Mesylates for Use in Stereospecific Cross-Coupling Reactions


Glenn Ralph^{†,‡} and Mark R. Biscoe^{*,†,‡,§}

[†]Department of Chemistry, The City College of New York (CCNY), 160 Convent Avenue, New York, New York 10031, United States

[‡]Ph.D. Program in Chemistry, The Graduate Center of The City University of New York (CUNY), 365 Fifth Avenue, New York, New York 10016, United States

Supporting Information

ABSTRACT: We report the preparation of enantioenriched secondary alkylcarbastannatranes via a stereoinvertive S_N2 reaction of enantioenriched alkyl mesylates and carbastannatranyl anion equivalents. Using this process, enantioenriched secondary alcohols may be converted into highly enantioenriched alkylcarbastannatranes, which are useful in stereospecific cross-coupling reactions.



The use of configurationally stable enantioenriched organometallic nucleophiles in stereospecific transformations constitutes a powerful approach to the preparation of complex, nonracemic molecules.^{1,2} Accordingly, enantioenriched organoboron and organotin compounds have been extensively used in metal-catalyzed cross-coupling reactions. These transformations rely on stereospecific mechanisms that effectively translate chiral information from the starting materials to final products. Thus, it is vital that such precursors can be prepared with high enantiopurity in a straightforward manner.

Building upon the pioneering work of Jurkschat³ and Vedejs,⁴ in 2013, our laboratory developed a Stille cross-coupling reaction of enantioenriched secondary alkylcarbastannatrane nucleophiles and aryl electrophiles (Figure 1).^{5–7}




Figure 1. Stereospecific Stille cross-coupling reaction using enantioenriched secondary alkylcarbastannatranes.

This method afforded arylation products in high yields and with high enantiospecificity. The unique reactivity of alkylcarbastannatranes is attributed to the selective labilization of its apical alkyl substituent as a consequence of the dative N–Sn interaction in the “atrane” tin backbone.⁴ This approach enabled the first example of a stereospecific cross-coupling reaction using an unactivated secondary organotin species.⁸

Subsequently, we extended this chemistry to stereospecific acylation reactions, as well as stereospecific fluorination reactions.^{9,10}

The lack of an efficient route to prepare enantioenriched carbastannatrane nucleophiles has been a major bottleneck to their broader synthetic application. Because unactivated secondary alkylstannanes typically exhibit low reactivity, there has previously been limited value in devising synthetic strategies to prepare enantioenriched variants. As a result, our laboratory has relied heavily on stereospecific lithiation/stannylation processes¹¹ and chiral preparatory HPLC separation of racemates to obtain enantioenriched alkylcarbastannatrane nucleophiles. We set out to address this limitation by developing a more general approach to the preparation of enantioenriched alkylcarbastannatranes. Herein, we report a study of carbastannatranyl anion equivalents in stereoinvertive S_N2 reactions with enantioenriched alkyl mesylates. We have found that carbastannatranyl lithium (2) readily undergoes substitution reactions with secondary alkyl mesylates. This strategy enables the preparation of highly enantiopure secondary alkylcarbastannatranes from commercially available single-enantiomer alcohols. With access to more diverse alkylcarbastannatrane nucleophiles, the scope of stereospecific Stille cross-coupling reactions is significantly broadened.

Special Issue: Asymmetric Synthesis Enabled by Organometallic Complexes

Received: July 10, 2019

In 2015, Wang and Uchiyama reported the highly efficient lithiation of trialkyltin chlorides using *in situ* generated naphthalide radical anions.¹² Using this method, clean substitution of methyl and primary allylic/benzylic halides was observed, which enabled preparation of the corresponding organostannanes in excellent yield. We hypothesized that carbastannatryl lithium (2) could be produced under similar reductive conditions and might be employed in S_N2 reactions with enantioenriched secondary alkyl electrophiles to afford enantioenriched secondary alkylcarbastannatranes via stereoinversion. To circumvent potential racemization via a background radical pathway, we opted to investigate the use of secondary alkyl sulfonates, which have a lower propensity to form alkyl radicals than do alkyl halides. We quickly found that *sec*-butyl mesylate underwent efficient substitution with tributylstannyl lithium prepared by the method of Wang and Uchiyama. However, an analogous approach using carbastannatryl chloride failed to generate the corresponding secondary alkylcarbastannatrane. Ensuing studies revealed that carbastannatryl lithium (2) could be generated through sonication of the reaction mixture at 45 °C (see the Supporting Information).

Having developed effective conditions for the *in situ* generation of tin lithium species 2, we investigated the enantiospecificity¹³ of the substitution reaction using enantioenriched alkyl mesylate 5 (Table 1). The reaction of 2 with

Table 1. Optimization of Enantiospecificity in the Reaction of Carbastannatryl Anion Equivalents and Alkyl Mesylate 5

Entry	Tin Nucleophile	Solvent	Yield (%) ^a	% es ^b
1	2	THF	74	74
2	2 (with TMEDA)	THF	65	98
3	2	ether	63	99
4	2	hexane	68	99
5	5	hexane	30	98
6	6	hexane	< 5	--
7	7	hexane	65	98

2

5

6

7

^aCalibrated ^1H NMR yields. ^bDetermined by chiral HPLC.

3 predissolved in THF resulted in eroded enantiopurity in the substitution product. However, we found that excellent enantiospecificity could be achieved when TMEDA was added to 2 or when ether or hexane was employed as the solvent. Access to carbastannatryl lithium (2) not only was found to be useful in forming tin–carbon bonds directly but could also be used as a precursor to other metalated tin nucleophiles.¹⁴ Using 5 as a model substrate for analysis, we examined the reactivity of carbastannatryl anion equivalents 5–7 in this substitution reaction. Reactions involving 5 and 7 showed excellent enantiospecificity. Though none of these

nucleophiles proved superior to the use of 2 in hexane or ether, the ability of attenuated nucleophiles such as 5 and 7 to undergo stereoinvertive substitution with high enantiospecificity suggests that alkyl mesylates bearing functional groups not compatible with tin lithium reagents will be still viable electrophiles in this process.

Using the conditions developed in Table 1, a series of secondary alkylcarbastannatranes was prepared from their corresponding mesylates (Table 2). Yields of ca.

Table 2. Preparation of Enantioenriched Secondary Alkylcarbastannatranes from Their Corresponding Mesylates


	$\xrightarrow[\text{THF, 45 } \text{°C, 3 h}]{\text{Li (2.2 equiv), naph. (10 mol %)}}$		$\xrightarrow[\text{hexane}]{\text{OMs, R1-R2}}$	
(<i>R</i>)-9: 47% (99% es) (<i>S</i>)-9: 52% (99% es)				
(<i>R</i>)-4: 42% (99% es) (<i>S</i>)-4: 48% (99% es)				
(±)-10: 37% (<i>S</i>)-10: 35% (97% es)				
(±)-11: 54% (<i>S</i>)-11: 51% (99% es)				
(±)-12: 46% (<i>S</i>)-12: 45% (99% es)				
(±)-13: 51%				

50% were typically obtained alongside high levels of enantioenrichment, with separable elimination products accounting for the remaining mass balance. These unactivated enantioenriched secondary alkylcarbastannatranes (8) are completely tolerant of air and water. They are also inert to neutral, basic, and reductive reaction conditions, which enables synthetic modifications to be performed following installation of the carbastannatrane unit. For instance, desilylation of 12 was readily achieved by treatment with TBAF. Previously, we have demonstrated that diborylation of olefins and LiAlH_4 reduction of esters do not affect the integrity of the carbastannatrane unit.^{10,15,16} The majority of the enantioenriched alkylcarbastannatranes prepared using this method contain a methyl group on the stereogenic carbon center. Unfortunately, analytical chiral HPLC conditions could not be found to separate enantiomers of carbastannatrane 13, in which an ethyl group replaces the methyl group. Though the enantio purity of 13 could not be determined, we feel that the enantiospecificity would likely be very high from the enantioenriched alkyl mesylate, as was observed when other mesylates were employed alongside our standard conditions.

Previously, we had only reported two examples of stereospecific arylation reactions using unactivated secondary alkylcarbastannatranes (prepared using preparatory chiral

HPLC).⁵ Now having broader access to enantioenriched alkylcarbastannatranes nucleophiles, we conducted an investigation of their use in stereospecific Stille reactions. The cross-coupling products in Table 3 were all generated with high

Table 3. Stereospecific Stille Cross-Coupling Reactions Using Enantioenriched Secondary Alkylcarbastannatranes^a



^aIsolated yields of duplicate runs. ^bUsing racemic 13. ^cUsing the thioester of naproxen as electrophile; no KF added.

enantiospecificity. In contrast to our observations for stereospecific Suzuki reactions,² the electronic properties of the electrophilic component did not influence the transfer of stereochemistry during the cross-coupling reaction. Both electron-rich and electron-deficient electrophiles underwent coupling with nearly perfect transfer of the initial stereochemistry. The presence of heteroatoms and acidic protons

was well tolerated in these reactions. The free alcohol formed from desilylation of (S)-12 underwent arylation with high yield and excellent enantiospecificity. Use of (rac)-13 as a nucleophile resulted in the formation of arylation product 14h in high yield. Thus, transmetalation of a secondary alkylcarbastannatrane still occurs efficiently from the bulkier, non-methyl-containing alkylcarbastannatrane nucleophile. Finally, when (S)-11 was employed in a coupling reaction with the thioester of (S)-naproxen, acylation product 14j was obtained with completely reagent controlled diastereoselectivity.

In summary, we have developed a reliable synthetic method to access enantioenriched secondary alkylcarbastannatranes from their corresponding alkyl mesylates. Treatment of enantioenriched alkyl mesylates with stannatranylolithium (2) results in the formation of highly enantioenriched secondary alkylcarbastannatranes via a stereoinvertive S_N2 pathway. Other carbastannatranyl anion equivalents (e.g., 5 and 7) were also shown to undergo substitution reactions with high enantiospecificity. We subsequently demonstrated that these enantioenriched alkylcarbastannatrane nucleophiles readily undergo cross-coupling reactions with aryl bromides, with highly general preference for retention of stereochemistry. We expect that this method to access enantioenriched alkylcarbastannatranes will facilitate the future development of new stereospecific transformations of alkylcarbastannatrane compounds.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.organomet.9b00467](https://doi.org/10.1021/acs.organomet.9b00467).

Synthetic procedures, spectral data (¹H and ¹³C NMR), and HPLC data ([PDF](#))

AUTHOR INFORMATION

Corresponding Author

*E-mail for M.R.B.: mbiscoe@ccny.cuny.edu.

ORCID

Mark R. Biscoe: [0000-0003-1257-6288](https://orcid.org/0000-0003-1257-6288)

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the City College of New York, the National Institutes of Health (R01GM131079), and the National Science Foundation (CHE-1665189) for support of this work.

REFERENCES

- (1) For review articles on stereospecific cross-coupling reactions, see: (a) Rygus, J. P. G.; Cradden, C. M. Enantiospecific and Iterative Suzuki-Miyaura Cross-Couplings. *J. Am. Chem. Soc.* **2017**, *139*, 18124–18137. (b) Sandford, C.; Aggarwal, V. K. Stereospecific Functionalizations and Transformations of Secondary and Tertiary Boronic Esters. *Chem. Commun.* **2017**, *53*, 5481–5494. (c) Cherney, A. H.; Kadunce, N. T.; Reisman, S. E. Enantioselective and Enantiospecific Transition-Metal-Catalyzed Cross-Coupling Reactions of Organometallic Reagents to Construct C–C Bonds. *Chem. Rev.* **2015**, *115*, 9587–9652. (d) Wang, C. Y.; Derosa, J.; Biscoe, M. R. Configurationally Stable, Enantioenriched Organometallic Nucleophiles in Stereospecific Pd-Catalyzed Cross-Coupling Reactions: An

Alternative Approach to Asymmetric Synthesis. *Chem. Sci.* **2015**, *6*, 5105–5113.

(2) (a) Zhao, S.; Gensch, T.; Murray, B.; Niemeyer, Z. L.; Sigman, M. S.; Bischof, M. R. Enantiodivergent Pd-Catalyzed C–C Bond Formation Enabled through Ligand Parameterization. *Science* **2018**, *362*, 670–674. (b) Li, L.; Zhao, S.; Joshi-Pangu, A.; Diane, M.; Bischof, M. R. Stereospecific Pd-Catalyzed Cross-Coupling Reactions of Secondary Alkylboron Nucleophiles and Aryl Chlorides. *J. Am. Chem. Soc.* **2014**, *136*, 14027–14030.

(3) (a) Jurkschat, K.; Tzschach, A.; Meunier-Piret, J.; Van Meerssche, M. Crystal and Molecular Structure of 1-Aza-5-Stanna-5-Chlorotricyclo[3.3.0^{1,5}]undecane, a 2,8,9-Tricarbostannatran. *J. Organomet. Chem.* **1985**, *290*, 285–289. (b) Jurkschat, K.; Tzschach, A.; Meunier-Piret, J. Crystal and Molecular Structure of 1-Aza-5-Stanna-5-Methyltricyclo[3.3.0^{1,5}]undecane. Evidence for a Transannular Donor-Acceptor Interaction in a Tetraorganotin Compound. *J. Organomet. Chem.* **1986**, *315*, 45–49.

(4) (a) Vedejs, E.; Haight, A. R.; Moss, W. O. Internal Coordination at Tin Promotes Selective Alkyl Transfer in the Stille Coupling Reaction. *J. Am. Chem. Soc.* **1992**, *114*, 6556–6558. (c) Theddu, N.; Vedejs, E. Stille Coupling of an Aziridinyl Stannatran. *J. Org. Chem.* **2013**, *78*, 5061–5066.

(5) Li, L.; Wang, C. Y.; Huang, R.; Bischof, M. R. Stereoretentive Pd-Catalyzed Stille Cross-Coupling Reactions of Secondary Alkyl Azastannatranes and Aryl Halides. *Nat. Chem.* **2013**, *5*, 607–12.

(6) For other examples of carbastannatran use, see: (a) Jensen, M. S.; Yang, C.; Hsiao, Y.; Rivera, N.; Wells, K. M.; Chung, J. Y. L.; Yasuda, N.; Hughes, D. L.; Reider, P. J. Synthesis of an Anti-Methicillin-Resistant *Staphylococcus aureus* (MRSA) Carbapenem via Stannatran-Mediated Stille Coupling. *Org. Lett.* **2000**, *2*, 1081–1084. (b) Sebahar, H. L.; Yoshida, K.; Hegedus, L. S. Effect of Adjacent Chiral Tertiary and Quaternary Centers on the Metal-Catalyzed Allylic Substitution Reaction. *J. Org. Chem.* **2002**, *67*, 3788–3795. (c) Fillion, E.; Taylor, N. J. Cine-Substitution in the Stille Coupling: Evidence for the Carbenoid Reactivity of *sp*³-*gem*-Organodimetalllic Iodopalladio-Trialkylstannylalkane Intermediates. *J. Am. Chem. Soc.* **2003**, *125*, 12700–12701. (d) Kavoosi, A.; Fillion, E. Synthesis and Characterization of Tricarbostannatranes and Their Reactivity in B(C₆F₅)₃-Promoted Conjugate Additions. *Angew. Chem., Int. Ed.* **2015**, *54*, 5488–5492. (e) Srivastav, N.; Singh, R.; Kaur, V. Carbastannatranes: Powerful Coupling Mediators in Stille Coupling. *RSC Adv.* **2015**, *5*, 62202–62213. (f) Fillion, E.; Kavoosi, A.; Nguyen, K.; Ieritano, C. B(C₆F₅)₃-Catalyzed Transfer 1,4-Hydrostannylation of α,β -Unsaturated Carbonyls Using iPr-tricarbostannatran. *Chem. Commun.* **2016**, *52*, 12813–12816. (g) Simidzija, P.; Lecours, M. J.; Marta, R. A.; Steinmetz, V.; McMahon, T. B.; Fillion, E.; Hopkins, W. S. Changes in Tricarbostannatran Transannular N–Sn Bonding upon Complexation Reveals Lewis Base Donicities. *Inorg. Chem.* **2016**, *55*, 9579–9585.

(7) For carbagermanatran use, see: Xu, M.-Y.; Jiang, W.-T.; Li, Y.; Xu, Q.-H.; Zhou, Q.-L.; Yang, S.; Xiao, B. Alkyl Carbagermatranes Enable Practical Palladium-Catalyzed *sp*²–*sp*³ Cross-Coupling. *J. Am. Chem. Soc.* **2019**, *141*, 7582–7588.

(8) Activated organotin nucleophiles feature the presence of a C(sp²) α -carbon, heteroatomic α -carbon, and/or strongly coordinating β -carbonyl group.

(9) Wang, C. Y.; Ralph, G.; Derosa, J.; Bischof, M. R. Stereospecific Palladium-Catalyzed Acylation of Enantioenriched Alkylcarbastannatranes: A General Alternative to Asymmetric Enolate Reactions. *Angew. Chem., Int. Ed.* **2017**, *56*, 856–860.

(10) Ma, X.; Diane, M.; Ralph, G.; Chen, C.; Bischof, M. R. Stereospecific Electrophilic Fluorination of Alkylcarbastannatran Reagents. *Angew. Chem., Int. Ed.* **2017**, *56*, 12663–12667.

(11) Beak, P.; Basu, A.; Gallagher, D. J.; Park, Y. S.; Thayumanavan, S. Regioselective, Diastereoselective, and Enantioselective Lithiation–Substitution Sequences: Reaction Pathways and Synthetic Applications. *Acc. Chem. Res.* **1996**, *29*, 552–560.

(12) Wang, D. Y.; Wang, C.; Uchiyama, M. Stannyli-Lithium: A Facile and Efficient Synthesis Facilitating Further Applications. *J. Am. Chem. Soc.* **2015**, *137*, 10488–10491.

(13) Enantiospecificity (%es) = (%ee_{product} / %ee_{starting material}) \times 100%.

(14) Hibino, J.-I.; Matsubara, S.; Morizawa, Y.; Oshima, K.; Nozaki, H. Regioselective Stannylimetalation of Acetylenes in the Presence of Transition-Metal Catalyst. *Tetrahedron Lett.* **1984**, *25*, 2151–2154.

(15) We have found secondary alkylcarbastannatranes to be incompatible with oxidative conditions, though it is plausible that very mild oxidants may be tolerated.

(16) Cyclohexyl mesylate could be efficiently converted to the corresponding cyclohexylcarbastannatran. However, 4-*tert*-butylcyclohexyl mesylate failed to react under similar conditions.