

REGULAR PAPER

Life history, distribution and molecular phylogenetics of the Upward-Mouth Spikefish *Atrophacanthus japonicus* (Teleostei: Tetraodontiformes: Triacanthodidae)

Katherine E. Bemis^{1,2}  | James C. Tyler³ | Dahiana Arcila⁴

¹Department of Fisheries Science, Virginia Institute of Marine Science, College of William & Mary, Gloucester Point, Virginia, USA

²Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, District of Columbia, USA

³Department of Paleobiology, National Museum of Natural History, Smithsonian Institution, Washington, District of Columbia, USA

⁴Sam Noble Oklahoma Museum of Natural History and Department of Biology, University of Oklahoma, Norman, Oklahoma, USA

Correspondence

Katherine E. Bemis, Department of Fisheries Science, Virginia Institute of Marine Science, College of William & Mary, P.O. Box 1346, Gloucester Point, VA, 23062, USA.

Email: kebemis@vims.edu

Funding information

Research support was provided by the National Science Foundation to KEB (GRFP), JCT (DEB-1541552) and DA (DBI-1811748) and the Virginia Institute of Marine Science Office of Academic Studies, Clyde D., & Lois W. Marlatt, Jr. Fellowship.

Ninety-six juvenile specimens (37–54 mm standard length; L_S) of the rarely collected Upward-Mouth Spikefish *Atrophacanthus japonicus* (Triacanthodidae) were obtained from the stomachs of three Yellowfin Tuna *Thunnus albacares* collected off Guam in the Mariana Islands in the central Pacific Ocean. These specimens extend the range of *A. japonicus* eastward into Oceania. We review the systematic characters of the monotypic genus *Atrophacanthus* and present colour photographs of freshly collected specimens. The diet of the juvenile specimens of *A. japonicus* consisted of thecosome pteropods and foraminiferans. We present a range map of *A. japonicus* based on all known specimens and show that specimen size is related to whether specimens were collected in the pelagic zone or on the bottom. Our results support that, compared to all other Triacanthodidae, *A. japonicus* has an unusually extended pelagic larval and juvenile period, up to 54 mm L_S , before settling to the bottom as adults. Lastly, we provide a multilocus phylogeny addressing the phylogenetic placement of *Atrophacanthus* based on eight of 11 triacanthodid genera and six genetic markers. Our results reveal that *Atrophacanthus* is the sister group of *Macrorhamphosodes* and they provide new insights about the evolutionary history of the family.

KEY WORDS

Atrophacanthus japonicus, biogeography, Guam, Mariana Islands, Oceania, systematics

1 | INTRODUCTION

The rarely collected Upward-Mouth Spikefish *Atrophacanthus japonicus* (Kamohara 1941) is the only species of its triacanthodid spikefish genus. Unlike all other triacanthodids, which settle into benthic habitats at small sizes, larval and juvenile *A. japonicus* have longer pelagic periods (Tyler, 1968). Most known specimens are larval or juvenile stages that were collected pelagically by the *Dana* Expedition in the early 1900s and only six benthic adult specimens are known. Herein, we report on 96 new juvenile specimens collected from the stomachs of three Yellowfin Tuna *Thunnus albacares* (Bonnaterre 1788) caught off Guam in the western Pacific Ocean. We provide a review of the systematic characters of the genus, diet data, colour photographs of the fresh new specimens from Guam and a distribution map based on

all known specimens. We also present a molecular phylogeny of the eight genera of Triacanthodidae for which genetic data presently is available.

2 | MATERIALS AND METHODS

2.1 | Collection and curation of new specimens from Guam

Ninety-six juvenile specimens of *A. japonicus* were collected off Guam, Mariana Islands (Figure 1). The specimens were taken from the stomachs of three *T. albacares* weighing 12.2 to 20.4 kg. The *T. albacares* were caught by trolling surface lures from the F.V. *Stonfish* (Captains




FIGURE 1 Map showing distribution of examined specimens of *Atrophacanthus japonicus* in this study and Tyler (1968, 1970a). L = Larvae, J = Juveniles, A = Adult. Most larvae and juveniles were collected in the pelagic zone, whereas adults were collected using benthic gear. The new specimens collected off Guam represent the most eastward occurrence of *A. japonicus*. All lengths standard length

James and Ken Borja) about 2.4 km offshore from the southeast coast of Guam at 13° 12.829' N, 144° 43.178' E, depth c. 850 m on 16 April 2016.

Many of the specimens from the *T. albacares* stomachs were in relatively good condition and were photographed on the deck worktable (Figure 2) before the specimens were frozen aboard the ship. During freezing, the specimens became crowded together in a bundle, whereupon many of the slender dorsal and pelvic spines were broken and fin rays and regions of the bodies became distorted in alignment. The frozen fish were sent to the Division of Fishes, National Museum of Natural History, Smithsonian Institution, by B. Mundy, NOAA National Marine Fisheries Service, Hawaii, and his cooperating colleagues, the Captains J. and K. Borja and K. Kawamoto, M. Mitsuyasu, E. Cruz, C. Tam and R. Humphreys. The specimens were thawed and preserved in 95% ethanol by D. Pitassy. Eighteen specimens (38–54 mm standard length; L_s) in relatively good condition were selected for individual cataloguing (USNM 440413–440430) as vouchers for tissue samples. These 18 voucher specimens were individually radiographed by S. Raredon. The remaining 78 specimens (37–50 mm L_s) were placed in one jar as a lot catalogued as USNM 440431. All specimens were stored in 95% ethanol.

We recorded meristic and morphological data for the 18 individually catalogued voucher specimens from Guam to compare with the large dataset presented in Tyler (1968) for *A. japonicus* (larval, juvenile and adults), with emphasis on diagnostic features. We report L_s for



FIGURE 2 (a) Anterior right and (b) Anterior left views of *Atrophacanthus japonicus* from the stomachs of three *Thunnus albacares* caught near the surface 2.4 km off the southeast coast of Guam (13° 12.829' N, 144°, 43.178' E) in about 850 m of water on 16 April 2016. Specimens are unspecified individuals photographed on the deck of the F.V. Stonefish and catalogued among USNM 440413–440431

the Guam specimens rounded to the nearest millimetre because many specimens sustained length distortion and other damage.

2.2 | Stomach contents of new specimens from Guam

Stomach contents were initially surveyed for the 18 individually catalogued specimens (USNM 440413–440430) using radiographs. We removed the stomach contents from four of these specimens (USNM 440413, USNM 440416, USNM 440418 and USNM 440419) for study under a dissecting microscope to confirm initial identifications made from the radiographs.

2.3 | Other material

Institutional abbreviations follow Sabaj (2016). In addition to the new specimens from Guam, we studied 10 specimens of *A. japonicus* from the ANSP, BMNH, MNHN and USNM collected since Tyler (1968). This material included juveniles and adults (Figure 3). The localities of all records of *A. japonicus* are summarized in Figure 1.

2.4 | Tissue samples and DNA extractions

Tissue samples were collected from two of the new specimens of *A. japonicus* (USNM 440413 and USNM 440414). DNA was extracted using the DNeasy tissue kit (Qiagen; www.qiagen.com) according to the manufacturer's protocol. Double-stranded DNA was synthesized via PCR using universal primers for the mitochondrial 16 s subunit (Kocher *et al.*, 1989) marker. The PCR products were electrophoresed in 1.5% low-melting-point agarose gels using a Tris-acetate buffer for

quality control and submitted for purification and sequencing in both directions to the sequencing and genomics facility of the University of Puerto Rico (www.sgf.hpcf.upr.edu). Sequences were assembled and aligned manually with Geneious 10.0.9 (www.geneious.com).

2.5 | Taxon sampling and phylogenetic methods

Molecular data were newly generated only for *A. japonicus*. Additional sequence data for 224 tetraodontiform species were retrieved from the National Center for Biotechnology Information (NCBI) database, including nine of the 24 valid species of Triacanthodidae. Valid species of triacanthodids are summarized in Matsuura (2015: table 1), except that there are only 23 species in that table because *Mephisto albomaculatus* Matsuura, Psomadakis & Than Tun 2018 was described subsequently by Matsuura *et al.* (2018). One taxon included, *Triacanthodes* sp., presumably is one of the two species of this genus not included in our dataset; *i.e.*, *Triacanthodes indicus* Matsuura 1982 or *Triacanthodes intermedius* Matsuura & Fourmanoir 1984. Our dataset includes eight of the 11 valid genera in Triacanthodidae (all except *Hollardia* Poey 1861, *Johnsonina* Myers 1934 and *Mephisto* Tyler 1966). Six genetic markers were examined: two mitochondrial genes (*col* and *16 s*) and four nuclear markers (ENC1, GLYT, MYH6 and PLAGL2). The data for each species and marker are summarised in Supporting Information Table S1. We analysed the concatenated dataset using maximum likelihood (ML) in RAxML 8.2.10 (Stamatakis, 2006) partitioning by codon position and selecting the best tree inferred from 10 independent replicates generated by the GTRGAMMA model. We assessed branch support using 1000 replicates using RAxML's rapid bootstrapping algorithm, with the collection of sample trees used to draw the bipartition frequencies on the optimal tree. We performed all RAxML analyses using CIPRES portal 3.3. Sequence data produced by this study are deposited in GenBank (MH796904).

3 | RESULTS

3.1 | Synonomy

Atrophacanthus japonicus (Kamohara 1941)

Ukeguchi-kawamuki (Japanese name)

Upward-Mouth Spikefish (new English name based upon that of Kamohara, 1941)

Tydemania japonica Kamohara, 1941: p. 166 (original description; Japanese name: Ukeguchi-kawamuki; type locality: Kochi Prefecture, Shikoku, Japan, on the bottom at > 100 fathoms (c. 183 m); illustration). Kamohara, 1952: p. 60 (description; illustration; type locality given as Mimase (Kochi Prefecture)). Tomiyama and Abe, 1958: p. 30 (description; colour illustration; Kochi Prefecture).

Atrophacanthus danae Fraser-Brunner, 1950: p. 3 (original description; type species of *Atrophacanthus*; illustration; type localities: several *Dana* stations in the Celebes Sea, bathypelagic nets between 600 and 5000 m of wire out over depths of 2160–4950 m). de Beaufort and Briggs, 1962: pp. 262, 266–268 (included in key; illustrations and description after Fraser-Brunner, 1950). George and Dayanandan, 1966: pp. 220–221 (description and illustration of




FIGURE 3 (a) Alcohol preserved and (b) Radiograph of adult *Atrophacanthus japonicus* ANSP 103654, 88.0 mm standard length, collected in a benthic trawl off the coast of Tanzania (06° 48' S, 39° 51' E), bottom depth 300–400 m, by the R.V. Anton Bruun on 20 November 1964

CMFRI-F.179/497, 98 mm total length (L_T), caught in bottom trawl off southwest coast of India. Murty, 1969: p. 34 (listing of CMFRI-F.179/497 at Central Marine Fisheries Institute, India).

***Atrophacanthus* [sic] *japonicus*.** Kamohara, 1958: p. 51 (listed with *danae* as synonym; Kochi Prefecture, Japan to Celebes). Kamohara, 1961: p. 5 (synonymy; references; type destroyed and no other specimens available for neotype designation). Kamohara, 1964: p. 67 (listed with *danae* as synonym; illustration; Kochi Prefecture to Celebes).

***Atrophacanthus japonicus*.** Tyler, 1968: pp. 163–175 (synonymy; description; illustration; photograph; drawing of lateral view of entire skeleton; discussion of relationships; discussion and figure about depth range). Tyler, 1970a: p. 3 (description of an adult specimen from a bottom trawl in Arabian Sea off India (CMFRI-F.179/497) and of four juvenile specimens from $07^{\circ} 00' N$, $139^{\circ} 56' E$). Tyler, 1970b: pp. 8–9, 43 (description and illustration of caudal skeleton). Tyler, 1980: pp. 70–72 (illustration of bones of head; discussion of relationships condensed from Tyler, 1968). Aboussouan and Leis, 1984: pp. 452–454, fig. 247 (eggs unknown; larvae probably unpigmented; composite illustration of specimens 2.6–2.7 mm L_T). Gloerfelt-Tarp and Kailola, 1984: pp. 281 (colour photo and description of 48 mm standard length (L_S) specimen, USNM 265190, from Sumatra, Indonesia). Matsuura, 1984: p. 357 (listed; illustration; distribution from Kochi Prefecture to Philippines, Celebes Sea, and east coast of Africa). Tyler, 1986: p. 887 (description; distribution includes Tanzania). Adam et al., 1998: p. 16 (62 mm L_T specimen, BMNH 1996.9.25.1, from the Maldives Islands, $04^{\circ} 05' N$, $73^{\circ} 20' E$). Matsuura, 2000: p. 647 (listed; South China Sea). Shinohara et al., 2001: p. 337 (listed; Tosa Bay, Japan (off Kochi Prefecture, near type locality); Japanese name: Ukeguchi-kawamuki). Bijukumar and Deepthi, 2009: p. 153 (caught in trawl bycatch off Kerala, India; size range of specimens of *A. japonicus* 54–97 mm L_T). Konstantinidis and Johnson, 2012a: pp. 98–114, fig. 3C–F (photographs of cleared and stained specimens and descriptions of caudal skeleton in three specimens, 14.5 mm, 18 mm and 58 mm L_S). Konstantinidis and Johnson, 2012: pp. 351–366, fig. 1C, D (photographs of cleared and stained specimens and descriptions of jaw and suspensorium of two specimens, 18 mm and 58 mm L_S). Matsuura, 2015: pp. 74–76 (valid species; distribution given as off Tanzania, Maldives Islands, Japan and Celebes Sea).

3.2 | Colouration of new specimens from Guam

The new specimens of *A. japonicus* from Guam have pinkish red bodies except for strong silvery overtones in the abdominal region and lower half of the head, beginning at about the middle of the eye and continuing ventrally (Figure 2). In alcohol, the new specimens are uniformly light in colour except for the webbing between the first, second and third dorsal-fin spines, where there are small dark spots (not visible in Figure 3).

3.3 | Diagnostic features of new specimens from Guam

The following 12 features are diagnostic for *A. japonicus* and present in the new specimens from Guam. Average values for the

18 individually catalogued specimens from Guam are given in parentheses where applicable: (1) only the first three dorsal-fin spines well developed and prominently exposed, with the fourth spine much shorter than the first three but frequently exposed and easily seen externally and the fifth and sixth spines reduced in size and barely protruding to the surface or not protruding at all; (2) first dorsal-fin spine relatively short, not reaching much beyond fourth spine when depressed; (3) dorsal-fin rays 13 or 14 (14.0); (4) anal-fin rays 11–13 (12.8); (5) pectoral-fin rays 13–15 (14.0); (6) a single, short pelvic-fin ray protruding to the exterior just behind the locking flange on the posterior region of the pelvic-fin-spine base (single, short pelvic ray present); (7) short conical teeth in a single row, 19–23 in upper jaw and 22–27 in lower jaw (short conical teeth in a single row, 22 in upper jaw and 24 in lower jaw); (8) profile of head straight; (9) mouth distinctly supraterminal; (10) snout short, about equal to or somewhat longer than the postorbital distance, snout length 9.4%–12% L_S (10.9% L_S); (11) gill opening moderately long, usually extending down about one-half to four-fifths of the pectoral-fin base, gill-opening length 6.1%–8.0% L_S (7.2% L_S); and (12) pelvis narrow, basin-like, being flat ventrally with low dorsally upturned lateral edges, usually about five times in pelvic length, width into length 4.6–5.7 times (5.1 times).

3.4 | Diet

The stomach contents of the new juvenile specimens from Guam contained predominantly two species of pelagic thecosome pteropods and undetermined species of foraminiferans. Among the 18 radiographed specimens, only one individual had an empty stomach, whereas one individual had more than 40 pelagic thecosome pteropods and foraminiferans in its stomach and intestine (Figure 4).

3.5 | Molecular phylogeny

Our analysis (Figure 5) suggests that there are three major lineages among triacanthodids: (1) a clade composed of *Parahollandia* Fraser-Brunner 1941; (2) a clade including *Halimochirurgus* Alcock 1899, *Tydemania* Weber 1913, *Paratriacanthodes* Fowler 1934 and




FIGURE 4 Radiograph of the gut of *Atrophacanthus japonicus* USNM 440416, containing more than 40 pelagic thecosome pteropods and foraminiferans




FIGURE 5 Molecular phylogenetic relationships among genera of Triacanthodidae for which tissue samples are available (no tissue samples available for *Hollardia*, *Mephisto*, and *Johnsonina*)

Bathyphylax Myers 1934; and (3) a clade including *Triacanthodes* Bleeker 1857, *Atrophacanthus* and *Macrorhamphosodes* Fowler 1934. Of the 10 triacanthodid species (including *Triacanthodes* sp.) included in this analysis, *A. japonicus* is most closely related to *Macrorhamphosodes uradoi* (Kamohara 1933).

4 | DISCUSSION

4.1 | Distribution

The new specimens from Guam are conspecific with *A. japonicus* from Japan, Indonesia, the Philippines (Celebes Sea), Maldives, India and East Africa (Figure 1; for locality records see: Adam *et al.*, 1998; Matsuura, 1984, 2000, 2015; Murty, 1969; Shinohara *et al.*, 2001; Tyler, 1968, 1970a). The specimens of *A. japonicus* from Guam represent a range extension eastward into Oceania from a station sampled by the R.V. Vityaz between the Caroline Islands and New Guinea ($7^{\circ} 00' N$, $139^{\circ} 56' E$ (Tyler, 1970a: p. 3).

4.2 | Depth records of life stages

Atrophacanthus japonicus has been collected infrequently but occasionally in large numbers and more than 250 specimens are now available. Larval and juvenile stages ($\leq c. 55 \text{ mm } L_S$) are bathy and mesopelagic and have been collected at depths of c. 200–2500 m (Tyler, 1968: p. 171); *n.b.*, depth provided is depth collected in water column, not bottom depth. Juvenile stages have twice been taken in relatively large numbers: (1) in the fine mesh nets of many stations of the *Dana* Expedition in the Celebes Sea, with one station (3739 VI) obtaining 145 specimens (Tyler, 1968: p. 174); and (2) the new

96 specimens from the stomachs of three *Thunnus albacares* (Bonnaterre 1788) off Guam that we describe herein. Based upon *Dana* Expedition material, Fraser-Brunner (1950) noted that the majority of *A. japonicus* were collected in water with a bottom depth of 4000 m and no specimens were collected in water less than 2000 m deep; however, none of the *A. japonicus* studied by Fraser-Brunner were adults. Only six adult specimens of *A. japonicus* ($c. 60 \text{ mm to } 117.2 \text{ mm } L_S$) are known; all were collected from benthic bottom trawls at relatively shallower depths of 121–491 m (all collected as single individuals except the holotype and paratype of *Tydemania japonica* (Kamohara 1941), which were collected together). Juveniles that have been collected benthically range from 46.9 to 54.9 mm L_S , so while there is some variation in the size at which *A. japonicus* settle, only large juveniles have been collected benthically. This suggests that larvae and juveniles have an extended pelagic period over deep water and subsequently settle as young adults into shallower benthic shelf and slope habitats.

We interpret that the juvenile specimens of *A. japonicus* from the *T. albacares* stomachs had been ingested recently because of their bright coloration and only superficial digestion. *Thunnus albacares* usually feed in the top 100 m of the water column, so it is likely that they encountered the specimens of *A. japonicus* in the pelagic zone; however, *T. albacares* are known to dive occasionally to $> 1000 \text{ m}$ (Schaefer *et al.*, 2007), so it is possible, albeit unlikely, that the new specimens had settled into benthic habitats when the *T. albacares* ate them. We emphasize that the gut contents of the specimens of *A. japonicus* from Guam contained prey that included pelagic taxa (two species of thecosome pteropods and undetermined foraminiferans). These observations strongly suggest to us that the *T. albacares* ingested these large juvenile *A. japonicus* in the pelagic environment. Fraser-Brunner (1950) reported a similar diet for the *A. japonicus* he studied (pteropods and foraminiferans, as well as ostracods and *Sagitta* spp.) that were collected in the water column by the *Dana* Expedition.

4.3 | Other Triacanthodidae known from the Mariana Islands

Two other species of triacanthodids are known from the Mariana Islands: *Halimochirurgus alcocki* Weber 1913 (Myers, 1988; Myers & Donaldson, 2003; Tenorio, 2014) and, based upon underwater photographs taken aboard the NOAA R.V. *Okeanos Explorer*, *Hollardia* cf. *goslinei* Tyler 1968 also occurs in the region. The photographs show *H. cf. goslinei* feeding on the bottom at depths of 359–525 m on the Zealandia Bank of the Marianas on 21, 22 April and 3 May 2016 (photographs available through NOAA Office of Ocean Exploration and Research (OER), 2016 Deepwater Exploration of the Marianas: D2-EX1605-L1, images 01-01:51:17, 02-00:47:53, 12-00:22:48 and 12-00:23:36. Four photographs taken at locations between $12^{\circ} 43' 57'' N$ to $16^{\circ} 54' 06'' N$ and $144^{\circ} 16' 21'' E$ to $145^{\circ} 53' 53'' E$). R.V. *Okeanos Explorer* subsequently photographed another individual of *H. cf. goslinei* on the bottom at 448 m on 13 August 2016 off Wake Island (see NOAA OER Deepwater Wonders of Wake: Exploring the Pacific Remote Islands Marine National Monument: D2-EX1606-11, image 03:00:31. The photo was taken at $19^{\circ} 17' 21.0'' N$, $166^{\circ} 39'$

55.5° E). *Hollardia goslinei* is known from nine specimens and *in situ* observations from the Hawaiian Archipelago (Chave & Malahoff, 1998; Chave & Mundy, 1994; Matsuura, 2015), as well as several other Pacific localities where R.V. *Okeanos Explorer* has observed them.

4.4 | Phylogenetic placement

Morphological studies have proposed that *Atrophacanthus* and a clade containing *Bathyphylax* + *Halimochirurgus* and another including *Tydemania* + *Macrorhamphosodes* were related lineages derived from a *Paratriacanthodes*-like clade (Tyler, 1968, 1980). Our molecular analysis supports three major clades among triacanthodids: (1) a clade composed of *Parahollardia*; (2) a clade including *Halimochirurgus*, *Tydemania*, *Paratriacanthodes* and *Bathyphylax*; and (3) a clade including *Triacanthodes*, *Atrophacanthus* and *Macrorhamphosodes* (Figure 5). The close relationship of *Atrophacanthus* and *Macrorhamphosodes* is contrary to the interpretation of Tyler (1968: pp. 164–166) and Tyler (1980: p. 70), in which *Tydemania* was thought to have the closest relationship to *Macrorhamphosodes*, but those studies were based only on osteological features. We do not attempt to resolve that issue herein because further taxonomic and genetic sampling are needed to obtain a better understanding of the relationships among triacanthodid genera. We note that the support evidenced in our molecular phylogeny clearly challenges the non-cladistic conclusions of Tyler (1968, 1980) concerning these genera of triacanthodids.

To date, neither a cladistically based morphological analysis nor a comprehensive molecular phylogenetic analysis has assessed the relationships within the Triacanthodidae. The deep habitats of Triacanthodidae (continental shelf, slope and mesopelagic for juveniles of *A. japonicus*), comparatively infrequent collection and relatively low number of specimens available for study have challenged such research. Our ML analysis provides strong support for the sister-group relationship of *Atrophacanthus* and *Macrorhamphosodes* (bootstrap support = 99%), but we still lack DNA sequence data for three genera considered to be among the earliest branching lineages (*Hollardia*, 3 species; *Mephisto*, 2 species; and the monotypic *Johnsonina*; Tyler, 1968, 1980).

Our molecular results imply two independent origins of elongated snouts within triacanthodids (*Halimochirurgus* and *Macrorhamphosodes*). It also implies that lepidophagy (scale feeding) has evolved twice, once in *Tydemania* (Mok, 1978) and also in *Macrorhamphosodes* (Nakae & Sasaki, 2002). Future studies addressing instances of morphological and ecological convergence among triacanthodids may provide insights into the evolutionary mechanisms responsible for convergence.

5 | OTHER MATERIAL EXAMINED

Each lot only has one specimen unless otherwise indicated.

Atrophacanthus japonicus: 105 specimens examined: ANSP 103654, 88.0 mm L_S , Tanzania; BMNH 1987.1.23.51–52, two specimens, est. 30 mm, est. 35 mm L_S , South Indonesia; BMNH 1987.1.23.54–56, three specimens, est. 32 mm, est. 20 mm, est. 23 mm L_S , South Indonesia; BMNH 1996.9.25.1, 60.5 mm L_S , Maldives; MNHN 2002–2990, 54.9 mm L_S , Philippines.

MNHN 2014-2112, 117.2 mm L_S , Madagascar; USNM 265190, 46.9 mm L_S , Sumatra, Indonesia; USNM 440413, 51 mm L_S , Guam; USNM 440414, 47 mm L_S , Guam; USNM 440415, 54 mm L_S , Guam; USNM 440416, 44 mm L_S , Guam; USNM 440417, 51 mm L_S , Guam; USNM 440418, 42 mm L_S , Guam; USNM 440419, 51 mm L_S , Guam; USNM 440420, 49 mm L_S SL, Guam; USNM 440421, 48 mm L_S , Guam; USNM 440422, 51 mm L_S , Guam; USNM 440423, 46 mm L_S , Guam; USNM 440424, 49 mm L_S , Guam; USNM 440425, 44 mm L_S , Guam; USNM 440426, 38 mm L_S , Guam; USNM 440427, 47 mm L_S , Guam; USNM 440428, 48 mm L_S , Guam; USNM 440429, 46 mm L_S , Guam; USNM 440430, 51 mm L_S , Guam; USNM 440431, 78 specimens, 37–50 mm L_S , Guam.

ACKNOWLEDGEMENTS

We are grateful to J. and K. Borja for their keen interest in marine sciences and their practice of preserving interesting organisms they come across while fishing from their vessel F.V. *Stonfish*. J. Borja froze the specimens of *Atrophacanthus japonicus* he obtained from the *T. albacares* stomachs and began the process leading to the specimens making their way to Honolulu through the good offices of B. Mundy, NOAA National Marine Fisheries Service, Hawaii and his many cooperating fishery biologists in Guam and Hawaii. The frozen materials were sent to the Division of Fishes, National Museum of Natural History, Smithsonian Institution, where they were cared for and tissue sampled by D. Pitassy and radiographed and photographed by S. Raredon. The efforts of all the above are greatly appreciated.

We thank M. Arce-H. and M. Sabaj at ANSP, J. MacLaine BMNH and Z. Gabsi MNHN, for providing loans of triacanthodids for our study. For their help in obtaining images and associated data of *Hollardia* cf. *goslinei* photographed by the NOAA R.V. *Okeanos Explorer* we thank C. Kelley and M. Putts at the University of Hawaii and M. Vecchione and K. Cantwell at NOAA. We benefited from discussions about tuna feeding with B. Collette, Smithsonian Institution. M. Vecchione, NOAA and R. Mann and J. Conroy, Virginia Institute of Marine Science, advised us on pteropod identification. K. Matsuura provided a translation for the Japanese common name of *A. japonicus*, which we offer as the English name. W. Bemis helped prepare the map. We thank D. Tyler for editorial comments and for hosting a productive writing week. B. Mundy provided a thorough review of the manuscript and an anonymous reviewer was also helpful. This paper is contribution No. 3805 of the Virginia Institute of Marine Science, College of William & Mary.

ORCID

Katherine E. Bemis  <https://orcid.org/0000-0002-7471-9283>

REFERENCES

Aboussouan, A., & Leis, J. M. (1984). Balistoidei: Development. In H. G. Moser, W. J. Richards, D. M. Cohen, M. P. Fahay, A. W. Kendall, & S. L. Richardson (Eds.), *Ontogeny and systematics of fishes*, Special Publication Number 1, American Society of Ichthyologists and Herpetologists (pp. 450–459). Lawrence, KS: Allen Press.
 Adam, M. S., Merrett, N. R., & Anderson, R. C. (1998). Additions to the fish fauna of the Maldivian Islands. Part 1: An Annotated checklist of the

deep demersal fishes of the Maldivian Islands. *Ichthyological Bulletin of the J.L.B. Smith Institute of Ichthyology*, 67, 1–19.

de Beaufort, L. F., & Briggs, J. C. (1962). Scleroparei, Hypostomides, Pediculati, Plectognathi, Opisthomii, Discocephali, Xenopterygii. In *The fishes of the Indo-Australian Archipelago* (Vol. 11: i–xi +, pp. 1–481). Leiden: E. J. Brill.

Bijukumar, A., & Deepthi, G. R. (2009). Mean trophic index of fish fauna associated with trawl bycatch of Kerala, southwest coast of India. *Journal of Marine Biology Association of India*, 51, 145–157.

Chave, E. H., & Malahoff, A. (1998). *In deeper waters: Photographic studies of Hawaiian deep-sea habitats and lifeforms*. Honolulu, HI: University of Hawai'i Press.

Chave, E. H., & Mundy, B. C. (1994). Deep-sea benthic fish of the Hawaiian Archipelago, Cross Seamount and Johnston Atoll. *Pacific Science*, 48, 367–409.

Fraser-Brunner, A. (1950). *Studies in plectognath fishes from the "Dana" Expeditions. I: An interesting new genus of triacanthodid fishes from the Celebes Sea* (Dana Report No. 35) pp. 1–8. Copenhagen: Andr. Fred. Host & Son.

George, K. C., & Dayanandan, M. G. (1966). *Atrophacanthus danae* Fraser Brunner and *Chaetodon jayakari* Norman, new records of fishes from the Arabian Sea off the South-West coast of India. *Journal of the Marine Biological Association of India*, 8, 220–222.

Gloerfelt-Tarp, T., & Kailola, P. J. (1984). *Trawled fishes of southern Indonesia and northwestern Australia*. Australia and Singapore: Australian Development Assistance Bureau, Australia, Directorate General of Fishes, Indonesia and German Agency for Technical Cooperation, Federal Republic of Germany.

Kamohara, T. (1941). Descriptions of one new and two rare fishes from Japan. *Annotations Zoologicae Japonenses*, 20, 166–168.

Kamohara, T. (1952). Revised descriptions of the offshore bottom-fishes of Prov. Tosa, Shikoku, Japan. *Research Reports of the Kochi University Natural Science No. 3*. pp. 1–122. Kochi, Japan: Kochi University.

Kamohara, T. (1958). A catalogue of fishes of Kochi Prefecture (Province Tosa), Japan. *Reports of the USA Marine Biological Station*, 5, 1–76.

Kamohara, T. (1961). Notes on the type specimens of fishes in my laboratory. *Reports of the USA Marine Biological Station*, 8, 1–9, Pls. 1–7.

Kamohara, T. (1964). Revised catalogue of fishes of Kochi Prefecture, Japan. *Reports of the USA Marine Biological Station*, 11, 1–99.

Kocher, T. D., Thomas, W. K., Meyer, A., Edwards, S. V., Pääbo, S., Villablanca, F. X., & Wilson, A. C. (1989). Dynamics of mitochondrial DNA evolution in animals: Amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences*, 86, 6196–6200. <https://doi.org/10.1073/pnas.86.16.6196>.

Konstantinidis, P., & Johnson, G. D. (2012a). A comparative ontogenetic study of the tetraodontiform caudal complex. *Acta Zoologica*, 93, 98–114. <https://doi.org/10.1111/j.1463-6395.2010.00490.x>.

Konstantinidis, P., & Johnson, G. D. (2012b). Ontogeny of the jaw apparatus and suspensorium of the Tetraodontiformes. *Acta Zoologica*, 93, 351–366. <https://doi.org/10.1111/j.1463-6395.2011.00509.x>.

Matsuura, K. (1984). Tetraodontiformes. In H. Masuda, K. Amaoka, C. Araga, T. Uyeno, & T. Yoshino (Eds.), *The fishes of the Japanese Archipelago* (pp. 357, 1–366). Shinjuku-ku Tokyo, Japan: Tokai University Press.

Matsuura, K. (2000). Tetraodontiformes. In J. E. Randall & K. K. P. Lim (Eds.), *A checklist of the fishes of the South China Sea*, The Raffles Bulletin of Zoology, Supplement 8 (pp. 647, 569–649, 667). Singapore: National University of Singapore.

Matsuura, K. (2015). Taxonomy and systematics of tetraodontiform fishes: A review focusing primarily on progress in the period from 1980 to 2014. *Ichthyological Research*, 62, 72–113. <https://doi.org/10.1007/s10228-014-0444-5>.

Matsuura, K., Psomadakis, P., & Than Tun, M. (2018). *Mephisto albomaculosus*, a new spikefish (Actinopterygii: Tetraodontiformes: Triacanthodidae) collected off Myanmar, Indian Ocean. *Ichthyological Research*, 66, 1–4. <https://doi.org/10.1007/s10228-018-0642-7>.

Mok, H. K. (1978). Scale-feeding in *Tydemania navigatoris* (Pisces: Triacanthodidae). *Copeia*, 1978(2), 338–340. <https://doi.org/10.2307/1443577>.

Murty, V. S. (1969). Catalogue of fishes (excluding from the Laccadives) in the reference collections of the Central Marine Fisheries Research Institute. *Central Marine Fisheries Research Institute, Bulletin*, 10, 1–36.

Myers, R. F. (1988). An annotated checklist of the fishes of the Mariana Islands. *Micron*, 21, 115–180.

Myers, R. F., & Donaldson, T. J. (2003). The fishes of the Mariana Islands. *Micron*, 35–36, 594–648.

Nakae, M., & Sasaki, K. (2002). A scale-eating triacanthodid, *Macrorhamphosodes uradoi*: prey fishes and mouth "handedness" (Tetraodontiformes, Triacanthoidei). *Ichthyological Research*, 49, 7–14. <https://doi.org/10.1007/s102280200001>.

Sabaj, M. H. (2016, August 16). Standard Symbolic Codes for Institutional Resource Collections in Herpetology and Ichthyology: An Online Reference (Version 6.5). Washington, DC: American Society of Ichthyologists and Herpetologists. Retrieved from <http://www.asih.org/>

Schaefer, K. M., Fuller, D. W., & Block, B. A. (2007). Movements, behavior and habitat utilization of Yellowfin Tuna (*Thunnus albacares*) in the northeastern Pacific Ocean, ascertained through archival tag data. *Marine Biology*, 152, 503–525.

Shinohara, G., Endo, H., Matsuura, K., Machida, Y., & Honda, H. (2001). Annotated checklist of the Deepwater fishes from Tosa Bay, Japan. *National Science Museum, Monograph*, 20, 283–343.

Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics*, 22, 2688–2690.

Tenorio, M. C. (2014). *Updates to the checklist of fishes of the CNMI*. Division of Fish and Wildlife, Commonwealth of the Northern Mariana Islands. viii+5 pp.+34 pp. Retrieved from http://www.cnmi-dfw.com/docs/Checklist_of_fishes_2014_11714.pdf

Tomiyama, I., & Abe, T. (1958). *Encyclopaedia zoological illustrated in colours*. Tokyo: Pisces and cyclostoma.

Tyler, J. C. (1968). A monograph on plectognath fishes of the superfamily Triacanthoidea. *Academy of Natural Sciences of Philadelphia, Monograph*, 16, 1–364.

Tyler, J. C. (1970a). New records of triacanthoid plectognath fishes. *Academy of Natural Sciences of Philadelphia, Notulae Naturae*, 435, 1–7.

Tyler, J. C. (1970b). The progressive reduction in number of elements supporting the caudal fin of fishes of the order Plectognathi. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 122, 1–85.

Tyler, J. C. (1980). Osteology, phylogeny and higher classification of the fishes of the order Plectognathi (Tetraodontiformes). *NOAA Technical Report, NMFS Circular*, 434, 1–422.

Tyler, J. C. (1986). Family No. 265: Triacanthodidae, Spikefishes, Spyker-visse. In M. M. Smith & P. C. Heemstra (Eds.), *Smith's sea fishes* (pp. 887–890). Grahamstown, South Africa: Smith Institute of Ichthyology, Macmillan.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Bemis KE, Tyler JC, Arcila D. Life history, distribution and molecular phylogenetics of the Upward-Mouth Spikefish *Atrophacanthus japonicus* (Teleostei: Tetraodontiformes: Triacanthodidae). *J Fish Biol*. 2019;1–7. <https://doi.org/10.1111/jfb.13923>