

Immune System Promiscuity in Human and Nonhuman Primate Evolution

Jessica F. Brinkworth^{1,2*} and Courtney C. Babbitt^{3*}

ABSTRACT

Many genes that respond to infection have functions outside of immunity and have been found to be under natural selection. Pathogens may therefore incidentally alter nonimmune physiology through engagement with immune system genes. This raises a logical question of how genetically promiscuous the immune system is, here defined as how heavily cross-referenced the immune system is into other physiological systems. This work examined immune gene promiscuity across physiological systems in primates by assessing the baseline (unperturbed) expression of key tissue and cell types for differences, and primate genomes for signatures of selection. These efforts revealed “immune” gene expression to be cross-referenced extensively in other physiological systems in primates. When immune and nonimmune tissues diverge in expression, the differentially expressed genes at baseline are enriched for cell biological activities not immediately identifiable as immune function based. Individual comparisons of immune and nonimmune tissues in primates revealed low divergence in gene expression between tissues, with the exception of whole blood. Immune gene promiscuity increases over evolutionary time, with hominoids exhibiting the most cross-referencing of such genes among primates. An assessment of genetic sequences also found positive selection in the coding regions of differentially expressed genes between tissues functionally associated with immunity. This suggests that, with increasing promiscuity, divergent gene expression between the immune system and other physiological systems tends to be adaptive and enriched for immune functions in hominoids.

The human immune system is a complex, multitiered system of cellular, tissue, and protein components that comprises the primary defense of our species against invading microorganisms. It is an ancient, energetically expensive biological system that is fundamental to the survival of living things. Given its importance, it should not be surprising that the immune system is under tremendous selective pressure and that the genes involved can rapidly diversify (Barreiro et al. 2008; Cortez et al. 2017; Ferrer-Admetlla et al. 2008; Kosiol et al. 2008; Novembre and Han 2012; Pickrell et al. 2009; Wang et al. 2017). Pathogen-mediated selection tends to be viewed as the primary force influencing human immune system evolution (Ferrer-Admetlla et al. 2008; Fumagalli

¹Evolutionary Immunology and Genomics Laboratory, Department of Anthropology, University of Illinois Urbana–Champaign, Urbana, Illinois, USA.

²Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana–Champaign, Urbana, Illinois, USA.

³Department of Biology, University of Massachusetts Amherst, Amherst, Massachusetts, USA.

*Jessica F. Brinkworth, Department of Anthropology, University of Illinois Urbana–Champaign, 109, 607 S. Mathews Ave., Urbana, IL 61801 USA. E-mail: jfbrinkw@illinois.edu.

Courtney C. Babbitt, Department of Biology, University of Massachusetts Amherst, 221 Morrill Science Center III, 611 North Pleasant St., Amherst, MA 01003 USA. E-mail: cbabbitt@bio.umass.edu.

KEY WORDS: EVOLUTIONARY IMMUNOLOGY, IMMUNE SYSTEM, PLEIOTROPY, PRIMATE IMMUNITY, IMMUNE PROMISCUITY, PRIMATE IMMUNOGENOMICS.

et al. 2011). Eukaryotic life, after all, was born into an environment where prokaryotes, viruses, and archaea were already competing for resources via infection (Casanova 2015). Moreover, humans have evolved in a diverse microbial landscape, under profound, routine assault by pathogens, and have carried a heavy disease burden for most of our existence (Fumagalli et al. 2009; Harper and Armelagos 2010, 2013; Karlsson et al. 2014). Prior interactions with microorganisms appear to be reflected in our genomes. Genes with clear immune functions tend to comprise the largest functional categories of positively selected genes in human populations and across mammalian species (Kosiol et al. 2008; Vallender and Lahn 2004). While pathogen-mediated selection is no doubt an extremely important factor driving the evolution of the species, this framework of immune system evolution prioritizes microbial conflict as *the* factor shaping immunity and obscures the importance and impact of the system being highly cross-referenced physiologically on genomic evolution (Brinkworth 2017).

The immune system is very physiologically promiscuous, stretching into every tissue, lending and co-opting genes to and from the developmental and cellular programs of “nonimmune” systems. Indeed, most genes that fulfill immune functions are pleiotropic and also engaged in activities traditionally considered to be “nonimmune” (Goselin et al. 2005; Hattermann et al. 2008; Johnson et al. 2009; Mori et al. 2016; Piccinin et al. 2010; Sferruzzi-Perri et al. 2009; Sheridan and Murphy 2013). The ancient origins of immunity and the increasing complexity of interactions between microorganisms and hosts over evolutionary time likely partially explain the multifunctional and cross-system infiltrative nature of immune system components. The “borrowing” and “lending” of immune system building blocks has led to multifunctionality in many of the components of the system, with associated genes appearing to have evolved in parallel with other systems, functions, and processes (Adler and Rogers 2005; Ambrosi et al. 2014; Kitaya et al. 2007; Réaux-Le Goazigo et al. 2013; Takizawa et al. 2008). System genetics approaches over the last decade have well established that most phenotypes, including characteristics of immunity, are controlled by pleiotropic networks of many genes (reviewed in Boyle et

al. 2017; Wagner and Zhang 2011; most recently referred to as the “omnigenic model of complex traits”: Boyle et al. 2017). While the evolution of single pleiotropic genes may sometimes be constrained due to their role in multiple phenotypes (Wagner and Zhang 2011), selection on phenotypes expressed via genes networks means that adaptive shifts in target genes are accompanied by small functional shifts in many other networked genes to compensate for changes at one physiological interface (Pavlicev and Wagner 2012). An important aspect of human immune gene pleiotropy is that many of the genes in question are involved in the development of tissues that manifest features considered to be particularly human and hominoid, for example, forebrain development and brain volume (Kee et al. 2003; Luo et al. 2012; Tsai et al. 2002), cognitive function (Ben Menachem-Zidon et al. 2008; Bliss and Collingridge 1993; Meola et al. 2013), and embryogenesis and placental invasion (Chatterjee et al. 2013; Das et al. 2002; Kitaya et al. 2007; Piccinni et al. 1998; Sferruzzi-Perri et al. 2009; Takahashi et al. 2006). In this context, factors driving the adaptive evolution of immunity may be contributing to the divergence and diversification of other traits and vice versa. More specifically, it should be expected that evolutionary interactions with pathogens could mediate changes in heavily immune cross-referenced physiology considered important to the human evolutionary story.

In this article, we illustrate the cross-referencing of the immune system in other physiological systems and, broadly, the degree to which immune tissues transcriptionally diverge from nonimmune tissues, using evolutionary immunology and genomic approaches to analyze previously published data sets (Clark et al. 2015; Derrien et al. 2012; Lin et al. 2014; Linsley et al. 2014; Peng et al. 2015). We contrast genome-wide gene expression of primate immune and nonimmune tissues by species and tissue type, examine differentially expressed (DE) genes to identify divergent functions and expression specificity across species, and assess DE genes for signatures of positive selection. We show that “immune” gene expression is extensively cross-referenced in other physiological systems in primates. These genes show high overlap in expression between immune and nonimmune tissue types and are enriched for a mix of cell biological activities both immune and nonimmune in nature.

Our analysis indicates that this cross-referencing increases over evolutionary time in primates, with positively selected divergent gene expression in hominoids enriched for immune function.

Multiple Human Nonimmune Tissues Have Been Immune Tissues in the Past

The highly networked and promiscuous nature of the immune system is at once a strong indication of the importance of host defense in species survival and an outcome of a very ancient system expanding through extensive genomic change over time. With novel species-specific changes in expanded and restricted immune gene families and differences in immune tissue development and immune cell content, the human immune system is a relatively new configuration of mammalian immunity (Boehm et al. 2012a; Brinkworth and Thorn 2013). Over the course of vertebrate evolution, for example, organs and tissues not considered to be involved in human immunity have been key generators of immune cells and components. B cell development in mammals, for instance, is completed almost entirely in the bone marrow, with minor activity occurring in the spleen. In fish and amphibians, B cells extensively develop in other locations, such as the kidneys (cartilaginous and bony fish), spleen (cartilaginous fish, amphibians), and liver (amphibians; Boehm et al. 2012a, 2012b; Du Pasquier et al. 2000; Greenhalgh et al. 1993; Pickel et al. 1993). Secondary lymphoid tissues and organs are usually differentiated structures that have entirely different primary functions but are at the interface of the host's internal and external environments. Gut-associated lymphoid tissue extensively develops in the gut tube, the central component of the digestive system. The spleen is primarily a blood filter, though it is also a site for lymphocyte development, and the action of filtering has been cited as having an immune-like effect, as a possible reason for malaria resistance in sickle-cell anemia carriers (Kwiatkowski 2005). Similarly, the genes the immune system have undergone co-option during immune system evolution. Toll-like receptors, some of the most intensely studied genes that initiate immune responses against invading pathogens, are engaged in determining the ventral-dorsal axis in invertebrates (Ambrosi et al. 2014; Brennan et al. 2017). Chemokine and cytokine genes, which are key to the first line of defense in immune

responses, including cell trafficking, inflammation, and immune cell differentiation, are extensively cross-referenced in forebrain/brain development and neurogenesis (e.g., *CXCL12*, *CXCR4*, *CXCL16*, *CXCR6*, *IL3*; Hattermann et al. 2008; Klein et al. 2001; Luo et al. 2012; Réaux-Le Goazigo et al. 2013; Reiss et al. 2002). A host of chemokines engaged in antimicrobial activities are also key to normal neurological function (*CXCL12*, *CXCR4*, *CXCL8*/*IL-8*, *CCL2*, *CCR2*, *CXCL31*; Adler and Rogers 2005; Giovannelli et al. 1998; Gosselin et al. 2005; Piccinin et al. 2010) and cognitive memory (*CX3CL1*, *CX3CR1*; reviewed in Sheridan and Murphy 2013). Similarly, cytokines are very actively engaged in reproduction, including embryo implantation (reviewed in Mori et al. 2016), trophoblast differentiation (*CSF2*; Sferruzzi-Perri et al. 2009), placental invasion and development (*IRF1*, *STAT1*; Johnson et al. 2009; Kitaya et al. 2007), decidual/maternal blood pressure regulation (*IL-4*, *IL-40*, *CSF1*, *LIF*; Chatterjee et al. 2013; Piccinin et al. 1998, 2001), and platelet architecture and embryonic hematopoiesis (cytokine regulator *SH2B3*, *IL-7*; Dravid et al. 2011; Takizawa et al. 2008). This cross-referencing of immune genes with “nonimmune” tissues means that evolutionary interactions with pathogens may be altering nonimmune phenotypes through direct interactions with genes in the responsible gene network.

Primates Respond to Bacteria with Genes Engaged in Nonimmune Activities

From an evolutionary standpoint the net effect of immune promiscuity may be that pathogens interacting with the immune system exert direct selective pressure on genes responsible for both immune and nonimmune phenotypes. To illustrate that possibility, we assessed the top 10 upregulated genes that responded in human, chimp, and rhesus macaque monocytes stimulated with lipopolysaccharide (LPS) derived from *Escherichia coli* by Barreiro et al (2010), against the GeneCards (<https://www.genecards.org>) and OMIM (Online Mendelian Inheritance in Man; <https://omim.org/>) databases (Amberger et al. 2015; Stelzer et al. 2016), as well as an extensive literature search for known functions. Here, “top 10 upregulated” refers to the 10 genes most strongly upregulated across all three species. This simple examination revealed that most of these 10 genes (70%) that were strongly

upregulated in response to challenge by gram-negative bacteria are involved in two physiological systems that significantly inform reproductive success, the reproductive tract, and the central nervous system. These findings broadly agree with our prior examination of multiple human cell whole genome responses to bacterial (*Mycobacterium tuberculosis*), viral (influenza A), and parasitic (*Trypanosoma cruzi*) pathogens, in which we examined the top five upregulated responding genes shared across cell infections and found they were highly multifunctional and played important roles in other physiological systems such as digestive and reproductive systems (Brinkworth 2017). Similarly, we have previously found that immune genes that both are associated with a chronic inflammatory and autoimmune diseases and are signatures of positive selection in humans are also involved in important non-immune-related processes, such as uterine remodeling, decidua regulation, respiration, brain development, and sensory motor gating (Brinkworth and Barreiro 2014). In the present analysis, at least 5 of the 10 genes responding to LPS appear to be under mild to strong purifying selection in primates ($dN/dS = 0.01\text{--}0.85$, marked with * in Table 1) and are cross-referenced in central nervous system function (Table 1). In combination with previous results, the present examination suggests that, despite pleiotropy, genes responding to pathogen stimulus and involved in nonimmune functions are not always evolutionarily constrained by their multiple roles. The significant upregulation of genes involved in nonimmune activities in response to pathogen stimulus that are also under stronger purifying selection supports the possibility that pathogen-immune system evolutionary interactions can affect the function of nonimmune systems.

A Large Proportion of Genes Expressed in Nonimmune Tissues at Physiological Baseline Have a Role in Immunity: Genetic Promiscuity

In combination, the multifunctional nature of genes responding to immune challenge and evidence that these responding genes are under natural selection support the hypothesis that pathogens may incidentally alter nonimmune physiology through engagement with immune system genes. This raises a logical question of how genetically promiscuous the immune system is in primates,

here defined as how heavily cross-referenced into other physiological systems the immune system is in the primate order. We compared the whole-genome gene expression (RNA-Seq) of eight immunological and nonimmunological tissue types in five primate species at physiological baseline (i.e., tissues not known to be infected, injured, or otherwise unperturbed) using transcriptomes made publicly available by the Nonhuman Primate Reference Transcriptome Resource (NHPRTR) or published in peer-reviewed journals and archived the Genome Expression Omnibus (GEO) database (Clark et al. 2015; Derrien et al. 2012; Lin et al. 2014; Linsley et al. 2014; Peng et al. 2015). Primates included in the study were human (*Homo sapiens*), chimp (*Pan troglodytes*), rhesus macaque (*Macaca mulatta*), common marmoset (*Callithrix jacchus*), and the mouse lemur (*Microcebus murinus*). Tissues were categorized as immune versus nonimmune based on current known function in cellular immunity. Under this scheme, colon, lymph node, spleen, thymus, and whole blood represent primary and secondary lymphoid sites and were categorized as immune, and kidney, liver, and brain were categorized as nonimmune (Supplementary Table S1). As the evolutionary distance between species grows, the number of genes expressing species-level programs increases (Lin et al. 2014; Pishesha et al. 2014). Several previous studies have used transcriptomic data from multiple organs and multiple species (e.g., organ differences among amniotes) to look at relative rates of change in organs over time. Those studies have either directionally scaled their data or limited the data set to tissue-specific gene expression (Barbosa-Morais et al. 2012; Brawand et al. 2011; Breschi et al. 2016; Merkin et al. 2012). Here we are doing something different: since we seek to highlight genes that are expressed across physiological systems in primates, we have not directionally scaled the transcriptomic data, as doing so would mask this information. A principal components analysis (PCA) on Spearman's rank correlations of normalized expression of 8,869 genetic orthologs for all tissues (full data set), as well as a reduced matrix of only tissues available for all five species, reveals that 75.9–82.1% of the variation expressed (PC1) is explained by membership in the family Hominidae (Supplementary Figures S1 and S2). By contrast, very little gene expression variation (2.6%; PC4) in the full data set

Table 1. Immune and Nonimmune Functions of the Top 10 Genes Most Strongly Responding to Lipopolysaccharide (LPS) in Human, Chim, and Rhesus Macaque Monocytes

LPS-Responding Gene	Immune Functions (Selected)	Other Physiological Systems	Other Functions
<i>IL6*</i>	Acute-phase responses (Zilberstein et al. 1986) B-cell differentiation (Ikebuchi et al. 1987) Inflammation (Zilberstein et al. 1986) T17 and Treg cell development (Zheng et al. 2008)	Central nervous system Musculoskeletal Circulatory	Neurogenesis (Bauer et al. 2007; Deverman and Patterson 2009) Neuronal function (Fann and Patterson 1994; Marz et al. 1998; Oh and O'Malley 1994) Myokine/fat breakdown (Gudiksen et al. 2017; Wueest et al. 2014) Osteoclast activity (Ota et al. 1999 2001) Angiogenesis (Wei et al. 2003)
<i>CCL20</i>	Cell trafficking of DC, T, and B cells (Anderle et al. 2005; Ito et al. 2011) Antimicrobial activity toward <i>E. coli</i> (Hoover et al. 2002) Inflammation (Hromas et al. 1997)	Reproductive Genomic	Regulates sperm motility (Caballero-Campo et al. 2014; Diao et al. 2014) Induces DNA repair (Schwarz et al. 2002)
<i>IL12B*</i>	Th1 cell development (Cua et al. 2003) NK cell growth factor (Oppmann et al. 2000)	—	—
<i>IL1A</i>	Apoptosis (Hogquist et al. 1991) Fever generator (Cannon and Dinarello 1985) Antimicrobial activity toward <i>E. coli</i> (March et al. 1985)	Reproductive Musculoskeletal Central nervous system	Spermatogenesis (Sarkar et al. 2008) Ovulation (luteal phase; Cannon and Dinarello 1985) Osteoclast activity (Lord et al. 1991; Sabatini et al. 1988) Astroglial growth factor (Giulian et al. 1988)
<i>TNFAIP6</i>	Cell migration (Dyer et al. 2014, 2016)	Reproductive	Ovulation (Joyce et al. 2001; Ochsner et al. 2003)
<i>GCH1*</i>	Response to LPS (Werner-Felmayer et al. 1993)	Central nervous system	Dopamine synthesis (Duan et al. 2005)
<i>TNF</i>	Inflammation (Wu et al. 2011) Apoptosis (Obeid et al. 1993) Fever generator (Dinarello et al. 1986)	Respiratory Musculoskeletal Central nervous system	Lung morphogenesis (Zhu et al. 2007) Osteoclast differentiation (Komine et al. 2001) Smooth muscle proliferation (Barath et al. 1990; Ohta et al. 2005) Preserves synaptic strength (Beattie et al. 2002)
<i>PDE4B*</i>	Regulates interferon-gamma, interleukin-2 (Peter et al. 2007)	Central nervous system	Mood regulation (O'Donnell and Zhang 2004)
<i>G0S2</i>	Apoptosis (Welch et al. 2009)	Growth/metabolism	Lipolysis (Bednarski et al. 2016; Heier et al. 2015)
<i>ELOVL7*</i>	—	Growth/metabolism	Lipid metabolism and membrane development (Ohno et al. 2010)

* dN/dS = 0.01–0.85.

is explained by a tissue belonging to the immune or nonimmune categories, a finding consistent with promiscuous gene expression (Figure 1). This ordination on the PCA plot is due to the choice of scaling mentioned above.

To identify how many genes broadly categorized as immune genes are expressed in tissues considered to be nonimmune, we used the genes identified by the Gene Ontology consortium (GO) as “immune system process” (3,553 genes). We then compared these genes in humans against three nonimmune” tissues expressed in the five primate species. In the grouped nonimmune tissues, 1,352 of




FIGURE 1. Principal components analysis of baseline gene expression in eight immune and nonimmune tissues from five primate species. Brain, kidney, and liver were categorized as nonimmune, whereas colon, lymph node, whole blood, bone marrow, spleen, and thymus were categorized as immune. fr, brain frontal lobe; tc, temporal cortex. Principal components PC1 and PC4 are shown.

8,869 (15.2%) orthologous immune system process genes were consistently expressed above a counts per million normalized value (cpm) of ≥ 1 (Supplementary Table S2). *Homo-Pan* and *Homo-Macaca* comparison of coding regions revealed 44 of these cross-referenced genes with a dN/dS ratio > 1.2 suggesting they may be under positive selection in humans. A more refined comparison seeking possible cross-referencing of genes that are engaged in immune responses (GO: “activation of immune response”) in humans (826 genes) found that 301 genetic orthologs (3.6% orthologs and 36.4% of the GO category) are consistently expressed (cpm ≥ 1) in these nonimmune tissues as a group. Twelve of these genes appear to be under positive selection in humans (dN/dS > 1.2) and not in the other comparisons. These data lend support to the notion that a significant number of genes active in response to infection play a role in the function and maintenance of nonimmune tissues and that evolutionary change in genes categorized as immune may affect function in nonimmune tissues.




FIGURE 2. Overrepresentation analysis of GO biological processes in the immune-nonimmune significantly differentially expressed (DE) gene list, in semantic space. Bubbles represent collapsed Gene Ontology (GO) terms (REVIGO) that immune-nonimmune DE genes populate. Bubble size indicates frequency that genes in the DE list populate a GO term in the underlying Gene Ontology Annotation (GOA) database. Bubble color represents significance of GO term overrepresentation given the DE list and a background of 8,868 genes, with increasingly cool colors indicating lower FDR values ($-1/\text{FDR}$). Bubbles are arranged in semantic space according to relatedness of the GO terms. Clusters represent GO categories of similar biological processes.


Divergent Gene Expression between Immune and Nonimmune Tissue Is Associated with Basic Cell Function, Not Immunity

That the entire “immune system process” gene list is not cross-referenced in genes expressed at baseline in primate nonimmune tissues is not surprising. Logically, immune and nonimmune tissues must differ in gene expression as well. Given that pathogens play a strong role in the evolution of the immune system, it would be reasonable to expect that strong differences in baseline gene expression between immune and nonimmune tissues occur in genes with known immune functions. When nonimmune tissue gene expression is contrasted against gene expression in immune tissues across all five species, 16.3% (1,448) of genes are significantly differentially expressed (DE; 20%-fold change and a Benjamini-Hochberg false discovery rate [FDR] < 0.1 ; Supplementary Table S3). However, when these genes are examined for GO biological process category enrichment (overrepresentation analysis, GOrilla/REVIGO), they appear to be mainly engaged in activities other than immune function (allowed similarity of 0.7, SimRel algorithm, FDR < 0.001 ; Eden et al. 2009; Supek et al. 2011). The most strongly enriched GO biological process categories were found to be semantically similar to *ion transmembrane transport* (q -value = 8.1×10^{-9}) a biological process of membrane transport that is thought to have diversified significantly by cell type and transported substrate (Attwood et al. 2017), and three categories that could be lumped into cellular signaling, including *regulation of biological quality* (q -value = 5.13×10^{-3}), *single organism signaling* (q -value = 5.37×10^{-4}), and *protein activation cascade* (q -value = 2.6×10^{-5} ; Figure 2, Supplementary Table S4; GOrilla/REVIGO results). This is true also of the 838 (9.4%) genes that are upregulated in immune tissues versus nonimmune tissues (20%-fold change, FDR < 0.1), where the most strongly enriched GO biological process categories were found to be semantically similar to the *response to metal ion* category and include organic acid metabolic process, carboxylic acid metabolic process, and oxoacid metabolic process (q -values = 8.15×10^{21} , 1.15×10^{20} , 4.70×10^{20} ; Supplementary Table S4; GOrilla/REVIGO results). While cell signaling often includes cytokines, which complete antimicrobial activities directly and through cell communication, few other DE

genes in these categories have known antimicrobial functions (Selmaj et al. 1991). It appears that gene expression between immune and nonimmune tissues has primarily diverged in functions such as membrane transport and cell signaling that may impact immune activities but are not restricted to or necessarily immediately identifiable as immune function based.

Nonimmune Tissues Exhibit Low Divergent Gene Expression from Almost All Other Tissues

A by-product of the cross-referencing of genes between the immune system and other physiological systems should be homogeneity in differently expressed genes between immune tissues and nonimmune tissues. A more granular analysis of tissue-type gene expression reveals a gene transcription pattern supporting this notion. When the gene expression values of each tissue are contrasted pairwise across species (20%-fold change, FDR <0.1), individual nonimmune tissues minimally diverge in expression from immune and nonimmune tissues (Figure 3, Supplementary Table S3). The exceptions to this pattern were mainly nonimmune tissue contrasts against whole blood, an immune compartment that maintains a very high concentration and diversity of immune cell types. Those contrasts produced a low to moderate number of DE genes (kidney-whole blood, 559; brain-whole blood, 964; liver-whole blood, 639 genes). Low to moderate divergence was also noted for whole blood contrasted against colon (492 genes) and liver contrasted against brain (771 genes). When the DE genes of the whole blood contrasts were examined for gene enrichment across a broad range of function (GO biological processes), phenotype/disease (Jensen Disease and dbGaP databases), and cell (Human Gene Atlas) ontogenies via the program Enrichr, a distinct pattern of enrichment for a mix of immune and nonimmune traits emerges (Kuleshov et al. 2016). All nonimmune tissue-whole blood contrasts produced DE genes enriched for the phenotypes of body height (mainly growth factors used throughout life) and potentially antimicrobial cholesterol type, along with the immune biological process of neutrophil degranulation and the disease trait carcinoma (Supplementary Table S5; Chen et al. 2013; Kuleshov et al. 2016). Contrasts with the brain or kidney and thymus produced strong enrichments for T-cell activity,



highlighting the mammalian thymus specialization in T-cell development. A mixed pattern was noted for the brain-liver contrast, rather than an immune versus nonimmune tissue contrast: DE genes tended to be enriched for categories associated with immunity (carcinoma) as well as more general cell function such as synaptogenesis and signal peptide processing. Interestingly, when the remaining 25 tissue contrasts that exhibited DE genes were examined for DE gene enrichment, 64% (16/25) were enriched for carcinomas, suggesting that the regulation of genes associated with cancer has evolved differently in different tissues, and across human and nonhuman primates; both notions have been explored experimentally (Babbitt et al. 2017; Pizzollo et al. 2018). With the exception of the colon-whole blood contrast, immune tissues exhibited very minor if any divergent gene expression compared to other immune tissues.

FIGURE 3. Number of differentially expressed (DE) genes between tissues. DE genes from pairwise contrasts between tissue types across primates (edgeR) were assessed as genes with a >20%-fold change (fc) and a Benjamini-Hochberg false discovery rate (fdr) of <0.1.

Gene Expression across Immune and Nonimmune Tissues Becomes More Promiscuous over Evolutionary Time, and Is Most Promiscuous in the Hominoid Lineage

Though immune system genes are promiscuous in their expression across tissues in primates, there is ample evidence that primate immune systems have diverged in genomic responses to infection and cell physiology (Advani et al. 2016; Barreiro et al. 2010; Brawand et al. 2011; Brinkworth et al. 2012; Pizzollo et al. 2018). This suggests that, along with the multifunctional nature of immune system genes,




FIGURE 4. Density plot of specificity to either immune or nonimmune tissues over evolutionary time. In this analysis, the input values are the log fold change in immune versus nonimmune tissues. There is one distribution for each species, allowing for species-to-species contrast. The distributions represent specificity values (0 = even expression in all tissues; 1 = gene expression specific to a given set of tissues).

there is species specificity in immunogenomic activity in primates. Species specificity in immune gene activity tends to be described as novel elements (Chuong et al. 2016; Gombart et al. 2009) and highly divergent gene code or copy number (Kehrer-Sawatzki and Cooper 2008; Perry et al. 2008; Romero et al. 2017). It is also, on occasion, described as genomic responses to immune challenge that are significantly up- or downregulated over or between particular FDRs in one species, but not over or between particular FDRs of a second or multiple species in the study in question (Barreiro et al. 2010; Tian et al. 2018; Valenzuela-Muñoz et al. 2017). From a functional standpoint, evolving nonimmune phenotypes involve multifunctional genes also associated with immunity. Species may evolve unique immune responses through subtle, networked alterations in the promiscuity of genes associated with both the immune system and other physiological systems. These changes cannot be detected using the first two methods mentioned above, and their exploration would benefit from analytical tools that can complement the use of multiple *p*-value thresholds as a means of assessment. One method for addressing the possibility of species-specific gene expression promiscuity is to compare the specificity of gene expression across species representing major clades of primates using

a scoring system that has been used to examine tissue specificity of gene expression across tissues and species, which controls for the magnitude of expression and so makes head-to-head comparisons possible (Babbitt et al. 2017; Haygood et al. 2007, 2010). For each gene and tissue, we computed a specificity score between 0 and 1 representing the specificity (Haygood et al. 2007; Kosiol et al. 2008) of the gene's expression to the tissue (immune vs. nonimmune), which can also be expanded across species pairs (Babbitt et al. 2017). The specificity score is high if the expression is very specific to the contrast across species, and is low even for the gene's tissue of maximal expression if the gene nearly as highly expressed in other tissues.

Specificity of gene expression changes the distribution of genes specific to either immune or nonimmune tissues over evolutionary time. In this analysis, the input values are the log fold change in immune versus nonimmune tissues. There is one distribution for each species, allowing for species-to-species contrast. The distributions represent specificity values (where 0 shows even expression in all tissues and 1 shows gene expression specific to a given set of tissues). There is less contrast-specific expression (immune vs. nonimmune) in the human-chimpanzee comparison, compared to human and *Macaca mulatta*, *Callithrix jacchus*, and *Microcebus murinus* (Figure 4). The number of contrast-specific genes increases with evolutionary time. This suggests there is more promiscuous gene expression across nonimmune and immune leading to the hominoid lineage, compared to the other primate species.

Conversely, the genes that are more evenly expressed across all the tissues can be examined. With these genes (specificity <0.2), GO biological process enrichments in housekeeping categories such as *protein phosphorylation*, *spliceosomal complex assembly*, and *histone phosphorylation* are apparent—as might be expected, categories essential to cellular function and not necessary to specific tissue biological processes.

There Are Correlations with Positively Selected Genes and Differences in Expression

The high promiscuity of gene expression across immune and nonimmune tissue leads to low divergence in gene expression between such tissues in primates. That such gene sharing is common

and possibly adaptive suggests that, for gene expression to significantly depart from sharing, such singular gene expression patterns are beneficial. To intersect these patterns with signatures of selection, we examined the overlap between DE and signatures of positive selection in immune versus nonimmune tissues. Signatures of positive selection were defined by dN/dS ratios across species comparisons taken from Ensembl (Zerbino et al. 2018). In both sets of tissues we see a weakly positive, but significant, correlation when looking at human-chimpanzee dN/dS ratios. For the immune tissues there is a weak but significant correlation (permutations p -value < 0.0001 , $\rho = 0.0967$, $n = 147$, 20,000 simulations); a similar pattern is seen in nonimmune tissues (permutation p -value $= 3 \times 10^{-4}$, $\rho = 0.0996$, $n = 84$, 20,000 simulations). Enrichment analyses on protein-coding regions showing evidence ($dN/dS > 1$, $n = 430$ orthologs) of accelerated change show enrichments in diffuse immune and transport categories, such as *cytokine-cytokine receptor interaction*. In contrast, enrichments using the human-macaque comparison ($dN/dS > 1$, $n = 81$ orthologs) show a stronger enrichment for immune response categories and their roles in human disease (Supplementary Table S6). This suggests that these differences are more readily apparent at slightly longer evolutionary distances and that those rates of change may inform how global adaptive immune changes have impacts over evolutionary time scales.

Discussion

The degree to which multiple traits are affected by pleiotropy has come into focus recently due to more precise measurements of both genotypic and phenotypic changes (Cannell et al. 2015; De et al. 2014; Li et al. 2017). It has become progressively apparent that pleiotropy is modular and somewhat constrained, allowing for evolvability (reviewed in Wagner and Zhang 2011). The extent of pleiotropic interactions between tissue functions, such as immune/nonimmune compartments, has not been explored in the study of human evolutionary biology. In this review of immune promiscuity, we offer an analysis to explore how tightly associated these different functions are across tissues and evolutionary time.

The nature of the underlying gene networks, however, is still unclear. There can be tissue-specific effects for genes, even genes that are widely expressed (Bossi and Lehner 2009). The degree of pleiotropy for most genes has been estimated to be rather small (reviewed in Wagner and Zhang 2011). However, such assessments suffer from a lack of information about gene expression throughout an organism's development. No ontological series of gene expression is available for all of these primate species and tissues. Acquisition of such data could enhance our understanding of the interplay of these genes over life histories, as many of the cross-referenced genes might be turned on and off during development. A better understanding of how cross-referenced genes are transcribed and silenced over the life course can provide important information on evolutionary trade-offs between the immune and other physiological systems. Importantly, such data can reveal how experiences in early life that impact immune gene expression can alter physiological function of other body systems and health later in life.

As we have shown here, the immune system is very promiscuous, and in primates this increases over the evolution of the order. We also see evidence of accelerated changes between immune and nonimmune gene expression in the hominoid lineage. Immune and nonimmune tissues show low to moderate divergence in expression, yet the functional enrichment of those genes is for a mix of immune and nonimmune characteristics, suggesting that this interplay between systems is driving a host of phenotypes to diverge. When they do diverge, there is strong enrichment for disease susceptibility genes underlying a common human disease: cancer.

As for possible adaptive consequences, we also see that the DE genes between immune and nonimmune tissues under positive selection in hominoids appear to be enriched mainly for immune function. This suggests that immune functions are shaping other phenotypes. An exciting prospect of increasingly accessible gene editing technology is that very controlled assessments of the impact of immune divergence on nonimmune phenotypes and the many specific mutations that may be involved can be readily initiated in animal or cell models.

Taken together, the net effect of immune promiscuity may be that pathogens interacting

with the immune system exert direct selective pressure on genes responsible for both immune and nonimmune phenotypes. In our view, this notion necessitates that future studies examining the evolution of human gene expression and physiology explore what the term “immune” really means. If the immune system is so promiscuous as to be cross-referenced into such diverse functions as brain development, neurological function, placental implantation, spermatogenesis, and bone development, is it fair to describe a cytokine, for example, that is engaged in such pathways as “immune” alone? Though study areas concerned with evolutionary trade-offs and life history frequently consider the energetic requirements of immunity versus other body systems, as a whole human evolutionary biology and biological anthropology have not fully incorporated a view of the immune system as a consideration for the evolution of other physiological systems. If we are not considering the role genes cross-referenced into the immune system may play in the development and evolution of any other system, the view of human and eukaryotic biology, broadly, is incomplete. We will also have a missed opportunity to reiterate the connectedness of human evolution to the extant human condition and to leverage studies concerned with human evolution to better inform fields associated with human health and immunity.

This is especially true in biological fields that readily interface with clinical research, which tend to transact in the language of immunity even though the physiological compartment of focus is ostensibly not the immune system. It is critical that human evolution studies continue to develop in this manner, as key characteristics of the human species appear to be strongly tied to the immune system. Human stature, for example, is connected to immune function, with body height associated B cell responses to infection, and pro-inflammatory cytokines slowing long bone development at the growth plate (Krams et al. 2014; MacRae et al. 2006; Sederquist et al. 2014). Human neurological function and behavior appear inextricably tied to immune activity (Piccinni et al. 1998; Réaux-Le Goazigo et al. 2013; Sheridan and Murphy 2013), and evolutionary interactions with pathogens are hypothesized to affect current cytokine expression in mental health conditions such as depression and anxiety disorders (e.g., PATHOS-D hypothesis;

Raison and Miller 2013). The process of deep inter-digitation of the placenta, a factor that influences maternal mortality and is characteristic of humans and other catarrhines, appears to be influenced by immune gene expression (Benirschke and Kaufmann 2006; Capellini 2012; Crosley et al. 2013; Pudney et al. 2016; Yang et al. 2017). We posit that the genetic promiscuity of the immune system is such that its overlap with other body system makes it, at times, inseparable from other body functions. When considering what defines immune versus nonimmune functions, it is also important that studies consider seemingly nonimmune body components can be integral to the act of eliminating invading foreign matter or dying and cancerous cells. That many of our most famous examples of immune adaptation (i.e., HBS sickle cell alleles, thalassemias) involve the canonical immune tissues and cells very indirectly suggests that maybe the descriptor “immune” can be very broadly applied to any genetic or physiological alterations that can impede invading microorganisms or cancerous and dying cells, regardless of expressing tissue or component (Brinkworth 2017). Minimally, the promiscuity of “immune” genes and the immune activities of nonimmune tissues support the assertive consideration of the immune system and the function of immunity in the evolutionary modeling of all other body systems.

Materials and Methods

Data Sources and Tissue Designations

Data sources for expression analysis are listed in Supplementary Tables S1 and S2 and are otherwise noted in text. For the purposed of our analyses, we designate kidneys, liver and brain as “nonimmune” because they do not serve as major sites of leukocyte development and education, compared to colon, lymph nodes, spleen, thymus, and whole blood/marrow, which are key in the development and maintenance of cellular immunity.

RNA-Seq Mapping and Differential Expression (DE) Analyses

RNA-Seq data were downloaded from the NHPRT (Peng et al. 2015), with human RNA-Seq sequences for the same tissues obtained from the Illumina Body Map2 project (Derrien et al. 2012) stored by

ENCODE and three other studies of human tissues (Clark et al. 2015; Lin et al. 2014; Linsley et al. 2014) archived in the GEO database. RNA isolation and library preparation for the NHPRT data were completed by the same groups across species, and sequencing was completed on an Illumina HiSeq 2000. The human data were collected across projects and sequenced on the HiSeq 2000, with the exception of whole blood, which was sequenced on an Illumina HiScanSQ. PCA analysis was completed on the complete data set, and no batch effects were noted for human (see Figure 1, Supplementary Figures S1 and S2). FASTQ sequences were mapped to the species-specific genome (hg38, panTro4, calJac3.2.1, rheMac8, and micMurl) using the STAR aligner (Dobin et al. 2013). Counts per orthologous genes for all tissues were generated using HTSeq (Anders et al. 2015). Orthology was assessed using Ensembl Genes 86 (Zerbino et al. 2018). Orthologous gene expression across the primate order were assessed in this study, as opposed to novel genes in a lineage, as this is a more conservative approach to understanding human immunity. Orthologs were identified via BiomaRt in the R environment (biomaRt; Durinck et al. 2009; Zerbino et al. 2018). Gene expression was normalized using edgeR (Lun et al. 2016; Robinson et al. 2010), and differential expression (DE) was assessed as minimum 20%-fold change in expression at a false discovery rate (FDR) of 10% or less.

Signatures of Positive Selection in Protein-Coding Regions

dN/dS values from human, chimpanzee, and rhesus macaque were obtained from Ensembl using biomaRt (Durinck et al. 2009), with correlations performed in R, where $1 - dN/dS$ values were compared with p -values of DE. Enrichment analyses were performed using Enrichr (Chen et al. 2013; Kuleshov et al. 2016) and GOrilla/REVIGO (Eden et al. 2009).

Specificity Scores

Specificity scores were calculated as in Haygood et al. (2007) and Kosiol et al. (2008), and species-specific shifts in specificity as in (Babbitt et al. 2017). Briefly, the input values are the log fold change in immune versus nonimmune tissues. There is one distribution for each species, allowing for species-to-species contrast. Here, 0 = even expression in

tissues and 1 = gene expression specific to a given set of tissues.

ACKNOWLEDGMENTS

We thank Ripan Malhi for his editorial oversight, and Jeremy Sykes, Trisha Zintel, and our reviewers for their helpful comments. We also acknowledge our funding provided by the National Science Foundation through grants BCS-1750675 (JFB) and BCS-1750377 (CCB).

Received 1 June 2018; accepted for publication 14 May 2019.

LITERATURE CITED

- Adler, M. W., and T. J. Rogers. 2005. Are chemokines the third major system in the brain? *J. Leukoc. Biol.* 78:1,204–1,209.
- Advani, A. S., A. Y. Chen, and C. C. Babbitt. 2016. Human fibroblasts display a differential focal adhesion phenotype relative to chimpanzee. *Evol. Med. Public Health* 2016:110–116.
- Amberger, J. S., C. A. Bocchini, F. Schiettecatte et al. 2015. OMIM.org: Online Mendelian Inheritance in Man (OMIM®), an online catalog of human genes and genetic disorders. *Nucleic Acids Res.* 43:D789–D798.
- Ambrosi, P., J. S. Chahda, H. R. Koslen et al. 2014. Modeling of the dorsal gradient across species reveals interaction between embryo morphology and Toll signaling pathway during evolution. *PLoS Comput. Biol.* 10:1–17.
- Anderle, P., M. Rumbo, F. Sierro et al. 2005. Novel markers of the human follicle-associated epithelium identified by genomic profiling and microdissection. *Gastroenterology* 129:321–327.
- Anders, S., P. T. Pyl, and W. Huber. 2015. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166–169.
- Attwood, M. M., A. Krishnan, M. S. Almen et al. 2017. Highly diversified expansions shaped the evolution of membrane bound proteins in metazoans. *Sci. Rep.* 7:1–15.
- Babbitt, C. C., R. Haygood, W. J. Nielsen et al. 2017. Gene expression and adaptive noncoding changes during human evolution. *BMC Genomics* 18:435–445.
- Barath, P., M. C. Fishbein, J. Cao et al. 1990. Tumor necrosis factor gene expression in human vascular intimal smooth muscle cells detected by in situ hybridization. *Am. J. Pathol.* 137:503–509.
- Barbosa-Morais, N. L., M. Irimia, Q. Pan et al. 2012. The evolutionary landscape of alternative splicing in vertebrate species. *Science* 338:1,587–1,593.

- Barreiro, L. B., G. Laval, H. Quach et al. 2008. Natural selection has driven population differentiation in modern humans. *Nat. Genet.* 40:340–345.
- Barreiro, L. B., J. C. Marioni, R. Blekhman et al. 2010. Functional comparison of innate immune signaling pathways in primates. *PLoS Genet.* 6:1–13.
- Bauer, S., B. J. Kerr, and P. H. Patterson. 2007. The neuropoietic cytokine family in development, plasticity, disease and injury. *Nat. Rev. Neurosci.* 8:221–232.
- Beattie, E. C., D. Stellwagen, W. Morishita et al. 2002. Control of synaptic strength by glial TNF α . *Science* 295:2,282–2,285.
- Bednarski, T., A. Olichwier, A. Opasinska et al. 2016. Stearoyl-CoA desaturase 1 deficiency reduces lipid accumulation in the heart by activating lipolysis independently of peroxisome proliferator-activated receptor α . *Biochim. Biophys. Acta.* 1861:2,029–2,037.
- Benirschke, K., and P. Kaufmann. 2006. *Pathology of the Human Placenta*. New York: Springer.
- Ben Menachem-Zidon, O., I. Goshen, T. Kreisel et al. 2008. Intrahippocampal transplantation of transgenic neural precursor cells overexpressing interleukin-1 receptor antagonist blocks chronic isolation-induced impairment in memory and neurogenesis. *Neuropsychopharmacology* 33:2,251–2,262.
- Bliss, T. V., and G. L. Collingridge. 1993. A synaptic model of memory: Long-term potentiation in the hippocampus. *Nature* 361:31–39.
- Boehm, T., I. Hess, and J. B. Swann. 2012a. Evolution of lymphoid tissues. *Trends Immunol.* 33:315–321.
- Boehm, T., N. Iwanami, and I. Hess. 2012b. Evolution of the immune system in the lower vertebrates. *Annu. Rev. Genomics Hum. Genet.* 13:127–149.
- Bossi, A., and B. Lehner. 2009. Tissue specificity and the human protein interaction network. *Mol. Syst. Biol.* 5:1–7.
- Boyle, E. A., Y. I. Li, and J. K. Pritchard. 2017. An expanded view of complex traits: From polygenic to omnigenic. *Cell* 169:1,177–1,186.
- Brawand, D., M. Soumillon, A. Necsulea et al. 2011. The evolution of gene expression levels in mammalian organs. *Nature* 478:343–348.
- Brennan, J. J., J. L. Messerschmidt, L. M. Williams et al. 2017. Sea anemone model has a single Toll-like receptor that can function in pathogen detection, NF- κ B signal transduction, and development. *Proc. Natl. Acad. Sci. U. S. A.* 114:E10122–E10131.
- Breschi, A., S. Djebali, J. Gillis et al. 2016. Gene-specific patterns of expression variation across organs and species. *Genome Biol.* 17:1–13.
- Brinkworth, J. F. 2017. Infectious disease and the diversification of the human genome. *Hum. Biol.* 89:47–65.
- Brinkworth, J. F., and L. B. Barreiro. 2014. The contribution of natural selection to present-day susceptibility to chronic inflammatory and autoimmune disease. *Curr. Opin. Immunol.* 31:66–78.
- Brinkworth, J. F., E. A. Pechenkina, J. Silver et al. 2012. Innate immune responses to TLR2 and TLR4 agonists differ between baboons, chimpanzees and humans. *J. Med. Primatol.* 41:388–393.
- Brinkworth, J. F., and M. Thorn. 2013. Vertebrate immune system evolution and comparative primate immunity. In *Primates, Pathogens and Evolution*, J. F. Brinkworth and K. Pechenkina, eds. New York: Springer, 17–64.
- Caballero-Campo, P., M. G. Buffone, F. Benencia et al. 2014. A role for the chemokine receptor CCR6 in mammalian sperm motility and chemotaxis. *J. Cell Physiol.* 229:68–78.
- Cannell, I. G., K. A. Merrick, S. Morandell et al. 2015. A pleiotropic RNA-binding protein controls distinct cell cycle checkpoints to drive resistance of p53-defective tumors to chemotherapy. *Cancer Cell* 28:623–637; erratum in 28:831.
- Cannon, J. G., and C. A. Dinarello. 1985. Increased plasma interleukin-1 activity in women after ovulation. *Science* 227:1,247–1,249.
- Capellini, I. 2012. The evolutionary significance of placental interdigitation in mammalian reproduction: Contributions from comparative studies. *Placenta* 33:763–768.
- Casanova, J.-L. 2015. Human genetic basis of interindividual variability in the course of infection. *Proc. Natl. Acad. Sci. U. S. A.* 112:E7118–E7127.
- Chatterjee, P., S. E. Kopriva, V. L. Chiasson et al. 2013. Interleukin-4 deficiency induces mild preeclampsia in mice. *J. Hypertens.* 31:1,414–1,423.
- Chen, E. Y., C. M. Tan, Y. Kou et al. 2013. Enrichr: Interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* 14:1–14.
- Chuong, E. B., N. C. Elde, and C. Feschotte. 2016. Regulatory evolution of innate immunity through co-option of endogenous retroviruses. *Science* 351:1,083–1,087.
- Clark, M. B., T. R. Mercer, G. Bussotti et al. 2015. Quantitative gene profiling of long noncoding RNAs with targeted RNA sequencing. *Nat. Methods* 12:339–342.
- Cortez, M. J. V., J. F. Rabajante, J. M. Tubay et al. 2017. From epigenetic landscape to phenotypic fitness landscape: Evolutionary effect of pathogens on host traits. *Infect. Genet. Evol.* 51:245–254.
- Crosley, E. J., M. G. Elliot, J. K. Christians et al. 2013. Placental invasion, preeclampsia risk and adaptive molecular

- evolution at the origin of the great apes: Evidence from genome-wide analyses. *Placenta* 34:127–132.
- Cua, D. J., J. Sherlock, Y. Chen et al. 2003. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 421:744–748.
- Das, C., V. S. Kumar, S. Gupta et al. 2002. Network of cytokines, integrins and hormones in human trophoblast cells. *J. Reprod. Immunol.* 53:257–268.
- De, I., M. Nikodemova, M. D. Steffen et al. 2014. CSF1 overexpression has pleiotropic effects on microglia in vivo. *Glia* 62:1,955–1,967.
- Derrien, T., R. Johnson, G. Bussotti et al. 2012. The GENCODE v7 catalog of human long noncoding RNAs: Analysis of their gene structure, evolution, and expression. *Genome Res.* 22:1,775–1,789.
- Deverman, B. E., and P. H. Patterson. 2009. Cytokines and CNS development. *Neuron* 64:61–78.
- Diao, R., K. L. Fok, H. Chen et al. 2014. Deficient human beta-defensin 1 underlies male infertility associated with poor sperm motility and genital tract infection. *Sci. Transl. Med.* 6:249a108.
- Dinareello, C. A., J. G. Cannon, S. M. Wolff et al. 1986. Tumor necrosis factor (cachectin) is an endogenous pyrogen and induces production of interleukin 1. *J. Exp. Med.* 163:1,433–1,450.
- Dobin, A., C. A. Davis, F. Schlesinger et al. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.
- Dravid, G., Y. Zhu, J. Scholes et al. 2011. Dysregulated gene expression during hematopoietic differentiation from human embryonic stem cells. *Mol. Ther.* 19:768–781.
- Duan, C. L., Y. Su, C. L. Zhao et al. 2005. The assays of activities and function of TH, AADC, and GCH1 and their potential use in ex vivo gene therapy of PD. *Brain Res. Brain Res. Protoc.* 16:37–43.
- Du Pasquier, L., J. Robert, M. Courtet et al. 2000. B-cell development in the amphibian *Xenopus*. *Immunol. Rev.* 175:201–213.
- Durinck, S., P. T. Spellman, E. Birney et al. 2009. Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* 4:1,184–1,191.
- Dyer, D. P., C. L. Salanga, S. C. Johns et al. 2016. The anti-inflammatory protein TSG-6 regulates chemokine function by inhibiting chemokine/glycosaminoglycan interactions. *J. Biol. Chem.* 291:12,627–12,640.
- Dyer, D. P., J. M. Thomson, A. Hermant et al. 2014. TSG-6 inhibits neutrophil migration via direct interaction with the chemokine CXCL8. *J. Immunol.* 192:2,177–2,185.
- Eden, E., R. Navon, I. Steinfeld et al. 2009. GOrilla: A tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* 10:1–7.
- Fann, M. J., and P. H. Patterson. 1994. Neurotrophic cytokines and activin A differentially regulate the phenotype of cultured sympathetic neurons. *Proc. Natl. Acad. Sci. U. S. A.* 91:43–47.
- Ferrer-Admetlla, A., E. Bosch, M. Sikora et al. 2008. Balancing selection is the main force shaping the evolution of innate immunity genes. *J. Immunol.* 181:1,315–1,322.
- Fumagalli, M., U. Pozzoli, R. Cagliani et al. 2009. Parasites represent a major selective force for interleukin genes and shape the genetic predisposition to autoimmune conditions. *J. Exp. Med.* 206:1,395–1,408.
- Fumagalli, M., M. Sironi, U. Pozzoli et al. 2011. Signatures of environmental genetic adaptation pinpoint pathogens as the main selective pressure through human evolution. *PLoS Genet.* 7:1–14.
- Giovannelli, A., C. Limatola, D. Ragozzino et al. 1998. CXC chemokines interleukin-8 (IL-8) and growth-related gene product alpha (GROalpha) modulate Purkinje neuron activity in mouse cerebellum. *J. Neuroimmunol.* 92:122–132.
- Giulian, D., D. G. Young, J. Woodward et al. 1988. Interleukin-1 is an astroglial growth factor in the developing brain. *J. Neurosci.* 8:709–714.
- Gombart, A. F., T. Saito, and H. P. Koeffler. 2009. Exaptation of an ancient Alu short interspersed element provides a highly conserved vitamin D-mediated innate immune response in humans and primates. *BMC Genomics* 10:1–11.
- Gosselin, R. D., C. Varela, G. Banisadr et al. 2005. Constitutive expression of CCR2 chemokine receptor and inhibition by MCP-1/CCL2 of GABA-induced currents in spinal cord neurones. *J. Neurochem.* 95:1,023–1,034.
- Greenhalgh, P., C. E. Olesen, and L. A. Steiner. 1993. Characterization and expression of recombination activating genes (RAG-1 and RAG-2) in *Xenopus laevis*. *J. Immunol.* 151:3,100–3,110.
- Gudiksen, A., L. Bertholdt, M. B. Vingborg et al. 2017. Muscle interleukin-6 and fasting-induced PDH regulation in mouse skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 312:E204–E214.
- Harper, K., and G. Armelagos. 2010. The changing disease-scape in the third epidemiological transition. *Int. J. Environ. Res. Public Health* 7:675–697.
- Harper, K. N., and G. J. Armelagos. 2013. Genomics, the origins of agriculture, and our changing microbe-scape: Time to revisit some old tales and tell some new ones. *Am. J. Phys. Anthropol.* 152:135–152.
- Hattermann, K., A. Ludwig, V. Gieselmann et al. 2008. The

- chemokine CXCL16 induces migration and invasion of glial precursor cells via its receptor CXCR6. *Mol. Cell Neurosci.* 39:133–141.
- Haygood, R., C. C. Babbitt, O. Fedrigo et al. 2010. Contrasts between adaptive coding and non-coding changes during human evolution. *Proc. Natl. Acad. Sci. U. S. A.* 107:7,853–7,857.
- Haygood, R., O. Fedrigo, B. Hanson et al. 2007. Promoter regions of many neural- and nutrition-related genes have experienced positive selection during human evolution. *Nat. Genet.* 39:1,140–1,144.
- Heier, C., F. P. Radner, T. Moustafa et al. 2015. G0/G1 switch gene 2 regulates cardiac lipolysis. *J. Biol. Chem.* 290:26,141–26,150.
- Hogquist, K. A., M. A. Nett, E. R. Unanue et al. 1991. Interleukin 1 is processed and released during apoptosis. *Proc. Natl. Acad. Sci. U. S. A.* 88:8,485–8,489.
- Hoover, D. M., C. Boulegue, D. Yang et al. 2002. The structure of human macrophage inflammatory protein-3alpha / CCL20: Linking antimicrobial and CC chemokine receptor-6-binding activities with human beta-defensins. *J. Biol. Chem.* 277:37,647–37,654.
- Hromas, R., P. W. Gray, D. Chantry et al. 1997. Cloning and characterization of exodus, a novel beta-chemokine. *Blood* 89:3,315–3,322.
- Ikebuchi, K., G. G. Wong, S. C. Clark et al. 1987. Interleukin 6 enhancement of interleukin 3-dependent proliferation of multipotential hemopoietic progenitors. *Proc. Natl. Acad. Sci. U. S. A.* 84:9,035–9,039.
- Ito, T., W. F. Carson, K. A. Cavassani et al. 2011. CCR6 as a mediator of immunity in the lung and gut. *Exp. Cell Res.* 317:613–619.
- Johnson, G. A., F. W. Bazer, R. C. Burghardt et al. 2009. Conceptus-uterus interactions in pigs: Endometrial gene expression in response to estrogens and interferons from conceptuses. *Soc. Reprod. Fertil. Suppl.* 66:321–332.
- Joyce, I. M., F. L. Pendola, M. O'Brien et al. 2001. Regulation of prostaglandin-endoperoxide synthase 2 messenger ribonucleic acid expression in mouse granulosa cells during ovulation. *Endocrinology* 142:3,187–3,197.
- Karlsson, E. K., D. P. Kwiatkowski, and P. C. Sabeti. 2014. Natural selection and infectious disease in human populations. *Nat. Rev. Genet.* 15:379–393.
- Kee, H. J., J. T. Koh, S. Y. Yang et al. 2003. A novel murine long-chain acyl-CoA synthetase expressed in brain participates in neuronal cell proliferation. *Biochem. Biophys. Res. Commun.* 305:925–933.
- Kehrer-Sawatzki, H., and D. N. Cooper. 2008. Comparative analysis of copy number variation in primate genomes. *Cytogenet. Genome Res.* 123:288–296.
- Kitaya, K., T. Yasuo, T. Yamaguchi et al. 2007. Genes regulated by interferon-gamma in human uterine microvascular endothelial cells. *Int. J. Mol. Med.* 20:689–697.
- Klein, R. S., J. B. Rubin, H. D. Gibson et al. 2001. SDF-1 alpha induces chemotaxis and enhances Sonic hedgehog-induced proliferation of cerebellar granule cells. *Development* 128:1,971–1,981.
- Komine, M., A. Kukita, T. Kukita et al. 2001. Tumor necrosis factor-alpha cooperates with receptor activator of nuclear factor kappaB ligand in generation of osteoclasts in stromal cell-depleted rat bone marrow cell culture. *Bone* 28:474–483.
- Kosiol, C., T. Vinař, R. R. da Fonseca et al. 2008. Patterns of positive selection in six Mammalian genomes. *PLoS Genet.* 4:1–17.
- Krams, I. A., I. Skrinda, S. Kecko et al. 2014. Body height affects the strength of immune response in young men, but not young women. *Sci. Rep.* 4:1–3.
- Kuleshov, M. V., M. R. Jones, A. D. Rouillard et al. 2016. Enrichr: A comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* 44:W90–W97.
- Kwiatkowski, D. P. 2005. How malaria has affected the human genome and what human genetics can teach us about malaria. *Am. J. Hum. Genet.* 77:171–192.
- Li, J. J., F. X. Ma, Y. W. Wang et al. 2017. Knockdown of IL-8 provoked premature senescence of placenta-derived mesenchymal stem cells. *Stem Cells Dev.* 26:912–931.
- Lin, S., Y. Lin, J. R. Nery et al. 2014. Comparison of the transcriptional landscapes between human and mouse tissues. *Proc. Natl. Acad. Sci. U. S. A.* 111:17,224–17,229.
- Linsley, P. S., C. Speake, E. Whalen et al. 2014. Copy number loss of the interferon gene cluster in melanomas is linked to reduced T cell infiltrate and poor patient prognosis. *PLoS One* 9:1–9.
- Lord, P. C., L. M. Wilmoth, S. B. Mizel et al. 1991. Expression of interleukin-1 alpha and beta genes by human blood polymorphonuclear leukocytes. *J. Clin. Invest.* 87:1,312–1,321.
- Lun, A. T., Y. Chen, and G. K. Smyth. 2016. It's DE-licious: A recipe for differential expression analyses of RNA-seq experiments using quasi-likelihood methods in edgeR. *Methods Mol. Biol.* 1418:391–416.
- Luo, X. J., M. Li, L. Huang et al. 2012. The interleukin 3 gene (*IL3*) contributes to human brain volume variation by regulating proliferation and survival of neural progenitors. *PLoS One* 7:1–17.
- MacRae, V. E., C. Farquharson, and S. F. Ahmed. 2006. The restricted potential for recovery of growth plate chondrogenesis and longitudinal bone growth following

- exposure to pro-inflammatory cytokines. *J. Endocrinol.* 189:319–328.
- March, C. J., B. Mosley, A. Larsen et al. 1985. Cloning, sequence and expression of two distinct human interleukin-1 complementary DNAs. *Nature* 315:641–647.
- Marz, P., J. G. Cheng, R. A. Gadiant et al. 1998. Sympathetic neurons can produce and respond to interleukin 6. *Proc. Natl. Acad. Sci. U. S. A.* 95:3,251–3,256.
- Meola, D., Z. Huang, and J. M. Petitto. 2013. Selective neuronal and brain regional expression of IL-2 in IL2P 8-GFP transgenic mice: Relation to sensorimotor gating. *J. Alzheimers Dis. Parkinsonism* 3:1–6.
- Merkin, J., C. Russell, P. Chen et al. 2012. Evolutionary dynamics of gene and isoform regulation in Mammalian tissues. *Science* 338:1,593–1,599.
- Mori, M., A. Bogdan, T. Balassa et al. 2016. The decidua-the maternal bed embracing the embryo-maintains the pregnancy. *Semin. Immunopathol.* 38:635–649.
- Novembre, J., and E. Han. 2012. Human population structure and the adaptive response to pathogen-induced selection pressures. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 367:878–886.
- Obeid, L. M., C. M. Linardic, L. A. Karolak et al. 1993. Programmed cell death induced by ceramide. *Science* 259:1,769–1,771.
- Ochsner, S. A., D. L. Russell, A. J. Day et al. 2003. Decreased expression of tumor necrosis factor- α -stimulated gene 6 in cumulus cells of the cyclooxygenase-2 and EP2 null mice. *Endocrinology* 144:1,008–1,019.
- O'Donnell, J. M., and H. T. Zhang. 2004. Antidepressant effects of inhibitors of cAMP phosphodiesterase (PDE4). *Trends Pharmacol. Sci.* 25:158–163.
- Oh, Y. J., and K. L. O'Malley. 1994. IL-6 increases choline acetyltransferase but not neuropeptide transcripts in sympathetic neurons. *Neuroreport* 5:937–940.
- Ohno, Y., S. Suto, M. Yamanaka et al. 2010. ELOVL1 production of C24 acyl-CoAs is linked to C24 sphingolipid synthesis. *Proc. Natl. Acad. Sci. U. S. A.* 107:18,439–18,444.
- Ohta, H., H. Wada, T. Niwa et al. 2005. Disruption of tumor necrosis factor- α gene diminishes the development of atherosclerosis in ApoE-deficient mice. *Atherosclerosis* 180:11–17.
- Oppmann, B., R. Lesley, B. Blom et al. 2000. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 13:715–725.
- Ota, N., S. C. Hunt, T. Nakajima et al. 1999. Linkage of interleukin 6 locus to human osteopenia by sibling pair analysis. *Hum. Genet.* 105:253–257.
- Ota, N., T. Nakajima, I. Nakazawa et al. 2001. A nucleotide variant in the promoter region of the interleukin-6 gene associated with decreased bone mineral density. *J. Hum. Genet.* 46:267–272.
- Pavlicev, M., and G. P. Wagner. 2012. A model of developmental evolution: Selection, pleiotropy and compensation. *Trends Ecol. Evol.* 27:316–322.
- Peng, X., J. Thierry-Mieg, D. Thierry-Mieg et al. 2015. Tissue-specific transcriptome sequencing analysis expands the non-human primate reference transcriptome resource (NHPRTR). *Nucleic Acids Res.* 43:D737–D742.
- Perry, G. H., F. Yang, T. Marques-Bonet et al. 2008. Copy number variation and evolution in humans and chimpanzees. *Genome Res.* 18:1,698–1,710.
- Peter, D., S. L. Jin, M. Conti et al. 2007. Differential expression and function of phosphodiesterase 4 (PDE4) subtypes in human primary CD4+ T cells: Predominant role of PDE4D. *J. Immunol.* 178:4,820–4,831.
- Piccinin, S., S. Di Angelantonio, A. Piccioni et al. 2010. CX3CL1-induced modulation at CA1 synapses reveals multiple mechanisms of EPSC modulation involving adenosine receptor subtypes. *J. Neuroimmunol.* 224:85–92.
- Piccinni, M. P., L. Beloni, C. Livi et al. 1998. Defective production of both leukemia inhibitory factor and type 2 T-helper cytokines by decidual T cells in unexplained recurrent abortions. *Nat. Med.* 4:1,020–1,024.
- Piccinni, M. P., C. Scaletti, A. Vultaggio et al. 2001. Defective production of LIF, M-CSF and Th2-type cytokines by T cells at fetomaternal interface is associated with pregnancy loss. *J. Reprod. Immunol.* 52:35–43.
- Pickel, J. M., W. T. McCormack, C. H. Chen et al. 1993. Differential regulation of V(D)J recombination during development of avian B and T cells. *Int. Immunol.* 5:919–927.
- Pickrell, J. K., G. Coop, J. Novembre et al. 2009. Signals of recent positive selection in a worldwide sample of human populations. *Genome Res.* 19:826–837.
- Pishesha, N., P. Thiru, J. Shi et al. 2014. Transcriptional divergence and conservation of human and mouse erythropoiesis. *Proc. Natl. Acad. Sci. U. S. A.* 111:4,103–4,108.
- Pizzollo, J., W. J. Nielsen, Y. Shibata et al. 2018. Comparative serum challenges show divergent patterns of gene expression and open chromatin in human and chimpanzee. *Genome Biol. Evol.* 10:826–839.
- Pudney, J., X. He, Z. Masheeb et al. 2016. Differential expression of toll-like receptors in the human placenta across early gestation. *Placenta* 46:1–10.
- Raison, C. L., and A. H. Miller. 2013. The evolutionary significance of depression in Pathogen Host Defense (PATHOS-D). *Mol. Psychiatry* 18:15–37.

- Réaux-Le Goazigo, A., J. Van Steenwinckel, W. Rostène et al. 2013. Current status of chemokines in the adult CNS. *Prog. Neurobiol.* 104:67–92.
- Reiss, K., R. Mentlein, J. Sievers et al. 2002. Stromal cell-derived factor 1 is secreted by meningeal cells and acts as chemotactic factor on neuronal stem cells of the cerebellar external granular layer. *Neuroscience* 115:295–305.
- Robinson, M. D., D. J. McCarthy, and G. K. Smyth. 2010. edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26:139–140.
- Romero, V., K. Hosomichi, H. Nakaoka et al. 2017. Structure and evolution of the flaggrin gene repeated region in primates. *BMC Evol. Biol.* 17:1–13.
- Sabatini, M., B. Boyce, T. Aufdemorte et al. 1988. Infusions of recombinant human interleukins 1 alpha and 1 beta cause hypercalcemia in normal mice. *Proc. Natl. Acad. Sci. U. S. A.* 85:5,235–5,239.
- Sarkar, O., P. P. Mathur, C. Y. Cheng et al. 2008. Interleukin 1 alpha (IL1A) is a novel regulator of the blood-testis barrier in the rat. *Biol. Reprod.* 78:445–454.
- Schwarz, A., S. Ständer, M. Berneburg et al. 2002. Interleukin-12 suppresses ultraviolet radiation-induced apoptosis by inducing DNA repair. *Nat. Cell Biol.* 4:26–31.
- Sederquist, B., P. Fernandez-Vojvodich, F. Zaman et al. 2014. Recent research on the growth plate: Impact of inflammatory cytokines on longitudinal bone growth. *J. Mol. Endocrinol.* 53:T35–T44.
- Selmaj, K., C. S. Raine, M. Farooq et al. 1991. Cytokine cytotoxicity against oligodendrocytes. Apoptosis induced by lymphotoxin. *J. Immunol.* 147:1,522–1,529.
- Sferruzzi-Perri, A. N., A. M. Macpherson, C. T. Roberts et al. 2009. Csf2 null mutation alters placental gene expression and trophoblast glycogen cell and giant cell abundance in mice. *Biol. Reprod.* 81:207–221.
- Sheridan, G. K., and K. J. Murphy. 2013. Neuron-glia crosstalk in health and disease: Fractalkine and CX₃CR1 take centre stage. *Open Biol.* 3:1–14.
- Stelzer, G., N. Rosen, I. Plaschkes et al. 2016. The GeneCards suite: From gene data mining to disease genome sequence analyses. *Curr. Protoc. Bioinformatics* 54:1.30.1–1.30.33.
- Supek, F., M. Bosnjak, N. Skunca et al. 2011. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* 6:1–9.
- Takahashi, Y., M. Dominici, J. Swift et al. 2006. Trophoblast stem cells rescue placental defect in SOCS3-deficient mice. *J. Biol. Chem.* 281:11,444–11,445.
- Takizawa, H., K. Eto, A. Yoshikawa et al. 2008. Growth and maturation of megakaryocytes is regulated by Lnk/Sh2b3 adaptor protein through crosstalk between cytokine- and integrin-mediated signals. *Exp. Hematol.* 36:897–906.
- Tian, L., S. Shi, F. Nasir et al. 2018. Comparative analysis of the root transcriptomes of cultivated and wild rice varieties in response to *Magnaporthe oryzae* infection revealed both common and species-specific pathogen responses. *Rice* 11:1–15.
- Tsai, H. H., E. Frost, V. To et al. 2002. The chemokine receptor CXCR2 controls positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. *Cell* 110:373–383.
- Valenzuela-Muñoz, V., S. Boltaña, and C. Gallardo-Escárate. 2017. Uncovering iron regulation with species-specific transcriptome patterns in Atlantic and coho salmon during a *Caligus rogercresseyi* infestation. *J. Fish Dis.* 40:1,169–1,184.
- Vallender, E. J., and B. T. Lahn. 2004. Positive selection on the human genome. *Hum. Mol. Genet.* 13:R245–R254.
- Wagner, G. P., and J. Zhang. 2011. The pleiotropic structure of the genotype-phenotype map: The evolvability of complex organisms. *Nat. Rev. Genet.* 12:204–213.
- Wang, S., C. Liu, A. B. Wilson et al. 2017. Pathogen richness and abundance predict patterns of adaptive major histocompatibility complex variation in insular amphibians. *Mol. Ecol.* 26:4,671–4,685.
- Wei, L. H., M. L. Kuo, C. A. Chen et al. 2003. Interleukin-6 promotes cervical tumor growth by VEGF-dependent angiogenesis via a STAT3 pathway. *Oncogene* 22:1,517–1,527.
- Welch, C., M. K. Santra, W. El-Asaad et al. 2009. Identification of a protein, G0S2, that lacks Bcl-2 homology domains and interacts with and antagonizes Bcl-2. *Cancer Res.* 69:6,782–6,789.
- Werner-Felmayer, G., E. R. Werner, D. Fuchs et al. 1993. Pteridine biosynthesis in human endothelial cells: Impact on nitric oxide-mediated formation of cyclic GMP. *J. Biol. Chem.* 268:1,842–1,846.
- Wu, F., W. Zhang, L. Li et al. 2011. Inhibitory effects of honokiol on lipopolysaccharide-induced cellular responses and signaling events in human renal mesangial cells. *Eur. J. Pharmacol.* 654:117–121; erratum in 658:284–285.
- Wueest, S., F. Item, C. N. Boyle et al. 2014. Interleukin-6 contributes to early fasting-induced free fatty acid mobilization in mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 306:R861–R867.
- Yang, S. W., E. H. Cho, S. Y. Choi et al. 2017. DC-SIGN expression in Hofbauer cells may play an important role in


immune tolerance in fetal chorionic villi during the development of preeclampsia. *J. Reprod. Immunol.* 124:30–37.

Zerbino, D. R., P. Achuthan, W. Akanni et al. 2018. Ensembl 2018. *Nucleic Acids Res.* 46:D754–D761.


Zheng, S. G., J. Wang, and D. A. Horwitz. 2008. Cutting edge: Foxp3+CD4+CD25+ regulatory T cells induced by IL-2 and TGF-beta are resistant to Th17 conversion by IL-6. *J. Immunol.* 180:7,112–7,116.

Zhu, N. L., C. Li, H. H. Huang et al. 2007. TNF-alpha represses transcription of human Bone Morphogenetic Protein-4 in lung epithelial cells. *Gene* 393:70–80.

Zilberstein, A., R. Ruggieri, J. H. Korn et al. 1986. Structure and expression of cDNA and genes for human interferon-beta-2, a distinct species inducible by growth-stimulatory cytokines. *EMBO J.* 5:2,529–2,537.



SUPPLEMENTARY FIGURE S1. Principal component analysis of normalized expression values of tissues represented by each primate (kidney, brain, liver, spleen).



SUPPLEMENTARY FIGURE S2. Principal component analysis of normalized expression values of all tissues in this study (brain, kidney, liver, colon, lymph node, whole blood, spleen, thymus, bone marrow).

Supplementary Table S1. Sequence Data (FASTQ) Used in the Analysis

Species	Whole Blood	Spleen	Thymus	Frontal Cortex (Brain)
<i>Homo sapiens</i>	Linsey et al., 2014 http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM1479433	Lin et al., 2014 https://www.encodeproject.org/experiments/ENCSR448VSW/	Clark et al., 2015 http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM1505612	E-MTAB-513-HCT20160 (PRJEB2445) SAMEA962344 ERS025085 ERX011186 ERR030890) https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-513/ ★
<i>Pan troglodytes</i>	Chimpanzee_Wholeblood_HCT22009_S3_L002_R2_001.fastq.gz ★	Chimpanzee_Spleen_364_D2GJ2ACXX-5-ID25_2_sequence.txt.bz2 ★	Chimpanzee_Thymus_363_D2E8TACXX-8-ID27_2_sequence.txt.bz2 ★	Chimpanzee_BrainFrontalCortex_352_D2F1YACXX-2-ID10_2_sequence.txt.bz2 ★
<i>Papio anubis</i>	Baboon_Wholeblood_HCT22016_S10_L002_R2_001.fastq.gz ★	Baboon_Spleen_85_C238CACXX-2-ID01_2_sequence.txt.bz2 ★	Baboon_Thymus_86_C2Y0UACXX-6-ID03_2_sequence.txt.bz2 ★	Baboon_BrainFrontalCortex_72_D23L3ACXX-4-ID05_2_sequence.txt.bz2 ★
<i>Macaca mulatta</i>	RhesusmacaqueIndian_Wholeblood_HCT22013_S7_L001_R1_001.fastq ★	RhesusmacaqueIndian_Spleen_325_R1.fastq.gz ★	RhesusmacaqueIndian_Thymus_326_R1.fastq.gz ★	RhesusmacaqueIndian_Brainfrontalcortex_315_R1.fastq.gz ★
<i>Callithrix jacchus</i>	(bone marrow) Marmoset_BoneMarrow_250_C2B3BACXX-7-ID18_2_sequence.txt.bz2 ★	Marmoset_Spleen_246_C2CB0ACXX-6-ID23_1_sequence.txt ★	N/A	Marmoset_BrainLeftHemisphere_253_C2CG6ACXX-6-ID19_1_sequence.txt ★
<i>Microcebus murinus</i>	N/A	MouseLemur_Spleen_186_H0V2UADXX-1-ID16_1_sequence.txt ★	N/A	MouseLemur_BrainFrontalCortex_173_C2GGMACXX-7-ID04_1_sequence.txt ★

All data RNA-seq, sequenced on Illumina HiSeq 2000, except human whole blood which was sequenced on Illumina HiScanSQ.
 ★ Peng et al., 2015, downloaded from: <http://nhprtr.org>. ★ Illumina Body Map ● Encode sourced

Supplementary Table S2. Consistently Expressed Immune Genes (counts per million normalized value of ≥ 10) in Primate Nonimmune Tissues

NI_genes_immune_process and NI_genes_immune_activation = normalized genes expressed in nonimmune tissues that overlap with the Gene Ontology biological process categories “immune process” and “immune activation,” respectively.

Go to the following link for Supplementary Table S2: https://digitalcommons.wayne.edu/cgi/viewcontent.cgi?filename=1&article=1142&context=humbiol_preprints&type=additional

Supplementary Table S3. Differentially Expressed Genes

Contrasts were completed using edgeR. Full results are presented here. Significantly differentially expressed (DE) genes are defined here as 20%-fold change and FDR < 0.1. Tabs are labeled with the contrast. The second tissue/group is contrasted with the first. I_NI_contrast = immune-nonimmune contrast. All other tabs are pairwise tissue contrasts,

with the second named tissue contrasted against the first named tissue (e.g., Brain_liver_contrast is liver gene expression for all species contrasted with brain gene expression for all species).

Go to the following link for Supplementary Table S3: https://digitalcommons.wayne.edu/cgi/viewcontent.cgi?filename=2&article=1142&context=humbiol_preprints&type=additional

Supplementary Table S4. Results from Enrichment Analysis of Significantly Differentially Expressed (DE) Genes Generated by Immune versus Nonimmune Tissue Contrast

Enrichment analysis was completed using the program GOrilla and revised via the program REVIGO (Eden et al. 2009). “Significantly DE genes” is defined as genes with differential expression of 20%-fold change at an FDR < 0.1. Categories were revised using an allowed similarity of 0.7, SimRel algorithm, FDR < 0.001. Enrichment results for the Gene Ontology biological process (GO_Biological_Process) and revised categories (REVIGO_of_GO_BP) are listed by page.

Temporal Lobe (Brain)	Kidney	Liver	Colon	Lymph Node
N/A	Encode ENCLB3471NC-ENCFF640PYL https://www.encodeproject.org/experiments/ENCSR071ZMQ/ ●	Encode ENCLB828MEI-ERR030895 (PRJEB2445) SAMEA962335 ERS025096 ERX011211 ERR030895) ●	E-MTAB-513-HCT20162 run ERR030882 (PRJEB2445) SAMEA962341 ERS025089 ERX011192 ERR030892) https://www.ebi.ac.uk/ena/data/view/ERR030892 ✚	E-MTAB-513-HCT20146 -ERR030878 (run PRJEB2445) SAMEA962345 ERS025086 ERX011193 ERR030878) ✚
N/A	Chimpanzee_Kidney_358_C2MTPACXX-7-ID23_2_sequence.txt.bz2 ★	Chimpanzee_Liver_355_D2GJ2ACXX-5-ID02_2_sequence.txt.bz2 ★	Chimpanzee_Colon_367_D2EKWACXX-8-ID21_2_sequence.txt.bz2 ★	Chimpanzee_Lymphnode_366_D2E8TACXX-4-ID05_2_sequence.txt.bz2 ★
Baboon_BrainTemporallobe_75_D2FVJACXX-7-ID07_1_sequence.txt.bz2 ★	Baboon_Kidney_78_C23H9ACXX-6-ID14_1_sequence.txt.bz2 ★	Baboon_Liver_79_C238CACXX-2-ID15_2_sequence.txt.bz2 ★	Baboon_Colon_76_C23H9ACXX-6-ID12_1_sequence.txt.bz2 ★	Baboon_Lymphnode_81_C238CACXX-2-ID18_2_sequence.txt.bz2 ★
N/A	RhesusmacaqueIndian_Kidney_319_R1.fastq.gz ★	RhesusmacaqueIndian_Liver_320_R1.fastq.gz ★	N/A	N/A
N/A	Marmoset_Kidney_240_D2CC5ACXX-2-ID10_1_sequence.txt ★	Marmoset_Liver_237_D2CC5ACXX-2-ID11_1_sequence.txt ★	N/A	Marmoset_Lymphnode_248_C2CD5ACXX-3-ID21_1_sequence.txt ★
MouseLemur_BrainTemporallobe_176_D2FVCACXX-7-ID05_1_sequence.txt ★	MouseLemur_Kidney_180_H0V2UADXX-1-ID12_1_sequence.txt ★	MouseLemur_Liver_181_C298AACXX-6-ID13_1_sequence.txt ★	MicMur_colon ★	N/A

Go to the following link for Supplementary Table S4: https://digitalcommons.wayne.edu/cgi/viewcontent.cgi?filename=3&article=1142&context=humbiol_preprints&type=additional

Supplementary Table S5. Results from Enrichment Analysis of Significantly Differentially Expressed (DE) Genes Generated by Pairwise Tissue Contrasts

Enrichment analysis was completed using the program Enrichr (Chen et al. 2013; Kuleshov et al. 2016) on 1 May 2018. “Significantly DE genes” is defined as genes with differential expression of 20%-fold change at an FDR <0.1. The following tissue contrasts did not produce either DE genes or significant enrichment results and are therefore not represented in this file: thymus–whole blood, lymph node–thymus, lymph node–spleen, colon–spleen, colon–lymph node. Enrichment results for the Gene Ontology, Jensen Disease, dbGaP, and Human Gene Atlas databases were completed and are stored here in single pages by tissue contrast. Database is noted in the column “Database.”

Go to the following link for Supplementary Table S5: https://digitalcommons.wayne.edu/cgi/viewcontent.cgi?filename=4&article=1142&context=humbiol_preprints&type=additional

Supplementary Table S6. dN/dS Values and Enrichments for Primates, Hominoids, and Humans

dN/dS values were obtained from Ensembl using biomaRt (Durinck et al. 2009), with correlations performed in R, where 1 – dN/dS values were compared with *p*-values of DE. Homo_Mus_dn_ds column = primate calculation; Homo_Macaca_dn_ds column = hominoid calculation; Homo_Pan_dn_ds column = human calculation. Enrichment analyses were performed using Enrichr (Chen et al. 2013; Kuleshov et al. 2016) and GOrilla/REVIGO (Eden et al. 2009).

Go to the following link for Supplementary Table S6: https://digitalcommons.wayne.edu/cgi/viewcontent.cgi?filename=5&article=1142&context=humbiol_preprints&type=additional

Copyright of Human Biology is the property of Wayne State University Press and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.