

Near-Wall Vorticity Dynamics in Channel-Flow Turbulence: A Stochastic Lagrangian Analysis

G. L. Eyink^{1,2,3}, A. Gupta¹, and T. A. Zaki³

¹Department of Applied Mathematics & Statistics, The Johns Hopkins University, Baltimore, MD 21218, USA

²Department of Physics & Astronomy, The Johns Hopkins University, Baltimore, MD 21218, USA

³Department of Mechanical Engineering, The Johns Hopkins University, Baltimore, MD 21218, USA

(Received)

We use an online database of a turbulent channel-flow simulation at $Re_\tau = 1000$ (Graham *et al.* 2016) to determine the origin of vorticity in the near-wall buffer layer. Following an experimental study of Sheng *et al.* (2009), we identify typical “ejection” and “sweep” events in the buffer layer by local minima/maxima of the wall-stress. In contrast to their conjecture, however, we find that vortex-lifting from the wall is not a discrete event requiring ~ 1 viscous time and ~ 10 wall units, but is instead a distributed process over a space-time region at least $1 \sim 2$ orders of magnitude larger in extent. To reach this conclusion, we exploit a rigorous mathematical theory of vorticity dynamics for Navier-Stokes solutions, in terms of stochastic Lagrangian flows and stochastic Cauchy invariants, conserved on average backward in time. This theory yields exact expressions for vorticity inside the flow domain in terms of vorticity at the wall, as transported by viscous diffusion and by nonlinear advection, stretching and rotation. We show that Lagrangian chaos observed in the buffer layer can be reconciled with saturated vorticity magnitude by “virtual reconnection”: although the Eulerian vorticity field in the viscous sublayer has a single sign of spanwise component, opposite signs of Lagrangian vorticity evolve by rotation and cancel by viscous destruction. Our analysis reveals many unifying features of classical fluids and quantum superfluids. We argue that “bundles” of quantized vortices in superfluid turbulence will also exhibit stochastic Lagrangian dynamics and satisfy stochastic conservation laws resulting from particle relabelling symmetry.

Key words: vortex dynamics, turbulent boundary layers, quantum fluids

1. Introduction

Vorticity is well-recognized to play a fundamental role in turbulent flows and its ultimate origin lies quite frequently at solid walls or flow boundaries. It therefore becomes a basic question to understand how vorticity in the interior of the flow evolved from vorticity generated at the wall. Recent mathematical work of Constantin & Iyer (2008, 2011) has provided new exact tools to answer this question in terms of stochastic Lagrangian particle trajectories evolved backward in time. In the previous work of Eyink *et al.* (2020a)—hereafter referred to as paper I—it was shown that the mathematical

representations of Constantin & Iyer (2008, 2011) have a simple fluid-dynamical interpretation in terms of the “vortex-momentum density” associated to a continuous distribution of infinitesimal vortex-rings, which is the basis of the Kuz’min (1983)-Oseledets (1989) formulation of the incompressible Navier-Stokes equation. For smooth ideal Euler solutions, the vortex-momentum density is Lie-transported by the fluid flow as a 1-form and its curl, the vorticity, is transported as a 2-form (Oseledets 1989; Tur & Yanovsky 1993). This Lie-transport leads naturally to the vorticity invariants of Cauchy (1815) for incompressible Euler solutions and to generalized Cauchy invariants for the vortex-momentum density (Tur & Yanovsky 1993; Besse & Frisch 2017). The mathematical theory of Constantin & Iyer (2008, 2011) provides corresponding “stochastic Cauchy invariants” for Navier-Stokes solutions that are conserved on average by the stochastic Lagrangian flow backward in time and these invariants provide an explicit representation of the vortex-momentum density and the vorticity in terms of boundary data. We furthermore discussed in paper I some relations of the Constantin & Iyer (2008, 2011) stochastic Lagrangian representation with the Eulerian theory of Lighthill (1963)-Morton (1984) for vorticity-generation at solid walls and with an exact statistical result of Taylor (1932)-Huggins (1994) for the “vorticity flux tensor”. The latter is an anti-symmetric tensor Σ_{ij} , which represents the flux of the j th-component of the vorticity in the i th coordinate direction and which was observed by Huggins (1994) (following Taylor (1932) for 2D pipe flow) to have a mean value directly proportional to the mean pressure-gradient in the k th direction, with i, j, k all distinct. This relation applies to drag generally for any flows driven by imposed pressure-gradients and/or freestream velocity, such as turbulent shear layers and wakes (Brown & Roshko 2012). The “vorticity source” of Lighthill (1963) and Morton (1984) is defined at the wall by the vector $\sigma_i = \Sigma_{ij}\hat{n}_j$, summed over repeated index j , where $\hat{n}$ is the outward-pointing unit normal vector on the boundary (Lyman 1990; Eyink 2008).

The main result that we exploit in the present work is the relation [paper I, eq.(2.51)]

$$\boldsymbol{\omega}(\mathbf{x}, t) = \mathbb{E}[\tilde{\boldsymbol{\omega}}_s(\mathbf{x}, t)], \quad s < t \quad (1.1)$$

which expresses the vorticity vector at space-time point $(\mathbf{x}, t)$ as an expectation $\mathbb{E}$ of the *stochastic Cauchy invariant* $\tilde{\boldsymbol{\omega}}_s(\mathbf{x}, t)$ [paper I, eq.(2.55)]. This is a random vector that can be evaluated along stochastic Lagrangian particle trajectories satisfying

$$\hat{d}\tilde{\mathbf{A}}_t^s(\mathbf{x}) = \mathbf{u}(\tilde{\mathbf{A}}_t^s(\mathbf{x}), s) ds + \sqrt{2\nu} \hat{d}\tilde{\mathbf{W}}(s), \quad s < t; \quad \tilde{\mathbf{A}}_t^t(\mathbf{x}) = \mathbf{x}, \quad (1.2)$$

which are released at $(\mathbf{x}, t)$ and move backward in time s . Here $\tilde{\mathbf{W}}(s)$ is a random Wiener process that represents diffusion by molecular viscosity. For every realization of the process there is a largest time $s = \tilde{\sigma}_t(\mathbf{x})$ at which the stochastic particle first hits the wall, backward in time. Each particle is stopped at the wall where it first hits and in that particular realization $\tilde{\boldsymbol{\omega}}_s(\mathbf{x}, t) = \tilde{\boldsymbol{\omega}}_{\tilde{\sigma}_t(\mathbf{x})}(\mathbf{x}, t)$ for $s < \tilde{\sigma}_t(\mathbf{x})$, thus remaining fixed at earlier times. If one considers $s \ll t$, then the inequality $s < \tilde{\sigma}_t(\mathbf{x})$ will hold with near certainty and, in that case, the formula (1.1) represents the interior vorticity in terms of wall vorticity which is transported to $(\mathbf{x}, t)$ by advection, diffusion and stretching. A numerical Monte Carlo Lagrangian algorithm was also developed in paper I to calculate realizations of the stochastic Cauchy invariants and their statistics, given an Eulerian space-time solution of the incompressible Navier-Stokes equation.

Here we shall exploit that approach to make a first-of-its-kind numerical study of stochastic Lagrangian dynamics of vorticity in a turbulent channel-flow at high Reynolds number. If we denote Cartesian coordinate directions as streamwise x , wall-normal y , and spanwise z , then statistical homogeneity in x and z and reflection-symmetry in y provide




Figure 1: A typical array of vortex lines pointing spanwise and lifting in an arch over a low-speed streak at a wall-stress minimum during an “ejection” event in the buffer-layer of a turbulent boundary layer. The wide arrow indicates the direction of the mean flow.

simplifications in long-time averages. In particular, the mean flux of spanwise vorticity vertically from the wall becomes independent of wall-normal location as a consequence of conservation, $\partial_y \bar{\Sigma}_{yz} = 0$, and is given by

$$\bar{\Sigma}_{yz} = \overline{v'\omega'_z - w'\omega'_y} - \nu \frac{\partial \bar{\omega}_z}{\partial y} = \frac{\partial \bar{p}}{\partial x} = -\frac{u_*^2}{h} < 0. \quad (1.3)$$

where $f' := f - \bar{f}$ defines fluctuation away from the long-time mean value, where velocity vector $\mathbf{u} = (u, v, w)$, where u_* is the friction velocity, and where h is the channel half-height (Huggins 1994; Eyink 2008). According to eq.(1.3) the constant flux of spanwise vorticity away from the wall is numerically equal to the rate of mean downstream pressure drop, which characterizes turbulent drag and dissipation. In this paper we shall study the Lagrangian dynamics of vorticity in the buffer layer of turbulent channel-flow, which is conventionally taken to be the layer of fluid at heights y^+ over the range $5 < y^+ < 30$ (Tennekes & Lumley 1972), with y^+ in dimensionless wall units (see section 2). We have both pragmatic and fundamental reasons to focus on the buffer layer. Since all interior flow vorticity traces its origin back to a solid wall, the “youngest” vorticity must lie closest to the wall. This makes the near-wall region numerically easiest to study by our approach. There is also motivation to understand the turbulent physics close to the wall, since one can expect that control of vorticity creation and transport at the earliest stage will be most efficient in reducing drag (Koumoutsakos 1999; Zhao *et al.* 2004). On the other hand, nonlinear contributions to the dynamics are subleading (on average) in the viscous sublayer $y^+ < 5$, and nonlinear contributions to mean vorticity transport first become substantial in the buffer layer.

Our study is directly motivated by laboratory experiments of Sheng *et al.* (2009), who investigated buffer-layer structures in a turbulent square-duct channel flow at $Re_\tau = 1470$ using a technique of digital holographic microscopy that yields well-resolved measurements of three-dimensional velocity and velocity-gradient fields. Conditional sampling based on local wall shear-stress maxima and minima revealed two types of structures that generate such extreme stress events. In accord with a common terminology, these may be called “sweeps” and “ejections”, respectively. The latter type of flow event generates arrays of vortex lines with a simple “hairpin” structure that rise in an arch above the location of the wall-stress minimum. See Fig. 1, which represents well the typical

geometry of vortex lines observed by Sheng *et al.* (2009) during an “ejection”. These raised vortex lines with a non-vanishing vertical component of vorticity are the signature closest to the wall of a contribution by nonlinear stretching and rotation to transport of spanwise vorticity upward from the wall. Such an orderly spatial array of lines as illustrated in Fig. 1 invites interpretation in terms of a similarly simple temporal progression, with each line apparently “moving” forward and evolving into its downstream neighbor. Indeed, Sheng *et al.* (2009) on the basis of such spatial arrays of lines (see their Figures 7 & 21) have proposed an “abrupt lifting” of vorticity in just one viscous time or, spatially, in a short distance of 10 wall units, above the location of a local stress-minimum.

As we shall show by detailed Lagrangian analysis exploiting the stochastic Cauchy invariants, the deceptively simple Eulerian picture of vortex-lines in Fig. 1 is in fact the outcome of a hidden, violent conflict between intense nonlinear stretching and rotation of vorticity vector elements on the one hand, and vigorous destruction of vorticity by viscosity on the other. Previous work of Johnson *et al.* (2017) has demonstrated existence of Lagrangian chaos in channel-flow turbulence, which leads to rapid exponential stretching of material line elements. This poses a theoretical puzzle, however, because such stretching should lead to unbounded growth of vorticity but the channel flow nevertheless attains a statistical steady state with a saturated magnitude of vorticity. The obvious mechanism that limits vortex-stretching is viscous destruction (Taylor 1938), but advection and diffusion of same-sign vorticity cannot quench the growth due to stretching. We find that vorticity vector elements in the buffer layer are not only exponentially magnified, but also strongly rotated, so that they often point opposite to the negative spanwise direction pictured in Fig. 1. Cancellation of this oppositely-directed vorticity by viscous diffusion leads to the regular geometry in Fig. 1. Furthermore, we find that the vortex lifting is not an “abrupt” discrete event, but is instead a highly distributed process spread over more than 100 viscous times and 1000 wall units.

These results are obtained by a numerical study using an online computational dataset of channel-flow turbulence at $Re_\tau = 1000$ (Graham *et al.* 2016). The accuracy of this database to study buffer-layer physics has been carefully evaluated and documented in paper I, as briefly summarized in section 2. Our study not only yields new insights into Lagrangian vorticity dynamics of pressure-driven wall-bounded flows, but also reveals as well many common features of classical fluids and quantum superfluids, especially for wall-bounded turbulent flows through pipes and channels. To mention here just a few of these similarities, the Kuz'min (1983)-Oseledets (1989) representation of the classical fluid by the vortex-momentum density is closely related to intuitive pictures of “vortex-tangles” in superfluids as superpositions of small vortex-rings (Campbell 1972; Schwarz 1982; Kuz'min 1999). Perhaps most importantly, Huggins (1994) has emphasized that eq.(1.3) is the exact analogue of the “Josephson-Anderson relation” in quantum fluids; see Josephson (1962); Anderson (1966), and Packard (1998); Varoquaux (2015) for reviews. This relation provides the accepted explanation for effective drag in an otherwise non-dissipative superfluid by cross-stream motion of quantized vortex lines. For further discussions of this analogy, see also Huggins (1970) and Eyink (2008). We argue briefly in the conclusion of the present paper that such similarity should extend to Lagrangian dynamics of vorticity and that motion of “bundles” of quantized vortex lines in turbulent superfluids should be also intrinsically stochastic.

The contents of this paper are outlined as follows. Section 2 summarizes the essentials of our numerical methods, which are described more completely in paper I. Section 3 explains how the two specific flow events were selected for examination (3.1) and then describes both the ejection (3.2) and sweep (3.3) events in detail in conventional Eulerian terms. Our novel Lagrangian analysis is presented in section 4, where we first

Table 1: Simulation Parameters for Turbulent Channel-Flow Dataset

N_x	N_y	N_z	Re_τ	dp/dx	ν	u_*	U_{bulk}	Δx^+	Δz^+	Δt
2048	512	1536	1000	-2.5×10^{-3}	5×10^{-5}	5×10^{-2}	1.00	12.3	6.1	1.3×10^{-3}

choose specific vorticity vectors for quantitative study (4.1) and then determine their dynamical origin at the wall for both the ejection (4.2) and the sweep (4.3). The final section 5 summarizes our results and conclusions, especially on common features of wall-bounded turbulence in classical fluids and quantum superfluids. Additional material that complements the discussion in the main text is provided as Supplemental Materials (SM).

2. Numerical Methods

We review here for completeness some necessary information about our computational approach from paper I, which should be consulted for full details. The Johns Hopkins Turbulence Databases (JHTDB) channel-flow dataset (Graham *et al.* 2016) is exploited for the empirical study in this paper. This data was generated from a Navier-Stokes simulation in a channel using a pseudospectral method in the plane parallel to the walls and a seventh-order B-splines collocation method in the wall-normal direction (Lee *et al.* 2013). For numerical solution, the Navier-Stokes equations were formulated in wall-normal velocity-vorticity form (Kim *et al.* 1987). Pressure was computed by solving the pressure Poisson equation only when writing to disk, which was every five time steps for 4000 snapshots, enough for about one domain flow-through time. The simulation domain $[0, 8\pi h] \times [-h, h] \times [0, 3\pi h]$, $h = 1$, was discretized using a spatial grid of $2048 \times 512 \times 1536$ points. Time advancement was made with a third-order low-storage Runge-Kutta method and dealiasing was performed with 2/3 truncation (Orszag 1971). A constant pressure head was applied to drive the flow at $Re_\tau = 1000$ ($Re_{bulk} = \frac{2hU_{bulk}}{\nu} = 40\,000$) with bulk velocity near unity. As is common, we shall indicate with a superscript “+” non-dimensionalized quantities in viscous wall units, with velocities scaled by friction velocity u_* and lengths by viscous length $\delta_\nu = \nu/u_* = 10^{-3}$. Also as usual, we define $y^+ = (h \mp y)/\delta_\nu$ near $y = \pm h$. In these units, the first y -grid point in the simulation is located at distance $\Delta y_1^+ = 1.65199 \times 10^{-2}$ from the wall, while in the center of the channel $\Delta y_c^+ = 6.15507$. Other numerical parameters are summarized in Table 1.

In paper I we developed and tested a numerical Monte Carlo Lagrangian algorithm to calculate the stochastic Cauchy invariants and their statistics, by discretizing the stochastic ODE (1.2) with a step-size Δs and by averaging over N independent realizations $\widetilde{\mathbf{W}}^{(n)}(s)$, $n = 1, \dots, N$ of the Wiener process. We showed in that work for the two specific cases from the JHTDB channel-flow dataset studied in the present paper that $\Delta s = 10^{-3}$ and $N = 10^7$ sufficed to give converged results for the mean and variance of the stochastic Cauchy invariant over a time interval $-100 < \delta s^+ < 0$ with $\delta s := s - t$. In particular, it was shown that the mean conservation law (1.1) holds for those two cases to within a few percent over that time interval, which is a quite stringent test of validity of our numerics. The residual errors in the mean conservation can be explained by some near-wall under-resolution of the numerical JHTDB data, indicated by the sizable Δx^+ and Δz^+ values in Table 1, and by errors in the finite-difference approximation of velocity-gradients within the database. To test that hypothesis, we also

spatially coarse-grained the relation (1.1) over n_i grid-spacings in each of the coordinate directions $i = x, y, z$, since such coarse-grained fields from the JHTDB dataset should represent better a coarse-grained Navier-Stokes solution. We verified that such coarse-graining noticeably improves mean conservation, in particular for $(n_x, n_y, n_z) = (4, 0, 4)$.

In sections 3.1 and 4.1 below, we describe the criteria that we used to select the two test cases for study in this work. In particular, we discuss in section 3.1 how a pair of events, an “ejection” and a “sweep”, were identified in the Eulerian dataset, following the work of Sheng *et al.* (2009). In section 4.1 we then describe how we selected a space-time point $(\mathbf{x}, t)$ in each event for comparative study by stochastic Lagrangian analysis. We also show there that coarse-graining the JHTDB fields with $(n_x, n_y, n_z) = (4, 0, 4)$ does not change qualitatively the Eulerian and Lagrangian picture of the two events. These results, together with those presented in paper I, validate both our Monte Carlo numerical method to calculate the stochastic Cauchy invariant and also the adequacy of the JHTDB channel-flow database to resolve the physics of the turbulent buffer layer.

3. Eulerian Vorticity Dynamics in the Buffer Layer

3.1. Identification of Ejections and Sweeps

Following the approach of Sheng *et al.* (2009), we selected events where the viscous shear stress $\tau_{xy} = \nu(\partial u / \partial y)$ at the wall has local minima and local maxima with magnitudes satisfying a threshold condition. For this purpose, we downloaded the stress field at the entire top wall of the channel-flow database at the final archived time $t_f = 25.9935$ and searched for local minima and maxima. Note that we used the data at the top wall because the bottom-wall data was temporarily unavailable when we began our study; in order to present our results below we always rotate the figures 180° around the streamwise axis, so that the top wall is exchanged with the bottom wall. In SM we provide a plot of the normalized stress field $\tau_{xy}^+ = \tau_{xy} / u_*^2$ over the entire channel wall and a PDF of its values, which range from -2.55 to $+7.54$ and have mean value unity. To find local minima and maxima, we used a fast peak-finder for 2D scalar data (Natan 2013). We found that the points identified by this code for field τ_{xy} were indeed local maxima and for $-\tau_{xy}$ were local minima, but that weaker maxima/minima were often missed. We therefore do not regard the output of this algorithm to be completely reliable to obtain statistics of the local extrema, but it suffices for our purpose of identifying specific local maxima/minima. Nevertheless, we do provide in the SM for the interest of readers the obtained PDF’s of the stress values at the positions both of the local minima and also of the local maxima. The PDF of the local-minimum stress values shows a large peak at $\tau_p^+ \doteq 0.6$, while the PDF of the local-maximum stress values shows a large peak at $\tau_p^+ \doteq 1.8$. Interestingly, the condition that Sheng *et al.* (2009) applied to identify “extreme stress events” was $\tau_{xy}^+ < 0.6$ for local-minima and $\tau_{xy}^+ > 1.8$ for local-maxima, in good agreement with these peaks values. We therefore searched for two typical events of this type, namely, for a local-minimum of stress with value $\tau_{xy}^+ \doteq 0.6$ and for a local-maximum of stress with value $\tau_{xy}^+ \doteq 1.8$. We also looked for such local-extremum points which were relatively isolated from others. After examining several candidates, we selected as representative the two local-extrema with database space-time coordinates given in Table 2. The reader will note that these coordinates correspond, as stated above, to points on the top wall of the channel. The reader should transform results in paper I according to $(\omega_x, \omega_y, \omega_z) \rightarrow (\omega_x, -\omega_y, -\omega_z)$ for consistency with visualizations in the present paper. In particular, mean spanwise vorticity $\bar{\omega}_z$ under this transformation becomes negative.

We shall refer to the local-minimum stress event as an “ejection” and to the local-

Table 2: Coordinates of Local-Minimum and Local-Maximum Wall-Stress Events

	x	y	z	t
local-minimum	21.107576	1.000000	7.565593	25.9935
local-maximum	0.711767	1.000000	0.724039	25.9935

maximum stress event as a “sweep.” This terminology is in agreement with the classification of Sheng *et al.* (2009) in their Table 2, but it differs somewhat from the most common characterization of such structures based on quadrant analysis in the (u', v') velocity plane, with connected regions of Q2 fluctuations designated as “ejections” and regions of Q4 fluctuations as “sweeps” (Jiménez 2013). As we shall see from a detailed Eulerian description of these two selected events in the following sections 3.2-3.3, our use of the terms “ejection” and “sweep” is not unrelated with the traditional quadrant analysis. We have purposely avoided using the term “burst” to describe either of our two events, although “ejections” have sometimes in the past been equated with “bursts”. In more current understanding, however, buffer-layer “bursting” is believed to be associated with quasi-periodic breakdown of unstable coherent structures presumably described by traveling-wave solutions of Navier-Stokes equations (Jiménez (2013), sections IV.A-B; Park *et al.* (2018)). The quasi-period of this bursting is expected to be $\simeq 400$ viscous times $t_\nu = \nu/u_*^2$ with duration $\simeq 100$ viscous times, during which the traveling structure evolves from a low wall-stress to high wall-stress state. E.g. see Jiménez (2013), Figure 10 or Park *et al.* (2018), Figure 6. Therefore “ejections” and “sweeps” in our sense may both be associated with buffer-layer bursting, at different stages in the evolution of the burst. Our interest here is mainly in analyzing the Lagrangian dynamics of vorticity within these two buffer-layer events and not in determining their relation with “bursting.”

In subsections 3.2-3.3 below we first provide a detailed description of these events in Eulerian terms. This does illuminate some connections with “bursting”, but our primary purpose is to describe these two events in terms of standard Eulerian theory of vorticity generation at walls and to compare with prior numerical and experimental results. Among these, we wish to compare our chosen events with those selected in Sheng *et al.* (2009) by identical criteria and to verify that our events have the same characteristic features. In particular, we shall show that our “ejection” event is quite typical of those studied by Sheng *et al.* (2009) and used by them to postulate the “abrupt lifting” of vortex-lines. Here it is appropriate to make a remark on the relative importance of nonlinear dynamics and viscous diffusion for vorticity transport in the buffer layer. As stressed in the Introduction, an “abrupt lifting” event requires nonlinear stretching and rotation of spanwise vorticity in order to create a vertical arch. However, linear viscous diffusion dominates the mean vertical flux of spanwise vorticity not only in the viscous sublayer and buffer layer but even into the log-layer! In fact, the simple relation

$$\overline{v'\omega'_z - w'\omega'_y} = \frac{\partial}{\partial y} (\overline{-u'v'}) \quad (3.1)$$

implies that the nonlinear contribution to spanwise vorticity flux is positive for $y^+ < y_p^+$, the location of the peak Reynolds shear stress $\overline{-u'v'}$ (Klewicki *et al.* 2007; Eyink 2008). In the JHTDB channel-flow dataset at $Re_\tau = 1000$, this peak occurs at about $y_p^+ \doteq 50$ (Graham *et al.* 2016). The net effect of the nonlinear terms for $y^+ < y_p^+$ is thus to transport spanwise vorticity *opposite* to the conserved total flux $\overline{\Sigma}_{yz}$ in eq.(1.3), and viscous




Figure 2: Field of viscous shear stress $\tau_{xy}^+ = (\partial u^+ / \partial y^+)$ at the wall $y^+ = 0$, in wall units so that the average is unity. Black arrows represent the two-dimensional in-wall stress vector τ_W . The asterisk “*” marks the location of the selected stress local-minimum.




Figure 3: (a) Isosurface $\lambda_2^+ = -0.0163$, with magnitude 4 times the local box-average value $\langle |\lambda_2^+| \rangle = 4.09 \times 10^{-3}$. The shear-stress field from Fig. 2 is replotted in the bottom x - z -plane for reference. (b) Field of streamwise vorticity ω_x^+ in plane $x^+ = 85.9$ (transparent in panel (a)). The black arrows represent cross-stream velocity vectors (w, v) .

transport must be even more negative to compensate. We shall see also in the Lagrangian description that viscous diffusion plays an essential role in buffer-layer vorticity transport, even for extreme-stress events where nonlinear terms are enhanced.

3.2. Eulerian Description of Ejection Event

The stress local-minimum that we selected for study is located within a long low-speed streak of the type commonly observed in near-wall turbulence, with typical spanwise separations $\delta z^+ \simeq 100$ between streaks (Jiménez 2013). This environment is illustrated in Fig. 2, which plots the viscous shear-stress $\tau_{xy} = \nu(\partial u / \partial y)$ at the wall, with magnitudes represented by the color (or shade, in greyscale), together with the location of the stress local-minimum as an asterisk “*”. This figure also plots the two-dimensional wall stress field $\tau_W = \nu(\partial u / \partial y, \partial w / \partial y)$ with black arrows. The arrows indicate a near-wall, in-plane

flow which is converging toward the streak. This convergence is consistent with a vertical flow that is upward, away from the wall, at the streak and it agrees with results of Sheng *et al.* (2009) for the conditional average stress-field in the vicinity of such local-minima (see their Fig. 6(f)). More insight into the local flow conditions is provided by Fig. 3, which in panel (a) visualizes the coherent vortices in the vicinity of the local-minimum using isosurfaces of λ_2 , or the second eigenvalue of the $(\nabla_x \mathbf{u})^2$ matrix (Jeong & Hussain 1995). Somewhat different choices of λ_2 -levels and different vortex visualization criteria yield similar results. Clearly observed are two long, equal-strength, streamwise vortices located on each side of the low-speed streak and inclined away from the wall. Measurement of ω_x reveals that these vortices are counter-rotating, generating a lifting flow above the low-speed streak. This is illustrated in Fig. 3(b), which provides a color-plot of ω_x in the transverse y - z -plane through the middle of the visualized box and which also plots as arrows the two-dimensional cross-stream flow vectors (w, v) within that plane. This type of counter-rotating vortex-pair generating a lifting flow between them is a quite common buffer-layer configuration, encountered in about 16% of all of samples in the study of Sheng *et al.* (2009) and in 98% of their realizations satisfying the condition $\tau_{xy}^+ < 0.6$. Our selected event thus appears to be quite typical of such stress-minima and similar features were observed in many other local-minima stress events that we identified in the JHTDB channel-flow dataset satisfying the criteria $\tau_{xy}^+ \simeq 0.6$.

This typicality is confirmed by Fig. 4 which plots vortex-lines crossing the visualized domain in the spanwise direction, initialized at evenly spaced streamwise locations and at initial elevations $y^+ = 2, 4, 6, 8$. These vortex lines are lifted in arches above the low-speed streak, with a typical “hairpin” geometry. The arches are nearly vertical for lines initiated at $y^+ = 2, 4$ and also for $y^+ = 6, 8$ at points well upstream of the stress minimum. For lines initiated at $y^+ = 6$ the arches rise and bend downstream approaching the stress-minimum, while the lines initiated at $y^+ = 8$ near the stress-minimum have instead an “ Ω -vortex” geometry and the uppermost tips are bent back slightly upstream. These arrays of vortex lines are typical of those observed in the vicinity of stress local-minima in the study of Sheng *et al.* (2009), as illustrated in their Figures 4(a), 8(a) for individual realizations and in their Figure 7 for lines of the conditionally averaged vorticity field given $\tau_{xy}^+ < 0.6$. Note Fig. 4(c) plots the same vortex lines shown in Fig. 1 in the Introduction. One of the primary goals of this work is to elucidate the surprisingly complex and violent Lagrangian dynamics underlying this simple vortex-line structure.

Further insight into the local vorticity dynamics from the Eulerian perspective is provided by results on the in-wall pressure distribution $p(x, z)$, spanwise vorticity source $\sigma_z(x, z)$, and selected in-wall vortex-lines, as plotted in Fig. 5 for the vicinity of our stress local-minimum. Panel (a) of that figure reveals that our selected stress minimum is very close to a local pressure minimum, which in turn is flanked upstream and downstream at distances $\delta x^+ \simeq \pm 60$ by a pair of local pressure maxima. The pressure isolines or isobars in this plot are the lines of instantaneous generation of tangential vorticity in the Lighthill-Morton theory, with positive (counterclockwise) sense of rotation around pressure maxima and negative (clockwise) rotation around pressure minima. Of course, these isobars align only with the direction of generation of vorticity and the instantaneous vortex-lines within the wall are instead pointed mainly in the spanwise direction with small streamwise deviations, as shown in Fig. 5(b). The bending of these lines is explained in detail by the relation $\boldsymbol{\tau}_W = \nu \hat{\mathbf{n}} \times \boldsymbol{\omega}_W$ between the in-wall stress and vorticity fields (Lighthill 1963; Morton 1984). As a consequence, the stress vectors $\boldsymbol{\tau}_W$ plotted in Fig. 2 are locally perpendicular to the in-wall vortex-lines in Fig. 5(b) and the concavity of the lines is exactly that required to produce a converging flow at the low-speed streak. We plot in Fig. 5(b) as well the *negative* spanwise vorticity source $-\sigma_z^+ = -\partial p^+ / \partial x^+ = \partial \omega_z^+ / \partial y^+$




Figure 4: Vortex lines initiated at points with streamwise separations $\Delta x^+ = 12.3$, with $z^+ = 128.8$ and (a) $y^+ = 2$, (b) $y^+ = 4$, (c) $y^+ = 6$, (d) $y^+ = 8$. The shear-stress field from Fig. 2 is replotted in the bottom x - z -planes for reference.




Figure 5: (a) Pressure field p^+ at the wall, with selected isolines in black. (b) The source field $-\sigma_z^+$ of the negative spanwise vorticity and selected in-wall vortex lines. The asterisk “*” in both panels marks the location of the selected stress local-minimum.




Figure 6: (a) Field of spanwise vorticity fluctuation $\omega_z^{+'}$ in the x - y -plane at $z^+ = 85.9$. The black arrows represent the vectors (u', v') of the cross-section velocity fluctuation. (b) Same as (a), but with fluctuations calculated relative to local planar averages at constant y^+ and plotted in the near-wall region $y^+ < 10$.

in wall units. We included the minus sign since drag evidenced by a streamwise drop in pressure is associated to flux of negative spanwise vorticity away from the wall. Thus, the color/shading schemes in Figs. 2 and Fig. 5(b) are consistent, with yellow/light corresponding to increased drag (high stress, pressure drop) and blue/dark corresponding to reduced drag (low stress, pressure rise). Note however that the mean pressure drop associated to dissipative turbulent drag is $-\partial\bar{p}^+/\partial x^+ = 1/Re_\tau = 10^{-3}$, whereas the instantaneous streamwise pressure-gradients plotted in Fig. 5(b) are 100 times larger in magnitude, spanning a range from -0.15 to +0.15. It is consistent with the results in Fig. 5(b) that instantaneous pressure-gradients at the wall scale as u_*^2/δ_ν and thus remain $O(1)$ in wall-units. The average streamwise pressure drop is thus the result of near-cancellation between large instantaneous gradients of both signs.

It is interesting to observe in Fig. 5(b) that a region of negative streamwise pressure-gradient occurs just upstream of the stress local-minimum and a corresponding region of positive gradient occurs just downstream. This seems to agree with experimental observations of “bipolar” spanwise vorticity generation by Andreopoulos & Agui (1996) and Klewicki *et al.* (2008), based on conditionally-averaged time series of pressure and vorticity-flux, and on time-correlation functions of pressure and pressure-gradients. Andreopoulos & Agui (1996) proposed a conceptual model of ejections as rising “mushroom vortices” that would produce exactly such a bipolar pattern of spanwise vorticity source at the wall. See Andreopoulos & Agui (1996), Figure 28(b). However, a plot in Fig. 6 of the spanwise vorticity fluctuation ω_z' and velocity vector fluctuation (u', v') in the plane $z^+ = 85.9$ for our event is not consistent with such a picture. Panel (a) of that figure shows the entire y^+ -range, for context, and panel (b) zooms into the near-wall region $y^+ < 10$. To make the flow pattern more clear in Fig. 6(b) we have calculated fluctuations with respect to local planar averages at fixed distances y^+ from the wall (see SM for details). We observe that the “bipolar” source is produced by a fluid layer with $\omega_z' < 0$ being lifted up and to the right from the bottom wall, while replacement fluid with $\omega_z' > 0$ is advected in from the right and downward toward the wall. This dynamics is very similar to that postulated by Jimenez *et al.* (1988), Fig.6, as a mechanism of sublayer ejections. Those authors also observed thin, low-inclined layers of ω_z' like those




Figure 7: Field of viscous shear stress $\tau_{xy}^+ = (\partial u^+ / \partial y^+)$ at the wall $y^+ = 0$, in wall units so that the average is unity. Black arrows represent the two-dimensional in-wall stress vector τ_W . The asterisk “*” marks the location of the selected stress local-maximum.

in our Fig. 6(a) and interpreted them as Tollmien-Schlichting waves. Such shear layers are also inferred for coherent, nonlinear travelling waves (Waleffe (1998), Fig. 1). Finally, Andreopoulos & Agui (1996) have argued that “ejections which carry fluid of negative ω_z away from the wall ... are expected to be characterized by positive $\partial \omega_z / \partial y$. Negative $\partial \omega_z / \partial y$ is expected to be the distinguishing feature of sweeps...”. The ejection event that we consider has near $y^+ \simeq 5 - 10$ an upward flux of negative spanwise vorticity associated with $v' \omega'_z < 0$. However, there is no *instantaneous* balance between this advective flux and the viscous flux $\sigma_z = -\nu(\partial \omega_z / \partial y)$ at the wall. Thus it is not clear that the region with $\partial \omega_z / \partial y > 0$ upstream of the stress local-minimum should be regarded as the “source” of the advective spanwise vorticity flux at $y^+ \simeq 5 - 10$. Our Lagrangian analysis in section 4 shall show indeed that there is no causal connection.

3.3. Eulerian Description of Sweep Event

The stress local-maximum that we selected for study is likewise located within a high-speed streak of a type also commonly observed in near-wall turbulence, generally shorter than the low-speed streaks in streamwise extent and flanking them (Jiménez 2013). This neighborhood is illustrated in Fig. 7, which plots the viscous shear-stress $\tau_{xy} = \nu(\partial u / \partial y)$ at the wall and the location of the stress local-maximum as an asterisk “*”. The arrows representing the two-dimensional wall stress field τ_W indicate a near-wall, in-plane flow which is diverging from the streak. This divergence is consistent with a vertical flow that is downward toward the wall at the streak and it agrees with results of Sheng *et al.* (2009) for the conditional average stress-field in the vicinity of such local-maxima (see their Fig. 6(e)). Vortex visualization in Fig. 8(a) via λ_2 -isosurfaces shows a more complex environment than for the preceding local-minimum event. There are two or three large quasi-streamwise vortices at heights $y^+ > 20$, but these do not seem to influence strongly the near-wall physics. Instead at elevations $y^+ < 20$ there is a pair of counter-rotating almost streamwise vortices, one on each side of the observed high-speed streak. The plot in Fig. 8(b) of streamwise vorticity ω_x and cross-stream velocity vectors (w, v) , in the transverse y - z -plane cutting through the middle of the visualized box, shows clearly that this low-lying pair generate a downward, splatting flow between them. Unlike the pair observed near the stress-minimum, however, this pair is asymmetrical in strength,




Figure 8: (a) Isosurface $\lambda_2^+ = -0.0107$, with magnitude 3 times the local box-average value $\langle |\lambda_2^+| \rangle = 3.57 \times 10^{-3}$. The shear-stress field from Fig. 7 is replotted in the bottom x - z -plane for reference. (b) Field of streamwise vorticity ω_x^+ in plane $x^+ = 85.9$ (transparent in panel (a)). The black arrows represent cross-stream velocity vectors (w, v) .




Figure 9: Vortex lines initiated at points with streamwise separations $\Delta x^+ = 12.3$, with $z^+ = 128.8$ and (a) $y^+ = 4$, (b) $y^+ = 8$, (c) $y^+ = 12$, (d) $y^+ = 16$. The shear-stress field from Fig. 7 is replotted in the bottom x - z -planes for reference.




Figure 10: (a) Pressure field p^+ at the wall, with selected isolines in black. (b) The source field $-\sigma_z^+$ of the negative spanwise vorticity and selected in-wall vortex lines. The asterisk “*” in both panels marks the location of the selected stress local-maximum.

with the rightmost member of the pair distinctly weaker. If we increase the magnitude of the threshold value of λ_2 by even 33% to $\lambda_2^+ = -0.0143$, then no isosurface appears for this weaker vortex and only the single stronger vortex is observed at $y^+ < 20$. This is consistent with the findings of Sheng *et al.* (2009), who did encounter counter-rotating vortex pairs generating a splatting flow between them, in 11% of all of the samples in their study. However, under the condition $\tau_{xy}^+ > 1.8$, only about 8% of the realizations were of this type and all of these vortex pairs were quite asymmetrical in strength. Instead, 55% of the realizations in the study of Sheng *et al.* (2009) that satisfied the condition $\tau_{xy}^+ > 1.8$ had the stress maximum generated by a single low-lying vortex. Our selected event thus exhibits typical features for such stress-maxima. Similar features were observed also in other local-maxima stress events that we identified in the JHTDB channel-flow dataset satisfying the criteria $\tau_{xy}^+ \simeq 1.8$.

The vortex lines that we observe near this stress local-maximum likewise show expected features. See Fig. 9, which plots vortex-lines crossing the visualized domain in the spanwise direction, initialized at evenly spaced streamwise locations and at initial elevations $y^+ = 4, 8, 12, 16$. These lines are clearly squashed or depressed toward the wall by the downward splatting flow at the high-speed streak. Such arrays of vortex lines are typical of those observed in the vicinity of stress local-maxima in the study of Sheng *et al.* (2009), as illustrated in their Figure 16 for individual realizations and in their Figure 17 for lines of the conditionally averaged vorticity field given $\tau_{xy}^+ > 1.8$. As they also observed, the “troughs” of depressed lines are wider than the corresponding “hairpins” above low-speed streaks. Also, the asymmetry in strength of the streamwise vortices is clearly visible, with the ridge of lines located at the strong vortex obviously twisted higher than those at the weak vortex. In section 4 we shall study in depth the Lagrangian dynamics of the illustrated squashed lines at the local-maximum of stress.

First, however, we consider the vorticity dynamics for this event in more detail from the Eulerian perspective. The plot in Fig. 10(a) of the pressure field and its isolines at the wall shows that the stress maximum is close to a local pressure maximum, with a pair of local pressure minima upstream and downstream at distances $\delta x^+ \simeq \pm 80$. In contrast to the instantaneous generation of vorticity along the isobars, the actual vortex lines at the wall plotted in Fig. 10(b) are aligned mainly in the spanwise direction. The lines




Figure 11: Field of spanwise vorticity fluctuation $\omega_z^{+'}$ in the x - y -plane at $z^+ = 79.7$. The black arrows represent the vectors (u', v') of the cross-section velocity fluctuation. (b) Same as (a), but with fluctuations calculated relative to local planar averages at constant y^+ and plotted in the near-wall region $y^+ < 10$.

bend in the streamwise direction so that the locally-perpendicular stress vectors $\boldsymbol{\tau}_W$, as plotted in Fig. 7, correspond to a near-wall flow diverging away from the high-speed streak. The negative spanwise vorticity source $-\sigma_z$ also plotted in Fig. 10(b) again shows a bipolar pattern of the type inferred by Andreopoulos & Agui (1996) and Klewicki *et al.* (2008) from experimental data, with a region of positive streamwise pressure-gradient occurring just upstream of the stress local-maximum and a region of negative gradient just downstream. In the conceptual model of Andreopoulos & Agui (1996), Figure 28(a), sweeps correspond to inverted “mushroom vortices” moving toward the wall, producing just such a pattern of positive spanwise vorticity source upstream and negative spanwise source downstream. However, we do not observe such a mushroom vortex here. The plot in Fig. 11(a) of the spanwise vorticity fluctuation ω_z' and velocity vector fluctuation (u', v') in the plane $z^+ = 79.7$ does exhibit a fluid layer near $y^+ \simeq 10$ with $\omega_z' > 0$ and $v' < 0$, associated with a spanwise vorticity flux $v'\omega_z' < 0$. However, this occurs mainly upstream from the stress local-maximum, which is exactly opposite to what is proposed in the mushroom-vortex model. Fig. 11(b) shows instead underneath the primary vortex with $\omega_z' > 0$ a layer of strong secondary vorticity with $\omega_z' < 0$ just above the wall. (Unlike for the ejection case earlier, the local plane-average of streamwise velocity u , used to define the fluctuation u' in Fig. 11(b), is noticeably larger than the global average. See Fig. 11(a) and the SM.) This secondary layer is apparently produced by strong interaction of the primary vortex with the wall, as illustrated in Figure 26 of Andreopoulos & Agui (1996), but the primary and secondary layers are not rolled up to form the head of a mushroom vortex. It is the presence of the secondary layer with $\omega_z' < 0$ just above the wall that makes $\partial\omega_z/\partial y < 0$ upstream of the stress local-maximum, while the absence of such a secondary layer makes $\partial\omega_z/\partial y > 0$ downstream. The thin, inclined layers of ω_z' observed in Fig. 11(a), although now for a sweep rather than an ejection, are again quite similar to those reported by Jimenez *et al.* (1988), who also emphasized the important role of opposite-sign vorticity induced at the wall in producing such structures. These observations suggest some possible common features in viscous sublayer dynamics of ejections and sweeps. In the following section 4 we shall attempt to further explicate the physics of both types of events by stochastic Lagrangian analysis.

Table 3: Coordinates of Analyzed Vorticity Vectors

	x	y	z	t	ω_x	ω_y	ω_z
ejection	21.094707	0.994647	7.563944	25.9935	-2.24948978	-0.110395804	22.1811829
sweep	0.715000	0.995100	0.725900	25.9935	0.05745593	-0.1597188	47.2303467

4. Stochastic Lagrangian Dynamics of Vorticity in the Buffer Layer

4.1. Selection of Vorticity Vectors

We now select for analysis specific vortex lines and specific vorticity vectors lying upon them. In this initial study, we shall compare the Lagrangian dynamics of two vorticity vectors both pointing approximately in the negative spanwise direction and located at points at a similar distance from the wall, $y^+ \simeq 5$, at the bottom of the buffer layer. For the ejection event in particular, we want to investigate the initial nonlinear transfer of spanwise vorticity from the wall, which was described by Sheng *et al.* (2009) as the process where “spanwise vorticity lifts abruptly from the wall, creating initially a vertical arch”. We have therefore considered for Lagrangian analysis in the ejection event the lowest-lying vortex lines in Fig. 4(a), which are replotted in Fig. 12(a). These lines all start on the right at height $y^+ = 2$ and then pass left in the negative spanwise direction, rising up in a nearly vertical arch over the low-speed streak. We chose the middle of these lines, colored solid black, and in particular the vorticity vector at the top of the arch on that line at height $y^+ = 5.35$, indicated by the arrow. For comparison, we considered in the sweep event the vortex lines which are squashed down to a comparable height above the wall by the splatting flow. This is the set of lines in Fig. 4(c), replotted in Fig. 12(b), which all start at $y^+ = 12$ and as they pass leftward are depressed in a trough above the high-speed streak. We chose the middle of these lines, colored solid black, and in particular the vorticity vector at the bottom of the trough on that line at height $y^+ = 4.90$, indicated by the arrow. The components of the two vorticity vectors and their position coordinates in the channel-flow database are recorded in Table 3. Note that the vorticity vector for the sweep has a magnitude around twice that for the ejection. This is consistent with the argument of Lighthill (1963) that sweeps to the wall should stretch and magnify spanwise vorticity, while ejections should attenuate it (see his Fig.II.22).

Although the two vectors have different magnitudes, they are otherwise similar in orientation and in distance from the wall. This pair of vectors is thus well-suited to illuminate differences in Lagrangian dynamics that arise solely from the different flow conditions that exist in the two events. In particular, our expectation is that the vorticity in the ejection event should have arisen recently from the wall and should be “younger”, while the vorticity at the same distance from the wall in the sweep event should be “older” vorticity that entered from the wall at an earlier time, was processed by the flow, and was then returned by the splatting motion toward the wall. We shall see whether these expectations are borne out by our stochastic Lagrangian analysis.

We must first consider, however, the effect of filtering/coarse-graining on the structure and dynamics of the two events pictured in Fig. 12. As discussed in section 2, the mean conservation law (1.1) obtained from the JHTDB simulation data would be expected to have improved validity when spatially coarse-grained over n_i grid-spaces in the three coordinate directions $i = x, y, z$. This expectation was verified in paper I with a box-filter; see Fig. 3 there for the ejection case and Fig. 5 for the sweep case. However, to justify the JHTDB channel-flow data as a Navier-Stokes solution accurate enough to




Figure 12: The selected vortex lines, solid black, and the two selected vorticity vectors represented as arrows for (a) the ejection and (b) the sweep event. Other vortex lines in the two flow events are plotted as dotted lines.

investigate buffer-layer physics, it must also be shown that the filtering does not smear out the motions of interest. We have thus attempted to choose (n_x, n_y, n_z) as small as possible, sufficient to verify that mean conservation is improved but not so large that the coarse-graining obscures the essential physics. We have found that coarse-graining in the y -direction does not help to improve mean conservation, presumably since the numerical resolution in the channel-flow simulation was sufficient in that direction, so that we take $n_y = 0$. By trial, we have found that a good choice of filtering lengths in the wall-parallel directions are $n_x = n_z = 4$, although somewhat smaller or larger values give similar results. Fig. 13 verifies that the coarse-grained fields for $(n_x, n_y, n_z) = (4, 0, 4)$ in the ejection event preserve the basic flow features. Plotted in Fig. 13(a) is the field-line of the coarse-grained vorticity field which passes through the same space point as the field-line of the fine-grained vorticity that was plotted in Fig. 12(a) (see Table 3 for the spatial coordinates). This line starts at the right of the figure at vertical height $y^+ = 2.7039$ and Fig. 13(a) also plots an array of lines that start at that same height and pass in the spanwise direction. As in Fig. 12(a), the vortex lines rise up nearly vertically in an arch over the low-speed streak, and the main effect of the coarse-graining is that the arch is broadened in the spanwise direction. The coarse-grained vorticity field at the selected point is $\hat{\omega}(\mathbf{x}, t) = (-1.2871, 0.3523, -29.8522)$ with magnitude increased by mixing with adjacent stronger vorticity. Fig. 13(b) plots isosurfaces of the λ_2 -invariant for the coarse-grained field, which reveals a pair of quasi-streamwise, counter-rotating vortices flanking the low-speed streak. Compared with Fig. 3(a) for the fine-grained field, the vortices are somewhat smoother and weaker, but otherwise are quite similar. The relatively mild effects of the coarse-graining are explainable by the well-known long-range coherence in the streamwise direction of buffer-layer, slow-speed streaks, which extend with some meander for $10^3 - 10^4$ wall units (Jiménez 2013).

Similar observations are presented in Fig. 14 for the sweep event, considering also fields coarse-grained over a box with sides of $(n_x, n_y, n_z) = (4, 0, 4)$ grid-lengths. At the same space point as shown in Fig. 12(b), the coarse-grained vorticity vector is $\hat{\omega}(\mathbf{x}, t) = (0.099327, -1.5828, 49.04330)$, which is not much changed from the value of the fine-grained vorticity reported in Table 3. We have plotted through this point the integral line of the coarse-grained vorticity, which starts at the right of the figure at a height




Figure 13: (a) Vortex line (solid black line) and vorticity vector (arrow), for the coarse-grained vorticity field at the selected point in the ejection event. Other vortex lines are plotted as dotted lines. (b) Isosurface $\lambda_2^+ = -0.00815$ of the coarse-grained field, with magnitude 4 times the local box-average value $\langle |\lambda_2^+| \rangle = 2.038 \times 10^{-3}$.




Figure 14: (a) Vortex line (solid black line) and vorticity vector (arrow), for the coarse-grained vorticity field at the selected point in the sweep event. Other vortex lines are plotted as dotted lines. (b) Isosurface $\lambda_2^+ = -0.005338$ of the coarse-grained field, with magnitude 3 times the local box-average value $\langle |\lambda_2^+| \rangle = 1.780 \times 10^{-3}$.

$y^+ = 10.5622$. This is just a bit lower than the height $y^+ = 12$ of the corresponding line for the fine-grained field. An evenly spaced array of lines of coarse-grained vorticity that start also at height $y^+ = 10.5622$ is shown in Fig. 12(b). These are depressed into a trough over the high-speed streak, which is shallower but just slightly wider than for the similar trough of lines in Fig. 12(b) for the fine-grained vorticity. Fig. 14(b) plots isosurfaces of the λ_2 -invariant for the coarse-grained field, which have a somewhat simpler structure but are qualitatively very similar to the isosurfaces in Fig. 8(a) for fine-grained vorticity. The effects of coarse-graining are even less pronounced here than for the ejection event, presumably because the sweep is a bit broader in the spanwise direction.

We believe that these results provide sufficient *a priori* justification for our use of the JHTDB dataset to study buffer-layer vortex dynamics. We shall present below the results of our stochastic Lagrangian analysis of the raw/unfiltered vorticity, where mean conser-

vation of the stochastic Cauchy invariant already holds to within a few percent. In the SM we present the corresponding Lagrangian analysis for the coarse-grained stochastic Cauchy invariant, with $(n_x, n_y, n_z) = (4, 0, 4)$, demonstrating improved conservation. All results are slightly changed quantitatively, but not qualitatively, by the filtering.

4.2. Lagrangian Description of Ejection Event

4.2.1. Origin of Vorticity at the Wall

We now turn to the central question of the paper, the origin at the wall of the buffer-layer vorticity. A regular spatial array of vortex-lines such as plotted in Fig. 12(a) might suggest a simple dynamical process of “abrupt lifting” of the vorticity away from the wall within a few viscous times. There are, however, *a priori* theoretical reasons to expect that the process is much slower and more complex. We have already discussed in paper I the strong Lagrangian chaos in the buffer-layer (Johnson *et al.* 2017), which argues against a simple ideal lifting motion. Furthermore, we note that stochastic Lagrangian particles, moving backward in time, cannot ever reach the wall by fluid advection alone but only through viscous diffusion, because the wall-normal velocity drops to zero rapidly with decreasing distance to the wall. Diffusion is an intrinsically slow process. If wall-normal velocity were exactly zero, then a Brownian particle with diffusivity ν released at height y^+ would reach the wall at $y^+ = 0$ in a random time $\tilde{\sigma}^+ > 0$ (in wall units) with probability density

$$p(\sigma^+) = \frac{y^+}{\sqrt{4\pi\sigma^{+3}}} \exp\left(-\frac{y^{+2}}{4\sigma^+}\right), \quad \sigma^+ > 0 \quad (4.1)$$

which is a special case of an “inverse Gaussian distribution”; see Chhikara & Folks (1988), Borodin & Salminen (2015), Part II, formula 2.02 (p.295). The mean value of the hitting time with distribution Eq. (4.1) is infinite, which implies that it takes a very long time to reach the wall, with a high probability. This result does not contradict the natural expectation that vorticity created at the wall will diffuse across the viscous sublayer to $y^+ \sim 5$ in a time of order $t_\nu = \delta_\nu^2/\nu$, because the diffusion process is asymmetric in time. Under time-reversal, a Brownian particle that reaches the wall is described by a “3-dimensional Bessel process” forward in time (Borodin & Salminen (2015), Part I, Chapter II.5, p.35), which is strongly repelled from the wall at $y = 0$, because it cannot return there again. The stochastic trajectory released at $y^+ = 5$ and moving backward in time will take much longer to reach the wall, because it will cross and recross the level $y^+ = 5$ many times before ultimately reaching the wall located at $y^+ = 0$.

Numerical results on stochastic Lagrangian dynamics of vorticity from our Monte Carlo approach confirm the above theoretical expectations. The percentage of particles released at the selected point at $y^+ = 5.35$ which have hit the wall going backward in time is plotted in Fig. 15(a) versus $\delta s = s - t$. The percentage grows faster in time for this fluid ejection event than it does for pure diffusion, because the lifting flow advects particles toward the wall backward in time. Nevertheless, hundreds of viscous times are required for nearly the entire ensemble of particles to hit the wall. After an initial fairly abrupt rise to $\sim 75\%$ in the interval $-50 < \delta s^+ < 0$, less than 90% of the particles have hit the wall even after 150 viscous times. An even more salient issue is the relative contributions to the stochastic Cauchy invariant which arise from particles that have arrived at the wall and from those still in the flow interior. We can define such partial contributions by

$$\mathbb{E}_C[\tilde{\omega}_{s i}(\mathbf{x}, t)] := \mathbb{E}[\mathbf{1}_C \tilde{\omega}_{s i}(\mathbf{x}, t)]$$




Figure 15: (a) Fraction of particles at the wall, (b) partial means of the parallel component of the stochastic Cauchy invariant, and (c) partial means of the perpendicular component of the stochastic Cauchy invariant. All quantities in wall units plotted versus δs^+ .

where C is a subset of stochastic trajectories satisfying conditions such as

$$\text{particle at wall at time } s: \quad W = \{|\tilde{b}_t^s(\mathbf{x})| = h\}$$

$$\text{particle in interior at time } s: \quad I = \{|\tilde{b}_t^s(\mathbf{x})| < h\}$$

and $\mathbf{1}_C$ is the indicator function of this subset, = 1 on the set and = 0 on its complement. The stochastic Cauchy invariants are vector quantities, and we consider here the intrinsic parallel and perpendicular components as defined in paper I, Eq.(3.14), or:

$$\tilde{\omega}_{s\parallel}(\mathbf{x}, t) := \tilde{\omega}_s(\mathbf{x}, t) \cdot \hat{\mathbf{n}}_\omega(\mathbf{x}, t), \quad \tilde{\omega}_{s\perp}(\mathbf{x}, t) := \tilde{\omega}_s(\mathbf{x}, t) - \tilde{\omega}_{s\parallel}(\mathbf{x}, t)\hat{\mathbf{n}}_\omega(\mathbf{x}, t), \quad (4.2)$$

where $\hat{\mathbf{n}}_\omega(\mathbf{x}, t) = \boldsymbol{\omega}(\mathbf{x}, t)/|\boldsymbol{\omega}(\mathbf{x}, t)|$. In Fig. 15(b)-(c) we plot partial means for subsets $C = W, I$ and for components $i = \parallel, \perp$. We plot also in Fig. 15(b)-(c) the conserved total means of the stochastic Cauchy invariants, reproducing the results for $i = \parallel, \perp$ in Fig. 2 of Paper I. As can be observed, the conserved means are the summed results of strongly time-dependent contributions separately from particles at the wall and in the interior and conservation for $i = \perp$, in particular, involves complete vector cancellations between the two contributions. (To make this cancellation visually obvious, we have plotted $+\mathbb{E}_I[\tilde{\omega}_{s\perp}(\mathbf{x}, t)]$ and $-\mathbb{E}_W[\tilde{\omega}_{s\perp}(\mathbf{x}, t)]$ in Fig. 15(c). The complete cancellation is more perfectly exhibited by the results on the coarse-grained Cauchy invariant presented in the SM, since the total perpendicular component remains closer to vanishing there.) For the parallel component plotted in Fig. 15(b) there is a gradual crossover from the conserved mean arising from interior particles to instead arising from wall particles going backward in time. The contribution from the wall particles is notably larger than the fraction of the particles located at the wall. Indeed, at $\delta s^+ \simeq -100$, almost 100% of the conserved parallel component arises from the wall contribution, even though only about 85% of the particles have reached the wall.

Even so, however, the ~ 100 viscous times to get the entire parallel vorticity component originating from the wall is much longer than the few viscous times postulated by Sheng *et al.* (2009). Lifting of vorticity from the wall is not an “abrupt” event but is instead a very prolonged process. To quantify the time required or the mean “age” of the vorticity vector $\boldsymbol{\omega}(\mathbf{x}, t)$, we can introduce an integral formation time for the parallel component

$$T_{\parallel}(\mathbf{x}, t) := \int_{-\infty}^t ds f_{\parallel}(s; \mathbf{x}, t), \quad f_{\parallel}(s; \mathbf{x}, t) := \frac{\mathbb{E}_I[\tilde{\omega}_{s\parallel}(\mathbf{x}, t)]}{|\boldsymbol{\omega}(\mathbf{x}, t)|}. \quad (4.3)$$

Here $f_{\parallel}(s; \mathbf{x}, t)$ is the fractional contribution of the parallel vorticity arising at times $s < t$ from interior particles, satisfying $f_{\parallel}(t; \mathbf{x}, t) = 1$ and $f_{\parallel}(-\infty; \mathbf{x}, t) = 0$. Fig. 16 plots as a function of δs the fraction $f_{\parallel}(s; \mathbf{x}, t)$ obtained from our numerical Monte Carlo method,




Figure 16: Fractional contribution of the parallel vorticity arising at times $s < t$ from interior particles, plotted versus δs^+ , and its discrete approximation as “abrupt lifting”.

together with the discrete approximation that corresponds to a Heaviside step function with jump at $\delta s = -T_{\parallel}^+(\mathbf{x}, t)$. Numerical quadrature gives $T_{\parallel}^+(\mathbf{x}, t) \simeq 20.7$ in wall units. The step function represents graphically the “abrupt lifting” proposed by Sheng *et al.* (2009), pictured as a discrete event. This is not an unreasonable caricature of the actual vortex-lifting process, except that the integral time is about an order of magnitude larger than the heuristic time estimate by Sheng *et al.* (2009) and the discrete picture misses entirely the long, slowly decaying tail. As a matter of fact, the true “age” of the vorticity $\boldsymbol{\omega}(\mathbf{x}, t)$ in our selected point is even much larger than estimated by $T_{\parallel}(\mathbf{x}, t)$ because, as shown in Fig. 15(c), the interior particles contribute also a very slowly vanishing *perpendicular* component to the vorticity. One could define an integral time for decay of this perpendicular component to zero, for example, by

$$T_{\perp}(\mathbf{x}, t) := \int_{-\infty}^t ds f_{\perp}(s; \mathbf{x}, t), \quad f_{\perp}(s; \mathbf{x}, t) := \frac{|\mathbb{E}_I[\tilde{\omega}_{s\perp}(\mathbf{x}, t)]|}{\max_s |\mathbb{E}_I[\tilde{\omega}_{s\perp}(\mathbf{x}, t)]|}. \quad (4.4)$$

We shall not attempt a quantitative evaluation of this quantity, because our numerical Monte Carlo scheme with $N = 10^7$ particles does not yield converged results for the perpendicular Cauchy invariant at times $\delta s^+ < -100$. However, Fig. 15(c) shows that the perpendicular component at $\delta s^+ = -100$ still remains about $0.1|\boldsymbol{\omega}|$. It is thus clear from Fig. 15(b)-(c) at least that $T_{\perp}(\mathbf{x}, t) \gg T_{\parallel}(\mathbf{x}, t)$.

The broad space-time distribution and intricate Lagrangian dynamics of the vorticity generation process is further revealed by Fig. 17. Plotted at the channel wall are realizations of the parallel component of the stochastic Cauchy invariant, $\tilde{\omega}_{s\parallel}(\mathbf{x}, t)$, with values encoded by color/shade, at the position $(\tilde{a}_*(\mathbf{x}, t), \tilde{c}_*(\mathbf{x}, t))$ where the particle hits the wall and grouped in intervals of hitting time $-5(5k+1) < \delta\tilde{\sigma}_t^+(\mathbf{x}) < -25k$ for $k = 0, 1, 2, 3, 4, 5$, with $\delta\tilde{\sigma}_t(\mathbf{x}) := \tilde{\sigma}_t(\mathbf{x}) - t$. This figure was created with a sub-ensemble of 10^6 stochastic Lagrangian particles and each panel plots Cauchy vorticity contributions for all of the particles in this sub-ensemble that hit the wall in the given 5-viscous-time interval. In the SM, we provide a video with similar plots as frames, at a larger set of backward times δs , with greater time resolution, and employing all 10^7 available particles. Averaging over all of these wall contributions yields the resultant vorticity magnitude $|\boldsymbol{\omega}(\mathbf{x}, t)|$ at the selected space-time point $(\mathbf{x}, t)$ in the ejection event. For spatial reference, Fig. 17 plots also in each frame three points: (1) the wall-parallel position (x, z) of the selected point, taken as the coordinate origin $(0, 0)$, plotted as a black asterisk “*”; (2) the mean position of all particles in the entire ensemble at the given time s , plotted as




Figure 17: Scatterplot of $\tilde{\omega}_{s\parallel}^+(\mathbf{x}, t)$ values for particles hitting the wall in 5-unit time intervals ending at: (a) $\delta s^+ = -5$, (b) $\delta s^+ = -30$, (c) $\delta s^+ = -55$, (d) $\delta s^+ = -80$, (e) $\delta s^+ = -105$, (f) $\delta s^+ = -130$. Mean position of all particles (*) and only in interior (*). The resultant vorticity magnitude $|\omega^+(\mathbf{x}, t)| \doteq 0.45$.

a red/dark-grey asterisk “*”; and, (3) the mean position of all interior particles at the given time s , plotted as a green/light-grey asterisk “*”. The green point drifts quickly upstream backward in time, since the “living” particles are generally at higher elevations where the streamwise velocity is larger. The red point also drifts upstream, but more and more slowly as particles hit the wall and stop at their “birth place.” These points provide useful context in the figure.

The wall-plots in Fig. 17 display clearly the space-time origin of the resultant vorticity. The particles newly arrived to the wall in each frame land roughly between the mean positions of all particles in the ensemble and of interior “living” particles. Over the range of times $-100 < \delta s^+ < 0$ that contribute substantially to $\omega(\mathbf{x}, t)$, the stochastic particles hit the wall in a region extending ~ 1000 wall units in the streamwise direction and ~ 200 wall units in the spanwise direction. The “cloud” of points instantaneously hitting the wall in each frame also expands going backward in time, faster in the streamwise direction than in the spanwise. The faster growth of the particle cloud in the streamwise direction is explained by the dispersive effect of the mean-shear, which produces a super-ballistic $\sim (\delta s)^3$ growth of the mean-square streamwise extent of the cloud (Corrsin 1953). Such super-ballistic growth was previously verified for stochastic Lagrangian particles in the buffer-layer of this same channel-flow database (Drivas & Eyink 2017b) and it is also observed here (see SM). The mean-square spanwise extent of the cloud also grows backward in time, but at a slower diffusive rate $\sim \delta s$ (see SM). We conclude that the vorticity in the vertical arch at $y^+ = 5.35$ does not arise from a location ~ 10 wall units upstream, as conjectured by Sheng *et al.* (2009), but instead from a region at the wall at least 1-2 orders of magnitude larger in extent.

The magnitude of the final vorticity in the arch is $|\omega^+| \doteq 0.45$, but the individual contributions of the stochastic Cauchy invariant are much larger. The maximum values observed will, of course, increase with the number of samples N employed in our calculation. For the $N = 10^6$ ensemble used to prepare Fig. 17, the largest values of $\tilde{\omega}_{s||}^+(\mathbf{x}, t)$ were seen to grow with increasing $|\delta s|$ at a slightly less than exponential rate over the interval $-150 < \delta s^+ < 0$, from values near ± 1 at small δs^+ to around ± 500 at $\delta s^+ = -150$. There are substantial fluctuations from smooth (sub)exponential growth, however, and the largest values of the stochastic invariant $\tilde{\omega}_{s||}^+$ encountered over the interval $-150 < \delta s^+ < 0$ with $N = 10^6$ were $\pm 10^4$. It should be emphasized that these large values do not correspond to the vorticity magnitudes sampled by the particles when they hit the wall. The wall-vorticity is pointed mainly in the spanwise direction, as illustrated in Fig. 5(b), and has magnitude $\doteq 1$ in wall units. The large values are instead the consequence of exponential growth of the wall-vorticity as it is transported forward in time along the stochastic Lagrangian trajectories, consistent with the growth of variances observed in Paper I, Fig. 2(b) and with the strong Lagrangian chaos reported in the buffer-layer (Johnson *et al.* 2017). The mean value of the realizations $\tilde{\omega}_{s||}^+$ arising from the wall can yield the order-unity value $|\omega^+| \doteq 0.45$ only if there is nearly complete cancellation between contributions from opposite sign. This is clearly exhibited in Fig. 17, where yellow/light indicates large positive values and blue/dark large negative values. The negative values arise from vorticity elements that start at the wall aligned in the negative spanwise direction with the mean vorticity, but whose parallel component is rotated 180° as the vector is transported from its “birth place” at the wall to the final point $(\mathbf{x}, t)$ on the arch. The extensive cancellation between oppositely signed contributions is the representation in our stochastic Lagrangian framework of strong viscous destruction of vorticity, which counterbalances the exponential growth of vorticity by strong Lagrangian chaos.

The plots in Fig. 17 exhibit interesting and nontrivial structure of the parallel Cauchy invariant plotted against wall-position, especially for intermediate values of δs^+ . The particles that hit the wall in the earliest time interval $-5 < \delta s^+ < 0$ (panel (a)) contribute only positive values of order unity. The wall-vorticity in this early time is transported essentially by pure diffusion and without stretching or rotation. However, for more negative values of δs^+ , large opposite signs of $\tilde{\omega}_{\parallel}^+$ develop at the wall, with clear spatial organization. This order presumably reflects in part the well-known Eulerian vortex structures in the flow, such as the counter-rotating pair of streamwise vortices pictured in Fig. 3(a). However, these patterns involve also the Lagrangian evolution over time and become progressively more complex and fine-grained as δs^+ grows more negative. The complex, intertwined positive and negative values enhance the amount of cancellation. Eventually, for $\delta s^+ < -100$, the scatter of positive and negative values of $\tilde{\omega}_{\parallel}^+$ becomes essentially random and the cancellation is nearly complete (panel (f)). Particles continue to hit the wall backward in time for $\delta s^+ \ll -100$ and the r.m.s. values of $\tilde{\omega}_{\parallel}^+$ grow larger, but these very early contributions become less and less probable and cancel almost entirely, giving no net contribution to the resultant magnitude $|\omega(\mathbf{x}, t)|$ at the top of the arch.

4.2.2. Relation to the Eulerian Vorticity Source

To make connection with the Eulerian theory of Lighthill (1963) and Morton (1984), we present in Fig. 18 a plot of the negative spanwise vorticity source $-\sigma_z(\mathbf{x}, s)$ at time $\delta s^+ = -5k$, together with a scatterplot of particles landing at the wall in the interval of hitting time $-(5k+1) < \delta \tilde{\sigma}_t^+(\mathbf{x}) < -5k$, for values $k = 0, 1, 2, 3, 4, 5$ in successive panels. To render more clear the pattern of the source underneath the particle markers we have added isolines $-\sigma_z^+ = k/30$ for $|k| \leq 4$. We go back in time by only about $T_{\parallel}^+ \doteq 20.7$, since Fig. 16 shows that more than 50% of the final parallel vorticity is generated from the wall in that interval. We provide in the SM a movie whose frames are plots of the same format, but with more frames and going back 50 viscous times. The first panel, Fig. 18(a), is essentially the same as Fig. 5(b) but showing a larger domain. The two regions with $\sigma_z < 0$ just upstream of the stress-minimum and with $\sigma_z > 0$ just downstream in Fig. 5(b) now appear as parts of larger “band” structures. The spanwise vorticity source plotted in Fig. 18 exhibits an alternating pattern of such bands, each with streamwise thickness $L_x^+ \sim 25 - 50$ and spanwise length $L_z^+ \sim 50 - 100$. The reverse sign in successive bands is the manifestation of the “bipolarity” of the vorticity source. Since it has been proposed in the literature that ejections should be associated with values $\sigma_z < 0$ (Andreopoulos & Agui 1996), we would like to investigate whether the region with $\sigma_z < 0$, just upstream of the point marked with an asterisk “*”, can be the source of the vertical vortex arch at height $y^+ = 5.35$.

A striking feature in Fig. 18 which is even more apparent in the associated movie (see SM) is that the band patterns seem to travel in the negative streamwise direction, backward in time. In fact, it is well-known that velocity and pressure structures at the wall consist of travelling waves with streamwise velocities $\sim 10 - 15$ measured in units of the friction velocity u_* (Kim & Hussain 1993), agreeing well with the velocity inferred by eye from Fig. 18. This means in particular that propagation speeds of the waves exceed the mean flow velocity $\bar{u}(y)$ for $y^+ < 15$ and increasingly so as y^+ decreases. A consequence is that the band with $\sigma_z < 0$ just upstream of the “*” moves further upstream (backward in time) at a considerably faster speed than does the cloud of particles released at $y^+ = 5.35$, and thus particles in that ensemble hit this moving band with very low probability. It therefore seems ruled out that the upstream band with $\sigma_z < 0$ is the “source” of the vorticity in the arch at $y^+ = 5.35$. Only particles released at very small heights $y^+ \lesssim 1$,




Figure 18: Color plots of negative spanwise vorticity source together with scatterplots of particles hitting the wall in unit time intervals ending at: (a) $\delta s^+ = -1$, (b) $\delta s^+ = -6$, (c) $\delta s^+ = -11$, (d) $\delta s^+ = -16$, (e) $\delta s^+ = -21$, (f) $\delta s^+ = -26$. The light grey lines are contour levels $-\sigma_z^+ = k/30$ for integers $|k| \leq 4$. Also shown are the mean position of all particles (*) and of only those in the interior (*).




Figure 19: (a) Fraction of particles at the wall, (b) partial means of the parallel component of the stochastic Cauchy invariant, and (c) partial means of the perpendicular component of the stochastic Cauchy invariant. All quantities in wall units plotted versus δs^+ .

well below the arch, hit that band with substantial probability and at those heights the vortex lines are flat, with almost no vertical component.

Instead, the particles released from the vortex arch at $y^+ = 5.35$ and going backward in time hit structures that were originally *downstream* of “*” in Fig. 18(a) but which rapidly moved upstream of “*” in subsequent panels. In particular, there is a band with large negative values $\sigma_z^+ \simeq -0.1$ about 100 wall units downstream of “*” in Fig. 18(a) which particles hit with very high probability at times $-15 < \delta s^+ < -10$ (Fig. 18(c-d)). At this same interval in Fig. 16 one sees a very sharp increase in the contribution from the wall to the final vorticity magnitude. The intense band 100 wall units downstream of “*” in Fig. 18(a) is thus a more likely source of the enhanced negative spanwise vorticity in the vertical arch (Fig. 6), or at least the part associated to “abrupt lifting”. Of course, the negative spanwise vorticity injected by this source is not all delivered to this one arch but is instead distributed more generally throughout the flow. Fig. 18(e-f) shows that the particles for $-26 < \delta s^+ < -21$ sample a broad region with smaller negative values of source σ_z^+ , corresponding to the turnover to a slowly decaying tail in Fig. 16.

4.3. Lagrangian Description of Sweep Event

We now discuss the results for the vorticity at the bottom of the trough in the “sweep” event, as pictured in Fig. 12(b). In particular, we investigate how the vorticity in the trough originates at the wall. We are especially interested to compare with the previously discussed results for the low arch in the “ejection” event pictured in Fig. 12(a).

4.3.1. Origin of Vorticity at the Wall

Plotted in Fig. 19(a) is the percentage of particles residing at the wall as a function of $\delta s = s - t$, in wall units. The percentage is more slowly rising (backward in time) than for the ejection case pictured in Fig. 15(a) and, indeed, at $\delta s^+ = -150$ has risen to only 50%. This is consistent with the expectation that the vorticity is “older” in the sweep than it is in the ejection, at the same height above the wall. Since the wall-normal velocity is downward in the sweep, the ensemble of particles moves generally upward going backward in time and fewer particles hit the wall over the same time interval than for the ejection.

Plotted next in Fig. 19 are the partial contributions to the mean Cauchy invariants arising from the wall ($C = W$) and from interior ($C = I$) as functions of δs^+ , for parallel component ($i = \parallel$) in panel (b) and perpendicular component ($i = \perp$) in panel (c). Just as for the ejection case, we see that the conservation of the mean invariant is non-trivial and involves detailed cancellations between contributions from the wall and from the interior. Indeed, for the perpendicular component of the stochastic Cauchy invariant in




Figure 20: Fractional contribution of the parallel vorticity arising at times $s < t$ from interior particles, plotted versus δs^+ , and its discrete approximation as “abrupt lifting”.

Fig. 19(c) the two separate contributions are both *increasing* backward in time, roughly linearly in δs^+ over the range $-150 < \delta s^+ < 0$. The entire perpendicular contribution $\omega_{\perp, \text{int-near}}^+$ from the interior at time $\delta s^+ = -150$ has a magnitude near $|\omega^+|$, the value of the ultimate parallel component. It is not surprising to see such a larger perpendicular contribution for the sweep. The cloud of interior particles rises steadily backward in time from $y^+ \simeq 5$ at $\delta s^+ = 0$ to $y^+ \simeq 42.5$ at $\delta s^+ = -150$ (see plot in SM), while simultaneously the number of interior particles as a percentage of the total drops from 100% to 50%. This subcloud of particles represents the vorticity brought down from the interior of the flow by the splatting motion, forward in time. Since the vorticity high in the buffer layer is more variable, it is natural that this interior contribution $\omega_{\perp, \text{int-near}}^+$ to the resultant vorticity is not mainly spanwise but points instead in an orthogonal direction. This orthogonal component is exactly cancelled by an equal and opposite contribution $\omega_{\perp, \text{wall-near}}^+ = -\omega_{\perp, \text{int-near}}^+$ from the other 50% of the particles that hit the wall in the near-time interval $-150 < \delta s^+ < 0$.

A first conclusion of the results plotted in Fig. 19(c) is that the vorticity in the sweep is indeed very “old” and arose from the wall in the distant past, as expected. The quantity $|\mathbb{E}_I[\tilde{\omega}_{s\perp}(\mathbf{x}, t)]|$ that appears in the definition (4.4) of the perpendicular integral time $T_{\perp}(\mathbf{x}, t)$ is apparently increasing past $\delta s^+ = -150$ and to values $> |\omega|$, before finally decaying to zero. We do not have results well enough converged, even with $N = 10^7$, in order to evaluate this time accurately, but it is clear at least that $T_{\perp}^+(\mathbf{x}, t) \gg 100$ for the sweep. A more surprising and puzzling conclusion follows, however, from the exact anti-correlation $\omega_{\perp, \text{wall-near}}^+ = -\omega_{\perp, \text{int-near}}^+$. The interior contribution $\omega_{\perp, \text{int-near}}^+$ at time $\delta s^+ = -150$ was itself the result of vorticity shed from the wall in the far distant past. Going far backward to distant times where nearly 100% of the particles have hit the wall must reproduce it. Thus, $\omega_{\perp, \text{int-near}}^+ = \omega_{\perp, \text{wall-dist}}^+$, where the latter is the contribution from particles that hit the wall in the distant past $\delta s^+ < -150$. The immediate implication is that there is also a perfect anti-correlation $\omega_{\perp, \text{wall-dist}}^+ = -\omega_{\perp, \text{wall-near}}^+$. In other words, the vorticity shed from the wall in the far distant past $\delta s^+ < -150$ is making a very sizable contribution to streamwise and wall-normal components of vorticity in the trough, with magnitude $|\omega_{\perp, \text{wall-dist}}^+| > 1$, but this is *exactly* cancelled by a large, exactly anti-parallel contribution $\omega_{\perp, \text{wall-near}}^+$ from the wall in the near past $-150 < \delta s^+ < 0$! This very long-range temporal correlation is required by mean conservation of the value 0 of the perpendicular Cauchy invariant, but it seems a bit counterintuitive, fluid dynamici-

cally. A possible explanation is that $\omega_{\perp,wall-near}^+$ arises from secondary vorticity induced by the strong interaction with the wall of the primary interior vorticity $\omega_{\perp,int-near}^+$ as it is advected downward. This picture may help make plausible the exact anti-correlation $\omega_{\perp,wall-near}^+ = -\omega_{\perp,int-near}^+$. In any case, our findings emphasize not only the great “age” of the vorticity vector in the “trough” but also its very complex origin at the wall.

A surprise in the opposite direction is that the final parallel component of the vorticity vector in the “trough”, which is pointed almost spanwise, is just about as “old” as the vorticity vector at the same height on the “arch” in the ejection case. Indeed, the history of formation of the parallel component out of vorticity shed from the wall is remarkably similar for the “ejection” and the “sweep”, as can be seen by comparing the results in Fig. 15(b) and Fig. 19(b). Except for the different magnitudes of the parallel components in the two cases, the plots otherwise agree quite closely. This fact is underlined by Fig. 20, which plots for the vorticity vector in the “trough” the fractional contribution to the parallel component arising from the interior. This plot is almost identical to that in Fig. 16 for the ejection case. The integral time of formation of the parallel component calculated from Eq.(4.3) must therefore be similar also for the two cases. Indeed the value obtained by numerical quadrature for the sweep, $T_{\parallel}^+(\mathbf{x}, t) \doteq 24.6$, is just slightly larger than the value $T_{\parallel}^+(\mathbf{x}, t) \doteq 20.7$ for the ejection.

To further explore this issue we have made for the sweep case a figure of the same type as Fig. 17 for the ejection, again plotting realizations of the parallel component of the stochastic Cauchy invariant. See Fig. 21, which uses the same time intervals and the same size subensemble of 10^6 particles as in the earlier plot. In the SM, we provide a video with greater time resolution and employing all 10^7 available particles. This plot only deepens the mystery, however. The plots in Fig. 21 are broadly similar to those in Fig. 17 for the ejection, but also show significant differences. The clouds of particles are clearly more compact for the sweep case and disperse less quickly going backward in time. Furthermore, the spatial pattern at the wall of the parallel Cauchy invariant values is strikingly “bipartite” for the sweep at times $-60 < \delta s^+ < 0$, with large positive values in the lower half of the particle cluster and large negative values in the upper half. This pattern is presumably due to the rotation of vorticity vectors by the pair of low-lying quasi-streamwise vortices pictured in Fig. 8(a), and it is much more ordered than the pattern for the ejection case in Fig. 17. Despite these clear differences in the spatial patterns in the two plots, the summed results yield the time variations plotted in Fig. 16 and Fig. 20, which are remarkably similar.

4.3.2. Relation to the Eulerian Vorticity Source

Some further insight may be gained by considering the Eulerian picture. In Fig. 22 we plot the negative spanwise vorticity source $-\sigma_z(\mathbf{x}, s)$ together with a scatterplot of particles landing at the wall. We consider the same set of times $\delta s^+ = -(5k + 1)$, $k = 0, 1, 2, 3, 4, 5$ as in Fig. 18 for the ejection, going backward by 26 viscous times. This is close in value to the integral shedding time $T_{\parallel}^+ \doteq 24.6$ for the sweep. A movie with greater time-resolution and going back further in time is provided in the SM. As before, the first panel, Fig. 22(a), is essentially the same as Fig. 10(b) but over a larger spatial domain. It was conjectured by Andreopoulos & Agui (1996) that positive wall sources $\sigma_z > 0$ should be associated with sweeps. Indeed, just upstream of the point marked with an asterisk “*” in Fig. 22(a) there is a band with large values $\sigma_z \doteq 0.1$. However, as shown by the subsequent panels (or by the movie, in more detail) the particles hit that region of the wall with negligible probability backward in time, since this band with $\sigma_z > 0$ retreats upstream with high velocity $\sim 10u_*$. Instead, the particles hit mainly




Figure 21: Scatterplot of $\tilde{\omega}_{s\parallel}^+(\mathbf{x}, t)$ values for particles hitting the wall in 5-unit time intervals ending at: (a) $\delta s^+ = -5$, (b) $\delta s^+ = -30$, (c) $\delta s^+ = -55$, (d) $\delta s^+ = -80$, (e) $\delta s^+ = -105$, (f) $\delta s^+ = -130$. Mean position of all particles (*) and only in interior (*). The resultant vorticity magnitude $|\omega^+(\mathbf{x}, t)| = 0.95$.




Figure 22: Color plots of negative spanwise vorticity source together with scatterplots of particles hitting the wall in unit time intervals ending at: (a) $\delta s^+ = -1$, (b) $\delta s^+ = -6$, (c) $\delta s^+ = -11$, (d) $\delta s^+ = -16$, (e) $\delta s^+ = -21$, (f) $\delta s^+ = -26$. The light grey lines are contour levels $-\sigma_z^+ = k/30$ for integers $|k| \leq 4$. Also shown are the mean position of all particles (*) and of only those in the interior (*).

regions with $\sigma_z < 0$ over the time interval shown in Fig. 22. In particular, there is a space band with moderately negative values originally about 200 wall units downstream of the point marked “*” which travels rapidly upstream backward in time and which intensifies to values $\sigma_z \doteq -0.1$. The particles released from the depressed vortex line at height $y^+ = 4.90$ are sampling mainly points in this band of strong negative source $\sigma_z \doteq -0.1$ over the time interval $-25 < \delta s^+ < -16$. In fact, the pattern of vorticity source sampled by the particles in this sweep case is rather similar to that sampled in the ejection case, pictured in Fig. 18. This seems to contradict the proposal of Andreopoulos & Agui (1996). Here it should be noted that nonlinear advection contributes also in the sweep case a *negative* flux of spanwise vorticity away from the wall over much of the region $y^+ < 15$. As illustrated in Fig. 11, positive fluctuations of spanwise vorticity $\omega'_z > 0$ are here often associated pointwise with downward fluctuations $v' < 0$ of wall-normal velocity.

These results suggest a possible explanation for the surprising agreement of Fig. 16 for the ejection and Fig. 20 for the sweep. This agreement could be expected if viscous diffusion dominates the transport of spanwise vorticity in the viscous sublayer, not only on average (Klewicki *et al.* 2007; Eyink 2008), but also in instantaneous realizations of the flow. The structure of in-wall vortex lines is very similar for the ejection and sweep, with magnitudes $|\omega^+| \doteq 1$ and pointed mainly in the spanwise direction (or negative spanwise direction for rotated visualizations). Of course, these nearly uni-directional wall vorticity vectors are strongly stretched and rotated by the stochastic Lagrangian flow and yield both large positive and large negative values of the parallel stochastic Cauchy invariant, as illustrated in Figs. 17, 21 for the ejection and sweep, respectively. It is possible however that these large values almost completely cancel and leave only the contributions of viscous diffusion and advection by the mean velocity. Here we note that the viscous fluxes/wall sources of spanwise vorticity sampled by the stochastic particles are also quite large. The values $\sigma_z^+ \doteq -0.1$ sampled in both the ejection and sweep events for this $Re_\tau = 1000$ simulation are ~ 100 times larger than the mean value $\langle \sigma_z^+ \rangle \doteq -1 \times 10^{-3}$. The very similar temporal profiles in Fig. 16 and Fig. 20 may just reflect a common origin of the spanwise vorticity at height $y^+ \doteq 5$ in the parallel-transport of initially spanwise wall-vorticity by viscous diffusion and by advection due to the mean shear-velocity.

In summary, our results for the sweep rather dramatically contradict the naive idea that vortex lines are approximately “frozen-in” and advected by the flow. If that idea were correct, the spanwise vorticity at the bottom of the depressed vortex lines over the high-speed streak would have been swept down from the interior of the flow. We have found that there is indeed a large non-spanwise vorticity contribution swept down from the interior in the near past, but this contribution is exactly cancelled by an equal and opposite vorticity originating from the wall in that same period. This cancelling contribution from the wall can be plausibly explained to arise from opposite-signed vorticity induced by interactions of the solid wall and the downswept interior vorticity. In contrast to this “old” interior vorticity, the resultant (mainly spanwise) vorticity at the bottom of the depressed vortex line at $y^+ = 5$ is much “younger”, arising from vorticity shed by the wall in the near past and perhaps transported primarily by viscous diffusion.

5. Conclusions and Prospects

We have presented a first concrete application of our Monte Carlo numerical Lagrangian method to channel-flow turbulence, using an online database of a high- Re channel-flow simulation (Graham *et al.* 2016). We have analyzed the Lagrangian vorticity dynamics for two specific events in the near-wall buffer layer, a pair selected as generic examples of an “ejection” and a “sweep”. We find that the growth of vorticity

magnitude due to nonlinear Lagrangian chaos is compensated by viscous cancellation of oppositely-signed vector components, or viscous reconnection in a generalized sense. We may refer to this as “virtual reconnection”, because spanwise anti-parallel components of the Eulerian vorticity field almost never appear in the viscous sublayer but arise by non-trivial Lagrangian dynamics in an average over virtual, stochastic processes. We find also that vortex-lifting from the wall is a highly distributed development in space-time, not an abrupt, discrete event. Consistent with expectations, we find that, at the same distance from the wall, the vorticity in the “sweep” event is older than in the “ejection” event, being birthed at the wall in the more distant past. Surprisingly, however, the greater age is evidenced only by a persistent orthogonal contribution from the wall, which requires many hundreds of viscous times to decay, whereas the vorticity in the ultimate parallel direction is assembled over ~ 100 viscous times in a similar manner for both events.

Future numerical studies should examine many more such events, to verify whether such features are typical and to reach statistically relevant conclusions. Such studies should illuminate the detailed Lagrangian mechanisms of the organized transport of spanwise vorticity away from the wall, which is required for turbulent drag and dissipation. More empirical studies are required even of the Eulerian turbulent vorticity flux, which has been examined much less than momentum transport and Reynolds stress, in order to understand which flow structures sustain the required flow of spanwise vorticity. As emphasized by Brown & Roshko (2012), “The subject stands at the beginning of a new era in which both LES and DNS calculations can provide details of the vorticity field and the fluxes of vorticity (vortex force).” The stochastic Lagrangian methods developed in our work can provide even deeper insight into the dynamics, especially when tightly integrated with the Eulerian picture. We take some further steps in this direction in the following paper of Eyink *et al.* (2020b), which develops a stochastic Lagrangian representation in which the Eulerian vorticity source of Lighthill (1963)-Morton (1984) is incorporated as Neumann boundary conditions for the Helmholtz equation via stochastic particle trajectories that are reflected from the wall.

A remarkable aspect of the stochastic Lagrangian theory is the many unifying features that emerge naturally between classical and quantum fluids. Even based upon our preliminary results, we can make some relevant comparisons with quantum turbulence in superfluids (Barenghi *et al.* 2014). The regime with the closest correspondence to the classical case is two-fluid turbulence in ^4He . Forced flows of superfluid ^4He in the two-fluid regime through smooth-wall tubes and square channels at high Reynolds numbers suffer a pressure-drop in reasonable agreement with classical friction laws (Swanson *et al.* 2000; Fuzier *et al.* 2001) and velocity profiles determined from particle-imaging velocimetry exhibit a near-wall turbulent boundary layer (Xu & Van Sciver 2007). A factor in favor of such classical correspondences is the locking or coupling of the two fluid components (Vinen 2000; Kivotides 2007), but complete understanding of the similar behaviors is still lacking. We believe that the theory developed in this paper and paper I may assist in developing such explanations, because of the several connections it exposes between classical and quantum fluids.

To emphasize this point, we briefly summarize here some of the common features. The Kuz'min (1983)-Oseledets (1989) formulation of Navier-Stokes dynamics in terms of a continuous distribution of infinitesimal vortex-rings is very similar to the intuitive picture of a quantized vortex tangle proposed by Campbell (1972), as a superposition of small vortex rings, which was invoked by Schwarz (1982) to explain intuitively the phase-slip process in superfluid turbulence. See also Kuz'min (1999). Huggins (1994) and Eyink (2008) have already emphasized that constant flux of a conserved vorticity current is necessary for dissipative drag in both classical and superfluid turbulent channel-flows.

The mechanism in quantum turbulence proposed by Schwarz (1988, 1990), based on his vortex-filament simulations, was the cross-stream “ballooning” of ring vortices, which are ultimately driven to annihilate at the wall. As observed by Huggins (1994), an equivalent flux in the classical case results from vorticity creation at the wall and subsequent transport to the channel center, where opposite orientations cancel. Last but not least, Eyink (2010) showed that mean conservation of the stochastic Cauchy invariants and of stochastic circulations (Kelvin Theorem) for incompressible Navier-Stokes solutions arises from particle relabelling symmetry in a stochastic least-action principle. These conservation laws hold, in close analogy to those for ideal Euler equations, although viscosity vitiates the standard “frozen-in” property of vorticity and permits vortex reconnection. Similarly, it has been shown for superfluids, both in the zero-temperature Gross-Pitaevskii model (Nilsen *et al.* 2006) and in strongly rotating, chiral flows (Wiegmann 2019), that the Kelvin theorem holds while simultaneously quantized vortices are *not* frozen into the flow. For the chiral flows the Kelvin theorem is derived as a consequence of particle relabelling symmetry (Wiegmann 2019), while this issue seems open for Gross-Pitaevskii. The contrary finding of Kedia *et al.* (2018) with the Thomas-Fermi approximation neglects the quantum pressure, which is crucial to determine the correct motion of quantized vortices (Nilsen *et al.* 2006) and to permit vortex reconnection (Koplik & Levine 1993).

This underlying unity has importance because fewer differences between classical and quantum fluids appear starker than the differences in reconnection physics. Vortex reconnection is believed to be a crucial element of superfluid turbulence, as discussed in works of Schwarz (1982, 1988, 1990), and we have argued that reconnection in some generalized sense is an essential feature also of classical channel-flow turbulence. In a classical fluid, vortex lines are continuously distributed in space and cannot be unambiguously tracked in time. Vorticity may be attributed a stochastic law of motion and viscous reconnection then results from cancellations in averaging random contributions, just as for the similar case of resistive magnetic reconnection in plasmas (Eyink *et al.* 2013). The vorticity of a superfluid is quantized, on the other hand, and individual segments of vortex-lines are topological defects that may be followed objectively and deterministically. We believe, however, that stochastic laws of motion similar to those for classical viscous fluids will hold also in superfluid turbulence for coherent “bundles” of quantized vortex lines (L’vov *et al.* 2007; Baggaley *et al.* 2012). There seems no reason to doubt that individual vortex lines in a superfluid regime with a Kolmogorov energy spectrum (Nore *et al.* 1997; Barenghi *et al.* 2014) will exhibit “spontaneous stochasticity” due to turbulent Richardson dispersion (Bernard *et al.* 1998; Drivas & Eyink 2017*a*). Because of such explosive dispersion effects and ubiquitous microscopic reconnection, the motion and collective reconnection of vortex bundles in superfluids (Alamri *et al.* 2008) should appear stochastic just as in classical viscous fluids. Stochastic Lagrangian invariants would naturally arise in a dissipative effective action for coarse-grained fields preserving relabelling symmetry (Crossley *et al.* 2017).

Acknowledgements

We are grateful to Joseph Katz, Joseph Klewicki, Demosthenes Kivotides, Charles Meneveau, and Pablo Mininni for useful discussions. We thank Paul Weigmann for informing us of his unpublished work. We acknowledge the NSF grant BigData:OCE-1633124 for support and G.E. also acknowledges the Simons Foundation through Targeted Grant in MPS-663054 for partial support. This research project was conducted using simulation data from the Johns Hopkins Turbulence Database (JHTDB) and scientific computing

services at the Maryland Advanced Research Computing Center (MARCC). Declaration of Interests: The authors report no conflict of interest.

REFERENCES

- ALAMRI, SULTAN Z, YOUNG, ANTHONY J & BARENGHI, CARLO F 2008 Reconnection of superfluid vortex bundles. *Physical review letters* **101** (21), 215302.
- ANDERSON, P. W. 1966 Considerations on the flow of superfluid helium. *Reviews of Modern Physics* **38** (2), 298.
- ANDREOPOULOS, J & AGUI, JH 1996 Wall-vorticity flux dynamics in a two-dimensional turbulent boundary layer. *Journal of Fluid Mechanics* **309**, 45–84.
- BAGGALEY, ANDREW W, BARENGHI, CARLO F, SHUKUROV, ANVAR & SERGEEV, YURI A 2012 Coherent vortex structures in quantum turbulence. *EPL (Europhysics Letters)* **98** (2), 26002.
- BARENGHI, CARLO F, SKRBEK, LADISLAV & SREENIVASAN, KATEPALLI R 2014 Introduction to quantum turbulence. *Proceedings of the National Academy of Sciences* **111** (Supplement 1), 4647–4652.
- BERNARD, DENIS, GAWEDZKI, KRZYSZTOF & KUPIAINEN, ANTTI 1998 Slow modes in passive advection. *Journal of Statistical Physics* **90** (3-4), 519–569.
- BESSE, N. & FRISCH, U. 2017 Geometric formulation of the Cauchy invariants for incompressible Euler flow in flat and curved spaces. *Journal of Fluid Mechanics* **825**, 412–478.
- BORODIN, A. N. & SALMINEN, P. 2015 *Handbook of Brownian Motion - Facts and Formulae*. Birkhäuser Basel.
- BROWN, GARRY L & ROSHKO, ANATOL 2012 Turbulent shear layers and wakes. *Journal of Turbulence* **13**, N51.
- CAMPBELL, LJ 1972 A critical look at a class of critical velocity theories for superfluid he. *Journal of Low Temperature Physics* **8** (1-2), 105–113.
- CAUCHY, A. L. 1815 Sur l'état du fluide à une époque quelconque du mouvement. Mémoires extraits des recueils de l'Académie des sciences de l'Institut de France, Théorie de la propagation des ondes à la surface d'un fluide pesant d'une profondeur indéfinie. (*Extraits des Mémoires présentés par divers savants à l'Académie royale des Sciences de l'Institut de France et imprimés par son ordre*). *Sciences mathématiques et physiques*. Tome I, 1827 Seconde Partie, pp. 33–73.
- CHHIKARA, R. & FOLKS, J. L. 1988 *The Inverse Gaussian Distribution: Theory: Methodology, and Applications*. Taylor & Francis.
- CONSTANTIN, P. & IYER, G. 2008 A stochastic Lagrangian representation of the 3-dimensional incompressible Navier-Stokes equations. *Commun. Pure Appl. Math.* **61**, 330–345.
- CONSTANTIN, P. & IYER, G. 2011 A stochastic-Lagrangian approach to the Navier-Stokes equations in domains with boundary. *The Annals of Applied Probability* **21.4**, 1466–1492.
- CORRSIN, S. 1953 Remarks on turbulent heat transfer. In *Proceedings of the Iowa Thermodynamics Symposium*, American Society for Engineering Education and Society for the Promotion of Engineering Education (U.S.), , vol. 60, pp. 5–30. State University of Iowa, Iowa City.
- CROSSLEY, M., GLORIOSO, P. & LIU, H. 2017 Effective field theory of dissipative fluids. *Journal of High Energy Physics* **2017** (9), 95.
- DRIVAS, THEODORE D & EYINK, GREGORY L 2017a A lagrangian fluctuation–dissipation relation for scalar turbulence. part i. flows with no bounding walls. *Journal of Fluid Mechanics* **829**, 153–189.
- DRIVAS, THEODORE D & EYINK, GREGORY L 2017b A lagrangian fluctuation–dissipation relation for scalar turbulence. part ii. wall-bounded flows. *Journal of Fluid Mechanics* **829**, 236–279.
- EYINK, GREGORY, VISHNIAC, ETHAN, LALESCU, CRISTIAN, ALUIE, HUSSEIN, KANOV, KALIN, BÜRGER, KAI, BURNS, RANDAL, MENEVEAU, CHARLES & SZALAY, ALEXANDER 2013 Flux-freezing breakdown in high-conductivity magnetohydrodynamic turbulence. *Nature* **497** (7450), 466–469.
- EYINK, G. L. 2008 Turbulent flow in pipes and channels as cross-stream inverse cascades of vorticity. *Physics of Fluids* **20** (12), 125101.

- EYINK, G. L. 2010 Stochastic least-action principle for the incompressible Navier–Stokes equation. *Physica D: Nonlinear Phenomena* **239** (14), 1236–1240.
- EYINK, G. L., GUPTA, A. & ZAKI, T. A. 2020a Stochastic Lagrangian dynamics of vorticity for viscous, incompressible fluids. *J. Fluid Mech.* (submitted).
- EYINK, G. L., ZAKI, T. A. & WANG, M. 2020b A stochastic Lagrangian representation of the Lighthill–Morton theory. in preparation.
- FUZIER, S., BAUDOUY, B. & VAN SCIVER, SW 2001 Steady-state pressure drop and heat transfer in he ii forced flow at high reynolds number. *Cryogenics* **41** (5-6), 453–458.
- GRAHAM, J., KANOV, K., YANG, X. I. A., LEE, M., MALAYA, N., LALESCU, C. C., BURNS, R., EYINK, G., SZALAY, A., MOSER, R. D. & MENEVEAU, C. 2016 A web services accessible database of turbulent channel flow and its use for testing a new integral wall model for LES. *Journal of Turbulence* **17** (2), 181–215.
- HUGGINS, E. R. 1970 Energy-dissipation theorem and detailed Josephson equation for ideal incompressible fluids. *Physical Review A* **1** (2), 332.
- HUGGINS, E. R. 1994 Vortex currents in turbulent superfluid and classical fluid channel flow, the Magnus effect, and Goldstone boson fields. *Journal of low temperature physics* **96** (5-6), 317–346.
- JEONG, J. & HUSSAIN, F. 1995 On the identification of a vortex. *Journal of fluid mechanics* **285**, 69–94.
- JIMÉNEZ, JAVIER 2013 Near-wall turbulence. *Physics of Fluids* **25** (10), 101302.
- JIMENEZ, JAVIER, MOIN, P, MOSER, R & KEEFE, L 1988 Ejection mechanisms in the sublayer of a turbulent channel. *The Physics of fluids* **31** (6), 1311–1313.
- JOHNSON, P. L., HAMILTON, S. S., BURNS, R. & MENEVEAU, C. 2017 Analysis of geometrical and statistical features of lagrangian stretching in turbulent channel flow using a database task-parallel particle tracking algorithm. *Physical Review Fluids* **2** (1), 014605.
- JOSEPHSON, B. D. 1962 Possible new effects in superconductive tunnelling. *Physics letters* **1** (7), 251–253.
- KEDIA, HRIDESH, KLECKNER, DUSTIN, SCHEELER, MARTIN W & IRVINE, WILLIAM TM 2018 Helicity in superfluids: Existence and the classical limit. *Physical Review Fluids* **3** (10), 104702.
- KIM, JOHN & HUSSAIN, FAZLE 1993 Propagation velocity of perturbations in turbulent channel flow. *Physics of Fluids A: Fluid Dynamics* **5** (3), 695–706.
- KIM, J., MOIN, P. & MOSER, R. 1987 Turbulence statistics in fully developed channel flow at low Reynolds number. *Journal of fluid mechanics* **177**, 133–166.
- KIVOTIDES, DEMOSTHENES 2007 Relaxation of superfluid vortex bundles via energy transfer to the normal fluid. *Physical Review B* **76** (5), 054503.
- KLEWICKI, JOE, FIFE, PAUL, WEI, TIE & MCMURTRY, PAT 2007 A physical model of the turbulent boundary layer consonant with mean momentum balance structure. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **365** (1852), 823–840.
- KLEWICKI, JC, PRIYADARSHANA, PJA & METZGER, MM 2008 Statistical structure of the fluctuating wall pressure and its in-plane gradients at high reynolds number. *Journal of Fluid Mechanics* **609**, 195–220.
- KOPLIK, JOEL & LEVINE, HERBERT 1993 Vortex reconnection in superfluid helium. *Physical Review Letters* **71** (9), 1375.
- KOUMOUTSAKOS, P. 1999 Vorticity flux control for a turbulent channel flow. *Physics of Fluids* **11** (2), 248–250.
- KUZ’MIN, GA 1999 Vortex momentum density and invariants of the hydrodynamic equations of superfluidity and superconductivity. *Low Temperature Physics* **25** (1), 1–4.
- KUZ’MIN, G. A. 1983 Ideal incompressible hydrodynamics in terms of the vortex momentum density. *Physics Letters A* **96** (2), 88–90.
- LEE, M., MALAYA, N. & MOSER, R. D. 2013 Petascale direct numerical simulation of turbulent channel flow on up to 786K cores. In *SC ’13, Proceedings of the International Conference on High Performance Computing, Networking, Storage and Analysis, Denver, Colorado November 17 - 21, 2013*, Article 61. ACM, New York.
- LIGHTHILL, M. J. 1963 Boundary layer theory. In *Laminar Boundary Layers* (ed. L. Rosenhead), pp. 46–113. Oxford University Press, Oxford.

- L'VOV, VICTOR S, NAZARENKO, SERGEI V & RUDENKO, OLEKSII 2007 Bottleneck crossover between classical and quantum superfluid turbulence. *Physical Review B* **76** (2), 024520.
- LYMAN, FA 1990 Vorticity production at a solid boundary. *Appl. Mech. Rev* **43** (8), 157–158.
- MORTON, B. R. 1984 The generation and decay of vorticity. *Geophysical & Astrophysical Fluid Dynamics* **28** (3-4), 277–308.
- NATAN, A. 2013 Fast 2d peak finder. Matlab File Exchange, <https://www.mathworks.com/matlabcentral/fileexchange/37388-fast-2d-peak-finder>.
- NILSEN, HALVOR M, BAYM, GORDON & PETHICK, CJ 2006 Velocity of vortices in inhomogeneous bose–einstein condensates. *Proceedings of the National Academy of Sciences* **103** (21), 7978–7981.
- NORE, C., ABID, M. & BRACHET, M. E. 1997 Decaying Kolmogorov turbulence in a model of superflow. *Physics of Fluids* **9** (9), 2644–2669.
- ORSZAG, S. A. 1971 On the elimination of aliasing in finite-difference schemes by filtering high-wavenumber components. *Journal of the Atmospheric sciences* **28** (6), 1074–1074.
- OSELEDETS, V. I. 1989 On a new way of writing the Navier-Stokes equation. The Hamiltonian formalism. *Russian Mathematical Surveys* **44** (3), 210.
- PACKARD, R. E. 1998 The role of the Josephson-Anderson equation in superfluid helium. *Reviews of Modern Physics* **70** (2), 641.
- PARK, JAE SUNG, SHEKAR, ASHWIN & GRAHAM, MICHAEL D 2018 Bursting and critical layer frequencies in minimal turbulent dynamics and connections to exact coherent states. *Physical Review Fluids* **3** (1), 014611.
- SCHWARZ, KW 1982 Generation of superfluid turbulence deduced from simple dynamical rules. *Physical Review Letters* **49** (4), 283.
- SCHWARZ, KW 1988 Three-dimensional vortex dynamics in superfluid he 4: Homogeneous superfluid turbulence. *Physical Review B* **38** (4), 2398.
- SCHWARZ, KW 1990 Phase slip and turbulence in superfluid he 4: A vortex mill that works. *Physical review letters* **64** (10), 1130.
- SHENG, J, MALKIEL, E & KATZ, J 2009 Buffer layer structures associated with extreme wall stress events in a smooth wall turbulent boundary layer. *Journal of Fluid Mechanics* **633**, 17–60.
- SWANSON, CHRIS J, DONNELLY, RUSSELL J & IHAS, GARY G 2000 Turbulent pipe flow of he i and he ii. *Physica B: Condensed Matter* **284**, 77–78.
- TAYLOR, G. I. 1932 The transport of vorticity and heat through fluids in turbulent motion. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* **135** (828), 685–702.
- TAYLOR, G. I. 1938 Production and dissipation of vorticity in a turbulent fluid. *Proceedings of the Royal Society of London. Series A-Mathematical and Physical Sciences* **164** (916), 15–23.
- TENNEKES, H. & LUMLEY, J.L. 1972 *A First Course in Turbulence*. Cambridge, MA: MIT Press.
- TUR, A. V. & YANOVSKY, V. V. 1993 Invariants in dissipationless hydrodynamic media. *Journal of Fluid Mechanics* **248**, 67–106.
- VAROQUAUX, ERIC 2015 Anderson's considerations on the flow of superfluid helium: Some offshoots. *Reviews of Modern Physics* **87** (3), 803.
- VINEN, WF 2000 Classical character of turbulence in a quantum liquid. *Physical Review B* **61** (2), 1410.
- WALEFFE, FABIAN 1998 Three-dimensional coherent states in plane shear flows. *Physical Review Letters* **81** (19), 4140.
- WIEGMANN, P 2019 Quantization of hydrodynamics: Rotating superfluid and gravitational anomaly. *arXiv preprint arXiv:1906.03788*.
- XU, T & VAN SCIVER, SW 2007 Particle image velocimetry measurements of the velocity profile in he ii forced flow. *Physics of fluids* **19** (7), 071703.
- ZHAO, H., WU, J.-Z. & LUO, J.-S. 2004 Turbulent drag reduction by traveling wave of flexible wall. *Fluid Dynamics Research* **34** (3), 175.