

Looking down on NF- κ B

Leah M. Williams, Thomas D. Gilmore

Department of Biology, Boston University, Boston, MA 02215, USA

*Corresponding author: Thomas D. Gilmore, Biology Department, Boston University, 5
Cummington Mall, Boston, MA 02215, USA; 617-353-5444 (phone); 617-353-6340
(fax); E-mail: gilmore@bu.edu

24

25 **Abstract**

26 The diversified NF- κ B transcription factor family has been extensively characterized in
27 organisms from flies to humans. However, homologs of NF- κ B and many upstream
28 signaling components have recently been characterized in basal phyla, including Cnidaria
29 (sea anemones, corals, hydras, jellyfish), Porifera (sponges), and single-celled protists
30 including *Capsaspora owczarzaki* and some choanoflagellates. Herein, we review what is
31 known about basal NF- κ Bs and how that knowledge informs on the evolution and
32 conservation of key sequences and domains in NF- κ B, as well as the regulation of NF- κ B
33 activity. The structures and DNA-binding activities of basal NF- κ B proteins resemble
34 mammalian NF- κ B p100 proteins, and their post-translational activation appears to have
35 aspects of both canonical and non-canonical pathways in mammals. Several studies
36 suggest that the single NF- κ B proteins found in some basal organisms have dual roles in
37 development and immunity. Further research on NF- κ B in invertebrates will reveal
38 information about the evolutionary roots of this major signaling pathway, shed light on
39 the origins of regulated innate immunity, and may have relevance to our understanding of
40 the responses of ecologically important organisms to changing environmental conditions
41 and emerging pathogen-based diseases.

42

43

44 Keywords: *NF-kappaB; evolution; development; innate immunity; signal transduction*

45 Running title: NF- κ B in basal invertebrates

INTRODUCTION

The Nuclear Factor- κ B (NF- κ B) superfamily comprises a group of related transcription factors that have been intensively studied for their involvement in development and immunity since their near simultaneous discovery in a retrovirus, flies, and mice almost 35 years ago (1–5). Indeed, there are now approximately 100,000 papers on NF- κ B, the vast majority focusing on mammals, flies, and viruses. However, with the proliferation of genomic and transcriptomic sequencing over the past decade, it has been discovered that many organisms ostensibly less complex than insects have NF- κ B-like genes (6–13). Herein, we critically review what is known about NF- κ B in these more basal organisms, namely cnidarians, poriferans, and protists.

All NF- κ B proteins are related by an N-terminal DNA-binding and dimerization region called the Rel Homology Domain (RHD) containing a nuclear localization sequence (NLS), and the RHD allows them to enter the nucleus, bind to specific DNA sites (“ κ B sites”), and activate or repress transcription of target genes for specified biological outcomes (14). In vertebrates and flies, the NF- κ B superfamily can be divided into two subfamilies: the NF- κ B proteins that consist of vertebrate p100 and p105 and *Drosophila* Relish; and the Rel proteins, which include vertebrate RelA, RelB, and c-Rel, as well as *Drosophila* Dif and Dorsal (14). Thus, flies and vertebrates all contain multiple NF- κ B proteins that, for the most part, show complete combinatorial diversity by forming homodimers and heterodimers, which have distinct DNA target site specificities. The two subfamilies can be distinguished phylogenetically by sequence alignment of their RHDs, and by sequences C-terminal to the RHD. That is, NF- κ B proteins contain C-terminal inhibitory Ankyrin (ANK) repeat domains and Rel proteins contain C-terminal

transactivation domains (Fig. 1). The ANK repeats, either within the NF- κ B proteins themselves or in a separate family of NF- κ B inhibitors (I κ Bs), regulate the subcellular localization of NF- κ Bs by binding to the RHD and sequestering them in the cytoplasm. Activation of the pathway by an appropriate upstream signal results in the degradation of the ANK repeat inhibitor, thus allowing the NF- κ B dimer to enter the nucleus and bind DNA (14, 15). NF- κ B p100 and p105 proteins also contain a C-terminal death domain (DD) that is important for protein-protein interactions with other members of the DD superfamily, which serve as adaptors in signaling pathways and/or to recruit other proteins into signaling complexes (15).

Many evolutionarily conserved receptors can elicit downstream signals to activate NF- κ B translocation to the nucleus. These receptors include Toll-like receptors (TLRs), tumor necrosis factor (TNF) receptors, and interleukin-1 receptors (IL-1Rs) (15), which interact with cytoplasmic adaptor proteins (e.g., MYD88 and MAL) to initiate downstream signaling. Once the pathway is activated, a series of phosphorylation events leads to the degradation of the ANK repeat inhibitor sequences, freeing NF- κ B to translocate from the cytoplasm to the nucleus. In vertebrates, activation of NF- κ B generally happens in one of two ways. In the canonical pathway, activation of an I κ B kinase β (IKK β) complex results in phosphorylation of an independent I κ B protein, and phosphorylated I κ B then undergoes ubiquitination and proteasomal degradation to liberate the NF- κ B dimer. In the non-canonical pathway, activation of the related kinase IKK α leads to phosphorylation of a serine cluster located C-terminal to the ANK repeats on the p100 protein (16). These phosphorylations result in proteasomal processing of the inhibitory ANK repeats on p100 until the proteasome reaches a glycine-rich region

(GRR). At the GRR the proteasome falls off, generating p52, a C-terminally truncated version of p100 that can enter the nucleus (16). Similar pathways exist in flies, except the p100 homolog Relish has no GRR and the truncated, active form of Relish is generated by a site-specific proteolytic cleavage that removes the ANK repeat region (17).

Genomic and transcriptomic sequencing data strongly suggest that NF- κ B was also pervasive through early evolution, based on the presence of NF- κ B-like homologs in many extant organisms basal to flies and vertebrates. NF- κ B homologs have been identified in single-celled organisms such as the protists *Capsaspora owczarzaki* (11) and choanoflagellates (12), which are hypothesized to be the closest living relatives to multicellular animals. NF- κ Bs have also been identified in a variety of basal marine organisms, including poriferans (sponges) and cnidarians (corals, sea anemones, hydras, and jellyfish) (7–10, 13). In most of these basal organisms, there are single NF- κ B proteins, which, as discussed below, most closely resemble NF- κ B subfamily proteins such as Relish and p100. Curiously, no NF- κ B homologs have been found in nematodes (e.g., *Caenorhabditis elegans*), where the pathway appears to have been lost.

As discussed in this review, only recently have several studies investigated how transcription factor NF- κ B and its upstream and downstream pathways function in early-branching organisms. We provide insight into how this highly conserved transcription factor and its regulatory pathway arose, describe domains that have appeared at pivotal points of evolution and biological processes likely controlled by NF- κ B, and discuss areas of basal NF- κ B knowledge that remain unanswered.

STRUCTURES OF BASAL NF- κ B PROTEINS

Many basal eukaryotes have NF- κ B proteins that are structurally and phylogenetically most similar to vertebrate NF- κ B proteins rather than Rel proteins. In most cases, these organisms have single p100-like RHD-ANK repeat NF- κ B proteins. For example, the protist *Capsaspora*, the sponge *Amphimedon queenslandica*, and several cnidarians all have NF- κ Bs with an RHD and C-terminal ANK repeats on the same transcript (8, 18–23). Many ANK repeat-containing proteins are found throughout the genomes of archaea, bacteria, and all plants and animals (24, 25), and therefore, we think it likely that during the evolution of eukaryotes there arose a primitive RHD-only protein, which developed the ability to interact with a pre-existing ANK repeat-containing protein. At some point, these interacting RHD and ANK-repeat genes fused to create the more modern single RHD-ANK-repeat protein.

Nevertheless, some basal organisms have NF- κ B-like proteins without C-terminal ANK repeat domains. Among protists, several choanoflagellates have NF- κ B-like proteins that do not have C-terminal ANK repeat domains (12). Choanoflagellates comprise a sister group to all metazoans, and are part of a diverse group of single-celled, colony-forming organisms that live in waters around the world. Although the genomes of the classically studied choanoflagellates *Monosiga brevicollis* and *Salpingoeca rosetta* do not contain NF- κ B homologs (26), a large-scale transcriptomic sequence analysis of 19 choanoflagellates (12) showed that 12 had transcripts that encode RHD-containing NF- κ B-like sequences (12). However, none of these 12 choanoflagellate NF- κ B-like proteins appears to have C-terminal ANK repeat domains (Fig. 1). Moreover, several choanoflagellate NF- κ B-like proteins also have extended N-terminal domains that are not present in any other NF- κ B proteins (Fig. 1, Table S1). Overall, these 19 choanoflagellate

transcriptomes encode zero to three NF- κ B-like proteins, none of which has C-terminal ANK repeats. Indeed, it is not yet known whether choanoflagellate NF- κ Bs are regulated by an ANK repeat-containing I κ B-like inhibitor. The lack of ANK repeats in choanoflagellate NF- κ B proteins and the wide diversification of NF- κ Bs among choanoflagellates are likely a result of 600 million years of independent evolution among the diverse species of choanoflagellates. In contrast, the protist *Capsaspora* does contain a bipartite RHD-ANK protein (11). Thus, the earliest NF- κ B with an RHD-ANK fusion likely arose in the protist lineage that led to *Capsaspora*, but was excluded or lost in the lineage that led to choanoflagellates.

Although the choanoflagellate and *Capsaspora* NF- κ Bs differ from one another in their overall structures (i.e., especially the absence of a C-terminal ANK-repeat region in choanoflagellates), they share two motifs within their RHDs (Fig. 2, Motif 7 and 10 in teal and yellow) that are not present in any metazoans. Among choanoflagellate NF- κ Bs, there is a great deal of sequence diversity, which is consistent with the overall genetic differences among choanoflagellates wherein the average phylogenetic distance between any pair of choanoflagellates is greater than the phylogenetic distance between sponges and mammals (12). Nevertheless, a Bayesian tree analysis of the 12 choanoflagellate RHDs identified to date clusters them as an outgroup to several fly and vertebrate NF- κ Bs and Rels, which demonstrates a clear divergence of the choanoflagellate NF- κ Bs from the metazoan NF- κ B superfamily (Fig. 2). In general, one finds that NF- κ Bs within a single choanoflagellate species that has multiple NF- κ B proteins (e.g., *Salpingoeca helianthicam*) are highly related (Fig. 2), suggesting that they arose by gene duplication events.

Among poriferans, the fully sequenced genome of the demisponge *Amphimedon queenslandica* encodes a single full-length NF- κ B protein that contains clear features of the human p100 NF- κ B protein: that is, it has an RHD, NLS, GRR, six ANK repeats, and a DD (21, 27) (Fig. 1). However, between the GRR and the ANK repeats, *A. queenslandica* NF- κ B contains a region that has no known function and no homology to any other protein (21, 27) (Fig. 1, yellow). More limited sequencing data indicate that other sponges also contain NF- κ B-like transcripts, which, to date, only contain RHD sequences (28). Many of these sponge NF- κ B-like proteins appear much shorter than the prototypical NF- κ B proteins, but this may simply be due to lack of sequence coverage; for example, *Corticium candelabrum* is reported to have a transcript for NF- κ B that consists of only 86 amino acids that are similar to the beginning of the RHD (28).

All characterized cnidarians appear to contain single NF- κ B proteins, but the overall structures of these NF- κ B proteins have diverged, in some cases, likely by gene-splitting events. Thus, many cnidarians (the anemones *Edwardsella lineata* (23) and *Exaiptasia pallida* (22) and the corals *Acropora digitifera*, *Stylophora pistillata*, *Orbicella faveolata*, and *Pocillopora damicornis* (18–20) have prototypical RHD-ANK repeat bipartite proteins, whereas others have separate RHD and ANK repeat proteins. In the best studied example of the latter, the sea anemone *Nematostella vectensis* has a single 440 amino acid NF- κ B-like protein that is most similar to the processed vertebrate p50/p52 NF- κ B proteins based on phylogenetic analysis of RHD amino acid sequences, intron-exon structure of the RHD, and the presence of a GRR after the RHD (29). Moreover, a separate *N. vectensis* gene encodes an I κ B-like ANK-repeat protein with substantial homology to C-terminal sequences of the ANK-repeat domains of

vertebrate p100/p105 (29). That the separate *N. vectensis* NF- κ B and I κ B genes arose due to a gene splitting event is supported by four lines of evidence: 1) the existence of RHD-ANK repeat bipartite proteins in more basal organisms (e.g., *Capsaspora*, *A. queenslandica*); 2) the presence of remnant GRR sequences at the C terminus of *N. vectensis* NF- κ B; 3) the homology of the separate *N. vectensis* I κ B protein to mammalian NF- κ B protein C-terminal ANK repeats; and 4) the presence of an intact RHD-ANK protein in the closely related anemone *E. lineata* (23). Similar to *N. vectensis*, the hydras *Hydractinia symbiolongicarpus* (30) and *Hydra magnipapillata* (31) and the jellyfish *Aurelia* (13) have separate RHD-only NF- κ B proteins and I κ B-like genes, all of which likely came about due to gene-splitting events, which are common in cnidarians (32, 33).

ACTIVITY AND REGULATION OF BASAL NF- κ Bs

The activity and regulation of NF- κ B proteins in vertebrates are now known in great detail. Many of these properties are conserved in basal NF- κ Bs, however, there are clear exceptions.

Although not formally shown, either biochemically or structurally, it is likely that all basal NF- κ B-like proteins bind conserved DNA sites as homodimers. For example, the NF- κ B- κ B site protein-DNA complexes seen in electrophoretic mobility shift assays using cnidarian and sponge NF- κ B proteins migrate in a manner suggesting that the NF- κ B proteins are dimers (18, 22, 27, 34). Furthermore, based on protein-binding microarrays (PBMs), the NF- κ B proteins of *Capsaspora*, the sponge *A. queenslandica*, and two sea anemones bind to a set of κ B sites that are more similar to the sites bound by

mammalian NF- κ B proteins than by Rel proteins (22, 35) a biochemical finding that is consistent with their overall structural organization and phylogenetic data.

Basal NF- κ B proteins appear to be activators of κ B site-containing promoters. That is, the RHD sequences of sponge and cnidarian NF- κ B proteins function as activators of transcription when expressed in reporter gene assays in yeast and human cells (18, 22, 27, 34). The ability of these basal NF- κ Bs to function as independent activators of transcription is more similar to Relish, which can act as a homodimeric transcriptional activator (17), than to mammalian p50/p52 proteins, which are generally activators of transcription when in heterodimers with a Rel protein (14, 15). No DNA-binding or transcriptional regulatory studies have been performed with choanoflagellate NF- κ Bs.

The most extensive research on basal NF- κ B activity has been done with the NF- κ B protein of the sea anemone *N. vectensis* (Nv-NF- κ B). Curiously, it was found that wild populations of these anemones have two major alleles of Nv-NF- κ B that differ at 10 residues, six of which are in the RHD (36). Two of these variable residues are ones that are predicted to contact DNA, and, as a consequence, one of the allelic Nv-NF- κ B proteins (Nv-NF- κ B-C) binds DNA with an approximately two-fold higher affinity than the other variant (Nv-NF- κ B-S) (37). Despite its reduced DNA-binding activity, Nv-NF- κ B-S activates transcription in reporter gene assays in human cells more effectively than Nv-NF- κ B-C (37). Overall, the PBM-based binding site profiles of both Nv-NF- κ B alleles still largely resemble that of mammalian p50 (22, 35). Interestingly, the DNA-binding site profiles of Nv-NF- κ B-C and Nv-NF- κ B-S are as different from each other as each is from human p50 (22), suggesting that within a given species (i.e., *N. vectensis*)

there can be considerable flexibility in DNA binding site recognition that does not have an obvious effect on the organism's phenotype.

In the mammalian non-canonical pathway, human p100 is phosphorylated by IKK α at a cluster of serine residues that are located C-terminal to the ANK repeats, and this phosphorylation induces proteasomal processing of the C-terminal sequences up to the GRR (16). This cluster of serine residues is conserved in many cnidarian NF- κ Bs, including those of the corals *Pocillopora damicornis*, *Orbicella faveolata*, *Stylophora pistillata*, and *Acropora millepora*, and the sea anemones *Actinia tenebrosa*, *Aulactinia veratra*, and *Aiptasia*, as well as one sponge (*A. queenslandica*) (7, 18–22, 27, 38, 39). Furthermore, phosphorylation and proteasomal processing of NF- κ Bs from two cnidarians (*Aiptasia* and *O. faveolata*) and the sponge *A. queenslandica* can be induced by co-expression of human IKK α and IKK β or the single *Aiptasia* or *O. faveolata* IKKs (18, 22, 27), suggesting that IKK-dependent processing of cnidarian and sponge NF- κ Bs can occur in their natural settings. However, no IKK homolog has yet been identified in the sponge *A. queenslandica*. On the other hand, the *Capsaspora* NF- κ B protein clearly has C-terminal ANK repeats and a GRR, but it does not have any apparent C-terminal IKK target site serine residues and no IKK can be found in the *Capsaspora* genome. Overall, the emergence of IKK proteins appears generally coincident with the presence of regulatory serine residues in the ANK repeat domains of basal NF- κ Bs. Taken together, these findings suggest that phosphorylation-dependent processing of the RHD-ANK repeat proteins was first invented in sponges and that a different form of regulated processing occurs with *Capsaspora* NF- κ B. Of note, the *Drosophila* Relish protein also lacks C-terminal serine residues and a GRR and is processed by a site-specific protease

cleavage event (17), and in some conditions, shortened human p105 NF- κ B proteins can be generated by ribosomal stalling (40).

For the *N. vectensis* NF- κ B protein, which lacks C-terminal ANK repeats, it was shown that the independent Nv-I κ B could directly bind to Nv-NF- κ B and that this Nv-I κ B could be phosphorylated by an Nv-IKK-related kinase (34). Thus, aspects of canonical signaling may also be present in cnidarians.

Proteasomal-mediated processing has not been directly demonstrated to induce nuclear translocation of any basal NF- κ B in its natural setting. However, in a reconstituted mammalian cell system, IKK-induced processing of the anemone *Aiptasia* NF- κ B can be blocked by a proteasome inhibitor (22). In addition, when expressed in chicken tissue culture cells, the full-length RHD-ANK proteins of the coral *O. faveolata*, the anemone *Aiptasia*, and the sponge *A. queenslandica* are located in the cytoplasm, and deletion of the ANK repeat domains of these NF- κ Bs causes them to localize to the nucleus (18, 22, 27), suggesting that C-terminal processing would do the same in their host organisms. Similarly, in these same assays, the naturally truncated anemone Nv-NF- κ B protein localizes to the nucleus, but it is sequestered in the cytoplasm when Nv-I κ B is co-expressed (34). Of note, although the C-terminal ANK repeat sequences block nuclear translocation and reporter gene activation by the sponge NF- κ B protein, the ANK repeats do not inhibit NF- κ B's *in vitro* DNA-binding activity (27), unlike what is seen with NF- κ Bs of cnidarians and other higher metazoans. Nothing is known about the subcellular localization or regulation of the choanoflagellate NF- κ B proteins, which lack a clear NLS as well as C-terminal ANK repeats.

Notwithstanding the above-described reconstitution experiments conducted with cnidarian and sponge NF- κ Bs in vertebrate cell systems, it is not known whether or how regulated processing of basal NF- κ Bs occurs in any natural setting. Indeed, in both the anemone *Aiptasia* (22) and one demosponge (27), most NF- κ B protein is largely processed and nuclear in the absence of any known stimulus. Furthermore, certain treatments (e.g., bleaching in *Aiptasia* and LPS treatment of *O. faveolata* coral tissue) may induce NF- κ B activation by inducing transcriptional upregulation of NF- κ B pathway genes rather than C-terminal processing (18, 22).

BASAL NF- κ Bs LIKELY HAVE ROLES IN DEVELOPMENT AND IMMUNITY

One of the most important questions regarding NF- κ B is what biological processes are regulated by this transcription factor in cnidarians, sponges, and protists? At least in cnidarians and sponges, there is evidence that NF- κ B has roles in both early development and adult immunity (Table 1).

Role of NF- κ B in sponge and cnidarian development

In the sponge *A. queenslandica* and several cnidarians, NF- κ B transcripts are expressed in the early embryo. In *A. queenslandica*, fluorescent *in situ* staining has shown that NF- κ B transcripts are broadly expressed throughout the embryo after cleavage, and are particularly strong in granular cells, which eventually become the outer layer of the developing embryo (21). Similarly, some cnidarians express NF- κ B transcripts in the early embryo and juvenile larvae (41-46). In the anemone *N. vectensis*, NF- κ B mRNA expression is seen as early as one hour post fertilization (hpf) (43-46), and nuclear NF- κ B

protein was detected as early as the late gastrula stage (42). Morpholino-based knockdown of NF- κ B in the developing *N. vectensis* embryo led to a failure to develop cnidocytes at the juvenile polyp stage (42). Cnidocytes are a Cnidaria-specific cell type that is involved in a variety of sensing, prey capture, and perhaps defense roles (47). In addition, morpholino knockdown of the single NF- κ B-inducing Toll-like receptor (TLR) transcript also led to an early developmental defect in *N. vectensis* (48).

Role of NF- κ B in juvenile and adult sponges and cnidarians

NF- κ B transcripts and proteins have also been detected in several cell types in developed sponge tissues. Whole-mount *in situ* hybridization of juvenile sponge tissue showed that NF- κ B transcripts are present in flask cells, which are large ciliated cells that express a range of genes whose orthologues play roles in eumetazoan neurons (21). Two recent studies (49, 50) performed single cell sequencing on different adult sponge cell types, and they reported that NF- κ B transcripts were primarily expressed in two cell types: choanocytes, which are cells that reside in the chambers that allow the sponge to filter feed, and archaeocytes, which are motile phagocytic cells that are inside the mesoglea. In a black encrusting sponge (*Cliona sp.*), nuclear NF- κ B protein was detected in several scattered cells throughout the animal (27). Furthermore, anti-NF- κ B Western blots of whole tissue extracts from this *Cliona sp.* contained proteins similar in size to the NF- κ B proteins of *A. queenslandica*. Treatment of this sponge tissue with lipopolysaccharide (LPS, a potent TLR-to-NF- κ B inducer in mammals) produced further processing of the putative full-length NF- κ B protein and an increase in κ B-site binding activity. Similarly, treatment of two Mediterranean species of sponges, *Aplysina*

aerophoba and *Dysidea avara* with an immunogenic cocktail of LPS and peptidoglycan led to the upregulation of immune-related receptors involved in signaling to NF- κ B (51).

Taken together, the above studies suggest that NF- κ B plays a role in a select number of cells involved in sponge immunity. In one sponge, NF- κ B protein appears to be constitutively in its processed active state and in the nucleus of select cells (27). Whether this constitutive nuclear localization of NF- κ B is the result of a chronic processing of this protein in the sponge or represents a naturally truncated isoform that is preferentially expressed has not been determined.

Several studies have suggested that NF- κ B also has a role in immunity in adult cnidarians. In the anemone *N. vectensis*, cytoplasmic NF- κ B and I κ B are expressed in a subset of cnidocytes in the body column of juvenile and adult anemones (42). NF- κ B protein is also highly expressed in cnidocytes that are present in circulating multicellular bodies called nematosomes, which also express high levels of TLR and c-GAS-STING innate immune signaling components that are upstream of NF- κ B (48, 52). Those results, along with the ability of nematosomes to take up bacteria (48, 52), suggest that the nematosome is a primitive circulating immune organ in some anemones.

Aiptasia is a tropical anemone that has been used as a model for cnidarian symbiosis with algal dinoflagellates in the family Symbiodiniaceae (53). Infection of both larval and adult *Aiptasia* with algal symbionts of certain strains results in down-regulation of NF- κ B transcripts, protein, and DNA-binding activity (22, 54, 55).

Conversely, induction of loss-of-symbiosis with either chemical or heat treatment results in increased levels of NF- κ B expression and activity (22), and *Aiptasia* lacking symbionts are more resistant to bacterial infection than ones with symbionts (55). These results have

led to the hypothesis that suppression of NF- κ B-directed immunity is required for the establishment of algal symbiosis in some cnidarians, similar to what has been found in amphibians (56). It is noteworthy that the symbiont-modulated changes in NF- κ B activity in *Aiptasia* occurred primarily at the level of expression (22), and not by post-translational induction as generally seen in vertebrate systems. That is, both symbiotic and aposymbiotic *Aiptasia* had nuclear NF- κ B-staining cells, but NF- κ B-staining cells were substantially increased in *Aiptasia* in which symbiont loss had occurred (22).

The research on NF- κ B in *Aiptasia* is of high interest because corals host the same family of algal symbionts and because ocean warming-induced loss of symbiosis (“bleaching”), as well as microbial pathogen infections, are causing large-scale loss of coral reef health (53). Several studies have also investigated NF- κ B and NF- κ B pathway gene expression in corals, and such studies further suggest a role for NF- κ B in immunity and symbiosis/dysbiosis (Table 1). For example, transcripts encoding NF- κ B and its signaling components have been identified in transcriptomes from bleached *O. faveolata* (57), and treatment of *O. faveolata* tissue with LPS resulted in differential expression of genes in a manner that indicated activation of the NF- κ B pathway (18). Furthermore, NF- κ B mRNA was substantially upregulated in the coral *Acropora palmata* following extended exposure to elevated water temperature (58). While another study found that NF- κ B transcripts were only transiently induced shortly after heat treatment in the coral *Acropora hyacinthus* (59). Thus, NF- κ B may be affected in different ways in different cnidarians undergoing heat stress-induced changes. Additionally, some studies have shown an increase in transcripts encoding members of the NF- κ B pathway in corals with

microbial diseases (18, 60), further suggesting that NF- κ B has an immune-related role in cnidarians.

Several lines of evidence suggest that TLR and TLR-like pathways are upstream activators of NF- κ B in cnidarians. First, homologs of TLR-to-NF- κ B pathway signaling are present in most cnidarians (6-8, 13, 18, 19, 29, 32, 34). Second, *Hydra* and *N. vectensis* TLR-like proteins have a conserved ability to activate NF- κ B when ectopically expressed in human cells (31, 48). Third, as mentioned above, TLR and NF- κ B proteins are expressed in many of the same cnidocytes in adult *N. vectensis* (48). Fourth, treatment of *O. faveolata* coral tissue with the mammalian TLR ligand LPS can induce NF- κ B pathway gene expression (18). Indeed, TLR-to-NF- κ B signaling has been proposed to play a role in embryonic development (42, 48), immunity (18, 31, 38, 48, 60), and regeneration (61) in cnidarians. Nevertheless, TLR is not the sole receptor that can signal to NF- κ B in mammals that is also present in cnidarians. For example, cnidarians also possess homologs to TNF receptors and their downstream components (29), which are extensively characterized activators of NF- κ B in vertebrates. However, the ability of cnidarian TNF receptors to activate NF- κ B in cnidarians has not been investigated.

The biological role of NF- κ B in single-celled organisms remains unknown (Table 1). However, one may speculate that biological processes controlled by NF- κ B in protists are quite different than in multicellular animals, given that protists have no innate or adaptive immunity. Thus, if NF- κ B plays a role in immunity in protists, it may do so in an unexpected manner. Or, in the case of *Capsaspora*, which is a symbiont of the snail *B. galbrata*, NF- κ B may modulate either protist or host immunity to facilitate symbiosis, similar to what has been proposed for *Aiptasia*-*Symbiocinicae* symbiosis. Alternatively,

NF- κ B could play a “developmental” role in the solitary vs. colony states seen in many protists, including choanoflagellates. Finally, NF- κ B may be involved in some protist-specific process, for example, one species of choanoflagellate has been shown to be induced to sexually reproduce upon stimulation with bacterial components (62).

CONCLUSIONS AND FUTURE PROSPECTS

Since its discovery, NF- κ B has been studied as a prominent player in many biological processes in vertebrate and fly systems. Nonetheless, until a decade ago, NF- κ B was not known to be present in the genomes of organisms basal to insects. The discovery of NF- κ Bs in single-celled protists has led to improved hypotheses for how this transcription factor evolved (Fig. 3). That is, we propose that prior to the rise of holozoan life, ANK repeats were present in bacteria and archaea genomes. Some ancestral organism developed a primordial RHD-only protein that eventually fused to an ANK repeat protein, resulting in the modern-day bipartite NF- κ B protein. RHDs then diversified in choanoflagellates, while metazoans retained the full-length NF- κ B fusion protein and also developed separate NF- κ B/Rel and I κ B proteins, likely from gene splitting and duplication events (Fig. 3).

Several studies have now demonstrated that the NF- κ B proteins of evolutionarily basal organisms have many of the same structural features, activities, modes of regulation, and biological effects that are found in the expanded set of NF- κ B and Rel proteins of more complex organisms. For example, ANK repeat regulation of the NF- κ B RHD is even found in protists, and NF- κ B is likely to have dual roles in development and immunity in organisms from sponges to humans. Nevertheless, there are clear differences

in basal NF- κ Bs. Particularly intriguing is the finding that NF- κ B proteins appear to be constitutively processed and active in some sponges and anemones, and that induced activation of the NF- κ B pathway may be at the transcriptional level in some of these same basal organisms.

Studies on the evolution of NF- κ B have only scratched the surface of what is yet to be discovered. Future studies will likely seek to identify the most primitive biological processes controlled by NF- κ B in protists. Moreover, the identification of NF- κ B target genes and the development of genetic systems to study gene function will certainly reveal much about the function of NF- κ B in cnidarians and sponges. In particular, the identification of immune response gene targets of NF- κ B in basal organisms may provide insights into novel microbials, as well as reveal information about molecular processes underlying global ecological crises of marine invertebrates, such as coral bleaching and microbial pathogenesis. Looking backwards at the evolutionary history of NF- κ B will no doubt bring knowledge to our understanding of ancient immunity, especially in those organisms that are environmentally sensitive and are situated at the base of multicellular life.

Materials and Methods

Phylogenetic analysis.

The RHD sequences of NF- κ B from cnidarians, poriferans, and choanoflagellate organisms were compared phylogenetically to annotated vertebrate NF- κ B and Rel proteins and rooted with the predicted RHD of *Capsaspora owczarzaki* NF- κ B. Details on databases and sequence acquisition can be found in Table S2. Conserved motifs from

MEME analysis were truncated based on motif predictions (Table S2), were aligned by Clustal Omega (63) for Bayesian Analysis (Fig. 1). Other phylogenetic analyses showed similar results (data not shown).

Acknowledgments

We thank Anvitha Addanki for help with the analysis of choanoflagellate NF- κ B proteins. The authors' research on the evolution and basal functions of NF- κ B was supported by the following National Science Foundation grants (to T.D.G.): MCB-0924749, IOS-1557804, and IOS-1937650. L.M.W. was supported by an NSF Graduate Research Fellowship.

The authors declare no conflict of interest.

References

1. Gilmore TD, Temin HM. 1986. Different localization of the product of the v-rel oncogene in chicken fibroblasts and spleen cells correlates with transformation by REV-T. *Cell* 44:791–800.
2. Gilmore TD. 1990. NF- κ B, KBF1, dorsal, and related matters. *Cell* 62:841–843.
3. Steward R. 1987. Dorsal, an embryonic polarity gene in *Drosophila*, is homologous to the vertebrate proto-oncogene, c-rel. *Science* 238:692–694.
4. Ghosh S, Gifford AM, Riviere LR, Tempst P, Nolan GP, Baltimore D. 1990. Cloning of the p50 DNA binding subunit of NF- κ B: homology to rel and dorsal. *Cell* 62:1019–1029.
5. Kieran M, Blank V, Logeat F, Vandekerckhove J, Lottspeich F, Le Bail O, Urban MB, Kourilsky P, Baeuerle PA, Israël A. 1990. The DNA binding subunit of NF- κ B is identical to factor KBF1 and homologous to the *rel* oncogene product. *Cell* 62:1007–1018.
6. Gilmore TD, Wolenski FS. 2012. NF- κ B: where did it come from and why? *Immunol Rev* 246:14–35.
7. Baumgarten S, Simakov O, Esherrick LY, Liew YJ, Lehnert EM, Michell CT, Li Y, Hambleton EA, Guse A, Oates ME, Gough J, Weis VM, Aranda M, Pringle JR, Voolstra CR. 2015. The genome of *Aiptasia*, a sea anemone model for coral symbiosis. *Proc Natl Acad Sci USA* 112:11893–11898.

- 468 8. Shinzato C, Shoguchi E, Kawashima T, Hamada M, Hisata K, Tanaka M, Fujie M,
469 Fujiwara M, Koyanagi R, Ikuta T, Fujiyama A, Miller DJ, Satoh N. 2011. Using the
470 *Acropora digitifera* genome to understand coral responses to environmental change.
471 Nature 476:320–323.
- 472 9. Putnam NH, Srivastave M, Hellsten U, Dirks B, Chapman J, Salamov A, Terry A,
473 Shapiro H, Lindquist E, Kapitono VV, Jurka J, Genikhovich G, Grigoriev IV, Lucas
474 SM, Steele RE, Finnerty JR, Technau U, Martindale MQ, Rokhsar DS. 2007. Sea
475 anemone genome reveals ancestral eumetazoan gene repertoire and genomic
476 organization. Science 317:86-94.
- 477 10. Srivastava M, Simakov O, Chapman J, Fahey B, Gauthier MEA, Mitros T, Richards
478 GS, Conaco C, Dacre M, Hellsten U, Larroux C, Putnam NH, Stanke M, Adamska
479 M, Darling A, Degnan SM, Oakley TH, Plachetzki DC, Zhai Y, Adamski M,
480 Calcino A, Cummins SF, Goodstein DM, Harris C, Jackson DJ, Leys SP, Shu S,
481 Woodcroft BJ, Vervoort M, Kosik KS, Manning G, Degnan BM, Rokhsar DS.
482 2010. The *Amphimedon queenslandica* genome and the evolution of animal
483 complexity. Nature 466:720–726.
- 484 11. Suga H, Chen Z, Mendoza A de, Sebé-Pedrós A, Brown MW, Kramer E, Carr M,
485 Kerner P, Vervoort M, Sánchez-Pons N, Torruella G, Derelle R, Manning G, Lang
486 BF, Russ C, Haas BJ, Roger AJ, Nusbaum C, Ruiz-Trillo I. 2013. The *Capsaspora*
487 genome reveals a complex unicellular prehistory of animals. Nat Commun 4:1–9.
- 488 12. Richter DJ, Fozouni P, Eisen MB, King N. 2018. Gene family innovation,
489 conservation and loss on the animal stem lineage. eLife 7:e34226.

- 490 13. Gold DA, Katsuki T, Li Y, Yan X, Regulski M, Ibberson D, Holstein T, Steele RE,
491 Jacobs DK, Greenspan RJ. 2019. The genome of the jellyfish *Aurelia* and the
492 evolution of animal complexity. *Nat Ecol Evol* 3:96–104.
- 493 14. Gilmore TD. 2006. Introduction to NF- κ B: players, pathways, perspectives.
494 *Oncogene* 25:6680–6684.
- 495 15. Hayden MS, Ghosh S. 2008. Shared principles in NF- κ B signaling. *Cell* 132:344.362.
- 496 16. Sun S-C. 2011. Non-canonical NF- κ B signaling pathway. *Cell Res* 21:71–85.
- 497
- 498 17. Stöven S, Silverman N, Junell A, Hedengren-Olcott M, Erturk D, Engström Y,
499 Maniatis T, Hultmark D. 2003. Caspase-mediated processing of the *Drosophila* NF-
500 κ B factor Relish. *Proc Natl Acad Sci USA* 100:5991–5996.
- 501 18. Williams LM, Fuess LE, Brennan JJ, Mansfield KM, Salas-Rodriguez E, Welsh J,
502 Awtry J, Banic S, Chacko C, Chezian A, Dowers D, Estrada F, Hsieh Y-H, Kang J,
503 Li W, Malchiodi Z, Malinowski J, Matuszak S, McTigue T, Mueller D, Nguyen B,
504 Nguyen M, Nguyen P, Nguyen S, Njoku N, Patel K, Pellegrini W, Pliakas T, Qadir
505 D, Ryan E, Schiffer A, Thiel A, Yunes SA, Spilios KE, Pinzón C JH, Mydlarz LD,
506 Gilmore TD. 2018. A conserved toll-like receptor-to-NF- κ B signaling pathway in
507 the endangered coral *Orbicella faveolata*. *Dev Comp Immunol*.
- 508 19. Voolstra CR, Li Y, Liew YJ, Baumgarten S, Zoccola D, Flot J-F, Tambutté S,
509 Allemand D, Aranda M. 2017. Comparative analysis of the genomes of *Stylophora*
510 *pistillata* and *Acropora digitifera* provides evidence for extensive differences
511 between species of corals. *Sci Rep* 7:1–14.

- 512 20. Cunning R, Bay RA, Gillette P, Baker AC, Traylor-Knowles N. 2018. Comparative
513 analysis of the *Pocillopora damicornis* genome highlights role of immune system in
514 coral evolution. *Sci Rep* 8:1–10.
- 515 21. Gauthier M, Degnan BM. 2008. The transcription factor NF- κ B in the demosponge
516 *Amphimedon queenslandica*: insights on the evolutionary origin of the Rel
517 homology domain. *Dev Genes Evol* 218:23–32.
- 518 22. Mansfield KM, Carter NM, Nguyen L, Cleves PA, Alshanbayeva A, Williams LM,
519 Penvose AR, Crowder C, Finnerty JR, Gilmore TD, Siggers TW, Weis VM. 2017.
520 Transcription factor NF- κ B is modulated by symbiotic status in a sea anemone
521 model of cnidarian bleaching. *Sci Rep* 7:16025.
- 522 23. Stefanik DJ, Lubinski TJ, Granger BR, Byrd AL, Reitzel AM, DeFilippo L, Lorenc
523 A, Finnerty JR. 2014. Production of a reference transcriptome and transcriptomic
524 database (EdwardsiellaBase) for the lined sea anemone, *Edwardsiella lineata*, a
525 parasitic cnidarian. *BMC Genomics* 15:71.
- 526 24. Al-Khodori S, Price CT, Kalia A, Abu Kwaik Y. 2010. Functional diversity of
527 ankyrin repeats in microbial proteins. *Trends Microbiol* 18:132–139.
- 528 25. Jernigan KK, Bordenstein SR. 2014. Ankyrin domains across the Tree of Life. *PeerJ*
529 2:e264.
- 530 26. Hoffmeyer TT, Burkhardt P. 2016. Choanoflagellate models - *Monosiga brevicollis*
531 and *Salpingoeca rosetta*. *Curr Opin Genet Dev* 39:42–47.

- 532 27. Williams LM, Inge MM, Mansfield KM, Rasmussen A, Afghani J, Agrba M, Albert
533 C, Andersson C, Babaei M, Babaei M, Bagdasaryants A, Bonilla A, Browne A,
534 Carpenter S, Chen T, Christie B, Cyr A, Dam K, Dulock N, Erdene G, Esau L,
535 Esonwune S, Hanchate A, Huang X, Jennings T, Kasabwala A, Kehoe L, Kobayashi
536 R, Lee M, LeVan A, Liu Y, Murphy E, Nambiar A, Olive M, Patel D, Pavesi F,
537 Petty CA, Samofalova Y, Sanchez S, Stejskal C, Tang Y, Yapo A, Cleary JP, Yunes
538 SA, Siggers T, Gilmore TD. 2020. Transcription factor NF- κ B in a basal metazoan,
539 the sponge, has conserved and unique sequences, activities, and regulation. *Dev*
540 *Comp Immunol* 103559.
- 541 28. Riesgo A, Farrar N, Windsor PJ, Giribet G, Leys SP. 2014. The analysis of eight
542 transcriptomes from all poriferan classes reveals surprising genetic complexity in
543 sponges. *Mol Biol Evol* 31:1102–1120.
- 544 29. Sullivan JC, Kalaitzidis D, Gilmore TD, Finnerty JR. 2007. Rel homology domain-
545 containing transcription factors in the cnidarian *Nematostella vectensis*. *Dev Genes*
546 *Evol* 217:63–72.
- 547 30. Zárate-Potes A, Ocampo ID, Cadavid LF. 2019. The putative immune recognition
548 repertoire of the model cnidarian *Hydractinia symbiolongicarpus* is large and
549 diverse. *Gene* 684:104–117.
- 550 31. Franzenburg S, Fraune S, Künzel S, Baines JF, Domazet-Lošo T, Bosch TCG. 2012.
551 MyD88-deficient Hydra reveal an ancient function of TLR signaling in sensing
552 bacterial colonizers. *Proc Natl Acad Sci* 109:19374–19379.

- 553 32. Chapman JA, Kirkness EF, Simakov O, Hampson SE, Mitros T, Weinmaier T, Rattei
554 T, Balasubramanian PG, Borman J, Busam D, Disbennett K, Pfannkoch C, Sumin
555 N, Sutton GG, Viswanathan LD, Walenz B, Goodstein DM, Hellsten U, Kawashima
556 T, Prochnik SE, Putnam NH, Shu S, Blumberg B, Dana CE, Gee L, Kibler DF, Law
557 L, Lindgens D, Martinez DE, Peng J, Wigge PA, Bertulat B, Guder C, Nakamura Y,
558 Ozbek S, Watanabe H, Khalturin K, Hemmrich G, Franke A, Augustin R, Fraune S,
559 Hayakawa E, Hayakawa S, Hirose M, Hwang JS, Ikeo K, Nishimiya-Fujisawa C,
560 Ogura A, Takahashi T, Steinmetz PRH, Zhang X, Aufschnaiter R, Eder M-K, Gorny
561 A-K, Salvenmoser W, Heimberg AM, Wheeler BM, Peterson KJ, Böttger A,
562 Tischler P, Wolf A, Gojobori T, Remington KA, Strausberg RL, Venter JC,
563 Technau U, Hobmayer B, Bosch TCG, Holstein TW, Fujisawa T, Bode HR, David
564 CN, Rokhsar DS, Steele RE. 2010. The dynamic genome of *Hydra*. *Nature*
565 464:592–596.
- 566 33. Gacesa R, Chung R, Dunn SR, Weston AJ, Jaimes-Becerra A, Marques AC,
567 Morandini AC, Hranueli D, Starcevic A, Ward M, Long PF. 2015. Gene
568 duplications are extensive and contribute significantly to the toxic proteome of
569 nematocysts isolated from *Acropora digitifera* (Cnidaria: Anthozoa: Scleractinia).
570 *BMC Genomics* 16:774.
- 571 34. Wolenski FS, Garbati MR, Lubinski TJ, Traylor-Knowles N, Dresselhaus E, Stefanik
572 DJ, Goucher H, Finnerty JR, Gilmore TD. 2011. Characterization of the core
573 elements of the NF- κ B signaling pathway of the sea anemone *Nematostella*
574 *vectensis*. *Mol Cell Biol* 31:1076–1087.

- 575 35. Ryzhakov G, Teixeira A, Saliba D, Blazek K, Muta T, Ragoussis J, Udalova IA.
576 2013. Cross-species analysis reveals evolving and conserved features of the nuclear
577 factor κ B (NF- κ B) proteins. J Biol Chem 288:11546–11554.
- 578 36. Sullivan JC, Wolenski FS, Reitzel AM, French CE, Traylor-Knowles N, Gilmore TD,
579 Finnerty JR. 2009. Two alleles of NF- κ B in the sea anemone *Nematostella vectensis*
580 are widely dispersed in nature and encode proteins with distinct activities. PloS One
581 4:e7311.
- 582 37. Wolenski FS, Chandani S, Stefanik DJ, Jiang N, Chu E, Finnerty JR, Gilmore TD.
583 2011. Two polymorphic residues account for the differences in DNA binding and
584 transcriptional activation by NF- κ B proteins encoded by naturally occurring alleles
585 in *Nematostella vectensis*. J Mol Evol 73:325–336.
- 586 38. Anderson DA, Walz ME, Weil E, Tonellato P, Smith MC. 2016. RNA-Seq of the
587 Caribbean reef-building coral *Orbicella faveolata* (Scleractinia-Merulinidae) under
588 bleaching and disease stress expands models of coral innate immunity. PeerJ
589 4:e1616.
- 590 39. Ying H, Hayward DC, Cooke I, Wang W, Moya A, Siemering KR, Sprungala S, Ball
591 EE, Forêt S, Miller DJ. 2019. The whole-genome sequence of the coral *Acropora*
592 *millepora*. Genome Biol Evol 11:1374–1379.
- 593 40. Lin L, DeMartino GN, Greene WC. 1998. Cotranslational biogenesis of NF- κ B p50
594 by the 26S proteasome. Cell 92:819–828.

- 595 41. Siboni N, Abrego D, Motti CA, Tebben J, Harder T. 2014. Gene expression patterns
596 during the early stages of chemically induced larval metamorphosis and settlement
597 of the coral *Acropora millepora*. PLOS ONE 9:e91082.
- 598 42. Wolenski FS, Bradham CA, Finnerty JR, Gilmore TD. 2013. NF- κ B is required for
599 cnidocyte development in the sea anemone *Nematostella vectensis*. Dev Biol
600 373:205–215.
- 601 43. Warner JF, Guerlais V, Amiel AR, Johnston H, Nedoncelle K, Röttinger E. 2018.
602 NvERTx: a gene expression database to compare embryogenesis and regeneration in
603 the sea anemone *Nematostella vectensis*. Development 145:dev162867.
- 604 44. Fischer AH, Mozzherin D, Eren AM, Lans KD, Wilson N, Cosentino C, Smith J.
605 2014. SeaBase: a multispecies transcriptomic resource and platform for gene
606 network interface. Integr Comp Biol 54:250-263.
- 607 45. Tulin S, Aguiar D, Istrail S, Smith J. 2013. A quantitative reference transcriptome for
608 *Nematostella vectensis* early embryonic development: a pipeline for *de novo*
609 assembly in emerging model systems. Evodevo 4:16.
- 610 46. Helm RR, Siebert S, Tulin S, Smith J, Dunn CW. 2013. Characterization of
611 differential transcript abundance through time during *Nematostella vectensis*
612 development. BMC Genomics 14:266.
- 613 47. Babonis LS, Martindale MQ. 2014. Old cell, new trick? Cnidocytes as a model for
614 the evolution of novelty. Integr Comp Biol 54:714–722.

- 615 48. Brennan JJ, Messerschmidt JL, Williams LM, Matthews BJ, Reynoso M, Gilmore
616 TD. 2017. Sea anemone model has a single Toll-like receptor that can function in
617 pathogen detection, NF- κ B signal transduction, and development. Proc Natl Acad
618 Sci USA 201711530.
- 619 49. Sogabe S, Hatleberg WL, Kocot KM, Say TE, Stoupin D, Roper KE, Fernandez-
620 Valverde SL, Degnan SM, Degnan BM. 2019. Pluripotency and the origin of animal
621 multicellularity. Nature 570:519-522.
- 622 50. Musser JM, Schippers KJ, Nickel M, Mizzon G, Kohn AB, Pape C, Hammel JU,
623 Wolf F, Liang C, Hernández-Plaza A, Achim K, Schieber NL, Francis WR, R SV,
624 Kling S, Renkert M, Feuda R, Gaspar I, Burkhardt P, Bork P, Beck M, Kreshuk A,
625 Wörheide G, Huerta-Cepas J, Schwab Y, Moroz LL, Arendt D. 2019. Profiling
626 cellular diversity in sponges informs animal cell type and nervous system evolution.
627 bioRxiv <https://doi.org/10.1101/758276>.
- 628 51. Pita L, Hoepfner M, Ribes M, Hentschel U. 2018. Differential expression of immune
629 receptors in two marine sponges upon exposure to microbial-associated molecular
630 patterns. Sci Rep 8:16081.
- 631 52. Babonis LS, Martindale MQ, Ryan JF. 2016. Do novel genes drive morphological
632 novelty? An investigation of the nematosomes in the sea anemone *Nematostella*
633 *vectensis*. BMC Evol Biol 16:114.
- 634 53. Weis VM. 2008. Cellular mechanisms of cnidarian bleaching: stress causes the
635 collapse of symbiosis. J Exp Biol 211:3059–3066.

- 636 54. Wolfowicz I, Baumgarten S, Voss PA, Hambleton EA, Voolstra CR, Hatta M, Guse
637 A. 2016. Aiptasia sp. larvae as a model to reveal mechanisms of symbiont selection
638 in cnidarians. *Sci Rep* 6:1–12.
- 639 55. Mansfield KM, Cleves PA, Vlack EV, Kriefall NG, Benson BE, Camacho DJ,
640 Hemond O, Pedroza M, Siggers T, Pringle JR, Davies SW, Gilmore TD. 2019.
641 Varied effects of algal symbionts on transcription factor NF- κ B in a sea anemone
642 and a coral: possible roles in symbiosis and thermotolerance. *bioRxiv*
643 <https://doi.org/10.1101/640177>.
- 644 56. Burns JA, Zhang H, Hill E, Kim E, Kerney R. 2017. Transcriptome analysis
645 illuminates the nature of the intracellular interaction in a vertebrate-algal symbiosis.
646 *eLife* 6:e22054.
- 647 57. Pinzón JH, Kamel B, Burge CA, Harvell CD, Medina M, Weil E, Mydlarz LD. 2015.
648 Whole transcriptome analysis reveals changes in expression of immune-related
649 genes during and after bleaching in a reef-building coral. *R Soc Open Sci* 2:140214.
- 650 58. DeSalvo MK, Sunagawa S, Voolstra CR, Medina M. 2010. Transcriptomic responses
651 to heat stress and bleaching in the elkhorn coral *Acropora palmata*. *Mar Ecol Prog*
652 *Ser.* 17:3952-3971
- 653 59. Traylor-Knowles N, Rose NH, Sheets EA, Palumbi SR. 2017. Early transcriptional
654 responses during heat stress in the coral *Acropora hyacinthus*. *Biol Bull* 232:91–
655 100.

- 656 60. Fuess LE, C JHP, Weil E, Grinshpon RD, Mydlarz LD. 2017. Life or death: disease-
657 tolerant coral species activate autophagy following immune challenge. Proc R Soc B
658 284:20170771.
- 659 61. Wenger Y, Buzgariu W, Reiter S, Galliot B. 2014. Injury-induced immune responses
660 in *Hydra*. Semin Immunol 26:277–294.
- 661 62. Woznica A, Gerdt JP, Hulett RE, Clardy J, King N. 2017. Mating in the closest living
662 relatives of animals is induced by a bacterial chondroitinase. Cell 170:1175-
663 1183.e11.
- 664 63. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
665 Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation
666 of high-quality protein multiple sequence alignments using Clustal Omega. Mol
667 Syst Biol 7:539.
- 668 64. Jacobovitz MR, Rupp S, Voss PA, Gornik SG, Guse A. 2019. Dinoflagellate
669 symbionts escape vomocytosis by host cell immune suppression.
670 bioRxiv <https://doi.org/10.1101/864579>.
- 671

672 **Table 1. Functions of NF-κB in basal organisms.**

Organism	Biological Function	Data Available	References
<i>Aiptasia</i>	Immunity	protein, transcriptomic, mRNA, DNA binding	22, 54, 55, 64
<i>N. vectensis</i>	Immunity (?) and Development	protein, transcriptomic, mRNA, DNA-binding	34, 36, 37, 42, 48, 52
<i>Hydra</i>	Immunity	transcriptomic, protein	30, 31
Corals	Immunity	Transcriptomic	18, 38, 57–60
Sponge	Immunity and Development	protein, transcriptomic, mRNA, DNA-binding	21, 27, 50, 51
<i>Capsaspora</i>	?	genomic	11
Choanoflagellates	?	transcriptomic	12

673
674
675
676
677
678
679
680
681
682
683
684

Figure Legends


Figure 1. The structure of NF- κ Bs in organisms basal to Bilateria is most similar to human NF- κ B proteins rather than Rel proteins. The basic domain structures of homologs to NF- κ B in *Capsaspora*, choanoflagellates, poriferans (sponges), and cnidarians compared to the fly and vertebrate NF- κ Bs (p100/p105, Relish), Rels (RelA, RelB, c-Rel, Dif, Dorsal), and I κ B (Cactus). Green, Rel Homology Domain (RHD); pink, Nuclear Localization Signal (NLS); blue, Glycine Rich Region (GRR); black bars, Ankyrin repeats; red, Death Domain; SSS/SS, conserved serines important for phosphorylation and degradation of the protein (red SSS indicate homology to p100); and rainbow, N-terminal domains of choanoflagellates not typically seen in NF- κ B proteins.

Figure 2. Phylogenetic analysis places Choanoflagellate NF- κ Bs as an outgroup of vertebrate and fly NF- κ Bs. A Bayesian analysis of holozoan RHDs, including the recently identified choanoflagellate NF- κ Bs. MEME analysis was performed on each sequence to identify shared motifs. NF- κ B proteins and Rel proteins cluster separately from each other, and choanoflagellate NF- κ Bs cluster as a single outgroup. Conserved motifs are highlighted, and two choanoflagellate specific sequences are shown in teal and yellow.

Figure 3. The evolution of NF- κ B. Prior to the rise of holozoan life, Ankyrin repeats (ANK, red) were present in bacteria and archaea. The appearance of an RHD-containing protein (green) likely led to RHD-only proteins that diversified in choanoflagellates and to an RHD-ANK fusion in an ancestor of *Capsaspora*. Metazoans retained the full-length


NF- κ B fusion protein or in some cnidarians (e.g., Hydra and Nematostella) underwent gene splitting events to create separate RHD and ANK repeat proteins.

Fig. 1



731

Fig. 2



733

734

735

Fig. 3

