

HARMONIC GRADIENTS ON HIGHER DIMENSIONAL SIERPIŃSKI GASKETS

LUKE BROWN, GIOVANNI FERRER, GAMAL MOGRABY, LUKE G. ROGERS,
AND KARUNA SANGAM

ABSTRACT. We consider criteria for the differentiability of functions with continuous Laplacian on the Sierpiński Gasket and its higher-dimensional variants SG_N , $N > 3$, proving results that generalize those of Teplyaev [1]. When SG_N is equipped with the standard Dirichlet form and measure μ we show there is a full μ -measure set on which continuity of the Laplacian implies existence of the gradient ∇u , and that this set is not all of SG_N . We also show there is a class of non-uniform measures on the usual Sierpiński Gasket with the property that continuity of the Laplacian implies the gradient exists and is continuous everywhere, in sharp contrast to the case with the standard measure.

1. INTRODUCTION

In analysis on fractals the basic differential operator is a Laplacian obtained either by probabilistic methods [2] or as a renormalized limit of graph Laplacians [3]. There are then various approaches to defining a gradient, or first derivative, such as those in [4, 5, 6, 7, 8], and related questions remain an active area of research [9, 10, 11]. The results of this paper are a contribution to understanding what connection there is between smoothness measured using the Laplacian and the pointwise existence of a gradient, as the situation is very different than in the setting of Euclidean spaces or manifolds.

A fundamental result relating the regularity of the Laplacian and existence of a gradient was proven by Teplyaev in [1], who gave an example in which functions with continuous Laplacian are differentiable a.e. but can fail to be differentiable at a countable dense set of points. The innovative idea was not the example itself, which was just the standard Sierpiński gasket with its usual Laplacian and Bernoulli measure, but a concrete description of the gradient which allowed the points of differentiability to be described fairly precisely. It should be noted that, on the Sierpiński gasket, Teplyaev's gradient can be identified with that of Kusuoka [4].

In Section 2 we introduce the N -vertex Sierpiński Gasket SG_N and its analytic structure, and in Section 3 we review Teplyaev's gradient and some basic results from [1]. In Section 4 we then build on Teplyaev's work to show how the structure of the measure affects the connection between Laplacian regularity and existence of a gradient. Theorem 4.1 shows that, in contrast with the previously mentioned results from [1], if we equip the Sierpiński gasket with a suitably chosen self-similar measure

Date: January 25, 2021.

2010 Mathematics Subject Classification. Primary 31C25, Secondary 28A80.

Key words and phrases. Sierpinski gasket, harmonic gradient, harmonic structure, Dirichlet form, Laplacian, fractal.

Research supported in part by the NSF Grants DMS-1659643 and DMS-1613025.

having unequal weights, then we find that functions with continuous Laplacian are not only differentiable everywhere but the gradient is continuous. The discontinuity of natural gradients in the case of the standard measure is well-known, and it is rather unexpected that a continuous gradient can be obtained with such a simple modification of the measure.

Section 5 is concerned with differentiability results on the gaskets SG_N . We show that if the self-similar Laplacian and measure are symmetric under the symmetries of the underlying simplex then the results proved in [1] can be generalized to SG_N , though the description of the points of differentiability is less explicit and some proofs are correspondingly more complicated.

2. HIGHER DIMENSIONAL SIERPIŃSKI GASKETS

Let $N \in \mathbb{N}$ with $N \geq 3$. We largely follow [6, 12] in the following definitions and basic results. Note that, in the definition below, SG_3 is the usual Sierpiński Gasket.

Definition 2.1. Let $\{p_i\}_{i=0}^{N-1}$ be the vertices of a regular N simplex in $\mathbb{R}^{N-1}$ such that $|p_j - p_k| = 1$ if $j \neq k$. Let $\{F_i\}_{i=0}^{N-1}$, with $F_i : \mathbb{R}^{N-1} \rightarrow \mathbb{R}^{N-1}$, be the iterated function system defined by $F_j(x) = \frac{1}{2}(x - p_j) + p_j$. Then the N -dimensional Sierpiński Gasket, denoted SG_N , is the unique non-empty compact set such that $SG_N = \bigcup_{j=0}^N F_j(SG_N)$.

Definition 2.2. Let $S_N = \{0, 1, \dots, N-1\}$ and $\Omega_N = S_N^{\mathbb{N}}$ be the collection of one-sided infinite words over S_N . Similarly, a finite word of length $m \in \mathbb{N}$ is an element of the n -fold product S_N^m .

For simplicity, we often omit the index N in Ω_N and S_N and write Ω, S respectively. SG_N is post-critically finite with post-critical set $V_0 = \{p_0, \dots, p_{N-1}\}$. We write $F_w = F_{w_1} \circ \dots \circ F_{w_m}$, where $w = w_1 \dots w_m$ is a finite word of length m over the alphabet S . Let $V_m = \bigcup_{w \in S^m} F_w(V_0)$ and consider these points as vertices of a graph in which adjacency $x \sim_m y$ means there is a word w of length m such that $x, y \in F_w(V_0)$. A non-negative definite, symmetric, quadratic form on SG_N may be defined as a limit of graph energies as follows.

Definition 2.3. Let u, v be continuous functions on SG_N . The bilinear form

$$\mathcal{E}_m(u, v) = \left(\frac{N+2}{N} \right)^m \sum_{x \sim_m y} (u(x) - u(y))(v(x) - v(y))$$

defines the graph energy of level m .

We write $\mathcal{E}_m(u) = \mathcal{E}_m(u, u)$. Then $\{\mathcal{E}_m(u)\}$ is a nondecreasing sequence of graph energies, so $\mathcal{E}(u) = \lim_{m \rightarrow \infty} \mathcal{E}_m(u)$ is well-defined; setting its domain to be $\text{dom } \mathcal{E} = \{u : SG_N \rightarrow \mathbb{R} \mid \mathcal{E}(u) < \infty\}$ one obtains a non-negative definite, symmetric quadratic form that extends to the completion of $\bigcup_m V_m$, which can be shown to be SG_N , and the domain is uniform-norm dense in the continuous functions. For proofs of these facts see [3]. By construction this form is also self-similar in the sense that

$$(2.1) \quad \mathcal{E}(f, f) = \left(\frac{N+2}{N} \right) \sum_{i=0}^{N-1} \mathcal{E}(f \circ F_i, f \circ F_i)$$

We equip SG_N with a Bernoulli measure μ with weights $\{0 < \mu_i < 1\}_{i=0}^{N-1}$, $\sum_i \mu_i = 1$, at which point $(\mathcal{E}, \text{dom } \mathcal{E})$ is a Dirichlet form and we may define the Dirichlet μ -Laplacian as follows (see [3, 13]):

Definition 2.4. Let $u \in \text{dom } \mathcal{E}$, and let f be continuous. Then $u \in \text{dom } \Delta_\mu$ with $\Delta_\mu u = f$ if

$$\mathcal{E}(u, v) = - \int_{SG_N} f v d\mu \text{ for all } v \in \text{dom}_0 \mathcal{E},$$

where $\text{dom}_0 \mathcal{E}$ is the subspace of $\text{dom } \mathcal{E}$ consisting of functions that vanish at V_0 .

A function $h \in \text{dom } \mathcal{E}$ is called harmonic if it has specified values on V_0 and minimizes the graph energies $\mathcal{E}_n(u)$ for all $n \geq 1$. We can calculate $h|_{V_{m+1}}$ from $h|_{V_m}$ using harmonic extension matrices.

Definition 2.5. Let h be a harmonic function on SG_N . The harmonic extension matrices $\{A_i\}_{i=0}^{N-1}$ are defined by

$$\begin{pmatrix} h(F_i(p_0)) \\ \vdots \\ h(F_i(p_{N-1})) \end{pmatrix} = A_i \begin{pmatrix} h(p_0) \\ \vdots \\ h(p_{N-1}) \end{pmatrix}.$$

The harmonic extension matrices for SG_N are derived in [12], and given by

$$(2.2) \quad A_0 = \frac{1}{N+2} \begin{pmatrix} N+2 & \mathbf{0} \\ \mathbf{2} & I_{N-1} + J_{N-1} \end{pmatrix}$$

where I_{N-1} is the $(N-1) \times (N-1)$ identity matrix, J_{N-1} is the $(N-1) \times (N-1)$ matrix with all entries equal 1, and $\mathbf{0}$ and $\mathbf{2}$ are the $(N-1)$ size vectors with all entries 0 and 2 respectively. All other harmonic extension matrices can be found with cyclic row and column permutations.

Let A_i be a harmonic extension matrix of SG_N and $\sigma(A_i)$ the set of eigenvalues of A_i . Then the eigenspace of $\lambda \in \sigma(A_i)$, denoted by $E_i[\lambda]$, is the subspace of $\mathbb{R}^N$ spanned by the eigenvectors of A_i corresponding to λ . We need an elementary lemma.

Lemma 2.6. Let A_i be a harmonic extension matrix for SG_N . Then, the eigenvalues of A_i are $\sigma(A_i) = \{1, \frac{N}{N+2}, \frac{1}{N+2}\}$. The corresponding eigenspaces have the following dimensions:

$$\begin{aligned} \dim E_i[1] &= 1 \\ \dim E_i[\frac{N}{N+2}] &= 1 \\ \dim E_i[\frac{1}{N+2}] &= N-2. \end{aligned}$$

Proof. It can be easily verified that $(1, \dots, 1)^T$ is a simple eigenvector with eigenvalue 1, $(0, 1, \dots, 1)^T$ is a simple eigenvector with eigenvalue $\frac{N}{N+2}$ and that $(\dots, 0, 1, -1, 0, \dots)^T$ are $N-2$ eigenvectors corresponding to the eigenvalue $\frac{1}{N+2}$. $\square$

Let $\mathcal{H}$ denote the space of harmonic functions. Since these are determined by their values on V_0 this space is N -dimensional. Let $W \subset \mathcal{H}$ be the subspace of constant functions and $P : \mathcal{H} \rightarrow \mathcal{H}/W =: \tilde{\mathcal{H}}$ be the quotient map. As W is the eigenspace for the eigenvalue 1 we have $A_i(W) = \{A_i w \mid w \in W\} \subseteq W$ for $i \in \{0, \dots, N-1\}$ and thus there is $\tilde{A}_i : \tilde{\mathcal{H}} \rightarrow \tilde{\mathcal{H}}$, such that $\tilde{A}_i \circ P = P \circ A_i$.

We call $\{\tilde{A}_i\}_{i=0}^{N-1}$ the induced harmonic extension matrices. The energy $\mathcal{E}(\cdot, \cdot)$ is a bilinear form on $\mathcal{H}$ and $\mathcal{E}(u, u) = 0$ if and only if u is a constant function on SG_N , so the restriction of $\mathcal{E}(\cdot, \cdot)$ on $\tilde{\mathcal{H}}$ is a well-defined inner product that makes $\tilde{\mathcal{H}}$ a Hilbert space. The following is an immediate consequence of Lemma 2.6.

Corollary 2.7. *Let $\tilde{A}_i$ be a induced harmonic extension matrix for SG_N . Then, the eigenvalues of $\tilde{A}_i^{-1}$ are $\sigma(\tilde{A}_i^{-1}) = \{\frac{N+2}{N}, N+2\}$. The corresponding eigenspaces have the following dimensions:*

$$\begin{aligned} \dim E_i\left[\frac{N+2}{N}\right] &= 1 \\ \dim E_i[N+2] &= N-2. \end{aligned}$$

Remark 2.8. *A simple calculation shows that the eigenspaces $E_i\left[\frac{N+2}{N}\right]$ and $E_i[N+2]$ of $\tilde{A}_i^{-1}$ are orthogonal subspaces in $(\tilde{\mathcal{H}}, \mathcal{E})$.*

3. HARMONIC GRADIENTS IN THE SENSE OF TEPLYAEV

We define a harmonic gradient on SG_N following the approach and notation of Teplyaev [1], which is closely related to work of Kusuoka [4]. We require some notation for a cell containing a point described by an infinite word and for a harmonic approximation to the function on such a cell.

Definition 3.1. *For $\omega = \omega_1\omega_2\cdots \in \Omega_N$ the truncated word $[\omega]_n \in S_N^n$ is $[\omega]_n = \omega_1\omega_2\cdots\omega_n$.*

Definition 3.2. *The n -level harmonic approximation at word $\omega \in \Omega$ is*

$$\nabla_n f(\omega) = \tilde{A}_{[\omega]_n}^{-1} \tilde{H}(f \circ F_{[\omega]_n}),$$

where $\tilde{H}(g) = PH(g)$, and $H(g)$ is the unique harmonic function that coincides with g on the boundary of SG_N . The harmonic gradient at ω is defined to be

$$\nabla f(\omega) = \lim_{n \rightarrow \infty} \nabla_n f(\omega)$$

if the limits exist in $\tilde{\mathcal{H}}$.

Observe that the preceeding is analogous to the way in which secants converge to a tangent in elementary calculus, with harmonic functions playing the role of linear functions (because the latter are harmonic on $\mathbb{R}$). The n -level harmonic gradient of f at a point x is akin to a secant modulo constant functions because it is the unique globally harmonic function modulo constants that agrees with f at the boundary points of a cell containing the point. The matrices $\tilde{A}_{[\omega]_n}^{-1}$ are used simply to find the boundary values of this harmonic function from data on the cell at scale n . Then the limit of the harmonic approximations as the scale goes to zero is the harmonic gradient.

The following theorems are essential to our treatment of the topic, and were proved for a resistance form satisfying the identity

$$(3.1) \quad \mathcal{E}(f, f) = \sum_{i=0}^{N-1} r_i^{-1} \mathcal{E}(f \circ F_i, f \circ F_i)$$

and a Bernoulli measure with weights $0 < \mu_i < 1$ in [1].




FIGURE 1. SG_3 Level 1 (left) and 2 (right) with respect to the IFS $\{F_{ij}\}_{i,j \in S}$. Cells shaded according to their measure weights.

Theorem 3.3 ([1, Theorem 1]). *Suppose $f \in \text{dom } \Delta_\mu$. Then, $\nabla f(\omega)$ exists for every $\omega \in \Omega$ such that*

$$\sum_{n \geq 1} r_{[\omega]_n} \mu_{[\omega]_n} \|\tilde{A}_{[\omega]_n}^{-1}\| < \infty$$

Corollary 3.4 ([1, Corollary 5.1]). *Suppose that $f \in \text{dom } \Delta_\mu$. Then, $\nabla f(\omega)$ exists for all $\omega \in \Omega$ if*

$$r_j \mu_j \|\tilde{A}_j^{-1}\| < 1$$

For $j = 1, \dots, N$. Moreover, in this case, $\nabla f(\omega)$ is continuous in Ω .

Teplyaev [1] points out that Corollary 3.4 is not applicable to the Sierpiński Gasket when μ is the standard (uniform) Bernoulli measure. The same is true for SG_N for any $N \geq 3$ because one may readily compute that $\|A_j^{-1}\| = N + 2$ for each $j = 0, \dots, N - 1$, while each $\mu_j = N^{-1}$ and, as previously noted (see (2.1)), each $r_j = \frac{N}{N+2}$, so that $r_j \mu_j \|\tilde{A}_j^{-1}\| = 1$. Teplyaev shows that one can apply Theorem 3.3 to certain points on the Sierpiński Gasket, but their description is rather complicated.

4. A MEASURE ON SG_3 FOR WHICH FUNCTIONS WITH CONTINUOUS LAPLACIAN HAVE CONTINUOUS GRADIENT

On the standard Sierpiński Gasket SG_3 with its usual self-similar resistance form (as defined in Section 2) we consider the Laplacian associated to a non-uniform Bernoulli measure defined using the iterated function system of the second level, meaning that the similarities are compositions $F_{ij} = F_i \circ F_j$ of the usual three contractions on SG_3 . The following theorem gives a condition on the measure sufficient to ensure functions with continuous Laplacian have continuous gradients.

Theorem 4.1. *Let μ be the Bernoulli measure on SG_3 with the weights $\{\mu_{ij}\}_{i,j \in S}$ corresponding to the iterated function system $\{F_{ij} = F_i \circ F_j\}_{i,j \in S}$, so*

$$\mu(A) = \sum_{i,j \in S} \mu_{ij} \mu(F_{ij}^{-1}(A))$$

for any Borel set $A \subset SG_3$. If $\{\mu_{ij}\}_{i,j \in S}$ satisfy

$$(4.1) \quad \mu_{ij} < \begin{cases} \frac{1}{9} & i = j \\ \frac{1}{\sqrt{17+4\sqrt{13}}} & i \neq j \end{cases},$$

then $u \in \text{dom } \Delta_\mu$ implies that $\nabla u(\omega)$ exists and is continuous for all $\omega \in \Omega$.

Remark 4.2. The theorem provides examples because $\sqrt{17+4\sqrt{13}} < 9$, so there are many choices of μ_{ij} satisfying both (4.1) and $\sum_{i,j \in S} \mu_{ij} = 1$.

Proof. We apply Corollary 3.4, for which purpose we need the values of r_{ij} , μ_{ij} and an estimate of the norms of the matrices $\tilde{A}_{ij}^{-1}$ where $\tilde{A}_{ij}$ is the reduced harmonic extension matrix of the composition $F_{ij} = F_i \circ F_j$. The μ_{ij} values are given in (4.1) and $r_{ij} = (\frac{3}{5})^2$ because the energy scaling for any F_i is $\frac{3}{5}$, as is apparent by comparing equations (2.1) and (3.1).

We can calculate the spectral radius $\rho((\tilde{A}_{ij}^{-1})^* \tilde{A}_{ij}^{-1})$ for each i and j , using the description in Corollary 2.7 and Mathematica, to obtain

$$\sqrt{\rho[(\tilde{A}_{ij}^{-1})^* \tilde{A}_{ij}^{-1}]} = \sqrt{\rho[(\tilde{A}_i^{-1} \tilde{A}_j^{-1})^* (\tilde{A}_i^{-1} \tilde{A}_j^{-1})]} = \begin{cases} 25 & i = j \\ \frac{25}{9} \sqrt{17+4\sqrt{13}} & i \neq j \end{cases}.$$

Thus, we see that

$$r_{ij} \mu_{ij} \sqrt{\rho[(\tilde{A}_{ij}^{-1})^* \tilde{A}_{ij}^{-1}]} = \begin{cases} 9\mu_{ij} & i = j \\ \mu_{ij} \sqrt{17+4\sqrt{13}} & i \neq j \end{cases},$$

and hence from (4.1) that Corollary 3.4 is applicable because the spectral radius dominates the norm. It follows that if $u \in \text{dom } \Delta_\mu$ then $\nabla u(\omega)$ exists and is continuous for all $\omega \in \Omega$. In Figure 4, we illustrate one such μ . $\square$

A similar argument works on SG_N for any $N \geq 3$, though we do not know a convenient procedure for determining the optimal weights (corresponding to those in (4.1)) if $N > 3$.

5. GRADIENTS ON SG_N WITH THE STANDARD BERNOULLI MEASURE

As we noted at the end of Section 3, Corollary 3.4 is not applicable to the Sierpiński gasket or any SG_N , $N \geq 3$ when they are equipped with the standard (fully symmetric) measure and Dirichlet form. Hence in this setting there may be functions u with continuous Laplacian but for which ∇u fails to exist, at least at some points. Indeed, in [1], Teplyaev gives an example which may be used to construct a function u with continuous Laplacian such that ∇u is undefined on a countable dense set. This example may readily be generalized to SG_N , $N > 3$. However, Teplyaev also proves there is a full μ -measure set of words $\omega \in \Omega$ for which continuity of Δu implies existence of $\nabla u(\omega)$. The purpose of this section is to prove a generalization of this result to SG_N , $N > 3$.

The key idea in Teplyaev's proof of the result mentioned above is that harmonic functions have an improved scaling behavior near points defined by words that are asymptotically sufficiently non-constant. An appropriate generalization to our context uses the following concept.

Definition 5.1. A **k -block** for an alphabet $S_N = \{0, 1, \dots, N-1\}$ is a length k word w with k distinct letters, meaning $w = w_1 w_2 \dots w_k$ such that each $w_i \in S_N$ and $w_i \neq w_j$ for all $i \neq j$.

The key scaling behavior for an $(N-1)$ -block is the following estimate.

Lemma 5.2. Fix N . There is $\beta_N < 1$ such that for any $(N-1)$ -block $w \in S_N^{N-1}$

$$\frac{1}{N+2} \left\| \tilde{A}_w^{-1} \right\|^{1/(N-1)} \leq \beta_N.$$

Proof. Since the set of $(N-1)$ -blocks is finite it suffices to show the estimate for an arbitrary $(N-1)$ -block $w = w_1 \dots w_{N-1}$. Then $\tilde{A}_w^{-1} = \tilde{A}_{w_{N-1}}^{-1} \dots \tilde{A}_{w_1}^{-1}$ and the maximal eigenvalue for each $\tilde{A}_{w_j}^{-1}$ is $N+2$, so the result is true unless there is a vector common to the $(N+2)$ -eigenspaces of all of the $\tilde{A}_{w_j}^{-1}$. However we determined these eigenspaces explicitly in the proof of Lemma 2.6. Recalling that passage from A_i to $\tilde{A}_i$ eliminated the constant eigenspace (which was common to all A_i), we see that the eigenvectors of $\tilde{A}_i^{-1}$ with eigenvalue $N+2$ correspond to vectors in $\mathbb{R}^N$ that are orthogonal to the constants and to the unit vector in the i^{th} direction. A vector common to the eigenspaces of all $\tilde{A}_{w_j}^{-1}$ would then need to be orthogonal to the constants and to the unit vector in the w_j direction for $j = 1, \dots, N-1$, thus to all of $\mathbb{R}^N$. This shows the estimate for an arbitrary $(N-1)$ -block and proves the lemma. $\square$

Remark 5.3. One can compute β_N explicitly, but we do not know an elementary way to do this for general N . In [1] it is shown that $\beta_3 = \sqrt{\frac{7+\sqrt{13}}{18}}$.

The significance of a $(N-1)$ -block from our perspective is that the harmonic gradient exists at the point $F_w(X)$ if w has sufficient asymptotic density of $(N-1)$ -blocks. The density is counted using the following.

Definition 5.4. The **block counting function** $C_N: \Omega \times \mathbb{N} \rightarrow \mathbb{N} \cup \{0\}$ is defined for $\omega \in \Omega$ by

$$C_N(\omega, n) = \#\{i \in \mathbb{N} \mid i \leq n - (N-2), [i, i+N-2] \text{ is an } (N-1)\text{-block}\}.$$

The following theorem now provides a criterion sufficient for existence of the harmonic gradient.

Theorem 5.5. Let $u: SG_N \rightarrow \mathbb{R}$ and suppose $\Delta_\mu u$ is continuous, where μ is the standard Bernoulli measure. Then $\nabla u(\omega)$ is defined at every $\omega \in \Omega$ such that

$$(5.1) \quad \liminf_{n \rightarrow \infty} \frac{C_N(\omega, n)}{\log n} > \frac{1}{|\log \beta_N|}.$$

Lemma 5.6. Let $\omega \in \Omega_N$. Then

$$\left\| \tilde{A}_{[\omega]_n}^{-1} \right\| \leq (N+2)^n \beta_N^{C_N(\omega, n)}.$$

Proof. The proof is inductive with base case $n = 1$, for which $C_N(\omega, 1) = 0$ (by definition) and we are bounding the norm of $\tilde{A}_{w_1}^{-1}$ by its maximal eigenvalue. For the inductive step we consider two cases.

If the last $N - 1$ letters of $[\omega]_{n+1}$ do not form a $(N - 1)$ -block then $C_N(\omega, n) = C_N(\omega, n + 1)$ and bounding the norm of $\tilde{A}_{w_{n+1}}^{-1}$ by the maximal eigenvalue $(N + 2)$ we have from the induction hypothesis

$$\left\| \tilde{A}_{[\omega]_{n+1}}^{-1} \right\| \leq (N + 2) \left\| \tilde{A}_{[\omega]_n}^{-1} \right\| \leq (N + 2)^{n+1} \beta_N^{C_N(\omega, n)} = (N + 2)^{n+1} \beta_N^{C_N(\omega, n+1)}.$$

In the other case, where the last $N - 1$ letters form a $(N - 1)$ -block, we instead use Lemma 5.2 on this block and the inductive bound on for $n + 1 - (N - 1)$ to obtain

$$\begin{aligned} \left\| \tilde{A}_{[\omega]_{n+1}}^{-1} \right\| &\leq \left\| \tilde{A}_{[\omega]_{n-N+2}}^{-1} \right\| \left\| \tilde{A}_{\omega_{n-N+3}}^{-1} \cdots \tilde{A}_{\omega_{n+1}}^{-1} \right\| \\ &\leq (N + 2)^{n-N+2} \beta_N^{C_N(\omega, n-N+2)} (N + 2)^{N-1} \beta_N^{N-1} \\ &\leq (N + 2)^{n+1} \beta_N^{C_N(\omega, n-N+2)+N-1} \\ &\leq (N + 2)^{n+1} \beta_N^{C_N(\omega, n+1)}, \end{aligned}$$

where we also used the fact that $C_N(\omega, n + 1) - C_N(\omega, n - N + 2) \leq N - 1$, which is immediate from the definition. $\square$

Proof of Theorem 5.5. For the standard Bernoulli measure and Dirichlet form on the higher dimensional Sierpiński Gasket we have $r_{[\omega]_n} \mu_{[\omega]_n} = \frac{1}{(N+2)^n}$, as noted after Corollary 3.4. Inserting this and the result of Lemma 5.6 we compute, using that $0 < \beta_N < 1$ that

$$\sum_{n=1}^{\infty} r_{[\omega]_n} \mu_{[\omega]_n} \left\| \tilde{A}_{[\omega]_n}^{-1} \right\| \leq \sum_{n=1}^{\infty} \beta_N^{C_N(\omega, n)} \leq \sum_{n=1}^{\infty} n^{-C_N(\omega, n) |\log \beta_N| / \log n}$$

which is convergent because

$$\frac{C_N(\omega, n) \log |\beta_N|}{\log n} > \frac{1}{2} \left(1 + \liminf_n \frac{C_N(\omega, n) \log |\beta_N|}{\log n} \right) > 1$$

for all sufficiently large n . This gives the result by Theorem 3.3. $\square$

Theorem 5.7. *Let $u: SG_N \rightarrow \mathbb{R}$ and suppose $\Delta_\mu u$ is continuous, where μ is the standard Bernoulli measure. Then $\nabla u(\omega)$ is defined μ -a.e.*

Proof. We give a crude but sufficient lower bound on the set of words for which $C(\omega, n)$ satisfies the estimate in Theorem 5.5. Suppose we split a word of length $n - (N - 2)$ up into disjoint intervals of length $N - 1$. Evidently there are at least $k = \frac{n}{N-1} - 2$ of these. The probability of any one such interval being an $N - 1$ block is $p_N = (N!)N^{-N}$, so the probability that $C_N(\omega, n) < \frac{\log n}{|\log \beta_N|} := l$ does not exceed that of having l successes in k binomial trials where success has probability p_N . Using Chernoff's inequality to bound the stated probability by $\exp(-(kp_N - l)^2 / (2kp_N))$ and taking n large enough that $kp_N > 2l$ we have probability less than $\exp(-kp_N/8) = \exp(-nq_N)$ for a $q_N > 0$ that does not depend on n . The latter is summable over n , so the bound required in Theorem 5.5 follows from the first Borel-Cantelli lemma. $\square$

It is perhaps interesting to note that the preceding reasoning allows one to bound the Hausdorff dimension of the set of points at which ∇u is undefined by a value strictly less than the Hausdorff dimension of SG_N ; we omit the details.

6. ACKNOWLEDGEMENTS

The authors are grateful to Alexander Teplyaev and Daniel Kelleher for helpful discussions.

REFERENCES

- [1] Alexander Teplyaev. Gradients on fractals. *J. Funct. Anal.*, 174(1):128–154, 2000.
- [2] Martin T. Barlow. Diffusions on fractals. In *Lectures on probability theory and statistics (Saint-Flour, 1995)*, volume 1690 of *Lecture Notes in Math.*, pages 1–121. Springer, Berlin, 1998.
- [3] Jun Kigami. *Analysis on fractals*, volume 143 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2001.
- [4] Shigeo Kusuoka. Dirichlet forms on fractals and products of random matrices. *Publ. Res. Inst. Math. Sci.*, 25(4):659–680, 1989.
- [5] Robert S. Strichartz. Taylor approximations on Sierpinski gasket type fractals. *J. Funct. Anal.*, 174(1):76–127, 2000.
- [6] Jun Kigami. Measurable Riemannian geometry on the Sierpinski gasket: the Kusuoka measure and the Gaussian heat kernel estimate. *Math. Ann.*, 340(4):781–804, 2008.
- [7] Anders Pelander and Alexander Teplyaev. Products of random matrices and derivatives on p.c.f. fractals. *J. Funct. Anal.*, 254(5):1188–1216, 2008.
- [8] Masanori Hino. Energy measures and indices of Dirichlet forms, with applications to derivatives on some fractals. *Proc. Lond. Math. Soc. (3)*, 100(1):269–302, 2010.
- [9] Masanori Hino. Measurable Riemannian structures associated with strong local Dirichlet forms. *Math. Nachr.*, 286(14-15):1466–1478, 2013.
- [10] Naotaka Kajino. Analysis and geometry of the measurable Riemannian structure on the Sierpiński gasket. In *Fractal geometry and dynamical systems in pure and applied mathematics. I. Fractals in pure mathematics*, volume 600 of *Contemp. Math.*, pages 91–133. Amer. Math. Soc., Providence, RI, 2013.
- [11] Fabrice Baudoin and Daniel J. Kelleher. Differential one-forms on Dirichlet spaces and Bakry-Émery estimates on metric graphs. *Trans. Amer. Math. Soc.*, 371(5):3145–3178, 2019.
- [12] Sara Chari, Joshua Frisch, Daniel J. Kelleher, and Luke G. Rogers. Measurable riemannian structure on higher dimensional harmonic sierpiński gaskets. *arXiv:1703.03380*, 9 Mar 2017.
- [13] Robert S. Strichartz. *Differential equations on fractals*. Princeton University Press, Princeton, NJ, 2006. A tutorial.

(Luke Brown) DREXEL UNIVERSITY
DEPARTMENT OF MATHEMATICS
Email address: `lcb94@drexel.edu`

(Giovanni Ferrer) UNIVERSITY OF PUERTO RICO (MAYAGÜEZ)
DEPARTMENT OF MATHEMATICS
Email address: `giovanni.ferrer@upr.edu`

(Gamal Mograby) UNIVERSITY OF CONNECTICUT
DEPARTMENT OF MATHEMATICS
Email address: `gamal.mograby@uconn.edu`

(Luke Rogers) UNIVERSITY OF CONNECTICUT
DEPARTMENT OF MATHEMATICS
Email address: `luke.rogers@uconn.edu`

(Karuna Sangam) RUTGERS UNIVERSITY
DEPARTMENT OF MATHEMATICS
Email address: `karuna.sangam@rutgers.edu`