

Age and intraspecific diversity of resilient *Acropora* communities in Belize

Adele Irwin¹, Lisa Greer², Robert Humston¹, Meghann Devlin-Durante³, Paul Cabe¹, Halard Lescinsky⁴, Karl Wirth⁵, H. Allen Curran⁶, and Iliana B Baums³

¹Department of Biology, Washington & Lee University, Lexington, VA ²Department of Geology, Washington & Lee University, Lexington, VA ³Department of Biology, Pennsylvania State University, University Park, PA ⁴Department of Biology and Earth Science, Otterbein University, Westerville, OH ⁵Department of Geology, Macalester College, St. Paul, MN ⁶Department of Geosciences, Smith College, Northampton, MA

Adele Irwin

Biology Department, Washington & Lee University
204 W. Washington Street
Lexington, VA 24450

Lisa Greer

Professor of Geology, Washington & Lee University
204 W. Washington Street
Lexington, VA 24450
540-458-8870
greerl@wlu.edu

Robert Humston

Associate Professor of Biology, Washington & Lee University
204 W. Washington Street
Lexington, VA 24450

Meghann Devlin-Durante

Research Technologist III, Pennsylvania State University
213 Mueller
University Park, PA 16802

Paul Cabe

Professor of Biology, Washington and Lee University
204 W. Washington Street
Lexington, VA 24450

Halard Lescinsky

Professor of Biology and Earth Science, Otterbein University
Science Center
1 South Grove Street
Westerville, OH 43081

Karl Wirth

Associate Professor of Geology, Macalester College

Olin-Rice Science Center, 113
1600 Grand Avenue
Saint Paul, MN 55105

H. Allen Curran

Kenan Professor Emeritus of Geosciences, Smith College
Clark Science Center
44 College Lane
Northampton, MA 01063

Ilana B Baums

Associate Professor of Biology, Pennsylvania State University
213 Mueller
University Park, PA 16802

Keywords: *Acropora*; microsatellite; Belize; genotypic diversity; coral reef; Caribbean

abstract

The corals *Acropora palmata* and *Acropora cervicornis* are important Caribbean reef-builders that have faced significant mortality in recent decades. While many studies have focused on the recent demise of these species, data from areas where *Acropora* spp. have continued to thrive are limited. Understanding the genetic diversity, recruitment, and temporal continuity of healthy populations of these threatened *Acropora* spp. and the hybrid they form ("*Acropora prolifera*") may provide insights into the demographic processes governing them. We studied three reef sites with abundant *Acropora cervicornis*, *Acropora palmata*, and hybrid *Acropora* populations offshore of Ambergris Caye, Belize at Coral Gardens, Manatee Channel, and Rocky Point. Samples were collected from all three *Acropora* taxa. We used microsatellite markers to determine: 1) genotypic diversity; 2) dominant reproductive mode supporting local recruitment; 3) minimum and maximum genet age estimates of all three acroporids; and 4) the history of hybrid colonization at these sites. We found *Acropora* populations to be highly clonal with local recruitment primarily occurring through asexual fragmentation. We also estimated the ages of 10 *Acropora* genets using recent methodology based on somatic mutation rates from genetic data. Results indicate minimum ages between 62 – 409 years old for *A. cervicornis*, between 187-561 years for *A. palmata*, and between 156-281 years old for the *Acropora* hybrids at these sites. Our data indicate that existing *A. cervicornis*, *A. palmata*, and *Acropora* hybrid genets persisted during the 1980s Caribbean-wide *Acropora* spp. collapse, suggesting that these sites have been a refuge for Caribbean *Acropora* corals. Additionally, our data suggest that formation of extant hybrid *Acropora* genets predates the widespread collapse of its parent taxa.

introduction

The global decline of shallow coral populations in response to increasing sea-surface temperatures, overfishing, rising sea level, disease, eutrophication, sedimentation, and ocean acidification has been well documented (Hughes 1994; Hughes et al. 2003; Bellwood et al. 2004; Hoegh-Guldberg et al. 2007; Kuffner et al. 2015). Caribbean populations of *Acropora cervicornis* and *Acropora palmata* (Fig. 1), two of the most important Caribbean reef framework-builders, have experienced up to 98% mortality over just a few decades in response to increasing environmental stress and disease prevalence (Gladfelter 1982; Aronson and Precht 2001; Pandolfi 2002; Gardner et al. 2003; Eakin et al. 2010; Randall and van Woesik 2015). Consequently, these were the first two coral species listed as threatened under the US Endangered Species Act (NOAA 2005).

Despite Caribbean-wide declines, relatively abundant *A. cervicornis* and *A. palmata* populations have been reported in Honduras (Keck et al. 2005; Purkis et al., 2006), the Dominican Republic (Lirman et al. 2010), Mexico (Larson et al. 2014), Florida (Fort Lauderdale area; Vargas-Angel et al. 2003), and Belize (Peckol et al. 2003; Brown-Saracino et al. 2007; Macintyre and Toscano 2007; Busch et al. 2016). These studies have variably included general habitat surveys, quantification of percent of algal and live coral coverage, *Acropora* colony sizes, number of *Acropora* colonies, fish and invertebrate population densities, image-based spatial characterization, and disease prevalence, but none included genetic data.

Genotypic diversity (the number of distinct genotypes or clones per species per site) is a critical factor when assessing the long-term population trends and evolutionary potential of these sites because Caribbean acroporids do not self-fertilize and thus sexual reproduction requires the presence of several genotypes (Szmant 1986; Baums et al. 2005a; Baums et al. 2006; Fogarty et al. 2012). Clonal reproduction is also an important process in many populations. Caribbean

acroporids experience frequent breakage due to physical impacts such as storms and anchor damage. Branches are genetically identical to their donor colonies (ie. ramets of the same genet) and can re-attach to the benthos. Once re-attached they grow into new colonies, usually within 10s of meters of the donor colony (Highsmith 1982; Lirman 2000, Baums et al. 2006; Williams et al. 2008). This sometimes results in large areas of a reef dominated by a single genotype (Baums et al. 2006). Genotypic diversity of foundation fauna has been associated with overall population health and has been predicted to have a positive relationship with persistence during adverse conditions by providing a diverse set of alleles to the population (Altizer et al. 2003; Reed and Frankham 2003; Reusch et al. 2005, Downing et al. 2012; Williams et al. 2014). However, recent findings show that genotypic diversity in *Acropora* populations does not necessarily increase their sustainability over short time scales as predicted (Williams et al. 2014). Thus, the role of genotypic diversity in persistence remains unclear.

A. cervicornis and *A. palmata* hybridize to form “*Acropora prolifera*”, the only known scleractinian hybrid in the Caribbean (Fig. 1; van Oppen et al. 2000; Vollmer and Palumbi 2002; Willis et al. 2006). The hybrid has not been found in the fossil record (Budd et al. 1994), and has been reported as rare compared to its parents (Lang et al. 1998; Willis et al. 2006; Fogarty, 2007; Fogarty 2012). However, recently it has been observed more frequently and in higher abundance in Florida (Wheaton et al. 2006), Curaçao (Fogarty 2010), Honduras (Keck et al. 2005), the Lesser Antilles (Japaud et al. 2014), and Belize (Fogarty 2010; Fogarty et al. 2012; Richards and Hobbs 2015). The apparent recent increase in the hybrid may suggest either that hybridization is now occurring more frequently, or that hybridization has always occurred, but environmental conditions now favor maturation of hybrid larvae over those of the diminishing parent species (Willis et al. 2006).

In this study, we analyzed genetic data of *A. cervicornis*, *A. palmata*, and the hybrid from Coral Gardens, Manatee Channel, and Rocky Point, sites offshore of Ambergris Caye, Belize, where acroporids are highly abundant (Table 1, Fig. 2). At these sites our specific objectives were to: 1) determine genotypic diversity of *Acropora* taxa; 2) calculate the degree to which sexual vs. asexual recruitment has occurred; 3) estimate the age of *Acropora* genets; and 4) determine the history and nature of hybrid colonization. To do this, we analyzed data from microsatellite loci to identify genets and ramets and subsequently draw inference about the genotypic diversity, as well as the degree of asexual versus sexual reproduction in recruitment to these sites (Baums et al. 2005b). We will refer to a “genet” as an assemblage of genetically identical colonies (clones) that are descendants of a single zygote (Harper 1977, Hughes 1989, Carvalho 1994), and a genet’s component colonies as “ramets” (Kays and Harper 1974) or clone mates. A method based on accumulation of somatic mutations in clonal organisms (Weiher et al. 1999; Eriksson 2000; Ally et al. 2008; de Witte and Stocklin 2010), recently adapted to study *Acropora* genets (Devlin-Durante et al. 2016), was applied to estimate clonal age and assess the history of *Acropora* spp. and *A. prolifera* hybrid persistence at these sites.). We analyzed clonal age data for all *Acropora* taxa to determine persistence and, for the hybrid at our study sites, to elucidate the history of its emergence and spread.

materials and methods

Study Site

The three reef sites sampled lie within roughly a 50-km distance between Caye Caulker to the south and the northern end of Ambergris Caye in Belize (Fig. 2). Coral Gardens

(17°50'00.36"N, 87°59'32.45"W, Table 1, Fig. 1) is a shallow-water back reef setting with a maximum depth of ~7m. It is comprised of lagoonal patch reefs dominated by interconnected thickets of *A. cervicornis*, large peripheral colonies of *A. palmata*, and small hybrid colonies. The *Acropora* thickets are interspersed with mixed coral stands dominated by *Orbicella*, *Agaricia*, *Porites*, and *Millepora* species, and areas of sandy bottom. Peckol et al. (2003) and Brown-Saracino et al. (2007) reported live coral cover (all species) in this area exceeding 43%. More recently, Busch et al. (2016) reported an average value of ~30% live *A. cervicornis* coral cover (species specific) in the *Acropora* thickets sampled in this study. The extent of live *Acropora* thickets in the greater Coral Gardens area exceeded 7.5 hectares in 2015, making it one of the largest documented sites of its kind in the Caribbean (Busch et al. 2016).

Manatee Channel (17°47'58.08"N, 87°59'45.57"W; Table 1, Fig. 2) is approximately 1-2 m deep and located just inside the reef crest. It is comprised of patch reefs dominated by thickets of the hybrid and scattered stands of *A. cervicornis* and *A. palmata*. Acroporid thickets and colonies are surrounded by mixed coral stands dominated by *Orbicella*, *Agaricia*, *Porites*, and *Millepora* species and sandy bottom. Manatee Channel has been previously described in the literature by Fogarty (2012) as “north Caye Caulker.”

Rocky Point, (18°12'37.95"N, 87°82'60.64"W; Table 1, Fig. 2) approximately 2-3 m depth, is dominated by individual colonies and small patches of the hybrid, both parent species, and some colonies of *Orbicella*, *Porites*, and *Siderastrea* species (Figures 2 and 3). Rocky Point lies within the Bacalar Chico National Park and Marine Reserve and is the only site among our study locations with marine protected area (MPA) status.

Sampling

Live coral samples approximately 1 cm³ were collected from *Acropora* branch tips using surgical bone cutters. Samples were preserved in 95% ethanol and refrigerated prior to DNA extractions. Coral tissue samples were collected from Coral Gardens in June of 2013 and 2014, and from Manatee Channel and Rocky Point in June 2014. In 2013, coral tissue was collected at Coral Gardens from haphazardly selected *A. cervicornis* colonies (n=60) within the reef boundary, and each observed colony of *A. palmata* (n=78) and the hybrid (n=48) was sampled. Additional sampling in 2014 was designed to increase the likelihood of sampling ramets of the same genet: At Coral Gardens, *A. cervicornis* samples (n=158) were collected from 5-m radius plots in which a center point was designated and samples were collected along 4 randomly generated headings (at least 5° apart) every 1 m for a total of 20 samples per plot. The likelihood that the same colonies were sampled in 2013 and 2014 is small. At Manatee Channel, hybrid samples (n=80) were collected using the same circle plot method, with 4 plots (20 samples per plot) placed at areas of largest abundance and never overlapping. All observed colonies of *A. cervicornis* (n=20) and *A. palmata* (n=15) across the entire site were also sampled. We defined a colony as a continuous, upright entity of living coral attached to a base or seafloor. At Rocky Point the hybrid was sampled haphazardly (n=32) due to irregular patch shape. *A. cervicornis* and *A. palmata* were not sampled at this site.

Genotypic Analysis

Nuclear DNA was extracted from tissue of 3-5 polyps using the Qiagen DNEasy kit (Qiagen, Germany) and protocol with the following modifications: 1) we performed proteinase K digestion overnight at 56°C in a stationary water bath; and, 2) precipitates were dissolved in Buffer AL for 10 minutes 56°C in a stationary water bath. We amplified five microsatellite loci (166, 181, 182, 192, 207) via polymerase chain reaction in two multiplexes following methods

described by Baums et al (2005a). Locus 192 does not amplify in *A. cervicornis* and amplifies only the *A. palmata* allele in the hybrid (making the marker appear homozygous). Fragment lengths were determined using an ABI 3730 (Gene Scan 500-Liz, Applied Biosystems) and electropherograms were scored using GeneMapper Software 3.0 (Applied Biosystems).

Data Analyses

Genotypic diversity:

We used data from highly polymorphic microsatellite loci to identify unique genotypes (genets). Samples were considered ramets of the same genet, i.e. genotypically identical colonies arisen from fragmentation, if they have identical genotypes across all five or four loci, or if they have identical genotypes across all five or four (*A. cervicornis*) loci with an additional allele(s). Because of the smaller number of loci retrieved for *A. cervicornis* the power to distinguish clonemates from closely related individuals is lower for *A. cervicornis* (average $P_{gen} = 1.11^{-04}$) than the other taxa (*A. palmata* average $P_{gen} = 2.99^{-07}$, hybrid average $P_{gen} = 2.74^{-05}$). Nevertheless, using MLGSim 2.0 (Stenberg et al. 2003) we determined that each of the identified genets, regardless of taxon, was highly likely to be the product of random mating within the population (P_{sex} , $p < 0.05$ for each genet).

In addition, samples with matching genotypes across four or three (*A. cervicornis*) loci were also considered to be ramets of a single genet if the fifth or fourth (*A. cervicornis*) locus only differed by single- or double-step mutations. Such genotypic differences among ramets were assumed to arise via mutations in somatic cell lines that can accumulate with age (Devlin-Durante et al. 2016; see below). We calculated diversity metrics following methods described in

Baums et al (2006) as adapted from Stoddart and Taylor (1988). Observed genotypic diversity (G_o) within and among sampling sites for each species was calculated as:

$$\frac{1}{\sum_i^k g_i^2}$$

Where g_i is the relative frequency of an observed genotype among all (k) genotypes. Expected genotypic diversity under the condition of exclusively sexual reproduction (G_e) is assumed equal to the number of colonies sampled (i.e. sample size, n), since only one sample was collected from each separate colony. Thus, the ratio G_o/G_e provides an index of genotypic diversity within the population that captures the relative degree to which sexual reproduction contributes to recruitment. Genotypic evenness is represented as the ratio of observed genotypic diversity versus genotypic richness (G_o/N_g), where N_g is the number of unique genets in the sample. Finally, we calculated standardized genotypic richness among and within sites relative to sample size (N_g/n). Baums et al. (2006) use these metrics to draw conclusions about the demography of local recruitment, classifying sites with high clonality as “asexual” (most colony recruitment arising from fragmentation) and those with the low clonality as “mostly sexual” (recruitment arising from sexual reproduction and settlement of planulae). We used values for these metrics from this study and those from Baums et al. (2006) to make relative inferences about clonal structure and recruitment of *Acropora* at our sites compared to others in the Caribbean.

Somatic mutations:

Genotypes accumulate somatic mutations over time. Assuming a constant mutation rate, the number of somatic mutations can be used to estimate the age of the genome by applying coalescent methods (Ally et al. 2008). Microsatellites are ideal for estimating genetic divergence because they have high mutation rates and low technical error rates. Hence, they can provide adequate resolution of somatic mutations. A ramet was identified as having a somatic mutation if there was an amplification of an additional allele/s, but alleles were otherwise identical at all 5 or 4 (*A. cervicornis*) loci (Devlin-Durante et al. 2016). While the appearance of an additional allele at a locus (rendering an individual polyploid instead of diploid at this locus) may seem odd, it has been deemed common in *A. palmata*, and predicted to occur due to genomes accumulating somatic duplications over time.

As somatic gene duplications accumulate, multiple copies of the microsatellite locus become available for replication slippage (Devlin-Durante et al. 2016). For example, consider the case where an unmutated ramet had the ancestral genotype of allele 1 = 160 bp, allele 2 = 172 bp (in short: 160/172) at locus 1, whereas a mutated ramet had alleles of 160/172/175 at locus 1. In this instance, the ramet mutated when its second allele duplicated, and the new third allele underwent a single-step mutation, adding a tri-nucleotide repeat. Alternatively, one of the alleles may have mutated in some of the cells or polyps without a prior gene duplication event (Devlin-Durante et al. 2016). Because DNA was extracted from multiple polyps this could also lead to the appearance of three alleles in the electropherograms. A ramet was also identified as having a somatic mutation if it matched a genotype across four or three (*A. cervicornis*) loci, but at the fifth or fourth (*A. cervicornis*) locus only differed by single- or double-step mutations.

Next we required that a genet have at least five ramets, a minimum requirement set by Devlin-Durante et al. (2016) to improve accuracy of aging. The mutation rates established by Devlin-Durante et al. (2016) (lowest=1.542-05 per locus per year, highest=1.195-04 per locus per year) were then applied to the equation described in Ally et al. (2008) to obtain minimum and maximum ages of each genet. The methods for calculating clonal age utilizing genetic divergence are described in Ally et al. (2008). In brief, there are two statistics, π_k and S_k , that describe genetic divergence within a clone (Slatkin 1996). We calculated the average number of pairwise differences per locus for the k th clone

$$\pi_k = \frac{1}{\binom{n}{2}} \sum_{i=1}^{n-1} \sum_{j=i+1}^n s_{ij}$$

where n is the number of sampled ramets and s_{ij} is the number of genetic differences between ramet i and j averaged across loci (Ally et al. 2008). We chose π_k to measure the level of genetic divergence because it has been shown to be more robust to deviations from a starlike phylogeny than S_k (the observed proportion of polymorphic loci) (Ally et al. 2008).

results

Genotypic diversity

Collectively we identified 18 genets among $n = 227$ individuals of *A. cervicornis* sampled, 31 genets among $n = 92$ *A. palmata*, and 14 genets among $n = 147$ hybrid samples (Table 2). Acroporids at our study sites showed a range in genotypic diversity suggesting that local recruitment in some locations was predominantly asexual, i.e. supported by fragmentation, while at other sites recruitment was supported to a greater degree by sexual reproduction (Table 2; Fig.3). At Manatee Channel, all hybrid colonies ($n=70$) shared the same genotype and were therefore ramets of a single genet (Table 2, Fig. 3). Thus, recruitment of existing hybrid colonies at Manatee Channel had occurred only through fragmentation from existing colonies. By contrast, *A. cervicornis* and *A. palmata* at Manatee Channel showed the highest genotypic diversity ($G_o/G_e = 0.26$ and 0.37 respectively) and evenness ($G_o/N_g = 0.62$ and 0.69 respectively) among locations¹, indicating that a substantial proportion of local recruitment was via sexual reproduction (following Baums et al. 2006; Table 2, Fig. 3). Coral Gardens was characterized by low genotypic diversity across all three taxa, indicating a high degree of asexual recruitment (Table 2, Fig. 3). It is possible but unlikely that a few individual colonies were re-sampled from 2013 to 2014. If so, this could partially contribute to the low genotypic diversity seen at Coral Gardens. Rocky Point, at which only hybrids were sampled, was characterized by low genotypic diversity and predominantly asexual recruitment as well (Table 2, Fig. 3).

When samples were pooled across all three sites, both diversity and evenness metrics were relatively low and indicated predominantly asexual recruitment (Table 2, Fig. 3). Overall, *A. palmata* showed the highest genotypic diversity ($G_o/G_e=0.43$), evenness ($G_o/N_g=0.15$), and

¹ Note that although the calculated value for evenness is highest for *A. prolifera* hybrid at Manatee Channel ($G_o/N_g = 1$), this value is not meaningful as all individuals shared the same genotype.

richness ($N_g/N=0.34$) across all three species (Table 2, Fig. 3). *A. cervicornis* and the hybrid were comparatively much lower in genotypic diversity and richness (Table 2, Fig. 3).

Somatic Ages

From our 466 *Acropora* samples, we were able to calculate preliminary minimum and maximum age estimates for 10 of the 63 genets (Table 3). We were unable to estimate ages for the remaining 53 genets using this method either because we sampled fewer than five ramets of that genet (44 genets) or because there were no observed somatic mutations in the loci amplified (9 genets). In the latter scenario, we concluded these genets lack somatic mutations because they are of recent origin.

At Coral Gardens, *A. cervicornis* genets were estimated to be between 62-460 years old (two genets), 179-1337 years old (one genet), 393-2931 years old (one genet), and 409-3052 years old (one genet) (Table 3). *A. palmata* genets at this site were estimated to be between 187-1397 years old (one genet), 219-1636 years old (one genet), and 561-4191 years old (one genet) (Table 3). A single hybrid genet at this site was estimated to be between 281-2096 years old (Table 3). At Manatee Channel, no somatic mutations were observed at these loci for any *A. cervicornis*, *A. palmata*, and hybrid genets. At Rocky Point, we were able to estimate the age for one hybrid genet to be between 156-1164 years old (Table 3).

discussion

In this study we had the rare opportunity to analyze large living stands of ecologically significant *Acropora* spp. and the hybrid they form, at a time when the existence of acroporids in the Caribbean is in peril. From our collection and analyses of 466 samples, we determined that acroporids offshore of Ambergris Caye, Belize were mostly clonal, with little sexual recruitment. Although genetic diversity was low, we found that both veteran and more recent genets were represented in all three acroporid taxa at these sites. Furthermore, our data suggested that hybrid colonization at some of these sites pre-dated the widespread Caribbean collapse of its parent acroporid species.

The genotypic diversity and evenness of acroporid populations at our study sites was low overall, with the exception of *A. palmata* and *A. cervicornis* at Manatee Channel, which exhibited a relatively high degree of genotypic diversity and evenness. From this information, we deduced that there is minimal input from local sexual reproduction or from ‘upstream’ source populations for all three Caribbean acroporids at Coral Gardens and the hybrid at Manatee Channel and Rocky Point. Only the Manatee Channel *A. palmata* and *A. cervicornis* samples in our study indicated substantive inputs from sexual reproduction. This may be due to localized variation in current patterns or subsequent differences in delivery or retention of planulae at these sites. A similar prevalence of asexual recruitment was observed by Baums et al. (2006) at a nearby *A. palmata* population in Chinchorro, Mexico (~49 km north of our northernmost site, Rocky Point), as well as in most other *A. palmata* populations at Western Caribbean sites (Baums et al. 2005b). Our results further validate their claim that Western Caribbean *Acropora* populations exhibit little sexual reproduction.

Previous studies from other abundant *Acropora* sites predicted that large populations were critical sources of propagules that might contribute to genetic diversity and coral coverage at neighboring sites (Keck et al. 2005; Zubillaga et al. 2008; Lirman et al. 2010, Vargas-Angel et al. 2003). However, because Caribbean acroporids usually do not self-fertilize, production of sexual offspring is a function of the number of genotypes present and not just a function of colony density or size (Levitan and McGovern 2005; Fogarty et al. 2012; Williams et al. 2014). Our study suggests that while Coral Gardens may be the largest extant acroporid site in the Caribbean (Busch et al., 2016), this may not translate into successful downstream sexual recruitment. Only observation of spawning and gamete-crossing experiments at these sites would provide additional insight into the production of gametes by these stands.

Although the genotypic diversity is generally low, our age estimates of acroporid genets vary. Using a new genetic aging technique, we were able to estimate age ranges for 10 genets in this study. While the uncertainty in the age range estimates may be considerable (Devlin-Durante et al. 2016), we believe that the more conservative minimum age estimates may be valuable at least in distinguishing new from ‘older’ genets. Our data suggested that both new (minimal somatic mutations) and veteran (minimum ages from 62-561 years) genets co-exist at these sites and that veteran Belize acroporids pre-date the widespread Caribbean collapse of acroporids. Future determination of microsatellite mutation rates in *Acropora* coral would help to narrow the currently rather wide range for age estimates.

Two of the veteran genets we identified are from hybrid samples. We estimated that these hybrid genets have existed for at least 156 and 281 years, respectively. It is thought that the hybrid only recently expanded its presence in Caribbean reef areas (Willis et al. 2006, Fogarty 2010, Richards and Hobbs 2015); however, no data on the ages of extant hybrid genets exist in

the literature. Here we show that at least a few hybrid genets originated before the Caribbean wide acroporid die-off event in the 1980s. It is unclear whether the persistence of *A. cervicornis*, *A. palmata*, and hybrid veteran genets at our sites is a function of: 1) their innate ability to rebound from disturbances (Eriksson 2000; Riegl et al. 2009); 2) whether certain colonies simply got lucky and survived environmental changes at random; or 3) whether environmental or oceanographic conditions at this site have been particularly conducive to acroporid persistence at these sites.

Our study sites are home to acroporid populations observed in high abundances in a time when they are facing Caribbean-wide decline. The existence of *Acropora* populations like these provides hope that perhaps these species and their hybrid can continue to persist. However, without analyzing the demographic processes of such populations, their potential resilience cannot be known. Here, we studied genotypic diversity and age in unison. The genotypic diversity of acroporids at our study sites was relatively low, which when considered alone does not bode well with respect to the potential of these populations to combat future environmental stresses (Schmid 1994; Steinger et al. 1996; Reusch et al. 2005; Garcia et al. 2008; Williams et al. 2008). However, we also know that some of the genets in these populations are relatively old and may have survived past environmental stresses, perhaps due to increased (and unknown) fitness attributes. Such veteran populations have hypothesized potential to expand (Noss 2001; Taberlet and Cheddadi 2002; Loarie et al. 2008; Keppel et al. 2012). Thus, we are left with two contradicting conclusions: the populations have low genotypic diversity so they are vulnerable to environmental change, and the populations have veteran genets whose past fitness may indicate that they will continue to survive in the future. In either scenario, these sites have potential to be valuable resources to *Acropora* spp. recovery initiatives, and should be investigated further. We

hope that the methods applied here can be useful to future studies that aim to identify sites of conservation priorities.

acknowledgments

This work was supported by the National Science Foundation under Grant No. NSF-REU #1358987 to LG, OCE-1537959 to IB, and the Keck Geology Consortium, the Howard Hughes Medical Institute through the Precollege and Undergraduate Science Education Program, as well as the Washington and Lee University Department of Geology, Office of the Provost, Johnson Opportunity Grant, and the Evans Fund for International Student Experiences to AI and LG. We wish to thank Kelly Bezold for laboratory work as well as Ken Mattes and Maureen Gannon at Belize Marine TREC for logistical support of this project. We are grateful for the support of the Belize Fisheries Department and Mr. James Azueta of the Belize Ecosystems Management Unit and Mr. Miguel Alamilla of the Hol Chan Marine Reserve for research permits to facilitate this work.

references

- Ally D, Ritland K, Otto SP (2008) Can clone size serve as a proxy for clone age? An exploration using microsatellite divergence in *Populus tremuloides*. *Mol Ecol* 17:4897-491
- Altizer S, Harvell D, Friedle E (2003) Rapid evolutionary dynamics and disease threats to biodiversity. *Trends Ecol Evol* 11:589-596
- Aronson RB, Precht WF (2001) White-band disease and the changing face of Caribbean coral reefs. *Hydrobiologia* 460:25-38
- Baums IB, Hughes CR, Hellberg MH (2005a) Mendelian microsatellite loci for the Caribbean coral *Acropora palmata*. *Mar Ecol Prog Series* 288:115-127

419 Baums IB, Miller MW, Hellberg ME (2005b) Regionally isolated populations of an imperiled
420 Caribbean coral, *Acropora palmata*. *Mol Ecol* 14(5): 1377-1390

421 Baums IB, Miller MW, Hellberg ME (2006) Geographic variation in clonal structure in a reef-
422 building Caribbean coral, *Acropora palmata*. *Ecol Monogr* 76(4): 503-519

423 Baums IB, Devlin-Durante MK, Polato NR, Xu D, Giri S, Altman NS, Ruiz D, Parkinson JE,
424 Boulay JN (2013) Genotypic variation influences reproductive success and thermal stress
425 tolerance in the reef building coral, *Acropora palmata*. *Coral Reefs* 32:703-717

426 Bellwood DR, Hughes TP, Folke C, Nystrom M (2004) Confronting the coral reef crisis. *Nature*
427 429:827-833

428 Brown-Saracino C, Peckol P, Curran HA, Robbart ML (2007) Spatial variation in sea urchins, fish
429 predators, and bioerosion rates on coral reefs of Belize. *Coral Reefs* 26: 71-78

430 Budd AF, Stemann TA, Johnson KG (1994) Stratigraphic distributions of genera and species of
431 Neogene to recent Caribbean reef corals. *J Paleont* 68:951-957

432 Busch J, Greer L, Harbor D, Wirth K, Lescinsky H, Curran HA (2016) Quantifying unusually large
433 populations of *Acropora* spp. corals in Belize using sub-meter multispectral satellite
434 imagery classification. *Bull Mar Sci* 92:265-283.

435 Carvalho GR (1994) Genetics of aquatic clonal organisms. In: *Genetics and Evolution of Aquatic*
436 *Organisms* (ed. Beaumont AR), pp. 291-323. Chapman and Hall, London

437 Devlin-Durante MK, Miller MW, Caribbean Acropora Research, Precht WF, Baums IB (2016)
438 How old are you? Genet age estimates in a clonal animal. *Mol Ecol* 25(22): 5628-5646

439 de Witte LC, Stocklin J (2010) Longevity of clonal plants: why it matters and how to measure it.
440 *AoB Plants* 106:859-870

441 Downing A, van Nes E, Mooij WM, Scheffer M (2012) The resilience and resistance of an
442 ecosystem to a collapse of diversity. *PLoS ONE* 7(9)

443 Eakin CM, Morgan JA, Heron SF, Smith TB, Liu G, Alvarez-Filip L, Baca B, Bartels E,
444 Bastidas C, Boucon C, Brandt M, Bruckner AW, Bunkley-Williams L, Cameron A,
445 Causey BD, Chiappone M, Christensen TRL, Crabbe MJC, Day O, Guardia E, Diaz-
446 Pulido G, DiResta D, Gil-Agudelo DL, Gilliam DS, Ginsburg RN, Gore S, Guzman HM,
447 Hendee JC, Hernandez-Delgado EA, HusainE, Jeffrey CFG, Jones RJ, Jordan-Dahlgren
448 E, Kaufman LS, Kline DI, Kramer PA, Lang JC, Lirman D, Mallela J, Manfrino C,
449 Marechal JP, Marks K, Mihaly J, Miller WJ, Mueller EM, Muller EM, Toro CAO,
450 Oxenford HA, Ponce-Taylor D, Quinn N, Ritchie KB, Rodriguez S, Ramirez AR,
451 Romano S, Samhuri JF, Sanchez JA, Schmal GP, Shank BV, Skirving WJ, Steiner SCC,
452 Villamizar E, Walsh SM, Walter C, Weil E, Williams EH, Roberson KW, Yusuf Y
453 (2010) Caribbean Corals in Crisis: Record Thermal Stress, Bleaching, and Mortality in
454 2005. *PLoS ONE* 5(11)

455 Eriksson O (2000) Functional roles of remnant plant populations in communities and
456 ecosystems. *Glob Ecol Biogeog* 9(6):443-449

457 Fogarty ND (2010) Reproductive Isolation and Hybridization Dynamics in Threatened
 458 Caribbean Acroporid Corals. Doctoral dissertation. Nova Southeastern University.
 459 Retrieved from NSUWorks. (4)

460 Fogarty ND (2012) Caribbean acroporid coral hybrids are viable across life history stages. *Mar*
 461 *Ecol Prog Ser* 446:145-159

462 Fogarty ND, Vollmer SV, Levitan DR (2012) Weak prezygotic isolating mechanisms in
 463 threatened Caribbean *Acropora* corals. *PLoS ONE* 7(2): 1-11

464 Garcia MB, Pico FX, Ehrlén J (2008) Life span correlates with population dynamics in perennial
 465 herbaceous plants. *Am J Bot* 95(2):258-262

466 Gardner TA, Cote IM, Gill JA, Grant A, Watkinson AR (2003) Long-term region-wide declines
 467 in Caribbean corals. *Science* 301:958-960

468 Gladfelter WB (1982) White-band disease in *Acropora palmata*: implications for the structure
 469 and growth of shallow reefs. *Bull Mar Sci* 32:639-643

470 Harper JL (1977) *Population Biology of Plants*. Academic Press, London.

471 Highsmith RC (1982) Reproduction by fragmentation in corals. *Mar Ecol Prog Series* 7: 207-226

472 Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD,
 473 Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N,
 474 Bradbury RH, Dubi A, Hatziolos ME (2007) Coral reefs under rapid climate change and
 475 ocean acidification. *Science* 318(5857):1737-1742

476 Hughes RN (1989) *A Functional Biology of Clonal Animals*. Chapman and Hall, London and
 477 New York.

478 Hughes TP (1994) Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral
 479 reef. *Science* 265:1547-1551

480 Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-
 481 Guldberg O, Jackson JBC, Kleypas J, Lough JM, Marshall P, Nystrom M, Palumbi SR,
 482 Pandolfi JM, Rosen B, Roughgarden J (2003) Climate Change, Human Impacts, and the
 483 Resilience of Coral Reefs. *Science* 301:929-933

484 Japaud A, Bouchon C (2014) Unexpected high densities of the hybrid coral *Acropora prolifera*
 485 (Lamarck 1816) in Guadeloupe Island, Lesser Antilles. *Coral Reefs* 33:593

486 Kays S, Harper JL (1974) The regulation of plant and tiller density in a grass sward. *J Ecol*
 487 63:97-105

488 Keck J, Houston RS, Purkis S, Riegl BM (2005) Unexpectedly high cover of *Acropora*
 489 *cervicornis* on offshore reefs in Roatán (Honduras). *Coral Reefs* 24:509

490 Keppel G, Val Niel KP, Wardell-Johnson GW, Yates CJ, Byrne M, Mucina L, Schut AGT,
 491 Hopper SD, Franklin SE (2012) Refugia: identifying and understanding safe havens for
 492 biodiversity under climate change. *Glob Ecol Biogeogr* 21:393-404

493 Kuffner IB, Lidz BH, Hudson JH, Anderson JS (2015) A century of ocean warming on Florida
 494 Keys coral reefs: historic in situ observations. *Estuaries and Coasts* 38: 1085-1096

495 Lang JC, Alcolando P, Carricart-Ganivet JP, Chippone M, Curran A (1998) Status of coral reefs
 496 in the northern areas of the world, Vol. Australian Institute of Marine Science,
 497 Townsville

498 Larson EA, Gilliam DS, Padierna ML, Walker BK (2014) Possible recovery of *Acropora*
 499 *palmata* (Scleractinia: Acroporidae) within the Veracruz Reef System, Gulf of Mexico: a
 500 survey of 24 reefs to assess the benthic communities. *Rev Biol Trop* 62 (3): 75-84

501 Levitan DR, McGovern TM (2005) The Allee effect in the Sea. Pages 47-57 in E.A. Norse and
 502 I.B. Crowder, eds. *Mar Cons Biol. The Science of Maintaining the Sea's Biodivers Isl*
 503 Press

504 Lirman D (2000) Fragmentation in the branching coral *Acropora palmata* (Lamarck): growth,
 505 survivorship, and reproduction of colonies and fragments. *J Exp Mar Bio Ecol* 25: 41-57

506 Lirman D, Bowden-Kerby A, Schopmeyer S, Huntington B, Thyberg T, Gough M, Gough T,
 507 Gough R, Gough Y (2010) A window to the past: documenting the status of one of the
 508 last remaining 'megapopulations' of the threatened staghorn coral *Acropora cervicornis*
 509 in the Dominican Republic. *Aquat Conserv* 20(7):773-781

510 Loarie SR, Carter BE, Hayhoe K, McMahon S, Moe R, Knight CA, Ackerly DD (2008) Climate
 511 change and the future of California's endemic flora. *PLoS ONE* 3(6)

512 Macintyre IG, Toscano MA (2007) The Elkhorn Coral *Acropora palmata* is coming back to the
 513 Belize Barrier Reef. *Coral Reefs* 26:757

514 NOAA (National Oceanic and Atmospheric Administration) (2005) Endangered and Threatened
 515 Wildlife and Plants; 12-Month Finding on a Petition to List Elkhorn Coral, Staghorn
 516 coral, and Fused-staghorn coral as Threatened or Endangered. Code Fed Regul 70:13151-
 517 13152

518 Noss RF (2001) Beyond Kyoto: Forest management in a time of rapid climate change. *Conserv*
 519 *Biol* 15(3):578-590

520 van Oppen MJH, Willis BL, Van Vugt H, Miller DJ (2000) Examination of species boundaries in
 521 the *Acropora cervicornis* group (Scleractinia, Cnidaria) using nuclear DNA sequence
 522 analyses. *Mol Ecol* 9:1363-1373

523 Pandolfi J (2002) Coral community dynamics at multiple scales. *Coral Reefs* 21:13-23

524 Peckol PM, Curran HA, Floyd EY, Robbart ML, Greenstein BJ, Buckman KL (2003)
 525 Assessment of selected reef sites in northern and southern-central Belize, including
 526 recovery from bleaching and hurricane disturbances (stony corals, algae and fish). *Atoll*
 527 *Res Bull* 496: 146-171

528 Purkis SJ, Myint SW, Riegl BM (2006) Enhanced detection of the coral *Acropora cervicornis*
 529 from satellite imagery using a textural operator. *Remote Sens Environ* (101):82-94

530 Randall CJ, van Woesik R (2015) Contemporary white-band disease in Caribbean corals driven
 531 by climate change. *Nat Clim Chang* 5:375-379

532 Reed DH, Frankham R (2003) Correlation between fitness and genetic diversity. *Conserv Biol*
533 17:230-237

534 Reusch TBH, Ehlers A, Hammerli A, Worm B (2005) Ecosystem recovery after climatic
535 extremes enhanced by genotypic diversity. *Proc Natl Acad Sci USA* 102:2826-2931

536 Richards ZT, Hobbs JPA (2015) Hybridisation on coral reefs and the conservation of
537 evolutionary novelty. *Curr Zool* 2015

538 Riegl B, Purkis SJ, Keck J, Rowlands GP (2009) Monitored and modeled coral population
539 dynamics and the refuge concept. *Mar Poll Bul* 58:24-38

540 Schmid B (1994) Effects of genetic diversity in experimental stands of *Solidago altissima*:
541 evidence for the potential role of pathogens as selective agents in plant-populations. *J*
542 *Ecol* 82:165–175

543 Steinger T, Korner C, Schmid B (1996) Long-term persistence in a changing climate: DNA
544 analysis suggest very old ages of Alpine *Carex curvula*. *Oecologia* 105(1):94-99

545 Stenberg P, Lundmark M, Saura A (2003) MLGsim: a program for detecting clones using a
546 simulation approach. *Molecular Ecology Notes* 3, 329-331.

547 Szmant AM (1986) Reproductive ecology of Caribbean reef corals. *Coral Reefs* 5:43-53

548 Stoddart JA, Taylor JF (1988) Genotypic diversity: estimation and prediction in samples.
549 *Genetics* 118:705-711

550 Taberlet P, Cheddadi R (2002) Quaternary refugia and persistence of biodiversity. *Science*
551 297:2009-2010

552 Vargas-Angel B, Thomas JD, Hoke SM (2003) High-latitude *Acropora cervicornis* thickets off
553 Fort Lauderdale, Florida, USA. *Coral Reefs* 22:464-473

554 Vollmer SV, Palumbi SR (2002) Hybridization and the evolution of reef coral diversity. *Science*
555 296:2023-2025

556 Weiher E van der Werf A, Thompson K, Roderick M, Garnier E, Eriksson O (1999) Challenging
557 Theophrastus: A common core list of plant traits for functional ecology. *Journal of*
558 *Vegetation Science* 10(5):609-620

559 Wheaton J, Callahan M, Brooke S, Beaver C, Wade S, Johnson D, Kupfner S, Kidney J, Bertin
560 M (2006) Dry Tortugas National Park (DRTO) Long term monitoring and assessment
561 project annual report 2005-2006. Fish and Wildlife Research Institute, St. Petersburg, FL

562 Williams DE, Miller MW, Kramer KL (2008) Recruitment failure in Florida Keys *Acropora*
563 *palmata*, a threatened Caribbean coral. *Coral Reefs* 27:697-705

564 Williams DE, Miller MW, Baums IB (2014). Cryptic changes in the genetic structure of a highly
565 clonal coral population and the relationship with ecological performance. *Coral Reefs*
566 33:595-606

567 Willis BL, van Oppen M.J.H, Miller DJ, Vollmer SJ, Ayre DJ (2006) The role of hybridization
568 in the evolution of reef corals. *Annu Rev Ecol Syst* 37:489-517

Zubillaga AL, Marquez LM, Croquer A, Bastidas C (2008) Ecological and genetic data indicate recovery of the endangered coral *Acropora palmata* in Los Roques, Southern Caribbean. Coral Reefs 27:63-72

figure captions

Fig. 1 Caribbean *Acropora* species: A) *Acropora cervicornis*, or Staghorn coral; B) *A. palmata*, or Elkhorn coral; C) “*A. prolifera*”, the hybrid of *A. cervicornis* and *A. palmata*, also known as “Fused Staghorn” coral; all from Coral Gardens, Belize.

Fig. 2 Map of Belize, inlaid with map of study sites Coral Gardens, Manatee Channel, and Rocky Point. Reef area is depicted in blue.

Fig. 3 Comparison of genotypic diversity and evenness metrics calculated in our study with those calculated by Baums et al (2006) for Caribbean populations of *A. palmata*, with inference on demography of local recruitment. Values for our locations are labeled by site (MC, *Manatee Channel*; CG, *Coral Gardens*; RP, *Rocky Point*; BZ, Belize i.e. all sites combined) and color coded by species (blue, *A. cervicornis*; red, *A. palmata*; green, hybrid). Data from Baums et al. (2006) are shown as unfilled circles. Ellipses grouping sites according to relative degree of sexual versus asexual recruitment are redrawn from Figure 4 in Baums et al. (2006).