

Floating topological phases

Trithip Devakul  and S. L. Sondhi*Department of Physics, Princeton University, Princeton, New Jersey 08544, USA*S. A. Kivelson *Department of Physics, Stanford University, Stanford, California 94305, USA*

Erez Berg

Department of Physics, Weizmann Institute of Science, Rehovot 76100, Israel

(Received 13 July 2020; accepted 31 August 2020; published 21 September 2020)

While quasi-two-dimensional (layered) materials can be highly anisotropic, their asymptotic long-distance behavior generally reflects the properties of a fully three-dimensional phase of matter. However, certain topologically ordered quantum phases with an emergent $(2 + 1)$ -dimensional gauge symmetry can be asymptotically impervious to interplane couplings. We discuss the stability of such “floating topological phases,” as well as their diagnosis by means of a nonlocal order parameter. Such a phase can produce a divergent ratio $\rho_{\perp}/\rho_{\parallel}$ of the interlayer to intralayer resistivity as $T \rightarrow 0$, even in an insulator where both ρ_{\perp} and ρ_{\parallel} individually diverge. Experimental observation of such a divergence would constitute proof of the existence of a topological (e.g., spin-liquid) phase.

DOI: [10.1103/PhysRevB.102.125136](https://doi.org/10.1103/PhysRevB.102.125136)

I. INTRODUCTION

Although bulk materials are, obviously, three dimensional, it is common to encounter materials with highly anisotropic electronic structure, either quasi-one-dimensional (quasi-1D) or quasi-two-dimensional (quasi-2D) structures. Naturally, macroscopic thermodynamic correlation functions and linear response properties of such materials are highly anisotropic, reflecting the microscopic anisotropy. What is less obvious, but an intriguing possibility, is that such anisotropic systems might exhibit one form or another of a “floating phase,” i.e., a phase that exhibits asymptotic properties characteristic of a lower-dimensional system [1–13]. For instance, the ground state of a one-dimensional electron gas (1DEG) generically exhibits power-law correlations (e.g., Luttinger liquid or Luther-Emery liquid behavior [14,15]); a quantum floating phase of a quasi-1D array of weakly coupled wires, were such a phase to exist, would be a $T = 0$ phase which exhibits power-law correlations along the wires, but exponentially falling correlations in the transverse directions [7,8,10]. A classical ($T > 0$) floating phase that has been considered [3–6] in a quasi-2D material of stacked planes, would exhibit power-law charge-density-wave (CDW) or superconducting (SC) correlations in plane that fall exponentially with distance perpendicular to the planes.

Disappointingly, for the most part whenever a lower-dimensional system exhibits power-law correlations, arbitrarily weak higher-dimensional couplings cause the onset of long-range order rather than a floating phase. This reflects the fact that power-law phases are often critical in the sense that the susceptibility to one or another form of broken symmetry

is infinite. Even in more mundane phases, either disordered or long-range ordered, in which the connected correlation functions fall exponentially with distance, the correlation length characterizing that falloff can be anisotropic, reflecting the microscopic anisotropy of the system, but it is ultimately governed by a single three-dimensional (3D) Ornstein-Zernike form at long enough distances.

There do appear to be special extreme circumstances in which such floating phases may be stable phases of matter [16]. In most cases, however, the best one can hope for in the standard phases of matter is to find circumstances in which the simplest (and largest) higher-dimensional couplings are in some way frustrated and rendered irrelevant, so that $(3+1)$ D behavior is only apparent beyond some extremely long (emergent) length scale.

The situation is more promising for topological phases. Consider a stack of initially decoupled planes, each characterized by some form of topological order. If we were to turn on interplane couplings, this can affect all sorts of local correlations, but cannot effect the topological order. Thus, as we will discuss (formalizing and extending ground-breaking work of Senthil and Fisher [17]), if the couplings are weak enough, they cannot couple the topological order associated with each plane. The topologically nontrivial properties of the state in the absence of interplane coupling, and in particular the fact that there is an emergent gauge symmetry associated with each plane individually, survive in 3D, provided the microscopic couplings are sufficiently anisotropic. In short, “floating topological phases” (as defined below) are predicted to be a generic feature of topological phases when (and if) they exist in the lower-dimensional context.

From a more direct empirical perspective, a standard measure of the macroscopic anisotropy of a system is the ratio of transport coefficients in various directions, for instance, the ratio of the values of the resistivity tensor $\rho_{\alpha\beta}$ along the different principal axes of a crystal. In all conventional phases of matter, including a band insulator, a diffusive metal, and a disordered insulator (in the regime where it exhibits variable range hopping), even when the resistivity tensor is highly anisotropic, the ratio ρ_{cc}/ρ_{aa} approaches a finite constant as $T \rightarrow 0$. (Here and henceforth we take c to be the most resistive direction and a the least.) In contrast, in a floating phase, when the in-plane and interplane currents are carried by different combinations of elementary excitations, this ratio can diverge as $T \rightarrow 0$. Note that this distinction can apply in an insulator; even when both ρ_{aa} and ρ_{cc} diverge as $T \rightarrow 0$, it is still possible to distinguish cases in which their ratio approaches a constant (as in conventional anisotropic phases of matter), or diverges. As was pointed out previously [18], divergent anisotropies in thermal transport are also expected, although a typically large phonon contribution may mask the effect.

A diverging anisotropy ratio would certainly require some form of dynamical dimensional decoupling for which a more or less exotic explanation would be needed. In the early days of cuprate high-temperature superconductivity, Anderson and Zou [19] proposed that just such a divergent resistivity anisotropy should be expected based on ideas of spin-charge separation. Experiments carried out at high magnetic fields (to suppress the superconductivity) indeed were initially suggestive that this indeed occurs [20]. More recently [21,22], in the stripe-ordered cuprate, LBCO, a form of dynamical layer decoupling has been observed, in which the in-plane resistivity ρ_{aa} drops rapidly below a well-defined onset temperature T_{onset} , becoming immeasurably small below an apparent in-plane superconducting transition temperature T_{2d} , while ρ_{cc} , the resistivity in the out-of plane direction, remains large and only vanishes at a substantially lower critical temperature $T_{3d} < T_{2d}$. The resistivity anisotropy increases by at least three orders of magnitude between T_{onset} and T_{2d} , and is equal to infinity within experimental error for $T_{2d} > T > T_{3d}$. For reasons that we will review and expand upon, it was suggested that a search for a related divergent anisotropy, in this case of the thermal conductivity tensor, could be used as a clear experimental signature of the existence of certain kinds of spin-liquid phases. Similar ideas were explored in the context of stacked quantum Hall layers by Balents and Fisher [23], Naud *et al.* [24,25], and Levin and Fisher [26].

In this paper we offer a precise theoretical definition of a topological floating phase. We propose that a form of absolute stability, even in the presence of (weak) higher-dimensional couplings, may be considered as a defining feature of certain sorts of spin-liquid phases. Floating topological phases are also closely related to fracton topological ordered phases [27–30], which have attracted much theoretical attention in recent years [31,32]. Fracton topologically ordered phases host fractionalized quasiparticles with restricted mobility and a subextensive ground-state degeneracy, all properties shared by floating topological phases. The phases studied in this paper are therefore very basic versions of fracton phases.

This paper is organized as follows. We define and discuss gapped floating topological phase in Sec. II, as well as their diagnostic in terms of nonlocal correlation functions in Sec. III. In Sec. IV, we extend this analysis to various gapless floating topological phases, and discuss their stability to interlayer couplings. In Sec. V, we demonstrate that a diverging conductivity ratio may serve as an experimentally accessible signature of a floating topological phase. Finally, we end with some concluding remarks in Sec. VI.

II. GAPPED TOPOLOGICAL FLOATING PHASES

We begin with the simplest possible case: a floating phase of gapped topological orders in two spatial dimensions [i.e., (2+1)D including the time dimension]. In such models, topological properties such as the ground-state degeneracy and statistical properties of fractionalized quasiparticle excitations are stable to arbitrary interlayer couplings. As the models we will consider later take the form of stacked gauge theories coupled to matter, we will first begin with a simplest such example: the stack of (2+1)D Ising gauge theories coupled to Ising matter. This model is equivalent to a stack of (2+1)D Kitaev toric codes [33].

The (2+1)D Ising gauge theory (IGT) on the square lattice is described [34,35] by a model with matter degrees of freedom τ_r living on the sites r , and gauge degrees of freedom σ_ℓ living on the links $\ell = (rr')$ connecting nearest-neighbor sites r and r' . The IGT Hamiltonian is

$$H_{\text{IGT}} = -K \sum_{\square} \prod_{\ell \in \square} \sigma_\ell^z - \Gamma \sum_{\ell} \sigma_\ell^x - J \sum_{(rr')} \tau_r^z \sigma_{(rr')}^z \tau_{r'}^z - \Gamma_M \sum_r \tau_r^x, \quad (1)$$

where the first sum is over square plaquettes, and $\sigma^{x,y,z}$ and $\tau^{x,y,z}$ are Pauli matrices acting on the σ and τ degrees of freedom, respectively. Here the first two terms are the ‘‘Maxwell’’ terms, and the remaining terms are the gauge-invariant matter terms. This Hamiltonian is invariant under local gauge transformations $G_r^\dagger H G_r = H$, generated by

$$G_r = \tau_r^x \prod_{r'} \sigma_{(rr')}^x \quad (2)$$

on each site r , where the product is over the four nearest neighbors. We take the physical subspace to be the one with $G_r = 1$ on every site. $K > 0$ favors a zero-flux ground state, and Γ makes the gauge field dynamical.

This model describes a deconfined \mathbb{Z}_2 gauge theory when $K/\Gamma \gg 1$ and $\Gamma_M/J \gg 1$. In this limit, H_{IGT} describes a perturbed version of Kitaev’s toric-code model [33]; this can be seen by working in the gauge $\tau_r^z = 1$, in which H_{IGT} can be expressed entirely in terms of the gauge fields

$$\begin{aligned} \tilde{H}_{\text{IGT}} &= -K \sum_{\square} \prod \sigma^z - \Gamma_M \sum_{+} \prod \sigma^x \\ &\quad - J \sum_{\ell} \sigma_\ell^z - \Gamma \sum_{\ell} \sigma_\ell^x \\ &= H_{\text{TC}} - J \sum_{\ell} \sigma_\ell^z - \Gamma \sum_{\ell} \sigma_\ell^x, \end{aligned} \quad (3)$$

where the K and Γ_M terms are the stabilizers of the toric-code Hamiltonian H_{TC} , and J and Γ are small σ^z and σ^x perturbations that make the model dynamical.

The toric-code Hamiltonian H_{TC} possesses nontrivial topological order which is stable to arbitrary local perturbations [36]. When placed on a torus, H_{TC} has four exactly degenerate ground states in the thermodynamic limit. These ground states may be distinguished via nonlocal Wilson and 't Hooft operators. Define the Wilson loop operator

$$W_C = \prod_{l \in C} \sigma_l^z, \quad (4)$$

where C denotes a closed loop on the square lattice, and $l \in C$ are all the links involved. We may also define the dual Wilson loop (or 't Hooft operator)

$$V_{\bar{C}} = \prod_{l \in \bar{C}} \sigma_l^x, \quad (5)$$

where \bar{C} denotes a loop on the dual square lattice, and $l \in \bar{C}$ are all the links cut by \bar{C} . These operators commute with H_{TC} . Let W_1 and V_1 denote the noncontractible Wilson loops going around the torus in the x direction, and similarly W_2 and V_2 along y . The operators (W_1, V_2) and (W_2, V_1) generate Pauli algebras on the four-dimensional ground-state manifold. The ground-state degeneracy is stable to J and Γ perturbations due to the fact that these noncontractible Wilson loop operators only appear at high order $O(L)$ in perturbation theory where L is the circumference of the torus; any lifting of the degeneracy is thus exponentially suppressed at large L , going as $\sim (J/\Gamma_M)^L$ or $\sim (\Gamma/K)^L$.

Next, consider the bilayer of two such systems, with some weak coupling between them

$$H = H_{\text{IGT}}^{(1)} + H_{\text{IGT}}^{(2)} + \lambda H_{\text{inter}}, \quad (6)$$

where H_{inter} contains local terms coupling the two layers. On a torus, this bilayer now has a 4^2 -fold degenerate ground-state manifold, which is only split perturbatively by the interlayer couplings at order λ^L as before. This simply describes a new topological order, which inherits all its topological properties from the stack of two decoupled toric codes. Indeed, this is simply the $\mathbb{Z}_2 \times \mathbb{Z}_2$ generalization of the toric code (which describes the gauge theory of a bilayer Ising model in which each layer has a separate \mathbb{Z}_2 symmetry).

Now, let us consider a $(3+1)$ D system on a 3-torus obtained by stacking L such models in the xy plane along the z direction, allowing for small (but arbitrary) local perturbations. This model will have a robust 4^L ground-state degeneracy which is stable to any arbitrary small interlayer interactions. We define a $(D+1)$ -dimensional system to be in a nontrivial gapped floating topological phase if it can be smoothly connected (via a finite depth local unitary transformation) to a decoupled stack of $(d+1)$ -dimensional topologically ordered systems, where $0 < d < D$. In the cases we consider, $D = 3$ and $d = 2$. The stack of toric-code models, with weak interlayer coupling terms, realizes a nontrivial floating topological phase by this definition.

Quasiparticle excitations of this model are constrained to move within a single $(2+1)$ D xy plane. As an aside, we note that decoupled stacks of topologically ordered planes,

exactly as we have formulated, have appeared multiple times [37,38] in the literature of fracton topological order [31,32]. If we take the definition of a fracton topological order to be a subextensive ground-state degeneracy $\ln \text{GSD} \sim L$ on a 3-torus and subdimensional quasiparticle excitations, then the stack of $(2+1)$ D topological orders are indeed (very simple versions of) fracton models. These may also be obtained as gauge theories of models with planar subsystem symmetries along each xy plane (a 1-foliated planar subsystem symmetry [39]).

The low-energy field theory description of the single-layer toric code is given by the $(2+1)$ D BF theory, or equivalently the Chern-Simons theory [40,41]

$$\mathcal{L}_{\text{CS}} = \frac{K_{IJ}}{4\pi} \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J \quad (7)$$

with the 2×2 \mathbf{K} matrix $\mathbf{K}^{\text{TC}} = 2\sigma^x$. The ground-state degeneracy of such a model on a manifold of genus g is then given by $\text{GSD} = |\det \mathbf{K}|^g$. The bilayer toric code described by the Hamiltonian (6) then admits a similar low-energy description, except with the 4×4 \mathbf{K} matrix given by the direct sum $\mathbf{K}^{2\text{TC}} = \mathbf{K}^{\text{TC}} \oplus \mathbf{K}^{\text{TC}}$. The floating topological phase of toric codes discussed above may then be characterized by the extensively large \mathbf{K} matrix $\mathbf{K}^{\text{float}} = \mathbf{K}^{\text{TC}} \oplus \dots \oplus \mathbf{K}^{\text{TC}}$ (such ‘‘giant’’ \mathbf{K} matrices also appear in the classification of fracton phases [42]). This can be generalized to stacks of general Abelian topological phases characterized by the matrix \mathbf{K} [43].

We also note the interesting possibility of off-diagonal elements in the large- \mathbf{K} matrix which couples different layers. These types of systems have been studied [24,25,44,45] and found to exhibit interesting behavior (such as an irrational braiding statistic) which cannot be found in $(2+1)$ D systems. Such systems are not floating topological phases by our definition as they cannot be deformed to the decoupled limit, but nevertheless have an emergent decoupled gauge symmetry.

Finally, most of our discussion can also be extended to stacks of non-Abelian topological orders.¹ Such a phase can be characterized by the topological properties of the quasiparticle excitations, such as the fusion coefficients and topological spin [47]. Stacking two layers results in a new phase whose quasiparticles are directly inherited from the individual layers (see Ref. [47]), and survive interlayer couplings as long as the gap is not closed. The floating phase of gapped non-Abelian topological orders is therefore also stable.

III. FLOATING PHASES VIA THE FREDENHAGEN-MARCU ORDER PARAMETER

The goal of this section is to differentiate a floating topological phase from either the $(3+1)$ D topological order or the trivial phase by means of a correlation function.

¹More specifically, we concern ourselves with only *non-invertible* topological orders, which possess fractionalized quasiparticles. Stacks of invertible topological phases [46] need not form stable floating phases.

A. Usual deconfinement diagnostic

We first review how this is done in the usual case of diagnosing deconfinement in the Ising gauge theory [48]. Let us take $W(L)$ to be the Wilson loop defined in Eq. (4) along the contour C taken to be an $L \times L$ square. For a pure gauge theory without dynamical matter [$J = 0$ in Eq. (3)], the scaling of the expectation value of the Wilson loop is sufficient to diagnose deconfinement: for large L , $\ln\langle W(L) \rangle \sim -L$ scales linearly with the perimeter of the loop in the deconfined phase, but in the confined phase scales with the area $\ln\langle W(L) \rangle \sim -L^2$. However, as soon as $J \neq 0$, the Wilson loop scales with the perimeter in both the deconfined and confined phases, and therefore fails to distinguish between the two.

To correct this shortcoming, consider (an equal time formulation of) the Fredenhagen-Marcu order parameter [48,49] (FMOP). Define

$$W_{\frac{1}{2}}(L) \equiv \tau_r^z \tau_{r'}^z \prod_{l \in C_{\frac{1}{2}}} \sigma_l^z \quad (8)$$

to be a gauge-invariant open Wilson line (or horseshoe) operator, where $C_{\frac{1}{2}}$ is the $L \times L/2$ horseshoe terminated at sites r, r' , obtained by cutting C in half. Similarly, let $W_{-\frac{1}{2}}$ be the other half of the Wilson loop. We note that a more general geometry is possible; we simply choose horseshoe shape for simplicity. It can be shown (see below) that the ratio

$$R(L) = \langle W_{\frac{1}{2}}(L) \rangle \langle W_{-\frac{1}{2}}(L) \rangle / \langle W(L) \rangle \quad (9)$$

goes to 0 in the deconfined phase, while $\lim_{L \rightarrow \infty} R(L) = R_0 > 0$ in the confined phase. Thus, other than in the special case $J = 0$, this provides a suitable signature of a deconfined phase.

This behavior can be understood in many ways. If we adopt the same gauge choice as in Eq. (3), we can express the ground state of the Hamiltonian in the σ^x basis as

$$|\psi\rangle = \sum_c \alpha_c |c\rangle, \quad (10)$$

where $c = \{\sigma_l^x\}$ label all the configurations. The ground state of the toric code is the equal amplitude superposition of all configurations of closed $\sigma^x = -1$ loops, with zero amplitude for all other c ; it is a loop condensate. With perturbations, the weights of each configuration in the ground state are no longer exactly equal and configurations with open strings now exist, albeit with weights that are exponentially small in the separation between the two end points ℓ going as $\sim (J/\Gamma_M)^\ell$.

The expectation value of the horseshoe operator [Eq. (8)]

$$\langle W_{\frac{1}{2}} \rangle = \sum_c \alpha_c^* \alpha_{c'}, \quad (11)$$

where $|c'\rangle = W_{\frac{1}{2}} |c\rangle$ is the configuration c with σ^x flipped along the support of $W_{\frac{1}{2}}$. [The terminal factors of τ_r^z in Eq. (8) are set equal to 1 by the choice of gauge.] There is an analogous expression for the expectation value of W . To see how these considerations distinguish the two phases, we use this expression to compute $R(L)$ at points deep inside the respective phases.

Manifestly, $\langle W_{\frac{1}{2}}(L) \rangle$ vanishes in the toric-code ground state since $W_{1/2}$ generates a string from r to r' , meaning that for

any c such that $\alpha_c \neq 0$, $\alpha_{c'} = 0$. In the perturbed problem, $\langle W_{\frac{1}{2}} \rangle$ is not identically zero, but decays exponentially with L . The form of its decay can be derived using perturbation theory in both J and Γ : $\langle W_{\frac{1}{2}}(L) \rangle \sim e^{-(2a+b)L}$. Here, $a \sim (\Gamma/K)$ comes from the sides of the horseshoe and $b \sim -\ln(J/\Gamma_M)$ comes from the string from r to r' . The Wilson loop scales as $\langle W(L) \rangle \sim e^{-4aL}$. Thus, $R(L) \sim e^{-2bL} \rightarrow 0$ in the deconfined phase.

To characterize the confined phase, consider the ground state of Eq. (3) in the large- Γ limit. Here, the significant configurations are those which are mostly polarized with $\sigma^x = +1$, plus small loop fluctuations [suppressed by factors of $(K/\Gamma)^A$ where A is the enclosed area] and open line fluctuations [exponentially suppressed in their length as $(J/\Gamma)^\ell$]. For large L , $\langle W_{1/2}(L) \rangle \sim (J/\Gamma)^{2L}$ and $\langle W(L) \rangle \sim (J/\Gamma)^{4L}$. Hence, $R(L) \rightarrow \text{const}$ in the large- L limit in the confined phase. Some intuition can be gained along the special $\Gamma = 0$ axis, where $R(L)$ is (the square of) the original Ising $\sigma^x \sigma^z$ correlation function, and the approach to a constant can be understood in terms of spontaneous symmetry breaking. The generalization of this FMOP construction beyond \mathbb{Z}_2 is possible and discussed by Gregor *et al.* [48].

B. Floating deconfinement diagnostic

We now turn our discussion to the problem of diagnosing a floating phase of (2+1)D topological orders from either a confining phase or a fully (3+1)D topological ordered phase. To do this we consider an anisotropic 3D generalization of the gauge-fixed version of the Ising gauge theory [Eq. (3)] with qubits defined on the nearest-neighbor bonds of a tetragonal lattice, with couplings K, J, Γ for the in-plane terms, and $K_\perp, J_\perp, \Gamma_\perp$ for those involving bonds in the interplane direction. This system now has the same Hilbert space as that of the (3+1)D toric code. Indeed, for $J = J_\perp = \Gamma = \Gamma_\perp = 0$ and $K = K_\perp$, this model reduces to the (3+1)D toric code, and to stacks of (2+1)D toric codes if we then take the limit $\Gamma_\perp \rightarrow \infty, K_\perp \rightarrow 0$.

It is thus clear that in different limits, this one model can support all three of the possible phases in question. Our diagnostic for floating topological order is inspired by the usual deconfinement diagnostic just discussed.

Let $W(z; L)$ be the Wilson loop on a $L \times L$ loop in the z th plane. We further split the loop into two equal horseshoes such that $W(z; L) = W_{\frac{1}{2}}(z; L) W_{-\frac{1}{2}}(z; L)$, where $W_{\frac{1}{2}}$ is defined on the left horseshoe and $W_{-\frac{1}{2}}$ on the right. We then consider the ratio

$$R_2(L) = \frac{W_{\frac{1}{2}}(z; L) W_{-\frac{1}{2}}(z+1; L)}{\sqrt{W(z; L) W(z+1; L)}}. \quad (12)$$

As we will see, in analogy with the previous analysis,

$$\lim_{L \rightarrow \infty} R_2(L) = \begin{cases} 0, & \text{deconfined floating} \\ \text{const}, & \text{confined or (3+1)D deconfined} \end{cases} \quad (13)$$

distinguishes between the deconfined floating topological phase from a confined or fully (3+1)D deconfined topological phase.

First, consider the decoupled limit with all the interplane couplings set equal to 0: now, $R_2(L)$ factors as $R_2(L) = R(L)$,

where $R(L)$ is the FMOP for a single (2+1)D plane defined in Eq. (9). Thus, in this case, for the deconfined phase, $R_2(L) \rightarrow 0$, while for the confining phase $R_2(L) \rightarrow \text{const} > 0$. It is easy to see that these results remain true even in the presence of arbitrary small perturbations.

Finally, consider the case of an isotropic (3+1)D phase. The numerator is now the expectation value of a Wilson loop minus only the two vertical bonds: the full Wilson loop thus differs from this by a factor which is order $O(1)$ (i.e., independent of L). Moreover, this Wilson loop can be viewed as a slightly distorted relative of the the Wilson loop in the denominator other except for two ‘‘kinks’’ where it changes over between planes z and $z + 1$. These kinks also make an $O(1)$ correction to the total expectation value of the Wilson loop. The scaling of numerator and denominator again cancel, so independent of whether the (3+1)D system is confining or not, $R_2(L) \rightarrow \text{const}$. In short, the vanishing of $R_2(L)$ at large L is a signature of a topological floating phase.

Owing to the emergent subdimensional Lorentz symmetry of the floating topological phase, the order parameter may be oriented in various space-time directions (within the x, y, τ subspace), each of which have a physical interpretation [48]. The order parameter (12) corresponds to an equal time-slice orientation. We will now discuss another orientation of this order parameter (which in the usual case corresponded to the order parameter discovered by Fredenhagen and Marcu [49]). Let us denote by $|\psi_z\rangle$ a trial state with two (spinon) excitations located at \vec{r} and $\vec{r}' = \vec{r} + L\hat{x}$ on plane $r_z = z$, defined by

$$|\psi_z\rangle = \tau_r^z \tau_{r'}^z V_{rr'}(-T/2)|\text{GS}\rangle, \quad (14)$$

where $V_{rr'}(-T/2) \equiv e^{-HT/2} V_{rr'} e^{HT/2}$, $V_{rr'} = \prod_l \sigma_l^z$ is the (non-gauge-invariant) Wilson line operator connecting the points r and r' , and $|\text{GS}\rangle$ is the ground state. $V(-T/2)$ acts on the ground state by creating two ‘‘defects’’ at r and r' where $G_r = -1$, and (in the limit of large- T) projects to the lowest-energy state with such defects.

The order parameter is given by

$$\tilde{R}_2(L, T) = \frac{\langle \psi_z | \psi_{z+1} \rangle}{\sqrt{\langle \psi_z | \psi_z \rangle \langle \psi_{z+1} | \psi_{z+1} \rangle}} \quad (15)$$

which will exhibit the same asymptotic behavior as $L, T \rightarrow \infty$ as $F(L)$. In this picture, we see that \tilde{R}_2 is probing the orthogonality of the trial spinon states on plane z and $z + 1$. In the deconfined floating phase, $|\psi_z\rangle$ and $|\psi_{z+1}\rangle$ will be orthogonal since spinons cannot tunnel between planes. However, in a fully (3+1)D deconfined phase a spinon may move between the two planes and so $\langle \psi_z | \psi_{z+1} \rangle \neq 0$. While \tilde{R} is useful conceptually, in condensed matter systems where the gauge symmetry is emergent, the equal-time formulation R_2 [Eq. (12)] should be used [48].

Finally, all of this discussion can be extended beyond \mathbb{Z}_2 . One simply replaces V with the appropriate Wilson line operator and τ by the appropriate charged matter operator for gauge invariance [48].

IV. GAPLESS FLOATING TOPOLOGICAL PHASES

In this section, we analyze the stability of gapless floating topological phases. We first consider the floating phase of gapless Dirac matter coupled to a gapped \mathbb{Z}_2 gauge field. We

then go on to consider the floating phase of Dirac fermions coupled to a $U(1)$ gauge field, in which both the matter and gauge sectors are gapless. We show that these gapless floating phases are stable to interlayer couplings in the renormalization group (RG) sense. That is, the interlayer couplings are irrelevant perturbations: at long distances and low energies, the system flows back to the decoupled limit.

A. Gapless matter, gapped gauge

Let us first consider the case with gapless Dirac matter coupled to a gapped gauge field. At low energies, a single (2+1)D layer is described by Dirac fermions hopping in a background static gauge field. This describes, for example, the gapless phase of Kitaev’s honeycomb model [50], in which case the minimum energy configuration for the gauge field is equivalent to the trivial (flux-free) configuration.

The low-energy continuum Hamiltonian for a single layer l is

$$H_l = \int d^2r \psi_l^\dagger(\vec{r}) [-iv\vec{\sigma} \cdot \vec{\partial}] \psi_l(\vec{r}), \quad (16)$$

where $\psi_l^\dagger(\vec{r})$ [$\psi_l(\vec{r})$] is a two-component spinor which creates (annihilates) a complex fermion at position $\vec{r} = (x, y)$ on layer l , v is the Fermi velocity, $\vec{\partial} = (\partial_x, \partial_y)$, and $\vec{\sigma} = (\sigma^x, \sigma^y)$ is a vector of Pauli matrices acting on the spinor indices.²

The corresponding Euclidean action for the single layer is

$$S_l = \int d^2r d\tau \bar{\psi}_l(\vec{r}, \tau) \gamma^\mu \partial_\mu \psi_l(\vec{r}, \tau), \quad (17)$$

where summation over $\mu = \tau, x, y$ is implied, $\gamma^\mu = (\sigma^z, \sigma^y, -\sigma^x)$, $\bar{\psi} = \psi^\dagger \sigma^z$, and we have rescaled coordinates to set $v = 1$.

The gauge fields are fully gapped out, and hence do not appear in the low-energy description. Nevertheless, they are important as they restrict the terms which are allowed to appear to only those which are gauge invariant. When we have multiple layers, gauge invariance within each layer implies that any interlayer term must consist of operators which are individually gauge invariant on each layer. Crucially, this forbids quadratic interlayer hopping terms, which are not gauge invariant within a single layer. The simplest gauge-invariant terms coupling two layers are four-body interaction terms such as $(\bar{\psi}_l \psi_l)(\bar{\psi}_{l'} \psi_{l'})$ between two layers l, l' . However, all such quartic terms are irrelevant at the (2+1)D Dirac fermion Gaussian fixed point with $S = \sum_l S_l$. This can be seen by simple power counting: the field ψ has length dimensions $[\psi] = L^{-1}$, and so the quartic term has dimensions $[(\bar{\psi} \psi)^2] = L^{-4}$. Under an RG transformation in which we rescale time and the two continuous spatial dimensions, the quartic term therefore flows to zero. If the system were to flow to a fully (3+1)D phase, we would instead expect that the interlayer couplings would increase under this RG flow. In this case, however, the system flows back to the decoupled layers limit at large distances. This system is therefore an example of a stable gapless floating phase.

²This low-energy form is obtained from Kitaev’s honeycomb model by first combining the two Majorana cones into a single Dirac cone.

Finally, we note that there are single-layer terms which are relevant. These include quadratic terms of the form $\bar{\psi}\gamma^\mu\psi$, which may open up a gap or create a Fermi surface. In assessing the stability of these phases, we have implicitly assumed that these terms are forbidden by symmetries of the microscopic Hamiltonian, and that the microscopic interlayer couplings also respect such symmetries.

B. Gapless matter, gapless gauge

Let us now consider the case of Dirac fermions coupled to a gapless U(1) gauge field. The pure U(1) gauge theory is confining in $(2+1)D$, but a stable deconfined phase can exist when coupled to a large number of Dirac fermions [51]. In this situation, the low-energy continuum description of each $(2+1)D$ layer l is simply large- N quantum electrodynamics: N Dirac fermion flavors $\psi_{i,l}$ ($i = 1, \dots, N$) coupled to an emergent gauge field $a_{\mu,l}$. The Euclidean Lagrangian is

$$\mathcal{L}_l^{\text{QED}_3} = \sum_{i=1}^N \bar{\psi}_{i,l} \gamma^\mu (\partial_\mu + i a_{\mu,l}) \psi_{i,l} + \frac{1}{4e^2} f_{\mu\nu,l} f_l^{\mu\nu}, \quad (18)$$

where $f_{\mu\nu,l} = \partial_\mu a_{\nu,l} - \partial_\nu a_{\mu,l}$ is the field strength. The gauge transformation sends $\psi_l \rightarrow e^{i\alpha_l} \psi_l$ and $a_{\mu,l} \rightarrow a_{\mu,l} - \partial_\mu \alpha_l$ for an arbitrary space-time function $\alpha_l(\vec{r}, \tau)$ on each layer l .

The Maxwell term, although included, is irrelevant at large N . This is exemplified by the fact that with a clever choice of nonlocal gauge-fixing term, the gauge photon propagator can be written in such a way that the $e^2 \rightarrow \infty$ limit can be taken at the beginning of a calculation [52]. Indeed, as written, we have $[a] = L^{-1}$ and so $[f_{\mu\nu} f^{\mu\nu}] = L^{-4}$ is irrelevant, while the coupling $[\bar{\psi} a \psi] = L^{-3}$ is marginal.

As in the previous case, the simplest interlayer coupling terms that are gauge invariant are either quartic in the fermion operators or pure gauge (of the form $f_{\mu\nu,l} f_l^{\mu\nu}$), both of which are strictly irrelevant perturbations at large N . However, as before, we have implicitly assumed that, for reasons of symmetry, relevant single-layer terms are not present. These now include, for example, a Chern-Simons term.

Like in the gapped case, rather than the low-energy model with a separate gauge symmetry on each individual layer, a more natural starting point is a single anisotropic emergent $(3+1)D$ gauge field with interlayer couplings much weaker than intralayer. In this case, one has a gauge fields $a_{z,l}$, and gauge-invariant interlayer hopping terms of the form $\bar{\psi}_{l+1} e^{i a_{z,l}} \psi_l$ are allowed. In this limit, $a_{z,l}$ are strongly fluctuating (and therefore gapped), so they can be integrated out resulting in a local effective action with a separate gauge symmetry on each individual layer. We go through this explicitly in Appendix for an anisotropic U(1) lattice gauge theory with fermions. Such a system is partially confined: fractionalized quasiparticles are confined along z , but deconfined within each layer.

Such ‘‘layered’’ phases of the U(1) gauge theory have been studied previously [53,54] and found to be stable in higher than $(3+1)D$. These are examples of stable higher-dimensional gapless floating topological phases.

C. Gapless matter with a Fermi surface

Finally, we turn to the case where the emergent (gauged) fermions form a Fermi surface with a finite density of states [55,56]. This situation may arise in a layered, gapless spin liquid. We first consider the simpler case of a \mathbb{Z}_2 gauge field, where the gauge degrees of freedom are gapped, and then comment on the more complex case of a U(1) gapless gauge field.

1. Fermi surface with \mathbb{Z}_2 gauge field

In the case of gapless fermions coupled to a \mathbb{Z}_2 gauge field, the only low-energy degrees of freedom are the fermions. At a single layer l , the effective Hamiltonian is

$$H_{\mathbb{Z}_2 \text{ FS}} = \sum_l \int \frac{d^2k}{(2\pi)^2} \varepsilon(\vec{k}) \psi_l^\dagger(\vec{k}) \psi_l(\vec{k}) + H_{\text{pert}}, \quad (19)$$

where $\varepsilon(\vec{k})$ is the dispersion of the fermions, that vanishes at the Fermi surface specified by $\varepsilon(\vec{k}) = 0$, and H_{pert} includes various possible perturbations, to be discussed in the following. We have assumed for simplicity that the gauged fermions, created by the operator $\psi_l^\dagger(\vec{k})$, do not carry any quantum number other than the layer index l and momentum \vec{k} (e.g., spin is not conserved).

Before turning to the effects of interlayer interactions, we first discuss stability of the Fermi surface in the limit where the layers are decoupled. It is not immediately obvious that the Fermi surface is stable, even in this limit, since the number of fermions is only conserved mod (2) . Thus, pairing terms of the form $\Delta(\vec{k}) \psi_l^\dagger(\vec{k}) \psi_l^\dagger(-\vec{k}) + \text{H.c.}$ are generally allowed in H_{pert} , and one may conclude that (except under fine-tuned circumstances [57,58]) the Fermi surface is always gapped, with the exception of a discrete set of nodal points where both $\varepsilon(\vec{k}) = 0$ and $\Delta(\vec{k}) = 0$. Nevertheless, it turns out that stable Fermi surfaces are, in fact, possible in certain circumstances [59–62]: (i) If both time-reversal and inversion symmetries are broken, either spontaneously or explicitly, such that $\varepsilon(\vec{k}) \neq \varepsilon(-\vec{k})$, then the Fermi surface remains even for a nonzero generic $\Delta(\vec{k})$, as long as the maximum of $|\Delta(\vec{k})|$ is below a certain critical value. (ii) Even in the presence of time-reversal symmetry, there are situations where the Fermi surface is stable. This happens if the action of time reversal \mathcal{T} on the gauged fermions is such that $\mathcal{T} \psi_l(\vec{k}) \mathcal{T}^{-1} = \psi_l(-\vec{k} + \vec{Q})$ with a certain nonzero wave vector \vec{Q} . In this case, $\varepsilon(\vec{k}) \neq \varepsilon(-\vec{k})$ even in the presence of time reversal, and the Fermi surface is again stable.³

Note that the excitations near the Fermi surface are actually ‘‘Bogoliubov-type’’ quasiparticles, consisting of superpositions of ψ_l and ψ_l^\dagger . Nevertheless, we can always diagonalize the quadratic part of the Hamiltonian including the pairing

³In this case, one can always choose a gauge where the action of time reversal is $\mathcal{T} \psi(\vec{k}) \mathcal{T}^{-1} = \psi(-\vec{k})$ by shifting the momentum of the fermions. However, in this case, the ‘‘pairing term’’ becomes modulated in space, as in a pair density wave (PDW) state, since it includes pairs of electrons with a nonzero momentum. Pair density wave states are known to support stable Fermi surfaces [63,64].

terms, bringing it to the form (19) with only quartic and higher-order terms in H_{pert} .

For a generic dispersion $\varepsilon(\vec{k})$, short-range quartic terms, both intralayer and interlayer, are marginal by power counting, as in the RG analysis of a Fermi liquid [65]. Higher-order terms are irrelevant. Taking into account the fact that the number of fermions must be conserved mod (2) in each layer, the most general quartic interaction is of the form

$$H_{\text{pert}} = \sum_{l,l'} \sum_{\{\eta_i=\pm 1\}} \int_{\{\vec{k}_i\}} \delta\left(\sum_{i=1}^4 \eta_i \vec{k}_i\right) \times W_{l,l'}(\{\eta_i\}, \{\vec{k}_i\}) \psi_l^{\eta_1}(\vec{k}_1) \psi_l^{\eta_2}(\vec{k}_2) \psi_l^{\eta_3}(\vec{k}_3) \psi_l^{\eta_4}(\vec{k}_4), \quad (20)$$

where we have introduced the notation $\psi_l^{\eta=+1} \equiv \psi_l$, $\psi_l^{\eta=-1} \equiv \psi_l^\dagger$. From the Hermiticity of H_{int} , the interaction function must satisfy $W_{l,l'}^*(\{\eta_1 \dots \eta_4\}, \{\vec{k}_1 \dots \vec{k}_4\}) = W_{l,l'}(\{-\eta_4 \dots -\eta_1\}, \{\vec{k}_4 \dots \vec{k}_1\})$.

The stability analysis of the Fermi surface including ‘‘anomalous’’ (particle-number-nonconserving) terms of the form (20) can be done along similar lines of the RG analysis for a Fermi liquid [65]. We shall not go through the details of this analysis here, and only point out the main results. Most importantly, in the absence of a Kramers-type degeneracy $\varepsilon(\vec{k}) \neq \varepsilon(-\vec{k})$, the interaction function (20) is exactly marginal. The imaginary part of the single-fermion self-energy on the Fermi surface, to order $|W|^2$, is $\Sigma''(\omega, T) \propto \max(\omega^2, T^2) \times \log[\frac{\varepsilon_F}{\max(|\omega|, T)}]$ (where ε_F is the Fermi energy), similar to that of an ordinary two-dimensional Fermi liquid. These results indicate that the resulting stable phase is a ‘‘floating Fermi liquid’’ with long-lived low-energy fermionic quasiparticles that can propagate coherently within each layer, but not between layers.

Note that the interaction (20) includes terms that do not conserve the quasiparticle number $\psi_l^\dagger(\vec{k})\psi_l(\vec{k})$ at a particular point on the Fermi surface. Such terms are not included in the standard Fermi-liquid effective Hamiltonian, but nevertheless do not change its low-energy properties. They are analogous to the umklapp terms in a two-dimensional Fermi liquid, which are marginal under RG [66] but do not destabilize the Fermi liquid.

Since the layers are coupled by forward-scattering density-density interactions [included in Eq. (20)], the collective mode spectrum of the system may include ‘‘zero sound’’ modes that propagate in all three directions, depending on the nature of the interlayer interactions.

2. Fermi surface with U(1) gauge field

Finally, we comment on the case of a layered system with a Fermi surface coupled to an emergent U(1) gauge field in each layer. The Lagrangian density of an individual layer is given by

$$\mathcal{L}_l^{\text{FS}} = \sum_{\alpha=1}^N \bar{\psi}_{i,l} [\partial_\tau + ia_{0,l} + \varepsilon(-i\nabla + \vec{a}_l)] \psi_{i,l} + \frac{1}{4e^2} f_{\mu\nu,l} f_l^{\mu\nu}. \quad (21)$$

As in the \mathbb{Z}_2 case, we must first address the stability of the Fermi surface in a single layer, before considering the effects

of interlayer coupling. Unfortunately, even this is at present an unsolved problem. The traditional expansion in $1/N$ turns out to be problematic, due to a proliferation of divergences in terms that are naively of high order in $1/N$, making the expansion unreliable at asymptotically low energies [67–69].

A possible way to cure the problem [70,71] is to generalize the bare ‘‘photon dispersion,’’ coming from the last (Maxwell) term in Eq. (21), to $\omega^2 \propto |\vec{q}|^{1+\epsilon}$. The physical case is $\epsilon = 1$. The problem can be solved in a double expansion in both $1/N$ and ϵ . To lowest order, one recovers the self-consistent one-loop approximation for the fermionic and gauge field self-energies [71,72]. Note that the price of this approach is that for $\epsilon < 1$ the action is nonlocal in space, which entails potential subtleties in an RG analysis (which generally assumes a local form for the action, a property preserved by the RG transformation).

Proceeding nevertheless along these lines, Metlitski *et al.* [73] showed that the single-layer action (21) is stable in the presence of arbitrary quartic interactions, as long as they are sufficiently weak. In particular, the Fermi surface does not have a BCS instability toward Cooper pairing [74].

With the above caveats in mind, we may assess the effects of interlayer interactions within the fixed point described by the $(\epsilon, 1/N)$ expansion. The lowest-order gauge-invariant interlayer terms are of the form⁴

$$\mathcal{L}_{\text{inter}} = \sum_{l,l'} [u_{l,l'} n_l n_{l'} + g_{l,l'} f_{\mu\nu,l} f_l^{\mu\nu}]. \quad (22)$$

Here, $n_l = \sum_i \psi_{i,l}^\dagger \psi_{i,l}$ is the density of fermions in layer l . The first term is irrelevant at the decoupled layer limit, by the same argument that makes the single-layer fixed point stable with respect to Cooper pairing. The second term is marginal by scaling, and renormalizes the dispersion of the (overdamped) photon mode, which now becomes dependent on the component of the wave vector perpendicular to the planes. Hence, at least within the $(\epsilon, 1/N)$ expansion, a floating phase of Fermi liquids coupled to U(1) gauge fields remains stable in the presence of interplane coupling. Whether this conclusion extends beyond this limit remains to be determined.

V. EXPERIMENTAL DIAGNOSTIC: DIVERGING CONDUCTIVITY ANISOTROPY

Having discussed the stability of floating phases, we now turn to possible experimental diagnostics of such phases. The correlation function diagnostic in Sec. III is useful conceptually, but is unlikely to be measurable experimentally. In this section, we discuss an experimentally accessible signature of a floating topological phase in the form of the electrical conductivity anisotropy at low temperatures $T \rightarrow 0$.

In a floating topological phase, it is possible that the transport coefficients become *parametrically* anisotropic in the limit $T \rightarrow 0$. In particular, the ratio of the in-plane and out-of-plane conductivities diverges as $T \rightarrow 0$. As far as we know,

⁴For simplicity, we omit here some terms that are allowed by symmetry, e.g., a term of the form $(\nabla \times \vec{a}_l)(\nabla \times \vec{a}_{l'})$ [which is linearly independent from the term that appears at the end of Eq. (22)]. These terms obey the same scaling as the term we have retained.

this is impossible in ordinary, three-dimensional (nonfloating) phases, where we expect the conductivities in all directions to scale in the same way as a function of T , even when the microscopic parameters are strongly anisotropic.

In gapless floating quantum spin liquids (QSLs), the anisotropy of the thermal conductivity has been studied by Werman *et al.* [18]. However, in gapped spin liquids, phonons always dominate the thermal transport at low temperatures, and the signatures of fractionalization of the spin excitations are masked. Even in gapless QSLs, where the magnetic excitations dominate at low temperatures, one may need to go to exceedingly low T to make the phonon contribution negligible. Here, we consider the *electrical* conductivity, which may be mediated by gapped “holon” (or chargon) excitations (in both gapped and gapless QSLs). Since there is no phonon contribution to the electrical conductivity, the electronic anisotropy may be more easily measurable.

In weakly disordered systems, we find conditions under which the anisotropy in the conductivity indeed diverges at low temperatures, in both gapped and gapless layered QSLs. The functional form of the divergence of the anisotropy depends on the type of QSL and the properties of its lowest-energy excitations.

Interestingly, if the charge carriers are localized and gapless, and hence the conductivity at low temperature occurs through variable-range hopping, it turns out that the resistivity anisotropy *saturates*. Thus, a rapidly growing resistivity anisotropy can serve as a signature of floating topological phases, although in the presence of disorder, localization may ultimately preempt a true divergence.

We illustrate these principles by computing the resistivity anisotropy in two examples: a gapped QSL and a gapless \mathbb{Z}_2 Dirac QSL, both with weak disorder. We then discuss possible nonperturbative effects of quenched disorder that may appear at low temperature, such as a modification of the low-energy spinon spectrum and Anderson localization.

A. Gapped QSL

Let us consider a layered, gapped \mathbb{Z}_2 QSL. The mechanism for conductivity at low temperature depends on the character of the lowest-energy charge excitations. Crucially, the transport of charge within the planes can be carried by fractionalized excitations (that carry a nontrivial gauge charge or flux), whereas interlayer transport must be carried by gauge-neutral, charged excitations (such as electrons). This is the source of the parametric anisotropy in the limit $T \rightarrow 0$.

As a concrete example, let us assume that the lowest-energy charged excitation in each layer is a holon that carries charge $+e$ and is “electrically charged” under the gauge field. The self-statistics of the lowest-energy holon excitation may either be fermionic or bosonic, depending on energetics [75]. Let us denote the holon creation operator by $h_l^\dagger(\vec{r})$. Then, we can define a “spinon” excitation $\psi_{l,\sigma}(\vec{r}) \sim h_l^\dagger(\vec{r})c_{l,\sigma}^\dagger(r)$, where $c_{l,\sigma}^\dagger(r)$ creates an electron in layer l with spin $\sigma = \uparrow, \downarrow$. The spinon carries spin $\frac{1}{2}$, zero electric charge, and a nontrivial gauge charge. Note that if the holon has fermionic self-statistics, then the spinon must have bosonic self-statistics and vice versa, such that the electron is a fermion.

The low-energy effective Hamiltonian is then given by

$$H = \sum_{\vec{k},l} \varepsilon_{\vec{k}}^h h_{\vec{k},l}^\dagger h_{\vec{k},l} + \sum_{\vec{k},\sigma,l} \varepsilon_{\vec{k}}^\psi \psi_{\vec{k},\sigma,l}^\dagger \psi_{\vec{k},\sigma,l} + H_{\text{dis}} + H_{\text{inter}}, \quad (23)$$

where \vec{k} is the in-plane momentum, and at low energies we may expand $\varepsilon_{\vec{k}}^h = \Delta_h + \frac{k^2}{2m_h} + \dots$, with Δ_h and m_h the holon gap and effective mass, respectively. Similarly, $\varepsilon_{\vec{k}}^\psi = \Delta_\psi + \frac{k^2}{2m_\psi} + \dots$. H_{dis} and H_{inter} are disorder and interplane coupling terms, to be discussed below. Crucially, we assume that the holon and spinon do not form an electronlike bound state, i.e., the gap for creating an electron excitation Δ_c is larger than $\Delta_\psi + \Delta_h$. Under these conditions, both in-plane and out-of-plane conductivity are carried by spinons and holons, and electronic excitations can be ignored at low temperatures. We will discuss situations where this condition is violated below.

Note that spinon number is only conserved mod 2 so terms of the form $\psi_{l,\uparrow}^\dagger(\vec{r})\psi_{l,\downarrow}^\dagger(\vec{r})$ are allowed in the Hamiltonian. However, such terms can be eliminated by a Bogoliubov transformation. Quartic terms that describe interactions between spinons or between spinons and holons include, in general, terms that do not conserve the spinon number. Terms of the form $h_l^\dagger(\vec{r})c_{l,\sigma}^\dagger(\vec{r})\psi_{l,\sigma}(\vec{r})$ are also allowed. However, at low temperatures, excitations of all types are very dilute, and intralayer interactions do not play an important role, as long as the spinon and holon interact repulsively and do not form a bound state.

To describe interlayer transport, we include an interlayer coupling of the form

$$H_{\text{inter}} = g_\perp \int d^2r \sum_{l,\sigma} h_{l+1}^\dagger(\vec{r})h_{l+1}(\vec{r})\psi_{l,\sigma}^\dagger(\vec{r})\psi_{l+1,\sigma}(\vec{r}) + \text{H.c.} \quad (24)$$

In order to render the in-plane conductivity finite, we need to take into account terms that break translational symmetry in the plane. To this end, we include also a weak disorder potential H_{dis} , that may couple to both the spinons and holons. The explicit form of H_{int} is not important for our present purpose. We will assume that the disorder is sufficiently weak that its effects can be considered perturbatively; nonperturbative effects leading to localization will be discussed in Sec. V C 3.

The frequency-dependent conductivity perpendicular to the layers $\sigma_\perp(\Omega)$ can be computed perturbatively in g_\perp . Assuming that the holons obey Bose statistics, whereas the spinons are fermions, σ_\perp is given by

$$\begin{aligned} \sigma_\perp(\Omega) = & \frac{4\pi e^2 |g_\perp|^2}{\hbar \Omega d} \int_{\vec{k},\vec{k}',\vec{q}} [\delta(\Omega - \varepsilon_{\vec{k}}^h + \varepsilon_{\vec{k}+\vec{q}}^h - \varepsilon_{\vec{k}'}^\psi + \varepsilon_{\vec{k}'-\vec{q}}^\psi) \\ & - (\Omega \rightarrow -\Omega)] n_B(\varepsilon_{\vec{k}}^h) n_B(-\varepsilon_{\vec{k}+\vec{q}}^h) n_F(\varepsilon_{\vec{k}'}^\psi) \\ & \times n_F(-\varepsilon_{\vec{k}'-\vec{q}}^\psi). \end{aligned} \quad (25)$$

Here, $n_{F,B}(\varepsilon)$ are the Fermi and Bose functions, respectively, and $\int_{\vec{k}} \equiv \int \frac{d^2k}{(2\pi)^2}$.

At low temperature, we may replace $n_{F,B} \approx e^{-\varepsilon/T}$. Taking the dc ($\Omega \rightarrow 0$) limit, we obtain in a gapped QSL

$$\begin{aligned} \sigma_{\perp} &= \frac{4\pi e^2 |g_{\perp}|^2}{Td} \int_{\vec{k}, \vec{k}', \vec{q}} \delta\left(\varepsilon_{\vec{k}}^h - \varepsilon_{\vec{k}+\vec{q}}^h + \varepsilon_{\vec{k}}^{\psi} - \varepsilon_{\vec{k}'-\vec{q}}^{\psi}\right) e^{-\frac{\varepsilon_{\vec{k}}^h + \varepsilon_{\vec{k}'}^{\psi}}{T}} \\ &\propto e^{-\frac{\Delta_h + \Delta_{\psi}}{T}}. \end{aligned}$$

In contrast, the in-plane conductivity is proportional to the density of holons, and hence $\sigma_{\parallel} \propto e^{-\frac{\Delta_h}{T}}$ (with a prefactor proportional to the elastic mean-free time of a holon in the plane). The anisotropy ratio

$$\left(\frac{\sigma_{\parallel}}{\sigma_{\perp}}\right)_{\text{gapped QSL}} \propto \exp\left[\frac{\Delta_{\psi}}{T}\right] \quad (26)$$

diverges at low T .

Importantly, as noted above, we assumed that there is no attraction between spinons and holons in a layer. Hence, the energy of an electron or a hole excitation is larger than $\Delta_h + \Delta_{\psi}$. If the minimal excitation energy of an electron Δ_c is smaller than $\Delta_h + \Delta_{\psi}$, then the interplane conductivity is mostly carried by electronic excitations, so $\sigma_{\perp} \propto e^{-\Delta_c/T}$. Still, so long as $\Delta_c > \Delta_h$, the in-plane current is carried by holons and the anisotropy still diverges as $\sigma_{\parallel}/\sigma_{\perp} \propto \exp[(\Delta_c - \Delta_h)/T]$. However, if $\Delta_c < \Delta_h$, then both the in-plane and out-of-plane currents are carried by electrons, and the anisotropy ratio does not diverge in the $T \rightarrow 0$ limit. Thus, a divergent anisotropy implies that the system is in a floating phase, but the converse is not necessarily true: a gapped floating phase may or may not have a divergent anisotropy.

B. Gapless Dirac QSL

Next, we consider the case of a gapless Dirac QSL coupled to a \mathbb{Z}_2 gauge field in each layer, considered in Sec. IV A. The spinons are fermionic, with an in-plane action given by Eq. (16), supplemented (assuming that the system is spin rotationally invariant) by a summation over the spin index. The system is electrically insulating, and the bosonic holons are assumed to be gapped with a minimal gap Δ_h . The interlayer coupling is taken to be of the form (24).

As in the gapped case, if the gap to create electron or hole excitations is smaller than Δ_h , then the conductivity anisotropy ratio saturates at low temperature. In contrast, if the holon is the lowest-energy charged excitation, then the out-of-plane conductivity is of the form of Eq. (25), with the spinon dispersion given by $\varepsilon_{\vec{k}}^{\psi} = v|\vec{k}|$. We explore this case henceforth.

At low temperature, we approximate $n_B(\varepsilon_{\vec{k}}^h) \approx e^{-\varepsilon_{\vec{k}}^h/T}$ and $n_B(-\varepsilon_{\vec{k}+\vec{q}}^h) \approx -1$. The dc limit $\Omega \rightarrow 0$ is then taken. We may evaluate the integral in (25) as follows. Due to the Fermi and Bose functions and the energy conservation condition, $|\vec{k}'|$ and $|\vec{q}|$ are both of order $\frac{T}{v}$, whereas $|\vec{k}| \sim \sqrt{2m_h T}$. Therefore, $|\varepsilon_{\vec{k}}^h - \varepsilon_{\vec{k}+\vec{q}}^h| \approx \frac{\vec{k}\cdot\vec{q}}{m_h} \sim \sqrt{\frac{2T}{m_h}} \frac{T}{v} \ll T$, and we may neglect the $\varepsilon_{\vec{k}}^h - \varepsilon_{\vec{k}+\vec{q}}^h$ term relative to $\varepsilon_{\vec{k}'}^{\psi} - \varepsilon_{\vec{k}'-\vec{q}}^{\psi}$ in the δ function. The integral over \vec{k} can then be done independently of the integrals

over \vec{k}' and \vec{q} . This results in

$$\sigma_{\perp} \propto T^3 e^{-\frac{\Delta_h}{T}}. \quad (27)$$

The in-plane conductivity can be estimated from the Einstein relation: $\sigma_{\parallel} = \kappa D$, where κ is the compressibility and D is the diffusion constant. The compressibility is $\kappa \sim m_h e^{-\frac{\Delta_h}{T}}$. The diffusion constant is $D = \frac{1}{2}\langle v^2 \rangle \tau$, where $\langle v^2 \rangle \sim 2T/m_h$ is the average velocity of the holons, and τ is the mean-free time of the holon due to impurity scattering (assumed to be temperature independent). We obtain

$$\sigma_{\parallel} \sim T e^{-\frac{\Delta_h}{T}}, \quad (28)$$

and hence

$$\left(\frac{\sigma_{\parallel}}{\sigma_{\perp}}\right)_{\mathbb{Z}_2 \text{ Dirac QSL}} \sim \frac{1}{T^2}. \quad (29)$$

The conductivity anisotropy ratio diverges algebraically as $T \rightarrow 0$, unlike the gapped case where it diverges exponentially. As in the gapped case, if $\Delta_c < \Delta_h$ then the anisotropy ratio does not diverge since both the in-plane and out-of-plane currents are carried by electrons.

C. Nonperturbative disorder effects

So far, we have treated the disorder potential in the layers as weak; basically, its only effect was to provide the holons with a finite in-plane transport lifetime. Here, we comment on possible nonperturbative effects of disorder, that couple either to the holons or the spinons.

1. Spinon disorder

In the case of a Dirac QSL, disorder may modify the nature of Dirac fermions qualitatively, depending on the symmetry of the problem. In certain cases, such as in the Kitaev honeycomb model with time-reversal symmetry, the disorder couples to the spinons as a random vector potential [76]. Then, the DOS scales as a nonuniversal power law with energy, with a power that depends continuously on the disorder strength [77]. The latter behavior may result in a nonuniversal power-law dependence of the conductivity anisotropy ratio on temperature. Such behavior has been predicted for the thermal conductivity anisotropy in a layered Dirac QSL [18]. If the disorder couples to the density of the Dirac spinons, then the density of states (DOS) approaches a nonzero constant at low energy [78]. We leave a detailed prediction of the conductivity anisotropy in this case, as well as a discussion of the anisotropy in the case of a QSL with a disordered spinon Fermi surface, to future studies.

2. Screw dislocations

Interestingly, in a floating topological phase, the transport properties are qualitatively affected by the presence of lattice screw dislocations along the axis perpendicular to the layers. This is because such dislocations allow fractionalized excitations, e.g., spinons and holons in a layered QSL, to move coherently between layers. Thus, in the presence of a finite

density of screw dislocations, the resistivity anisotropy saturates at sufficiently low temperature, with a saturation value that depends on the density of screw dislocations.

A screw dislocation in a layered system is a version of a “twist defect” discussed in multilayered systems [79]. Depending on the type of topological order in each layer, screw dislocations may also carry propagating gapless modes. Similar phenomena have been discussed in weak topological insulators [80]. If present, such gapless modes may further affect the resistivity anisotropy.

3. Localization and variable range hopping

Ignoring interactions, the motion of a low-energy holon is two dimensional, and hence quenched disorder is generally expected to localize it. Moreover, for a generic disorder distribution, the holon excitations are then gapless (even if there is a finite gap in the absence of disorder) since at any given energy within the original gap there is a nonzero probability to find a localized state. At finite temperature, the dc conductivity is then dominated by variable range hopping of either holons or electrons. Moreover, as we shall now argue, at low enough T , the conductivity, even for currents in the in-plane direction, is always dominated by interplane hopping processes.

To see this, consider hopping of charge between two localized states separated by a displacement \vec{R} . The hopping rate, generically, depends exponentially on \vec{R} as $\nu \propto \exp[-S(\vec{R})]$. If the relevant charged excitations are fractionalized, then the behavior of $S(\vec{R})$ can be different depending on whether the two localized states are in the same plane or in different planes. For two localized states in the same plane, $S = S_{\text{intra}} = |\vec{R}|/\xi_{\text{intra}}$ where ξ_{intra} is the in-plane localization length. By contrast for two localized states in different planes, $S = S_{\text{inter}} = \sqrt{(R_{\parallel}/\xi_{\parallel})^2 + (R_{\perp}/\xi_{\perp})^2}$ where the subscripts \parallel and \perp refer to the in-plane and out-of-plane directions, and ξ_{\parallel} is not necessarily equal to ξ_{intra} . The important point is that for a given concentration of active states c , the typical distance between neighboring states in a given plane grows in proportion to $c^{-1/2}$ while if interplane processes are considered it grows in proportion to $c^{-1/3}$.

Thus, since the concentration of thermally accessible states vanishes as $T \rightarrow 0$, even in the limit $\xi_{\text{intra}} \gg \xi_{\parallel} > \xi_{\perp}$, at low enough T interplane hopping is always preferred. In this limit, the system can be considered as an anisotropic continuum, with the result that the anisotropy has no effect on the T dependence of the resistivity (it has the usual form for variable range hopping) and the resistivity anisotropy $\rho_{\perp}/\rho_{\parallel} = (\xi_{\parallel}/\xi_{\perp})^2$. Thus, while there could arise an intermediate T regime in which the anisotropy in the hopping conductivity grows with decreasing T (reflecting an intermediate regime in which hopping of fractionalized particles within a given plane dominates the transport in the \parallel direction), at low enough T the anisotropy will always saturate.

VI. CONCLUDING REMARKS

One of the more remarkable recent advances in our understanding of phases of matter is the discovery of “absolute stability,” the stability of a phase to absolutely any perturbation. The primary class of such absolutely stable phases are

topological phases which exhibit topological order characterized by emergent gauge fields and fractionalization. Such phases are well known to be stable to all weak perturbations to their Hamiltonians [81]. Another class of absolutely stable phases which was discovered even more recently is that of discrete-time crystals [82] which are nonequilibrium phases of Floquet systems. Here the stability is with respect to arbitrary weak perturbations of the drive that preserve the period.

In this paper we have examined the stability of topological phases to weak perturbations in dimensionality, which take place when weakly coupled stacks of d -dimensional systems are in the $(d + 1)$ st direction. Remarkably, as first observed by Senthil and Fisher [17] and studied at length in this paper, topological phases are stable to such perturbations too, in the sense that the resulting “floating” phases are connected perturbatively to the strictly decoupled stacks of lower dimensional systems and share their universal properties. This stability is extraordinarily useful from the viewpoint of realizing topological phases in materials. While in cases of broken symmetry at long wavelength the higher, physical, dimension always wins and thus any interesting lower-dimensional phenomena are always ultimately obscured, for topological phases such as spin liquids the opposite is true. Given that spin liquids are more likely to be found in two dimensions on account of stronger quantum fluctuations, it is highly encouraging that their weakly three-dimensional continuations will, in a sharp sense, continue to exhibit the universal properties of the planar phases regardless of the details of the couplings in the third direction just as long as they are weak. In addition to this general observation, we have presented a fairly general tool for identifying such floating spin liquids, which is to study the low-temperature anisotropy in their charge transport. We trust that this combination of the general and the particular will encourage and inform the ongoing search for spin liquids.

Indeed, an analogous strategy for probing the presence of a candidate topological phase has recently been applied in the cuprate high-temperature superconductors. There, the observation [83] of an anomalously large nonelectronic component of the in-plane thermal Hall conductivity κ_{xy} led to speculative interpretations [84–86] in terms of a heat current carried by spinons associated with the presence, or near proximity of a chiral topological phase in the individual Cu-O planes. However, a followup study [87] found an effect of comparable magnitude in κ_{xz} , i.e., when the thermal current is in the direction perpendicular to the plane. This observation is generally taken to rule out such an exotic explanation, and suggests instead that the thermal Hall response is associated with a still notable, and not fully understood aspect of the phonon dynamics [88,89].

Before concluding, we briefly list some items worthy of further investigation:

(i) *Frequency dependent conductivity.* We have focused on the anisotropy in the T dependence of the conductivity. However, it may well be that in an insulator it is much more convenient to look at the variation of the electrical response with frequency. The optical conductivity of a gapless spin liquid is generically finite at frequencies below the charge gap [90]. The physics discussed in this paper will again imply that this response becomes infinitely anisotropic as $\omega \rightarrow 0$. The

precise form of the anisotropy will require computation in various cases of interest.

(ii) *Chiral phases.* Two-dimensional spin liquids also come in a chiral variant where the long-wavelength action contains a Chern-Simons term; indeed, there is strong numerical evidence that they can be found in relatively simple Hamiltonians. These phases, which break time-reversal symmetry spontaneously, can be coupled three dimensionally in more interesting ways in which the topological order floats, while the discrete broken symmetry orders with various periods. Now, we can end up with transport which proceeds either via the bulk or via a potentially gapless surface of the kind which has been studied in the context of layered quantum Hall systems. We note that the bulk response will have both longitudinal and transverse components of which the in-plane components (σ_{xy} and κ_{xy}) will be nontrivial and especially interesting. We intend to return to this in future work. The more general observation here is that such considerations will apply to other stacked phases where the lower-dimensional phase exhibits both topological order and broken symmetries, such as quantum Hall ferromagnets. Indeed, even stacked topological phases with unbroken symmetries, symmetry-enriched topological phases, can potentially give rise to distinct floating phases wherein the same floating topological order is joined to different realizations of the now three-dimensional symmetry group.

(iii) *Metallic phases.* We have focused on insulating phases but the basic intuition also carries over to metallic phases that also exhibit fractionalization and emergent gauge fields, e.g., the “orthogonal metal” [91,92]. A related phenomenon was explored in Ref. [93] where the $T \rightarrow 0$ resistance anisotropy in a conventional (Fermi-liquid) metallic phase was shown to diverge on approach to a topological insulating phase with a spinon Fermi surface.

ACKNOWLEDGMENTS

We thank T. Senthil, J. Tranquada, and V. Calvera for helpful comments. S.A.K. was supported, in part, by NSF Grant No. DMR-1608055 at Stanford. E.B. was supported by the European Research Council (ERC) under grant HQ-MAT (Grant No. 817799), by the U. S.-Israel Binational Science Foundation (BSF), and by CRC 183 of the Deutsche Forschungsgemeinschaft. T.D. acknowledges support from the Charlotte Elizabeth Procter Fellowship at Princeton University.

APPENDIX: LAYERED GAUGE INVARIANCE FROM ANISOTROPIC LATTICE QED

In this Appendix, we argue that a model for an anisotropic lattice QED in $(3+1)$ D (which has the usual gauge symmetry) can be described at low energy as a model with a separate gauge symmetry on each layer such as those discussed in the main text.

Let us first describe the lattice $U(1)$ gauge theory [94]. On each bond in of the $(3+1)$ D hypercubic lattice, we define the $U(1)$ variable $U_\mu(\vec{x}) = \exp i \int_{\vec{x}}^{\vec{x}+\hat{\mu}} A_\mu(\vec{x}') dx'$, where $\mu = 0, 1, 2, 3$ corresponding to τ, x, y, z . The anisotropic $U(1)$

gauge theory is described by the Euclidean action

$$\begin{aligned} S_g &= - \sum_{\vec{x}, \mu, \nu < \mu} K_{\mu\nu} U_\mu(\vec{x}) U_\nu(\vec{x} + \hat{\mu}) U_\mu^\dagger(\vec{x} + \hat{\nu}) U_\nu^\dagger(\vec{x}) + \text{H.c.} \\ &\equiv - \sum_{\vec{x}, \mu, \nu < \mu} K_{\mu\nu} F_{\mu\nu}(\vec{x}) + \text{H.c.} \end{aligned} \quad (\text{A1})$$

which is the discretization of $\cos(\text{curl}A)$, with a direction-dependent coefficient $K_{\mu\nu}$ which encodes the anisotropy. The isotropic model $K_{\mu\nu} = K$ has a deconfined phase at weak coupling $K \ll 1$, and a confined phase at strong coupling $K \gg 1$, with a phase transition in-between. The anisotropic limit we wish to consider has $K_{\mu\nu} = K$ for $\mu, \nu < 3$, and $K_{3\mu} = K'$, in the limit $K \gg q$ and $K' \ll 1$. It was shown in Ref. [53] that in $(3+1)$ D in the pure gauge theory, this limit is always confined and therefore there is no stable layered phase [this reflects the fact that the $(2+1)$ D $U(1)$ pure gauge theory is always confined].

We now discuss fermions, which for simplicity we take to be the naive discretization of $(3+1)$ D (isotropic) Dirac fermions given by

$$\begin{aligned} S_f &= \frac{1}{2} \sum_{\vec{x}, \mu, n} \bar{\psi}_n(\vec{x}) \gamma_\mu [U_\mu^\dagger(\vec{x}) \psi_n(\vec{x} + \hat{\mu}) \\ &\quad - U_\mu(\vec{x} - \hat{\mu}) \psi_n(\vec{x} - \hat{\mu})], \end{aligned} \quad (\text{A2})$$

where $\psi, \bar{\psi} = \psi^\dagger \gamma_0$ are four-component Grassmann spinors, $n = 1, \dots, N$ label fermion flavors, and γ_μ satisfy $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$. Note that if we ignore the gauge field by setting $U = 1$, this actually leads to a low-energy description with $2^3 N$ Dirac fermions (due to fermion doubling). Since we are working at large N (and are interested in a layered phase which will have Dirac fermions associated with each layer anyway), this does not pose a problem.

The full action is

$$S = S_g + S_f \quad (\text{A3})$$

and has the gauge symmetry that involves sending

$$\psi_n(\vec{x}) \rightarrow e^{i\alpha(\vec{x})} \psi_n(\vec{x}), \quad (\text{A4})$$

$$\bar{\psi}_n(\vec{x}) \rightarrow e^{-i\alpha(\vec{x})} \bar{\psi}_n(\vec{x}), \quad (\text{A5})$$

$$U_\mu(\vec{x}) \rightarrow e^{i(\alpha(\vec{x}+\hat{\mu})-\alpha(\vec{x}))} U_\mu(\vec{x}). \quad (\text{A6})$$

We now argue that in the anisotropic limit $K' \ll 1 \ll K$, this theory describes a gapless layered phase with a separate gauge invariance in each plane.

In the limit $K' \ll 1$, U_z is strongly fluctuating and can be integrated out explicitly via a “high-temperature” expansion in K' . The effective action after integrating out U_z ,

$$S_{\text{eff}} = - \ln \int \mathcal{D}U_z e^{-S}, \quad (\text{A7})$$

can be obtained as a power series in K' . We use the convention

$$\int dU U^n = \delta_{n0} \quad (\text{A8})$$

for integrating over the $U(1)$ group element U .

Let us write $S_g = S_{g,\perp} + S_{g,z}$ and $S_f = S_{f,\perp} + S_{f,z}$ where $S_{g/f,z}$ involves terms with U_z , and $S_{g/f,\perp}$ are all other terms.

Since $S_{g,z}$ is proportional to K' , we may write $S = S_0 + K'S_1$ where $S_0 = S_{g,\perp} + S_f$ and $S_1 = S_{g,z}/K'$. Then,

$$S_{\text{eff}} = -\ln \int \mathcal{D}U_z e^{-S_0} \left(1 - K'S_1 + \frac{1}{2!}(K'S_1)^2 - \dots \right) \quad (\text{A9})$$

we see that the only terms which survive are those that contain no factors of any U_z . Furthermore, the logarithm means that only the connected terms survive.

At zeroth order in K' , we have

$$S_{\text{eff}}^{(0)} = S_{g,\perp} + S_{f,\perp} + \frac{1}{4} \sum_{\vec{x},n} [\bar{\psi}_n(\vec{x})\gamma_3\psi_n(\vec{x} + \hat{z})][\bar{\psi}_n(\vec{x} + \hat{z})\gamma_3\psi_n(\vec{x})], \quad (\text{A10})$$

a density-density type interaction term between neighboring layers.

At first order in K' , we have

$$S_{\text{eff}}^{(1)} = \frac{K'}{4} \sum_{\vec{x},\mu < 3,n,m} \bar{\psi}_n(\vec{x})\gamma_3\psi_n(\vec{x} + \hat{z})U_\mu^\dagger(\vec{x})U_\mu(\vec{x} + \hat{z}) \times \bar{\psi}_m(\vec{x} + \hat{z} + \hat{\mu})\gamma_3\psi_m(\vec{x} + \hat{\mu}) + (\psi \leftrightarrow \bar{\psi}, U \leftrightarrow U^\dagger), \quad (\text{A11})$$

which has the form of a gauge-invariant nearest-neighbor pair-hopping term involving two adjacent layers.

At second order, keeping only terms at fourth power in $\bar{\psi}, \psi$,

$$S_{\text{eff}}^{(2)} = \frac{(K')^2}{4} \sum_{\langle x,x' \rangle} \sum_{\text{paths } C} \sum_{m,n} \bar{\psi}_n(\vec{x})\gamma_3\psi_n(\vec{x} + \hat{z}) \times \left(\prod_{x'',\mu \in C} U_\mu^\dagger(\vec{x}'')U_\mu(\vec{x}'' + \hat{z}) \right) \times \bar{\psi}_m(\vec{x}' + \hat{z})\gamma_3\psi_m(\vec{x}') + (\psi \leftrightarrow \bar{\psi}, U \leftrightarrow U^\dagger), \quad (\text{A12})$$

where the first sum over $\langle x, x' \rangle$ is over all next-nearest-neighbor pairs with the same z coordinate, the second sum is over all paths C of length $\text{len}(C) = 2$ connecting x and x' , and the product of U is along the path connecting x and x' . This has the form of a gauge-invariant next-nearest-neighbor pair-hopping term.

At higher orders, we will have longer-range pair-hopping terms of this, made gauge invariant by the Wilson line operators summed over all paths C . At small K' , these terms decay exponentially with distance as $(K')^{\text{len}(C)}$, reflecting the fact that the U_z degrees of freedom are gapped and strongly fluctuating.

We may now take the continuum limit along τ, x, y , in which the exponentially decaying term becomes essentially local. The continuum model describes a stack of U(1) gauge theories coupled to Dirac fermions with a separate gauge invariance on each layer, and local quartic couplings between layers. As discussed in the main text, such quartic terms are irrelevant at low energies, for large N . By construction, S_{eff} does not depend on U_z and so therefore has a separate gauge invariance on each layer. The key result here is that the resulting theory is local; had K' been too large, highly nonlocal terms would have been generated upon integrating out U_z .

-
- [1] K. B. Efetov, Zh. Eksp. Teor. Fiz. **76**, 1781 (1979) [Sov. Phys.–JETP **49**, 905 (1979)].
- [2] S. E. Korshunov and A. I. Larkin, Phys. Rev. B **46**, 6395 (1992).
- [3] B. Horowitz, Phys. Rev. B **45**, 12632 (1992).
- [4] L. Golubovic and M. Golubovic, Phys. Rev. Lett. **80**, 4341 (1998).
- [5] C. S. O'Hern and T. C. Lubensky, Phys. Rev. Lett. **80**, 4345 (1998).
- [6] C. S. O'Hern, T. C. Lubensky, and J. Toner, Phys. Rev. Lett. **83**, 2745 (1999).
- [7] V. J. Emery, E. Fradkin, S. A. Kivelson, and T. C. Lubensky, Phys. Rev. Lett. **85**, 2160 (2000).
- [8] A. Vishwanath and D. Carpentier, Phys. Rev. Lett. **86**, 676 (2001).
- [9] S. L. Sondhi and K. Yang, Phys. Rev. B **63**, 054430 (2001).
- [10] R. Mukhopadhyay, C. L. Kane, and T. C. Lubensky, Phys. Rev. B **64**, 045120 (2001).
- [11] S. Tewari, J. Toner, and S. Chakravarty, Phys. Rev. B **72**, 060505(R) (2005).
- [12] S. Tewari, J. Toner, and S. Chakravarty, Phys. Rev. B **73**, 064503 (2006).
- [13] K. Shtengel, C. Nayak, W. Bishara, and C. Chamon, J. Phys. A: Math. Gen. **38**, L589 (2005).
- [14] V. Emery, in *Highly Conducting One-Dimensional Solids*, edited by J. Devreese *et al.* (Plenum, New York, 1979).
- [15] J. Solyom, Adv. Phys. **28**(2), 201 (1979).
- [16] C. Murthy and C. Nayak, Phys. Rev. Lett. **124**, 136801 (2020).
- [17] T. Senthil and M. P. A. Fisher, Phys. Rev. B **63**, 134521 (2001); Phys. Rev. Lett. **86**, 292 (2001).
- [18] Y. Werman, S. Chatterjee, S. C. Morampudi, and E. Berg, Phys. Rev. X **8**, 031064 (2018).
- [19] P. W. Anderson and Z. Zou, Phys. Rev. Lett. **60**, 132 (1988).
- [20] Y. Ando, G. S. Boebinger, A. Passner, N. L. Wang, C. Geibel, and F. Steglich, Phys. Rev. Lett. **77**, 2065 (1996).
- [21] Q. Li, M. Hucker, G. D. Gu, A. M. Tsvelik, and J. M. Tranquada, Phys. Rev. Lett. **99**, 067001 (2007).
- [22] E. Berg, E. Fradkin, E.-A. Kim, S. A. Kivelson, V. Oganesyan, J. M. Tranquada, and S. C. Zhang, Phys. Rev. Lett. **99**, 127003 (2007).
- [23] L. Balents and M. P. A. Fisher, Phys. Rev. Lett. **76**, 2782 (1996).
- [24] J. D. Naud, L. P. Pryadko, and S. L. Sondhi, Phys. Rev. Lett. **85**, 5408 (2000).

- [25] J. D. Naud, L. P. Pryadko, and S. L. Sondhi, *Nucl. Phys. B* **594**, 713 (2001).
- [26] M. Levin and M. P. A. Fisher, *Phys. Rev. B* **79**, 235315 (2009).
- [27] C. Chamon, *Phys. Rev. Lett.* **94**, 040402 (2005).
- [28] J. Haah, *Commun. Math. Phys.* **342**, 771 (2016).
- [29] B. Yoshida, *Phys. Rev. B* **88**, 125122 (2013).
- [30] S. Vijay, J. Haah, and L. Fu, *Phys. Rev. B* **92**, 235136 (2015).
- [31] R. M. Nandkishore and M. Hermele, *Annu. Rev. Condens. Matter Phys.* **10**, 295 (2019).
- [32] M. Pretko, X. Chen, and Y. You, *Int. J. Mod. Phys. A* **35**, 2030003 (2020).
- [33] A. Kitaev, *Ann. Phys.* **303**, 2 (2003).
- [34] J. B. Kogut, *Rev. Mod. Phys.* **51**, 659 (1979).
- [35] E. Fradkin and S. H. Shenker, *Phys. Rev. D* **19**, 3682 (1979).
- [36] S. Bravyi, M. Hastings, and S. Michalakis, *J. Math. Phys.* **51**, 093512 (2010).
- [37] K. Slagle and Y. B. Kim, *Phys. Rev. B* **96**, 195139 (2017).
- [38] W. Shirley, K. Slagle, and X. Chen, *Ann. Phys.* **410**, 167922 (2019).
- [39] T. Devakul, W. Shirley, and J. Wang, *Phys. Rev. Research* **2**, 012059 (2020).
- [40] T. H. Hansson, V. Oganesyan, and S. L. Sondhi, *Ann. Phys.* **313**, 497 (2004).
- [41] M. Freedman, C. Nayak, K. Shtengel, K. Walker, and Z. Wang, *Ann. Phys.* **310**, 428 (2004).
- [42] W. Shirley, K. Slagle, and X. Chen, *Phys. Rev. B* **102**, 115103 (2020).
- [43] X. G. Wen and A. Zee, *Phys. Rev. B* **46**, 2290 (1992).
- [44] X. Qiu, R. Joynt, and A. H. MacDonald, *Phys. Rev. B* **42**, 1339 (1990).
- [45] X. Qiu, R. Joynt, and A. H. MacDonald, *Phys. Rev. B* **40**, 11943 (1989).
- [46] L. Kong and X.-G. Wen, [arXiv:1405.5858](https://arxiv.org/abs/1405.5858); D. S. Freed, [arXiv:1406.7278](https://arxiv.org/abs/1406.7278).
- [47] T. Lan, L. Kong, and X. G. Wen, *Phys. Rev. B* **94**, 155113 (2016).
- [48] K. Gregor, D. A. Huse, R. Moessner, and S. L. Sondhi, *New J. Phys.* **13**, 025009 (2011).
- [49] K. Fredenhagen and M. Marcu, *Phys. Rev. Lett.* **56**, 223 (1986); *Nucl. Phys. Proc. Suppl.* **4**, 352 (1988).
- [50] A. Kitaev, *Ann. Phys.* **321**, 2 (2006).
- [51] M. Hermele, T. Senthil, M. P. A. Fisher, P. A. Lee, N. Nagaosa, and X.-G. Wen, *Phys. Rev. B* **70**, 214437 (2004).
- [52] S. M. Chester and S. S. Pufu, *J. High Energ. Phys.* **08** (2016) 069.
- [53] Y. K. Fu and H. B. Nielsen, *Nucl. Phys. B* **236**, 167 (1984); **254**, 127 (1985).
- [54] A. Hulsebos, C. P. Korthals-Altes, and S. Nicolis, *Nucl. Phys. B* **450**, 437 (1995).
- [55] P. W. Anderson, *Science* **235**, 1196 (1987).
- [56] G. Baskaran, Z. Zou, and P. W. Anderson, *Solid State Commun.* **63**, 973 (1987).
- [57] H. Yao, S.-C. Zhang, and S. A. Kivelson, *Phys. Rev. Lett.* **102**, 217202 (2009).
- [58] C. Wu, D. Arovas, and H.-H. Hung, *Phys. Rev. B* **79**, 134427 (2009).
- [59] G. Baskaran, G. Santhosh, and R. Shankar, [arXiv:0908.1614](https://arxiv.org/abs/0908.1614).
- [60] H.-H. Lai and O. I. Motrunich, *Phys. Rev. B* **84**, 085141 (2011).
- [61] M. Barkeshli, H. Yao, and S. A. Kivelson, *Phys. Rev. B* **87**, 140402(R) (2013).
- [62] M. Hermanns and S. Trebst, *Phys. Rev. B* **89**, 235102 (2014).
- [63] S. Baruch and D. Orgad, *Phys. Rev. B* **77**, 174502 (2008).
- [64] E. Berg, E. Fradkin, S. A. Kivelson, and J. M. Tranquada, *New J. Phys.* **11**, 115004 (2009).
- [65] R. Shankar, *Rev. Mod. Phys.* **66**, 129 (1994).
- [66] D. Kumar and R. Rajaraman, *Phys. Rev. B* **54**, 4379 (1996).
- [67] S.-S. Lee, *Phys. Rev. B* **80**, 165102 (2009).
- [68] M. A. Metlitski and S. Sachdev, *Phys. Rev. B* **82**, 075127 (2010).
- [69] T. Holder and W. Metzner, *Phys. Rev. B* **92**, 041112(R) (2015).
- [70] C. Nayak and F. Wilczek, *Nucl. Phys. B* **417**, 359 (1994).
- [71] D. F. Mross, J. McGreevy, H. Liu, and T. Senthil, *Phys. Rev. B* **82**, 045121 (2010).
- [72] B. L. Altshuler, L. B. Ioffe, and A. J. Millis, *Phys. Rev. B* **50**, 14048 (1994).
- [73] M. A. Metlitski, D. F. Mross, S. Sachdev, and T. Senthil, *Phys. Rev. B* **91**, 115111 (2015).
- [74] A non-Fermi liquid with no superconducting instability can also be constructed in a different setting: see S. Raghu, G. Torroba, and H. Wang, *Phys. Rev. B* **92**, 205104 (2015).
- [75] S. A. Kivelson, *Phys. Rev. B* **39**, 259 (1989).
- [76] A. J. Willans, J. T. Chalker, and R. Moessner, *Phys. Rev. Lett.* **104**, 237203 (2010).
- [77] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, *Phys. Rev. B* **50**, 7526 (1994).
- [78] E. Fradkin, *Phys. Rev. B* **33**, 3263 (1986).
- [79] M. Barkeshli, C.-M. Jian, and X.-L. Qi, *Phys. Rev. B* **87**, 045130 (2013).
- [80] Y. Ran, Yi Zhang, and A. Vishwanath, *Nat. Phys.* **5**, 298 (2009).
- [81] X.-G. Wen and Q. Niu, *Phys. Rev. B* **41**, 9377 (1990).
- [82] C.-W. von Keyserlingk, V. Khemani, and S. L. Sondhi, *Phys. Rev. B* **94**, 085112 (2016).
- [83] G. Grissonnanche, A. Legros, S. Badoux, E. Lefrançois, V. Zatkó, M. Lizaïre, F. Laliberté, A. Gourgout, J.-S. Zhou, S. Pyon *et al.*, *Nature (London)* **571**, 376 (2019).
- [84] R. Samajdar, S. Chatterjee, S. Sachdev, and M. S. Scheurer, *Phys. Rev. B* **99**, 165126 (2019).
- [85] J. H. Han, J.-H. Park, and P. A. Lee, *Phys. Rev. B* **99**, 205157 (2019).
- [86] R. Samajdar, M. S. Scheurer, S. Chatterjee, H. Guo, C. Xu, and S. Sachdev, *Nat. Phys.* **15**, 1290 (2019).
- [87] G. Grissonnanche, S. Theriault, A. Gourgout, M.-E. Boulanger, E. Lefrançois, A. Ataei, F. Laliberté, M. Dion, J.-S. Zhou, S. Pyon, T. Takayama, H. Takagi, N. Doiron-Leyraud, and L. Taillefer, *Nat. Phys.* (2020).
- [88] J.-Y. Chen, S. A. Kivelson, and X.-Q. Sun, *Phys. Rev. Lett.* **124**, 167601 (2020).
- [89] X. Li, B. Fauque, Z. Zhu, and K. Behnia, *Phys. Rev. Lett.* **124**, 105901 (2020).
- [90] T.-K. Ng and P. A. Lee, *Phys. Rev. Lett.* **99**, 156402 (2007).
- [91] R. Nandkishore, M. A. Metlitski, and T. Senthil, *Phys. Rev. B* **86**, 045128 (2012).
- [92] R. K. Kaul, Y. B. Kim, S. Sachdev, and T. Senthil, *Nat. Phys.* **4**, 28 (2008).
- [93] L. Zou and T. Senthil, *Phys. Rev. B* **94**, 115113 (2016).
- [94] K. G. Wilson, *Phys. Rev. D* **10**, 2445 (1974).