

Molecular Phylogenetics, Phylogenomics, and Phylogeography

Higher-Level Phylogeny and Reclassification of Lampyridae (Coleoptera: Elateroidea)

Gavin J. Martin^{1,10,*}, Kathrin F. Stanger-Hall², Marc A. Branham³, Luiz F. L. Da Silveira², Sarah E. Lower^{4,5}, David W. Hall⁴, Xue-Yan Li⁶, Alan R. Lemmon⁷, Emily Moriarty Lemmon⁸, Seth M. Bybee^{1,9}

¹Department of Biology, 4102 LSB, Brigham Young University, Provo, UT 84602, ²Department of Plant Biology, University of Georgia, Athens, GA 30602, ³Department of Entomology & Nematology, University of Florida, Gainesville, FL 32611, ⁴Department of Genetics, University of Georgia, Athens, GA 30602, ⁵Department of Biology, Bucknell University, Lewisburg, PA 17837, ⁶Max-Planck Junior Scientist Group on Evolutionary Genomics, Kunming Institute of Zoology, Chinese Academy of Sciences, 32 Jiaochang Donglu Kunming, Yunnan 650223 ⁷Department of Scientific Computing, Florida State University, Tallahassee, FL 32306, ⁸Department of Biological Science, Florida State University, Tallahassee, FL 32306, ⁹Monte L. Bean Museum of Natural History, Brigham Young University, Provo, UT 84602, and ¹⁰Corresponding author, email: gavin.jon.martin@gmail.com

Subject Editor: Bjarte Jordal

Received 1 May, 2019; Editorial decision 18 October, 2019

Abstract

Fireflies (Lampyridae Rafinesque) are a diverse family of beetles which exhibit an array of morphologies including varying antennal and photic organ features. Due in part to their morphological diversity, the classification within the Lampyridae has long been in flux. Here we use an anchored hybrid enrichment approach to reconstruct the most extensive molecular phylogeny of Lampyridae to date (436 loci and 98 taxa) and use this phylogeny to evaluate the higher-level classification of the group. None of the currently recognized subfamilies were recovered as monophyletic with high support. We propose several classification changes supported by both phylogenetic and morphological evidence: 1) *Pollaclasis* Newman, Vestini McDermott (incl. *Vesta* Laporte, *Dodacles* Olivier, *Dryptelytra* Laporte, and *Ledocas* Olivier), *Phototus* McDermott, and *Araucariocladus* Silveira & Mermudes are transferred to Lampyridae *incertae sedis*, 2) *Psilocladinae* McDermott, 1964 *status novum* is reestablished for the genus *Psilocladus* Blanchard, 3) *Lamprohizini* Kazantsev, 2010 is elevated to *Lamprohizinae* Kazantsev, 2010 *status novum* and *Phausis* LeConte is transferred to *Lamprohizinae*, 4) *Memoan* Silveira and Mermudes is transferred to *Amydetinae* Olivier, and 5) *Scissicauda* McDermott is transferred to *Lampyrinae* Rafinesque.

Key words: Firefly, phylogenomics, target capture, *Amydetinae*, *Lamprohizinae*

Fireflies (Lampyridae Rafinesque, 1815) are a cosmopolitan group of beetles with approximately 2,200 species (Branham 2010) and a varied and confounded history of classification (Olivier 1907, 1910a; Green 1959; McDermott 1966; Crowson 1972; Nakane 1991; Jeng 2008). Higher-level firefly classification schemes (e.g., Olivier 1907) have traditionally been framed in the context of morphological structures related to the production and detection of sexual signals (e.g., photic organs, eyes, and antennae). The morphological evolution of these features appears to have been driven by species-specific sexual communication, such as pheromones, glows, and flashes (Buck 1937, Lloyd 1971, Ohba 1983, Branham and Wenzel 2003, Stanger-Hall and Lloyd 2015, Stanger-Hall et al. 2018). Recent work suggests that bioluminescent sexual signals have evolved multiple times during firefly evolution and have been repeatedly lost, with a likely reversal to the use of pheromones for a

sexual signal (Branham and Wenzel 2003, Stanger-Hall et al. 2007, Martin et al. 2017). Therefore, morphologies related to these signals (production/reception) may represent instances of convergent evolution and the continued dependence on these morphologies to support firefly classification and taxonomy (Branham and Wenzel 2003) needs to be reconsidered.

The first catalogue of Lampyridae was authored by Ernst Olivier in 1907 and updated in 1910. In this initial taxonomic effort, Olivier included 9 subfamilies (based almost entirely on antennal morphology) and approximately 1,000 species. Drawing on the earlier work of Green (1948, 1959), McDermott (1964) rearranged the taxa into seven subfamilies and added a tribal classification for the Lampyrinae. However, he noted that his classification, while logical, did not reflect phylogeny and was ‘more or less arbitrary’, as it relied heavily on characters such as antennae and photic organ

morphology. This work leads to McDermott's (1966) supplement to the Olivier catalogue. Since McDermott, there have been two revised classification schemes for Lampyridae (Crowson 1972, Nakane 1991; see Table 1). The classification of the family by Crowson (1972) was incomplete as it did not include all lampyrid genera, and the work of Nakane (1991; published in Japanese) has largely been overlooked by current taxonomic workers. Recent phylogenetic efforts have provided deeper insight into the classification of fireflies by expanding morphological data sets (Branham and Wenzel 2001, 2003, Jeng 2008); molecular data sets (Stanger-Hall et al. 2007); or both (Martin et al. 2017). All of these studies highlighted the need to update the higher-level classification of this group within a phylogenetic framework. While focused on the evolution of specific traits, each of these studies recovered nonmonophyly of at least one subfamily. We are presenting here the first classification of Lampyridae that uses extensive taxon sampling and explicitly takes phylogenetic history into account.

Materials and Methods

Taxon Sampling

Our ingroup sample included 53 (of ~145) lampyrid genera and 88 species, representing seven of the eight subfamilies Ssensu Nakane 1991 (Table 2). However, the Otoretadriinae has since been synonymized under the Otoretinae (Janisova and Bocakova 2013). DNA-grade material for the Cheguevarinae was not available. The appropriate sister taxon to Lampyridae has been a topic of debate (Branham and Wenzel 2001, 2003, Stanger-Hall et al. 2007, Jeng 2008, Bocakova et al. 2007, Kundrata et al. 2014, Martin et al. 2017); therefore, 10 outgroup taxa from several closely related elateroid families were sampled: Omethidae, Eucnemidae, Cantharidae, Lycidae, Elateridae, Rhagophthalmidae, and Phengodidae (Table 2).

DNA Extraction

Specimens were collected in the field, preserved in 95% ethanol and stored at -80°C for long-term storage. Each specimen was independently identified by at least two coauthors. Muscle tissue was removed from a single metacoxae, when possible, for each specimen. In some instances, the small size of the specimen necessitated more of the specimen to be used. DNA was then extracted using a Qiagen DNeasy extraction kit. DNA samples were sent to the Center for Anchored Phylogenomics at Florida State University, Tallahassee, FL for anchored hybrid enrichment sequencing. The specimens and

remaining DNA samples were deposited as vouchers in the BYU Insect Genomics Collection in the Monte L. Bean Museum (BYU) of Natural History, Provo, UT, USA, the Lampyridae cryo-collection in the Stanger-Hall lab (UGA), Athens, GA, and the Coleção Entomológica Prof. José Alfredo Pinheiro Dutra, Departamento de Zoologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil (DZRJ); see Table 2.

Probe Design

In collaboration with the Center for Anchored Phylogenomics (www.anchoredphylogeny.com), an AHE kit representing the diversity of Lampyridae was developed, targeting a subset of the low-copy AHE 941 loci (exons) that Haddad et al. (2018) developed for Coleoptera. To this end, genomic resources, assembled genomes, assembled transcriptomes, and unassembled low-coverage genomic reads (unpublished data, Kathrin Stanger-Hall (K.S.H.); see Supp Table 1 [online only] for details), for 14 species of Lampyridae were gathered. In order to collect the low coverage genomic reads, Illumina libraries were prepared following Lemmon et al. (2012) and each was sequenced at 5x coverage on an Illumina HiSeq 2500 with a 150bp paired-end protocol with an 8-bp indexing read. After demultiplexing with no mismatches tolerated, overlapping reads were merged following Rokyta et al. (2012).

Target regions appropriate for Lampyridae were identified by following the general methodologies described in Hamilton et al. (2016). The assembled genomes and transcriptomes were scanned using all of the 941 *Tribolium castaneum* reference sequences developed by Haddad et al. (2018). After reads were mapped and extended to the same references, the reads were extended up to 2,000 bp in each direction to produce longer sequences. The resulting sequences were then aligned in Mafft v. 7.407 (Katoh and Standley 2013) and visually inspected in Geneious. Well-aligned regions were selected for downstream analysis. Of the 941 beetle exons identified by Haddad et al. (2018), only 586 regions of >180 bp in which at least 16 of 19 taxa were present were identified. Seventy overlapping loci were removed. Repetitive regions were masked using analyses of kmer distributions (see Hamilton et al. 2016 for details).

The final target regions were comprised of 516 target loci covering 578,322 bp. Probes were tiled at 2x coverage across each of the 19 lineages represented in the alignments. After removing redundant probes, 56,979 probes were submitted to Agilent Technologies for synthesis (SureSelect XP kit).

Library Preparation

From extracted DNA, libraries with standard Illumina adapters were prepared following Lemmon et al. (2012) and Prum et al. (2015) using a Beckmann Coulter FXp liquid-handling robot. After adding 8 bp indexes, libraries were pooled in groups of ~16 and enriched using the Lampyridae-specific enrichment kit described above. Enriched library pools were then pooled and sequenced on three Illumina 2500 sequencing lanes with C-bot clustering and a paired-end 150 bp protocol (~100 Gb of raw data were collected).

Read Assembly and Orthology Assessment

Raw reads passing the CASAVA high-chastity filter were merged following Rokyta et al. (2012) and assembled using the quasi-de novo approach described by Hamilton et al. (2016), with probe region sequences from *Phausis reticulata*, *Pyrractomena borealis*, *Photinus pyralis*, *Photinus scintillans*, *Phausis reticulata* and a soldier beetle (Cantharidae), serving as references. Assembly clusters derived from less than 42 reads were removed from downstream analyses in order to

Table 1. Comparison of major classification schemes in Lampyridae

Olivier 1910	McDermott 1966	Crowson 1972	Nakane 1991
Amydetinae	Amydetinae	Amydetinae	Amydetinae
Luciolinae	Luciolinae	Luciolinae	Luciolinae
Photurinae	Photurinae	Photurinae	Photurinae
Lampyrinae	Lampyrinae	Lampyrinae	Lampyrinae
Dadophorinae			
Lamprocerinae			
Photinae			
Lucidotinae			
Megalophthalminae			
	Pterotinae	Pterotinae	Pterotinae
	Matheteinae		
	Rhagophthalminae		
		Ototretinae	Ototretinae
		Cyphonocerinae	Psilodadinae
		Ototretadriinae	Ototretadriinae

Table 2. Taxa included in current study

Code	Family	Subfamily	Species	Coll.
CO1548	Cantharidae		Cantharidae sp. 1	BYU
KSH_12271	Cantharidae		Cantharidae sp. 2	UGA
CO1310	Elateridae		<i>Ampedus pomonae</i>	BYU
CO1077	Elateridae		<i>Pyrophorus</i> sp.	BYU
CO1309	Eucnemidae		<i>Microhagus pygmaeus</i>	BYU
CO1545	Lycidae		<i>Calopteron reticulatum</i>	BYU
CO1286	Omethidae		<i>Telegeusis nubifer</i>	BYU
CO1544	Phengodidae		<i>Phengodes</i> sp.	BYU
CO1285	Phengodidae		<i>Zarhipis integripennis</i>	BYU
CO1304	Rhagophthalmidae		<i>Rhagophthalmus obbai</i>	BYU
CO1212	Lampyridae	Amydetinae	<i>Amydetes fastigiata</i>	DZRJ
CO1211	Lampyridae	Amydetinae	<i>Cladodes illigeri</i>	DZRJ
CO1213	Lampyridae	Amydetinae	<i>Ethra axillaris</i>	DZRJ
CO1214	Lampyridae	Amydetinae	<i>Memoan ciceroi</i>	DZRJ
CO1216	Lampyridae	Amydetinae	<i>Psilocladus sigillatus</i>	DZRJ
CO1245	Lampyridae	Amydetinae	<i>Psilocladus</i> sp. 1	BYU
CO1218	Lampyridae	Amydetinae	<i>Psilocladus</i> sp. 2	BYU
CO1215	Lampyridae	Amydetinae	<i>Scissicauda disjuncta</i>	DZRJ
CO1305	Lampyridae	Amydetinae	<i>Vesta impressicollis</i>	BYU
CO1299	Lampyridae	Amydetinae	<i>Vesta saturnalis</i>	BYU
CO1219	Lampyridae	Amydetinae	<i>Vesta</i> sp.	BYU
CO1281	Lampyridae	Cyphonocerinae	<i>Pollaclasis bifaria</i>	BYU
CO1069	Lampyridae	Lampyrinae	<i>Aspisoma</i> sp.	BYU
CO1233	Lampyridae	Lampyrinae	<i>Aspisoma sticticum</i>	BYU
CO1234	Lampyridae	Lampyrinae	<i>Cratomorphus</i> sp.	BYU
CO1292	Lampyridae	Lampyrinae	<i>Diaphanes pectinealis</i>	BYU
CO1223	Lampyridae	Lampyrinae	<i>Diaphanes</i> sp. 1	BYU
CO1516	Lampyridae	Lampyrinae	<i>Diaphanes</i> sp. 2	BYU
CO1524	Lampyridae	Lampyrinae	<i>Diaphanes</i> sp. 3	BYU
CO1523	Lampyridae	Lampyrinae	<i>Diaphanes</i> sp. 4	BYU
KSH_6091	Lampyridae	Lampyrinae	<i>Ellychnia corrusca</i>	UGA
CO1197	Lampyridae	Lampyrinae	<i>Ellychnia</i> sp.	BYU
CO1067	Lampyridae	Lampyrinae	<i>Heterophotinus</i> sp.	BYU
CO1294	Lampyridae	Lampyrinae	<i>Lamprigera yunnanensis</i>	BYU
CO1060	Lampyridae	Lampyrinae	<i>Lamprocera</i> sp. 1	BYU
CO1459	Lampyridae	Lampyrinae	<i>Lamprocera</i> sp. 2	BYU
CO1457	Lampyridae	Lampyrinae	<i>Lamprocera</i> sp. 3	BYU
CO1231	Lampyridae	Lampyrinae	<i>Lamprocera</i> sp. 4	BYU
CO1287	Lampyridae	Lampyrinae	<i>Lamprohiza splendidula</i>	BYU
CO1071	Lampyridae	Lampyrinae	<i>Lampyrus noctiluca</i>	BYU
CO1551	Lampyridae	Lampyrinae	<i>Lucidota atra</i>	BYU
CO1232	Lampyridae	Lampyrinae	<i>Lucio blattinum</i>	BYU
CO1226	Lampyridae	Lampyrinae	<i>Lychnacris</i> sp.	BYU
CO1092	Lampyridae	Lampyrinae	<i>Microphotus</i> sp.	BYU
CO1227	Lampyridae	Lampyrinae	<i>Petalacmis</i> sp.	BYU
CO1549	Lampyridae	Lampyrinae	<i>Phausis reticulata</i>	BYU
CO1289	Lampyridae	Lampyrinae	<i>Phosphaenopterus metzneri</i>	BYU
KSH_17651	Lampyridae	Lampyrinae	<i>Photinus ardens</i>	UGA
KSH_1028	Lampyridae	Lampyrinae	<i>Photinus australis</i>	UGA
KSH_9013	Lampyridae	Lampyrinae	<i>Photinus brimleyi</i>	UGA
KSH_9670	Lampyridae	Lampyrinae	<i>Photinus carolinus</i>	UGA
KSH_10061	Lampyridae	Lampyrinae	<i>Photinus floridanus</i>	UGA
KSH_14611	Lampyridae	Lampyrinae	<i>Photinus granulatus</i>	UGA
CO1074	Lampyridae	Lampyrinae	<i>Photinus macdermotti</i> 1	BYU
KSH_87151	Lampyridae	Lampyrinae	<i>Photinus macdermotti</i> 2	UGA
CO1547	Lampyridae	Lampyrinae	<i>Photinus pyralis</i>	BYU
CO1224	Lampyridae	Lampyrinae	<i>Photinus</i> sp. 1	BYU
CO1528	Lampyridae	Lampyrinae	<i>Photinus</i> sp. 2	BYU
CO1057	Lampyridae	Lampyrinae	<i>Photinus stellaris</i>	BYU
CO1230	Lampyridae	Lampyrinae	<i>Photinini</i> sp.	BYU
CO1307	Lampyridae	Lampyrinae	<i>Pleotomodes needhami</i>	BYU
KSH_4081	Lampyridae	Lampyrinae	<i>Pyractomena borealis</i>	UGA
CO1063	Lampyridae	Lampyrinae	<i>Pyractonema</i> sp. 1	BYU

Table 2. Continued

Code	Family	Subfamily	Species	Coll.
COARG	Lampyridae	Lampyrinae	<i>Pyractonema</i> sp. 2	BYU
CO1293	Lampyridae	Lampyrinae	<i>Pyrocoelia pygidialis</i>	BYU
CO1552	Lampyridae	Lampyrinae	<i>Pyropyga decipiens</i>	BYU
CO1080	Lampyridae	Lampyrinae	<i>Pyropyga nigricans</i>	BYU
CO1306	Lampyridae	Lampyrinae	<i>Tenaspis angularis</i>	BYU
CO1291	Lampyridae	Luciolinae	<i>Abscondita cerata</i>	BYU
CO1297	Lampyridae	Luciolinae	<i>Asymmetricata circumdata</i>	BYU
CO1221	Lampyridae	Luciolinae	<i>Atyphella flammulans</i>	BYU
CO1520	Lampyridae	Luciolinae	<i>Australoluciola nigra</i>	BYU
CO1222	Lampyridae	Luciolinae	<i>Australoluciola</i> sp.	BYU
CO1298	Lampyridae	Luciolinae	<i>Curtos bilineatus</i>	BYU
CO1303	Lampyridae	Luciolinae	<i>Curtos obscuricolor</i>	BYU
CO1255	Lampyridae	Luciolinae	<i>Curtos</i> sp.	BYU
CO1295	Lampyridae	Luciolinae	<i>Emeia pseudosanteri</i>	BYU
CO1220	Lampyridae	Luciolinae	<i>Lloydella uberia</i>	BYU
CO1518	Lampyridae	Luciolinae	<i>Luciola</i> sp. 1	BYU
CO1225	Lampyridae	Luciolinae	<i>Luciola</i> sp. 2	BYU
CO1505	Lampyridae	Luciolinae	Luciolinae nr. <i>Luciola</i> 1	BYU
CO1525	Lampyridae	Luciolinae	Luciolinae nr. <i>Luciola</i> 2	BYU
CO1510	Lampyridae	Luciolinae	Luciolinae nr. <i>Luciola</i> 3	BYU
CO1509	Lampyridae	Luciolinae	Luciolinae nr. <i>Luciola</i> 4	BYU
CO1534	Lampyridae	Luciolinae	Luciolinae sp.	BYU
CO1517	Lampyridae	Luciolinae	<i>Pteroptyx</i> sp.	BYU
CO1296	Lampyridae	Luciolinae	<i>Pygoluciola qingyu</i>	BYU
CO1229	Lampyridae	Luciolinae	<i>Trisinuata</i> sp.	BYU
CO1308	Lampyridae	Ototretinae	<i>Drilaster</i> sp.	BYU
CO1302	Lampyridae	Ototretinae	<i>Stenocladus shirakii</i>	BYU
CO1290	Lampyridae	Photurinae	<i>Bicellonycha</i> sp. 1	BYU
CO1228	Lampyridae	Photurinae	<i>Bicellonycha</i> sp. 2	BYU
CO1546	Lampyridae	Photurinae	<i>Photuris congener</i>	BYU
CO1049	Lampyridae	Photurinae	<i>Photuris divisia</i>	BYU
KSH_406	Lampyridae	Photurinae	<i>Photuris frontalis</i>	UGA
KSH_8870	Lampyridae	Photurinae	<i>Photuris quadrifulgens</i>	UGA
CO1062	Lampyridae	Photurinae	<i>Pyrogaster</i> sp.	BYU
CO1550	Lampyridae	Pterotinae	<i>Pterotus obscuripennis</i>	BYU

mitigate the effects of any low-level contamination (clusters were typically comprised of 100–1,000 reads). Homologous consensus sequences were derived from the remaining clusters, with ambiguous base calls being made for each site with a base pattern that could not be explained by sequencing error, which was assumed to be no more than 1%. For each of the 516 target loci, orthologous sets of homologs were determined using a distance matrix (constructed by comparing kmer distributions) and a neighbor-joining approach for clustering (with at most one sequence per individual allowed in each ortholog cluster). After removing ortholog sets with less than 50% species representation (<51 of 98), 436 loci remained (Supp Table 2 [online only]).

Alignment

The sequences obtained for individual loci were aligned via Mafft v. 7.407 (Katoh and Standley 2013) using the default automatic alignment option. For maximum likelihood (ML) analyses, the alignments of the individual loci included flanking regions (head, tail, or both) of each locus and were concatenated before analysis and resulted in an alignment length of 1,833,533 base pairs. Breinholt et al. (2017) showed that trimming data from the probe or flanking region have little effect on the final topology; consequently, we elected not to trim these regions (Lemmon et al. 2012). However, it is possible that missing data may bias phylogenetic reconstruction (Lemmon et al. 2009; see also Wiens and Morrill 2011); therefore, we generated a second data set utilizing a portion of the pipeline of Breinholt

et al. (2017) (alignment_DE_trim.py) to remove sites from the alignment with <75% density (i.e., completeness), and with entropy >1.5 (to account for sites estimated to be nearly random). This resulted in a trimmed alignment of 401,423 base pairs.

Phylogenetic Reconstruction

ML reconstruction was performed in IQ-Tree v 1.6.8 (Nguyen et al. 2015, Chernomor et al. 2016) with 1,000 ultrafast bootstrap iterations (UFBoot; Hoang et al. 2018). Model estimation for concatenated locus data sets was carried out using ModelFinder (Kalyaanamoorthy et al. 2017) implemented within IQ-Tree, using BIC criteria. GTR+F+R9 was selected as the best-fit model for the untrimmed data set and GTR+R9 was selected for the trimmed data set. To account for the potential effect of incomplete lineage sorting on our phylogenetic reconstruction, we also used a coalescent approach to estimate a species tree via ASTRAL-III (Zhang et al. 2018) for each dataset (trimmed and untrimmed). ML gene tree topologies for each locus were inferred via IQ-Tree (Sensu Breinholt et al. 2017) before estimating the ASTRAL topology for each data set. We used localized posterior probabilities (Sayyari and Mirarab 2016) to assess nodal support for the coalescent species trees as assessed in ASTRAL-III.

Nomenclature

This paper has been registered in Zoobank (www.zoobank.org), the official register of the International Commission

on Zoological Nomenclature. The LSID (Life Science Identifier) number of the publication is: urn:lsid:zoobank.org:pub:D43452DB-0140-4863-84A8-94BFA6FD39EE

Results

Phylogenetic Reconstruction

ML Reconstruction

Within the ML reconstruction of the untrimmed data set (Topology 1 (T1); Fig. 1), we find a strong support for a sister group relationship between the Elateridae, Rhagophthalmidae, and Phengodidae with the Lampyridae (Fig. 1). The Luciolineae + *Lamprigera* Motschulsky were recovered as the sister lineage to all other firefly subfamilies with strong support (UFboot 100%). Within the lineage of the remaining subfamilies, a clade comprised of (*Pterotus* LeConte + *Pollaclasis* Newman) + the monophyletic Otoretinae forms the sister group to the remaining fireflies. However, neither this clade, nor the Otoretinae clade are well supported (UFboot: 72% for both nodes). The clade (*Lamprohiza* Motschulsky + *Phausis* LeConte) is recovered as sister to the clade ((*Psilocladus* Blanchard + (Amydetinae + Photurinae)) + Lampyrinae) with UFboot support of 100% at each node.

The ML reconstruction of the trimmed data set (Topology 2 (T2); Supp Fig. 1 [online only]) was largely identical to T1, with the exception of the placement of the paraphyletic Otoretinae and *Pollaclasis* in T2. In T2, Otoretinae is rendered paraphyletic, as *Stenocladus* Fairmaire in Deyrolle and Fairmaire is recovered as the sister taxon to all other Lampyridae, while *Drilaster* Kiesenwetter is recovered as the sister taxon to a clade comprised of *Pterotus* + *Pollaclasis*, albeit with low support (UFboot 77%). Compared with T1, there is also lower support along the backbone of T2 (Supp Fig. 1 [online only]).

Coalescent Reconstruction

The coalescent reconstruction of the untrimmed data set (Topology 3 (T3); Supp Fig. 2 [online only]) was largely congruent with the ML topology of the same data set, except for the placement of the Otoretinae (incl. *Pterotus*). In the coalescent analysis (T3), the Otoretinae are rendered polyphyletic by inclusion of *Pterotus*, and this clade is recovered as the sister group to all other fireflies, while in the ML analysis (T1) this position was held by Luciolineae + *Lamprigera* (Fig. 1; localized posterior probability (LPP): 0.95).

The coalescent reconstruction of the trimmed data set (Topology 4 (T4); Supp Fig. 3 [online only]) was identical in respect to subfamilial relationships in the untrimmed data set (T3), with only minor variation in species relationships within the major clades (i.e., placement of the lucioline clade comprised of *Abscondita cerata* and *Curtos* sp.; Supp Figs. 3 and 4 [online only]).

Consensus Among Analysis

In all analyses, Lampyridae was reconstructed as monophyletic with support values of UFboot: 100% or LPP: 1) With the exception of the Otoretinae (discussed separately below), our four different analyses yielded no topological differences that impact the monophyly of the subfamilies among Lampyridae. Therefore, we will use Topology 1 (Fig. 1) to discuss the phylogenetic relationships among the seven subfamilies sampled. Topology 1 (T1) is also the best supported (via ultrafast bootstrap analysis) and is based on the full, untrimmed data set.

In T1, the Luciolineae are reconstructed as paraphyletic with the inclusion of *Lamprigera*.

The Otoretinae were represented by two genera (*Drilaster* and *Stenocladus*) in our analyses, and these formed a monophyletic clade, albeit with low support (72%). T1 is the only topology to recover the Otoretinae as monophyletic and sister to a clade comprised of *Pterotus* and *Pollaclasis*. In both T2 and T4 (Supp Figs. 1 and 3 [online only]) *Stenocladus* was recovered as the sister lineage to all other fireflies with low support (UFboot: 77% & LPP: 0.75), while in T3 (Supp Fig. 2 [online only]) a clade consisting of *Stenocladus* + (*Drilaster* + *Pterotus*) is recovered as sister to the remaining fireflies (LPP: 0.95). Given this incongruence, the exact position of the Otoretinae among firefly subfamilies remains uncertain. However, in all phylogenetic reconstructions, *Pterotus* is either sister to, or a lineage within, the Otoretinae. The subfamily Amydetinae is recovered as polyphyletic and its taxa can be found closely associated with diverse subfamilies: *Psilocladus* is recovered as sister to the Amydetinae + Photurinae, *Pollaclasis* forms a clade with *Pterotus* (Pterotinae), *Vesta* Laporte is recovered within the photurine clade, and *Cladodes* Solier, *Ethra* Laporte, and *Scissicauda* McDermott are recovered within the Lampyrinae. The Photurinae are rendered paraphyletic by inclusion of the old-world *Vesta* (see Martin et al. 2017). Lampyrinae is recovered as polyphyletic: *Lamprigera* is recovered within the Luciolineae, and there is high support (UFboot: 100%) for a monophyletic *Lamprohiza* + *Phausis* clade outside the Lampyrinae (see above). In addition, several amydetine taxa (*Cladodes*, *Ethra*, *Scissicauda*) cluster within the remaining Lampyrinae.

As such, in T1, the only subfamily recovered as monophyletic, as currently classified, is the Otoretinae. However, across all other topologies, even the Otoretinae are also rendered nonmonophyletic.

Discussion

The higher-level classification of Lampyridae is in need of revision. Our molecular analyses confirm that a few morphological characters related to sexual communication, many likely convergent, cannot be central to diagnosing a natural classification of Lampyridae. The use of these characters in subfamilial classification needs to be reassessed in light of a more complete phylogenetic history of the group. Our phylogenetic analyses based on 436 loci and 98 taxa provide a robust framework for a revision of the classification of Lampyridae.

Luciolineae

The Luciolineae are a subfamily of fireflies that have received much attention over the years (for a taxonomic treatment, see Ballantyne et al. 2015) and seem to share stable morphological characters such as a reduced number of abdominal ventrites in the males (see discussion of *Lamprigera* below).

Our analyses largely agree with recent morphological analyses in terms of the major groupings of the Luciolineae (Ballantyne and Lambkin, 2013). Thanks in large part to the effort of Ballantyne and Lambkin (Ballantyne 1987a, Ballantyne and Lambkin 2009, 2013), there also exists a discussion of the aedeagal sheath, which has provided evidence in support of the current taxonomic framework for these taxa. Many of the morphological characters used in these analyses (Ballantyne and Lambkin 2009, 2013, Ballantyne et al. 2015, 2016) and the phylogenetic patterns they support, correlate well with the phylogenetic hypotheses generated from our molecular data. However, some minor differences between previous phylogenetic work and our overall topology do exist. For example, the *Atyphella* complex is not recovered as the most highly derived lineage, as it is in Ballantyne and Lambkin (2013). Our molecular data set recovers this lineage as more basal compared with the other




Fig. 1. ML reconstruction of full, untrimmed dataset aligned in MAFFT; log-likelihood = -26,321,164.240. The numbers at nodes indicate UFBoot support. Unlabeled nodes have UFBoot values of 100%, colors represent the newly proposed classification. Star at node depicts Lampyridae; A.-I. Dorsal habitus of genus representatives: A. *Lamprigera yunnana*; B. *Luciola cyathigera*; C. *Stenocladus davidi*; D. *Amydetes fastigiata*; E. *Vesta chevrolati*; F. *Diaphanes* sp.; G. *Aspisoma gentile*; H. *Ellychnia aurora*; I. *Photinus laticollis*;

Luciolinae. In addition, the lampyrine genus *Lamprigera* is recovered as a lineage within the lucioline clade.

Jeng et al. (2000) remarked that the systematic position of *Lamprigera* was uncertain and would hopefully be more accurately defined through future phylogenetic investigation. Previous analyses have recovered *Lamprigera* in various positions within the Lampyridae. In 2006, on the basis of the 16S mitochondrial marker, Li et al. performed a phylogenetic reconstruction of the Lampyridae. In this analysis, *Stenocladus* was recovered as the basal lineage, while *Lamprigera* was recovered in a surprising clade together with the amydetine genus *Vesta* sister to the remaining fireflies (Li et al. 2006), albeit with low support. In his thesis, Jeng (2008) recovered *Lamprigera* sister to the *Phausis* + *Lamprohiza* using >400 morphological characters. Wang et al. (2017) recovered *Lamprigera* as a member of the Lampyrinae on the basis of 13 mitochondrial genes. It should be noted, however, that the Wang et al. (2017) analysis suffered from a limited taxon sampling (six lucioline taxa and two additional lampyrine taxa from a single genus). Martin et al. (2017) recovered *Lamprigera* as sister to the monophyletic Luciolinae with high support and placed it as Lampyridae *incertae sedis*.

Here we recover *Lamprigera* as a member of the Luciolinae for the first time with both strong support and congruence between all of our analyses. However, this placement is based on a single *Lamprigera* species and major morphological differences between *Lamprigera* and the luciolines, e.g., the number of abdominal ventrites, need to be addressed. A major, long-standing morphological synapomorphy for Luciolinae has been males with six abdominal ventrites, whereas all other Lampyridae have seven or eight abdominal ventrites. Males of *Lamprigera* exhibit the 'typical' abdominal morphology of Lampyridae with seven ventrites.

In certain elateroid lineages, ventrite number is known to vary greatly, even within recently derived tribes/subfamilies (Kundrata and Bocak 2019). In contrast, ventrite number has not been shown to vary in Luciolinae. To rigorously test the classification of *Lamprigera* relative to the Luciolinae, an expanded taxon sampling including deeper species coverage within the *Lamprigera*, combined with an in-depth morphological investigation, including the plasticity of ventrite number across these taxa will be needed. Until these analyses can be done, we elect to keep *Lamprigera* as Lampyridae *incertae sedis* (Martin et al., 2017).

Pterotinae, Cyphonocerinae, and Psilocladinae (McDermott, 1964) stat. nov.

Jeng et al. 1998, 2006, summarizing the work of McDermott (1966), Crowson (1972), and Nakane (1991), were the first to formally delineate the Psilocladinae (previously known as Cyphonocerinae), by identifying the constituents of the group (*Cyphonocerus* Kiesenwetter, *Psilocladus*, and *Pollaclasis*) and laying out nine morphological features uniting the group. Jeng et al. (1998) also recognized Cyphonocerinae as a subjective synonym of Psilocladinae based on priority. In 2016, Silveira et al. treated *Scissicauda* as a member of the amydetine subtribe Psilocladina sensu McDermott 1964, distinguishing *Scissicauda* from the other members of the group (*Ethra*, *Photocetus* McDermott, *Psilocladus*, and *Pollaclasis*). In 2017, another genus (*Araucariocladus* Silveira & Mermudes) was added to the Psilocladina sensu McDermott. As a consequence of recognizing Psilocladina sensu McDermott instead of Psilocladinae sensu Jeng/Nakane, *Cyphonocerus* was left as the sole member of the Cyphonocerinae.

Our phylogenetic analyses supports in part the classification presented by Crowson (1972) and Nakane (1991) and supports the classification of *Psilocladus* as a separate lineage from

the Amydetinae. Without *Cyphonocerus* in our taxon sample, the placement of *Psilocladus* within the Cyphonocerinae cannot truly be tested; however, given that *Psilocladus* did not form a monophyletic lineage with *Pollaclasis*, we formally recognize the subfamily Psilocladinae, distinguished by the antennae with 11 articles, articles 2–11 with two weak, ciliate branches, with the only constituent genus being *Psilocladus*.

Following McDermott (1966) and Silveira et al. (2016), *Pollaclasis* is classified as a member of the Amydetinae. However, both our ML and coalescent analyses challenge this classification. The ML analysis places *Pollaclasis* as a sister taxon to the North American *Pterotus*, the sole member of the Pterotinae (Fig. 1). In contrast, the coalescent analysis places *Pollaclasis* as sister to a *Phausis* + *Lamprohiza* clade (Supp Fig. 2). Hypothesized to be a close relative of *Cyphonocerus*, *Pollaclasis* has previously been classified in the Psilocladinae based on the morphology of antennae, mandibles, and abdominal segmentation (see Jeng et al. 1998). The present analyses do not support the monophyly of *Pollaclasis* and *Psilocladus*. On the basis of this evidence, as well as the absence of *Cyphonocerus* in our taxon sample, we transfer *Pollaclasis* to Lampyridae *incertae sedis*. Future efforts need to be made to ascertain whether *Pollaclasis* is indeed a member of the Cyphonocerinae, or as the ML analyses suggest, more closely related to *Pterotus*.

Ototretinae

Our analyses indicate that there may be a close relation between the Ototretinae and the Pterotinae. Recently, the Ototretinae were revised, and their taxonomic placement was addressed (Janisova and Bocakova 2013). In that study, the monotypic Ototretadrilinae were synonymized under the Ototretinae, and the ototretine lineage was transferred from Elateriformia *incertae sedis* to Lampyridae. In addition, the genus *Stenocladus* was transferred from Elateriformia *incertae sedis* to Ototretinae. Our analyses corroborate these transfers; however, as our taxon sample included only 2 of the 18 genera from the Ototretinae, *Stenocladus* and *Drilaster*, a wider generic sampling is needed to address the phylogenetic history of this lineage.

Two other genera, *Anadrilus* Kirsch, 1875 and *Pachytarsus* Motschulsky, 1861, were both excluded from the Ototretinae by Janisova and Bocakova (2013) on the basis of lack of type material to examine. As we were also unable to sample these genera, they remain Lampyridae *incertae sedis*. Of note, when he described *Pachytarsus*, Motschulsky (1861) was apparently unaware of the true bug genus by the same name, having been described the year before: *Pachytarsus* Fieber, 1860 (for current usage, see Ballal et al. 2018). Due to the name *Pachytarsus* being preoccupied, we propose the replacement name *Crassitarsus* Martin.

Crassitarsus Martin

New Replacement Name

Pachytarsus Motschulsky, 1861

Included species: *Crassitarsus basalis* (Motschulsky, 1861) NEW COMBINATION, *Crassitarsus bicolor* (Pic, 1929) NEW COMBINATION, *Crassitarsus bryanti* (Wittmer, 1940) NEW COMBINATION, *Crassitarsus lateralis* (Motschulsky, 1861) NEW COMBINATION, *Crassitarsus longicornis* (Pic, 1921) NEW COMBINATION, *Crassitarsus minutus* (Pic, 1933) NEW COMBINATION, *Crassitarsus obscurus* (Pic, 1927) NEW COMBINATION, *Crassitarsus testaceus* (Motschulsky, 1861) NEW COMBINATION.

Etymology: *Crassus*, a synonym for *pachy*, both meaning thick, seems an apt replacement to conserve the original thoughts of Motschulsky.

Lamprohizinae Kazantsev, 2010 stat. nov.

There is strong support for a *Lamprohiza* + *Phausis* clade in all our topologies. This is supported by a high degree of morphological similarity between these two genera (Fig. 2). *Phausis* was erected for *Lampyrus reticulata* Say by LeConte (1852) and *Lamprohiza* was erected in 1853 by Motschulsky for *Lampyrus splendidula* Linnaeus. *Lamprohiza* was later synonymized with *Phausis* by Lacordaire in 1857. In contrast, Mulsant, 1862 treated *Lamprohiza* as an independent genus, but in 1881 LeConte wrote '[*Phausis*] is not sufficiently distinct from the European *Lamprohiza*, and in fact the European species seems to have been naturalized in Maryland and Illinois' and included *L. splendidula* within *Phausis*. This classification was accepted until 1964 when McDermott separated *Lamprohiza* from *Phausis* by the 'minute appendage on the 11th antennal article' of the latter. However, Fender (1966) treated the two genera as one in his treatment on the '*Phausis*' of North America, while Miksic (1969) followed McDermott in treating them as separate genera. From a phylogenetic perspective, Stanger-Hall et al. (2007) found *Phausis* as sister to Photurinae + Lampyrinae, similar to our results. In 2008, Jeng found support for these genera as members of Lampyrinae, however, he noted that they differed from the traditional Lampyrinae in the 'unmodified mandibles *sensu* Green (1949), dorsal abdominal spiracles, and a symmetrical

aedeagal sheath.' Our analyses, based on the type species of each genus, strongly support the rank of subfamily and we herein elevate Lamprohizini Kazantsev, 2010 to Lamprohizinae Kazantsev, 2010 stat. nov.

Diagnosis

The Lamprohizinae are distinguished from all other subfamilies with the following combination of characters, based on adult males: mandibles unmodified (i.e., not reduced in size); antennae filiform, 11-segmented, with or without terminal sensorium, if without then posterior margin of ventrite 7 with weak to strong medial projection, projection emarginate at midline; tarsal claws simple, not bifid; abdomen with seven–eight ventrites; abdominal spiracles dorsal; aedeagal sheath symmetrical.

Key to the genera of Lamprohizinae

Terminal antennomere with small, beadlike sensorium; abdominal ventrite 7 unmodified; ventrite 8 apparent*Phausis*
 Terminal antennomere without small, beadlike sensorium; abdominal ventrite 7 with weak to strong medial projection; ventrite 8 not apparent*Lamprohiza*

Amydetinae

McDermott (1964) recognized the Amydetini as a tribe within the Lampyrinae based on the mandibles being 'normal, arcuate, regularly narrowing to tips', antennae bearing rami or branches, and lack of secondary elytral pubescence of the males. To further classify the rather heterogeneous group of genera included in the Amydetini, McDermott (1964) recognized three subtribes based on variation in antennal morphology: Amydetina ('Antennae with more than 14 articles, uniramous'), Vestina ('Antennae with 11 articles, Antennae uniramous, rami remiform, relatively broad, folding like a fan'), and Psilocladina ('Antennae with 11 articles, Antennae bi- or uniramous, rami either long and diffuse, or if short and straight, not fan-folding'). In his 1966 update to the classification, McDermott elevated the Amydetini to the subfamily level. Nakane (1991) reduced the Amydetinae to *Amydetes* Hoffmannsegg in Illiger and *Magnoculus* McDermott; however, Silveira and Mermudes (2013) described the new genus *Memoan* Silveira and Mermudes and, based on it having features of both the Amydetinae and the Lampyrinae, placed it in Lampyridae *incertae sedis*.

Owing to the phylogenetic position of *Memoan* as sister to *Amydetes*, and the morphological features shared by *Memoan* with other taxa within the Amydetinae (as identified by Silveira and Mermudes, 2013), including a 'continuous glow, pleuralventral suture, ventral approximate eyes, deep punctures on pronotum and scutellum, and absence of tibial spurs', we formally transfer *Memoan* to the Amydetinae, thus the Amydetinae now include *Amydetes*, *Magnoculus*, and *Memoan*.

Cheguevariinae

Kazantsev (2006) described two enigmatic species in the genus *Cheguevaria* Kazantsev, 2006, one from Puerto Rico, and one from Dominican Republic. In the description, he remarks on the similarity of *Cheguevaria* to the Phengodidae; however, he does note the differences, and allies *Cheguevaria* with the Lampyridae based on the structure of the genital capsule. While he suspected that the genus represented a unique lampyrid subfamily, given the peculiar morphology of the genus, as well as the lack of phylogenetic stability at the time, Kazantsev placed the Cheguevariini as Lampyridae




Fig. 2. Lamprohizinae habitus images; A: *Phausis reticulata* dorsal; B: *Phausis reticulata* ventral; C: *Lamprohiza delarouzei* dorsal; D: *Lamprohiza delarouzei* ventral.

incertae sedis (Kazantsev, 2006). Recent phylogenetic analyses have corroborated the placement of *Cheguevaria* within the Lampyridae and, confirming the suspicion of Kazantsev, elevated the tribe to subfamilial status (Ferreira, et al. 2019). We were unable to sample the Cheguevariinae for the present analyses.

Photurinae and Lampyrinae

All photurine genera in our analyses (we were unable to sample *Presbyolampis* Buck, 1947) were recovered in a monophyletic lineage with the old-world *Vesta*. Known only from the New World, the Photurinae are typically characterized by bifurcate claws in males, and eyes expanding beyond the hypomeron. Nakane (1991) transferred the members of the Vestina (*Vesta*, *Cladodes*, *Ledocas* Olivier, *Dodacles* Olivier, and *Dryptelytra* Laporte) to the Lampyrinae, and moved *Psilocladus* to the Cyphonocerinae (see above) and *Ethra* to the Lampyrinae; he did not mention *Scissicauda* or *Photoctus*. Our analyses agree with Nakane (1991) in the placement of *Cladodes* and *Ethra*; however, the old-world *Vesta* sampled for our analyses formed a monophyletic lineage within the Photurinae, a new-world clade. *Vesta* is a speciose genus with representatives in both the Old and New World. McDermott (1964) recognized that *Vesta* might not be a natural group. In his 1966 catalogue he continued to treat the new-world taxa as potentially independent from the old-world taxa. As we were unable to sample the new-world *Vesta*, or the other members of the Vestini, our analyses are inconclusive with regard to the subfamilial placement of the *Vesta*. Therefore, we transfer *Vesta* and the other members of the Vestini (*Dodacles*, *Dryptelytra*, and *Ledocas*) to Lampyridae *incertae sedis* pending further morphological and phylogenetic analyses.

The Lampyrinae are a diverse assemblage of genera lacking any stable morphological characters (McDermott 1964; Jeng 2008). It is in part due to this that we believe so many ‘lampyrine’ genera are/have been mis-classified. Based on our present phylogenetic analyses, we transfer the genus *Scissicauda* from Amydetinae into Lampyrinae. Since neither *Photoctus* nor *Araucariocladus* were included in the present analyses we transfer these to Lampyridae *incertae sedis*, pending further phylogenetic analyses.

Conclusions

With 88 lampyrid taxa in 53 genera for seven of the eight previously recognized subfamilies, these analyses represent the largest molecular phylogeny of the group, thereby providing a unique opportunity to robustly test the monophyly of each subfamily. The results of our four phylogenetic reconstructions were largely congruent, with the only major difference concerning the placement of the Otoretinae, which were under-sampled in this study. The inclusion of additional otoretine taxa will likely aid the resolution of their placement. Aside from the monotypic Pterotinae, among the subfamilies in the current analyses, only the Otoretinae (as currently defined) were reconstructed as monophyletic (T1). As a result, we propose several changes to the higher-level classification of Lampyridae, with an updated higher-level classification scheme for the family. *Pollaclasis*, *Vesta* (as well as the other constituents of the Vestini: *Dodacles*, *Dryptelytra*, and, *Ledocas*), *Photoctus*, and *Araucariocladus* are transferred to Lampyridae *incertae sedis*. *Psilocladinae* is re-established for *Psilocladus*. Amydetinae is limited to *Amydetes*, *Magnoculus*, and *Memoan* with *Scissicauda* being transferred to the Lampyrinae. *Lamprohizini* is elevated to the rank of subfamily, *Phausis* is transferred to Lamprohizinae. This comprehensive classification of firefly lineages will provide the phylogenetic scaffold in support

of future studies into the evolution of the fascinating morphology and behavior of fireflies.

List of Genera

Subfamily Luciolinae Lacordaire, 1857

- Abscondita* Ballantyne, Lambkin, & Fu in Ballantyne et al., 2013
Luciola terminalis Olivier, 1883: 330
Aquatica Fu, Ballantyne, and Lambkin, 2010
Aquatica wuhana Fu, Ballantyne, and Lambkin, 2010: 8
Aquilonia Ballantyne in Ballantyne and Lambkin, 2009
Luciola costata Lea, 1921: 66
Asymmetricata Ballantyne in Ballantyne and Lambkin, 2009
Luciola circumdata Motschulsky, 1854b: 50
Atyphella Olliff, 1890
Atyphella lychnus Olliff, 1890: 647
Australoluciola Ballantyne in Ballantyne and Lambkin, 2013
Lampyris australis Fabricius, 1775: 201
Colophotia Motschulsky, 1853
Lampyris praeusta Eschscholtz, 1822: 57
Convexa Ballantyne in Ballantyne and Lambkin, 2009
Luciola wolffi Olivier, 1910b: 343
Curtos Motschulsky, 1845
Curtos mongolicus Motschulsky, 1845: 36
Emarginata Ballantyne in Ballantyne et al., 2019
Luciola trilucida Jeng and Lai in Jeng et al., 2003: 248
Emeia Fu, Ballantyne and Lambkin, 2012
Curtos pseudosauteri Geisthardt, 2004: 1
Inflata Boontop in Ballantyne et al., 2015
Luciola indica Motschulsky, 1854b: 53
Kuantana Ballantyne in Ballantyne et al., 2019
Kuantana menayah Ballantyne in Ballantyne et al., 2019: 82
Lampyroidea Costa 1875
Lampyroidea syriaca Costa 1875: CLXIX
Lloydidiella Ballantyne in Ballantyne and Lambkin, 2009
Luciola majuscula Lea, 1915: 495
Luciola Laporte, 1833
Luciola pedemontana Motschulsky, 1853: 53
Magnalata Ballantyne in Ballantyne and Lambkin, 2009
Luciola limbata Blanchard, 1853: 73
Medeopteryx Ballantyne in Ballantyne and Lambkin, 2013
Pteropteryx effulgens Ballantyne, 1987b: 141
Missimia Ballantyne in Ballantyne and Lambkin, 2009
Missimia flavida Ballantyne in Ballantyne and Lambkin, 2009
Pacifica Ballantyne in Ballantyne and Lambkin, 2013
Atyphella salomonis Olivier, 1911c: 172
PhoturoLuciola Pic, 1931
PhoturoLuciola deplanata Pic, 1931: 12
Pteropteryx Olivier, 1902
Luciola testacea Motschulsky, 1854b: 48
Pygatyphella Ballantyne, 1968
Atyphella obsoleta Olivier, 1911c: 174
Pygoluciola Wittmer, 1939
Pygoluciola styliifer Wittmer, 1939: 22
Pyrophanes Olivier, 1885
Pyrophanes similis Olivier, 1885: 370
Sclerotia Ballantyne in Ballantyne et al., 2016
Luciola aquatilis Thancharoen in Thancharoen et al., 2007: 56
Serratia Ballantyne in Ballantyne et al. 2019
Serratia sibuyaniana Ballantyne in Ballantyne et al., 2019: 147
Triangulara Pimpasalee in Ballantyne et al., 2016

Triangulara frontoflava Pimpasalee in Ballantyne et al., 2016: 242

Trisinuata Ballantyne in Ballantyne and Lambkin, 2013

Trisinuata caudabifurca Ballantyne in Ballantyne and Lambkin, 2013: 117

Tribe *Pristolycini* Kazantsev, 2010

Pristolycus Gorham, 1883

Pristolycus sagulatus Gorham, 1883: 407

Subfamily *Pterotinae* LeConte, 1861

Pterotus LeConte, 1859

Pterotus obscuripennis LeConte, 1859: 86

Subfamily *Ototretinae* McDermott, 1964

Baolacus Pic, 1915

Baolacus lajoyei Pic, 1915: 21

Brachylampis Van Dyke, 1939

Brachylampis sanguinicollis Van Dyke, 1939: 16

Brachypterodrilus Pic, 1918

Brachypterodrilus pallidipes Pic, 1918: 1

Ceylanidrilus Pic, 1911

Ceylanidrilus bipartitus Pic, 1911a: 187

Drilaster Kiesenwetter, 1879

Drilaster axillaris Kiesenwetter, 1879: 311

Emasia Bocakova and Janisova, 2010

Emasia dentata Bocakova and Janisova, 2010: 61

Eugeusis Westwood, 1853

Eugeusis palpator Westwood, 1853: 239

Falsophaeopterus Pic, 1911

Falsophaeopterus fruhstorferi Pic, 1911a: 187

Flabellopalpodes Bocakova and Bocak, 2016

Flabellopalpodes flavus Bocakova and Bocak, 2016: 372

Flabellototreta Pic, 1911

Flabellototreta fruhstorferi Pic, 1911b: 156

Gorhamia Pic, 1911

Gorhamia compressicornis Pic, 1911a: 187

Harmatelia Walker, 1858

Harmatelia bilinea Walker, 1858: 281

Hydaspidoides Bocakova and Janisova, 2013

Hydaspidoides kanarensis Janisova and Bocakova, 2013: 7

Hyperstoma Wittmer, 1979

Hyperstoma marginata Wittmer, 1979: 86

Lamellipalpodes Maulik, 1921

Lamellipalpodes annandalei Maulik, 1921: 584

Lamellipalpus Maulik, 1921

Eugeusis nigripennis Pascoe, 1887: 10

Ototretadrilus Pic, 1921

Ototretadrilus atritarsis Pic, 1921: 13

Picodrilus Wittmer, 1938

Ototreta drescheri Pic, 1937: 138

Stenocladus Fairmaire in Deyrolle and Fairmaire, 1878

Stenocladus davidis Fairmaire in Deyrolle and Fairmaire, 1878: 113

Subfamily *Lamprohizinae* Kazantsev, 2010

Phausis LeConte, 1852

Lampyrus reticulata Say, 1825: 163

Lamprohiza Motschulsky, 1853

Lampyrus splendidula Linnaeus, 1767: 644

Subfamily *Cyphonocerinae* Crowson, 1972

Cyphoncerus Kiesenwetter, 1879

Cyphoncerus ruficollis Kiesenwetter, 1879: 312

Subfamily *Psilocladinae* McDermott, 1964

Psilocladus Blanchard, 1846

Psilocladus miltoderus Blanchard, 1846: 122

Subfamily *Amydetinae* Olivier in Wytsman, 1907

Tribe *Amydetini* Olivier in Wytsman, 1907

Amydetes Hoffmannsegg in Illiger, 1807

Homalisus apicalis Germar, 1824: 67

Magnoculus McDermott, 1964

Meglophthalmus bennetti Gray, 1832: 371

Memoan Silveira and Mermudes, 2013

Memoan ciceroi Silveira and Mermudes, 2013: 84

Subfamily *Cheguevariinae* Kazantsev, 2006

Tribe *Cheguevariini* Kazantsev, 2006

Cheguevaria Kazantsev, 2006

Cheguevaria taino Kazantsev, 2006: 370

Subfamily *Photurinae* Lacordaire, 1857

Bicellonycha Motschulsky, 1853

Bicellonycha deleta Motschulsky, 1854b: 58

Photuris Dejean, 1833

Lampyrus versicolor Fabricius, 1798: 123

Presbyolampis Buck, 1947

Presbyolampis immigrans Buck, 1947: 75

Pyrogaster Motschulsky, 1853

Pyrogaster grylloides Motschulsky, 1853: 53

Subfamily *Lampyrinae* Rafinesque, 1815

Tribe *Cratomorphini* Green, 1948

Aspisoma Laporte, 1833

Lampyrus ignita Linnaeus, 1767: 645

Aspisomoides Zaragoza-Caballero, 1995

Aspidosma bilineatum Gorham, 1880b: 86

Cassidomorphus Motschulsky, 1853

Cassidomorphus silphoides Motschulsky, 1853: 36

Cratomorphus Motschulsky, 1853

Photinus fabricii Laporte, 1840: 268

Micronaspis Green, 1948

Micronaspis floridana Green, 1948: 63

Paracratomorphus Zaragoza-Caballero, 2013

Paracratomorphus reyesi Zaragoza-Caballero, 2013: 145

Pyractomena Melsheimer, 1846

Pyractomena lucifera Melsheimer, 1846: 304

Tribe *Lamprocerini* Olivier, 1907

Alecton Laporte, 1833

Alecton discoidalis Laporte, 1833: 135

Lamprocera Laporte, 1833
Homalilus grandis Sturm, 1826: 58
Lucernuta Laporte, 1833
Lampyrus savignii Kirby, 1818: 388
Lucio Laporte, 1833
Lucio abdominalis Laporte, 1833: 136
Lychnacris Motschulsky, 1853
Lychnacris triguttula Motschulsky, 1853: 33
Tenaspis LeConte, 1881
Hyas angularis Gorham, 1880: 7

Tribe Lampyrini Rafinesque, 1815

Afrodiaphanes Geisthardt, 2007
Lampyrus marginipennis Boheman, 1851: 439
Diaphanes Motschulsky, 1853
Diaphanes luniger Motschulsky, 1853: 45
Lampyrus Geoffroy, 1762
Lampyrus noctiluca Linnaeus, 1767: 643
Lychnobius Geisthardt, 1983
Lampyrus conspicua Gyllenhal in Schönherr, 1817: 20
Microlampyrus Pic, 1956
Microlampyrus basilewski Pic, 1956: 193
Microphotus LeConte, 1866
Microphotus dilatatus LeConte, 1866: 90
Nelsonphotus Cicero, 2006
Nelsonphotus aridus Cicero, 2006: 201
Nyctophila Olivier, 1884
Lamprotomus caucasica Motschulsky, 1854a: 19
Ovalampis Fairmaire, 1898
Ovalampis crispaticollis Fairmaire, 1898: 404
Paraphausis Green, 1949
Paraphausis eximia Green, 1949: 4
Pelania Mulsant, 1860
Lampyrus mauritanica Linnaeus, 1767: 645
Petalacmis Olivier, 1908
Petalacmis praeclarus Olivier, 1908: 187
Prolutacea Cicero, 2006
Lampyrus pulsator Cicero, 1984: 322
Pyrocoelia Gorham, 1880
Lampyrus bicolor Fabricius, 1801: 100

Tribe Photinini LeConte, 1881

Ankonophallus Zaragoza-Caballero and Navarrete-Heredia, 2014
Ankonophallus zuninoi Zaragoza-Caballero and Navarrete-Heredia, 2014: 126
Aorphallus Zaragoza-Caballero and Gutierrez-Carranza, 2018
Aorphallus cibriani Zaragoza-Caballero and Gutierrez-Carranza, 2018: 160
Callopisma Motschulsky, 1853
Lampyrus rufa G.A. Olivier, 1790: 28
Calotrechelum Pic, 1930
Calotrechelum olivieri Pic, 1930: 88
Dadophora Olivier in Wytsman, 1907
Dadophora hyalina Olivier in Wytsman, 1907: 27
Dilychnia Motschulsky, 1853
Dilychnia basalis Motschulsky, 1853: 30
Ellychnia Blanchard, 1845
Lampyrus corrusca Linnaeus, 1767: 644
Erythrolychnia Motschulsky, 1853

Erythrolychnia dimidiatipennis Motschulsky, 1853: 29
Heterophotinus Olivier, 1894
Photinus limbipennis du Val, 1857: 86
Jamphotus Barber, 1941
Jamphotus tuberculatus Barber, 1941: 11
Lucidina Gorham, 1883
Lucidina accensa Gorham, 1883: 408
Lucidota Laporte, 1833
Lucidota banoni Laporte, 1833: 137
Lucidotopsis McDermott, 1960
Lucidota cruenticollis Fairmaire, 1889: 38
Luciuranus Silveira, Khattar, and Mermudes in Silveira et al., 2016
Luciuranus josephi Silveira, Khattar, and Mermudes in Silveira et al., 2016: 377
Macrolampis Motschulsky, 1853
Macrolampis longipennis Motschulsky, 1853: 37
Microdiphot Barber, 1941
Microdiphot cavernarum Barber, 1941: 13
Mimophotinus Pic, 1935
Mimophotinus angustatus Pic, 1935: 9
Oliviereus Pic, 1930
Oliviereus flavus Pic, 1930: 88
Phosphaenopterus Schaufuss, 1870
Phosphaenopterus metzneri Schaufuss, 1870: 61
Phosphaenus Laporte, 1833
Lampyrus hemiptera Geoffroy in Fourcroy, 1785: 58
Photinoides McDermott, 1963
Photinoides penai McDermott, 1963: 87
Photinus Laporte, 1833
Lampyrus pallens Fabricius, 1798: 124
Platylampis Motschulsky, 1853
Platylampis latiuscula Motschulsky, 1853: 44
Pseudolychnuris Motschulsky, 1853
Pseudolychnuris vittata Motschulsky, 1853: 32
Pyraetionema Solier in Gay, 1849
Pyraetionema compressicornis Solier in Gay, 1849: 446
Pyropyga Motschulsky, 1852
Lampyrus nigricans Say, 1823: 179
Pyropygodes Zaragoza-Caballero, 2000
Pyropygodes huautlae Zaragoza-Caballero, 2000: 20
Robopus Motschulsky, 1853
Robopus roseicollis Motschulsky, 1853: 42
Rufolychnia Kazantsev, 2006
Callopisma boreconca Leng and Mutchler, 1922: 440
Uanauna Campello-Gonçalves, Souto, Mermudes, and Silveira, 2019
Uanauna angaporan Campello-Gonçalves, Souto, Mermudes, and Silveira, 2019: 67
Ybytyramoan Silveira and Mermudes, 2014
Ybytyramoan praeclarum Silveira and Mermudes, 2014: 328

Tribe Pleotomini Summers, 1875

Calyptocephalus Gray in Griffith and Pidgeon, 1832
Calyptocephalus fasciatus Gray in Griffith and Pidgeon, 1832: 371
Ophoelis Olivier, 1911
Ophoelis impura Olivier, 1911: 48
Phaenolis Gorham, 1880
Phaenolis laciniatus Gorham, 1880: 10
Pleotomodes Green, 1948
Pleotomodes needhami Green, 1948: 65
Pleotomus LeConte, 1861

Pleotomus pallens LeConte, 1866: 88
Roleta McDermott, 1962
Roleta coracina McDermott, 1962: 69

Lampyrinae incertae sedis

Cladodes Solier in Gay, 1849
Cladodes flabellatus Solier in Gay, 1849: 445
Ethra Laporte, 1833
Cladophorus marginatus Gray in Griffith and Pidgeon, 1832: 371
Scissicauda McDermott, 1964
Lucidota disjuncta Olivier, 1896: 1

Lampyridae incertae sedis

Anadrilus Kirsch, 1875
Anadrilus indus Kirsch, 1875: 37
Araucariocladus Silveira and Mermudes, 2017
Araucariocladus hiems Silveira and Mermudes, 2017: 209
Crassitarsus Martin, nom. nov.
Pachytarsus basalis Motschulsky, 1861: 135
Lamprigera Motschulsky, 1853
Lamprigera boyei Motschulsky, 1853: 48
Oculogryphus Jeng, Engel, and Yang, 2007
Oculogryphus fulvus Jeng in Jeng, Engel, and Yang, 2007: 7
Photocetus McDermott, 1961
Photocetus boliviae McDermott, 1961: 174
Pollaclasis Newman, 1838
Pollaclasis ovatus Newman, 1838: 385

Subtribe Vestini McDermott, 1964

Dodacles Olivier, 1885
Dodacles elegans Olivier, 1885: 141
Dryptelytra Laporte, 1833
Dryptelytra cayennensis Laporte, 1833: 139
Ledocas Olivier, 1885
Ledocas parallelus Olivier, 1885: 140
Vesta Laporte, 1833
Vesta chevrolati Laporte, 1833: 133

Supplementary Data

Supplementary data are available at *Insect Systematics and Diversity* online.

Data Availability Statement

Data from this study are available from the Dryad Digital Repository: doi:10.5061/dryad.737c8t8.

Acknowledgments

We thank the members of the Bybee and Whiting labs at Brigham Young University, the Stanger-Hall lab at the University of Georgia, and the Branham lab at the University of Florida for specimen acquisition. We also specifically thank Brian Cassell, Joe Cicero, Raphael DeCock, E. Evans, Lynn Faust, Jim Lloyd, Arwin Provonsha, P. Smith, Fred Vencel, and Ming-Luen Jeng for collecting specimens (Stanger-Hall lab collection). We thank Michelle Kortyna, Kirby Birch, and Sean Holland at the Center for Anchored Phylogenomics for assistance with data collection and analyses. We also thank Jesse Breinholt, Robert Erickson, Anton Suvorov, and Kyle Warren for assistance and advice on analyses, as well as the Fulton Supercomputing Group at BYU where all analyses

were performed. We want to thank Michael Ivie, Oliver Keller, Nathan Lord, Yelena Pacheco, Gareth Powell, and Caroline Storer for general comments on the manuscript. We thank Floyd Shockley (USNM), Azadeh Taghavian (MNHN), Isabelle Zuercher (MNB), and Max Barclay and Michael Geiser (NHM) for access to museums for specimen examination and for imaging of *Phausis reticulata* (USNM) and *Lamprohiza delarouzei* (MNHN). This project was funded via the by the National Science Foundation (DEB-1655981 S.M.B.; DEB-1655908 K.S.H.; and DEB-1655936 M.A.B) and by the Brigham Young University Mentoring Environment Grant (S.M.B.).

References Cited

- Ballal, C. R., S. A. Akbar, K. Yamada, A. A. Wachkoo, and R. Varshney. 2018. Annotated catalogue of the flower bugs from India (Heteroptera: Anthrenidae, Lasiochilidae). *Acta Entomol. Mus. Natl. Pragae*. 58(1): 207–26.
- Ballantyne, L. A. 1968. Revisional studies of Australian and Indomalayan Luciolini (Coleoptera: Lampyridae: Luciolinae). *Univ. Queensl. Papers* 2(6): 105–139.
- Ballantyne, L. A. 1987a. Luciline morphology, taxonomy and behaviour: a reappraisal (Coleoptera: Lampyridae). *Trans. Am. Entomol. Soc.* 113(2): 171–188.
- Ballantyne, L. A. 1987b. Further revisional studies on the firefly genus *Pteroptyx* Olivier (Coleoptera: Lampyridae: Luciolinae: Luciolini). *Trans. Am. Entomol. Soc.* 113(2): 117–170.
- Ballantyne, L. A. and C. L. Lambkin. 2009. Systematics of Indo-Pacific fireflies with a redefinition of Australasian *Atypheila* Olliff, Madagascan *Photurolociola* Pic, and description of seven new genera from the Luciolinae (Coleoptera: Lampyridae). *Zootaxa*. 1997: 1–188.
- Ballantyne, L. A. and C. L. Lambkin. 2013. Systematics and phylogenetics of Indo-Pacific Luciolinae fireflies (Coleoptera: Lampyridae) and the description of new genera. *Zootaxa*. 3653(1): 1–162.
- Ballantyne, L. A., X.-H. Fu, C. Lambkin, M.-L. Jeng, L. Faust, W. M. C. D. Wijekoon, D. Li, and T. Zhu. 2013. Studies on south-east Asian fireflies: *Abscondita*, a new genus with details of life history, flashing patterns and behaviour of *Abscondita chinensis* (L.) and *Abscondita terminalis* (Olivier) (Coleoptera: Lampyridae: Luciolinae). *Zootaxa*. 3721(1): 1–48.
- Ballantyne, L. A., C. L. Lambkin, Y. Boontop, and W. F. A. Jusoh. 2015. Revisional studies on the Luciolinae fireflies of Asia (Coleoptera: Lampyridae): 1. The genus *Pyrophanes* Olivier with two new species. 2. Four new species of *Pteroptyx* Olivier and 3. A new genus *Inflata* Boontop, with redescription of *Luciola indica* (Motsch.) as *Inflata indica* comb. nov. *Zootaxa*. 3959(1): 1–84.
- Ballantyne, L. A., C. L. Lambkin, X. Luan, Y. Boontop, S. Nak-Eiam, S. Pimpasalee, S. Silalom, and A. Thancharoen. 2016. Further studies on south eastern Asian Luciolinae: 1. *Sclerotia* Ballantyne, a new genus of fireflies with back swimming larvae 2. *Triangulata* Pimpasalee, a new genus from Thailand (Coleoptera: Lampyridae). *Zootaxa*. 4170(2): 201–249.
- Ballantyne, L. A., C. L. Lambkin, J.-Z. Ho, W. F. A. Jusoh, B. Nada, S. Nak-Eiam, A. Thancharoen, W. Wattanachaiyingcharoen, and V. Yiu. 2019. The Luciolinae of S.E. Asia and the Australopacific region: a revisionary checklist (Coleoptera: Lampyridae) including description of three new genera and 13 new species. *Zootaxa*. 4687(1): 001–174.
- Barber, H. S. 1941. Species of fireflies in Jamaica (Coleoptera, Lampyridae). *Proc. Rochester Acad. Sci.* 8(1): 1–13.
- Blanchard, C. E. 1853. Description des insectes, pp. 1842–54. In: J. S. C. Dumont-D'Urville. Voyage au Pole sud et dans l'Océanie sur les corvettes l'Astrolabe et la Zélée. Paris, Zoologie, t.4, 73,74, plates V,VI.
- Blanchard, É. 1845. Histoire naturelle des insectes leurs mœurs, leurs, métamorphoses et leur classification ou traité élémentaire d'entomologie. Tome II. F. Savy, Paris, France. 524 pp.
- Blanchard, É. and A. Brullé. 1846. Voyage dans l'Amérique Méridionale (le Brésil, la République orientale de l'Uruguay, la République Argentine, la Patagone, la République du Chili, la République de Bolivie, la République du Pérou), exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833, par Alcide D'Orbigny. Tome Sixième. 2.^e partie: insectes. P. Bertrand, Paris & Ve. Lerrault, Strasbourg, France. 222 pp.

- Bocakova, M. and L. Bocak. 2016. A new genus of ototretine firefly endemic to Indian subcontinent (Coleoptera: Lampyridae). *Ann. Zool.* 66(3): 371–380.
- Bocakova, M. and K. Janisova. 2010. A new genus and species of ototretine firefly from Borneo (Coleoptera: Lampyridae). *Zootaxa*. 2347: 59–63.
- Bocakova, M., L. Bocak, T. Hunt, M. Teräväinen, and A. P. Vogler. 2007. Molecular phylogenetics of Elateriformia (Coleoptera): evolution of bioluminescence and neoteny. *Cladistics* 23: 477–496.
- Boheman, C. H. 1851. Fam. XI. Lampyrides. *Insecta Caffrariae annis 1838–1845 a J. A. Wahlberg collecta, amici auxilio suffultus, descripsit Carolus H. Boheman* 1(2): 439–451.
- Branham, M. A. and J. W. Wenzel. 2001. The evolution of bioluminescence in cantharoids (Coleoptera: Elateroidea). *Fla. Entomol.* 84(4): 565–586.
- Branham, M. A. and J. W. Wenzel. 2003. The origin of photic behavior and the evolution of sexual communication in fireflies (Coleoptera: Lampyridae). *Cladistics*. 19: 1–22.
- Branham, M. A. 2010. Lampyridae Latreille, 1817, pp. 141–149. *In*: R. A. B. Leschen, R. G. Beutel, and J. F. Lawrence (Eds.), *Coleoptera, Beetles. Volume 2: Morphology and Systematics (Elateroidea, Bostrichiformia, Cucujiformia partim)*. Walter de Gruyter, Berlin, Germany. 786 pp.
- Breinholz, J. W., C. Earl, A. R. Lemmon, E. M. Lemmon, L. Xiao, and A. Y. Kawahara. 2017. Resolving relationships among the megadiverse butterflies and moths with a novel pipeline for Anchored Phylogenomics. *Syst. Biol.* 67(1): 78–93.
- Buck, J. B. 1937. Studies on the firefly. II. The signal system and color vision in *Photinus pyralis*. *Physiol. Zool.* 10(4): 412–419.
- Buck, J. B. 1947. Studies on the firefly, IV: Ten new lampyrids from Jamaica. *Proc. U. S. Nat. Mus.* 97(3210): 59–79.
- Campello-Gonçalves, L., P. M. Souto, J. R. M. Mermudes, and L. F. L. Silveira. 2019. *Uanauna* gen. nov., a new genus of fireflies endemic to the Brazilian Atlantic forest (Coleoptera: Lampyridae), with key to Brazilian genera of Lucidolina. *Zootaxa*. 4585 (1): 59–72.
- Chernomor, O., A. von Haeseler, and B. Q. Minh. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Syst. Biol.* 65(6): 997–1008.
- Cicero, J. M. 1984. A new species of *Lampyris* from southern Arizona and Mexico (Coleoptera, Lampyridae). *Coleopt. Bull.* 38(4): 322–324.
- Cicero, J. M. 2006. Two new genera and one new species of firefly (Coleoptera: Lampyridae: Lampyrinae: Lampyrini), with notes on their biology. *Pan-Pac. Entomol.* 82(2): 200–207.
- Costa, A. 1875. M. A. Costa prend la parole et fait la communication suivante. *Bull. Séances Bull. Bibliograph. Soc. Entomol. France* 1875: clxix–clxx.
- Crowson, R. A. 1972. A review of the classification of Cantharoidea (Coleoptera), with the definition of two new families, Cneoglossidae and Omethidae. *Rev. Univ. Madrid* 21(82): 35–77.
- Dejean, P. F. M. A. 1833. Catalogue des Coléoptères de la Collection de M. le Comte Dejean. Livraisons 1–2. Méquignon-marvis Père et Fils, Paris, France. 1–176.
- Deyrolle, H. and L. Fairmaire. 1878. Descriptions de Coléoptères recueillis par M. l'abbé David dans la Chine centrale. *Ann. Soc. Entomol. Fr.* 8: 87–140.
- Fabricius, J. C. 1775. *Systema Entomologiae, sistens insectorum classes, ordines genera, species, adiectus synonymus, locis, descriptionibus, observationibus*. Officina Libraria Kortii, Flensburgi (=Flensburg) and Lipsiae (=Leipzig), Germany. 832 pp.
- Fabricius, J. C. 1798. *Supplementum Entomologiae Systematicae*. Proft et Storch, Hafniae (=Kopenhagen), Denmark. 572 pp.
- Fabricius, J. C. 1801. *Systema Eleutheratorum Secundum Ordines, Genera, Species Adiecticis Synonymis, Locis, Observationibus, Descriptionibus*. Tomus II. Impensis Bibliopolii Academici Novi, Kiliae (=Kiel), Germany. 506 pp.
- Fairmaire, L. 1898. Matériaux pour la faune Coléoptérique de la region Malgache. *Ann. Soc. Entomol. Belg.* 42: 390–439.
- Fender, K. M. 1966. The genus *Phausis* in America North of Mexico (Coleoptera-Lampyridae). *Northwest Sci.* 40(3): 83–95.
- Ferreira, V. S., O. Keller, M. A. Branham, and M. A. Ivie. 2019. Molecular data support the placement of the enigmatic *Cheguevaria* as a subfamily of Lampyridae (Insecta: Coleoptera). *Zool. J. Linn. Soc.* XX: 1–6.
- Fieber, F. X. 1860. Exegesen in Hemipteren. *Wiener Entomol. Monatschr.* 4: 257–272.
- Fu, X.-H., L. A. Ballantyne, and C. L. Lambkin. 2010. *Aquatica* gen. nov. from mainland China with a description of *Aquatica wuhana* sp. nov. (Coleoptera: Lampyridae: Luciolinae). *Zootaxa*. 2530: 1–18.
- Fu, X.-H., L. A. Ballantyne, and C. L. Lambkin. 2012. *Emeia* gen. nov., a new genus of Luciolinae fireflies from China (Coleoptera: Lampyridae) with an unusual trilobite-like larva, and a redescription of the genus *Curtos* Motschulsky. *Zootaxa*. 3403: 1–53.
- Geisthardt, M. 1983. Zur Kenntnis der Gattung *Nyctophila* Olivier 1884 und Aufstellung einer neuen Gattung *Lychnobius* gen. nov. (Coleoptera: Lampyridae: Lampyrinae). *Entomol. Blätter Biol. Syst. Käfer* 79(1): 37–60.
- Geisthardt, M. 2004. New and known fireflies from Mount Emei (China) (Coleoptera: Lampyridae). *Mitteilungen des Internationalen Entomologischen Vereins e. V., Frankfurt a. M.* 29(1–2): 1–10.
- Geisthardt, M. 2007. *Afrodiaphanes* a new genus from Africa (Coleoptera: Lampyridae: Lampyrini). *Mitteilungen des Internationalen Entomologischen Vereins e. V., Frankfurt a. M.* 32(1–2): 45–50.
- Geoffroy, M. 1762. *Histoire abrégée des Insectes, qui se trouvent aux environs de Paris; dans laquelle ces Animaux sont rangés suivant un ordre méthodique*. Tome 1. Chez Durand, Paris, France. 523 pp.
- Geoffroy, M. 1785. *Lampyris*, pp. 58–59. *In*: A. F. Fourcroy, (Ed.), *Entomologia Parisiensis; sive, catalogus insectorum quae in agro Parisiensi reperiuntur*. Paris, France. 231 pp.
- Gorham, H. S. 1880a. I. Materials for a revision of the Lampyridae. *Trans. Entomol. Soc. Lond.* 1880: 1–36.
- Gorham, H. S. 1880b. VIII. Materials for a revision of the Lampyridae. *Trans. Entomol. Soc. Lond.* 1880: 83–112.
- Gorham, H. S. 1883. Revision of the genera and species of Malacoderm Coleoptera of the Japanese fauna. *Trans. Entomol. Soc. Lond.* 1883: 393–411.
- Gray, G. 1832. Supplement on the Serricornes. 356–383, pp. 570 pp. *In*: E. Griffith, and E. Pidgeon, (Eds.), *The animal kingdom*. Vol. 14. Gilbert and Rivington, London, United Kingdom.
- Green, J. W. 1948. Two new species of Lampyridae from southern Florida, with a generic revision of the Nearctic fauna (Coleoptera). *Trans. Am. Entomol. Soc.* 74(2): 61–73.
- Green, J. W. 1949. A new genus and new species of American Lampyrini, and other notes (Coleoptera: Lampyridae, Cantharidae). *Trans. Am. Entomol. Soc.* 75(1): 1–6.
- Green, J. W. 1959. Revision of the species of *Microphotus*, with an emendation of the Lampyrini (Lampyridae). *Coleopt. Bull.* 13(3): 80–96.
- Gyllenhal, L. 1817. *Lampyris*, 20–25 *In*: C. J. Schönherr, (Ed.) *Appendix ad C. J. Schönherr Synonymiam Insectorum sistens Descriptiones Novarum Specierum*. Tom I Part 3. Scaris in Officina Lewerentziana, Stockholm, Sweden. 266 pp.
- Haddad, S., S. Shin, A. R. Lemmon, E. Moriarty Lemmon, P. Svacha, B. Farrell, A. Sliński, D. Windsor, and D. D. McKenna. 2018. Anchored hybrid enrichment provides new insights into the phylogeny and evolution of longhorned beetles (Cerambycidae). *Syst. Ent.* 43:68–89.
- Hamilton, C. A., A. R. Lemmon, E. Moriarty Lemmon, and J. E. Bond. 2016. Expanding anchored hybrid enrichment to resolve both deep and shallow relationships within the spider tree of life. *BMC Evol. Biol.* 16(1): 212.
- Hoang, D. T., O. Chernomor, A. von Haeseler, B. Q. Minh, and L. S. Vinh. 2018. UFBoot2: improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* 35(2): 518–522.
- Illiger, J. K. W. 1807. Vorschlag zur Aufnahme im Fabricischen Systeme fehlender Käfergattungen. *Magazin Insek.* 6: 318–349.
- Jacquelin du Val, P. N. C. 1857. Coléoptères. 1–328, *In*: R. Sagra de la, (Ed.), *Historia Fisica, Politica y natural de la Isla de Cuba. Segunda parte: Historia natural*. Tomo VII: Crustaceos, Aragnides é Insectos. Arthus Bertrand, Paris, France. lxxxvii + 868 pp.
- Janisova, K. and M. Bocakova. 2013. Revision of the subfamily Ototretinae (Coleoptera: Lampyridae). *Zool. Anz.* 252: 1–19.
- Jeng, M.-L. 2008. Comprehensive phylogenetics, systematics, and evolution of neoteny of Lampyridae (Insecta: Coleoptera). Ph.D. dissertation. University of Kansas, Lawrence, KS. 388 pp.

- Jeng, M.-L., P.-S. Yang, and M. Satô. 1998. The genus *Cyphonocerus* (Coleoptera, Lampyridae) from Taiwan and Japan, with notes on the subfamily Cyphonocerinae. *Elytra*, Tokyo. 26(2): 379–398.
- Jeng, M.-L., J. Lai, P.-S. Yang, and M. Satô. 2000. Notes on the taxonomy of *Lamprigera yunnana* (Fairmaire) and the genus *Lamprigera* Motschulsky (Coleoptera, Lampyridae). *Jpn. J. Syst. Entomol.* 6(2): 313–319.
- Jeng, M.-L., P.-S. Yang, and J. Lai. 2003. Notes on the genus *Luciola* (Coleoptera, Lampyridae, Luciolinae) of Taiwan. *Spec. Bull. Japan. Soc. Coleop.* Tokyo 6: 247–262.
- Jeng, M.-L., P.-S. Yang, and M. Satô. 2006. Synopsis of *Cyphonocerus* (Coleoptera: Lampyridae) with the description of four new species and a key to the genus. *Zool. Stud.* 45(2): 157–167.
- Jeng, M.-L., M. S. Engel, and P.-S. Yang. 2007. *Oculogryphus*, a remarkable new genus of fireflies from Asia (Coleoptera: Lampyridae). *Am. Mus. Novit.* 3600: 1–19.
- Kalyaanamoorthy, S., B. Q. Minh, T. K. Wong, A. von Haeseler, and L. S. Jermin. 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* 14(6): 587.
- Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30(4): 772–780.
- Kazantsev, S. V. 2006. New firefly taxa from Hispaniola and Puerto Rico (Coleoptera: Lampyridae), with notes on biogeography. *Russ. Entomol. J.* 15(4): 367–392.
- Kazantsev, S. V. 2010. Fireflies of Russia and adjacent territories (Coleoptera: Lampyridae). *Russ. Entomol. J.* 19(3): 187–208.
- Kiesenwetter, H. von. 1879. Coleoptera Japoniae collecta a Domino Lewis et aliis. *Deut. Entomol. Z.* 23(2): 305–320.
- Kirsch, T. 1875. Neue Käfer aus Malacca. *Mittheilungen aus dem Königlichen Zool. Mus. Dresden.* 1: 25–58.
- Kirby, W. 1818. A century of insects, including several new genera described from his cabinet. *Transactions of the Linnean Society of London* 12(2): 375–453.
- Kundrata, R., M. Bocakova, and L. Bocak. 2014. The comprehensive phylogeny of the superfamily Elateroidea (Coleoptera: Elateriformia). *Mol. Phylogenet. Evol.* 76: 162–171.
- Lacordaire, M. T. 1857. Tribu II. Lampyrides. *Histoire Naturelle des Insectes. Genres des Coléoptères ou exposé Méthodique et Critique de tous les Genres Proposés jusqu'ici dans cet Ordre d'Insectes* 4: 304–341.
- Laporte, [=Castelnau] F. L. N. C. de. 1833. Essai d'une révision du genre *Lampyre*. *Ann. Soc. Entomol. Fr.* 2: 122–153.
- Lea, A. M. 1915. Descriptions of new species of Australian Coleoptera. Part XI. *Proc. Linn. Soc. N. S. W.* 40: 490–521.
- Lea, A. M. 1921. On Australian Coleoptera of the family Malacodermidae. *Trans. R. Soc. S. Aust.* 45: 50–135.
- LeConte, J. L. 1852. Synopsis of the Lampyrides of temperate North America [1851]. *Proc. Acad. Nat. Sci. Phila.* 5: 331–347.
- LeConte, J. L. 1859. Catalogue of the Coleoptera of Fort Tejon, California. *Proc. Acad. Nat. Sci. Phila.* 11: 69–90.
- LeConte, J. L. 1861. Classification of the Coleoptera of North America. Part 1. *Smithson. Misc. Collect.* 3: xxv + 1–208.
- LeConte, J. L. 1866. Descriptions of new species. *Smithson. Misc. Collect.* 167: 1–177.
- LeConte, J. L. 1881. Synopsis of the Lampyridae of the United States. *Trans. Am. Entomol. Soc.* 9: 15–72.
- Lemmon, A. R., J. M. Brown, K. Stanger-Hall, and E. M. Lemmon. 2009. The effect of ambiguous data on phylogenetic estimates obtained by maximum likelihood and Bayesian inference. *Syst. Biol.* 58(1): 130–145.
- Lemmon, A. R., S. Emme, and E. M. Lemmon. 2012. Anchored hybrid enrichment for massively high-throughput phylogenetics. *Syst. Biol.* 61: 721–744.
- Leng, C. W. and A. J. Mutchler. 1922. The Lycidae, Lampyridae, and Cantharidae (Telephoridae) of the West Indies. *Bull. Am. Mus. Nat. Hist.* 4: 413–499.
- Li, X.-Y., S. Yang, and X.-C. Liang. 2006. Phylogenetic relationship of the firefly, *Diaphanes pectinealis* (Insecta, Coleoptera, Lampyridae) based on DNA sequence and gene structure of Luciferase. *Zool. Res.* 27(4): 367–374.
- Linnaeus, C. 1767. *Systema Naturae*. Tome I, Pars II. Editio Duodecima Reformata. Impensis Direct. Laur. Salvii, Stockholm, Sweden. 794 pp.
- Lloyd, J. E. 1971. Bioluminescent communication in insects. *Annu. Rev. Entomol.* 16: 97–122.
- Martin, G. J., M. A. Branham, M. F. Whiting, and S. M. Bybee. 2017. Total evidence phylogeny and the evolution of adult bioluminescence in fireflies (Coleoptera: Lampyridae). *Mol. Phylogenet. Evol.* 107: 564–575.
- Maulik, S. 1921. New Indian drilid beetles. *Proc. Zool. Soc. Lond.* 1921(3): 579–586.
- McDermott, F. A. 1960. *Lucidotopsis* genus novum. (Coleoptera: Lampyridae). *Coleopt. Bull.* 14(1): 18.
- McDermott, F. A. 1961. A new genus and species of firefly: *Photocus boliviae* (Coleoptera: Lampyridae). *Entomol. News* 72(7): 174–176.
- McDermott, F. A. 1962. *Roleta coracina* gen. nov., sp. nov. (Coleoptera: Lampyridae, Pleotomini). *Entomol. News* 73(3): 69–73.
- McDermott, F. A. 1963. *Photinoides*, a new genus, with three new species (Coleoptera: Lampyridae). *Coleopt. Bull.* 17(3): 84–89.
- McDermott, F. A. 1964. The taxonomy of the Lampyridae (Coleoptera). *Trans. Am. Entomol. Soc.* 90(1): 1–72.
- McDermott, F. A. 1966. Lampyridae, pp. 1–149. In: Steel, W. O. (Ed.), *Coleopterorum catalogus supplementa, pars 9*. Uitgeverij Dr. W. Junk, Gravenhage, Netherlands.
- Melsheimer, F. E. 1846. Descriptions of new species of Coleoptera of the United States. *Proc. Acad. Nat. Sci. Phila.* 2(12): 302–318.
- Mikšić, R. 1969. The composition and the distribution of the fauna of Lampyridae in the Palaearctic region. In *Proceedings of the Third Kongres Biologov Jugoslavije*: 182.
- Motschulsky, V. 1845. Demarques sur la collection de Coléoptères Russes. *Bull. Soc. Nat. Moscou.* 18(1): 3–127.
- Motschulsky, V. 1853. Lampyrides. *Etud. Entomol.* 1: 25–58.
- Motschulsky, V. 1854a. Lampyrides. *Etud. Entomol.* 3: 15–26.
- Motschulsky, V. 1854b. Lampyrides. *Etud. Entomol.* 3: 47–62.
- Motschulsky, V. 1861. Essai d'un Catalogue des Insectes de l'île Ceylan. *Bull. Soc. Nat. Moscou.* 34(1): 95–155.
- Mulsant, E. 1860. Observations sur les Lampyrides. *Opus. Entomol.* 11: 113–133.
- Mulsant, E. 1862. *Histoire Naturelle des Coléoptères de France*. Mollipennes. Magnin, Blanchard et C^{ie}, successeurs de Louis Janet, Paris, France. 440 pp.
- Nakane, T. 1991. Lampyrid insects of the world, pp. 3–11. In: *The Association of Nature Restoration of Japan*. (Ed.), The reconstruction of firefly environments. *Reconquista SP. No. 1*. Siedeku, Tokyo, Japan. 136 pp.
- Newman, E. 1938. Entomological notes. *Entomol. Mag.* 6: 372–402.
- Nguyen, L. T., H. A. Schmidt, A. von Haeseler, and B. Q. Minh. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* 32(1): 268–274.
- Ohba, N. 1983. Studies on the communication system of Japanese fireflies. *Sci. Rep. Yokosuka City Mus.* 30: 1–62.
- Olivier, G. A. 1790. Lampyre. *Lampyris*, pp. 476–190. In: G. A. Olivier (Ed.), *Encyclopédie Méthodique. Histoire Naturelle des Insectes*. Tome Septième. Chez Panckoucke, Paris, France. 827 pp.
- Olivier, E. 1883. Lampyrides nouveaux ou peu connus. *Deuxième Mémoire. Rev. Entomol.* 2: 326–333.
- Olivier, E. 1884. Essai d'une révision des espèces européennes & circuméditerranéennes de la famille des Lampyrides. *L'Abeille* 22: 1–54.
- Olivier, E. 1885a. Catalogue des Lampyrides faisant partie des collections du Musée civique de Gênes. *Ann. Mus. Civ. Stor. Nat. Genova Serie 2* 22: 333–374.
- Olivier, E. 1885b. Études sur les Lampyrides I. *Ann. Soc. Entomol. Fr.* 5: 125–154.
- Olivier, E. 1894. Descriptions d'espèces nouvelles de Lampyrides. *Ann. Soc. Entomol. Fr.* 63: 23–24.
- Olivier, E. 1896. Descriptions de nouvelles espèces de Lampyrides du Musée de Tring. *Novit. Zool.* 3(1): 1–3.
- Olivier, E. 1902. Catalogue des espèces de *Luciola* et genres voisins décrits jusqu'à ce jour. *Rev. Sci. Bourbon. France.* 15: 69–88.
- Olivier, E. 1907. 53me fascicule. Coleoptera. Fam. Lampyridae, 1–74. In: Wytman P. A. (Ed.), *Genera Insectorum*. Vol. VIII. Verteneuil & Desmet, Bruxelles.
- Olivier, E. 1908. Description d'un nouveau genre et d'une nouvelle espèce de Lampyrides (Col.). *Bull. Soc. Entomol. Fr.* 1908(10): 186–187.

- Olivier, E. 1910a. Lampyridae. 1–68. In: S. Schenkling, (Ed.), *Coleopterorum catalogus*, pars 9. W. Junk, Berlin, Germany. 68 pp.
- Olivier, E. 1910b. Description de deux Lucioles Nouvelles de la Nouvelle-Guinée [Col. Lampyridae]. *Bull. Soc. Entomol. Fr.* 1910: 343–345.
- Olivier, E. 1911a. Révision des Lampyrides. *Rev. Sci. Bourbon.* 24(3): 63–85.
- Olivier, E. 1911b. Lampyrides rapportés de Formose par M. Hans Sauter. *Ann. Mus. Civ. Stor. Nat. Genova* 45: 145–148.
- Olivier, E. 1911c. Le genre *Atyphella* Oll. et descriptions d'espèces Nouvelles [Col. Lampyridae]. *Bull. Soc. Entomol. Fr.* 1911: 171–175.
- Olliff, A. S. 1890. New species of Lampyridae, including a notice of the Mt. Wilson fire-fly. *P. Linn. Soc. N. S. W. (Second Series)* 4: 643–653; 1297.
- Pascoe, F. P. 1887. Notes on Coleoptera, with descriptions of new genera and species. Part IV. *Ann. Magazine Nat. Hist.* 20(5th series) 8–20.
- Pic, M. 1911a. Quatre nouveaux genres de Malacodermes. *L'Échange* 27(324): 186–188.
- Pic, M. 1911b. Coléoptères exotiques nouveaux ou peu connus. *L'Échange* 27(320): 155–159.
- Pic, M. 1915. Nouvelles espèces de diverses familles. *Mélang. Exot.-Ent.* 15: 2–24.
- Pic, M. 1918. Courtes descriptions diverse. *Mélang. Exot.-Ent.* 27: 1–24.
- Pic, M. 1921. Nouveautés diverses. *Mélang. Exot.-Ent.* 33: 1–32.
- Pic, M. 1927. Malacodermes exotiques. *L'Échange* 43(hors-texte 430): 49–52.
- Pic, M. 1929. Nouveautés diverses. *Mélang. Exot.-Ent.* 53: 1–36.
- Pic, M. 1930. Malacodermes exotiques. *L'Échange* 46(hors-texte 441): 85–88.
- Pic, M. 1931. Nouveautés diverses. *Mélang. Exot.-Ent.* 57: 1–36.
- Pic, M. 1933. Nouveautés diverses. *Mélang. Exot.-Ent.* 62: 1–36.
- Pic, M. 1935. Nouveautés diverses. *Mélang. Exot.-Ent.* 66: 1–36.
- Pic, M. 1937. Malacodermes exotiques. *L'Échange* 53(hors-texte 468): 137–140.
- Pic, M. 1956. Cantharidae et Lampyridae nouveaux du Congo Belge (Coléoptères Malacodermes). *Rev. Zool. Afric.* 53(1–2): 179–194.
- Prum, R. O., J. S. Berv, A. Dornburg, D. J. Field, J. P. Townsend, E. C. Lemmon, and A. R. Lemmon. 2015. A fully resolved, comprehensive phylogeny of birds (Aves) using targeted next generation DNA sequencing. *Nature* 526:569–573.
- Rafinesque, C. S. 1815. *Analyse de la nature ou tableau de l'univers et des corps organisés*. C. S. Rafinesque, Palermo, 224 pp.
- Rokyta, D. R., A. R. Lemmon, M. J. Margres, and K. Arnow. 2012. The venom-gland transcriptome of the eastern diamondback rattlesnake (*Crotalus adamanteus*). *BMC Genomics* 13:312.
- Say, T. 1825. Descriptions of new species of coleopterous insects inhabiting the United States. *J. Acad. Nat. Sci. Phila.* 5: 160–204.
- Sayyari, E. and S. Mirarab. 2016. Fast coalescent-based computation of local branch support from quartet frequencies. *Mol. Biol. Evol.* 33(7): 1654–1668.
- Schaufuss, L. W. 1870. *Phosphaenopterus* nov. gen. malacodermata. *Nunq. Ot.* 1: 60–61.
- Silveira, L. F. L., and J. R. M. Mermudes. 2013. *Memoan ciceroi* gen. et sp. nov., a remarkable new firefly genus and species from the Atlantic Rainforest (Coleoptera: Lampyridae). *Zootaxa*. 3640(1): 79–87.
- Silveira, L. F. L., and J. R. M. Mermudes. 2014. *Ybytyramoan*, a new genus of fireflies (Coleoptera: Lampyridae, Lampyrinae, Photinini) endemic to the Brazilian Atlantic Rainforest, with description of three new species. *Zootaxa*. 3835(3): 325–337.
- Silveira, L. F. L., and J. R. M. Mermudes. 2017. A new tropical montane firefly genus and species, active during winter and endemic to the Southeastern Atlantic Rainforest (Coleoptera: Lampyridae). *Zootaxa*. 4221(2): 205–214.
- Silveira, L. F. L., G. Khattar, P. Souto, J. R. M. Mermudes, D. M. Takiya, and R. F. Monteiro. 2016. Integrative taxonomy of new firefly taxa from the Atlantic Rainforest. *Syst. Biodivers.* 14(4): 371–384.
- Silveira, L. F. L., J. R. M. Mermudes, and M. Bocáková. 2016. Systematic review of the firefly genus *Scissicauda* (Coleoptera, Lampyridae, Amydetinae) from Brazil. *ZooKeys*. 558: 55–75.
- Solier, A. J. J. 1849. Coleopteres, 105–508. In: C. Gay, (Ed.), *Historia fisica y politica de Chile segun documentos adquiridos en esta republica durante doce aos de residencia en ella y publicada bajo los auspicios del supremo gobierno*. Zoologia. Tomo cuarto. Museo de Historia Natural de Santiago, Paris, France and Santiago, Chile. 511 pp.
- Stanger-Hall, K. F. and J. E. Lloyd. 2015. Flash signal evolution in *Photinus* fireflies: character displacement and signal exploitation in a visual communication system. *Evolution*. 69(3): 666–682.
- Stanger-Hall, K. F., J. E. Lloyd, and D. M. Hillis. 2007. Phylogeny of North American fireflies (Coleoptera: Lampyridae): implications for the evolution of light signals. *Mol. Phylogenet. Evol.* 45: 33–49.
- Stanger-Hall, K. F., S. E. Sander Lower, L. Lindberg, A. Hopkins, J. Pallansch, and D. W. Hall. 2018. The evolution of sexual signal modes and associated sensor morphology in fireflies (Lampyridae, Coleoptera). *Proc. R. Soc. B.* 285: 20172384.
- Sturm, J. 1826. *Catalog meiner Insecten-Sammlung. Erster Theil. Käfer*. Self-published, Nürnberg, Germany. viii + 207 pp.
- Summers, S. V. 1875. Catalogue of the Coleoptera from the region of Pontchartrain, La. *Bull. Buffalo Soc. Nat. Sci.* 2(1874–1875): 78–99.
- Thancharoen, A., L. A. Ballantyne, M. A. Branham, and M.-L. Jeng. 2007. Description of *Luciola aquatilis* sp. nov., a new aquatic firefly (Coleoptera: Lampyridae: Luciolinae) from Thailand. *Zootaxa*. 1611: 55–62.
- Van Dyke, E. C. 1939. New species and subspecies of West American Coleoptera. *Pan-Pac. Entomol.* 15(1): 15–20.
- Walker, F. 1858. Characters of some apparently undescribed Ceylon insects. *The Annals and Magazine of Natural History, including Zoology, Botany, and Geology Third Series*. 2: 280–286.
- Wang, K., W. Hong, H.-W. Jiao, and H.-B. Zhao. 2017. Transcriptome sequencing and phylogenetic analysis of four species of luminescent beetles. *Sci. Rep.-UK*. 7(1814): 1–11.
- Westwood, J. O. 1853. Descriptions of some new species of Coleoptera from China and Ceylon. *Trans. Entomol. Soc. Lond. New Ser.* 2: 232–241.
- Wiens, J. J., and M. C. Morrill. 2011. Missing data in phylogenetic analysis: reconciling results from simulations and empirical data. *Syst. Biol.* 60(5): 719–731.
- Wittmer, W. 1938. 4. Beitrag zur Kenntnis der indo-malayischen Malacodermata (Coleoptera). *Mitt. Schweiz. Entomol. Ges.* 17: 209–210.
- Wittmer, W. 1939. 5. Beitrag zur Kenntnis der indo-malayischen Malacodermata (Col.). *Treubia*. 17: 21–32.
- Wittmer, W. 1940. Neue Malacodermata aus der Sammlung des British Museum. – 9. Beitrag zur Kenntnis der indo-malayischen Malacodermata. *Ann. Mag. Nat. Hist.* 6: 274–277.
- Wittmer, W. 1979. Zur Kenntnis der Lampyridae der orientalischen Region (Coleoptera). *Entomol. Arb. Mus. G. Frey Tutzing bei Münch.* 28: 83–92.
- Zaragoza-Caballero, S. 1995. La familia Lampyridae (Coleoptera) en la estación de biología tropical “Los Tuxtlas”, Veracruz, México. *Publ. Espec. Inst. Biol.* 14: 1–93.
- Zaragoza-Caballero, S. 2000. Cantharoidea (Coleoptera) de México. VI. Un nuevo género y una nueva especie de Lampyridae del Estado de Morelos, México. *Dugesiana*. 7(1): 19–22.
- Zaragoza-Caballero, S. 2013. *Paracratomorphus* nuevo género (Coleoptera: Lampyridae: Cratomorphini) de México. *Dugesiana*. 20(2): 145–148.
- Zaragoza-Caballero, S. and I. G. Gutiérrez-Carranza. 2018. *Aorphallus cibriani* gen. nov., sp. nov., y otros Photinini de México (Coleoptera: Lampyridae). *Dugesiana*. 25(2): 159–166.
- Zaragoza-Caballero, S. and J. L. Navarrete-Heredia. 2014. Descripción de cuatro especies de *Ankonophallus* gen. nov. (Coleoptera: Lampyridae: Photinini). *Dugesiana*. 21(2): 125–130.
- Zhang, C., M. Rabiee, E. Sayyari, and S. Mirarab. 2018. ASTRAL-III: polynomial time species tree reconstruction from partially resolved gene trees. *BMC Bioinformatics*. 19(6): 153.