

Parallel Social Information Processing Circuits Are Differentially Impacted in Autism

Highlights

- Cellular and circuit features distinguish magnocellular and parvocellular OT neurons
- Peer-peer, but not filial, social attachments are disrupted in *Fmr1*-KO ASD model mice
- *Fmr1* is required in parvocellular OT neurons for peer-peer social reward learning
- ASD risk and FMRP target genes are enriched in parvocellular OT neurons

Authors

Eastman M. Lewis,
Genevieve L. Stein-O'Brien,
Alejandra V. Patino, ..., Connie Jiang,
Loyal A. Goff, Gül Dölen

Correspondence

loyalgoff@jhmi.edu (L.A.G.),
gul@jhu.edu (G.D.)

In Brief

Analyzing magnocellular and parvocellular oxytocin neurons in detail, Lewis et al. provide evidence that pathocllisis of parvocellular neurons plays an important role in the pathogenesis of social impairments in fragile X and autism.



Article

Parallel Social Information Processing Circuits Are Differentially Impacted in Autism

Eastman M. Lewis,^{1,2,3,4,9} Genevieve L. Stein-O'Brien,^{1,3,5,6,9} Alejandra V. Patino,^{1,2,3,4,6,9} Romain Nardou,^{1,2,3,4} Cooper D. Grossman,^{1,2,3} Matthew Brown,¹ Bidli Bangamwabo,¹ Ndeye Ndiaye,¹ Daniel Giovinnazzo,¹ Ian Dardani,⁷ Connie Jiang,⁸ Loyal A. Goff,^{1,3,6,*} and Gül Dölen^{1,2,3,4,10,*}

¹The Solomon H. Snyder Department of Neuroscience, Johns Hopkins University, School of Medicine, Baltimore, MD 21205, USA

²The Brain Science Institute, Johns Hopkins University, School of Medicine, Baltimore, MD 21205, USA

³The Kavli Neuroscience Discovery Institute, Johns Hopkins University, School of Medicine, Baltimore, MD 21205, USA

⁴The Wendy Klag Institute for Autism and Developmental Disabilities, Johns Hopkins University, School of Medicine, Baltimore, MD 21205, USA

⁵Department of Oncology, Division of Biostatistics and Bioinformatics, Sidney Kimmel Comprehensive Cancer Center, Johns Hopkins University, School of Medicine, Baltimore, MD 21205

⁶McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University, School of Medicine, Baltimore, MD 21205, USA

⁷Department of Bioengineering, University of Pennsylvania, Philadelphia, PA 19104, USA

⁸Cell and Molecular Biology Group, University of Pennsylvania, Perelman School of Medicine, Philadelphia, PA 19104, USA

⁹These authors contributed equally

¹⁰Lead Contact

*Correspondence: loyalgoff@jhmi.edu (L.A.G.), gul@jhu.edu (G.D.)

<https://doi.org/10.1016/j.neuron.2020.10.002>

SUMMARY

Parallel processing circuits are thought to dramatically expand the network capabilities of the nervous system. Magnocellular and parvocellular oxytocin neurons have been proposed to subserve two parallel streams of social information processing, which allow a single molecule to encode a diverse array of ethologically distinct behaviors. Here we provide the first comprehensive characterization of magnocellular and parvocellular oxytocin neurons in male mice, validated across anatomical, projection target, electrophysiological, and transcriptional criteria. We next use novel multiple feature selection tools in *Fmr1*-KO mice to provide direct evidence that normal functioning of the parvocellular but not magnocellular oxytocin pathway is required for autism-relevant social reward behavior. Finally, we demonstrate that autism risk genes are enriched in parvocellular compared with magnocellular oxytocin neurons. Taken together, these results provide the first evidence that oxytocin-pathway-specific pathogenic mechanisms account for social impairments across a broad range of autism etiologies.

INTRODUCTION

Parallel processing pathways have been identified in both sensory (Dijkerman and de Haan, 2007; Haberly, 2001; Nassi and Callaway, 2009; Recanzone and Cohen, 2010; Roper, 2009; Zwergal et al., 2009) and striatal (Cui et al., 2013; Kravitz et al., 2012; Lobo and Nestler, 2011) circuits and are thought to dramatically expand the network capabilities of the nervous system. Interestingly, oxytocin (OT) neurons are composed of magnocellular and parvocellular subtypes (Gurdjian, 1927), raising the possibility that this organizational principle can be extended to social information processing. Indeed, magnocellular and parvocellular OT neuronal subtypes can be differentiated by a number of characteristic features (van den Pol, 2012), which suggest that they are specialized to subserve distinct behaviors. For example, because magnocellular OT neurons are specialized to release large quantities of OT in both the central and peripheral nervous system, this adaptation may enable coordination

of central and systemic responses during lactation and parturition (Ludwig and Leng, 2006). In contrast, parvocellular OT neurons release smaller quantities of OT that are restricted to the central brain, an adaptation that may subserve the measured reward associated with social cognition and peer-peer attachments (Dölen and Malenka, 2014). At this time this view is largely supported by indirect evidence from comparative studies across species (Dölen, 2015). Nevertheless, the existence of parallel circuits for social information processing will have significant implications for understanding the pathogenesis of autism spectrum disorder (ASD), a neurodevelopmental disease characterized by domain-specific impairments in social interactions (Vivanti and Nuske, 2017).

Selective vulnerability of neuronal subtypes to brain injury is called “pathocllisis” and has been proposed to explain how despite exposure to a common insult, disease pathogenesis may disproportionately reflect disrupted function of a subset of brain cells (Klatzo, 2003). For example, in Huntington’s disease,



even though the genetic lesion exists throughout the brain, characteristic motor impairments and striatal degeneration have been proposed to be the consequence of selective affinity of mutant Huntingtin (*HTT*) for Rhes, the Ras homolog enriched in the striatum (Paul et al., 2014). The etiology of ASD is also primarily genetic (Sullivan et al., 2012), with more than 1,000 genes implicated to date. Nevertheless, at this time little is known about the pathogenic mechanisms underlying social behavioral abnormalities in ASD and whether these are the consequence of selective vulnerability of specific cell types to genetic injury.

In 1943, Julia Bell and James Martin first described the genetic disorder now known as fragile X (Martin and Bell, 1943), the same year that Leo Kanner published his seminal psychiatric description of autism (Kanner, 1943). Using narrowly defined diagnostic criteria, early studies revealed the exceptionally high heritability of autism (Bailey et al., 1995; Folstein and Rutter, 1977; Freitag, 2007; Steffenburg et al., 1989), and identified fragile X as the first known genetic cause of the disorder (August, 1983; Brown et al., 1982, 1986). Broader modern clinical definitions have given rise to somewhat reduced heritability estimates (Sullivan et al., 2012); nevertheless, 50% of male patients and 20% of female patients with fragile X meet full diagnostic criteria for ASD (Kaufmann et al., 2017), making monogenic mutation of *FMR1*, the gene disrupted in fragile X (Verkerk et al., 1991), still the most common identified cause of the disease (Suhl and Warren, 2015). Moreover, FMRP, the protein product of *FMR1*, regulates roughly 20% of ASD susceptibility genes (Darnell et al., 2011; Van Driesche et al., 2019), thus this gene is thought to constitute a “node” of ASD etiology that can be studied to understand how mutations in a diverse array of genes produces the complex behavioral phenotypes that clinically unify the disorder (Dölen and Sahin, 2016).

Unlike the strong evidence implicating *FMR1* mutations in the etiology of ASD, evidence supporting a role for the genes encoding OT (*OXT*) and the OT receptor (*OXTR*) has been more controversial (Allen-Brady et al., 2009; Hovey et al., 2014; Satterstrom et al., 2020; Willsey et al., 2013; Zhang et al., 2017). At the same time, we and others have demonstrated that the nucleus accumbens (NAc), dorsal raphe nucleus (dRph), and ventral tegmental area (VTA) receive direct or indirect projections from hypothalamic OT neurons (Beier et al., 2015; Dölen et al., 2013; Hung et al., 2017; Nardou et al., 2019), which together constitute a neural circuit encoding social reward behaviors that may be especially relevant to autism (Chevallier et al., 2012). Interestingly, of these brain regions, only the hypothalamus, where OT neurons reside, exhibits the exaggerated protein synthesis phenotype seen in *Fmr1*-knockout (KO) mice (Qin et al., 2005). Furthermore, hypothalamic disruptions of OT neurons have been reported in human patients with ASD-related disorders such as Prader-Willi syndrome (Swaab et al., 1995), as well as in KO mice for ASD risk genes including *Fmr1* (Francis et al., 2014), *Cntnap2* (Peñagarikano et al., 2015), and *Pten* (Page et al., 2009). Nevertheless, social behavioral abnormalities seen in constitutive KO mice are not recapitulated following deletion of *Pten* in OT neurons (Clipperton-Allen et al., 2016), underscoring an important gap in our understanding of how pathognomonic features of the disease arise. Here we sought to bridge this gap by examining the interplay between genetic

disruption and molecularly defined circuits that subserve social behaviors that are disrupted in ASD.

RESULTS

Distribution of OT Neuronal Subtypes across the Rostral-Caudal Axis of the PVN

Magnocellular and parvocellular subtypes of OT neurons were first observed nearly a century ago (Gurdjian, 1927; Krieg, 1932; van den Pol, 2012), and early reports suggested that these subtypes can be segregated by their anatomical distribution (Armstrong et al., 1980; Hosoya and Matsushita, 1979; Sawchenko and Swanson, 1982; Swanson and Kuypers, 1980). Nevertheless, to date a quantitative analysis of the proportion and distribution of these OT neuron subtypes within the paraventricular nucleus (PVN) in mice has not been carried out. Here we used a well-established approach for differentiating between magnocellular and parvocellular OT neuronal subtypes, which is based on the observation that only magnocellular OT neurons send axonal projections outside of the blood-brain barrier (BBB) to the posterior pituitary (van den Pol, 2012). Specifically, intravenous (i.v.) injections of the BBB-impermeable retrograde tracer Fluoro-Gold (FG) into the lateral tail vein enable selective labeling of magnocellular but not parvocellular OT neurons (Merchenthaler, 1991). Combining this retrograde tracing approach with OT antibody (Ab) labeling (Ben-Barak et al., 1985; Whitnall et al., 1985) allowed us to conduct comprehensive quantification of the anatomical distribution and proportion of subtypes of OT neurons in the PVN of the hypothalamus in male mice (postnatal days 30–40). OT-Ab-labeled neurons were classified as magnocellular if they were FG positive and parvocellular if they were FG negative (Figures 1A–1F; Figure S1). Quantification of these subtypes ($n = 3$ mice, every other section quantified) revealed that in the PVN, 66% of OTergic neurons (952 ± 69 neurons) are magnocellular, while 34% of OTergic neurons (488 ± 25 neurons) are parvocellular (Figure 1G). Next we examined the distribution of all OT neurons according to their location along the rostral-caudal axis of the PVN. As shown in Figure 1H, magnocellular OT neurons are localized predominantly to the rostral division of the PVN, while parvocellular OT neurons are concentrated in the caudal division of the PVN. These data provide the first comprehensive quantification of the proportion and rostral-caudal distribution of magnocellular and parvocellular OT neurons and highlight the importance of examining the entire rostral-caudal axis of the PVN when assessing the contribution of OT neuronal subtypes to functional circuitry.

Electrophysiological Signatures of Magnocellular and Parvocellular OT Neurons

Within the PVN, magnocellular and parvocellular OT cells are intermixed with a wide variety of non-OTergic magnocellular and parvocellular neurons (Biag et al., 2012). Recent studies using electrophysiological circuit mapping of OTergic projections (Eliava et al., 2016; Xiao et al., 2017) relied on the implicit assumption that OT neurons can be classified as magnocellular or parvocellular using criteria established for PVN neurons of unspecified peptidergic identity (Hoffman et al., 1991; Luther et al., 2000, 2002; Tasker and Dudek, 1991). In order to explicitly test




Figure 1. Distribution of OT Neuronal Subtypes across the Rostral-Caudal Axis of the PVN

(A) Diagram illustrating labeling strategy. (B–E) Example images of rostral (B and C) and caudal (D and E) sections of the PVN at low magnification (B and D; scale bar, 200 μ m) and high magnification (C and E; scale bar, 50 μ m) showing OT-Ab (green, left), FG (magenta, center), and merged (right) images. (F) Drawing of magnocellular (left) and parvocellular (right) OT neurons identified by i.v. FG labeling. Scale bar indicates 20 μ m. (G) Magnocellular OT neurons account for 65.9% $\pm$ 2.6% and parvocellular OT neurons account for 34.1% $\pm$ 2.6% of PVN OT neurons ($n = 3$ animals; 1,439.3 $\pm$ 48.3 OT neurons per animal). (H) Magnocellular OT neurons are located predominantly in the rostral PVN, while parvocellular OT neurons are located predominantly in the caudal PVN. Data are represented as mean $\pm$ SEM. See also Figure S1.

this assumption, here we examined magnocellular and parvocellular electrophysiological signatures (Luther and Tasker, 2000) in identified OT neurons using our recently generated OT-Flp recombinase driver line (Nardou et al., 2019), crossed to a Flp-dependent GFP reporter line (fdGFP; Figure S1; Miyoshi et al., 2010). OT-2A-Flp::fdGFP mice ($n = 9$) were injected with i.v. FG and, blind to FG labeling status, GFP+ neurons were targeted for whole-cell current-clamp recording in acute slices of the PVN (Figure 2A). During recording, neurons were labeled with neurobiotin (Nb) to enable post hoc classification of cells as FG positive (FG+; Figure 2B) or FG negative (FG–; Figure 2C). Neurons were depolarized from a hyperpolarized membrane potential (approximately -100 mV; Figure S2), and the shape of the membrane depolarization preceding action potential (AP) initiation was classified as shoulder positive (Sh+; Figure 2D, blue) or shoulder negative (Sh–; Figure 2D, orange). OT neurons were examined for latency to first AP (Figures 2E and 2F), AP

duration (Figures 2H and 2I), and sensitivity to the voltage-gated K⁺ antagonist 4-aminopyridine (4-AP; Figures 2J–2O). These studies confirm that latency to first AP is longer in Sh+ compared with Sh– OT neurons (Figure 2E), as well in FG+ compared with FG– OT neurons when neurons are grouped by FG labeling (Figure 2F). Furthermore, AP duration (Figure 2G) is longer in Sh+ compared with Sh– OT neurons (Figure 2H) as well as longer in FG+ versus FG– OT neurons (Figure 2I). Application of the voltage-gated K⁺ channel blocker 4-AP (5 mM; Figures 2J and 2M) reduced the latency to first AP and increased AP duration in magnocellular (Sh+) OT neurons but had no effect on latency in parvocellular (Sh–) OT neurons (Figures 2K, 2L, 2N, and 2O).

To quantify whether an individual OT neuron's magnocellular or parvocellular identity can be predicted from its electrophysiological features, we used unsupervised k-means clustering analysis on the Z-scored values for AP frequency, duration, and latency to first AP for each neuron. The resulting best fit to these data are two clusters (Figure S2), which corresponded completely (100% [18 of 18 OT neurons]) with the qualitative magnocellular or parvocellular categorization on the basis of the presence or absence of a shoulder in the electrophysiological trace (Figure 2P) and in 94% (17 of 18 OT neurons) when defined by FG labeling (Figure 2Q). Consistent with the proportions determined in Figure 1, 61% of sampled neurons were Sh+ and 39% were Sh–. Similarly, 56% of sampled neurons were FG+ and 44% were FG– (Figure 2R). Further supporting our cross-validation, among Sh+ OT neurons, 91% were FG+, and 100% of Sh– neurons were FG– (Figure 2S). Similarly, 100% of FG+ neurons were Sh+ and 88% of FG– neurons were Sh– (Figure 2T). These results demonstrate that the electrophysiological signature of magnocellular and parvocellular neurons applies specifically to the subset of these cells that are oxytocinergic (Figures 2D–2O). Furthermore, these electrophysiological parameters can be used to build a mathematical model to predict magnocellular and parvocellular OT cell identity (Figures 2P and 2Q). As the proportion of electrophysiologically defined subtypes (Figures 2R–2T) is consistent with anatomical estimates (Figure 1G), these observations provide cross-validation of magnocellular and parvocellular OT neuronal subtype identities by two independently derived measures and demonstrate near perfect agreement of electrophysiological and anatomical classification systems (94% [17 of 18 OT neurons]).

scRNA-Seq of OT-Expressing Neurons in the PVN

Next we sought to determine whether magnocellular and parvocellular OT neurons exhibit distinct transcriptional profiles. Although previous studies have performed single-cell RNA sequencing (scRNA-seq) analysis of the entire hypothalamus (Chen et al., 2017; Romanov et al., 2017), low recovery rates of OT neurons in these studies limited their power for identifying magnocellular and parvocellular subtypes by transcriptional profiling, leading to confusion about existing classification systems (Althammer and Grinevich, 2017). Others have used modern molecular strategies to selectively enrich samples for OT neurons, but these studies used relatively low-throughput qRT-PCR to focus on a handful of target genes and bulk RNA-seq analyses that can mask important sources of heterogeneity across cell subtypes and states (Maynard




Figure 2. Electrophysiological Signatures of Magnocellular and Parvocellular OT Neurons

(A) Validation strategy using i.v. FG and Nb-filled recording pipette.

(B and C) FG+ (B) and FG- (C) neurons recovered after electrophysiological recording. Nb (blue, left), GFP (green, center-left), FG (magenta center-right), and merge (right). Scale bars, 20 μ m.

(D and G) Representative traces from magnocellular (blue) and parvocellular (orange) OT neurons.

(E, F, H, and I) Latency to first AP and AP duration are shorter in Sh- ($n = 7$) compared to Sh+ ($n = 11$) OT neurons ($U = 0$, $p > 0.001$ [latency]; $U = 0$, $p > 0.001$ [duration]) (E and H) and FG- ($n = 8$) compared with FG+ ($n = 10$) OT neurons ($U = 0$, $p < 0.001$ [latency]; $U = 5$, $p > 0.001$ [duration]) (F and I).

(J and M) Representative traces (blue, magnocellular [Sh+] OT; orange, parvocellular [Sh-] OT; black, following 5 mM 4-AP).

(K and L) 4-AP reduces latency to first AP in magnocellular (Sh+, $n = 10$, sum of signed ranks = 55, $p = 0.002$) (K) but not parvocellular (Sh-, $n = 7$, sum of signed ranks = 10, $p = 0.469$) (L) OT neurons.

(N and O) 4-AP increases AP duration in magnocellular (Sh+, $n = 10$, sum of signed ranks = -51, $p = 0.006$) (N) and parvocellular (Sh-, $n = 7$, sum of signed ranks = -28, $p = 0.016$) (O) OT neurons.

(P and Q) K-means cluster analysis using Z-scored values for AP frequency, duration, and latency to first AP for each neuron. (P) Clusters coincided completely (100% [18 of 18 neurons]) with qualitative magnocellular or parvocellular categorization on the basis of shoulder. (Q) Cluster assignment matched magnocellular or parvocellular identity of 94% (17 of 18) neurons defined by FG labeling.

(R) Left: 61% of sampled neurons ($n = 11$) were Sh+ and 39% ($n = 7$) were Sh-. Right: 56% of sampled neurons ($n = 10$) were FG+ and 44% ($n = 8$) were FG-.

(S) Left: 91% of Sh+ neurons ($n = 10$ of 11) were FG+. Right: 100% of Sh- neurons were FG- ($n = 7$ of 7).

(T) Left: 100% of FG+ neurons ($n = 10$ of 10) were also Sh+. Right: 88% of FG- neurons were Sh- ($n = 7$ of 8).

Data are represented as mean $\pm$ SEM. * $p < 0.05$; Mann-Whitney U test for unpaired (E, F, H, and I) or Wilcoxon rank-sum test for paired (K, L, N, and O) comparisons (both two-tailed). See also Figures S1 and S2.

et al., 2018). To overcome these technical hurdles, here we combined full-length cDNA scRNA-seq (SMART-Seq2) with molecular targeting of OT neuron subtypes in OT-2A-Flp::fdGFP mice and additionally used projection target mapping (Chev  e et al., 2018) to enable cross-validation of transcriptional profiles against OT neuronal subtype features with established ground truth (Merchenthaler, 1991; Figures 1 and

2). In total, 194 OT neurons were collected by fluorescence-activated cell sorting (FACS) and used as input for a modified SMART-Seq2 scRNA-Seq library preparation (Figure 3A). We next used the Uniform Manifold Approximation and Projection (UMAP) embedding method (Becht et al., 2018) to visualize the transcriptional landscape. We performed a silhouette analysis to quantitatively determine the optimal number of clusters




Figure 3. scRNA-seq of OT-Expressing Neurons in the PVN

(A) Schematic overview of experiment. (B) Plot of the average silhouette width versus number of clusters for scRNA-seq data indicating that the optimal number of clusters is 2. (C) UMAP of scRNA-seq data with the two clusters indicated by the silhouette analysis. (D) Cluster 1 is majority FG+ (91 of 125 cells [72.8% FG+]) and cluster 2 is devoid of Fluoro-Gold positive cells (21 of 21 cells [100% FG-]). (E) Magnocellular (blue) and parvocellular (orange) neuron identity assigned to cluster 1 and cluster 2, respectively. (F) Empirical cumulative distribution functions for all expressed genes in the two cell types indicated greater intra-cell type heterogeneity within magnocellular neurons ($p = 1.665e^{-15}$). Kolmogorov-Smirnov (K-S) test for statistical comparison. (G) Comparison of the density of mitochondrial RNA in the two clusters as calculated via the distributions of the per cell ratio of reads mapped to mitochondrial RNA versus total number of mapped reads. One-tailed t test for statistical comparison ($p = 0.0007972$). See also [Figures S1](#) and [S3](#).

in the UMAP embedding (Figure 3B). This calculation yielded a best fit of two clusters, which are plotted in Figure 3C.

To determine whether these two cell type clusters map onto magnocellular and parvocellular OT neuronal subtypes, we then assessed the distribution of FG labeling across the two clusters (Figure 3D). As 72.8% of cells in cluster 1 were FG positive (FG+), while cluster 2 contained only cells that were FG negative (FG-; 100%) across all three replicates (Figure S3), we deduce that cluster 1 cells are magnocellular, while cluster 2 cells are parvocellular OT neurons (Figures 3E and 3F). The FG tail vein injection labeling technique, like all labeling techniques, has a non-zero false-negative rate. In our assay, these false-negative magnocellular neurons would be represented in the FG- population of cluster 1. Additionally, as the emission wavelength of FG is known to be altered

at acidic pH (Schmued and Fallon, 1986), the absence of FG labeling in a subset of cells in the magnocellular cluster may represent acidification of the intracellular environment during dissociation and FACS sorting. Nevertheless, as the FG- cells in the magnocellular cluster have the same characteristics (e.g., mRNA levels, number of genes expressed) as the FG+ cells in that cluster (Figure S3), and as all cells had to pass strict quality metrics to be included in the analysis, it is unlikely that diminished cell health or quality accounts for this finding. Taken together with our observation that only one of the two transcriptionally defined OT populations demonstrated any FG+

staining, these results enabled us to use FG labeling as a supervisory metric to discriminate between the two populations.

To cross-validate this observation, we next quantified the density of mitochondrial RNA in the two clusters. Previous reports indicate that magnocellular neurons have increased mitochondrial density (Dayanithi et al., 2012; van den Pol, 1982), a specialization thought to support the increased metabolic demands of magnocellular neurons that synthesize and release significantly larger quantities of OT peptide compared with parvocellular neurons (Dreifuss, 1975; Ludwig and Leng, 2006; Nordmann and Morris, 1984; van den Pol, 1982, 2012; Pow and Morris, 1989). Consistent with these reports, magnocellular neurons had a greater ratio of mitochondrial reads to genomic reads than that of parvocellular neurons (Figure 3G). Moreover, within the magnocellular cluster, where FG+ neurons (purple

circles) dominate, the FG– neurons (green circles) are evenly distributed across mitochondrial density measurements (Figure 3G), consistent with our interpretation that FG– cells in this cluster represent a technical artifact, rather than a biologically significant subcategory of magnocellular neuron. Together, these studies identify two distinct transcriptional subtypes of OT neurons using high-resolution full-length sequencing techniques, in samples enriched for OT neurons by molecular techniques. By cross-validating these transcriptional profiles across two independent distinguishing features (projection target and mitochondrial density), these studies are the first to establish a transcriptional signature for classifying magnocellular and parvocellular OT neuronal subtypes.

Differential Expression Analysis of Clusters Determines Unique Molecular Markers of Magnocellular and Parvocellular OT Neurons

Having identified anatomical, circuit, electrophysiological, and transcriptional characteristics that differentiate magnocellular and parvocellular OT neuronal subtypes, next we made use of our high-resolution transcriptional data to identify differentially expressed genes across magnocellular and parvocellular OTergic neurons. Using the Monocle2 likelihood ratio test (Trapnell et al., 2014), we identified 181 genes (Table S1) with significant (0.1% false discovery rate [FDR]; Monocle2; Benjamini-Hochberg [BH] corrected) differential expression (DE) between magnocellular and parvocellular OT cells (Figure 4A), representing novel discriminating marker genes for these two subtypes. Among the most significantly differentially expressed genes was the calcium-binding protein Calbindin (*Calb1*) and a large conductance calcium-activated potassium channel subunit (*Kcnmb4*), both enriched for expression in the magnocellular OT neuronal population (Figure 4B), as well as the extracellular matrix serine protease Reelin (*Reln*) and the cannabinoid receptor 1 (*Cnr1*) genes, both significantly enriched in parvocellular OT neurons (Figure 4C). These results were validated with fluorescent *in situ* hybridization chain reaction version 3.0 (HCR 3.0; Table S2) (Choi et al., 2018) combined with anatomical criteria (Figure 1), which demonstrated that *Oxt*⁺ neurons in the rostral portion of the PVN were specifically colocalized with *Calb1* and not *Cnr1* mRNA (Figure 4D), while in more caudal sections, *Oxt*⁺ neurons colocalized with *Cnr1* and not *Calb1* (Figure 4E). In addition, 18 differentially expressed genes were identified as potentially contributing to the electrophysiological signature of magnocellular and parvocellular OT neurons (Figure S3). These results are the first to establish a set of discriminating molecular identifiers that can distinguish between magnocellular and parvocellular OT neuronal subtypes. Although many other differentially expressed genes likely also exist, this conservative list of 181 significantly differentially expressed genes provides a high-value candidate gene list for future studies examining the molecular mechanisms contributing to the functional, physiological, and connectivity differences between these two OT neuronal subtypes. Moreover, as described below, identification of differentially expressed genes paves the way for functional genomics in magnocellular and parvocellular OT neurons.

Fmr1-KO Mice Exhibit Social Domain-Specific Impairment in Reward Learning

For more than 20 years, ASD researchers have taken advantage of the high heritability of the disease to generate transgenic mice with construct validity for ASD (Nestler and Hyman, 2010) and used these model mice to examine how ASD risk genes might converge at the level of biochemical pathways and synapses (Dölen and Bear, 2009; Dölen and Sahin, 2016). This “bottom-up” approach has led to the discovery of some very interesting fundamental mechanisms, but today we are arguably no closer to understanding how disruption of these synaptic and biochemical functions gives rise to the behavioral phenotypes that unify the disease (Antoine et al., 2019). Because ASD remains a diagnosis that is defined by behavioral abnormalities, taking instead the “top-down” approach offers a significant opportunity to reconcile the etiological diversity of ASD with the convergence of the clinical presentation (Diester et al., 2015; Dölen et al., 2015). Combining this approach with the *Fmr1*-KO mouse model, which has established construct validity for the disease (The Dutch-Belgian Fragile X Consortium, 1994), here we focused on behaviors with face validity for social impairments in ASD (Nestler and Hyman, 2010).

The social motivation theory of autism proposes that a primary deficit in peer-peer social reward learning could account for all of the characteristic behavioral impairments that define the clinical picture in ASD (Chevallier et al., 2012). To test this theory, we began by characterizing three forms of reward learning in the constitutive *Fmr1* KO using peer-peer social, cocaine, and alloparent social conditioned place preference (CPP) assays (Bardo and Bevins, 2000; Fang and Wang, 2017; Panksepp and Lahvis, 2007). These results demonstrate a significant impairment in peer-peer social CPP (sCPP) in *Fmr1*-KO mice compared with wild-type (WT; Figures 5A–5D; Figure S4), while both WT and *Fmr1*-KO mice exhibit significant cocaine CPP (cCPP) (Figures 5E–5H), as well alloparent CPP (aCPP) (Figures 5I–5L, S4). Our finding that *Fmr1*-KO mice exhibit impairments in social reward but not drug reward (Figures 5A–5H) is consistent with human studies that have identified social (Demurie et al., 2011; Stavropoulos and Carver, 2014) but not food or patient salient object (Assaf et al., 2013; Cascio et al., 2012) reward deficits in ASD (but see Damiano et al., 2012; Smith et al., 2014). Furthermore, our finding that *Fmr1*-KO mice exhibit no significant impairment in filial attachments (Figures 5I–5L) is consistent with the human literature (Baron et al., 2020; Naber et al., 2007, 2008; Rutgers et al., 2004; Stokes et al., 2007; Vivanti and Nuske, 2017). Previously, we showed that changes in locomotor activity are unlikely to account for differences in sCPP across a variety of experimental manipulations (Dölen et al., 2013; Nardou et al., 2019). Although previous reports raise the possibility that *Fmr1*-KO mice exhibit altered locomotor behavior, these have been inconsistent across labs (Bernardet and Crusio, 2006); furthermore, as aCPP and cCPP were not different (Figures 5E–5L), it is unlikely that changes in locomotor activity could account for the observed differences in sCPP (Figures 5A–5D). Taken together, these findings support the view that social reward learning impairments in *Fmr1*-KO mice are domain specific and restricted to peer-peer social attachments. Furthermore, they provide improved face validity for social impairments in the *Fmr1*-KO




Figure 4. Differential Expression Analysis of Clusters Determines Unique Molecular Markers of Magnocellular and Parvocellular OT Neurons

(A) Heatmap of significantly differentially expressed genes (Monocle likelihood ratio test, FDR = 0.1%) between magnocellular and parvocellular OT neurons.

(B) UMAPs (left) and violin plots (right) for genes expressed more highly in magnocellular OT neurons: *Calb1* and *Kcnmb4* (identified using FDR of 0.00001%; $p = 1.28 \times 10^{-30}$ and $p = 5.46 \times 10^{-7}$, respectively).

(C) UMAPs (left) and violin plots (right) for genes expressed more highly in parvocellular OT neurons: *Reln* and *Cnr1* (identified using FDR of 0.00001%; $p = 1.48 \times 10^{-15}$ and $p = 6.67 \times 10^{-11}$, respectively).

(D and E) Validation of differential expression via fluorescent *in situ* hybridization chain reaction version 3.0 (HCR 3.0) within the PVN. *Calb1* but not *Cnr1* is colocalized with *Oxt* within the rostral PVN (D). *Cnr1* but not *Calb1* is colocalized with *Oxt* in the caudal PVN (E). Scale bars, 20 μ m. Distribution of *Calb1* (rostral) and *Cnr1* (caudal) expression in OT neurons recapitulates differential localization of magnocellular (rostrally enriched) and parvocellular (caudally enriched) neurons respectively.

* $p < 0.05$; Monocle2 likelihood ratio test for statistical comparisons. See also Figure S3 and Tables S1 and S2.




Figure 5. *Fmr1*-KO Mice Exhibit Social Domain-Specific Impairment in Reward Learning

(A) Timeline for sCPP. (B and C) Individual (top) and average (bottom) data indicate that WT mice (B; $t_{[23]} = 3.586$, $p = 0.002$) and not *Fmr1*-KO mice (C; $t_{[15]} = 1.324$, $p = 0.205$) exhibit sCPP. (D) Normalized ($t_{[38]} = 2.917$, $p = 0.006$) and subtracted ($t_{[38]} = 3.23$, $p = 0.003$) comparisons of social preference reveal a difference in the magnitude of sCPP between WT and *Fmr1*-KO mice.

mouse model of ASD, as well as additional support for the social motivation theory of autism.

Parvocellular OT Neurons Innervate NAC

The impairment in peer-peer sCPP in the *Fmr1*-KO mouse (Figures 5A–5D) recapitulates previously reported consequences of OT receptor blockade or ablation in the NAC (Dölen et al., 2013), raising the possibility that these phenotypes share overlapping mechanisms. To test this possibility, we began by taking an anatomical approach that we have used previously to qualitatively identify both magnocellular and parvocellular OT neuronal projections from the PVN to the VTA (Hung et al., 2017). Example images of immunohistochemically identified OT neurons from a WT mouse that received an i.v. FG injection and NAC injection of retrogradely transported fluorescent microspheres (retrobeads [RtBs]; Katz et al., 1984), indicated that unlike the VTA, the NAC receives an exclusively parvocellular OTergic projection (Figure S5). To quantify this finding, we next injected RtBs into the NAC (Figure S6) of OT-2A-Flp::fdGFP mice ($n = 6$ mice) and performed current-clamp recordings from RtB/GFP double-positive neurons in acute slices of the PVN (Figures 6A and 6B). All RtB-labeled, OT-positive neurons recorded (nine of nine) exhibited a parvocellular electrophysiological phenotype by latency to first AP, AP duration, and 4-AP sensitivity (Figures 6C–6J), providing quantitative electrophysiological cross-validation of our immunohistochemical observations.

Next, we used unsupervised k-means clustering, trained with electrophysiological data from Figure 2, to provide unbiased quantitative cross-validation of our electrophysiological findings (Figure 6K). We further used the soft k-means method to calculate the probability that a cell belongs to a given cluster on the basis of the Euclidean distance between that cell and the centers of each cluster in feature space. The probability that a cell belongs to an individual cluster is given by the following softmax equation:

$$p(x_i, c_j) = \frac{1}{\sum_{n=1}^N \left(\frac{\|x_i - c_j\|}{\|x_i - c_n\|} \right)^2}$$

where x_i is a cell's coordinates, c_j is the coordinates of the center of the cluster j , $p(x_i, c_j)$ is the probability that x_i belongs to cluster j ,

(E) Protocol and timeline for cocaine CPP.

(F and G) Individual (top) and average (bottom) data indicate that WT (F; $t_{[14]} = 6.454$, $p < 0.0001$) and *Fmr1*-KO mice (G; $t_{[10]} = 4.297$, $p = 0.002$) exhibit cocaine CPP.

(H) Normalized ($t_{[24]} = 0.609$, $p = 0.548$) and subtracted ($t_{[24]} = 0.644$, $p = 0.525$) comparisons of cocaine preference reveal a similar magnitude of cocaine CPP in WT and *Fmr1*-KO mice.

(I) Protocol and timeline for alloparent CPP.

(J and K) Individual (top) and average (bottom) data indicate that WT (J; $t_{[11]} = 2.931$, $p = 0.014$) and *Fmr1*-KO mice (K; $t_{[8]} = 2.488$, $p = 0.038$) exhibit alloparent CPP.

(L) Normalized ($t_{[19]} = 0.234$, $p = 0.817$) and subtracted ($t_{[19]} = 0.354$, $p = 0.727$) comparisons of alloparent preference reveal a similar magnitude of alloparent CPP in WT and *Fmr1*-KO mice.

Data are represented as mean $\pm$ SEM. * $p < 0.05$, Student's t test (two-tailed, paired [B, C, F, G, J, and K] or unpaired [D, H, and L]) for statistical comparisons.




Figure 6. Parvocellular OT Neurons Innervate NAc

(A) RtB labeling and recording strategy in OT-2A-Flp::fdGFP mice.

(B) Example of a GFP/RtB double-positive OT neuron targeted for electrophysiological recording. Left to right: DIC (gray), GFP (green), RtB (red), and merge. Scale bar, 10 μ m.

(C and E) Traces recorded from a NAc-projecting OT neuron.

(D) Latency to first AP of NAc-projecting OT neurons ($n = 9$; 21.2 ± 4.2 ms).

(F) AP duration of NAc-projecting OT neurons ($n = 9$; 1.4 ± 0.07 ms).

(G and I) Traces (red, RtB+; black, following 5 mM 4-AP).

(H and J) 4-AP does not change latency to first AP ($n = 8$, sum of signed ranks = 6, $p = 0.742$) (H) but increases AP duration ($n = 8$, sum of signed ranks = -34, $p = 0.016$) (J) in NAc-projecting OT neurons.

(K) Using the k-means clustering algorithm trained using data from Figure 2, 100% (nine of nine) sampled RtB/GFP double-positive cells were assigned to the parvocellular cluster.

Data are represented as mean $\pm$ SEM. * $p < 0.05$, Wilcoxon rank-sum test (two-tailed, paired [H and J]). See also Figures S1, S2, S5, and S6.

and N is the total number of clusters. Using this analysis, we determined that NAc-projecting OT neurons belong to the parvocellular cluster with a mean probability of $92.8\% \pm 2.7\%$. This is similar ($H_{[2]} = 0.762$, $p = 0.6833$; Kruskal-Wallis test) to the probabilities for FG- neurons and Sh- neurons ($87.2\% \pm 8.9\%$ [Figure 2Q] and $96\% \pm 1.9\%$ [Figure 2P], respectively). Although we and others have identified a number of brain regions involved in the regulation of social behaviors that receive both magnocellular and parvocellular OTergic projections (Hung et al., 2017; Knobloch et al., 2012; Xiao et al., 2017), until technological advancements enable functional genomic interrogation of magnocellular and parvocellular OT neurons separately, identification of the NAc as a brain region that receives an exclusively parvocellular OTergic input in male mice provides a unique opportunity to selectively interrogate the functional role of *Fmr1* in parvocellular OT neurons for social reward learning.

Parvocellular OT Neurons Require *Fmr1* to Mediate Social Reward Learning

To directly examine the behavioral consequence of disrupting the parvocellular OT input to the NAc, we began by testing whether ongoing *Fmr1* function is required for peer-peer social reward learning. We used conditional (Cre-dependent) *Fmr1*-KO mice (Mientjes et al., 2006) (*cdFmr1* KO) receiving injections

of canine adenovirus-expressing Cre (CAV2-Cre-GFP) or GFP (CAV2-GFP) (Hung et al., 2017; Kremer et al., 2000; Uematsu et al., 2017) bilaterally into the NAc. As we have previously used the CAV2-Cre-GFP to manipulate gene expression for examining the interplay between genetic and circuit mechanisms of sCPP (Hung et al., 2017), here we controlled for genetic and rearing effects in *cdFmr1*-KO mice and examined sCPP behaviors following injection of control virus (CAV2-GFP). Injection sites and viral function were confirmed, as shown in Figure S6. Seven days later, sCPP was measured, and we observed impaired sCPP in *cdFmr1*-KO mice injected with CAV2-Cre-GFP but not CAV2-GFP (Figures 7A–7D). Importantly, as acute knockdown of *Fmr1* in adolescence is sufficient to recapitulate peer-peer sCPP impairments seen in the constitutive *Fmr1* KO, these results demonstrate an ongoing requirement of *Fmr1* in regulating these behaviors and suggest that previous findings that hysteresis plays an important role in the pathogenesis of social impairments in ASD (Orefice et al., 2019) do not generalize across etiologies.

To further circumscribe the OT cell-type specific role of *Fmr1* in regulating social reward learning, we next crossed *cdFmr1*-KO mice to OT-2A-Flp mice and generated retrograde adeno-associated virus expressing Flp-dependent-Cre (rgAAV-fdCre) (Penzo et al., 2015; Schneeberger et al., 2019; Tervo et al.,




Figure 7. Parvocellular OT Neurons Require *Fmr1* to Mediate Social Reward Learning

(A) Protocol for sCPP following acute knockdown of *Fmr1* in neurons projecting to NAC.

2016) (Figure 7E). As our experiments above indicated that control *cdFmr1* mice injected with CAV-GFP (Figure 7B) exhibit sCPP that is comparable with that seen in WT mice (Figure 5B), here we decided to control for potential off-target effects of rgAAV-fdCre, which although well validated for selectivity and efficiency, had not been previously tested in this behavioral and injection target context. As shown in Figures 7F–7H, sCPP was absent in OT-2A-Flp::*cdFmr1* KO, while WT animals injected with rgAAV-fdCre exhibited normal social reward learning. Injection sites and viral function were confirmed, as shown in Figure S6. Taken together with the studies above identifying the source of these OTergic inputs to the NAC as parvocellular (Figure 6), these results demonstrate that *Fmr1* in OT neurons projecting to the NAC specifically, is required for peer-peer social reward learning. Furthermore, as previously parvocellular OT neurons had only been directly implicated in non-social functions (e.g., feeding, yawning, erection, cardiovascular function, pain; Althammer and Grinevich, 2017; Eliava et al., 2016; Melis et al., 1986; Petersson, 2002; Valassi et al., 2008), these studies identify the first social function directly encoded by this subtype of OT neuron.

Although the studies above provide direct evidence that normal function of OT neurons is necessary for social reward learning, in order to test the specificity of this function, we next sought to determine the consequences of *Fmr1* deletion in magnocellular OT neurons. To date we have yet to identify a brain region that receives an exclusively magnocellular OTergic projection (Hung et al., 2017), so the projection targeting method used above could not be used to selectively knock down *Fmr1* in magnocellular OT neurons. Instead, we capitalized on our discovery that *Calb1* is a discriminatory marker of magnocellular compared with parvocellular OT neurons. Crossing *Calb1*-IRES-Cre mice (Daigle et al., 2018; Nigro et al., 2018) with *cdFmr1*-KO mice allowed us to selectively knock down *Fmr1* in magnocellular OT neurons expressing *Calb1*. Although this manipulation also knocked down

(B and C) Individual (top) and average (bottom) data indicating that *cdFmr1*-KO mice injected with CAV2-GFP (B; $t_{118} = 3.95$, $p = 0.001$) but not CAV2-Cre-GFP (C; $t_{119} = 0.69$, $p = 0.499$) exhibit sCPP.

(D) Normalized ($t_{137} = 2.453$, $p = 0.019$) and subtracted ($t_{137} = 2.566$, $p = 0.015$) comparisons reveal reduced magnitude sCPP in *cdFmr1*-KO mice injected with CAV2-Cre-GFP compared with CAV2-GFP.

(E) Protocol for sCPP following acute knockdown of *Fmr1* in OT neurons projecting to NAC.

(F and G) Individual (top) and average (bottom) data indicating that WT (F; $t_{431} = 4.947$, $p > 0.0001$) but not OT-2A-Flp::*cdFmr1* KO (G; $t_{351} = 0.349$, $p = 0.729$) mice injected with rgAAV-fdCre exhibit sCPP.


(H) Normalized ($t_{78} = 3.295$, $p = 0.002$) and subtracted ($t_{78} = 2.937$, $p = 0.004$) comparisons reveal reduced magnitude sCPP in OT-2A-Flp::*cdFmr1* KO compared with WT mice injected with rgAAV-fdCre.

(I) Protocol and timeline for sCPP.

(J and K) Individual (top) and average (bottom) data indicate that WT (J; $t_{191} = 3.901$, $p = 0.004$) and *Calb1*-IRES-Cre::*cdFmr1*-KO mice (K; $t_{191} = 7.146$, $p > 0.0001$) exhibit social CPP.

(L) Normalized ($t_{17.941} = 1.642$, $p = 0.118$) and subtracted ($t_{17.874} = -1.965$, $p = 0.065$) comparisons of social preference reveal that WT and *Calb1*-IRES-Cre::*cdFmr1*-KO mice exhibit similar magnitude sCPP.

Data are represented as mean $\pm$ SEM. * $p < 0.05$, Student's *t* test (two-tailed, paired [B, C, F, G, J, and K]; two-tailed, unpaired [D, H, and L]) for statistical comparisons. Welch's correction was used to account for unequal variance and small sample size in (L). See also Figure S6.



Fmr1 in other, non-OTergic *Calb1*-expressing neuronal populations elsewhere in the brain (see also Figures 4D and 4E), remarkably, sCPP was not impaired in *Calb1*-IRES-Cre::cd*Fmr1*-KO mice (Figures 7I–7L), consistent with the selective requirement of *Fmr1* in parvocellular OT neurons for peer-peer social reward learning. Furthermore, as *Calb1* is also expressed in the Purkinje cells of the cerebellum, these results suggest that previous reports implicating the cerebellum in the pathogenesis of social impairments in the *Tsc1*-KO model of ASD (Kelly et al., 2020; Tsai et al., 2012) do not generalize across ASD etiologies. Taken together, these anatomical, electrophysiological, and functional genomic data demonstrate that social reward learning deficits seen in constitutive *Fmr1*-KO mice, are not the consequence of hysteresis but rather the ongoing requirement of *Fmr1* in parvocellular, but not magnocellular, OT neurons and directly implicate these neurons in the pathogenesis of a peer-peer social reward learning behavior that has clear face validity for social impairments that define ASD.

ASD Risk and FMRP Target Genes Are Enriched in Parvocellular OT Neurons

Having identified a role for *Fmr1* in parvocellular OT neuronal regulation of domain-specific social reward learning, we next sought to determine whether this pathogenic mechanism in

Figure 8. ASD Risk and FMRP Target Genes Are Enriched in Parvocellular OT Neurons

(A) Density of the log₂ fold change for (black) all expressed genes, (orange) all ASD risk genes, and (red) all significantly differentially expressed (FDR = 0.1%) ASD risk genes. ASD risk genes (orange versus black, $p = 0.0009279$, one-sided Wilcoxon rank-sum test) and differentially expressed ASD risk genes (red versus black, $p = 8.589 \times 10^{-9}$) are enriched in parvocellular OT neurons. Individually labeled genes are the union of ASD risk genes and FMRP binding partners that are differentially expressed between neuron types.

(B) Density of the log₂ fold change for (black) all expressed genes, (light blue) all FMRP binding partners, and (dark blue) all significantly differentially expressed (FDR = 0.1%) FMRP binding partners. FMRP binding partners (light blue versus black, $p = 6.83 \times 10^{-16}$, one-sided Wilcoxon rank-sum test) and differentially expressed FMRP binding partners (dark blue versus black, $p = 1.751 \times 10^{-9}$) are enriched in parvocellular OT neurons. Individually labeled genes are the union of ASD risk genes and FMRP binding partners that are differentially expressed between neuron types (FDR = 0.1%).

(C) Density of the log₂ fold change for (black) all expressed genes, (light green) all schizophrenia risk genes, and (dark green) all significantly differentially expressed (FDR = 0.1%) schizophrenia risk genes. Schizophrenia risk genes are not enriched in magnocellular or parvocellular OT neurons (light green versus black, $p = 0.98$) and differentially expressed schizophrenia risk genes (dark green versus black, $p = 0.21$, one-sided Wilcoxon rank-sum test) are not enriched in parvocellular OT neurons.

(D) Density of the log₂ fold change for (black) all expressed genes, (light yellow) all Alzheimer's disease (AD) risk genes, and (dark yellow) all significantly differentially expressed (FDR = 0.1%) AD risk genes. AD risk genes are not enriched in magnocellular or parvocellular OT neurons ($p = 0.99$).

(E) Density of the log₂ fold change for (black) all expressed genes, (light pink) all type 2 diabetes risk genes, and (dark pink) all significantly differentially expressed (FDR = 0.1%) type 2 diabetes risk genes. Type 2 diabetes risk genes are not enriched in magnocellular or parvocellular OT neurons ($p = 0.33$). Vertical tick marks below density plots indicate the individual differentially expressed ASD risk genes (A), FMRP binding partners (B), schizophrenia risk genes (C), and the single differentially expressed AD risk gene (D). Note the lack of differentially expressed type 2 diabetes risk genes (E). See also Figure S7 and Tables S1, S3, and S4.

fragile X may generalize to other ASD etiologies. Given the diversity of ASD risk gene functions, we hypothesized that the pathogenic overlap might occur at the level of selective vulnerability of parvocellular OT neurons to genetic injury, even if cellular phenotypes are not shared across all ASD etiologies. If so, we predict that this overlap would be evident in the transcriptional profile of these cells. In order to test this hypothesis, we performed gene set enrichment analysis for ASD risk genes (Abrahams et al., 2013) and FMRP binding partners (Van Driesche et al., 2019). In both cases, genes were significantly enriched in parvocellular relative to magnocellular neurons (Figures 8A and 8B; Tables S3 and S4). Additionally, 10 of the 12 differentially expressed genes in the intersection of the 2 groups (*Dlgap2*, *Cnr1*, *Pacs1*, *Tanc2*, *Anks1b*, *Slc6a1*, *Synj1*, *Tspan7*, *Camk2b*, *Stxbp1*, *Fbxo11*, and *Foxp1*; Figures 8A and 8B; individually labeled genes; Figure S7) were also enriched in parvocellular cells. As OT neurons in both the rostral and caudal PVN (Figure S7) express FMRP, these results indicate that subtype-specific expression of FMRP itself is unlikely to account for selective enrichment of ASD risk genes and FMRP binding partners in parvocellular OT neurons.

Next we repeated this analysis on risk genes for schizophrenia, Alzheimer's disease, and type 2 diabetes (Kanehisa and Goto, 2000; Scelsi et al., 2018; Sullivan et al., 2012). Although a handful of schizophrenia risk genes were differentially expressed in OT neuronal subtypes, these were not significantly enriched in either magnocellular or parvocellular OT neurons ($p = 0.98$; Figure 8C). Moreover, Alzheimer's disease had only a single differentially expressed gene (*Lhx5*), and type 2 diabetes risk genes were completely absent among differentially expressed genes in OT cells (Figures 8D and 8E). These results support the view that cell type-specific vulnerability of parvocellular OT neurons to genetic injury generalizes across multiple ASD etiologies but is not apparent in unrelated genetic disorders that are not defined by impairments in social behavior.

DISCUSSION

In the present studies, we took a comprehensive approach to provide validation of 8 of the 14 characteristics identified as distinguishing between magnocellular and parvocellular OT neuron, having discovered two of these characteristic features here (Figure S8, center; Figures 1, 2, 3, 4, and 8). To date, OT's role as a regulator of social behaviors has largely focused on reproductive behaviors such as pair bonding and parental behaviors. Here we provide evidence that these filial social behaviors (Figure S8, left) are governed by mechanisms distinct from those governing peer-peer social interactions (Figure S8, right). Specifically, peer-peer but not filial social attachments are disrupted in the *Fmr1* KO (Figure 5). Furthermore, in parvocellular OT neurons (Figure 6) but not magnocellular OT neurons (Figure 4), *Fmr1* is required for peer-peer social reward learning (Figure 7). Taken together these results delineate a novel example of parallel processing, which enables a single peptide to encode a wide diversity of social behaviors, significantly expanding the network capabilities of the social brain (Figure S8).

For more than a decade, research into the pathogenesis of ASD has focused largely on understanding unifying biochem-

ical mechanisms on the basis of the hypothesis that ASD is a "synaptopathy" (Bear et al., 2008; Dölen and Bear, 2009; Grabrucker et al., 2011). One implicit assumption of this hypothesis is that identification of synaptic phenotypes anywhere in the brain provides insight into disease pathogenesis (Dölen et al., 2007). Our present findings provide evidence that the pathogenesis of social impairments in ASD can be better understood using the framework of "pathocllisis" (Klatzo, 2003), whereby subtype-specific differences render a specific neuronal population vulnerable to genetic lesion. Furthermore, by examining mechanisms at the intersection of both construct and face validity (Dölen and Sahin, 2016; Dölen et al., 2015), the present studies have identified a critical role of parvocellular OT in the pathogenesis of ASD. Importantly, this mechanistic link has been overlooked by genetic association studies, underscoring the importance of taking a neurobiological approach to the disease that does not forfeit etiological considerations to genetics alone.

Clinical trials examining core symptoms of ASD have failed to consistently demonstrate efficacy of intranasal OT treatment (Anagnostou et al., 2012; Dadds et al., 2014; Guastella et al., 2015; Hollander et al., 2007), likely because intranasal OT targets magnocellular but not parvocellular OT neurons (Dölen, 2015; Leng and Ludwig, 2016). Furthermore, although much of the understanding of OT as a regulator of social behaviors is based on the importance of magnocellular OT neurons for pair bonding, we found no evidence to support the view that the magnocellular subtype of OT neurons is involved in the pathogenesis of ASD. Rather, our findings implicate the parvocellular OT neurons as the appropriate therapeutic target for correcting social impairments in ASD. Moreover, the only study to date that has examined the therapeutic potential of intranasal OT specifically in the fragile X population (Hall et al., 2012) found a correlation between improvements in a measure of social anxiety (a syndromic symptom that is seen more frequently in fragile X than ASD; Thurman et al., 2014) and decreased serum cortisol levels (Hall et al., 2012). As intranasally delivered OT also reaches systemic OT receptors in the adrenal cortex (Hammock and Levitt, 2013), whose activation suppresses cortisol release (Chiodera and Legros, 1981; Legros et al., 1984; de Oliveira et al., 2007; Stachowiak et al., 1995), these anxiolytic effects likely reflect indirect activity of OT in the periphery.

Future studies dissecting input and output connections, defining developmental programs, and computational modeling of this circuitry will be enabled by the technical achievements of this work, not the least of which is the identification of novel molecular markers, which can be used in combination with our OT-Flp driver line to specify (by multiple feature selection) circuit elements for functional genomics (including optogenetics and calcium imaging). Multiple feature selection-based approaches will be particularly useful in the study of OTergic projections to brain regions such as the VTA, which are innervated by mixed populations of magnocellular and parvocellular neurons, as well as to enable studies to strengthen the mechanistic link between cellular and behavioral consequences of *Fmr1* deletion. Ultimately, such studies will lay the foundation for understanding the universal motifs governing the brain circuit organization of sociality.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
 - Lead Contact
 - Materials Availability
 - Data and Code Availability
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Animals
 - Genotyping
- **METHOD DETAILS**
 - Social Conditioned Place Preference
 - Cocaine Conditioned Place Preference
 - Alloprenant Conditioned Place Preference
 - Stereotaxic and tail-vein injections
 - Electrophysiology
 - Cell Isolation, Enrichment, and cDNA Library Preparation
 - *In situ* hybridization chain reaction
 - Immunohistochemistry
 - Histology
- **QUANTIFICATION AND STATISTICAL ANALYSIS**

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.neuron.2020.10.002>.

ACKNOWLEDGMENTS

We thank members of the Dölen and Goff labs for comments on the manuscript, Elizabeth Vincent for histological consultation, and Ben Emert and Arjun Raj (UPenn) for assistance with the HCR 3.0 protocol. PS38 and VA10 antibodies were kind gifts from Dr. Harold Gainer. This work was supported by funds to G.D. (National Institute of Mental Health [NIMH] grant R56MH115177 and NIMH grant R01MH117127; National Institute of Child Health and Human Development [NICHD] grant R01HD98184) and L.A.G. (Chan - Zuckerberg Initiative grant SVCF 2018-183445 and National Science Foundation [NSF] grant IOS-1656592).

AUTHOR CONTRIBUTIONS

E.M.L., A.V.P., G.L.S.-O., L.A.G., and G.D. designed the experiments and wrote the manuscript. E.M.L. and C.D.G. conducted electrophysiology and analyzed the data. E.M.L., R.N., A.V.P., and N.N. conducted behavioral experiments and analyzed the data. E.M.L., A.V.P., and B.B. conducted immunoblotting experiments and analyzed the data. A.V.P. and G.L.S.-O. conducted the scRNA-seq experiments. G.L.S.-O. and L.A.G. analyzed the scRNA-seq data. A.V.P. and G.L.S.-O. performed the HCR smFISH under the direction of I.D. C.J. and I.D. performed smFISH imaging.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: May 6, 2020

Revised: August 21, 2020

Accepted: October 3, 2020

Published: October 27, 2020

SUPPORTING CITATIONS

The following references appear in the Supplemental Information: Castel and Morris (1988); Kamm et al. (1928); Lambert et al. (1993); Ott and Scott (1911); Pedersen et al. (1982); Zhang and Hernández (2013); Zhang et al. (2009); Zhang et al. (2011).

REFERENCES

- Abrahams, B.S., Arking, D.E., Campbell, D.B., Mefford, H.C., Morrow, E.M., Weiss, L.A., Menashe, I., Wadkins, T., Banerjee-Basu, S., and Packer, A. (2013). SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). *Mol. Autism* 4, 36.
- Allen-Brady, K., Miller, J., Matsunami, N., Stevens, J., Block, H., Farley, M., Krasny, L., Pingree, C., Lainhart, J., Leppert, M., et al. (2009). A high-density SNP genome-wide linkage scan in a large autism extended pedigree. *Mol. Psychiatry* 14, 590–600.
- Althammer, F., and Grinevich, V. (2017). Diversity of oxytocin neurons: beyond magno- and parvocellular cell types? *J. Neuroendocrinol.* 140, 874–888.
- Altstein, M., and Gainer, H. (1988). Differential biosynthesis and posttranslational processing of vasopressin and oxytocin in rat brain during embryonic and postnatal development. *J. Neurosci.* 8, 3967–3977.
- Anagnostou, E., Soorya, L., Chaplin, W., Bartz, J., Halpern, D., Wasserman, S., Wang, A.T., Pepa, L., Tanel, N., Kushki, A., and Hollander, E. (2012). Intranasal oxytocin versus placebo in the treatment of adults with autism spectrum disorders: a randomized controlled trial. *Mol. Autism* 3, 16.
- Antoine, M.W., Langberg, T., Schnepel, P., and Feldman, D.E. (2019). Increased excitation-inhibition ratio stabilizes synapse and circuit excitability in four autism mouse models. *Neuron* 101, 648–661.e4.
- Armstrong, W.E., Warach, S., Hatton, G.I., and McNeill, T.H. (1980). Subnuclei in the rat hypothalamic paraventricular nucleus: a cytoarchitectural, horse-radish peroxidase and immunocytochemical analysis. *Neuroscience* 5, 1931–1958.
- Assaf, M., Hyatt, C.J., Wong, C.G., Johnson, M.R., Schultz, R.T., Hendler, T., and Pearlson, G.D. (2013). Mentalizing and motivation neural function during social interactions in autism spectrum disorders. *Neuroimage Clin.* 3, 321–331.
- August, G.J. (1983). A genetic marker associated with infantile autism. *Am. J. Psychiatry* 140, 813.
- Bailey, A., Le Couteur, A., Gottesman, I., Bolton, P., Simonoff, E., Yuzda, E., and Rutter, M. (1995). Autism as a strongly genetic disorder: evidence from a British twin study. *Psychol. Med.* 25, 63–77.
- Bardo, M.T., and Bevins, R.A. (2000). Conditioned place preference: what does it add to our preclinical understanding of drug reward? *Psychopharmacology (Berl.)* 153, 31–43.
- Baron, D., Holland, C.M., Carlson, K., Wolfrum, E., and Thompson, B.L. (2020). Adapting social conditioned place preference for use in young children. *Neurobiol. Learn. Mem.* 172, 107235.
- Bear, M.F., Dölen, G., Osterweil, E., and Nagarajan, N. (2008). Fragile X: translation in action. *Neuropsychopharmacology* 33, 84–87.
- Becht, E., McInnes, L., Healy, J., Dutertre, C.-A.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP. *Nat. Biotechnol.* 37, 38–47.
- Beier, K.T., Steinberg, E.E., DeLoach, K.E., Xie, S., Miyamichi, K., Schwarz, L., Gao, X.J., Kremer, E.J., Malenka, R.C., and Luo, L. (2015). Circuit architecture of VTA dopamine neurons revealed by systematic input-output mapping. *Cell* 162, 622–634.
- Ben-Barak, Y., Russell, J.T., Whitnall, M.H., Ozato, K., and Gainer, H. (1985). Neuropeptide in the hypothalamo-neurohypophyseal system. I. Production and characterization of monoclonal antibodies. *J. Neurosci.* 5, 81–97.
- Bernardet, M., and Crusio, W.E. (2006). *Fmr1* KO mice as a possible model of autistic features. *ScientificWorldJournal* 6, 1164–1176.
- Biag, J., Huang, Y., Gou, L., Hintiryan, H., Askarinam, A., Hahn, J.D., Toga, A.W., and Dong, H.W. (2012). Cyto- and chemoarchitecture of the

- p hypothalamic paraventricular nucleus in the C57BL/6J male mouse: a study of immunostaining and multiple fluorescent tract tracing.
- J. Comp. Neurol.*
- 520, 6–33.
- Brown, W.T., Friedman, E., Jenkins, E.C., Brooks, J., Wisniewski, K., Raguthu, S., and French, J.H. (1982). Association of fragile X syndrome with autism. *Lancet* 1, 100.
- Brown, W.T., Jenkins, E.C., Cohen, I.L., Fisch, G.S., Wolf-Schein, E.G., Gross, A., Waterhouse, L., Fein, D., Mason-Brothers, A., Ritvo, E., et al. (1986). Fragile X and autism: a multicenter survey. *Am. J. Med. Genet.* 23, 341–352.
- Cascio, C.J., Foss-Feig, J.H., Heacock, J.L., Newsom, C.R., Cowan, R.L., Benningfield, M.M., Rogers, B.P., and Cao, A. (2012). Response of neural reward regions to food cues in autism spectrum disorders. *J. Neurodev. Disord.* 4, 9.
- Castel, M., and Morris, J.F. (1988). The neurophysin-containing innervation of the forebrain of the mouse. *Neuroscience* 24, 937–966.
- Chen, R., Wu, X., Jiang, L., and Zhang, Y. (2017). Single-cell RNA-seq reveals hypothalamic cell diversity. *Cell Rep.* 18, 3227–3241.
- Chevallier, C., Kohls, G., Troiani, V., Brodtkin, E.S., and Schultz, R.T. (2012). The social motivation theory of autism. *Trends Cogn. Neurosci.* 16, 231–239.
- Chev  e, M., Robertson, J.J., Cannon, G.H., Brown, S.P., and Goff, L.A. (2018). Variation in activity state, axonal projection, and position define the transcriptional identity of individual neocortical projection neurons. *Cell Rep.* 22, 441–455.
- Chiodera, P., and Legros, J.J. (1981). [Intravenous injection of synthetic oxytocin induces a decrease of cortisol plasma level in normal man]. *C. R. Seances Soc. Biol. Fil.* 175, 546–549.
- Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha, A., and Pierce, N.A. (2018). Third-generation *in situ* hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development* 145, 1–10.
- Clipperton-Allen, A.E., Chen, Y., and Page, D.T. (2016). Autism-relevant behaviors are minimally impacted by conditional deletion of Pten in oxytocinergic neurons. *Autism Res.* 9, 1248–1262.
- Cui, G., Jun, S.B., Jin, X., Pham, M.D., Vogel, S.S., Lovinger, D.M., and Costa, R.M. (2013). Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature* 494, 238–242.
- Dadds, M.R., MacDonald, E., Cauchi, A., Williams, K., Levy, F., and Brennan, J. (2014). Nasal oxytocin for social deficits in childhood autism: a randomized controlled trial. *J. Autism Dev. Disord.* 44, 521–531.
- Daigle, T.L., Madisen, L., Hage, T.A., Valley, M.T., Knoblich, U., Larsen, R.S., Takeno, M.M., Huang, L., Gu, H., Larsen, R., et al. (2018). A suite of transgenic driver and reporter mouse lines with enhanced brain-cell-type targeting and functionality. *Cell* 174, 465–480.e22.
- Damiano, C.R., Alois, J., Treadway, M., Bodfish, J.W., and Dichter, G.S. (2012). Adults with autism spectrum disorders exhibit decreased sensitivity to reward parameters when making effort-based decisions. *J. Neurodev. Disord.* 4, 13.
- Darnell, J.C., Van Driesche, S.J., Zhang, C., Hung, K.Y.S., Mele, A., Fraser, C.E., Stone, E.F., Chen, C., Fak, J.J., Chi, S.W., et al. (2011). FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* 146, 247–261.
- Dayanithi, G., Forostyak, O., Ueta, Y., Verkhatsky, A., and Toescu, E.C. (2012). Segregation of calcium signalling mechanisms in magnocellular neurons and terminals. *Cell Calcium* 51, 293–299.
- de Oliveira, L.F., Camboim, C., Diehl, F., Consiglio, A.R., and Quillfeldt, J.A. (2007). Glucocorticoid-mediated effects of systemic oxytocin upon memory retrieval. *Neurobiol. Learn. Mem.* 87, 67–71.
- Demurie, E., Roeyers, H., Baeyens, D., and Sonuga-Barke, E. (2011). Common alterations in sensitivity to type but not amount of reward in ADHD and autism spectrum disorders. *J. Child Psychol. Psychiatry* 52, 1164–1173.
- Diester, I., Hefti, F., Mansuy, I., Pascual-Leone, A., Robbins, T.W., Rubin, L.L., Sawa, A., Wernig, M., D  len, G., Hyman, S.E., et al. (2015). Bridging the gap between patients and models. In *Translational Neuroscience: Toward New Therapies*, K. Nikolich, S.E. Hyman, and J. Lupp, eds. (MIT Press), pp. 209–244.
- Dijkerman, H.C., and de Haan, E.H.F. (2007). Somatosensory processes subserving perception and action. *Behav. Brain Sci.* 30, 189–201, discussion 201–239.
- D  len, G. (2015). Oxytocin: parallel processing in the social brain? *J. Neuroendocrinol.* 27, 516–535.
- D  len, G., and Bear, M.F. (2009). Fragile X syndrome and autism: from disease model to therapeutic targets. *J. Neurodev. Disord.* 1, 133–140.
- D  len, G., and Malenka, R.C. (2014). The emerging role of nucleus accumbens oxytocin in social cognition. *Biol. Psychiatry* 76, 354–355.
- D  len, G., and Sahin, M. (2016). Editorial: essential pathways and circuits of autism pathogenesis. *Front. Neurosci.* 10, 182.
- D  len, G., Osterweil, E., Rao, B.S.S., Smith, G.B., Auerbach, B.D., Chattarji, S., and Bear, M.F. (2007). Correction of fragile X syndrome in mice. *Neuron* 56, 955–962.
- D  len, G., Darvishzadeh, A., Huang, K.W., and Malenka, R.C. (2013). Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin. *Nature* 501, 179–184.
- D  len, G., Malenka, R., Perlumutter, J., Brose, N., Frackowiak, R., Cuthbert, B., Diester, I., Mansuy, I., Kroker, K., Boekers, T., et al. (2015). Pathophysiological toolkit: genes to circuits. In *Translational Neuroscience: Toward New Therapies*, K. Nikolich, S.E. Hyman, and J. Lupp, eds. (MIT Press), pp. 139–163.
- Dreifuss, J.J. (1975). A review on neurosecretory granules: their contents and mechanisms of release. *Ann. N Y Acad. Sci.* 248, 184–201.
- The Dutch-Belgian Fragile X Consortium (1994). Fmr1 knockout mice: a model to study fragile X mental retardation. *Cell* 78, 23–33.
- Eliava, M., Melchior, M., Knobloch-Bollmann, H.S., Wahis, J., da Silva Gouveia, M., Tang, Y., Ciobanu, A.C., Triana Del Rio, R., Roth, L.C., Althammer, F., et al. (2016). A new population of parvocellular oxytocin neurons controlling magnocellular neuron activity and inflammatory pain processing. *Neuron* 89, 1291–1304.
- Fang, Q., and Wang, J. (2017). Place preferences associated with pups or cocaine change the expression of D2R, V1aR and OTR in the NAcc and MeA and the levels of plasma AVP, OT, T and E2 in mandarin vole fathers. *Psychoneuroendocrinology* 80, 147–154.
- Folstein, S., and Rutter, M. (1977). Genetic influences and infantile autism. *Nature* 265, 726–728.
- Francis, S.M., Sagar, A., Levin-Decanini, T., Liu, W., Carter, C.S., and Jacob, S. (2014). Oxytocin and vasopressin systems in genetic syndromes and neurodevelopmental disorders. *Brain Res.* 1580, 199–218.
- Freitag, C.M. (2007). The genetics of autistic disorders and its clinical relevance: a review of the literature. *Mol. Psychiatry* 12, 2–22.
- Furler, S., Paterna, J.C., Weibel, M., and B  eler, H. (2001). Recombinant AAV vectors containing the foot and mouth disease virus 2A sequence confer efficient bicistronic gene expression in cultured cells and rat substantia nigra neurons. *Gene Ther.* 8, 864–873.
- Gabel, L.A., Won, S., Kawai, H., McKinney, M., Tartakoff, A.M., and Fallon, J.R. (2004). Visual experience regulates transient expression and dendritic localization of fragile X mental retardation protein. *J. Neurosci.* 24, 10579–10583.
- Grabrucker, A.M., Schmeisser, M.J., Schoen, M., and Boeckers, T.M. (2011). Postsynaptic ProSAP/Shank scaffolds in the cross-hair of synaptopathies. *Trends Cell Biol.* 21, 594–603.
- Guastella, A.J., Gray, K.M., Rinehart, N.J., Alvares, G.A., Tonge, B.J., Hickie, I.B., Keating, C.M., Cacciotti-Sajja, C., and Einfeld, S.L. (2015). The effects of a course of intranasal oxytocin on social behaviors in youth diagnosed with autism spectrum disorders: a randomized controlled trial. *J. Child Psychol. Psychiatry* 56, 444–452.
- Gurdjian, E.S. (1927). The diencephalon of the albino rat: studies on the brain of the rat. No. 2. *J. Comp. Neurol.* 43, 1–114.

- Haberly, L.B. (2001). Parallel-distributed processing in olfactory cortex: new insights from morphological and physiological analysis of neuronal circuitry. *Chem. Senses* 26, 551–576.
- Hall, S.S., Lightbody, A.A., McCarthy, B.E., Parker, K.J., and Reiss, A.L. (2012). Effects of intranasal oxytocin on social anxiety in males with fragile X syndrome. *Psychoneuroendocrinology* 37, 509–518.
- Hammock, E.A.D., and Levitt, P. (2013). Oxytocin receptor ligand binding in embryonic tissue and postnatal brain development of the C57BL/6J mouse. *Front. Behav. Neurosci.* 7, 195.
- Hoffman, N.W., Tasker, J.G., and Dudek, F.E. (1991). Immunohistochemical differentiation of electrophysiologically defined neuronal populations in the region of the rat hypothalamic paraventricular nucleus. *J. Comp. Neurol.* 307, 405–416.
- Hollander, E., Bartz, J., Chaplin, W., Phillips, A., Sumner, J., Soorya, L., Anagnostou, E., and Wasserman, S. (2007). Oxytocin increases retention of social cognition in autism. *Biol. Psychiatry* 61, 498–503.
- Hosoya, Y., and Matsushita, M. (1979). Identification and distribution of the spinal and hypophyseal projection neurons in the paraventricular nucleus of the rat. A light and electron microscopic study with the horseradish peroxidase method. *Exp. Brain Res.* 35, 315–331.
- Hovey, D., Zettergren, A., Jonsson, L., Melke, J., Anckarsäter, H., Lichtenstein, P., and Westberg, L. (2014). Associations between oxytocin-related genes and autistic-like traits. *Soc. Neurosci.* 9, 378–386.
- Hung, L.W., Neuner, S., Polepalli, J.S., Beier, K.T., Wright, M., Walsh, J.J., Lewis, E.M., Luo, L., Deisseroth, K., Dölen, G., and Malenka, R.C. (2017). Gating of social reward by oxytocin in the ventral tegmental area. *Science* 357, 1406–1411.
- Kamm, O., Aldrich, T.B., Grote, I.W., Rowe, L.W., and Bugbee, E.P. (1928). The active principles of the pituitary gland. *J. Am. Chem. Soc.* 50, 573–598.
- Kanehisa, M., and Goto, S. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.* 28, 27–30.
- Kanner, L. (1943). Autistic disturbances of affective contact. *Nerv. Child J. Psychopathol. Psychother. Ment. Hyg. Guid. Child* 2, 217–250.
- Katz, L.C., Burkhalter, A., and Dreyer, W.J. (1984). Fluorescent latex microspheres as a retrograde neuronal marker for in vivo and in vitro studies of visual cortex. *Nature* 310, 498–500.
- Kaufmann, W.E., Kidd, S.A., Andrews, H.F., Budimirovic, D.B., Esler, A., Haas-Givler, B., Stackhouse, T., Riley, C., Peacock, G., Sherman, S.L., et al. (2017). Autism spectrum disorder in fragile X syndrome: cooccurring conditions and current treatment. *Pediatrics* 139 (Suppl 3), S194–S206.
- Kelly, E., Meng, F., Fujita, H., Morgado, F., Kazemi, Y., Rice, L.C., Ren, C., Escamilla, C.O., Gibson, J.M., Sajadi, S., et al. (2020). Regulation of autism-relevant behaviors by cerebellar-prefrontal cortical circuits. *Nat. Neurosci.* 23, 1102–1110.
- Klatzo, I. (2003). Cécile & Oskar Vogt: the significance of their contributions in modern neuroscience. *Acta Neurochir. Suppl. (Wien)* 86, 29–32.
- Knobloch, H.S., Charlet, A., Hoffmann, L.C., Eliava, M., Khrulev, S., Cetin, A.H., Osten, P., Schwarz, M.K., Seeburg, P.H., Stoop, R., and Grinovich, V. (2012). Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* 73, 553–566.
- Kravitz, A.V., Tye, L.D., and Kreitzer, A.C. (2012). Distinct roles for direct and indirect pathway striatal neurons in reinforcement. *Nat. Neurosci.* 15, 816–818.
- Kremer, E.J., Boutin, S., Chillon, M., and Danos, O. (2000). Canine adenovirus vectors: an alternative for adenovirus-mediated gene transfer. *J. Virol.* 74, 505–512.
- Krieg, W.J.S. (1932). The hypothalamus of the albino rat. *J. Comp. Neurol.* 55, 19–89.
- Lambert, R.C., Moos, F.C., and Richard, P. (1993). Action of endogenous oxytocin within the paraventricular or supraoptic nuclei: a powerful link in the regulation of the bursting pattern of oxytocin neurons during the milk-ejection reflex in rats. *Neuroscience* 57, 1027–1038.
- Lengros, J.J., Chiodera, P., Geenen, V., Smits, S., and von Frenckell, R. (1984). Dose-response relationship between plasma oxytocin and cortisol and adrenocorticotropin concentrations during oxytocin infusion in normal men. *J. Clin. Endocrinol. Metab.* 58, 105–109.
- Leng, G., and Ludwig, M. (2016). Intranasal oxytocin: myths and delusions. *Biol. Psychiatry* 79, 243–250.
- Lobo, M.K., and Nestler, E.J. (2011). The striatal balancing act in drug addiction: distinct roles of direct and indirect pathway medium spiny neurons. *Front. Neuroanat.* 5, 41.
- Ludwig, M., and Leng, G. (2006). Dendritic peptide release and peptide-dependent behaviours. *Nat. Rev. Neurosci.* 7, 126–136.
- Luther, J.A., and Tasker, J.G. (2000). Voltage-gated currents distinguish parvocellular from magnocellular neurones in the rat hypothalamic paraventricular nucleus. *J. Physiol.* 523, 193–209.
- Luther, J.A., Halmos, K.C., and Tasker, J.G. (2000). A slow transient potassium current expressed in a subset of neurosecretory neurons of the hypothalamic paraventricular nucleus. *J. Neurophysiol.* 84, 1814–1825.
- Luther, J.A., Daftary, S.S., Boudaba, C., Gould, G.C., Halmos, K.C., and Tasker, J.G. (2002). Neurosecretory and non-neurosecretory parvocellular neurones of the hypothalamic paraventricular nucleus express distinct electrophysiological properties. *J. Neuroendocrinol.* 14, 929–932.
- Martin, J.P., and Bell, J. (1943). A pedigree of mental defect showing sex-linkage. *J. Neurol. Psychiatry* 6, 154–157.
- Maynard, K.R., Hobbs, J.W., Phan, B.N., Gupta, A., Rajpurohit, S., Williams, C., Rajpurohit, A., Shin, J.H., Jaffe, A.E., and Martinowich, K. (2018). BDNF-TrkB signaling in oxytocin neurons contributes to maternal behavior. *eLife* 7, e33676.
- Melis, M.R., Argiolas, A., and Gessa, G.L. (1986). Oxytocin-induced penile erection and yawning: site of action in the brain. *Brain Res.* 398, 259–265.
- Merchenthaler, I. (1991). Neurons with access to the general circulation in the central nervous system of the rat: a retrograde tracing study with Fluoro-Gold. *Neuroscience* 44, 655–662.
- Mientjes, E.J., Nieuwenhuizen, I., Kirkpatrick, L., Zu, T., Hoogeveen-Westerveld, M., Severijnen, L., Rifé, M., Willemsen, R., Nelson, D.L., and Oostra, B.A. (2006). The generation of a conditional Fmr1 knock out mouse model to study Fmrp function in vivo. *Neurobiol. Dis.* 27, 549–555.
- Miyoshi, G., Hjerling-Leffler, J., Karayannis, T., Sousa, V.H., Butt, S.J., Battiste, J., Johnson, J.E., Machold, R.P., and Fishell, G. (2010). Genetic fate mapping reveals that the caudal ganglionic eminence produces a large and diverse population of superficial cortical interneurons. *J. Neurosci.* 30, 1582–1594.
- Naber, F.B.A., Swinkels, S.H.N., Buitelaar, J.K., Dietz, C., van Daalen, E., Bakermans-Kranenburg, M.J., van Ijzendoorn, M.H., and van Engeland, H. (2007). Joint attention and attachment in toddlers with autism. *J. Abnorm. Child Psychol.* 35, 899–911.
- Naber, F.B.A., Bakermans-Kranenburg, M.J., van Ijzendoorn, M.H., Swinkels, S.H.N., Buitelaar, J.K., Dietz, C., van Daalen, E., and van Engeland, H. (2008). Play behavior and attachment in toddlers with autism. *J. Autism Dev. Disord.* 38, 857–866.
- Nardou, R., Lewis, E.M., Rothhaas, R., Xu, R., Yang, A., Boyden, E., and Dölen, G. (2019). Oxytocin-dependent reopening of a social reward learning critical period with MDMA. *Nature* 569, 116–120.
- Nassi, J.J., and Callaway, E.M. (2009). Parallel processing strategies of the primate visual system. *Nat. Rev. Neurosci.* 10, 360–372.
- Nestler, E.J., and Hyman, S.E. (2010). Animal models of neuropsychiatric disorders. *Nat. Neurosci.* 13, 1161–1169.
- Nigro, M.J., Hashikawa-Yamasaki, Y., and Rudy, B. (2018). Diversity and connectivity of layer 5 somatostatin-expressing interneurons in the mouse barrel cortex. *J. Neurosci.* 38, 1622–1633.
- Nordmann, J.J., and Morris, J.F. (1984). Method for quantitating the molecular content of a subcellular organelle: hormone and neurophysin content of newly formed and aged neurosecretory granules. *Proc. Natl. Acad. Sci. U S A* 81, 180–184.

- Orefice, L.L., Mosko, J.R., Morency, D.T., Wells, M.F., Tasnim, A., Mozeika, S.M., Ye, M., Chirila, A.M., Emanuel, A.J., Rankin, G., et al. (2019). Targeting peripheral somatosensory neurons to improve tactile-related phenotypes in ASD models. *Cell* 178, 867–886.e24.
- Ott, I., and Scott, J.C. (1911). The action of infundibulin upon the mammary secretion. *Proc. Soc. Exp. Biol. Med.* 8, 48–49.
- Page, D.T., Kuti, O.J., Prestia, C., and Sur, M. (2009). Haploinsufficiency for Pten and Serotonin transporter cooperatively influences brain size and social behavior. *Proc. Natl. Acad. Sci. U S A* 106, 1989–1994.
- Panksepp, J.B., and Lahvis, G.P. (2007). Social reward among juvenile mice. *Genes Brain Behav.* 6, 661–671.
- Paul, B.D., Sbodio, J.I., Xu, R., Vandiver, M.S., Cha, J.Y., Snowman, A.M., and Snyder, S.H. (2014). Cystathionine γ -lyase deficiency mediates neurodegeneration in Huntington's disease. *Nature* 509, 96–100.
- Pedersen, C.A., Ascher, J.A., Monroe, Y.L., and Prange, A.J., Jr. (1982). Oxytocin induces maternal behavior in virgin female rats. *Science* 216, 648–650.
- Peñagarikano, O., Lázaro, M.T., Lu, X.-H., Gordon, A., Dong, H., Lam, H.A., Peles, E., Maidment, N.T., Murphy, N.P., Yang, X.W., et al. (2015). Exogenous and evoked oxytocin restores social behavior in the Cntnap2 mouse model of autism. *Sci. Transl. Med.* 7, 271ra8.
- Penzo, M.A., Robert, V., Tucciarone, J., De Bundel, D., Wang, M., Van Aelst, L., Darvas, M., Parada, L.F., Palmiter, R.D., He, M., et al. (2015). The paraventricular thalamus controls a central amygdala fear circuit. *Nature* 519, 455–459.
- Petersson, M. (2002). Cardiovascular effects of oxytocin. *Prog. Brain Res.* 139, 281–288.
- Picelli, S., Faridani, O.R., Björklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R. (2014). Full-length RNA-seq from single cells using Smart-seq2. *Nat. Protoc.* 9, 171–181.
- Pow, D.V., and Morris, J.F. (1989). Dendrites of hypothalamic magnocellular neurons release neurohypophysial peptides by exocytosis. *Neuroscience* 32, 435–439.
- Qin, M., Kang, J., Burlin, T.V., Jiang, C., and Smith, C.B. (2005). Postadolescent changes in regional cerebral protein synthesis: an in vivo study in the FMR1 null mouse. *J. Neurosci.* 25, 5087–5095.
- Qiu, X., Hill, A., Packer, J., Lin, D., Ma, Y.-A., and Trapnell, C. (2017). Single-cell mRNA quantification and differential analysis with Censur. *Nat. Methods* 14, 309–315.
- Raj, A., van den Bogaard, P., Rifkin, S.A., van Oudenaarden, A., and Tyagi, S. (2008). Imaging individual mRNA molecules using multiple singly labeled probes. *Nat. Methods* 5, 877–879.
- Recanzone, G.H., and Cohen, Y.E. (2010). Serial and parallel processing in the primate auditory cortex revisited. *Behav. Brain Res.* 206, 1–7.
- Romanov, R.A., Zeisel, A., Bakker, J., Girach, F., Hellysaz, A., Tomer, R., Alpár, A., Mulder, J., Clotman, F., Keimpema, E., et al. (2017). Molecular interrogation of hypothalamic organization reveals distinct dopamine neuronal subtypes. *Nat. Neurosci.* 20, 176–188.
- Roper, S.D. (2009). Parallel processing in mammalian taste buds? *Physiol. Behav.* 97, 604–608.
- Rutgers, A.H., Bakermans-Kranenburg, M.J., van Ijzendoorn, M.H., and van Berckelaer-Onnes, I.A. (2004). Autism and attachment: a meta-analytic review. *J. Child Psychol. Psychiatry* 45, 1123–1134.
- Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.Y., Peng, M., Collins, R., Grove, J., Klei, L., et al.; Autism Sequencing Consortium; iPSYCH-Broad Consortium (2020). Large-scale exome sequencing study implicates both developmental and functional changes in the neurobiology of autism. *Cell* 180, 568–584.e23.
- Sawchenko, P.E., and Swanson, L.W. (1982). Immunohistochemical identification of neurons in the paraventricular nucleus of the hypothalamus that project to the medulla or to the spinal cord in the rat. *J. Comp. Neurol.* 205, 260–272.
- Saxena, A., Wagatsuma, A., Noro, Y., Kuji, T., Asaka-Oba, A., Watahiki, A., Gurnot, C., Fagiolini, M., Hensch, T.K., and Carninci, P. (2012). Trehalose-enhanced isolation of neuronal sub-types from adult mouse brain. *Biotechniques* 52, 381–385.
- Scelsi, M.A., Khan, R.R., Lorenzi, M., Christopher, L., Greicius, M.D., Schott, J.M., Ourselin, S., and Altmann, A. (2018). Genetic study of multimodal imaging Alzheimer's disease progression score implicates novel loci. *Brain* 141, 2167–2180.
- Schmued, L.C., and Fallon, J.H. (1986). Fluoro-Gold: a new fluorescent retrograde axonal tracer with numerous unique properties. *Brain Res.* 377, 147–154.
- Schneeberger, M., Parolari, L., Das Banerjee, T., Bhawe, V., Wang, P., Patel, B., Topilko, T., Wu, Z., Choi, C.H.J., Yu, X., et al. (2019). Regulation of energy expenditure by brainstem GABA neurons. *Cell* 178, 672–685.e12.
- Smith, L.N., Jedynak, J.P., Fontenot, M.R., Hale, C.F., Dietz, K.C., Taniguchi, M., Thomas, F.S., Zirlin, B.C., Birnbaum, S.G., Huber, K.M., et al. (2014). Fragile X mental retardation protein regulates synaptic and behavioral plasticity to repeated cocaine administration. *Neuron* 82, 645–658.
- Stachowiak, A., Macchi, C., Nussdorfer, G.G., and Malendowicz, L.K. (1995). Effects of oxytocin on the function and morphology of the rat adrenal cortex: in vitro and in vivo investigations. *Res. Exp. Med. (Berl.)* 195, 265–274.
- Stavropoulos, K.K.M., and Carver, L.J. (2014). Reward anticipation and processing of social versus nonsocial stimuli in children with and without autism spectrum disorders. *Psychiatry Allied Discip.* 55, 1398–1408.
- Steffenburg, S., Gillberg, C., Hellgren, L., Andersson, L., Gillberg, I.C., Jakobsson, G., and Bohman, M. (1989). A twin study of autism in Denmark, Finland, Iceland, Norway and Sweden. *J. Child Psychol. Psychiatry* 30, 405–416.
- Stokes, M., Newton, N., and Kaur, A. (2007). Stalking, and social and romantic functioning among adolescents and adults with autism spectrum disorder. *J. Autism Dev. Disord.* 37, 1969–1986.
- Suhl, J.A., and Warren, S.T. (2015). Single-nucleotide mutations in FMR1 reveal novel functions and regulatory mechanisms of the fragile X syndrome protein FMRP. *J. Exp. Neurosci.* 9 (Suppl 2), 35–41.
- Sullivan, P.F., Daly, M.J., and O'Donovan, M. (2012). Genetic architectures of psychiatric disorders: the emerging picture and its implications. *Nat. Rev. Genet.* 13, 537–551.
- Swaab, D.F., Purba, J.S., and Hofman, M.A. (1995). Alterations in the hypothalamic paraventricular nucleus and its oxytocin neurons (putative satiety cells) in Prader-Willi syndrome: a study of five cases. *J. Clin. Endocrinol. Metab.* 80, 573–579.
- Swanson, L.W., and Kuypers, H.G.J.M. (1980). The paraventricular nucleus of the hypothalamus: cytoarchitectonic subdivisions and organization of projections to the pituitary, dorsal vagal complex, and spinal cord as demonstrated by retrograde fluorescence double-labeling methods. *J. Comp. Neurol.* 194, 555–570.
- Tasker, J.G., and Dudek, F.E. (1991). Electrophysiological properties of neurons in the region of the paraventricular nucleus in slices of rat hypothalamus. *J. Physiol.* 434, 271–293.
- Tervo, D.G.R., Hwang, B.Y., Viswanathan, S., Gaj, T., Lavzin, M., Ritola, K.D., Lindo, S., Michael, S., Kuleshova, E., Ojala, D., et al. (2016). A designer AAV variant permits efficient retrograde access to projection neurons. *Neuron* 92, 372–382.
- Thurman, A.J., McDuffie, A., Hagerman, R., and Abbeduto, L. (2014). Psychiatric symptoms in boys with fragile X syndrome: a comparison with non-syndromic autism spectrum disorder. *Res. Dev. Disabil.* 35, 1072–1086.
- Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J., Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat. Biotechnol.* 32, 381–386.
- Tsai, P.T., Hull, C., Chu, Y., Greene-Colozzi, E., Sadowski, A.R., Leech, J.M., Steinberg, J., Crawley, J.N., Regehr, W.G., and Sahin, M. (2012). Autistic-like

behaviour and cerebellar dysfunction in Purkinje cell Tsc1 mutant mice. *Nature* 488, 647–651.

Uematsu, A., Tan, B.Z., Ycu, E.A., Cuevas, J.S., Koivumaa, J., Junyent, F., Kremer, E.J., Witten, I.B., Deisseroth, K., and Johansen, J.P. (2017). Modular organization of the brainstem noradrenergic system coordinates opposing learning states. *Nat. Neurosci.* 20, 1602–1611.

Valassi, E., Scacchi, M., and Cavagnini, F. (2008). Neuroendocrine control of food intake. *Nutr. Metab. Cardiovasc. Dis.* 18, 158–168.

van den Pol, A.N. (1982). The magnocellular and parvocellular paraventricular nucleus of rat: intrinsic organization. *J. Comp. Neurol.* 206, 317–345.

van den Pol, A.N. (2012). Neuropeptide transmission in brain circuits. *Neuron* 76, 98–115.

Van Driesche, S.J., Sawicka, K., Zhang, C., Hung, S.K.Y., Park, C.Y., Fak, J.J., Yang, C., Darnell, R.B., and Darnell, J.C. (2019). FMRP binding to a ranked subset of long genes is revealed by coupled CLIP and TRAP in specific neuronal cell types. *bioRxiv*. <https://doi.org/10.1101/762500>.

Verkerk, A.J.M.H., Pieretti, M., Sutcliffe, J.S., Fu, Y.H., Kuhl, D.P.A., Pizzuti, A., Reiner, O., Richards, S., Victoria, M.F., Zhang, F.P., et al. (1991). Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 65, 905–914.

Vivanti, G., and Nuske, H.J. (2017). Autism, attachment, and social learning: three challenges and a way forward. *Behav. Brain Res.* 325 (Pt B), 251–259.

Whitnall, M.H., Key, S., Ben-Barak, Y., Ozato, K., and Gainer, H. (1985). Neurophysin in the hypothalamo-neurohypophyseal system. II. Immunocytochemical studies of the ontogeny of oxytocinergic and vasopressinergic neurons. *J. Neurosci.* 5, 98–109.

Willsey, A.J., Sanders, S.J., Li, M., Dong, S., Tebbenkamp, A.T., Muhle, R.A., Reilly, S.K., Lin, L., Fertuzinhos, S., Miller, J.A., et al. (2013). Coexpression networks implicate human midfetal deep cortical projection neurons in the pathogenesis of autism. *Cell* 155, 997–1007.

Wu, Z., Xu, Y., Zhu, Y., Sutton, A.K., Zhao, R., Lowell, B.B., Olson, D.P., and Tong, Q. (2012). An obligate role of oxytocin neurons in diet induced energy expenditure. *PLoS ONE* 7, e45167.

Xiao, L., Priest, M.F., Nasenbeny, J., Lu, T., and Kozorovitskiy, Y. (2017). Biased oxytocinergic modulation of midbrain dopamine systems. *Neuron* 95, 368–384.e5.

Zhang, L., and Hernández, V.S. (2013). Synaptic innervation to rat hippocampus by vasopressin-immuno-positive fibres from the hypothalamic supraoptic and paraventricular nuclei. *Neuroscience* 228, 139–162.

Zhang, Z., Bhalla, A., Dean, C., Chapman, E.R., and Jackson, M.B. (2009). Synaptotagmin IV: a multifunctional regulator of peptidergic nerve terminals. *Nat. Neurosci.* 12, 163–171.

Zhang, G., Bai, H., Zhang, H., Dean, C., Wu, Q., Li, J., Guariglia, S., Meng, Q., and Cai, D. (2011). Neuropeptide exocytosis involving synaptotagmin-4 and oxytocin in hypothalamic programming of body weight and energy balance. *Neuron* 69, 523–535.

Zhang, R., Zhang, H.F., Han, J.S., and Han, S.P. (2017). Genes related to oxytocin and arginine-vasopressin pathways: associations with autism spectrum disorders. *Neurosci. Bull.* 33, 238–246.

Zwergal, A., Strupp, M., Brandt, T., and Büttner-Ennever, J.A. (2009). Parallel ascending vestibular pathways: anatomical localization and functional specialization. *Ann. N Y Acad. Sci.* 1164, 51–59.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse anti-OT neurophysin	Whitnall et al., 1985 Ben-Barak et al., 1985 ; gift from Harold Gainer, PhD	PS38; RRID: AB_2315026488
Alexa 488 donkey anti-mouse	Life Technologies	A-21202; RRID: AB_141607647
Alexa 647 donkey anti-mouse	Life Technologies	A-31571; RRID: AB_2535804350
Alexa 647 goat anti-mouse	Life Technologies	A-21235; RRID: AB_2535804350
Alexa 350 goat anti-mouse	Life Technologies	A-11045; RRID: AB_2534100
Rabbit anti-FluoroGold	Fluorochrome	Cat# FG; RRID: AB_2314407647
Alexa 647 donkey anti-rabbit	Life Technologies	A-31573; RRID: AB_2536183
Chicken anti-GFP	ABCAM	Ab13970; RRID: AB_300798488
Alexa 488 goat anti-chicken	Life Technologies	A-11039; RRID: AB_2534096
Goat anti-mCherry	SICGEN	AB0040; RRID: AB_2333093555
Alexa 555 donkey anti-goat	Life Technologies	A-21432; RRID: AB_2535853
Mouse anti-FMRP	Developmental Studies Hybridoma Bank	2F5-1; RRID: AB_10805421
Alexa 647 goat anti-mouse IgG2b	Jackson ImmunoResearch	115-607-187; RRID: AB_2632546
Rabbit anti-OT	Altstein and Gainer, 1988 ; gift from Harold Gainer, PhD	VA10, Gainer; not found on scicrunch
Alexa 488 donkey anti-rabbit	Life Technologies	A-21206; RRID: AB_2535792
Bacterial and Virus Strains		
rgAAV-EF1a-fDIO-Cre	Schneeberger et al., 2019	Addgene 121675-AAVrg
rgAAV-CAG-fDIO-GFP-Cre	Stanford University Neuroscience Gene Vector and Virus Core	N/A
CAV-2-GFP	Plateforme de Vectorologie de Montpellier	N/A
CAV-2-Cre-GFP	Plateforme de Vectorologie de Montpellier	N/A
Chemicals, Peptides, and Recombinant Proteins		
Neurobiotin	Vector Laboratories	SP-1120
Alexa 350 streptavidin conjugate	Life Technologies	S11249
FluoroGold	Fluorochrome	N/A
Red Retrobeads	Lumafluor	N/A
4-Aminopyridine	Sigma-Aldrich	275875
Cocaine hydrochloride	Sigma-Aldrich	C5776
Critical Commercial Assays		
Worthington Papain Dissociation System	Worthington Biochemical Corporation	Cat#LK003182
Illumina TruSeq kit	Illumina Cat#RS-122-2001	Cat#RS-122-200
Deposited Data		
Accession number GSE147092	GEO	
Experimental Models: Organisms/Strains		
C57BL6/J	Jackson Laboratories	Stock # 000664
<i>Calb1</i> -IRES-Cre	Jackson Laboratories	Stock # 028532
Flp-dependent GFP reporter mice	Mutant Mouse Resource Center	Stock # 32038

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Ai9 Cre-dependent TdTomato reporter mice	Jackson Laboratories	Stock # 007909
Constitutive <i>Fmr1</i> knockout	Jackson Laboratories	Stock # 003025
Conditional <i>Fmr1</i> knockout	Mientjes et al., 2006	N/A
Oxytocin-2A-Flp-optimized	Nardou et al., 2019	N/A
Oligonucleotides		
see supplemental tables (HCR)		
Recombinant DNA		
fdIO-Cre-GFP	Penzo et al., 2015	Dr. Bo Li and Dr. Linda Van Aelst
Software and Algorithms		
R version 3.5	The R project	https://www.r-project.org/
MATLAB	Mathworks	https://www.mathworks.com/
Prism 5	GraphPad	https://www.graphpad.com/
Igor Pro	WaveMetrics	https://www.wavemetrics.com/
Recording Artist (plugin for Igor Pro)	Richard C Gerkin, PhD	https://github.com/rgerkin/recording-artist

RESOURCE AVAILABILITY**Lead Contact**

Further requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Gül Dölen (gul@jhu.edu).

Materials Availability

OT-2A-Flp mice will be made available with a materials transfer agreement.

Data and Code Availability

The accession number for the scRNA-Seq data reported in this paper is NCBI GEO: GSE147092). Additionally, raw and processed scRNA-Seq data, and code to reproduce the complete analysis, are also publicly available at https://github.com/gofflab/OT_neuron_study_2020.

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Animals**

Experiments were conducted in 3 - 7 week-old male mice. Wild-type C57BL6/J (Stock # 000664; 3 - 4 weeks old) and *Calb1*-IRES-Cre (Stock # 028532) mice were obtained from Jackson Laboratories. Flp-dependent GFP reporter mice (Stock # 32038) were obtained from the Mutant Mouse Resource Center. Transgenic mice were bred in-house and weaned at 3 weeks of age. OT-2A-Flp-optimized (OT-2A-Flp) knockin mice were generated by Cyagen Biosciences (Santa Clara, CA) (Figure S1) (Nardou et al., 2019). Similar to the strategy employed by the OT-IRES-Cre knockin mouse (Wu et al., 2012) the stop codon in exon 3 of the endogenous OT gene was replaced with a 2A-Flp-optimized (FlpO) construct. The co-translational cleavage 2A peptide strategy (known to have a more robust expression of its downstream gene compared to IRES (Furler et al., 2001)) has been used to separate Flp and OT, since Flp is directed to the nucleus, while OT is directed to the cytoplasm. OT-2A-Flp mice were kept on a C57BL/6 background. Despite these designed-in safeguards against disrupting native OT function, mice homozygous for the OT-2A-Flp knockin, like the OT-IRES-Cre mice, exhibit an impaired lactation phenotype. The mice are otherwise phenotypically normal. Therefore, in order to control for possible effects of knockin on *Oxt* gene function, all OT-Flp mice used in these experiments were heterozygous, leaving one WT copy of the *Oxt* gene.

To generate hemizygous *Fmr1* KO males for behavior, homozygous *Fmr1* KO females were bred to hemizygous *Fmr1* KO males. The same strategy was used to generate Cre-dependent conditional *Fmr1* KO (*cdFmr1* KO) males. To generate hemizygous *Fmr1* KO males for cocaine CPP, homozygous *Fmr1* KO females were bred to OT-2A-Flp::fdGFP homozygous males. To generate OT-2A-Flp::Ai9 and OT-2A-Flp::cd*Fmr1* KO mice, homozygous OT-2A-Flp males were crossed with homozygous Ai9 or *cdFmr1* KO females respectively. To generate *Calb1*-Cre::cd*Fmr1* KO males for behavior, homozygous *Calb1*-IRES-Cre males were crossed with homozygous *cdFmr1* KO females. Homozygous or heterozygous OT-2A-Flp and fdGFP mice were crossed to generate OT-2A-Flp::fdGFP mice. All mice were maintained on a 12:12h natural light dark cycle, starting at

7:30am with food and water *ad libitum*. All procedures were conducted in accordance with protocols approved by the Johns Hopkins Animal Care and Use Committee

Genotyping

OT-2A-Flp knockin mice, fdGFP, *Fmr1* KO, and *cdFmr1* KO mice were genotyped using polymerase chain reaction (PCR) analysis of DNA isolated from tail snips taken before weaning. For OT-2A-Flp mice, the wild-type allele was identified by a 622-bp PCR product and the mutant allele by a 308-bp PCR product using the primers mOxtR1: TCCGACAATTAGACACCAGTCAA; mOxtF1: CTACCTG AGCAGCTACATCAACAG; mOxtF2: AGGGCTTTGGGAAGTGTAGGCT. The reactions were run under the following conditions: 94°C × 3 min, (94°C × 30 s, 60°C × 35 s, 72°C × 35 s) × 38 cycles, 72°C × 5 min. For fdGFP mice, the wild-type allele was identified by a 603-bp PCR product, and the mutant allele by a 320-bp PCR product using the following primers; mutant: CCA GGC GGG CCA TTT ACC GTA AG; common: AAA GTC GCT CTG AGT TGT TAT; wild-type: GGA GCG GGA GAA ATG GAT ATG. The reactions were run using the Gt(ROSA)26Sor^{tm(CAG)} protocol published by Jackson Laboratories. For the *Fmr1* KO mice, the wild-type allele was identified by a 131-bp PCR product, and the mutant allele by a 400-bp PCR product using the following primers; wild-type: TGT GAT AGA ATA TGC AGC ATG TGA; common: CTT CTG GCA CCT CCA GCT T; mutant: CAC GAG ACT AGT GAG ACG TG. The reactions were run using the *Fmr1*^{tm1Cgr} protocol published by Jackson Laboratories. For *cdFmr1* KO mice, the wild-type allele was identified by a 120-bp PCR product, and the mutant allele was detected by a 220-bp PCR product using the following primers: CCC ACT GGG AGA GGA TTA TTT GGG and GTT GAG CGG CCG AGT TTG TGA G. Reactions were run under the following conditions: 95°C × 3 min, (95°C × 30 s, 55°C × 30 s, 72°C × 60 s) × 35 cycles, 72°C × 7 min.

METHOD DETAILS

Social Conditioned Place Preference

The sCPP assay (Figure S4) was conducted as previously described (Nardou et al., 2019). Briefly, animals were weaned and socially housed (3–5 same sex cage mates based on unpublished observations that group number does not impact sCPP magnitude within this range) in a cage containing corncob bedding (Anderson Cob, 1/4" cob or 1/8" cob; Animal Specialties and Provisions). At approximately 6 weeks of age, animals were placed individually in one of five open field activity chambers (ENV-510, Med Associates) employing infrared beams and a software interface (Activity Monitor, Med Associates) to monitor mouse position. The chamber was partitioned into two equally sized zones using a clear Plexiglas wall with a hole at floor level to allow the mouse to pass freely between zones. Each zone contained one type of novel bedding (Alpha-Dri; Animal Specialties and Provisions, Kaytee Soft Granule; Petco, Anderson Cob, 1/4" cob; Animal Specialties and Provisions, or Aspen Chip; Northeastern Products). To establish each animal's baseline preference for the bedding cues, the amount of time each mouse spent exploring each zone was recorded during a 30 min pre-conditioning trial. Immediately after this trial, mice received social conditioning with cage mates for 24 hr on one type of bedding used in the pre-conditioning trial, followed by 24 hr isolation conditioning on the other bedding used in the pre-conditioning trial. Immediately following isolation conditioning, a 30 min post-conditioning trial was conducted to establish preference for the two conditioned cues. Note that clean bedding was used for each phase of testing. Chamber assignments were counterbalanced for side and bedding cues. Exclusion criteria for this behavior were defined as a pre-conditioning preference score $\left(\frac{\text{time side 1}}{\text{total time}} \right)$ of > 1.5 or < 0.5. Experimental conditions were compared using normalized social preference scores $(\text{social zone post} / \text{social zone pre})$, and subtracted social preference scores $((\text{social zone post} - \text{social zone pre}) / 900)$ (Dölen et al., 2013; Nardou et al., 2019).

Cocaine Conditioned Place Preference

The protocol for cocaine CPP was conducted as previously described (Nardou et al., 2019). Experiments were performed in the same open field activity chambers as sCPP using an identical configuration. After 3 days of habituation to i.p. saline injections in the home cage, the pre-conditioning trial was performed as stated above. After 24 h, mice received an i.p. injection of cocaine (20 mg/kg) immediately followed by 30 min conditioning on one bedding (Soft Granule or Alpha Dri), which was randomly assigned in a counter-balanced fashion. A second 30 min conditioning session was conducted 24 h later on the other bedding after an i.p. injection of saline (equal volume to cocaine). A 30 min post-conditioning test session was conducted 24 h later to determine each mouse's preference for the cocaine versus saline associated beddings.

Alloparent Conditioned Place Preference

The alloparent CPP assay was adapted from a previously used assay (Fang and Wang, 2017), and experiments were conducted in the same open field activity chambers using an identical configuration to the sCPP assay. To generate familiar alloparent stimulus animals, adult (> 9 weeks old) virgin wild-type C57BL6/J females were pair housed with pregnant WT or *Fmr1* KO females. This housing configuration remained intact following the birth of WT or *Fmr1* KO pups until alloparent CPP testing began. Alloparent conditioning was conducted as follows. Beginning at postnatal day 19–20, and continuing until each male littermate was tested, male mice were individually weaned from their home cage and placed directly in an activity chamber with two novel beddings. To establish each an-

imal's baseline preference for the novel bedding cues, the amount spent exploring each zone of the activity chamber was recorded during a 30 min pre-conditioning trial. Immediately after this trial, mice (postnatal day 19-24) individually received alloparent conditioning with their virgin female alloparent for 24 hr on one type of bedding, followed by 24 hr isolation conditioning on the other bedding. Immediately following the isolation conditioning, a 30 min post-conditioning trial was conducted to establish preference for the two conditioned cues. This process was repeated individually until each male littermate had been tested. Identical to the methods used for sCPP, chamber assignments were counterbalanced for side and bedding cues. Exclusion criteria for this behavior were defined as a pre-conditioning preference score $\left(\frac{\text{time side 1}}{\text{total time}} \right)$ of > 1.5 or < 0.5 . Experimental conditions were compared using normalized social preference scores $(\text{social zone post} / \text{social zone pre})$, and subtracted social preference scores $((\text{social zone post} - \text{social zone pre}) / 900)$ (Dölen et al., 2013; Nardou et al., 2019).

Stereotaxic and tail-vein injections

All stereotaxic injections into the nucleus accumbens (NAc; distance from bregma: anterior +1.54 mm; lateral ± 1.065 mm; ventral -4.1 mm) were performed under general ketamine-medetomidine anesthesia using a stereotaxic instrument (David Kopf). For electrophysiological analysis of NAc-projecting OT neurons, a small volume (~ 30 nl) of diluted Rtb solution (1:4; Lumafluor; Red Retrobeads) was injected unilaterally or bilaterally into NAc core at a slow rate (20 nl/min) using a syringe pump (Harvard Apparatus, MA). For conditional knockdown of *Fmr1*, 1 μ L viral suspension (0.1 μ L/min; CAV2-GFP, CAV2-Cre-GFP, rgAAV-fDIO-Cre-GFP, or rgAAV-fDIO-Cre-HA) was injected bilaterally into NAc 7 days before behavioral testing. CAV2-GFP and CAV2-Cre-GFP viruses were obtained from Plateforme de Vectorologie de Montpellier. The fDIO-Cre-GFP plasmid was a gift of Dr. Bo Li and Dr. Linda Van Aelst and viral packaging was conducted by Stanford University's Neuroscience Gene Vector and Virus Core. rgAAV-fDIO-Cre-HA was obtained from Addgene (Catalog # 121675-AAVrg). To confirm viral-mediated Cre function, 1 μ L CAV2-Cre-GFP or 1.5 μ L rgAAV-fDIO-Cre-HA was injected into NAc of Ai9 (tdTomato Cre-reporter) or OT-2A-Flp::Ai9 mice respectively. After all injections, the injection pipette was left in place for at least 5 min prior to removal from the brain. Injection sites were confirmed post hoc by preparing sections (20-50 μ m) containing the NAc. For i.v. labeling of magnocellular OT neurons with FG, 1 or 2 lateral tail-vein injections of 15 μ L 4% FG (Fluorochrome) diluted in ~ 100 μ L sterile saline were performed using a 1 mL syringe and 25-gauge needle at least 24 hours prior to sacrifice. To facilitate i.v. injections, mice were placed under a warm lamp for 3-5 min and were briefly restrained using a tail vein restrainer (TV-150; Braintree Scientific). Prior to injection, the tail was rubbed firmly with 70% ethanol to clean the injection area and enhance visibility of the tail vein.

Electrophysiology

Using a Leica VT 1200s vibrating microtome, coronal sections (250 μ m) containing the paraventricular nucleus were cut in an ice-cold sucrose solution containing the following (in mM): Sucrose 228; NaHCO₃ 26; Glucose 11; KCl 2.5; NaH₂PO₄ 1; MgSO₄ 7; CaCl₂ 0.5. Immediately after cutting, slices were transferred into 32°C aCSF containing (in mM): NaCl 119; KCl 2.5; NaH₂PO₄ 1; NaHCO₃ 26.2; Glucose 11; MgCl₂ 1.3; CaCl₂ 2.5. Slices were allowed to recover at 29°C for at least 1 h prior to recording. GFP+ and Rtb+ neurons were visualized using 470 nm and 535 nm LEDs respectively (Cool LED). LED excitation was delivered through a 40x microscope objective, (Olympus) and fluorescence was detected using a camera and visualized using SliceScope Pro software (Scientifica). Recordings were conducted in current clamp using patch pipettes (2-4 M Ω) filled with internal solution containing the following (in mM): 130 K-Gluconate, 10 HEPES, 1 NaCl, 1 CaCl₂, 10 EGTA, 1 MgCl, 2 Mg-ATP, 0.5 Na-GTP, and 0.125% Neurobiotin) and the liquid junction potential (13 mV) was corrected after recording. Recording was discontinued for neurons that failed to maintain a hyperpolarized resting membrane potential in the absence of current injection, and neurons that failed to produce APs > 50 mV were excluded from further analysis. Data were acquired and analyzed using the Recording Artist plugin in Igor Pro and custom software written in MATLAB (Mathworks). AP initiation was determined as the first point where the membrane potential accelerated past 40 ms/s². This threshold was confirmed with comparison to manual determination of AP initiation. AP peaks were identified prior to correction of the liquid junction potential by finding local maxima with a minimum peak of 0 mV and a minimum separation of 5 ms and absolute membrane potential values were subsequently corrected. Nine mice, with a maximum of 4 neurons per animal were used to cross-validate FG and electrophysiological differentiation of magnocellular and parvocellular neurons. Neurons were targeted and recorded blind to FG labeling status. Six mice, with a maximum of 2 neurons per animal were used for electrophysiological characterization of OTergic projections to NAc.

Cell Isolation, Enrichment, and cDNA Library Preparation

OT-2A-Flp::fGFP mice were weaned at P21 and given i.v. injections of FluoroGold on that and the following day. At P23 mice were sacrificed. Brains were rapidly removed and 250 μ m thick coronal slices (n = 7 mice) containing the paraventricular nucleus of the hypothalamus were sectioned using a Leica VT-1200s vibrating microtome in ice-cold ACSF solution containing the following (in mM): 124 NaCl; 2.5 KCl; 1.2 NaH₂PO₄; 24 NaHCO₃; 5 HEPES; 13 glucose; 2 MgSO₄; 2 CaCl₂, oxygenated with carbogen gas (95% O₂ and 5% CO₂) to pH 7.3 - 7.4. To microdissect the PVN, slices were placed in a Petri-dish containing ice-cold, oxygenated ACSF solution and the PVN was identified using the 3rd ventricle and other structural markers. Following dissection, tissue was placed into 2.6 mL equilibrated Papain DNase-I solution. The dissociation protocol was similar to Chev  e et al., 2018, itself an adaptation from the trehalose-enhanced

neuronal isolation protocol (Saxena et al., 2012) using the Worthington Papain Dissociation System (Worthington Biochemical Corporation). The following modifications from Chevée et al., 2018 were made: a single low speed centrifugation (300xg for 5 min) was performed after dissociation, and the pellet was resuspended in 250 μ L media (DMEM, 5% trehalose (w/v), 25 μ M AP-V, 0.4 mM kynurenic acid, 6 μ L of 40 U/ μ L RNase inhibitor) and 250 μ L of EBSS#2 (EBSS, 25 mM AP-V, 100 mM Kynurenic acid, ovomucoid protease inhibitor with BSA, DNase-I, 5% w/v Trehalose, 40 U/ μ L RNase inhibitor) at room temperature. Following resuspension, single-cell suspensions were introduced into a FACS machine (Beckman Coulter MoFlo Cell Sorter).

Neurons were sorted based on fluorescence (GFP+/FG- or GFP+/FG+) directly into individual wells of a 96-well plate containing 2 μ L Smart-Seq2 lysis buffer + RNAase inhibitor, 1 μ L oligo-dT primer, and 1 μ L dNTPs (Picelli et al., 2014). Three plates were collected across 2 separate dates (Plate 1: n = 1 mouse, 32 GFP+/FG- neurons and 47 GFP+/FG+ neurons; Plate 2: n = 1 mouse, 32 GFP+/FG- neurons and 33 GFP+/FG+ neurons; Plate 3: n = 5 mice, 25 GFP+/FG- neurons and 48 GFP+/FG+ neurons). Following the sort, plates were briefly spun-down in a table-top microcentrifuge and immediately placed on dry ice. Single-cell lysates were kept at -80°C until cDNA library preparation. Library preparation and amplification of single-cell samples was performed using a modified version of the Smart-Seq2 protocol (Chevée et al., 2018).

Individual libraries were quality controlled, pooled, and sequenced on two lanes of an Illumina HiSeq 2500 sequencer to an average depth of $1,424,499 \pm 77,731$ paired-end 50bp reads per neuron. Reads were aligned to the reference mouse genome (mm10) using the Hisat 2 aligner and transcript abundances were estimated using CuffNorm (Qiu et al., 2017) against the mouse GENCODE reference transcriptome (vM8) (Qiu et al., 2017). Relative abundances were converted to absolute estimates of gene expression using the Monocle2 CENSUS approach and used as input for the dpFeature workflow (Qiu et al., 2017). 146 neurons passed our quality assessment, and we identified a total of 10,172 genes with detectable expression in at least 5 neurons. For those cells that passed QC, the mean mRNA-copies per cell was 27,551 and mean number of genes detected across all cells was 3,713. A Uniform Manifold Approximation and Projection (UMAP) (Becht et al., 2018) embedding was obtained by collapsing the top 50 principal components learned on the top 1000 genes with highest residuals to the Monocle model fit. All data are archived in the Gene Expression Omnibus (GEO Accession number GSE147092). Raw and processed scRNA-Seq data and code to reproduce the complete analysis are also publicly available at https://github.com/gofflab/OT_neuron_study_2020.

In situ hybridization chain reaction

Experiments were performed on 3 week-old FG injected male wild-type C57BL6/J mice obtained from Jackson Laboratories. Brains were extracted and immediately frozen in OCT with liquid nitrogen. Brains were sectioned on a cryostat at 7 μ m. Serial sections were collected onto charged slides, briefly washed in PBS and then fixed in 4% PFA for 10 mins. Slides were then washed twice in PBS, transferred to 70% ethanol, and kept at 4°C . *In situ* HCR v3.0 was performed using the protocols previously detailed (Choi et al., 2018). Probe hybridization and amplification steps were both performed overnight (12–16 h). *In situ* probes (Table S2) were design using publicly available software (Raj et al., 2008).

Immunohistochemistry

Prior to immunostaining, sections were mounted on slides and rinsed 4×10 min in PBS followed by 1 hr in blocking solution (0.5% Triton X-100, 10% horse serum, 0.2% bovine serum albumin in PBS). Antibodies were diluted in PBS containing the following: 0.5% Triton X-100, 1% horse serum, 0.2% bovine serum albumin. Primary antibodies were applied overnight at room temperature (RT), and after rinsing slides in PBS, secondary antibodies were applied for 2 hr at RT. With the exception of FMRP immunohistochemistry, OT neurons were labeled using the antibody PS38 (gift of Dr. Harold Gainer; 1:150). PS38 was detected using Alexa 488, (magnocellular and parvocellular distribution; Life Technologies; donkey anti-mouse; 1:1000) Alexa 647 (GFP or tdTomato colocalization; Life Technologies; goat-anti mouse or donkey-anti mouse respectively; 1:1000), or Alexa 350 (RtB colocalization; Life Technologies; goat-anti mouse; 1:200). FluoroGold was labeled with an anti-fluorogold antibody, (Fluorochrome; 1:100), and detected using Alexa 647 (Life Technologies; donkey anti-rabbit; 1:1000). This approach was chosen instead of visualizing the native FG fluorescence to maximize the detectable signal with our microscope configuration. GFP signal was amplified using the anti-GFP antibody, ab13970, (ABCAM; 1:1000) and visualized using Alexa 488 (Life Technologies; goat-anti chicken; 1:1000). tdTomato signal was amplified using the anti-mCherry antibody AB0040 (SICGEN; 1:4000) and visualized using Alexa 555 (Life Technologies; donkey-anti goat; 1:1000). For post hoc identification of electrophysiologically recorded neurons, Neurobiotin was detected using streptavidin-conjugated Alexa 350 (Life Technologies; 1:800).

FMRP co-labeling experiments were conducted as described above with the following exceptions. After rinsing in PBS, slides were transferred to a hot (90°C) sodium citrate (10 mM Sodium Citrate, 0.05% Tween 20, pH 6.0) bath for antigen retrieval (30 min) followed by 1 hr in blocking solution at RT. Sections were labeled with anti-FMRP 2F5-1 ((Gabel et al., 2004) Developmental Studies Hybridoma Bank; 1:1) which was detected using Alexa 647 (Jackson ImmunoResearch; goat anti-mouse IgG2b; 1:500). OT neurons were labeled with the anti-OT antibody VA10 (Altstein and Gainer, 1988) (gift of Dr. Harold Gainer; 1:1000) and detected using Alexa 488 (Life Technologies; donkey anti-rabbit; 1:1000).

Histology

Following transcardial perfusion with 1M PBS and 10% formalin, brains were kept in formalin at 4°C overnight and then transferred to PBS. To prepare them for re-sectioning, acute brain slices used for electrophysiology were placed in 10% formalin immediately after

recording, and then transferred to PBS. For all experiments in which fluorescent colocalization was analyzed, serial brain sections (20 μm thickness) were cut using a cryostat following cryoprotection of tissue with a 30% sucrose solution containing 0.01% azide. Following immunostaining, images were acquired using an EVOS or Olympus BX41 microscope with 4x, 10x, and 40x objectives and analyzed using ImageJ.

QUANTIFICATION AND STATISTICAL ANALYSIS

All behavioral and electrophysiological analyses were performed with MATLAB (Mathworks), Prism (GraphPad), or R (The R Project). Electrophysiological comparisons were made using a two-tailed Mann-Whitney U Test or Wilcoxon Signed-Rank Test for unpaired and paired comparisons, respectively. $p < 0.05$ was considered significant. Behavioral comparisons were made using a two-tailed, Student's *t* test (paired or unpaired and with or without Welch's correction as appropriate per the Central Limit Theorem), with $p < 0.05$ considered significant. Statistical parameters can be found in figure legends. All data were subjected to the Shapiro-Wilk normality test and F test to compare variances. For k-means clustering analysis, two clusters were specified, as this number of clusters was indicated by the elbow method (Figure S2). Electrophysiological feature values used for the clustering analysis were z-scored across all neurons, while the clustering algorithm was trained only on the values from neurons from Figure 2. *Calb1-Cre::cdFmr1KO* experiments were done blind, as were a subset of *OT-Flip::cdFmr1KO* experiments. For electrophysiological analysis, neurons were targeted for recording blind to FG labeling status.

All scRNAseq analysis was performed in R/Bioconductor following alignment with Hisat2. Log2 expression estimates (with a pseudocount of 1) of high-variance genes were used as input for PCA analysis and UMAP clustering of individual cells. After cluster assignment, differential expression testing was performed using the Monocle2 VGAM model comparison test (Trapnell et al., 2014) (FDR 0.1%, Monocle2 test, Benjamini-Hochberg corrected). Gene set enrichment was computed using the hypergeometric test unless stated otherwise. Using the full dataset with a *q* value of 0.001 (0.1% FDR), *Post hoc* power calculations were made for observing 181 differentially expressed genes. Based on these calculations, there is 99.84% power (probability of true positives) to observe 181 genes with a significant two-fold change ($\log_2 \text{Magno/Parvo} \geq 1$ or $\log_2 \text{Magno/Parvo} \leq -1$) in gene expression. Given that the observed log2-fold changes (magnocellular versus parvocellular) for the mean expression of *Calb1*, *Kcnmb4*, *Reln*, *Cnr1* are 6.59, 2.91, -7.38 , and -3.92 respectively, it was determined that there was more than sufficient (greater than 99.84%) power for this analysis. Limiting the power calculation to a balanced design by using only enough magnocellular cells to match the reduced number of parvocellular cells (thereby disregarding the observed ratio of cell types), results in 96.94% power for and FDR of 0.001 to observe 181 genes with a significant two-fold change ($\log_2 \text{Magno/Parvo} \geq 1$ or $\log_2 \text{Magno/Parvo} \leq -1$). Code to reproduce the complete analysis is publicly available at https://github.com/gofflab/OT_neuron_study_2020.