

Lower and Upper Bounds for Positive Bases of Skein Algebras

Thang T. Q. Lê^{1,*}, Dylan P. Thurston² and Tao Yu³

¹School of Mathematics, 686 Cherry Street, Georgia Tech, Atlanta, GA 30332, USA, ²Department of Mathematics, Indiana University, Bloomington, IN 47405-7106, and ³School of Mathematics, 686 Cherry Street, Georgia Tech, Atlanta, GA 30332, USA

**Correspondence to be sent to: e-mail: letu@math.gatech.edu*

We show that if a sequence of normalized polynomials gives rise to a positive basis of the skein algebra of a surface, then it is sandwiched between the two types of Chebyshev polynomials. For the closed torus, we show that the normalized sequence of Chebyshev polynomials of type one ($\hat{T}_n$) is the only one that gives a positive basis.

1 Introduction

1.1 Results

Let R be a commutative integral domain with a distinguished invertible element $q \in R$. The main examples are $R = \mathbb{Z}[q^{\pm 1}]$ and $R = \mathbb{Z}$ with $q = 1$. Let $\Sigma = \Sigma_{g,p}$ be the oriented surface of genus g with p points removed. The skein algebra $\mathcal{S}(\Sigma; R)$ is the R -algebra spanned by isotopy classes of framed links in $\Sigma \times (-1, 1)$ modulo the skein relation and the trivial loop relation in Figure 1. The product is given by superposition. For details, see Section 2.1.

We assume that the ring R has a positive part R_+ , see Section 2.4. When $R = \mathbb{Z}$ its positive part is $R_+ = \mathbb{Z}_+$, the set of non-negative integers, and when $R = \mathbb{Z}[q^{\pm 1}]$ its positive part is $R_+ = \mathbb{Z}_+[q^{\pm 1}]$. A basis B of an R -algebra is *positive* if the structure

Received August 8, 2019; Revised February 24, 2020; Accepted March 26, 2020
 Communicated by Prof. Anton Alekseev



Fig. 1. Defining relations for $\mathcal{S}(M)$.

constants are in R_+ , that is, for any $x, y \in B$ the product xy is a linear combination of elements in B with coefficients in R_+ . An important conjecture [4] in cluster algebra theory is that the skein algebra $\mathcal{S}(\Sigma)$ has a positive basis.

A sequence of polynomials $(P_n(x))_{n=0}^\infty$ with R coefficients is *normalized* if $P_n(x)$ is monic with degree n for each $n \geq 0$. Note $P_0(x) = 1$ by definition. A normalized sequence (P_n) defines a basis B_P of $\mathcal{S}(\Sigma; R)$ as in Section 2.3. We say a sequence of normalized polynomials (P_n) is *positive on Σ over R* if the corresponding basis B_P is a positive basis of $\mathcal{S}(\Sigma; R)$.

Two important sequences considered here are the normalized Chebyshev polynomials of type one $(\hat{T}_n)$ and type two (S_n) , defined inductively by

$$\begin{aligned} \hat{T}_0(x) &= 1, & \hat{T}_1(x) &= x, & \hat{T}_2(x) &= x^2 - 2, & \hat{T}_n(x) &= x\hat{T}_{n-1}(x) - \hat{T}_{n-2}(x), & n &\geq 3, \\ S_0(x) &= 1, & S_1(x) &= x, & & S_n(x) &= xS_{n-1}(x) - S_{n-2}(x), & n &\geq 2. \end{aligned}$$

Note that the polynomials $\hat{T}_n(x)$ coincide with the Chebyshev polynomials of type one except for $\hat{T}_0$. The reason is we want $\hat{T}_n$ to be monic.

The 2nd named author [13] showed that when $R = \mathbb{Z}$ and $q = 1$, the sequence $(\hat{T}_n)$ is positive on any surface and made a conjecture, refining an earlier one of Fock and Goncharov [4], that $(\hat{T}_n)$ is positive when $R = \mathbb{Z}[q^{\pm 1}]$ as well.

The 1st named author [7] showed that if the sequence (P_n) is positive on a surface Σ having genus ≥ 1 , then $(P_n) \geq (\hat{T}_n)$. Here, $(P_n) \geq (Q_n)$ means each P_n is an R_+ -linear combination of $Q_0, Q_1, \dots, Q_n$.

We extend the lower-bound result in [7] to surfaces of genus 0, and at the same time give an upper bound for all surfaces.

Theorem 1. Suppose $R = \mathbb{Z}[q^{\pm 1}]$ and (P_n) is normalized with integer coefficients. Let Σ be a surface with genus at least 1 or with at least 4 punctures. If (P_n) is positive on Σ , then $(\hat{T}_n) \leq (P_n) \leq (S_n)$.

In the case of the closed torus $\Sigma_{1,0}$, our result is much more precise.

Theorem 2. A normalized sequence (P_n) is positive on the closed torus $\Sigma_{1,0}$ over $\mathbb{Z}$ or $\mathbb{Z}[q^{\pm 1}]$ if and only if $(P_n) = (\hat{T}_n)$.

See Section 2.5 for more refined statements.

Theorem 2 is somewhat surprising. The sequence (S_n) gives simple objects in the ring of modules over the quantum group $U_q(sl_2)$. From this point of view, (S_n) is more natural than the sequence $(\hat{T}_n)$. However, it does not give a positive basis on the torus. On the other hand, if the surface has negative Euler characteristic, it is conjectured that the (S_n) is positive [13].

Conjecture 1. Both $(\hat{T}_n)$ and (S_n) are positive on any surface Σ with negative Euler characteristic.

If the skein algebra of a surface Σ has a positive basis, then one can ask whether the skein algebra can be categorified. A categorification would give the skein algebra a richer structure and hopefully will lead to an interesting 4D topological quantum field theory. The (positive) basis of a normalized sequence is functorial in the sense that if $f : \Sigma \rightarrow \Sigma'$ is a strict embedding (see the definition in Section 2.3), then f maps injectively the (positive) basis of Σ to the (positive) basis of Σ' . For some initial results and an extensive discussion of strategies to categorify the conjectural positive bases of the sequences $(\hat{T}_n)$ and (S_n) , see [12].

Remark 1.1. In Theorem 1, we exclude the case when (g, p) is one of $(0, 0)$, $(0, 1)$, $(0, 2)$, $(0, 3)$ because in these cases the skein algebra $\mathcal{S}(\Sigma_{g,p}; R)$ is a commutative polynomial algebra and hence obviously has a positive basis; for example, the monomial basis.

Remark 1.2. For other types of positivity, see for example [2, 3, 8, 10].

2 Skein Algebras

2.1 Basic definitions

Let R be a commutative integral domain with unit and with a distinguished invertible element $q \in R$. Suppose M is an oriented 3-manifold, not necessarily closed. A *framed link* L in M is a smooth, unoriented, closed one-dimensional submanifold equipped with a normal vector field. By convention, the empty set is also considered as a framed link. The skein module $\mathcal{S}(M; R)$ of M , introduced independently by Przytycki [11] and Turaev

[14, 15], is defined as the quotient of the R -module freely generated by isotopy classes of framed links in M by the skein relation and the trivial loop relation in Figure 1.

In the figure, the diagrams represent links that are identical outside of a ball in the manifold M . The shaded part is the projection onto the equatorial plane of the ball where the difference is. The framings on the diagrams are vertical, that is, pointing out to the reader. The skein relations were introduced by Kauffman [6].

For $\Sigma = \Sigma_{g,p}$, the oriented surface with genus g and p punctures, let

$$\mathcal{S}(\Sigma; R) := \mathcal{S}(\Sigma \times (-1, 1); R),$$

which has a product structure given by stacking. More precisely, the product of two framed links L and K in $\mathcal{S}(\Sigma; R)$ is given by the union $i_+(L) \cup i_-(K)$, where $i_{\pm} : M \rightarrow M$ are the embeddings defined by $i_{\pm}(x, t) = (x, \frac{t \pm 1}{2})$. It is easy to see that this is a well-defined product that turns $\mathcal{S}(\Sigma; R)$ into an R -algebra. It should be noted that there are surfaces $\Sigma \neq \Sigma'$ such that $\Sigma \times (-1, 1)$ and $\Sigma' \times (-1, 1)$ are diffeomorphic as 3-manifolds, but $\mathcal{S}(\Sigma; R)$ and $\mathcal{S}(\Sigma'; R)$ have different product structures, hence different as R -algebras.

An orientation preserving embedding of surfaces $i : \Sigma \rightarrow \Sigma'$ induces an R -algebra homomorphism $i_* : \mathcal{S}(\Sigma; R) \rightarrow \mathcal{S}(\Sigma'; R)$ by applying $(x, t) \mapsto (i(x), t)$ to links. In particular, the mapping class group of Σ acts on the skein algebra $\mathcal{S}(\Sigma; R)$.

2.2 Multicurves

A *multicurve* γ on a surface $\Sigma = \Sigma_{g,p}$ is a closed unoriented 1-submanifold of Σ none of the components of which bounds a disk in Σ . A *simple closed curve* is a multicurve with one component. A *peripheral curve* is simple closed curve bounding a once-punctured disk. By convention, the empty set is a multicurve.

A multicurve γ of Σ defines an element of $\mathcal{S}(\Sigma; R)$ by the embedding $\gamma \subset \Sigma \cong \Sigma \times \{0\}$. The framing on γ is vertical, that is, parallel to the $(-1, 1)$ direction and pointing towards 1.

If α and β are disjoint multicurves, then $\alpha \cup \beta$ is also a multicurve, and $\alpha\beta = \beta\alpha = \alpha \cup \beta$ as elements of $\mathcal{S}(\Sigma; R)$. It follows that peripheral curves are in the center of the skein algebra.

By grouping together isotopic components, every multicurve γ can be uniquely written as

$$\gamma = \gamma_1^{n_1} \dots \gamma_r^{n_r}, \quad (1)$$

where $\gamma_1, \dots, \gamma_r$ are disjoint and pairwise non-isotopic simple closed curves, and n_i are positive integers.

2.3 Bases

Let $B = B(\Sigma)$ denote the set of all isotopy classes of multicurves. The module structure of the skein algebra is very simple.

Theorem 2.1 ([11]). As an R -module, $\mathcal{S}(\Sigma; R)$ is free with basis $B(\Sigma)$.

From this description of free bases, it is easy to see that, as an R -algebra $\mathcal{S}(\Sigma_{g,p}; R)$ is isomorphic to a commutative polynomial algebra for the case when $g = 0$ and $p \leq 3$. Namely $\mathcal{S}(\Sigma_{0,0}; R) \cong \mathcal{S}(\Sigma_{0,1}; R) \cong R$. When $(g, p) = (0, 2)$, one has $\mathcal{S}(\Sigma_{0,2}; R) \cong R[x]$ where x is the only peripheral curve. Finally, $\mathcal{S}(\Sigma_{0,3}; R) \cong R[x, y, z]$, where x, y, z are the three peripheral curves. These surfaces will not be considered below. For all other (g, p) , over $R = \mathbb{Z}[q^{\pm 1}]$, the skein algebra $\mathcal{S}(\Sigma_{g,p}; R)$ is non-commutative.

An embedding $\iota : \Sigma \hookrightarrow \Sigma'$ is *strict* if the induced map from $B(\Sigma)$ to $B(\Sigma')$ is injective. From Theorem 2.1, we get the following.

Corollary 2.2. If $\iota : \Sigma \hookrightarrow \Sigma'$ is a strict embedding, then $\iota_* : \mathcal{S}(\Sigma; R) \rightarrow \mathcal{S}(\Sigma'; R)$ is an algebra embedding.

The basis $B(\Sigma)$ can be twisted by polynomial sequences as follows. Let $P = (P_n)$ be a normalized sequence of polynomials. If $\gamma = \gamma_1^{n_1} \dots \gamma_r^{n_r}$ as in Equation (1), define

$$P(\gamma) = P_{n_1}(\gamma_1) \dots P_{n_r}(\gamma_r).$$

Let

$$B_P(\Sigma) := \{P(\gamma) : \gamma \in B(\Sigma)\}.$$

Then, $B_P(\Sigma)$ is also a free R -basis of $\mathcal{S}(\Sigma; R)$. When $P_n(x) = x^n$ one recovers $B_P(\Sigma) = B(\Sigma)$.

2.4 Positivity

Let $R_+ = \mathbb{Z}_+$ if $R = \mathbb{Z}$ and $R_+ = \mathbb{Z}_+[q^{\pm 1}]$ if $R = \mathbb{Z}[q^{\pm 1}]$. More generally, when R is an arbitrary commutative domain with a distinguished invertible element q , a *positive part* of R is any subset R_+ satisfying

- (1) $q, q^{-1} \in R_+$,
- (2) R_+ is closed under addition and multiplication,
- (3) $R_+ \cap (-R_+) = \{0\}$.

Fix such a positive part of R . A basis of $\mathcal{S}(\Sigma; R)$ is *positive* if the structure constants are in R_+ , that is, for any basis elements x, y , the product xy is an R_+ -linear combination of the basis elements. A normalized sequence of polynomials $P = (P_n)$ is *positive* on Σ over R if the basis B_P is positive for $\mathcal{S}(\Sigma; R)$.

Recall that given two normalized sequences of polynomials (P_n) and (Q_n) , one defines $(P_n) \leq (Q_n)$ if each Q_n is an R_+ -linear combination of $P_0, P_1, \dots, P_n$.

Lemma 2.3. The binary relation $(\leq)$ is a partial order on the set of normalized sequences of polynomials.

Proof. It is clear that $(\leq)$ is reflexive and transitive, and we need to show that it is anti-symmetric. Assume that $(P_n) \leq (Q_n)$ and $(Q_n) \leq (P_n)$. Writing each sequence (P_n) and (Q_n) as an infinite column vectors, then $(Q_n) = A \times (P_n)$, where A is a $\mathbb{Z}_+ \times \mathbb{Z}_+$ matrix that is upper triangular, having 1 on the diagonal, and having entries R_+ . We can write $A = I + N$, where I is the identity matrix and N is a strictly upper triangular matrix. Suppose the matrix $N = (N_{ij})$ is not 0. Among all the non-zero entries of N , let N_{ij} be the one with the smallest pair $(j - i, i)$ in the lexicographic order. Then, it is easy to see that $(A^{-1})_{ij} = -N_{ij}$. Since $(Q_n) \leq (P_n)$, all the entries of A^{-1} are in R_+ . It follows that both N_{ij} and $-N_{ij}$ are in R_+ , a contradiction. Thus, $N = 0$, and $(P_n) = (Q_n)$. ■

The Chebyshev polynomials of type one (T_n) and type two (S_n) are defined by the recurrence relations

$$\begin{aligned} T_0(x) &= 2, & T_1(x) &= x, & T_n(x) &= xT_{n-1}(x) - T_{n-2}(x), & n &\geq 2, \\ S_0(x) &= 1, & S_1(x) &= x, & S_n(x) &= xS_{n-1}(x) - S_{n-2}(x), & n &\geq 2. \end{aligned}$$

They can be characterized by

$$T_n(t + t^{-1}) = t^n + t^{-n}, \quad S_n(t + t^{-1}) = t^n + t^{n-2} + \dots + t^{-n}.$$

Thus, $S_n(x) - S_{n-2}(x) = T_n(x)$ for $n \geq 2$.

While (S_n) is a normalized sequence, (T_n) is not. We normalize T_n by setting $\hat{T}_0(x) = 1$ and $\hat{T}_n(x) = T_n(x)$ for $n > 0$, as in the Introduction. Then, $(\hat{T}_n)$ is a normalized sequence, and

$$S_n(x) = \sum_{i=0}^{\lfloor n/2 \rfloor} \hat{T}_{n-2i}(x).$$

Thus, $(\hat{T}_n) \leq (S_n)$.

2.5 Results

Here are fuller, more refined versions of Theorems 1 and 2. For the convenience of proofs, we formulated them in three statements.

Theorem 2.4. Suppose R is a commutative domain with a distinguished invertible element q and a positive part R_+ . Assume either $g \geq 1$ or $p \geq 4$. Let (P_n) be a normalized sequence of polynomials that is positive on $\Sigma_{g,p}$ over R .

- (a) One has $(P_n) \geq (\hat{T}_n)$ and $P_1(x) = x$.
- (b) If $P_n = \hat{T}_n$ for $n \leq 3$ then $(P_n) = (\hat{T}_n)$.

Theorem 2.5. Suppose R is a commutative domain with a distinguished invertible element q and a positive part R_+ . A normalized sequence polynomial (P_n) is positive on the torus $\Sigma_{1,0}$ over R if and only if $(P_n) = (\hat{T}_n)$.

Theorem 2.6. Suppose $R = \mathbb{Z}[q^{\pm 1}]$ and (P_n) are normalized sequences of polynomials with integer coefficients. Assume either $g \geq 1$ or $p \geq 4$. If (P_n) is positive on $\Sigma_{g,p}$ over R , then $(P_n) \leq (S_n)$.

To prove these theorems, it suffices to consider three *basic surfaces*: the closed torus $\Sigma_{1,0}$, the once-punctured torus $\Sigma_{1,1}$, and the sphere with four punctures $\Sigma_{0,4}$. This can be seen as follows. If $\Sigma_{g,p}$ has at least four punctures, then there is a strict embedding of $\Sigma_{0,4}$ into $\Sigma_{g,p}$, and by Corollary 2.2, the skein algebra $\mathcal{S}(\Sigma_{0,4}; R)$ embeds in $\mathcal{S}(\Sigma; R)$. The results for $\Sigma_{0,4}$ implies the corresponding results for $\Sigma_{g,p}$. If Σ has genus at least 1, then either $\Sigma = \Sigma_{1,0}$ or $\Sigma_{1,1}$ strictly embeds into $\Sigma_{g,p}$. In the latter case, the results for $\Sigma_{1,1}$ imply those for $\Sigma_{g,p}$.

Theorem 2.4 is proved in Section 4. Theorem 2.5 is proved in Section 5.1. Theorem 2.6 is proved in Section 5.2 for $\Sigma_{1,1}$ and Section 5.3 for $\Sigma_{0,4}$. The case of $\Sigma_{1,0}$ is a corollary of Theorem 2.5.

3 Parameterization of Curves on Basic Surfaces

A non-oriented simple closed curve on the torus $\Sigma_{1,0}$ is determined by the homology class up to sign. After choosing a basis of homology on $\Sigma_{1,0}$, every simple closed curve is represented by a pair of coprime integers (r, s) , which is identified with $(-r, -s)$.




Fig. 2. The involution of the torus.



(A) a , b , and puncture numbering

(B) The $(1, 1)$ curve


Fig. 3. Curves on $\Sigma_{0,4}$.

The isotopy classes of simple curves on $\Sigma_{1,1}$ except the peripheral curve are in one-to-one correspondence with those on $\Sigma_{1,0}$. Thus, the same notations can be used to represent essential simple closed curves on $\Sigma_{1,1}$.

In both cases, the mapping class group is $SL_2(\mathbb{Z})$ and the action of a mapping class on the curves is the standard linear action.

The sphere with four punctures $\Sigma_{0,4}$ can be regarded as the quotient of the torus $\Sigma_{1,0}$ by the involution in Figure 2. The action has four fixed points that corresponds to the punctures. This quotient also identifies the essential simple closed curves on $\Sigma_{1,0}$ with the non-peripheral ones on $\Sigma_{0,4}$. Thus, coprime integers (r, s) , with the identification $(r, s) = (-r, -s)$, also represent curves on $\Sigma_{0,4}$.

More concretely, represent $\Sigma_{0,4}$ as in Figure 3. Choose the curves a and b , and number the punctures as in the figure. The curve surrounding puncture i is denoted by γ_i . To obtain the (r, s) curve on $\Sigma_{0,4}$, take $|r|$ parallel copies of a and $|s|$ parallel copies of b , and resolve each intersection such that one would turn left from a to b if $rs > 0$ and turn right if $rs < 0$. Thus, $a = (1, 0)$, $b = (0, 1)$. The $(1, 1)$ curve is demonstrated in Figure 3. Thus, the (r, s) curve is the Luo product of a^r and b^s [9] if $r \geq 0$ and $s \geq 0$ and is the Luo product of a^{-r} and b^s if $r < 0$ and $s \geq 0$. The Luo product of two multicurves α and β can be defined as the unique multicurve γ such that $\alpha\beta = q^{I(\alpha, \beta)}\gamma + x$, where $I(\alpha, \beta)$ is the geometric intersection index of α and β , and x is a $\mathbb{Z}[q, q^{-1}]$ -linear combination of multicurves with coefficients being Laurent polynomials in q of highest degrees $< I(\alpha, \beta)$.

Fig. 4. Four resolutions of ab .

For $\Sigma_{0,4}$, the mapping class group is $(\mathbb{Z}/2)^2 \rtimes PSL_2(\mathbb{Z})$. The mapping classes that fix puncture 4 forms a subgroup isomorphic to $PSL_2(\mathbb{Z}) = SL_2(\mathbb{Z})/\{\pm 1\}$. The action of this subgroup on (r, s) curves is the projective linear action.

4 Lower Bound, Proof of Theorem 2.4

Note that Theorem 2.4 about lower bounds does not assume $R = \mathbb{Z}[q^{\pm 1}]$.

Proof of Theorem 2.4. (1) The case when Σ has genus at least 1 is already proved in [7]. Now, assume $\Sigma = \Sigma_{0,4}$. Let σ be the counterclockwise half twist along a , fixing punctures 3 and 4 and exchanging punctures 1 and 2. Then, σ can be represented by the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. For convenience, define $b_n = (n, 1) = \sigma^n(b)$.

By resolving both crossings between a and b as in Figure 4, we get

$$ab = q^2 b_1 + q^{-2} b_{-1} + c_0,$$

where

$$c_0 = \gamma_1 \gamma_3 + \gamma_2 \gamma_4.$$

Applying σ^n to both sides, we have

$$ab_n = q^2 b_{n+1} + q^{-2} b_{n-1} + c_n, \quad (2)$$

where we defined

$$c_n = \sigma^n(c_0) = \begin{cases} c_0 = \gamma_1 \gamma_3 + \gamma_2 \gamma_4, & n \text{ even,} \\ c_1 = \gamma_1 \gamma_4 + \gamma_2 \gamma_3, & n \text{ odd.} \end{cases}$$

The following can be proved easily using induction.

Lemma 4.1. For $n \geq 0$,

$$T_n(a)b = q^{2n}b_n + q^{-2n}b_{-n} + c_0f_n + c_1g_n,$$

where

$$f_n(a) = \sum_{\substack{0 < i \leq n \\ i \text{ odd}}} [i] \hat{T}_{n-i}(a) \quad \text{and} \quad g_n(a) = \sum_{\substack{0 < i \leq n \\ i \text{ even}}} [i] \hat{T}_{n-i}(a)$$

are polynomials of a with R coefficients. Here, $[i] = q^{2i-2} + q^{2i-6} + \dots + q^{2-2i}$ is the quantum integer. ■

It is also possible to express f_n and g_n in terms of S_n , (see e.g., [1] Theorem 3.5). The expressions of f_n and g_n are not needed in the following.

First, we show $P_1(x) = x$. Write $P_1(x) = x + \delta$. Then,

$$\begin{aligned} P_1(a)P_1(b) &= (a + \delta)(b + \delta) \\ &= (q^2b_1 + q^{-2}b_{-1} + c_0) + \delta(a + b) + \delta^2 \\ &= q^2(P_1(b_1) - \delta) + q^{-2}(P_1(b_{-1}) - \delta) + (P_1(\gamma_1) - \delta)(P_1(\gamma_3) - \delta) + \\ &\quad + (P_1(\gamma_2) - \delta)(P_1(\gamma_4) - \delta) + \delta(P_1(a) + P_1(b) - 2\delta) + \delta^2. \end{aligned}$$

The positivity of (P_n) implies that $-\delta$ and δ , the coefficients of $P_1(\gamma_1)$ and $P_1(a)$, respectively, are both in R_+ . Thus, $\delta = 0$, that is $P_1(x) = x$.

Now, consider $P_n(x) = T_n(x) + \delta_{n-1}T_{n-1}(x) + \dots + \delta_1T_1(x) + \delta_0$ with $n \geq 2$. Then,

$$\begin{aligned} P_n(a)P_1(b) &= (T_n(a) + \delta_{n-1}T_{n-1}(a) + \dots + \delta_0)b \\ &= q^{2n}b_n + q^{-2n}b_{-n} + \delta_{n-1}(q^{2n-2}b_{n-1} + q^{2-2n}b_{1-n}) + \\ &\quad + \dots + \delta_1(q^2b_1 + q^{-2}b_{-1}) + \delta_0b + c_0F + c_1G, \end{aligned}$$

where F and G are polynomials of a with R coefficients. Then, by the positivity of (P_n) , the coefficients of $P_1(b_i) = b_i$ are in R_+ , which implies $\delta_i \in R_+$. Thus, $(P_n) \geq (\hat{T}_n)$.

(2) Choose an essential simple closed curve z on Σ and a regular neighborhood N of z , which is an annulus. Then, $\mathcal{S}(N) \cong R[z]$ is a subalgebra of $\mathcal{S}(\Sigma)$, and the positivity of (P_n) on Σ implies that $(P_n(z))$ is a positive basis for $R[z]$.

Assume $P_i(x) = \hat{T}_i(x)$ for $i < k$ where $k \geq 4$. Since $(P_n) \geq (\hat{T}_n)$, we can write

$$P_k(x) = T_k(x) + \sum_{i=0}^{k-1} \delta_i \hat{T}_i(x).$$

for some $\delta_i \in R_+$. Consider

$$P_1(z)P_{k-1}(z) = T_1(z)T_{k-1}(z) = T_k(z) + T_{k-2}(z).$$

This should be an R_+ -linear combination of $P_0(z), \dots, P_k(z)$. The coefficient of $P_k(z)$ is 1 by the monic condition. Thus,

$$P_1(z)P_{k-1}(z) - P_k(z) = -\delta_{k-1}\hat{T}_{k-1}(z) + (1 - \delta_{k-2})\hat{T}_{k-2}(z) + \sum_{i=0}^{k-3} (-\delta_i)\hat{T}_i(z)$$

is an R_+ -linear combination of $P_0(z) = \hat{T}_0(z), \dots, P_{k-1}(z) = \hat{T}_{k-1}(z)$. This shows $\delta_i = 0$ for $i < k$ except $i = k - 2$. A similar argument with $P_2(z)P_{k-2}(z)$ shows $\delta_{k-2} = 0$ as well. Thus, $P_k(x) = \hat{T}_k(x)$. By induction, $(P_n) = (\hat{T}_n)$.

5 Upper Bound

The strategy to obtain an upper bound on positive polynomials is to compute the product of simple closed curves with more and more intersections.

5.1 Proof of Theorem 2.5

Proof. For any pair of integers $(r, s) \neq (0, 0)$, define $(r, s)_T = T_d((r/d, s/d))$ where $d = \gcd(r, s)$. For convenience, let $(0, 0)_T = 2$. In this notation, the basis $B_{\hat{T}}$ is

$$B_{\hat{T}} = \{(r, s)_T : (r, s) \neq (0, 0)\} \cup \{1\}.$$

Note $(r, s)_T = (-r, -s)_T$. The structure constants of the skein algebra of the torus in the basis $B_{\hat{T}}$ were computed by Frohman and Gelca [5],

$$(r, s)_T(u, v)_T = q^{rv-us}(r+u, s+v)_T + q^{-(rv-us)}(r-u, s-v)_T,$$

which shows that $(\hat{T}_n)$ is positive.

By Theorem 2.4, one has $(P_n) \geq (\hat{T}_n)$ and $P_1(x) = x = T_1(x)$. To show the opposite inequality, consider

$$P_1((n, 1))P_1((0, 1)) = (n, 1)_T(0, 1)_T = q^n(n, 2)_T + q^{-n}(n, 0)_T.$$

First, let $n = 2$. Write $T_2(x) = P_2(x) + \delta_1 P_1(x) + \delta_0$. Then,

$$\begin{aligned} P_1((2, 1))P_1((0, 1)) &= q^2(2, 2)_T + q^{-2}(2, 0)_T \\ &= q^2(P_2((1, 1)) + \delta_1 P_1((1, 1)) + \delta_0) \\ &\quad + q^{-2}(P_2((1, 0)) + \delta_1 P_1((1, 0)) + \delta_0). \end{aligned}$$

By the positivity of $(P_n(x))$, δ_1 and $(q^2 + q^{-2})\delta_0$ are in R_+ . On the other hand, $P_2(x) = T_2(x) - \delta_1 T_1(x) - \delta_0$ implies $-\delta_1, -\delta_0 \in R_+$. Thus, $\delta_1 = 0 = \delta_0$ and $P_2(x) = T_2(x)$.

For $n > 2$, $(n, 2)_T$ is either $P_1((n, 2))$ or $P_2((n/2, 1))$. Then, by the positivity of (P_n) , $(n, 0)_T = T_n((1, 0))$ is an R_+ -linear combination of $\{P_k((1, 0))\}$. Thus, $(P_n) \leq (\hat{T}_n)$. ■

5.2 Theorem 2.6, punctured torus case

Assume $\Sigma = \Sigma_{1,1}$. Define $T_{r,s} = T_d((r/d, s/d))$, $S_{r,s} = S_d((r/d, s/d))$ for $(r, s) \neq (0, 0)$, where $d = \gcd(r, s)$. Let $T_{0,0} = S_{0,0} = 1$. The products on $\Sigma_{1,1}$ are more complicated than the $\Sigma_{1,0}$ case. No general formula is available yet. However, to prove the theorem, only special cases are needed.

Lemma 5.1. If the curves $(r/d, s/d)$ and (u, v) intersect once, where $d = \gcd(r, s)$, then

$$T_{r,s}T_{u,v} = q^{rv-su}T_{r+u,s+v} + q^{-(rv-su)}T_{r-u,s-v}. \quad (3)$$

Proof. There is a diffeomorphism of $\Sigma_{1,1}$ sending the curves $(r/d, s/d)$ and (u, v) to the curves $(1, 0)$ and $(0, 1)$, respectively. Then, (3) becomes

$$T_{d,0}T_{0,1} = q^dT_{d,1} + q^{-d}T_{d,-1}.$$

This is essentially a reformulation of Proposition 3.1 in [7]. ■

Lemma 5.2. On $\Sigma_{1,1}$, let U be the peripheral curve. Then,

$$T_{1,0}T_{n,2} = q^2T_{n+1,2} + q^{-2}T_{n-1,2} + (U + q^2 + q^{-2})A_n,$$

where $A_n = 0$ if n is even, and $A_n = 1$ if n is odd.




Fig. 5. Four resolutions of $T_{1,0}T_{1,2}$.

Proof. If n is even, then $(n/2, 1)$ and $(1, 0)$ intersect once. In this case, the formula is a specialization of Lemma 5.1. When $n = 1$, the curves $(1, 0)$ and $(1, 2)$ intersect twice as in Figure 5.

Opposite sides of the squares are identified, and the corners represent the puncture. Resolving both crossings, we get

$$\begin{aligned} T_{1,0}T_{1,2} &= q^2(1, 1)^2 + U + (-q^2 - q^{-2}) + q^{-2}(0, 1)^2 \\ &= q^2T_{2,2} + q^{-2}T_{0,2} + U + q^2 + q^{-2}. \end{aligned}$$

For a general odd n , apply $(n-1)/2$ Dehn twists along $(1, 0)$ to the equation above gives

$$T_{1,0}T_{n,2} = q^2T_{n+1,2} + q^{-2}T_{n-1,2} + (U + q^2 + q^{-2}).$$

■

Lemma 5.3. On $\Sigma_{1,1}$,

$$T_{n,1}T_{0,1} = q^nT_{n,2} + q^{-n}T_{n,0} + (U + q^2 + q^{-2})G_n,$$

where $G_0 = G_1 = 0$, and

$$G_n = \sum_{i=1}^{\lfloor n/2 \rfloor} q^{4i-n-2} S_{n-2i,0}$$

if $n > 1$.

Proof. When $n = 0, 1$, the result follows from direct calculation. To use induction, note $(n, 1)$ and $(1, 0)$ intersect at one point. Thus, $T_{1,0}T_{n,1} = qT_{n+1,1} + q^{-1}T_{n-1,1}$.

$$\begin{aligned}
 T_{n+1,1}T_{0,1} &= (q^{-1}T_{n,1}T_{1,0} - q^{-2}T_{n-1,1})T_{0,1} \\
 &= q^{-1} \left(q^n T_{n,2} + q^{-n} T_{n,0} + (U + q^2 + q^{-2})G_n \right) T_{1,0} \\
 &\quad - q^{-2} \left(q^{n-1} T_{n-1,2} + q^{1-n} T_{n-1,0} + (U + q^2 + q^{-2})G_{n-1} \right) \\
 &= q^{n-1} (q^2 T_{n+1,2} + q^{-2} T_{n-1,2} + (U + q^2 + q^{-2})A_n) \\
 &\quad + q^{-n-1} (T_{n+1,0} + T_{n-1,0}) + (U + q^2 + q^{-2})q^{-1}G_n T_{1,0} \\
 &\quad - q^{n-3} T_{n-1,2} - q^{-n-1} T_{n-1,0} - (U + q^2 + q^{-2})q^{-2}G_{n-1} \\
 &= q^{n+1} T_{n+1,2} + q^{-n-1} T_{n+1,0} + (U + q^2 + q^{-2})(q^{-1}G_n T_{1,0} - q^{-2}G_{n-1} + q^{n-1}A_n) \\
 &= q^{n+1} T_{n+1,2} + q^{-n-1} T_{n+1,0} + (U + q^2 + q^{-2})G_{n+1},
 \end{aligned}$$

where the last equality can be directly verified using the expression of G_n . Thus, the formula holds by induction. $\blacksquare$

Proof of Theorem 2.6, punctured torus case. Since (P_n) is positive, $P_1(t) = t$. Thus,

$$P((n, 1))P((0, 1)) = T_{n,1}T_{0,1}$$

is an R_+ -linear combination of basis elements in B_P . Since (P_n) has integer coefficients, terms with different exponents of q are separately $\mathbb{Z}_+$ -linear combinations of basis elements in B_P . Rearranging the product in terms of the exponents of q , we have

$$P((n, 1))P((0, 1)) = q^{-n}S_n((1, 0)) + (\text{higher degree in } q).$$

Therefore, $(P_n) \leq (S_n)$. $\blacksquare$

5.3 Theorem 2.6, sphere with four punctures case

The proof is similar to the punctured torus case. Define $S_{r,s} = S_d((r/d, s/d))$ for $(r, s) \neq (0, 0)$, where $d = \gcd(r, s)$. Let $S_{0,0} = 1$.

Recall that γ_i is the curve surrounding puncture i , and $c_0 = \gamma_1\gamma_3 + \gamma_2\gamma_4$. $\sigma = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ represents the counterclockwise half twist along a , fixing punctures 3 and 4 and exchanging punctures 1 and 2. $c_k = \sigma^k(c_0)$.

Lemma 5.4. On $\Sigma_{0,4}$,

$$\begin{aligned} S_{1,0}S_{2k,2} &= q^4S_{2k+1,2} + q^{-4}S_{2k-1,2} + c_kS_{k,1} + [S_{1,0} + (q^2 + q^{-2})(\gamma_1\gamma_2 + \gamma_3\gamma_4)], \\ S_{1,0}S_{2k+1,2} &= q^4S_{2k+2,2} + q^{-4}S_{2k,2} + (q^2c_kS_{k+1,1} + q^{-2}c_{k+1}S_{k,1}) + \Gamma. \end{aligned}$$

where $\Gamma = \gamma_1\gamma_2\gamma_3\gamma_4 + \gamma_1^2 + \gamma_2^2 + \gamma_3^2 + \gamma_4^2 - 2$.

Proof. By direct computation,

$$\begin{aligned} S_{1,0}S_{0,2} &= q^4S_{1,2} + q^{-4}S_{-1,2} + c_0S_{0,1} + [S_{1,0} + (q^2 + q^{-2})(\gamma_1\gamma_2 + \gamma_3\gamma_4)], \\ S_{1,0}S_{1,2} &= q^4S_{2,2} + q^{-4}S_{0,2} + (q^2c_0S_{1,1} + q^{-2}c_1S_{0,1}) + \Gamma. \end{aligned}$$

The equations follow by applying σ^k . ■

Lemma 5.5 (see also [1] Theorem 3.6). On $\Sigma_{0,4}$, for $n \geq 0$,

$$S_{n,1}S_{0,1} = q^{2n}S_{n,2} + q^{-2n}S_{n,0} + g_n + h_n.$$

Here, $g_0 = g_1 = 0$, and for $n \geq 2$,

$$g_n = \sum_{i=1}^{\lfloor n/2 \rfloor} q^{4i-2} \sum_{j=i}^{n-i} c_{n-j+1}S_{j,1}.$$

h_n is a polynomial of q , $S_{1,0}$ and γ_i 's only. $h_0 = 0$, and for $n \geq 1$, the exponents of q in h_n are between $-2n + 2$ and $2n - 2$.

Proof. When $n = 0$, the equation clearly holds. When $n = 1$, resolving both crossings of $S_{1,1}$ and $S_{0,1}$, yields

$$S_{1,1}S_{0,1} = q^2S_{1,2} + q^{-2}S_{1,0} + (\gamma_1\gamma_2 + \gamma_3\gamma_4).$$

Thus, the equation holds with $h_1 = \gamma_1\gamma_2 + \gamma_3\gamma_4$.

When $n > 1$,

$$\begin{aligned}
S_{n+1,1}S_{0,1} &= q^{-2}S_{1,0}(S_{n,1}S_{0,1}) - q^{-4}S_{n-1,1}S_{0,1} - q^{-2}c_nS_{0,1} \\
&= q^{-2}S_{1,0}(q^{2n}S_{n,2} + q^{-2n}S_{n,0} + g_n + h_n) \\
&\quad - q^{-4}(q^{2n-2}S_{n-1,2} + q^{2-2n}S_{n-1,0} + g_{n-1} + h_{n-1}) - q^{-2}c_nS_{0,1}. \\
&= q^{2n-2}S_{1,0}S_{n,2} + q^{-2n-2}(S_{n+1,0} + S_{n-1,0}) + q^{-2}S_{1,0}g_n + q^{-2}S_{1,0}h_n \\
&\quad - q^{2n-6}S_{n-1,2} - q^{-2n-2}S_{n-1,0} - q^{-4}g_{n-1} - q^{-4}h_{n-1} - q^{-2}c_nS_{0,1} \\
&= (q^{2n-2}S_{1,0}S_{n,2} - q^{2n-6}S_{n-1,2}) + q^{-2n-2}S_{n+1,0} \\
&\quad + (q^{-2}S_{1,0}g_n - q^{-4}g_{n-1} - q^{-2}c_nS_{0,1}) + (q^{-2}S_{1,0}h_n - q^{-4}h_{n-1}).
\end{aligned}$$

To continue, apply Lemma 5.4 to the 1st term. For the product $S_{1,0}g_n$, we can use Equation 2, written in the notations of this section

$$S_{1,0}S_{n,1} = q^2S_{n+1,1} + q^{-2}S_{n-1,1} + c_n.$$

After a routine reduction, the product $S_{n,1}S_{0,1}$ has the desired form. ■

Proof of Theorem 2.6, sphere with 4 punctures case. In the product $(n, 1)(0, 1) = S_{n,1}S_{0,1}$, the terms with the lowest q -exponent is

$$q^{-2n}S_{n,0} = q^{-2n}S_n((1, 0))$$

for $n > 0$. This shows that $S_n((1, 0))$ is a $\mathbb{Z}_+$ -linear combination of $\{P_k((1, 0))\}$. Thus, $(P_n) \leq (S_n)$. This completes the proof of Theorem 2.6. ■

Funding

This work was supported in part by NSF grants. T. Le is supported by the National Science Foundation, Grant DMS-1811114. D. Thurston is supported by the National Science Foundation, Grant DMS-1507244.

Acknowledgments

The 1st author would like to thank the CIMI Excellence Laboratory, Toulouse, France, for inviting him on a Excellence Chair during the period of January–July 2017 when part of this work was done.

References

- [1] Bakshi, R. P., S. Mukherjee, J. H. Przytycki, M. Silvero, and X. Wang. "On multiplying curves in the Kauffman bracket skein algebra of the thickened four-holed sphere." (2018): preprint arXiv:1805.06062.
- [2] Cho, S. Y., H. Kim, H. Kim, and D. Oh. "Laurent positivity of quantized canonical bases for quantum cluster varieties from surfaces." *Comm. Math. Phys.* 373, no. 2 (2020): 655–705.
- [3] Davison, B. "Positivity for quantum cluster algebras." *Ann. of Math. (2)* 187, no. 1 (2018): 157–219.
- [4] Fock, V. and A. Goncharov. "Moduli spaces of local systems and higher Teichmüller theory." *Publ. Math. Inst. Hautes Études Sci. No.* 103 (2006): 1–211.
- [5] Frohman, C. and R. Gelca. "Skein modules and the noncommutative torus." *Trans. Amer. Math. Soc.* 352, no. 10 (2000): 4877–88.
- [6] Kauffman, L. "States models and the Jones polynomial." *Topology* 26, no. 3 (1987): 395–407.
- [7] Lê, T. T. Q. "On positivity of Kauffman bracket skein algebras of surfaces." *Int. Math. Res. Not. IMRN* 2018, no. 5 (2018): 1314–28.
- [8] Lee, K. and R. Schiffler. "Positivity for cluster algebras." *Ann. of Math. (2)* 182, no. 1 (2015): 73–125.
- [9] Luo, F. "Simple loops on surfaces and their intersection numbers." *J. Differential Geom.* 85, no. 1 (2010): 73–116.
- [10] Musiker, G., R. Schiffler, and L. Williams. "Bases for cluster algebras from surfaces." *Compos. Math.* 149, no. 2 (2013): 217–63.
- [11] Przytycki, J. H. "Fundamentals of Kauffman bracket skein modules." *Kobe J. Math.* 16, no. 1 (1999): 45–66.
- [12] Queffelec, H. and P. Wedrich. "Khovanov homology and categorification of skein modules." (2018): preprint arXiv:1806.03416.
- [13] Thurston, D. "Positive basis for surface skein algebras." *Proc. Natl. Acad. Sci. USA* 111, no. 27 (2014): 9725–32.
- [14] Turaev, V. "Conway and Kauffman modules of a solid torus." *J. Soviet. Math.* 52, no. 1 (1990): 2799–805.
- [15] Turaev, V. "Skein quantization of Poisson algebras of loops on surfaces." *Ann. Sci. Sc. Norm. Sup.* 24, no. 6 (1991): 635–704.