

LIGHT GROUPS OF ISOMORPHISMS OF BANACH SPACES AND INVARIANT LUR RENORMINGS

L. ANTUNES, V. FERENCZI, S. GRIVAUX, AND C. ROSENDAL

ABSTRACT. Megrelishvili defines in [15] *light groups* of isomorphisms of a Banach space as the groups on which the Weak and Strong Operator Topologies coincide, and proves that every bounded group of isomorphisms of Banach spaces with the Point of Continuity Property (PCP) is light. We investigate this concept for isomorphism groups G of classical Banach spaces X without the PCP, specially isometry groups, and relate it to the existence of G -invariant LUR or strictly convex renormings of X .

1. INTRODUCTION

The general objective of this note is to determine conditions on a bounded group of isomorphisms of Banach spaces that ensure the existence of a locally uniformly rotund (LUR) renorming invariant under the action of this group. In particular, we will be interested in this context in the notion of lightness for such groups.

1.1. Light groups. A frequent problem in functional analysis is to determine under which conditions weak convergence and norm convergence coincide. For example, it is well-known that conditions of convexity of the norm of a Banach space ensure that weak and strong convergence are equivalent on its unit sphere. The corresponding problem for isomorphisms of Banach spaces (or more generally of locally convex spaces) was studied by Megrelishvili in [15] in the context of group representations, using the concept of fragmentability.

Given a (real) Banach space X , we denote by $L(X)$ the set of bounded linear operators on X , and by $GL(X)$ the group of bounded isomorphisms of X . We also denote by $\text{Isom}(X)$ the group of surjective linear isometries of X . If G is a subgroup of $GL(X)$, we write $G \leq GL(X)$. Recall that given a Banach space X , the Strong Operator Topology on $L(X)$ is the topology of pointwise convergence, i.e. the initial topology generated by the family of functions $f_x: L(X) \rightarrow X$, $x \in X$, given by $f_x(T) = Tx$, $T \in L(X)$, and the Weak Operator Topology on $L(X)$ is generated by the family of functions $f_{x,x^*}: L(X) \rightarrow \mathbb{R}$, $x \in X$, $x^* \in X^*$, given by $f_{x,x^*}(T) = x^*(Tx)$, $T \in L(X)$.

Megrelishvili gives the following definition.

Date: October 30, 2017.

1991 Mathematics Subject Classification. 22F50, 46B03.

Key words and phrases. groups of isomorphisms of Banach spaces, isometry groups, light groups, renormings of Banach spaces, LUR renormings, invariant renormings.

The first author was fully supported by UTFPR (Process 23064.004102/2015-40). The second author was supported by CNPq, grant 303034/2015-7 and Fapesp, grants 2013/11390-4 and 2016/25574-8. The third author was supported in part by EU IRSES grant AOS (PIRSES-GA-2012-318910) and by the Labex CEMPI (ANR-11-LABX-0007-01). The fourth author was supported by the NSF (DMS #1464974).

Definition 1.1 ([15]). *A group $G \leq GL(X)$ of isomorphisms on a Banach space X is light if the Weak Operator Topology (WOT) and the Strong Operator Topology (SOT) coincide on G .*

Observe that since the two operator topologies are independent of the specific choice of norm on X , the same holds for lightness of G .

Obvious examples of light groups are the group $U(H)$ of unitary operators on a Hilbert space H , as well as the bounded amenable subgroups of $GL(H)$. However, the main result of [15] concerning light groups indicates that the restriction of amenability is not needed. Recall that a Banach space X has the Point of Continuity Property (PCP) if for every norm-closed non-empty bounded subset C of X , the identity on C has a point of continuity from the weak to the norm topology:

Theorem 1.2 ([15]). *If X is a Banach space with the Point of Continuity Property (PCP) and if $G \leq GL(X)$ is bounded in norm, then G is light.*

In particular, if X has the Radon-Nikodym Property (RNP) (e.g. if X is reflexive or is a separable dual space), then every bounded subgroup of $GL(X)$ is light.

We note here that in the literature (and indeed in [15]) PCP sometimes appears as the formally weaker condition “every weakly-closed non-empty bounded subset has a weak-to-norm point of continuity for the identity”. However, as was pointed out to us by G. Godefroy, if X satisfies the second definition and F is norm-closed and bounded, it is readily seen that any point of continuity of the weak closure $\overline{F}^w$ belongs to F , so the two definitions are equivalent.

1.2. Bounded non-light groups. A natural question that arises from Megrelishvili’s result is to investigate in which respect his result is optimal, and whether “smallness” assumptions on G or weaker assumptions than the PCP on X could imply that G is light. After observing that SOT-compact groups G of isomorphisms are light (Proposition 2.3), we show (Theorem 4.6) that any separable space containing an isomorphic copy of c_0 admits a bounded cyclic group of isomorphisms which is not light. This shows that we cannot really expect further general results in this direction.

Megrelishvili gives as example of a non-light group the group $\text{Isom}(C([-1, 1]^2))$. His proof uses a construction of Helmer [12] of a separately continuous group action on $[-1, 1]^2$ that is not jointly continuous, and the equivalence of pointwise compactness and weak compactness of bounded subsets of $C(K)$. This leads us to look for further examples of non-light isometry groups, for instance within the class of spaces $C(K)$, where K is an infinite compact space. We first prove (Proposition 4.3) that the isometry group of c , the space of real convergent sequences, is not light. Neither is the isometry group of $C(\{0, 1\}^{\mathbb{N}})$ (Proposition 5.5). On the other hand, as a consequence of Theorem 5.7, we show that the isometry group of $C[0, 1]$ is light, while those of the spaces $C([0, 1]^n)$, $n \geq 2$, are not light. These constructions simplify the initial example of Megrelishvili.

1.3. Light groups and LUR renormings. In another direction, we study the relation between light groups and the existence of LUR renormings invariant under the action of the group. Recall that a norm $\|\cdot\|$ on X is *rotund* or *strictly convex* if whenever the vectors x, y belong to the unit sphere S_X of X and $\|x + y\| = 2$, $x = y$. It is *locally uniformly convex* (LUC) or *locally uniformly rotund* (LUR) at a vector $x_0 \in X$ if whenever $(x_n)_{n \in \mathbb{N}}$ is a sequence of vectors of X such that $\lim \|x_n\| = \|x_0\|$ and $\lim \|x_0 + x_n\| = 2\|x_0\|$,

$\lim \|x_n - x_0\| = 0$, and LUR if it is LUR at every point x_0 of X (or, equivalently, of S_X). The property of the dual norm $\|\cdot\|_*$ on X^* being strictly convex or LUR is closely related to the differentiability of the norm $\|\cdot\|$ on X , in the senses of Fréchet and Gateaux respectively. All (real) separable Banach spaces admit an equivalent LUR renorming. For this and much more on renormings of Banach spaces, see [4].

A fundamental result in the study of LUR renorming is the following theorem, due to Lancien (see [14]):

Theorem 1.3 ([14]). *If X is a separable Banach space with the RNP, X admits an isometry invariant LUR renorming.*

If $G \leq \text{GL}(X, \|\cdot\|)$ is a bounded group of isomorphisms on X , the norm on X defined by

$$\|x\| = \sup_{g \in G} \|gx\|, \quad x \in X,$$

is a G -invariant renorming of X . In other words, $G \leq \text{Isom}(X, \|\cdot\|)$. So a consequence of Lancien's Theorem 1.3 is that whenever X is a separable space with the RNP and G is a bounded group of isomorphisms on X , there exists a G -invariant LUR renorming of X . The existence of G -invariant LUR renormings for general groups of isomorphisms G was first investigated by Ferenczi and Rosendal in [7]. In this paper, problems of maximal symmetry in Banach spaces were studied, analyzing the structure of subgroups of $\text{GL}(X)$ when X is a separable reflexive Banach space. An example of a super-reflexive space with no maximal bounded group of isomorphisms was also exhibited in [7].

The relation between light groups and G -invariant LUR renormings is given by Proposition 2.2. We observe that if a Banach space X admits a G -invariant LUR renorming, then G is light. In fact, this is true even if the norm is LUR only on a dense subset of S_X . We also show that the converse assertion is false: although the isometry group of $C[0, 1]$ is light, $C[0, 1]$ admits no strictly convex isometry invariant renorming (Corollary 5.10).

1.4. Light groups and distinguished families. In [6], Ferenczi and Galego investigated groups that may be seen as the group of isometries of a Banach space under some renorming. Among other results, they prove that if X is a separable Banach space and G is a finite group of isomorphisms of X such that $-\text{Id} \in G$, X admits an equivalent norm $\|\cdot\|$ such that $G = \text{Isom}(X, \|\cdot\|)$. They also prove that if X is a separable Banach space with LUR norm $\|\cdot\|$ and if G is an isometry group of X such that $-\text{Id} \in G$ and such that G admits a point $x \in X$ with $\inf_{g \neq \text{Id}} \|gx - x\| > 0$, then $G = \text{Isom}(X, \|\cdot\|)$ for some equivalent norm $\|\cdot\|$ on X . A point x satisfying the condition

$$\inf_{g \neq \text{Id}} \|gx - x\| > 0$$

is called in [8] a *distinguished point* of X for the group G .

In [8], Ferenczi and Rosendal generalized results of [6] to certain uncountable Polish groups, and also defined the concept of *distinguished family* for the action of a group G on a Banach space X . It is clear that if G is an isometry group with a distinguished point, G is SOT-discrete. However, the following question remained open: if G is an isomorphism group of X which is SOT-discrete, should X have a distinguished point for G ? In Proposition 6.1 we will see that the answer to this question is negative, and will give an example of an infinite countable group of isomorphisms G of c_0 which is SOT-discrete but does not admit a distinguished point for G . In addition, this group is also not light.

1.5. Light groups on quasi-normed spaces. Although Megrelishvili has defined the concept of light group only for groups of isomorphisms on locally convex spaces, we can extend the definition to quasi-normed spaces, even if they are not locally convex. We finish this article by investigating whether the isometry groups of the quasi-normed spaces ℓ_p and $L_p[0, 1]$, $0 < p < 1$, are light.

2. LUR RENORMINGS AND LIGHT GROUPS

Let G be a bounded group of isomorphisms on a Banach space $(X, \|\cdot\|)$. In this section we are interested in the existence of a G -invariant LUR renorming of X , i.e. in the existence of an equivalent norm $\|\cdot\|$ on X which is both invariant under the action of G and is LUR; or simply in the existence of a G -invariant *dense LUR renorming*, meaning a renorming which is invariant under the action of G and is LUR on a dense subset of S_X . When $G = \text{Isom}(X, \|\cdot\|)$ we shall talk about *isometry invariant* renormings.

Proposition 2.1. *Let X be a Banach space and let $G \leq \text{GL}(X)$. If G is SOT-compact and if X admits an LUR renorming, X admits a G -invariant LUR renorming.*

Proof. Suppose that $\|\cdot\|$ is an equivalent LUR norm on X . The formula

$$\|x\| = \sup_{T \in G} \|Tx\|, \quad x \in X,$$

defines a G -invariant LUR renorming of X . Indeed, suppose that x_n , $n \in \mathbb{N}$, and x are vectors of X such that $\|x_n\| = \|x\| = 1$ for every $n \in \mathbb{N}$ and $\lim \|x_n + x\| = 2$. Then we can find elements T_n , $n \in \mathbb{N}$, of G such that $\lim \|T_n x_n + T_n x\| = 2$. By SOT-compactness of G we can assume without loss of generality that T_n tends SOT to some element T of G , from which it follows that $\|T_n x_n + Tx\|$ converges to 2. Since $\|T_n x_n\| \leq \|x_n\| = 1$ for every $n \in \mathbb{N}$ and $\|Tx\| \leq \|x\| = 1$, we deduce that $\|Tx\| = 1$ and that $\|T_n x_n\|$ converges to 1. In particular, if we set, for every $n \in \mathbb{N}$, $y_n = \frac{T_n x_n}{\|T_n x_n\|}$, then y_n belongs to the unit sphere of $(X, \|\cdot\|)$ and $\|y_n + Tx\|$ converges to 2. By the LUR property of $\|\cdot\|$ at the point Tx , we deduce that y_n converges to Tx . This means that $T_n x_n$ converges to Tx . So $\|x_n - x\| = \|T_n x_n - T_n x\|$ converges to 0 since both $T_n x_n$ and $T_n x$ converge to Tx . This shows that $\|\cdot\|$ is LUR. $\square$

In [8], Ferenczi and Rosendal investigate LUR renormings in the context of transitivity of norms. Recall that a norm $\|\cdot\|$ on X is called *transitive* if the orbit $\mathcal{O}(x)$ of every point $x \in S_X$ under the action of the isometry group $\text{Isom}(X)$ is the whole sphere S_X . If for every $x \in X$ the orbit $\mathcal{O}(x)$ is dense in S_X we say that $\|\cdot\|$ is *almost transitive*, and if the closed convex hull of $\mathcal{O}(x)$ is the unit ball B_X , we say that $\|\cdot\|$ is *convex transitive*. Ferenczi and Rosendal proved that if an almost transitive norm on a Banach space is LUR at some point of the unit sphere, it is uniformly convex. They also proved that if a convex transitive norm on a Banach space is LUR on a dense subset of the unit sphere, it is almost transitive and uniformly convex.

In the next proposition, we exhibit a relation between the existence of LUR renormings and light groups.

Proposition 2.2. *Let $G \leq \text{GL}(X)$ be a group of isomorphisms of a Banach space X . If X admits a G -invariant renorming which is LUR on a dense subset of S_X , G is light.*

Proof. Let $\|\cdot\|$ be a G -invariant renorming of X which is LUR on a dense subset of S_X . Let (T_α) be a net of elements of G which converges WOT to the identity operator Id on X , and assume that T_α does not converge SOT to Id . Let $x \in S_X$ be such that $T_\alpha x$ does

not converge to x . Without loss of generality, we can suppose that the norm is LUR at x , and that there exists $\delta > 0$ such that, for every α , $\|T_\alpha x - x\| \geq \delta$. By the LUR property, there exists $\varepsilon > 0$ such that $\|T_\alpha x + x\| \leq 2 - \varepsilon$ for every α . Let $\phi \in X^*$ with $\|\phi\| = 1$ be such that $\phi(x) = 1$. Since T_α converges WOT to Id, $\phi(T_\alpha x) \rightarrow 1$. On the other hand,

$$2 - \varepsilon \geq \|T_\alpha x + x\| = \max_{\substack{\psi \in X^* \\ \|\psi\|=1}} |\psi(T_\alpha x + x)| \geq |\phi(T_\alpha x) + 1| \quad \text{for every } \alpha,$$

which contradicts the WOT convergence of T_α to Id. $\square$

Since every separable space admits an equivalent LUR renorming, it follows by combining Propositions 2.1 and 2.2 that every SOT-compact group $G \leq \text{GL}(X)$ of a separable Banach space X is light. In fact, this result holds true even for non-separable spaces, as observed in the next proposition:

Proposition 2.3. *Let $G \leq \text{GL}(X)$ be a group of isomorphisms of a Banach space X . If G is SOT-compact, G is light.*

Proof. The assumption implies that G is also WOT compact, since the WOT is weaker than the SOT. However, the WOT is also Hausdorff, and so the two topologies must agree on G . In other words, G is light. $\square$

3. LIGHT GROUPS AND DISTINGUISHED POINTS

As recalled in the introduction, Lancien proved in [14] that if X is separable with the RNP, X admits an isometry invariant LUR renorming. Although separable spaces always admit LUR renormings, the generalization of Lancien's result to all separable spaces is false. For example, if $X = C([-1, 1]^2)$ and $G = \text{Isom}(X)$ then, since G is not light [15], there is by Proposition 2.2 no equivalent G -invariant (not even dense) LUR renorming. Another example mentioned in [8] is the case where $X = L_1[0, 1]$ and $G = \text{Isom}(L_1[0, 1])$. In this case there is not even a strictly convex G -invariant renorming.

Here we discuss conditions which clarify the relations between the two properties of a group $G \leq \text{GL}(X)$ being light and X having a G -invariant LUR renorming, in the case when G is SOT-discrete. The following notion was defined in [8].

Definition 3.1. *Let X be a Banach space, let $G \leq \text{GL}(X)$ be a bounded group of isomorphisms of X , and let $x \in X$. We say that x is distinguished for G (or for the action of G on X) if*

$$\inf_{T \neq \text{Id}} \|Tx - x\| > 0.$$

If $\{x_1, \dots, x_n\}$ is a finite family of vectors of X , then it is distinguished for G if

$$\inf_{T \neq \text{Id}} \max_{1 \leq i \leq n} \|Tx_i - x_i\| > 0,$$

or, equivalently, if the n -tuple $(x_1, \dots, x_n)$ is distinguished for the canonical action of G on X^n .

This notion does not depend on the choice of an equivalent norm on X . Note also that G is SOT-discrete exactly when it admits a distinguished finite family of vectors. We also have, considering the adjoint action of G on X^* :

Lemma 3.2. *Assume that $G \leq \text{GL}(X)$ is light. If G acts as an SOT-discrete group on X , then G acts as an SOT-discrete group on X^* .*

Proof. Define $\psi : G \rightarrow \text{GL}(X^*)$ by setting $\psi(T)(x^*) = x^* \circ T^{-1}$ for every $T \in G$ and $x^* \in X^*$. We want to show that $\psi(G)$ is an SOT-discrete subgroup of $\text{GL}(X^*)$. For this it suffices to show the existence of $\varepsilon > 0$ and $x_1^*, \dots, x_n^* \in S_{X^*}$ such that the only element T of G such that $\|\psi(T)(x_i^*) - x_i^*\| < \varepsilon$ for every $1 \leq i \leq n$ is the identity operator Id_X on X . Since G is light and acts as an SOT-discrete group on X , it is WOT-discrete. So there exist $\varepsilon > 0$, $x_1, \dots, x_m \in S_X$ and $x_1^*, \dots, x_n^* \in S_{X^*}$ such that the only element T of G such that $|x_i^*(T^{-1}x_j - x_j)| < \varepsilon$ for every $1 \leq i \leq n$ and $1 \leq j \leq m$ is $T = \text{Id}_X$. The conclusion follows immediately. $\square$

Lemma 3.3. *Let X be a Banach space, let G be a bounded subgroup of $\text{GL}(X)$, and let $\{x_1, \dots, x_n\}$ be a distinguished family of vectors for the action of G on X . Let $\|\cdot\|$ be a G -invariant norm on X which is LUR at x_i for every $1 \leq i \leq n$. For any functional $x_i^* \in S_{X^*, \|\cdot\|}^*$ such that $x_i^*(x_i) = \|x_i\|$ for every $1 \leq i \leq n$, the family $\{x_1^*, \dots, x_n^*\}$ is distinguished for the action of G on X^* .*

Proof. Assume that $\|x_i\| = 1$ for every $1 \leq i \leq n$. Let $\alpha = \inf_{T \neq \text{Id}_X} \max_{1 \leq i \leq n} \|Tx_i - x_i\| > 0$. For every $T \neq \text{Id}_X$, choose $1 \leq i \leq n$ such that $\|T^{-1}x_i - x_i\| \geq \alpha$. By the LUR property of the norm at x_i , there exists $\varepsilon > 0$ depending on α but not on i such that $\|T^{-1}x_i + x_i\| \leq 2 - \varepsilon$. So $x_i^*(T^{-1}x_i) \leq 1 - \varepsilon$. From this it follows, using the notation introduced in the proof of Lemma 3.2 above, that $\psi(T)(x_i^*)(x_i) - x_i^*(x_i) \leq -\varepsilon$, so that $\|\psi(T)(x_i^*) - x_i^*\| \geq \varepsilon$. This being true for every $T \neq \text{Id}_X$, $\{x_1^*, \dots, x_n^*\}$ is distinguished for the action of G on X^* . $\square$

As a direct corollary, we obtain:

Corollary 3.4. *Let X be a Banach space, let $G \leq \text{GL}(X)$ be SOT-discrete, and assume that X admits a G -invariant dense LUR renorming. If there exists a distinguished family of cardinality n for the action of G on X , there also exists a distinguished family of cardinality n for the action of G on X^* .*

4. BOUNDED GROUPS WHICH ARE NOT LIGHT

Isometry groups are especially relevant to our study. We introduce the following definition:

Definition 4.1. *A Banach space X is light if $\text{Isom}(X)$ is a light subgroup of $\text{GL}(X)$.*

Observe that since the isometry group of a Banach space $(X, \|\cdot\|)$ is not invariant by equivalent renorming, the notion of lightness for a Banach space depends very much on the choice of the norm. In our terminology, Megrelishvili proves in [15] that all spaces with the PCP are light but that $C([0, 1]^2)$ is not light. Also, it is easy to observe the following fact:

Fact 4.2. *The space c_0 is light.*

Proof. Every isometry T of c_0 (endowed with the usual supremum norm) has the form

$$T((x_k)_{k \in \mathbb{N}}) = (\varepsilon_k x_{\sigma(k)})_{k \in \mathbb{N}}, \quad (x_k)_{k \in \mathbb{N}} \in c_0,$$

where $(\varepsilon_k)_{k \in \mathbb{N}} \in \{-1, 1\}^{\mathbb{N}}$ and σ is a permutation of $\mathbb{N}$. Using this, it is not difficult to check that whenever $(T_\alpha)_{\alpha \in I}$ is a net of isometries of c_0 which converges to Id in the WOT, it automatically converges to Id in the SOT. $\square$

Another proof of Fact 4.2 is based on the observation that c_0 admits a particular LUR renorming, namely the Day's renorming given by

$$\|x\|_D = \sup \left\{ \left(\sum_{k=1}^n \frac{x_{\sigma(k)}^2}{4^k} \right)^{\frac{1}{2}} \right\}, \quad x \in c_0,$$

where the supremum is taken over all $n \in \mathbb{N}$ and all permutations σ of $\mathbb{N}$ (see [4, p. 69]). Since this renorming is isometry invariant, it follows from Proposition 2.2 that c_0 is light.

Note that the Day's renorming is actually defined on ℓ_∞ , and therefore on the space c of convergent real sequences. In view of Proposition 4.3 below, it may be amusing to observe that Day's renorming is not strictly convex on c (not even on a dense subset of S_c). In fact, it is not strictly convex at the point $(1, 1, \dots)$ since for every $x = (x_k)_{k \in \mathbb{N}} \in c$ such that $\|x\|_\infty = 1$ and $|x_k| = 1$ for infinitely many indices k , we have $\|x\|_D = \|(1, 1, \dots)\|_D$.

We now provide an elementary example of a space which is not light.

Proposition 4.3. *There exists a subgroup $G \leq \text{Isom}(c)$ which has a distinguished point, but whose dual action on ℓ_1 is not SOT-discrete. In particular the space c is not light.*

Proof. Define G as the subgroup of isometries T of c of the form

$$T((x_k)_{k \in \mathbb{N}}) = (\varepsilon_k x_k)_{k \in \mathbb{N}}, \quad (x_k)_{k \in \mathbb{N}} \in c,$$

where the sequence $(\varepsilon_k)_k \in \{-1, 1\}^{\mathbb{N}}$ is eventually constant. One easily sees that $(1, 1, \dots)$ is a distinguished point for G . On the other hand, the dual space of c identifies isomorphically with ℓ_1 , where $\varphi = (y_k)_{k \in \mathbb{N}} \in \ell_1$ acts on an element $x = (x_k)_{k \in \mathbb{N}} \in c$ by the formula

$$\varphi(x) = y_1 \lim_{k \rightarrow \infty} x_k + \sum_{k=2}^{\infty} y_k x_{k-1}.$$

For every $n \in \mathbb{N}$, define the operator $T_n \in G$ by setting, for every $(x_k)_{k \in \mathbb{N}} \in c$,

$$T_n(x_1, x_2, \dots, x_{n-1}, x_n, x_{n+1}, \dots) = (x_1, x_2, \dots, x_{n-1}, -x_n, x_{n+1}, \dots).$$

Obviously $T_n \xrightarrow{\text{SOT}} \text{Id}$, but for every $n \in \mathbb{N}$ and every $x \in c$ we have

$$\varphi(T_n(x)) = y_1 \lim_{k \rightarrow \infty} (T_n(x))_k + \sum_{k=2}^{\infty} y_k (T_n(x))_{k-1} = \left(y_1 \lim_{k \rightarrow \infty} x_k + \sum_{k=2}^{\infty} y_k x_{k-1} \right) - 2y_n x_n$$

which tends to $\varphi(x)$ as n tends to infinity. Hence $T_n \xrightarrow{\text{WOT}} \text{Id}$ and G is not light, which implies that $\text{Isom}(c)$ itself is not light. Actually the inequality $|(T_n^* \varphi - \varphi)(x)| = 2|y_n x_n| \leq 2|y_n| \|x\|$, $x \in c$, $\varphi \in \ell_1$, implies that T_n^* tends SOT to Id , so the dual action of G on ℓ_1 is not SOT-discrete. $\square$

Remark 4.4. Note that the non-light subgroup G of $\text{Isom}(c)$ constructed in the proof above has the property that all its elements belong to the group $\text{Isom}_f(c)$ of isometries which are finite rank perturbations of the identity.

We observe the following relation between groups acting on a space and on a complemented subspace.

Lemma 4.5. *Assume Y embeds complementably in X . If every bounded group of isomorphisms on X is light, then every bounded group of isomorphisms on Y is light.*

Proof. Let Z be a closed subspace of X such that $X \simeq Y \oplus Z$. Let $G \leq GL(Y)$ be a bounded subgroup and for each $T \in G$, consider the operator $\tilde{T} = T \oplus \text{Id}_Y \in GL(X)$. These operators form a bounded subgroup $\tilde{G}$ of $GL(X)$ which is therefore light.

Let $(T_\alpha)_{\alpha \in I}$ be a net in G such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}_Y$. Then $\tilde{T}_\alpha \xrightarrow{\text{WOT}} \text{Id}_X$, and since $\tilde{G}$ is light, $\tilde{T}_\alpha \xrightarrow{\text{SOT}} \text{Id}_X$. Since for every $y \in Y$,

$$\|T_\alpha(y) - y\|_Y = \|\tilde{T}_\alpha(y, 0) - (y, 0)\|_X \rightarrow 0,$$

it follows that $T_\alpha \xrightarrow{\text{SOT}} \text{Id}_Y$. $\square$

Assume that X is separable and that $G \leq GL(X)$ is a bounded group of isomorphisms on X . As we have seen, if X either has the RNP or G is SOT-compact, then X admits a G -invariant LUR-renorming. It is natural to wonder whether the assumption on G may be weakened somewhat and, in particular, whether a similar result holds true for cyclic groups G . We show that it is not the case.

Theorem 4.6. *Let X be a separable Banach space containing an isomorphic copy of c_0 . Then $GL(X)$ contains a WOT-indiscrete bounded cyclic subgroup G with a distinguished point in X . In particular, G is not light.*

Proof. Consider the space $c(\mathbb{R}^2)$ of convergent sequences in the euclidean space $\mathbb{R}^2$ with the supremum norm. We define an isometry T of $c(\mathbb{R}^2)$ by setting

$$T((x_n)_{n \in \mathbb{N}}) = (R_n x_n)_{n \in \mathbb{N}} \quad \text{for every } x = (x_n)_{n \in \mathbb{N}} \in c(\mathbb{R}^2),$$

where, for every $n \in \mathbb{N}$,

$$R_n = \begin{pmatrix} \cos(\frac{2\pi}{n}) & -\sin(\frac{2\pi}{n}) \\ \sin(\frac{2\pi}{n}) & \cos(\frac{2\pi}{n}) \end{pmatrix}$$

is the rotation of $\mathbb{R}^2$ of angle $\frac{2\pi}{n}$. Observe that, since $\lim_n \frac{2\pi}{n} = 0$, we have

$$\lim_n T((x_n)_{n \in \mathbb{N}}) = \lim_n (R_n x_n)_{n \in \mathbb{N}} = \lim_n (x_n)_{n \in \mathbb{N}} \quad \text{for every } x = (x_n)_{n \in \mathbb{N}} \in c(\mathbb{R}^2).$$

As also $R_n^{k!} = \text{Id}_{\mathbb{R}^2}$ whenever $k \geq n$, we deduce that $T^{k!} \xrightarrow{\text{WOT}} \text{Id}$. So the cyclic subgroup $\langle T \rangle$ of $GL(c(\mathbb{R}^2))$ generated by T is indiscrete in the WOT.

On the other hand, if we define $x = (x_n)_{n \in \mathbb{N}} \in c(\mathbb{R}^2)$ by setting $x_n = (1, 0)$ for every $n \in \mathbb{N}$, we find that, for every $k \in \mathbb{N}$,

$$\|T^k x - x\|_{c(\mathbb{R}^2)} \geq \|R_{2k}^k x_{2k} - x_{2k}\|_2 = \|(-1, 0) - (1, 0)\|_2 = 2.$$

So x is a distinguished point for $\langle T \rangle$.

Observe now that $c(\mathbb{R}^2) \simeq c \oplus c \simeq c_0 \oplus c_0 \simeq c_0$, so T can be seen as an automorphism of c_0 . Also, if X is a separable Banach space containing c_0 , c_0 is complemented in X by Sobczyk's Theorem, i.e. X can be written as $X = c_0 \oplus Y$ for some subspace Y of X . Then Lemma 4.5 applies. Actually the group G generated by $S = T \oplus \text{Id}$ on X is not light, since $S^{k!} \xrightarrow{\text{WOT}} \text{Id}$, while G has a distinguished point in X . $\square$

Remark 4.7. It follows from Theorem 4.6 that any separable Banach space X containing an isomorphic copy of c_0 admits a renorming $\|\cdot\|$ such that $(X, \|\cdot\|)$ is not light.

We finish this section with the following observation:

Lemma 4.8. *Suppose G is an abelian group acting by isometries on a metric space (X, d) without isolated points, and inducing a dense orbit $G \cdot x$ for some element $x \in X$. Then, for every $\epsilon > 0$, there exists $g \in G \setminus \{1\}$ such that $\sup_{z \in X} d(gz, z) < \epsilon$.*

Proof. Indeed, since X has no isolated points and the orbit $G \cdot x$ is dense, we may pick $g \in G$ so that $0 < d(gx, x) < \epsilon$. For any y in $G \cdot x$, written $y = hx$ for $h \in G$, we have

$$d(gy, y) = d(ghx, hx) = d(hgx, hx) = d(gx, x) < \epsilon.$$

The result follows by density. $\square$

As a particular instance, note that if G is an SOT-discrete group of isometries of a Banach space X of dimension > 1 with a dense orbit on S_X , then G cannot be abelian.

5. LUR AND STRICTLY CONVEX ISOMETRY INVARIANT RENORMINGS

Proposition 2.2 leads to the following question:

Question 5.1. *Does there exist a light Banach space X which admits no isometry invariant LUR renorming?*

It was observed in [8] that $X = L_1[0, 1]$ does not admit any isometry invariant dense LUR renorming. In fact, since the norm of $L_1[0, 1]$ is almost transitive and is not strictly convex, any equivalent renorming is just a multiple of the original norm, so it is not strictly convex either, and hence is not LUR. Thus $L_1[0, 1]$ could be a natural example of a light Banach space which admits no isometry invariant LUR renorming. However,

Proposition 5.2. *The space $L_1[0, 1]$ is not light.*

Proof. For every $n \in \mathbb{N}$, define $\varphi_n : [0, 1] \rightarrow [0, 1]$ by setting

$$\varphi_n(t) = t + \frac{1 - \cos(2^n \pi t)}{2^n \pi}, \quad t \in [0, 1],$$

and $T_n : L_1[0, 1] \rightarrow L_1[0, 1]$ by

$$T_n(f)(t) = \varphi'_n(t) f(\varphi_n(t)), \quad f \in L_1[0, 1], \quad t \in [0, 1].$$

Note that φ_n is a differentiable bijection from $[0, 1]$ into itself. So T_n is a surjective linear isometry of $L_1[0, 1]$. Moreover, $T_n \xrightarrow{\text{SOT}} \text{Id}$, since for $f \equiv 1$ we have

$$\|T_n(1) - 1\|_1 = \|\sin(2^n \pi x)\|_1 = \frac{2}{\pi} \quad \text{for every } n \in \mathbb{N}.$$

On the other hand, $T_n \xrightarrow{\text{WOT}} \text{Id}$. To prove this, we need to check that

$$\int_0^1 T_n(f)(t) g(t) dt \longrightarrow \int_0^1 f(t) g(t) dt \quad \text{for every } f \in L_1[0, 1] \text{ and } g \in L_\infty[0, 1].$$

By the linearity of T_n and the density of step functions in $L_1[0, 1]$, it is sufficient to consider the case where f is the indicator function of a segment $I_{m,k} = [\frac{2k}{2^m}, \frac{2(k+1)}{2^m}]$, where $m \geq 1$ and $0 \leq k \leq 2^{m-1} - 1$. In this case the function φ_n is a bijection from $I_{m,k}$ into itself for every $n \geq m$. Thus $f \circ \varphi_n = f$, and

$$\begin{aligned} \int_0^1 T_n(f)(t) g(t) dt &= \int_0^1 \varphi'_n(t) f(\varphi_n(t)) g(t) dt = \int_0^1 \varphi'_n(t) f(t) g(t) dt \\ &= \int_0^1 f(t) g(t) dt + \int_0^1 \sin(2^n \pi t) f(t) g(t) dt. \end{aligned}$$

The result then follows from the Riemann-Lebesgue Lemma. $\square$

Remark 5.3. Another space of which it is well known that it does not admit any LUR renorming is ℓ_∞ . Indeed ℓ_∞ does not admit any equivalent norm with the Kadec-Klee property ([4, Ch. 2, Th. 7.10]), while every LUR norm satisfies the Kadec-Klee property ([4, Ch. 2, Prop. 1.4]). However, ℓ_∞ does admit a strictly convex renorming (see [5, p. 120]). We note here that it does not admit any isometry invariant strictly convex renorming. To see this, consider the points $x = (1, 1, 0, 1, 0, 1, 0, \dots)$ and $y = (-1, 1, 0, 1, 0, 1, 0, \dots)$. Setting $z = (x + y)/2 = (0, 1, 0, 1, 0, 1, 0, \dots)$, it is readily seen that there exist two isometries T and S of ℓ_∞ such that $Tx = y$ and $Sx = z$. So, for any isometry invariant renorming $\|\cdot\|$ of ℓ_∞ we have $\|x\| = \|y\| = \|z\|$, and therefore $\|\cdot\|$ cannot be strictly convex.

We now prove

Proposition 5.4. *The space ℓ_∞ is not light.*

Proof. Consider the sequence of isometries $T_n : \ell_\infty \rightarrow \ell_\infty$, $n \in \mathbb{N}$, defined by

$$T_n(x_1, \dots, x_{n-1}, x_n, x_{n+1}, \dots) = (x_1, \dots, x_{n-1}, -x_n, x_{n+1}, \dots), \quad x = (x_k)_{k \in \mathbb{N}} \in \ell_\infty.$$

Notice that $T_n \not\xrightarrow{\text{SOT}} \text{Id}$, since $\|T_n(1, 1, \dots) - (1, 1, \dots)\|_\infty = 2$ for every $n \in \mathbb{N}$. On the other hand, observe that $T_n \xrightarrow{\text{WOT}} \text{Id}$. Indeed, if $(e_j)_{j \in \mathbb{N}}$ denotes the canonical basis of ℓ_∞ , the sequence $(\beta(e_j))_{j \in \mathbb{N}}$ belongs to ℓ_1 for every $\beta \in \ell_\infty^*$. In particular, $\beta(e_j) \rightarrow 0$. Thus $\beta(T_n x - x) = -2x_n \beta(e_n) \rightarrow 0$ for every $x \in \ell_\infty$ and $\beta \in \ell_\infty^*$, showing that $T_n \xrightarrow{\text{WOT}} \text{Id}$. $\square$

A similar proof shows that the space of continuous functions on the Cantor space is not light.

Proposition 5.5. *The space $C(\{0, 1\}^\mathbb{N})$ is not light.*

Proof. For each integer $n \in \mathbb{N}$, define $T_n \in \text{Isom}(C(\{0, 1\}^\mathbb{N}))$ by setting, for every $f \in C(\{0, 1\}^\mathbb{N})$ and every $x \in \{0, 1\}^\mathbb{N}$,

$$T_n(f)(x) = \begin{cases} -f(x), & \text{if } x \in N_n, \\ f(x), & \text{otherwise,} \end{cases}$$

where N_n is the basic open subset of $\{0, 1\}^\mathbb{N}$ defined by

$$N_n = \{(x_k)_{k \in \mathbb{N}} \in \{0, 1\}^\mathbb{N}; \quad x_1 = \dots = x_{n-1} = 1, \quad x_n = 0\}.$$

Then $T_n \not\xrightarrow{\text{SOT}} \text{Id}$, since taking the constant function $f \equiv 1$ we have $\|T_n(f) - f\|_\infty = 2$ for every $n \in \mathbb{N}$. On the other hand, the same proof as that of Proposition 5.4 shows that $T_n \xrightarrow{\text{WOT}} \text{Id}$. Indeed, any functional $\Phi \in C(\{0, 1\}^\mathbb{N})^*$ extends to a continuous functional on $L^\infty(\{0, 1\}^\mathbb{N})$, also denoted by Φ . If χ_n denotes for each $n \in \mathbb{N}$ the indicator function of the set N_n , $\Phi(\chi_n) \rightarrow 0$, and this proves our claim. $\square$

So far we know that the spaces $C([0, 1]^2)$ and $C(\{0, 1\}^\mathbb{N})$ are not light. It may seem reasonable to conjecture that none of the spaces $C(K)$, where K is any infinite compact metric space, is light. However, our next result shows that this is not the case.

Proposition 5.6. *Let K be an infinite compact connected space. Then $C(K)$ is light if and only if the topologies of pointwise and uniform convergence coincide on the group $\text{Homeo}(K)$ of homeomorphisms of K .*

Proof. Suppose first that the topologies of pointwise and uniform convergence coincide on $\text{Homeo}(K)$. Let $(T_\alpha)_{\alpha \in I}$ be a net of isometries of $C(K)$ such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. By the Banach-Stone Theorem and the connectedness of K , each isometry T_α of $C(K)$ has the form

$$T_\alpha(f) = \varepsilon_\alpha f \circ \varphi_\alpha \quad \text{for every } f \in C(K),$$

where $\varepsilon_\alpha \in \{-1, 1\}$ and $\varphi_\alpha \in \text{Homeo}(K)$. Since $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$, $\varepsilon_\alpha \rightarrow 1$, so we can suppose without loss of generality that $\varepsilon_\alpha = 1$ for every $\alpha \in I$. Moreover, the fact that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$ also implies that the net $(\varphi_\alpha)_{\alpha \in I}$ converges pointwise to the identity function id_K on K . Our assumption then implies that $(\varphi_\alpha)_{\alpha \in I}$ converges uniformly to id_K on K , from which it easily follows that $T_\alpha \xrightarrow{\text{SOT}} \text{Id}$. Thus $C(K)$ is light.

Conversely, suppose that $C(K)$ is light. Let $(\varphi_\alpha)_{\alpha \in I}$ be a net of elements of $\text{Homeo}(K)$ which converges pointwise to $\varphi \in \text{Homeo}(K)$. Consider the isometries T_α and T of $C(K)$ defined by

$$T_\alpha(f) = f \circ \varphi_\alpha \quad \text{and} \quad T(f) = f \circ \varphi \quad \text{for every } f \in C(K).$$

Then $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. Since $C(K)$ is light, $T_\alpha \xrightarrow{\text{SOT}} \text{Id}$ and thus $(\varphi_\alpha)_{\alpha \in I}$ converges to φ uniformly on K . $\square$

Birkhoff studied in his paper [2] various topologies on so-called “transformation spaces”, in particular on the groups of homeomorphisms of topological spaces. The notions of A -, B - and C -convergence of sequences of homeomorphisms on a given space X introduced there correspond respectively to pointwise convergence, continuous convergence, and continuous convergence in both directions. Since on compact spaces continuous convergence and uniform convergence coincide, Proposition 5.6 can be rephrased, using Birkhoff’s language, as saying that for compact connected spaces K , $C(K)$ is light if and only if A - and B -convergence coincide on $\text{Homeo}(K)$. Now, it is observed in [2, Th. 18] that A -convergence implies B - and C -convergence for homeomorphisms of finite unions of segments of the real line (this is essentially the content of Dini’s second convergence theorem), while if K contains an n -dimensional region with $n \geq 2$ (i.e. an open set homeomorphic to an open subset of $\mathbb{R}^n$), A -convergence implies neither B - nor C -convergence for homeomorphisms of K ([2, Th. 19]). In a more modern language, there exists under this assumption a sequence $(\varphi_n)_{n \in \mathbb{N}}$ of homeomorphisms of K such that φ_n converges pointwise but not uniformly on K to the identity function on K . Combined with Proposition 5.6 above, this yields:

Theorem 5.7. *Let K be an infinite compact connected space.*

- (a) *If K is homeomorphic to a finite union of segments of $\mathbb{R}$, $C(K)$ is light.*
- (b) *If K contains an n -dimensional region for some $n \geq 2$, $C(K)$ is not light.*

For instance, the space $C[0, 1]$ is light, while spaces $C([0, 1]^n)$, $n \geq 2$, are not light. We thus retrieve in a natural way the original example of Megrelishvili of a non-light space.

Theorem 5.7 allows us to answer Question 5.1 in the negative. Although $C[0, 1]$ is light, it does not admit any isometry invariant LUR renorming. In fact, $C[0, 1]$ does not admit any isometry invariant strictly convex renorming. In order to prove this, we need the following lemma:

Lemma 5.8. *Let $f \in C[0, 1]$ be such that there exists an interval $[a, b] \subset [0, 1]$, $a < b$, on which f is strictly monotone. Then there exists $g \in C[0, 1]$ with the following three properties:*

- (a) $\|f\|_\infty = \|g\|_\infty = \left\| \frac{f+g}{2} \right\|_\infty$;
 (b) $\|f - g\|_\infty > 0$;
 (c) *there exist two homeomorphisms φ and ψ of $[0, 1]$ such that $g = f \circ \varphi$ and $\frac{f+g}{2} = f \circ \psi$.*

Proof. Let $0 \leq a < b \leq 1$ be such that f is strictly monotone on $[a, b]$. Without loss of generality, suppose that f is strictly increasing on $[a, b]$. Let $\xi : [a, b] \rightarrow [f(a), f(b)]$ be an increasing homeomorphism such that $\xi \neq f|_{[a, b]}$. Define $g \in C[0, 1]$ and an homeomorphism $\varphi : [0, 1] \rightarrow [0, 1]$ by

$$g(x) = \begin{cases} \xi(x) & \text{if } x \in [a, b]; \\ f(x) & \text{otherwise} \end{cases} \quad \varphi(x) = \begin{cases} f^{-1}(\xi(x)) & \text{if } x \in [a, b]; \\ x & \text{otherwise.} \end{cases}$$

Then, $g = f \circ \varphi$, $\|g\|_\infty = \|f\|_\infty = \left\| \frac{f+g}{2} \right\|_\infty$ and $\|f - g\|_\infty > 0$. Moreover, $f \circ \psi = \frac{f+g}{2}$, where $\psi : [0, 1] \rightarrow [0, 1]$ is the homeomorphism defined by

$$\psi(x) = \begin{cases} f^{-1}\left(\frac{\xi(x) + f(x)}{2}\right) & \text{if } x \in [a, b]; \\ x & \text{otherwise.} \end{cases}$$

□

Proposition 5.9. *Let $\|\cdot\|$ be an isometry invariant renorming of $C[0, 1]$. Then there exists a dense subset of $C[0, 1]$ where $\|\cdot\|$ is not strictly convex.*

Proof. Let $f \in C[0, 1]$ be a non-constant and affine function, and take g, φ and ψ as in Lemma 5.8. Since $f \mapsto f \circ \varphi$ and $f \mapsto f \circ \psi$ define surjective linear isometries of $C[0, 1]$,

$$\|g\| = \|f \circ \varphi\| = \|f\| = \|f \circ \psi\| = \left\| \frac{f+g}{2} \right\|.$$

So $\|\cdot\|$ is not strictly convex at the point f . The result then follows from the fact that the set of piecewise linear functions is dense in $C[0, 1]$. □

Combining Theorem 5.7 and Proposition 5.9, we obtain:

Corollary 5.10. *The space $C[0, 1]$ is light, but does not admit any isometry invariant LUR renorming.*

Remark 5.11. Using the same arguments as in the proofs of Proposition 5.6, Theorem 5.7 and Proposition 5.9, one can prove that $C_0(\mathbb{R})$ is light, but does not admit a strictly convex isometry invariant renorming either.

Remark 5.12. The examples presented in this section show that there is no general relation between closed subspaces and their respective isometry groups, in terms of being light, apart from Lemma 4.5. In fact:

- (1) c_0 is a closed subspace of ℓ_∞ , c_0 is light, but ℓ_∞ is not;
- (2) c is isometrically isomorphic to a closed subspace of $C[0, 1]$, c is not light but $C[0, 1]$ is light.

6. AN EXAMPLE OF A GROUP WITH A DISCRETE ORBIT BUT NO DISTINGUISHED POINT

In this section we solve a problem of [8], mentioned in the introduction, by exhibiting an SOT-discrete group of isomorphisms of c_0 which admits no distinguished point. More generally, we show the following:

Proposition 6.1. *For any integer $r \geq 2$, there exists a bounded infinite SOT-discrete group of isomorphisms of c_0 of the form $\text{Id} + F$, $F \in L(c_0)$ of finite rank, admitting a distinguished family of cardinality r , but none of cardinality $r - 1$.*

Proof. Since $c_0 \simeq \ell_1^r \oplus_\infty c_0$ it is enough to define the group G as an infinite bounded SOT-discrete group of isomorphisms on $\ell_1^r \oplus_\infty c_0$.

Let $(e_n)_{n \in \mathbb{N}}$ be the canonical basis of c_0 , and let $(U_n)_{n \in \mathbb{N}}$ be the sequence of isometries of c_0 defined by setting, for every $n, m \in \mathbb{N}$, $U_n(e_n) = -e_n$ and $U_n(e_m) = e_m$ whenever $m \neq n$. Let $(\phi_n)_{n \in \mathbb{N}}$ be dense in the unit sphere of ℓ_∞^r , and define the rank-one operator $R_n : \ell_1^r \rightarrow c_0$ by $R_n(x) = \phi_n(x)e_n$, $x \in \ell_1^r$. We then define an operator T_n on $\ell_1^r \oplus_\infty c_0$ in matrix form as

$$T_n = \begin{pmatrix} \text{Id} & 0 \\ R_n & U_n \end{pmatrix}.$$

It is readily checked that $T_n^2 = \text{Id}$ for every $n \in \mathbb{N}$ and that for every distinct integers $n_1, \dots, n_k$,

$$T_{n_1} \dots T_{n_k} = \begin{pmatrix} \text{Id} & 0 \\ R_{n_1} + \dots + R_{n_k} & U_{n_1} \dots U_{n_k} \end{pmatrix}.$$

Therefore the group G generated by the operators T_n is abelian. Furthermore, since for every $x \in \ell_1^r$

$$\|(R_{n_1} + \dots + R_{n_k})x\| = \|\phi_{n_1}(x)e_{n_1} + \dots + \phi_{n_k}(x)e_{n_k}\| \leq \max_i |\phi_{n_i}(x)| \cdot \|x\|$$

it follows that $\|T_{n_1} + \dots + T_{n_k}\| \leq 2$, and thus G is a bounded subgroup of $\text{GL}(\ell_1^r \oplus_\infty c_0)$.

We claim that no family $\{x_1, \dots, x_{r-1}\}$ of $\ell_1^r \oplus_\infty c_0$ is distinguished for G . Indeed, writing each vector x_i , $1 \leq i \leq r-1$, as (y_i, z_i) with $y_i \in \ell_1^r$ and $z_i \in c_0$, we note that $U_n z_i \rightarrow z_i$ for every $1 \leq i \leq r-1$. Since the vectors $y_1, \dots, y_{r-1}$ generate a subspace of dimension strictly less than r of ℓ_1^r , there exists a norm 1 functional $\phi \in \ell_\infty^r$ such that $\phi(y_i) = 0$ for every $1 \leq i \leq r-1$. Let $D \subset \mathbb{N}$ be such that $\phi_n \rightarrow \phi$ in ℓ_∞^r as n tends to infinity along D . Then $R_n(y_i) \rightarrow 0$ as n tends to infinity along D , and therefore $T_n(x_i) \rightarrow x_i$ as n tends to infinity along D for every $1 \leq i \leq r-1$. So the family $\{x_1, \dots, x_{r-1}\}$ is not distinguished for G .

On the other hand, if we denote by $(f_1, \dots, f_r)$ the canonical basis of ℓ_1^r , then the family $\{f_1 \oplus 0, \dots, f_r \oplus 0\}$ is distinguished for G . To check this, note that for any operator $T \in \text{GL}(\ell_1^r \oplus_\infty c_0)$ of the form

$$T = \begin{pmatrix} \text{Id} & 0 \\ \sum_{k \in F} R_k & U \end{pmatrix},$$

where F is a non-empty subset of $\mathbb{N}$, and U is an isometry of c_0 , we have

$$\|T(f_s \oplus 0) - f_s \oplus 0\| = \max_{k \in F} |\phi_k(f_s)| \quad \text{for every } 1 \leq s \leq r.$$

Since, for each $k \in F$, ϕ_k is normalized in ℓ_∞^r , $|\phi_k(f_s)| \geq 1$ for at least one index s . It follows that

$$\max_{1 \leq s \leq r} \|T(f_s \oplus 0) - f_s \oplus 0\| \geq 1,$$

and so

$$\inf_{\substack{T \in G \\ T \neq \text{Id}}} \left\{ \max_{1 \leq s \leq r} \|T(f_s \oplus 0) - f_s \oplus 0\| \right\} \geq 1.$$

Hence $\{f_1, \dots, f_r\}$ is a distinguished family for G . $\square$

We immediately deduce

Corollary 6.2. *The group of isomorphisms of c_0 constructed in the proof of Proposition 6.1 is not light.*

Proof. For every $x \in \ell_1^r$, the sequence $(R_n(x))_{n \in \mathbb{N}}$ tends weakly to 0 in c_0 . We also know that the sequence $(U_n)_{n \in \mathbb{N}}$ tends WOT to Id. Therefore $(T_n)_{n \in \mathbb{N}}$ also tends WOT to Id. On the other hand, we have for every $x \in \ell_1^r$ and every $n \in \mathbb{N}$ that

$$\|T_n(x \oplus 0) - x \oplus 0\| = \|R_n(x)\| = \|\phi_n(x)\|.$$

By the density of the sequence $(\phi_n)_{n \in \mathbb{N}}$ in the unit sphere of ℓ_∞^r , this implies that the sequence $(T_n(x \oplus 0))_{n \in \mathbb{N}}$ does not tend to x in norm, and thus $(T_n)_{n \in \mathbb{N}}$ does not tend SOT to Id. $\square$

We have thus proved:

Corollary 6.3. *There exists a bounded group G of isomorphisms of c_0 which is infinite, not light, SOT-discrete, and does not admit a distinguished point.*

Proof. Take $r = 2$ in Proposition 6.1. $\square$

7. QUASI-NORMED SPACES

Although Megrelishvili has defined the concept of light group of isomorphisms only for locally convex spaces, we can extend the definition to quasi-normed spaces, even if these spaces are not locally convex. One could ask if there is a general answer for the isometry groups of non-locally convex spaces, in terms of being light. The spaces ℓ_p and $L_p[0, 1]$, $0 < p < 1$, are examples that give a negative answer to this question.

Recall that for $0 < p < 1$, $(L_p[0, 1])^* = \{0\}$, i.e., the only linear continuous functional $f : L_p[0, 1] \rightarrow \mathbb{R}$ is the constant function $f \equiv 0$ (see [13, p. 18]). Considering the sequence $(T_n)_{n \in \mathbb{N}}$ constantly equal to $-\text{Id}$, we observe that $T_n \not\rightarrow^{\text{SOT}} \text{Id}$ while $T_n \xrightarrow{\text{WOT}} \text{Id}$. So $L_p[0, 1]$ is trivially non-light for every $0 < p < 1$. On the other hand,

Proposition 7.1. *For $0 < p < 1$, the space ℓ_p is light.*

Proof. Let $0 < p < 1$, and let $(T_\alpha)_{\alpha \in I}$ be a net in $\text{Isom}(\ell_p)$ such that $T_\alpha \xrightarrow{\text{WOT}} \text{Id}$. Each T_α acts on vectors $(x_n)_{n \in \mathbb{N}} \in \ell_p$ as $T_\alpha((x_n)_{n \in \mathbb{N}}) = (\varepsilon_n^{(\alpha)} x_{\sigma_\alpha(n)})_{n \in \mathbb{N}}$, where σ_α is a permutation of $\mathbb{N}$ and $(\varepsilon_n^{(\alpha)})_{n \in \mathbb{N}}$ is a sequence of elements of $\{-1, 1\}$ (the proof of this fact is similar of the case where $p > 1$ and $p \neq 2$, found in [1, p. 178]). Assume, by contradiction, that $T_\alpha \not\rightarrow^{\text{SOT}} \text{Id}$. Then there exist $x \in \ell_p$, $\varepsilon > 0$ and an infinite sequence $(\alpha_i)_{i \in \mathbb{N}}$ of indices in I such that $\|T_{\alpha_i} x - x\|_p^p > \varepsilon$ for every $i \in \mathbb{N}$. Since $x \in \ell_p$, there exists $N \in \mathbb{N}$ such that $\sum_{k=N+1}^\infty |x_k|^p < \varepsilon/2$. The dual space of ℓ_p identifies isomorphically with ℓ_∞ , where $\Phi = (y_k)_{k \in \mathbb{N}}$ acts on an element $x = (x_k)_{k \in \mathbb{N}} \in \ell_p$ by the formula $\Phi(x) = \sum_{k \in \mathbb{N}} y_k x_k$ (see [13, p. 21]). Considering for $1 \leq j \leq N$ the functionals Φ_j identified with the vectors of

the canonical basis $e_j \in \ell_\infty$, as well as the vectors $e_k \in \ell_p$ for $1 \leq k \leq N$, we obtain by the WOT convergence of T_α to Id that

$$\Phi_j(T_\alpha(e_k)) - \Phi_j(e_k) = \varepsilon_j^{(\alpha)} \delta_{\sigma_\alpha(k),j} - \delta_{k,j} \rightarrow 0,$$

where $\delta_{k,j} = 1$ if $k = j$ and $\delta_{k,j} = 0$ if $k \neq j$. In particular, $\varepsilon_k^{(\alpha)} \delta_{\sigma_\alpha(k),k} \rightarrow 1$ for every $1 \leq k \leq N$. So we may assume that the permutations σ_α fix the first N integers and that $\varepsilon_k^{(\alpha)} = 1$ for every $1 \leq k \leq N$. Hence we have for every $i \in \mathbb{N}$

$$\sum_{k=N+1}^{\infty} |(T_{\alpha_i}(x))_k|^p = \sum_{k=N+1}^{\infty} |x_k|^p < \frac{\varepsilon}{2}.$$

However, taking $z = (z_k)_{k \in \mathbb{N}} \in \ell_p$ defined by $z_k = 0$ if $1 \leq k \leq N$ and $z_k = 1$ if $k > N$, we have $\|z\|_p^p = \sum_{k=N+1}^{\infty} |x_k|^p < \varepsilon/2$ and

$$\|T_{\alpha_i}x - x\|_p^p = \|T_{\alpha_i}z - z\|_p^p \leq \|T_{\alpha_i}z\|_p^p + \|z\|_p^p = \varepsilon$$

for every $i \in \mathbb{N}$, which is a contradiction. $\square$

We finish the paper with a few related questions or comments.

8. QUESTIONS AND COMMENTS

Our first question concerns renormings of the space c . Since it c is not light, as proved in Proposition 4.3, it does not admit any isometry invariant LUR renorming. But it may still admit an isometry invariant strictly convex renorming.

Question 8.1. *Does c admits an isometry invariant strictly convex renorming?*

We have observed in Section 4 that if the isometry group $\text{Isom}(X)$ of a Banach space X of dimension > 1 acts almost transitively on S_X and is SOT-discrete, it is not abelian.

Question 8.2. *Suppose X is a separable Banach space of dimension > 1 and $G \leq \text{Isom}(X)$ is an SOT-discrete amenable subgroup. Can G have a dense orbit on S_X ?*

We have seen in Corollary 6.3 that there exists a bounded group G of isomorphisms of c_0 which is infinite, not light, SOT-discrete, and does not admit a distinguished point. One may wonder about the role of the space c_0 in this construction. For example, one can ask:

Question 8.3. *Does there exist a reflexive space X with an SOT-discrete bounded group $G \leq \text{GL}(X)$ that does not admit a distinguished point?*

Of course such a group G , if it exists, must be light, as all reflexive spaces are light. Noting that the example of Proposition 4.3 is a group of finite rank perturbations of the identity on the space c_0 , a question in the same vein is:

Question 8.4. *Does there exist a reflexive space X with an SOT-discrete infinite bounded group $G \leq \text{GL}(X)$ such that all elements of G are finite rank perturbations of the identity?*

This question is relevant to [7], where isometry groups on complex, reflexive, separable, hereditarily indecomposable spaces are studied. A negative answer to this question would imply that all isometry groups on such spaces act almost trivially, i.e., there would exist an isometry invariant decomposition $F \oplus H$ of the space where F is finite dimensional and all elements of the group act as multiple of the identity on H , [7] Theorem 6.9.

Another natural space which could be investigated in this context is the universal space of Gurarij, whose isometry group possesses a very rich structure (see [11] for its definition and [10] for a recent survey).

Question 8.5. *Is the isometry group of the Gurarij space light?*

Finally it remains open whether the converse to Megreleshvili's result holds:

Question 8.6. *Does a Banach space X have the PCP if and only if all bounded subgroups of $GL(X)$ are light?*

The answer is positive when X has an unconditional basis: this follows from Theorem 4.6, the fact that an unconditional basis whose span does not contain c_0 must be boundedly complete, and the fact that separable dual spaces have the RNP and therefore the PCP.

REFERENCES

- [1] S. Banach, *Théorie des opérations linéaires* [Theory of linear operators], Monografie Matematyczne, Tom I., Seminar. Matem. Univ. Warszawa, Warszawa (1932).
- [2] G. Birkhoff, *The topology of transformation-sets*, Ann. of Math. 35 (1934), 861875.
- [3] J. M. F. Castillo, M. González, *Three-space problems in Banach space theory*, Lecture Notes in Mathematics, 1667, Springer-Verlag, Berlin (1997).
- [4] R. Deville, G. Godefroy, V. Zizler, *Smoothness and renormings in Banach spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 64, Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York (1993).
- [5] J. Diestel, *Geometry of Banach spaces: selected topics*, Lecture Notes in Mathematics, Vol. 485. Springer-Verlag, Berlin (1975).
- [6] V. Ferenczi, E. M. Galego, *Countable groups of isometries on Banach spaces*, Trans. Amer. Math. Soc. 362 (2010), no. 8, 4385–4431.
- [7] V. Ferenczi, C. Rosendal, *On isometry groups and maximal symmetry*, Duke Math. J. 162 (2013), no. 10, 1771–1831.
- [8] V. Ferenczi, C. Rosendal, *Displaying Polish groups on separable Banach spaces*, Extracta Math. 26 (2011) 195–233.
- [9] R. J. Fleming, J. E. Jamison, *Isometries on Banach spaces: function spaces*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, 129, Chapman & Hall/CRC, Boca Raton, FL (2003).
- [10] J. Garbulińska, W. Kubiś, *Remarks on Gurarij spaces*, Extracta Math. 26(2) (2011/2012), 235–269.
- [11] V. I. Gurarij, *Spaces of universal placement, isotropic spaces and a problem of Mazur on rotations of Banach spaces*, Sibirsk. Mat. Z. 7 (1966), 1002–1013.
- [12] D. Helmer, *Joint continuity of affine semigroup actions*, Semigroup Forum 21 (1980), no. 2-3, 153–165.
- [13] N. J. Kalton, N. T. Peck, J. W. Roberts, *An F -space sampler*, London Mathematical Society Lecture Note Series, 89, Cambridge University Press, Cambridge (1984).
- [14] G. Lancien, *Dentability indices and locally uniformly convex renormings*, Rocky Mountain J. Math. 23 (1993), no. 2, 635–647.
- [15] M. Megreleshvili, *Operator topologies and reflexive representability*, Nuclear groups and Lie groups (Madrid, 1999), 197–208, Res. Exp. Math., 24, Heldermann, Lemgo (2001).

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE TECNOLÓGICA FEDERAL DO PARANÁ, CAMPUS TOLEDO, 85902-490 TOLEDO, PR, BRAZIL

E-mail address: `leandroantunes@utfpr.edu.br`

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, 05311-970 SÃO PAULO, SP, BRAZIL

E-mail address: `ferenczi@ime.usp.br`

CNRS, LABORATOIRE PAUL PAINLEVÉ, UMR 8524, UNIVERSITÉ DE LILLE, CITÉ SCIENTIFIQUE, BÂTIMENT M2, 59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

E-mail address: `grivaux@math.univ-lille1.fr`

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE (M/C 249), UNIVERSITY OF ILLINOIS AT CHICAGO, 851 S. MORGAN ST., CHICAGO, IL 60607-7045, USA

E-mail address: `rosendal.math@gmail.com`