

Terrestrial and marine floral response to latest Eocene and Oligocene events on the Antarctic Peninsula

Sophie Warny, C. Madison Kymes, Rosemary Askin, Krzysztof P. Krajewski & Andrzej Tatur

To cite this article: Sophie Warny, C. Madison Kymes, Rosemary Askin, Krzysztof P. Krajewski & Andrzej Tatur (2019) Terrestrial and marine floral response to latest Eocene and Oligocene events on the Antarctic Peninsula, *Palynology*, 43:1, 4-21, DOI: [10.1080/01916122.2017.1418444](https://doi.org/10.1080/01916122.2017.1418444)

To link to this article: <https://doi.org/10.1080/01916122.2017.1418444>



Published online: 16 Feb 2018.



Submit your article to this journal [↗](#)



Article views: 181



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)



Terrestrial and marine floral response to latest Eocene and Oligocene events on the Antarctic Peninsula

Sophie Warny^{a,b}, C. Madison Kymes^a, Rosemary Askin^c, Krzysztof P. Krajewski^{d,†} and Andrzej Tatur^e

^aLSU Department of Geology and Geophysics E-235 Howe-Russell, Baton Rouge, Louisiana 70803; ^bLSU Museum of Natural Science, 109 Foster Hall, Baton Rouge, Louisiana 70803; ^cConsultant, Jackson, Wyoming, United States; ^dInstitute of Geological Sciences, Polish Academy of Sciences, Twarda 51/55, 00-818 Warszawa, Poland; ^eFaculty of Geology, University of Warsaw, Żwirki i Wigury 93, 02-089 Warszawa, Poland

ABSTRACT

Palynological results from opposite sides of the northernmost Antarctic Peninsula provide insight on terrestrial vegetation and sea-surface conditions immediately before the Eocene–Oligocene transition (EOT), through Early Oligocene glacial conditions and the subsequent Late Oligocene interglacial interval. A latest Eocene sample set from the uppermost La Meseta Formation on Seymour Island, James Ross (back-arc) Basin, records a low-diversity *Nothofagus* (southern beech)-dominated vegetation with some podocarp conifers similar to Valdivian-type forest found today in Chile and Argentina. Marine organic-walled phytoplankton include leiospheres and Eocene dinoflagellate cysts such as *Vozzhennikovia rotunda*, *V. apertura*, *Senegalinium asymmetricum* and *Spinidinium macmurdoense*. Immediately before the EOT near the top of the section the decrease in terrestrial palynomorphs, increase in reworked specimens, disappearance of key dinocysts, and overwhelming numbers of sea-ice-indicative leiospheres plus the small dinoflagellate cyst *Impletosphaeridium* signal the onset of glacial conditions in a subpolar climate. Early to Late Oligocene samples from the Polonez Cove and Boy Point formations on King George Island, South Shetland Islands (magmatic arc), yielded an extremely depauperate terrestrial flora, likely resulting in part from poor vegetation cover during the Polonez Glaciation but also because of destruction of vegetation due to continued regional volcanism. The prevalence of sea-ice-indicative leiospheres in the marine palynomorph component is consistent with polar to subpolar conditions during and following the Polonez Glaciation.

KEYWORDS

Antarctica; Seymour Island; King George Island; palynology; Eocene; Oligocene

1. Introduction

The interval from the early Eocene through to the Eocene–Oligocene Transition (EOT, ca. 34 Ma) represents a global cooling trend marked by a high-latitude sea-surface temperature decrease from $\sim 18^\circ\text{C}$ in early Eocene to $\sim 6^\circ\text{C}$ transitioning into the Oligocene (Stott et al. 1990; Liu et al. 2009). The $\sim 12^\circ\text{C}$ temperature change has been correlated with increasing deep-sea benthic foraminiferal oxygen isotope ($\delta^{18}\text{O}$) values, and decreasing atmospheric carbon dioxide concentrations (Zachos et al. 1996, 2001, 2008; Prothero & Berggren 2014). This major cooling trend was accompanied by substantial ice growth in Antarctica. Ice growth continued with some fluctuations as temperatures decreased from ‘greenhouse’ conditions approaching the EOT, to the Oligocene–present day ‘icehouse’ or glacial state in Antarctica (Prothero et al. 2003; Lear et al. 2008).

This paper reports palynological results from a master’s of science (MS) project by Madison Kymes (2015), originally based on two sample sets provided by Krzysztof P. Krajewski and Andrzej Tatur from King George Island in the South Shetland Islands (Figure 1). These included the current Lower to Upper Oligocene set from the Polonez Cove and the Boy Point formations, as well as a Lower Miocene set from the Cape Melville

Formation, reported earlier in Warny et al. (2016). For comparison, an additional sample set was selected from the uppermost Eocene sediments on Seymour Island in the Weddell Sea (Figure 1). Together these studies cover three small but significant slices of time during the Cenozoic shift to icehouse conditions in West Antarctica. They are compared with earlier results in the region from well-dated SHALDRIL cores (Anderson et al. 2011; Warny & Askin 2011a, 2011b; Griener et al. 2013; Feakins et al. 2014). This MS project is a small part within the broader objectives of the authors and their colleagues of clarifying Antarctica’s Cretaceous and Cenozoic vegetation and climate history.

2. Geological setting

2.1. La Meseta Formation, Seymour Island

The older (Eocene) sample set discussed here is from Seymour Island, which lies within the James Ross (back-arc) Basin, south-east of the tip of the Antarctic Peninsula (Figure 1(B)) at $64^\circ 17'\text{S}$ latitude and $56^\circ 45'\text{W}$ longitude. The island is roughly 20.5 km long and 9.6 km wide, showing slopes from sea level up to a pronounced plateau or mesa at roughly 200 m above sea level (Elliot et al. 1975). Seymour Island outcrops lack permanent ice




Figure 1. (A) Map of Antarctica showing the location of the Antarctic Peninsula; the red rectangle is the part enlarged in (B). (B) Locations of Seymour Island, King George Island, and SHALDRIL II sites 3C and 12A in the northern Antarctic Peninsula region.

and are thus well exposed, and contain a rich assortment of well-preserved marine and terrestrial macro- and microfossils.

The selected samples are from the uppermost sediments of the La Meseta Formation (Rinaldi et al. 1978; Elliot & Trautman 1982) (Figure 2). La Meseta Formation siliciclastic strata crop out on the northern third of the island, filling a valley incised into the upper Palaeocene Cross Valley Formation, itself incised into Maastrichtian to lower Palaeocene sediments of the López de Bertodano and Sobral formations, which are exposed in the south-western two-thirds, and also the northern tip of the island. Following the initial three-fold subdivision of the formation by Elliot & Trautman (1982), various authors have described and subdivided this complex back-arc shelfal succession of

shallow marine, estuarine and deltaic sands, silts, muddy sediments and shellbeds. Seven lithofacies (designated Tertiary Eocene La Meseta/Telm 1–7) were mapped in detail by Sadler (1988). Porębski (1995, 2000) recognised three major eustatically controlled depositional sequences, with subdivisions that are more tectonically influenced. Marenssi & Santillana (1994) and Marenssi et al. (1998a, 1998b) divided the formation into six primarily eustatically controlled depositional sequences or allomembers. The stratigraphic scheme of Marenssi and colleagues is followed here.

The Seymour Island sample set was collected from the top of the La Meseta Formation (Submeseta Allomember according to Marenssi et al. 1998a, 1998b) along the side of a gully




Figure 2. Simplified geological map of northern Seymour Island, showing the La Meseta Formation with the distribution of the Submeseta Allomember in relationship to other La Meseta allomembers. The location of the D6 section is marked by a red line. The relationship of the sample location with the lithology sampled is presented in the inset to the right.

downcutting the central/south-eastern side of the mesa (Figures 2 and 3), from monotonous unconsolidated sands and silts with shell beds and concretionary horizons. Strontium-isotope analyses conducted on *Cucullaea* specimens collected throughout the La Meseta Formation by Dutton et al. (2002 and references therein; Ivany et al. 2008) had supported the previous palaeontologically assigned age range of late Early Eocene to Late Eocene; however, more recent results suggest the oldest La Meseta sediments may be Mid rather than late Early Eocene (Douglas et al. 2014), with the entire formation ranging in age from 45 to 34 Ma. Specifically, for the uppermost beds sampled in this study (upper Submeseta Allomember, or upper Telm7), Dutton et al. (2002), Ivany et al. (2008) and Douglas et al. (2014) place these sediments between 36 and 34 Ma. These upper beds underlie the Miocene glaciogenic Hobbs Glacier Formation (Marensi et al. 2010) and the Pliocene–Pleistocene till of the Weddell Sea Formation (Zinsmeister and deVries 1983; Gaździcki et al. 2004) that tops the mesa. The discovery, off the opposite northern side of the mesa, by Ivany et al. (2006) of three thin layers of sediments (pebbly mudstone; diamict; pebbly mudstone) between the uppermost La Meseta Formation and the overlying Weddell Sea Formation provides more certain age and palaeoclimatic control. Ivany et al. (2006) report $^{87}\text{Sr}/^{86}\text{Sr}$ ratios of bivalves yielding ages of 33.57–34.78 Ma from the uppermost La Meseta Formation at their locality, confirming an EOT age at the top of the formation (E–O boundary 33.9 Ma, Gradstein et al. 2012). Dinoflagellate cysts indicate a late Eocene and earliest Oligocene age, respectively, for the lower and upper pebbly mudstone (Ivany et al. 2006), with the intervening diamict suggesting glacial ice at the EOT. Our sample set from a section on the opposite side of the mesa is likely of similar latest Eocene age to the uppermost La Meseta beds dated by Ivany et al. (2006; see also Dutton et al. 2002; Ivany et al. 2008; Douglas et al. 2014).

2.2. Polonez Cove Formation and Boy Point Formation, King George Island

The younger (Oligocene) sample set discussed here is from King George Island, located approximately 120 km off the northern coast of the Antarctic Peninsula at 62°01'S latitude and 58°33'W longitude (Figure 1(B)). King George Island is roughly 95 km long and 25 km wide, making it the largest of the South Shetland Islands. The South Shetland Islands are part of a magmatic arc, detached from the northern Antarctic Peninsula magmatic arc by the Bransfield Strait (Figure 1(B)). The Bransfield Rift is less than 4 million years old, with oceanic crust dating from about 1.3 Ma (Barker & Austin 1998). Thus, the studied sedimentary units representing the back-arc basin (Seymour Island) and the magmatic arc (King George Island) were deposited in relatively close proximity to each other on opposite sides of the Antarctic Peninsula in the latest Eocene and Oligocene.

The geological history of King George Island is substantially more complicated than that of the Seymour Island sediments. The Eocene through Oligocene of King George Island reflects a complex history of volcanism, glaciation and marine transgressions. The studied sample set is from part of the Chopin Ridge Group (Birkenmajer 1980) exposed on the western side of King George Bay (Figure 4). The Chopin Ridge Group includes porphyritic lava flows and pyroclastics, marine tillites, associated glaciogenic sediments and sandstones, divided into five formations (in ascending order): the Mazurek Point Formation or Lions Cove Formation, Polonez Cove Formation, Boy Point Formation and Wesele Cove Formation (Birkenmajer 2001). The sample set was collected from the Polonez Cove Formation and basal Boy Point Formation along a section at Linton Knoll (Figures 4 and 5(A)). Supplementary samples from the Polonez Cove Formation were collected at Conglomerate Bluff, located approximately 2 km to the south-east from Linton Knoll




Figure 3. Photograph looking to the south-east on Seymour Island of section D6 towards a person (black arrow) standing on the resistant bed near the section base (between samples 1 and 2). The red arrow indicates the direction of the D6 section.



Figure 4. Geological map of King George Island, showing the distribution of the Chopin Ridge Group and the location of Linton Knoll and Conglomerate Bluff sections.

(Figures 4 and 5(B)). Their stratigraphic position is compiled on the Linton Knoll section (Figure 6).

The Mazurek Point Formation and the Lions Cove Formation (Eocene) comprise basaltic/andesitic substratum for the overlying glacio-marine and volcanogenic formations along the outcrop belt of the Chopin Ridge Group (Birkenmajer 2001; Pańczyk & Nawrocki 2011). The overlying Polonez Cove Formation is subdivided into six members, though not all are present at every location (Birkenmajer 1980, 1982; Porębski & Gradziński 1987; Troedson & Smellie 2002). These are (in ascending order): the Krakowiak Glacier, Bayview, Low Head, Siklawa, Oberek and Chlamys Ledge members. Lodgement till at the base of the Polonez Cove Formation and the associated glacio-marine sediments, together making up the Krakowiak Glacier Member, represent the Polonez Glaciation (Birkenmajer 1987). The name Krakowiak Glacier Member was based on the presence of a small glacial body that existed along the Chopin Ridge, and it was introduced by K. Birkenmajer in 1980. It is sad to note that this ice body has now fully vanished as a result of the warming trend in West Antarctica (K. Krajewski, pers. comm.). A late Early Oligocene age is indicated from Sr dating of carbonate material from the Krakowiak Glacier Member and the Low Head Member (ca. 30–28 Ma, Dingle et al. 1997; Dingle & Lavelle 1998). Unpublished strontium isotope results point to even older geological ages (Early Oligocene, ca. 32–30 Ma) of parts of the glacio-marine deposits of the Krakowiak Glacier Member (K. Krajewski, pers. comm.). Younger Ar–Ar dates were derived from the inter-fingering and overlying basaltic lavas and hyaloclastic rocks in the Low Head Member and Oberek Cliff Member, although precision of these measurements is low (29–22 Ma, Smellie et al. 1998; Troedson & Smellie 2002). The upper part of the Polonez Cove Formation, which comprises shallow marine basaltic

sandstones (Chlamys Ledge Member), gave a single late Oligocene strontium age (27.1 ± 0.3 Ma) on the basis of a *Chlamys* shell analysis (K. Krajewski, pers. comm.). The whole-rock K–Ar ages obtained from the overlying terrestrial Boy Point Formation (ca. 24–22 Ma) indicate an age of youngest Oligocene/oldest Miocene, though they might in part reflect a reset by younger thermal events (Birkenmajer & Gaździcki 1986). The Boy Point Formation consists mostly of dacitic rocks, with the lower informal member (Loud Waterfall member) dominated by dacitic clastic deposits, and the upper informal member (Linton Knoll member) dominated by agglomerates and lavas. A Late Oligocene age for the formation is suggested here on the basis of geological correlation.

3. Studied materials

Thirty samples were analysed for palynology: 14 from Seymour Island (Table 1), and 16 from King George Island (Table 2). The 14 Seymour Island samples were collected from the uppermost La Meseta Formation for R. Askin in December 1986 and were obtained for this study from the Polar Rock Repository, Byrd Polar and Climate Research Center, The Ohio State University. These 14 samples are from the top of the Submeseta Allomember (or upper Telm7 in the terms of Sadler 1988) and were collected at 3-m intervals through a 39-m section of monotonous fine sands and silts (Figure 2).

The 16 samples from King George Island included sandstones to mudstones and were collected from the Linton Knoll (11 samples) and Conglomerate Bluff (five samples) sections (Figures 4–6). Fourteen of these samples were taken from some members of the Polonez Cove Formation, and two samples from the base of the Boy Point Formation. They were collected




Figure 5. (A) Photograph of the section at Linton Knoll. (B) Photograph of the section at Conglomerate Bluff. King George Bay, King George Island. Red arrows indicate directions of the sections. KGM – Krakowiak Glacier Member; LHM – Low Head Member; CLM – Chlamys Ledge Member.

during the Polish Academy of Sciences (PAS) two-part expedition, which took place in January 2007 and January 2009.

4. Methods

All samples were processed using standard chemical palynological processing techniques. Dry sediment was weighed and spiked with a known quantity of *Lycopodium* spores to allow for calculation of palynomorph concentrations. Dry sediment was successively treated with hydrochloric acid, hydrofluoric acid and heavy liquid separation (e.g. Brown 2008). Samples were sieved between a 10- and 250- μm fraction, and the remaining residue was mounted on microscope slides using glycerin jelly.

Palynological analysis was conducted in the Louisiana State University's Center for Excellence in Palynology (CENEX) lab. When possible, 300 palynomorphs were tabulated per sample, using an Olympus BX41 microscope. Palynomorphs were identified to the lowest taxonomic level possible. After palynomorphs were tallied, palynomorph concentration was calculated for

each specimen using the equation from Benninghoff (1962): $C = (P_c \times L_t \times T) / (L_c \times W)$, where C = concentration (per gramme of dried sediment, gdw^{-1}), P_c = the number of palynomorphs counted, L_t = the number of *Lycopodium* spores per tablet, T = the total number of *Lycopodium* tablets added per sample, L_c = the number of *Lycopodium* spores counted and W = the weight of dried sediment.

All specimens analysed from the assemblages were identified at the time of scanning and counting as being either reworked or *in situ* (considered penecontemporaneous with deposition), before palaeoenvironmental conditions could be considered. High numbers of reworked specimens are commonplace in Antarctic Cenozoic assemblages, in large part because of glacial and/or fluvial scouring and redeposition of mainly unconsolidated sediments. When age-restricted species older than Eocene or Oligocene were found (e.g. Cretaceous species), they could be easily recognised as reworked. However, for species with longer age ranges, it is often difficult to differentiate between reworked and *in situ* in Antarctic successions,




Figure 6. Detailed section of the Polonez Cove Formation at Linton Knoll showing samples collected during the PAS expeditions in 2007 and 2009. * indicates stratigraphic location of samples from the adjacent Conglomerate Bluff section. Polonez Cove Formation: KGM – Krakowiak Glacier Member; BM – Bayview Member; LHM – Low Head Member; SM – Siklawa Member; OCM – Oberek Member; CLM – Chlamys Ledge Member. Boy Point (BP) Formation: LWM – Loud Waterfall member; LKM – Linton Knoll member.

because of minimal time and depths of burial of eroded strata and subsequent lack of differences in preservation and thermal maturities between reworked and *in situ* grains. Where possible, differentiation of reworked specimens was done on the basis of subtle to more overt differences in the preservation of the

grains (e.g. battered specimens, presence of corrosion) as well as the thermal maturity of their walls. Some thick-walled cryptogam spores were counted as reworked. Some specimens, if differentiation was not possible, were counted as *in situ*, resulting in taxa such as *Nothofagidites* spp. (*brassii* group) being

Table 1. Palynomorph concentration for the uppermost Eocene D6 section of the La Meseta Formation.

Sample name	Elevation (m)	Palynological counts	Concentration of palynomorphs (gdw ⁻¹)
D6-14	39	300	753
D6-13	36	300	634
D6-12	33	300	759
D6-11	30	300	812
D6-10	27	300	652
D6-09	26	300	557
D6-08	21	300	669
D6-07	18	300	760
D6-06	15	300	873
D6-05	12	300	844
D6-04	9	300	776
D6-03	6	300	987
D6-02	3	86	373
D6-01	0	98	119

included as part of the penecontemporaneous flora, though these 'warmer-climate' species are typically considered as becoming scarce and possibly disappearing from the region in the latest Eocene (e.g. Chen 2000; R. Askin, pers. comm.).

5. Results

5.1. Palynological results from the La Meseta Formation

The 14 La Meseta samples yielded well-preserved palynomorphs, but with a fairly low diversity. The overall numbers of palynomorphs counted are shown in Figure 7, and our interpretations of *in situ* vs. reworked species in Figure 8. Table 3 provides raw counts. Note that specimens considered to be *in situ* (interpreted as derived from the penecontemporaneous terrestrial and marine flora) are differentiated in blue, and reworked specimens in red, throughout the figures and tables. Selected *in situ* specimens are illustrated in Plate 1.

Palynomorph concentrations are low in these sandy sediments and range from 119 to 987 per gramme of dried sediments (Table 1), which is far less than the concentrations (varying from 700 to 100,000) of recovered palynomorphs from the late Eocene 3C section (Figure 1(B)) sampled by the

SHALDRIL programme offshore and north-east of Seymour Island (Warny & Askin 2011a), and from many older samples of the La Meseta Formation. Both of the latter sample sets generally have a higher mud content and thus tend to be more productive. Interestingly, the preparations recovered in this study contain higher concentrations of palynomorphs than most of the same uppermost La Meseta samples summarised by Askin (1997). Several factors may explain this mismatch in palynomorph concentration between the two studies: a larger amount of sample was processed in the current study, recent improvements in processing techniques may have resulted in more abundant organic matter being recovered, or the processed sample fraction may have contained a slightly higher proportion of finer (muddy) sediment.

Age-indicative dinoflagellate cysts occur throughout most of the section, from D6-01 to D6-12 (Figure 7; Table 3). They are absent in the uppermost part (samples D6-13, D6-14). The presence of *Vozzhennikovia rotunda* (Lentin & Williams 1981), *V. apertura* (Lentin & Williams 1981), *Senegalinium asymmetricum* (Stover & Evitt 1978) and *Spinidinium macmurdoense* (Wilson 1967) in the La Meseta samples is consistent with Eocene age, based on age ranges from Ocean Drilling Program's (ODP) Publication 189 (Sluijs et al. 2003; see also the Supporting Information discussion in Douglas et al. 2014 with their reference to the recent biostratigraphic age model of Bijl et al. 2013). It is noted that even though *Vozzhennikovia* spp. and *S. macmurdoense* are shown by Bijl et al. (2013) to range into the Early Oligocene, other dating methods restrict the La Meseta Formation to the Eocene (see Geological setting, Section 2).

Reworked palynomorphs occur throughout the section, becoming progressively more common above the basal samples, and after some fluctuations again increasing in the uppermost portion of the section in samples D6-13 and especially D6-14. Reworked species of palynomorphs range from Permian to Palaeocene in age and have been well documented on Seymour Island (e.g. Askin & Elliot 1982; Bowman et al. 2012). Identified reworked taxa are mostly Cretaceous to Paleogene in age.

Presumed penecontemporaneous or '*in situ*' specimens predominate throughout section D6, with terrestrial pollen of *Nothofagidites* spp. the most abundant in the lower samples (D6-01 to D6-07) and marine leiospheres dominating the upper samples (D6-08 to D6-14), illustrating a trend to a more marine assemblage upsection. Some other observations within these broad trends are summarised below.

D6-01 to D6-07. This lower part of the section contained the most varied (though hardly diverse) terrestrial palynoflora. The predominant *Nothofagidites* (southern beech) pollen considered *in situ* are mainly *N. spp.* (*fusca* group), with *Nothofagidites* species of the *brassii* and *menziesii* groups. Other angiosperm pollen include *Myricipites harrisii* and *M. parvus*, *Peninsulapollis* spp. and *Proteacidites* spp. Some podocarpaceous conifer pollen, *Phyllocladites* spp. and *Podocarpidites* spp. (Table 3), occur in these lower samples. Other conifer pollen were observed throughout the section, but they were interpreted at the time of scanning as reworked specimens (see Section 4, Methods).

Table 2. Palynomorph concentrations of reworked (Rw) and *in situ* palynomorphs from Polonez Cove and Boy Point formation samples.

Sample name	Elevation (m)	Palynological counts	Concentration of palynomorphs (gdw ⁻¹) (Rw)	Concentration of palynomorphs (gdw ⁻¹) (<i>in situ</i>)
L-12	55	11	41	9
L-13	48	7	5	4
G-13	46	11	1	1
L-11	43	46	9	20
LHMB	26	101	4	20
LHMB-cb	26*	40	7	10
G8	25	37	73	44
LRx	25	46	8	21
L-6	21	73	11	18
L-5	14	54	4	10
K-1	14	64	15	21
KGLMB-4	7*	95	29	33
KGLMB-3	7*	45	4	18
KGLMB-2	7*	10	7	1
KGLMB-1	7*	0	0	0
G3	5	40	44	117

* Samples from Conglomerate Bluff, with their stratigraphic position shown on the Linton Knoll section.

Seymour La Meseta




Figure 7. Overall frequency (raw counts) of palynomorph species observed in the La Meseta Formation of Seymour Island; blue = *in situ* and red = reworked.




Figure 8. Relative abundance of reworked (Rw) and *in situ* palynomorphs in La Meseta Formation samples.

In the marine dinoflagellate cyst component, taxa typical of the Antarctic Eocene are most common in this part of the section, and a small peak in abundance in *Impletosphaeridium* spp. occurs in sample D6-05.

D6-08 to D6-12. The marine component, which dominates in this part of the section, is marked by increasing

numbers of leiospheres, included as the sphaeromorph acritarch *Leiosphaeridia* spp. Dinoflagellate cysts show slight decreases in numbers compared with the underlying part of the section. Diversity has dropped off drastically in the terrestrial assemblage. Apart from the last conifer pollen considered *in situ* in this study in sample D6-08, only pollen of *Nothofagidites* (mainly *fusca* group)

SAMPLE #	DEPTH IN M	Dinoflagellate cysts				Algae	Spores				Gymnosperms	Angiosperms				Reworking								
		<i>Asterionum distichum</i> <i>Dicodanidium</i> sp. <i>Impatiopsisphaeridium</i> spp. <i>Manumiella seelandica</i> <i>Paleocystodinium</i> sp. <i>Phelodinium</i> sp. <i>Senegalinium asymmetricum</i> <i>Spinidinium lanterna</i> <i>Spinidinium macmurdoense</i> <i>Spiniferites</i> spp. <i>Vozzhennikovia apertura</i> <i>Vozzhennikovia rotunda</i>	<i>Leiosphaeridia</i> spp. <i>Pterospirillum australe</i> <i>Acantholittorites teretangulatus</i> <i>Clavifera triplex</i> <i>Cycladites minor</i> <i>Ditythophyllites</i> spp. <i>Foraminiparis dailyi</i> <i>Laevigatosporites ovatus</i> <i>Lentolittorites directus</i> <i>Osmundacites wellmannii</i> <i>Punctatisporites</i> sp. <i>Stereosporites antiquasporites</i> <i>Trilobosporites</i> spp.	<i>Microacantholittorites antarcticus</i> <i>Phyllocladites</i> spp. <i>Phyllocladites</i> spp. <i>Podocarpites</i> spp. <i>Podocarpites</i> spp. <i>Myricipites harrisi</i> <i>Myricipites parvus</i> <i>Nothofagidites</i> spp. (<i>brassii</i> gp.) <i>Nothofagidites</i> spp. (<i>brassii</i> gp.) <i>Nothofagidites</i> spp. (<i>jusca</i> gp.) <i>Nothofagidites</i> spp. (<i>jusca</i> gp.) <i>Nothofagidites</i> spp. (<i>menziesii</i> gp.) <i>Nothofagidites</i> spp. (<i>menziesii</i> gp.) <i>Peninsulapollis</i> spp. <i>Peninsulapollis</i> spp. <i>Proteacidites</i> spp. <i>Proteacidites</i> spp. <i>Tricolpites</i> spp. <i>Triparapollenites</i> spp.	<i>Reworked Counts Marine</i> <i>In Situ Counts Marine non sea ice</i> <i>In Situ Marine sea-ice indicator</i> <i>Reworked Counts Terrestrial</i> <i>In Situ Counts Terrestrial</i>																			
D6-14	39	19	2			111	3				10	17		38	88	7		5	2	0	130	168	0	
D6-13	36	28	1	1		179	4			2	22	13		9	32		8		3	0	207	90	0	
D6-12	33	19				172	2				4	8			50	25	3	5	0	2	191	54	53	
D6-11	30	1	9		7	2	4		4		7	13		8	14	70	20	1	3	4	4	7	6	
D6-10	27	2			2	2			4	193	9	7		0	19	21	30		1	2	3	4	2	
D6-9	24	7	4		7	1		1	3	106	6	12		0	19	75	37	1		3	4	2	3	
D6-8	21	1	2	5	8	1	1		7	102	1	2	1	1	11	22	76	29		5	5	1	5	
D6-7	18	4	1		7	3			1	65	1	4	2	2	18	29	87	44	6	3	4	5	0	
D6-6	15	2	3		6	2			5	66	3	1			3	1	100	27	8	4	5	6	3	
D6-5	12	2	21	2	7	6			10	51	5	3			24	8	81	19	15	8	2	9	13	
D6-4	9	3	1	6	12	11		4	13	68	4	1		1	24	6	73	11	18	6	2	6	1	
D6-3	6		3	4	11	1	8		3	69	9				26	71	21		11	9	2	2	4	
D6-2	3				2	3			1	27	2	2	2		3					1	3	5	0	6
D6-1	0								3	17	3	2	4	1	2					3	4	10	0	3

are the reworked species.

D6-13 and D6-14. These upper two assemblages are entirely marine and composed mainly of leiospheres along with the small chorate dinoflagellate cyst *Implettosphaeridium* spp. This latter form fluctuates in abundance through the section above sample D6-02, reaching a peak in sample D6-13. No other *in situ* dinocysts or any *in situ* terrestrial components were observed. There is no change in total frequencies of specimens recovered compared with the underlying part of the section (Table 1).

Unlike most of the La Meseta Formation samples, Polonez Cove and Boy Point Formation samples contain very few palynomorphs (Table 2). The overall number of identifiable palynomorphs observed is summarised in Figure 9, our interpretations of *in situ* vs. reworked specimens are given in Figure 10, and counts are given in Table 4. The listed samples are from the Krakowiak Glacier Member (KGM), Bay View Member (BM), Low Head Member (LHM), and the Chlamys Ledge Member (CLM) of the Polonez Cove Formation (listed from oldest to youngest), and from the lower Boy Point (BP) Formation (Figure 6).

sampled by the SHALDRIL programme off the Antarctic Peninsula (Warny & Askin 2011b). Many of the latter samples contained higher proportions of fine-grained carbonaceous material and thus likely more palynomorphs, as well as being deposited in a marine environment more conducive to palynomorph deposition than the diamictites and volcanobreccias of the Polonez Cove samples.

There were no age-diagnostic marine *in situ* palynomorphs observed in these samples. Radiometric dating for these beds is outlined in [Section 2.2](#) (Geological setting). Leiospheres dominate in these very sparse assemblages, typically in overwhelming abundance. Of the non-leiosphere fraction, most specimens are reworked (~85% of the non-leiosphere fraction). Many reworked specimens are of such high thermal maturity that they are unidentifiable, and thus are not included in any counts. The identifiable reworked assemblages, readily differentiated from *in situ* by their darker exines, are of possible Jurassic but mainly Cretaceous and Paleogene age, and include rare dinoflagellate cysts and fluctuating proportions of terrestrial cryptogam spores, podocarp conifer pollen and various angiosperm pollen ([Figures 9 and 10](#); [Table 4](#)).

The *in situ* assemblage throughout the Polonez Cove Formation and Boy Point Formation composite section is notable for its extreme lack of diversity, being composed almost entirely of the marine sphaeromorph acritarchs *Leiosphaeridia* spp. The few exceptions to this observation in the Polonez Cove Formation (see Table 4) include the following.

- A few specimens of terrestrial angiosperm pollen *Chenopodipollis* sp. occur in the basal KGM (sample G-3, from a marine lodgement till, [Figure 6](#)), and in the LHM (samples G-8, LHMb-cb (Conglomerate Bluff), LHMb),




Plate 1. Light photomicrographs of palynomorphs considered *in situ*. Images 1–11 are from the La Meseta Fomation, Seymour Island, and 12 is from the Polonez Cove Formation, King George Island. 1–4. **Various beech pollen grains:** 1–2. *Fusca* group: *Nothofagidites rocaensis-saraensis* complex (1. sample D6-09 16.2×139.1 ; 2 sample D6-08 15.9×141.7); 3. *Menziesii* group: *Nothofagidites* sp. cf. *N. asperus* (sample D6-11 15.5×117.5), 4. *Brassii* group: *Nothofagidites mataurensis* (sample D6-05 4.1×143.5). 5–8. **Other angiosperms:** 5. *Myricipites harrisii* (sample D6-01 14.2×119.4), 6. *Proteacidites* sp. cf. *P. parvus* (sample D6-01 13.4×143.9), 7. *Peninsulapollis askiniae* (sample D6-03 16.2×147.3), 8. *Peninsulapollis gillii* (sample D6-03 3.4×144.4). 9. **Gymnosperm:** *Phyllocladites* sp. cf. *P. exiguus* (sample D6-08 13.3×147.6). 10–12. **Marine palynomorphs:** 10. *Vozzhennikovia rotunda* (sample D6-01 6.3×117.8), 11. *Spinidinium macmurdoense* (sample D6-12 14.6×144.9), 12. *Leiosphaeridia* sp. (Sample G-13 12.1×141.0). All images were taken at $60\times$ under oil immersion. Scale bar = $20\ \mu\text{m}$.

- There are rare occurrences of podocarp conifer pollen (*Phyllocladites* spp. and *Podocarpidites* spp.) in the lower part of the section (samples G3 and L-6),
- Scattered occurrences of *Nothofagidites* spp. (*fusca* group) were observed in the basal KGM (sample G-3), in the upper glaci-marine KGM (samples K1, L5), and in some of

- the volcanoclastic marine samples of the BM (sample L-6), LHM (sample G-8) and CLM (sample L-11),
- The highest abundance of leiospheres was encountered in the LHM (sample LHMb), and the dinocyst *Impletosphaeridium* spp. occurs near the top of the Polonez Cove Formation in the CLM (sample L-11).

Table 4. Palynomorph counts from Polonez Cove and Boy Point formation samples. Reworked specimens are listed in pink and in situ specimens are listed in blue.

MEMBER/FORMATION	SAMPLE NAME	Sample #	Dinoflagellate cysts					Spores		Gymnosperms	Angiosperms										Reworking									
			<i>Deflandrea</i> spp.	<i>Dicodinium</i> sp.	<i>Impletosphaeridium</i> spp.	<i>Odontochitina</i> spp.	<i>Leiosphaeridia</i> spp.	<i>Calamospira</i> spp.	<i>Cyathidites minor</i>	<i>Dictyophyllidites</i> spp.	<i>Stereosporites antiquasporites</i>	<i>Alisporites</i> sp.	<i>Microcachrydites antarcticus</i>	<i>Phyllocladites</i> spp.	<i>Podocarpidites</i> spp.	<i>Chenopodipollis</i> spp.	<i>Haloragacidites</i> spp.	<i>Myricipites</i> spp.	<i>Nothofagidites</i> spp. (<i>brassii</i> gp.)	<i>Nothofagidites</i> spp. (<i>fusca</i> gp.)	<i>Nothofagidites</i> spp. (<i>fusca</i> gp.)	<i>Peninsulapollis</i> spp.	<i>Proteacidites</i> spp.	<i>Tricolpites</i> spp.	<i>Tripolipollenites</i> spp.	Reworked Counts Marine	In Situ Counts Marine non sea ice	In Situ Marine sea-ice indicator	Reworked Counts Terrestrial	In Situ Counts Terrestrial
BP	L-12	16						1		1	2			1											4	0	0	2	9	0
	L-13	15					3					2	2								1	1		1	1	0	0	3	4	0
CLM	G-13	14					4		2				2							1	1		1	1	1	0	0	4	7	0
	L-11	13		3			27					4	4		2				2	7					1	0	0	30	17	2
LHM	LHMB	12					83		1			3	4	1	1					6			2	2	2	0	0	83	17	1
	LHMB-cb	11					22		1			2	3	3	1	2	1			2			1	5	0	0	0	22	17	1
	G8	10			1		13						6	6	3	2			1	7			4	4	1	0	13	19	4	
	LRx	9					34		2				2	4						3	1				0	0	34	12	0	
BM	L-6	8	1				42		2			3	5					5	2	8		3		2	1	0	42	28	2	
KGM	L-5	7					39		2	1			6		2				1					3	0	0	39	14	1	
	K-1	6					31	1	9	2			3							2	11		4	0	0	0	31	30	2	
	KGLMb-4	5	1				51					3	21					3		11			4	1	1	0	51	43	0	
	KGLMb-3	4					37					2		1						5					0	0	37	8	0	
	KGLMb-2	3					1						5							2					0	0	1	9	0	
	KGLMb-1	2					0																		0	0	0	0	0	0
	G3	1				2	10	1			2		7		3	2	2		5				4	2	2	2	0	10	20	8

are the in situ species.

are the reworked species.

KGM = Krakowiak Glacier Member; BM = Bay View Member; LHM = Low Head Member; CLM = Chlamys Ledge Member; BP = Bay Point Formation.

The two assemblages from the basal Boy Point Member are very sparse, consisting of a few leiospheres. No *in situ* terrestrial material was observed.

6. Discussion

6.1. Uppermost La Meseta Formation

6.1.1. Terrestrial vegetation

Amongst the terrestrial component, *Phyllocladites* spp. and *Podocarpidites* spp. represent southern podocarp conifers which, together with the southern beech *Nothofagidites* spp. (*fusca* gp.) and *Nothofagidites* spp. (*menziesii* gp.), thrive in cool temperate humid climates (Veblen et al. 1996; Farjon 2010). As noted in Section 4 (Methods), the penecontemporaneous nature of *Nothofagidites* spp. (*brassii* gp.) through this section is equivocal. Thus, for the lower D6 section of this La Meseta dataset, we can infer the environment was comparable to the cool temperate Valdivian-type forest found in parts of modern-day Chile, as described in Veblen et al. (1996), although the diversity of the recovered La Meseta assemblages is somewhat lower. Podocarp conifer pollen interpreted as *in situ* were not observed above sample D6-08, and above this level the only woody taxa were *Nothofagidites* spp. These too disappeared before the upper two samples. This conclusion, however, assumes correct reworked vs. *in situ* interpretations. We note that Chen (2000) reported *Nothofagidites* spp. in sample D6-13, and Askin (unpublished data) recorded apparently *in situ* podocarp conifer and *Nothofagidites* pollen in both samples D6-13 and D6-14, albeit very sparse in the upper sample. Overall, the frequency and relative abundance of terrestrial pollen decrease towards the top of the section, which suggests deteriorating climatic conditions (also considering the other palynomorph data

noted below) and possibly a shift from forest to more stunted woodland-shrubby trees surviving in colder temperatures. The composition of the latest Eocene pollen record is ambiguous, but if penecontemporaneous, the presence of woody plants suggests that land temperatures remained above the ~10 °C January mean that delimits the modern austral polar-alpine treeline (Raine 1998; Körner and Paulsen 2004).

6.1.2. Marine and sea-ice influence

Small numbers of the typical Eocene dinocysts *Senegalinium asymmetricum*, *Spinidinium macmurdoense* and *Vozzhennikovia* spp. occur through much of the section, becoming less frequent upsection. These *in situ* dinoflagellates died out near the top of the section, coincident with an increase in the small dinocyst *Impletosphaeridium* spp. and great abundance of leiospheres. The leiospheres, assigned to *Leiosphaeridia* spp., a group usually associated with sphaeromorph acritarchs, are known to be abundant at the limit between pack ice and sea ice (Mudie 1992; Troedson & Riding 2002; Warny et al. 2006). Similar to *Leiosphaeridia* spp., *Impletosphaeridium* spp. are believed to be present during sea-ice formation (Warny et al. 2007).

6.1.3. Environmental trends

Several simultaneous trends in environmental indicators are evident in the upper part of the section, from sample D6-08 to the top. Approaching the end of the Eocene, the data include a dearth of penecontemporaneous terrestrial palynomorphs, the overwhelming abundance of *in situ* sea-ice-indicative leiospheres, increased numbers of sea-ice-indicative *Impletosphaeridium* spp., decreasing numbers of other *in situ* dinoflagellates, and towards the top an increasing relative abundance of reworked specimens. Together these imply a less

Polonez Cove




Figure 9. Overall frequency (raw counts) of palynomorph species observed in the Polonez Cove and Boy Point samples from King George Island. For details about the position of samples in relation to the formations and sampled members, see [Figure 6](#). Blue = *in situ* and red = reworked.

vegetated, more glacially dominated environment for the uppermost part of the La Meseta Formation.

Our data corroborate the marine latest Eocene temperature record for Seymour Island (e.g. Dutton et al. 2002; Ivany et al. 2006, 2008), and echo the marked cooling at the EOT in the Zachos et al. (2008) $\delta^{18}\text{O}$ curve. This trend was also evidenced by the microtexture analysis of sand grains (Anderson et al. 2011) from D6 samples (D6-03, D6-05, D6-07) which revealed high-stress glacially induced surface textures from relatively

low (<33%) occurrence, compared with younger, more glacially influenced sediments from the area, to medium abundance higher in the section. This analysis supported encroaching glacial activity in the vicinity in the latest Eocene. By the EOT, the presence of glacial ice was also indicated by the Ivany et al. (2006) report of diamict at the Eocene–Oligocene boundary on Seymour Island.

Interestingly, the Late Eocene palynomorph assemblages described from SHALDRIL II 3C sediment cores offshore and




Figure 10. Relative abundance of reworked (Rw) and *in situ* palynomorphs in Polonez Cove and Boy Point Formation samples. For details about the position of samples in relation to the formations and sampled members, see Figure 6.

north-east of Seymour Island (Warny & Askin 2011a) may actually predate at least the upper D6 samples, rather than postdate the youngest La Meseta Formation sediments as concluded by Anderson et al. (2011) based on seismic evidence. Diatom stratigraphy placed the 3C cores between 37 and 32 Ma, and a strontium date halfway through the 3C composite section provided an age of 35.9 Ma (Bohaty et al. 2011). Both the marine and terrestrial 3C assemblages were more diverse and more abundant (see Section 5.1) than those encountered in the latest Eocene Seymour D6 samples, and the more refined chronostratigraphy for the top of the La Meseta Formation, as discussed in Section 2, supports this contention. Furthermore, although the upper 3C

assemblages record significant cooling and a sea-level drop with an increase in erosion and reworking (Warny & Askin 2011a), these 3C samples lack the abundant sea-ice-indicative leiosphere acritarchs, and thus suggest sea-ice was not yet widespread during their deposition. This is in agreement with temperature data reconstructed by Feakins et al. (2014), who modeled leaf wax hydrogen isotopic evidence from the SHAL-DRIL II 3C sediments and combined their results with the pollen data of Warny & Askin (2011a). Their δD and modeling show cooling and drying conditions during that latest Eocene time, but with temperatures remaining above freezing, from ca. 7 to 2 °C and precipitation around 700 to 600 mma^{-1} with an isotopic shift in δD of about -15‰ .

6.2. Polonez Cove Formation and Boy Point Formation

6.2.1. Terrestrial vegetation

Our sampling on King George Island covers a period from Early to Late Oligocene. Farther west on King George Island, the *Cytisella* flora of the Point Thomas Formation, previously thought to be of Early Oligocene age (e.g. Birkenmajer & Zastawniak 1989), is now believed to represent preglacial Eocene vegetation (Mozer 2012, 2013). In addition, the various Point Hennequin floras in the Mount Wawel Formation, previously thought to be of Late Oligocene age (Birkenmajer & Zastawniak 1989), are now included in the Eocene (Hunt & Poole 2003; Nawrocki et al. 2011; Mozer 2013). This is also the case for several other flora sites on King George Island (Mozer et al. 2015). Thus, the current study provides important new information on the Oligocene vegetation.

The Oligocene sections studied from the Polonez Cove Formation indicate that by that time, on the west side of the Antarctic Peninsula, vegetation was reduced to a sparse tundra flora consisting of a few herbaceous plants with rare podocarp conifers in the lower section and some southern beech. The woody beech plants were probably of prostrate habit, surviving in sheltered locations in severe conditions (e.g. Francis & Hill 1996; Raine 1998; Askin & Raine 2000). Although occasional *in situ* terrestrial pollen were observed, the vast majority of the terrestrial palynomorphs in the Polonez Cove samples were reworked, which is consistent with advanced ice sheet expansion, and ice erosion and transport on the Peninsula related to the Polonez Glaciation.

6.2.2. Sea-ice influence

As evidenced by the overwhelming dominance of leiospheres in most of our samples, plus some *Impletosphaeridium* spp. in one sample, sea ice persisted in the waning stages of the Polonez Glaciation, during deposition of the upper KGM and overlying shallow marine, glacially influenced (at least for the lower samples), and basaltic volcanism-dominated BM, LHM and CLM, and lowermost BP Formation.

6.2.3. Environmental trends and volcanism

Figures 9 and 10 highlight how the overall abundance, diversity and preservation of palynomorphs have diminished compared to those of La Meseta Formation. The lower Polonez Cove samples from the KGM represent sediments deposited during the Polonez Glaciation, with palynomorphs from the sea-ice flora and a few pollen from a very sparse periglacial terrestrial flora. The Antarctic periglacial tundra vegetation that survived from the EOT, and through the Oligocene and Miocene, in non-glaciated locations has been described from SHALDRIL cores off Seymour Island (Anderson et al. 2011; Warny & Askin 2011b) and from as far away as the southern Victoria Land margin of the Ross Sea (e.g. Raine 1998; Askin & Raine 2000; Raine & Askin 2001; Feakins et al. 2012; Greiner et al. 2015). Compared to these other records, the vegetation recovered from the Polonez Cove Formation was even sparser. Our data imply that local terrestrial vegetation barely survived through the Polonez Glaciation. The palynological record suggests conifers are present only in the lower part of the section. Southern

beech fared a little better with their interpreted *in situ* record extending up into the late Oligocene top of the Polonez Cove Formation. Because of the scarcity of unambiguous terrestrial remains, we refrain from making comparisons to modern climate parameters (temperatures, etc.). The only recovered record of herbaceous plants considered *in situ* is pollen of Chenopodiaceae, a group that evidently survived well into the Neogene in the Antarctic Peninsula area (Warny & Askin 2011b). This great dearth of local vegetation is hardly surprising, considering the plants had to contend with numerous episodes of local volcanism as well as glacial climates and ice. The presence of Chenopodiaceae and very little else is consistent with these harsh conditions, as today these are typically weedy plants that are well adapted to disturbed and chemically challenging soils.

Farther to the east on King George Island, it seems that at least some vegetation regained a foothold in the region as it emerged from the worst of the volcanism. From the Destruction Bay Formation, of Late Oligocene age (25.3 Ma, Dingle & Lavelle 1998), Troedson & Riding (2002) described a 'moderately diverse' assemblage with abundant pollen of *Nothofagidites* spp. (*fusca* group), with, among other taxa, some podocarp pollen and *Cyathidites* (fern) spores, plus dinoflagellate cysts. Sparse, low-diversity assemblages of mainly podocarp pollen, with a few *Nothofagidites* pollen, *Cyathidites* and indeterminate spores, plus dinocysts, were reported by Troedson & Riding (2002) from the early Miocene Cape Melville Formation. Also from the early Miocene Cape Melville Formation, the *in situ* palynoflora described by Warny et al. (2016) contained *Nothofagidites* spp. (*fusca* group), rare podocarp pollen, moss spores (*Coptospora*), and pollen of Asteraceae, Caryophyllaceae (*Colobanthus*-type) and Chenopodiaceae. These assemblages, from sediments deposited during the Melville Glaciation, are somewhat richer in diversity than the Polonez assemblages and represent a periglacial tundra flora with components common to other Antarctic Miocene tundra assemblages (Warny et al. 2016).

Nearby, to the east of the tip of the Antarctic Peninsula, there are palynomorph assemblages of Late Oligocene age (24.0–28.6 Ma from diatom biostratigraphy, Bohaty et al. 2011) from the SHALDRIL II 12A sediment cores. Palynomorph concentrations, however, are substantially less at Polonez Cove and the terrestrial flora is far less diverse than that described from the 12A cores (Warny & Askin 2011b). Probably, as noted above, the intermittent volcanism has caused this dearth of terrestrial vegetation on the northern Antarctic Peninsula magmatic arc. On both sides of the northernmost peninsula the marine component is dominated by a sea-ice acritarch flora, but with a difference. The Polonez Cove samples are dominated by *Leiosphaeridia* spp., and the 12A samples by a *Michrystidium*/*Leiosphaeridia* association, with the former acanthomorph acritarchs typically more common except for the uppermost two 12A samples where the leiospheres predominate. We note that the dinocyst *Impletosphaeridium* and the acritarch *Michrystidium* are morphologically similar and likely occupied similar ecological niches. It is possible that the difference comes from the fact that the Polonez Cove samples are from a nearshore depositional site with closer proximity to glaciers and their freshwater

outwash, while the SHALDRIL II 12A samples were from a more marine palaeoenvironment, albeit within a partly land-locked back-arc basin.

7. Conclusions

Palynomorph assemblages from the La Meseta Formation on Seymour Island, and from the Polonez Cove and Boy Point formations on King George Island, provide insight into latest Eocene and Oligocene climatic evolution in the northern Antarctic Peninsula. La Meseta Formation assemblages capture a small slice of time at the end of the Eocene, recording the shift from cool temperate, humid Valdivian-type forest to a more depauperate vegetation. This was accompanied by a decrease in typical Eocene dinoflagellate cysts, an increase in sea-ice-indicative marine phytoplankton, and an increase in reworked palynomorphs. We suggest these palynofloras record a shift from cool temperate to periglacial conditions and a subpolar climate just before the EOT boundary in the back-arc James Ross Basin.

By the early to late Oligocene, on the other side of the Antarctic Peninsula, land vegetation had been decimated by both glaciation and volcanism in the northern Antarctic Peninsula magmatic arc. The terrestrial palynomorph assemblage is mostly reworked and penecontemporaneous traces of terrestrial vegetation are sparse. We believe the apparent record of extremely reduced vegetative cover at this time is overprinted by regional volcanism and cannot be accurately used as an indicator of climate. The marine palynoflora does, however, indicate the presence of sea ice and thus a polar to subpolar climate.

Acknowledgments

The authors would like to thank the Institute of Geological Sciences, Polish Academy of Sciences, for providing samples from the Polonez Cove and Boy Point formations on King George Island. Field assistance of Michał Guźniczak, Marcin Klisz, Mariusz Potocki and Mateusz Zabłocki during 2007 and 2009 expeditions is greatly appreciated. Fieldwork on King George Island was financially supported by the Polish Ministry of Science and Higher Education (Grant No. DWM/N8IPY/2008). Samples from the section on Seymour Island were collected for R. Askin by Fred Barbis and Tim Kelley as part of the 1986–1987 US Antarctic Program, and splits were subsequently obtained for this study from the Polar Rock Repository. This is a contribution to the Antarctic Climate Evolution (ACE) programme. We are grateful to the NSF Polar and CAREER programme for funding this research. Funding for this research was provided by the US National Science Foundation ANT-1048343 to SW. We appreciate our reviewers' suggestions, which greatly improved this paper. Thanks are extended to the HESS corporation for funding the Stratabugs software license for CENEX and to HESS staff members (D. Pocknall, P. Griggs and M. Thomas) for Stratabugs training, and to M. Thomas for help with some of the graphics. This article is dedicated to the memory of our colleague Dr Krzysztof Krajewski who passed away suddenly during the final stage of preparation of this manuscript. His knowledge and contribution to Antarctic geology will be greatly missed.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the Polish Ministry of Science and Higher Education (to KPK and AT) [grant number DWM/N8IPY/2008]; and the US National Science Foundation (to SW) [grant number ANT-1048343].

Notes on contributors



SOPHIE WARNY is an associate professor of palynology in the Department of Geology and Geophysics and a curator at the Museum of Natural Science at Louisiana State University in Baton Rouge. She has a long history with AASP as she won the AASP Student Award in 1996, served as Director-At-Large on the AASP board from 2006 to 2007, was the AASP newsletter editor from 2006 to 2015, and is now the AASP Chair in Palynology.

She received her PhD degree from the Université Catholique de Louvain, in Belgium working with Dr Jean-Pierre Suc on the Messinian Salinity Crisis. In 2011, she received a NSF CAREER award to conduct palynological research in Antarctica. Since being hired at LSU in 2008, she directed 18 graduate students' theses on various Cretaceous to Cenozoic sections. Her students are now employed with the oil and gas industry (HESS, BP, DEVON, CHEVRON, BHP Billiton Petroleum, and EOG), with the US Department of Homeland Security, with environmental companies, with IODP, or as instructor.



C. MADISON KYMES earned an MS degree in palynology at CENEX (the Center for Excellence in Palynology) in the Department of Geology and Geophysics at Louisiana State University, Louisiana, USA, under the direction of S. Warny and R. Askin in 2015. Madison was born in Shreveport, Louisiana, but spent the majority of his life growing up in Madison, Mississippi. During his MS degree, Madison interned as an environmental

geologist with a consulting company and was hired, after his graduation, by the Mississippi Department of Environmental Quality, in their office of Land and Water, in Jackson, Mississippi.



ROSEMARY ASKIN is currently a palynological consultant in Jackson, Wyoming. She earned her PhD degree from Victoria University of Wellington, New Zealand. She has conducted research and taught in several US universities, most recently at Byrd Polar Research Center, The Ohio State University. Her most recent project there was establishing the US Polar Rock Repository. Her specialty is the use of palynology as a tool in biostratigraphy

and paleoenvironmental reconstruction. Current research projects primarily involve taxonomy, evolution and distribution of Cretaceous–Neogene terrestrial palynomorphs from the Ross Sea region, Siple Coast and northern Antarctic Peninsula; and on Permian–Triassic palynomorphs of the Transantarctic Mountains. She served as Director-At-Large on the AASP Board from 1994 to 1996, and was elected a fellow of the Geological Society of America in 1997.



KRZYSZTOF P. KRAJEWSKI was a professor of earth sciences at the Institute of Geological Sciences, Polish Academy of Sciences in Warszawa. He obtained his PhD degree in 1986, and continued his studies and research at the Polish Academy of Sciences, University of Oslo, and University of Oldenburg. He completed his DSc degree in 2001 in the fields of sedimentary petrology and geochemistry. His scientific interests focused on the interactions

between biological activity and mineral deposition in carbonate, phosphate, and organic carbon-rich sedimentary systems. For more than 20 years, Krajewski carried out geological research in the Arctic, where he

concentrated on the Mesozoic phosphorites and petroleum source beds. He was also involved in the research on the evolution of Cenozoic sedimentary systems in Antarctica. Krzysztof passed away on 21 November 2017, during the final stages of revision of this paper. His field expertise and knowledge of the Antarctic Peninsula will be greatly missed.



ANDRZEJ TATUR graduated from the Department of Geology of Warsaw University in the field of petrography, geochemistry and mineralogy in 1969. From 1970 to 2002, Tatur worked for the Institute of Ecology, at the Polish Academy of Sciences. In 2011, he became the director of the Department of Polar Biology, and then was hired as a full professor in the Department of Geology at University of Warsaw where he still works today. He obtained his PhD degree in 1978, habilitation in 2002 and professor degree in 2011. He took part in more than 10 geological and ecological expeditions with Polish, Argentinian and Russian teams to Antarctica, Patagonia and to the Arctic. He participated in several international EU and Antarctic Scientific projects and published over 100 publications that are available in Google Scholar. From 2005 to 2011, he served as the expert in Natural Environmental Protection for the Polish Delegation of the Ministry of Foreign Affairs at annual ATCM/CEP Meetings.

References

- Anderson JB, Warny S, Askin RA, Wellner JS, Bohaty SM, Kirshner AE, Majewski W. 2011. Progressive Cenozoic cooling and the demise of Antarctica's last refugium. *Proceedings of the National Academy of Sciences* 239; 108 (28):11356–11360.
- Askin RA. 1997. Eocene–earliest Oligocene terrestrial palynology of Seymour Island, Antarctica. In: Ricci CA, editor. *The Antarctic region: Geological evolution and processes*. Siena (Italy): Terra Antarctica Publication; p. 993–996.
- Askin RA, Elliot DH. 1982. Geologic implications of recycled Permian and Triassic palynomorphs in Tertiary rocks of Seymour Island, Antarctic Peninsula. *Geology* 10(4):547–551.
- Askin RA, Raine JL. 2000. Oligocene and early Miocene terrestrial palynology of the Cape Roberts drillhole CRP-2/2A, Victoria Land Basin, Antarctica. *Terra Antarctica* 7(4):493–501.
- Barker DHN, Austin JA. 1998. Rift propagation, detachment faulting, and associated magmatism in Bransfield Strait, Antarctic Peninsula. *Journal of Geophysical Research* 103:24017–24043.
- Benninghoff WS. 1962. Calculation of pollen and spore density in sediments by addition of exotic pollen in known quantities. *Pollen et Spores* 4 (2):332–333.
- Bijl PK, Sluijs A, Brinkhuis H. (2013). A magneto- and chemostratigraphically calibrated dinoflagellate cyst zonation of the early Palaeogene South Pacific Ocean. *Earth-Science Reviews* 124:1–31.
- Birkenmajer K. 1980. Discovery of Pliocene glaciations on King George Island, South Shetland Islands (west Antarctica). *Académie Polonaise des Sciences, Bulletin, Série Science de la Terre* 29:331–340.
- Birkenmajer K. 1982. Pliocene tillite-bearing succession of King George Island (South Shetland Islands, Antarctica). *Studia Geologica Polonica* 74:7–72.
- Birkenmajer K. 1987. Oligocene–Miocene glacio-marine sequences of King George Island (South Shetland Islands), Antarctica. *Palaeontologia Polonica* 239; 49(1):9–36.
- Birkenmajer K. 2001. Mesozoic and Cenozoic stratigraphic units in parts of the South Shetland Islands and Northern Antarctic Peninsula (as used by the Polish Antarctic Programmes). *Studia Geologica Polonica* 118:5–188.
- Birkenmajer K, Gaździcki A. 1986. Age of the Pecten conglomerate on King George Island, West Antarctica. *Bulletin of the Polish Academy of Sciences, Earth Sciences* 34:219–226.
- Birkenmajer K, Zastawniak E. 1989. Late Cretaceous–early Tertiary floras of King George Island, West Antarctica: their stratigraphic distribution and palaeoclimatic significance. In: Crame JA, editor. *Origins and evolution of the Antarctic Biota*. Geological Society Special Publication 47:277–240.
- Bohaty SM, Kulhanek DK, Wise SW Jr, Jennison K, Warny S, Sjunneskog C. 2011. Age assessment of Eocene–Pliocene drill cores recovered during the SHALDRIL II expedition, Antarctic Peninsula. In: Anderson JB, Wellner JS, editors. *Tectonic, climatic and Cryospheric evolution of the Antarctic Peninsula*. AGU Special Publication 63:63–114.
- Bowman VC, Francis JE, Riding JB, Hunter SJ, Haywood AM. 2012. A latest Cretaceous to earliest Paleogene dinoflagellate cyst zonation from Antarctica, and implications for phytoprovincialism in the high southern latitudes. *Review of Palaeobotany and Palynology* 171:40–56.
- Brown C. 2008. Palynological techniques. In: Riding JB, Warny S, editors. *AASP Special Publications*; p. 137. ISBN 9780931871078.
- Chen B. 2000. Evolutionary, biogeographic and paleoclimatic implications of Eocene Nothofagidites pollen from Seymour Island, Antarctica [Unpubl. MS Thesis]. The Ohio State University; p. 105.
- Dingle RV, Lavelle M. 1998. Antarctic Peninsula cryosphere: Early Oligocene (ca. 30 Ma) initiation and a revised glacial chronology. *Geological Society (London) Journal* 155:433–437.
- Dingle RV, McArthur JM, Vroon P. 1997. Oligocene and Pliocene interglacial events in the Antarctic Peninsula dated using strontium isotope stratigraphy. *Journal of the Geological Society* 239; 154(2):257–264.
- Douglas PMJ, Affek HP, Ivany LC, Houben AJP, Sijp WP, Sluijs A, Schouten S, Pagani M. 2014. Pronounced zonal heterogeneity in Eocene southern high-latitude sea surface temperatures. *Proceedings of the National Academy of Sciences* 239; 111(18):6582–6587.
- Dutton AL, Lohmann KC, Zinsmeister WJ. 2002. Stable isotope and minor element proxies for Eocene climate of Seymour Island, Antarctica. *Paleoceanography* 239; 17(2):6–13.
- Elliot DH, Rinaldi C, Zinsmeister WJ, Trautman TA, Bryant WA, Valle R. 1975. Geological investigations on Seymour Island, Antarctic Peninsula. *Antarctic Journal of the United States* 239; 10(4):182–186.
- Elliot DH, Trautman TA. 1982. Lower Tertiary strata on Seymour Island, Antarctic Peninsula. In: Craddock C, editor. *Antarctic geosciences*. Madison (WI): University of Wisconsin Press; p. 287–297.
- Farjon A. 2010. *A handbook of the World's Conifers*, 2 vols. Leiden & Boston: J. Brill; p. 1111.
- Feakins SJ, Warny S, DeConto R. 2014. Snapshot of cooling and drying before onset of Antarctic Glaciation. *Earth and Planetary Science Letters* 404:154–166.
- Feakins SJ, Warny S, Lee J-E. 2012. Hydrologic cycling over Antarctica during the middle Miocene warming. *Nature Geoscience* 5:557–560.
- Francis JE, Hill RS. 1996. Fossil plants from the Pliocene Sirius Group, Transantarctic Mountains: evidence for climate from growth rings and fossil leaves. *Palaios* 11:389–396.
- Gaździcki A, Tatur A, Hara U, Valle RAD. 2004. The Weddell Sea Formation: post-late Pliocene terrestrial glacial deposits on Seymour Island, Antarctic Peninsula. *Polish Polar Research* 25:189–204.
- Gradstein FM, Ogg JG, Schmitz MD, Ogg GM. 2012. *The geologic time scale 2012*. Amsterdam: Elsevier BV; p. 1144.
- Greiner KW, Warny S, Askin R, Acton G. 2015. Early to middle Miocene vegetation history of Antarctica supports eccentricity-paced warming intervals during the Antarctic icehouse phase. *Global and Planetary Change* 127:67–78.
- Griener KW, Nelson DM, Warny S. 2013. Declining moisture availability on the Antarctic Peninsula during the Late Eocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 239; 383:72–78.
- Hunt RJ, Poole I. 2003. Revising Paleogene West Antarctic climate and vegetation history in light of new data from King George Island. In: Wing SL, Gingerich PD, Schmitz B, Thomas E, editors. *Causes and consequences of globally warm climates in the Early Paleogene*. Boulder (CO): GSA Special Paper 369; p. 395–412.
- Ivany LC, Lohmann KC, Hasiuk F, Blake DB, Glass A, Aronson RB, Moody RM. 2008. Eocene climate record of a high southern latitude continental shelf: Seymour Island, Antarctica. *Geological Society of America Bulletin* 120(5/6):659–678.
- Ivany LC, Van Simaeys S, Domack EW, Samson SD. 2006. Evidence for an earliest Oligocene ice sheet on the Antarctic Peninsula. *Geology* 34(5):377–380.
- Körner, C, Paulsen J. 2004. A world-wide study of high altitude treeline temperatures. *Journal of Biogeography* 31:713–732.
- Kymes CM. 2015. A Palynological analysis of Seymour Island and King George Island off the Antarctic Peninsula: A Dating and Climatic

- Reconstruction [LSU Master's Theses Series. 392]. Baton Rouge (LA): Louisiana State University. p. 55.
- Lear CH, Bailey TR, Pearson PN, Coxall HK, Rosenthal Y. 2008. Cooling and ice growth across the Eocene-Oligocene transition. *Geology* 239; 36(3):251–254.
- Lentin JK, Williams GL. 1981. Fossil dinoflagellates: index to genera and species, 1981 edition. Dartmouth: Bedford Institute of Oceanography Report Series. BI.
- Liu Z, Pagani M, Zinniker D, DeConto R, Huber M, Brinkhuis H, Pearson A. 2009. Global cooling during the Eocene-Oligocene climate transition. *Science* 239; 323(5918):1187–1190.
- Marensi A, Santillana SN. 1994. Unconformity bounded units within the La Meseta Formation, Seymour Island, Antarctica: a preliminary approach. Warsaw: XXI Polar Symposium; p. 33–37.
- Marensi SA, Casadio S, Santillana SN. 2010. Record of Late Miocene glacial deposits on Isla Marambio (Seymour Island), Antarctic Peninsula. *Antarctic Science* 22:93–198.
- Marensi SA, Santillana SN, Rinaldi CA. 1998a. Paleoambientes sedimentarios de la Aloformación La Meseta (Eoceno), Isla Marambio (Seymour), Antártida (No. 464). Buenos Aires: Dirección Nacional del Antártico, Instituto Antártico Argentino.
- Marensi SA, Santillana SN, Rinaldi CA. 1998b. Stratigraphy of the La Meseta Formation (Eocene), Marambio (Seymour) Island, Antarctica. *Paleógeno de América del Sur y de la Península Antártica* 239; 5:137–146.
- Mozer A. 2012. Pre-glacial sedimentary facies of the point Thomas formation (Eocene) at Cyadela, Admiralty Bay, King George Island, West Antarctica. *Polish Polar Research* 33(1):41–62.
- Mozer A. 2013. Eocene sedimentary facies in a volcanogenic succession on King George Island, South Shetland Islands: a record of pre-ice sheet terrestrial environments in West Antarctica. *Geological Quarterly* 57:385–394.
- Mozer A, Pécskay Z, Krajewski KP. 2015. Eocene age of the Baranowski Glacier Group at Red Hill, King George Island, West Antarctica. *Polish Polar Research* 36:307–324.
- Mudie PJ. 1992. Circum-arctic Quaternary and Neogene marine palynofloras: paleoecology and statistical analysis. In: Head MJ, Wrenn JH, editors. *Neogene and quaternary Dinoflagellate cysts and acritarchs*. Dallas: American Association of Stratigraphic Palynologists Foundation; p. 347–390.
- Nawrocki J, Pańczyk M, Williams IS. 2011. Isotopic ages of selected magmatic rocks from King George Island (West Antarctica) controlled by magnetostratigraphy. *Geological Quarterly* 55:301–322.
- Pańczyk M, Nawrocki J. 2011. Geochronology of selected andesitic lavas from the King George Bay area (SE King George Island). *Geological Quarterly* 55:323–334.
- Porębski SK. 1995. Facies architecture in a tectonically-controlled incised-valley estuary: La Meseta Formation (Eocene) of Seymour Island, Antarctic Peninsula. *Studia Geologica Polonica* 107:7–97.
- Porębski SK. 2000. Shelf-valley compound fill produced by shelf subsidence and eustatic sea-level changes, Eocene La Meseta Formation, Seymour Island, Antarctica. *Geology* 28:147–150.
- Porębski SK, Gradzinski R. 1987. Depositional history of the Polonez Cove Formation (Oligocene), King George Island, West Antarctica: a record of continental glaciations, shallow marine sedimentation and contemporaneous volcanism. *Studia Geologica Polonica* 93:7–62.
- Prothero DR, Berggren WA, editors. 2014. *Eocene-Oligocene climatic and biotic evolution*. Princeton (NJ): Princeton University Press; p. 584.
- Prothero DR, Ivany LC, Nesbitt ER. 2003. From greenhouse to icehouse: the marine Eocene-Oligocene transition. New York (NY): Columbia University Press; p. 560.
- Raine JI. 1998. Terrestrial palynomorphs from Cape Roberts drillhole CRP-1, Ross Sea, Antarctica. *Terra Antarctica* 5(3):539–548.
- Raine JI, Askin RA. 2001. Terrestrial palynology of Cape Roberts Project drill-hole CRP-3, Victoria Land Basin, Antarctica. *Terra Antarctica* 8(4):389–400.
- Rinaldi CA, Massabie A, Morelli J, Rosenmann LH, del Valle RA. 1978. *Geología de la isla Vicecomodoro Marambio, Antártida*. Buenos Aires (Argentina): Instituto Antártico Argentino Contribucion 217; p. 37.
- Sadler PM. 1988. Geometry and stratification of uppermost Cretaceous and Paleogene units on Seymour Island, northern Antarctic Peninsula. *Geological Society of America Memoirs* 239; 16:303–320.
- Sluijs A, Brinkhuis H, Stickley CE, Warnaar J, Williams GL, Fuller M. 2003. Dinoflagellate cysts from the Eocene/Oligocene transition in the Southern Ocean; results from ODP Leg 189. *Proceedings of the Ocean Drilling Program-Scientific Results* 189:1–42.
- Smellie JL, Millar IL, Rex DC, Butterworth PJ. 1998. Subaqueous, basaltic lava dome and carapace breccia on King George island, South Shetland Islands, Antarctica. *Bulletin of Volcanology* 239; 59(4):245–261.
- Stott LD, Kennett JP, Shackleton NJ, Corfield RM. 1990. The Evolution of Antarctic surface waters during the Paleogene: Inferences from the stable isotopic composition of planktonic foraminifers, ODP LEG 1131. *Proceeding of the Ocean Drilling Project Scientific Results* 239; 113:849–863.
- Stover LE, Evitt WR. 1978. Analyses of pre-Pleistocene organic-walled dinoflagellates. *Stanford University Publications, Geological Sciences* 15:1–300.
- Troedson AL, Riding JB. 2002. Upper Oligocene to lowermost Miocene strata of King George Island, South Shetland Islands, Antarctica: stratigraphy, facies analysis, and implications for the glacial history of the Antarctic Peninsula. *Journal of Sedimentary Research* 239; 72(4):510–523.
- Troedson AL, Smellie JL. 2002. The Polonez Cove Formation of King George Island, Antarctica: stratigraphy, facies and implications for mid-Cenozoic cryosphere development. *Sedimentology* 49:277–301.
- Veblen TT, Hill RS, Read J. 1996. *The Ecology and biogeography of nothofagus forests*. New Haven (CT): Yale University Press; p. 403.
- Warny S, Anderson JB, Londeix L, Bart PJ. 2007. Analysis of the dinoflagellate cyst genus *Impletosphaeridium* as a marker of sea-ice conditions off Seymour Island: An ecomorphological approach. *Denver (CO): US Geological Survey, No. 2007-1047-SRP-079*; p. 4.
- Warny S, Askin R. 2011a. Vegetation and organic-walled phytoplankton at the end of the Antarctic greenhouse world: latest cooling events. In: Anderson JB, Wellner JS, editors. *Tectonic, Climatic and Cryospheric Evolution of the Antarctic Peninsula*. Washington (DC): AGU Special Publication 63:193–210. doi:10.1029/2010SP000965.
- Warny S, Askin R. 2011b. Last remnants of Cenozoic vegetation and organic-walled phytoplankton in the Antarctic Peninsula's icehouse world. In: Anderson JB, Wellner JS, editors. *Tectonic, Climatic and Cryospheric Evolution of the Antarctic Peninsula*. Washington (DC): AGU Special Publication 63:167–192. doi:10.1029/2010SP000996.
- Warny S, Kymes CM, Askin RA, Krajewski KP, Bart PJ. 2016. Remnants of Antarctic vegetation on King George Island during the early Miocene Melville Glaciation. *Palynology* 40(1):66–82.
- Warny S, Wrenn JH, Bart PJ, Askin RA. 2006. Palynology of the NBP03-01A transect in Northern Basin, western Ross Sea, Antarctica: a Late Pliocene record. *Palynology* 30:151–182.
- Wilson GJ. 1967. Some new species of lower Tertiary dinoflagellates from McMurdo Sound, Antarctica. *New Zealand Journal of Botany* 239; 5(1):57–83.
- Zachos JC, Dickens GR, Zeebe RE. (2008). An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* 451(7176):279–283.
- Zachos J, Pagani M, Sloan L, Thomas E, Billups K. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to Present. *Science* 292:686–693.
- Zachos JC, Quinn TM, Salamy KA. 1996. High-resolution (104 years) deep-sea foraminiferal stable isotope records of the Eocene–Oligocene climate transition. *Paleoceanography* 239; 11(3):251–266.
- Zinsmeister WJ, DeVries TJ. 1983. Quaternary glacial marine deposits on Seymour Island. *Antarctic Journal U.S.* 18:64–65.