

Journal of Stochastic Analysis

Volume 1 | Number 4

Article 8

December 2020

An Asymptotic Formula for Integrals of Products of Jacobi Polynomials

Maxim Derevyagin

University of Connecticut, Storrs, CT 06269-1009, USA, maksym.derevyagin@uconn.edu

Nicholas Juricic

University of Connecticut, Storrs, CT 06269-1009, USA, nicholas.juricic@uconn.edu

Follow this and additional works at: <https://digitalcommons.lsu.edu/josa>

 Part of the [Analysis Commons](#), and the [Other Mathematics Commons](#)

Recommended Citation

Derevyagin, Maxim and Juricic, Nicholas (2020) "An Asymptotic Formula for Integrals of Products of Jacobi Polynomials," *Journal of Stochastic Analysis*: Vol. 1 : No. 4 , Article 8.

DOI: 10.31390/josa.1.4.08

Available at: <https://digitalcommons.lsu.edu/josa/vol1/iss4/8>

AN ASYMPTOTIC FORMULA FOR INTEGRALS OF PRODUCTS OF JACOBI POLYNOMIALS

MAXIM DEREVYAGIN* AND NICHOLAS JURICIC

Dedicated to Professor Leonard Gross on the occasion of his 88th birthday

ABSTRACT. We recast Byerly's formula for integrals of products of Legendre polynomials. Then we adopt the idea to the case of Jacobi polynomials. After that, we use the formula to derive an asymptotic formula for integrals of products of Jacobi polynomials. The asymptotic formula is similar to an analogous one recently obtained by the first author and Jeff Geronimo for a different case. Thus, it suggests that such an asymptotic behavior is rather generic for integrals of products of orthogonal polynomials.

1. Introduction

In [4] the Alpert multiresolution analysis was studied and important in this study was the integral

$$f_{n,m} = \int_0^1 \hat{p}_n(t) \hat{p}_m(2t - 1) dt,$$

where \hat{p}_n is the orthonormal Legendre polynomial to be precisely defined in the next section. These coefficients are entries in the refinement equation associated with the multiresolution analysis and they have been shown to satisfy a variety of difference equations. As a matter of fact, it turns out that more general coefficients of the form

$$u_{n,m} = \int_{\mathbb{R}} P_n(t) Q_m(\alpha t + \beta) d\sigma(t),$$

where $\{P_n\}_{n=0}^{\infty}$ and $\{Q_n\}_{n=0}^{\infty}$ are two families of orthonormal polynomials, σ is a measure on \mathbb{R} with finite moments, and the numbers $\alpha \neq 0, \beta$ are complex, satisfy a generalized wave equation on the two dimensional lattice

$$a_{n+1}u_{n+1,m} + b_nu_{n,m} + a_nu_{n-1,m} = \frac{c_{m+1}}{\alpha}u_{n,m+1} + \frac{d_m - \beta}{\alpha}u_{n,m} + \frac{c_m}{\alpha}u_{n,m-1}$$

for $n, m = 0, 1, 2, \dots$ (see [3] for details). It was also observed in [3] that a damped oscillatory behavior took place for such coefficients. In particular, for the

Received 2020-10-24; Accepted 2020-11-19; Communicated by A. Stan et al.

2010 *Mathematics Subject Classification.* Primary 33C45; Secondary 39A05; 39A14; 34D05.

Key words and phrases. Jacobi polynomials, asymptotic formulas, Darboux transformation, difference equations.

* Corresponding author.

coefficients

$$f_{n,m}^{(\lambda)} = \int_0^1 \hat{p}_n^{(\lambda)}(t) \hat{p}_m^{(\lambda)}(2t-1) (t(1-t))^{\lambda-1/2} dt,$$

where $\hat{p}_n^{(\lambda)}$ are the orthonormal ultraspherical polynomials and $\lambda > -1/2$, it was shown that

$$f_{n,m}^{(\lambda)} = k_m \frac{\cos(\pi(m + \frac{\lambda}{2} - \frac{n}{2} + \frac{1}{4}))}{\sqrt{\pi} n^{\lambda+1/2}} + O\left(\frac{1}{n^{\lambda+3/2}}\right), \quad (1.1)$$

as $n \rightarrow \infty$, where the coefficient k_m can be explicitly found in terms of m .

The goal of the present paper is to obtain an asymptotic formula similar to (1.1) for the integrals

$$\int_x^1 P_n^{(\alpha,\beta)}(t) P_m^{(\alpha,\beta)}(t) (1-t)^\alpha (1+t)^\beta dt,$$

as $n \rightarrow \infty$, where x, m are fixed, and $P_n^{(\alpha,\beta)}(x)$ are Jacobi polynomials. First, we recast a useful formula by Byerly in the case of Legendre polynomials and then we will derive the desired formula.

2. Legendre Polynomials

Recall that the Legendre polynomial P_n of degree n is a polynomial solution to the second order differential equation

$$(1-x^2)y''(x) - 2xy'(x) + n(n+1)y(x) = 0. \quad (2.1)$$

Actually, it is not so hard to check that for every nonnegative integer n equation (2.1) has a unique polynomial solution up to a multiplicative constant. To be definite, throughout this section we assume that P_n is a monic polynomial.

In his book published in 1893, W.E. Byerly noted the following relation for Legendre polynomials that showed itself to be useful in a few instances where Legendre polynomials appear.

Proposition 2.1 (See p. 172 in [2]). *Let n and m be nonnegative integers such that $n \neq m$. Then we have that*

$$\int_x^1 P_n(t) P_m(t) dt = \frac{(1-x^2) [P_m(x)P'_n(x) - P_n(x)P'_m(x)]}{n(n+1) - m(m+1)}, \quad (2.2)$$

where x is any real number.

Proof. Evidently, the polynomials P_n and P_m satisfy the differential equations

$$\frac{d}{dt} \left[(1-t^2) \frac{dP_n(t)}{dt} \right] = -n(n+1)P_n(t) \quad (2.3)$$

and

$$\frac{d}{dt} \left[(1-t^2) \frac{dP_m(t)}{dt} \right] = -m(m+1)P_m(t). \quad (2.4)$$

Multiplying (2.3) by P_m and (2.4) by P_n , respectively, and then subtracting them and integrating the result gives

$$\begin{aligned} & [n(n+1) - m(m+1)] \int_x^1 P_n(t) P_m(t) dt \\ &= \int_x^1 P_n(t) \frac{d}{dt} \left[(1-t^2) \frac{dP_m(t)}{dt} \right] dt - \int_x^1 P_m(t) \frac{d}{dt} \left[(1-t^2) \frac{dP_n(t)}{dt} \right] dt. \end{aligned} \quad (2.5)$$

After integrating each of the integrals on the right-hand side of (2.5) by parts we arrive at (2.2). \square

Formula (2.2) immediately implies the orthogonality of Legendre polynomials.

Corollary 2.2. *The polynomials P_n and P_m are orthogonal with respect to the Lebesgue measure on $[-1, 1]$ provided that $n \neq m$, that is,*

$$\int_{-1}^1 P_n(t) P_m(t) dt = 0.$$

Proof. It directly follows from (2.2) if we set $x = -1$. \square

Another aspect of formula (2.2) is that it shows a relation between integrals of products of orthogonal polynomials and the “augmented Wronskian” introduced by Karlin and Szegő, who were motivated by some probabilistic problems [6]. Namely, they studied the polynomials

$$\varphi_n(m; x) = \begin{vmatrix} P_m(x) & P_n(x) \\ P'_m(x) & P'_n(x) \end{vmatrix}.$$

In these notations, formula (2.2) reads that

$$\varphi_n(m; x) = \frac{n(n+1) - m(m+1)}{1-x^2} \int_x^1 P_n(t) P_m(t) dt$$

which gives an integral representation of polynomials $\varphi_n(m; x)$. In addition, Karlin and Szegő showed that the polynomials $\varphi_n(m; x)$ satisfy a second order differential equation [6, Chapter 4]. At the same time, in the literature on integrable systems the family of polynomials $\varphi_n(m; x)$ is called the Darboux transformation of the system $P_n(x)$ (for instance, see [7]) and is known to produce solvable equations if the original equation is solvable.

3. Jacobi Polynomials

In this section we will apply Byerly’s idea to the Jacobi differential equation. To this end, let $P_n^{(\alpha, \beta)}(x)$ be a polynomial solution of the Jacobi differential equation

$$(1-x^2) \frac{d^2y}{dx^2} + (\beta - \alpha - (\alpha + \beta + 2)x) \frac{dy}{dx} + n(n + \alpha + \beta + 1)y = 0, \quad (3.1)$$

where we assume that $\alpha, \beta > -1$. It is clear that such a definition determines a polynomial up to a constant factor. So, to be more precise, one can check that

$$P_n^{(\alpha, \beta)}(x) = \frac{(\alpha+1)_n}{n!} \sum_{k=0}^n \frac{(-n)_k (\alpha + \beta + n + 1)_k}{(\alpha+1)_k} \left(\frac{1-x}{2} \right)^k \quad (3.2)$$

is a solution to (3.1), where $(a)_k$ is the Pochhammer symbol, that is, $(a)_0 = 1$ and $(a)_k = a(a+1)\dots(a+k-1)$ for $k = 1, 2, 3, \dots$. Besides, formula (3.2) defines the Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ uniquely.

Next, following Karlin and Szegő [6] let us introduce the “augmented Wronskian” of the form

$$\varphi_n(m; \alpha, \beta, x) = \begin{vmatrix} P_m^{(\alpha, \beta)}(x) & P_n^{(\alpha, \beta)}(x) \\ \frac{dP_m^{(\alpha, \beta)}(x)}{dx} & \frac{dP_n^{(\alpha, \beta)}(x)}{dx} \end{vmatrix}.$$

Remark 3.1. The polynomials $\varphi_n(m; \alpha, \beta, x)$ are in fact the Darboux transform of $P_n^{(\alpha, \beta)}$ (for the definition of the Darboux transform see [7]). Also, the polynomials $\varphi_n(m; \alpha, \beta, x)$ are a particular case of the generalized Jacobi polynomials introduced in [1].

Theorem 3.2. *Let $\alpha, \beta > -1$. Then for any nonnegative integers n and m such that $n \neq m$, we have*

$$\int_x^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt = \frac{w(x)(1-x^2)\varphi_n(m; \alpha, \beta, x)}{n(n+\alpha+\beta+1) - m(m+\alpha+\beta+1)}, \quad (3.3)$$

where $w(t) = (1-t)^\alpha(1+t)^\beta$ and x is any real number.

Proof. By definition, the polynomials $P_n^{(\alpha, \beta)}$ and $P_m^{(\alpha, \beta)}$ satisfy the differential equations

$$(1-t^2) \frac{d^2 P_n^{(\alpha, \beta)}(t)}{dt^2} + (\beta - \alpha - (\alpha + \beta + 2)t) \frac{dP_n^{(\alpha, \beta)}(t)}{dt} + n(n + \alpha + \beta + 1) P_n^{(\alpha, \beta)}(t) = 0 \quad (3.4)$$

and

$$(1-t^2) \frac{d^2 P_m^{(\alpha, \beta)}(t)}{dt^2} + (\beta - \alpha - (\alpha + \beta + 2)t) \frac{dP_m^{(\alpha, \beta)}(t)}{dt} + m(m + \alpha + \beta + 1) P_m^{(\alpha, \beta)}(t) = 0. \quad (3.5)$$

We multiply (3.4) by $P_m^{(\alpha, \beta)}$ and (3.5) by $P_n^{(\alpha, \beta)}$, respectively, and then subtract the results to get a differential equation for $P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t)$. After using the product rule on $\varphi_n(m; \alpha, \beta, x)$ and rewriting the differential equation becomes

$$-CP_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) = (1-t^2)\varphi'_n(m; \alpha, \beta, t) + (\beta - \alpha - (\alpha + \beta + 2)t)\varphi_n(m; \alpha, \beta, t), \quad (3.6)$$

where $C = n(n + \alpha + \beta + 1) - m(m + \alpha + \beta + 1)$. We now multiply (3.6) by $w(t)$ and integrate over $[x, 1]$ to obtain

$$\begin{aligned} -C \int_x^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt &= \int_x^1 (1-t^2)\varphi'_n(m; \alpha, \beta, t) w(t) dt \\ &\quad + \int_x^1 (\beta - \alpha - (\alpha + \beta + 2)t)\varphi_n(m; \alpha, \beta, t) w(t) dt. \end{aligned}$$

After handling the left integral on the right-hand side by parts, we are left with

$$\begin{aligned} -C \int_x^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt &= -(1-x^2) w(x) \varphi_n(m; \alpha, \beta, x) \\ &- \int_x^1 \varphi_n(m; \alpha, \beta, t) (1-t^2) w'(t) dt + \int_x^1 (\beta - \alpha - (\alpha + \beta)t) \varphi_n(m; \alpha, \beta, t) w(t) dt. \end{aligned} \quad (3.7)$$

Then one can check that

$$\int_x^1 \varphi_n(m; \alpha, \beta, t) (1-t^2) w'(t) dt = \int_x^1 (\beta - \alpha - (\alpha + \beta)t) \varphi_n(m; \alpha, \beta, t) w(t) dt,$$

which completes the proof. \square

As in the case of Legendre polynomials, formula (3.3) immediately leads to orthogonality.

Corollary 3.3. *Let $\alpha, \beta > -1$. Then the polynomials $P_n^{(\alpha, \beta)}$ and $P_m^{(\alpha, \beta)}$ are orthogonal with respect to the weight $w(x) = (1-x)^\alpha (1+x)^\beta$ on $[-1, 1]$ provided that $n \neq m$, that is,*

$$\int_{-1}^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) (1-t)^\alpha (1+t)^\beta dx = 0.$$

The following result is another consequence of formula (3.3), and a well-known form of this result will be used in the next section.

Corollary 3.4. *Let $\alpha, \beta > -1$ and let $w(t) = (1-t)^\alpha (1+t)^\beta$. Then for any $n, m \geq 1$ and $n \neq m$, we have*

$$\int_{-1}^1 \left(\int_x^1 P_n^{(\alpha, \beta)}(t) w(t) dt \int_x^1 P_m^{(\alpha, \beta)}(t) w(t) dt \right) \frac{dx}{(1-x^2) w(t)} = 0. \quad (3.8)$$

Remark 3.5. It follows from (3.3) that

$$\int_x^1 P_n^{(\alpha, \beta)}(t) w(t) dt = \frac{(1-x^2) w(x) \varphi_n(0; \alpha, \beta, x)}{n(n+\alpha+\beta+1)}$$

and thus (3.8) can be rewritten in the following manner

$$\int_{-1}^1 \varphi_n(0; \alpha, \beta, t) \varphi_m(0; \alpha, \beta, t) (1-t^2) w(t) dt = 0, \quad (3.9)$$

which in particular shows that the integral from (3.8) exists.

Proof. To apply formula (3.3), note that $n > 0$ and $P_0^{(\alpha, \beta)} \equiv 1$ to get the identity

$$\int_x^1 P_n^{(\alpha, \beta)}(t) w(t) dt = \frac{(1-x^2) w(x) (P_n^{(\alpha, \beta)}(x))'}{n(n+\alpha+\beta+1)}. \quad (3.10)$$

Substituting this expression into (3.8) yields a cancellation, leaving

$$\frac{1}{n(n+\alpha+\beta+1)} \int_{-1}^1 \left((P_n^{(\alpha, \beta)}(x))' \int_x^1 P_m^{(\alpha, \beta)}(t) w(t) dt \right) dx.$$

After integrating by parts and using the fundamental theorem of calculus on the term

$$\int_x^1 P_m^{(\alpha, \beta)}(t) w(t) dt,$$

the integral in (3.8) becomes (up to a constant factor which is not important since we have 0 on the right-hand side)

$$\left(P_n^{(\alpha, \beta)}(x) \int_x^1 P_m^{(\alpha, \beta)}(t) w(t) dt \right) \Big|_{x=-1}^{x=1} + \int_{-1}^1 P_n^{(\alpha, \beta)}(x) P_m^{(\alpha, \beta)}(x) w(x) dx = 0.$$

Observe that $\int_{-1}^1 P_m^{(\alpha, \beta)}(t) w(t) dt = 0$ and $\int_{-1}^1 P_n^{(\alpha, \beta)}(x) P_m^{(\alpha, \beta)}(x) w(x) dx = 0$ using Corollary 3.3. \square

Remark 3.6. Since

$$\varphi_n(0; \alpha, \beta, x) = (P_n^{(\alpha, \beta)}(x))'$$

formula (3.9) implies the following well-known formula

$$\frac{d}{dx} \left[P_n^{(\alpha, \beta)}(x) \right] = \left(\frac{n + \alpha + \beta + 1}{2} \right) P_{n-1}^{(\alpha+1, \beta+1)}(x). \quad (3.11)$$

Here we are taking into account the normalization given by (3.2).

4. Asymptotic Formula for the Integrals in Question

In this section we will obtain an asymptotic formula for the integral

$$\int_x^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt$$

when x and m are fixed and $n \rightarrow \infty$. To begin with, let us recall a classical asymptotic result for Jacobi polynomials.

Proposition 4.1 ([5]). *Let $\alpha, \beta > -1$ and set*

$$N = n + \frac{\alpha + \beta + 1}{2}, \quad \gamma = -\frac{(\alpha + 1/2)\pi}{2}.$$

Then for a fixed $\theta \in (0, \pi)$, we have

$$P_n^{(\alpha, \beta)}(\cos \theta) = \frac{k(\theta)}{\sqrt{n}} \cos(N\theta + \gamma) + O(n^{-3/2}), \quad (4.1)$$

as $n \rightarrow \infty$, where

$$k(\theta) = \frac{1}{\sqrt{\pi}} [\sin(\theta/2)]^{-\alpha-1/2} [\cos(\theta/2)]^{-\beta-1/2}. \quad (4.2)$$

To apply the above statement to our integrals, note that combining (3.3) and (3.11) gives

$$\begin{aligned} & \frac{n(n + \alpha + \beta + 1) - m(m + \alpha + \beta + 1)}{(1 - x^2)w(x)} \int_x^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt \\ &= \frac{n + \alpha + \beta + 1}{2} P_{n-1}^{(\alpha+1, \beta+1)}(x) P_m^{(\alpha, \beta)}(x) - \frac{m + \alpha + \beta + 1}{2} P_{m-1}^{(\alpha+1, \beta+1)}(x) P_n^{(\alpha, \beta)}(x). \end{aligned} \quad (4.3)$$

Then the latter formula and Proposition 4.1 lead to the following result.

Theorem 4.2. *Let α , β , N , and γ be as in Proposition 4.1. Suppose n and m are nonnegative integers, m is fixed, and $n \neq m$. Then for a fixed $\theta \in (0, \pi)$ we have*

$$\int_{\cos \theta}^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt = \frac{\ell(\theta)}{n^{3/2}} \sin(N\theta + \gamma) + O(n^{-5/2}) \quad (4.4)$$

as $n \rightarrow \infty$, where

$$\ell(\theta) = \frac{2^{\alpha+\beta+1}}{\sqrt{\pi}} [\sin(\theta/2)]^{\alpha+1/2} [\cos(\theta/2)]^{\beta+1/2} P_m^{(\alpha, \beta)}(\cos \theta).$$

Proof. For convenience, let us set

$$I_n = \int_{\cos \theta}^1 P_n^{(\alpha, \beta)}(t) P_m^{(\alpha, \beta)}(t) w(t) dt.$$

Next, applying Proposition 4.1 to $P_{n-1}^{(\alpha+1, \beta+1)}$ and $P_n^{(\alpha, \beta)}$, we arrive at

$$P_{n-1}^{(\alpha+1, \beta+1)}(x) = \frac{[\sin(\theta/2)]^{-\alpha-3/2} [\cos(\theta/2)]^{-\beta-3/2}}{\sqrt{(n-1)\pi}} \cos(N\theta + \gamma - \pi/2) + O(n^{-3/2}) \quad (4.5)$$

and

$$P_n^{(\alpha, \beta)}(x) = \frac{[\sin(\theta/2)]^{-\alpha-1/2} [\cos(\theta/2)]^{-\beta-1/2}}{\sqrt{n\pi}} \cos(N\theta + \gamma) + O(n^{-3/2}), \quad (4.6)$$

as $n \rightarrow \infty$ and where we set $x = \cos \theta$. Observe that $\cos(N\theta + \gamma - \pi/2) = \sin(N\theta + \gamma)$ and so (4.5) takes the form

$$P_{n-1}^{(\alpha+1, \beta+1)}(x) = \frac{[\sin(\theta/2)]^{-\alpha-3/2} [\cos(\theta/2)]^{-\beta-3/2}}{\sqrt{(n-1)\pi}} \sin(N\theta + \gamma) + O(n^{-3/2}). \quad (4.7)$$

Now, combining (4.6) and (4.7) with (4.3), we get

$$\begin{aligned} & \frac{(n(n+\alpha+\beta+1) - m(m+\alpha+\beta+1))I_n}{(1-x^2)w(x)} [\sin(\theta/2)]^{\alpha+3/2} [\cos(\theta/2)]^{\beta+3/2} \\ &= \frac{n+\alpha+\beta+1}{2\sqrt{(n-1)\pi}} P_m^{(\alpha, \beta)}(x) \sin(N\theta + \gamma) + O(n^{-1/2}). \end{aligned} \quad (4.8)$$

Then since $x = \cos \theta$, the weight function can be rewritten in the following way

$$w(x) = (1 - \cos \theta)^\alpha (1 + \cos \theta)^\beta = 2^{\alpha+\beta} [\sin(\theta/2)]^{2\alpha} [\cos(\theta/2)]^{2\beta}.$$

Also, we have $1 - x^2 = \sin^2 \theta = 4[\sin(\theta/2)]^2 [\cos(\theta/2)]^2$. As a result, (4.8) reduces to

$$\begin{aligned} & \frac{(n(n+\alpha+\beta+1) - m(m+\alpha+\beta+1))I_n}{2^{\alpha+\beta+2} [\sin(\theta/2)]^{\alpha+1/2} [\cos(\theta/2)]^{\beta+1/2}} \\ &= \frac{\sqrt{n}}{2\sqrt{\pi}} P_m^{(\alpha, \beta)}(x) \sin(N\theta + \gamma) + O(n^{-1/2}), \end{aligned}$$

which yields (4.4). \square

Acknowledgment. M.D. was supported by the NSF DMS grant 2008844 and by the University of Connecticut Research Excellence Program.

References

1. Bonneux, N.: Exceptional Jacobi polynomials, *J. Approx. Theory* **239** (2019), 72–112.
2. Byerly, W. E.: *An elementary treatise on Fourier's series and spherical, cylindrical, and ellipsoidal harmonics, with applications to problems in mathematical physics*, Dover Publications Inc., New York, 1959.
3. Derevyagin, M.; Geronimo, J. S.: The entries of a refinement equation and a generalization of the discrete wave equation, arXiv:2001.10650.
4. Geronimo, J.S.; Marcellán, F.: On Alpert multiwavelets, *Proc. Amer. Math. Soc.* **143** (2015), no. 6, 2479–2494.
5. Ismail, M. E. H.: *Classical and quantum orthogonal polynomials in one variable*, Cambridge University Press, Cambridge, 2009.
6. Karlin, S.; Szegő, G.: On certain determinants whose elements are orthogonal polynomials, *J. Analyse Math.* **8** (1960/61), 1–157.
7. Matveev, V. B.; Salle, M. A.: *Darboux transformations and solitons*, Springer Series in Nonlinear Dynamics, Springer-Verlag, Berlin, 1991.

M. DEREVYAGIN: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-1009, USA

E-mail address: maksym.derevyagin@uconn.edu

NICHOLAS JURICIC: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CT 06269-1009, USA

E-mail address: nicholas.juricic@uconn.edu