

Physical Violations of the Bulk-Edge Correspondence in Topological Electromagnetics

S. Ali Hassani Gangaraj^{*} and Francesco Monticone[†]*School of Electrical and Computer Engineering, Cornell University, Ithaca, New York 14853, USA*

(Received 4 November 2019; accepted 25 March 2020; published 13 April 2020)

In this Letter, we discuss two general classes of apparent violations of the bulk-edge correspondence principle for continuous topological photonic materials, associated with the asymptotic behavior of the surface modes for diverging wave numbers. Considering a nonreciprocal plasma as a model system, we show that the inclusion of spatial dispersion (e.g., hydrodynamic nonlocality) formally restores the bulk-edge correspondence by avoiding an unphysical response at large wave numbers. Most importantly, however, our findings show that, for the considered cases, the correspondence principle is *physically* violated for all practical purposes, as a result of the unavoidable attenuation of highly confined modes even if all materials are assumed perfect, with zero intrinsic bulk losses, due to confinement-induced Landau damping or nonlocality-induced radiation leakage. Our work helps clarifying the subtle and rich topological wave physics of continuous media.

DOI: 10.1103/PhysRevLett.124.153901

Introduction.—The bulk-edge correspondence is a widely used principle of topological wave physics, which allows determining the number of unidirectional edge modes from the topological properties of the bulk modes [1–9]. In topological photonic insulators with broken time-reversal symmetry (nonreciprocal), the relevant topological invariant is the gap Chern number, i.e., the sum of the Chern numbers of all bulk modes below the bulk-mode bandgap, $C_{\text{gap}} = \sum_i C_i$. The bulk-edge correspondence then states that the difference in gap Chern numbers between the materials forming an interface is equal to the net number of unidirectional surface modes propagating along the interface [7–11]. While this principle works well for topological photonic insulators based on periodic structures, difficulties arise in the case of continuous topological materials with no intrinsic periodicity, due to the absence of a finite Brillouin zone, which may lead to an ill-behaved response for diverging wave numbers if spatial dispersion is not included [12–16]. A proof of the bulk-edge correspondence principle in topological photonics has been recently published [17], which does require the inclusion of spatial dispersion (nonlocality), but hinges on the assumption of an ideally dissipationless structure. As discussed in the following, however, even if all materials are assumed lossless (i.e., with no bulk damping), dissipation channels may still be present in a physical scenario. Thus, questions still remain regarding the possibility of breaking the bulk-edge correspondence in a physical system.

In this Letter, we present and discuss two general classes of violations of the bulk-edge correspondence in continuous topological systems, and we carefully assess the role of spatial dispersion and dissipation. These two classes are illustrated in Fig. 1: (i) The number of unidirectional edge

modes is inconsistent with the gap Chern number difference between the two materials at the interface [Fig. 1(a)]; (ii) The dispersion curve of the unidirectional edge mode does not span the entire bulk-mode band gap, due to a flat asymptotic dispersion of the mode at a certain frequency within the gap [Fig. 1(b)]. Examples of these violations are easily realized using the simplest possible continuous topological photonic insulator, namely, a nonreciprocal (magnetized) plasma. Indeed, it has been known for decades that, at the interface between a magnetized plasma and a conductor, unidirectional transverse-magnetic (TM)




FIG. 1. Apparent violations of the bulk-edge correspondence. (a),(b) Illustration of the typical dispersion diagram for the two classes of violations studied here. Solid red curves indicate the surface modes, and solid black curves denote the bulk modes along the x direction, delimiting the region of the projected bulk bands (light blue areas). Gray areas highlight the bulk-mode band gap. (c) Configurations that exhibit class-I violations of the bulk-edge correspondence. The dispersion diagram in (a) corresponds to the geometry on the left. (d) Configuration that implements a class-II violation.

surface waves appear within the common bulk-mode band gap [18,19]. Such surface waves are an example of unidirectional surface plasmon polaritons (SPPs) [15]. Recently, it has been shown that their unidirectional nature is indeed rooted in the different topological properties of the trivial conductor and the biased plasma [12,20–28]. Specifically, considering a plasma magnetized along the $\pm z$ axis, the TM bulk modes in the plane orthogonal to the bias exhibit nontrivial topological properties, yielding a gap Chern number $\mathcal{C}_{\text{gap}} = \mp 1$. Here, we use this model system to study physical configurations that provide apparent counterexamples to the bulk-edge correspondence.

Class-I violations.—To realize the simplest possible class-I violation, consider an interface between a biased plasma and a perfect magnetic conductor (PMC), a configuration considered for example in [15,19] and illustrated in Fig. 1(c, left). Since a PMC is a topologically trivial insulator, the difference in gap Chern numbers is $\Delta\mathcal{C}_{\text{gap}} = +1$. Thus, according to the bulk-edge correspondence principle, one topological TM surface mode should emerge across the bulk-mode band gap. However, no TM surface mode is allowed on this interface, because the PMC boundary “short-circuits” the magnetic field [29]. This absence of a surface mode within the band gap directly violates the predictions of the bulk-edge correspondence. One might argue that, since PMCs are not natural materials, but have to be realized in the form of metasurfaces [30], this may be a rather artificial scenario. However, a closely related configuration that does not involve a PMC wall, but still violates the bulk-edge correspondence, is an interface between two oppositely-biased plasmas [Fig. 1(c, right)]. Indeed, as we recently showed in [31], the difference in gap Chern numbers is now $\Delta\mathcal{C}_{\text{gap}} = +2$, so one would expect two surface states, but only one appears, a mode with even-symmetric magnetic field distribution with respect to the interface. Interestingly, due to the nature of a magnetic mirror, the surface modes of the PMC-terminated structure are identical to the odd-symmetric surface modes of the configuration with oppositely-biased plasmas. This suggests that the origin of the bulk-edge-correspondence violation and the nature of the missing mode are the same in both cases.

To elucidate the physical mechanism behind these violations, we consider again the configuration in Fig. 1(c, left), but we now include a thin spacer layer, filled with an isotropic material with permittivity ϵ_m , between the magnetized plasma and the PMC boundary, as shown in the inset of Fig. 2(a). The following analysis concerns TM modes, with $e^{-i\omega t}$ time-harmonic dependence. As usually done, the magnetized plasma can be modeled by a gyrotropic permittivity tensor $\epsilon = \epsilon_0[\epsilon_{11}\mathbf{I}_t + \epsilon_{33}\hat{\mathbf{z}}\hat{\mathbf{z}} - i\epsilon_{12}\hat{\mathbf{z}} \times \mathbf{I}]$, where $\mathbf{I}_t = \mathbf{I} - \hat{\mathbf{z}}\hat{\mathbf{z}}$. ϵ_{11} , ϵ_{12} , and ϵ_{33} follow a standard magnetized Drude model (see, e.g., [32]). As a realistic example, we consider the case of a magnetized n -doped semiconductor in the low THz regime, namely, n -type InSb, a material that has been studied in several recent works [15,16,33–35]. A typical




FIG. 2. Class-I violations of the bulk-edge correspondence. (a) Trajectory of the surface-mode wave number, for the configuration in the inset, as the separation d is reduced to zero. Blue and red curves are the trajectories for the local and the nonlocal case ($\beta = 1.07 \times 10^6$ m/s), respectively, at a frequency within the bulk-mode band gap, $\omega/\omega_p = 1.05$. (b) Self-consistent iterative calculation of Landau damping for the configuration in (a) with $d = 0$. The inset shows the distribution (time snapshot) of the SPP electric field normal to the interface, with zero bulk damping and zero Landau damping (blue) and with the calculated Landau damping (red).

sample of this material has plasma frequency $\omega_p/2\pi = 2$ THz, electron density $N_e = 1.1 \times 10^{22}/\text{m}^3$, and bound-charge contribution to the permittivity function $\epsilon_\infty = 15.6$. We consider a moderate dc magnetic field of 0.2 T, which is sufficient to produce a moderately large cyclotron frequency, $\omega_c/\omega_p = 0.2$. We first consider a lossless and local scenario. Solving Maxwell’s equations with suitable boundary conditions, it is straightforward to show that the SPPs supported by the interface between the isotropic spacer and the biased plasma satisfy the following dispersion equation,

$$\frac{\epsilon_{11}k_x + \epsilon_{12}\alpha_p}{\alpha_p k_x - k_0^2 \epsilon_{12}} \cos(k_y d) - \frac{\epsilon_m k_y}{k_x^2 - k_0^2 \epsilon_m} \sin(k_y d) = 0, \quad (1)$$

where d is the isotropic spacer thickness, $k_0 = \omega/c$, $k_y = \sqrt{k_0^2 \epsilon_m - k_x^2}$, $\alpha_p = \sqrt{k_x^2 - k_0^2 \epsilon_{\text{eff}}}$, and $\epsilon_{\text{eff}} = (\epsilon_{11}^2 - \epsilon_{12}^2)/\epsilon_{11}$. The blue curve in Fig. 2(a) represents the trajectory of the SPP wave number, calculated with (1) at a frequency within the bulk-mode band gap, as the spacer thickness is reduced, which corresponds to transforming the geometry in Fig. 2(a) to the configuration in Fig. 1(c, left). From this plot, one can clearly see that, as $d \rightarrow 0$, the SPP wave number tends to infinity. Hence, this analysis shows that, although the PMC-plasma configuration in Fig. 1(c, left) does not seem to support any mode, one topological surface mode does exist in the asymptotic part of the spatial spectrum for $k \rightarrow +\infty$.

As mentioned above, an interface between a magnetized plasma and a PMC is closely related to the case of two oppositely-biased plasmas. We have recently studied this configuration in a different context [31], where we showed that the wave number of the supported odd-symmetric mode rapidly diverges, as the spacing between the two plasmas is reduced, fully consistent with the blue curve trajectory in Fig. 2(a). This asymptotic behavior is indeed

the reason for all class-I violations of the bulk-edge correspondence.

At this point, it is crucial to stress that, since we are dealing with a mode with very large wave number (in principle infinite), a simple material model that neglects dissipative and nonlocal effects, while mathematically consistent, is not physically accurate [36,37]. We start by assessing the impact of nonlocal effects, which make the permittivity a function of the wave vector. Specifically, here we consider a well-established hydrodynamic treatment of plasma nonlocalities [15,36]; however, we stress that our considerations are general since any model of nonlocality leads to qualitatively similar predictions [38–41]. Most importantly, nonlocal effects generically lead to a high spatial-frequency cutoff for the material response [12,36].

Using the hydrodynamic model, the induced free-electron currents \mathbf{J} are governed by the equation [36]:

$$\beta^2 \nabla(\nabla \cdot \mathbf{J}) + \omega(\omega + i\Gamma)\mathbf{J} = i\omega(\omega_p^2 \epsilon_0 \epsilon_\infty \mathbf{E} - \mathbf{J} \times \omega_c \hat{z}), \quad (2)$$

where β is the nonlocal parameter and Γ is the damping rate due to absorption losses. By simultaneously solving Maxwell's equations and the hydrodynamic equation in the magnetized plasma, the dispersion equation of the TM bulk modes can be found as $k^4 \beta^2 \omega^2 - k^2 [\Omega_b(\omega_p^2 + \beta^2 k_0^2 \epsilon_\infty) + \Omega_0] + k_0^2 \epsilon_\infty \Omega_0 = 0$, where $\Omega_b = \omega^2 - \omega_p^2$ and $\Omega_0 = \Omega_b^2 - \omega^2 \omega_c^2$. If $\beta \neq 0$, the first term of this equation leads to the emergence of an additional bulk mode with respect to the local case [15], which implies that an additional boundary condition should be included to solve for the fields in the presence of interfaces. A physically meaningful choice is $\mathbf{J} \cdot \hat{y} = 0$ [36], which forces the normal component of \mathbf{J} to vanish at the boundary. Following this approach, we can assess the impact of nonlocality on the surface-wave dispersion for the configuration in Fig. 1(c, left) as $d \rightarrow 0$. The solid red line in Fig. 2(a) shows the modified trajectory of the SPP wave number in the presence of nonlocality. While nonlocalities have little effect for large separations and small wave numbers, they become very important at larger values of k : in contrast to the local case, the SPP wave number does not diverge, but it stops at a finite value, $k \approx 116.6k_p$, for $d \rightarrow 0$. This result shows that an SPP with finite wave number is indeed supported by the PMC-plasma interface. Hence, the bulk-edge correspondence is formally restored if nonlocal effects are properly included in the material model.

Interestingly, the presence of this unidirectional surface mode at a PMC-biased-plasma interface (or between oppositely biased plasmas) has so far gone unnoticed (e.g., [15,19]). This is, however, not surprising. Indeed, even though the SPP wave number does not diverge in the nonlocal case, it is still very large for any realistic values of nonlocality [42]. This implies that, first, the SPP mode has very large wave impedance; therefore, it is very difficult to excite due to the large impedance mismatch with any

source or feeding mechanism. Most importantly, this mode is so highly confined to the interface that it becomes extremely sensitive to absorption. Thus, the topological mode is quickly damped in the presence of a loss channel of any type, including phonon and defect scattering, electron-electron interactions, and interface roughness (or even just the effective loss due to numerical error and mesh granularity in numerical simulations). Even in the absence of any scattering loss in the bulk of the considered materials, it is physically impossible for a surface plasmon-polariton to experience zero dissipation. Indeed, no matter the intrinsic quality of the materials and interfaces, the smallest possible loss is determined by surface-collision-induced Landau damping, i.e., by the direct excitation of electron-hole pairs in the plasma by the highly confined electric field on the interface. Following [44,45], we determine the loss rate due to Landau damping for the geometry in Fig. 1(c, left) as

$$\Gamma_L = \frac{3\pi\omega}{2} \frac{\int_1^\infty q^{-3} |F_y(q)|^2 dq}{\int_0^\infty (|F_x(q)|^2 + |F_y(q)|^2) dq}, \quad (3)$$

where $F_y(q)$ and $F_x(q)$ are the Fourier transforms of the electric-field components, normal and parallel to the interface, and $q = k/(\omega/v_F)$ is a normalized wave number, where v_F is the Fermi velocity, proportional to the nonlocal parameter $\beta^2 = v_F^2(\frac{3}{5}\omega + \frac{1}{3}i\Gamma)/(\omega + i\Gamma)$ [46,47]. To investigate the effect of Landau damping, we consider an ideal sample of InSb with zero intrinsic bulk loss, $\Gamma = 0$, and solve for Γ_L and the resulting SPP fields. The correct solution needs to be found in a self-consistent manner [48]. Figure 2(b) shows the iterative solution steps for Γ_L up to convergence. It can be seen that, even in the absence of bulk loss, a significant level of Landau damping is present, $\Gamma_L \approx 0.13\omega_p$, which represents the smallest possible physical level of loss for this configuration. Surface-induced Landau damping strongly affects the highly confined topological surface mode. The inset of Fig. 2(b) compares the SPP field distributions with and without Landau damping. The difference is striking: the surface mode dies out very quickly in the presence of Landau damping, over a distance of less than $0.03\lambda_p$ (λ_p is the free-space wavelength at ω_p).

We would like to summarize here the main message of this section: while the inclusion of nonlocal effects formally restores the bulk-edge correspondence in the considered lossless configurations, the correspondence principle is physically violated since, due to confinement-induced damping, the surface mode is attenuated almost immediately even if the considered materials are assumed perfect, with zero intrinsic bulk losses.

Class-II violations.—To realize a class-II violation of the bulk-edge correspondence, as illustrated in Fig. 1(b), we consider an interface between a magnetized plasma and a trivial opaque medium, as shown in Fig. 1(d). A large body of work [12,15–19,21–23,28,31] has shown that the




FIG. 3. Class-II violations of the bulk-edge correspondence. (a),(b) Dispersion diagrams (dashed blue lines) of the surface modes supported by the configuration in Fig. 1(d). The dispersion diagrams are also plotted as density plots of the inverse determinant of the boundary-condition matrix. The bright bands correspond to the SPP poles, with the band becoming blurred if the pole moves to a complex value of wave number. (a) Local and lossless case, with $\beta = \beta^m = 0$, $\omega_p^m = 1.2\omega_p$ and $\epsilon_\infty^m = \epsilon_\infty$. (b) Nonlocal and lossless case with $\beta = \beta^m = 1.07 \times 10^6$ m/s. The dashed white lines indicate the edges of the bulk-mode band gap. (c),(d) Spatial distributions (time snapshots) of the SPP electric field normal to the interface, for the local (c) and nonlocal (d) cases, calculated at the frequency indicated by the white dot in panels (a),(b).

bulk-edge correspondence appears to correctly predict the emergence of one topological surface state within the common bulk-mode band gap of the two materials, consistent with $\Delta\mathcal{C}_{\text{gap}} = +1$. If the opaque material is frequency dispersive, following, for instance, a classical Drude dispersion with plasma frequency ω_p^m , the interface supports a surface-plasmon resonance at a frequency $\omega_{\text{SP}} < \omega_p^m$ at which $\epsilon_m = -(\epsilon_{11} \pm \epsilon_{12})$. Approaching this frequency, the surface-mode band flattens out and tends to infinite wave number, as recognized in [15]. Surface-wave propagation is not allowed at frequencies $\omega_{\text{SP}} < \omega < \omega_p^m$. Thus, if we tune ω_p^m such that ω_{SP} falls within the bulk-mode band gap of the magnetized plasma, as illustrated in Fig. 1(b), we obtain a common band gap for the two materials whose lower-frequency portion supports exactly one topological surface mode, whereas no surface mode can propagate in the higher-frequency portion of the band gap.

As a realistic example of this scenario, we consider an interface between magnetized n -type InSb (same parameters as above) and a dispersive isotropic metal with parameters given in the caption of Fig. 3. We first consider a local scenario, and we choose the metal plasma frequency close to the upper edge of the InSb bulk-mode band gap, e.g., $\omega_p^m = 1.2\omega_p$. Figure 3(a) shows the dispersion curve

of the unidirectional surface mode supported by this configuration, revealing its flat asymptotic dispersion within the band gap. This behavior indeed violates the bulk-edge correspondence, which would predict the presence of one unidirectional surface mode spanning the entire band gap. As for class-I violations, this form of violation is due to the asymptotic behavior of the surface mode for large wave numbers, which suggests that nonlocality may again restore the correspondence. To verify this, we include hydrodynamic nonlocalities both in the metal and the magnetized plasma, following the approach discussed in [15,49].

If nonlocal effects are properly included, the nonphysical flat asymptotes in the modal dispersion disappear, as discussed in [15,16] for different scenarios. Indeed, as shown in Fig. 3(b), in the nonlocal case under consideration, the dispersion curve of the unidirectional SPP bends upward and monotonically grows with frequency, spanning the entire band gap. This implies that, for each frequency within the gap, there exists exactly one unidirectional surface mode, which is now consistent with the bulk-edge correspondence. Most importantly, however, we also note that the dispersion band in Fig. 3(b) becomes significantly “blurred” at higher frequencies, which is a direct indication of the migration of the dispersion-equation root (pole of the system’s Green function) into the complex wave number plane. Since we assumed that all the materials are lossless, a complex modal wave number indicates *radiation loss*: the bound surface mode has become a “leaky mode” that gradually loses energy as radiation [50]. This form of nonlocality-induced radiation leakage was originally predicted in [49] and recently discussed in [15], where it was shown that the additional nonlocality-induced bulk mode of the magnetized plasma provides a continuum of radiation modes, propagating at different angles, which the surface mode can couple to, thus leaking energy into the bulk. Figures 3(c) and 3(d) compare the field distribution of the SPP mode, in the local and nonlocal cases, at a frequency near the middle of the band gap: in the nonlocal case, the unidirectional mode is quickly attenuated over a distance of less than $0.5\lambda_p$ due to radiation leakage. Much higher attenuation is obtained at higher frequencies. We also note that, because of such a strong attenuation, the degree of confinement of the SPP mode is low (especially compared to the SPPs in Fig. 2); therefore, the effect of Landau damping in this configuration is negligible compared to radiation damping. However, the main finding of this analysis is essentially the same as in the previous section: while the inclusion of nonlocal effects formally restores the bulk-edge correspondence, the correspondence principle is violated for all practical purposes because the surface mode is attenuated almost immediately even if the considered materials are assumed lossless.

Discussion and conclusion.—Both classes of apparent violations discussed here arise due to the asymptotic behavior of the photonic bands for diverging wave

numbers. This is possible only for continuous media, with an infinite Brillouin zone. We note that the presence of topological surface modes in the asymptotic part of the spatial spectrum has also been recently recognized in [14], based on a different approach. Topological photonic insulators based on periodic structures, e.g., [7,51], are not expected to exhibit the bulk-edge-correspondence violations discussed here. For continuous media, we have shown that the correspondence principle is formally restored if nonlocal effects are included, consistent with the correct definition of Chern number in topological continua, which requires a high spatial-frequency cutoff as demonstrated in [12,13]. Most importantly, however, we have shown that, even if the considered materials are assumed perfect, with zero bulk losses, the “recovered” surface mode that would satisfy the bulk-edge correspondence is very strongly attenuated due to either confinement-induced Landau damping, or nonlocality-induced radiation leakage. We, therefore, conclude that, in the considered cases, the bulk-edge correspondence principle is violated for all practical purposes.

Our findings also help clarify the behavior of some extreme nonreciprocal configurations, such as terminated one-way waveguides. In a junction between two structures, one supporting one or more unidirectional modes, and one that does not support any mode (as in the case of an interface between a magnetized plasma and a metal terminated by an orthogonal PMC wall, forming the T-like junction studied in several recent papers [15,16,23,34,52–54]), the energy incident on the junction will not be able to escape. We argue, however, that this situation does not pose any physical problem (the energy does not build up indefinitely at the junction or termination), even assuming media with vanishing losses, since the incident wave will be dissipated into a so-called “wedge mode” while the wavelength shrinks to zero, as discussed in [15,35], and as a result of Landau damping or radiation leakage, as discussed here.

To conclude, our work identifies, for the first time, extreme configurations in which the bulk-edge correspondence principle is physically violated. Our findings are expected to help making more accurate predictions in topological systems involving uniform media.

F.M. and S.A.H.G. acknowledge support from the National Science Foundation (NSF) with Grant No. 1741694, and the Air Force Office of Scientific Research with Grant No. FA9550-19-1-0043.

*ali.gangaraj@gmail.com

†francesco.monticone@cornell.edu

[1] Y. Hatsugai, Chern Number and Edge States in the Integer Quantum Hall Effect, *Phys. Rev. Lett.* **71**, 3697 (1993).

[2] S. A. Skirlo, L. Lu, and M. Soljacic, Multimode One-Way Waveguides of Large Chern Numbers, *Phys. Rev. Lett.* **113**, 113904 (2014).

[3] S. A. Skirlo, L. Lu, Y. Igarashi, Q. Yan, J. Joannopoulos, and M. Soljacic, Experimental Observation of Large Chern Numbers in Photonic Crystals, *Phys. Rev. Lett.* **115**, 253901 (2015).

[4] W. Gao, M. Lawrence, B. Yang, F. Liu, F. Fang, B. Bri, J. Li, and S. Zhang, Topological Photonic Phase in Chiral Hyperbolic Metamaterials, *Phys. Rev. Lett.* **114**, 037402 (2015).

[5] F. D. M. Haldane and S. Raghu, Possible Realization of Directional Optical Waveguides in Photonic Crystals with Broken Time-Reversal Symmetry, *Phys. Rev. Lett.* **100**, 013904 (2008).

[6] S. Raghu and F. D. M. Haldane, Analogs of quantum-Hall effect edge states in photonic crystals, *Phys. Rev. A* **78**, 033834 (2008).

[7] L. Lu, J. D. Joannopoulos, and M. Soljacic, Topological photonics, *Nat. Photonics* **8**, 821 (2014).

[8] S. Q. Shen, *Topological Insulators: Dirac Equation in Condensed Matters* (Springer, Berlin, 2012).

[9] B. A. Bernevig and T. L. Hughes, *Topological Insulators and Topological Superconductors* (Princeton University Press, Princeton, NJ, 2013).

[10] K. Y. Bliokh, D. Leykam, M. Lein, and F. Nori, Topological non-Hermitian origin of surface Maxwell waves, *Nat. Commun.* **10**, 580 (2019).

[11] S. A. R. Horsley, Topology and the optical Dirac equation, *Phys. Rev. A* **98**, 043837 (2018).

[12] M. G. Silveirinha, Chern invariants for continuous media, *Phys. Rev. B* **92**, 125153 (2015).

[13] M. G. Silveirinha, Bulk-edge correspondence for topological photonic continua, *Phys. Rev. B* **94**, 205105 (2016).

[14] C. Tauber, P. Delplace, and A. Venaille, Anomalous bulk-edge correspondence in continuous media, *Phys. Rev. Research* **2**, 013147 (2020).

[15] S. A. Hassani Gangaraj and F. Monticone, Do truly unidirectional surface plasmon-polaritons exist?, *Optica* **6**, 1158 (2019).

[16] S. Buddhiraju, Y. Shi, A. Song, C. Wojcik, M. Minkov, I. A. D. Williamson, A. Dutt, and S. Fan, Absence of unidirectionally propagating surface plasmon-polaritons at nonreciprocal metal-dielectric interfaces, *Nat. Commun.* **11**, 674 (2020).

[17] M. G. Silveirinha, Proof of the Bulk-Edge Correspondence Through a Link Between Topological Photonics and Fluctuation-Electrodynamics, *Phys. Rev. X* **9**, 011037 (2019).

[18] S. R. Seshadri, Excitation of surface waves on a perfectly conducting screen covered with anisotropic plasma, *IRE Trans. Microwave Theory Tech.* **10**, 573 (1962).

[19] A. Ishimaru, Unidirectional waves in anisotropic media and the resolution of the thermodynamic paradox, Technical Reports, Air Force Cambridge Research Laboratories, Bedford, Massachusetts, 1962.

[20] D. Jin, L. Lu, Z. Wang, C. Fang, J. D. Joannopoulos, M. Soljacic, L. Fu, and N. X. Fang, Topological magnetoplasmons, *Nat. Commun.* **7**, 13486 (2016).

- [21] L. Shen, Y. You, and X. Deng, Backscattering-immune one-way surfacemagnetoplasmons at terahertz frequencies, *Opt. Express* **23**, 950 (2015).
- [22] Z. Yu, G. Veronis, Z. Wang, and S. Fan, One-Way Electromagnetic Waveguide Formed at the Interface Between a Plasmonic Metal Under a Static Magnetic Field and a Photonic Crystal, *Phys. Rev. Lett.* **100**, 023902 (2008).
- [23] D. E. Fernandes and M. G. Silveirinha, Topological Origin of Electromagnetic Energy Sinks, *Phys. Rev. Applied* **12**, 014021 (2019).
- [24] D. Jin, T. Christensen, M. Soljacic, N. X. Fang, L. Lu, and X. Zhang, Infrared Topological Plasmons in Graphene, *Phys. Rev. Lett.* **118**, 245301 (2017).
- [25] S. A. Hassani Gangaraj, M. G. Silveirinha, and G. W. Hanson, Berry phase, Berry connection, and Chern number for a continuum bianisotropic material from a classical electromagnetics perspective, *IEEE J. Multiscale Multiphys. Comput. Tech.* **2**, 3 (2017).
- [26] S. A. Hassani Gangaraj, A. Nemilentsau, and G. W. Hanson, The effects of three-dimensional defects on one-way surface plasmon propagation for photonic topological insulators comprised of continuum media, *Sci. Rep.* **6**, 30055 (2016).
- [27] T. V. Mechelen and J. Zubin, Unidirectional Maxwellian spin waves, *Nanophotonics* **8**, 1399 (2019).
- [28] A. Davoyan and N. Engheta, Theory of wave propagation in magnetized near-zero-epsilon metamaterials: Evidence for One-Way Photonic States and Magnetically Switched Transparency and Opacity, *Phys. Rev. Lett.* **111**, 257401 (2013).
- [29] A PMC boundary, being the dual a perfect electric conductor, requires zero tangential magnetic field on its surface.
- [30] D. Sievenpiper, L. Zhang, R. F. J. Broas, N. G. Alexopolous, and E. Yablonovitch, High-impedance electromagnetic surfaces with a forbidden frequency band, *IEEE Trans. Microwave Theory Tech.* **47**, 2059 (1999).
- [31] S. A. Hassani Gangaraj and F. Monticone, Coupled topological surface modes in gyrotropic structures: Green's function analysis, *IEEE Antennas Wireless Propag. Lett.* **17**, 1993 (2018).
- [32] J. A. Bittencourt, *Fundamentals of Plasma Physics* (Springer-Verlag, New York, 2004), 3rd ed.
- [33] E. Palik, R. Kaplan, R. Gammon, H. Kaplan, R. Wallis, and J. Quinn, Coupled surface magnetoplasmon-optic-phonon polariton modes on InSb, *Phys. Rev. B* **13**, 2497 (1976).
- [34] K. Tsakmakidis, L. Shen, S. Schulz, X. X. Zheng, J. Upham, X. Deng, H. Altug, A. Vakakis, and R. Boyd, Breaking Lorentz reciprocity to over-come the time-bandwidth limit in physics and engineering, *Science* **356**, 1260 (2017).
- [35] S. A. Mann, D. L. Sounas, and A. Alu, Nonreciprocal cavities and the time-bandwidth limit, *Optica* **6**, 104 (2019).
- [36] S. Raza, S. I. Bozhevolnyi, M. Wubs, and N. A. Mortensen, Nonlocal optical response in metallic nanostructures, *J. Phys. Condens. Matter* **27**, 183204 (2015).
- [37] V. M. Agranovich and V. Ginzburg, *Crystal Optics with Spatial Dispersion, and Excitons*, Springer Series in Solid-State Sciences (Springer-Verlag, Berlin Heidelberg, 1984), 2nd ed.
- [38] C. Ciraci, R. T. Hill, J. J. Mock, Y. Urzhumov, A. I. Fernandez-Dominguez, S. A. Maier, J. B. Pendry, A. Chilkoti, and D. R. Smith, Probing the ultimate limits of plasmonic enhancement, *Science* **337**, 1072 (2012).
- [39] P. J. Feibelman, Microscopic calculation of surface-plasmon dispersion and damping, *Phys. Rev. B* **9**, 5077 (1974).
- [40] R. Esteban, A. G. Borisov, P. Nordlander, and J. Aizpurua, Bridging quantum and classical plasmonics with a quantum-corrected model, *Nat. Commun.* **3**, 825 (2012).
- [41] N. A. Mortensen, Nonlocal formalism for nanoplasmonics: Phenomenological and semi-classical considerations, *Photonics Nanostruct. Fundam. Appl.* **11**, 303 (2013).
- [42] InSb has one of the largest naturally available values of β due to its small effective mass [43].
- [43] J. R. Maack, N. A. Mortensen, and M. Wubs, Size-dependent nonlocal effects in plasmonic semiconductor particles, *Europhys. Lett.* **119**, 17003 (2017).
- [44] J. B. Khurgin, Hot carriers generated by plasmons: Where are they generated and where do they go from there?, *Faraday Discuss.* **214**, 35 (2019).
- [45] J. Khurgin, W. Y. Tsai, D. P. Tsai, and G. Sun, Landau damping and limit to field confinement and enhancement in plasmonic dimers, *ACS Photonics* **4**, 2871 (2017).
- [46] P. Halevi, Hydrodynamic model for the degenerate free-electron gas: Generalization to arbitrary frequencies, *Phys. Rev. B* **51**, 7497 (1995).
- [47] D. Vasileska and S. M. Goodnick, *Nano-Electronic Devices: Semiclassical and Quantum Transport Modeling* (Springer, New York, NY, 2011).
- [48] At each iteration after the first one, we calculate the fields with the loss rate obtained with Eq. (3) at the previous step, which modifies the field intensity, the degree of mode confinement, and, therefore, the value of Γ_L , as further explained in Refs. [44,45].
- [49] A. D. Boardman and R. Ruppini, The boundary conditions between spatially dispersive media, *Surf. Sci.* **112**, 153 (1981).
- [50] F. Monticone and A. Alu, Leaky-wave theory, techniques and applications: From microwaves to visible frequencies, *Proc. IEEE* **103**, 793 (2015).
- [51] A. B. Khanikaev, S. H. Mousavi, W.-K. Tse, M. Kargarian, A. H. MacDonald, and G. Shvets, Photonic topological insulators, *Nat. Mater.* **12**, 233 (2013).
- [52] U. K. Chettiar, A. R. Davoyan, and N. Engheta, Hotspots from nonreciprocal surface waves, *Opt. Lett.* **39**, 1760 (2014).
- [53] L. Shen, X. Zheng, and Z. Deng, Backscattering-immune one-way surface magnetoplasmons at terahertz frequencies, *Opt. Express* **23**, 950 (2015).
- [54] M. Marvasti and B. Rejaei, Formation of hotspots in partially filled ferrite-loaded rectangular waveguides, *J. Appl. Phys.* **122**, 233901 (2017).