




JAYADEV ATHREYA

STEVE LALLEY

JENYA SAPIR

MATTHEW WROTEN

LOCAL GEOMETRY OF RANDOM GEODESICS ON NEGATIVELY CURVED SURFACES

GÉOMÉTRIE LOCALE DES GÉODÉSIQUES
ALÉATOIRES SUR LES SURFACES DE
COURBURE STRICTEMENT NÉGATIVE

ABSTRACT. — We show that the tessellation of a compact, negatively curved surface induced by a long random geodesic segment, when properly scaled, looks locally like a *Poisson line process*. This implies that the global statistics of the tessellation – for instance, the fraction of triangles – approach those of the limiting Poisson line process.

Keywords: self-intersection, random tessellation, geodesic, hyperbolic surface, Poisson line process.
2020 Mathematics Subject Classification: 37D40, 37E35, 37B10.

DOI: <https://doi.org/10.5802/ahl.70>

(*) Athreya is supported by National Science Foundation CAREER grant DMS 1559860, and grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network), and National Science Foundation grant DMS 2003528. Lalley is supported by National Science Foundation grant DMS 1612979. Athreya and Sapir thank the Mathematical Sciences Research Institute for its hospitality in Fall 2019.

RÉSUMÉ. — Nous montrons que la tessellation d'une surface compacte de courbure strictement négative induite par un long segment géodésique aléatoire ressemble localement à un processus de Poisson en droites, après rééchelonnement. Ceci implique que les statistiques globales de la tessellation (par exemple, la proportion de triangles) convergent vers celles du processus de Poisson en droites limite.

1. Main Results: Intersection Statistics of Random Geodesics

1.1. Local Statistics

Any sufficiently long geodesic segment γ on a compact, negatively curved surface S partitions S into a finite number of non-overlapping geodesic polygons of various shapes and sizes, whose vertices⁽¹⁾ are the self-intersection points of γ . If a geodesic segment γ of length T is chosen by selecting its initial tangent vector v at random, according to (normalized) Liouville measure μ_L on the unit tangent bundle T^1S , then with probability 1, as $T \rightarrow \infty$ the maximal diameter of a polygon in the induced partition will converge to 0, and hence the number of polygons in the partition will become large. The goal of this paper is to elucidate some of the statistical properties of this random polygonal partition for large T . Our main result will be a *local* geometric description of the partition: roughly, this will assert that in a neighborhood of any point $x \in S$ the partition will, in the large- T limit, look as if it were induced by a *Poisson line process* [Mil64a, Mil64b]. We will also show that this result has implications for the *global* statistics of the partition: for instance, it will imply that with probability ≈ 1 the fraction of polygons in the partition that are *triangles* will stabilize near a non-random limiting value $\tau_3 > 0$.

DEFINITION 1.1. — A Poisson line process $\mathcal{L}$ of intensity $\kappa > 0$ is a random collection $\mathcal{L} = \{L_n\}_{n \in \mathbb{Z}}$ of lines in $\mathbb{R}^2$ constructed as follows. Let $\{(R_n, \Theta_n)\}_{n \in \mathbb{Z}}$ be the points of a Poisson point process⁽²⁾ of intensity κ/π on the infinite strip $\mathbb{R} \times [0, \pi)$. For each $n \in \mathbb{Z}$ let L_n be the line

$$(1.1) \quad L_n := \{(x, y) \in \mathbb{R}^2 : R_n = x \cos \Theta_n + y \sin \Theta_n\}.$$

That is, we consider the line through the origin of angle Θ_n to the horizontal, and L_n is the line orthogonal to this line passing through it at distance R_n from the origin. Observe that the mapping (1.1) of points (r, θ) to lines is a bijection from the strip $\mathbb{R} \times [0, \pi)$ to the space of all lines in $\mathbb{R}^2$. For any convex region $\Omega \subset \mathbb{R}^2$,

⁽¹⁾ A long segment of a random geodesic ray doesn't quite induce a tessellation, as there will be two faces [triangles, quadrilaterals, or whatever] that contain the two ends of the geodesic segment. We ignore these, however, since they will not influence statistics when the length of the geodesic segment is large.

⁽²⁾ The ordering of the points doesn't really matter, but for definiteness take $\dots < R_{-1} < 0 < R_0 < R_1 < \dots$. The assumption that $\{(R_n, \Theta_n)\}_{n \in \mathbb{Z}}$ is a Poisson point process of intensity κ/π is equivalent to the assumption that $\{R_n\}_{n \in \mathbb{Z}}$ is a Poisson point process of intensity κ on $\mathbb{R}$ and that $\{\Theta_n\}_{n \in \mathbb{Z}}$ is an independent sequence of i.i.d. random variables with uniform distribution on $[0, \pi]$.

call the restriction to Ω of a Poisson line process a Poisson line process in Ω . It is not difficult to show (see Lemma 2.4 below) that, with probability one, if Ω is a bounded domain with piecewise smooth boundary then the Poisson line process in Ω will consist of only finitely many line segments, and that at most two line segments will intersect at any point of Ω . For any realization of the process, the line segments will uniquely determine (and be determined by) their intersection points with $\partial\Omega$, grouped in (unordered) pairs.

In order to formulate our main result, we must explain how geodesic segments in a small neighborhood of a point $x \in S$ are associated with line segments in the tangent space $T_x S$. We shall assume throughout that the Riemannian metric ρ on S is C^∞ ; therefore, geodesics are C^∞ curves that depend smoothly on their initial tangent vectors. Furthermore, we will only consider geodesics of unit speed. Fix $x \in S$, and consider a small disk $D(x, r)$ on S of radius r centered at x . A (unit-speed) geodesic ray $\gamma_t(v)$ with initial tangent vector $v \in T^1 S$ distributed according to normalized Liouville measure μ_L (that is, $v = (y, \theta)$ where $y \in S$ is distributed according to normalized surface area measure and θ is distributed according to the uniform distribution on the set $[0, 2\pi]$ of directions based at x) will, with probability one, eventually enter $D(x, r)$, at a time roughly of order $1/r$ (this will follow from our main results). Thus, if we wish to study the local intersection statistics of a random geodesic segment of (large) length T in a neighborhood of x , we should focus on the intersections of the geodesic segment with neighborhoods of x of diameters proportional to $1/T$.

For any $\alpha > 0$ and $T > 0$, set $\mathcal{D}_T(x, \alpha) := D(x, \alpha T^{-1})$ to be the ball of radius α/T about x in S . Let $\exp_x : T_x S \rightarrow S$ be the exponential mapping. Then,

$$(1.2) \quad \mathcal{D}_T(x, \alpha) = \left\{ \exp_x(v) : \|v\| \leq \alpha T^{-1} \right\}.$$

The boundary $\partial(x, \alpha T^{-1})$ is a smooth closed curve. Consequently, the intersection of $\mathcal{D}_T(x, \alpha)$ with any geodesic segment will consist of (i) finitely many geodesic crossings of $\mathcal{D}_T(x, \alpha)$; (ii) up to two incomplete geodesic crossings; and (iii) a finite number of isolated points on $\partial\mathcal{D}_T(x, \alpha)$, the latter coming from tangencies of the geodesic with the boundary. Since the set of all unit tangent vectors tangent to the curve $\partial\mathcal{D}_T(x, \alpha)$ has Liouville measure 0, tangent intersections will have probability zero if the initial vector of the geodesic is chosen randomly; hence, we shall henceforth ignore these. Furthermore, incomplete geodesic crossings will occur if and only if the initial or terminal point of the geodesic segments lies in the interior of $D(x, \alpha T^{-1})$; this will occur with probability of order $O(\text{area}(D(x, \alpha T^{-1}))) = O(T^{-2})$, and so can also be ignored in the $T \rightarrow \infty$ limit. Thus, with probability $\rightarrow 1$, the intersection consists of finitely many geodesic crossings. Now any geodesic crossing of $D(x, \alpha T^{-1})$ pulls back, via the scaled exponential mapping $v \mapsto \exp_x(v/T)$, to a smooth curve in the ball $B(0, \alpha)$ with endpoints on the circle $\partial B(0, \alpha)$. When T is large, such a curve will closely approximate the *chord* of the circle with the same endpoints on $\partial B(0, \alpha)$.

Suppose now that $v \in T^1 S$ is a unit vector chosen randomly according to normalized Liouville measure μ_L . Define $\mathcal{I}_T = \mathcal{I}_T(v; x, \alpha)$ to be the intersection of the geodesic segment $\gamma_{[0, T]}(v)$ with the set $D(x, \alpha T^{-1})$, and define $\mathcal{L}_T = \mathcal{L}_T(v; x, \alpha)$ to

be the finite set of chords in $B(0, \alpha) \subset T_x S$ obtained by pulling back the geodesic crossings from $\mathcal{I}_T$ and then replacing the resulting curves by the corresponding chords.

THEOREM 1.2. — *Let S be a compact surface of genus $g \geq 2$, and assume that S is endowed with a C^∞ Riemannian metric ρ of negative curvature. Fix $x \in S$ and $\alpha > 0$, and let $\mathcal{L}_T = \mathcal{L}_T(v; x, \alpha)$ be the random chord process corresponding to the intersection $\mathcal{I}_T = \mathcal{I}_T(v; x, \alpha)$ of a random geodesic segment (i.e., one whose initial tangent vector v is chosen randomly according to the normalized Liouville measure) of length T with the neighborhood $D(x; \alpha T^{-1})$ of x . As $T \rightarrow \infty$, the random chord process $\mathcal{L}_T$ converges in distribution to a Poisson line process in $B(0; \alpha)$ of intensity*

$$(1.3) \quad \kappa = \kappa_S = \frac{1}{\text{area}(S)}.$$

Because the elements of the random processes here live in somewhat unusual spaces (finite unions of chords), we now elaborate on the meaning of convergence in distribution. In general, we say that a sequence of random elements of a complete metric space $\mathcal{X}$ converge in distribution if their distributions (the induced probability measures on $\mathcal{X}$) converge weakly. Weak convergence is defined as follows [Bil68]: if μ_n, μ are Borel probability measures on a complete metric space $\mathcal{X}$, then $\mu_n \rightarrow \mu$ weakly if for every bounded, continuous function $f : \mathcal{X} \rightarrow \mathbb{R}$,

$$(1.4) \quad \lim_{n \rightarrow \infty} \int f d\mu_n = \int f d\mu.$$

In Theorem 1.2, the appropriate metric space is

$$\mathcal{X} = \bigcup_{n=0}^{\infty} \mathcal{X}_n$$

where $\mathcal{X}_n$ is the set of all collections of n *unordered* pairs $y_i, z_i \in \partial B(0; \alpha)$. For any two such unordered pairs $\{y, z\}, \{y', z'\}$, set

$$d(\{y, z\}, \{y', z'\}) = \min(d(y, y') + d(z, z'), d(y, z') + d(z, y'));$$

and for any two elements $F, F' \in \mathcal{X}$, define

$$\begin{aligned} d(F, F') &= \min_{\pi \in \mathcal{S}_n} d(\{y_i, z_i\}, \{y'_{\pi(i)}, z'_{\pi(i)}\}) \quad \text{if } F, F' \in \mathcal{X}_n, \\ &= \infty \quad \text{otherwise} \end{aligned}$$

where $\mathcal{S}_n$ is the group of permutations of the set $[n]$. Henceforth, we will refer to this space $\mathcal{X}$ as *configuration space* (the dependence on the parameter $\alpha > 0$ will be suppressed).

The proof of Theorem 1.2 will also show that the limiting Poisson line processes in neighborhoods of distinct points of S are independent.

THEOREM 1.3. — *Fix two distinct points $x, x' \in S$ and $\alpha > 0$, and let $\mathcal{L}_T$ and $\mathcal{L}'_T$ be the chord processes induced by intersections of a random geodesic of length T with the neighborhoods $D(x; \alpha T^{-1})$ and $D(x'; \alpha T^{-1})$, respectively. Then as $T \rightarrow \infty$, the random chord processes $\mathcal{L}_T$ and $\mathcal{L}'_T$ converge jointly in distribution to a pair of independent Poisson line processes in $B(0; \alpha)$, both of intensity $\kappa = \frac{1}{\text{area}(S)}$, as in (1.3).*

1.2. Heuristics

There is an explanation for the convergence to Poisson line processes that falls short of being a complete proof. This heuristic argument is, in essence, the same as that used in [Lal96] to guess the limiting frequency of self-intersections of a random geodesic segment. It rests on the fact that the (normalized) Liouville measure μ_L on the unit tangent bundle T^1S is a mixing invariant measure for the geodesic flow.

Let $\tilde{\gamma} : [0, \infty) \rightarrow T^1S$ be a random geodesic ray with distribution μ_L , viewed as a (random) curve in the unit tangent bundle T^1S , and let γ be its projection to the surface S . Since μ_L is invariant for the geodesic flow, for any fixed time $t > 0$ the random point $\gamma(t)$ will be uniformly distributed on S (according to normalized surface area measure), and the tangent angle $\gamma'(t)$ will be uniformly distributed on $[0, 2\pi]$ (according to normalized Lebesgue measure). Fix $\varepsilon > 0$, and let $T = N\varepsilon$ be a large integer multiple of ε ; then the geodesic segment $\tilde{\gamma}[0, T)$ can be partitioned into N nonoverlapping segments $\Gamma_j := \tilde{\gamma}[j\varepsilon, (j+1)\varepsilon)$, each of whose initial tangent vectors $\tilde{\gamma}(j\varepsilon)$ is uniformly distributed according to μ_L . If ε is sufficiently small then any pair of segments $\Gamma_j, \Gamma_{j'}$ will intersect at most once. Moreover, as $\varepsilon \rightarrow 0$ the segments Γ_j approximate straight line segments of length ε in the tangent plane at the initial point.

Now we appeal to the fact that the geodesic flow is mixing relative to μ_L . This implies that for any two integers j, j' such that $|j - j'|$ is large, the random vectors $\tilde{\gamma}(j\varepsilon)$ and $\tilde{\gamma}(j'\varepsilon)$ of the random segments Γ_j and $\Gamma_{j'}$ are *approximately* independent. This suggests that the pattern and number of self-intersections in $\gamma[0, T]$ should not differ appreciably from those of a random sample of N independent random geodesic segments Γ'_j of length ε , each of whose initial tangent vectors is randomly chosen from μ_L .




Figure 1.1. Γ_j and $\Gamma_{j'}$ intersect in an angle close to θ .

Consider, in particular, the number of self-intersections of $\gamma[0, T)$. For $\theta \in [0, \pi]$ and for any pair of indices j, j' , the event $F_{j,j'}(\theta; d\theta)$ that the projections to S of the segments Γ_j and $\Gamma_{j'}$ cross at angle between $\theta - d\theta$ and $\theta + d\theta$ is, up to an error of size $O(d\theta)$, the same as the event that (i) the point $\gamma(j'\varepsilon)$ lies in a rhombus on S whose sides meet at angle θ and whose “top” side is the projection to S of Γ_j , and (ii) the tangent angle of $\tilde{\gamma}(j'\varepsilon)$ differs from that of $\tilde{\gamma}(j\varepsilon)$ by $\theta \pm d\theta$ (Figure 1.1). (Similarly, for Γ_j and $\Gamma_{j'}$ to cross at angle $\pi + (\theta \pm d\theta)$ the “bottom” side of the rhombus

should be the projection of Γ_j .) Since Γ_j and $\Gamma_{j'}$ are approximately independent, the probability of this event is (approximately) the relative area of this rhombus times $2d\theta$ divided by 2π . Summing over θ and taking $d\theta \rightarrow 0$ now shows that the probability of intersection is about

$$\frac{2\varepsilon^2 \int_0^\pi \sin \theta d\theta}{2\pi \text{area}(S)} = \frac{2\varepsilon^2 \kappa}{\pi}$$

where $\kappa = (1/\text{area}(S))$. This fails, of course, if $|j - j'|$ is small, but for most pairs j, j' the difference will be large. Consequently, by the law of large numbers, the number of self-intersections of the segment $\gamma[0, T]$, when divided by T^2 , should satisfy

$$\frac{1}{T^2} \sum_{j=1}^N \sum_{j'=j+1}^N \mathbf{1} \{ \Gamma_{j'} \text{ crosses } \Gamma_j \} \approx \frac{\kappa}{\pi}.$$

Amplification of this argument “explains” the local convergence of the induced tessellation to the Poisson line process. Consider, for instance, the number of distinct geodesic arcs that cross the disk $D(x, \alpha/T)$, for some fixed point $x \in S$: we will argue that this should have approximately a Poisson distribution. Choose ε small, and let $T = N\varepsilon$ be an integer multiple of ε large enough that $1/T \ll \varepsilon$. For each $j \leq N$, the probability that (the projection of) Γ_j crosses the disk $D(x, \alpha T^{-1})$ is, for small ε , about $C\varepsilon\alpha/T = C\alpha/N$ for a suitable geometric constant $C > 0$. (This follows by a simple geometric argument similar to that given above for self-intersections.) Thus, if the random segments Γ_j were actually independent, the number that would cross the disk $D(x, \alpha/T)$ would be the sum of N independent Bernoulli random variables each with mean $C\alpha/N$. For N large, the distribution of this count would therefore converge to Poisson with mean $C\alpha$ (cf. Proposition 2.11 below).

The sticky point, of course, is that the random segments Γ_j are not independent. What is worse, the events of interest (for instance, the event that Γ_j crosses the disk $D(x, \alpha T^{-1})$) are events whose probabilities become small as N becomes large; thus, the mixing property of the geodesic flow does not by itself imply that

$$\frac{P(\{\Gamma_j \text{ crosses } D(x, \alpha T^{-1})\} \cap \{\Gamma_{j'} \text{ crosses } D(x, \alpha T^{-1})\})}{P\{\Gamma_j \text{ crosses } D(x, \alpha T^{-1})\} P\{\Gamma_{j'} \text{ crosses } D(x, \alpha T^{-1})\}} \approx 1$$

even for $|j - j'|$ large. The rigorous arguments to be given below are largely designed to circumvent the failure of mixing at this level by exploiting the Gibbsian structure of the Liouville measure.

Mixing problems in which the events of interest have probabilities tending to zero are known as “shrinking target” problems. Such problems occur naturally in hyperbolic dynamics: see, for instance, [Sul82], where the “target” is one of the cusps of a non-compact hyperbolic surface of finite area, or Kleinbock–Margulis [KM99], who consider related problems for diagonal flows on finite-volume homogeneous spaces. For shrinking target problems where the targets lie in the compact part of the space, see Dolgopyat [Dol04], and Maucourant [Mau06]. Unfortunately none of these results is easily adapted to the problems we consider here.

1.3. Global Statistics

Theorems 1.2–1.3 describe the “local” structure of the random tessellation $\mathcal{T}_T$ of the surface S induced by a long segment $\gamma[0, T]$ of a random geodesic. The tessellation $\mathcal{T}_T$ will consist of geodesic polygons, typically of diameter of order T^{-1} , since the $O(T^2)$ self-intersections will subdivide the length T geodesic segment into sub-segments of length $O(T^{-1})$. Thus, it is natural to look at the statistics of the scaled tessellation $T\mathcal{T}_T$, which we view as consisting of a random number of triangles, quadrilaterals, etc., each with its own set of side-lengths and interior angles.

The empirical frequencies of triangles, quadrilaterals, etc. and the empirical distribution of side-length and interior-angle sets in a Poisson line process of intensity κ on the ball $B(0; \alpha)$ of radius α converge as $\alpha \rightarrow \infty$. (These results are evidently due to R. E. Miles [Mil64a, Mil64b]; proofs are given in Section 2 below.) Theorem 1.2 asserts that when L is large, then for any point $x \in S$ the statistics of the polygonal partition in $B(x; \alpha^{-1}L)$ induced by a random geodesic segment of length L should approach those of a Poisson line process. From this observation we will deduce the following assertion regarding *global* statistics.

THEOREM 1.4. — *Let $\mathcal{T}_T$ be the tessellation of S induced by a random geodesic of length T . Then with probability approaching 1 as $T \rightarrow \infty$, the empirical frequencies of triangles, quadrilaterals, etc. and the empirical distribution of side-length and interior-angle sets in $\mathcal{T}_T$ approach the corresponding theoretical frequencies for a Poisson line process.*

For example, for each $v \in T^1(S)$, let $\mathcal{T}_T(v)$ be the tessellation induced by the length T arc with initial direction v . Let f_3 be the function that returns the frequency of triangles in a tessellation. Suppose the expected value of f_3 is τ_3 for a Poisson line process. Then we show that for any $\epsilon > 0$,

$$\lim_{T \rightarrow \infty} \mu_L \left\{ v \in T^1(S) : |f_3(\mathcal{T}_T(v)) - \tau_3| > \epsilon \right\} = 0$$

where μ_L is the Liouville measure on $T^1(S)$.

Plan of the paper.

The proofs of Theorems 1.2–1.3 will occupy most of the paper. The strategy will be to reduce the problem to a corresponding counting problem in symbolic dynamics. Preliminaries on Poisson line processes will be collected in Section 2, and preliminaries on symbolic dynamics for the geodesic flow in Section 3. Section 4 will be devoted to heuristics and a reformulation of the problem; the proofs of Theorems 1.2–1.3 will then be carried out in Sections 5–8. Theorem 1.4 will be proved in Section 9. Finally, in Section 10, we give a short list of conjectures, questions, and possible extensions of our main results.

2. Preliminaries: Poisson line processes

The Poisson line process and its generalizations have a voluminous literature, with notable early contributions by Miles [Mil64a, Mil64b]. See [SKM87] for an extended

discussion and further pointers to the literature. In this section we will record some basic facts about these processes. These are mostly known – some of them are stated as theorems in [Mil64a] without proofs – but proofs are not easy to track down, so we shall provide proof sketches in Appendix A.

2.1. Statistics of a Poisson line process

LEMMA 2.1. — *A Poisson line process of constant intensity κ is rotationally and translationally invariant, that is, if A is any orientation preserving isometry of $\mathbb{R}^2$ then the configuration $\{AL_n\}_{n \in \mathbb{Z}}$ has the same joint distribution as the configuration $\{L_n\}_{n \in \mathbb{Z}}$.*

Remark 2.2. — This result is stated without proof in [Mil64a]. A proof of the corresponding fact for the intensity measure can be found in [San04], and another in [SKM87, Chapter 8]. A short, elementary proof is given in Appendix A. The following Corollary 2.3, which is stated without proof as [Mil64a, Theorem 2], follows easily from isometry-invariance.

COROLLARY 2.3. — *Let $\mathcal{L}$ be a Poisson line process of intensity $\kappa > 0$. For any fixed line ℓ in $\mathbb{R}^2$, the point process of intersections of ℓ with lines in $\mathcal{L}$ is a Poisson point process of intensity $2\kappa/\pi$.*

LEMMA 2.4. — *Let $\mathcal{L}$ be a Poisson line process of intensity $\kappa > 0$, and for each point $x \in \mathbb{R}^2$ and each real $r > 0$ let $N(B(x;r))$ be the number of lines in $\mathcal{L}$ that intersect the ball $B(x;r)$ of radius r centered at x . Then the random variable $N(B(x;r))$ has the Poisson distribution with mean $2\kappa r$. Consequently, with probability one, for any compact set $K \subseteq \mathbb{R}^2$ the set of lines L_n in $\mathcal{L}$ that intersect K is finite.*

Proof. — Without loss of generality, take $K = B(0; R)$ to be the closed ball of radius R centered at the origin. Then the line L_n intersects K if and only if $|R_n| \leq R$. Since a Poisson point process on $\mathbb{R}$ of constant intensity has at most finitely many points in any finite interval, the result follows. $\square$

The next result characterizes the Poisson line process (see also Proposition 2.10 below). Fix a bounded, convex region $D \subset \mathbb{R}^2$ with C^∞ boundary $\Gamma = \partial D$, and let A, B be non-intersecting closed arcs on Γ . For any line process $\mathcal{L}$, let

$$(2.1) \quad N_{\{A,B\}} = \# \{\text{lines that cross both } A \text{ and } B\}.$$

For any angle $\theta \in [-\pi/2, \pi/2]$, the set of lines that intersect both A and B and meet the x -axis at angle $\theta + \pi/2$ constitute an infinite strip that intersects the line $\{re^{i\theta}\}_{r \in \mathbb{R}}$ in an interval; see Figure 2.1 below. Let $\psi(\theta) = \psi_{A,B}(\theta)$ be the length of this interval, and define

$$(2.2) \quad \beta_{A,B} = \frac{1}{\pi} \int_{-\pi/2}^{\pi/2} \psi(\theta) d\theta.$$




Figure 2.1. Lines that cross A and B at angle θ .

PROPOSITION 2.5. — *A line process $\mathcal{L}$ in D is a Poisson line process of rate $\kappa > 0$ if and only if*

- (i) *for any two non-intersecting arcs $A, B \subset \Gamma$, the random variable $N_{\{A, B\}}$ has the Poisson distribution with mean $\kappa\beta_{A, B}$, and*
- (ii) *for any finite collection $\{A_i, B_i\}_{i \leq m}$ of pairwise disjoint boundary arcs, the random variables $N_{\{A_i, B_i\}}$ are mutually independent.*

See Appendix A for the proof of the forward implication, along with that of the following corollary. The converse implication in Proposition 2.5 will follow from Proposition 2.10 in Section 2.3 below.

COROLLARY 2.6. — *Let $D \subset \mathbb{R}^2$ be a compact, convex region, and let $\mathcal{L}$ be a Poisson line process with intensity κ . The number $V(D)$ of intersection points (vertices) of $\mathcal{L}$ in D has expectation*

$$EV(D) = \kappa^2 |D| / \pi$$

where $|D|$ is the Lebesgue measure of D .

2.2. Ergodic theorem for Poisson line processes

The configuration space $\mathcal{C}$ in which a Poisson line process takes values is the set of all countable, locally finite collections of lines in $\mathbb{R}^2$. This space has a natural metric topology, specifically, the weak topology generated by the Hausdorff topologies on the restrictions to balls in $\mathbb{R}^2$. Moreover, $\mathcal{C}$ admits an action (by translations) of $\mathbb{R}^2$. Denote by ν_κ the distribution of the Poisson line process with intensity κ . By Lemma 2.1, the measure ν_κ is translation-invariant.

PROPOSITION 2.7. — *The probability measure ν_κ is mixing (and therefore ergodic) with respect to the translational action of $\mathbb{R}^2$ on $\mathcal{C}$.*

Remark 2.8. — Ergodicity of the measure ν_κ is asserted in Miles' papers [Mil64a, Mil64b], and proved in his unpublished Ph. D. dissertation. We have been unable to locate a proof in the published literature, so we have provided one in the Appendix.

COROLLARY 2.9. — *Let $\Phi_{n, k}$ be the fraction of k -gons, F_n (for “faces”) the total number of polygons, and V_n (for “vertices”) the number of intersection points in the tessellation of the square $[-n, n]^2$ induced by a Poisson line process $\mathcal{L}$ of intensity κ .*

There exist constants $\phi_k > 0$ such that with probability 1,

$$(2.3) \quad \lim_{n \rightarrow \infty} F_n / (2n)^2 = \kappa^2 / \pi,$$

$$(2.4) \quad \lim_{n \rightarrow \infty} V_n / (2n)^2 = \kappa^2 / \pi, \quad \text{and}$$

$$(2.5) \quad \lim_{n \rightarrow \infty} \Phi_{n,k} = \phi_k.$$

Integral formulas for the quantities ϕ_k are given in [Cal03].

The ergodic theorem can also be used to prove that a variety of other statistical properties stabilize in large squares. Consider, for example, the number $N_n(A, B, C)$ of triangles contained in $[-n, n]^2$ whose side lengths α, β, γ lie in the intervals A, B, C ; then as $n \rightarrow \infty$,

$$N_n(A, B, C) / (2n)^2 \longrightarrow E \mathbf{1}_{G(A, B, C)}(\mathcal{L})$$

where $G(A, B, C)$ is the event that the polygon containing the origin is a triangle with side lengths in A, B, C .

2.3. Weak convergence to a Poisson line process

For any unordered pair $\{A, B\}$ of non-overlapping boundary arcs of the disk $B(0, \alpha)$, let $L_{\{A, B\}}$ be the set of lines in $\mathbb{R}^2$ that intersect both A and B . This set can be identified with the set of point pairs $\{x, y\}$ where $x \in A$ and $y \in B$. This allows us to view any random collection of unordered point pairs $\{x, y\}$ as a *line process* in $B(0, \alpha)$, even when the collection consists of endpoints of arcs across $B(0, \alpha)$ that are not line segments (in particular, when they are pullbacks of geodesic arcs to the tangent space). For any line process $\mathcal{L}$ in $B(0, \alpha)$ let $N_{\{A, B\}}$ be the cardinality of $\mathcal{L} \cap L_{\{A, B\}}$ (cf. equation (2.1)).

PROPOSITION 2.10. — *Let $\mathcal{L}_n$ be a sequence of line processes in $B(0; \alpha)$, and let μ_n be the distribution of $\mathcal{L}_n$ (i.e., the probability measure on $\mathcal{X}$ induced by $\mathcal{L}_n$). In order that $\mu_n \rightarrow \mu$ weakly, where μ is the law of a rate- κ Poisson line process, it suffices that the following condition holds. For any finite collection $\{\{A_i, B_i\}\}_{i \leq m}$ of unordered pairs of non-overlapping boundary arcs of $B(0; \alpha)$ such that the sets $L_{\{A_i, B_i\}}$ are pairwise disjoint, the joint distribution of the counts $N_{\{A_i, B_i\}}$ under μ_n converges to the joint distribution under μ , that is, for any choice of nonnegative integers k_i ,*

$$(2.6) \quad \lim_{n \rightarrow \infty} \mu_n \left\{ N_{\{A_i, B_i\}} = k_i \quad \forall \quad i \leq m \right\} = \prod_{i=1}^m \frac{(\kappa \beta_{A_i, B_i})^{k_i}}{k_i!} e^{-\kappa \beta_{A_i, B_i}}.$$

Proof Sketch. — Recall that the configuration space $\mathcal{X}$ is the disjoint union of the sets $\mathcal{X}_k$, where $\mathcal{X}_k$ is the set of all finite sets $F = \{\{x_i, y_i\}\}_{1 \leq i \leq k}$ consisting of k unordered pairs of points on $\partial B(0, \alpha)$. Since each set $\mathcal{X}_k$ is both open and closed in $\mathcal{X}$, to prove weak convergence $\mu_n \rightarrow \mu$ it suffices to establish the convergence (1.4) for every continuous function f supported by just one of the sets $\mathcal{X}_k$.

For each k , the space $\mathcal{X}_k$ is a quotient of $(\partial B(0, \alpha))^k$ with the usual topology, and so every continuous function $f : \mathcal{X}_k \rightarrow \mathbb{R}$ can be uniformly approximated by “step functions”, that is, functions g of configurations $F = \{\{x_i, y_i\}\}_{1 \leq i \leq k}$ that depend

only on the counts N_{A_i, B_i} for arcs A_i, B_i in some partition of $\partial B(0, \alpha)$. If (2.6) holds, then it follows by linearity of expectations that for any such step function g ,

$$\lim_{n \rightarrow \infty} \int g d\mu_n = \int g d\mu,$$

and hence (1.4) follows. $\square$

2.4. The “law of small numbers”

A elementary theorem of discrete probability theory states that for large n , the Binomial— $(n, \kappa/n)$ distribution is closely approximated by the Poisson distribution with mean κ . The following is a generalization that we will find useful.

PROPOSITION 2.11. — *Let $X_1, X_2, \dots, X_n$ be independent Bernoulli random variables with success parameters $EX_i = p_i$. Let $\alpha = \max_i p_i$ and $\kappa = \sum_i p_i$. Then there is a constant $C < \infty$ not depending on $p_1, p_2, \dots, p_n$ such that*

$$\sum_{k=0}^{\infty} \left| P \left\{ \sum_i X_i = k \right\} - \frac{\kappa^k}{k!} e^{-\kappa} \right| \leq C\alpha.$$

Note that for all $k > n$, the probability $P \{ \sum_i X_i = k \}$ is zero, but the elements of the sum are not.

See [LC60] for a proof. The important feature of the proposition for us is not the explicit bound, but the fact that the closeness of the approximation depends only on $\max p_i$.

A similar result holds for multinomial variables.

PROPOSITION 2.12. — *Let $X_1, X_2, \dots, X_n$ be independent random variables each taking values in the finite set $\{0, 1, 2, \dots, K\} = \{0\} \cup [K]$, and for each pair i, j set $p_{i,j} = P\{X_i = j\}$. Let $\alpha = \max_{j \geq 1} \max_i p_{i,j}$ and $\kappa_j = \sum_i p_{i,j}$, and for each j define*

$$T_j = \sum_{i=1}^n \mathbf{1}\{X_i = j\}.$$

Then there is a function $C_K(\alpha)$ satisfying $\lim_{\alpha \downarrow 0} C(\alpha) = 0$ such that

$$\sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \cdots \sum_{m_K=0}^{\infty} \left| P \{ T_j = m_j \ \forall j \in [K] \} - \prod_{j=1}^K \kappa_j^{m_j} e^{-\kappa_j} / m_j! \right| \leq C(\alpha).$$

3. Preliminaries: Symbolic Dynamics

3.1. Shifts and suspension flows

The geodesic flow on the unit tangent bundle $T^1 S$ of a compact, negatively curved surface S has a concrete representation as a suspension flow over a shift of finite type. In describing this representation, we shall follow (for the most part) the terminology and notation of [Bow75, Lal14, Ser81]. Let $\mathcal{A}$ be a finite alphabet and $\mathcal{F}$ a finite set of finite words on the alphabet $\mathcal{A}$, and define $\Sigma = \Sigma_{\mathcal{F}}$ to be the set of doubly infinite

sequences $\omega = (\omega_n)_{n \in \mathbb{Z}}$ such that no element of $\mathcal{F}$ occurs as a subword of ω . The sequence space Σ is given the metric $d(\omega, y) = \exp\{-n(\omega, y)\}$ where $n(\omega, y)$ is the minimum nonnegative integer n such that $\omega_j \neq y_j$ for $j = n$ or $j = -n$. For each nonnegative integer m and each $\omega \in \Sigma$, define the *cylinder set* $\Sigma_m(\omega)$ to be the set of all $\omega' \in \Sigma$ that agree with ω in all coordinates j such that $|j| \leq m$; equivalently,

$$(3.1) \quad \Sigma_m(\omega) = \{\omega' \in \Sigma : d(\omega, \omega') < e^{-m}\}.$$

The forward shift $\sigma : \Sigma \rightarrow \Sigma$ is known as a (two-sided) *shift of finite type*.⁽³⁾

For any continuous function $F : \Sigma \rightarrow (0, \infty)$ on Σ , define the *suspension space* Σ_F by

$$\Sigma_F := \{(\omega, t) : \omega \in \Sigma \text{ and } 0 \leq t \leq F(\omega)\},$$

with points $(\omega, F(\omega))$ and $(\sigma\omega, 0)$ identified. The metric d on the sequence space Σ induces a metric d_{TAXI} on Σ_F , the “taxicab” metric. (Roughly, the distance between any two points (ω, t) and (ω', t') in Σ_F is the length of the shortest “path” between them consisting of alternating “horizontal” and “vertical” segments. See [BW72] for the formal definition.) The *suspension flow* with height function F is the flow ϕ_t on Σ_F whose orbits proceed up vertical fibers

$$\mathcal{F}_\omega := \{(\omega, s) : 0 \leq s \leq F(\omega)\}$$

at speed 1, and upon reaching the ceiling at $(\omega, F(\omega))$ jump instantaneously to $(\sigma\omega, 0)$. If the height function $F : \Sigma \rightarrow \mathbb{R}$ is Hölder continuous with respect to the metric d , then the suspension flow ϕ_t is Hölder continuous with respect to the metric d_{TAXI} : in particular, there exists $\alpha > 0$ such that

$$(3.2) \quad d_{\text{TAXI}}(\phi_t(\omega, 0), \phi_t(\omega', 0)) \leq e^{\alpha|t|}d(\omega, \omega') \quad \text{for all } \omega, \omega' \in \Sigma \text{ and } t \in \mathbb{R}.$$

There is a bijective correspondence between invariant probability measures μ^* for the flow ϕ_t and shift-invariant measures μ on Σ . This correspondence can be specified as follows: for any continuous function $g : \Sigma_F \rightarrow \mathbb{R}$,

$$(3.3) \quad \int g d\mu^* = \int_{\Sigma} \int_0^{F(\omega)} g(\omega, s) ds d\mu(\omega) / \int_{\Sigma} F d\mu.$$

If μ is ergodic for the shift (Σ, σ) then μ^* is ergodic for the flow (Σ_F, ϕ_t) . By Birkhoff’s theorem, for any ergodic probability measure μ on Σ ,

$$(3.4) \quad \lim_{n \rightarrow \infty} n^{-1} \sum_{j=0}^{n-1} F \circ \sigma^j = \int_{\Sigma} F d\mu \quad \text{almost surely};$$

thus, under μ^* , almost every orbit makes roughly $T / \int F d\mu$ visits to the base $\Sigma \times \{0\}$ by time T , when T is large.

⁽³⁾Bowen [Bow75] requires that the elements of the set $\mathcal{F}$ all be of length 2. However, any shift of finite type can be “recoded” to give a shift of finite type obeying Bowen’s convention, by replacing the original alphabet $\mathcal{A}$ by $\mathcal{A}^m$, where m is the length of the longest word in $\mathcal{F}$, and then replacing each sequence ω by the sequence $\bar{\omega}$ whose entries are the successive length- m subwords of ω . In Series’ [Ser81] symbolic dynamics for the geodesic flow on a closed hyperbolic surface, the alphabet $\mathcal{A}$ is a set of natural generators for the fundamental group $\pi_1(S)$ of the surface S , and the forbidden subwords $\mathcal{F}$ are gotten from the relators of $\pi_1(S)$.

3.2. Symbolic dynamics for the geodesic flow

The following Proposition 3.1 is a special case of the main result of [Rat73] (see also [Bow73]), as the geodesic flow on a compact, negatively curved surface is an Anosov flow.

PROPOSITION 3.1. — *For any compact, negatively curved surface (S, ϱ) with C^∞ Riemannian metric ϱ , there exist a topologically mixing shift (Σ, σ) of finite type, a suspension flow (Σ_F, ϕ_t) over the shift with Hölder continuous height function F , and a surjective, Hölder-continuous mapping $\pi : \Sigma_F \rightarrow T^1S$ such that π is a semi-conjugacy with the geodesic flow γ_t on T^1S , i.e.,*

$$(3.5) \quad \pi \circ \phi_t = \gamma_t \circ \pi \quad \text{for all } t \in \mathbb{R}.$$

In the special case where ϱ is a hyperbolic (constant curvature) Riemannian metric, a much more explicit symbolic dynamics was constructed by Series: see especially [Ser81, Theorem 3.1] and also [BS79]. In this symbolic dynamics, the sequence space Σ is mapped to a subset of $\partial\mathbb{D} \times \partial\mathbb{D}$, where $\partial\mathbb{D}$ is the ideal boundary of the Poincaré disk, in such a way that every vertical fiber $\mathcal{F}_\omega$ of the suspension flow is mapped to a segment of the hyperbolic geodesic in $\mathbb{D}$ whose endpoints are gotten from the boundary correspondence. Series' symbolic dynamics can be extended to the variable curvature case using the *Conformal Equivalence Theorem* ([SY94, Theorem V.1.3]) and the structural stability theorem for Anosov flows. This more explicit symbolic dynamics will not be needed in the analysis below. However, we will need the following fact (see [PPS15, Chapter 7]).

PROPOSITION 3.2. — *Under the hypotheses of Proposition 3.1, the pullback $\lambda^* := \mu_L \circ \pi^{-1}$ of the normalized Liouville measure μ_L on T^1S is Gibbs, that is, it corresponds to a Gibbs state λ for the shift via the identity (3.3).*

3.3. Regenerative representation of Gibbs states

When the underlying shift (Σ, σ) is topologically mixing, Gibbs states with Hölder continuous potentials enjoy strong exponential mixing properties (e.g., [Bow75, the “exponential cluster property” 1.26 in Chapter 1]). We shall make use of an even stronger property, the *regenerative representation* of a Gibbs state established in [Lal86].⁽⁴⁾ This representation is most usefully described in terms of the stationary process governed by the Gibbs state. Let μ be a Gibbs state with Hölder continuous potential function $f : \Sigma \rightarrow \mathbb{R}$, where $\sigma : \Sigma \rightarrow \Sigma$ is a topologically mixing shift of finite type, and let $X_n : \Sigma \rightarrow \mathcal{A}$ be the coordinate projections on Σ , for $n \in \mathbb{Z}$. The sequence $(X_n)_{n \in \mathbb{Z}}$, viewed as a stochastic process on the probability space (Σ, μ) , is a stationary process that we will henceforth call a *Gibbs process*.

The regenerative representation relates the class of Gibbs processes to another class of stationary processes, called *list processes* (the term used by [Lal86]). A list

⁽⁴⁾The article [Lal86] proves the regenerative representation only for Gibbs states defined on a full shift, but the arguments are easily modified for topologically mixing shifts of finite type. See also [CFF02] for a somewhat different approach.

process is a stationary, positive-recurrent Markov chain $(Z_n)_{n \in \mathbb{Z}}$ with state space $\cup_{k \geq 1} \mathcal{A}^k$ and stationary distribution ν that obeys the following transition rules: first,

$$(3.6) \quad P(Z_{n+1} = (\omega_1, \omega_2, \dots, \omega_m) \mid Z_n = (\omega'_1, \omega'_2, \dots, \omega'_k)) = 0$$

unless either $m = 1$ or $m = k + 1$ and $\omega_i = \omega'_i$ for each $1 \leq i \leq k$; and second, for every letter ω_1 and every word $\omega'_1 \omega'_2 \dots \omega'_m$,

$$(3.7) \quad P(Z_{n+1} = \omega_1 \mid Z_{n+1} \in \mathcal{A}^1 \text{ and } Z_n = (\omega'_1, \omega'_2, \dots, \omega'_m)) = \nu((\omega_1)) / \nu(\mathcal{A}^1).$$

Thus, the process $(Z_n)_{n \in \mathbb{Z}}$ evolves by either adding one letter to the end of the list or erasing the entire list and beginning from scratch. Furthermore, by (3.7), at any time when the list is erased, the new 1-letter word chosen to begin the next list is independent of the past history of the entire process.

For any list process define the *regeneration times* $0 = \tau_0 < \tau_1 < \tau_2 < \dots$ by

$$\begin{aligned} \tau_1 &= \min \{n \geq 1 : Z_n \in \mathcal{A}^1\}; \\ \tau_{m+1} &= \min \{n \geq 1 + \tau_m : Z_n \in \mathcal{A}^1\}. \end{aligned}$$

By condition (3.7), the random variables $\tau_{m+1} - \tau_m$ are independent, and for $m \geq 1$ are identically distributed, as are the *excursions*

$$(Z_{\tau_m+1}, Z_{\tau_m+2}, \dots, Z_{\tau_{m+1}}).$$

Denote by $\pi : \cup_{k \geq 1} \mathcal{A}^k \rightarrow \mathcal{A}$ the projection onto the *last* letter.

PROPOSITION 3.3. — *If $(X_n)_{n \in \mathbb{Z}}$ is a Gibbs process then there is a list process $(Z_n)_{n \in \mathbb{Z}}$ such that the projected process $(\pi(Z_n))_{n \in \mathbb{Z}}$ has the same joint distribution as the Gibbs process $(X_n)_{n \in \mathbb{Z}}$. Thus, the random sequence obtained by concatenating the successive excursions $W_m := Z_{\tau_m}$, i.e.,*

$$W_1 \cdot W_2 \cdot W_3 \cdot \dots,$$

has the same distribution as the sequence $\{X_n\}_{n \geq 0}$. Moreover, the list process can be chosen in such a way that the excursion lengths $\tau_{m+1} - \tau_m$ satisfy

$$(3.8) \quad P(\tau_{m+1} - \tau_m \geq n) \leq C\alpha^n$$

for some $0 < \alpha < 1$ and $C < \infty$ not depending on either m or n .

See [Lal86, Theorem 1], or [CFF02, Theorem 4.1.], (The article [Lal86] uses the (older) term *chain with complete connections* for a Gibbs process, and a different (but equivalent) definition than that given in [Bow75]. Moreover, [Lal86] considers only the case where the underlying shift is the full shift on the symbol set $\mathcal{A}$, although the proof extends routinely to the general case. See [LNP19] for details.)

4. Theorem 1.2 Proof: Strategy

We shall use the symbolic dynamics outlined in Section 3.2 to translate the weak convergence problem to a problem involving the Gibbs state λ corresponding to the pullback λ^* of Liouville measure to the suspension space Σ_F . Recall (cf. Proposition 3.1) that the projection $\pi : \Sigma_F \rightarrow T^1S$ provides a semi-conjugacy (3.5) between

the suspension flow ϕ_t and the geodesic flow γ_t ; thus, each segment of the suspension flow projects (via the mapping $p \circ \pi$, where $p : T^1S \rightarrow S$ is the natural projection) to a geodesic segment of the same length, and in particular, each fiber $\mathcal{F}_\omega$ of the suspension space Σ_F projects to a geodesic segment of length $F(\omega)$.

By Birkhoff's ergodic theorem (cf. equation (3.4)), for λ -almost every $\omega \in \Sigma$, the number of returns to the base $\Sigma \times \{0\}$ made by the suspension flow by time T , divided by T , converges to $1/E_\lambda F$. We will show (cf. Proposition 5.1 in Section 5 below) that for any $\alpha > 0$, the probability that a random geodesic enters the disk $D(x, \alpha T^{-1})$ between times $(1 - \varepsilon)T$ and $(1 + \varepsilon)T$ is bounded by $C\varepsilon$ for some $C = C(\alpha) < \infty$ not depending on T . Consequently, with λ -probability approaching 1 as $T \rightarrow \infty$, the intersection of the random geodesic segment $p \circ \pi(\phi_{[0, T]}(\omega))$ with $D(x, \alpha T^{-1})$ is identical to the intersection of $D(x, \alpha T^{-1})$ with the geodesic segment

$$(4.1) \quad p \circ \pi(\cup_{i=0}^{n(T)-1} \mathcal{F}_{\sigma^i \omega}) \quad \text{where} \quad n(T) = [T/E_\lambda F].$$

Henceforth, we will use the abbreviation $n = n(T)$.

Denote by $I_n(x, \omega)$ be the intersection of the geodesic segment $p \circ \pi(\cup_{i=0}^{n-1} \mathcal{F}_{\sigma^i \omega})$ with the neighborhood $D(x; \alpha T^{-1})$, and by $J_n(x, \omega)$ the pullback to a finite collection of chords of the ball $B(0, \alpha)$ in the tangent space $T_x S$ (cf. the discussion preceding the statement of Theorem 1.2). Our goal is to prove that, for any fixed $x \in \mathcal{S}$, the sequence of line processes J_n converges in law to a Poisson line process on $B(0, \alpha)$. For this we will use the criterion of Proposition 2.10.

For any pair A, B of non-overlapping boundary arcs of $\partial B(0, \alpha)$, define $L_{A, B}$ to be the set of *oriented* line segments from A to B , and let $N_{A, B}(\omega)$ be the number of oriented chords in $J_n(x, \omega)$ from boundary arc A to boundary arc B in $B(0, \alpha)$, equivalently, the number of oriented geodesic segments in the collection $I_n(x; \omega)$ that cross the target neighborhood $D(x; \alpha T^{-1})$ from (the image of) arc A to (the image of) arc B . (Recall that $B(0, \alpha)$ is identified with the neighborhood $D(x, \alpha T^{-1})$ by the scaled exponential mapping. Henceforth, for any pair of arcs A, B in $\partial B(0, \alpha)$ we shall denote by A^T, B^T the corresponding boundary arcs of $D(x, \alpha T^{-1})$.) The counts $N_{A, B}$ depend on $n = [T/E_\lambda F]$ and ω , but to reduce notational clutter we shall suppress this dependence. Observe that the number of *undirected* crossings $N_{\{A, B\}}$ (cf. equation (2.1)) is given by

$$N_{\{A, B\}} = N_{A, B} + N_{B, A},$$

and consequently $EN_{\{A, B\}} = EN_{A, B} + EN_{B, A}$. Since the sum of independent Poisson random variables is Poisson, to prove that in the $n \rightarrow \infty$ limit the random variable $N_{\{A, B\}}$ becomes Poisson, it suffices to show that the *directed* crossing counts $N_{A, B}$ become Poisson. Thus, our objective now is to prove the following assertion, which, by Proposition 2.10, will imply Theorem 1.2.

PROPOSITION 4.1. — *For any finite collection $\{(A_i, B_i)\}_{i \leq r}$ of pairs of non-overlapping closed boundary arcs of $B(0, \alpha)$ such that the sets L_{A_i, B_i} are pairwise disjoint, and for any choice of nonnegative integers k_i ,*

$$(4.2) \quad \lim_{n \rightarrow \infty} \lambda \{ \omega : N_{A_i, B_i}(\omega) = k_i \ \forall \ i \} = \prod_{i=1}^r \frac{(\kappa \beta_{A_i, B_i} / 2)^{k_i}}{k_i!} e^{-\kappa \beta_{A_i, B_i} / 2}$$

where $\beta_{A,B}$ is defined by equation (2.2) (with $D = B(0, a)$) and $\kappa = 1/\text{area}(S)$.

Note that for fixed boundary arcs A, B the constants $\beta_{A,B} = \beta_{A,B}(\alpha)$ are proportional to α , because the function $\psi = \psi_{A,B}$ in (2.2) is proportional to α . See Figure 2.1.

The proof of Proposition 4.1 will be accomplished in four stages, as follows.

First, we will prove in Section 5 that for any positive function $f(T)$ satisfying

$$\lim_{T \rightarrow \infty} f(T)/T = 0,$$

- (a) the probability that a random geodesic ray enters the neighborhood $D(x, \alpha T^{-1})$ before time $f(T)$ converges to 0 as $T \rightarrow \infty$ (meaning there are no quick entries), and
- (b) the probability that a random geodesic ray enters $D(x, \alpha T^{-1})$ before time T and then re-enters (after having exited) within time $f(T)$ also converges to 0 (meaning there are no quick re-entries).

This will justify the replacement of the random length- T geodesic segment in the statement of Theorem 1.2 by the geodesic segment (4.1) above, and will also ultimately be used to partition this segment into nearly independent blocks.

Second, define $\Sigma(A, B; T)$ to be the set of all sequences $\omega \in \Sigma$ such that the geodesic segment $p \circ \pi(\mathcal{F}_\omega)$ intersects $D(x, \alpha T^{-1})$ in a geodesic segment with terminal endpoint in the boundary arc B^T , and either coincides with or extends to a geodesic crossing from boundary arc A^T to boundary arc B^T . (Note that if the image $p \circ \pi(\omega, 0)$ of the base point $(\omega, 0)$ lies in the interior of $D(x, \alpha T^{-1})$ then the intersection will only be a partial crossing.) By assertions (a) and (b) above, the event $\{\omega : N_{A,B}(\omega) = k\}$ coincides (up to a set of measure $\rightarrow 0$ as $T \rightarrow \infty$) with the set of sequences $\omega \in \Sigma$ such that

$$\sum_{i=0}^{n-1} \mathbf{1}_{\Sigma(A, B; T)}(\sigma^i \omega) = k,$$

that is, sequences whose forward σ -orbits $(\sigma^i \omega)_{i \geq 0}$ make exactly k visits to the set $\Sigma(A, B; T)$ for $i \leq n-1$. We will prove, in Section 6, that the set $\Sigma(A, B; T)$ has λ -measure satisfying

$$(4.3) \quad \lim_{T \rightarrow \infty} T \lambda(\Sigma(A, B; T)) = \frac{1}{2} \kappa \beta_{A,B} E_\lambda F.$$

Third, in Section 7, we will show that the set $\Sigma(A, B; T)$ can be represented approximately as a finite union of cylinder sets $\Sigma_m(\omega)$. This will be done in such a way that the lengths of the words defining the cylinder sets satisfy $m = (\log n)^2 = C'(\log T)^2$. It will then follow that the set $\{\omega : N_{A,B}(\omega) = k\}$ is (approximately) the set of all sequences $\omega \in \Sigma$ whose first n letters contain exactly k occurrences of one of the length- $2m+1$ sub-words

$$(4.4) \quad \omega_{-m} \omega_{-m+1} \cdots \omega_m$$

that define the cylinder sets $\Sigma_m(\omega)$.

Finally, in Section 8, we will use the results of steps 1, 2, and 3 to show that the number $N_{A,B}$ of crossings through arcs A, B on $\partial D(x; \alpha T^{-1})$ equals (with high probability) the number of length- $(\log T)^2$ blocks that contain one of the magic

subwords, and (using the regeneration theorem of Section 3.3) that these occurrence events are independent small-probability events. Furthermore, we will show that for distinct pairs (A_i, B_i) of boundary arcs the counts N_{A_i, B_i} are (approximately) independent. The desired result (4.2) will then follow from the Poisson convergence criterion of Section 2.3.

The strategy just outlined is easily adapted to Theorem 1.3. Fix distinct points $x, x' \in S$. For any pair A, B of non-overlapping boundary arcs of $\partial B(0, \alpha)$, denote by $N_{A, B}(\omega)$ and $N'_{A, B}(\omega)$ the numbers of geodesic arcs in the collections $I_n(x)$ and $I_n(x')$, respectively, that cross the target regions $D(x; \alpha T^{-1})$ and $D(x'; \alpha T^{-1})$ from arc A to arc B . To prove Theorem 1.3 it suffices to prove the following.

PROPOSITION 4.2. —

For any finite collections $\{(A_i, B_i)\}_{1 \leq i \leq r}$ and $\{(A'_i, B'_i)\}_{1 \leq i \leq r'}$ and any choice of nonnegative integers k_i, k'_i ,

$$(4.5) \quad \lim_{n \rightarrow \infty} \lambda \left\{ \omega : N_{A_i B_i}(\omega) = k_i \text{ and } N'_{A'_i B'_i}(\omega) = k'_i \quad \forall i \right\} \\ = \left(\prod_{i=1}^r \frac{1}{k_i!} (\kappa \beta_{A_i, B_i} / 2)^{k_i} e^{-\kappa \beta_{A_i, B_i} / 2} \right) \left(\prod_{i=1}^{r'} \frac{1}{k'_i!} (\kappa \beta_{A'_i, B'_i} / 2)^{k'_i} e^{-\kappa \beta_{A'_i, B'_i} / 2} \right).$$

Note: To avoid notational clutter, here and throughout Sections 5, 6, and 7 we will use the abbreviation κ for $\kappa_S = 1/\text{area}(S)$.

5. No Quick Entries or Re-entries of Small Disks

PROPOSITION 5.1. — *For any $\alpha > 0$ there exists $C < \infty$ such that if the geodesic ray γ has initial tangent vector chosen at random according to normalized Liouville measure, then for any $\varepsilon > 0$ the probability that γ enters the disk $D(x, \alpha T^{-1})$ before time εT is less than $C\varepsilon$.*

Proof. — For any unit vector $v \in T^1 S$, denote by $\tau(v)$ the smallest nonnegative time t (possibly $+\infty$) at which the geodesic ray $\gamma_t(v)$ with initial tangent vector v enters $D(x, \alpha T^{-1})$. Since any geodesic that enters $D(x, \alpha T^{-1})$ must spend at least $2\alpha T^{-1}$ units of time in the surrounding disk $D(x, 2\alpha T^{-1})$, we have, by the invariance of the Liouville measure,

$$\begin{aligned} \mu_L \{v : \tau(v) \leq \varepsilon T\} &\leq (2\alpha)^{-1} T \int_{T^1 S} \int_0^{\varepsilon T} \mathbf{1} \{ \gamma_t(v) \in D(x; 2\alpha T^{-1}) \} dt d\mu_L(v) \\ &= (2\alpha)^{-1} T \int_0^{\varepsilon T} \int_{T^1 S} \mathbf{1} \{ \gamma_t(v) \in D(x; 2\alpha T^{-1}) \} d\mu_L(v) dt \\ &= (2\alpha)^{-1} T \int_0^{\varepsilon T} \text{area} \left(D(x; 2\alpha T^{-1}) \right) dt / \text{area}(S) \\ &\leq C\varepsilon \end{aligned}$$

for a suitable $C < \infty$. □

PROPOSITION 5.2. — Let γ be a geodesic ray whose initial tangent vector is chosen at random according to normalized Liouville measure. If $f(T) = o(T)$ as $T \rightarrow \infty$ then the probability that γ enters (or begins in) $D(x, \alpha T^{-1})$ before time T and then re-enters within time $f(T)$ converges to 0 as $T \rightarrow \infty$.

The proof will be based on the following estimate Lemma 5.3.

LEMMA 5.3. — Fix $y \in D(x, 2\alpha T^{-1})$ and $0 < \beta < 1$, and define $R_T = R_T(y)$ to be the set of unit tangent vectors v based at y such that the geodesic ray with initial tangent vector v enters the ball $D(x, \alpha T^{-1})$ before time $f(T)$ after having first exited $D(x, 2\alpha T^{-1})$. There is a constant $K = K(\alpha) < \infty$ not depending on y such that for all $T \geq 1$ the Lebesgue measure of R_T satisfies

$$(5.1) \quad \text{Lebesgue}(R_T) \leq K f(T)/T.$$

The proof is deferred until after the proof of Proposition 5.2. Given the Lemma 5.3, Proposition 5.2 follows by an argument similar to that used in the proof of Proposition 5.1.

Proof of Proposition 5.2. — Denote by B_T the set of all unit tangent vectors v with base point in $D(x, \alpha T^{-1})$ such that the geodesic ray $\gamma_t(v)$ re-enters $D(x, \alpha T^{-1})$ before time $f(T)$ (after having first exited), and by B_T^* the set of all unit tangent vectors v with base point in the enlarged disk $D(x, 2\alpha T^{-1})$ such that the geodesic ray $\gamma_t(v)$ enters B_T before leaving $D(x, 2\alpha T^{-1})$. Clearly, $B_T \subset B_T^*$, and if $v \in B_T^*$, then the geodesic ray must spend at least αT^{-1} units of time in the disk $D(x, 2\alpha T^{-1})$ before exiting. Moreover, Lemma 5.3 implies that

$$\mu_L(B_T^*) \leq K T^{-1} f(T) \times \frac{\text{area}(D(x, 2\alpha T^{-1}))}{\text{area}(S)} \leq K' \alpha^2 T^{-3} f(T).$$

For any $v \in T^1(S)$ let $\tau^*(v)$ be the first time t that the geodesic ray $\gamma_t(v)$ enters the set B_T . Then by the invariance of Liouville measure,

$$\begin{aligned} \mu_L \{v : \tau^*(v) \leq T\} &\leq \alpha^{-1} T \int_{T^1 S} \int_0^T \mathbf{1} \{\gamma_t \in B_T^*\} dt d\mu_L(v) \\ &= \alpha^{-1} T \int_0^T \int_{T^1 S} \mathbf{1} \{\gamma_t \in B_T^*\} d\mu_L(v) dt \\ &= \alpha^{-1} T \int_0^T \mu_L(B_T^*) dt \\ &\leq K' \alpha^2 T^{-1} f(T) \longrightarrow 0 \quad \text{as } T \rightarrow \infty. \quad \square \end{aligned}$$

Proof of Lemma 5.3. — Let $\tilde{S}$ be the universal cover of S , viewed as the (open) unit disk $\mathbb{D}$ endowed with Riemannian metric $\tilde{\varrho}$, the natural lift of the Riemannian metric ϱ on S . The metric $\tilde{\varrho}$ is invariant by the fundamental group $\pi_1(S)$. Furthermore, the action of $\pi_1(S)$ on $\tilde{S}$ is discrete, and there is a *fundamental polygon* $\mathcal{P}$ for this action, bounded by geodesic segments, such that $\tilde{S}$ is tiled by the isometric images $g\mathcal{P}$, where g ranges over $\pi_1(S)$. Since the surface S is compact, the fundamental polygon $\mathcal{P}$ can be chosen so that it has finite diameter δ . Fix pre-images $\tilde{x}, \tilde{y} \in \tilde{S}$ of the points $x, y \in S$ in such a way that $\tilde{x} \in \mathcal{P}$ and $\tilde{y} \in D(\tilde{x}, 2\alpha T^{-1})$; then for all sufficiently large T the pre-image of the disk $D(x, \alpha T^{-1})$ is the disjoint union of

isometric disks $D(g\tilde{x}, \alpha T^{-1})$ where g ranges over $\pi_1(S)$. Clearly, since these disks are non-overlapping, only finitely many can intersect the fundamental polygon.

The set $R_T(y)$ lifts to a set $\tilde{R}_T(\tilde{y})$ of the same Lebesgue measure in the unit tangent space $T_{\tilde{y}}^1(\tilde{S})$; this lift $\tilde{R}_T(\tilde{y})$ contains all direction vectors $v \in T_{\tilde{y}}^1(\tilde{S})$ such that the geodesic ray $\tilde{\gamma}_t(v)$ in $\tilde{S}$ with initial tangent vector v intersects one of the balls $D(g\tilde{x}, \alpha T^{-1})$ with $g \neq 1$ at distance $\leq f(T)$ from $\tilde{y}$. To estimate the size of $\tilde{R}_T(\tilde{y})$, we decompose it by grouping the target disks $D(g\tilde{x}, \alpha T^{-1})$ in concentric shells by distance from the point $\tilde{y}$: thus, in particular,

$$\tilde{R}_T(\tilde{y}) \subset \bigcup_{m=1}^{\lfloor f(T) + 1 \rfloor / (3\delta)} \bigcup_{g \in \mathcal{A}_m} \Theta_{g,T}(\tilde{y})$$

where $\Theta_{g,T}(\tilde{y})$ is the set of all unit tangent vectors $v \in T_{\tilde{y}}^1(\tilde{S})$ such that the geodesic ray $\tilde{\gamma}_t(v)$ intersects the disk $D(g\tilde{x}, \alpha T^{-1})$, and

$$\mathcal{A}_m := \{g \in \pi_1(S) : 3(m-1)\delta \leq \text{distance}(g\tilde{x}, \tilde{y}) < 3m\delta\}.$$

(Recall that δ is the diameter of the fundamental polygon. Consequently, every point on the circle $\Gamma_{3m\delta}$ of radius $3m\delta$ centered at $\tilde{y}$ is within distance $5\delta/2$ of $g\tilde{x}$ for some $g \in \mathcal{A}_m$.) To prove inequality (5.1) it will suffice to prove that for some constant K independent of m, T and the choice of $\tilde{y} \in D(\tilde{x}, \alpha T^{-1})$,

$$(5.2) \quad \sum_{g \in \mathcal{A}_m} \text{Lebesgue}(\Theta_{g,T}) \leq KT^{-1}.$$

For each unit tangent vector $v \in T_{\tilde{y}}^1\tilde{S}$ and each real $t > 0$, define the *expansion factor* $\eta_t(v)$ for the geodesic flow at time t in direction v to be the amount by which the exponential map $\exp_{\tilde{y}}$ expands distances at the tangent vector tv in the direction $w = w(v) \in T^1S$ orthogonal to v , that is,

$$\eta_t(v) = \|(d\exp_{\tilde{y}}(tv))w\|$$

where $w = w(y) \in T_{\tilde{y}}^1\tilde{S}$ is the unit vector orthogonal to v (the choice of sign is irrelevant). Note that since our surface is compact, this quantity is bounded away from both 1 and ∞ (since the curvature is bounded away from zero). Thus, if $\Gamma_t = \Gamma_t(\tilde{y})$ is the circle of radius t centered at $\tilde{y}$ in $\tilde{S}$, then

$$(5.3) \quad \text{circumference}(\Gamma_t) = \int_{T_{\tilde{y}}^1\tilde{S}} \eta_t(v) d \text{Lebesgue}(v),$$

and more generally, for any interval $\Theta \subset T_{\tilde{y}}^1\tilde{S}$,

$$(5.4) \quad \text{arc-length}(\exp_{\tilde{y}}(t\Theta)) = \int_{\Theta} \eta_t(v) d \text{Lebesgue}(v).$$

CLAIM 1. — *There exists a constant $C < \infty$ not depending on the choice of $\tilde{y} \in \tilde{S}$ such that for any two unit vectors $v_1, v_2 \in T_{\tilde{y}}^1\tilde{S}$ satisfying the condition $\text{distance}(\tilde{\gamma}_t(v_1), \tilde{\gamma}_t(v_2)) \leq 3\delta$, any $t > 3\delta$, and any $0 \leq s \leq 3\delta$,*

$$(5.5) \quad C^{-1} \leq \frac{\eta_t(v_1)}{\eta_{t-s}(v_2)} \leq C.$$

Before proving the claim, we show how it implies inequality (5.2). For each deck transformation $g \in \mathcal{A}_m$, let $\Theta_{g,m}^*$ be the set of all direction vectors v based at $\tilde{y}$ for which the geodesic ray $\tilde{\gamma}_t(v)$ approaches the point $g\tilde{x}$ at least as closely as it approaches any other $g'\tilde{x}$, where $g' \in \mathcal{A}_m$. These sets overlap either in isolated points or not at all, and their union is the entire set $T_{\tilde{y}}^1 \tilde{S}$. Thus, the sets Θ_m^* , where $g \in \mathcal{A}_m$, form (up to a set of measure 0) a partition of $T_{\tilde{y}}^1 \tilde{S}$. Furthermore, since the action of $\pi_1(S)$ on the universal cover $\tilde{S}$ is discrete, there exists an integer $k \geq 1$ such that none of the sets Θ_m^* intersects more than k of the arcs $\Theta_{g,T}$.

Claim 1, together with the identity (5.4), implies that for a suitable constant $1 < C_1 < \infty$ not depending on $T, \tilde{y}, m$, or $g \in \mathcal{A}_m$,

$$\begin{aligned} \text{Lebesgue}(\Theta_{g,T}) &\leq C_1 T^{-1} \eta_{3m\delta}(v_g)^{-1} \quad \text{and} \\ \text{Lebesgue}(\Theta_{g,m}^*) &\geq C_1^{-1} \eta_{3m\delta}(v_g)^{-1}. \end{aligned}$$

where $v_g \in T_{\tilde{y}}^1 \tilde{S}$ is the unique direction such that the geodesic ray $\tilde{\gamma}_t(v_g)$ goes through the point $g\tilde{x}$. Since each arc $\Theta_{g,T}$ is contained in the union (over $g' \in \mathcal{A}_m$) of the sets $\Theta_{g',m}^*$, and since no $\Theta_{g',m}^*$ intersects more than k of the arcs $\Theta_{g,T}$, it follows that

$$\sum_{g \in \mathcal{A}_m} \text{Lebesgue}(\Theta_{g,T}) \leq k C_1^2 T^{-1} \sum_{g \in \mathcal{A}_m} \text{Lebesgue}(\Theta_{g,m}^*).$$

The desired result (5.2) now follows, because the sets $\Theta_{g,m}^*$ partition (up to a set of measure 0) the unit tangent space $T_{\tilde{y}}^1 \tilde{S}$.

Proof of Claim 1. — The expansion factor $\eta_t(v)$ can be calculated by integrating the infinitesimal expansion rates along the geodesic:

$$\begin{aligned} \eta_t(v) &= \exp \left\{ \int_0^t \zeta_s(v) ds \right\} \quad \text{where} \\ \zeta_t(v) &= \log \frac{d}{ds} \left(\left\| d \exp_{\tilde{\gamma}_t(v)}(s \tilde{\gamma}'_t(v)) w_t \right\| \right)_{s=0} \end{aligned}$$

and $w_t \in T_{\tilde{\gamma}_t(v)}^1 \tilde{S}$ is the unit vector tangent to the circle C_t (or equivalently, orthogonal to the direction $\tilde{\gamma}'_t(v)$ of the geodesic) at the point $\tilde{\gamma}_t(v)$. Because the curvature of the Riemannian metric ϱ is everywhere negative, the infinitesimal expansion rate $\zeta_t(v)$ is strictly positive. Moreover, because the Riemannian structure is C^∞ , so is the dependence of $\zeta_t(v)$ on both t and $v \in T^1 \tilde{S}$. Consequently, there exist constants $C_2 < \infty$ and $a > 0$ such that for any two unit vectors $v_1, v_2 \in T_{\tilde{y}}^1 \tilde{S}$ and any $t > 0$,

$$\begin{aligned} \text{distance}(\tilde{\gamma}_t(v_1), \tilde{\gamma}_t(v_2)) &\leq 3\delta \\ \implies \text{distance}(\tilde{\gamma}_{t-s}(v_1), \tilde{\gamma}_{t-s}(v_2)) &\leq C_2 e^{-as} \quad \text{for all } 0 \leq s \leq t. \end{aligned}$$

Therefore, the integral formula for the expansion rate $\eta_t(v)$ implies that for an appropriate constant $C_3 < \infty$,

$$\begin{aligned} \text{distance}(\tilde{\gamma}_t(v_1), \tilde{\gamma}_t(v_2)) &\leq 1 \\ \implies C_3^{-1} &\leq \frac{\eta_t(v_1)}{\eta_{t-s}(v_2)} \leq C_3 \quad \text{for all } t > 3\delta \quad \text{and} \quad 0 \leq s \leq 3\delta. \quad \square \end{aligned}$$

6. Measure of the Crossing Sets

Fix $x \in S$ and $\alpha > 0$. Let A, B be any two disjoint closed arcs, each with nonempty interior, on the boundary of the ball $B(0, \alpha)$ in the unit tangent space $T_x^1 S$. Recall that we have agreed to identify the arcs A, B with their images A^T, B^T on the closed curve $\partial D(x, \alpha T^{-1})$ under the scaled exponential mapping $v \mapsto \exp_x(v/T)$. Recall also that $\Sigma(A, B; T)$ is the set of all sequences $\omega \in \Sigma$ such that the intersection of the geodesic segment $p \circ \pi(\mathcal{F}_\omega)$ with the neighborhood $D(x, \alpha T^{-1})$ is a geodesic segment that either coincides with or extends to a geodesic crossing of $D(x, \alpha T^{-1})$ from boundary arc A^T to boundary arc B^T .

PROPOSITION 6.1. — For all A, B , and $\alpha > 0$,

$$(6.1) \quad \lim_{T \rightarrow \infty} T \lambda(\Sigma(A, B; T)) = \frac{1}{2} \kappa_{A, B} E_\lambda F$$

where $\beta_{A, B}$ is as defined by equation (2.2) (with $D = B(0, \alpha)$) and $\kappa = 1/\text{area}(S)$.

Proof. — By definition, if $\omega \in \Sigma(A, B; T)$ then there exist unique times $s_A(\omega) < s_B(\omega) < F(\omega)$ such that the segment $\phi([s_A(\omega), s_B(\omega)])$ projects via $p \circ \pi$ to a geodesic crossing of $D(x, \alpha T^{-1})$ from boundary arc A^T to boundary arc B^T . It is possible that $s_A(\omega) < 0$; this will occur if and only if $p \circ \pi(\omega, 0)$ is an interior point of $D(x, \alpha T^{-1})$. Now the surface area of $D(x, \alpha T^{-1})$ is of order T^{-2} ; hence, since λ^* is the pullback of the normalized Liouville measure μ_L , the λ -measure of the set of $\omega \in \Sigma(A, B; T)$ such that $s_A(\omega) < 0$ is also of order T^{-2} . Consequently, in proving (6.1) we may ignore the contribution of the set

$$\Sigma(A, B; T)_- = \{\omega \in \Sigma(A, B; T) : s_A(\omega) < 0\}.$$

Set $\Sigma(A, B; T)_+ = \Sigma(A, B; T) \setminus \Sigma(A, B; T)_-$.

Denote by $\Upsilon(A, B; T)$ the set of all $u \in T^1 S$ that are tangents to geodesic segments from arc A^T to arc B^T . This set nearly coincides with the projections of those points $(\omega, s) \in \Sigma_F$ such that $\omega \in \Sigma(A, B; T)$ and $0 \leq s_A(\omega) < s < s_B(\omega)$, the difference being accounted for by the set $\Sigma(A, B; T)_-$. Consequently, by equation (3.3),

$$\begin{aligned} \lambda(\Sigma(A, B; T)) &= \int_{\Sigma(A, B; T)} \frac{s_B - s_A}{s_B - s_A} d\lambda \\ &= \int_{\pi^{-1}\Upsilon(A, B; T)} \frac{1}{s_B(\omega) - s_A(\omega)} d\lambda^*(\omega, s) \times \int_\Sigma F d\lambda + O(T^{-2}) \\ &= \int_{\Upsilon(A, B; T)} \frac{1}{\tau(u)} d\mu_L(u) \times \int_\Sigma F d\lambda + O(T^{-2}), \end{aligned}$$

where $\tau(u)$ is the length of the geodesic segment from A to B on which u lies, for any $u \in \Upsilon(A, B; T)$.

Now we exploit the defining property of the Liouville measure μ_L , specifically, that locally μ_L is the product of normalized surface area with the Haar measure on the circle. For large T , the exponential mapping $v \mapsto \exp\{v/T\}$ maps the ball $B(0, \alpha)$ in the tangent space $T_x S$ onto $D(x, \alpha T^{-1})$ nearly isometrically (after scaling by the factor T^{-1}), so rescaled surface area on $D(x, \alpha T^{-1})$ is nearly identical with the pushforward of Lebesgue measure on $B(0, \alpha)$, scaled by T^{-2} . Furthermore, the inverse

images of geodesic segments across $D(x; \alpha T^{-1})$ are nearly straight line segments crossing $B(0; \alpha)$; those that cross from arc A^T to arc B^T in $\partial \mathcal{D}(x; \alpha T^{-1})$ will pull back to straight line segments from arc A to arc B in $\partial B(0; \alpha)$. These can be parametrized by the angle at which they meet the x -axis, as in Figure 2.1; for each angle θ , the integral of $1/\text{length}$ over the region in $B(0; \alpha)$ swept out by line segments crossing from arc A to arc B at angle θ is $\psi(\theta)$, as in Figure 2.1 (where the convex region is now $B(0; \alpha)$). Therefore, as $T \rightarrow \infty$,

$$\int_{\Upsilon(A, B; T)} \frac{1}{\tau(u)} d\mu_L(u) \sim T^{-1} \frac{1}{2\pi \text{area}(S)} \int_{-\pi/2}^{\pi/2} \psi(\theta) d\theta = T^{-1} \kappa \beta_{A, B}/2.$$

□

Similar calculations can be used to show there is vanishingly small probability that one of the first n geodesic segments $p \circ \pi(\mathcal{F}_{\sigma^i \omega})$ will hit both $D(x; \alpha T^{-1})$ and $D(x'; \alpha T^{-1})$, where $x \neq x'$ are distinct point of S . Define $H(x, x'; \alpha T^{-1})$ to be the set of all $\omega \in \Sigma$ such that the vertical fiber $\mathcal{F}_\omega$ over $(\omega, 0)$ in Σ_F projects to a geodesic segment that intersects both $D(x; \alpha T^{-1})$ and $D(x'; \alpha T^{-1})$.

PROPOSITION 6.2. — *For any two distinct points $x, x' \in S$ and each $\alpha > 0$,*

$$\lim_{T \rightarrow \infty} T \lambda \left(H \left(x, x'; \alpha T^{-1} \right) \right) = 0.$$

Proof. — Assume that T is sufficiently large that the closed disks $\bar{\mathcal{D}}(x; 2\alpha T^{-1})$ and $\bar{\mathcal{D}}(x'; 2\alpha T^{-1})$ do not intersect. For any ω such that the fiber $\mathcal{F}_\omega$ projects to a geodesic segment that enters $D(x; 2\alpha T^{-1})$ there will be unique times $0 < s_0(\omega) < s_1(\omega) < F(\omega)$ of entry and exit (except, as in the proof of Proposition 6.1, for a set of size $O(T^{-2})$); for those ω such that the projection of $\mathcal{F}_\omega$ enters the smaller disk $D(x; \alpha T^{-1})$, the sojourn time $s_1(\omega) - s_0(\omega)$ will be at least αT^{-1} .

Denote by $\Upsilon(x, x'; \alpha T^{-1})$ the set of all tangent vectors $u \in T^1 S$ based at points in $D(x; \alpha T^{-1})$ such that u lies on the directed geodesic segment $\pi(\mathcal{F}_{\sigma^i \omega})$ for some sequence $\omega \in H(x, x'; \alpha T^{-1})$. Since x and x' are distinct points of S , the neighborhoods $D(x; \alpha T^{-1})$ and $D(x'; \alpha T^{-1})$ are separated by at least $\text{dist}(x, x')/2$ (for large T), so there is a constant $C = C(x, x', \alpha) < \infty$ such that for every point $y \in D(x, \alpha T^{-1})$ the set of angles θ such that $(y, \theta) \in \Upsilon(x, x'; \alpha T^{-1})$ has Lebesgue measure less than CT^{-1} . Now

$$\begin{aligned} \lambda(H(x, x'; \alpha T^{-1})) &= \int_{H(x, x'; \alpha T^{-1})} \frac{s_1 - s_0}{s_1 - s_0} d\lambda \\ &= \int_{\pi^{-1}\Upsilon(x, x'; \alpha T^{-1})} \frac{1}{s_1(\omega) - s_0(\omega)} d\lambda^*(\omega, s) \times E_\lambda F \\ &= \int_{\Upsilon(x, x'; \alpha T^{-1})} \frac{1}{\tau(u)} dL(u) \times E_\lambda F \\ &\leq T^{-1} L \left(\Upsilon \left(x, x'; \alpha T^{-1} \right) \right) E_{\lambda F} \end{aligned}$$

where $\tau(u)$ is the crossing time of $D(x; 2\alpha T^{-1})$ by the geodesic with initial tangent vector u . Using once again the fact that (normalized) Liouville measure is the product of normalized hyperbolic area with Lebesgue angular measure, we see that

for a suitable constant $C' < \infty$,

$$L\left(\Upsilon\left(x, x'; \alpha T^{-1}\right)\right) \leq C'(\alpha T^{-1})^2 \times CT^{-1};$$

thus, $\lambda(H(x, x'; \alpha T^{-1})) = O(T^{-2})$. $\square$

7. Decomposition of the Events $N_{A,B} = k$

In this section we show that the events $\{\omega : N_{A,B}(\omega) = k\}$ can be approximated by sets consisting of those sequences $\omega \in \Sigma$ whose first n letters contain exactly k occurrence of certain “magic subwords” each of length $m = (\log n)^2 (\approx \log T)^2$. As in Section 6, let A, B be any two disjoint closed arcs, each with nonempty interior, on the boundary of the ball $B(0, \alpha)$ in the unit tangent space $T_x S$. We identify the neighborhood $D(x, \alpha T^{-1})$ in $S (= \mathcal{P})$ with the ball $B(0, \alpha)$ via the scaled exponential mapping (cf. equation (1.2)), so the arcs A, B are identified with arcs in $\partial D(x, \alpha T^{-1})$, denoted by A^T, B^T , whose arc-lengths are roughly proportional to T^{-1} . Recall that $N_{A,B}(\omega)$ is the number of crossings of the neighborhood $D(\omega, \alpha T^{-1})$ from boundary arc A^T to boundary arc B^T by the geodesic segment

$$p \circ \pi \left(\bigcup_{i=0}^{n-1} \mathcal{F}_{\sigma^i \omega} \right),$$

where $n = \lceil T/E_\lambda F \rceil$, π is the map from the suspension space Σ_F to $T^1(S)$, and p is the natural projection down from $T^1(S)$ to S . Recall also (Section 4) that $\Sigma(A, B; T)$ is the set of all sequences $\omega \in \Sigma$ such that the geodesic segment $p \circ \pi(\mathcal{F}_\omega)$ intersects $D(x, \alpha T^{-1})$ in a geodesic segment with terminal endpoint in the boundary arc B^T that extends to a geodesic crossing from boundary arc A^T to boundary arc B^T .

LEMMA 7.1. — *The set $\{\omega : N_{A,B} = k\}$ differs from the set*

$$\left\{ \omega \in \Sigma : \sum_{i=0}^{n-1} \mathbf{1}_{\Sigma(A, B; T)}(\sigma^i \omega) = k \right\}$$

by a set of λ -measure tending to 0 as $T \rightarrow \infty$.

Proof. — The symmetric difference of the two sets is contained in the set of all $\omega \in \Sigma$ such that either (a) $\omega \in \Sigma(A, B; T)$ or $\sigma^{n-1}\omega \in \Sigma(A, B; T)$, or (b) at least one of the geodesic segments $p \circ \pi(\mathcal{F}_{\sigma^i \omega})$, where $0 \leq i < n$, makes more than one visit to the neighborhood $D(x; \alpha T^{-1})$. Propositions 5.1 and 5.2 ensure that this set has measure $\rightarrow 0$ as $T \rightarrow \infty$. $\square$

Next, recall that the sequence space Σ is equipped with the metric $d(\omega, \omega') = e^{-n(\omega, \omega')}$, where $n(\omega, \omega')$ is the minimum nonnegative integer j such that the sequences ω, ω' differ in the $\pm j$ entry, and that the suspension space Σ_F inherits from d an induced “taxicab” metric satisfying the inequality (3.2). Cylinder sets $\Sigma_m(\omega)$ are open balls in Σ relative to the metric d (cf. equation (3.1)). Since the semi-conjugacy $\pi : \Sigma_F \rightarrow T^1 S$ is Hölder, it follows that if $\omega, \omega' \in \Sigma$ are at distance $< \varepsilon$, then the geodesics $p \circ \pi(\phi_t(\omega))$ and $p \circ \pi(\phi_t(\omega'))$ remain at distance $< e^{\max F} \varepsilon$ for all $|t| \leq \max_\Sigma F$. Thus, if one of the geodesic segments crosses from arc A^T to B^T without coming sufficiently near one of the endpoints of either A^T or B^T , then so

will the other; and similarly, if one stays sufficiently far away from the arcs A^T, B^T then so will the other. In particular, if

$$(7.1) \quad m = (\log n)^2 (\approx (\log T)^2),$$

then for every $\omega' \in \Sigma_m(\omega)$ the geodesic segments $p \circ \pi(\phi_t(\omega))_{|t| \leq \max F}$ and $p \circ \pi(\phi_t(\omega'))_{|t| \leq \max F}$ remain at distance less than $n^{-C \log n}$ for a suitable constant $C > 0$, and hence, for every $\omega \in \Sigma$ one of the following will hold:

- (i) $\Sigma_m(\omega) \subset \Sigma(A, B; T)$;
- (ii) $\Sigma_m(\omega) \subset \Sigma(A, B; T)^c$; or
- (iii) for every $\omega' \in \Sigma_m(\omega)$ the geodesic segment $p \circ \pi(\phi_t(\omega'))_{|t| \leq \max F}$ will pass within distance $C'n^{-C \log n}$ of one of the endpoints of arc A^T or arc B^T .

PROPOSITION 7.2. — *For each pair A, B of non-overlapping closed arcs of $\partial B(0, \alpha)$ and each $T \geq 1$ there exist finite subsets $\mathcal{J}_1 \subset \mathcal{J}_2$ of Σ such that*

- (A) *for each $\omega \in \mathcal{J}_1$ the cylinder set $\Sigma_m(\omega)$ is of type (i);*
- (B) *for each $\omega \notin \mathcal{J}_2$ the cylinder set $\Sigma_m(\omega)$ is of type (ii); and*
- (C) *the set $\cup_{\omega \in \mathcal{J}_2 \setminus \mathcal{J}_1} \Sigma_m(\omega)$ has λ -measure less than $o(n^{-r})$ for all $r > 0$.*

Proof. — The sets $\mathcal{J}_1$ and $\mathcal{J}_2 \setminus \mathcal{J}_1$ are gotten by selecting representatives of each cylinder $\Sigma_m(\omega)$ of type (i) and type (iii), respectively. What must be proved is assertion (C).

By construction, for every ω' not in $\cup_{\omega \in \mathcal{J}_2} \Sigma_m(\omega)$ the geodesic segment

$$p \circ \pi(\phi_t(\omega, 0))_{t \leq \max F}$$

must pass within distance $C'n^{-C \log n} = C''T^{-C'' \log T}$ of one of the four endpoints of A^T or B^T . Proposition 5.1 implies that the normalized Liouville measure of the set of geodesic rays that enter one of these four regions by time $\max F$ is of order $O(T^{-C'' \log T}) = O(n^{-C \log n})$. $\square$

DEFINITION 7.3. — *Given arcs A, B as in Proposition 7.2 and $T \geq 1$, define the magic subwords for the triple $(A, B; T)$ to be the words*

$$\omega_{-m}\omega_{-m+1} \cdots \omega_m$$

where $\omega \in \mathcal{J}_1$.

COROLLARY 7.4. — *The symmetric difference between the sets $\{\omega : N_{A, B} = k\}$ and the set of $\omega \in \Sigma$ with exactly k occurrences of one of the magic subwords in the segment $\omega_1\omega_2 \cdots \omega_n$ has λ -measure $\rightarrow 0$ as $T \rightarrow \infty$.*

Remark 7.5. — The set of magic subwords for a particular value of T will in general have no clear relationship to the magic subwords for a different value of T .

PROPOSITION 7.6. — *For each T let $\mathcal{M} = \mathcal{M}_T$ be the set of magic subwords for a fixed pair A, B of boundary arcs and fixed $\alpha > 0$. Then*

$$(7.2) \quad \lim_{T \rightarrow \infty} T \lambda\{\omega : (\omega_{-m}\omega_{-m+1} \cdots \omega_m) \in \mathcal{M}\} = \frac{1}{2} \kappa \beta_{A, B} E_\lambda F$$

where $\beta_{A, B}$ is defined by equation (2.2).

Proof. — This follows directly from Propositions 7.2 and 6.1. $\square$

COROLLARY 7.7. — *If $\omega \in \Sigma$ is chosen randomly according to λ , then the probability that the initial segment $\omega_1\omega_2 \cdots \omega_{\lfloor \log T \rfloor^\kappa}$ contains a magic subword converges to 0 as $T \rightarrow \infty$. Similarly, the probability that the segment $\omega_1\omega_2 \cdots \omega_n$ contains magic subwords separated by fewer than $(\log n)^\kappa$ letters converges to zero as $n \rightarrow \infty$.*

Proof. — This follows directly from Propositions 5.1 and 5.2. $\square$

8. Proof of Propositions 4.1–4.2

Proof of (4.2) for $r = 1$. — Consider first the case $r = 1$. In this case we are given a single pair (A, B) of non-overlapping boundary arcs of $\partial B(0, \alpha)$; we must show that for any integer $k \geq 0$,

$$(8.1) \quad \lim_{n \rightarrow \infty} \lambda\{\omega \in \Sigma : N_{A,B}(\omega) = k\} = \frac{(\kappa\beta_{A,B}/2)^k}{k!} e^{-\kappa\beta_{A,B}/2},$$

where $\beta_{A,B}$ is defined by equation (2.2). Recall that $N_{A,B}(\omega)$ is the number of geodesic segments in the collection $I_n(\omega)$ that cross the target disk $D(x; \alpha T^{-1})$ from arc A to arc B . By Corollary 7.4, $N_{A,B}(\omega)$ is well-approximated by the number $N'_{A,B}$ of magic subwords in the word $\omega_1\omega_2 \cdots \omega_n$; in particular, for any $k \geq 0$, the symmetric difference between the events $\{N_{A,B} = k\}$ and $\{N'_{A,B} = k\}$ has λ -measure tending to 0. Consequently, it suffices to prove that (8.1) holds when $N_{A,B}$ is replaced by $N'_{A,B}$.

Recall (Section 3.3) that any Gibbs process is the natural projection of a list process. Thus, on some probability space there exists a sequence $W_1, W_2, W_3, \dots$ of independent random words of random lengths τ_i , such that the infinite sequence obtained by concatenating $W_1, W_2, W_3, \dots$ has distribution λ , that is, for any Borel subset B of Σ^+ ,

$$P\{W_1 \cdot W_2 \cdot W_3 \cdots \in B\} = \lambda(B).$$

All but the first word W_1 have the same distribution, and the lengths τ_i have exponentially decaying tails (cf. inequality (3.8)). Since the magic subwords are of length $[\log n]^2$, any occurrence of one will typically straddle a large number of consecutive words in the sequence W_i . Thus, to enumerate occurrences of magic subwords, we shall break the sequence $\{W_i\}_{i \geq 1}$ into blocks of length $m = [\log n]^3$, and count magic subwords block by block. Set

$$\begin{aligned} \widetilde{W}_1 &= W_1 W_2 \cdots W_m, \\ \widetilde{W}_2 &= W_{m+1} W_{m+2} \cdots W_{2m}, \\ \widetilde{W}_3 &= W_{2m+1} W_{2m+2} \cdots W_{3m}, \end{aligned}$$

etc., and denote by $\tilde{\tau}_k = \sum_{i=mk-k+1}^{mk} \tau_i$ the length (in letters) of the word $\widetilde{W}_k$.

CLAIM 2. — *For each $C > E\tau_2$, there exists $\Lambda(C) > 0$ such that for any integer $k \geq 1$*

$$(8.2) \quad P\left\{\sum_{i=1}^k \tau_i \geq Ck\right\} \leq e^{-k\Lambda(C)},$$

and for all sufficiently large $C < \infty$,

$$(8.3) \quad \lim_{n \rightarrow \infty} P \left\{ \max_{k \leq n} \tilde{\tau}_k \geq Cm \right\} = 0.$$

The function $C \mapsto \Lambda(C)$ is convex and satisfies $\liminf_{C \rightarrow \infty} \Lambda(C)/C > 0$.

Proof of Claim 2. — These estimates follow from the exponential tail decay property (3.8) by standard results in the elementary large deviations theory, in particular, *Cramér's theorem* (cf. [DZ98, Section 2.2]) for sums of independent, identically distributed random variables with exponentially decaying tails. The block lengths $\tilde{\tau}_k$ are gotten by summing the lengths τ_i of their m constituent words W_i ; for all but the first block $\tilde{W}_1$, these lengths are i.i.d. and satisfy (3.8). Hence, Cramér's theorem guarantees⁽⁵⁾ the existence of a convex rate function $C \mapsto \Lambda(C)$ and constants $C' = C'(C) < \infty$ such that inequality (8.2) holds for all $k \geq 1$. Applying this inequality with $k = m = [\log n]^3$ yields

$$P \left\{ \sum_{i=2}^{m+1} \tau_i \geq Cm \right\} \leq e^{-m\Lambda(C)} = n^{-3\Lambda(C)}.$$

Cramér's theorem also implies that $\Lambda(C)$ grows at least linearly in C , so by taking C sufficiently large we can ensure that $\Lambda(C) \geq 2/3$, which makes the probability above smaller than n^{-2} . Since there are only n blocks, it follows that the probability that $\tilde{\tau}_k \geq Cm$ for one of them is smaller than n^{-1} . $\square$

CLAIM 3. — *The probability that a magic subword occurs in the concatenation of the first two blocks $\tilde{W}_1\tilde{W}_2$ converges to 0 as $T \rightarrow \infty$.*

Proof of Claim 3. — The event that one of the first two blocks has length $\geq C[\log n]^3$ can be ignored, by Claim 2. On the complementary event, an occurrence of a magic subword in $\tilde{W}_1\tilde{W}_2$ would require that the magic subword occurs in the first $2C[\log n]^3$ letters. By Corollary 7.7, the probability of this event tends to 0 as $T \rightarrow \infty$. $\square$

It follows from Claim 2 and Corollary 7.7 that with probability tending to 1 as $n \rightarrow \infty$, no block $\tilde{W}_k$ among the first n will contain more than one magic subword. On this event, then, the number $N_{A,B}$ of magic subwords that occur in the first n letters can be obtained by counting the number of blocks $\tilde{W}_k$ that contain magic subwords and then adding the number of magic subwords that straddle two consecutive blocks.

CLAIM 4. — *As $n \rightarrow \infty$, the probability that a magic subword straddles two consecutive blocks $\tilde{W}_k, \tilde{W}_{k+1}$ among the first $n/[\log n]^3$ blocks converges to 0.*

Proof of Claim 4. — A magic subword, since it has length $[\log n]^2$, can only straddle consecutive blocks $\tilde{W}_k, \tilde{W}_{k+1}$ if it begins in one of the last $[\log n]^2$ word W_i of the $m = [\log n]^3$ words that constitute $\tilde{W}_k$. The words W_i are i.i.d. (except for

⁽⁵⁾The length of the initial block has a different distribution than the subsequent blocks, because the first excursion of the list process has a different law than the rest. However, the length of the first excursion also has an exponentially decaying tail, by Proposition 3.3, so the upper bounds given by Cramér's theorem still apply.

W_1 , and by Claim 3 we can ignore the possibility that a magic subword begins in $\widetilde{W}_1\widetilde{W}_2$, so the probability that a magic subword begins in W_i does not depend on i . Since only $[\log n]^2$ of the $[\log n]^3$ words in each block $\widetilde{W}_k$ would produce straddles, it follows that the *expected* number of magic subwords in $\widetilde{W}_1\widetilde{W}_2\cdots\widetilde{W}_{n/m}$ is at least $[\log n]$ times the probability that a magic subword straddles two consecutive blocks. Therefore, the Claim 4 will follow if we can show that the expected number of magic subwords in $\widetilde{W}_1\widetilde{W}_2\cdots\widetilde{W}_{n/m}$ remains bounded as $T \rightarrow \infty$. Denote the number of such magic subwords by $N''_{A,B}$.

The number of letters in the concatenation $\widetilde{W}_1\widetilde{W}_2\cdots\widetilde{W}_{n/m}$ is $\sum_{i=1}^n \tau_i$, which by Claim 2 obeys the large deviation bound (8.2). Fix $K < \infty$, and let G be the event that $\sum_{i=1}^n \tau_i \leq nK$. On this event, $N''_{A,B}$ is bounded by the number of magic subwords in the first nK letters of the concatenation $W_1W_2\cdots$. Since the concatenation $W_1W_2\cdots$ is, by Proposition 3.3, a version of the Gibbs process associated with the Gibbs state λ , which by shift-invariance is stationary, it follows that the expected number of magic subwords in the first nK letters is $nK \times$ the probability that a magic subword begins at the very first letter of $W_1W_2\cdots$. But by Proposition 6.1, this probability is asymptotic to $T^{-1}\alpha\beta_{A,B}E_\lambda F$; thus, for large T ,

$$EN''_{A,B}\mathbf{1}_G \leq nKT^{-1}\alpha\beta_{A,B}E_\lambda F = K\alpha\beta_{A,B}.$$

It remains to bound the contribution to the expectation from the complementary event G^c . For this, we use the large deviation bound (8.2). On the event that $\sum_{i=1}^n \tau_i \leq n(K+k)$, the count $N''_{A,B}$ cannot be more than $n(K+k)$; hence,

$$EN''_{A,B}\mathbf{1}_{G^c} \leq \sum_{k=1}^{\infty} n(K+k)e^{-n\Lambda(K+k)}.$$

Since $\Lambda(C)$ grows at least linearly in C , this sum remains bounded provided K is sufficiently large. $\square$

Recall that $N'_{A,B}$ is the number of magic subwords in the first n letters of the sequence $\widetilde{W}_1\widetilde{W}_2\cdots$ obtained by concatenating the words in the regenerative representation. The blocks $\widetilde{W}_k$ are independent, and except for the first all have the same distribution, with common mean length $mE\tau_2$. Let $N^*_{A,B}$ be the number of magic subwords in the segment $\widetilde{W}_2\widetilde{W}_3\cdots\widetilde{W}_\nu$, where $\nu = \nu(n) = n/[mE\tau_2]$. By the central limit theorem, with probability approaching 1 the length $\sum_{i=1}^{\nu m} \tau_i$ of the segment $\widetilde{W}_2\widetilde{W}_3\cdots\widetilde{W}_\nu$ differs by no more than $\sqrt{n}\log n$ from n , and by the same argument as in the proof of Claim 4, the probability that a magic subword occur within the stretch of $2\sqrt{n}\log n$ letters surrounding the n th letter converges to 0. Thus, as $T \rightarrow \infty$,

$$P\{N'_{A,B} \neq N^*_{A,B}\} \rightarrow 0.$$

By Claim 2 and Corollary 7.7, with probability approaching 1 no block $\widetilde{W}_k$ will contain more than 1 magic subword, and by Claim 4 no magic subword will straddle two blocks $\widetilde{W}_k, \widetilde{W}_{k+1}$. Therefore, with probability $\rightarrow 1$,

$$N'_{A,B} = N^*_{A,B} = \sum_{k=2}^{\nu} Y(\widetilde{W}_k),$$

where $Y(\widetilde{W}_k)$ is the indicator of the event that the block $\widetilde{W}_k$ contains a magic subword. These indicators are independent, identically distributed Bernoulli random variables; by Proposition 7.6,

$$EY(\widetilde{W}_k)mE\tau_2 \sim T^{-1}\frac{1}{2}\kappa\beta E_\lambda F$$

and so

$$E \sum_{k=2}^{\nu} Y(\widetilde{W}_k) \longrightarrow \frac{1}{2}\kappa\beta_{A,B}.$$

Now Proposition 2.11 implies that for any integer $J \geq 0$,

$$P \left\{ \sum_{k=2}^{\nu} Y(\widetilde{W}_k) = J \right\} \longrightarrow \frac{(\kappa\beta_{A,B}/2)^J}{J!} e^{-\kappa\beta_{A,B}/2},$$

proving (8.1). $\square$

Proof of (4.2) for $r \geq 1$. — (Sketch) In general, given $r \geq 1$, we are given a set $\{A_i, B_i\}$ of pairwise non-overlapping boundary arcs of $\partial B(0, \alpha)$; we must show that the counts N_{A_i, B_i} converge *jointly* to independent Poissons with means $\alpha\beta_{A_i, B_i}$, respectively. The key to this is that the sets $\mathcal{M}_i$ of magic words for the different pairs (A_i, B_i) are pairwise disjoint, because the arcs A_i, B_i are non-overlapping (a geodesic segment crossing of $D(x; \alpha T^{-1})$ has unique entrance and exit points on $\partial D(x, \alpha T^{-1})$, so at most one of the pairs (A_i, B_i) can contain these).

By the same argument as in the case $r = 1$, the counts N_{A_i, B_i} can be replaced by the sums

$$N_{A_i, B_i}^* = \sum_{k=2}^{\nu} Y_i(\widetilde{W}_k)$$

where $Y_i(\widetilde{W}_k)$ is the indicator of the event that the block $\widetilde{W}_k$ contains a magic subword for the pair A_i, B_i . Since the sets $\mathcal{M}_i$ of magic subwords are non-overlapping, the vector of these sums follows a multinomial distribution; hence, by Proposition 2.12, the vector

$$(N_{A_i, B_i}^*)_{1 \leq i \leq r}$$

converges in distribution to the product of r Poisson distributions, with means $\frac{1}{2}\kappa\alpha\beta_{A_i, B_i}$. $\square$

Proof of Proposition 4.2. — The argument is virtually the same as that for the case $r \geq 2$ of Proposition 4.1; the only new wrinkle is that the sets $\mathcal{M}_i$ and $\mathcal{M}'_{i'}$ of magic words for the pairs (A_i, B_i) and $(A'_{i'}, B'_{i'})$ need not be disjoint, because it is possible for a geodesic segment across the fundamental polygon $\mathcal{P}$ to enter both $D(x, \alpha T^{-1})$ and $D(x', \alpha T^{-1})$. However, Proposition 6.2 implies that the expected number of such double-hits in the first n crossings of $\mathcal{P}$ converges to 0 as $T \rightarrow \infty$, and consequently the probability that there is even one double-hit tends to zero. Thus, the magic subwords for pairs $A'_{i'}, B'_{i'}$ that also occur as magic subwords for pairs A_i, B_i can be deleted without affecting the counts (at least with probability

$\rightarrow 1$ as $T \rightarrow \infty$), and so the counts N_{A_i, B_i} and $N'_{A'_{i'}, B'_{i'}}$ may be replaced by

$$N_{A_i, B_i}^* = \sum_{k=2}^{\nu} Y_i(\widetilde{W}_k) \quad \text{and} \\ N_{A'_{i'}, B'_{i'}}^{**} = \sum_{k=2}^{\nu} Y'_{i'}(\widetilde{W}_k)$$

where $Y_i(\widetilde{W}_k)$ and $Y'_{i'}$ are the indicators of the events that the block $\widetilde{W}_k$ contains a magic subword for the appropriate pair (with deletions of any duplicates). Since the adjusted sets of magic subwords are non-overlapping, the vector of these counts N_{A_i, B_i}^* and $N_{A'_{i'}, B'_{i'}}^{**}$ follows a multinomial distribution, and so the convergence (4.5) holds, by Proposition 2.12. $\square$

9. Global Statistics

In this section we show how Theorem 1.4, which describes the “global” statistics of the tessellation $\mathcal{T}_T$ induced by a random geodesic segment of length T , follows from the “local” description provided by Theorem 1.2 and the ergodicity of the Poisson line process with respect to translations. Theorem 1.2 and Proposition 2.7 (cf. also Corollary 2.9) imply that locally – in balls $D(x; \alpha T^{-1})$, where α is large – the empirical distributions of polygons, their angles and side lengths (after scaling by T) stabilize as $T \rightarrow \infty$. Since this is true in neighborhoods of all points $x \in S$, it is natural to expect that these empirical distributions also converge globally. To prove this, we must show that in those small regions of S where empirical distributions behave atypically the counts are not so large as to disturb the global averages. The key is the following proposition, which limits the numbers of polygons, edges, and vertices in $\mathcal{T}_T$.

PROPOSITION 9.1. — *Let $f = f_T$, $v = v_T$, and $e = e_T$ be the number of polygons, vertices and edges in the tessellation $\mathcal{T}_T$. With probability one, as $T \rightarrow \infty$,*

$$(9.1) \quad \lim_{T \rightarrow \infty} v_T/T^2 = \kappa/\pi,$$

$$(9.2) \quad \lim_{T \rightarrow \infty} e_T/T^2 = (2\kappa)/\pi, \quad \text{and}$$

$$(9.3) \quad \lim_{T \rightarrow \infty} f_T/T^2 = \kappa/\pi.$$

Moreover, there exists a (nonrandom) constant $C = C_S < \infty$ such that for every tessellation $\mathcal{T}_T$ induced by a geodesic segment of length T ,

$$(9.4) \quad v_T + e_T + f_T \leq CT^2.$$

For the proof we will need to know that multiple intersection points (points of S that a geodesic ray passes through more than twice) do not occur in typical geodesics. We have the following:

LEMMA 9.2. — *For almost every unit tangent vector $v \in T^1S$, there are no multiple intersection points on the geodesic ray $(\gamma_t(v))_{t \geq 0}$.*

Proof. — Suppose $v \in T^1S$ gives rise to triple intersection. Let γ denote a lift of the geodesic ray $(\gamma_t(v))_{t \geq 0}$ to the universal cover $\mathbb{H}^2$, we have that there must be deck transformations A, B so that the geodesic rays $A\gamma$ and $B\gamma$ have a triple intersection. In [EJM91], it is shown that the set of such geodesics is a positive codimension subvariety for any fixed A, B , and therefore, a set of measure 0. Taking the (countable) union over all possible pairs A, B , we have our result. $\square$

Proof of Proposition 9.1. — The number v_T of vertices is the number of self-intersections of the random geodesic segment $\gamma_T := (\gamma_t(\cdot))_{0 \leq t \leq T}$ (unless one counts the beginning and end points of γ_T as vertices, in which case the count is increased by 2). It is an easy consequence of Birkhoff's ergodic theorem (see [Lal14, Section 2.3] for the argument, but beware that [Lal14] seems to be off by a factor of 4 in his calculation of the limit) that the number of self-intersections satisfies (9.1). Following is a brief resume of the argument.

Fix $\epsilon > 0$ small, and partition the segment γ_T into non-overlapping geodesic segments γ_T^i of length ϵ (if necessary, extend or delete the last segment; this will not change the self-intersection count by more than $O(T)$). If ϵ is smaller than the injectivity radius then

$$(9.5) \quad v_T = \sum_{i \neq j} \sum \mathbf{1}(\gamma_T^i \cap \gamma_T^j \neq \emptyset)$$

is the number of pairs (i, j) such that γ_T^i and γ_T^j cross. Birkhoff's theorem implies that for each i , the fraction of indices j such that γ_T^j crosses γ_T^i converges, as $T \rightarrow \infty$, to the normalized Liouville measure of that region R_ϵ of T^1S where the geodesic flow will produce a ray that crosses γ_T^i by time ϵ . This implies that the limit on the left side of (9.1) exists. To calculate the limit, let $\epsilon \rightarrow 0$: if $\epsilon > 0$ is small, then for each angle θ the set of points $x \in S$ such that $(x, \theta) \in R_\epsilon$ is approximately a rhombus of side ϵ with interior angle θ . Integrating the area of this rhombus over θ , one obtains a sharp asymptotic approximation to the normalized Liouville measure of R_ϵ :

$$L(R_\epsilon) \sim 2\epsilon^2 \int_0^\pi \sin \theta d\theta / (2\pi \text{ area}(S)) = 2\epsilon^2 \kappa / \pi.$$

Since the number of terms in the sum (9.5) is $\frac{1}{2}[T/\epsilon^2]$, it follows that $v_T/T^2 \rightarrow \kappa/\pi$.

The limiting relations (9.2) and (9.3) follow easily from (9.1). With probability one, the geodesic segment γ_T has no multiple intersection points, by Lemma 9.2. Consequently, as one traverses the segment γ_T from beginning to end, one visits each vertex twice, and immediately following each such visit encounters a new edge of $\mathcal{T}_T$ (except for the initial edge), so $e_T = 2v_T \pm 2$, and hence (9.2) follows from (9.1). Finally, by Euler's formula, $v - e + f = -\chi(S)$, and therefore (9.3) follows from (9.1)–(9.2).

No geodesic ray can intersect itself before time ϱ , where ϱ is the injectivity radius of S , so for every geodesic segment γ_T to length T the corresponding tessellation must satisfy $v_T \leq T^2/\varrho^2$. The inequality (9.4) now follows by Euler's formula and the relation $e = v \pm 2$. $\square$

Proof of Theorem 1.4. — We will prove only the assertion concerning the empirical frequencies of k -gons in the induced tessellation. Similar arguments can be used to prove that the empirical distributions of scaled side-lengths, interior angles, etc. converge to the corresponding theoretical frequencies in a Poisson line process. Denote by $\mathcal{T}_T$ the tessellation of the surface S induced by a random geodesic segment of length T .

We first give a heuristic argument that explains how Theorem 1.2, Corollary 2.9, and Proposition 9.1 together imply the convergence of empirical frequencies. Suppose that, for large T , the surface S could be partitioned into non-overlapping regions R_i each nearly isometric, by the scaled exponential mapping from the tangent space based at its center x_i , to a square of side αT^{-1} . (Of course this is not possible, because it would violate the fact that S has non-zero scalar curvature.) The hyperbolic area of R_i would be $\sim \alpha^2/T^2$, and so the number of squares R_i in the partition would be $\sim T^2/(\alpha^2\kappa)$.

Assume that α is sufficiently large that with probability at least $1 - \epsilon$, the absolute errors in the limiting relations (2.3), (2.4), and (2.5) (for some fixed k) of Corollary 2.9 are less than ϵ . By Theorem 1.2, for any point $x \in S$ and any α , the restriction of the geodesic tessellation $\mathcal{T}_T$ to the disk $D(x, 2\alpha T^{-1})$, when pulled back to the ball $B(0, 2\alpha)$ of the tangent space $T_x S$, converges in distribution, as a line process, to the Poisson line process of intensity κ . Since this holds for every x , it follows that for all sufficiently large T , with probability at least $1 - 2\epsilon$, in all but a fraction ϵ of the regions R_i the counts $V_T(R_i)$ and $F_T(R_i)$ of vertices and faces in the regions R_i (in the tessellation $\mathcal{T}_T$) and the fractions $\Phi_{k,T}(R_i)$ of k -gons will satisfy

$$(9.6) \quad |V_T(R_i)/\alpha^2 - \kappa^2/\pi| < 2\epsilon,$$

$$(9.7) \quad |F_T(R_i)/\alpha^2 - \kappa^2/\pi| < 2\epsilon, \quad \text{and}$$

$$(9.8) \quad |\Phi_{k,T}(R_i) - \phi_k| < 2\epsilon.$$

Call the regions R_i where these inequalities hold *good*, and the others *bad*. Since all but an area of size $\epsilon \times \text{area}(S)$ is covered by good squares R_i , relations (9.7) and (9.3) imply that the total number of faces of $\mathcal{T}_T$ in the bad squares satisfies

$$\sum_{i \text{ bad}} F_T(R_i) \leq 4\epsilon T^2 \times \text{area}(S).$$

Consequently, regardless of how skewed the empirical distribution of faces in the bad regions might be, it cannot affect the overall fraction of k -gons by more than 8ϵ . Since $\epsilon > 0$ can be made arbitrarily small, it follows from (9.8) that

$$(9.9) \quad \lim_{T \rightarrow \infty} \Phi_{k,T}(S) = \phi_k.$$

To provide a rigorous argument, we must explain how the partition into “squares” R_i can be modified. Fix $\delta > 0$ small, and let Δ be a triangulation of S whose triangles τ all (a) have diameters less than δ and (b) have geodesic edges. If $\delta > 0$ is sufficiently small, the triangles of Δ will all be contained in coordinate patches nearly isometric, by the exponential mapping, to disks $B(0, 2\delta)$ in the tangent space TS_{x_τ} , where x_τ is a distinguished point in the interior of τ . In each such ball $B(0, 2\delta)$, use an orthogonal coordinate system to foliate $B(0, 2\delta)$ by lines parallel to the coordinate

axes, and then use the exponential mapping to project these foliations to foliations of the triangles τ ; call these foliations $\mathcal{F}_x(\tau)$ and $\mathcal{F}_y(\tau)$. If $\delta > 0$ is sufficiently small then the curves in $\mathcal{F}_x(\tau)$ will cross curves in $\mathcal{F}_y(\tau)$ at angles $\theta \in [\frac{\pi}{2} - \epsilon, \frac{\pi}{2} + \epsilon]$, where $\epsilon > 0$ is small.

The foliations $\mathcal{F}_x(\tau)$ and $\mathcal{F}_y(\tau)$ can now be used as guidelines to partition τ into regions $R_i(\tau)$ whose boundaries are segments of curves in one or the other of the foliations. In particular, each boundary $\partial R_i(\tau)$ should consist of four segments, two from $\mathcal{F}_x(\tau)$ and two from $\mathcal{F}_y(\tau)$, and each should be of length $\sim \alpha T^{-1}$; thus, for large T each region $R_i(\tau)$ will be nearly a “parallelogram” (more precisely, the image of a parallelogram in the tangent space $TS_{x_i(\tau)}$ at a central point $x_i(\tau) \in R_i(\tau)$) whose interior angles are within ϵ of $\pi/2$. The collection of all regions $R_i(\tau)$, where τ ranges over the triangulation Δ , is nearly a partition of S into rhombi; only at distances $O(\alpha T^{-1})$ of the boundaries $\partial\tau$ are there overlaps. The total area in these boundary neighborhoods is $o(1)$ as $T \rightarrow \infty$.

Corollary 2.9, as stated, applies only to squares. However, any rhombus R whose interior angles are within ϵ of $\pi/2$ can be bracketed by squares $S_- \subset R \subset S_+$ in such a way that the area of $S_+ \setminus S_-$ is at most $C\epsilon \text{ area}(S_+)$, for some $C < \infty$ not depending on ϵ . Since Corollary 2.9 applies for each of the bracketing squares, it now follows as in the heuristic argument above that with probability $\geq 1 - C'\epsilon$, in all but a fraction $C\epsilon$ of the regions $R_i(\tau)$ the inequalities (9.6), (9.7), and (9.8) will hold. The limiting relation (9.9) now follows as before. $\square$

10. Extensions, Generalizations, and Speculations

A. Finite-area hyperbolic surfaces with cusps.

We expect also that Theorems 1.2–1.4 extend to finite-area hyperbolic surfaces with cusps. For this, however, genuinely new arguments would seem to be needed, as our analysis for the compact case relies heavily on the symbolic dynamics of Proposition 3.1 and the regenerative representation of Gibbs states (Proposition 3.3). The geodesic flow on the modular surface has its own very interesting symbolic dynamics (cf. for example [Ser85] and [AF82]), but this uses a countably infinite alphabet (the natural numbers) rather than a finite alphabet. At present there seems to be no analogue of the regenerative representation theorem (Proposition 3.3) for Gibbs states on sequence spaces with infinite alphabets.

B. Tessellations by closed geodesics.

It is known that statistical regularities of “random” geodesics (where the initial tangent vector is chosen from the maximal-entropy invariant measure for the geodesic flow) mimic those of typical long closed geodesics. This correspondence holds for first-order statistics (cf. [Bow72]), but also for second-order statistics (i.e., “fluctuations”): see [Lal14, Lal87, Lal89]. Thus, it should be expected that Theorems 1.2–1.4 have analogues for long closed geodesics. In particular, we conjecture the following.

CONJECTURE 10.1. — *Let S be a closed hyperbolic surface, and let $x \in S$ be a fixed point on S . From among all closed geodesics of length $\leq T$ choose one – call it γ_T – at random, and let A_T be the intersection of γ_T with the ball $D(x; \alpha T^{-1})$. Then as $T \rightarrow \infty$ the random collection of arcs A_T converge in distribution to a Poisson line process on $B(0; \alpha)$ of intensity κ .*

We do not expect that this will be true on a surface of variable negative curvature, because the maximal-entropy invariant measure for the geodesic flow coincides with the Liouville measure only in constant curvature.

C. Tessellations by several closed geodesics.

Given Conjecture 10.1, it is natural to expect that if two (or more) closed geodesics γ_T, γ'_T are chosen at random from among all closed geodesics of length $\leq T$, the resulting tessellations should be independent. Thus, the intersections of these tessellations with a ball $D(x, \alpha T^{-1})$ should converge jointly in law to independent Poisson line processes of intensity κ .

Appendix A. Poisson Line Processes

Proof of Lemma 2.1. — Rotational invariance is obvious, since the angles Θ_n are uniformly distributed, so it suffices to establish invariance by translations along the x -axis. To accomplish this, we will exhibit a sequence $\mathcal{L}_m$ of line processes that converge pointwise to a Poisson line process $\mathcal{L}$, and show by elementary means that each $\mathcal{L}_m$ is translationally invariant.

Let $\{(R_n, \Theta_n)\}_{n \in \mathbb{Z}}$ be the Poisson point process used in the construction (1.1) of $\mathcal{L}$. For each $m = 3, 5, 7, \dots$, let $\mathcal{A}_m = \{k\pi/m\}_{0 \leq k < m}$ (the restriction to odd m prevents $\pi/2$ from occurring in $\mathcal{A}_m$). For each $n \geq 1$, let $\Theta_n^m = [m\Theta_n]/m$ be the nearest point in $\mathcal{A}_m$ less than Θ_n . By construction, for each m the random variables Θ_n^m are independent and identically distributed, with the uniform distribution on the finite set $\mathcal{A}_m$. Now define $\mathcal{L}_m$ to be the line process constructed in the same manner as $\mathcal{L}$, but using the discrete random variables Θ_n^m instead of the continuous random variables Θ_n . Clearly, as $m \rightarrow \infty$ the sequence $\mathcal{L}_m$ of line processes converges to $\mathcal{L}$.

It remains to show that each of the line processes $\mathcal{L}_m$ is invariant by translations along the x -axis. For this, observe that for each $\theta_k \in \mathcal{A}_m$ the thinned process $\mathcal{R}_{m,k}$ consisting of those R_n such that $\Theta_n^m = \theta_k$ is itself a Poisson point process on $\mathbb{R}$ of intensity κ/m , and that these thinned Poisson point processes are mutually independent.⁽⁶⁾ Consequently, the line process $\mathcal{L}_m$ is the superposition of m independent line processes $\mathcal{L}_m^k$, with $k = 1, 2, \dots, m$, where $\mathcal{L}_m^k$ is the subset of all lines in $\mathcal{L}_m$ that meet the x -axis at angle $\pi/2 - \theta_k$. Since the constituent processes $\mathcal{L}_m^k$ are independent, it suffices to show that for each k the line process $\mathcal{L}_m^k$ is translation-invariant. But this is elementary: the points where the lines in $\mathcal{L}_m^k$ meet the x -axis form a

⁽⁶⁾The thinning and superposition laws are elementary properties of Poisson point processes. The thinning law follows from the superposition property; see Kingman [Kin93] for a proof of the latter.

Poisson point process on the real line, and Poisson point processes on the real line of constant intensity are translation-invariant. $\square$

Proof of Corollary 2.3. — By rotational invariance, it suffices to show this for the x -axis. Let $\mathcal{L}_m$ and $\mathcal{L}_m^k$ be as in the proof of Lemma 2.1; then by an easy calculation, the point process of intersections of the lines in $\mathcal{L}_m^k$ with the x -axis is a Poisson point process of intensity $(\kappa/m) \sin \theta_k$. Summing over k and then letting $m \rightarrow \infty$, one arrives at the desired conclusion. $\square$

Proof of Proposition 2.5. — The hypothesis that Γ encloses a strictly convex region guarantees that if a line intersects both A and B then it meets each in at most one point. Denote by $L_{\{A,B\}}$ the set of all lines that intersect both A and B . If A and B are partitioned into non-overlapping sub-arcs A_i and B_j then $L_{\{A,B\}}$ is the *disjoint* union $\cup_{i,j} L_{\{A_i,B_j\}}$. Since the sets $L_{\{A_i,B_j\}}$ are piecewise disjoint, the corresponding regions of the strip $\mathbb{R} \times [0, \pi)$ (in the standard parametrization (1.1)) are non-overlapping, and so, by a defining property of the Poisson point process $\{(R_n, \Theta_n)\}_{n \in \mathbb{Z}}$, the counts $N_{\{A_i,B_j\}}$ are independent Poisson random variables. Since the sum of independent Poisson random variables is Poisson, to finish the proof it suffices to show that for arcs A, B of length $< \varepsilon$ the random variables $N_{\{A,B\}}$ are Poisson, with means $\kappa\beta_{A,B}$.

If $\varepsilon > 0$ is sufficiently small then any line L that intersects two boundary arcs A, B of length $\leq \varepsilon$ must intersect the two straight line segments $\tilde{A}, \tilde{B}$ connecting the endpoints of A and B , respectively; conversely, any line that intersects both $\tilde{A}, \tilde{B}$ will intersect both A, B . Therefore, we may assume that the arcs A, B, A_i, B_i are straight line segments of length $\leq \varepsilon$. Because Poisson line processes are rotationally invariant, we may further assume that A is the interval $[-\varepsilon/2, \varepsilon/2] \times \{0\}$.

We now resort once again to the discretization technique used in the proof of Lemma 2.1. For each $m = 3, 5, 7, \dots$, let $N_{\{A,B\}}^m$ be the number of lines in the line process $\mathcal{L}_m$ that cross the segments A, B . Clearly, $N_{\{A,B\}}^m \rightarrow N_{\{A,B\}}$ as $m \rightarrow \infty$, so it suffices to show that for each m the random variable $N_{\{A,B\}}^m$ has a Poisson distribution with mean $\mu_m \rightarrow \kappa\beta_{A,B}$.

Recall that the line process $\mathcal{L}_m$ is a superposition of m independent line processes $\mathcal{L}_m^k$, and that for each k the lines in $\mathcal{L}_m^k$ all meet the x -axis at a fixed angle $|\pi/2 - \theta_k|$. Hence, $N_{\{A,B\}}^m = \sum_k N_{\{A,B\}}^{m,k}$, where $N_{\{A,B\}}^{m,k}$ is the number of lines in $\mathcal{L}_m^k$ that cross both A and B . The random variables $N_{\{A,B\}}^{m,k}$ are independent; thus, to show that $N_{\{A,B\}}^m$ has a Poisson distribution it suffices to show that each $N_{\{A,B\}}^{m,k}$ is Poisson. By construction, the lines in $\mathcal{L}_m^k$ meet the line $(s \cos \theta_k, s \sin \theta_k)_{s \in \mathbb{R}}$ at the points of a Poisson point process of intensity κ/m ; consequently, they meet the x -axis at the points of a Poisson point process of intensity $\kappa |\cos \theta_k|/m$. Now a line that meets the x -axis at angle θ_k will cross both $A = [-\varepsilon/2, \varepsilon/2] \times \{0\}$ and B if and only if its point of intersection with the x -axis lies in the θ_k -shadow J_k of B on A . Therefore, $N_{\{A,B\}}^{m,k}$ has the Poisson distribution with mean $\kappa |J_k \cos \theta_k|/m = \kappa \psi_{A,B}(\theta_k)$. It follows

that $N_{\{A,B\}}^m$ has the Poisson distribution with mean

$$\begin{aligned} EN_{\{A,B\}}^m &= m^{-1} \sum_{k=0}^{m-1} \kappa |J_k \cos \theta_k| \\ &= m^{-1} \sum_{k=0}^{m-1} \kappa \psi_{A,B}(\theta_k) \\ &\longrightarrow \frac{\kappa}{\pi} \int_{-\pi/2}^{\pi/2} \psi_{A,B}(\theta) d\theta. \end{aligned} \quad \square$$

Proof of Corollary 2.6. — It suffices to prove this for disks of small radius, because by the translation-invariance of $\mathcal{L}$,

$$EV(D) \sim \frac{1}{\pi \varrho^2} \int_D EV(B(x, \varrho)) dx = EV(B(0, \varrho)) |D| / (\pi \varrho^2)$$

as $\varrho \rightarrow 0$. Let γ be a chord of $B(0, \varrho)$, and H_γ the event that $\gamma \in \mathcal{L} \cap B(0, \varrho)$. Conditional on H_γ , the number of intersection points on γ is Poisson with mean $2\kappa|\gamma|/\pi$, by Corollary 2.3 and Proposition 2.5.⁽⁷⁾ Therefore,

$$EV(B(0, \varrho)) = \frac{1}{2} \frac{2\kappa}{\pi} E \left(\sum_{\gamma \in \mathcal{L} \cap B(0, \varrho)} |\gamma| \right) := \frac{\kappa}{\pi} E\Psi(\mathcal{L} \cap B(0, \varrho)).$$

(The factor of $1/2$ accounts for the fact that each intersection point lies on two chords.)

The expectation $E\Psi(\mathcal{L} \cap B(0, \varrho))$ is easily evaluated using the standard construction of the Poisson line process (Definition 1.1). The lines of $\mathcal{L}$ that cross $B(0, \varrho)$ are precisely those corresponding to points R_n such that $-\varrho < R_n < \varrho$. For any such R_n , the length of the chord $\gamma = \gamma_n$ is $|\gamma_n| = 2\sqrt{\varrho^2 - R_n^2}$. Therefore,

$$E\Psi(\mathcal{L} \cap B(0, \varrho)) = \kappa \int_{r=-\varrho}^{\varrho} 2\sqrt{\varrho^2 - r^2} dr = \kappa \pi \varrho^2. \quad \square$$

Proof of Proposition 2.7. — Let $\mathcal{L}$ be the Poisson line process with intensity κ , and denote by τ_z the translation by $z \in \mathbb{R}^2$. It suffices to prove that for any two bounded, continuous functions $f, g : \mathcal{C} \rightarrow \mathbb{R}$,

$$(A.1) \quad \lim_{|z| \rightarrow \infty} Ef(\mathcal{L})g(\tau_z \mathcal{L}) = Ef(\mathcal{L})Eg(\mathcal{L}).$$

Since the Poisson line process is rotationally invariant, it suffices to consider only translations τ_z for $z = (x, 0)$ on the x -axis. Moreover, since continuous functions that depend only on the restrictions of configurations to balls are dense in the space of all bounded, continuous functions, it suffices to establish (A.1) for functions f, g that depend only on configurational restrictions to the ball of radius $r > 0$ centered at the origin.

⁽⁷⁾The event H_γ has probability 0, but it is the limit of the positive-probability events that $\mathcal{L}$ has a line which intersects small boundary arcs centered at the endpoints of γ . The conditional distribution of $\mathcal{L}$ given H_γ can be interpreted as the limit of the conditional distributions given these approximating events. The independence assertion of Proposition 2.5 guarantees that, conditional on H_γ , the distribution of $\mathcal{L} \cap B(0, \varrho)$ is the same as the unconditional distribution of $(\mathcal{L} \cap B(0, \varrho)) \cup \{\gamma\}$.

To prove (A.1), we will show that on some probability space there are Poisson line processes $\mathcal{L}, \mathcal{L}', \mathcal{L}''$, each with intensity κ , such that

- (a) the line processes $\mathcal{L}'$ and $\mathcal{L}''$ are independent;
- (b) $f(\mathcal{L}) = f(\mathcal{L}')$ with probability one; and
- (c) $g(\tau_z \mathcal{L}) = g(\tau_z \mathcal{L}'')$ with probability $\rightarrow 1$ as $|z| \rightarrow \infty$.

It will then follow, by translation invariance, that

$$\begin{aligned} |Ef(\mathcal{L})g(\tau_z \mathcal{L}) - Ef(\mathcal{L})Eg(\mathcal{L})| &= |Ef(\mathcal{L})g(\tau_z \mathcal{L}) - Ef(\mathcal{L}')g(\mathcal{L}'')| \\ &= |Ef(\mathcal{L})g(\tau_z \mathcal{L}) - Ef(\mathcal{L}')g(\tau_z \mathcal{L}'')| \\ &\leq 2\|f\|_\infty \|g\|_\infty P\{g(\tau_z \mathcal{L}) \neq g(\tau_z \mathcal{L}'')\} \rightarrow 0. \end{aligned}$$

The line processes $\mathcal{L}, \mathcal{L}', \mathcal{L}''$ can be built on any probability space that supports independent Poisson point processes $\{R'_n\}_{n \in \mathbb{Z}}$ and $\{R''_n\}_{n \in \mathbb{Z}}$ on $\mathbb{R}$ of intensity κ , and independent sequences $\{\Theta'_n\}_{n \in \mathbb{Z}}$ and $\{\Theta''_n\}_{n \in \mathbb{Z}}$ of random variables uniformly distributed on the interval $[-\pi, \pi]$. Let $\mathcal{L}'$ be the line process obtained by using the “standard construction” (that is, the construction explained in Definition 1.1) with the point process $\{R'_n\}_{n \in \mathbb{Z}}$ and the accompanying uniform random variables $\{\Theta'_n\}_{n \in \mathbb{Z}}$, and let $\mathcal{L}''$ be the line process obtained by the standard construction using the point process $\{R''_n\}_{n \in \mathbb{Z}}$ and the random variables $\{\Theta''_n\}_{n \in \mathbb{Z}}$. Clearly, $\mathcal{L}'$ and $\mathcal{L}''$ are independent.

The line process $\mathcal{L}$ is constructed by splicing the marked Poisson point processes $\mathcal{R}' = \{(R'_n, \Theta'_n)\}_{n \in \mathbb{Z}}$ and $\mathcal{R}'' = \{(R''_n, \Theta''_n)\}_{n \in \mathbb{Z}}$ as follows: in the interval $(-r, r)$, use the marked points of $\{(R'_n, \Theta'_n)\}_{n \in \mathbb{Z}}$; but in $\mathbb{R} \setminus (-r, r)$, use the marked points of $\{(R''_n, \Theta''_n)\}_{n \in \mathbb{Z}}$. Thus, the resulting marked point process $\mathcal{R} = \{(R_n, \Theta_n)\}_{n \in \mathbb{Z}}$ consists of (i) all pairs (R'_n, Θ'_n) such that $-r < R'_n < r$, and (ii) all pairs (R''_n, Θ''_n) such that $R''_n \notin (-r, r)$. By standard results in the elementary theory of Poisson processes, the marked point process $\mathcal{R}$ has the same distribution as $\mathcal{R}'$ and $\mathcal{R}''$, in particular, $\{R_n\}_{n \in \mathbb{Z}}$ is a rate- κ Poisson point process on $\mathbb{R}$, and the random variables $\{\Theta_n\}_{n \in \mathbb{Z}}$ are independent and uniformly distributed on $[-\pi, \pi]$. Let $\mathcal{L}$ be the Poisson line process constructed using $\mathcal{R}$.

It remains to show that the Poisson line processes $\mathcal{L}, \mathcal{L}', \mathcal{L}''$ satisfy properties (b) and (c) above. Observe first that in the standard construction (Definition 1.1), only those pairs (R_n, Θ_n) such that $R_n \in (-r, r)$ will produce lines that intersect the ball $B(0, r)$ of radius r centered at the origin. Consequently, the restrictions of $\mathcal{L}$ and $\mathcal{L}'$ to $B(0, r)$ are equal; since f depends only on the configuration in $B(0, r)$, it follows that $f(\mathcal{L}) = f(\mathcal{L}')$.

Next, consider the configurational restrictions of $\mathcal{L}$ and $\mathcal{L}''$ to the ball $B((x, 0), r)$ for $x \gg 2r$. In the standard construction, a pair (R_n, Θ_n) such that $R_n \in (-r, r)$ will produce a line of $\mathcal{L}$ that intersects $B((x, 0), r)$ only if $|\tan \Theta_n| \leq r/(x - 2r)$. The probability that there is such a pair, in either $\mathcal{R}$ or $\mathcal{R}''$, tends to 0 as $x \rightarrow \infty$; hence, with probability $\rightarrow 1$, the restrictions of $\mathcal{L}$ and $\mathcal{L}''$ agree in $B((x, 0), r)$, and on this event $g(\mathcal{L}) = g(\mathcal{L}'')$. $\square$

Proof of Corollary 2.9. — The number of lines in a Poisson line process $\mathcal{L}$ that intersect a given line segment of length m has the Poisson distribution with mean

$C\kappa m$, where C is a finite positive constant not depending on either m or κ . Consequently, the probability that the number of polygons in the induced tessellation of the plane intersecting one of the four sides of $[-n, n]^2$ exceeds $n^{3/2}$ is exponentially small in n .

Given a line configuration $\mathcal{L}$, let $1/f(\mathcal{L})$ be the area of the polygon containing the origin in the induced tessellation. (This is well-defined and positive with probability 1.) Let A_n^- be the union of all polygons of the tessellation that lie entirely in the open square $(-n, n)^2$, and let A_n^+ be the union of the polygons that intersect $[-n, n]^2$. Then

$$\int_{A_n^-} f(\tau_z \mathcal{L}) dz \quad \text{and} \quad \int_{A_n^+} f(\tau_z \mathcal{L}) dz$$

count the number of polygons in A_n^- and A_n^+ , respectively; since the difference between these is less than $n^{3/2}$, except with exponentially small probability, it follows that except with small probability

$$\left| F_n - \int_{[-n, n]^2} f(\tau_z \mathcal{L}) dz \right| \leq n^{3/2}.$$

Hence, by the multi-parameter ergodic theorem (see, for example, [Cal53]), $F_n/n^2 \rightarrow Ef(\mathcal{L})$ almost surely.

The proof of the assertion regarding empirical frequencies of k -gons is similar. If G_k is the event that the polygon containing the origin is a k -gon, then the total number of k -gons in the region $A_n^\pm$ is

$$\int_{A_n^\pm} (f \mathbf{1}_{G_k})(\tau_z \mathcal{L}) dz.$$

Hence, the ergodic theorem implies that the number of k -gons divided by n^2 converges to $E(f \mathbf{1}_{G_k}(\mathcal{L}))$, and it follows that the fraction of k -gons converges to

$$\phi_k = \frac{E(f \mathbf{1}_{G_k}(\mathcal{L}))}{Ef(\mathcal{L})}.$$

Now consider the number of vertices V_n . Because there is probability 0 that three distinct lines of a Poisson line process meet at a point, all interior vertices are shared by exactly 4 edges, and each edge is incident to two vertices; thus, since the number of vertices on the boundary of the square is $O(n^{3/2})$, we have $\mathcal{E}_n = 2V_n + O(n^{3/2})$. By Euler's formula, $V_n - \mathcal{E}_n + F_n = 1$, so $V_n = F_n + O(n^{3/2})$; hence,

$$\lim_{n \rightarrow \infty} V_n/n^2 = \lim_{n \rightarrow \infty} F_n/n^2.$$

The value of the limit is determined by Corollary 2.6, which implies that $EV_n = 4\kappa^2 n^2/\pi$. $\square$

BIBLIOGRAPHY

- [AF82] Roy L. Adler and Leopold Flatto, *Cross section map for the geodesic flow on the modular surface*, Conference in modern analysis and probability (New Haven, Conn., 1982), Contemporary Mathematics, vol. 26, American Mathematical Society, 1982, pp. 9–24. $\uparrow 218$
- [Bil68] Patrick Billingsley, *Convergence of probability measures*, John Wiley & Sons, 1968. $\uparrow 190$

- [Bow72] Rufus Bowen, *The equidistribution of closed geodesics*, Am. J. Math. **94** (1972), 413–423. ↑218
- [Bow73] ———, *Symbolic dynamics for hyperbolic flows*, Am. J. Math. **95** (1973), 429–460. ↑199
- [Bow75] ———, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture Notes in Mathematics, vol. 470, Springer, 1975. ↑197, 198, 199, 200
- [BS79] Rufus Bowen and Caroline Series, *Markov maps associated with Fuchsian groups*, Publ. Math., Inst. Hautes Étud. Sci. **50** (1979), 153–170. ↑199
- [BW72] Rufus Bowen and Peter Walters, *Expansive one-parameter flows*, J. Differ. Equations **12** (1972), 180–193. ↑198
- [Cal53] Alberto P. Calderon, *A general ergodic theorem*, Ann. Math. **58** (1953), 182–191. ↑223
- [Cal03] Pierre Calka, *An explicit expression for the distribution of the number of sides of the typical Poisson–Voronoi cell*, Adv. Appl. Probab. **35** (2003), no. 4, 863–870. ↑196
- [CFF02] Francis M. Comets, Roberto Fernández, and Pablo A. Ferrari, *Processes with long memory: regenerative construction and perfect simulation*, Ann. Appl. Probab. **12** (2002), no. 3, 921–943. ↑199, 200
- [Dol04] Dmitry Dolgopyat, *Limit theorems for partially hyperbolic systems*, Trans. Am. Math. Soc. **356** (2004), no. 4, 1637–1689. ↑192
- [DZ98] Amir Dembo and Ofer Zeitouni, *Large deviations techniques and applications*, corrected reprint of the second ed., Applications of Mathematics, vol. 38, Springer, 1998. ↑212
- [EJM91] Charles Epstein, Troels Jorgensen, and Curt McMullen, *Multiple intersections on negatively curved surfaces*, J. Differ. Geom. **33** (1991), no. 1, 253–261. ↑216
- [Kin93] John F. C. Kingman, *Poisson processes*, Oxford Studies in Probability; Oxford Science Publications, vol. 3, Clarendon Press; Oxford University Press, 1993. ↑219
- [KM99] Dmitry Y. Kleinbock and Grigoriĭ A. Margulis, *Logarithm laws for flows on homogeneous spaces*, Invent. Math. **138** (1999), no. 3, 451–494. ↑192
- [Lal86] Steven P. Lalley, *Regenerative representation for one-dimensional Gibbs states*, Ann. Probab. **14** (1986), no. 4, 1262–1271. ↑199, 200
- [Lal87] ———, *Distribution of periodic orbits of symbolic and Axiom A flows*, Adv. Appl. Math. **8** (1987), no. 2, 154–193. ↑218
- [Lal89] ———, *Closed geodesics in homology classes on surfaces of variable negative curvature*, Duke Math. J. **58** (1989), no. 3, 795–821. ↑218
- [Lal96] ———, *Self-intersections of closed geodesics on a negatively curved surface: statistical regularities*, Convergence in ergodic theory and probability (Columbus, OH, 1993), Ohio State University Mathematical Research Institute Publications, vol. 5, Walter de Gruyter, 1996, pp. 263–272. ↑191
- [Lal14] ———, *Statistical regularities of self-intersection counts for geodesics on negatively curved surfaces*, Duke Math. J. **163** (2014), no. 6, 1191–1261. ↑197, 216, 218
- [LC60] Lucien Le Cam, *An approximation theorem for the Poisson binomial distribution*, Pac. J. Math. **10** (1960), 1181–1197. ↑197
- [LNP19] Jialun Li, Frédéric Naud, and Wenyu Pan, *Kleinian Schottky groups, Patterson–Sullivan measures, and Fourier decay, with an appendix on stationarity of Patterson–Sullivan measures*, <https://arxiv.org/abs/1902.01103>, to appear in *Duke Mathematical Journal*, 2019. ↑200
- [Mau06] François Maucourant, *Dynamical Borel–Cantelli lemma for hyperbolic spaces*, Isr. J. Math. **152** (2006), 143–155. ↑192
- [Mil64a] Roger E. Miles, *Random polygons determined by random lines in a plane*, Proc. Natl. Acad. Sci. USA **52** (1964), 901–907. ↑188, 193, 194, 195

- [Mil64b] ———, *Random polygons determined by random lines in a plane. II*, Proc. Natl. Acad. Sci. USA **52** (1964), 1157–1160. ↑188, 193, 195
- [PPS15] Frédéric Paulin, Mark Pollicott, and Barbara Schapira, *Equilibrium states in negative curvature*, Astérisque, vol. 373, Société Mathématique de France, 2015. ↑199
- [Rat73] Marina Ratner, *Markov partitions for Anosov flows on n -dimensional manifolds*, Isr. J. Math. **15** (1973), 92–114. ↑199
- [San04] Luis A. Santaló, *Integral geometry and geometric probability*, 2nd ed., Cambridge Mathematical Library, Cambridge University Press, 2004, with a foreword by Mark Kac. ↑194
- [Ser81] Caroline Series, *Symbolic dynamics for geodesic flows*, Acta Math. **146** (1981), no. 1-2, 103–128. ↑197, 198, 199
- [Ser85] ———, *The modular surface and continued fractions*, J. Lond. Math. Soc. **31** (1985), no. 1, 69–80. ↑218
- [SKM87] Dietrich K. Stoyan, Wilfrid S. Kendall, and Joseph Mecke, *Stochastic geometry and its applications*, Wiley Series in Probability and Mathematical Statistics, John Wiley & Sons, 1987, With a foreword by D. G. Kendall. ↑193, 194
- [Sul82] Dennis Sullivan, *Disjoint spheres, approximation by imaginary quadratic numbers, and the logarithm law for geodesics*, Acta Math. **149** (1982), no. 3, 3–4. ↑192
- [SY94] Richard Schoen and Shing-Tung Yau, *Lectures on differential geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology, vol. 1, International Press, 1994, Lecture notes prepared by Wei Yue Ding, Kung Ching Chang [Gong Qing Zhang], Jia Qing Zhong and Yi Chao Xu, Translated from the Chinese by Ding and S. Y. Cheng, Preface translated from the Chinese by Kaising Tso. ↑199

Manuscript received on 12th February 2019,
revised on 19th April 2020,
accepted on 27th April 2020.

Recommended by Editor S. Gouëzel.
Published under license CC BY 4.0.



This journal is a member of Centre Mersenne.



Jayadev ATHREYA
Department of Mathematics,
University of Washington,
PO Box 354350, Seattle,
WA, 98195-4350 (USA)
jathreya@uw.edu

Steve LALLEY
Department of Statistics,
University of Chicago,
5734 University Avenue,
Chicago, IL 60637 (USA)
lalley@galton.uchicago.edu

Jenya SAPIR
Department of Mathematical Sciences,
Binghamton University,
PO Box 6000 Binghamton,
New York 13902-6000 (USA)
sapis@math.binghamton.edu

Matthew WROTEN
Cold Spring Harbor Laboratory,
One Bungtown Road
Cold Spring Harbor,
NY 11724 (USA)
mwroten@cshl.edu