



# Metagenomes from Experimental Hydrologic Manipulation of Restored Coastal Plain Wetland Soils (Tyrell County, North Carolina)

 Ariane L. Peralta,<sup>a</sup> Regina B. Bledsoe,<sup>a</sup>  Mario E. Muscarella,<sup>b</sup> Marcel Huntemann,<sup>c</sup> Alicia Clum,<sup>c</sup> Brian Foster,<sup>c</sup> Bryce Foster,<sup>c</sup> Simon Roux,<sup>c</sup> Krishnaveni Palaniappan,<sup>c</sup> Neha Varghese,<sup>c</sup> Supratim Mukherjee,<sup>c</sup>  T. B. K. Reddy,<sup>c</sup> Chris Daum,<sup>c</sup> Alex Copeland,<sup>c</sup> I-Min A. Chen,<sup>c</sup> Natalia N. Ivanova,<sup>c</sup> Nikos C. Kyrpides,<sup>c</sup> Tijana Glavina del Rio,<sup>c</sup> Emiley A. Eloë-Fadrosch<sup>c</sup>

<sup>a</sup>Department of Biology, East Carolina University, Howell Science Complex, Greenville, North Carolina, USA

<sup>b</sup>Institute of Arctic Biology and Department of Biology and Wildlife, University of Alaska Fairbanks, Fairbanks, Alaska, USA

<sup>c</sup>Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, California, USA

**ABSTRACT** Hydrologic changes modify microbial community structure and ecosystem functions, especially in wetland systems. Here, we present 24 metagenomes from a coastal freshwater wetland experiment in which we manipulated hydrologic conditions and plant presence. These wetland soil metagenomes will deepen our understanding of how hydrology and vegetation influence microbial functional diversity.

Microbial community structure, soil physicochemical properties, and the abundance and composition of vegetation interact to influence biogeochemical functions (1). Changes in wetland hydrology due to drought, draining, and rewetting cause shifts in soil redox potential, microbial community composition, and associated ecosystem processes such as greenhouse gas (GHG) production (2–6). Vegetation also affects microbial processes by facilitating the transport of oxygen into the root zone and the transport of methane from the root zone into the atmosphere (7, 8). Here, we present wetland soil metagenomes from a mesocosm experiment in which we manipulated hydrology and plant presence to examine microbial community responses. These data complement targeted amplicon sequencing data, GHG fluxes, and soil physicochemical properties from the wetland mesocosm experiment (9).

We collected wetland soils from a restored coastal freshwater wetland at the Timberlake Observatory for Wetland Restoration on the Albemarle Peninsula in Tyrell County, North Carolina (latitude, 35.8959; longitude, –76.1658). We collected soils from three locations with different water table levels (–20 cm, –10 cm, and 0 cm) (10). We altered redox conditions by manipulating hydrology over 8 weeks. The experimental design and sampling details were published by Bledsoe and Peralta (9). We collected and combined six soil cores (3-cm diameter, 10-cm depth) from plant or no-plant areas. We completed 16S rRNA amplicon sequencing (NCBI BioProject [PRJNA636184](https://doi.org/10.5281/zenodo.4042110)), GHG flux measurements, and soil environmental analyses (9). We sequenced metagenomes that represented the most distinct microbial communities based on amplicon sequencing, and we chose the following samples: wetland soils sampled from field sites at which the water table measured –20 cm and 0 cm, to capture “mesocosm baseline” ( $n = 8$ ) functional diversity, and a subset of samples at the end of the 8-week hydrologic manipulation (i.e., prolonged drying or wetting only) in the presence or absence of vegetation ( $n = 16$ ) (9).

We used the Qiagen DNeasy PowerMax soil kit to extract genomic DNA from freeze-dried soils. Purified DNA products were sent to the U.S. Department of Energy (DOE) Joint Genome Institute (JGI) for sequencing and analysis. Metagenomes were sequenced at the DOE JGI (GitHub [see SupplementalTableS1\_MetagenomeSummary\_Peralta\_et\_al.csv at <https://doi.org/10.5281/zenodo.4042110>]), and project information

**Citation** Peralta AL, Bledsoe RB, Muscarella ME, Huntemann M, Clum A, Foster B, Foster B, Roux S, Palaniappan K, Varghese N, Mukherjee S, Reddy TBK, Daum C, Copeland A, Chen I-MA, Ivanova NN, Kyrpides NC, Glavina del Rio T, Eloë-Fadrosch EA. 2020. Metagenomes from experimental hydrologic manipulation of restored coastal plain wetland soils (Tyrell County, North Carolina). *Microbiol Resour Announc* 9:e00882-20. <https://doi.org/10.1128/MRA.00882-20>.

**Editor** J. Cameron Thrash, University of Southern California

**Copyright** © 2020 Peralta et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ariane L. Peralta, [peraltaa@ecu.edu](mailto:peraltaa@ecu.edu).

**Received** 29 July 2020

**Accepted** 15 September 2020

**Published** 8 October 2020

**TABLE 1** Annotation and gene-calling summary for the 24 metagenomes

Gene type and database used <sup>a</sup>	No. (mean ± SD)
CDSs	992,951 ± 305,299
rRNAs	775 ± 173
Other genes	4,878 ± 2,340
CDSs with COGs	633,925 ± 194,834
CDSs with SMART	127,732 ± 38,536
CDSs with SUPFAM	641,238 ± 189,914
CDSs with CATH FunFam	509,413 ± 154,168
CDSs with Pfam	603,115 ± 187,921

<sup>a</sup> CDSs, coding sequences; COGs, Clusters of Orthologous Groups; SUPFAM, superfamily; FunFam, functional families.

can be accessed under GOLD (11) study project accession number [Gs0142547](https://doi.org/10.5281/zenodo.4042110) and NCBI BioProject accession number [PRJNA641216](https://doi.org/10.5281/zenodo.4042110). Plate-based DNA library preparation for Illumina sequencing was performed according to published protocols in GitHub (see Supplemental\_Methods\_Details\_MRA\_WetlandSoilMetagenomes.txt at <https://doi.org/10.5281/zenodo.4042110>). The sequencing project resulted in 583.2 Gbp of raw sequence data. The average read length for each metagenome is found in GitHub (see SupplementalTableS1\_MetagenomeSummary\_Peralta\_et\_al.csv at <https://doi.org/10.5281/zenodo.4042110>). These data were processed using the DOE JGI Metagenome Annotation Pipeline using IMG/M v.5.0.9 (12–14). Initial sequence quality control details can be found in GitHub (see Supplemental\_Methods\_Details\_MRA\_WetlandSoilMetagenomes.txt at <https://doi.org/10.5281/zenodo.4042110>). Illumina reads were quality control filtered according to the protocol described in GitHub (see Supplemental\_Methods\_Details\_MRA\_WetlandSoilMetagenomes.txt at <https://doi.org/10.5281/zenodo.4042110>).

Annotation and gene calling resulted in  $600,507 \pm 172,049$  annotated contigs per sample (mean ± standard deviation [SD]), with a mean of 998,604 gene features (SD, 307,431 gene features) identified in each (Table 1). Based on phylogenetic associations (determined on the basis of bidirectional best hits to genes in other genomes), the estimated alpha diversity across all metagenomes is  $10,365 \pm 837$  operational taxonomic units (OTUs) (mean ± SD) (GitHub [see SupplementalTableS1\_MetagenomeSummary\_Peralta\_et\_al.csv at <https://doi.org/10.5281/zenodo.4042110>]).

**Data availability.** Metagenomes were sequenced at the DOE JGI, and the study information can be found under GOLD study project accession number [Gs0142547](https://doi.org/10.5281/zenodo.4042110) and NCBI BioProject accession number [PRJNA641216](https://doi.org/10.5281/zenodo.4042110). Additional sample-specific metagenome statistics can be found in GitHub (see SupplementalTableS1\_MetagenomeSummary\_Peralta\_et\_al.csv at <https://doi.org/10.5281/zenodo.4042110>). Details on metagenomic library preparation and sequence filtering can be found in GitHub (see Supplemental\_Methods\_Details\_MRA\_WetlandSoilMetagenomes.txt at <https://doi.org/10.5281/zenodo.4042110>).

## ACKNOWLEDGMENTS

We thank J. LeCrone, L. Armstrong, M. Stillwagon, G. Gunderson, C. Eakins, J. Basco, and C. Bledsoe for field and laboratory assistance. We thank J. Gill and the East Carolina University grounds crew for their efforts in maintaining the grounds surrounding the shade house.

This work was supported by the National Science Foundation (Graduate Research Fellowship Program [GRFP] grant to R.B.B. and grant DEB 1845845 to A.L.P.). The metagenomes were produced by the DOE JGI under the Community Science Program (CSP) (JGI CSP grant 503952). The work conducted by the DOE JGI, a DOE Office of Science User Facility, is supported under contract DE-AC02-05CH11231.

## REFERENCES

- Jansson JK, Hofmockel KS. 2020. Soil microbiomes and climate change. 1. *Nat Rev Microbiol* 18:35–46. <https://doi.org/10.1038/s41579-019-0265-7>.
- Peralta AL, Ludmer S, Matthews JW, Kent AD. 2014. Bacterial community response to changes in soil redox potential along a moisture gradient in restored wetlands. *Ecol Eng* 73:246–253. <https://doi.org/10.1016/j.ecoleng.2014.09.047>.

3. Lazcano C, Deol AS, Brummell ME, Strack M. 2020. Interactive effects of vegetation and water table depth on belowground C and N mobilization and greenhouse gas emissions in a restored peatland. *Plant Soil* 448: 299–313. <https://doi.org/10.1007/s11104-020-04434-2>.
4. Bansal S, Johnson OF, Meier J, Zhu X. 2020. Vegetation affects timing and location of wetland methane emissions. *J Geophys Res Biogeosci* 125: e2020JG005777.
5. Ou Y, Rousseau AN, Wang L, Yan B, Gumiere T, Zhu H. 2019. Identification of the alteration of riparian wetland on soil properties, enzyme activities and microbial communities following extreme flooding. *Geoderma* 337: 825–833. <https://doi.org/10.1016/j.geoderma.2018.10.032>.
6. Peralta AL, Ludmer S, Kent AD. 2013. Hydrologic history influences microbial community composition and nitrogen cycling under experimental drying/wetting treatments. *Soil Biol Biochem* 66:29–37. <https://doi.org/10.1016/j.soilbio.2013.06.019>.
7. Chanton JP. 2005. The effect of gas transport on the isotope signature of methane in wetlands. *Org Geochem* 36:753–768. <https://doi.org/10.1016/j.orggeochem.2004.10.007>.
8. Carmichael MJ, Bernhardt ES, Bräuer SL, Smith WK. 2014. The role of vegetation in methane flux to the atmosphere: should vegetation be included as a distinct category in the global methane budget? *Biogeochemistry* 119:1–24. <https://doi.org/10.1007/s10533-014-9974-1>.
9. Bledsoe RB, Peralta AL. 2020. Dynamic hydrology and plant-mediated effects reduce greenhouse gas emissions and alter wetland microbial communities. *bioRxiv* <https://doi.org/10.1101/2020.06.29.178533>.
10. Ardón M, Morse JL, Doyle MW, Bernhardt ES. 2010. The water quality consequences of restoring wetland hydrology to a large agricultural watershed in the southeastern coastal plain. *Ecosystems* 13:1060–1078. <https://doi.org/10.1007/s10021-010-9374-x>.
11. Mukherjee S, Stamatis D, Bertsch J, Ovchinnikova G, Katta HY, Mojica A, Chen I-MA, Kyrpides NC, Reddy TBK. 2019. Genomes Online Database (GOLD) v.7: updates and new features. *Nucleic Acids Res* 47:D649–D659. <https://doi.org/10.1093/nar/gky977>.
12. Markowitz VM, Chen I-MA, Chu K, Szeto E, Palaniappan K, Pillay M, Ratner A, Huang J, Pagani I, Tringe S, Huntemann M, Billis K, Varghese N, Tennesen K, Mavromatis K, Pati A, Ivanova NN, Kyrpides NC. 2014. IMG/M 4 version of the integrated metagenome comparative analysis system. *Nucleic Acids Res* 42:D568–D573. <https://doi.org/10.1093/nar/gkt919>.
13. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Tennesen K, Palaniappan K, Szeto E, Pillay M, Chen I-MA, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2016. The standard operating procedure of the DOE-JGI Metagenome Annotation Pipeline (MAP v.4). *Stand Genomic Sci* 11:17. <https://doi.org/10.1186/s40793-016-0138-x>.
14. Chen I-MA, Markowitz VM, Chu K, Palaniappan K, Szeto E, Pillay M, Ratner A, Huang J, Andersen E, Huntemann M, Varghese N, Hadjithomas M, Tennesen K, Nielsen T, Ivanova NN, Kyrpides NC. 2017. IMG/M: integrated genome and metagenome comparative data analysis system. *Nucleic Acids Res* 45:D507–D516. <https://doi.org/10.1093/nar/gkw929>.