

*Annual Review of Entomology*

# The Interplay Between Viruses and RNAi Pathways in Insects

Bryony C. Bonning<sup>1</sup> and Maria-Carla Saleh<sup>2</sup>

<sup>1</sup>Department of Entomology and Nematology, University of Florida, Gainesville, Florida 32611, USA; email: bbonning@ufl.edu

<sup>2</sup>Viruses and RNA Interference Unit, Institut Pasteur, CNRS UMR 3569, 75724 Paris CEDEX 15, France; email: carla.saleh@pasteur.fr

Annu. Rev. Entomol. 2021. 66:61–79

The *Annual Review of Entomology* is online at  
ento.annualreviews.org

<https://doi.org/10.1146/annurev-ento-033020-090410>

Copyright © 2021 by Annual Reviews.  
All rights reserved

ANNUAL REVIEWS **CONNECT**



- [www.annualreviews.org](http://www.annualreviews.org)
- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

## Keywords

RNAi pathways, antiviral immunity, endogenous viral elements, viral suppressors of RNAi

## Abstract

As an overarching immune mechanism, RNA interference (RNAi) displays pathogen specificity and memory via different pathways. The small interfering RNA (siRNA) pathway is the primary antiviral defense mechanism against RNA viruses of insects and plays a lesser role in defense against DNA viruses. Reflecting the pivotal role of the siRNA pathway in virus selection, different virus families have independently evolved unique strategies to counter this host response, including protein-mediated, decoy RNA-based, and microRNA-based strategies. In this review, we outline the interplay between insect viruses and the different pathways of the RNAi antiviral response; describe practical application of these interactions for improved expression systems and for pest and disease management; and highlight research avenues for advancement of the field.

## INTRODUCTION

**siRNA pathway:** small interfering RNAs of approximately 20–22 base pairs processed from long dsRNA by Dicer-2; this pathway is mainly antiviral

**RNA interference (RNAi):** comprised of three pathways; mechanism for sequence-specific gene, virus, or transposon silencing based on sRNA production

**Virus-induced gene silencing (VIGS):** use of a virus vector to deliver silencing RNAs to silence a target gene

**piRNA pathway:** PIWI-interacting RNAs of approximately 26–32 nucleotides that primarily function to silence transposons in the germline

**Dicer-2 (Dcr2):** RNase III nuclease that cleaves dsRNA into siRNAs

**Argonaute-2 (Ago2):** primary RISC component of the siRNA pathway mediating target recognition and cleavage of viral genome and transcripts

Like all organisms, insects are subject to infection by viruses, but only with the advent of next-generation sequencing have the abundance and diversity of insect-associated viruses been revealed (7, 63). The course and outcomes of viral infections rely on complex molecular host–virus interactions. Viruses hijack host cellular factors for infection and replication, while the host counters by restricting access to key factors and/or by mounting an antiviral immune response. A major player in the invertebrate antiviral immune response is the highly conserved small interfering RNA (siRNA) pathway of the RNA interference (RNAi) response (117). The siRNA pathway is the primary antiviral defense mechanism against insect viruses with RNA genomes (86) but also functions in defense against insect viruses with DNA genomes. Virus-derived double-stranded RNA (dsRNA) triggers the antiviral siRNA pathway in host cells, which can completely block virus replication.

Study of the interplay between viruses and the siRNA pathway has resulted in several tools, including methods for detection of covert viruses in insect cell lines (13, 14), optimized baculovirus expression systems (17, 101), and the potential use of virus-induced gene silencing (VIGS) for pest and disease management (106). The (re)emergence of insect vector-borne viruses such as dengue, Chikungunya, and Zika viruses highlights the importance of understanding insect–virus molecular interactions for providing solutions for global health.

## INSECT RNAi-DEPENDENT IMMUNE MECHANISMS AGAINST VIRAL INFECTIONS


The three primary small RNA (sRNA)-directed RNA silencing pathways in insects are the siRNA, microRNA (miRNA), and PIWI-interacting RNA (piRNA) pathways (139). While all three pathways center around the silencing of target RNA by Argonaute family proteins, which themselves are directed by the sRNAs in a sequence-specific manner, they are differentiated by the length and origin of the guiding sRNAs, enzymatic requirements, and functional outcomes.

### The siRNA Pathway

RNAi directed by siRNAs is a potent antiviral mechanism against both RNA and DNA viruses in a wide variety of eukaryotic organisms. This pathway begins with the recognition of dsRNA of viral origin in the cytoplasm by the RNase-III enzyme Dicer-2 (Dcr2) (5) (Figure 1). dsRNA is then cleaved by Dcr2 into approximately 21-nucleotide (nt) siRNA duplexes (109) containing 19 base-paired nts flanked by 2-nt 3' overhangs (23). One strand of the siRNA duplex, known as the guide strand, is incorporated into the RNA-induced silencing complex (RISC), while the other strand (the passenger strand) is degraded (72, 105). The guide strand is subsequently 2'-O-methylated at the 3' end by DmHen1 methyltransferase to form a mature RISC (28).

Argonaute-2 (Ago2) is the catalytic core of the RISC. Ago2 stably binds the guide strand within the protein complex through interactions between the 3' and 5' ends of the guide strand and the Ago2 PAZ and MID domains, respectively (15). siRNAs bound to Ago2 guide the RISC to complementary RNAs in the cytoplasm in a sequence-specific manner through Watson-Crick base pairing. Following target recognition, Ago2 catalyzes cleavage of target RNAs between nts 10 and 11 as measured from the 5' end of the guiding siRNA (24, 104). RNA molecules cleaved by Ago2 are then rapidly degraded through the activity of the XRN1 exoribonuclease and the exosome (94). Cleavage by Ago2 is sequence specific and requires perfect base pairing between nts 2–21 of the guide strand (70).

Single-stranded RNA (ssRNA), dsRNA, single-stranded DNA, and double-stranded DNA viruses are all targeted by siRNAs in insects (9, 85, 97, 128, 138). During the infection cycles



**Figure 1**

The antiviral small RNA pathways. While the antiviral role of the small interfering RNA (siRNA) pathway is well established, direct evidence for a role of the PIWI-interacting RNA (piRNA) pathway in antiviral immunity is lacking. We propose that the interaction of endogenous viral elements (EVEs) with exogenous cognate viruses could generate viral piRNAs with an antiviral role. These EVEs could originate from the integration of viral DNA forms of RNA viruses produced during viral infection in insects. In this way, there may be crosstalk between the two pathways to maintain host fitness during viral infection.

of RNA viruses, dsRNA is formed from the genomes of dsRNA viruses or by replication intermediates of ssRNA viruses (113). DNA viruses are thought to produce dsRNA by convergent transcription. For both DNA and RNA viruses, secondary structure within single-stranded viral RNA or transcripts can serve as a source of dsRNA (113). Notably, siRNAs can also be produced from endogenous sources of dsRNA (endo-siRNAs), and these are implicated in gene regulation and repression of transposable elements (TEs) (33).

Interaction between viral RNA and the siRNA pathway is indicated by the Dcr2-dependent accumulation of approximately 21-nt virus-derived siRNAs during virus infection in insects. The antiviral nature of the siRNA pathway is clearly demonstrated by the finding that knockout of Dcr2 or Ago2 results in enhanced virus replication and increased host mortality in a variety of systems (1, 52, 110, 128). This is further supported by the observation that both DNA and RNA viruses encode suppressors of siRNA-based RNAi (detailed below).

## The miRNA Pathway

Unlike siRNAs, which are usually produced from exogenous sources of dsRNA, the precursors of miRNAs are transcribed from endogenous loci by RNA polymerase II (63). These transcripts, known as primary miRNAs, are highly structured molecules that are cleaved by Drosha into approximately 60-nt hairpins known as precursor miRNAs (62). Precursor miRNAs are exported

to the cytoplasm, where they are recognized and cleaved by Dicer-1 to form a miRNA duplex (64, 69). One strand of the miRNA duplex is then loaded onto Argonaute-1 (Ago1) within the miRNA induced silencing complex (28). Mature miRNAs are approximately 22 nt, and in insects, they typically direct silencing of endogenous target RNAs through imperfect base pairing (3). Like Ago2, the endonuclease activity of Ago1 requires perfect base pairing along the guide miRNA (3). Because miRNA target recognition relies on imperfect base pairing, miRNA-directed RNA silencing usually involves translational repression through Ago1-mediated deadenylation and decapping (48). Relatively little work has explored the antiviral potential of miRNAs in insects. In mammals, miRNAs are known to exert antiviral effects by directing transcriptional changes that result in an unfavorable state for virus replication. More infrequently, mammalian miRNAs can directly target viral RNA (19). While virus infection in insects is known to alter the abundance of some host miRNAs in a temporal and tissue-specific manner, the effects that such changes might have on virus replication are unclear (81).

### The piRNA Pathway

The most recently discovered sRNA-based silencing pathway, the piRNA pathway, is the primary mechanism of TE repression in the animal germline (20). Insect piRNA pathway details are most clearly understood in *Drosophila melanogaster*, but the general mechanisms of piRNA biogenesis are conserved in most animals (30). piRNA biogenesis is classically divided into the primary and secondary pathways (7). In the *D. melanogaster* primary piRNA pathway, discrete genomic loci (piRNA clusters) give rise to long piRNA precursor transcripts, which are exported to the cytoplasm and fragmented by Zucchini endonuclease (Zuc) into precursor piRNAs (7, 91). Precursor piRNAs are then loaded onto Piwi or Aubergine (Aub), and their 3' ends are trimmed and 2'-O-methylated, resulting in 26–32-nt mature primary piRNAs (28, 41). Because Zuc cleavage preferentially occurs just upstream from uracil residues, and because the MID domains of PIWI-clade Argonaute proteins favor interactions with uracil residues, primary piRNAs have a strong bias for uracil as the fifth nt (1U bias) (18, 91). piRNA clusters contain a high density of TE-derived sequences (7). Thus, the primary piRNA pool contains an abundance of piRNAs specific to TEs.

The secondary piRNA pathway, also known as the ping-pong cycle, involves recognition of complementary RNA by Aub-bound primary piRNAs via nearly perfect Watson-Crick base pairing (7, 44). Like Ago2, Aub catalyzes the cleavage of target RNA between nts 10 and 11 with respect to the 5' end of the guiding primary piRNA (7). Cleaved target RNA with a 5' end defined by Aub cleavage is then loaded onto Argonaute-3 (Ago3) and processed into a 26–32-nt secondary piRNA with a 2'-O-methylated 3' end defined by Zuc cleavage or exonucleolytic degradation of cleaved RNA by nibbler (7, 36, 39, 41, 78, 132). Secondary piRNAs have a strong bias for adenine as the 10th nt (10A bias) and are antisense to the primary piRNA that directed the cleavage event resulting in their formation (7). Ago3-bound secondary piRNAs direct cleavage of complementary RNA targets, which are subsequently processed into Aub- or Piwi-bound primary piRNAs (7). Thus, the ping-pong cycle is a series of reciprocal cleavage events that serves as a post-transcriptional silencing mechanism and amplifies the piRNA-based response against active TEs (7, 20). piRNAs are distinguished from other types of sRNAs by their size (26–32 nt), and ping-pong-dependent piRNAs can be identified by 1U and 10A biases, as well as by a distance of 10 nt between the 5' ends of complementary piRNAs.

Interestingly, virus-derived, ping-pong-dependent piRNAs (vpiRNAs) are present during infection of *Aedes aegypti* and *Aedes albopictus* mosquitoes with several different ssRNA viruses (77), as well as in cell lines derived from these species. vpiRNAs derived from a single-stranded DNA virus were also observed during infection of *A. aegypti*-derived Aag2 cells (97). Unlike *D. melanogaster*, which expresses only three PIWI-clade Argonaute proteins (Aub, Ago3, and Piwi), *A. aegypti* and

*A. albopictus* express eight PIWI-clade Argonaute proteins (Ago3 and Piwi1–7) (12, 65). In Aag2 cells, Piwi5 and Ago3, but none of the other PIWI-clade Argonaute proteins, are required for biogenesis of vpiRNAs (76). In contrast, production of TE-derived piRNAs relies on a more diverse set of PIWI proteins (76). This suggests that the novel PIWI-clade Argonaute proteins in mosquitoes have undergone functional diversification to produce piRNAs from a variety of RNA sources. vpiRNAs are not produced during RNA or DNA virus infection in *D. melanogaster* and have not yet been observed outside of mosquitoes (98).

Despite their essential roles for vpiRNA biogenesis, knockdown of Piwi5 and Ago3 does not lead to enhanced virus replication in Aag2 cells (76, 111, 124, 129). Knockdown of Piwi4 does not significantly impact vpiRNA abundance in Aag2 cells or *A. aegypti* mosquitoes but does lead to increased replication of several RNA viruses in Aag2 cells, as well as increased replication of dengue virus *in vivo* in *A. aegypti* (76, 111, 124, 129, 130). While intriguing, the links among PIWI protein expression, vpiRNA abundance, and virus replication in mosquitoes are not yet clear, and direct evidence of an antiviral role for the piRNA pathway is lacking.

**Immune priming:**  
memory in insect  
immune response to a  
subsequent exposure  
to a pathogen

**Endogenous viral  
element (EVE):**  
complete or partial  
viral sequence  
integrated into host  
genome

## Immune Memory

A characteristic feature of the antiviral response in jawed vertebrates is the maintenance of long-lasting protection against future infections with the same virus (immune memory). Because insects lack antibodies, they have generally been considered incapable of preserving a functional record of prior infections. However, this view has been challenged by recent results demonstrating that DNA forms of RNA virus sequences are produced during viral infections, serve as sources of siRNAs, and are maintained even after clearance of the infection (34, 35, 79, 100, 123, 124). We are only beginning to understand the mechanisms underlying this immune memory or immune priming in insects.

The reverse transcriptases encoded by endogenous retrotransposons can reverse transcribe viral RNA in insects, resulting in the formation of chimeric viral DNAs (vDNAs) consisting partly of virus sequence and partly of TE sequence (34, 35, 100). Importantly, vDNAs are transcribed and serve as a source of the siRNAs that modulate virus infection (34, 35, 100, 123) (Figure 1). The prevention of vDNA formation during virus infection of *D. melanogaster* or *A. albopictus* decreases virus-specific siRNA abundance and increases insect mortality (34, 35). Remarkably, vDNAs generated from RNA viruses taken up by hemocytes in *D. melanogaster* give rise to virus-specific siRNAs that are secreted in exosome-like vesicles capable of priming an antiviral response in uninfected cells (123). Similarly, vDNAs purified from *D. melanogaster*-derived S2 cells infected with *Flock house virus* (FHV) confer virus-specific protective immunity following injection into naïve *D. melanogaster* flies, raising the possibility that vDNA may serve as a biologically relevant source of long-lasting and sequence-specific nucleic acid-based memory (100). Most studies of insect antiviral mechanisms have employed intrathoracic injection of virions, which generally results in death of the host or the development of a lifelong persistent infection (79). The persistent state in *D. melanogaster* and *A. albopictus* requires the formation of vDNA, highlighting its importance for long-term immune processes (34, 35). Because infections initiated by intrathoracic injection are not typically cleared, the potential role of vDNA in priming a protective response against future infections is difficult to assess.

vDNAs are episomal molecules produced during the course of virus infection (100). In contrast, the sequences of endogenous viral elements (EVEs) derive from full or partial integrations of viral sequence into the host genome (43, 51). EVEs from both DNA and RNA viruses have been described in a wide variety of eukaryotes (43, 51, 125). Nonretroviral RNA virus sequences within host genomes are surprising because these viruses do not undergo a DNA stage during replication.

### Viral suppressor of RNAi (VSR):

viral mechanism for countering host siRNA-mediated antiviral response

While the mechanisms of EVE generation are unclear, it is tempting to speculate that vDNAs are precursors of EVEs.

There are several well-known examples of EVEs being co-opted and playing important roles in the biology of the host (27, 75). Since their discovery, EVEs have been speculated to play an antiviral role (58) (Figure 1). In support of their antiviral potential, many EVEs are transcribed and contain uninterrupted open reading frames (ORFs). However, few reports have presented experimental evidence (29, 121). In honey bees, 30% of wild populations harbored an EVE derived from *Israeli acute paralysis virus* (IAPV) encoding a protein capable of cross-reacting with an antibody specific for the corresponding IAPV capsid protein (71). In vivo, individuals carrying this EVE exhibited reduced mortality following injection of IAPV compared to individuals without the EVE.

Given that nucleic acid-based defense mechanisms represent the primary antiviral strategy in arthropods, and that vDNA is capable of inducing immune priming via siRNAs, it is hypothesized that EVEs might similarly provide the specificity determinants of a long-lasting and heritable nucleic acid-based silencing system. In support of this, EVEs in *A. aegypti* and *A. albopictus* were found to be enriched within piRNA clusters and give rise to piRNAs (95, 135) (Figure 1). This finding was extended to arthropods in general by an analysis of the EVEs present within arthropods with sequenced genomes that have corresponding publicly available sRNA data sets (125).

So far, experiments on the antiviral potential of EVE-derived piRNAs have been limited to Aag2 cells (124, 135). An EVE derived from *Phasi Charoen-like virus* (PCLV) gives rise to a single piRNA mapping antisense to the PCLV genome (135). In Aag2 cells persistently infected with PCLV, the 5' end of this piRNA was separated by 10 nt from a piRNA mapping to the opposite strand, and knockdown of Piwi4 led to an approximately twofold increase in PCLV RNA levels (135). The Aag2 genome also contains EVEs derived from *Cell fusing agent virus* (CFAV) (124, 135). Aag2 cells persistently infected with CFAV show increased CFAV RNA levels, and knockdown of Piwi4 or Ago3 reduced CFAV-derived piRNA abundance (124). Although this evidence supports an antiviral role for these proteins, it is difficult to confirm an antiviral role for EVE-derived piRNAs based on CFAV-derived piRNA abundance without more detailed sRNA mapping profiles. Intriguingly, insertion of sequences corresponding to piRNA-producing EVEs within the *Sindbis virus* 3' UTR leads to diminished replication of the recombinant viruses in Aag2 cells following Piwi4 knockdown (124). The sRNA profiles during infection with such recombinant viruses should be the subject of future investigation.

## VIRAL COUNTER-DEFENSE MECHANISMS

### Viral Suppressors of siRNA-Based RNAi

Both RNA and DNA viruses encode viral suppressors of RNAi (VSRs) to counter the negative impact of the siRNA pathway on virus replication. VSRs from different virus families lack sequence or structural similarity, indicating independent evolution of different mechanisms to counter the siRNA pathway. VSRs act via four primary mechanisms: (a) binding to long dsRNA to prevent Dcr2-mediated cleavage, (b) binding to siRNA to prevent siRNA loading into RISC, (c) interfering with Dcr2 or Ago2 function, and (d) degrading siRNAs. Some VSRs combine multiple modes of action (Table 1).

The VSRs of viruses that infect *D. melanogaster* are particularly well studied. While homologous at the genetic level, the two dicistroviruses *Cricket paralysis virus* (CrPV) and *Drosophila C virus* (DCV) exhibit distinct virulence characteristics in wild-type *Drosophila*. CrPV results in paralytic, lethal (acute) infection, while DCV causes persistent, nonlethal (chronic) infection. The VSRs of CrPV and DCV map to the 1A protein at the N terminus of the ORF1 polypeptide but have low sequence similarity (approximately 9%) and lack homology to other proteins (89, 128, 133).

**Table 1** VSRs encoded by insect viruses and arboviruses

Family	Virus	VSR	Binds to			Inhibits		Decoy siRNA	Decoy RNA	References
			dsRNA	siRNA	Dcr2	Ago2	Dcr2			
<b>RNA virus</b>										
<i>Nodaviridae</i>	<i>Flleck house virus</i>	B2	X	X	X	—				2, 16, 66
	<i>Nodamura virus</i>	B2	X	X	—		X			120
	<i>Wuhan nodavirus</i>	B2	X	X	X					102, 103
	<i>Mosinavirus</i>	B2	X	—						114
<i>Dicistroviridae</i>	<i>Drosophila C virus</i>	1A	X	—	—					128
	<i>Cricket paralysis virus</i>	1A <sup>a</sup>	—	—	X					90, 134
<i>Birnaviridae</i>	<i>Drosophila X virus</i>	VP3	X	X	—					126
	<i>Culex Y virus</i>	VP3	X	X	—					126
Unassigned	<i>Drosophila melanogaster Nora virus</i>	VP1	—	—	X					127
	<i>Drosophila immigrans</i>	VP1			X					127
	<i>Nora-like virus</i>									
<b>DNA virus</b>										
<i>Flaviviridae</i>	<i>Dengue virus</i>	siRNA						X <sup>b</sup>		112
	<i>West Nile virus</i>	sRNA						X <sup>c</sup>		112
<i>Asfarviridae</i>	<i>Heliothis virescens asfarvirus</i>	RNase III					X			45
<i>Iridoviridae</i>	<i>Invertebrate iridescent virus</i>	IIV6-340R	X	X	—					8

<sup>a</sup>Also disrupts stress granule formation (56).

<sup>b</sup>Based on use of human Dicer. Inhibition of Dicer-2 in insects is assumed.

<sup>c</sup>Inhibits siRNA-based RNA interference in insect cells. This function presumed based on similarity to *West Nile virus* siRNA. Abbreviations: Ago2, Argonaute-2; Dcr2, Dicer-2; dsRNA, double-stranded RNA; siRNA, small interfering RNA; VSR, viral suppressor of RNA interference.

Expression of CrPV 1A or DCV 1A in *Drosophila* S2 cells blocked the siRNA pathway, increasing cell susceptibility to virus infection (89, 128). The pathogenicity traits of CrPV and DCV are reflected in the different efficacies of their respective VSRs in suppressing the siRNA pathway: The 166-amino-acid CrPV 1A prevents the Ago2 RISC from targeting RNA (134), while the 97-amino-acid DCV 1A binds to dsRNA, preventing processing by Dcr2 (128). Introduction of CrPV 1A into a recombinant *Sindbis* virus markedly increased virulence, while introduction of DCV 1A resulted in only a modest increase in virulence (89). Two elements within CrPV 1A that bind specific host proteins confer dual functionality in suppression of the siRNA pathway. In addition to acting as a competitive inhibitor to block Ago2, a BC box domain mediates ubiquitination of Ago2, thereby targeting Ago2 for degradation (90).

Stress granules form in the cytoplasm following inhibition of translation. They appear to function as a cellular antiviral response by sequestering viral proteins or RNA. Some viruses have evolved measures to counter stress granule formation. For example, CrPV and DCV, which inhibit S2 cell translation, do not result in formation of stress granules. CrPV 1A disrupts stress granule formation by a mechanism independent of its VSR function (56).

The 106-amino-acid B2 protein of FHV is a VSR that binds to dsRNA and siRNAs, inhibiting the siRNA pathway by sequestering siRNAs and preventing processing of dsRNA (2, 16, 66). B2 also binds to dsRNA regions in FHV RNA2 to prevent movement into cytoplasmic granules (which differ from stress granules) where translation is silenced. Deletion of B2 thus impairs translation of the capsid protein from RNA2 (99). FHV B2 is arguably the best studied of the insect virus VSRs due to the early availability of an infectious replicon that allows for strong genetic evidence. Indeed, other insect VSRs have been tested for their ability to rescue the FHV ΔB2 replicon (25). Transgenic *Drosophila* (6) show that, in addition to blocking antiviral defense, B2 and CrPV 1A also interfere with transposon silencing in somatic tissues by endo-siRNAs.

The Nora virus VP1 and Nora-like virus VP1 proteins function as VSRs by interacting with Ago2 and blocking cleavage of target sequences (127). As VSRs provide counter-defense against host antiviral RNAi, both VSRs and their targets in the siRNA pathway (Dcr2, R2D2, and Ago2) evolve rapidly (93). This rapid evolution can result in VSRs that are host specific. For example, the Nora-like virus VP1 binds *Drosophila immigrans* Ago2 but not *D. melanogaster* Ago2 (127). This observation highlights the importance of studying VSR activity in a relevant host species, rather than using a more tractable model species.

In addition to serving as hosts for mosquito-specific viruses, some mosquitoes also vector viruses (arthropod-borne viruses, or arboviruses) of medical and veterinary importance. VSRs have been identified from arboviruses in the genera *Alphaviridae* (120) and *Flaviviridae* (50). *Nodamura virus* (*Nodaviridae*) is a mosquito-transmitted arbovirus that, similar to FHV, encodes a B2 VSR protein that binds dsRNA and siRNAs (120). *Dengue virus* nonstructural protein 4B (NS4B) is reported to function as a VSR, although the mechanism is unclear (50).

Three VSRs have been identified in insect DNA viruses. *Heliothis virescens ascovirus* encodes an RNase III that degrades siRNAs (45). *Invertebrate iridescent virus 6* (IIV-6) encodes a VSR, 340R, that binds siRNAs and dsRNA. This VSR rescues FHV lacking its VSR, confirming the ability of this protein to antagonize antiviral RNAi (8). IIV-6 completely masks the impact of the siRNA pathway in *Drosophila* (10). The inhibitor of apoptosis, P35, derived from the baculovirus *Autographa californica multiple nucleopolyhedrovirus* (AcMNPV) also functions as a VSR acting downstream of cleavage by Dcr2 (74).

## Decoy RNAs

*West Nile virus* (WNV) and *Dengue virus* are mosquito-transmitted flaviviruses. These viruses produce noncoding subgenomic flavivirus RNAs (sfRNAs) as a result of incomplete degradation of

genomic RNA (112). The stem-loop structures of these sfRNAs may act as competitors or decoys for Dcr2 activity, thereby decreasing the efficiency of the antiviral RNAi response. Similarly, hot spots in the genome that were more frequently targeted were identified based on siRNA profiles of two mosquito cell lines following infection with the alphavirus *Semliki Forest virus*. These areas may serve as decoy RNA sequences, reducing the efficacy of the antiviral response by providing regions that are more susceptible to siRNA-mediated RNAi based on structure, but less damaging to the replicating virus (118). This phenomenon was also noted for the phlebovirus *Rift Valley fever virus*, whose defective interfering particle and S segment were heavily targeted for siRNA production but were not stabilized for Ago2 loading (108). This putative decoy strategy represents an alternative mechanism for evasion of RNAi-based antiviral defense.

### **Viral miRNA-Based Manipulation of the Host Genome**

miRNAs can also influence the outcome of virus infection (82). Some viruses encode miRNAs that target host or viral mRNA with proviral consequences. Viral miRNAs that target viral mRNA can modulate the timing of viral replication or gene expression. For example, *Heliothis virescens ascovirus* miR-1 reduces viral replication, thus delaying viral replication and promoting long-term virus production (46). Similarly, AcMNPV miR-1 targets multiple viral proteins, including ODV-E25, to reduce budded virus production (140, 141), and *Bombyx mori nucleopolyhedrovirus* (BmNPV) miR-3 reduces late gene expression to avoid early detection by the host (115).

Viral miRNAs that regulate host gene function include WNV kun-miR-1, which targets and upregulates *GATA4* to the benefit of the virus (47). BmNPV miR-1 impacts miRNA biogenesis, resulting in increased virus replication by targeting the host gene *Ran* that functions in export of pre-miRNAs from the nucleus into the cytoplasm (116). The reduced miRNAs seen in *D. melanogaster* infection with DCV could result from a similar impact on miRNA biogenesis (83).

## **PRACTICAL APPLICATIONS AND IMPLICATIONS**

### **Practical Applications for Viral Suppressors of RNA**

Knowledge of molecular factors contributing to insect virus–RNAi interactions provide opportunities for novel applications. VSRs such as CrPV 1A and FHV B2 can be used ectopically for suppression of the siRNA pathway to reveal latent infections in cell lines (13, 14) or to promote replication of other RNA or DNA viruses (57). VSRs also provide tools to investigate the molecular mechanisms of antiviral RNAi: A single-molecule approach has been employed to elucidate how VSRs discriminate between viral and cellular RNA (25). VSRs can also be used to track dsRNA in live cells (84).

### **RNAi for Optimization of Baculovirus Expression**

RNAi can be used to optimize protein expression from insect virus–based vectors such as the baculovirus expression system (17, 101). For example, insect cell lines stably transformed to downregulate *caspase-1* by siRNA-based RNAi had prolonged resistance to apoptosis following baculovirus infection, thereby increasing the duration of protein expression (131). Similarly, siRNA-based RNAi has been used to restrict baculovirus gp64 production, reducing the number of baculovirus budded virus contaminants present in candidate vaccines (61), and to improve glycosylation of baculovirus-expressed therapeutic proteins (87). Although investigation of insect virus–derived VSRs is relatively young (42), the potential value of VSRs for improved baculovirus expression of recombinant proteins was demonstrated by use of plant virus–derived VSRs (21, 67).

## Virus-Induced Gene Silencing

The efficacy of dsRNA for gene silencing in insect pests can be severely restricted by nucleases in the saliva or gut that degrade dsRNA. As the sRNA viruses of insects have evolved to overcome this enzymatic challenge, the viral particle could have potential use as an enzyme-resistant dsRNA delivery system. VIGS is the delivery of silencing RNAs from a virus. The use of insect virus-based VIGS vectors in pest management is in its infancy, largely due to the limited number of infectious clones of insect RNA viruses. However, multiple plant virus-based VIGS vectors have been developed, and some have been employed to silence essential genes in crop pests (106). In this system, the plant virus produces silencing RNAs in the host plant that are ingested by insects feeding on the plant, so-called host-induced gene silencing. For example, on infection of *Nicotiana benthamiana*, a *Tobacco rattle virus* VIGS vector produced silencing RNAs targeting three *Manduca sexta* gut cytochrome P450 genes (59). A *Citrus tristeza virus* vector was used to target the abnormal wing disc gene of the Asian citrus psyllid, *Diaphorina citri* (38); a *Tobacco mosaic virus* vector was used to target genes in the potato psyllid, *Bactericera cockerelli*, and citrus mealybug, *Planococcus citri* (55, 136); and a *Potato virus X* vector was used to silence genes in the cotton mealybug, *Phenacoccus solenopsis* (54).

## NEEDS FOR FUTURE RESEARCH

### Natural Pathosystems

Studies of antiviral mechanisms in insects have traditionally relied on a small number of model insects such as *D. melanogaster* and *A. aegypti*. These investigations have often focused on viruses with direct relevance to human health or agriculture, and in many cases, the viruses under study are not natural pathogens of the experimental host. Furthermore, most studies have delivered viruses via intrathoracic injection, bypassing the hosts' natural barriers to viral infection. While these approaches have revealed a wealth of knowledge regarding host–virus interactions, they provide an incomplete and biased picture. Indeed, several recent studies have highlighted the importance of the route of infection for the outcome of infection (4, 37, 73, 79). Different routes of infection are also known to trigger different immune responses. For example, the Toll pathway is required for resistance to oral infection with DCV, CrPV, or FHV in *D. melanogaster* but does not appear to play a role during systemic infections (26). The observation that *D. melanogaster* flies can clear DCV infection independent of RNAi underscores the fact that different antiviral mechanisms are required to control virus infection following different routes of exposure (79). More work is needed to understand both the antiviral responses triggered by different routes of infection and the mechanisms by which these responses restrict virus replication.

Natural host–pathogen systems are shaped by coevolution, and natural pathogens can be subject to more structured and effective immune responses (26, 73). Susceptibility to virus infection also varies among different strains of the same insect species; thus, results obtained from common laboratory strains may not necessarily reflect the situation in nature (96). Such differences highlight the importance of studying host–virus interactions in a more natural context to provide more robust and relevant biological data.

### Antiviral RNAi: Inducible or Constitutive?

A core feature of immune mechanisms is the recognition of pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors, resulting in the induction of innate immune responses. Several cell signaling pathways, including the JAK-STAT, Toll, and Imd pathways, are induced upon pathogen recognition as part of the insect immune system. dsRNA is known

to serve as a potent PAMP in insects, and administration of dsRNA, regardless of the sequence, induces the activation of an antiviral response (11, 22, 68). In some cases, recognition of dsRNA has been attributed to the DExD/H-box helicase of Dcr2, suggesting a possible link between components of the siRNA pathway and non-RNAi-based antiviral responses (22). Despite the central role of RNAi in controlling virus infections in insects, relatively little is known about the regulation of the core components of RNAi. The expression of Dcr2 and Ago2 mRNAs was recently shown to be upregulated in a sequence-independent manner by dsRNA in a variety of systems, and a similar increase in Ago2 and Dcr2 expression was observed upon viral infection in *B. terrestris*, *Apis mellifera*, and *D. melanogaster* (31, 32, 40, 68, 92, 137). Intriguingly, injection of heat-inactivated Zika virus also upregulates the expression of RNAi components in *D. melanogaster* (40). Together, these results suggest that virus presence may be sensed by insects independent of viral replication, and that RNAi is regulated as part of a response that is not virus specific. *D. melanogaster* flies constitutively expressing an active form of dFOXO exhibit elevated expression of Dcr2 and Ago2, establishing a link between stress response signaling and RNAi and providing further support for the inducible nature of RNAi (119). Despite these advances, our understanding of RNAi regulation remains limited. Future studies should address whether the route of infection alters the regulation of RNAi components and whether such regulation takes place at the transcriptional or translational levels. Finally, given the central role of RNAi in controlling virus infections, it will be important to determine whether RNAi is differentially regulated among different insect strains or during infection with different viruses.

### Long-Term Immune Memory Against Viruses

As discussed above, traces of RNA viral genomes persist in DNA form following clearance of orally acquired viral infections in *D. melanogaster* (79). While similar vDNAs are known to serve as potent sources of siRNAs and are required for the establishment of persistent infections in *D. melanogaster* and *A. albopictus*, it is unknown whether vDNAs play a role in immune priming during responses to previously encountered viruses (34, 79, 80). Future investigations should seek to clarify the role of vDNAs in immune priming, as well as how the routes of prior infections influence susceptibility to subsequent infection via the same or alternate routes.

Another key question is the duration of immune priming following initial pathogen exposure. Does transgenerational immune priming or even heritable immune priming occur? If so, what is the nature of the archived pathogen specificity determinant, and do the mechanisms of its mobilization in subsequent generations differ from those in the generation actually exposed to the pathogen?

While many questions remain regarding the ability of vDNAs to mediate immune memory, it is now clear that vDNAs play a central role in RNAi-based antiviral responses. Recent results indicate that the majority of vDNAs are derived from reverse transcription of defective viral genomes (DVGs), i.e., truncated or rearranged viral genomes produced during viral replication (100). Modulation of DVG production alters the antiviral response in *D. melanogaster*, with higher levels of DVGs being associated with increased vDNA synthesis and an enhanced antiviral response (100). Thus, future studies should examine the dynamics of DVG production during infection, with a particular focus on whether manipulation of DVGs plays a role in long-term immune processes.

### Reprogramming Endogenous Viral Elements as Antivirals

Additional studies are needed to evaluate the role of EVEs in antiviral immune responses, as well as the mechanisms of EVE generation. While genomic integration of vDNAs is a likely scenario for the endogenization of virus sequences, this has yet to be experimentally demonstrated. Furthermore, it is conceivable that different types of viruses (DNA versus RNA) may

become endogenized by one or more alternative mechanisms, and the route of endogenization may influence the functionality of integrated virus sequences. Beyond the mechanisms of EVE generation, the role of EVEs in antiviral responses is still unclear. Given the potential for EVEs to serve as heritable markers of pathogen specificity, it is important to clarify their role during infection with cognate viruses. If it is indeed possible for EVEs to mediate antiviral responses, then their heritable nature provides a potentially valuable means of manipulating insect genomes to modulate susceptibility to virus infection.

## Infectious Clones

While infectious clones for FHV and *Sindbis virus* have proven to be valuable research tools, their extensive host ranges preclude their use as VIGS vectors for pest management. Indeed, the first demonstration in a proof of concept study was for the use of an FHV-based vector in *Drosophila* (122). Infectious clones available for bona fide insect viruses include those for CrPV (53), which has a broad host range, and *Deformed wing virus* of honey bees (60, 107). The lack of insect virus infectious clones results in part from the lack of virus-free cell lines derived from key crop pests that would support replication of candidate viruses of interest. The development of infectious clones for the mosquito-specific flaviviruses *Niénokoué virus* (49) and *Eilat virus* (88) may ultimately allow for their use in combatting mosquito-borne disease (including flaviviruses dengue and Zika) or reducing mosquito populations through antiviral RNAi-mediated targeting.

## CONCLUSION

While significant progress has been made toward increased understanding of the interplay between insect viruses and the host immune response, significant challenges remain to be addressed, not only from the mechanistic point of view (how virus infection is recognized, how the host nutritional status and stress affect the antiviral RNAi-mediated response), but also from a translational perspective for future application of this knowledge in the control of agricultural pests and vectors of diseases, as well as the protection of beneficial insects. There is tremendous potential for powerful biotechnological applications to result from research on insect RNAi pathways.

### SUMMARY POINTS

1. The host siRNA pathway and, possibly, the piRNA pathway are involved in antiviral defense against insect viruses.
2. Both DNA and RNA viruses have evolved suppressors of the siRNA pathway. These suppressors are most important for RNA virus replication, reflecting the primary role of the siRNA pathway in antiviral defense against RNA viruses.
3. As a nucleic acid-based immune system, the siRNA pathway confers long-term immune memory against the same pathogen. This memory may be inherited by the progeny.
4. Virus-derived decoy RNAs may provide an alternative mechanism for evasion of siRNA-based antiviral defense.
5. Virus-derived miRNA can manipulate the host genome to benefit virus replication.
6. Virus-derived DNA sequence, from which transcripts feed into the siRNA pathway, provides insects with immune memory.

7. Virus-derived DNA sequences that are integrated into host genomes, inherited, and retained as endogenous viral elements produce piRNAs that may function in antiviral immunity.
8. The interplay between insect viruses and RNAi has been exploited for practical purposes, including optimization of the baculovirus expression vector system and use of VSRs to detect covert viruses in insect cell lines.
9. Infectious clones of insect viruses have proven invaluable for the study of virus–host molecular interactions and may have future utility for VIGS-based suppression of insect pests and arboviral disease.

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

The authors thank Jared Nigg, Vanesa Mongelli, Bryan Arias, and Grace Crummer for helpful comments. B.C.B. acknowledges support from grants from the National Science Foundation Industry-University Cooperative Research Center, the Center for Arthropod Management Technologies under grants IIP-1338775 and 1821914; from industry partners; and from the USDA/NIFA 2019-06119, Hatch, and State of Florida funds. M.-C.S. is funded by the European Research Council (FP7/2013–2019 ERC CoG 615220) and the French Government’s Investissement d’Avenir program Laboratoire d’Excellence Integrative Biology of Emerging Infectious Diseases (grant ANR-10-LABX-62-IBEID).

## LITERATURE CITED

1. Adelman ZN, Blair CD, Carlson JO, Beaty BJ, Olson KE. 2001. Sindbis virus-induced silencing of dengue viruses in mosquitoes. *Insect. Mol. Biol.* 10:265–73
2. Aliyari R, Wu Q, Li HW, Wang XH, Li F, et al. 2008. Mechanism of induction and suppression of antiviral immunity directed by virus-derived small RNAs in *Drosophila*. *Cell Host Microbe* 4:387–97
3. Bartel DP. 2009. MicroRNAs: target recognition and regulatory functions. *Cell* 136:215–33
4. Behrens S, Peuß R, Milutinović B, Eggert H, Esser D, et al. 2014. Infection routes matter in population-specific responses of the red flour beetle to the entomopathogen *Bacillus thuringiensis*. *BMC Genom.* 15:445
5. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* 409:363–66
6. Berry B, Deddouche S, Kirschner D, Imler JL, Antoniewski C. 2009. Viral suppressors of RNA silencing hinder exogenous and endogenous small RNA pathways in *Drosophila*. *PLOS ONE* 4:e5866
7. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, et al. 2007. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* 128:1089–103
8. Bronkhorst AW, van Cleef KW, Venselaar H, van Rij RP. 2014. A dsRNA-binding protein of a complex invertebrate DNA virus suppresses the *Drosophila* RNAi response. *Nucleic Acids Res.* 42:12237–48
9. Bronkhorst AW, van Cleef KWR, Vodovar N, İnce İA, Blanc H, et al. 2012. The DNA virus *Invertebrate iridescent virus 6* is a target of the *Drosophila* RNAi machinery. *PNAS* 109:E3604–13

10. Bronkhorst AW, Vogels R, Overheul GJ, Pennings B, Gausson-Dorey V, et al. 2019. A DNA virus-encoded immune antagonist fully masks the potent antiviral activity of RNAi in *Drosophila*. *PNAS* 116:24296–302
11. Brutscher LM, Daughenbaugh KF, Flenniken ML. 2017. Virus and dsRNA-triggered transcriptional responses reveal key components of honey bee antiviral defense. *Sci. Rep.* 7:6448
12. Campbell CL, Black WC, Hess AM, Foy BD. 2008. Comparative genomics of small RNA regulatory pathway components in vector mosquitoes. *BMC Genom.* 9:425
13. Carrillo-Tripp J, Dolezal AG, Goblirsch MJ, Miller WA, Toth AL, Bonning BC. 2016. In vivo and in vitro infection dynamics of honey bee viruses. *Sci. Rep.* 6:22265
14. Carrillo-Tripp J, Krueger EN, Harrison RL, Toth AL, Miller WA, Bonning BC. 2014. *Lymantria dispar iflavivirus 1* (LdIV1), a new model to study iflaviral persistence in lepidopterans. *J. Gen. Virol.* 95:2285–96
15. Cenik ES, Zamore PD. 2011. Argonaute proteins. *Curr. Biol.* 21:R446–49
16. Chao JA, Lee JH, Chapados BR, Debler EW, Schneemann A, Williamson JR. 2005. Dual modes of RNA-silencing suppression by *Flock House virus* protein B2. *Nat. Struct. Mol. Biol.* 12:952–57
17. Chavez-Pena C, Kamen AA. 2018. RNA interference technology to improve the baculovirus-insect cell expression system. *Biotechnol. Adv.* 36:443–51
18. Cora E, Pandey RR, Xiol J, Taylor J, Sachidanandam R, et al. 2014. The MID-PIWI module of Piwi proteins specifies nucleotide- and strand-biases of piRNAs. *RNA* 20:773–81
19. Cullen BR. 2011. Viruses and microRNAs: RISCy interactions with serious consequences. *Genes Dev.* 25:1881–94
20. Czech B, Hannon GJ. 2016. One loop to rule them all: the ping-pong cycle and piRNA-guided silencing. *Trends Biochem. Sci.* 41:324–37
21. de Oliveira VC, da Silva Morgado F, Ardisson-Araujo DM, Resende RO, Ribeiro BM. 2015. The silencing suppressor (NSs) protein of the plant virus *Tomato spotted wilt virus* enhances heterologous protein expression and baculovirus pathogenicity in cells and lepidopteran insects. *Arch. Virol.* 160:2873–79
22. Deddouche S, Matt N, Budd A, Mueller S, Kemp C, et al. 2008. The DExD/H-box helicase Dicer-2 mediates the induction of antiviral activity in *Drosophila*. *Nat. Immunol.* 9:1425–32
23. Elbashir SM, Lendeckel W, Tuschl T. 2001. RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes Dev.* 15:188–200
24. Elbashir SM, Martinez J, Patkaniowska A, Lendeckel W, Tuschl T. 2001. Functional anatomy of siRNAs for mediating efficient RNAi in *Drosophila melanogaster* embryo lysate. *EMBO J.* 20:6877–88
25. Fareh M, van Lopik J, Katechis I, Bronkhorst AW, Haagsma AC, et al. 2018. Viral suppressors of RNAi employ a rapid screening mode to discriminate viral RNA from cellular small RNA. *Nucleic Acids Res.* 46:3187–97
26. Ferreira ÁG, Naylor H, Esteves SS, Pais IS, Martins NE, Teixeira L. 2014. The Toll-dorsal pathway is required for resistance to viral oral infection in *Drosophila*. *PLOS Pathog.* 10:e1004507
27. Feschotte C. 2008. Transposable elements and the evolution of regulatory networks. *Nat. Rev. Genet.* 9:397–405
28. Förstemann K, Horwich MD, Wee L, Tomari Y, Zamore PD. 2007. *Drosophila* microRNAs are sorted into functionally distinct argonaute complexes after production by dicer-1. *Cell* 130:287–97
29. Fujino K, Horie M, Honda T, Merriman DK, Tomonaga K. 2014. Inhibition of Borna disease virus replication by an endogenous bornavirus-like element in the ground squirrel genome. *PNAS* 111:13175–80
30. Gainetdinov I, Colpan C, Arif A, Cecchini K, Zamore PD. 2018. A single mechanism of biogenesis, initiated and directed by PIWI proteins, explains piRNA production in most animals. *Mol. Cell* 71:775–90
31. Galbraith DA, Yang X, Nino EL, Yi S, Grozinger C. 2015. Parallel epigenomic and transcriptomic responses to viral infection in honey bees (*Apis mellifera*). *PLOS Pathog.* 11:e1004713
32. Garbutt JS, Reynolds SE. 2012. Induction of RNA interference genes by double-stranded RNA: implications for susceptibility to RNA interference. *Insect Biochem. Mol. Biol.* 42:621–28
33. Ghildiyal M, Seitz H, Horwich MD, Li C, Du T, et al. 2008. Endogenous siRNAs derived from transposons and mRNAs in *Drosophila* somatic cells. *Science* 320:1077–81

34. Goic B, Stapleford KA, Frangeul L, Doucet AJ, Gausson V, et al. 2016. Virus-derived DNA drives mosquito vector tolerance to arboviral infection. *Nat. Commun.* 7:12410
35. Goic B, Vodovar N, Mondotte JA, Monot C, Frangeul L, et al. 2013. RNA-mediated interference and reverse transcription control the persistence of RNA viruses in the insect model *Drosophila*. *Nat. Immunol.* 14:396–403
36. Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, et al. 2007. A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science* 315:1587–90
37. Gupta V, Vasanthakrishnan RB, Siva-Jothy J, Monteith KM, Brown SP, Vale PF. 2017. The route of infection determines Wolbachia antibacterial protection in *Drosophila*. *Proc. R. Soc. B* 284:20170809
38. Hajeri S, Killiny N, El-Mohtar C, Dawson WO, Gowda S. 2014. *Citrus tristeza virus*-based RNAi in citrus plants induces gene silencing in *Diaphorina citri*, a phloem-sap sucking insect vector of citrus greening disease (Huanglongbing). *J. Biotechnol.* 176:42–49
39. Han BW, Wang W, Li C, Weng Z, Zamore PD. 2015. piRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production. *Science* 348:817–21
40. Harsh S, Ozakman Y, Kitchen SM, Paquin-Proulx D, Nixon DF, Eleftherianos I. 2018. Dicer-2 regulates resistance and maintains homeostasis against *Zika virus* infection in *Drosophila*. *J. Immunol.* 201:3058–72
41. Hayashi R, Schnabl J, Handler D, Mohn F, Ameres SL, Brennecke J. 2016. Genetic and mechanistic diversity of piRNA 3'-end formation. *Nature* 539:588–92
42. Hodgson JJ, Wenger LW, Clem RJ, Passarelli AL. 2019. Inhibition of dicer activity in lepidopteran and dipteran cells by baculovirus-mediated expression of *Flock House virus* B2. *Sci. Rep.* 9:14494
43. Holmes EC. 2011. The evolution of endogenous viral elements. *Cell Host Microbe* 10:368–77
44. Huang XA, Yin H, Sweeney S, Raha D, Snyder M, Lin H. 2013. A major epigenetic programming mechanism guided by piRNAs. *Dev. Cell* 24:502–16
45. Hussain M, Abraham AM, Asgari S. 2010. An Ascovirus-encoded RNase III autoregulates its expression and suppresses RNA interference-mediated gene silencing. *J. Virol.* 84:3624–30
46. Hussain M, Taft RJ, Asgari S. 2008. An insect virus-encoded microRNA regulates viral replication. *J. Virol.* 82:9164–70
47. Hussain M, Torres S, Schnettler E, Funk A, Grundhoff A, et al. 2012. *West Nile virus* encodes a microRNA-like small RNA in the 3' untranslated region which up-regulates GATA4 mRNA and facilitates virus replication in mosquito cells. *Nucleic Acids Res.* 40:2210–23
48. Iwakawa HO, Tomari Y. 2015. The functions of microRNAs: mRNA decay and translational repression. *Trends Cell Biol.* 25:651–65
49. Junglen S, Korries M, Grasse W, Wieseler J, Kopp A, et al. 2017. Host range restriction of insect-specific flaviviruses occurs at several levels of the viral life cycle. *mSphere* 2:e00375-16
50. Kakumani PK, Ponia SS, S RK, Sood V, Chinnappan M, et al. 2013. Role of RNA interference (RNAi) in dengue virus replication and identification of NS4B as an RNAi suppressor. *J. Virol.* 87:8870–83
51. Katzourakis A, Gifford RJ. 2010. Endogenous viral elements in animal genomes. *PLOS Genet.* 6:e1001191
52. Keene KM, Foy BD, Sanchez-Vargas I, Beaty BJ, Blair CD, Olson KE. 2004. RNA interference acts as a natural antiviral response to O'nyong-nyong virus (Alphavirus; Togaviridae) infection of *Anopheles gambiae*. *PNAS* 101:17240–45
53. Kerr CH, Wang QS, Keatings K, Khong A, Allan D, et al. 2015. The 5' untranslated region of a novel infectious molecular clone of the dicistrovirus *Cricket paralysis virus* modulates infection. *J. Virol.* 89:5919–34
54. Khan AM, Ashfaq M, Khan AA, Naseem MT, Mansoor S. 2018. Evaluation of potential RNA-interference-target genes to control cotton mealybug, *Phenacoccus solenopsis* (Hemiptera: Pseudococcidae). *Insect Sci.* 25:778–86
55. Khan AM, Ashfaq M, Kiss Z, Khan AA, Mansoor S, Falk BW. 2013. Use of recombinant *Tobacco mosaic virus* to achieve RNA interference in plants against the citrus mealybug, *Planococcus citri* (Hemiptera: Pseudococcidae). *PLOS ONE* 8:e73657
56. Khong A, Kerr CH, Yeung CH, Keatings K, Nayak A, et al. 2017. Disruption of stress granule formation by the multifunctional *Cricket paralysis virus* 1A protein. *J. Virol.* 91:e01779-16

57. Khoo CC, Doty JB, Heersink MS, Olson KE, Franz AW. 2013. Transgene-mediated suppression of the RNA interference pathway in *Aedes aegypti* interferes with gene silencing and enhances *Sindbis virus* and *dengue virus* type 2 replication. *Insect Mol. Biol.* 22:104–14
58. Klenerman P, Hengartner H, Zinkernagel RM. 1997. A non-retroviral RNA virus persists in DNA form. *Nature* 390:298–301
59. Kumar P, Pandit SS, Baldwin IT. 2012. *Tobacco rattle virus* vector: a rapid and transient means of silencing *Manduca sexta* genes by plant mediated RNA interference. *PLOS ONE* 7:e31347
60. Lamp B, Url A, Seitz K, Eichhorn J, Riedel C, et al. 2016. Construction and rescue of a molecular clone of *Deformed wing virus* (DWV). *PLOS ONE* 11:e0164639
61. Lee HS, Lee HY, Kim YJ, Jung HD, Choi KJ, et al. 2015. Small interfering (Si) RNA mediated baculovirus replication reduction without affecting target gene expression. *Virus Res.* 199:68–76
62. Lee Y, Ahn C, Han J, Choi H, Kim J, et al. 2003. The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425:415–19
63. Lee Y, Kim M, Han J, Yeom KH, Lee S, et al. 2004. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.* 23:4051–60
64. Lee YS, Nakahara K, Pham JW, Kim K, He Z, et al. 2004. Distinct roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA silencing pathways. *Cell* 117:69–81
65. Lewis SH, Salmela H, Obbard DJ. 2016. Duplication and diversification of dipteran Argonaute genes, and the evolutionary divergence of Piwi and Aubergine. *Genome Biol. Evol.* 8:507–18
66. Lingel A, Simon B, Izaurralde E, Sattler M. 2005. The structure of the *flock house virus* B2 protein, a viral suppressor of RNA interference, shows a novel mode of double-stranded RNA recognition. *EMBO Rep.* 6:1149–55
67. Liu Y, Zhang L, Zhang Y, Liu D, Du E, Yang Z. 2015. Functional analysis of RNAi suppressor P19 on improving baculovirus yield and transgene expression in Sf9 cells. *Biotechnol. Lett.* 37:2159–66
68. Lozano J, Gomez-Orte E, Lee H-J, Belles X. 2012. Super-induction of Dicer-2 expression by alien double-stranded RNAs: an evolutionary ancient response to viral infection? *Dev. Genes Evol.* 222:229–35
69. Lund E, Güttinger S, Calado A, Dahlberg JE, Kutay U. 2004. Nuclear export of microRNA precursors. *Science* 303:95–98
70. Ma J-B, Yuan Y-R, Meister G, Pei Y, Tuschl T, Patel DJ. 2005. Structural basis for 5'-end-specific recognition of guide RNA by the *A. fulgidus* Piwi protein. *Nature* 434:666–70
71. Maori E, Tanne E, Sela I. 2007. Reciprocal sequence exchange between non-retro viruses and hosts leading to the appearance of new host phenotypes. *Virology* 362:342–49
72. Martinez J, Patkaniowska A, Urlaub H, Lührmann R, Tuschl T. 2002. Single-stranded antisense siRNAs guide target RNA cleavage in RNAi. *Cell* 110:563–74
73. Martins NE, Faria VG, Teixeira L, Magalhães S, Sucena É. 2013. Host adaptation is contingent upon the infection route taken by pathogens. *PLOS Pathog.* 9:e1003601
74. Mehrabadi M, Hussain M, Matindoost L, Asgari S. 2015. The baculovirus antiapoptotic p35 protein functions as an inhibitor of the host RNA interference antiviral response. *J. Virol.* 89:8182–92
75. Mi S, Lee X, Li X, Veldman GM, Finnerty H, et al. 2000. Syncytin is a captive retroviral envelope protein involved in human placental morphogenesis. *Nature* 403:785–89
76. Miesen P, Girardi E, van Rij RP. 2015. Distinct sets of PIWI proteins produce arbovirus and transposon-derived piRNAs in *Aedes aegypti* mosquito cells. *Nucl. Acids Res.* 43:6545–56
77. Miesen P, Joosten J, van Rij RP. 2016. PIWIs go viral: arbovirus-derived piRNAs in vector mosquitoes. *PLOS Pathog.* 12:e1006017
78. Mohn F, Handler D, Brennecke J. 2015. piRNA-guided slicing specifies transcripts for Zucchini-dependent, phased piRNA biogenesis. *Science* 348:812–17
79. Mondotte JA, Gausson V, Frangeul L, Blanc H, Lambrechts L, Saleh MC. 2018. Immune priming and clearance of orally acquired RNA viruses in *Drosophila*. *Nat. Microbiol.* 3:1394–403
80. Mondotte JA, Saleh M-C. 2018. Antiviral immune response and the route of infection in *Drosophila melanogaster*. *Adv. Virus Res.* 100:247–78
81. Monsanto-Hearne V, Johnson KN. 2018. miRNAs in insects infected by animal and plant viruses. *Viruses* 10:354

82. Monsanto-Hearne V, Johnson KN. 2019. miRNA modulation of insect virus replication. *Curr. Issues Mol. Biol.* 34:61–82
83. Monsanto-Hearne V, Tham ALY, Wong ZS, Asgari S, Johnson KN. 2017. *Drosophila* miR-956 suppression modulates Ectoderm-expressed 4 and inhibits viral replication. *Virology* 502:20–27
84. Monsion B, Incarbone M, Hleibieh K, Poignavent V, Ghannam A, et al. 2018. Efficient detection of long dsRNA in vitro and in vivo using the dsRNA binding domain from FHV B2 protein. *Front. Plant Sci.* 9:70
85. Mueller S, Gausson V, Vodovar N, Deddouche S, Troxler L, et al. 2010. RNAi-mediated immunity provides strong protection against the negative-strand RNA vesicular stomatitis virus in *Drosophila*. *PNAS* 107:19390–95
86. Mussabekova A, Daeffler L, Imler J-L. 2017. Innate and intrinsic antiviral immunity in *Drosophila*. *Cell. Mol. Life Sci.* 74:2039–54
87. Nagata Y, Lee JM, Mon H, Imanishi S, Hong SM, et al. 2013. RNAi suppression of beta-N-acetylglucosaminidase (BmFDL) for complex-type N-linked glycan synthesis in cultured silkworm cells. *Biotechnol. Lett.* 35:1009–16
88. Nasar F, Palacios G, Gorchakov RV, Guzman H, Da Rosa AP, et al. 2012. *Eilat virus*, a unique alphavirus with host range restricted to insects by RNA replication. *PNAS* 109:14622–27
89. Nayak A, Berry B, Tassetto M, Kunitomi M, Acevedo A, et al. 2010. *Cricket paralysis virus* antagonizes Argonaute 2 to modulate antiviral defense in *Drosophila*. *Nat. Struct. Mol. Biol.* 17:547–54
90. Nayak A, Kim DY, Trnka MJ, Kerr CH, Lidsky PV, et al. 2018. A viral protein restricts *Drosophila* RNAi immunity by regulating Argonaute activity and stability. *Cell Host Microbe* 24:542–57.e9
91. Nishimasu H, Ishizu H, Saito K, Fukuhara S, Kamatani MK, et al. 2012. Structure and function of Zucchini endoribonuclease in piRNA biogenesis. *Nature* 491:284–87
92. Niu J, Smagghe G, De Coninck DIM, Van Nieuwerburgh F, Deforce D, Meeus I. 2016. In vivo study of Dicer-2-mediated immune response of the small interfering RNA pathway upon systemic infections of virulent and avirulent viruses in *Bombyx terrestris*. *Insect Biochem. Mol. Biol.* 70:127–37
93. Obbard DJ, Gordon KH, Buck AH, Jiggins FM. 2009. The evolution of RNAi as a defence against viruses and transposable elements. *Philos. Trans. R. Soc. Lond. B* 364:99–115
94. Orban TI, Izaurralde E. 2005. Decay of mRNAs targeted by RISC requires XRN1, the Ski complex, and the exosome. *RNA* 11:459–69
95. Palatini U, Miesen P, Carballar-Lejarazu R, Ometto L, Rizzo E, et al. 2017. Comparative genomics shows that viral integrations are abundant and express piRNAs in the arboviral vectors *Aedes aegypti* and *Aedes albopictus*. *BMC Genom.* 18:512
96. Palmer WH, Varghese FS, Van Rij RP. 2018. Natural variation in resistance to virus infection in dipteran insects. *Viruses* 10:118
97. Parry R, Bishop C, De Hayr L, Asgari S. 2019. Density-dependent enhanced replication of a densovirus in Wolbachia-infected *Aedes* cells is associated with production of piRNAs and higher virus-derived siRNAs. *Virology* 528:89–100
98. Petit M, Mongelli V, Frangeul L, Blanc H, Jiggins F, Saleh M-C. 2016. piRNA pathway is not required for antiviral defense in *Drosophila melanogaster*. *PNAS* 113:E4218–27
99. Petrillo JE, Venter PA, Short JR, Gopal R, Deddouche S, et al. 2013. Cytoplasmic granule formation and translational inhibition of nodaviral RNAs in the absence of the double-stranded RNA binding protein B2. *J. Virol.* 87:13409–21
100. Poirier EZ, Goic B, Tomé-Poderti L, Frangeul L, Boussier J, et al. 2018. Dicer-2-dependent generation of viral DNA from defective genomes of RNA viruses modulates antiviral immunity in insects. *Cell Host Microbe* 23:353–65
101. Possee RD, Chambers AC, Graves LP, Aksular M, King LA. 2019. Recent developments in the use of baculovirus expression vectors. *Curr. Issues Mol. Biol.* 34:215–30
102. Qi N, Cai D, Qiu Y, Xie J, Wang Z, et al. 2011. RNA binding by a novel helical fold of b2 protein from *Wuhan nodavirus* mediates the suppression of RNA interference and promotes b2 dimerization. *J. Virol.* 85:9543–54
103. Qi N, Zhang L, Qiu Y, Wang Z, Si J, et al. 2012. Targeting of dicer-2 and RNA by a viral RNA silencing suppressor in *Drosophila* cells. *J. Virol.* 86:5763–73

104. Rand TA, Ginalski K, Grishin NV, Wang X. 2004. Biochemical identification of Argonaute 2 as the sole protein required for RNA-induced silencing complex activity. *PNAS* 101:14385–89
105. Rand TA, Petersen S, Du F, Wang X. 2005. Argonaute2 cleaves the anti-guide strand of siRNA during RISC activation. *Cell* 123:621–29
106. Rosa C, Kuo YW, Wuriyanghan H, Falk BW. 2018. RNA interference mechanisms and applications in plant pathology. *Annu. Rev. Phytopathol.* 56:581–610
107. Ryabov EV, Childers AK, Lopez D, Grubbs K, Posada-Florez F, et al. 2019. Dynamic evolution in the key honey bee pathogen *deformed wing virus*: novel insights into virulence and competition using reverse genetics. *PLOS Biol.* 17:e3000502
108. Sabin LR, Zheng Q, Thekkat P, Yang J, Hannon GJ, et al. 2013. Dicer-2 processes diverse viral RNA species. *PLOS ONE* 8:e55458
109. Santos D, Mingels L, Vogel E, Wang L, Christiaens O, et al. 2019. Generation of virus- and dsRNA-derived siRNAs with species-dependent length in insects. *Viruses* 11:738
110. Santos D, Wynant N, Van den Brande S, Verdonckt TW, Mingels L, et al. 2018. Insights into RNAi-based antiviral immunity in Lepidoptera: acute and persistent infections in *Bombyx mori* and *Trichoplusia ni* cell lines. *Sci. Rep.* 8:2423
111. Schnettler E, Donald CL, Human S, Watson M, Siu RWC, et al. 2013. Knockdown of piRNA pathway proteins results in enhanced *Semliki Forest virus* production in mosquito cells. *J. Gen. Virol.* 94:1680–89
112. Schnettler E, Sterken MG, Leung JY, Metz SW, Geertsema C, et al. 2012. Noncoding flavivirus RNA displays RNA interference suppressor activity in insect and mammalian cells. *J. Virol.* 86:13486–500
113. Schuster S, Miesen P, van Rij RP. 2019. Antiviral RNAi in insects and mammals: parallels and differences. *Viruses* 11:448
114. Schuster S, Zirkel F, Kurth A, van Cleef KW, Drosten C, et al. 2014. A unique nodavirus with novel features: Mosinovirus expresses two subgenomic RNAs, a capsid gene of unknown origin, and a suppressor of the antiviral RNA interference pathway. *J. Virol.* 88:13447–59
115. Singh CP, Singh J, Nagaraju J. 2014. bmnpv-miR-3 facilitates BmNPV infection by modulating the expression of viral P6.9 and other late genes in *Bombyx mori*. *Insect Biochem. Mol. Biol.* 49:59–69
116. Singh J, Singh CP, Bhavani A, Nagaraju J. 2010. Discovering microRNAs from *Bombyx mori* nucleopolyhedrosis virus. *Virology* 407:120–28
117. Siomi H, Siomi MC. 2009. On the road to reading the RNA-interference code. *Nature* 457:396–404
118. Siu RW, Frakoudis R, Simmonds P, Donald CL, Chase-Topping ME, et al. 2011. Antiviral RNA interference responses induced by *Semliki Forest virus* infection of mosquito cells: characterization, origin, and frequency-dependent functions of virus-derived small interfering RNAs. *J. Virol.* 85:2907–17
119. Spellberg MJ, Marr MT. 2015. FOXO regulates RNA interference in *Drosophila* and protects from RNA virus infection. *PNAS* 112:14587–92
120. Sullivan CS, Ganem D. 2005. A virus-encoded inhibitor that blocks RNA interference in mammalian cells. *J. Virol.* 79:7371–79
121. Suzuki Y, Kobayashi Y, Horie M, Tomonaga K. 2014. Origin of an endogenous bornavirus-like nucleoprotein element in thirteen-lined ground squirrels. *Genes Genet. Syst.* 89:143–48
122. Taning CNT, Christiaens O, Li X, Swevers L, Casteels H, et al. 2018. Engineered *Flock house virus* for targeted gene suppression through RNAi in fruit flies (*Drosophila melanogaster*) in vitro and in vivo. *Front. Physiol.* 9:805
123. Tassetto M, Kunitomi M, Andino R. 2017. Circulating immune cells mediate a systemic RNAi-based adaptive antiviral response in *Drosophila*. *Cell* 169:314–25
124. Tassetto M, Kunitomi M, Whitfield ZJ, Dolan PT, Sánchez-Vargas I, et al. 2019. Control of RNA viruses in mosquito cells through the acquisition of vDNA and endogenous viral elements. *eLife* 8:e41244
125. ter Horst AM, Nigg JC, Dekker FM, Falk BW. 2019. Endogenous viral elements are widespread in arthropod genomes and commonly give rise to PIWI-interacting RNAs. *J. Virol.* 93:e02124–18
126. van Cleef KW, van Mierlo JT, Miesen P, Overheul GJ, Fros JJ, et al. 2014. Mosquito and *Drosophila* entomobirnaviruses suppress dsRNA- and siRNA-induced RNAi. *Nucleic Acids Res.* 42:8732–44
127. van Mierlo JT, Overheul GJ, Obadia B, van Cleef KW, Webster CL, et al. 2014. Novel *Drosophila* viruses encode host-specific suppressors of RNAi. *PLOS Pathog.* 10:e1004256

128. van Rij RP, Saleh MC, Berry B, Foo C, Houk A, et al. 2006. The RNA silencing endonuclease Argonaute 2 mediates specific antiviral immunity in *Drosophila melanogaster*. *Genes Dev.* 20:2985–95
129. Varjak M, Dietrich I, Sreenu VB, Till BE, Merits A, et al. 2018. Spindle-e acts antivirally against alphaviruses in mosquito cells. *Viruses* 10:88
130. Varjak M, Maringer K, Watson M, Sreenu VB, Fredericks AC, et al. 2017. *Aedes aegypti* Piwi4 is a non-canonical PIWI protein involved in antiviral responses. *mSphere* 2:e00144-17
131. Wang Q, Zhou Y, Chen K, Ju X. 2016. Suppression of Bm-caspase-1 expression in BmN cells enhances recombinant protein production in a baculovirus expression vector system. *Mol. Biotechnol.* 58:319–27
132. Wang W, Han BW, Tipping C, Ge DT, Zhang Z, et al. 2015. Slicing and binding by Ago3 or Aub trigger Piwi-bound piRNA production by distinct mechanisms. *Mol. Cell* 59:819–30
133. Wang XH, Aliyari R, Li WX, Li HW, Kim K, et al. 2006. RNA interference directs innate immunity against viruses in adult *Drosophila*. *Science* 312:452–54
134. Watanabe M, Iwakawa HO, Tadakuma H, Tomari Y. 2017. Biochemical and single-molecule analyses of the RNA silencing suppressing activity of CrPV-1A. *Nucleic Acids Res.* 45:10837–44
135. Whitfield ZJ, Dolan PT, Kunitomi M, Tassetto M, Seetin MG, et al. 2017. The diversity, structure, and function of heritable adaptive immunity sequences in the *Aedes aegypti* genome. *Curr. Biol.* 27:3511–19
136. Wuriyanghan H, Falk BW. 2013. RNA interference towards the potato psyllid, *Bactericera cockerelli*, is induced in plants infected with recombinant *Tobacco mosaic virus* (TMV). *PLOS ONE* 8:e66050
137. Ye C, An X, Jiang Y-D, Ding B-Y, Shang F, et al. 2019. Induction of RNAi core machinery's gene expression by exogenous dsRNA and the effects of pre-exposure to dsRNA on the gene silencing efficiency in the pea aphid (*Acyrtosiphon pisum*). *Front. Physiol.* 9:1906
138. Zambon RA, Vakharla VN, Wu LP. 2006. RNAi is an antiviral immune response against a dsRNA virus in *Drosophila melanogaster*. *Cell. Microbiol.* 8:880–89
139. Zhu KY, Palli SR. 2020. Mechanisms, applications, and challenges of insect RNA interference. *Annu. Rev. Entomol.* 65:293–311
140. Zhu M, Wang J, Deng R, Wang X. 2016. Functional regulation of an *Autographa californica* nucleopolyhedrovirus-encoded microRNA, AcMNPV-miR-1, in baculovirus replication. *J. Virol.* 90:6526–37
141. Zhu M, Wang J, Deng R, Xiong P, Liang H, Wang X. 2013. A microRNA encoded by *Autographa californica* nucleopolyhedrovirus regulates expression of viral gene ODV-E25. *J. Virol.* 87:13029–34

# Contents

Volume 66, 2021

Preference Provides a Plethora of Problems (Don't Panic) <i>Michael C. Singer</i> .....	1
A Century of Synergy in Termite Symbiosis Research: Linking the Past with New Genomic Insights <i>Michael E. Scharf and Brittany F. Peterson</i> .....	23
Chemical Ecology, Biochemistry, and Molecular Biology of Insect Hydrocarbons <i>Gary J. Blomquist and Matthew D. Ginzel</i> .....	45
The Interplay Between Viruses and RNAi Pathways in Insects <i>Bryony C. Bonning and Maria-Carla Saleh</i> .....	61
Growing Up in a Changing World: Environmental Regulation of Development in Insects <i>Christen K. Mirth, Timothy E. Saunders, and Christopher Amourda</i> .....	81
Semiochemicals for Thrips and Their Use in Pest Management <i>William D.J. Kirk, Willem Jan de Kogel, Elisabeth H. Koschier, and David A.J. Teulon</i> .....	101
Mechanisms of Resistance to Insecticidal Proteins from <i>Bacillus</i> <i>thuringiensis</i> <i>Juan Luis Jurat-Fuentes, David G. Heckel, and Juan Ferré</i> .....	121
Emergence of <i>Maruca vitrata</i> as a Major Pest of Food Legumes and Evolution of Management Practices in Asia and Africa <i>Ramasamy Srinivasan, Manuele Tamò, and Periasamy Malini</i> .....	141
Survive a Warming Climate: Insect Responses to Extreme High Temperatures <i>Chun-Sen Ma, Gang Ma, and Sylvain Pincebourde</i> .....	163
Honey as a Functional Food for <i>Apis mellifera</i> <i>May R. Berenbaum and Bernarda Calla</i> .....	185
Population Dynamics of Chewing Lice (Phthiraptera) Infesting Birds (Aves) <i>Terry D. Galloway and Robert J. Lamb</i> .....	209

Spider Diversification Through Space and Time <i>Dimitar Dimitrov and Gustavo Hormiga</i> .....	225
How Dung Beetles Steer Straight <i>Marie Dacke, Emily Baird, Basil el Jundi, Eric J. Warrant, and Marcus Byrne</i> .....	243
Laboulbeniomycetes: Intimate Fungal Associates of Arthropods <i>Danny Haelewaters, Meredith Blackwell, and Donald H. Pfister</i> .....	257
Tree Diversity and Forest Resistance to Insect Pests: Patterns, Mechanisms, and Prospects <i>Hervé Jactel, Xoaquín Moreira, and Bastien Castagneyrol</i> .....	277
Symbiont-Mediated Digestion of Plant Biomass in Fungus-Farming Insects <i>Hongjie Li, Soleil E. Young, Michael Poulsen, and Cameron R. Currie</i> .....	297
Navigation Along Windborne Plumes of Pheromone and Resource-Linked Odors <i>Ring T. Cardé</i> .....	317
Behaviors and Interactions of Insects in Mid-Mesozoic Ecosystems of Northeastern China <i>Taiping Gao, Chungkun Shieh, and Dong Ren</i> .....	337
Transposable Elements and the Evolution of Insects <i>Clément Gilbert, Jean Peccoud, and Richard Cordaux</i> .....	355
The Impact of Climate Change on Ticks and Tick-Borne Disease Risk <i>Lucy Gilbert</i> .....	373
Insect Transmission of Plant Single-Stranded DNA Viruses <i>Xiao-Wei Wang and Stéphane Blanc</i> .....	389
Engineering the Composition and Fate of Wild Populations with Gene Drive <i>Bruce A. Hay, Georg Oberhofer, and Ming Guo</i> .....	407
Evolution of Insect Color Vision: From Spectral Sensitivity to Visual Ecology <i>Casper J. van der Kooi, Doekele G. Stavenga, Kentaro Arikawa, Gregor Belušić, and Almut Kelber</i> .....	435
Biological Control with <i>Trichogramma</i> in China: History, Present Status, and Perspectives <i>Lian-Sheng Zang, Su Wang, Fan Zhang, and Nicolas Desneux</i> .....	463
Advancing Undergraduate Laboratory Education Using Non-Model Insect Species <i>Christopher W. Beck and Lawrence S. Blumer</i> .....	485