



Cite this article: Ruiz C, Theobald JC. 2020 Ventral motion parallax enhances fruit fly steering to visual sideslip. *Biol. Lett.* **16**: 20200046. <http://dx.doi.org/10.1098/rsbl.2020.0046>

Received: 27 January 2020

Accepted: 21 April 2020

Subject Areas:

behaviour, neuroscience, cognition

Keywords:

parallax, insect vision, optomotor response, optic flow, *Drosophila melanogaster*

Author for correspondence:

Carlos Ruiz

e-mail: cruiz093@fiu.edu

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.4966274>.

Ventral motion parallax enhances fruit fly steering to visual sideslip

Carlos Ruiz and Jamie C. Theobald

Department of Biological Sciences, Florida International University, Miami, FL 33199, USA

CR, 0000-0001-7595-0180

Flies and other insects use incoherent motion (parallax) to the front and sides to measure distances and identify obstacles during translation. Although additional depth information could be drawn from below, there is no experimental proof that they use it. The finding that blowflies encode motion disparities in their ventral visual fields suggests this may be an important region for depth information. We used a virtual flight arena to measure fruit fly responses to optic flow. The stimuli appeared below ($n = 51$) or above the fly ($n = 44$), at different speeds, with or without parallax cues. Dorsal parallax does not affect responses, and similar motion disparities in rotation have no effect anywhere in the visual field. But responses to strong ventral sideslip (206° s^{-1}) change drastically depending on the presence or absence of parallax. Ventral parallax could help resolve ambiguities in cluttered motion fields, and enhance corrective responses to nearby objects.

1. Introduction

Flies execute extremely fast and precise aerial manoeuvres, requiring robust corrective responses to handle deviations from course. They use coherent motion from optic flow fields to countersteer against changes in direction and position during flight [1,2]. For this purpose, background motion is decomposed into translational and rotational components that are processed independently by large-field neurons in the lobula plate in flies [1–6], and arthropods in general [7].

Responding to translational self-motion requires nearby visual features, because image speed on the retina varies inversely with object distance [8]. Rotational image speeds, by contrast, are unaffected by distance. Some flies take advantage of this difference by increasing their sensitivity to translation in the frontolateral and subequatorial regions of their eyes, where perceived objects are usually closer during natural flight, while displacing the perception of rotation to the dorsal region [4]. *Drosophila*, for example, can use celestial cues above for evaluating changes in direction (reviewed by Warren *et al.* [9]), while positional tasks such as groundspeed control or responses to sudden changes in position are mostly based on optic flow below and near the horizon [10–12].

Translational optic flow additionally provides cues about the three-dimensional structure of the surroundings, which manifests as image speed being inversely proportional to object distances [8,13,14]. Flies can process motion patterns from the frontolateral regions of the visual field and use them to gauge distances [15,16], and separate objects from background [17–19], both critical tasks for navigating through cluttered environments. But motion depth cues are present outside of the frontal or lateral visual field. When flying low over patchy vegetation, for example, a wealth of information about the spatial distribution of features is available right underneath. Bees use this for altitude control [20–22], but flies, for some reason, do not [10,23]. Whether this is because they fail to integrate relative motion beneath, or shift attention to frontal areas during forward flight [24], remains unknown.




Figure 1. (a) Rear view of the projection arena showing the placement of the fly. The IR LED above illuminates the wings in motion casting a shadow on the sensor below the fly. (b) The stimulus projects onto the ventral or dorsal faces of a Perspex cube using mirrors. (c) Steering attempts are inferred from the difference in the size of shadows of the left and right wings captured by the dual sensor. Mean steering responses of the flies tested can be visualized as time series (solid lines), along with the standard error of the mean (s.e.m.) (shading). (d) Motion parallax in the stimuli is simulated by adding relative motion to the dot-field elements.

The recent finding of a neuron (VT1) in the blowfly *Calliphora vicina*, able to encode motion parallax in the forward and sideslip directions below the horizon [25], provides a partial answer to this question. It demonstrates that at least some groups of flies encode parallax in ventral optic flow, and suggests that this trait could be adaptive to flies traversing habitats with obstacles [26] or foraging for resources on the ground. This could be the case in *Drosophila melanogaster*, a slow flier that searches for fallen fruit.

Despite the abundance of derived traits associated with the lobula plate tangential cells (LPT) across groups of flies [27], horizontal system cells (HS) responsible for assessing yaw rotation are relatively conserved between blowflies and fruit flies [28]. It is therefore possible that they also share the ability to perceive and encode incoherent motion below during flight. We set out to test whether parallax affects the optomotor response of fruit flies during visual perturbations in the ventral or dorsal region of the flow field. We used a virtual flight arena to display perturbations, with and without depth cues, and measure optomotor responses of tethered fruit flies.

2. Material and methods

(a) Experimental subjects

We tested *D. melanogaster* females within 3–5 days of eclosion, reared in the laboratory under a 12 L:12 D cycle, kept at 21°C

and fed standard medium. Flies were cold anaesthetized, then glued to a fine tungsten rod by the mesonotum. They recovered in the dark for at least 30 min while holding a small piece of paper with their legs, preventing them from flapping their wings. We then removed the paper when suspending each fly in the centre of the arena (figure 1a). Each fly was tested only once in an experiment.

(b) Virtual flight arena

We projected visual stimuli onto the lower and upper surfaces of a 200 mm Perspex cube (figure 1b). Perspective-corrected stimuli displayed in a 90° diameter disc. Experiments took place in a dark room to increase contrast, and the sides of the cube prevented flies from getting light from any other direction. Further details of the arena are described in Cabrera & Theobald [29].

(c) Visual stimuli

Each experiment consisted of open-loop presentations of dot-fields moving either rightward (clockwise in rotation) or leftward (anticlockwise) projected to the ventral or dorsal visual region of the fly (see electronic supplementary material, video S1, for details). Dot-field motion was either rotational or sideslip, at one of four different angular speeds, and with or without parallax depth cues (figure 1c). We emulated depth cues by adding relative motion to a randomly distributed group of dots moving in the same direction, suggesting increased distance [12] (figure 1d). This ensured the number of dots was constant




Figure 2. (a,b) Mean response of *D. melanogaster* to unintended sideslip containing only coherent motion (blue), and with relative motion suggesting the presence of parallax (green). The stimuli were presented at four speeds and two directions on the dorsal and ventral regions of the visual field. Responses were averaged between 0.2 and 0.4 s after stimulus onset. (c,d) Time series showing the first 0.4 s of the mean response to the highest speed sideslip (206° s^{-1}) with and without depth, in the dorsal and ventral visual fields. (e,f) Mean responses to rotational stimuli with and without relative motion, presented in the dorsal and ventral visual fields. (g,h) Time series showing the response of the flies to both types of motion at high speed in their dorsal and ventral visual fields. Solid traces represent the mean responses obtained from n flies, shading represents s.e.m.

(113 dots/steradian), and allowed us to add differential speeds to rotational flow fields, which intrinsically have no such feature (see electronic supplementary material, videos S2–S5, for details on the stimuli used). The trials were presented in random order, and interspersed by segments of closed-loop bar fixation to standardize the behavioural state at the beginning of each test [30,31].

(d) Steering responses

Tethered flies were illuminated from above with an infrared light, while photodiodes below measured the shadow produced by each wing beat. Since flies steer by changing the relative amplitudes of left and right wing beats [32], attempts to turn produce a differential voltage by the sensor pair [33,34], which is reported as the voltage difference in wing beat amplitude (ΔWBA). Responses collected include roll and yaw attempts performed by the fly, as they both result from the same flight mechanics and are indistinguishable using a wing beat analyser [2].

3. Results

Flies responded to dorsal and ventral stimuli by steering in the direction of the flow, and increasing amplitude with flow speed. Coherent sideways flow with angular speeds up to 138° s^{-1} elicited responses of similar amplitude when presented dorsally and ventrally (figure 2a,b, blue lines). However, high speeds of sideways flow that suggested stronger disturbances with coherent motion (206° s^{-1}) produced significantly weaker ventral responses ($t = -2.365, p = 0.023$).

Due to the local optimization of the dorsal region of the eye for evaluating rotation [4], if flies are sensitive to parallax, they might respond to it only when it is present in ventral disturbances. Since motion parallax is exclusive to translation, we expected flies to be unresponsive to it when was added to rotation. For optic flow including relative motion (simulating parallax), response to strong ventral sideslip disturbances

increased significantly ($t = -2.313, p = 0.023$) (figure 2d, green trace). This was similar in amplitude to the response to a dorsal stimulus, either in the absence or presence of depth cues ($t = -0.001, p = 0.999$ and $t = -0.227, p = 0.821$, respectively) (figure 2c). Relative motion had no effect on steering response when added to dorsal sideslip (figure 2a, green) or rotation (figure 2e,f). This occurs even at high-speed translational disturbances presented dorsally ($t = 0.23, p = 0.819$) (figure 2a, green), or rotational in both regions of the visual field (figure 2g,h, green) [35].

4. Discussion

(a) Response to dorsal and ventral sideslip disturbances without depth cues

In contrast with hawkmoths *Manduca sexta*, that maintain flight control even with the ventral region of their eyes covered [36], steering responses to positional changes in flies may be strongly based on flow fields below the horizon, as demonstrated in blowflies [4]. In fact, flies respond weakly to translational cues present only in the upper visual hemisphere [12]. However with the narrower dorsal and ventral visual fields shown here, low-speed disturbances containing only coherent motion elicited similar steering responses in both of these regions. Only high-speed disturbances caused a strong difference in the weakening of ventral perturbations without parallax.

Similar steering response shifts are seen during forward flow. Flies shift attention to anterior regions of the ventral flow field as forward flow speed increases [24]. This may alleviate motion blur [37] by focusing attention on areas with slower optic flow, potentially reducing responsiveness to perturbations below. Fast optic flow can also induce spatial summation, forcing the fly to spatially pool information in

lateral regions. This increases the ability to respond to fast stimuli at the cost of spatial resolution [38]. It is unknown if sideways perturbations can trigger such neural strategies.

(b) Response to wide-field incoherent motion

As expected, relative motion cues affected steering responses only in sideslip, and not yaw rotation. Since rotational and translational components of motion are processed separately [1,3,4], encoding relative motion may be a property of large-field neurons, responsible for the translational components of self-motion only. Further, the similarity in the responses to translating dot-fields with and without relative motion in the dorsal region of the eye suggests the presence of a less specialized system for the perception of translation there, in contrast with regions near the horizon where translational cues are more relevant [12]. While incoherent motion is known to be informative in frontolateral regions [15,16], our results extend that range to the ventral region, showing that fruit flies perceive, encode and use depth cues below them. Due to the conservation of traits inherent to the perception of rotation between blowflies and fruit flies [28], we believe elements involved in the integration of depth cues in the ventral flow field in *D. melanogaster* may be homologous to those found by Longden *et al.* [25] in *C. vicina*. Due to strong selective pressures acting on LPT cells [27], the ability to perceive ventral parallax in *D. melanogaster* is a selective trait, with a variety of possibly adaptive roles.

(c) Height control

Although the depth cues from the ventral optic flow could be used for height control when flying over structured environments (bees are an example [20–22,39,40]), fruit flies seem to control their height during flight using information from frontolateral areas of the visual field [10,23], while referring to ventral optic flow in order to control groundspeed [10,41]. Because the effect of relative motion was notable only during strong sideslip, ventral parallax is probably not involved in altitude control, but further experiments with different levels of parallax would be required to rule it out completely.

(d) Dealing with ambiguity

Different types of self-motion can generate identical flow fields when perceived by small regions of the eye [4,42], and partial stimulation of wide-field neurons could therefore be ambiguous.

In our experiment, for example, moving dots in the small ventral visual field could be perceived by the fly as either sideslip or a roll. However, the presence of parallax in the ventral optic flow could confirm that an otherwise ambiguous stimulus results from translation, because incoherent motion is absent from rotation. However, this is complicated because VS neurons sensitive to roll branch out laterally instead of ventrally on the eye in blowflies [43–46], suggesting that lateral motion on a small region of the ventral field is possibly perceived only as translational, which could also apply to *Drosophila*.

(e) Navigating complex environments

Our results suggest the presence of relative motion ventrally, even in a narrow cone of vision, is enough to prevent attention from shifting forward, and keep the fly responsive to potential risks below during strong perturbations. The sense of nearness produced by motion parallax induces stronger corrective responses to sideslip disturbances in fruit flies [29]. The fact that fruit flies share the ability to encode parallax information from ventral flow fields with blow flies is not that surprising. The saprophagous nature of both flies forces them to move around in search of ephemeral resources that can be far apart and usually at ground level. While *C. vicina* is a fast flier that moves across patches of differently structured vegetation [26], *D. melanogaster* can forage longer distances and even migrate if necessary in search of resources [47]. With such a natural history, both species can certainly benefit from being aware of the dangers below when traversing unknown structured environments.

We have demonstrated that fruit flies respond to the presence of parallax during strong sideways disturbances in their ventral optic flow. The robustness of this response suggests that it is an adaptive trait, but its full significance is unresolved.

Data accessibility. Data are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.2jm63xskb> [35].

Authors' contributions. C.R. and J.C.T. designed the experiment and wrote the manuscript. C.R. collected and analysed the data. Both authors gave final approval for publication and agree to be held accountable for the work performed therein.

Competing interests. We declare we have no competing interests.

Funding. This research was supported by a grant from the National Science Foundation, IOS-1750833.

Acknowledgements. We thank Cleopatra Pimienta, Pablo Currea, Nick Palermo and Ravindra Palavalli-Nettimi for ideas and comments on the manuscript.

References

1. Hausen K, Egelhaaf M. 1989 Neural mechanisms of visual course control in insects. In *Facets of vision* (eds DG Stavenga, RC Hardie), pp. 391–424. Berlin, Germany: Springer.
2. Theobald JC, Ringach DL, Frye MA. 2010 Dynamics of optomotor responses in *Drosophila* to perturbations in optic flow. *J. Exp. Biol.* **213**, 1366–1375. (doi:10.1242/jeb.037945)
3. Duistermars BJ, Chow DM, Condron M, Frye MA. 2007 The spatial, temporal and contrast properties of expansion and rotation flight optomotor responses in *Drosophila*. *J. Exp. Biol.* **210**, 3218–3227. (doi:10.1242/jeb.007807)
4. Krapp HG, Hengstenberg R. 1996 Estimation of self-motion by optic flow processing in single visual interneurons. *Nature* **384**, 463–466. (doi:10.1038/384463a0)
5. Borst A, Haag J, Reiff DF. 2010 Fly motion vision. *Annu. Rev. Neurosci.* **33**, 49–70. (doi:10.1146/annurev-neuro-060909-153155)
6. Borst A, Euler T. 2011 Seeing things in motion: models, circuits, and mechanisms. *Neuron* **71**, 974–994. (doi:10.1016/j.neuron.2011.08.031)
7. Mauss AS, Borst A. 2019 Motion vision in arthropods. In *The Oxford handbook of invertebrate neurobiology* (ed. JH Byrne), pp. 318–344. Oxford, UK: Oxford University Press.
8. Srinivasan MV. 1992 Distance perception in insects. *Curr. Dir. Psychol. Sci.* **1**, 22–26. (doi:10.1111/1467-8721.ep10767830)
9. Warren TL, Giraldo YM, Dickinson MH. 2019 Celestial navigation in *Drosophila*. *J. Exp. Biol.* **222**, jeb186148. (doi:10.1242/jeb.186148)
10. David CT. 1979 Optomotor control of speed and height by free-flying *Drosophila*. *J. Exp. Biol.* **82**, 389–392.
11. David CT. 1982 Compensation for height in the control of groundspeed by *Drosophila* in a new,

- 'barber's pole' wind tunnel. *J. Comp. Physiol.* **147**, 485–493. (doi:10.1007/BF00612014)
12. Mazo C, Theobald JC. 2014 To keep on track during flight, fruit flies discount the skyward view. *Biol. Lett.* **10**, 20131103. (doi:10.1098/rsbl.2013.1103)
 13. Horridge GA. 1986 A theory of insect vision: velocity parallax. *Proc. R. Soc. Lond. B* **229**, 13–27. (doi:10.1098/rspb.1986.0071)
 14. Horridge GA. 1987 The evolution of visual processing and the construction of seeing systems. *Proc. R. Soc. Lond. B* **230**, 279–292. (doi:10.1098/rspb.1987.0020)
 15. Wehner R, Horn E. 1975 The effect of object distance on pattern preferences in the walking fly, *Drosophila melanogaster*. *Experientia* **31**, 641–643. (doi:10.1007/bf01944603)
 16. Heisenberg M, Wolf R. 1984 *Vision in Drosophila: genetics of microbehavior*. Berlin, Germany: Springer.
 17. Egelhaaf M. 1985. On the neuronal basis of figure-ground discrimination by relative motion in the visual system of the fly. *Biol. Cybern.* **52**, 123–140. (doi:10.1007/bf00364003)
 18. Egelhaaf M. 1985. On the neuronal basis of figure-ground discrimination by relative motion in the visual system of the fly II: figure-detection cells, a new class of visual interneurons. *Biol. Cybern.* **52**, 195–209. (doi:10.1007/BF00339948)
 19. Egelhaaf M. 1985. On the neuronal basis of figure-ground discrimination by relative motion in the visual system of the fly III: possible input circuitries and behavioural significance of the FD-cells. *Biol. Cybern.* **52**, 267–280. (doi:10.1007/BF00336983)
 20. Lehrer M, Srinivasan MV, Zhang SW, Horridge GA. 1988 Motion cues provide the bee's visual world with a third dimension. *Nature* **332**, 356–357. (doi:10.1038/332356a0)
 21. Srinivasan MV, Zhang SW, Chahl JS, Barth E, Venkatesh S. 2000 How honeybees make grazing landings on flat surfaces. *Biol. Cybern.* **83**, 171–183. (doi:10.1007/s004220000162)
 22. Srinivasan MV, Zhang S, Chahl JS. 2001 Landing strategies in honeybees, and possible applications to autonomous airborne vehicles. *Biol. Bull.* **200**, 216–221. (doi:10.2307/1543319)
 23. Straw AD, Lee S, Dickinson MH. 2010 Visual control of altitude in flying *Drosophila*. *Curr. Biol.* **20**, 1550–1556. (doi:10.1016/j.cub.2010.07.025)
 24. Palermo N, Theobald J. 2019 Fruit flies increase attention to their frontal visual field during fast forward optic flow. *Biol. Lett.* **15**, 20180767. (doi:10.1098/rsbl.2018.0767)
 25. Longden KD, Wicklein M, Hardcastle BJ, Huston SJ, Krapp HG. 2017 Spike burst coding of translatory optic flow and depth from motion in the fly visual system. *Curr. Biol.* **27**, 3225–3236. (doi:10.1016/j.cub.2017.09.044)
 26. Coyler CN, Hammond CO. 1951 *Flies of the British Isles*. London, UK: Warne & Co. Ltd.
 27. Buschbeck EK, Strausfeld NJ. 1997 The relevance of neural architecture to visual performance: phylogenetic conservation and variation in dipteran visual systems. *J. Comp. Neurol.* **383**, 282–304. (doi:10.1002/(SICI)1096-9861(19970707)383:3<282::AID-CNE2>3.0.CO;2-#)
 28. Schnell Á, Joesch M, Forstner F, Raghu SV, Otsuna H, Ito K, Borst A, Reiff DF. 2010 Processing of horizontal optic flow in three visual interneurons of the *Drosophila* brain. *J. Neurophysiol.* **103**, 1646–1657. (doi:10.1152/jn.00950.2009)
 29. Cabrera S, Theobald JC. 2013 Flying fruit flies correct for visual sideslip depending on relative speed of forward optic flow. *Front. Behav. Neurosci.* **7**, 76. (doi:10.3389/fnbeh.2013.00076)
 30. Reichardt W, Wenking H. 1969 Optical detection and fixation of objects by fixed flying flies. *Naturwissenschaften* **56**, 424. (doi:10.1007/bf00593644)
 31. Heisenberg M, Wolf R. 1979 On the fine structure of yaw torque in visual flight orientation of *Drosophila melanogaster*. *J. Comp. Physiol.* **130**, 113–130. (doi:10.1007/bf00611046)
 32. Götz KG. 1968 Flight control in *Drosophila* by visual perception of motion. *Kybernetik* **4**, 199–208. (doi:10.1007/bf00272517)
 33. Götz KG. 1987 Course-control, metabolism and wing interference during ultralong tethered flight in *Drosophila melanogaster*. *J. Exp. Biol.* **128**, 35–46.
 34. Frye MA, Dickinson MH. 2004 Motor output reflects the linear superposition of visual and olfactory inputs in *Drosophila*. *J. Exp. Biol.* **207**, 123–131. (doi:10.1242/jeb.00725)
 35. Ruiz C, Theobald JC. 2020 Data from: Ventral motion parallax enhances fruit fly steering to visual sideslip. Dryad Digital Repository. (<https://doi.org/10.5061/dryad.2jm63xskb>)
 36. Copley S, Parthasarathy K, Willis MA. 2018 Optomotor steering and flight control requires a specific sub-section of the compound eye in the hawkmoth, *Manduca sexta*. *J. Exp. Biol.* **221**, jeb178210. (doi:10.1242/jeb.178210)
 37. Srinivasan MV, Bernard GD. 1975 The effect of motion on visual acuity of the compound eye: a theoretical analysis. *Vision Res.* **15**, 515–525. (doi:10.1016/0042-6989(75)90029-2)
 38. Theobald J. 2017 Optic flow induces spatial filtering in fruit flies. *Curr. Biol.* **27**, R212–R213. (doi:10.1016/j.cub.2017.02.018)
 39. Portelli G, Ruffier F, Franceschini N. 2010 Honeybees change their height to restore their optic flow. *J. Comp. Physiol. A* **196**, 307–313. (doi:10.1007/s00359-010-0510-z)
 40. Portelli G, Serres JR, Ruffier F. 2017 Altitude control in honeybees: joint vision-based learning and guidance. *Sci. Rep.* **7**, 1–10. (doi:10.1038/s41598-017-09112-5)
 41. Kellogg FE, Frizel DE, Wright RH. 1962 The olfactory guidance of flying insects. IV. *Drosophila*. *Can. Ent.* **94**, 884–888. (doi:10.4039/ent94884-8)
 42. Dahmen HJ, Franz MO, Krapp HG. 2001 Extracting egomotion from optic flow: limits of accuracy and neural matched filters. In *Motion vision* (eds JM Zanker, J Zeil), pp. 143–168. Berlin, Germany: Springer.
 43. Eckert H, Bishop LG. 1978 Anatomical and physiological properties of the vertical cells in the third optic ganglion of *Phaenicia sericata* (Diptera, Calliphoridae). *J. Comp. Physiol.* **126**, 57–86. (doi:10.1007/bf01342651)
 44. Hengstenberg, R. 1982 Common visual response properties of giant vertical cells in the lobula plate of the blowfly *Calliphora*. *J. Comp. Physiol.* **149**, 179–193. (doi:10.1007/bf00619212)
 45. Hengstenberg R. 1984 Roll-stabilization during flight of the blowfly's head and body by mechanical and visual cues. In *Localization and orientation in biology and engineering* (eds D Varjú, HU Schnitzler) pp. 121–134. Berlin, Germany: Springer.
 46. Hengstenberg R, Hausen K, Hengstenberg B. 1982 The number and structure of giant vertical cells (VS) in the lobula plate of the blowfly *Calliphora erythrocephala*. *J. Comp. Physiol.* **149**, 163–177. (doi:10.1007/bf00619211)
 47. Coyne JA, Boussy IA, Prout T, Bryant SH, Jones JS, Moore JA. 1982 Long-distance migration of *Drosophila*. *Am. Nat.* **119**, 589–595. (doi:10.1086/283936)