

Weil representation and Arithmetic Fundamental Lemma

Author(s): W. Zhang

Source: *Annals of Mathematics*, May 2021, Vol. 193, No. 3 (May 2021), pp. 863-978

Published by: Mathematics Department, Princeton University

Stable URL: <https://www.jstor.org/stable/10.4007/annals.2021.193.3.5>

REFERENCES

Linked references are available on JSTOR for this article:

https://www.jstor.org/stable/10.4007/annals.2021.193.3.5?seq=1&cid=pdf-reference#references_tab_contents

You may need to log in to JSTOR to access the linked references.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

Mathematics Department, Princeton University is collaborating with JSTOR to digitize, preserve and extend access to *Annals of Mathematics*

Weil representation and Arithmetic Fundamental Lemma

By W. ZHANG

Abstract

We study a partially linearized version of the relative trace formula for the arithmetic Gan–Gross–Prasad conjecture for the unitary group $U(V)$. The linear factor in this relative trace formula provides an SL_2 -symmetry which allows us to prove by induction the arithmetic fundamental lemma over \mathbb{Q}_p when p is odd and $p \geq \dim V$.

Contents

1. Introduction	864
Part 1. Local theory	872
2. FL and variants	873
3. AFL and variants	879
4. Relation between the two versions of AFL	884
5. Local constancy of intersection numbers	896
Part 2. Global theory	900
6. Shimura varieties and their integral models	900
7. Kudla–Rapoport divisors and the derived CM cycles	906
8. Modular generating functions of special divisors	918
9. Local intersection: non-archimedean places	924
10. Local intersection: archimedean places	930
11. Weil representation and RTF	932
12. RTF with Gaussian test functions	940

Keywords: Arithmetic Fundamental Lemma, arithmetic Gan–Gross–Prasad conjecture, Kudla–Rapoport special divisor, relative trace formula, unitary Shimura variety, Weil representation

AMS Classification: Primary: 11F27, 11F67, 11G40, 14C25, 14G35.

Research of W. Zhang is partially supported by the NSF grant DMS #1838118 and #1901642.

© 2021 Department of Mathematics, Princeton University.

Part 3. Proof of the main theorems	953
13. The proof of FL	953
14. The comparison for arithmetic intersections	961
15. The proof of AFL	968
Appendix A. Weil representation commutes with smooth transfer	969
Appendix B. Grothendieck groups for formal schemes	971
References	975

1. Introduction

The theorem of Gross and Zagier [14] relates the Néron–Tate heights of Heegner points on modular curves to the central derivative of certain L -functions. The arithmetic Gan–Gross–Prasad conjecture [9], [46], [40] is a generalization of this theorem to higher-dimensional Shimura varieties. This conjecture is inspired by the (usual) Gan–Gross–Prasad conjecture relating period integrals on classical groups to special values of Rankin–Selberg tensor product L -functions. In [21] Jacquet and Rallis proposed a relative trace formula (RTF) approach to this last conjecture in the case of unitary groups, and there has been much progress along this direction in the past years. Inspired by their approach, in [46] the author proposed a relative trace formula approach to the arithmetic Gan–Gross–Prasad conjecture. This approach reduces the problem to certain local statements, notably the arithmetic fundamental lemma (AFL) conjecture formulated by the author in [46], and the arithmetic transfer (AT) conjecture formulated by Rapoport, Smithling, and the author [37], [38]. The AFL and AT conjectures relate the special values of the derivative of orbital integrals to arithmetic intersection numbers on a Rapoport–Zink formal moduli space (RZ space) of p -divisible groups,

$$\partial\text{Orb}(\gamma, \mathbf{1}_{S_n(O_{F_0})}) = -\text{Int}(g) \cdot \log q;$$

cf. the precise statement of [Conjecture 3.8](#) for the AFL conjecture.

The goal of this paper is to give a proof of the AFL conjecture over $F_0 = \mathbb{Q}_p$ when $p \geq n$ for an open dense subset of regular semisimple elements (i.e., the set of “strongly regular semisimple elements” in the sense of [45]); cf. [Theorem 15.1](#). This restriction is harmless for the relative trace formula approach to the arithmetic Gan–Gross–Prasad conjecture.

In fact, we also obtain a proof of the Jacquet–Rallis fundamental lemma (FL) conjecture over p -adic field, a theorem due to Yun [45] and Gordan [13] for p large, which is an identity between two orbital integrals

$$\text{Orb}(\gamma, \mathbf{1}_{S_n(O_{F_0})}) = \text{Orb}(g, \mathbf{1}_{K_0});$$

cf. the precise statement of [Conjecture 2.3](#). The idea is similar to the proof of the AFL and is easier to explain. For our proof of the FL, the main input is a

study of a “partially linearized” version of the Jacquet–Rallis RTF, which we call a semi-Lie algebra version. This is closely related to the RTF of Yifeng Liu to the Fourier–Jacobi periods/cycles [29], [30]. The advantage of the linearization is to gain more “symmetry,” i.e., there is an “action” on the RTF (changing test functions) by SL_2 under the Weil representation. The SL_2 -modularity plays the role in the global setting of the Fourier transform in the local harmonic analysis, a crucial ingredient in [47] to prove the smooth transfer conjecture of Jacquet–Rallis.

Now we give a little more detail of our approach. Let F_0 be a totally real number field, and F a CM quadratic extension of F_0 . Let V be an F/F_0 -hermitian space with $\dim_F V = n$, and let $\mathrm{U}(V)$ be the associated isometry group (a reductive group over F_0). Consider the (diagonal) action of $\mathrm{U}(V)$ on the product $\mathrm{U}(V) \times V$, where the two factors are viewed as affine varieties over F_0 endowed with the conjugation action and the standard action respectively. For unexplained notation, we refer the reader to [Section 1.2](#) and the main body of the paper. To any Schwartz function $\Phi \in \mathcal{S}((\mathrm{U}(V) \times V)(\mathbb{A}_0))$, we can associate a kernel function

$$\mathcal{K}_\Phi(g) = \sum_{(\delta, u) \in (\mathrm{U}(V) \times V)(F_0)} \Phi(g^{-1}(\delta, u)), \quad g \in \mathrm{U}(V)(\mathbb{A}_0),$$

which is left invariant under $\mathrm{U}(V)(F_0)$. Then, as one usually does in the theory of relative trace formula, one may study the distribution on $(\mathrm{U}(V) \times V)(\mathbb{A}_0)$ (at least for certain nice test functions Φ),

$$\mathbb{I}(\Phi) = \int_{[\mathrm{U}(V)]} \mathcal{K}_\Phi(g) dg.$$

Here $[G] := G(F_0) \backslash G(\mathbb{A}_0)$ for an algebraic group G over F_0 . Similarly, one can start with the (diagonal) action of GL_{n, F_0} on the product $S_n \times V'_n$, where $V'_n = \mathrm{M}_{1, n} \times \mathrm{M}_{n, 1}$ is the product of the space of column and row vectors; cf. [Section 2.1](#). To any Schwartz function $\Phi' \in \mathcal{S}((S_n \times V'_n)(\mathbb{A}_0))$, we have a similar kernel function $\mathcal{K}_{\Phi'}$ and a distribution (for nice test functions Φ')

$$\mathbb{J}(\Phi') = \int_{[\mathrm{GL}_{n, F_0}]} \mathcal{K}_{\Phi'}(g) \eta_{F/F_0} \circ \det(g) dg.$$

By the smooth transfer between Φ and Φ' through their orbital integrals (relative to the group actions here), one can match the distributions \mathbb{I} and \mathbb{J} .

Now, due to the presence of the linear factors V and V'_n respectively, the Weil representation ω of $\mathrm{SL}_2(\mathbb{A}_0)$ acts on $\mathcal{S}((\mathrm{U}(V) \times V)(\mathbb{A}_0))$ and $\mathcal{S}((S_n \times V'_n)(\mathbb{A}_0))$, hence on the distributions \mathbb{I} and \mathbb{J} ,

$$\mathbb{I}(h, \Phi) := \mathbb{I}(\omega(h)\Phi) \quad \text{and} \quad \mathbb{J}(h, \Phi') := \mathbb{J}(\omega(h)\Phi'),$$

where $h \in \mathrm{SL}_2(\mathbb{A}_0)$. Moreover, the action is “modular” in the sense that $h \mapsto \mathbb{I}(h, \Phi)$ and $\mathbb{J}(h, \Phi')$ are left invariant under $\mathrm{SL}_2(F_0)$, as an application of the Poisson summation formula. In other words, we may enrich the kernel function to a two-variable one:

$$\mathcal{K}_\Phi(g, h) = \sum_{(\delta, u) \in (\mathrm{U}(V) \times V)(F_0)} \omega(h) \Phi(g^{-1}(\delta, u)), \quad g \in \mathrm{U}(V)(\mathbb{A}_0), h \in \mathrm{SL}_2(\mathbb{A}_0).$$

The natural question now is how the Weil representation fits into the comparison of the two distributions. From [47] and [43] one can deduce that the Weil representation commutes with smooth transfer; cf. [Theorem A.1](#) in the appendix.

Both distributions \mathbb{I} and \mathbb{J} can be expanded as a sum over orbital integrals. Then the SL_2 -modularity amounts to certain recursive relations between the orbital integrals appearing in \mathbb{I} and \mathbb{J} . One may hope that the recursive relations are ample enough to allow us to extract identities such as the aforementioned fundamental lemma, starting from some simple identities that can be verified directly. This resembles the situation in the geometric approach (cf. [34], [45]) where one also needs to verify some simple cases directly as a starting point before applying the “perverse continuation principle.”

The idea does not work directly to yield a proof of the Jacquet–Rallis FL; however, it does work if we take two additional inputs. The first input is to consider a “slice” of the semi-Lie algebra version. For example, we fix a suitable monic polynomial α and denote by $\mathrm{U}(V)(\alpha)$ the subscheme of $\mathrm{U}(V)$ consisting of elements with characteristic polynomial equal to α . We then introduce a kernel function,

$$\mathcal{K}_{\Phi, \alpha}(g) = \sum_{(\delta, u) \in (\mathrm{U}(V)(\alpha) \times V)(F_0)} \Phi(g^{-1}(\delta, u)), \quad g \in \mathrm{U}(V)(\mathbb{A}_0).$$

Here the sum runs only over a subset of $\mathrm{U}(V)(F_0)$ -orbits on $(\mathrm{U}(V) \times V)(F_0)$. Similarly, we define a distribution

$$\mathbb{I}_\alpha(\Phi) = \int_{[\mathrm{U}(V)]} \mathcal{K}_{\Phi, \alpha}(g) dg.$$

This still keeps the action of $\mathrm{SL}_2(\mathbb{A}_0)$ under the Weil representation ω ,

$$(1.1) \quad \mathbb{I}_\alpha(h, \Phi) = \mathbb{I}_\alpha(\omega(h)\Phi), \quad h \in \mathrm{SL}_2(\mathbb{A}_0).$$

We have the similar construction for $S_n \times V'_n$. Clearly by varying α we have refined the relations between the orbital integrals appearing in \mathbb{I} and \mathbb{J} . In the local situation, this sliced version was utilized in [47] to prove the existence of smooth transfer by an induction argument. Here we are exploiting the global analog, i.e., the $\mathrm{SL}_2(F_0)$ -modularity of (1.1) and its counterpart for \mathbb{J} .

Another input is to impose that $\mathrm{U}(V)$ is compact at archimedean places, and at the same time to plug in a (weaker version of) Gaussian test function; cf.

[Section 12](#). This simplifies the spectra of the SL_2 -automorphic forms $\mathbb{I}_\alpha(\cdot, \Phi)$ and its counterpart on $S_n \times V'_n$, to the extent that the spectra are finite. In fact, in our case, they lie in a finite dimensional vector space corresponding to classical holomorphic modular forms with known levels and weights.

The two inputs allow us to deduce the Jacquet–Rallis fundamental lemma by induction on the dimension of V , at least when $p \geq \dim V$.

Now that we have explained our approach to the FL, let us move to the AFL conjecture. We have indicated that the extra symmetry is the SL_2 -modularity of the kernel function, which follows from the Poisson summation formula. In the arithmetic setting, the extra symmetry is a version of the modularity of generating series of special divisors in the arithmetic Chow groups of the integral models of unitary Shimura varieties (e.g., in the recent work of Bruinier–Howard–Kudla–Rapoport–Yang [5]).

To take advantage of the modularity, we consider the semi-Lie algebraic version of the AFL conjecture, which has appeared in Mihatsch’s thesis [31, §8] and in Liu’s work [30, Conj. 1.11]. In the semi-Lie algebraic version, we consider the intersection numbers of the Kudla–Rapoport divisors (KR divisors, for short) [24] and the (derived) fixed point locus of an automorphism of the RZ space. We show in [Section 3](#) that there is an inductive structure similar to the smooth transfer and the fundamental lemma. More precisely, it is possible to reduce the special case when the KR divisor is (formally) smooth to the AFL in one-dimension lower. This is still hardly useful if we only work on the local moduli space. Therefore we introduce a global version of the fixed point locus, called “the derived CM cycle,” or “the fat big CM cycle,” being a “thickened” variant of the “big CM cycle” in the work of Bruinier–Kudla–Yang [6] and Howard [18]. The naively defined CM cycle may have dimension larger than expected. However, we note that it is a union of connected components of the fixed point locus of a Hecke correspondence (over the integral model); cf. [Section 7.5](#). Therefore there is a natural derived structure on the naive CM cycle, and the derived CM cycle has virtual dimension one, as expected.

By the modularity of generating series of special divisors mentioned above, we obtain a modular form (with known level and weight) by taking the (arithmetic) intersection numbers (cf. [\(9.4\)](#)) of a fixed (derived) CM cycle with special divisors; cf. [Section 9.2](#). The rest is then similar to the proof of the FL conjecture. The resulting modular form is the arithmetic analog of [\(1.1\)](#). By induction, together with a special case of the AFL (cf. [Proposition 3.9](#)), one may assume that the ξ -th Fourier coefficients are known if ξ is prime to a certain finite set of places. The desired equality for *all* Fourier coefficients then follows from the modularity of the generating series and a density principle for the Fourier coefficients of holomorphic modular forms (cf. [Lemma 13.6](#)). Finally, one deduces the AFL conjecture from the global identity, together with a local constancy property of the intersection numbers on RZ spaces; cf. [Theorem 5.5](#).

In our approach, it is important to understand the archimedean local intersection (i.e., the values of Green functions; cf. Section 10), and correspondingly the derivatives of the archimedean orbital integrals for Gaussian test functions (cf. Section 12). After subtracting the archimedean terms, the intersection numbers and derivative of orbital integrals at non-archimedean places all lie in the \mathbb{Q} -linear span of $\log p$ for a finite set of primes p . One can then separate the contribution from different primes by the linear independence of logarithms of prime numbers.

We have restricted this paper to the case $F_0 = \mathbb{Q}$ since in a few places there are missing ingredients in the literature and some of them are subtle. However, we have tried to present most of the arguments in the general totally real field case, especially in the analytic side of RTF.

We would like to point out some earlier works related to the AFL conjecture. The author proved the AFL for low ranks of the unitary group ($n = 2$ and 3) in [46]. Rapoport, Terstiege and the author [41] proved it for arbitrary rank $n \leq p$ and *minuscule* group elements. A Lie algebraic version (in the case of artinian intersection) was studied by Mihatsch in [33], [31], simplifying the proof and generalizing the result in [46]. Finally, in the minuscule case, Li and Zhu in [27] have given a simplified proof of [41]; recently, He, Li, and Zhu [17] have also removed the restriction on the residue characteristic.

During the preparation of this paper, the author learned that Beuzart-Plessis [2] has given a purely local proof of the Jacquet–Rallis fundamental lemma for all p -adic fields with p odd, by induction and using a more precise version (i.e., a local relative trace formula) of the compatibility between the *local* Weil representation (mainly the Fourier transform) and smooth transfer. It is an interesting question whether there is a purely local proof of the AFL along the line of his proof.

1.1. Acknowledgements. We thank Chao Li, Andreas Mihatsch, Michael Rapoport, Chen Wan and the referee for their comments. An earlier version of the paper was circulated in the ARGOS seminar in the spring 2019 and in a seminar in Morningside center Beijing in the summer 2019. The author would like to thank Michael Rapoport and Ye Tian for communicating comments from their seminars, which have helped the author improve the paper.

1.2. Notation.

Notation on algebra

- \mathbb{R}_+ : the set of positive real numbers.
- Let F be a field of character zero. For a reductive group H acting on an affine variety X , we say that a point $x \in X(F)$ is
 - H -semisimple if Hx is Zariski closed in X (when F is a local field, equivalently, $H(F)x$ is closed in $X(F)$ for the analytic topology);

– H -regular if the stabilizer H_x of x is trivial.

And we say that x is *regular semisimple* if it is regular and semisimple. We denote by $X(F)_{\text{rs}}$ the set of regular semisimple elements and by $[X(F)]_{\text{rs}}$ the set of regular semisimple $H(F)$ -orbits. We denote the categorical quotient by $X_{//H}$ with the natural map $X \rightarrow X_{//H}$.

- For global fields, unless otherwise stated, F denotes a CM number field and F_0 denotes its (maximal) totally real subfield of index 2. We denote by $a \mapsto \bar{a}$ the nontrivial automorphism of F/F_0 . Let $F_{0,+}$ (resp. $F_{0,\geq 0}$) be the set of totally positive (resp. semi-positive) elements in F_0 .
- We denote $\mathbf{H} = \text{SL}_2$ as an algebraic group over F_0 . Denote by B the Borel subgroup of upper triangular matrices and by N its unipotent radical.
- We use the symbols v and v_0 to denote places of F_0 , and we use w and w_0 to denote places of F . We write $F_{0,v}$ for the v -adic completion of F_0 , and we set $F_v := F \otimes_{F_0} F_{0,v}$; thus F_v is isomorphic to $F_{0,v} \times F_{0,v}$ or to a quadratic field extension of $F_{0,v}$ accordingly as v is split or non-split in F . We write $\mathcal{O}_{F_{0,v}} \subset F_{0,v}$ for the ring of integers. We use analogous notation for other fields in place of F_0 and other finite places in place of v .
- Unless otherwise stated, we write \mathbb{A} , \mathbb{A}_0 , and \mathbb{A}_F for the adele rings of \mathbb{Q} , F_0 , and F , respectively. We systematically use a subscript f for the ring of finite adeles and a superscript p for the adeles away from the prime number p .
- For an abelian scheme A over a locally noetherian scheme S on which the prime number p is invertible, we write $\text{T}_p(A)$ for the p -adic Tate module of A (regarded as a smooth \mathbb{Z}_p -sheaf on S) and $\text{V}_p(A) := \text{T}_p(A) \otimes \mathbb{Q}$ for the rational p -adic Tate module (regarded as a smooth \mathbb{Q}_p -sheaf on S). When S is a $\mathbb{Z}_{(p)}$ -scheme, we similarly write $\widehat{\text{V}}^p(A)$ for the rational prime-to- p Tate module of A . When S is a scheme in characteristic zero, we write $\widehat{\text{V}}(A)$ for the full rational Tate module of A .
- We use a superscript \circ to denote the operation $- \otimes_{\mathbb{Z}} \mathbb{Q}$ on groups of homomorphisms of abelian schemes so that, for example,

$$\text{Hom}^\circ(A, A') := \text{Hom}(A, A') \otimes_{\mathbb{Z}} \mathbb{Q}.$$

- All Chow groups and K -groups have \mathbb{Q} -coefficients.
- Given a discretely valued field L , we denote the completion of a maximal unramified extension of it by \check{L} .
- We write 1_n for the $n \times n$ identity matrix. Let $\text{M}_{n,m}(R)$ denote the R -module of $n \times m$ -matrices with coefficients in a ring R .
- For a vector space V over a field F_0 (of characteristic not equal to 2), a quadratic form $\mathfrak{q} : V \rightarrow F$ has an associated symmetric bilinear pairing

defined by

$$(1.2) \quad \langle x, y \rangle = \mathfrak{q}(x + y) - \mathfrak{q}(x) - \mathfrak{q}(y), \quad x, y \in V.$$

In particular,

$$(1.3) \quad \langle x, x \rangle = 2\mathfrak{q}(x).$$

For a quadratic field extension F of F_0 , an F/F_0 -hermitian space is an F_0 -vector space V endowed with an F_0 -linear action of F and an “ F/F_0 -hermitian form,” i.e., a map $\langle \cdot, \cdot \rangle : V \times V \rightarrow F$ that is F -linear on the first factor, and conjugate-linear on the second factor. Its dimension will be the dimension as an F -vector space. It induces a symmetric bi- F_0 -linear pairing by $(x, y) \mapsto \text{tr}_{F/F_0}\langle x, y \rangle \in F_0$. In particular, the corresponding quadratic form on V is

$$(1.4) \quad \mathfrak{q}(x) = \langle x, x \rangle \in F_0.$$

We will treat V as an affine variety over F_0 , and for $\xi \in F_0$, we denote by V_ξ the subscheme defined by $\mathfrak{q}(x) = \xi$.

- For a F/F_0 -hermitian space V over a non-archimedean local field, and an O_F -lattice $\Lambda \subset V$ (of full rank), we denote by Λ^\vee its dual lattice under the hermitian form.
- Let R be a commutative ring. We denote by $(\text{LNSch})_{/R}$ the category of locally noetherian schemes over $\text{Spec } R$. We denote by $R[T]_{\deg=m}$ the set of monic polynomials with coefficients in R of degree m .

Notation on automorphic forms.

- Fix the non-trivial additive character $\psi = \psi_{\mathbb{Q}} \circ \text{tr}_{F_0/\mathbb{Q}} : F_0 \setminus \mathbb{A}_0 \rightarrow \mathbb{C}^\times$, where $\psi_{\mathbb{Q}}$ is the standard one and $\text{tr}_{F_0/\mathbb{Q}} : F_0 \setminus \mathbb{A}_0 \rightarrow \mathbb{Q} \setminus \mathbb{A}$ is the trace map. For $\xi \in F_0$, we denote by $\psi_\xi(x) = \psi(\xi x)$.
- For a smooth algebraic variety X over a local field F , we denote $\mathcal{S}(X(F))$ by the space of Schwartz functions on $X(F)$. When F is non-archimedean, this is the same as the space of locally constant functions with compact support. When F is archimedean, $\mathcal{S}(X(F))$ consists of smooth functions ϕ on $X(F)$ such that, for every algebraic differential operator D on X , the function $D\phi$ is bounded. Similarly, for a smooth algebraic variety X over a global field F , we denote $\mathcal{S}(X(\mathbb{A}))$ by the space of Schwartz functions on $X(\mathbb{A})$.
- $\mathcal{H} = \{\tau = b + ia \in \mathbb{C} \mid a > 0\}$: the complex upper half plane.
- For $\xi \in \mathbb{R}$ and $k \in \mathbb{Z}$, the weight k Whittaker function on $\text{SL}_2(\mathbb{R})$ is defined by

$$(1.5) \quad W_\xi^{(k)}(h) = |a|^{k/2} e^{2\pi i \xi(b+ai)} \chi_k(\kappa_\theta),$$

where we write $h \in \mathrm{SL}_2(\mathbb{R})$ according to the Iwasawa decomposition

$$(1.6) \quad h = \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix} \begin{pmatrix} a^{1/2} & \\ & a^{-1/2} \end{pmatrix} \kappa_\theta, \quad a \in \mathbb{R}_+, \quad b \in \mathbb{R}$$

and

$$(1.7) \quad \kappa(\theta) = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \in \mathrm{SO}(2, \mathbb{R}).$$

Here the weight k -character of $\mathrm{SO}(2, \mathbb{R})$, for $k \in \mathbb{Z}$, is defined by

$$(1.8) \quad \chi_k(\kappa_\theta) = e^{ik\theta}.$$

- The principal congruence subgroups of $\mathrm{SL}_2(\mathbb{Z})$ are

$$\Gamma(N) = \left\{ \gamma \in \mathrm{SL}_2(\mathbb{Z}) \mid \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{N} \right\}.$$

- $\mathcal{A}_{\mathrm{hol}}(\Gamma, k)$ is the space of holomorphic modular forms of level Γ , weight k , for Γ where $\Gamma(N) \subset \Gamma \subset \mathrm{SL}_2(\mathbb{Z})$. For any subfield $L \subset \mathbb{C}$, we denote by $\mathcal{A}_{\mathrm{hol}}(\Gamma, k)_L$ the L -vector space consisting of $f \in \mathcal{A}_{\mathrm{hol}}(\Gamma, k)$ whose Fourier coefficients in the q -expansion at the cusp $i\infty$ all lie in L . Fixing an embedding $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$, the \mathbb{C} -vector space $\mathcal{A}_{\mathrm{hol}}(\Gamma, k)$ has a $\overline{\mathbb{Q}}$ -structure via the q -expansion at the cusp $i\infty$, i.e., $\mathcal{A}_{\mathrm{hol}}(\Gamma, k) = \mathcal{A}_{\mathrm{hol}}(\Gamma, k)_{\overline{\mathbb{Q}}} \otimes_{\overline{\mathbb{Q}}} \mathbb{C}$. For any L -vector space W , we have an L -vector space

$$(1.9) \quad \mathcal{A}_{\mathrm{hol}}(\Gamma, k)_L \otimes_L W.$$

We will view this vector space as the space of formal power series in $q^{1/N}$ with coefficients in W

$$\sum_{\xi \geq 0, \xi \in \frac{1}{N}\mathbb{Z}} A_\xi q^\xi, \quad A_\xi \in W,$$

where there exist elements $f_i \in \mathcal{A}_{\mathrm{hol}}(\Gamma, k)_L$ indexed by a finite set I whose q -expansion at the cusp $i\infty$ are given by $\sum_{\xi \geq 0, \xi \in \frac{1}{N}\mathbb{Z}} a_\xi(f_i) q^\xi \in L[[q^{1/N}]]$, and elements $w_i \in W, i \in I$, such that

$$A_\xi = \sum_{i \in I} a_\xi(f_i) w_i \quad \text{for all } \xi.$$

- $\mathcal{A}_{\mathrm{hol}}(\mathbf{H}(\mathbb{A}_0), K, k)$ is the space of automorphic forms (with moderate growth) on $\mathbf{H}(\mathbb{A}_0)$, invariant under $K \subset \mathbf{H}(\mathbb{A}_{0,f})$, and parallel weight k under the action of $\prod_{v \in \mathrm{Hom}(F_0, \mathbb{R})} \mathrm{SO}(2, \mathbb{R})$, holomorphic (i.e., annihilated by the element $\frac{1}{2} \begin{pmatrix} i & 1 \\ 1 & -i \end{pmatrix}$ in the complexified Lie algebra of $\mathbf{H}(F_{0,v}) \simeq \mathrm{SL}_2(\mathbb{R})$ for every $v \in \mathrm{Hom}(F_0, \mathbb{R})$). This is a finite dimensional vector space over \mathbb{C} , and it has a $\overline{\mathbb{Q}}$ -structure via the q -expansion at the cusp $i\infty$. For any subfield

$L \subset \mathbb{C}$ and any L -vector space W , we define $\mathcal{A}_{\text{hol}}(\mathbf{H}(\mathbb{A}_0), K, k)_L$ similar to $\mathcal{A}_{\text{hol}}(\Gamma, k)_L$, and

$$(1.10) \quad \mathcal{A}_{\text{hol}}(\mathbf{H}(\mathbb{A}_0), K, k)_L \otimes_L W$$

similar to $\mathcal{A}_{\text{hol}}(\Gamma, k)_L \otimes_L W$ as above.

- To a (parallel) weight k function $\phi : \mathbf{H}(\mathbb{A}_0) \rightarrow \mathbb{C}$ and $h_f \in \mathbf{H}(\mathbb{A}_{0,f})$, we define $\phi_{h_f}^\flat$ to be the function:

$$(1.11) \quad \begin{aligned} \phi_{h_f}^\flat : \quad \prod_{v|\infty} \mathcal{H} &\longrightarrow \mathbb{C} \\ \tau = (\tau_v)_{v|\infty} &\longmapsto |a_\infty|^{-k/2} \phi(h_\infty, h_f), \end{aligned}$$

where $h_\infty = (h_v)_{v|\infty}$, $h_v = \begin{pmatrix} 1 & b_v \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a_v^{1/2} \\ a_v^{-1/2} \end{pmatrix}$, $\tau_v = b_v + a_v i \in \mathcal{H}$ and $|a_\infty| = \prod_{v|\infty} |a_v|$. When $h_f = 1$, we simply write it as ϕ^\flat . If $\phi \in \mathcal{A}_{\text{hol}}(\mathbf{H}(\mathbb{A}_0), K, k)$, then $\phi_{h_f}^\flat \in \mathcal{A}_{\text{hol}}(\Gamma, k)$, where $\Gamma = h_f K h_f^{-1} \cap \mathbf{H}(F_0)$.

- For a left $N(F_0)$ -invariant continuous function $\phi : \mathbf{H}(\mathbb{A}_0) \rightarrow \mathbb{C}$, its ξ -th Fourier coefficient for $\xi \in F_0$ is defined as the function

$$(1.12) \quad h \in \mathbf{H}(\mathbb{A}_0) \longmapsto W_\phi(h) := \int_{F_0 \backslash \mathbf{H}_0} \phi \left[\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} h \right] \psi_{-\xi}(b) db.$$

Then there is a Fourier expansion (by an absolute convergent sum): for $h \in \mathbf{H}(\mathbb{A}_0)$,

$$(1.13) \quad \phi(h) = \sum_{\xi \in F_0} W_{\phi, \xi}(h).$$

- The case $F_0 = \mathbb{Q}$: the \mathbb{C} -vector space $\mathcal{A}_{\text{exp}}(\mathbf{H}(\mathbb{A}), K, k)$ consists of smooth functions ϕ on $\mathbf{H}(\mathbb{Q}) \backslash \mathbf{H}(\mathbb{A})$ with at worst exponential growth (i.e., for every $h_f \in \mathbf{H}(\mathbb{A}_f)$, there exists a constant C such that $|\phi(h_\infty h_f)| \leq e^{Ca}$ when $a \rightarrow \infty$, where $h_\infty \in \text{SL}_2(\mathbb{R})$ denotes the matrix (1.6)), invariant under $K \subset \mathbf{H}(\mathbb{A}_f)$ and weight k under the action of $\text{SO}(2, \mathbb{R})$, such that $\frac{1}{2} \begin{pmatrix} i & 1 \\ 1 & -i \end{pmatrix} \phi \in \mathcal{A}_{\text{hol}}(\mathbf{H}(\mathbb{A}), K, k-2)$. This is related to the space $\mathcal{A}_k^!(\rho_L^\vee)$ in [8, Def. 2.8, pp. 2104], noting that the differential operator $\frac{1}{2} \begin{pmatrix} i & 1 \\ 1 & -i \end{pmatrix}$ is the Maass lowering operator. This is an infinite dimensional vector space over \mathbb{C} .

Part 1. Local theory

Throughout this part, F_0 is a field of characteristic zero and F is a quadratic étale F_0 -algebra.

2. FL and variants

2.1. Group-theoretic setup.

Let

$$e := (0, \dots, 0, 1) \in \mathrm{M}_{n,1}(F) = F^n$$

be a column vector, and let $e^* \in \mathrm{M}_{1,n}(F) \simeq \mathrm{M}_{n,1}(F)^* = (F^n)^*$ be the transpose of e . Consider the embedding of algebraic groups over F ,

$$(2.1) \quad \begin{aligned} \mathrm{GL}_{n-1} &\longrightarrow \mathrm{GL}_n \\ \gamma_0 &\longmapsto \mathrm{diag}(\gamma_0, 1); \end{aligned}$$

this identifies GL_{n-1} with the subgroup of points γ in GL_n such that $\gamma e = e$ and $e^* \gamma = e^*$.

We introduce the algebraic group G' over F_0 and its subgroups,

$$\begin{aligned} G' &:= \mathrm{Res}_{F/F_0}(\mathrm{GL}_{n-1} \times \mathrm{GL}_n), \\ H'_1 &:= \mathrm{Res}_{F/F_0} \mathrm{GL}_{n-1}, \\ H'_2 &:= \mathrm{GL}_{n-1} \times \mathrm{GL}_n. \end{aligned}$$

Here H'_1 is embedded diagonally, and H'_2 is embedded in the obvious way. We consider the natural right action of $H'_1 \times H'_2$ on G' ,

$$(h_1, h_2) \cdot \gamma = h_1^{-1} \gamma h_2.$$

Consider the symmetric space

$$(2.2) \quad S := S_n := \{ g \in \mathrm{Res}_{F/F_0} \mathrm{GL}_n \mid g\bar{g} = 1_n \}$$

and its tangent space at 1_n , called “the Lie algebra” of S_n ,

$$(2.3) \quad \mathfrak{s} := \mathfrak{s}_n := \{ y \in \mathrm{Res}_{F/F_0} \mathrm{M}_n \mid y + \bar{y} = 0 \}.$$

Set

$$H' := \mathrm{GL}_{n-1}.$$

Then H' acts on S_n and \mathfrak{s}_n by conjugation

$$h \cdot \gamma = h^{-1} \gamma h.$$

We also consider a variant (arising from the Fourier–Jacobi period [9], [29]). Let

$$(2.4) \quad V'_{n-1} = F_0^{n-1} \times (F_0^{n-1})^*,$$

and consider the (diagonal) action of H' on the product $S_{n-1} \times V'_{n-1}$,

$$h \cdot (\gamma, (u_1, u_2)) = (h^{-1} \gamma h, (h^{-1} u_1, u_2 h)).$$

The action of H' on the its Lie algebra $\mathfrak{s}_{n-1} \times V'_{n-1}$ is defined similarly.

Next let V^\sharp be an F/F_0 -hermitian space of dimension $n \geq 2$. We fix a non-isotropic vector $u_0 \in V^\sharp$, which we call the *special vector*. We denote by

V the orthogonal complement of u_0 in V^\sharp . We define the algebraic group G over F_0 and its subgroups,

$$(2.5) \quad \begin{aligned} G &:= \mathrm{U}(V^\sharp), \\ H &:= \mathrm{U}(V), \\ G_V &:= H \times G. \end{aligned}$$

We have the natural action of $H \times H$ on G_V and the conjugation action of H on G . We also consider the adjoint action of H on the Lie algebra $\mathfrak{g} = \mathfrak{u}(V^\sharp)$ of G . When $\dim V = 1$, the Lie algebra $\mathfrak{u}(V)$ is denoted by $\mathfrak{u}(1)$, which is canonically isomorphic to F^- , the (-1) -eigenspace of F under the Galois conjugation.

We also need the variant arising from the RTF for the Fourier–Jacobi period [29]): the (diagonal) action of $H = \mathrm{U}(V)$ on the product $\mathrm{U}(V) \times V$ and $\mathfrak{u}(V) \times V$, where the two factors are endowed with the adjoint action (on the group and the Lie algebra) and the standard action respectively.

2.2. Orbit matching. There is a natural bijection of orbit spaces of *regular semisimple* elements,

$$(2.6) \quad \coprod_V [(\mathrm{U}(V^\sharp)(F_0)]_{\mathrm{rs}} \xrightarrow{\sim} [S_n(F_0)]_{\mathrm{rs}}$$

and

$$(2.7) \quad \coprod_V [(\mathrm{U}(V) \times V)(F_0)]_{\mathrm{rs}} \xrightarrow{\sim} [(S_{n-1} \times V'_{n-1})(F_0)]_{\mathrm{rs}}$$

(cf. [46] and [29]), where the disjoint union runs over the set of isometry classes of F/F_0 -hermitian spaces V , and the larger space $V^\sharp = V \oplus F \cdot u_0$ is then determined uniquely by demanding the special vector u_0 to have norm one (or any fixed number in F_0^\times when varying V). Here the left-hand (resp. right-hand) sides denote the orbits under the action of the group $\mathrm{U}(V)$ (resp. GL_{n-1}). The bijections define a *matching relation* between regular semisimple orbits. In both cases, there are also similar injections for orbits on the Lie algebras:

$$(2.8) \quad \coprod_V [(\mathfrak{u}(V^\sharp)(F_0)]_{\mathrm{rs}} \xrightarrow{\sim} [\mathfrak{s}_n(F_0)]_{\mathrm{rs}}$$

and

$$(2.9) \quad \coprod_V [(\mathfrak{u}(V) \times V)(F_0)]_{\mathrm{rs}} \xrightarrow{\sim} [(\mathfrak{s}_{n-1} \times V'_{n-1})(F_0)]_{\mathrm{rs}}.$$

We recall how the map (2.7) is defined. Choose an F -basis for V and complete it to a basis for V^\sharp by adjoining u_0 . This identifies V with F^{n-1} and V^\sharp with F^n in such a way that u_0 corresponds to the column vector $e := (0, \dots, 0, 1)$ in F^n , and hence determines embeddings of groups $\mathrm{U}(V^\sharp) \hookrightarrow \mathrm{Res}_{F/F_0} \mathrm{GL}_n$. An element $g \in \mathrm{U}(V)(F_0)_{\mathrm{rs}}$ and an element $\gamma \in S_n(F_0)_{\mathrm{rs}}$ are said to *match* if these two elements, when considered as elements in

$\text{Res}_{F/F_0} \text{GL}_n(F_0)$, are conjugate under $\text{Res}_{F/F_0} \text{GL}_{n-1}$. The matching relation is independent of the choice of embeddings and induces a bijection [46, §2]. In a similar way, we view elements in $(S_{n-1} \times V'_{n-1})(F_0)$ as elements in $\text{Res}_{F/F_0} \text{M}_{n,n}(F_0)$ by

$$(\gamma, (u_1, u_2)) \longmapsto \begin{pmatrix} \gamma & u_1 \\ u_2 & 0 \end{pmatrix},$$

and we view elements $(g, u) \in (\text{U}(V) \times V)(F_0)$ as elements in $\text{Res}_{F/F_0} \text{M}_{n,n}(F_0)$

$$(g, u) \longmapsto \begin{pmatrix} g & u \\ u^* & 0 \end{pmatrix}.$$

Here we view $u \in V(F_0)$ as the corresponding element in $\text{Hom}(V^\perp, V)$ sending $u_0 \in V^\perp = F \cdot u_0$ to u , and u^* is the element in $\text{Hom}(V, V^\perp) = \text{Hom}(V, F \cdot u_0)$ defined by $u' \mapsto \langle u', u \rangle u_0$. Then, an element $(g, u) \in (\text{U}(V) \times V)(F_0)_{\text{rs}}$ and an element $(\gamma, (u_1, u_2)) \in (S_{n-1}(F_0) \times F_0^{n-1} \times (F_0^{n-1})^*)_{\text{rs}}$ are said to *match* if these two elements, when considered as elements in $\text{Res}_{F/F_0} \text{M}_{n,n}(F_0)$, are conjugate under $\text{Res}_{F/F_0} \text{GL}_{n-1}$.

Equivalently, $(g, u) \in (\text{U}(V) \times V)(F_0)_{\text{rs}}$ matches

$$(\gamma, (u_1, u_2)) \in (S_{n-1}(F_0) \times F_0^{n-1} \times (F_0^{n-1})^*)_{\text{rs}}$$

if and only if the following invariants are equal:

$$\det(T \mathbf{1}_{n-1} + g) = \det(T \mathbf{1}_{n-1} + \gamma) \quad \text{and} \quad \langle g^i u, u \rangle = u_2 \gamma^i u_1, \quad 0 \leq i \leq n-2.$$

Here $\det(T \mathbf{1}_{n-1} + g) \in F[T]_{\deg=n-1}$ is the characteristic polynomial of g . (We remind the reader that $F[T]_{\deg=n-1}$ denotes the set of monic polynomials with coefficients in F of degree $n-1$; cf. Section 1.2.) In fact, these invariants define natural identifications of the categorical quotients $(\text{U}(V) \times V)_{/\text{U}(V)}$ and $(S_{n-1} \times V'_{n-1})_{/\text{GL}_{n-1}}$ with an F_0 -subscheme of the affine space $\text{Res}_{F/F_0}(F[T]_{\deg=n-1} \times F^{n-1})$, and we denote this F_0 -subscheme by \mathcal{B}_{n-1} :

$$(2.10) \quad \mathcal{B}_{n-1} \hookrightarrow \text{Res}_{F/F_0}(F[T]_{\deg=n-1} \times F^{n-1}).$$

We refer to [47] for the analogous case $\text{U}(V^\sharp)_{/\text{U}(V)} \simeq S_{n-1} \text{GL}_{n-1}$. Similarly, the characteristic polynomial defines a natural identification of $\text{U}(V)_{/\text{U}(V)}$ and $S_{n-1} \text{GL}_{n-1}$ with an F_0 -subscheme of the affine space $\text{Res}_{F/F_0}(F[T]_{\deg=n-1})$, which will be denoted by \mathcal{A}_{n-1} :

$$(2.11) \quad \mathcal{A}_{n-1} \hookrightarrow \text{Res}_{F/F_0}(F[T]_{\deg=n-1}).$$

More precisely, \mathcal{A}_{n-1} is the F_0 -scheme of conjugate self-reciprocal monic polynomials $\alpha \in F[T]_{\deg=n-1}$, i.e.,

$$T^{\deg(\alpha)} \alpha(T^{-1}) = \alpha(0) \bar{\alpha}(T),$$

where $\bar{\alpha}$ is the coefficient-wise Galois conjugate of α (in particular, $\alpha(0)\bar{\alpha}(0) = 1$). Moreover, both \mathcal{A} and \mathcal{B} have natural integral models over \mathcal{O}_{F_0} , so we will freely talk about their points over any \mathcal{O}_{F_0} -algebra.

For $\alpha \in \mathcal{A}_{n-1}(F_0)$, we will denote by $S_{n-1}(\alpha)$ its preimage under the natural morphism $S_{n-1} \rightarrow \mathcal{A}_{n-1}$. For $\xi \in F_0$, we will denote by $V'_{n-1,\xi}$ the subscheme of V'_{n-1} defined by $u_2u_1 = \xi$. We denote by $[S_{n-1}(\alpha)(F_0)]$ (resp. $[(S_{n-1}(\alpha) \times V'_{n-1})(F_0)]$ and $[(S_{n-1}(\alpha) \times V'_{n-1,\xi})(F_0)]$) the set of $\mathrm{GL}_{n-1}(F_0)$ -orbits in $S_{n-1}(\alpha)(F_0)$ (resp. $(S_{n-1}(\alpha) \times V'_{n-1})(F_0)$ and $(S_{n-1}(\alpha) \times V'_{n-1,\xi})(F_0)$). Similar notation applies to unitary groups.

2.3. Orbital integral matching: smooth transfer. We recall orbital integrals [38, §2.2]. Now let F/F_0 be a quadratic extension of local fields of characteristic zero. (The split $F = F_0 \times F_0$ is similar and simpler.) Let

$$\eta = \eta_{F/F_0} : F_0^\times \longrightarrow \{\pm 1\}$$

be the quadratic character associated to F/F_0 by local class field theory.

To simplify the exposition we consider the non-archimedean case, though the archimedean case requires very little change. Then there are exactly two isometry classes of F/F_0 -hermitian spaces of dimension $n-1$, denoted by V_0 and V_1 . When F/F_0 is unramified, we will assume that V_0 has a self-dual lattice. Then the orbit bijections are now

$$[(\mathrm{U}(V_0^\sharp)(F_0)]_{\mathrm{rs}} \coprod [(\mathrm{U}(V_1^\sharp)(F_0)]_{\mathrm{rs}} \xrightarrow{\sim} [S_n(F_0)]_{\mathrm{rs}}$$

and

$$[(\mathrm{U}(V_0) \times V_0)(F_0)]_{\mathrm{rs}} \coprod [(\mathrm{U}(V_1) \times V_1)(F_0)]_{\mathrm{rs}} \xrightarrow{\sim} [(S_{n-1} \times V'_{n-1})(F_0)]_{\mathrm{rs}}.$$

For $\gamma \in S_n(F_0)_{\mathrm{rs}}$, $f' \in \mathcal{S}(S_n(F_0))$, and $s \in \mathbb{C}$, we define

$$(2.12) \quad \mathrm{Orb}(\gamma, f', s) := \int_{\mathrm{GL}_{n-1}(F_0)} f'(h^{-1}\gamma h) |\det h|^s \eta(h) dh,$$

where $|\cdot|$ denotes the normalized absolute value on F_0 , where we set

$$\eta(h) := \eta(\det h).$$

We define the special values

(2.13)

$$\mathrm{Orb}(\gamma, f') := \omega(\gamma) \mathrm{Orb}(\gamma, f', 0) \text{ and } \partial \mathrm{Orb}(\gamma, f') := \omega(\gamma) \left. \frac{d}{ds} \right|_{s=0} \mathrm{Orb}(\gamma, f', s),$$

where the transfer factor $\omega(\gamma)$ is to be explicated below by (2.16). Here, we have included the transfer factor in the special values of the orbital integrals, different from [38, §2.2].

For $(\gamma, u') \in (S_{n-1} \times V'_{n-1})(F_0)$, $\Phi' \in \mathcal{S}((S_{n-1} \times V'_{n-1})(F_0))$, and $s \in \mathbb{C}$, we define

$$(2.14) \quad \text{Orb}((\gamma, u'), \Phi', s) := \int_{\text{GL}_{n-1}(F_0)} \Phi'(h \cdot (\gamma, u')) |\det h|^s \eta(h) dh,$$

and we define their special values similar to (2.13), replacing the transfer factor $\omega(\gamma)$ by $\omega(\gamma, u')$ to be explicated below by (2.17).

On the unitary side, for $g \in \text{U}(V^\sharp)(F_0)_{\text{rs}}$ and $f \in \mathcal{S}(\text{U}(V^\sharp)(F_0))$, we define

$$\text{Orb}(g, f) := \int_{\text{U}(V)(F_0)} f(h^{-1}gh) dh.$$

For $(g, u) \in (\text{U}(V) \times V)(F_0)_{\text{rs}}$ and $\Phi \in \mathcal{S}((\text{U}(V) \times V)(F_0))$, we define

$$(2.15) \quad \text{Orb}((g, u), \Phi) := \int_{\text{U}(V)(F_0)} \Phi(h \cdot (g, u)) dh.$$

Finally, we define an explicit transfer factors; cf. [38, §2.4]. First fix an extension $\tilde{\eta}$ of the quadratic character η from F_0^\times to F^\times (not necessarily of order 2). If F is unramified, then we take the natural extension $\tilde{\eta}(x) = (-1)^{v(x)}$. For S_n , we take the transfer factor

$$(2.16) \quad \omega(\gamma) := \tilde{\eta}(\det(\gamma)^{-\lfloor n/2 \rfloor} \det(\gamma^i e)_{0 \leq i \leq n-1}), \quad \gamma \in S_n(F_0)_{\text{rs}}.$$

For $(\gamma, u') \in (S_{n-1} \times V'_{n-1})(F_0)_{\text{rs}}$ where $u' = (u_1, u_2) \in V_{n-1}(F_0) = F_0^{n-1} \times (F_0^{n-1})^*$, we take

$$(2.17) \quad \omega(\gamma, u') := \tilde{\eta}(\det(\gamma)^{-\lfloor (n-1)/2 \rfloor} \det(\gamma^i u_1)_{0 \leq i \leq n-2}).$$

Similarly, we define transfer factors on \mathfrak{s}_n and $\mathfrak{s}_{n-1} \times V'_{n-1}$.

Definition 2.1. A function $f' \in \mathcal{S}(S_n(F_0))$ and a pair of functions $(f_0, f_1) \in \mathcal{S}(\text{U}(V_0^\sharp)(F_0)) \times \mathcal{S}(\text{U}(V_1^\sharp)(F_0))$ are (smooth) *transfers* of each other if for each $i \in \{0, 1\}$ and each $g \in \text{U}(V_i^\sharp)(F_0)_{\text{rs}}$,

$$\text{Orb}(g, f_i) = \text{Orb}(\gamma, f')$$

whenever $\gamma \in S(F_0)_{\text{rs}}$ matches g .

Definition 2.2. A function $\Phi' \in \mathcal{S}((S_{n-1} \times V_{n-1})(F_0))$ and a pair of functions $(\Phi_0, \Phi_1) \in \mathcal{S}((\text{U}(V_0) \times V_0)(F_0)) \times \mathcal{S}((\text{U}(V_1) \times V_1)(F_0))$ are (smooth) *transfers* of each other if for each $i \in \{0, 1\}$ and each $(g, u) \in (\text{U}(V_i) \times V_i)(F_0)_{\text{rs}}$,

$$(2.18) \quad \text{Orb}((g, u), \Phi_i) = \text{Orb}((\gamma, u'), \Phi')$$

whenever $(\gamma, u') \in (S_{n-1} \times V'_{n-1})(F_0)_{\text{rs}}$ matches (g, u) .

The definitions made above easily extend verbatim to the setting of the full Lie algebras $\mathfrak{u}(V) \times V$ and $\mathfrak{s}_{n-1} \times V'_{n-1}$. Finally, we remark that the definitions extend to the archimedean local field extension $F/F_0 = \mathbb{C}/\mathbb{R}$, where one only needs to replace the pair of functions (Φ_0, Φ_1) by a tuple of functions $\{\Phi_V\}_V$

indexed by the set of isometry classes of F/F_0 -hermitian spaces V , as in (2.7) and (2.9). We will not repeat the detail here.

2.4. Review of the FL conjecture. We review the FL conjecture; cf. [21], [46], [38]. Let F/F_0 be an unramified quadratic extension of p -adic field for an *odd* prime p . Assume furthermore that the special vectors $u_i \in V_i$ have norm one (or any fixed unit in O_{F_0}). Then the hermitian space V_i^\sharp is again split for $i = 0$ and non-split for $i = 1$. We write $G_i = \mathrm{U}(V_i^\sharp)$, $\mathfrak{g}_i = \mathrm{Lie} G_i$, and $H_i = \mathrm{U}(V_i)$. Fix a self-dual O_F -lattice

$$\Lambda_0 \subset V_0,$$

which exists and is unique up to $H_0(F_0)$ -conjugacy. Let

$$\Lambda_0^\sharp := \Lambda_0 \oplus O_F u_0 \subset V_0^\sharp,$$

which is again self-dual. We denote by

$$K_0 \subset H_0(F_0)$$

the stabilizer of Λ_0 , and by

$$K_0^\sharp \subset G_0(F_0) \quad \text{and} \quad \mathfrak{k}_0^\sharp \subset \mathfrak{g}_0(F_0)$$

the respective stabilizers of Λ_0 . Then K_0 and K_0^\sharp are both hyperspecial maximal subgroups.

We normalize the Haar measures on the groups

$$\mathrm{GL}_{n-1}(F_0) \quad \text{and} \quad \mathrm{U}(V_0)(F_0)$$

by assigning measure one to each of the respective subgroups

$$\mathrm{GL}_{n-1}(O_{F_0}) \quad \text{and} \quad K_0.$$

With respect to these normalizations, the Jacquet–Rallis fundamental lemma conjecture is the following statement; cf. [38, §3]. Note that the semi-Lie algebra version below is essentially the Fourier–Jacobi case arising from the relative trace formula of Yifeng Liu [29].

CONJECTURE 2.3 (Jacquet–Rallis fundamental lemma conjecture).

(a) (The group version) *The characteristic function $1_{S_n(O_{F_0})} \in \mathcal{S}(S_n(F_0))$ transfers to the pair of functions $(1_{K_0}, 0) \in \mathcal{S}(G_0(F_0)) \times \mathcal{S}(G_1(F_0))$.*

(b) (The semi-Lie algebra version) *The characteristic function*

$$1_{(S_{n-1} \times V'_{n-1})(O_{F_0})} \in \mathcal{S}((S_{n-1} \times V'_{n-1})(F_0))$$

transfers to the pair of functions

$$(1_{K_0 \times \Lambda_0}, 0) \in \mathcal{S}((H_0 \times V_0)(F_0)) \times \mathcal{S}((H_1 \times V_1)(F_0)).$$

Remark 2.4. There is also a Lie algebra version: the characteristic function $1_{\mathfrak{s}_n(O_{F_0})} \in \mathcal{S}(\mathfrak{s}_n(F_0))$ transfers to the pair of functions $(1_{\mathfrak{k}_0}, 0) \in \mathcal{S}(\mathfrak{g}_0(F_0)) \times \mathcal{S}(\mathfrak{g}_1(F_0))$. This Lie algebra version is equivalent to the group version, at least when p is odd; cf. [45, §2.6].

Remark 2.5. We note that the equal characteristic analog of the FL conjecture was proved by Z. Yun for $p > n$; cf. [45]; J. Gordon deduced the p -adic case for p large, but unspecified; cf. [13].

It is straightforward to check a special case.

PROPOSITION 2.6. *The semi-Lie algebra version FL holds for $(g, u) \in (G_0 \times V_0)(F_0)_{\text{rs}}$ when g is regular semisimple (i.e., $F[g]$ is a product of fields with total degree equal to $\dim V$) and generates a maximal order $O_F[g]$ (in $F[g]$).*

Proof. This is easy to check (see, e.g., [45, Lem. 2.5.5] for the Lie algebra version), but the argument is the same for the semi-Lie algebra version. \square

PROPOSITION 2.7. *Fix F/F_0 . Assume that $q \geq n$, where q denotes the cardinality of the residue field of O_{F_0} . Then*

- (i) *in Conjecture 2.3, parts (a) and (b) are equivalent;*
- (ii) *in Conjecture 2.3, part (a) for S_{n-1} implies part (b) for $(g, u) \in (H_0 \times V_0)(F_0)_{\text{rs}}$ where the norm of u is a unit.*

Proof. We will prove a similar statement, namely Proposition 4.12 for the AFL conjecture, where the situation is more delicate; we omit the argument here and only point out that the proof also works here. \square

3. AFL and variants

For Sections 3 and 4, we let F be an unramified quadratic field extension of a p -adic local field F_0 for an odd prime p .

3.1. The AFL conjecture and variants. For any $n \geq 1$, we recall the construction of the Rapoport–Zink formal moduli scheme $\mathcal{N}_n = \mathcal{N}_{n, F/F_0}$ associated to unitary groups; cf. [38, §4]. For $\text{Spf } O_F$ -schemes S , we consider triples (X, ι, λ) , where

- X is a p -divisible group of absolute height $2nd$ and dimension n over S , where $d := [F_0 : \mathbb{Q}_p]$,
- ι is an action of O_F such that the induced action of O_{F_0} on $\text{Lie } X$ is via the structure morphism $O_{F_0} \rightarrow \mathcal{O}_S$, and
- λ is a principal (O_{F_0} -relative) polarization.

Hence $(X, \iota|_{O_{F_0}})$ is a formal O_{F_0} -module of relative height $2n$ and dimension n . We require that the Rosati involution Ros_λ on O_F is the non-trivial Galois automorphism in $\text{Gal}(F/F_0)$, and that the *Kottwitz condition* of signature $(n-1, 1)$ is satisfied, i.e.,

$$(3.1) \quad \text{char}(\iota(a) \mid \text{Lie } X) = (T - a)^{n-1}(T - \bar{a}) \in \mathcal{O}_S[T] \quad \text{for all } a \in O_F.$$

An isomorphism $(X, \iota, \lambda) \xrightarrow{\sim} (X', \iota', \lambda')$ between two such triples is an O_F -linear isomorphism $\varphi : X \xrightarrow{\sim} X'$ such that $\varphi^*(\lambda') = \lambda$.

Over the residue field \bar{k} of O_{F^\flat} , there is a triple $(\mathbb{X}_n, \iota_{\mathbb{X}_n}, \lambda_{\mathbb{X}_n})$ such that \mathbb{X}_n is supersingular, unique up to O_F -linear quasi-isogeny compatible with the polarization. We fix such a triple, which we call a *framing object* (for the functor \mathcal{N}_n). Then \mathcal{N}_n (pro-)represents the functor over $\text{Spf } O_{F^\flat}$ that associates to each S the set of isomorphism classes of quadruples $(X, \iota, \lambda, \rho)$ over S , where the final entry is an O_F -linear quasi-isogeny of height zero defined over the special fiber,

$$\rho : X \times_S \bar{S} \longrightarrow \mathbb{X}_n \times_{\text{Spec } \bar{k}} \bar{S},$$

such that $\rho^*((\lambda_{\mathbb{X}_n})_{\bar{S}}) = \lambda_{\bar{S}}$. Here ρ is called a *framing*. The formal scheme \mathcal{N}_n is smooth over $\text{Spf } O_{F^\flat}$ of relative dimension $n-1$.

For $n \geq 2$, define the product $\mathcal{N}_{n-1,n} := \mathcal{N}_{n-1} \times_{\text{Spf } O_{F^\flat}} \mathcal{N}_n$. It is a (locally Noetherian) formal scheme of (formal) dimension $2(n-1)$, formally smooth over $\text{Spf } O_{F^\flat}$.

When $n = 1$, we have the (unique up to isomorphism) formal O_F -module \mathbb{E} (with signature $(1, 0)$) over \bar{k} and its canonical lift \mathcal{E} over O_{F^\flat} , as well as the “conjugate” objects $\bar{\mathbb{E}}$ and $\bar{\mathcal{E}}$ (with signature $(0, 1)$). For $n \geq 2$, there is a natural closed embedding of formal schemes

$$(3.2) \quad \begin{aligned} \delta = \delta_{\mathcal{N}_{n-1}} : \mathcal{N}_{n-1} &\longrightarrow \mathcal{N}_n \\ (X, \iota, \lambda, \rho) &\longmapsto (X \times \mathcal{E}, \iota \times \iota_{\mathcal{E}}, \lambda \times \lambda_{\mathcal{E}}, \rho \times \rho_{\mathcal{E}}), \end{aligned}$$

where we set $\mathbb{X}_1 = \bar{\mathbb{E}}$ and inductively take

$$(3.3) \quad \mathbb{X}_n = \mathbb{X}_{n-1} \times \mathbb{E}$$

as the framing object for \mathcal{N}_n . Let

$$(3.4) \quad \Delta_{\mathcal{N}_{n-1}} : \mathcal{N}_{n-1} \xrightarrow{(\text{id}_{\mathcal{N}_{n-1}}, \delta)} \mathcal{N}_{n-1} \times_{\text{Spf } O_{F^\flat}} \mathcal{N}_n = \mathcal{N}_{n-1,n}$$

be the graph morphism of δ . Then

$$(3.5) \quad \Delta := \Delta_{\mathcal{N}_{n-1}}(\mathcal{N}_{n-1})$$

is a closed formal subscheme of half the formal dimension of $\mathcal{N}_{n-1,n}$. Note that

$$(3.6) \quad \text{Aut}^\circ(\mathbb{X}_n, \iota_{\mathbb{X}_n}, \lambda_{\mathbb{X}_n}) \cong \text{U}(\mathbb{V}_n)(F_0),$$

where the left-hand side is the group of quasi-isogenies of the framing object $(\mathbb{X}_n, \iota_{\mathbb{X}_n}, \lambda_{\mathbb{X}_n})$, and

$$\mathbb{V}_n := \text{Hom}_{O_F}^0(\mathbb{E}, \mathbb{X}_n)$$

is the hermitian space on which the hermitian form is induced by the principle polarizations on \mathbb{X}_n and \mathbb{E} . Note that \mathbb{V}_n does not contain a self-dual lattice.

More concretely,

$$U(\mathbb{V}_n)(F_0) = \{g \in \text{End}_F(\mathbb{V}_n) \mid gg^* = \text{id}\}.$$

Here we denote by $g^* = \text{Ros}_{\lambda_{\mathbb{X}_n}}(g)$ the Rosati involution. Then the group $U(\mathbb{V}_n)(F_0)$ acts naturally on \mathcal{N}_n by acting on the framing:

$$g \cdot (X, \iota, \lambda, \rho) = (X, \iota, \lambda, g \circ \rho).$$

Furthermore, \mathbb{V}_n contains a natural special vector u_0 given by the inclusion of \mathbb{E} in $\mathbb{X}_n = \mathbb{X}_{n-1} \times \mathbb{E}$ via the second factor. The norm of u_0 is 1. Then \mathbb{V}_n is a non-split hermitian space of dimension n . Therefore, in the setting of [Section 2.3](#), we can choose identifications $V_1^\sharp = \mathbb{V}_n$ and $V_1 = \mathbb{V}_{n-1}$ compatible with the natural inclusions on both sides. Hence we obtain an action of $H_1(F_0)$ on \mathcal{N}_{n-1} , of $G_1(F_0)$ on \mathcal{N}_n , and of $G_{V_1}(F_0) = (U(\mathbb{V}_{n-1}) \times U(\mathbb{V}_n))(F_0)$ on $\mathcal{N}_{n-1,n}$; cf. [\(2.5\)](#). Furthermore, the maps $\delta_{\mathcal{N}_{n-1}}$ and $\Delta_{\mathcal{N}_{n-1}}$ are equivariant with respect to the respective embeddings $H_1(F_0) \hookrightarrow G_1(F_0)$ and $H_1(F_0) \hookrightarrow G_{V_1}(F_0)$.

For $g \in G_{V_1}(F_0)_{\text{rs}}$, we denote by $\text{Int}(g)$ the intersection product on $\mathcal{N}_{n-1,n}$ of Δ with its translate $g\Delta$, defined through the derived tensor product of the structure sheaves (cf. [\(B.4\)](#)):

$$(3.7) \quad \text{Int}(g) := \langle \Delta, g\Delta \rangle_{\mathcal{N}_{n-1,n}} := \chi(\mathcal{N}_{n-1,n}, \mathcal{O}_\Delta \overset{\mathbb{L}}{\otimes} \mathcal{O}_{g\Delta}).$$

We similarly define $\text{Int}(g)$ for $g \in G_1(F_0)_{\text{rs}}$,

$$(3.8) \quad \text{Int}(g) := \langle \Delta, (1 \times g)\Delta \rangle_{\mathcal{N}_{n-1,n}}.$$

In both cases, when g is regular semisimple, the right-hand side of this definition is finite since the (formal) schematic intersection $\Delta \cap g\Delta$ is a proper *scheme* over $\text{Spf } O_{\mathbb{F}}$. We refer to [Appendix B](#) for the terminology regarding various K -groups of formal schemes, following the work of Gillet–Soulé for schemes in [\[11\]](#).

Now we introduce a new variant of the above intersection number $\text{Int}(g)$ via the Kudla–Rapoport special divisors [\[24\]](#). This variant is closely related to in the AFL conjecture in the context of Fourier–Jacobi cycles in the work of Yifeng Liu [\[30, Conj. 1.11\]](#). A special case has also appeared in Mihatsch’s thesis [\[31, §8\]](#).

Recall from [\[24\]](#) that for every non-zero $u \in \mathbb{V}_n$, Kudla and Rapoport have defined a special divisor $\mathcal{Z}(u)$ in \mathcal{N}_n . This is the locus where the quasi-homomorphism $u : \mathbb{E} \rightarrow \mathbb{X}_n$ lifts to a homomorphism from \mathcal{E} to the universal

object over \mathcal{N}_n . By [24, Prop. 3.5], $\mathcal{Z}(u)$ is a relative divisor (or empty). Then the morphism δ in (3.2) induces an obvious closed embedding

$$(3.9) \quad \mathcal{N}_{n-1} \xrightarrow{\sim} \mathcal{Z}(u_0)$$

for the unit norm special vector u_0 , which is an isomorphism by [24, Lem. 5.2]. It follows from the definition that if $g \in \mathrm{U}(\mathbb{V}_n)(F_0)$, then

$$(3.10) \quad g\mathcal{Z}(u) = \mathcal{Z}(gu).$$

For simplicity, we will write $\mathcal{N}_n \times \mathcal{N}_n$ for the fiber product $\mathcal{N}_n \times_{\mathrm{Spf} O_{\breve{F}}} \mathcal{N}_n$. For $g \in \mathrm{U}(\mathbb{V}_n)(F_0)$, let $\Gamma_g \subset \mathcal{N}_n \times \mathcal{N}_n$ be the graph of the automorphism of \mathcal{N}_n induced by g . We define *the (naive) fixed point locus*, denoted by \mathcal{N}_n^g , as the (formal) schematic intersection (i.e., fiber product of formal schemes)

$$(3.11) \quad \mathcal{N}_n^g := \Gamma_g \cap \Delta_{\mathcal{N}_n},$$

viewed as a closed formal subscheme of \mathcal{N}_n . We also form a *derived fixed point locus*, denoted by ${}^{\mathbb{L}}\mathcal{N}_n^g$, i.e., the derived tensor product

$$(3.12) \quad {}^{\mathbb{L}}\mathcal{N}_n^g := \Gamma_g \cap \Delta_{\mathcal{N}_n} := \mathcal{O}_{\Gamma_g} \xrightarrow{\mathbb{L}} \otimes_{\mathcal{O}_{\mathcal{N}_n \times \mathcal{N}_n}} \mathcal{O}_{\Delta_{\mathcal{N}_n}}$$

viewed as an element in $K_0^{\mathcal{N}_n^g}(\mathcal{N}_n)$; cf. Section B.1.

For a pair $(g, u) \in (\mathrm{U}(\mathbb{V}_n) \times \mathbb{V}_n)(F_0)_{\mathrm{rs}}$, we define (cf. (B.4))

$$(3.13) \quad \mathrm{Int}(g, u) := \langle {}^{\mathbb{L}}\mathcal{N}_n^g, \mathcal{Z}(u) \rangle_{\mathcal{N}_n} := \chi \left(\mathcal{N}_n, {}^{\mathbb{L}}\mathcal{N}_n^g \xrightarrow{\mathbb{L}} \otimes_{\mathcal{O}_{\mathcal{N}_n}} \mathcal{O}_{\mathcal{Z}(u)} \right).$$

Similar to (3.7) and (3.8), when (g, u) is regular semisimple, $\mathcal{N}_n^g \cap \mathcal{Z}(u)$ is a proper *scheme* over $\mathrm{Spf} O_{\breve{F}}$ and hence the right-hand side of this definition is finite. The number $\mathrm{Int}(g, u)$ depends only on its $\mathrm{U}(\mathbb{V}_n)(F_0)$ -orbit.

Remark 3.1. By the projection formula for the closed immersion $\Delta : \mathcal{N}_n \rightarrow \mathcal{N}_n \times \mathcal{N}_n$, we obtain an equality in $K_0^{\Gamma_g \cap \Delta(\mathcal{Z}(u))}(\mathcal{N}_n \times \mathcal{N}_n)$,

$$\mathrm{R}\Delta_*({}^{\mathbb{L}}\mathcal{N}_n^g \xrightarrow{\mathbb{L}} \otimes_{\mathcal{O}_{\mathcal{N}_n}} \mathcal{O}_{\mathcal{Z}(u)}) = \mathcal{O}_{\Gamma_g} \xrightarrow{\mathbb{L}} \otimes_{\mathcal{O}_{\mathcal{N}_n \times \mathcal{N}_n}} \mathcal{O}_{\Delta(\mathcal{Z}(u))},$$

where we have used $\mathrm{R}\Delta_*(\mathcal{O}_{\mathcal{Z}(u)}) = \mathcal{O}_{\Delta(\mathcal{Z}(u))}$ for a closed immersion. Therefore, an equivalent definition of the intersection number (3.13) is

$$\mathrm{Int}(g, u) = \chi \left(\mathcal{N}_n \times \mathcal{N}_n, \mathcal{O}_{\Gamma_g} \xrightarrow{\mathbb{L}} \otimes_{\mathcal{O}_{\mathcal{N}_n \times \mathcal{N}_n}} \mathcal{O}_{\Delta_{\mathcal{Z}(u)}} \right).$$

This also appears in the AFL in the context of Fourier–Jacobi cycles in [30].

CONJECTURE 3.2 (Arithmetic fundamental lemma conjecture).

(a) (The group version) *Suppose that $\gamma \in S_n(F_0)_{\mathrm{rs}}$ matches an element $g \in \mathrm{U}(\mathbb{V}_n)(F_0)_{\mathrm{rs}}$. Then*

$$\partial\mathrm{Orb}(\gamma, \mathbf{1}_{S_n(O_{F_0})}) = -\mathrm{Int}(g) \cdot \log q.$$

(b) (The semi-Lie algebra version) Suppose that $(\gamma, u') \in (S_n \times V'_n)(F_0)_{\text{rs}}$ matches an element $(g, u) \in (U(\mathbb{V}_n) \times \mathbb{V}_n)(F_0)_{\text{rs}}$. Then

$$\partial\text{Orb}((\gamma, u'), \mathbf{1}_{(S_n \times V'_n)(O_{F_0})}) = -\text{Int}(g, u) \cdot \log q.$$

Remark 3.3. We refer to [38, §4, Conj. 4.1(a)] for the homogeneous group version of AFL involving the intersection numbers (3.7) rather than (3.8); it is equivalent to part (a) of [Conjecture 3.2](#).

Remark 3.4. Mihatsch [33] has pointed out that a naive formulation of Lie algebra version of AFL is not well behaved (unless the formal schematic intersection is artinian), unlike the case of FL (cf. [Remark 2.4](#)). Therefore the semi-Lie algebraic version seems to be the best possible linearization of the AFL conjecture.

Definition 3.5.

- (a) A regular semisimple element $(g, u) \in (U(V) \times V)(F_0)$ is called strongly regular semisimple (“srs” for short) if $g \in U(V)(F_0)$ is semisimple with respect to the conjugation action of $U(V)$.
- (b) A regular semisimple element $g \in U(V^\sharp)(F_0)$ with respect to the conjugation action of $U(V)$ is called strongly regular semisimple (“srs” for short) if it is also semisimple with respect to the conjugation action of $U(V^\sharp)$.

Definition 3.6.

- (a) A regular semisimple element $(\gamma, u') \in (S_{n-1} \times V'_{n-1})(F_0)$ is called strongly regular semisimple (“srs” for short) if $\gamma \in S_{n-1}(F_0)$ is semisimple with respect to the conjugation action of GL_{n-1, F_0} .
- (b) A regular semisimple element $\gamma' \in S_n(F_0)$ with respect to the conjugation action of GL_{n-1, F_0} is called strongly regular semisimple (“srs” for short) if it is also semisimple with respect to the conjugation action of GL_{n, F_0} .

Remark 3.7. On the Lie algebras the notion of “strongly regular semisimple” has appeared in [45].

CONJECTURE 3.8 (Arithmetic fundamental lemma conjecture for strongly regular semisimple elements).

- (a) (The group version) Suppose that $\gamma \in S_n(F_0)_{\text{srs}}$ matches an element $g \in U(\mathbb{V}_n)(F_0)_{\text{srs}}$. Then

$$\partial\text{Orb}(\gamma, \mathbf{1}_{S_n(O_{F_0})}) = -\text{Int}(g) \cdot \log q.$$

- (b) (The semi-Lie algebra version) Suppose that $(\gamma, u') \in (S_{n-1} \times V'_{n-1})(F_0)_{\text{srs}}$ matches an element $(g, u) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$. Then

$$\partial\text{Orb}((\gamma, u'), \mathbf{1}_{(S_{n-1} \times V'_{n-1})(O_{F_0})}) = -\text{Int}(g, u) \cdot \log q.$$

3.2. A special case of AFL.

PROPOSITION 3.9. *Let $p > n$. Conjecture 3.8 part (b) (i.e., the semi-Lie algebra version AFL) holds for $(g, u) \in (\mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\mathrm{srs}}$ when $O_F[g]$ is a maximal order (in $F[g]$).*

Proof. This follows from [31, Cor. 9.9] (as well as the fact that the assertion holds when $n = 2$). When $F_0 = \mathbb{Q}_p$, this can also be deduced from [18]. \square

4. Relation between the two versions of AFL

In this section, we continue to let F be an unramified quadratic field extension of a p -adic local field F_0 for an odd prime p .

4.1. *Orbits in $\mathrm{U}(\mathbb{V}_n)$.* We recall that the Cayley map is the rational morphism

$$(4.1) \quad \begin{aligned} \mathfrak{c} = \mathfrak{c}_n : \quad \mathfrak{u}(\mathbb{V}_n) &\longrightarrow \mathrm{U}(\mathbb{V}_n) \\ x &\longmapsto -\frac{1-x}{1+x}. \end{aligned}$$

Here $\frac{1+x}{1-x}$ denotes $(1-x)^{-1}(1+x) = (1+x)(1-x)^{-1}$ since the two factors commute. Its inverse is

$$\mathfrak{c}^{-1}(g') = \frac{1+g'}{1-g'}.$$

By definitions $\mathbb{V}_n = \mathrm{Hom}_{O_F}^{\circ}(\mathbb{E}, \mathbb{X}_n)$ and $\mathbb{X}_n = \mathbb{X}_{n-1} \times \mathbb{E}$, we decompose

$$\mathbb{V}_n = \mathbb{V}_{n-1} \oplus \mathrm{End}_{O_F}^{\circ}(\mathbb{E}) = \mathbb{V}_{n-1} \oplus F u_0.$$

Accordingly, write $g' \in \mathrm{U}(\mathbb{V}_n)$ in the matrix form

$$(4.2) \quad \begin{aligned} g' = \begin{pmatrix} h & u \\ w^* & d \end{pmatrix} \\ \mathbb{X}_{n-1} \times \mathbb{E} \xrightarrow{\quad} \mathbb{X}_{n-1} \times \mathbb{E}, \end{aligned}$$

where $*$ denotes the map $\mathrm{Hom}_{O_F}^{\circ}(\mathbb{E}, \mathbb{X}_{n-1}) \rightarrow \mathrm{Hom}_{O_F}^{\circ}(\mathbb{X}_{n-1}, \mathbb{E})$ induced by polarizations on \mathbb{X}_{n-1} and \mathbb{E} , and

$$h \in \mathrm{End}_{O_F}^{\circ}(\mathbb{X}_{n-1}), \quad u, w \in \mathbb{V}_{n-1}, \quad d \in \mathrm{End}_{O_F}^{\circ}(\mathbb{E}).$$

LEMMA 4.1. *Let $g' \in \mathrm{U}(\mathbb{V}_n)$ be as in (4.2). Write*

$$(4.3) \quad x' = \mathfrak{c}_n^{-1}(g') = \begin{pmatrix} x & \tilde{u} \\ -\tilde{u}^* & e \end{pmatrix} \in \mathfrak{u}(\mathbb{V}_n),$$

and define

$$(4.4) \quad g := \mathfrak{c}_{n-1}(x) \in \mathrm{U}(\mathbb{V}_{n-1}).$$

Then

$$(4.5) \quad \begin{cases} g = h + (1-d)^{-1}u w^*, \\ \tilde{u} = 2(1-d)^{-1}(1-g)^{-1}u, \\ \det(1-g') = (1-d)\det(1-g), \\ gw = \epsilon_d u, \end{cases}$$

where we define

$$(4.6) \quad \epsilon_d := \frac{1-\bar{d}}{1-d}.$$

Proof. By definition of \mathfrak{c}_n^{-1} , we expand the equality $1+g' = (1-g')x'$,

$$\begin{pmatrix} 1+h & u \\ w^* & 1+d \end{pmatrix} = \begin{pmatrix} 1-h & -u \\ -w^* & 1-d \end{pmatrix} \begin{pmatrix} x & \tilde{u} \\ -\tilde{u}^* & e \end{pmatrix},$$

to obtain

$$\begin{cases} 1+h = (1-h)x + u\tilde{u}^*, \\ w^* = -w^*x - (1-d)\tilde{u}^*. \end{cases}$$

The second equality yields

$$\tilde{u}^* = -(1-d)^{-1}w^*(1+x),$$

which implies that

$$(4.7) \quad \tilde{u} = -(1-\bar{d})^{-1}(1-x)w.$$

Plug into the first equality

$$1+h = (1-h)x - (1-d)^{-1}uw^*(1+x),$$

and this implies that

$$1+h + (1-d)^{-1}uw^* = (1-h - (1-d)^{-1}uw^*)x.$$

It follows that

$$g = \mathfrak{c}_{n-1}(x) = h + (1-d)^{-1}uw^*,$$

and this proves the first equality in (4.5).

Now note that the condition for $g'g'^* = 1$ amounts to

$$(4.8) \quad hh^* + uu^* = 1, \quad hw + \bar{d}u = 0, \quad w^*w + d\bar{d} = 1.$$

The last equality in (4.5) now follows:

$$\begin{aligned} gw &= hw + (1-d)^{-1}uw^*w \\ &= (-\bar{d} + (1-d\bar{d}))(1-d)^{-1}u \\ &= \frac{1-\bar{d}}{1-d}u. \end{aligned}$$

Now we return to (4.7), noting that $1 - x = -2(1 - g)^{-1}g$,

$$\tilde{u} = 2(1 - \bar{d})^{-1}(1 - g)^{-1}gw = 2(1 - d)^{-1}(1 - g)^{-1}u.$$

This proves the second equality in (4.5).

Finally, by $1 - g' = \begin{pmatrix} 1-h & -u \\ -w^* & 1-d \end{pmatrix}$ and the first equality in (4.5), we have

$$\begin{aligned} \det(1 - g') &= (1 - d) \det((1 - h) - (1 - d)^{-1}uw^*) \\ &= (1 - d) \det(1 - g). \end{aligned}$$

This proves the third equality in (4.5) and completes the proof. \square

We now define a rational map by the formulas in Lemma 4.1,

$$\begin{aligned} (4.9) \quad \mathfrak{r} : \mathrm{U}(\mathbb{V}_n) &\longrightarrow \mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \\ g' &\longmapsto \left(g, \frac{u}{(1-d)\sqrt{\epsilon}} \right), \end{aligned}$$

where $\epsilon \in O_{F_0}^\times$ is chosen such that $F = F_0[\sqrt{\epsilon}]$. We also define a variant

$$\begin{aligned} (4.10) \quad \mathfrak{r}^\sharp : \mathrm{U}(\mathbb{V}_n) &\longrightarrow \mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \\ g' &\longmapsto \left(g, \frac{\tilde{u}}{\sqrt{\epsilon}} \right). \end{aligned}$$

Following the notation in Lemma 4.1, let $\mathrm{U}(\mathbb{V}_n)^\circ$ be the open sub-variety of $\mathrm{U}(\mathbb{V}_n)$ defined by

$$1 - d \neq 0 \quad \text{and} \quad \det(1 - g') \neq 0.$$

Let $(\mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ$ be the open sub-variety of $\mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1)$ defined by

$$\det(1 - g) \neq 0 \quad \text{and} \quad \det(1 + x') \neq 0.$$

Here $(g, \tilde{u}, e) \in \mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1)$ and x' is as in (4.3), where $x = \mathfrak{c}_{n-1}^{-1}(g)$.

LEMMA 4.2. *The map \mathfrak{r} together with $e \in \mathfrak{u}(1)$ (cf. (4.3)) induce an isomorphism, equivariant under the action of $\mathrm{U}(\mathbb{V}_{n-1})$,*

$$\begin{aligned} \tilde{\mathfrak{r}} = (\mathfrak{r}, e) : \mathrm{U}(\mathbb{V}_n)^\circ &\xrightarrow{\sim} (\mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ \\ g' &\longmapsto (\mathfrak{r}(g'), e). \end{aligned}$$

The same holds if we replace \mathfrak{r} by \mathfrak{r}^\sharp .

Proof. By (4.5) we have

$$\det(1 - g') = (1 - d) \det(1 - g),$$

and by $1 - d \neq 0$, it follows that $\det(1 - g) \neq 0$. Then the map $x \mapsto \mathfrak{c}(x)$ is well defined since $1 - g = \frac{1}{1+x}$. Therefore the rational map $\tilde{\mathfrak{r}} = (\mathfrak{r}, e)$ is defined on $\mathrm{U}(\mathbb{V}_n)^\circ$

To reverse the map $\tilde{\tau}$, let $(g, u, e) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ$. First we send g to $\mathfrak{c}^{-1}(g) = x$. (This is defined since $\det(1-g) \neq 0$.) Then we define \tilde{u} by $\tilde{u} = 2\sqrt{\epsilon}(1-g)^{-1}u$; cf. (4.5) and (4.9). Finally, we apply Cayley map \mathfrak{c} (4.1) to $(\begin{smallmatrix} x & \tilde{u} \\ -\tilde{u}^* & e \end{smallmatrix})$ to obtain g' . (The Cayley map is well-defined by the second condition $\det(1+x') \neq 0$ when defining $(U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ$.) It is easy to see that the composition of above maps is defined on $(U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ$ and defines an inverse to the rational map $\tilde{\tau}$. The desired assertion for $\tilde{\tau}$ follows. It is easy to see the assertion for τ^\sharp . \square

We may apply the same construction to $\xi g'$ for $\xi \in F^1 = \ker(\mathrm{Nm} : F^\times \rightarrow F_0^\times)$:

$$(4.11) \quad \begin{aligned} \tau_\xi : U(\mathbb{V}_n) &\longrightarrow U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \\ g' &\longmapsto \tau(\xi g'). \end{aligned}$$

We define the variant τ_ξ^\sharp similar to (4.10).

LEMMA 4.3.

- (i) An element $g' \in U(\mathbb{V}_n)^\circ(F_0)$ is regular semisimple (with respect to the conjugation action of $U(\mathbb{V}_{n-1})$ for $\mathbb{V}_n = \mathbb{V}_{n-1} \oplus F u_0$) if and only if $\tau(g') = (g, u)$ is regular semisimple as an element in $(U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)$.
- (ii) Let $g' \in U(\mathbb{V}_n)^\circ(F_0)_{\mathrm{srs}}$. Then, for all but finitely many $\xi \in F^1$, we have $\xi g' \in U(\mathbb{V}_n)^\circ(F_0)$ and $\tau_\xi(g') \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\mathrm{srs}}$.
- (iii) Let $(g, u) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\mathrm{srs}}$. Then, for all but finitely many $e \in \mathfrak{u}(1)$, we have $(g, u, e) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ(F_0)$ and $\tilde{\tau}^{-1}(g, u, e) \in U(\mathbb{V}_n)^\circ(F_0)_{\mathrm{srs}}$.

Proof. The regular semi-simplicity for $g' \in U(\mathbb{V}_n)(F_0)$ is equivalent to the vectors

$$\{g'^i u_0 \mid 0 \leq i \leq n-1\}$$

being a basis of \mathbb{V}_n (as an F -vector space). By the decomposition (4.2), this is equivalent to $\{h^i u, 0 \leq i \leq n-2\}$ being a basis of \mathbb{V}_{n-1} . By the first equality in (4.5), we can show inductively that, for all $1 \leq i \leq n-2$, $g^i u - h^i u$ lies in the span of $u, hu, \dots, h^{i-1}u$. This proves part (i).

Let $P(\lambda) = \det(\lambda + h)$ be the characteristic polynomial of h , and let

$$Q(\lambda) = \det(\lambda + h) \cdot w^*(\lambda + h)^{-1}u,$$

which is a polynomial in λ of degree $n-2$. Then the characteristic polynomial of g' can be written as

$$(4.12) \quad \det(\lambda + g') = (\lambda + d)P(\lambda) - Q(\lambda).$$

Since $g' \in U(\mathbb{V}_n)^\circ(F_0)_{\mathrm{srs}}$ (particularly, regular semisimple relative to the $U(\mathbb{V}_n)$ -conjugation action), this polynomial in λ has only simple roots.

Let $\tau_\xi(g') = (g_\xi, u_\xi)$. We now study how the characteristic polynomial of g_ξ (or equivalently, of $\xi^{-1}g_\xi$) depends on ξ . By the first equality in (4.5),

$$\det(\lambda + \xi^{-1}g_\xi) = \det\left(\lambda + h + \frac{\xi}{1-d\xi}uw^*\right).$$

Set

$$t = \frac{\xi}{1-d\xi}.$$

Then

$$\begin{aligned} \det(\lambda + \xi^{-1}g_\xi) &= \det(\lambda + h) \det(1 + t uw^*(\lambda + h)^{-1}) \\ &= \det(\lambda + h) (1 + t w^*(\lambda + h)^{-1}u) \\ &= \det(\lambda + h) + t \det(\lambda + h) w^*(\lambda + h)^{-1}u \\ &= P(\lambda) + t Q(\lambda). \end{aligned}$$

Here in the second equality we have used the fact that $uw^* \in \text{End}(\mathbb{V}_{n-1})$ is of rank at most one.

Let $R(\xi)$ be the GCD of $P(\lambda)$ and $Q(\lambda)$. By the semi-simplicity of g' , the polynomial $R(\lambda)$ is multiplicity free. Fix an algebraic closed field $\Omega \supset F$. Since there are only finitely many $t \in \Omega$ such that $P/R + t Q/R$ and R have common roots, the question is reduced to the case $R = 1$ (and possibly smaller n). Now assume $R = 1$. Then $P + t Q \in F[t, \lambda]$ is an irreducible (over Ω) polynomial in t, λ , hence defines an irreducible curve C in A_F^2 (the affine plane in t, λ), and t defines a non-constant rational morphism to the projective line $C \rightarrow \mathbf{P}_F^1$. The polynomial $P + t Q$ has a repeated root precisely when the rational morphism is ramified at t . Hence there are only finitely many such $t \in \Omega$. This proves part (ii).

Part (iii) is proved similarly to part (ii). \square

4.2. Reduction of the intersection numbers. We recall from (3.2) that $\delta : \mathcal{N}_{n-1} \rightarrow \mathcal{N}_n$ is the embedding whose image is the special divisor $\mathcal{Z}(u_0)$ for a unit $u_0 \in \text{End}_{O_F}^0(\mathbb{E})$; cf. (3.9). Consider

$$\mathcal{N}_{n-1} \times \mathcal{N}_{n-1} \xrightarrow{(\delta, \delta)} \mathcal{N}_n \times \mathcal{N}_n,$$

and let $\pi_2 : \mathcal{N}_{n-1} \times \mathcal{N}_{n-1} \rightarrow \mathcal{N}_n$ be the projection to the second factor. We have the following pull-back formula for the graph of an automorphism.

LEMMA 4.4. *Let $g' \in \text{U}(\mathbb{V}_n)^\circ(F_0)$ be such that $1-d \in O_F^\times$, and let $(g, u) = \tau(g') \in (\text{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)$. Then*

$$(4.13) \quad (\delta, \delta)^* \Gamma_{g'} \simeq \Gamma_g \cap \pi_2^* \mathcal{Z}(u),$$

where $(\delta, \delta)^*$ is the naive pull-back, i.e., the fiber product

$$\begin{array}{ccc} (\delta, \delta)^* \Gamma_{g'} & \longrightarrow & \Gamma_{g'} \\ \downarrow & \square & \downarrow \\ \mathcal{N}_{n-1} \times \mathcal{N}_{n-1} & \xrightarrow{(\delta, \delta)} & \mathcal{N}_n \times \mathcal{N}_n. \end{array}$$

Moreover, if u is non-zero, then

$$(4.14) \quad \mathcal{O}_{\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_n \times \mathcal{N}_n} \mathcal{O}_{\Gamma_{g'}} = \mathcal{O}_{\Gamma_g \cap \pi_2^* \mathcal{Z}(u)} = \mathcal{O}_{\Gamma_g} \xrightarrow{\mathbb{L}} \mathcal{O}_{\pi_2^* \mathcal{Z}(u)},$$

as elements in $K'_0(\Gamma_g \cap \pi_2^* \mathcal{Z}(u))$.

Remark 4.5. By (4.5), we have $gw = \epsilon_d u$. Since $d \neq 1$, $\epsilon_d = \frac{1-\bar{d}}{1-d}$ is a unit in O_F , and hence we may replace $\pi_2^* \mathcal{Z}(u)$ by $\pi_1^* \mathcal{Z}(u)$ in the above statements.

Proof. We prove that the natural map on S -points is the identity map. Let (X_1, X_2) be an S -point of $\mathcal{N}_{n-1} \times_{\text{Spf } O_F} \mathcal{N}_{n-1}$, and let $X'_i = X_i \times \mathcal{E}$. (In the notation we have omitted S and the obvious additional structure ι, λ etc.)

We start from (X_1, X_2) on the graph $\Gamma_{g'}$; i.e., there exists (uniquely) $\varphi' : X'_1 \rightarrow X'_2$ lifting g' . Write φ' in the matrix form

$$X_1 \times \mathcal{E} \xrightarrow{\varphi' = \begin{pmatrix} \varphi & \psi \\ \psi'^* & d \end{pmatrix}} X_2 \times \mathcal{E},$$

which lifts the diagram (4.2). We then need to construct elements in $\Gamma_g \cap \pi_2^* \mathcal{Z}(u)$. The subtle point is that X_1 and X_2 are different, whereas the \mathbb{X}_n in the target and the source in the map g' of (4.2) are (unfortunately) identified.

First we have $X_2 \in \mathcal{Z}(u)$. (Note that the u in $\mathfrak{r}(g') = (g, u)$ differs from the u in (4.2) only by a unit $(1-d)\sqrt{\epsilon}$, hence we ignore the difference in this proof.) Consider the homomorphism

$$\tilde{\varphi} := \varphi + \frac{\psi \psi'^*}{1-d} : X_1 \longrightarrow X_2.$$

This is a lifting of $g \in U(\mathbb{V}_n)$ by Lemma 4.1, hence we have constructed (X_1, X_2) on $\Gamma_g \cap \pi_2^* \mathcal{Z}(u)$. Again by Lemma 4.1, ψ' lifts $\epsilon_d g^{-1} u$ (and $\epsilon_d = \frac{1-\bar{d}}{1-d}$ is a unit), hence

$$\psi' = \epsilon_d \tilde{\varphi}^{-1} \psi = \epsilon_d \tilde{\varphi}^* \psi$$

can be recovered from $\tilde{\varphi}$ and ψ . The desired isomorphism follows.

Now we prove the second part of the lemma. We assume that u is non-zero. If $\mathcal{Z}(u)$ is empty, then clearly both sides vanish. Now we assume that $\mathcal{Z}(u)$ is a (non-empty) relative divisor. Now note that the dimension of the intersection is as expected. Since both $\Gamma_{g'}$ and $\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}$ are local complete intersections in the ambient $\mathcal{N}_n \times \mathcal{N}_n$, Lemma B.2 shows that higher Tor all vanish. This proves the first equality in (4.14); the second one is proved similarly. \square

Recall from (3.5) that Δ is the image of the closed embedding $\Delta_{\mathcal{N}_{n-1}} : \mathcal{N}_{n-1} \rightarrow \mathcal{N}_{n-1,n} = \mathcal{N}_{n-1} \times \mathcal{N}_n$; cf. (3.4).

PROPOSITION 4.6. *Let $g' \in \mathrm{U}(\mathbb{V}_n)^\circ(F_0)$ be such that $1 - d \in O_F^\times$, and let $(g, u) = \mathfrak{r}(g') \in (\mathrm{U}(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)$. Assume further the vector $u \neq 0$ in \mathbb{V}_{n-1} . Then*

$$\Delta \cap (\mathrm{id} \times g')\Delta = \mathbb{L}\mathcal{N}_{n-1}^g \cap \mathcal{Z}(u)$$

as elements in $K'_0(\mathcal{N}_{n-1}^g \cap \mathcal{Z}(u))$. In particular, if g' is regular semisimple (hence so is (g, u) by Lemma 4.3(i)), then

$$\mathrm{Int}(g') = \mathrm{Int}(g, u).$$

Proof. Consider the following two cartesian squares, where we have applied Lemma 4.4 (4.13) to the middle term in the top row,

$$\begin{array}{ccccc} \mathcal{N}_{n-1}^g \cap \mathcal{Z}(u) & \longrightarrow & \Gamma_g \cap \pi_2^* \mathcal{Z}(u) & \longrightarrow & \Gamma_{g'} \\ \downarrow & \square & \downarrow & \square & \downarrow \\ \mathcal{N}_{n-1} & \xrightarrow{\Delta} & \mathcal{N}_{n-1} \times \mathcal{N}_{n-1} & \xrightarrow{(\delta, \delta)} & \mathcal{N}_n \times \mathcal{N}_n. \end{array}$$

We obtain equalities as elements in $K'_0(\mathcal{N}_{n-1}^g \cap \mathcal{Z}(u))$,

$$\begin{aligned} & \mathcal{O}_{\mathcal{N}_{n-1}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_n \times \mathcal{N}_n} \mathcal{O}_{\Gamma_{g'}} \\ &= \mathcal{O}_{\mathcal{N}_{n-1}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}} (\mathcal{O}_{\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_n \times \mathcal{N}_n} \mathcal{O}_{\Gamma_{g'}}) \\ &= \mathcal{O}_{\mathcal{N}_{n-1}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}} \mathcal{O}_{\Gamma_g \cap \pi_2^* \mathcal{Z}(u)} && \text{(Lemma 4.4 (4.14))} \\ &= (\mathcal{O}_{\mathcal{N}_{n-1}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}} \mathcal{O}_{\Gamma_g}) \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{N}_{n-1} \times \mathcal{N}_{n-1}} \mathcal{O}_{\pi_2^* \mathcal{Z}(u)} && \text{(Lemma 4.4 (4.14))} \\ &= \mathbb{L}\mathcal{N}_{n-1}^g \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{Z}(u)} && \text{(by (3.12))}. \end{aligned}$$

Similarly, we have two cartesian squares

$$\begin{array}{ccccc} \Delta \cap (1 \times g')\Delta & \longrightarrow & (1 \times g')\Delta & \longrightarrow & \Gamma_{g'} \\ \downarrow & \square & \downarrow & \square & \downarrow \\ \mathcal{N}_{n-1} & \xrightarrow{(\mathrm{id}_{\mathcal{N}_{n-1}}, \delta)} & \mathcal{N}_{n-1} \times \mathcal{N}_n & \xrightarrow{(\delta, \mathrm{id}_{\mathcal{N}_n})} & \mathcal{N}_n \times \mathcal{N}_n, \end{array}$$

with similar equalities as elements in $K'_0(\Delta \cap (1 \times g')\Delta) = K'_0(\mathcal{N}_{n-1}^g \cap \mathcal{Z}(u))$, which lead to

$$\Delta \cap_{\mathcal{N}_{n-1,n}} (1 \times g')\Delta = \mathcal{N}_{n-1} \cap_{\mathcal{N}_n \times \mathcal{N}_n} \Gamma_{g'}.$$

This completes the proof. \square

4.3. *Reduction of orbital integrals.* We use the Cayley map for S_n (a rational morphism):

$$(4.15) \quad \begin{aligned} \mathfrak{c} = \mathfrak{c}_n : \mathfrak{s}_n &\longrightarrow S_n \\ y &\longmapsto -\frac{1-y}{1+y}. \end{aligned}$$

Its inverse is

$$\mathfrak{c}^{-1}(\gamma) = \frac{1+\gamma}{1-\gamma}.$$

Similar to $U(\mathbb{V}_n)$, we now write $\gamma' \in S_n$ according to the decomposition $F^n = F^{n-1} \oplus Fu_0$:

$$\gamma' = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

LEMMA 4.7. *Let*

$$(4.16) \quad y' = \mathfrak{c}_n^{-1}(\gamma') = \begin{pmatrix} y & \tilde{b} \\ \tilde{c} & e \end{pmatrix} \in \mathfrak{s}_n \quad \text{and} \quad \gamma = \mathfrak{c}_{n-1}(y) \in S_{n-1}.$$

Then

$$(4.17) \quad \begin{cases} \gamma = a + (1-d)^{-1}bc, \\ \tilde{b} = 2(1-d)^{-1}(1-\gamma)^{-1}b, \\ \tilde{c} = -2c(1-d)^{-1}(1-\gamma)^{-1}, \\ \gamma\bar{b} = \epsilon_d b, \end{cases}$$

where we recall that $\epsilon_d = \frac{1-\bar{d}}{1-d}$; cf. (4.6).

Proof. Similar to the proof of 4.1, we obtain

$$\begin{cases} 1+a = (1-a)y + b\tilde{c}, \\ c = -cy - (1-d)\tilde{c}. \end{cases}$$

We then obtain

$$\tilde{c} = -(1-d)^{-1}c(1+y)$$

and

$$1+a + (1-d)^{-1}bc = (1-a - (1-d)^{-1}bc)y.$$

It follows that

$$\gamma = \mathfrak{c}_{n-1}(y) = a + (1-d)^{-1}bc.$$

The remaining assertions follow similarly. \square

We now define a rational map by the formulas in Lemma 4.7:

$$(4.18) \quad \begin{aligned} \mathfrak{r} : S_n &\longrightarrow S_{n-1} \times V'_{n-1} \\ \gamma' &\longmapsto \left(\gamma, \left(\frac{\tilde{b}}{\sqrt{\epsilon}}, \frac{\tilde{c}}{\sqrt{\epsilon}} \cdot (1-y^2)^{-1} \right) \right). \end{aligned}$$

From (4.17), and the fact that $y \in \mathfrak{s}_{n-1} \implies y^2 \in M_{n,n}$, it follows that the last component of $\mathfrak{r}(\gamma')$ indeed lies in $V'_{n-1} = F_0^{n-1} \times (F_0^{n-1})^*$. We also define a variant:

$$(4.19) \quad \begin{aligned} \mathfrak{r}^\sharp : S_n &\longrightarrow S_{n-1} \times V'_{n-1} \\ \gamma' &\longmapsto \left(\gamma, \left(\frac{\tilde{b}}{\sqrt{\epsilon}}, \frac{\tilde{c}}{\sqrt{\epsilon}} \right) \right). \end{aligned}$$

Following the notation in Lemma 4.7, let S_n° be the open sub-variety of S_n defined by

$$1 - d \neq 0 \quad \text{and} \quad \det(1 - \gamma') \neq 0.$$

Let $(S_{n-1} \times V'_{n-1} \times \mathfrak{s}_1)^\circ$ be the open sub-variety of $S_{n-1} \times V'_{n-1} \times \mathfrak{s}_1$ defined by

$$\det(1 - \gamma) \neq 0 \quad \text{and} \quad \det(1 + y') \neq 0.$$

LEMMA 4.8. *The map \mathfrak{r} together with $e \in \mathfrak{s}_1$ (cf. (4.16)) induce an isomorphism (between two open sub-varieties), equivariant under the action of GL_{n-1} ,*

$$\begin{aligned} \tilde{\mathfrak{r}} = (\mathfrak{r}, e) : S_n^\circ &\xrightarrow{\sim} (S_{n-1} \times V'_{n-1} \times \mathfrak{s}_1)^\circ \\ \gamma' &\longmapsto (\mathfrak{r}(\gamma'), e). \end{aligned}$$

The same holds if we replace \mathfrak{r} by \mathfrak{r}^\sharp .

Proof. The proof of Lemma 4.2 still works, and we omit the detail. \square

We may apply the same construction to $\xi\gamma'$ for $\xi \in F^1 = \ker(\mathrm{Nm} : F^\times \rightarrow F_0^\times)$:

$$(4.20) \quad \begin{aligned} \mathfrak{r}_\xi : S_n &\longrightarrow S_{n-1} \times V'_{n-1} \\ \gamma' &\longmapsto \mathfrak{r}(\xi\gamma'). \end{aligned}$$

We define \mathfrak{r}_ξ^\sharp similar to (4.19).

LEMMA 4.9.

- (i) *An element $\gamma' \in S_n^\circ(F_0)$ is regular semisimple if and only if $\mathfrak{r}_\xi(\gamma') \in (S_{n-1} \times V'_{n-1})(F_0)$ is regular semisimple.*
- (ii) *Let $\gamma' \in S_n^\circ(F_0)_{\mathrm{srs}}$. Then, for all but finitely many $\xi \in F^1$, we have $\xi\gamma' \in S_n^\circ(F_0)$ and $\mathfrak{r}_\xi(\gamma') \in (S_{n-1} \times V'_{n-1})(F_0)_{\mathrm{srs}}$.*
- (iii) *Let $(\gamma, u') \in (S_{n-1} \times V'_{n-1})(F_0)_{\mathrm{srs}}$. Then, for all but finitely many $e \in \mathfrak{s}_1$, the element (γ, u', e) lies in $(S_{n-1} \times V'_{n-1} \times \mathfrak{s}_1)^\circ(F_0)$, and $\tilde{\mathfrak{r}}^{-1}(\gamma, u', e) \in S_n^\circ(F_0)$ is strongly regular semisimple.*

Proof. The same argument as the proof of Lemma 4.3 works here. Hence we omit the details. \square

LEMMA 4.10. *If $\gamma' \in S_n(F_0)_{\text{srs}}$ and $g' \in U(\mathbb{V}_n)(F_0)_{\text{srs}}$ match, then the following pairs also match (whenever they are well defined for $\xi \in F^1$ under the rational maps):*

- $\tau_\xi^\natural(\gamma') \in (S_{n-1} \times V'_{n-1})(F_0)_{\text{srs}}$ and $\tau_\xi^\natural(g') \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$;
- $\tau_\xi(\gamma') \in (S_{n-1} \times V'_{n-1})(F_0)_{\text{srs}}$ and $\tau_\xi(g') \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$.

Proof. We retain the notation in [Lemmas 4.3](#) and [4.7](#). We may assume $\xi=1$. By choosing a basis of \mathbb{V}_{n-1} and of $\mathbb{V}_n = \mathbb{V}_{n-1} \oplus F u_0$, we write $g' \in M_{n,n}(F)$ in matrix form; cf. the discussion on matching orbits in [Section 2.2](#). Since the inverse Cayley map (cf. [\(4.1\)](#), [\(4.15\)](#)) preserve the matching conditions, $\mathfrak{c}^{-1}(\gamma')$ and $\mathfrak{c}^{-1}(g')$ also match. It follows that the two elements denoted by e in their lower right corner are equal. Moreover, there exists $k \in \text{GL}_{n-1}(F)$ such that

$$\begin{pmatrix} x & \tilde{u} \\ -\tilde{u}^* & e \end{pmatrix} = \begin{pmatrix} k^{-1} & \\ & 1 \end{pmatrix} \begin{pmatrix} y & \tilde{b} \\ \tilde{c} & e \end{pmatrix} \begin{pmatrix} k & \\ & 1 \end{pmatrix},$$

or equivalently,

$$x = k^{-1} y k, \quad \tilde{u} = k^{-1} \tilde{b}, \quad -\tilde{u}^* = \tilde{c} k.$$

It follows that

$$g = \mathfrak{c}(x) = k^{-1} \mathfrak{c}(y) k = k^{-1} \gamma k,$$

and hence

$$\begin{pmatrix} g & \frac{\tilde{u}}{\sqrt{\epsilon}} \\ \left(\frac{\tilde{u}}{\sqrt{\epsilon}}\right)^* & e \end{pmatrix} = \begin{pmatrix} k^{-1} & \\ & 1 \end{pmatrix} \begin{pmatrix} \gamma & \frac{\tilde{b}}{\sqrt{\epsilon}} \\ \frac{\tilde{c}}{\sqrt{\epsilon}} & e \end{pmatrix} \begin{pmatrix} k & \\ & 1 \end{pmatrix}.$$

This proves the first part.

By [Lemma 4.1 \(4.5\)](#), $\tilde{u} = 2(1-d)^{-1}(1-g)^{-1}u$, we obtain

$$u = 2^{-1}(1-d)(1-g)\tilde{u} = (1-d)(1+x)^{-1}\tilde{u}.$$

We compute the invariants of (g, u) . For $0 \leq i \leq n-1$,

$$\begin{aligned} u^* g^i u &= (1-d)(1-\bar{d})\tilde{u}^*(1+x^*)^{-1}g^i(1+x)^{-1}\tilde{u} \\ &= (1-d)(1-\bar{d})\tilde{u}^*(1-x^2)^{-1}g^i\tilde{u}, \end{aligned}$$

where we have used that g and x commute, and $x^* = -x$. In terms of the invariants of $(\gamma, \tilde{b}, \tilde{c})$, this last quantity is equal to

$$\begin{aligned} u^* g^i u &= (1-d)(1-\bar{d})\tilde{u}^*(1-x^2)^{-1}g^i\tilde{u} \\ &= -(1-d)(1-\bar{d})\tilde{c} k (1-x^2)^{-1}g^i k^{-1}\tilde{b} \\ &= -(1-d)(1-\bar{d})\tilde{c} (1-y^2)^{-1}\gamma^i \tilde{b}. \end{aligned}$$

Obviously g and γ have the same characteristic polynomial. It follows that $\mathfrak{r}(g') = (g, \frac{u}{\sqrt{\epsilon(1-d)}})$ has the same set of invariants as

$$\left(\gamma, \left(\sqrt{\epsilon}^{-1} \tilde{b}, \sqrt{\epsilon}^{-1} \tilde{c} \cdot (1 - y^2)^{-1} \right) \right) = \mathfrak{r}(\gamma').$$

This completes the proof of the second part. \square

LEMMA 4.11. *Let $\gamma' \in S_n(F_0)_{\text{rs}}$ and $g' \in \text{U}(\mathbb{V}_n)(F_0)_{\text{rs}}$ be a matching pair, and let $\xi \in F^1$. Assume that*

$$(4.21) \quad 1 - \xi d \in O_F^\times \quad \text{and} \quad \det(1 - \xi \gamma') \in O_F^\times.$$

Then

$$\begin{aligned} \text{Orb}(\gamma', \mathbf{1}_{S_n(O_{F_0})}, s) &= \text{Orb} \left(\mathfrak{r}_\xi^\natural(\gamma'), \mathbf{1}_{(S_{n-1} \times V'_{n-1})(O_{F_0})}, s \right) \\ &= \text{Orb} \left(\mathfrak{r}_\xi(\gamma'), \mathbf{1}_{(S_{n-1} \times V'_{n-1})(O_{F_0})}, s \right). \end{aligned}$$

Proof. It suffices to prove the assertions for $\xi = 1$. We also consider the orbital integral on the Lie algebra \mathfrak{s}_n . Since $\det(1 - \gamma') \in O_F^\times$ by assumption (4.21), and since the Cayley map is equivariant under the $\text{GL}_{n-1}(F_0)$,

$$h \cdot \mathfrak{c}^{-1}(\gamma') \in \mathfrak{s}_n(O_{F_0}) \quad \text{if and only if} \quad h \cdot \gamma' \in S_n(O_{F_0}).$$

It follows that

$$\text{Orb}(\mathfrak{c}^{-1}(\gamma'), \mathbf{1}_{\mathfrak{s}_n(O_{F_0})}, s) = \text{Orb}(\gamma', \mathbf{1}_{S_n(O_{F_0})}, s).$$

Similarly, by $\det(1 + y) = (1 - d)^{-1} \det(1 - \gamma')$ and (4.21), we know that $\det(1 + y) \in O_F^\times$. Therefore,

$$h^{-1}yh \in \mathfrak{s}_{n-1}(O_{F_0}) \quad \text{if and only if} \quad h^{-1}\gamma h \in S_{n-1}(O_{F_0}).$$

It follows that (note that d and e are now in O_F and $\mathfrak{s}_1(O_{F_0})$ respectively)

$$\text{Orb}(\mathfrak{c}^{-1}(\gamma'), \mathbf{1}_{\mathfrak{s}_n(O_{F_0})}, s) = \text{Orb} \left(\mathfrak{r}^\natural(\gamma'), \mathbf{1}_{(S_{n-1} \times V'_{n-1})(O_{F_0})}, s \right).$$

This proves the first equality.

We now simply denote

$$\tilde{c}' := \tilde{c} \cdot (1 - y^2)^{-1}$$

so that

$$\mathfrak{r}(\gamma') = \left(\gamma, \left(\tilde{b}/\sqrt{\epsilon}, \tilde{c}'/\sqrt{\epsilon} \right) \right).$$

Note now that $\det(1 - y^2) = \text{Nm} \det(1 + y) \in O_{F_0}^\times$ under our assumption. Therefore, when $h^{-1}\gamma h \in S_{n-1}(O_{F_0})$, we have

$$\frac{\tilde{c}'}{\sqrt{\epsilon}} h \in O_{F_0}^n \quad \text{if and only if} \quad \frac{\tilde{c}}{\sqrt{\epsilon}} h \in O_{F_0}^n.$$

This immediately implies the second equality. \square

4.4. Relation between the two versions of AFL.

PROPOSITION 4.12. *Assume that $q \geq n$, where q denotes the cardinality of the residue field of O_{F_0} . Then*

- (i) *in Conjecture 3.8, part (a) for \mathbb{V}_n is equivalent to part (b) for \mathbb{V}_{n-1} .*
- (ii) *in Conjecture 3.8, part (a) for S_{n-1} implies part (b) for \mathbb{V}_{n-1} and $(g, u) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$, where the norm of u is a unit.*

Remark 4.13. Similar results hold for Conjecture 3.2 for regular semisimple elements.

Proof. For part (i), let $g' \in U(\mathbb{V}_n)(F_0)_{\text{srs}}$. We may assume that $d \in O_F$ and the characteristic polynomial of both g' and g have integral coefficients (otherwise both sides of part (a) vanish). Since $q + 1 > n$, there exists $\xi \in F^1$ such that $\det(1 - \xi g') \in O_F^\times$ is a unit (looking at the reduction of the characteristic polynomial modulo the uniformizer ϖ_F of O_F). Since both sides of part (a) for \mathbb{V}_n are invariant under the substitution $g' \mapsto \xi g'$, we may just assume that g' has the property that $d \in O_F$ and $\det(1 - g') \in O_F^\times$. From the third equality in (4.5) and the integrality of $\det(1 - g)$, it follows that $1 - d \in O_F^\times$. Now $g' \in U(\mathbb{V}_n)^\circ(F_0)_{\text{srs}}$, so that we may apply the map τ . By Lemma 4.3, we may adjust $\xi \in F^1$ within the same residue class mod ϖ_F such that the image $\tau(g') = (g, u)$ lies in $(U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$. Now, by Proposition 4.6,

$$\text{Int}(g') = \text{Int}(g, u).$$

Now we consider the orbital integral. By Lemma 4.11,

$$\partial\text{Orb}(\gamma', 1_{S_n(O_{F_0})}) = \partial\text{Orb}(\tau(\gamma'), 1_{(S_{n-1} \times V'_{n-1})(O_{F_0})}).$$

Here we refer to [37, Lem. 11.9] for the comparison of the transfer factors. By Lemma 4.10, $\tau(\gamma') \in (S_{n-1} \times V'_{n-1})(F_0)_{\text{srs}}$ and $\tau(g') \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$ match. This shows that part (b) for \mathbb{V}_{n-1} implies part (a) for \mathbb{V}_n .

For the inverse direction, we start from $(g, u) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$. Again it suffices to prove part (b) when the invariants of (g, u) are all integers. By multiplying a suitable $\xi \in F^1$, we may assume $\det(1 - g) \in O_F^\times$. Then $\det(1 + x) \in O_F^\times$. By Lemma 4.3 part (iii), there exists $e \in \mathfrak{u}(1)(O_{F_0})$ such that $\det(1 + x') \in O_F^\times$, (g, u, e) lies in $(U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1} \times \mathfrak{u}(1))^\circ$ and $g' = \tilde{\tau}^{-1}(g, u, e) \in U(\mathbb{V}_n)_{\text{srs}}^\circ$. Then $\det(1 - g') \in O_F^\times$ and hence $(1 - d) \in O_F^\times$ (by the third equality in (4.5)), and we may therefore apply Proposition 4.6. A similar procedure proves the desired identity between orbital integrals. This shows that part (a) for \mathbb{V}_n implies part (b) for \mathbb{V}_{n-1} .

For part (ii), we note that for $g \in U(\mathbb{V}_{n-1})(F_0)_{\text{srs}}$, the pair $(g, u_0) \in (U(\mathbb{V}_{n-1}) \times \mathbb{V}_{n-1})(F_0)_{\text{srs}}$, and it is easy to see

$$\text{Int}(g) = \text{Int}(g, u_0).$$

One can show that the orbital integrals are equal easily; we leave the detail to the reader. \square

5. Local constancy of intersection numbers

This section is not used until [Section 15](#).

5.1. Local constancy of the function $\text{Int}(g, \cdot)$. We recall the Bruhat–Tits stratification of the underlying reduced scheme $\mathcal{N}_{n,\text{red}}$ of \mathcal{N}_n , following the work of Vollaard–Wedhorn [42] (see also [24, §2.2]). The scheme $\mathcal{N}_{n,\text{red}}$ admits a stratification by Deligne–Lusztig varieties of dimensions $0, 1, \dots, \lfloor \frac{n-1}{2} \rfloor$, attached to unitary groups in an odd number of variables and to Coxeter elements, with strata parametrized by the vertices of the Bruhat–Tits complex of the special unitary group for the non-split n -dimensional F/F_0 -hermitian space \mathbb{V}_n . The vertices of the Bruhat–Tits complex are bijective to the vertex lattices in \mathbb{V}_n , where an O_F -lattice (of full rank) $\Lambda \subset \mathbb{V}_n$ is called a vertex lattice if $\Lambda \subset \Lambda^\vee \subset \varpi^{-1}\Lambda$. The parametrization of the strata by vertex lattices in \mathbb{V}_n is compatible with the action of the group $U(\mathbb{V}_n)$ on $\mathcal{N}_{n,\text{red}}$ (cf. (3.6)) and on \mathbb{V}_n . The type of a vertex lattice Λ is by definition the integer $t(\Lambda) := \dim_k \Lambda^\vee / \Lambda$. Denote by $\mathcal{V}(\Lambda)$ the corresponding (generalized) Deligne–Lusztig variety; it is smooth projective of dimension $\frac{t(\Lambda)-1}{2}$; cf. loc. cit. Note that the type $t(\Lambda)$ is necessarily odd because the F/F_0 -hermitian space \mathbb{V}_n is non-split.

LEMMA 5.1. *Let $n \geq 3$, and let $\Lambda \subset \mathbb{V}_n$ be a vertex lattice of maximal type (i.e., type $2[(n-1)/2] + 1$). Let $C \in \text{Ch}_{1,\mathcal{V}(\Lambda)}(\mathcal{N}_{n,\text{red}})$. Then the function*

$$\begin{aligned} \text{Int}_C : \mathbb{V}_n &\longrightarrow \mathbb{Q} \\ u &\longmapsto \chi(\mathcal{N}_n, C \overset{\mathbb{L}}{\cap} \mathcal{Z}(u)) \end{aligned}$$

is locally constant and compactly supported. Here, even though the function is only defined for $u \neq 0$, the local constancy around $u = 0$ is to be interpreted as that the function takes a constant value for all $u \neq 0$ in a neighborhood of $0 \in \mathbb{V}$.

Proof. The proof is essentially the same as [26, Cor. 6.2.2], noting that Int_C is a linear combination of $\text{Int}_{\mathcal{V}(\Lambda')}$ for vertex lattices $\Lambda' \supset \Lambda$ of type 3. \square

PROPOSITION 5.2. *Fix a regular semisimple element $g \in U(\mathbb{V}_n)$. Let*

$$\mathbb{V}_{n,g} := \{u \in \mathbb{V}_n \mid (g, u) \text{ is not regular semisimple}\}.$$

Then the function

$$\begin{aligned} \text{Int}(g, \cdot) : \mathbb{V}_n \setminus \mathbb{V}_{n,g} &\longrightarrow \mathbb{Q} \\ u &\longmapsto \text{Int}(g, u) = \chi(\mathcal{N}_n, {}^L\mathcal{N}_n^g \overset{\mathbb{L}}{\cap} \mathcal{Z}(u)) \end{aligned}$$

is locally constant.

Proof. We first observe that, when $u \in \mathbb{V}_n \setminus \mathbb{V}_{n,g}$, the formal scheme $\mathcal{N}_n^g \cap \mathcal{Z}(u) = \mathcal{N}_n^g \cap \mathcal{Z}(u) \cap \mathcal{Z}(gu) \cap \cdots \cap \mathcal{Z}(g^{n-1}u)$ is a noetherian scheme. It follows that, since g is fixed, the scheme $\mathcal{N}_n^g \cap \mathcal{Z}(u)$ depends only on the lattice spanned by $g^i u$, $i = 0, 1, \dots, n-1$. For any $u \in \mathbb{V}_n \setminus \mathbb{V}_{n,g}$, there is a small open neighborhood \mathcal{U} of u in $\mathbb{V}_n \setminus \mathbb{V}_{n,g}$ such that this lattice does not vary when varying $u \in \mathcal{U}$. Let us fix such an open neighborhood \mathcal{U} . Therefore, without changing the notation, we may and will simply work with the restriction of the relevant coherent sheaves to a suitable fixed noetherian open formal subscheme \mathcal{N}_n° of \mathcal{N}_n .

In the codimension filtration (B.1), the classes of \mathcal{O}_{Γ_g} and $\mathcal{O}_{\Delta_{\mathcal{N}_n}}$ belong to $F^{n-1}K_0^{\Gamma_g}(\mathcal{N}_n \times \mathcal{N}_n)$ and $F^{n-1}K_0^{\Delta_{\mathcal{N}_n}}(\mathcal{N}_n \times \mathcal{N}_n)$ respectively. Therefore, by (B.3) the class ${}^{\mathbb{L}}\mathcal{N}_n^g$ (cf. (3.12)) lies in the filtration $F^{2n-2}K_0^{\mathcal{N}_n^g}(\mathcal{N}_n \times \mathcal{N}_n) = F^{n-1}K_0^{\mathcal{N}_n^g}(\mathcal{N}_n)$. Since $\mathcal{Z}(u)$ is a Cartier divisor, the Euler–Poincaré characteristic $\chi(\mathcal{N}_n, Z \cap \mathcal{Z}(u))$ vanishes if Z is a (noetherian) zero dimensional subscheme of \mathcal{N}_n . Therefore, it suffices to consider ${}^{\mathbb{L}}\mathcal{N}_n^g \in \text{Gr}^{n-1}K_0^{\mathcal{N}_n^g}(\mathcal{N}_n)$; see (B.2) for the definition of the graded groups Gr^*K_0 . We may represent (the restriction to \mathcal{N}_n° of) ${}^{\mathbb{L}}\mathcal{N}_n^g$ by a finite sum $\sum_C \text{mult}_C \cdot [\mathcal{O}_C]$ where $\text{mult}_C \in \mathbb{Q}$ and all C are one dimensional, formally reduced (i.e., the sheaf \mathcal{O}_C has trivial nilradical), irreducible, and closed formal subschemes of \mathcal{N}_n^g . Fix such a C . It suffices to show that the following function is locally constant:

$$\begin{aligned} \text{Int}_C : \mathcal{U} &\longrightarrow \mathbb{Q} \\ u &\longmapsto \chi(\mathcal{N}_n, C \cap \mathcal{Z}(u)). \end{aligned}$$

There are the following (mutually exclusive) two cases for C :

- C is a closed formal subscheme of $\mathcal{N}_{n,\text{red}}$;
- C is not a closed formal subscheme of $\mathcal{N}_{n,\text{red}}$.

For the first case, we can assume that $C \subset \mathcal{V}(\Lambda)$ for some vertex lattice Λ of maximal type. Then we have proved an even stronger statement in Lemma 5.1.

Now we consider the second case. We let \widetilde{C} be the normalization of C and let $\pi : \widetilde{C} \rightarrow C$ be the normalization morphism. (This is a finite morphism by the excellence of C .) It suffices to show the local constancy of $u \in \mathcal{U} \mapsto \chi(\mathcal{N}_n, \pi_* \mathcal{O}_{\widetilde{C}} \otimes^{\mathbb{L}} \mathcal{O}_{\mathcal{Z}(u)})$. Note that $\mathcal{Z}(u)$ is a Cartier divisor on \mathcal{N}_n . Then $\widetilde{C} \times_{\mathcal{N}_n} \mathcal{Z}(u)$ has expected dimension. (Otherwise $C \subset \mathcal{Z}(u)$; then $C \subset C \cap \mathcal{Z}(u) \subset \mathcal{N}_n^g \cap \mathcal{Z}(u)$, and hence C is a closed subscheme of $\mathcal{N}_{n,\text{red}}$.) Therefore we have equalities in $K'_0(\mathcal{N}_n^g \cap \mathcal{Z}(u))$:

$$\begin{aligned} \pi_* \mathcal{O}_{\widetilde{C}} \otimes^{\mathbb{L}}_{\mathcal{O}_{\mathcal{N}_n}} \mathcal{O}_{\mathcal{Z}(u)} &= \pi_* \mathcal{O}_{\widetilde{C}} \otimes_{\mathcal{O}_{\mathcal{N}_n}} \mathcal{O}_{\mathcal{Z}(u)} \\ &= \pi_* \mathcal{O}_{\widetilde{C}} \otimes_{\mathcal{O}_{\mathcal{N}_n^g}} (\mathcal{O}_{\mathcal{N}_n^g} \otimes_{\mathcal{O}_{\mathcal{N}_n}} \mathcal{O}_{\mathcal{Z}(u)}) \\ &= \pi_* \mathcal{O}_{\widetilde{C}} \otimes_{\mathcal{O}_{\mathcal{N}_n^g}} \mathcal{O}_{\mathcal{N}_n^g \cap \mathcal{Z}(u)}. \end{aligned}$$

It follows that

$$(5.1) \quad \chi(\mathcal{N}_n, \pi_* \mathcal{O}_{\widetilde{C}} \overset{\mathbb{L}}{\otimes} \mathcal{O}_{\mathcal{Z}(u)}) = \chi(\mathcal{N}_n, \pi_* \mathcal{O}_{\widetilde{C}} \otimes_{\mathcal{O}_{\mathcal{N}_n^g}} \mathcal{O}_{\mathcal{N}_n^g \cap \mathcal{Z}(u)}).$$

We have seen in the beginning of the proof that the sheaf $\mathcal{O}_{\mathcal{N}_n^g \cap \mathcal{Z}(u)}$ does not change when varying u in \mathcal{U} . Therefore the Euler–Poincaré characteristic (5.1) is a constant for $u \in \mathcal{U}$. \square

Remark 5.3. We could avoid (B.3) in the proof of [Proposition 5.2](#) as follows. Now we cannot conclude that $\mathbb{L}\mathcal{N}_{n,\gamma}^g \in F^{n-1}K_0^{\mathcal{N}_{n,\gamma}^g}(\mathcal{N}_n)$. Therefore, in the two cases of the proof, the closed formal subscheme C may have dimension higher than one. For the first case, [Lemma 5.1](#) holds for any $C \in \text{Ch}_{i,\mathcal{V}(\Lambda)}(\mathcal{N}_n, \text{red})$ of arbitrary dimension i (see [26, §6.4]). For the second case, we still have (5.1), and therefore the proof above still works.

5.2. Local constancy of the function $\text{Int}(\cdot, \cdot)$.

LEMMA 5.4. *Fix $(g_0, u_0, e_0) \in (\text{U}(\mathbb{V}_n) \times \mathbb{V}_n \times \mathfrak{u}(1))^\circ$ such that*

$$g' = \tilde{\tau}^{-1}(g_0, u_0, e_0) \in \text{U}(\mathbb{V}_{n+1})_{\text{srs}};$$

cf. [Lemma 4.3](#) for the notation. Then the map (defined on some open subsets of F_0 -varieties)

$$\begin{aligned} \text{char}(g_0, \cdot, \cdot) : \mathbb{V}_n \times \mathfrak{u}(1) &\longrightarrow \text{U}(\mathbb{V}_{n+1}) // \text{U}(\mathbb{V}_{n+1}) \\ (u, e) &\longmapsto \text{char poly}(\tilde{\tau}^{-1}(g_0, u, e)) \end{aligned}$$

is submersive (i.e., the induced map on tangent spaces is surjective) at (u_0, e_0) .

Here $\text{U}(\mathbb{V}_{n+1}) // \text{U}(\mathbb{V}_{n+1})$ denotes the categorical quotient (with respect to the conjugation action) and char poly denotes the characteristic polynomial.

Proof. The question is local on the source. Tracing the definition back to (4.9) and [Lemma 4.2](#), we may reduce the question to the Lie algebra version: for a fixed $(\begin{smallmatrix} x_0 & u_0 \\ -u_0^* & e_0 \end{smallmatrix}) \in \mathfrak{u}(\mathbb{V}_{n+1})_{\text{srs}}$, the map

$$\mathbb{V}_n \times \mathfrak{u}(1) \longrightarrow \mathfrak{u}(\mathbb{V}_{n+1}) // \text{U}(\mathbb{V}_{n+1})$$

sending (u, e) to the characteristic polynomial of $(\begin{smallmatrix} x_0 & u \\ -u^* & e \end{smallmatrix}) \in \mathfrak{u}(\mathbb{V}_{n+1})$ is submersive at (u_0, e_0) .

Note that a complete set of generators of invariants relative to the $\text{U}(\mathbb{V}_n)$ -action on $\mathfrak{u}(\mathbb{V}_{n+1})$ is given by

$$\text{char poly}(x), \quad e, \quad u^* x^j u, \quad 0 \leq j \leq n-1,$$

where $x' = (\begin{smallmatrix} x & u \\ -u^* & e \end{smallmatrix}) \in \mathfrak{u}(\mathbb{V}_{n+1})$; cf. [46]. It is easy to see that an equivalent set is

$$\text{char poly}(x), \quad \text{char poly}(x').$$

Therefore, it suffices to show the analogous map

$$\mathbb{V}_n \longrightarrow \prod_{i=0}^{n-1} F^{(-1)^j}$$

sending $u \in \mathbb{V}_n$ to the invariants

$$u^* x^j u, \quad 0 \leq j \leq n-1,$$

is submersive at u_0 . Here $F^{(-1)^j}$ is the $(-1)^j$ -eigenspace of F under the Galois conjugation. Now the assertion follows from the regular semi-simplicity of x_0 , which reduces the question to the case $n = 1$, but for the product of field extensions of F . This is routine and we omit the detail. \square

THEOREM 5.5. *The function*

$$\begin{aligned} \text{Int}(\cdot, \cdot) : (\text{U}(\mathbb{V}_n) \times \mathbb{V}_n)(F_0)_{\text{srs}} &\longrightarrow \mathbb{Q} \\ (g, u) &\longmapsto \text{Int}(g, u) \end{aligned}$$

is locally constant. Its support is compact modulo the action of $\text{U}(\mathbb{V}_n)(F_0)$.

Remark 5.6. See the forthcoming work of Mihatsch [32] for a different proof, which also yields the local constancy on the regular semisimple locus.

Proof. We may assume that the invariants of (g, u) are all integers. We now fix such a pair (g, u) , and we want to show the local constancy near (g, u) .

First, by the argument in the proof of part (i) of [Proposition 4.12](#), there exists $g' = \tilde{\tau}^{-1}(g, u, e) \in \text{U}(\mathbb{V}_{n+1})^\circ(F_0)_{\text{srs}}$ such that

$$(5.2) \quad \text{Int}(g') = \text{Int}(g, u).$$

In fact, by the same argument the equality holds if we replace (g, u, e) by any element $(g^\sharp, u^\sharp, e^\sharp)$ near it, and g' by the respective image g'^\sharp under the map $\tilde{\tau}^{-1}$.

On the other hand, we may write

$$\text{Int}(g') = \text{Int}(g', u'_0),$$

where $u'_0 \in \mathbb{V}_{n+1}$ is the fixed unit normed vector that induces the embedding $\mathcal{N}_n \hookrightarrow \mathcal{N}_{n+1}$. We now apply [Proposition 5.2](#) to (g', u'_0) :

$$\text{Int}(g', u'_0) = \text{Int}(g', u'),$$

where $u' \in \mathbb{V}_{n+1}$ is close to u'_0 . In particular, the equality holds for $u' = hu'_0$ for $h \in \text{U}(\mathbb{V}_{n+1})$ in a small neighborhood of 1. By the invariance under $\text{U}(\mathbb{V}_{n+1})$, for $u' = hu'_0$ we have

$$\text{Int}(g', u') = \text{Int}(h^{-1}g'h, u'_0).$$

It follows that $\text{Int}(g', u'_0) = \text{Int}(h^{-1}g'h, u'_0)$ and hence

$$(5.3) \quad \text{Int}(g') = \text{Int}(h^{-1}g'h)$$

for $h \in U(\mathbb{V}_{n+1})(F_0)$ in a small neighborhood of 1. This shows that, as a function on the quotient $[U(\mathbb{V}_{n+1})/U(\mathbb{V}_n)](F_0)$, $\text{Int}(g')$ is constant on those elements near g' and having the same characteristic polynomial (as g').

Now we claim that the desired local constancy near $(g, u) \in (U(\mathbb{V}_n) \times \mathbb{V}_n)(F_0)_{\text{srs}}$ follows from the following two properties:

- (1) the local constancy in the u -variable (for a fixed g), by [Proposition 5.2](#);
- (2) the invariance [\(5.3\)](#) under conjugation by elements h near 1 $\in U(\mathbb{V}_{n+1})$.

To show the claim, let g^\sharp be an element in a small neighborhood of g' . By [Lemma 5.4](#), there exists a neighborhood $\Omega \subset \mathbb{V}_n \times \mathfrak{u}(1)$ of (u, e) such that g^\sharp is conjugate (by an element $h \in U(\mathbb{V}_{n+1})(F_0)$ near 1) to $\tilde{\tau}^{-1}(g, u^\sharp, e^\sharp)$ for some $(u^\sharp, e^\sharp) \in \Omega$. By the invariance [\(5.3\)](#), we have

$$\text{Int}(g^\sharp) = \text{Int}(\tilde{\tau}^{-1}(g, u^\sharp, e^\sharp)).$$

By [\(5.2\)](#) (and the remark following it),

$$\text{Int}(\tilde{\tau}^{-1}(g, u^\sharp, e^\sharp)) = \text{Int}(g, u^\sharp).$$

By [Proposition 5.2](#) for the local constancy in u ,

$$\text{Int}(g, u^\sharp) = \text{Int}(g, u).$$

Again by [\(5.2\)](#), $\text{Int}(g, u) = \text{Int}(g')$, we obtain $\text{Int}(g^\sharp) = \text{Int}(g')$. The desired local constancy of $\text{Int}(g, u)$ follows from [\(5.2\)](#) (and the remark following it).

To show the compactness of the support modulo $U(\mathbb{V}_n)(F_0)$, it suffices to show the *claim*: the support is contained in the union of compact subsets

$$K_\Lambda \times \Lambda \subset (U(\mathbb{V}_n) \times \mathbb{V}_n)(F_0),$$

where Λ runs over all vertex lattices, and K_Λ is the stabilizer of Λ . Then the desired compactness follows from the fact that the group $U(\mathbb{V}_n)(F_0)$ acts transitively on the set of vertex lattices of any given type t (and there are only finitely many possible types $t = 1, 3, \dots, 2[(n-1)/2] + 1$). Now we show the claim. If $\text{Int}(g, u) \neq 0$, then there exists a point $x \in \mathcal{N}_n(\bar{k})$ lying on $\mathcal{Z}(u)$ and \mathcal{N}_n^g . Let $\mathcal{V}(\Lambda)$ for some vertex lattice Λ be the smallest stratum containing the point x . Then $g\Lambda = \Lambda$ (otherwise the intersection $g\mathcal{V}(\Lambda) \cap \mathcal{V}(\Lambda)$ is non-empty and is a strictly smaller stratum), and $u \in \Lambda$ by [\[24, Prop. 4.1\]](#). Therefore $(g, u) \in K_\Lambda \times \Lambda$ as desired. \square

Part 2. Global theory

6. Shimura varieties and their integral models

In this section we recall the construction of the integral models of unitary Shimura varieties, following [\[5\]](#), [\[40\]](#), [\[39\]](#). In fact, rather than the full strength of loc. cit., we only need a regular integral model away from a suitable finite

set of primes: the key is to keep those primes where the relevant hermitian space locally contains a self-dual lattice.

6.1. *Shimura varieties.* Let F be a CM number field with maximal totally real subfield F_0 and nontrivial F/F_0 -automorphism $a \mapsto \bar{a}$. Let n be a positive integer. A *generalized CM type of rank n* is a function $r : \text{Hom}_{\mathbb{Q}}(F, \overline{\mathbb{Q}}) \rightarrow \mathbb{Z}_{\geq 0}$, denoted $\varphi \mapsto r_{\varphi}$, such that

$$(6.1) \quad r_{\varphi} + r_{\bar{\varphi}} = n \quad \text{for all } \varphi.$$

Here $\bar{\varphi}$ denotes the pre-composition of φ by the nontrivial F/F_0 -automorphism. When $n = 1$, a generalized CM type is the same as a usual CM type (i.e., a half-system Φ of complex embeddings of F), via $\Phi = \{\varphi \in \text{Hom}_{\mathbb{Q}}(F, \overline{\mathbb{Q}}) \mid r_{\varphi} = 1\}$.

Let $(V, (\cdot, \cdot))$ be an F/F_0 -hermitian vector space of dimension n . Fix a CM type Φ of F . Then the signatures of V at the archimedean places determine a generalized CM type r of rank n (and vice versa), by the following recipe:

$$(6.2) \quad \text{sig}V_{\varphi} = (r_{\varphi}, r_{\bar{\varphi}}), \quad \varphi \in \Phi, \quad V_{\varphi} := V \otimes_{F, \varphi} \mathbb{C}.$$

Let $G^{\mathbb{Q}}$ be the group of unitary similitudes of $(V, (\cdot, \cdot))$,

$$G^{\mathbb{Q}} := \{g \in \text{Res}_{F_0/\mathbb{Q}} \text{GU}(V) \mid c(g) \in \mathbb{G}_m\},$$

considered as an algebraic group over \mathbb{Q} (with similitude factor in \mathbb{G}_m), where c denotes the similitude map.

Given Φ , r and V , we define a Shimura datum $(G^{\mathbb{Q}}, \{h_{G^{\mathbb{Q}}}\})$ as follows (cf. [39, §2.2]). For each $\varphi \in \Phi$, choose a \mathbb{C} -basis of V_{φ} with respect to which the matrix of (\cdot, \cdot) is given by

$$(6.3) \quad \text{diag}(1_{r_{\varphi}}, -1_{r_{\bar{\varphi}}}).$$

Then $\{h_{G^{\mathbb{Q}}}\}$ is the $G^{\mathbb{Q}}(\mathbb{R})$ -conjugacy class of the homomorphism

$$h_{G^{\mathbb{Q}}} : \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m \longrightarrow G_{\mathbb{R}}^{\mathbb{Q}},$$

defined with respect to the inclusion

$$(6.4) \quad G^{\mathbb{Q}}(\mathbb{R}) \subset \text{GL}_{F \otimes \mathbb{R}}(V \otimes \mathbb{R}) \xrightarrow[\sim]{\Phi} \prod_{\varphi \in \Phi} \text{GL}_{\mathbb{C}}(V_{\varphi}),$$

by $h_{G^{\mathbb{Q}}} = (h_{G^{\mathbb{Q}}, \varphi})_{\varphi \in \Phi}$ with the component $h_{G^{\mathbb{Q}}, \varphi}$ (on the \mathbb{R} -points)

$$h_{G^{\mathbb{Q}}, \varphi} : z \in \mathbb{C}^{\times} \longmapsto \text{diag}(z \cdot 1_{r_{\varphi}}, \bar{z} \cdot 1_{r_{\bar{\varphi}}}).$$

Then the reflex field $E(G^{\mathbb{Q}}, \{h_{G^{\mathbb{Q}}}\})$ is the reflex field E_r of r , which is the subfield of $\overline{\mathbb{Q}}$ defined by

$$(6.5) \quad \text{Gal}(\overline{\mathbb{Q}}/E_r) = \{ \sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \mid \sigma^*(r) = r \}.$$

Now, in addition to the CM type Φ , we also fix a distinguished element $\varphi_0 \in \Phi$. From now on we assume that the generalized CM type r is of *strict fake Drinfeld type* relative to Φ and φ_0 , in the sense of [39], i.e.,

$$r_\varphi = \begin{cases} n-1, & \varphi = \varphi_0, \\ n, & \varphi \in \Phi \setminus \{\varphi_0\}. \end{cases}$$

The first special case is when $n = 1$ and V is totally positive definite, i.e., V has signature $(1, 0)$ at each archimedean place.¹ In this case, we write $Z^\mathbb{Q} := G^\mathbb{Q}$ (a torus over \mathbb{Q}) and $h_{Z^\mathbb{Q}} := h_{G^\mathbb{Q}}$. The reflex field of $(Z^\mathbb{Q}, \{h_{Z^\mathbb{Q}}\})$ is E_Φ , the reflex field of Φ .

For general n , we set

$$\widetilde{G} := Z^\mathbb{Q} \times_{\mathbb{G}_m} G^\mathbb{Q},$$

where the two maps are respectively given by Nm_{F/F_0} and the similitude character. We form a Shimura datum for \widetilde{G} by

$$h_{\widetilde{G}} : \mathbb{C}^\times \xrightarrow{(h_{Z^\mathbb{Q}}, h_{G^\mathbb{Q}})} \widetilde{G}(\mathbb{R}).$$

Then the reflex field $E \subset \overline{\mathbb{Q}}$ of $(\widetilde{G}, \{h_{\widetilde{G}}\})$ is the composite $E_\Phi E_r$ (cf. [39, §3.2]). In particular, the field F is a subfield of E via φ_0 .

In [40, Rem. 3.2 (iii)] (also [39, §2.3]) the authors also defined a Shimura datum $(\text{Res}_{F_0/\mathbb{Q}} G, \{h_G\})$, where G is the unitary group $G = \text{U}(V)$ (an algebraic group over F_0); this gives the Shimura variety in the Gan–Gross–Prasad conjecture; cf. [9, §27]. Note that there is a natural isomorphism

$$\begin{aligned} \widetilde{G} &\xrightarrow{\sim} Z^\mathbb{Q} \times \text{Res}_{F_0/\mathbb{Q}} G \\ (z, g) &\longmapsto (z, z^{-1}g) \end{aligned}$$

and, when $K_{\widetilde{G}} = K_{Z^\mathbb{Q}} \times K_G$ is a decomposable compact open subgroup of $\widetilde{G}(\mathbb{A}_f)$, we have a product decomposition of the Shimura varieties over E

$$(6.6) \quad \text{Sh}_{K_{\widetilde{G}}}(\widetilde{G}, \{h_{\widetilde{G}}\}) \simeq \text{Sh}_{K_{Z^\mathbb{Q}}}(Z^\mathbb{Q}, \{h_{Z^\mathbb{Q}}\}) \times \text{Sh}_{K_G}(\text{Res}_{F_0/\mathbb{Q}} G, \{h_G\}).$$

6.2. Integral models.

6.2.1. *The auxiliary moduli problem for $Z^\mathbb{Q}$.* We recall the moduli problem \mathcal{M}_0 over O_{E_Φ} of [40, §3.2]. For a scheme S in $(\text{LNSch})_{/O_{E_\Phi}}$, we define $\mathcal{M}_0(S)$ to be the groupoid of triples $(A_0, \iota_0, \lambda_0)$, where

- A_0 is an abelian scheme over S ;
- $\iota_0 : O_F \rightarrow \text{End}(A_0)$ is an O_F -action satisfying the Kottwitz condition

$$(6.7) \quad \text{char}(\iota(a) \mid \text{Lie } A_0) = \prod_{\varphi \in \Phi} (T - \varphi(a)) \quad \text{for all } a \in O_F;$$

¹Here we follow the convention of [39], which differs from [40] where the space V is totally negative definite.

- λ_0 is a principal polarization on A_0 such that the induced Rosati involution via ι_0 coincides with the Galois involution on O_F .

A morphism between two objects $(A_0, \iota_0, \lambda_0)$ and $(A'_0, \iota'_0, \lambda'_0)$ in this groupoid is an O_F -linear isomorphism $\mu_0 : A_0 \rightarrow A'_0$ under which λ'_0 pulls back to λ_0 . Then the functor \mathcal{M}_0 is represented by a Deligne–Mumford stack, finite and étale over $\mathrm{Spec} O_{E_\Phi}$; cf. [18, Prop. 3.1.2].

The generic fiber M_0 of \mathcal{M}_0 is a disjoint union of copies of the Shimura variety $\mathrm{Sh}_{K_{Z\mathbb{Q}}^\circ}(Z^\mathbb{Q}, \{h_{Z\mathbb{Q}}\})$, where $K_{Z\mathbb{Q}}^\circ$ is the unique maximal compact subgroup of $Z^\mathbb{Q}(\mathbb{A}_f)$; cf. [40, Lem. 3.4] specializing to the ideal $\mathfrak{a} = O_{F_0}$. To avoid the possible emptiness of M_0 , we assume that F/F_0 is ramified throughout this paper (cf. [40, Rem. 3.5 (ii)]). For our purpose, it suffices to work with a fixed copy of the Shimura variety $\mathrm{Sh}_{K_{Z\mathbb{Q}}^\circ}(Z^\mathbb{Q}, \{h_{Z\mathbb{Q}}\})$ in the disjoint union in M_0 , and by abuse of notation we will still denote it by M_0 and by \mathcal{M}_0 the corresponding smooth integral model for the rest of the paper.

We also introduce a level structure for \mathcal{M}_0 . We let $K_{Z\mathbb{Q}} = \prod_p K_{Z\mathbb{Q},p} \subset Z^\mathbb{Q}(\mathbb{A}_f)$ be an open subgroup such that the prime-to- \mathfrak{d} components remain maximal. Analogous to \mathcal{M}_0 , there is a moduli functor $\mathcal{M}_{0,K_{Z\mathbb{Q}}}$ with $K_{Z\mathbb{Q}}$ -level structure, whose generic fiber is $\mathrm{Sh}_{K_{Z\mathbb{Q}}}(Z^\mathbb{Q}, \{h_{Z\mathbb{Q}}\})$. The construction is not important to this paper and we omit the detail (cf. [30, §C.3]); it suffices to mention that an object in the groupoid $\mathcal{M}_{0,K_{Z\mathbb{Q}}}(S)$ will be denoted by $(A_0, \iota_0, \lambda_0, \bar{\eta}_0)$, where $\bar{\eta}_0$ denotes a $K_{Z\mathbb{Q}}$ -level structure.

6.2.2. The RSZ integral model for $(\widetilde{G}, \{h_{\widetilde{G}}\})$. We now follow [40, §5.1] and [39, §6.1] to define the moduli interpretation of our Shimura varieties associated to the Shimura datum $(\widetilde{G}, \{h_{\widetilde{G}}\})$ for a certain special level structure. When $F_0 = \mathbb{Q}$, this is closely related to [25], [5]. In fact, for this paper, we only need an integral model over a suitable Zariski open subscheme of $\mathrm{Spec} O_E$.

Let \mathcal{D}_0 denote the finite set consisting of all non-archimedean places v of F_0 such that

- the residue characteristic of v is 2, or
- v is ramified in F , or
- v is inert in F where V_v is non-split.

Let \mathcal{D} be a finite set of non-archimedean places containing \mathcal{D}_0 , such that \mathcal{D} is pull-back from a set of places $\mathcal{D}_\mathbb{Q}$ of \mathbb{Q} . Define

$$\mathfrak{d} = \prod_{p \mid \mathcal{D}_\mathbb{Q}} p.$$

We will consider the Shimura variety for $(\widetilde{G}, \{h_{\widetilde{G}}\})$ with level-structure at the finite set of places dividing \mathfrak{d} .

For every non-archimedean $v \notin \mathcal{D}$, we fix a self-dual O_{F_v} -lattice $\Lambda_v^\circ \subset V_v$, i.e.,

$$(6.8) \quad \Lambda_v^\circ = (\Lambda_v^\circ)^\vee,$$

where we recall that $(\Lambda_v^\circ)^\vee$ denotes the dual lattice with respect to the hermitian forms on V_v . Let

$$(6.9) \quad K_{G,v}^\circ \subset \mathrm{U}(V_v)(F_{0,v})$$

be the stabilizer of the lattice Λ_v° , a hyperspecial compact open subgroup of $\mathrm{U}(V_v)(F_{0,v})$. Let $K_G = \prod_v K_{G,v} \subset G(\mathbb{A}_{0,f})$ be a compact open subgroup such that $K_{G,v} = K_{G,v}^\circ$ for all $v \nmid \mathfrak{d}$. Let $K_{Z\mathbb{Q}} = \prod_p K_{Z\mathbb{Q},p} \subset Z^\mathbb{Q}(\mathbb{A}_f)$ be an open subgroup such that the prime-to- \mathfrak{d} components remain maximal. Accordingly we define

$$K_{\tilde{G}} = K_{Z\mathbb{Q}} \times K_G \subset \tilde{G}(\mathbb{A}_f).$$

Recall that $E = E_\Phi E_r$ is the reflex field of $(\tilde{G}, \{h_{\tilde{G}}\})$.

Definition 6.1. The functor $\mathcal{M}_{K_{\tilde{G}}}(\tilde{G})$ associates the groupoid of tuples $(A_0, \iota_0, \lambda_0, A, \iota, \lambda, \bar{\eta})$ to each scheme S in $(\mathrm{LNSch})_{/O_E[1/\mathfrak{d}]}$, where

- $(A_0, \iota_0, \lambda_0, \bar{\eta}_0)$ is an object of $\mathcal{M}_{0, K_{Z\mathbb{Q}}}(S)$;
- A is an abelian scheme over S ;
- $\iota : O_F[1/\mathfrak{d}] \rightarrow \mathrm{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}[1/\mathfrak{d}]$ is an action satisfying the Kottwitz condition of signature

$$((n-1, 1)_{\varphi_0}, (n, 0)_{\varphi \in \Phi \setminus \{\varphi_0\}})$$

on $O_F[1/\mathfrak{d}]$;

- $\lambda : A \rightarrow A^\vee$ is a prime-to- \mathfrak{d} principle polarization whose Rosati involution inducing the Galois involution on $O_F[1/\mathfrak{d}]$ with respect to ι ;
- $\bar{\eta}$ is a $\prod_{v \mid \mathfrak{d}} K_{G,v}$ -orbit of isometries of hermitian modules (as smooth $F_\mathfrak{d} = \prod_{v \mid \mathfrak{d}} F_v$ -sheaves on S endowed with its natural hermitian form induced by the polarization)

$$(6.10) \quad \eta : \mathrm{V}_\mathfrak{d}(A_0, A) \xrightarrow{\sim} V(F_{0,\mathfrak{d}}),$$

where

$$\mathrm{V}_\mathfrak{d}(A_0, A) := \prod_{p \mid \mathfrak{d}} \mathrm{V}_p(A_0, A), \quad \mathrm{V}_p(A_0, A) = \mathrm{Hom}_{F \otimes_{\mathbb{Q}} \mathbb{Q}_p}(V_p(A_0), V_p(A))$$

and

$$(6.11) \quad V(F_{0,\mathfrak{d}}) := \prod_{p \mid \mathfrak{d}} V \otimes_{\mathbb{Q}} \mathbb{Q}_p = \prod_{v \mid \mathfrak{d}} V \otimes_{F_0} F_{0,v}.$$

More precisely, this is understood in the sense of, e.g., [25, Rem. 4.2]. Fixing any geometric point \bar{s} of a connected scheme S , the rational Tate module

$V_{\mathfrak{d}}(A_0, A)$ is a smooth $F_{\mathfrak{d}} = \prod_{v \mid \mathfrak{d}} F_v$ -sheaf on S determined by the rational Tate module $V_{\mathfrak{d}}(A_{0, \bar{s}}, A_{\bar{s}})$ together with the action of the fundamental group $\pi_1(S, \bar{s})$. Moreover, the polarizations on A_0 and A defines an $F_{\mathfrak{d}}$ -valued hermitian forms $\langle \cdot, \cdot \rangle$ on $V_{\mathfrak{d}}(A_{0, \bar{s}}, A_{\bar{s}})$:

$$\langle x, y \rangle = \lambda_0^{-1} \circ y^{\vee} \circ \lambda \circ x \in \text{End}_{F_{\mathfrak{d}}}(V_{\mathfrak{d}}(A_{0, \bar{s}})) = F_{\mathfrak{d}}.$$

Then the level structure $\bar{\eta}$ is a $\prod_{v \mid \mathfrak{d}} K_{G, v}$ -orbit of isometries of hermitian modules

$$(6.12) \quad \eta : V_{\mathfrak{d}}(A_{0, \bar{s}}, A_{\bar{s}}) \xrightarrow{\sim} V(F_{0, \mathfrak{d}})$$

that is required to be stable under the action of $\pi_1(S, \bar{s})$. The notion of $\prod_{v \mid \mathfrak{d}} K_{G, v}$ -level structure is independent of the choice of the geometric point \bar{s} on S .

- Finally, we impose the Eisenstein condition (cf. [39, §5.2]) for every place $w \nmid \mathfrak{d}$ of F ².

A morphism between two objects $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta})$ and $(A'_0, \iota'_0, \lambda'_0, \bar{\eta}'_0, A', \iota', \lambda', \bar{\eta}')$ is an isomorphism $(A_0, \iota_0, \lambda_0, \bar{\eta}_0) \xrightarrow{\sim} (A'_0, \iota'_0, \lambda'_0, \bar{\eta}'_0)$ in $\mathcal{M}_{0, K_{Z^{\mathbb{Q}}}}(S)$ and an O_F -linear \mathfrak{d} -isogeny $A \rightarrow A'$, pulling λ' back to λ and $\bar{\eta}'$ back to $\bar{\eta}$.

THEOREM 6.2. *The functor $\mathcal{M}_{K_{\tilde{G}}}(\tilde{G})$ is represented by a Deligne–Mumford stack. The morphism $\mathcal{M}_{K_{\tilde{G}}}(\tilde{G}) \rightarrow \text{Spec } O_E[1/\mathfrak{d}]$ is separated of finite type, and smooth of relative dimension $n - 1$.*

Proof. This follows from [40, Th. 5.2] (when all $p \nmid \mathfrak{d}$ are unramified in F) and [39, Th. 6.2], except we note that here we have omitted the sign conditions in loc. cit. However, the sign conditions hold automatically for all places away from \mathfrak{d} (cf. [39, Rem. 6.5(i)]), and therefore these two theorems still apply to the current situation. \square

Note that when both $\prod_{p \mid \mathfrak{d}} K_{Z^{\mathbb{Q}}, p}$ and $\prod_{p \mid \mathfrak{d}} K_{G, p}$ are small enough, the functor $\mathcal{M}_{K_{\tilde{G}}}(\tilde{G})$ is represented by a quasi-projective scheme, smooth over $\text{Spec } O_E[1/\mathfrak{d}]$. We will make this smallness assumption for the rest of the paper.

By [40, Prop. 3.5], the generic fiber $M_{K_{\tilde{G}}}(\tilde{G})$ of $\mathcal{M}_{K_{\tilde{G}}}(\tilde{G})$ is isomorphic to the canonical model of $\text{Sh}_{K_{\tilde{G}}}(\tilde{G}, \{h_{\tilde{G}}\})$. We also recall the moduli functor $M_{K_{\tilde{G}}}(\tilde{G})$ over $\text{Spec } E$ for *any* compact open subgroup $K_{\tilde{G}} \subset \tilde{G}(\mathbb{A}_f)$ of the form $K_{\tilde{G}} = K_{Z^{\mathbb{Q}}} \times K_G$. Similar to [Definition 6.1](#), the functor $M_{K_{\tilde{G}}}(\tilde{G})$ associates to each scheme S in $(\text{LNSch})_E$ the groupoid of tuples $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta})$,

²The Eisenstein condition at a place w of F [39, §5.2] follows from the Kottwitz condition if we assume that w is unramified over p . Therefore we only have this condition at finitely many places w ramified over p .

where everything is the same as [Definition 6.1](#) with the following minor change. Now $\iota : F \rightarrow \text{End}^\circ(A)$ is an F -action, $\lambda : A \rightarrow A^\vee$ is a polarization, and $\bar{\eta}$ is a K_G -orbit of isometries of $\mathbb{A}_{F,f}/\mathbb{A}_{F_0,f}$ -hermitian modules

$$(6.13) \quad \eta : \widehat{V}(A_0, A) \xrightarrow{\sim} V(\mathbb{A}_{F_0,f}),$$

where

$$\widehat{V}(A_0, A) := \prod_p V_p(A_0, A)$$

and $V(\mathbb{A}_{F_0,f}) = V \otimes_{F_0} \mathbb{A}_{F_0,f}$. The rest is the same as [Definition 6.1](#), with the appropriate modification of the definition of morphisms in the groupoid; cf. [\[40, §3.2\]](#).

7. Kudla–Rapoport divisors and the derived CM cycles

In this section we consider two types of special cycles on the integral models of Shimura varieties introduced in the previous section:

- the Kudla–Rapoport special divisors [\[25\]](#), and
- the derived CM cycle, which is a variant of the (1-dimensional) “big CM cycle” of Bruinier–Kudla–Yang and Howard [\[6\]](#), [\[18\]](#).

The derived CM cycle is the main novel geometric construction of this paper.

We make the following notational assumption: in Part 2 of the paper, all Schwartz functions on totally disconnected topological spaces are \mathbb{Q} -valued. The reason for this assumption is to define elements in various Chow groups or K -groups with \mathbb{Q} -coefficients.

7.1. The global Kudla–Rapoport divisors on $M_{K_{\widetilde{G}}}(\widetilde{G})$ over $\text{Spec } E$. First we define the global Kudla–Rapoport divisors on the canonical model $M_{K_{\widetilde{G}}}(\widetilde{G})$ of $\text{Sh}_{K_{\widetilde{G}}}(\widetilde{G}, \{h_{\widetilde{G}}\})$ over $\text{Spec } E$, introduced at the end of [Section 6.2.2](#), for an arbitrary compact open subgroup of the form $K_{\widetilde{G}} = K_{Z^\mathbb{Q}} \times K_G$. We follow [\[25\]](#) when $F_0 = \mathbb{Q}$, and [\[30, Def. 4.21\]](#) and [\[39, §3.5\]](#) for a general totally real field F_0 .

Let $\xi \in F_{0,+}$, and let $\mu \in V(\mathbb{A}_{0,f})/K_G$ be a K_G -orbit.

Definition 7.1. For each scheme S in $(\text{LNSch})_{/E}$, the S -points of the KR cycle $Z(\xi, \mu)$ is the groupoid of tuples $(A_0, \iota_0, \lambda_0, A, \iota, \lambda, \bar{\eta}, u)$, where

- $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}) \in M_{K_{\widetilde{G}}}(\widetilde{G})(S)$.
- $u \in \text{Hom}_F^\circ(A_0, A)$ such that $\langle u, u \rangle = \xi$, and $\bar{\eta}(u)$ is a homomorphism in the K_G -orbit μ . Here $\langle \cdot, \cdot \rangle$ denotes the hermitian form on $\text{Hom}_F^\circ(A_0, A)$ induced by the polarization λ_0 and λ for $x, y \in \text{Hom}_F^\circ(A_0, A)$:

$$\langle x, y \rangle = \lambda_0^{-1} \circ y^\vee \circ \lambda \circ x \in \text{End}_F^\circ(A_0) \simeq F.$$

A morphism between two objects $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, u)$ and $(A'_0, \iota'_0, \lambda'_0, \bar{\eta}'_0, A', \iota', \lambda', \bar{\eta}', u')$ is an isomorphism $(A_0, \iota_0, \lambda_0, \bar{\eta}_0) \xrightarrow{\sim} (A'_0, \iota'_0, \lambda'_0, \bar{\eta}'_0)$ in $\mathcal{M}_0(S)$ and an F -linear isogeny $\varphi : A \rightarrow A'$, compatible with λ and λ' , with $\bar{\eta}$ and $\bar{\eta}'$, and such that $u' = u \circ \varphi$.

Forgetting u defines a natural morphism $i : Z(\xi, \mu) \rightarrow M_{K_{\bar{G}}}(\bar{G})$; we defer to [Proposition 7.3](#) for its properties. In particular, the push-forward defines a class in the Chow group $\text{Ch}^1(M_{K_{\bar{G}}}(\bar{G}))$. For $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))^{K_G}$, we define

$$(7.1) \quad Z(\xi, \phi) := \sum_{\mu \in V_{\xi}(\mathbb{A}_{0,f})/K_G} \phi(\mu) Z(\xi, \mu),$$

viewed as an element in the Chow group $\text{Ch}^1(M_{K_{\bar{G}}}(\bar{G}))$. Here V_{ξ} is defined in [Section 1.2](#) after [\(1.4\)](#). Note that [\(7.1\)](#) is a finite sum due to the compactness of the support of ϕ (and $G(\mathbb{A}_{0,f})$ acts transitively on $V_{\xi}(\mathbb{A}_{0,f})$ when $\xi \neq 0$).

7.2. The global Kudla–Rapoport divisors on the integral model $\mathcal{M}_{K_{\bar{G}}}(\bar{G})$.

We now consider the moduli function $\mathcal{M} = \mathcal{M}_{K_{\bar{G}}}(\bar{G})$ with level structure at primes dividing \mathfrak{d} ; cf. [Definition 6.1](#). Here $K_{\bar{G}}$ is of the form $K_{\bar{G}} = K_{Z\mathbb{Q}} \times K_G$ with $K_G = \prod_v K_{G,v} \subset G(\mathbb{A}_{0,f})$ such that $K_{G,v} = K_{G,v}^{\circ}$ for $v \nmid \mathfrak{d}$, where $K_{G,v}^{\circ}$ is the stabilizer of the self-dual lattice Λ_v° ; cf. [\(6.8\)](#) and [\(6.9\)](#).

Let $\xi \in F_{0,+}$ and $\mu \in V(F_{0,\mathfrak{d}})/K_{G,\mathfrak{d}}$. Here $V(F_{0,\mathfrak{d}})$ is as in [\(6.11\)](#).

Definition 7.2. For each scheme S in $(\text{LNSch})_{/O_E[1/\mathfrak{d}]}$, the S -points of the KR cycle $Z(\xi, \mu)$ is the groupoid of tuples $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, u)$ where

- $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}) \in \mathcal{M}_{K_{\bar{G}}}(\bar{G})(S)$.
- $u \in \text{Hom}_{O_F}(A_0, A) \otimes_{\mathbb{Z}} \mathbb{Z}[1/\mathfrak{d}]$ such that $\langle u, u \rangle = \xi$, and $\bar{\eta}(u)$ is a homomorphism in the $K_{G,\mathfrak{d}}$ -orbit μ . Here $\langle \cdot, \cdot \rangle$ denotes the hermitian form induced by the polarization λ_0 and λ :

$$\langle x, y \rangle = \lambda_0^{-1} \circ y^{\vee} \circ \lambda \circ x \in \text{End}_{O_F}(A_0) \otimes_{\mathbb{Z}} \mathbb{Z}[1/\mathfrak{d}] \simeq O_F[1/\mathfrak{d}].$$

A morphism between two objects $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, u)$ and $(A'_0, \iota'_0, \lambda'_0, \bar{\eta}'_0, A', \iota', \lambda', \bar{\eta}', u')$ is an isomorphism $(A_0, \iota_0, \lambda_0, \bar{\eta}_0) \xrightarrow{\sim} (A'_0, \iota'_0, \lambda'_0, \bar{\eta}'_0)$ in $\mathcal{M}_{0,K_{Z\mathbb{Q}}}(S)$ and an O_F -linear prime-to- \mathfrak{d} isogeny $\varphi : A \rightarrow A'$, compatible with λ and λ' , with $\bar{\eta}$ and $\bar{\eta}'$, and such that $u' = u \circ \varphi$.

Forgetting u defines a natural morphism $i : Z(\xi, \mu) \rightarrow \mathcal{M}_{K_{\bar{G}}}(\bar{G})$.

PROPOSITION 7.3.

- (a) *The morphism $i : Z(\xi, \mu) \rightarrow \mathcal{M}_{K_{\bar{G}}}(\bar{G})$ is representable, finite and unramified.*
- (b) *The morphism i defines étale locally a Cartier divisor. Moreover, the morphism $Z(\xi, \mu) \rightarrow \text{Spec } O_E[1/\mathfrak{d}]$ is flat.*

Proof. When $F_0 = \mathbb{Q}$, part (a) follows from [25, Prop. 2.9], and part (b) follows from [5, §2.5]. For a general totally real F_0 , both follow from [30, Prop. 4.22]. \square

To a function $\phi_{\mathfrak{d}} \in \mathcal{S}(V_{\mathfrak{d}})^{K_{G,\mathfrak{d}}}$, we associate $\phi = \mathbf{1}_{\Lambda^{\mathfrak{d}}} \otimes \phi_{\mathfrak{d}} \in \mathcal{S}(V(\mathbb{A}_{0,f}))$, where $\Lambda^{\mathfrak{d}} = \prod_{v \nmid \mathfrak{d}} \Lambda_v^{\mathfrak{d}}$ for the self-dual lattice $\Lambda_v^{\mathfrak{d}}$ in (6.8). Then we define

$$(7.2) \quad \mathcal{Z}(\xi, \phi) := \sum_{\mu \in V_{\xi}(F_0, \mathfrak{d})/K_{G,\mathfrak{d}}} \phi_{\mathfrak{d}}(\mu) \mathcal{Z}(\xi, \mu),$$

viewed as an element in the Chow group $\text{Ch}^1(\mathcal{M}_{K_{\widetilde{G}}}(\widetilde{G}))$. For such functions $\phi_{\mathfrak{d}}$ and the associated ϕ , the generic fiber of $\mathcal{Z}(\xi, \phi)$ is $Z(\xi, \phi)$ defined by (7.1) (specializing to the current level $K_{\widetilde{G}}$). In particular, the generic fiber of $\mathcal{Z}(\xi, \mu)$ is the union of the KR cycles $Z(\xi, \mu')$ in Definition 7.1 for suitable $\mu' \in V(\mathbb{A}_{0,f})/K_G$.

7.3. Special divisors in the formal neighborhood of the basic locus. We consider the restriction of the KR divisors to the formal completion of $\mathcal{M} = \mathcal{M}_{K_{\widetilde{G}}}(\widetilde{G})$ along the basic locus.

Let $\nu \nmid \mathfrak{d}$ be a non-archimedean place of E . Its restriction to F (resp. F_0) is a place denoted by w_0 (resp. v_0). Assume that v_0 is *inert*. We recall from [40, §8, in the proof of Th. 8.15] the non-archimedean uniformization along the basic locus:

$$(7.3) \quad \widehat{\mathcal{M}_{O_{\check{E}_{\nu}}}} := (\mathcal{M}_{(\nu)} \otimes_{O_{E,(\nu)}} O_{\check{E}_{\nu}})^{\widehat{}} = \widetilde{G}'(\mathbb{Q}) \setminus \left[\mathcal{N}' \times \widetilde{G}(\mathbb{A}_f^p)/K_{\widetilde{G}}^p \right].$$

Here the hat on the left-hand side denotes the completion along the basic locus in the geometric special fiber of $\mathcal{M}_{(\nu)}$. The group \widetilde{G}' is an inner twist of \widetilde{G} . More precisely, the group \widetilde{G}' is associated to the “nearby” hermitian space V' that is positive definite at all archimedean places, and isomorphic to V , locally at all non-archimedean places except at v_0 . Let $\mathcal{N} = \mathcal{N}_{n, F_{w_0}/F_{v_0}} \rightarrow \text{Spf } O_{F_{w_0}}$ be the RZ space introduced in Section 3.1, and take its base change $\mathcal{N}_{O_{\check{E}_{\nu}}} = \mathcal{N} \widehat{\otimes}_{O_{F_{w_0}}} O_{\check{E}_{\nu}}$. Then as in loc. cit.,³ we may rewrite (7.3) as

$$(7.4) \quad \widehat{\mathcal{M}_{O_{\check{E}_{\nu}}}} = \widetilde{G}'(\mathbb{Q}) \setminus \left[\mathcal{N}_{O_{\check{E}_{\nu}}} \times \widetilde{G}(\mathbb{A}_f^{v_0})/K_{\widetilde{G}}^{v_0} \right].$$

³The formal scheme in the Rapoport–Zink uniformization theorem is the “absolute” RZ space of PEL-type in [31] rather than the “relative” RZ space $\mathcal{N}_{n, F_{w_0}/F_{v_0}}$ in Section 3.1. These two RZ spaces coincide by [31, Th. 3.1], noting that the Eisenstein condition imposed in [31] for the signature $(r, s) = (n - 1, 1)$ reduces to the Eisenstein condition in [39, §5.2, case (2)] for the definition of the moduli space $\mathcal{M} = \mathcal{M}_{K_{\widetilde{G}}}(\widetilde{G})$ in this paper.

Here, by abuse of notation, we denote

$$\widetilde{G}(\mathbb{A}_f^{v_0})/K_G^{v_0} = \widetilde{G}(\mathbb{A}_f^p)/K_G^p \times (Z^\mathbb{Q}(\mathbb{Q}_p)/K_{Z^\mathbb{Q},p}) \times \prod_{v \in S_p \setminus \{v_0\}} G(F_{0,v})/K_{G,v},$$

where S_p denotes the set of places of F_0 above p . For the action of the group $\widetilde{G}'(\mathbb{Q})$ in (7.4), we fix an isomorphism $\widetilde{G}'(\mathbb{A}_f^{v_0}) \simeq \widetilde{G}(\mathbb{A}_f^{v_0})$.

Note that the uniformization (7.4) induces a projection to a discrete set (in fact an abelian group)

$$(7.5) \quad \mathcal{M}_{O_{\check{E}_\nu}} \longrightarrow Z^\mathbb{Q}(\mathbb{Q}) \backslash (Z^\mathbb{Q}(\mathbb{A}_f)/K_{Z^\mathbb{Q}}).$$

This gives a partition of the formal scheme $\mathcal{M}_{O_{\check{E}_\nu}}$, each fiber is naturally isomorphic to

$$(7.6) \quad \mathcal{M}_{O_{\check{E}_\nu},0} := G'(F_0) \backslash \left[\mathcal{N}_{O_{\check{E}_\nu}} \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0} \right].$$

Recall that we have the local KR divisors $\mathcal{Z}(u)$ on $\mathcal{N} = \mathcal{N}_{n,F_{w_0}/F_{0,v_0}}$ for each $u \in V' \otimes F_{0,v_0} \simeq \text{Hom}^\circ(\mathbb{E}, \mathbb{X}_n)$, the hermitian space of local special homomorphisms (for some fixed framing objects \mathbb{E} and \mathbb{X}_n in the uniformization (7.4) above). For a pair $(u, g) \in V'(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$ with $u \neq 0$, we define the product divisor on $\mathcal{N}_{O_{\check{E}_\nu}} \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$,

$$(7.7) \quad \mathcal{Z}(u, g)_{K_G^{v_0}} = \mathcal{Z}(u) \times \mathbf{1}_{g K_G^{v_0}}.$$

We then consider

$$\sum \mathcal{Z}(u', g')_{K_G^{v_0}},$$

where the sum is over (u', g') in the $G'(F_0)$ -orbit of the pair (u, g) for the diagonal action of $G'(F_0)$ on $V'(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$. Since the sum is $G'(F_0)$ -invariant, it descends to a divisor on the quotient $\mathcal{M}_{O_{\check{E}_\nu},0}$ in (7.6), which we denote by $[\mathcal{Z}(u, g)]_{K_G^{v_0}}$.

PROPOSITION 7.4. *Let $\xi \in F_{0,+}$. The restriction of the special divisor $\mathcal{Z}(\xi, \phi)$ to each fiber of the above projection (7.5) is the sum*

$$(7.8) \quad \sum_{(u,g) \in G'(F_0) \backslash (V'_\xi(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0})} \phi^{v_0}(g^{-1}u) \cdot [\mathcal{Z}(u, g)]_{K_G^{v_0}},$$

viewed as a divisor on (7.6). This is a finite sum.

Remark 7.5. This is similar to the description of the special divisors over the complex number; cf. (8.8).

Proof. This follows from the proof of [25, Prop. 6.3]; also cf. [30, §4.2]. \square

7.4. Fat big CM cycles. We introduce a fat variant of the “big CM cycle” in [6], [18] on our moduli space $\mathcal{M} = \mathcal{M}_{K_{\bar{G}}}(\bar{G})$ with level structure at primes dividing \mathfrak{d} (cf. Definition 6.1).

Fix an $\alpha \in \mathcal{A}_n(F_0) \subset F[T]_{\deg=n}$ (cf. the end of Section 2.2). If α has no repeated roots, then

$$(7.9) \quad F' = F[T]/(\alpha)$$

is a semi-simple F -algebra. There is a unique F_0 -linear involution on F' sending $T \rightarrow T^{-1}$ and extending the Galois involution for F/F_0 . Then the fixed subalgebra F'_0 is a product of totally real field extensions of F_0 , and $F' \simeq F \otimes_{F_0} F'_0$ with the involution of F'/F'_0 induced from that of F/F_0 .

Now let $\alpha \in \mathcal{A}_n(O_{F_0}[1/\mathfrak{d}])$ be irreducible over F . Then the algebra F' in (7.9) is a field. Throughout the rest of the paper we will assume F' is a CM extension of F'_0 . This is a necessary condition for the functor in Definition 7.6 below to be non-empty. We denote

$$(7.10) \quad R_\alpha = O_F[1/\mathfrak{d}][T]/(\alpha),$$

viewed as a sub-ring of the CM field F' .

Definition 7.6. The functor $\mathcal{CM}(\alpha) = \mathcal{CM}_{K_{\bar{G}}}(\alpha)$ associates to each scheme S in $(\text{LNSch})_{/O_E[1/\mathfrak{d}]}$ the groupoid of tuples $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, \varphi)$ where $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}) \in \mathcal{M}_{K_{\bar{G}}}(\bar{G})$ and $\varphi \in \text{End}_{O_F}(A) \otimes \mathbb{Z}[1/\mathfrak{d}]$ such that

- the polynomial α annihilates the endomorphism φ ;
- φ is compatible with λ , i.e., $\varphi^* \lambda = \lambda$, or equivalently, the Rosati involution sends φ to φ^{-1} ; and
- φ preserves the level structure $\bar{\eta}$, i.e., we have a commutative diagram

$$\begin{array}{ccc} V_{\mathfrak{d}}(A_0, A) & \xrightarrow{\eta_1} & V(F_{0,\mathfrak{d}}) \\ \downarrow \varphi & & \downarrow \text{id} \\ V_{\mathfrak{d}}(A_0, A) & \xrightarrow{\eta_2} & V(F_{0,\mathfrak{d}}) \end{array}$$

for some $\eta_1, \eta_2 \in \bar{\eta}$.

Morphisms in the groupoid are defined in the obvious way.

We have a natural forgetful map

$$\mathcal{CM}_{K_{\bar{G}}}(\alpha) \longrightarrow \mathcal{M}_{K_{\bar{G}}}(\bar{G}).$$

We call $\mathcal{CM}_{K_{\bar{G}}}(\alpha)$ the (naive) fat big CM cycle, or simply CM cycle.

Remark 7.7. The abelian scheme A in the moduli functor $\mathcal{CM}(\alpha)$ carries an R_α -action where the T in (7.10) acts by φ . Our moduli functor $\mathcal{CM}(\alpha)$ is analogous to the big CM cycle defined in [18], where R_α is replaced by the ring of integers $O_{F'}$ in $F' = F[T]/(\alpha)$. A minor difference is that we do not impose any Kottwitz signature condition in our Definition 7.6, while [18, Def. 3.11]

does. The main new feature of our moduli functor $\mathcal{CM}(\alpha)$ is that we allow R_α to be a non-maximal order in $O_{F'}[1/\mathfrak{d}]$. As a result, it could have very complicated structure in positive characteristic (e.g., with large dimensional components). A complete understanding of the geometric structure of $\mathcal{CM}(\alpha)$ seems a hard question (e.g., to determine all of its irreducible components in its special fibers), and the AFL type identity in this paper gives us a partial answer.

We also define a twisted variant of $\mathcal{CM}(\alpha)$.

Definition 7.8. Let $g \in G(F_{0,\mathfrak{d}})$. The functor $\mathcal{CM}(\alpha, g) = \mathcal{CM}_{K_{\bar{G}}}(\alpha, g)$ associates to each $O_E[1/\mathfrak{d}]$ -scheme S the groupoid of tuples $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, \varphi)$, where $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}) \in \mathcal{M}_{K_{\bar{G}}}(\bar{G})$ and $\varphi \in \text{End}_{O_F}(A) \otimes \mathbb{Z}[1/\mathfrak{d}]$ such that

- the polynomial α annihilates the endomorphism φ ;
- φ is compatible with λ , i.e., $\varphi^* \lambda = \lambda$, or equivalently, the Rosati involution sends φ to φ^{-1} ; and
- we have a commutative diagram

$$\begin{array}{ccc} V_{\mathfrak{d}}(A_0, A) & \xrightarrow{\eta_1} & V(F_{0,\mathfrak{d}}) \\ \downarrow \varphi & & \downarrow g \\ V_{\mathfrak{d}}(A_0, A) & \xrightarrow{\eta_2} & V(F_{0,\mathfrak{d}}) \end{array}$$

for some $\eta_1, \eta_2 \in \bar{\eta}$.

Morphisms are defined in the obvious way.

Denote by $\text{Ram}(\alpha)$ the set of non-archimedean places $v \nmid \mathfrak{d}$ of F_0 where $R_\alpha = O_F[1/\mathfrak{d}][T]/(\alpha)$ is non-maximal (i.e., $R_{\alpha,v} := R_\alpha \otimes_{O_{F_0}} O_{F_0,v}$ is not a product of DVRs).

PROPOSITION 7.9. *Let $\alpha \in \mathcal{A}_n(O_{F_0}[1/\mathfrak{d}])$ be irreducible over F . Let $g \in \prod_{v \mid \mathfrak{d}} G(F_v)$.*

- (a) *The morphism $\mathcal{CM}(\alpha, g) \rightarrow \mathcal{M}$ is representable, finite and unramified.*
- (b) *The morphism $\mathcal{CM}(\alpha, g) \rightarrow \text{Spec } O_E[1/\mathfrak{d}]$ is proper. Its restriction to the open sub-scheme $\text{Spec } O_E[1/\mathfrak{d}] \setminus \text{Ram}(\alpha)$ is finite étale.*

Proof. The first part follows similarly to [Proposition 7.3](#). (By the theory of Hilbert scheme, the morphism is representable by a disjoint union of schemes of finite type; it is of finite type by the first condition $\alpha(\varphi) = 0$; ⁴ it is then quasi-finite because there are only finitely many ways to endow an action of the order R_α to a given (A, ι, λ) over an arbitrary field; the unramifiedness

⁴Alternatively, one can argue using [Lemmas 7.11](#) and [7.15](#).

follows from the rigidity of quasi-isogeny; by the valuative criterion by the Néron property of abelian scheme, the morphism is proper, and hence finite.)

The properness of $\mathcal{CM}(\alpha, g) \rightarrow \text{Spec } O_E[1/\mathfrak{d}]$ follows by the valuative criterion. (The toric part of a semi-abelian scheme will have too small dimension to carry an action of R .) Finally, the argument of [18, Prop. 3.1.2(3)] still holds to show the finiteness and étaleness over $\text{Spec } O_E[1/\mathfrak{d}] \setminus \text{Ram}(\alpha)$: at every place above $v \notin \text{Ram}(\alpha)$, the local order $R_{\alpha, v}$ is maximal, and hence the p -divisible group has formal multiplication by a local maximal order. \square

7.5. Hecke correspondences and their fixed point loci. We first introduce the characteristic polynomial of an endomorphism of an abelian variety. Then we apply it to study the fixed point loci of Hecke correspondences.

Let k be an arbitrary field, and let A be an abelian variety over k . Then we define the characteristic polynomial of $\varphi \in \text{End}^\circ(A)$, denoted by $\text{char}_{\mathbb{Q}}(\varphi)$, as

$$\text{char}_{\mathbb{Q}}(\varphi) = \det(T - \varphi|V_\ell(A)) \in \mathbb{Q}_\ell[T]_{\deg=2\dim A},$$

where ℓ is any prime different from the characteristic of k , and $V_\ell(A)$ denotes the rational ℓ -adic Tate module of A . Similarly, if $\iota : F \rightarrow \text{End}^\circ(A)$ is an F -action, and $\varphi \in \text{End}_F^\circ(A)$, then $V_\ell(A)$ is a free $F \otimes_{\mathbb{Q}} \mathbb{Q}_\ell$ -module of rank $n := \frac{2\dim A}{[F:\mathbb{Q}]}$. We then define

$$\text{char}_F(\varphi) = \det_{F \otimes_{\mathbb{Q}} \mathbb{Q}_\ell}(T - \varphi|V_\ell(A)) \in F \otimes_{\mathbb{Q}} \mathbb{Q}_\ell[T]_{\deg=n},$$

viewing $V_\ell(A)$ as a free $F \otimes_{\mathbb{Q}} \mathbb{Q}_\ell$ -module.

LEMMA 7.10.

- (a) *The characteristic polynomial $\text{char}_{\mathbb{Q}}(\varphi)$ is in $\mathbb{Q}[T]_{\deg=2\dim A}$ and is independent of the choice of ℓ .*
- (b) *If $\iota : F \rightarrow \text{End}^\circ(A)$ is an F -action, and φ is in $\text{End}_F^\circ(A)$, then the characteristic polynomial $\text{char}_F(\varphi)$ is in $F[T]_{\deg=n}$ and is independent of the choice of ℓ .*

Proof. The characteristic polynomial $\text{char}_{\mathbb{Q}}(\varphi)$ is determined by its value at $T = m \in \mathbb{Q} \subset \text{End}^\circ(A)$, in which case we have

$$\text{char}_{\mathbb{Q}}(\varphi)(m) = \deg(m - \varphi) \in \mathbb{Q}_{\geq 0}.$$

This proves the first part.

Let $\text{tr}_{\mathbb{Q}}(\varphi)$ be the negation of the coefficient of $T^{2\dim A-1}$ in the polynomial $\text{char}_{\mathbb{Q}}(\varphi)$. Then we obtain a \mathbb{Q} -linear map $\text{tr}_{\mathbb{Q}} : \text{End}^\circ(A) \rightarrow \mathbb{Q}$. Then knowing $\text{tr}_{\mathbb{Q}}(\varphi^i)$ for all $i \geq 0$ is equivalent to knowing $\text{char}_{\mathbb{Q}}(\varphi)$. If φ commutes with the F -action $\iota : F \rightarrow \text{End}^\circ(A)$, we define $\text{tr}_F(\varphi) \in F$, characterized by

$$\text{tr}_{F/\mathbb{Q}}(a \text{tr}_F(\varphi)) = \text{tr}_{\mathbb{Q}}(\iota(a)\varphi) \quad \text{for all } a \in F.$$

From $\text{tr}_F(\varphi^i) \in F$ for all $i \geq 0$, there exists a unique polynomial in $F[T]_{\deg=n}$ recovering the characteristic polynomial $\text{char}_F(\varphi)$. This proves the second part. \square

LEMMA 7.11. *Let S be a connected locally noetherian scheme, and let $A \rightarrow S$ be an abelian scheme.*

- (a) *If $\varphi \in \text{End}^o(A)$, then the function $s \in S \mapsto \text{char}_{\mathbb{Q}}(\varphi) \in \mathbb{Q}[T]_{\deg=2\dim A}$ is constant.*
- (b) *Let $\iota : F \rightarrow \text{End}^o(A)$ and $\varphi \in \text{End}_F^o(A)$. Then the function $s \in S \mapsto \text{char}_F(\varphi) \in F[T]_{\deg=n}$ is constant.*

Proof. It suffices to show the assertion when some rational prime ℓ is invertible on S . (Otherwise, choose two primes $\ell_1 \neq \ell_2$, cover $\text{Spec } \mathbb{Z}$ by open sub-schemes $\text{Spec } \mathbb{Z}[1/\ell_1]$ and $\text{Spec } \mathbb{Z}[1/\ell_2]$, and then pull back to cover S .) Then the local constancy follows from the fact that the rational ℓ -adic Tate module $V_\ell(A)$ is a lisse étale sheaf on S . \square

We now define Hecke correspondences.

Definition 7.12. Let $g \in G(F_{0,\mathfrak{d}}) \subset G(\mathbb{A}_{0,f})$. The functor $\text{Hk}_{[K_G g K_G]}$ associates to each $O_E[1/\mathfrak{d}]$ -scheme S the groupoid of tuples

$$(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, A', \iota', \lambda', \bar{\eta}', \varphi),$$

where $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}), (A_0, \iota_0, \lambda_0, \bar{\eta}_0, A', \iota', \lambda', \bar{\eta}') \in \mathcal{M}_{K_{\bar{G}}}(\widetilde{G})(S)$, and a quasi-isogeny $\varphi \in \text{Hom}_{O_F}(A, A') \otimes \mathbb{Z}[1/\mathfrak{d}]$ such that

- φ is compatible with λ and λ' , i.e., $\varphi^* \lambda' = \lambda$;
- there exist $\eta \in \bar{\eta}$ and $\eta' \in \bar{\eta}'$ such that the following diagram commutes:

$$\begin{array}{ccc} V_{\mathfrak{d}}(A_0, A) & \xrightarrow{\eta} & V(F_{0,\mathfrak{d}}) \\ \downarrow \varphi & & \downarrow g \\ V_{\mathfrak{d}}(A_0, A') & \xrightarrow{\eta'} & V(F_{0,\mathfrak{d}}). \end{array}$$

Here the left vertical map on rational Tate modules is induced by φ . Note that this is to be understood similarly to the definition of level structure (cf. Definition 6.1).

Morphisms are defined in the obvious way.

We have a natural morphism

$$\text{Hk}_{[K_G g K_G]} \longrightarrow \mathcal{M} \times_{O_E[1/\mathfrak{d}]} \mathcal{M}.$$

This morphism is finite, and the projection to any one factor is a finite étale morphism.

Now consider the fiber product, called the “fixed point locus of the Hecke correspondence $\mathrm{Hk}_{[K_G g K_G]}$,”

$$\begin{array}{ccc} \mathcal{M}_{[K_G g K_G]} := \mathrm{Hk}_{[K_G g K_G]} \times_{\mathcal{M} \times \mathcal{M}} \Delta_{\mathcal{M}} & \longrightarrow & \mathrm{Hk}_{[K_G g K_G]} \\ \downarrow & & \downarrow \\ \mathcal{M} & \xrightarrow{\Delta} & \mathcal{M} \times_{O_E[1/\mathfrak{d}]} \mathcal{M}. \end{array}$$

Since \mathcal{M} is a scheme over $O_E[1/\mathfrak{d}]$ (under our smallness assumption on the compact open K_G), an object in $\mathcal{M}_{[K_G g K_G]}(S)$ can be represented by $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, \varphi)$.

By [Lemma 7.10](#) and [Lemma 7.11](#), we obtain a locally constant map (for the Zariski topology on the source)

$$(7.11) \quad \mathrm{char}_F : \mathcal{M}_{[K_G g K_G]} \longrightarrow F[T]_{\deg=n},$$

which sends a point $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, \varphi)$ in $\mathcal{M}_{[K_G g K_G]}$ to $\mathrm{char}_F(\varphi)$. The image is a finite set by [Lemma 7.11](#) because the source is of finite type and hence has only finitely many connected components. It follows that the fixed point locus $\mathcal{M}_{[K_G g K_G]}$ is a disjoint union of open and closed subschemes, indexed by the image under the map (7.11):

$$(7.12) \quad \mathcal{M}_{[K_G g K_G]} = \coprod_{\alpha \in \mathrm{Im}(\mathrm{char}_F)} \mathrm{char}_F^{-1}(\alpha).$$

LEMMA 7.13. *If $\alpha \in F[T]_{\deg=n}$ lies in the image of the map (7.11), then it is conjugate self-reciprocal, and all of its coefficients lie in $O_F[1/\mathfrak{d}]$ (i.e., $\alpha \in \mathcal{A}_n(O_F[1/\mathfrak{d}])$ in the notation (2.11)).*

Proof. Suppose that α is the image of a point $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, \varphi)$ over an algebraically closed field k . Since the endomorphism φ preserves the polarization λ , it preserves the hermitian form on $V_\ell(A_0, A)$ for any $\ell \nmid \mathfrak{d}$. Then the first assertion follows from the easy fact that the characteristic polynomial of an element preserving a hermitian form is conjugate self-reciprocal. To show that $\alpha \in O_F[1/\mathfrak{d}][T]$, it suffices to show that for every prime $\ell \nmid \mathfrak{d}$, we have $\alpha \in O_F \otimes_{\mathbb{Z}} \mathbb{Z}_\ell[T]$ when viewing $\alpha \in F \otimes_{\mathbb{Q}} \mathbb{Q}_\ell[T]$. If ℓ is different from the characteristic of the field k , the Tate module $T_\ell(A)$ is a free $O_F \otimes_{\mathbb{Z}} \mathbb{Z}_\ell$ -module. The endomorphism φ preserves $T_\ell(A)$, and hence its characteristic polynomial has coefficients in $O_F \otimes_{\mathbb{Z}} \mathbb{Z}_\ell$. If ℓ is equal to the characteristic of the field k , the desired integrality follows using the Dieudonné $M(A)$, a free $O_F \otimes_{\mathbb{Z}} W(k)$ -module of rank n , where $W(k)$ is the ring of Witt vectors of k . \square

Remark 7.14. Similarly, if $\varphi \in \mathrm{End}^\circ(A)$ preserves a polarization $\lambda : A \rightarrow A^\vee$ (i.e., $\varphi^* \lambda = \lambda$), then $\mathrm{char}_{\mathbb{Q}}(\varphi)$ is self-reciprocal (i.e., $T^{2 \dim A} \mathrm{char}_{\mathbb{Q}}(\varphi)(T^{-1}) = \mathrm{char}_{\mathbb{Q}}(\varphi)(T)$).

Finally, we relate the fixed point locus to the twisted CM cycle $\mathcal{CM}(\alpha, g)$ in [Definition 7.8](#).

LEMMA 7.15. *Let $\alpha \in \mathcal{A}_n(O_{F_0}[1/\mathfrak{d}])$ be irreducible over F . Let $g \in G(F_{0,\mathfrak{d}})$. Then the fiber of the map char_F (7.11) above the polynomial α is canonically isomorphic to the twisted CM cycle $\mathcal{CM}(\alpha, g)$ in [Definition 7.8](#).*

Proof. By [Definition 7.12](#), the fiber of the map char_F (7.11) above α is the functor whose S -points are the groupoid of tuples $(A_0, \iota_0, \lambda_0, \bar{\eta}_0, A, \iota, \lambda, \bar{\eta}, \varphi)$ satisfying the same conditions as in [Definition 7.8](#), except the first one, i.e., $\alpha(\varphi) = 0$. This condition is equivalent to the condition on the characteristic polynomial of φ by Cayley–Hamilton theorem and the assumption that α is irreducible. \square

7.6. Derived CM cycle ${}^{\mathbb{L}}\mathcal{CM}(\alpha, g)$. In [Section 7.5](#), the twisted CM cycle $\mathcal{CM}(\alpha, g)$ is recognized as a union of connected components of the fixed point locus $\mathcal{M}_{[K_G g K_G]}$ (cf. (7.11)):

$$(7.13) \quad \begin{array}{ccc} \mathcal{CM}(\alpha, g) & \longrightarrow & \mathcal{M}_{[K_G g K_G]} \longrightarrow \text{Hk}_{[K_G g K_G]} \\ & & \downarrow \\ & & \mathcal{M} \xrightarrow{\Delta} \mathcal{M} \times_{O_E[1/\mathfrak{d}]} \mathcal{M}. \end{array}$$

This allows us to define a derived CM cycle, by taking the restriction of the derived tensor product

$$(7.14) \quad {}^{\mathbb{L}}\mathcal{CM}(\alpha, g) := [\mathcal{O}_{\text{Hk}_{[K_G g K_G]}} \xrightarrow{\mathbb{L}} \mathcal{O}_{\mathcal{M}}] |_{\mathcal{CM}(\alpha, g)} \in K'_0(\mathcal{CM}(\alpha, g)).$$

Moreover, since Δ is a regular immersion of codimension $n - 1$, this element belongs to the filtration $F^{n-1}K'_0(\text{Hk}_{[K_G g K_G]})$, and hence by (B.3),⁵

$$(7.15) \quad {}^{\mathbb{L}}\mathcal{CM}(\alpha, g) \in F_1 K'_0(\mathcal{CM}(\alpha, g)).$$

We extend the derived CM cycle to a weighted version. Let $\mathcal{S}(G(F_{0,\mathfrak{d}}), K_{G,\mathfrak{d}})$ be the space of bi- $K_{G,\mathfrak{d}}$ -invariant Schwartz functions on $G(F_{0,\mathfrak{d}})$. For $\phi_{\mathfrak{d}} \in \mathcal{S}(G(F_{0,\mathfrak{d}}), K_{G,\mathfrak{d}})$, we denote $\phi_0 = \mathbf{1}_{K_G^{\mathfrak{d}}} \otimes \phi_{\mathfrak{d}} \in \mathcal{S}(G(\mathbb{A}_{0,f}), K_G)$; here $K_G^{\mathfrak{d}} = \prod_{v \nmid \mathfrak{d}} K_{G,v}^{\circ}$. We then define ${}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0)$ as the sum of above twisted CM cycles

$$(7.16) \quad {}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0) = \sum_{g \in K_G \backslash G(\mathbb{A}_{0,f}) / K_G} \phi_0(g) {}^{\mathbb{L}}\mathcal{CM}(\alpha, g),$$

⁵Strictly speaking the assertion (B.3) only applies to two closed subschemes Y, Z of X . Here, the right-most morphism in (7.13) is finite and hence preserves the dimension of any closed subscheme. Therefore we may apply (B.3) to the image of this morphism.

where we regard each summand ${}^L\mathcal{CM}(\alpha, g)$ as an element in

$$\bigoplus_{g \in K_{G,\mathfrak{d}} \backslash G(F_{0,\mathfrak{d}}) / K_{G,\mathfrak{d}}} K'_0(\mathcal{CM}(\alpha, g)).$$

Moreover, these elements lie in the filtration; cf. (7.15),

$$(7.17) \quad {}^L\mathcal{CM}(\alpha, \phi_0) \in \bigoplus_{g \in K_{G,\mathfrak{d}} \backslash G(F_{0,\mathfrak{d}}) / K_{G,\mathfrak{d}}} F_1 K'_0(\mathcal{CM}(\alpha, g)).$$

7.7. Hecke correspondences in the formal neighborhood of the basic locus.

We now consider the restriction of the Hecke correspondence $\mathrm{Hk}_{K_G g K_G}$ to the formal neighborhood of the basic locus at a non-archimedean place $v_0 \nmid \mathfrak{d}$ of F_0 , inert in F , via the RZ uniformization (7.4). We resume the notation there. We consider the fiber product (in the category of locally noetherian formal schemes)

$$\begin{array}{ccc} \widehat{\mathrm{Hk}}_{[K_G g K_G]} & \longrightarrow & \mathrm{Hk}_{[K_G g K_G]} \\ \downarrow & & \downarrow \\ \mathcal{M}_{O_{\check{E}_v}} \times_{\mathrm{Spf} O_{\check{E}_v}} \mathcal{M}_{O_{\check{E}_v}} & \longrightarrow & \mathcal{M} \times_{O_E[1/\mathfrak{d}]} \mathcal{M}. \end{array}$$

The commutative diagram in fact lives over the base $Z^{\mathbb{Q}}(\mathbb{Q}) \backslash (Z^{\mathbb{Q}}(\mathbb{A}_f) / K_{Z^{\mathbb{Q}}})$; cf. (7.5). Therefore it suffices to consider the fiber (cf. (7.6)) over any fixed element of $Z^{\mathbb{Q}}(\mathbb{Q}) \backslash (Z^{\mathbb{Q}}(\mathbb{A}_f) / K_{Z^{\mathbb{Q}}})$. It follows immediately that

PROPOSITION 7.16. *Let*

$$\mathrm{Hk}_{[K_G g K_G]}^{(v_0)} := \{(g_1, g_2) \in G(\mathbb{A}_{0,f}^{v_0}) / K_G^{v_0} \times G(\mathbb{A}_{0,f}^{v_0}) / K_G^{v_0} \mid g_1^{-1} g_2 \in K_G g K_G\}$$

with the two obvious projection maps, and the diagonal action by $G'(F_0)$ from the left multiplication. Then the fiber of the Hecke correspondence $\widehat{\mathrm{Hk}}_{[K_G g K_G]}$ over any fixed element of $Z^{\mathbb{Q}}(\mathbb{Q}) \backslash (Z^{\mathbb{Q}}(\mathbb{A}_f) / K_{Z^{\mathbb{Q}}})$ (cf. (7.6)) can be identified with

$$\begin{array}{ccc} \widehat{\mathrm{Hk}}_{[K_G g K_G], 0} & \xrightarrow{\sim} & G'(F_0) \backslash \left[\mathcal{N}_{O_{\check{E}_v}} \times \mathrm{Hk}_{[K_G g K_G]}^{(v_0)} \right] \\ \downarrow & & \downarrow \\ \left(\mathcal{M}_{O_{\check{E}_v}, 0} \right)^2 & \xrightarrow{\sim} & \left(G'(F_0) \backslash \left[\mathcal{N}_{O_{\check{E}_v}} \times G(\mathbb{A}_{0,f}^{v_0}) / K_G^{v_0} \right] \right)^2, \end{array}$$

where the right vertical map is induced by the diagonal $\mathcal{N}_{O_{\check{E}_v}} \rightarrow \mathcal{N}_{O_{\check{E}_v}} \times \mathcal{N}_{O_{\check{E}_v}}$ and the two projection maps from $\mathrm{Hk}_{[K_G g K_G]}^{(v_0)}$.

7.8. *CM cycles in the formal neighborhood of the basic locus.* We now consider the restriction of the fat big CM cycle and its derived version to the formal neighborhood of the basic locus at a non-archimedean place $v_0 \nmid \mathfrak{d}$ of F_0 , *inert* in F , via the RZ uniformization (7.4). We resume the notation there.

Let $\alpha \in \mathcal{A}_n(O_{F_0}[1/\mathfrak{d}])$ be irreducible over F . We denote by $\mathcal{CM}^\wedge(\alpha)$ (resp. $\mathcal{CM}^\wedge(\alpha, g)$) the formal completion along the basic locus of the CM cycle $\mathcal{CM}(\alpha)$ (resp. $\mathcal{CM}(\alpha, g)$ for $g \in G(F_{0,\mathfrak{d}})$). We denote the derived CM cycle

$${}^L\mathcal{CM}^\wedge(\alpha, g) \in K'_0(\mathcal{CM}^\wedge(\alpha, g)),$$

and for $\phi_0 = \mathbf{1}_{K_G^\mathfrak{d}} \otimes \phi_{\mathfrak{d}} \in \mathcal{S}(G(\mathbb{A}_{0,f}), K_G)$,

$${}^L\mathcal{CM}^\wedge(\alpha, \phi_0) \in \bigoplus_{g \in K_{G,\mathfrak{d}} \setminus G(F_{0,\mathfrak{d}})/K_{G,\mathfrak{d}}} K'_0(\mathcal{CM}^\wedge(\alpha, g)).$$

For $\delta \in G'(F_{0,v_0})$, let \mathcal{N}^δ be the fixed point locus of δ on the RZ space \mathcal{N} for $F_{w_0}/F_{0,v_0}$ (cf. Section 3.1), and let $\mathcal{N}_{O_{\check{E}_\nu}}^\delta$ be its base change to $O_{\check{E}_\nu}$. For $(\delta, h) \in G'(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$, we define a closed formal subscheme of $\mathcal{N}_{O_{\check{E}_\nu}} \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$:

$$(7.18) \quad \mathcal{CM}(\delta, h)_{K_G^{v_0}} = \mathcal{N}_{O_{\check{E}_\nu}}^\delta \times \mathbf{1}_{hK_G^{v_0}}.$$

We consider the sum

$$(7.19) \quad \sum \mathcal{CM}(\delta', h')$$

over all $(\delta', h') \in G'(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$ in the $G'(F_0)$ -orbit of (δ, h) . Here $G'(F_0)$ acts diagonally on $G'(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$ by $g \cdot (\delta, h) = (g\delta g^{-1}, gh)$. The sum is $G'(F_0)$ -invariant and hence descends to the quotient formal scheme (7.6), which we denote by $[\mathcal{CM}(\delta, h)]_{K_G^{v_0}}$.

Furthermore, we have a derived version of (7.18) and (7.19) by replacing the naive fixed point locus $\mathcal{N}_{O_{\check{E}_\nu}}^\delta$ in (7.18) by the derived fixed point locus ${}^L\mathcal{N}_{O_{\check{E}_\nu}}^\delta$ defined by (3.12).

We then have an analog of Proposition 7.4.

PROPOSITION 7.17. *Let $\alpha \in \mathcal{A}_n(O_{F_0}[1/\mathfrak{d}])$ be irreducible over F .*

(i) *The restriction of $\mathcal{CM}^\wedge(\alpha)$ to each fiber of the projection (7.5) is the disjoint union*

$$\coprod_{(\delta, h)} [\mathcal{CM}(\delta, h)]_{K_G^{v_0}},$$

where the index runs over the set

$$\{(\delta, h) \in G'(F_0) \setminus (G'(\alpha)(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}) \mid h^{-1}\delta h \in K_G^{v_0}\}.$$

(ii) Let $\phi_0 = 1_{K_G^v} \otimes \phi_v \in \mathcal{S}(G(\mathbb{A}_{0,f}), K_G)$, where $\phi_v \in \mathcal{S}(G(F_{0,v}), K_{G,v})$. The restriction of ${}^L\mathcal{CM}(\alpha, \phi_0)$ in (7.16) to each fiber of the projection (7.5) is the sum

$$\sum_{(\delta, h) \in G'(F_0) \setminus (G'(\alpha)(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0})} \phi_0^{v_0}(h^{-1}\delta h) \cdot [{}^L\mathcal{CM}(\delta, h)]_{K_G^{v_0}},$$

as an element in the group $\bigoplus_{g \in K_{G,v} \setminus G(F_{0,v})/K_{G,v}} K'_0(\mathcal{CM}(\alpha, g))$.

Remark 7.18. One can define an analog of the cycle ${}^L\mathcal{CM}(\alpha, \phi_0)$ on a semi-global integral model (i.e., over the localization $O_{E,(\nu)}$ of O_E at a place ν above v_0 ; cf. [40, §4]). Then we can allow more general level structure $K_G^{v_0}$ away from v_0 , and therefore allow $\phi_0 = 1_{K_{G,v_0}} \otimes \phi^{v_0} \in \mathcal{S}(G(\mathbb{A}_{0,f}^{v_0}), K_G^{v_0})$.

Proof. We only prove part (i); part (ii) concerning the derived version follows along the same line.

Over the formal scheme (7.6), $\mathcal{CM}(\alpha)$ consists of $G'(F_0)$ -cosets of $(X, hK_G^{v_0}) \in \mathcal{N}_{O_{E,\nu}} \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$ together with an isomorphism $\varphi_{v_0} : X \rightarrow X$ and $g \in G(\mathbb{A}_{0,f}^{v_0})$, satisfying the following conditions: there exists $\delta \in G'(F_0)$ such that the endomorphism of the framing object \mathbb{X}_n induced by φ_{v_0} is δ , and both g and δ fix $hK_G^{v_0}$ and induce the same automorphism of $hK_G^{v_0}$; the polynomial α annihilates g and φ_{v_0} (or equivalently δ by the rigidity of quasi-isogeny). In particular, $\delta \in G'(\alpha)(F_0)$ by the irreducibility of α .

Here we view $G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$ as a groupoid in which the automorphism group of $hK_G^{v_0}$ is isomorphic to $hK_G^{v_0}h^{-1}$. If both δ and g fix $hK_G^{v_0}$ and induce the same automorphism of $hK_G^{v_0}$, then $g = \delta$ (“rigidity away from v_0 ”). It follows that the condition g fixing $hK_G^{v_0}$ is equivalent to $\delta hK_G^{v_0} = hK_G^{v_0}$, i.e., $h^{-1}\delta h \in K_G^{v_0}$.

The condition on the existence of a quasi-isogeny φ_{v_0} lifting δ amounts to $X \in \mathcal{N}_{O_{E,\nu}}^\delta$.

Therefore, for a fixed $\delta \in G'(\alpha)(F_0)$, the desired pairs $(X, hK_G^{v_0})$ are exactly those lying on $\mathcal{N}_{O_{E,\nu}}^\delta \times 1_{hK_G^{v_0}}$ subject to the condition $h^{-1}\delta h \in K_G^{v_0}$. Then it remains to sum over all $\delta \in G'(\alpha)(F_0)$ to complete the proof of part (i). \square

8. Modular generating functions of special divisors

In this section we collect a few modularity results due to various authors for the generating functions of special divisors with values in Chow groups, and in a reduced version of arithmetic Chow groups.

8.1. Generating functions of special divisors on $M_{K_{\bar{G}}}(\widetilde{G})$. We first define the generating functions of special divisors on the canonical model $M_{K_{\bar{G}}}(\widetilde{G})$

over $\mathrm{Spec} E$. The moduli functor is introduced at the end of [Section 6](#) for an arbitrary compact open subgroup $K_{\tilde{G}}$ of the form $K_{\tilde{G}} = K_{Z\mathbb{Q}} \times K_G$. For $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))^{K_G}$ and $\xi \in F_{0,+}$, we have defined the divisor $Z(\xi, \phi) \in \mathrm{Ch}^1(M_{K_{\tilde{G}}}(\tilde{G}))$ by [\(7.1\)](#). When $\xi = 0$, we define

$$(8.1) \quad Z(0, \phi) = -\phi(0) c_1(\omega) \in \mathrm{Ch}^1(M_{K_{\tilde{G}}}(\tilde{G})),$$

where ω is the automorphic line bundle [\[22\]](#) and c_1 denotes the first Chern class.

In [Section 11.1](#) we will recall the Weil representation ω of $\mathbf{H}(\mathbb{A}_{0,f})$ on $\mathcal{S}(V(\mathbb{A}_{0,f}))^{K_G}$. We define the generating function on $\mathbf{H}(\mathbb{A}_0)$ by

$$(8.2) \quad Z(h, \phi) = Z(0, \omega(h_f)\phi) W_0^{(n)}(h_\infty) + \sum_{\xi \in F_{0,+}} Z(\xi, \omega(h_f)\phi) W_\xi^{(n)}(h_\infty),$$

where $h = (h_\infty, h_f) \in \mathbf{H}(\mathbb{A}_0)$, $h_\infty = (h_v)_{v|\infty} \in \prod_{v|\infty} \mathrm{SL}_2(F_{0,v})$ and

$$W_\xi^{(n)}(h_\infty) = \prod_{v|\infty} W_\xi^{(n)}(h_v);$$

cf. [\(1.5\)](#) for the weight n Whittaker function $W_\xi^{(n)}$ on $\mathrm{SL}_2(\mathbb{R})$.

THEOREM 8.1. *The generating function $Z(h, \phi)$ lies in*

$$\mathcal{A}_{\mathrm{hol}}(\mathbf{H}(\mathbb{A}_0), K, n)_{\overline{\mathbb{Q}}} \bigotimes_{\overline{\mathbb{Q}}} \mathrm{Ch}^1(M_{K_{\tilde{G}}}(\tilde{G}))_{\overline{\mathbb{Q}}},$$

where $K \subset \mathbf{H}(\mathbb{A}_{0,f})$ is a compact open subgroup that fixes $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))$ under the Weil representation.

For the definition of the vector space $\mathcal{A}_{\mathrm{hol}}(\mathbf{H}(\mathbb{A}_0), K, n)_{\overline{\mathbb{Q}}}$, we refer to [\(1.10\)](#) (and [\(1.9\)](#)). One can replace the field $\overline{\mathbb{Q}}$ by a number field, but it will not be more useful in this paper.

The result has an analog for orthogonal Shimura varieties, which is due to Borcherds when $F_0 = \mathbb{Q}$ (generalizing Gross–Kohnen–Zagier theorem), and [\[44\]](#) for totally real fields F_0 ; Bruinier also gave a proof in [\[4\]](#), where he also constructed the automorphic Green function we will use later. By the embedding trick [\[28, §3.2, Lem. 3.6\]](#), this result implies the analogous modularity for Shimura varieties $\mathrm{Sh}_{K_G}(\mathrm{Res}_{F_0/\mathbb{Q}} G, \{h_G\})$.⁶ Then the assertion in the theorem above follows from the fact that, after base change to \mathbb{C} , $M_{K_{\tilde{G}}}(\tilde{G})$ is a disjoint union of copies of $\mathrm{Sh}_{K_G}(\mathrm{Res}_{F_0/\mathbb{Q}} G, \{h_G\})$; cf. [\(6.6\)](#).

⁶In the unitary case, one expects to obtain a $\mathrm{U}(1, 1)$ -automorphic form. However, the SL_2 -automorphic form suffices for our purpose, and in fact the extra information in $\mathrm{U}(1, 1)$ -modularity is not useful for us at all because the analytic side only has SL_2 -modularity.

8.2. *Complex uniformization of special divisors.* We now study special divisors over the complex numbers. The situation is analogous to the description of the special divisors in the formal neighborhood of the basic locus; cf. Section 7.3.

We start with the complex uniformization of our Shimura varieties. This is very much similar to (7.4). Let $\nu : E \hookrightarrow \mathbb{C}$ be a complex place of the reflex field E . Its restriction to F (resp. F_0) is a place denoted by w_0 (resp. v_0). Let $M_{\nu, \mathbb{C}} = M_{K_{\widetilde{G}}}(\widetilde{G}) \otimes_{E, \nu} \mathbb{C}$ be the complex orbifold via ν . Let V' be the “nearby” hermitian space, i.e., the unique one that is positive definite at all archimedean places except v_0 where the signature is $(n-1, 1)$, and isomorphic to V locally at all non-archimedean places. Then let G' be the unitary group (viewed as a \mathbb{Q} -algebraic group) associated to V' . Let \mathcal{D}_{v_0} be the Grassmannian of negative definite \mathbb{C} -lines in $V' \otimes_{F, w_0} \mathbb{C}$. Then we have a complex uniformization

$$(8.3) \quad M_{\nu, \mathbb{C}} = \widetilde{G}'(\mathbb{Q}) \backslash \left[\mathcal{D}_{v_0} \times \widetilde{G}(\mathbb{A}_f) / K_{\widetilde{G}} \right].$$

When the embedding $\nu : E \hookrightarrow \mathbb{C}$ is the natural one for the reflex field E (recall that it is a subfield of \mathbb{C}), the uniformization is [40, Rem. 3.2, Prop. 3.5], and in general it follows from the proof of loc. cit.

Analogous to (7.5), we have a partition by the projection

$$(8.4) \quad M_{\nu, \mathbb{C}} \longrightarrow Z^{\mathbb{Q}}(\mathbb{Q}) \backslash (Z^{\mathbb{Q}}(\mathbb{A}_f) / K_{Z^{\mathbb{Q}}}),$$

where each fiber is naturally isomorphic to

$$(8.5) \quad M_{\nu, \mathbb{C}, 0} := G'(F_0) \backslash \left[\mathcal{D}_{v_0} \times G(\mathbb{A}_{0, f}) / K_G \right].$$

Here we fix an isomorphism $G'(\mathbb{A}_{0, f}) \simeq G(\mathbb{A}_{0, f})$.

Now we return to describe the complex uniformization of the special divisors. For each $u \in V'(F_0)$ with totally positive norm, let $\mathcal{D}_{v_0, u} \subset \mathcal{D}_{v_0}$ be the space of negative definite \mathbb{C} -lines perpendicular to u .⁷ For a pair $(u, g) \in V'(F_0) \times G(\mathbb{A}_{0, f}) / K_G$, we define

$$(8.6) \quad Z(u, g)_{K_G} = \mathcal{D}_{v_0, u} \times \mathbf{1}_{g K_G}.$$

We consider the sum

$$(8.7) \quad \sum Z(u', g')_{K_G},$$

over (u', g') in the $G'(F_0)$ -orbit of the pair (u, g) for the diagonal action of $G'(F_0)$ on $V'(F_0) \times G(\mathbb{A}_{0, f}) / K_G$. The sum is $G'(F_0)$ -invariant and hence descends to a divisor on the quotient (8.5), denoted by $[Z(u, g)]_{K_G}$.

⁷The codimension one analytic space $\mathcal{D}_{v_0, u}$ on \mathcal{D}_{v_0} is the archimedean analog of the local KR divisor $\mathcal{Z}(u)$ on \mathcal{N} in Section 7.3.

Then, we have an archimedean analog of [Proposition 7.4](#) for the special divisor $Z(\xi, \phi)$ defined by [\(7.1\)](#). In the case of $F_0 = \mathbb{Q}$ and a special level structure, this is proved in [\[25, §3.3\]](#); in general, the proof in loc. cit. works verbatim and hence we omit the detail.

PROPOSITION 8.2. *Let $\xi \in F_{0,+}$. Then the restriction of the special divisor $Z(\xi, \phi) \otimes_{E,\nu} \mathbb{C}$ to each fiber of the projection [\(7.5\)](#) is*

$$(8.8) \quad \sum_{(u,g) \in G'(F_0) \setminus (V'_\xi(F_0) \times G(\mathbb{A}_{0,f}) / K_G)} \phi(g^{-1}u) \cdot [Z(u, g)]_{K_G}.$$

Remark 8.3. We may rewrite the above result into a form that has appeared in the formula of special divisors in [\[44, §1\]](#). Let $G'_u \subset G'$ be the stabilizer of u under the action of G' on V' , viewed as an algebraic group over F_0 . Instead of [\(8.6\)](#), we define

$$\tilde{Z}(u, g)_{K_G} := \mathcal{D}_{v_0, u} \times 1_{G'_u(\mathbb{A}_{0,f}) g K_G}.$$

Similarly, we denote its image in the quotient [\(8.5\)](#) by $[\tilde{Z}(u, g)]_{K_G}$. Then we may rewrite the sum as [\(8.8\)](#):

$$\sum_{u \in G'(F_0) \setminus V'_\xi(F_0)} \sum_{g \in G'_u(\mathbb{A}_{0,f}) \setminus G(\mathbb{A}_{0,f}) / K_G} \phi(g^{-1}u) \cdot [\tilde{Z}(u, g)]_{K_G}.$$

This is exactly the formula in loc. cit.

8.3. Green functions. We recall the Green functions of Kudla [\[23\]](#) and the automorphic Green functions (cf. [\[36\]](#), [\[4\]](#)). The former is more convenient when comparing with the analytic side, while the latter is more suitable for proving (holomorphic) modularity of generating series. The difference between them is studied by Ehlen–Sankaran in [\[8\]](#) when $F_0 = \mathbb{Q}$.

We first recall Kudla's Green functions, defined for the orthogonal case in [\[23\]](#), which can be carried over easily to the unitary case (cf. [\[28, §4B\]](#)). Let $u \in V'(F_0)$ be as in the previous subsection. Let $z \in \mathcal{D}_{v_0}$. Let u_z be the orthogonal projection to the negative definite \mathbb{C} -line z of $V' \otimes_{F, w_0} \mathbb{C}$. Define

$$(8.9) \quad R(u, z) = \langle u_z, u_z \rangle = \frac{\langle u, \tilde{z} \rangle^2}{\langle \tilde{z}, \tilde{z} \rangle},$$

where \tilde{z} is any \mathbb{C} -basis of the line z .

We will need the exponential integral defined by

$$(8.10) \quad \text{Ei}(-r) = - \int_r^\infty \frac{e^{-t}}{t} dt, \quad r > 0.$$

This function has a logarithmic singularity around 0; more precisely, when $r \rightarrow 0^+$,

$$\text{Ei}(-r) = \gamma + \log r + \sum_{n=1}^{\infty} \frac{(-r)^n}{n \cdot n!}.$$

Here γ is the Euler constant.

Let $h_\infty = (h_v)_{v|\infty} \in \prod_{v|\infty} \text{SL}_2(F_{0,v})$ and $h_v = \begin{pmatrix} 1 & b_v \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \sqrt{a_v} \\ 1/\sqrt{a_v} \end{pmatrix} \kappa_v$ in the Iwasawa decomposition; cf. (1.6). For each *non-zero* vector $u \in V(F_0)$, Kudla [23] defined a Green function on \mathcal{D}_{v_0} , parametrized by h_∞ :

$$(8.11) \quad \mathcal{G}^{\mathbf{K}}(u, h_\infty)(z) = -\text{Ei}(2\pi a_{v_0} R(u, z)), \quad z \in \mathcal{D}_{v_0} \setminus \mathcal{D}_{v_0, u}.$$

It has logarithmic singularity along the divisor $\mathcal{D}_{v_0, u}$. Note that this is defined for *every* non-zero vector $u \in V'(F_0)$; in particular, u may have null-norm. If $\mathcal{D}_{v_0, u}$ is empty, the function is then smooth on \mathcal{D}_{v_0} . When $u = 0$, we set

$$(8.12) \quad \mathcal{G}^{\mathbf{K}}(0, h_\infty) = -\log |a_{v_0}|.$$

Now we descend the Green function on \mathcal{D}_{v_0} to the quotient (8.5): for all $\xi \in F_0$, define

$$(8.13) \quad \mathcal{G}^{\mathbf{K}}(\xi, h_\infty, \phi) = \sum \phi(g^{-1}u) \cdot (\mathcal{G}^{\mathbf{K}}(u, h_\infty) \times 1_{gK_G}),$$

where the sum is over $(u, g) \in V'_\xi(F_0) \times G(\mathbb{A}_{0,f})/K_G$. This defines a Green function for the divisor $Z(\xi, \phi)$; cf. [28, Prop. 4.9].

We now recall the automorphic Green function [36], [4], [5]. Since the role of those are indirect to this paper, we just say that there is a Green function $\mathcal{G}^{\mathbf{B}}(\xi, \phi)$ for each $\xi \in F_{0,+}$, and $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))$; cf. [5, §7.3].

We define the generating function of the difference of the two Green functions

$$(8.14) \quad \mathcal{Z}_{v_0, \text{corr}}(h, \phi) := \sum_{\xi \in F_0} (\mathcal{G}^{\mathbf{K}}(\xi, h_\infty, \omega(h_f)\phi) - \mathcal{G}^{\mathbf{B}}(\xi, \omega(h_f)\phi)) W_\xi^{(n)}(h_\infty),$$

where the notation is the same as in (8.2). We note that this definition depends on the archimedean place v_0 of F_0 , though it is omitted in the right-hand side of the equality.

The following theorem is due to Ehlen–Sankaran [8].

THEOREM 8.4. *Assume $F_0 = \mathbb{Q}$. The generating function $\mathcal{Z}_{\infty, \text{corr}}(h, \phi)$ lies in the space $\mathcal{A}_{\text{exp}}(\mathbf{H}(\mathbb{A}_0), K, n)$, in the sense that, for every point $[z, g] \in M_{\nu, \mathbb{C}}$, the value of the generating functions at $[z, g]$ lies in $\mathcal{A}_{\text{exp}}(\mathbf{H}(\mathbb{A}_0), K, n)$. Here $K \subset \mathbf{H}(\mathbb{A}_{0,f})$ is a compact open subgroup that fixes $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))$ under the Weil representation.*

Proof. In [8, Th. 3.6], the authors proved the assertion for orthogonal groups, from which the case of unitary groups follows (e.g, by [8, proof of Th. 4.13, p. 2131]). \square

8.4. *Modularity in the arithmetic Chow group $\widehat{\text{Ch}}_0^1(\mathcal{M})$.* We will use the Gillet–Soulé arithmetic intersection theory; cf. [12], [10]. (In the non-proper case, cf. [7].) We first recall the arithmetic Chow group $\widehat{\text{Ch}}^1(\mathcal{M})$ (with \mathbb{Q} -coefficient) for a regular flat scheme (possibly non-proper) $\mathcal{M} \rightarrow \text{Spec } O_E$. Elements are represented by arithmetic divisors, i.e., \mathbb{Q} -linear combinations of tuples $(Z, (g_{Z,w})_{w \in \text{Hom}_{\mathbb{Q}}(E, \overline{\mathbb{Q}})})$, where Z is a divisor on \mathcal{M} and $g_{Z,w}$ is a Green function of $Z_w(\mathbb{C})$ on the complex manifold $\mathcal{M}_w(\mathbb{C})$ via the embedding $w : E \hookrightarrow \overline{\mathbb{Q}} \subset \mathbb{C}$ (cf. [12, §3.3]). Principal arithmetic divisors are tuples associated to rational functions $f \in E(\mathcal{M})^\times$:

$$(\text{div}(f), (-\log |f|_w^2)_{w \in \text{Hom}_{\mathbb{Q}}(E, \overline{\mathbb{Q}})}).$$

(For example, when $E = \mathbb{Q}$, we have $\mathbf{V}_p = (0, 2\log |p|)$ in $\widehat{\text{Ch}}^1(\mathcal{M})$, where \mathbf{V}_p is the fiber of \mathcal{M} over a prime p .)

Now it is clear we can extend the same definition to a regular flat scheme $\mathcal{M} \rightarrow \text{Spec } O_E \setminus S$ for a finite set S of *non-archimedean* places. We still denote it by $\widehat{\text{Ch}}^1(\mathcal{M})$.

Remark 8.5. If we start with a regular flat scheme $\mathcal{M} \rightarrow \text{Spec } O_E$, and a finite set S , then two groups $\widehat{\text{Ch}}^1(\mathcal{M})$ and $\widehat{\text{Ch}}^1(\mathcal{M}^S)$ for $\mathcal{M}^S = \mathcal{M} \times_{\text{Spec } O_E} \text{Spec } O_E \setminus S$ are related as follows. We denote by $\text{Ch}_{|S|}^1(\mathcal{M})$ the subgroup of $\widehat{\text{Ch}}^1(\mathcal{M})$ consisting of elements supported at the fibers above $\nu \in S$. This is a finite dimensional vector space. Then there is a natural isomorphism

$$\widehat{\text{Ch}}^1(\mathcal{M}) / \text{Ch}_{|S|}^1(\mathcal{M}) \xrightarrow{\sim} \widehat{\text{Ch}}^1(\mathcal{M}^S).$$

Now we specialize to our interest, the moduli space $\mathcal{M} = \mathcal{M}_{K_{\widetilde{G}}}(\widetilde{G})$ introduced in [Definition 6.1](#). Let S be the set of places $\nu \mid \mathfrak{d}$. Recall that the morphism $\mathcal{M}_{K_{\widetilde{G}}}(\widetilde{G}) \rightarrow \text{Spec } O_E[1/\mathfrak{d}] = \text{Spec } O_E \setminus S$ is smooth.

Let $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))^{K_G}$ be of the form $\phi = 1_{\Lambda^0} \otimes \phi_0$; cf. [\(7.2\)](#). For $\xi \in F_{0,+}$, we endow the special divisor $\mathcal{Z}(\xi, \phi)$ (cf. [\(7.2\)](#)) with the automorphic Green function $\mathcal{G}^B(\xi, \phi)$. Denote by $\widehat{\mathcal{Z}}^B(\xi, \phi)$ the resulting element in $\widehat{\text{Ch}}^1(\mathcal{M})$. When $\xi = 0$, we define

$$(8.15) \quad \mathcal{Z}^B(0, \phi) = -\phi(0) c_1(\widehat{\omega}) \in \widehat{\text{Ch}}^1(\mathcal{M}),$$

where $\widehat{\omega} = (\omega, \|\cdot\|_{\text{Pet}})$ is the extension of the automorphic line bundle ω to the integral model \mathcal{M} , endowed with its Petersson metric [\[5, §7.2\]](#).

We define the generating series with coefficients in the arithmetic Chow group $\widehat{\text{Ch}}^1(\mathcal{M})$,

$$(8.16) \quad \widehat{\mathcal{Z}}^{\mathbf{B}}(\tau, \phi) = \sum_{\xi \in F_0, \xi \geq 0} \widehat{\mathcal{Z}}^{\mathbf{B}}(\xi, \phi) q^{\xi},$$

where

$$(8.17) \quad \tau = (\tau_v)_{v \mid \infty} \in \prod_{v \mid \infty} \mathcal{H}, \quad q^{\xi} := e^{2\pi i \text{tr}_{F_0/\mathbb{Q}}(\tau \xi)}.$$

The following theorem can be deduced from [5].

THEOREM 8.6 (Bruinier–Kudla–Howard–Rapoport–Yang). *Let $F_0 = \mathbb{Q}$. The generating series $\widehat{\mathcal{Z}}^{\mathbf{B}}(\cdot, \phi)$ lies in $\mathcal{A}_{\text{hol}}(\Gamma(N), n)_{\overline{\mathbb{Q}}} \otimes_{\overline{\mathbb{Q}}} \widehat{\text{Ch}}^1(\mathcal{M})_{\overline{\mathbb{Q}}}$, where N depends only on ϕ and all prime factors of N are contained in S .*

Proof. In [5] the authors proved a stronger version (i.e., Theorem B in loc. cit.) in a maximal level case (with principle polarization) over the full ring of integers of E . Since the arithmetic Chow group of \mathcal{M} considered here omits a finite set of bad places S (including primes ramified in F), the computation of divisors of the regularized theta lifts and Borcherds product on the integral models over $\text{Spec } O_E[1/\mathfrak{d}]$ of loc. cit. still applies to our (even simpler) situation. \square

9. Local intersection: non-archimedean places

9.1. Arithmetic intersection theory. We first recall an arithmetic intersection pairing on a pure dimensional flat (not necessarily proper) morphism $\mathcal{M} \rightarrow \mathcal{B} = \text{Spec } O_E$ of regular schemes with smooth generic fiber. Let $\widetilde{\mathcal{Z}}_{1,c}(\mathcal{M})$ be the group of proper (*over the base \mathcal{B}*) 1-cycles on \mathcal{M} (with \mathbb{Q} -coefficient). Then there is an arithmetic intersection pairing between two \mathbb{Q} -vector spaces (cf. [3, §2.3] when the ambient scheme is proper)

$$(9.1) \quad (\cdot, \cdot) : \widehat{\text{Ch}}^1(\mathcal{M}) \times \widetilde{\mathcal{Z}}_{1,c}(\mathcal{M}) \longrightarrow \mathbb{R}.$$

Now let S be a finite set of non-archimedean places of E , and let S_p be the subset of places above p . Let $\mathcal{M} \rightarrow \text{Spec } O_E \setminus S$, and let $\widehat{\text{Ch}}^1(\mathcal{M})$ be its arithmetic Chow group defined in Section 8.4. Consider the quotient of \mathbb{R} by a finite dimensional \mathbb{Q} -vector space,

$$(9.2) \quad \mathbb{R}_S := \mathbb{R} / \text{span}_{\mathbb{Q}}\{\log p : \#S_p \neq 0\},$$

which is an (infinite dimensional) \mathbb{Q} -vector space. Then the definition of [3] works directly if we replace the base $\text{Spec } O_E$ by $\text{Spec } O_E \setminus S$ (i.e., without an

integral model over the full ring of integers O_E), and it yields a pairing with values in \mathbb{R}_S :

$$(9.3) \quad (\cdot, \cdot) : \widehat{\text{Ch}}^1(\mathcal{M}) \times \widetilde{\mathcal{Z}}_{1,c}(\mathcal{M}) \longrightarrow \mathbb{R}_S.$$

Note that the cycles in $\widetilde{\mathcal{Z}}_{1,c}(\mathcal{M})$ are assumed to be *proper* over $\text{Spec } O_E \setminus S$.

We note that by [3, Prop. 2.3.1(ii)], for cycles in $\widetilde{\mathcal{Z}}_{1,c}(\mathcal{M})$ supported on special fibers, the pairing only depends on their rational equivalence classes. This motivates us to define a quotient group $\mathcal{Z}_{1,c}(\mathcal{M})$ of $\widetilde{\mathcal{Z}}_{1,c}(\mathcal{M})$ by the subgroup generated by 1-cycles that are supported on *proper* subschemes Y of the special fibers and are rationally equivalent to zero on Y . We have the resulting pairing

$$(9.4) \quad (\cdot, \cdot) : \widehat{\text{Ch}}^1(\mathcal{M}) \times \mathcal{Z}_{1,c}(\mathcal{M}) \longrightarrow \mathbb{R}_S.$$

We now let $\mathcal{M} = \mathcal{M}_{K_{\bar{G}}}(\widetilde{G})$ be the moduli stack introduced in [Definition 6.1](#). We apply the above pairing to $\mathcal{M} = \mathcal{M}_{K_{\bar{G}}}(\widetilde{G}) \rightarrow \mathcal{B} = \text{Spec } O_E \setminus S$ for any finite set S containing all places $\nu \mid \mathfrak{d}$. We define an element in $\mathcal{Z}_{1,c}(\mathcal{M})$ starting from the derived CM cycle ${}^L\mathcal{CM}(\alpha, g)$ ([7.14](#)), which is an element in $F_1 K'_0(\mathcal{CM}(\alpha, g))$, ([7.15](#)). The finite morphism $\mathcal{CM}(\alpha, g) \rightarrow \mathcal{M}$ induces a homomorphism

$$K'_0(\mathcal{CM}(\alpha, g)) \longrightarrow K'_{0, \mathcal{CM}(\alpha, g)}(\mathcal{M})$$

preserving the respective filtrations, where $K'_{0, \mathcal{CM}(\alpha, g)}(\mathcal{M})$ denotes the K -group of coherent sheaves with support on the image of $\mathcal{CM}(\alpha, g)$. Since $\mathcal{CM}(\alpha, g) \rightarrow \mathcal{B}$ is proper and the generic fiber of $\mathcal{CM}(\alpha, g)$ is zero dimensional (cf., [Proposition 7.9\(b\)](#)), there is a natural homomorphism $\text{Ch}_{1, \mathcal{CM}(\alpha, g)}(\mathcal{M}) \rightarrow \mathcal{Z}_{1,c}(\mathcal{M})$. We now consider the composition

$$F_1 K'_0(\mathcal{CM}(\alpha, g)) \longrightarrow \text{Gr}_1 K'_{0, \mathcal{CM}(\alpha, g)}(\mathcal{M}) \xrightarrow{\sim} \text{Ch}_{1, \mathcal{CM}(\alpha, g)}(\mathcal{M}) \longrightarrow \mathcal{Z}_{1,c}(\mathcal{M}),$$

where the isomorphism in the middle is [[11](#), Th. 8.2] and Gr_1 denotes the grading F_1/F_0 . By abuse of notation, we still denote by ${}^L\mathcal{CM}(\alpha, g)$ the image in $\mathcal{Z}_{1,c}(\mathcal{M})$ of the element ${}^L\mathcal{CM}(\alpha, g) \in F_1 K'_0(\mathcal{CM}(\alpha, g))$ (cf. ([7.15](#))) under the above composition.

9.2. Intersection of special divisors and CM cycles. For the rest of the article, we let $\Phi = \otimes_{v_0} \Phi_{v_0} \in \mathcal{S}((G \times V)(\mathbb{A}_{0,f}))$ be of the form $\phi_0 \otimes \phi$, where

- $\phi_0 = 1_{K_G^{\mathfrak{d}}} \otimes \phi_{0,\mathfrak{d}}$ and $\phi_{0,\mathfrak{d}} \in \mathcal{S}(G(F_{0,\mathfrak{d}}), K_{G,\mathfrak{d}})$ (cf. ([7.16](#))); and
- $\phi = 1_{\Lambda^{\mathfrak{d}}} \otimes \phi_{\mathfrak{d}}$ and $\phi_{\mathfrak{d}} \in \mathcal{S}(V(F_{0,\mathfrak{d}}))^{K_{G,\mathfrak{d}}}$ (cf. ([7.2](#))).

Recall from [Section 7.4](#) that we have also fixed a conjugate self-reciprocal polynomial $\alpha \in O_F[1/\mathfrak{d}][T]_{\deg=n}$, irreducible over F . We define a generating

series using the intersection pairing (9.4),

$$(9.5) \quad \text{Int}(\tau, \Phi) := \frac{1}{\tau(Z^{\mathbb{Q}}) \cdot [E : F]} \left(\widehat{\mathcal{Z}}^{\mathbf{B}}(\tau, \phi), \quad {}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0) \right),$$

where $\mathcal{Z}^{\mathbf{B}}(\tau, \phi)$ is (8.16), and

$$(9.6) \quad \tau(Z^{\mathbb{Q}}) := \#Z^{\mathbb{Q}}(\mathbb{Q}) \setminus (Z^{\mathbb{Q}}(\mathbb{A}_f)/K_{Z^{\mathbb{Q}}}).$$

Remark 9.1. By Theorem 8.6, when $F_0 = \mathbb{Q}$, this is a holomorphic modular form (of weight n , and level depending only on ϕ) with coefficients in the $\overline{\mathbb{Q}}$ -vector space $\mathbb{R}_{S, \overline{\mathbb{Q}}} := \mathbb{R}_S \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$, i.e.,

$$(9.7) \quad \text{Int}(\cdot, \Phi) \in \mathcal{A}_{\text{hol}}(\Gamma(N), n)_{\overline{\mathbb{Q}}} \otimes_{\overline{\mathbb{Q}}} \mathbb{R}_{S, \overline{\mathbb{Q}}}.$$

Our results in this and the next section are still valid for general totally real fields F_0 since they do not use the modularity.

Similarly, for each $\xi \in F_{0,+}$, we define

$$(9.8) \quad \text{Int}(\xi, \Phi) := \frac{1}{\tau(Z^{\mathbb{Q}}) \cdot [E : F]} \left(\widehat{\mathcal{Z}}^{\mathbf{B}}(\xi, \phi), \quad {}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0) \right).$$

When $\xi = 0$, this is by definition

$$(9.9) \quad \text{Int}(0, \Phi) = -\frac{1}{\tau(Z^{\mathbb{Q}}) \cdot [E : F]} \left(\widehat{\omega}, \quad {}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0) \right) \phi(0).$$

Then by (8.16),

$$(9.10) \quad \text{Int}(\tau, \Phi) = \sum_{\xi \in F_0, \xi \geq 0} \text{Int}(\xi, \Phi) q^{\xi}.$$

Now let $\xi \neq 0$. We will express the arithmetic intersection number (9.8) in terms of the local intersection numbers from the AFL over good places and the archimedean local intersection.

9.3. The support of the intersection. We first study the intersection of the special divisor $\mathcal{Z}(\xi, \phi)$ and the CM cycle ${}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0)$. First we have the following analog to [40, Th. 8.5].

THEOREM 9.2. *Let $\xi \neq 0$ and $\Phi = \otimes_{v_0} \Phi_{v_0} \in \mathcal{S}((G \times V)(\mathbb{A}_{0,f}))^{K_G}$. Let S be a finite set of places containing all places $\nu \mid \mathfrak{d}$ and such that at $v_0 \notin S$, $\Phi_{v_0} = \mathbf{1}_{K_{G,v_0}^{\circ}} \otimes \mathbf{1}_{\Lambda_{v_0}^{\circ}}$.*

Then the following statements on the support of the intersection of the special divisor $\mathcal{Z}(\xi, \phi)$ and the CM cycle $\mathcal{CM}(\alpha, \phi_0)$ on \mathcal{M} hold:

- (i) *The support does not meet the generic fiber.*
- (ii) *Let $\nu \notin S$ be a place of E lying over a place of F_0 that splits in F . Then the support does not meet the special fiber $\mathcal{M} \otimes_{O_E} \kappa_{\nu}$.*

(iii) Let $\nu \notin S$ be a place of E lying over a place of F_0 that does not split in F . Then the support meets the special fiber $\mathcal{M} \otimes_{O_E} \kappa_\nu$ only in its basic locus.

Proof. The proof of [40, Th. 8.5] goes through verbatim. (Since α is irreducible over F , the pair (g, u) is regular semisimple for any non-zero vector u in $V(F_0)$.) \square

Since their generic fibers do not intersect by [Theorem 9.2](#), the intersection pairing $\text{Int}(\xi, \Phi)$ localizes to a sum over all places of E . We define

$$(9.11) \quad \text{Int}_\nu^\natural(\xi, \Phi) := \langle \widehat{\mathcal{Z}}^B(\xi, \phi), {}^L\mathcal{CM}(\alpha, \phi_0) \rangle_\nu \log q_\nu,$$

where q_ν is the cardinality of the residue field of $O_{E,(\nu)}$ for non-archimedean ν (see below for the archimedean case). Here we recall that the local intersection number $\langle \cdot, \cdot \rangle_\nu$ is defined for a non-archimedean place ν through the Euler–Poincaré characteristic of a derived tensor product on $\mathcal{M} \otimes_{O_E} O_{E,(\nu)}$; cf. [12, 4.3.8(iv)]. For an archimedean place ν , the local intersection number is the value of the Green function at the complex point of the CM cycle:

$$(9.12) \quad \text{Int}_\nu^\natural(\xi, \Phi) := \langle \mathcal{G}_\nu^B(\xi, \phi), {}^L\mathcal{CM}(\alpha, \phi_0)_{\nu, \mathbb{C}} \rangle \log q_\nu,$$

where by definition $\log q_\nu = 2$ for complex places ν (and 1 if ν were a real place).

For a place v_0 of F_0 , we set

$$(9.13) \quad \text{Int}_{v_0}(\xi, \Phi) := \frac{1}{\tau(Z^\mathbb{Q}) \cdot [E : F]} \sum_{\nu|v_0} \text{Int}_\nu^\natural(\xi, \Phi).$$

Then we have a decomposition into a sum over places v_0 of F_0

$$(9.14) \quad \text{Int}(\xi, \Phi) = \sum_{v_0} \text{Int}_{v_0}(\xi, \Phi).$$

Combining (9.10), we obtain a decomposition of the generating function of arithmetic intersection numbers

$$(9.15) \quad \text{Int}(\tau, \Phi) = \text{Int}(0, \Phi) + \sum_{v_0} \text{Int}_{v_0}(\tau, \Phi),$$

where

$$(9.16) \quad \text{Int}_{v_0}(\tau, \Phi) := \sum_{\xi \in F_{0,+}} \text{Int}_{v_0}(\xi, \Phi) q^\xi.$$

COROLLARY 9.3 (to [Theorem 9.2](#)). *If v_0 is split in F/F_0 , then*

$$(9.17) \quad \text{Int}_{v_0}(\xi, \Phi) = 0.$$

9.4. *Local intersection: inert non-archimedean places.* Now let v_0 be a place of F_0 inert in F , and let w_0 be the unique place of F above v_0 . The notation here follows [Section 7.3](#).

THEOREM 9.4. *Assume that $v_0 \nmid \mathfrak{d}$ and $\Phi = \Phi_{v_0} \otimes \Phi^{v_0}$, where*

$$\Phi_{v_0} = \mathbf{1}_{K_{G,v_0}^\circ} \otimes \mathbf{1}_{\Lambda_{v_0}^\circ}.$$

Then

$$(9.18) \quad \text{Int}_{v_0}(\xi, \Phi) = 2 \log q_{v_0} \sum_{(\delta, u) \in [(G'(\alpha) \times V'_\xi)(F_0)]} \text{Int}_{v_0}(\delta, u) \cdot \text{Orb}((\delta, u), \Phi^{v_0}).$$

Here $\text{Int}_{v_0}(\delta, u)$ is the quantity defined in the AFL conjecture (semi-Lie algebra version) for the unramified quadratic extension $F_{w_0}/F_{0,v_0}$ (cf. (3.13)), and the orbital integral is the product of the local orbital integral defined by (2.15) with Haar measures on $G(F_{0,v})$ such that $\text{vol}(K_{G,v}) = 1$.

Proof. The proof follows a similar line to [46, Th. 3.11] and [40, Th. 8.15]. First, by [Theorem 9.2\(iii\)](#), the intersection only takes place in the basic locus. Hence it suffices to consider the question in the formal completion along the basic locus. We now fix a place ν of E above v_0 . Now by [Propositions 7.4](#) and [7.17](#), it suffices to consider the intersection number for each fiber of the projection (7.5) and multiply the result by the factor $\tau(Z^\mathbb{Q})$ (hence canceling the factor $\tau(Z^\mathbb{Q})$ in the denominator of (9.13)). Therefore we consider only the intersection on the fiber $\mathcal{M}_{O_{E_\nu}, 0}^\wedge$; cf. (7.6).

Recall that by [Proposition 7.4](#), the restriction to $\mathcal{M}_{O_{E_\nu}, 0}^\wedge$ of the special divisor $\mathcal{Z}(\xi, \phi)$ is

$$\sum_{(u, g') \in G'(F_0) \setminus (V'_\xi(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0})} \phi^{v_0}(g'^{-1}u) \cdot [\mathcal{Z}(u, g')]_{K_G^{v_0}},$$

and by [Proposition 7.17](#) the restriction of the derived CM cycle ${}^L\mathcal{CM}(\alpha, \phi_0)$ is the sum

$$\sum_{(\delta, h) \in G'(F_0) \setminus (G'(\alpha)(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0})} \phi_0^{v_0}(h^{-1}\delta h) \cdot [{}^L\mathcal{CM}(\delta, h)]_{K_G^{v_0}}.$$

We may compute the intersection number by pulling-back to the covering formal scheme $\mathcal{N}_{O_{E_\nu}} \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}$ in the uniformization (7.6). The intersection number ${}^L\mathcal{CM}(\alpha, \phi_0) \cap \mathcal{Z}(\xi, \phi) \log q_\nu$ (restricted to $\mathcal{M}_{O_{E_\nu}, 0}^\wedge$) is equal to a sum of

$$\phi_0^{v_0}(h^{-1}\delta h) \phi^{v_0}(g'^{-1}u) \cdot {}^L\mathcal{CM}(\delta, h)_{K_G^{v_0}} \cap \mathcal{Z}(u, g')_{K_G^{v_0}} \cdot \log q_\nu,$$

over $G'(F_0)$ -orbits (via diagonal action) of tuples (δ, h, u, g') :

$$(\delta, h) \in G'(\alpha)(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0} \quad \text{and} \quad (u, g') \in V'_\xi(F_0) \times G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}.$$

Here, we are abusing the notation $\overset{\mathbb{L}}{\cap}$ to denote the Euler–Poincaré characteristics of the corresponding derived tensor product.

By (7.7) and (7.18), we obtain

$$\mathbb{L}\mathcal{CM}(\delta, h)_{K_G^{v_0}} \overset{\mathbb{L}}{\cap} \mathcal{Z}(u, g')_{K_G^{v_0}} \cdot \log q_\nu = \mathbb{L}\mathcal{N}_{O_{E_\nu}}^\delta \overset{\mathbb{L}}{\cap} \mathcal{Z}(u)_{O_{E_\nu}} \log q_\nu \cdot \mathbf{1}_{K_G^{v_0}}(g'^{-1}h).$$

The first term is equal to

$$\begin{aligned} \mathbb{L}\mathcal{N}_{O_{E_\nu}}^\delta \overset{\mathbb{L}}{\cap} \mathcal{Z}(u)_{O_{E_\nu}} \log q_\nu &= [E_\nu : F_{w_0}] \cdot \left(\mathbb{L}\mathcal{N}^\delta \overset{\mathbb{L}}{\cap} \mathcal{Z}(u) \right) \log q_{w_0} \\ &= 2[E_\nu : F_{w_0}] \cdot \text{Int}_{v_0}(\delta, u) \log q_{v_0}. \end{aligned}$$

Here the factor 2 is due to $q_{w_0} = q_{v_0}$. In particular, it is invariant under the (diagonal) action of $G'(F_0)$ on the product $(G'(\alpha) \times V'_\xi)(F_0)$.

The second term $(g', h) \in (G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0})^2 \mapsto \mathbf{1}_{K_G^{v_0}}(g'^{-1}h)$ is also invariant under the (diagonal) $G'(F_0)$ -action. For a fixed pair (δ, u) , we obtain

$$\begin{aligned} \sum_{(g', h) \in (G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0})^2} \phi_0^{v_0}(h^{-1}\delta h) \phi^{v_0}(g'^{-1}u) \cdot \mathbf{1}_{K_G^{v_0}}(g'^{-1}h) \\ = \sum_{h \in G(\mathbb{A}_{0,f}^{v_0})/K_G^{v_0}} \phi_0^{v_0}(h^{-1}\delta h) \phi^{v_0}(h^{-1} \cdot u) \\ = \int_{G(\mathbb{A}_{0,f}^{v_0})} \phi_0^{v_0}(h^{-1}\delta h) \phi^{v_0}(h^{-1} \cdot u) \, dh \\ = \text{Orb}((\delta, u), \Phi^{v_0}), \end{aligned}$$

where we note that the Haar measure on $G'(\mathbb{A}_f^{v_0})$ is normalized such that $\text{vol}(K_G^{v_0}) = 1$.

To summarize, the intersection number $\mathbb{L}\mathcal{CM}(\alpha, \phi_0) \overset{\mathbb{L}}{\cap} \mathcal{Z}(\xi, \phi) \log q_\nu$ (restricted to $\mathcal{M}_{O_{E_\nu}, 0}$) is equal to

$$2[E_\nu : F_{w_0}] \sum_{(\delta, u)} \text{Orb}((\delta, u), \Phi^{v_0}) \cdot \text{Int}_{v_0}(\delta, u) \log q_{v_0},$$

where the sum is over $G'(F_0)$ -orbits of pairs $(\delta, u) \in (G'(\alpha) \times V'_\xi)(F_0)$.

Finally the sum over all places $\nu \mid v_0$ will cancel the factor $[E : F]$ in (9.13) by

$$\sum_{\nu \mid w_0} e_{\nu/w_0} f_{\nu/w_0} = \sum_{\nu \mid w_0} d_{\nu/w_0} = [E : F],$$

where e_{ν/w_0} (resp. $f_{\nu/w_0}, d_{\nu/w_0}$) denotes the ramification degree (resp. inert degree, degree) of the extension E_ν/F_{w_0} . This completes the proof. \square

10. Local intersection: archimedean places

The goal of this section is to compute the local intersection at ν of E above an archimedean place v_0 of F_0 . In fact, we will replace the automorphic Green function by Kudla's Green function; i.e., we consider the analog of (9.12):

$$(10.1) \quad \text{Int}_\nu^{\natural, \mathbf{K}}(\xi, \Phi) := \langle \mathcal{G}_\nu^{\mathbf{K}}(\xi, \phi), {}^{\mathbb{L}}\mathcal{CM}(\alpha, \phi_0)_{\nu, \mathbb{C}} \rangle \log q_\nu.$$

When $F_0 = \mathbb{Q}$, the difference is addressed by Theorem 8.4. Similar to (9.13), we set for $\xi \in F_0$,

$$(10.2) \quad \text{Int}_{v_0}^{\mathbf{K}}(\xi, \Phi) := \frac{1}{\tau(Z^\mathbb{Q}) \cdot [E : F]} \sum_{\nu|v_0} \text{Int}_\nu^{\natural, \mathbf{K}}(\xi, \Phi).$$

We note that by (8.11) and (8.12), there is a parameter $h_\infty \in \mathbf{H}(F_0 \otimes_{\mathbb{Q}} \mathbb{R})$ implicitly in the above expression.

The strategy is analogous to Theorem 9.4. We follow the notation in Sections 8.2 and 8.3.

THEOREM 10.1. *Let $\Phi \in \mathcal{S}((G \times V)(\mathbb{A}_{0,f}))$. Let $\xi \neq 0$. Then we have*

$$(10.3) \quad \text{Int}_{v_0}^{\mathbf{K}}(\xi, \Phi) = \sum_{(\delta, u) \in [(G'(\alpha) \times V'_\xi)(F_0)]} \text{Int}_{v_0}(\delta, u) \cdot \text{Orb}((\delta, u), \Phi).$$

Here $\text{Int}_{v_0}(\delta, u)$ is defined as the special value of the function

$$(10.4) \quad \text{Int}_{v_0}(\delta, u) = \mathcal{G}^{\mathbf{K}}(u, h_\infty)(z_\delta),$$

where z_δ is the unique fixed point of δ on \mathcal{D}_{v_0} . Moreover, the point z_δ does not lie on $\mathcal{D}_{v_0, u}$ for any non-zero vector $u \in V'(F_0)$.

Proof. The proof goes along the same line as that of Theorem 9.4, so we will not repeat the details, except to prove the claim on the point z_δ . Consider the n -dimensional \mathbb{C} -vector space $V' \otimes_{F, w_0} \mathbb{C}$ with the induced hermitian form. If a negative definite \mathbb{C} -line is fixed by δ , it must be an eigen-line for δ , which must be unique by the signature $(n-1, 1)$ condition on the hermitian form on $V' \otimes_{F, w_0} \mathbb{C}$. If z_δ lies on a divisor $\mathcal{D}_{v_0, u}$ for non-zero vector $u \in V'(F_0)$, it also lies on $\mathcal{D}_{v_0, \delta^i \cdot u}$, the translation of $\mathcal{D}_{v_0, u}$ under δ^i , for all $i \in \mathbb{Z}$. Equivalently, the line z_δ is perpendicular to all $\delta^i \cdot u \in V' \otimes_{F, w_0} \mathbb{C}$. Since $u \in V_0(F)$ is a non-zero vector and its characteristic polynomial α of δ is irreducible over F , the vectors $\delta^i \cdot u$ span V' over F , hence they also span $V' \otimes_{F, w_0} \mathbb{C}$ over F_{w_0} . Contradiction! \square

It remains to compute (10.4), or equivalently $R(u, z_\delta)$ defined by (8.9). The element $\delta \in G'(\alpha)(F_0)$ induces an action of the CM field F' (cf. (7.9)) on V' and makes V' into a one-dimensional F'/F'_0 -hermitian space $(W, \langle \cdot, \cdot \rangle_{F'_0})$

satisfying

$$(10.5) \quad (R_{F'/F}W, \text{tr}_{F'/F}\langle \cdot, \cdot \rangle_{F'_0}) \xrightarrow{\sim} (V', \langle \cdot, \cdot \rangle).$$

Here $R_{F'/F}W$ denotes the “restriction of scalar” of W , i.e., to view it as an F -vector space.

Remark 10.2. The above construction $\delta \in G'(\alpha)(F_0) \mapsto (W, \langle \cdot, \cdot \rangle_{F'_0})$ defines a bijection between the set of $G'(F_0)$ -conjugacy classes in $G'(\alpha)(F_0)$ and the set of one-dimensional F'/F'_0 -hermitian spaces (up to isometry) satisfying (10.5). In fact, fixing a $\delta_0 \in G'(\alpha)(F_0)$, we denote by W_0 the associated F'/F'_0 -hermitian space and by T the centralizer of δ_0 in G' . Then T is an anisotropic F_0 -torus isomorphic to $\text{Res}_{F'_0/F_0} \text{U}(W_0)$. Now the pointed set of $G'(F_0)$ -conjugacy classes in $G'(\alpha)(F_0)$ (with the conjugacy class of δ_0 as the distinguished element) is bijective to the pointed set $\ker(H^1(F_0, T) \rightarrow H^1(F_0, G'))$. Moreover, the pointed set $\ker(H^1(F_0, T) \rightarrow H^1(F_0, G'))$ is naturally isomorphic to the pointed set of one-dimensional F'/F'_0 -hermitian spaces (up to isometry) satisfying (10.5) (with the F'/F'_0 -hermitian space W_0 as the distinguished element). A similar remark applies to local fields rather than F/F_0 (except that the torus T may not be anisotropic).

It follows from (10.5) that the F'/F'_0 -hermitian space W has signatures $(1, 0)$ for all but one archimedean place v'_0 of F'_0 over v_0 . We define a refined invariant

$$(10.6) \quad \xi' = \mathfrak{q}'(u) \in F'_0,$$

where \mathfrak{q}' is the quadratic form attached to the F'/F'_0 -hermitian form on W ; cf. (1.4). In particular, $\text{tr}_{F'_0/F_0}(\xi') = \xi$.

According to the action of F'_0 , we have an orthogonal direct sum decomposition

$$V' \otimes_{F,w_0} \mathbb{C} = \bigoplus_{v' \in \text{Hom}(F'_0, \mathbb{R}), v'|_{F_0} = v_0} \mathbb{C}_{v'},$$

where F'_0 acts on the line $\mathbb{C}_{v'}$ through $v' : F'_0 \hookrightarrow \mathbb{R}$. Then there is a unique negative-definite summand, say $\mathbb{C}_{v'_0}$ for a place v'_0 above v_0 . It follows that

$$(10.7) \quad R(u, z_\delta) = v'_0(\mathfrak{q}'(u)) = -|\xi'|_{v'_0},$$

where the last equality is due to the fact $v'_0(\mathfrak{q}'(u)) < 0$.

COROLLARY 10.3. *Under the same assumptions as Theorem 10.1, we have*

$$(10.8) \quad \text{Int}_{v_0}^K(\xi, \Phi) = - \sum \text{Ei}(-2\pi|\xi'|_{v'_0}) \cdot \text{Orb}((\delta, u), \Phi),$$

where the sum runs over the $G'(F_0)$ -orbits (δ, u) in the product $(G'(\alpha) \times V'_\xi)(F_0)$, $\xi' = \mathfrak{q}'(u)$ is the refined invariant defined by (10.6), and $v'_0 \mid v_0$ is the unique archimedean place of F'_0 where ξ' is negative.

Finally, we address the difference between the two Green functions. Define, for any place $v \mid \infty$ of F_0 and $h \in \mathbf{H}(\mathbb{A}_0)$,

$$(10.9) \quad \text{Int}_v^{\mathbf{K}-\mathbf{B}}(h, \Phi) = \frac{1}{\tau(Z^\mathbb{Q}) \cdot [E : F]} \left(\mathcal{Z}_{v, \text{corr}}(h, \phi), \quad \mathbb{L}\mathcal{CM}(\alpha, \phi_0) \right)$$

(cf. (8.14)), and define

$$(10.10) \quad \text{Int}^{\mathbf{K}-\mathbf{B}}(h, \Phi) = \sum_{v \mid \infty} \text{Int}_v^{\mathbf{K}-\mathbf{B}}(h, \Phi).$$

We note that the definition works without any reference to the integral models \mathcal{M} , hence makes sense for all $\phi_0 \in \mathcal{S}(G(\mathbb{A}_{0,f}), K_G)$ and $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))^{K_G}$.

COROLLARY 10.4 (to Theorem 8.4). *Let $F_0 = \mathbb{Q}$. Then the function $h \in \mathbf{H}(\mathbb{A}_0) \mapsto \text{Int}^{\mathbf{K}-\mathbf{B}}(h, \Phi)$ belongs to $\mathcal{A}_{\text{exp}}(\mathbf{H}(\mathbb{A}_0), K, n)$, where $K \subset \mathbf{H}(\mathbb{A}_{0,f})$ is a compact open subgroup that fixes $\phi \in \mathcal{S}(V(\mathbb{A}_{0,f}))$ under the Weil representation.*

11. Weil representation and RTF

Starting from this section, we study a partially linearized version of the Jacquet–Rallis relative trace formula and the “action” on the RTF by $\text{SL}_2(\mathbb{A}_0)$ under the Weil representation (by changing testing functions on the linear factor of the RTF).

11.1. Weil representation and theta functions. For now we let F be a global field. Let (V, \mathfrak{q}) be a (non-degenerate) quadratic space over F of even dimension d , where $\mathfrak{q} : V \rightarrow F$ is the quadratic form with the associated symmetric bilinear pairing $\langle \cdot, \cdot \rangle : V \times V \rightarrow F$ by (1.2). Let $O(V) = O(V, \mathfrak{q})$ be the isometry group, viewed as an algebraic group over F .

Let $\mathcal{S}(V(\mathbb{A}_F))$ be the space of Schwarz functions. The product group $O(V)(\mathbb{A}_F) \times \text{SL}_2(\mathbb{A}_F)$ acts on $\mathcal{S}(V(\mathbb{A}_F))$ via the Weil representation denoted by ω : for $\phi \in \mathcal{S}(V(\mathbb{A}_F))$, the function $\omega(g, h)\phi$ is defined by

$$(\omega(g, h)\phi)(x) = (\omega(h))\phi(g^{-1}x), \quad (g, h) \in O(V)(\mathbb{A}_F) \times \text{SL}_2(\mathbb{A}_F),$$

where the action of $\text{SL}_2(\mathbb{A}_F)$ is defined as follows. Let $\chi_V = \prod_v \chi_{V_v}$ be the quadratic character of $F^\times \backslash \mathbb{A}_F^\times$ defined by

$$\chi_V(a) = (a, (-1)^{d/2} \det(V))_F,$$

where (\cdot, \cdot) is the Hilbert symbol over F and $\det(V) \in F^\times / (F^\times)^2$ is the determinant of the moment matrix $\frac{1}{2}(\langle x_i, x_j \rangle)_{1 \leq i, j \leq d}$ of any F -basis x_1, \dots, x_d of V .

For a place v of F , and $\phi_v \in \mathcal{S}(V(F_v))$, the action of $\mathrm{SL}_2(F_v)$ is determined by

$$(11.1) \quad \begin{aligned} \omega_v \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix} \phi_v(x) &= \chi_{V_v}(a) |a|_v^{d/2} \phi_v(ax), \\ \omega_v \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix} \phi_v(x) &= \psi_v(b\mathfrak{q}(x)) \phi_v(x), \\ \omega_v \begin{pmatrix} & 1 \\ -1 & \end{pmatrix} \phi_v(x) &= \gamma_{V_v} \widehat{\phi}_v(x), \end{aligned}$$

where γ_{V_v} is the Weil constant (a fourth root of unity under our assumption that $\dim V_v$ is even), and the Fourier transform is defined by

$$\widehat{\phi}_v(x) = \int_{V(F_v)} \phi_v(y) \psi_v(\langle x, y \rangle) dy.$$

Here dy is a self-dual Haar measure on $V(F_v)$.

For $\phi \in \mathcal{S}(V(\mathbb{A}_F))$, we define the theta function by the absolute convergent sum

$$\theta_\phi(g, h) = \sum_{\xi \in V} \omega(g, h) \phi(\xi), \quad (g, h) \in O(V)(\mathbb{A}_F) \times \mathrm{SL}_2(\mathbb{A}_F).$$

This is left invariant under $O(V)(F) \times \mathrm{SL}_2(F)$.

11.2. Automorphic kernel functions. In this subsection we work with a fairly general setting. It serves to explain the idea behind the more explicit setting in later sections.

Let G be a connected reductive algebraic group over F , acting on V and preserving the quadratic form \mathfrak{q} (i.e., the homomorphism $G \rightarrow \mathrm{GL}(V)$ factors through $O(V, \mathfrak{q})$). Let X be an affine variety over F with an action of G , and let $X_{/G}$ be the categorical quotient. Consider the diagonal action r of G on $X \times V$. Then $G(\mathbb{A}_F)$ acts on $\mathcal{S}((X \times V)(\mathbb{A}_F))$.

The group $\mathrm{SL}_2(\mathbb{A}_F)$ acts on $\mathcal{S}((X \times V)(\mathbb{A}_F))$ through the second factor V via the Weil representation. Note that now the formula (11.1) for the action of $\mathrm{SL}_2(\mathbb{A}_F)$ is only applied to the second coordinate; e.g., locally at v , the element $\begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$ acts on $\mathcal{S}((X \times V)(F_v))$ by (up to the Weil constant γ_{V_v}) the partial Fourier transform with respect to the V -component.

Let $\alpha \in X_{/G}(F)$ be a fixed semi-simple element and $X(\alpha)$ the preimage of α (under the quotient map $X \rightarrow X_{/G}$). Let $\phi_0 \in \mathcal{S}(X(\mathbb{A}_F))$ and $\phi \in \mathcal{S}(V(\mathbb{A}_F))$. We define the automorphic kernel function associated to $\Phi =$

$\phi_0 \otimes \phi \in \mathcal{S}((X \times V)(\mathbb{A}_F))$,

$$(11.2) \quad \begin{aligned} \mathcal{K}_{\Phi,\alpha}(g, h) &:= \sum_{(x,u) \in (X(\alpha) \times V)(F)} \phi_0(g^{-1} \cdot x) \omega(h) \phi(g^{-1} \cdot u) \\ &= \sum_{(x,u) \in (X(\alpha) \times V)(F)} \omega(h) \Phi(g^{-1} \cdot (x, u)), \end{aligned}$$

where $g \in G(\mathbb{A}_F)$, $h \in \mathrm{SL}_2(\mathbb{A}_F)$. This is again left invariant under $G(F) \times \mathrm{SL}_2(F)$. It follows that

$$(11.3) \quad h \in \mathrm{SL}_2(\mathbb{A}_F) \longmapsto \mathbb{J}(h, \Phi) := \int_{[G]} \mathcal{K}_{\Phi,\alpha}(g, h) dg,$$

when absolutely convergent, is left invariant under $\mathrm{SL}_2(F)$. The same applies if we replace the pure tensor $\phi_0 \otimes \phi$ by a more general function Φ in $\mathcal{S}((X \times V)(\mathbb{A}_F))$. (This does not make any essential difference at non-archimedean places, but does at archimedean places.)

Now we return to our earlier convention. Let F_0 be a totally real field, and let F/F_0 be a CM field extension. Let

$$\eta = \eta_{F/F_0} : F_0^\times \setminus \mathbb{A}_0^\times \longrightarrow \{\pm 1\}$$

be the quadratic character by class field theory. Note that now F_0 plays the role of the base field F in above discussion.

11.3. The case of unitary groups. Now we consider the Jacquet–Rallis RTF for unitary groups. Let V be a F/F_0 -hermitian space of dimension n . Let $G = \mathrm{U}(V)$ be the unitary group, and let $X = G$ with the conjugation action by G . Let $\alpha \in \mathcal{A}_n(F_0)$ be irreducible over F (cf. the end of [Section 2.2](#)). We rewrite the kernel function (11.2) according to $(\delta, u) \in (G(\alpha) \times V)(F_0)$ regular semisimple (equivalently $u \neq 0$ by the irreducibility of α) or not, and then (11.3) becomes

$$(11.4) \quad \begin{aligned} \mathbb{J}(h, \Phi) &= \int_{[G]} \sum_{(\delta,u) \in (G(\alpha) \times V)(F_0)} r(g) \omega(h) \Phi(\delta, u) dg \\ &= \int_{[G]} \sum_{\delta \in G(\alpha)(F_0)} \omega(h) \Phi(g^{-1} \cdot \delta, 0) dg \\ &\quad + \int_{[G]} \sum_{\substack{(\delta,u) \in (G(\alpha) \times V)(F_0) \\ u \neq 0}} \omega(h) \Phi(g^{-1} \cdot (\delta, u)) dg. \end{aligned}$$

The summands in (11.4) are related to the global Jacquet–Rallis (relative) orbital integral (for the G -action on $G \times V$) of $\omega(h)\Phi$. For $\Phi \in \mathcal{S}((G \times V)(\mathbb{A}_0))$

and a regular semisimple $(\delta, u) \in (G \times V)(F_0)$, we define

$$(11.5) \quad \text{Orb}((\delta, u), \Phi) := \int_{G(\mathbb{A}_0)} \Phi(g^{-1} \cdot (\delta, u)) dg.$$

For $\delta \in G(\alpha)(F_0)$, we define

$$(11.6) \quad \text{Orb}((\delta, 0), \Phi) := \text{vol}([G_\delta]) \int_{G_\delta(\mathbb{A}_0) \backslash G(\mathbb{A}_0)} \Phi(g^{-1} \cdot \delta, 0) dg,$$

where G_δ is the centralizer of δ in G , an anisotropic F_0 -torus.

The first summand in (11.4) is a sum over the set of $G(F_0)$ -conjugacy classes in $G(\alpha)(F_0)$ (cf. Section 2.2),

$$(11.7) \quad \sum_{\delta \in [G(\alpha)(F_0)]} \text{Orb}((\delta, 0), \omega(h)\Phi).$$

There are only finitely many non-zero terms (uniformly in $h \in \mathbf{H}(\mathbb{A}_0)$), and hence the sum is absolutely convergent. The second summand in (11.4) is a sum over the set, denoted by $[(G(\alpha) \times V)(F_0)]_{\text{rs}}$, of regular semisimple (equivalently, $u \neq 0$) $G(F_0)$ -orbits in $(G(\alpha) \times V)(F_0)$:

$$\sum_{(\delta, u) \in [(G(\alpha) \times V)(F_0)]_{\text{rs}}} \text{Orb}((\delta, u), \omega(h)\Phi).$$

We first justify the convergence. We recall that $\mathbf{H} = \text{SL}_2, F_0$.

LEMMA 11.1. (a) *For any $\Phi \in \mathcal{S}((G \times V)(\mathbb{A}_0))$, we have*

$$\sum_{(\delta, u) \in [(G \times V)(F_0)]_{\text{rs}}} \int_{G(\mathbb{A}_0)} |\Phi(g^{-1} \cdot (\delta, u))| dg < \infty.$$

In particular, the same holds if we only sum over $(\delta, u) \in [(G(\alpha) \times V)(F_0)]_{\text{rs}}$.

(b) *Assume that Φ is K_∞ -finite for the maximal compact $K_\infty = \prod_v \text{SO}(2, \mathbb{R})$ of $\mathbf{H}(F_{0,\infty})$. Then the sum in $\mathbb{J}(h, \Phi)$ converges absolutely and uniformly for h in any compact subset of $\mathbf{H}(\mathbb{A}_0)$. In particular, the function $h \in \mathbf{H}(\mathbb{A}_0) = \text{SL}_2(\mathbb{A}_0) \mapsto \mathbb{J}(h, \Phi)$ is smooth and left invariant under $\mathbf{H}(F_0)$.*

Proof. For part (a), we prove a stronger result:

$$\sum_{(\delta, u) \in [(G \times V)(F_0)]_{\text{rs}}} \int_{G(\mathbb{A}_0)} |\Phi(g^{-1} \cdot (\delta, u))| dg < \infty.$$

We first note that this is easy when Φ has compact support, since the sum would then have only finitely many non-zero terms. Next we may and do assume that $\Phi = \prod_v \Phi_v$ is a pure tensor (otherwise we can dominate Φ by a finite sum of pure tensors).

We refer to [1, §A.1] for the terminology in our proof. Consider $X = G \times V$, and the categorical quotient $\mathcal{B} = X // G$ with the natural map $\pi : X \rightarrow \mathcal{B}$. The regular semisimple locus in \mathcal{B} (resp. X), denoted by \mathcal{B}_{rs} (resp. X_{rs}), is defined

by $\Delta \neq 0$ (resp. $\Delta \circ \pi \neq 0$) for a regular function Δ on \mathcal{B} (resp. its pull-back to X). Then the restriction $\pi_{\text{rs}} : X_{\text{rs}} \rightarrow \mathcal{B}_{\text{rs}}$ is a G -torsor.

We fix a norm $\|\cdot\|_X$ on $X(\mathbb{A}_0)$, which is the product of local norms $\|\cdot\|_{X_v}$ on $X(F_{0,v})$. Similarly, we fix norms on $X_{\text{rs}}(\mathbb{A}_0)$ and $X_{\text{rs}}(F_{0,v})$. For the norm on the affine line, we write $\|t\|$ (resp. $\|t\|_v$) for $t \in \mathbb{A}_0$ (resp. $t \in F_{0,v}$), defined by $\|t\|_v = \max\{1, |t|_v\}$, cf. [1, §A.1, before (1)].

For $\Phi \in \mathcal{S}(X(\mathbb{A}_0))$, we have

$$(11.8) \quad |\Phi(x)| \ll \|x\|_X^{-d_1} \quad \text{for all } x \in X(\mathbb{A}_0)$$

for any constant $d_1 > 0$ ⁸. By [1, Prop. A.1.1(iii)], for all $x \in X_{\text{rs}}(\mathbb{A}_0)$,

$$\|x\|_{X_{\text{rs}}} \sim \|x\|_X \|\Delta \circ \pi(x)^{-1}\|.$$

In particular, there exists a constant $d_2 > 0$ such that, for all $x \in X_{\text{rs}}(\mathbb{A}_0)$,

$$\|x\|_{X_{\text{rs}}}^{d_2} \ll \|x\|_X \|\Delta \circ \pi(x)^{-1}\|.$$

By (11.8) and [1, Prop. A.1.1 (vii)], for any $d_3 > 0$, there exists $d_1 > 0$ large enough such that

$$(11.9) \quad \begin{aligned} \int_{G(\mathbb{A}_0)} |\Phi(g \cdot x)| dg &\ll \|\Delta \circ \pi(x)^{-1}\|^{d_1} \int_{G(\mathbb{A}_0)} \|g \cdot x\|_{X_{\text{rs}}}^{-d_1 d_2} \\ &\ll \|\Delta \circ \pi(x)^{-1}\|^{d_1} \|\pi_{\text{rs}}(x)\|_{\mathcal{B}_{\text{rs}}}^{-d_3} \end{aligned}$$

for all $x \in X_{\text{rs}}(\mathbb{A}_0)$. Note that this implies similar estimates: for any $d_3 > 0$, there exists $d_1 > 0$ such that

$$(11.10) \quad \int_{G(F_{0,v})} |\Phi_v(g \cdot x_v)| dg \ll \|\Delta \circ \pi(x_v)^{-1}\|_v^{d_1} \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d_3}$$

holds for every place v , and

$$(11.11) \quad \prod_{v \notin S} \int_{G(F_{0,v})} |\Phi_v(g \cdot x_v)| dg \ll \prod_{v \notin S} \|\Delta \circ \pi(x_v)^{-1}\|_v^{d_1} \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d_3}$$

holds for any finite set S of places, where $(x_v)_{v \notin S} \in X_{\text{rs}}(\prod_{v \notin S} F_{0,v})$. Here we emphasize that d_1, d_3 can be made independent of v .

Now we claim that for any $d_3 > 0$, there exists $d_1 > 0$ such that

$$(11.12) \quad \int_{G(F_{0,v})} |\Phi_v(g \cdot x_v)| dg \ll \|\Delta \circ \pi(x_v)\|_v^{-d_1} \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d_3}$$

holds for every place v . (Here we emphasize that d_1, d_3 can be made independent of v .) Indeed, if $|\Delta \circ \pi(x_v)|_v \leq 1$, then $\|\Delta \circ \pi(x_v)^{-1}\|_v = |\Delta \circ \pi(x_v)^{-1}|_v$, and

⁸Here, for two functions f_1, f_2 , the notation $f_1 \ll f_2$ means that there is a constant c such that $f_1 \leq c f_2$.

hence (11.12) follows from (11.10) in this case. Now suppose $|\Delta \circ \pi(x_v)|_v \geq 1$ and then $\|\Delta \circ \pi(x_v)^{-1}\|_v = 1$. It follows from (11.10) that

$$(11.13) \quad \int_{G(F_{0,v})} |\Phi_v(g \cdot x_v)| dg \ll \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d'_3}$$

for any constant $d'_3 > 0$. By [1, Prop. A.1.1(ii)] applied to the morphism $\Delta : \mathcal{B}_{\text{rs}} \rightarrow \mathbb{A}^1$ (here \mathbb{A}^1 denotes the affine line), there exists a constant $d_4 > 0$ such that

$$\|\Delta(b)\|_v \ll \|b\|_{\mathcal{B}_{\text{rs},v}}^{d_4}$$

for all $b \in \mathcal{B}_{\text{rs}}(F_{0,v})$.

Note that $|\Delta(b)|_v \leq \|\Delta(b)\|_v$. Choosing $d'_3 = d_3 + d_1 d_4$ in (11.13), we arrive at the estimate (11.12) (for any constants $d_1, d_3 > 0$).

Since the support of the non-archimedean component $\Phi^\infty = \prod_{v \nmid \infty} \Phi_v$ is compact, its image under $\Delta \circ \pi$ is also compact in $\mathbb{A}_{0,f}$. It follows that for $x \in X_{\text{rs}}(\mathbb{A}_0) \cap \text{supp}(\Phi)$,

$$(11.14) \quad \prod_{v \nmid \infty} \|\Delta \circ \pi(x)^{-1}\|_v \ll \prod_{v \nmid \infty} |\Delta \circ \pi(x)^{-1}|_v = \prod_{v \nmid \infty} |\Delta \circ \pi(x)|_v^{-1}.$$

It follows that when $x = (x_v)_v \in X_{\text{rs}}(\mathbb{A}_0)$, for any $d_3 > 0$, there exists $d_1 > 0$ such that

$$\begin{aligned} & \int_{G(\mathbb{A}_0)} |\Phi(g \cdot x)| dg \\ &= \prod_{v \nmid \infty} \int_{G(F_{0,v})} |\Phi_v(g \cdot x_v)| dg \prod_{v \nmid \infty} \int_{G(F_{0,v})} |\Phi_v(g \cdot x_v)| dg \\ &\ll \prod_{v \nmid \infty} |\Delta \circ \pi(x_v)|_v^{-d_1} \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d_3} \\ & \quad \cdot \prod_{v \nmid \infty} \|\Delta \circ \pi(x_v)^{-1}\|_v^{d_1} \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d_3} \quad (\text{by (11.12) and (11.11)}) \\ &\ll \prod_v |\Delta \circ \pi(x_v)|_v^{-d_1} \|\pi_{\text{rs}}(x_v)\|_{\mathcal{B}_{\text{rs},v}}^{-d_3} \quad (\text{by (11.14)}). \end{aligned}$$

Finally, let $x \in X_{\text{rs}}(F_0)$. By the product formula $\prod_v |\Delta \circ \pi(x)|_v = 1$, we obtain

$$\int_{G(\mathbb{A}_0)} |\Phi(g \cdot x)| dg \ll \|\pi_{\text{rs}}(x)\|_{\mathcal{B}_{\text{rs}}}^{-d_3}$$

for any constant $d_3 > 0$. The desired convergence then follows from [1, Prop. A.1.1 (v)] (applied to \mathcal{B}_{rs}):

$$\sum_{b \in \mathcal{B}_{\text{rs}}(F_0)} \|b\|_{\mathcal{B}_{\text{rs}}}^{-d_3} < \infty$$

for d_3 large enough.

To show part (b), it suffices to show that the constant implicit in \ll of (11.8) can be made uniform for $\omega(h)\Phi$ for $h \in H(\mathbb{A}_0)$ in a neighborhood of 1. The function Φ^∞ is invariant under a compact open of $H(\mathbb{A}_{0,f})$. So it suffices to consider $h \in H(F_{0,\infty})$. By the K_∞ -finiteness assumption, it suffices to consider upper triangular elements in $H(F_{0,\infty})$. Then it is easy to see the constant can be made uniform by the formula (11.1). \square

To summarize, we obtain

$$(11.15) \quad \mathbb{J}(h, \Phi) = \sum_{(\delta, u) \in [(G(\alpha) \times V)(F_0)]} \text{Orb}((\delta, u), \omega(h)\Phi).$$

When Φ_∞ is K_∞ -finite, it follows easily that for $\xi \in F_0^\times$, the ξ -th Fourier coefficient of $\mathbb{J}(\cdot, \Phi)$ is equal to

$$(11.16) \quad \sum_{(\delta, u) \in [(G(\alpha) \times V_\xi)(F_0)]} \text{Orb}((\delta, u), \omega(h)\Phi).$$

Here we refer to (1.12) for the definition of Fourier coefficients.

11.4. *The case of general linear groups.* Now we consider the Jacquet–Rallis RTF for general linear groups. Let $V_0 = F_0^n$ be the n -dimensional vector space of column vectors over F_0 . We identify the dual vector space $V_0^* = \text{Hom}_{F_0}(V_0, F_0)$ with the space of row vectors. Consider the natural quadratic form on $V' = V_0 \times V_0^*$:

$$(11.17) \quad \begin{aligned} \mathfrak{q} : \quad V_0 \times V_0^* &\longrightarrow F_0 \\ u' = (u_1, u_2) &\longmapsto u_2(u_1). \end{aligned}$$

Let

$$\langle \cdot, \cdot \rangle : V' \times V' \longrightarrow F_0$$

be the the associated symmetric bilinear pairing (so that $\langle u', u' \rangle = 2\mathfrak{q}(u')$). Let $G' = \text{GL}(V_0)$ act on V' by $(\text{std}, \text{std}^V)$. Then $G' \simeq \text{GL}_{n, F_0}$ via the given identification $V_0 = F_0^n$. Consider the diagonal action of G' on $S_n \times V'$; cf. (2.2).

Now let $\alpha \in \mathcal{A}_n(F_0)$ be irreducible over F . We rewrite the kernel function (11.2) according to $(\gamma, u') \in (S_n(\alpha) \times V')(F_0)$ regular semisimple or not. By the irreducibility of α , precisely three orbits are not regular semisimple, i.e., (γ, u') where u' are

$$(11.18) \quad \begin{cases} \{(0, 0)\}, \\ 0_+ := \{(u_1, 0) : u_1 \in V_0(F_0) \setminus \{0\}\}, \\ 0_- := \{(0, u_2) : u_2 \in V_0^*(F_0) \setminus \{0\}\}. \end{cases}$$

They will be called “(relative) nilpotent”; the last two are regular (i.e., with trivial stabilizers).

We define the (global) Jacquet–Rallis (relative) orbital integral (for the G' -action on $S_n \times V'$). For $\Phi' \in \mathcal{S}((S_n \times V')(\mathbb{A}_0))$ and a regular semisimple $(\gamma, u') \in (S_n \times V')(F_0)$, we define

$$(11.19) \quad \text{Orb}((\gamma, u'), \Phi', s) := \int_{G'(\mathbb{A}_0)} \Phi'(g^{-1} \cdot (\gamma, u')) |\det(g)|_{F_0}^s \eta(g) dg.$$

Here and thereafter we will simply denote by η the character $\eta \circ \det$ of $G'(\mathbb{A}_0)$. The global orbital integral is a product of local orbital integrals

$$(11.20) \quad \text{Orb}((\gamma, u'), \Phi'_v, s) := \int_{G'(F_{0,v})} \Phi'_v(g^{-1} \cdot (\gamma, u')) |\det(g)|_v^s \eta(g) dg.$$

We consider a one-parameter family of (11.3): for $\Phi' \in \mathcal{S}((S_n \times V')(\mathbb{A}_0))$ and $s \in \mathbb{C}$, we will define

$$\mathbb{J}(h, \Phi', s) = \int_{[G']} \left(\sum_{(\gamma, u') \in (S_n(\alpha) \times V')(F_0)} r(g) \omega(h) \Phi'(\gamma, u') \right) |\det(g)|_{F_0}^s \eta(g) dg.$$

Similar to the unitary case, we write it as a sum over orbits:

$$(11.21) \quad \mathbb{J}(h, \Phi', s) = \mathbb{J}(h, \Phi', s)_0 + \mathbb{J}(h, \Phi', s)_{\text{rs}},$$

where

$$\mathbb{J}(h, \Phi', s)_{\text{rs}} = \sum_{(\gamma, u') \in [(S_n(\alpha) \times V')(F_0)]_{\text{rs}}} \text{Orb}((\gamma, u'), \omega(h) \Phi', s),$$

and the term $\mathbb{J}(h, \Phi', s)_0$ is the sum over the two regular nilpotent orbits in (11.18), which will be defined in Section 12.6 by an analytic continuation. We have discarded the orbit $\{(0, 0)\}$ since η is then a non-trivial character on the stabilizer.

We first justify the convergence for the regular semisimple part $\mathbb{J}(h, \Phi', s)_{\text{rs}}$. We will defer the $H(F_0)$ -invariance to the next section; cf. Theorem 12.14.

LEMMA 11.2. (a) *For any $\Phi' \in \mathcal{S}((S_n \times V')(\mathbb{A}_0))$, the sum*

$$\sum_{(\gamma, u') \in [(S_n \times V')(F_0)]_{\text{rs}}} \int_{G'(\mathbb{A}_0)} |\Phi'(g^{-1} \cdot (\gamma, u'))| |\det(g)|_{F_0}^s dg < \infty$$

converges absolutely and uniformly for s in any compact subset in \mathbb{C} . In particular, the same holds if we only sum over $(\gamma, u') \in [(S_n(\alpha) \times V')(F_0)]_{\text{rs}}$.

(b) *Assume that Φ' is K_∞ -finite for the maximal compact $K_\infty = \prod_v \text{SO}(2, \mathbb{R})$ of $H(F_{0,\infty})$. Then the sum in $\mathbb{J}(h, \Phi', s)_{\text{rs}}$ converges absolutely and uniformly for (h, s) in any compact subset of $H(\mathbb{A}_0) \times \mathbb{C}$.*

Proof. This follows the same argument as the proof of Lemma 11.1. \square

12. RTF with Gaussian test functions

We now simplify (11.15) (resp. (11.21)) for a fixed $\alpha \in \mathcal{A}_n(F_0)$ by plugging in a Gaussian test function at every archimedean place.

12.1. Gaussian test functions: the compact unitary group case. Now let F/F_0 be the the archimedean local field extension \mathbb{C}/\mathbb{R} . Let V be an n -dimensional *positive definite* hermitian space with the unitary group $G = \mathrm{U}(V)$. We define a special test function, called the Gaussian test function (cf. [40, §7]) in the semi-Lie algebra setting,

$$(12.1) \quad \Phi(g, u) = \mathbf{1}_{G(\mathbb{R})}(g) \cdot e^{-\pi\langle u, u \rangle} \in \mathcal{S}((G \times V)(\mathbb{R})).$$

Since it is invariant under $G(\mathbb{R})$, its orbital integrals (2.15) take a very simple form:

$$(12.2) \quad \mathrm{Orb}((g, u), \Phi) = e^{-\pi\langle u, u \rangle}.$$

Here we normalize the Haar measure on $G(\mathbb{R})$ such that $\mathrm{vol}(G(\mathbb{R})) = 1$.

We explicate the action of $\mathrm{SL}_2(\mathbb{R})$ by the Weil representation (for the fixed additive character $\psi : x \in \mathbb{R} \mapsto e^{2\pi i x}$). Write $h \in \mathrm{SL}_2(\mathbb{R})$ according to the Iwasawa decomposition

$$(12.3) \quad h = \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix} \begin{pmatrix} a^{1/2} & \\ & a^{-1/2} \end{pmatrix} \kappa_\theta, \quad a \in \mathbb{R}_+, \quad b \in \mathbb{R},$$

where $\kappa(\theta)$ is as in (1.7). First of all, the Gaussian test functions above are eigen-vectors of weight $k = n$ under the action of the maximal compact $\mathrm{SO}(2, \mathbb{R})$ of $\mathrm{SL}_2(\mathbb{R})$, i.e.,

$$(12.4) \quad \omega(\kappa_\theta)\Phi = \chi_n(\kappa_\theta)\Phi,$$

where χ_n is the character (1.8). In general, for h of the form (12.3),

$$\omega(h)\Phi(g, u) = \chi_n(\kappa_\theta) \mathbf{1}_{G(\mathbb{R})}(g) \otimes |a|^{1/2} e^{\pi i(b+ia)\langle u, u \rangle}.$$

12.2. Gaussian test functions: the general linear group case. On the general linear group side, we define Gaussian test functions to be any smooth transfer of the Gaussian test functions on the unitary side (cf. [40, §7]). We recall the bijection of regular semisimple orbits (2.7) and (2.9). Note that in the disjoint union, one component is from the positive definite hermitian space V . We defined the notion of transfer at the end of Section 2.3.

Definition 12.1. We call $\Phi' \in \mathcal{S}((S_n \times V'_n)(\mathbb{R}))$ a Gaussian test function if it is a transfer of the tuple $\{\Phi_V\}_V$ where Φ_V is the Gaussian test functions (12.1) for the positive definite hermitian space V , and $\Phi_V = 0$ for all the other (isometry classes of) hermitian spaces V .

It is expected that Gaussian test functions exist. However, it seems very difficult to explicate the Gaussian test functions on $(S_n \times V')(\mathbb{R})$ (with one exception: the case $n = 1$). Fortunately a weaker version suffices for our purpose. We only need a partial matching, i.e., only Schwartz functions that have matching orbital integrals for elements with a fixed component on S_n ; we will name them “partial Gaussian test functions.”

We call the subset T_n of diagonal elements in $S_n(\mathbb{R})$ the compact Cartan subspace of $S_n(\mathbb{R})$. We have

$$T_n \xrightarrow{\sim} \mathrm{U}(1)(\mathbb{R})^n.$$

Let T_n^{rs} denote the open subset of the regular semisimple elements in the Cartan subspace T_n (i.e., those with distinct diagonal entries).

Definition 12.2. Let Ω be a compact subset of T_n^{rs} . We call $\Phi' \in \mathcal{S}((S_n \times V')(\mathbb{R}))$ a partial Gaussian test function (relative to Ω) if, for all regular semisimple $(\gamma, u') \in \Omega \times V'$ matching $(\delta, u) \in (\mathrm{U}(V) \times V)(\mathbb{R})$ for the positive definite hermitian space V , we have

$$(12.5) \quad \mathrm{Orb}((\gamma, u'), \Phi') = \mathrm{Orb}((\delta, u), \Phi),$$

where in the right-hand side Φ is the Gaussian test functions (12.1), and $\mathrm{Orb}((\gamma, u'), \Phi') = 0$ whenever a regular semisimple (γ, u') matches an orbit from non-positive-definite hermitian spaces in (2.7).

Now we construct “partial Gaussian test functions” explicitly for any compact subset Ω of T_n^{rs} . We first consider the case $n = 1$ and then reduce the general case to $n = 1$.

12.3. Gaussian test functions when $n = 1$. Assume $n = \dim V = 1$. Then $G'(\mathbb{R}) \simeq \mathbb{R}^\times$, and the symmetric space $S_1(\mathbb{R})$ is compact. The orbital integrals have been defined in Section 2.3; cf. (2.14). Since the G' -action on S_1 is trivial, we simply work with the vector space component and suppress the $\gamma \in S_1(\mathbb{R})$ in the orbital integrals.

Let $V = \mathbb{C}$ be 1-dimensional hermitian space (with the standard norm), and let

$$\phi(z) = e^{-\pi z\bar{z}} \in \mathcal{S}(V).$$

Then we have $\widehat{\phi} = \phi$.

Let $V' \simeq \mathbb{R} \times \mathbb{R}$, with \mathbb{R}^\times -action

$$t \cdot (x, y) = (t^{-1}x, ty).$$

Recall from (11.17) that the quadratic form on V' is $\mathfrak{q}(x, y) = xy$. We consider the following Schwartz function in the Fock model:

$$(12.6) \quad \phi'(x, y) = 2^{-3/2}(x + y)e^{-\frac{1}{2}\pi(x^2 + y^2)} \in \mathcal{S}(\mathbb{R} \times \mathbb{R}).$$

It has the symmetry

$$\phi'(x, y) = \phi'(y, x), \quad \phi'(-x, -y) = -\phi'(x, y).$$

Recall that the K -Bessel function is defined as

$$K_s(c) = \frac{1}{2} \int_{\mathbb{R}_+} e^{-\frac{1}{2}c(u+1/u)} u^s \frac{du}{u}, \quad c > 0, s \in \mathbb{C}.$$

LEMMA 12.3. *Let $\xi \in \mathbb{R}^\times$. Then*

$$\text{Orb}((1, \xi), \phi', s) = 2^{-1/2} |\xi|^{(-s+1)/2} (K_{(s+1)/2}(\pi|\xi|) + \eta(\xi) K_{(s-1)/2}(\pi|\xi|)).$$

In particular,

$$\text{Orb}((1, \xi), \phi') = \begin{cases} e^{-\pi\xi}, & \xi > 0, \\ 0, & \xi < 0, \end{cases}$$

and when $\xi < 0$,

$$\partial \text{Orb}((1, \xi), \phi') = \frac{1}{2} e^{-\pi\xi} \text{Ei}(-2\pi|\xi|).$$

Here Ei is the exponential integral (8.10).

Remark 12.4. Here the special value at $s = 0$ has taken into account the transfer factors; cf. Section 2.3.

Proof. By definition of orbital integrals (2.14) (except we have suppressed the \mathfrak{s}_1 and S_1 component), we have

$$\begin{aligned} \text{Orb}((1, \xi), \phi', s) &= 2^{-1/2} \int_{\mathbb{R}_+} (t + \eta(\xi)|\xi|/t) e^{-\frac{1}{2}\pi(t^2 + \xi^2/t^2)} t^{-s} \frac{dt}{t} \\ &= 2^{-1/2} |\xi|^{(-s+1)/2} \int_{\mathbb{R}_+} (t + \eta(\xi)/t) e^{-\frac{1}{2}\pi|\xi|(t^2 + 1/t^2)} t^{-s} \frac{dt}{t} \\ &= 2^{-1/2} |\xi|^{(-s+1)/2} \int_{\mathbb{R}_+} e^{-\frac{1}{2}\pi|\xi|(t^2 + 1/t^2)} (t^{-s+1} + \eta(\xi)t^{-s-1}) \frac{dt}{t} \\ &= 2^{-3/2} |\xi|^{(-s+1)/2} \int_{\mathbb{R}_+} e^{-\frac{1}{2}\pi|\xi|(u+1/u)} (u^{(-s+1)/2} + \eta(\xi)u^{(-s-1)/2}) \frac{du}{u} \\ &= 2^{-1/2} |\xi|^{(-s+1)/2} (K_{(-s+1)/2}(\pi|\xi|) + \eta(\xi) K_{(-s-1)/2}(\pi|\xi|)). \end{aligned}$$

To evaluate at $s = 0$, we note that

$$K_{1/2}(\xi) = \sqrt{\frac{\pi}{2}} \frac{e^{-\xi}}{\xi^{1/2}}.$$

Also we note that the transfer factor (2.17) takes value one at elements of the form $(1, \xi)$, applied to $F/F_0 = \mathbb{C}/\mathbb{R}$.

The assertion for the first derivative follows from the following identity [35]:

$$\frac{d}{ds} \Big|_{s=1/2} K_s(y) = -\sqrt{\frac{\pi}{2}} \frac{e^y}{y^{1/2}} \operatorname{Ei}(-2y), \quad y > 0. \quad \square$$

We now explicate the action of $\operatorname{SL}_2(\mathbb{R})$ by the Weil representation ω . Similar to the unitary case, the Gaussian test functions above are eigen-vectors of weight $k = n = 1$ under the action of the maximal compact $\operatorname{SO}(2, \mathbb{R})$; cf. (12.4), (1.8). Write $h \in \operatorname{SL}_2(\mathbb{R})$ according to the Iwasawa decomposition

$$h = \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix} \begin{pmatrix} a^{1/2} & \\ & a^{-1/2} \end{pmatrix} \kappa_\theta, \quad a \in \mathbb{R}_+, \quad b \in \mathbb{R},$$

where $\kappa_\theta \in \operatorname{SO}(2, \mathbb{R})$ is as in (1.7).

LEMMA 12.5. *Let $\xi \in \mathbb{R}^\times$. Then*

$$\begin{aligned} \operatorname{Orb}((1, \xi), \omega(h)\phi', s) \\ = 2^{-1/2} \chi_1(\kappa_\theta) a |\xi|^{(-s+1)/2} (K_{(-s+1)/2}(\pi a |\xi|) + \eta(\xi) K_{(-s-1)/2}(\pi a |\xi|)). \end{aligned}$$

In particular,

$$\operatorname{Orb}((1, \xi), \omega(h)\phi') = \begin{cases} a^{1/2} e^{\pi i \xi(b+ia)}, & \xi > 0, \\ 0, & \xi < 0, \end{cases}$$

and when $\xi < 0$,

$$\partial \operatorname{Orb}((1, \xi), \omega(h)\phi') = \frac{1}{2} \chi_1(\kappa_\theta) a^{1/2} e^{\pi i |\xi|(b-ia)} \operatorname{Ei}(-2\pi a |\xi|).$$

Here Ei is the exponential integral (8.10).

Proof. This follows by straightforward computation using Lemma 12.3, and the formulas (11.1) defining the Weil representation in Section 11.1. \square

12.4. *Partial Gaussian test functions: general n .* We will use the Iwasawa decomposition of the group $G'(\mathbb{R}) = \operatorname{GL}_n(\mathbb{R})$,

$$(12.7) \quad G'(\mathbb{R}) = ANK,$$

where $K = \operatorname{SO}(n, \mathbb{R})$, N is the group of unipotent upper triangular matrices, and $A \simeq (\mathbb{R}^\times)^n$ is the diagonal torus. We have a homeomorphism

$$(12.8) \quad G'(\mathbb{R}) \simeq AN \times_{\mu_2^{n-1}} K$$

as real manifolds, where the fiber product is over the intersection $AN \cap K$, which is equal to

$$K \cap A = \ker(\mu_2^n \rightarrow \mu_2) \simeq \mu_2^{n-1}.$$

We will take the natural Haar measure on each factor (e.g., the measure $\frac{dt}{|t|}$ on \mathbb{R}^\times and the product measure on $(\mathbb{R}^\times)^n \simeq A$) and take the induced measure on $G'(\mathbb{R})$ by the above product (12.8).

Note that the torus A is the stabilizer of a regular semisimple element in the Cartan subspace T_n . Then $NK \cdot T_n^{\text{rs}}$ (the conjugation action) defines an open subset $S_n^{c,\text{rs}}$ (“c” is for “compact”) in S_n :

$$\begin{aligned} NK \times T_n^{\text{rs}} &\xrightarrow{\sim} S_n^{c,\text{rs}} \subset S_n \\ (h, t) &\longmapsto h^{-1}th. \end{aligned}$$

The map is a $K \cap A$ -torsor and induces a $K \cap A$ -torsor:

$$\begin{aligned} (12.9) \quad NK \times T_n^{\text{rs}} \times (V_0 \times V_0^*) &\longrightarrow S_n^{c,\text{rs}} \times (V_0 \times V_0^*) \\ (h, t, u') &\longmapsto (h^{-1}th, h \cdot u'). \end{aligned}$$

Now let $\Omega \subset T_n^{\text{rs}}$ be any compact subset. We consider functions on $NK \times T_n^{\text{rs}} \times (V_0 \times V_0^*)$ of the form $\Psi = \phi_0 \otimes \phi'$, with $\phi' \in \mathcal{S}(V_0 \times V_0^*)$ and

$$(12.10) \quad \phi_0 = \varphi_N \otimes \varphi_K \otimes \varphi_{T_n},$$

where

- (1) the function $\varphi_{T_n} \in C_c^\infty(T_n^{\text{rs}})$ satisfies $\varphi_{T_n}|_\Omega = 1_\Omega$,
- (2) the function $\varphi_N \in C_c^\infty(N)$ satisfies $\int_N \varphi_N(n) dn = 1$,
- (3) the function φ_K is a constant multiple of 1_K such that $\int_K \varphi_K(k) dk = 1$,
- (4) the function ϕ' is invariant under the finite group $K \cap A$.

By the $K \cap A$ -invariance of ϕ_0 and ϕ' , the function $\Psi = \phi_0 \otimes \phi'$ descends along the map (12.9) to a Schwartz function Φ'^c on $S_n^{c,\text{rs}} \times (V_0 \times V_0^*)$. Then the extension-by-zero of Φ'^c , denoted by Φ' , is a Schwartz function on $S_n \times (V_0 \times V_0^*)$.

Finally we specify ϕ' on $V_0 \times V_0^*$. Identify $V_0 \times V_0^*$ with $\mathbb{R}^n \times \mathbb{R}^n \simeq (\mathbb{R} \times \mathbb{R})^n$, and we define

$$(12.11) \quad \phi' = 2^{-3n/2} \prod_{1 \leq i \leq n} (x_i + y_i) e^{-\frac{1}{2}\pi(x_i^2 + y_i^2)};$$

cf. (12.6) for the case $n = 1$. It is obviously invariant under $K \cap A$. Therefore by our recipe this function ϕ' (with any ϕ_0 above) gives us a Schwartz function Φ' on $S_n \times (V_0 \times V_0^*)$.

Now we define the orbital integral $\text{Orb}(u', \phi', s)$ for $u' \in V_0 \times V_0^*$, relative to the A -action on $V_0 \times V_0^*$, in the obvious way generalizing the case $n = 1$; cf. (2.14).

LEMMA 12.6. *Let $\gamma \in \Omega \subset T_n^{\text{rs}}$. Then for any regular semisimple (γ, u') , we have*

$$\text{Orb}((\gamma, u'), \Phi', s) = \text{Orb}(u', \phi', s),$$

where the left-hand side is the local orbital integral (11.20).

In particular, by Lemma 12.3 and (12.2), the function Φ' is a partial Gaussian test function (relative to the compact subset Ω).

Proof. By the Iwasawa decomposition (12.7), the local orbital integral (11.20) is equal to

$$\int_A \int_{NK} \Phi'((nk)^{-1} \cdot (\gamma, a^{-1} \cdot u')) |\det(a)|^s \eta(a) dn dk da.$$

By our choice of Φ' , we obtain

$$\begin{aligned} & \int_{NK} \Phi'((nk)^{-1} \cdot (\gamma, u')) dn dk \\ &= \left(\int_N \varphi_N(n) dn \int_K \varphi_K(k) dk \right) \varphi_{T_n}(\gamma) \phi'(u') \\ &= \phi'(u'). \end{aligned}$$

Therefore

$$\text{Orb}((\gamma, u'), \Phi', s) = \int_A \phi'(a^{-1} \cdot u') |\det(a)|^s \eta(a) da = \text{Orb}(u', \phi', s).$$

This completes the proof. \square

Remark 12.7. This result also holds if u' is a regular nilpotent orbit and the orbital integral is regularized by (12.18) and (12.27) below.

12.5. *Modular analytic generating functions when $n = 1$.* Now we return to the global situation Section 11.4. Assume that $n = 1$. Then we may identify $V' = F_0 \times F_0$, and the special orthogonal group $\text{SO}(V', \mathfrak{q})$ can be identified with the F_0 -group $G' := \text{GL}_{1, F_0}$, via the action on the V' by $g \cdot (u_1, u_2) = (g^{-1}u_1, gu_2)$. The map $u' = (u_1, u_2) \mapsto \xi = \mathfrak{q}(u') = u_1 u_2$ identifies the categorical quotient $V'_{/G'}$ with the affine line. Note that regular semisimple orbits (for the G' -action) are exactly the fibers over $\xi \neq 0$, and each fiber has exactly one G' -orbit.

Let $\phi' \in \mathcal{S}(V'(\mathbb{A}_0))$. Consider the integral,

$$(12.12) \quad \mathbb{J}(\phi', s) = \int_{[G']} \left(\sum_{u' \in V'(F_0)} \phi'(g^{-1} \cdot u') \right) |g|^s \eta(g) dg.$$

The integral is not necessarily convergent, and we define it by a regularization procedure as follows.

Recall from (11.21) that we can write the integrand as a sum over the $G'(F_0)$ -orbits in $V'(F_0)$. Then the regular semisimple part is

$$(12.13) \quad \sum_{\xi = \mathfrak{q}(u') \in F_0^\times} \text{Orb}(u', \phi', s),$$

where

$$(12.14) \quad \text{Orb}(u', \phi', s) := \int_{G'(\mathbb{A}_0)} \phi'(g^{-1} \cdot u') |g|^s \eta(g) dg.$$

By [Lemma 11.2](#), the sum in [\(12.13\)](#) converges absolutely and uniformly for s a compact set in \mathbb{C} .

The fiber over $\xi = 0$ breaks into three orbits:

$$\begin{cases} \{(0, 0)\}, \\ 0_+ = \{(u_1, 0) : u_1 \in F_0^\times\}, \\ 0_- = \{(0, u_2) : u_2 \in F_0^\times\}. \end{cases}$$

The stabilizer of the first one is G' , and the other two have a trivial stabilizer. Note that η is non-trivial on $G'(\mathbb{A}_0)$, and hence we *define* the integral for the first orbit to be zero. For the other two orbits, we define

$$(12.15) \quad \text{Orb}(0_+, \phi', s) := \int_{\mathbb{A}_0^\times} \phi'(g, 0) |g|^s \eta(g) dg$$

and

$$(12.16) \quad \text{Orb}(0_-, \phi', s) = \int_{\mathbb{A}_0^\times} \phi'(0, g^{-1}) |g|^s \eta(g) dg = \int_{\mathbb{A}_0^\times} \phi'(0, g) |g|^{-s} \eta(g) dg.$$

Both will be understood as Tate's global zeta integrals. More precisely,

$$(12.17) \quad \text{Orb}(0_+, \phi', s) = L(s, \eta) \prod_v \text{Orb}(0_+, \phi'_v, s),$$

where the local orbital integral for the regular nilpotent 0_+ is defined as (the analytic continuation of)

$$(12.18) \quad \text{Orb}(0_+, \phi'_v, s) := \frac{\int_{F_{0,v}^\times} \phi'_v(g, 0) |g|_v^s \eta_v(g) dg}{L(s, \eta_v)}.$$

We note that the local Tate integral [\(12.18\)](#) is absolutely convergent when $\text{Re}(s) > 0$ extends to an entire function of s (a polynomial in $q_v^{\pm s}$ when v is non-archimedean) and equal to one for unramified data. Here $L(s, \eta)$ is the *complete L-function* of the Hecke character η . Similarly, for 0_- , we have

$$(12.19) \quad \text{Orb}(0_-, \phi', s) = L(-s, \eta) \prod_v \text{Orb}(0_-, \phi'_v, -s),$$

where

$$\text{Orb}(0_-, \phi'_v, -s) = \frac{\int_{F_{0,v}^\times} \phi'_v(0, g) |g|_v^{-s} \eta_v(g) dg}{L(-s, \eta_v)}.$$

To summarize, we define [\(12.12\)](#) as the sum of [\(12.13\)](#), [\(12.15\)](#), and [\(12.16\)](#) (or rather, their analytic continuation to $s \in \mathbb{C}$):

$$(12.20) \quad \mathbb{J}(\phi', s) = \text{Orb}(0_+, \phi', s) + \text{Orb}(0_-, \phi', s) + \sum_{\xi = \mathfrak{q}(u') \in F_0^\times} \text{Orb}(u', \phi', s).$$

Define the analytic generating function on $\mathbf{H}(\mathbb{A}_0)$,

$$\mathbb{J}(h, \phi', s) = \mathbb{J}(\omega(h)\phi', s), \quad h \in \mathbf{H}(\mathbb{A}_0).$$

Remark 12.8. The function $\mathbb{J}(\cdot, \phi', s)$ may be viewed as the generating function of the above relative orbital integrals (12.14), (12.15), and (12.16), parametrized by $\xi \in F_0$. This is the analytic analog of the modular generating function of special divisors in Section 8.

THEOREM 12.9. *The function $\mathbb{J}(h, \phi', s)$ is smooth in $h \in H(\mathbb{A}_0)$ and entire in $s \in \mathbb{C}$. As a smooth function in $h \in H(\mathbb{A}_0)$, it is left invariant under $H(F_0)$.*

Proof. By Lemma 11.2, the smoothness and the entireness follow from the same property for each of (12.13), (12.15), and (12.16). To show the $H(F_0)$ -invariance, we first note that the invariance under the upper triangular elements follow from the definition of the Weil representation and that of the function $\mathbb{J}(h, \phi', s)$. It remains to show the invariance under $w = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$, i.e., the functional equation

$$(12.21) \quad \mathbb{J}(\phi', s) = \mathbb{J}(\widehat{\phi'}, s)$$

holds for all ϕ' .

By Poisson summation formula (note that the action of $G'(\mathbb{A}_0)$ commutes with the Weil representation),

$$\sum_{u' \in V'} \phi'(g^{-1} \cdot u') = \sum_{u' \in V'} \widehat{\phi'}(g^{-1} \cdot u'), \quad g \in G'(\mathbb{A}_0),$$

or equivalently,

$$(12.22) \quad \begin{aligned} & \sum_{u' \in V', \xi \neq 0} \phi'(g^{-1} \cdot u') - \sum_{u' \in V', \xi \neq 0} \widehat{\phi'}(g^{-1} \cdot u') \\ &= - \sum_{u' \in V'_{\xi=0}} \phi'(g^{-1} \cdot u') + \sum_{u' \in V'_{\xi=0}} \widehat{\phi'}(g^{-1} \cdot u'), \quad g \in G'(\mathbb{A}_0). \end{aligned}$$

We introduce a partial Fourier transform $\phi' \mapsto \mathcal{F}_1(\phi')$ for one of the two variables:

$$\mathcal{F}_1(\phi')(u_1, u_2) = \int_{\mathbb{A}_0} \phi'(u_2, w_2) \psi(-u_1 w_2) dw_2.$$

Apply Poisson summation formula to the line $u_1 = 0$:

$$\sum_{u_2 \in F_0} \phi'(0, g^{-1} u_2) = |g| \sum_{u_1 \in F_0} \mathcal{F}_1(\phi')(g u_1, 0).$$

We obtain an alternative expression of the right-hand side of (12.22) as the sum of

$$- \sum_{u_1 \in F_0} (\phi'(g u_1, 0) + |g| \mathcal{F}_1(\phi')(g u_1, 0)) + \phi'(0, 0)$$

and

$$\sum_{u'_1 \in F_0} (\widehat{\phi'}(g u_1, 0) + |g| \mathcal{F}_1(\widehat{\phi'})(g u'_1, 0)) - \widehat{\phi'}(0, 0).$$

Denote $[G']^1 = G'(F_0) \backslash G'(\mathbb{A}_0)^1$, where

$$G'(\mathbb{A}_0)^1 := \ker(|\det| : G'(\mathbb{A}_0) \longrightarrow \mathbb{R}_+).$$

We embed \mathbb{R}_+ into $G'(\mathbb{A}_0) = \mathbb{A}_0^\times$ by sending $t \in \mathbb{R}_+$ to (t_v) , where

$$t_v = \begin{cases} t^{1/[F_0:\mathbb{Q}]}, & v \mid \infty, \\ 1, & v \nmid \infty. \end{cases}$$

Then we have a direct product

$$(12.23) \quad \begin{aligned} G'(\mathbb{A}_0) &\xrightarrow{\sim} G'(\mathbb{A}_0)^1 \times \mathbb{R}_+ \\ g &\longmapsto (g_1, t). \end{aligned}$$

Since the quotient $[G']^1$ is compact, we may integrate (12.22) over $[G']^1$ first, and this kills the zero orbits (due to the non-triviality of $\eta|_{[G']^1}$). Then we integrate over \mathbb{R}_+ . (Now we use the alternative expression of the right-hand side.) Since the Tate integrals converges absolutely when $\text{Re}(s) > 1$, we obtain

$$\begin{aligned} &\text{Orb}(0_+, \phi', s) + \text{Orb}(0_+, \mathcal{F}_1(\phi'), 1+s) + \sum_{\xi \in F_0^\times} \text{Orb}(u', \phi', s) \\ &= \text{Orb}(0_+, \widehat{\phi}', s) + \text{Orb}(0_+, \mathcal{F}_1(\widehat{\phi}'), 1+s) + \sum_{\xi \in F_0^\times} \text{Orb}(u', \widehat{\phi}', s) \end{aligned}$$

when $\text{Re}(s) > 1$. Finally, we note that by (12.19) and the functional equation of Tate integrals,

$$\text{Orb}(0_-, \phi', s) = \text{Orb}(0_+, \mathcal{F}_1(\phi'), 1+s).$$

By analytic continuation, this completes the proof of (12.21) for all $s \in \mathbb{C}$. \square

Remark 12.10. The integral (12.12) can be viewed as the theta lifting for the pair

$$(\text{SO}(V', \mathfrak{q}), \text{SL}_2),$$

from the automorphic representation $\eta|\cdot|^s$ of $\text{SO}(V') \simeq \text{GL}_1$ to SL_2 . Therefore, the representation space spanned by $h \mapsto \mathbb{J}(h, \phi', s)$ is the space of degenerate Eisenstein series for the induced representation $\text{Ind}_{B(\mathbb{A}_0)}^{H(\mathbb{A}_0)}(\eta|\cdot|^s)$. (Here B is the Borel subgroup of upper triangular matrices). In this way, the two nilpotent orbital integrals become the constant terms of the associated Eisenstein series.

LEMMA 12.11. *Let $v \mid \infty$, and let ϕ'_v be the Gaussian test function (12.6). Then the local nilpotent orbital integral (12.18) is equal to*

$$\text{Orb}(0_+, \phi'_v, s) = 2^{\frac{s}{2}-1}.$$

The action of the group $\mathrm{SL}_2(\mathbb{R})$ is given as follows, for $h \in \mathrm{SL}_2(\mathbb{R})$ in the form of (1.6):

$$\mathrm{Orb}(0_+, \omega(h)\phi', s) = \chi_1(\kappa_\theta) a^{(-s+1)/2} 2^{\frac{s}{2}-1}.$$

Proof. By (12.6), we obtain $\phi'_v(x, 0) = 2^{-3/2} x e^{-\frac{1}{2}\pi x^2}$. Then $\mathrm{Orb}(0_+, \phi'_v, s)$ is the Tate's local zeta integral at an archimedean place:

$$\begin{aligned} 2 \int_{\mathbb{R}_+} e^{-\frac{1}{2}\pi x^2} |x|^{s+1} \frac{dx}{x} &= \int_{\mathbb{R}_+} e^{-\frac{1}{2}\pi x} |x|^{(s+1)/2} \frac{dx}{x} \\ &= (\pi/2)^{-(s+1)/2} \Gamma((s+1)/2). \end{aligned}$$

Note the local L-factor in (12.18) is by definition

$$L(s, \eta) = \pi^{-(s+1)/2} \Gamma\left(\frac{s+1}{2}\right).$$

We obtain

$$\mathrm{Orb}(0_+, \phi'_v, s) = 2^{\frac{s}{2}-1}.$$

The action of $\mathrm{SL}_2(\mathbb{R})$ is determined in a way similar to Lemma 12.5. \square

12.6. Modular analytic generating functions for general n . We now return to the setting of Section 11.4 for general n .

Recall from Section 7.4 that we have fixed $\alpha \in \mathcal{A}_n(F_0) \subset F[T]_{\deg=n}$ irreducible over F , the field $F' = F[T]/(\alpha)$ and its subfield F'_0 . Then $S_n(\alpha)(F_0)$ consists of exactly one $G'(F_0)$ -orbit; let us fix a representative $\gamma \in S_n(\alpha)(F_0)$. Denote by T' the stabilizer of γ , which is isomorphic to $\mathrm{Res}_{F'_0/F_0} \mathbb{G}_m$. It follows that the character $\eta \circ \det$ (of $G'(\mathbb{A}_0)$) is nontrivial on $T'(\mathbb{A}_0)$, which can be identified (via $T' \simeq \mathrm{Res}_{F'_0/F_0} \mathbb{G}_m$) with the quadratic character associated to F'/F'_0 by class field theory

$$\eta' = \eta_{F'/F'_0} : \mathbb{A}_{F'_0}^\times \longrightarrow \{\pm 1\}.$$

Similar to the F'/F'_0 -hermitian form (10.5), via the action of F'_0 , the vector space V_0 (hence V_0^*) carries a structure of a one-dimensional F'_0 -vector space. Furthermore, we can identify

$$\mathrm{Hom}_{F'_0}(V_0, F'_0) \simeq V_0^*$$

as one-dimensional F'_0 -vector spaces. There is a unique bi- F'_0 -linear symmetric pairing

$$(12.24) \quad \langle \cdot, \cdot \rangle_{F'_0} : V' \times V' \longrightarrow F'_0$$

such that

$$\langle u_1, u_2 \rangle = \mathrm{tr}_{F'_0/F_0} \langle u_1, u_2 \rangle_{F'_0}.$$

Let

$$(12.25) \quad \mathfrak{q}' : V_0 \times V_0^* \longrightarrow F'_0$$

be the associated quadratic form over F'_0 .

Definition 12.12. An element $\gamma \in S_n(F_0)$ is compact, if locally at all places $v \mid \infty$, it lies in the $G'(F_{0,v})$ -orbit of the compact Cartan subspace $T_n(F_{0,v})$.

Then $\gamma \in S_n(\alpha)(F_0)$ is compact if and only if the field F' is a CM extension of a totally real field F'_0 , which we have assumed since [Section 7.4](#).

Now, for every $v \mid \infty$, we fix the archimedean $\Phi'_v \in \mathcal{S}((S_n \times V')(F_{0,v}))$ to be the partial Gaussian test function constructed in [Section 12.4](#) (relative to a fixed compact neighborhood $\Omega_v \subset S_n(F_{0,v})$ of γ). Recall that Φ'_v is associated to the function ϕ'_v defined by [\(12.11\)](#).

There are two regular nilpotent orbits for the T' -action on $V'(F_0)$, denoted by 0_{\pm} in [\(11.18\)](#). We now define the constant term $\mathbb{J}(h, \Phi', s)_0$ in [\(11.21\)](#) as the sum of the two regular γ -nilpotent orbital integrals $\text{Orb}((\gamma, 0_{\pm}), \Phi', s)$ in a similar way to [\(12.15\)](#). More precisely, we define

$$(12.26) \quad \text{Orb}((\gamma, 0_+), \Phi', s) := L(s, \eta') \prod_v \text{Orb}((\gamma, 0_+), \Phi'_v, s),$$

where the local orbital integral is defined as

$$(12.27) \quad \text{Orb}((\gamma, 0_+), \Phi'_v, s) = \frac{\int_{G'(F_{0,v})} \Phi'_v(g^{-1} \cdot (\gamma, 0_+)) |\det(g)|_v^s \eta(g) dg}{L(s, \eta'_v)}.$$

Here the denominator is defined as

$$L(s, \eta'_v) = \prod_{v' \mid v} L(s, \eta'_{v'}),$$

where v' runs over all places of F'_0 above v . Note that $L(s, \eta') = L(s, \text{Ind}_{F'_0}^{F_0} \eta')$. We define $\text{Orb}((\gamma, 0_-), \Phi', s)$ similarly. Here we normalize the measure on $G'(F_{0,v})$ such that $\text{vol}(G'(O_{F_{0,v}})) = 1$ for all but finitely non-archimedean places v .

LEMMA 12.13. *The integral [\(12.27\)](#) is absolutely convergent when $\text{Re}(s) > 0$ extends to an entire function of s (a polynomial of $q_v^{\pm s}$ for non-archimedean v).*

Furthermore, for a fixed γ and a pure tensor $\Phi = \otimes_v \Phi_v$, where $\Phi'_v = \mathbf{1}_{(S_n \times V')(O_{F_{0,v}})}$ for all but finitely many v , the integral [\(12.27\)](#) is equal to one for all but finitely many places v (depending on γ and Φ).

Proof. When $v \mid \infty$, by [Lemma 12.6](#), the desired claim follows from (the product of n copies of) the same claim for $n = 1$.

Now let v be non-archimedean, and let Φ'_v be as in [Section 12.4](#). We fix a large compact subset Ω_v of $G'(F_{0,v})$ such that $\Phi'_v(g^{-1} \cdot \gamma, u') = 0$ unless

$g \in \Omega_v \cdot T'(F_{0,v})$. We introduce a Schwartz function (with a parameter $s \in \mathbb{C}$) on $V'(F_{0,v})$,

$$(12.28) \quad \phi'_{v,s}(u') := \int_{\Omega_v} \Phi'_v(g^{-1} \cdot (\gamma, u')) |\det(g)|_v^s \eta_v(g) dg.$$

It is easy to see that it is of the form

$$(12.29) \quad \phi'_{v,s} = \sum_{1 \leq i \leq m} a_i \lambda_i^s \phi_{v,i},$$

where

$$a_i \in \mathbb{Q}, \quad \lambda_i \in \mathbb{Q}_+^\times, \quad \phi_i \in \mathcal{S}(V'(F_{0,v})).$$

Then, for a suitable choice of measure dg on Ω_v in the integral (12.28),

$$(12.30) \quad \text{Orb}((\gamma, 0_+), \Phi'_v, s) = \text{Orb}(0_+, \phi'_{v,s}, s).$$

Here we view V_0 as a one-dimensional F'_0 -vector space and V_0^* as its F'_0 -dual vector space, and the right-hand side is (12.19) relative to the quadratic extension $F'_v/F'_{0,v}$ at v (i.e., F'_v is the product of $F'_{v'} = F' \otimes_{F'_0} F'_{0,v'}$ over all places v' of F'_0 over v). This shows that the local orbital integral for 0_+ is a polynomial of $q_{v'}^{\pm s}, v'|v$, particularly, an entire function in s .

Finally, let us assume that v is unramified in F' and $\Phi'_v = \mathbf{1}_{(S_n \times V')(\mathcal{O}_{F_{0,v}})}$. (Here we implicitly identified $V_0 = F_0^n$ and endow it with the natural integral structure.) For all but finitely many places v , the element γ belongs to $S_n(\mathcal{O}_{F_{0,v}})$ and generates the maximal order $\mathcal{O}_{F'_v}$ in F'_v . Then it is easy to see that $\phi'_{v,s} = \mathbf{1}_{V'(\mathcal{O}_{F_{0,v}})}$ in (12.30), and hence the integral is equal to one by the standard computation of Tate's local zeta integral for unramified data. \square

THEOREM 12.14. *The function $(h, s) \in \mathbf{H}(\mathbb{A}_0) \times \mathbb{C} \mapsto \mathbb{J}(h, \Phi', s)$ is entire in $s \in \mathbb{C}$ and left invariant under $\mathbf{H}(F_0)$.*

Proof. By the proof of Lemma 12.13, (12.29) and (12.30), there exists a finite collection of

$$a_i \in \mathbb{Q}, \quad \lambda_i \in \mathbb{Q}_+^\times, \quad \phi_i \in \mathcal{S}(V'(\mathbb{A}_0))$$

such that

$$\mathbb{J}(h, \Phi', s) = \sum_{1 \leq i \leq m} a_i \lambda_i^s \mathbb{J}(h, \phi'_i, s).$$

Here we note that for almost all places, the $\phi'_{v,s} = \mathbf{1}_{V'(\mathcal{O}_{F_{0,v}})}$ in (12.30), and the ϕ'_i are of the form $\otimes_v \phi'_{v,i_v}$ for ϕ'_{v,i_v} from (12.29). The desired claims follow now from Theorem 12.9 for $n = 1$, applied to the new quadratic extension F'/F'_0 . \square

For simplicity, we combine the two nilpotent orbital integrals into one:

$$(12.31) \quad \text{Orb}((\gamma, 0_{\pm}), \Phi', s) := \text{Orb}((\gamma, 0_+), \Phi', s) + \text{Orb}((\gamma, 0_-), \Phi', s).$$

Then we obtain an expansion as a sum of orbital integrals

$$(12.32) \quad \begin{aligned} \mathbb{J}(h, \Phi', s) &= \sum_{\substack{(\gamma, u') \in [(S_n(\alpha) \times V')(\mathbb{F}_0)] \\ u' \neq 0}} \text{Orb}((\gamma, u'), \omega(h)\Phi', s) \\ &= \text{Orb}((\gamma, 0_{\pm}), \omega(h)\Phi', s) \\ &\quad + \sum_{\substack{(\gamma, u') \in [(S_n(\alpha) \times V')(\mathbb{F}_0)]_{\text{rs}}}} \text{Orb}((\gamma, u'), \omega(h)\Phi', s). \end{aligned}$$

Moreover, for $\xi \in \mathbb{F}_0^{\times}$, the ξ -th Fourier coefficient of $\mathbb{J}(\cdot, \Phi', s)$ is equal to

$$(12.33) \quad \sum_{(\gamma, u') \in [(S_n(\alpha) \times V'_{\xi})(\mathbb{F}_0)]} \text{Orb}((\gamma, u'), \omega(h)\Phi', s).$$

This is the analog of (11.16) on the unitary side.

12.7. *The decomposition of the special value at $s = 0$.* We set

$$\begin{aligned} \mathbb{J}(h, \Phi') &:= \mathbb{J}(h, \Phi', 0), \\ \text{Orb}((\gamma, u'), \omega(h)\Phi') &:= \text{Orb}((\gamma, u'), \omega(h)\Phi', 0). \end{aligned}$$

Then the decomposition (12.32) specializes to

$$(12.34) \quad \mathbb{J}(h, \Phi') = \sum_{\substack{(\gamma, u') \in [(S_n(\alpha) \times V')(\mathbb{F}_0)] \\ u' \neq 0}} \text{Orb}((\gamma, u'), \omega(h)\Phi').$$

We set

$$(12.35) \quad \begin{aligned} \partial \mathbb{J}(h, \Phi') &:= \frac{d}{ds} \Big|_{s=0} \mathbb{J}(h, \Phi', s), \\ \partial \text{Orb}((\gamma, u'), \Phi'_v) &:= \frac{d}{ds} \Big|_{s=0} \text{Orb}((\gamma, u'), \Phi'_v, s). \end{aligned}$$

(The second equation also applies to the nilpotent orbit, in which case the local orbital integrals are defined by (12.27).)

Now we introduce

$$\partial \mathbb{J}_v(h, \Phi') := \partial \mathbb{J}_v(\omega(h)\Phi'), \quad \text{where}$$

$$(12.36) \quad \partial \mathbb{J}_v(\Phi') := \sum_{\substack{(\gamma, u') \in [(S_n(\alpha) \times V')(\mathbb{F}_0)] \\ u' \neq 0}} \partial \text{Orb}((\gamma, u'), \Phi'_v) \cdot \text{Orb}((\gamma, u'), \Phi'^v).$$

In the nilpotent case, $\text{Orb}((\gamma, 0_{\pm}), \Phi'^v)$ is interpreted as

$$L(0, \eta') \prod_{w \neq v} \text{Orb}((\gamma, 0_{\pm}), \Phi'^v);$$

cf. (12.26).

We define

$$\partial\text{Orb}(0_+, \Phi') = \frac{d}{ds} \Big|_{s=0} L(s, \eta) \prod_v \text{Orb}((\gamma, 0_+), \Phi'_v, 0),$$

and similarly for $\partial\text{Orb}(0_-, \omega(h)\Phi')$. Then we define

$$(12.37) \quad \partial\text{Orb}(0_\pm, \Phi') = \partial\text{Orb}(0_+, \Phi') + \partial\text{Orb}(0_-, \Phi').$$

Then by Leibniz's rule, we obtain a decomposition:

$$(12.38) \quad \partial\mathbb{J}(h, \Phi') = \partial\text{Orb}(0_\pm, \omega(h)\Phi') + \sum_v \partial\mathbb{J}_v(\omega(h)\Phi').$$

We call $\partial\text{Orb}(0_\pm, \omega(h)\Phi')$ the nilpotent term; it is part of the constant term (i.e., the 0-th Fourier coefficient).

Part 3. Proof of the main theorems

13. The proof of FL

13.1. *Smooth transfer: the global situation.* In [Section 2.3](#), we have defined the local transfer factor; cf. [\(2.17\)](#). The definition depends on a choice of an extension $\tilde{\eta}$ of the quadratic character η attached to the local quadratic extension. In the global case, we fix an extension of the quadratic character η_{F/F_0} of $F_0^\times \backslash \mathbb{A}_0^\times$ to a character $\tilde{\eta}$ of $F^\times \backslash \mathbb{A}^\times$ (not necessarily of order 2). The transfer factor for a global element then satisfies a product formula and transforms according to the desired rule; cf. [\[40, §7.3\]](#).

We are now in the setting of [Section 12.6](#); in particular, we have fixed an irreducible $\alpha \in \mathcal{A}_n(F_0) \subset F[T]_{\deg=n}$. Let $\Phi' = \otimes_v \Phi'_v \in \mathcal{S}((S_n \times V'_n)(\mathbb{A}_0))$ be a pure tensor such that for every $v \mid \infty$, Φ'_v is the partial Gaussian test function. We define a weaker notion of smooth transfer. Fix an $F_v/F_{0,v}$ -hermitian space V_v .

Definition 13.1. For a fixed $\alpha \in \mathcal{A}_n(F_{0,v})$, we say that Φ'_v partially (relative to α) transfers to $\Phi_v \in \mathcal{S}(\text{U}(V_v) \times V_v)(F_{0,v})$ if we only require the equality [\(2.18\)](#) in [Definition 2.2](#) to hold for matching orbits

$(\gamma, u') \in (S_n(\alpha) \times V'_n)(F_{0,v})_{\text{rs}}$ and $(\delta, u) \in (\text{U}(V_v)(\alpha) \times V_v)(F_{0,v})_{\text{rs}}$,
and $\text{Orb}((\gamma, u'), \Phi'_v) = 0$ for any other $(\gamma, u') \in (S_n(\alpha) \times V'_n)(F_{0,v})_{\text{rs}}$.

For $\Phi'^\infty = \otimes_{v \nmid \infty} \Phi'_v \in \mathcal{S}((S_n \times V'_n)(\mathbb{A}_{0,f}))$, we say that it partially transfers to (or matches) $\Phi^\infty = \otimes_{v \nmid \infty} \Phi_v \in \mathcal{S}((\text{U}(V) \times V)(\mathbb{A}_{0,f}))$ if Φ'_v partially transfers to Φ_v for every $v \nmid \infty$.

Remark 13.2. At those places of F_0 split in F , we will further demand Φ_v and Φ'_v to match in an elementary way analogous to [\[47\]](#).

13.2. *Comparison.* In this subsection, we compare $\mathbb{J}(h, \Phi')$ with $\mathbb{J}(h, \Phi)$ in the “coherent” case, i.e., $\Phi = \otimes_v \Phi_v \in \mathcal{S}((\mathrm{U}(V) \times V)(\mathbb{A}_0))$ for an n -dimensional F/F_0 -hermitian space V . We further assume that V is totally positively definite and Φ_v is the Gaussian test function for every $v \mid \infty$; cf. (12.1) in Section 12.1.

PROPOSITION 13.3. *The function*

$$h \in \mathbf{H}(\mathbb{A}_0) \longmapsto \mathbb{J}(h, \Phi'), \text{ resp. } \mathbb{J}(h, \Phi),$$

lies in $\mathcal{A}_{\mathrm{hol}}(\mathbf{H}(\mathbb{A}_0), K, n)$, where K is a compact open subgroup of $\mathrm{SL}_2(\mathbb{A}_{0,f})$ that acts trivially on both Φ and Φ' .

Proof. The K -invariance follows immediately from the definition of $\mathbb{J}(h, \Phi')$ and $\mathbb{J}(h, \Phi)$. By Theorem 12.14 (resp. Lemma 11.1) the function $\mathbb{J}(\cdot, \Phi')$ (resp. $\mathbb{J}(\cdot, \Phi)$) is invariant under $\mathbf{H}(F_0)$. The weight n condition follows from the action under $\mathrm{SO}(2, \mathbb{R})$ by (12.4) for Φ and by Lemma 12.5 for Φ' .

Finally we need to show the holomorphy on the complex upper half plane $\mathcal{H}^{[F_0:\mathbb{Q}]}$ and at all cusps. Equivalently, for any $h_f \in \mathbf{H}(\mathbb{A}_{0,f})$, the function $\mathbb{J}_{h_f}^b(\tau, \Phi')$ (resp. $\mathbb{J}_{h_f}^b(\tau, \Phi)$) defined by (1.11) is holomorphic in $\tau \in \prod_{v \mid \infty} \mathcal{H}$, and holomorphic at the cusp $i\infty$.

By (12.34) and Lemmas 12.3 and 12.6, the ξ -th Fourier coefficient of $\mathbb{J}_{h_f}^b(h_\infty, \Phi')$ vanishes unless $\xi \in F_0$ and $\xi \geq 0$ (i.e., totally semi-positive). Hence the Fourier expansion takes the form

$$\mathbb{J}_{h_f}^b(\tau, \Phi') = \sum_{\xi \in F_0, \xi \geq 0} A_\xi q^\xi, \quad A_\xi \in \mathbb{C},$$

where $A_\xi = 0$ unless ξ lies in a (fractional) ideal of F_0 depending on Φ'_f and h_f . This shows that $\mathbb{J}(\cdot, \Phi') \in \mathcal{A}_{\mathrm{hol}}(\mathbf{H}(\mathbb{A}_0), K, n)$. The assertion for Φ is proved similarly. \square

Now let us fix a regular elliptic compact element $\gamma \in S_n(\alpha)(F_0)$ (cf. Section 12.6). Let S be a finite set of non-archimedean places of F_0 such that

- S contains all places with residue characteristic 2;
- for all $v \in S$, Φ'_v partially (relative to α) transfers to $\Phi_v \in \mathcal{S}(\mathrm{U}(V_v) \times V_v)(F_{0,v})$;⁹ and
- for every non-archimedean $v \notin S$, the hermitian space V_v is split, $\Phi_v = \mathbf{1}_{(\mathrm{U}(V) \times V)(O_{F_{0,v}})}$ (with respect to a self-dual lattice in V_v), and $\Phi'_v = \mathbf{1}_{(S_n \times V')(O_{F_{0,v}})}$.

⁹Transfers exist by the result of [47]. Here we only need the weaker result of the existence of partial transfers for fixed α , which can be deduced easily from the $n = 1$ case.

We consider the difference

$$\mathcal{E}(h) = \mathbb{J}(h, \Phi') - \mathbb{J}(h, \Phi), \quad h \in \mathbf{H}(\mathbb{A}_0).$$

THEOREM 13.4. *Assume that [Conjecture 2.3](#) part (a) holds for all $F_{0,v}$ with $v \notin S$ and for S_n . Then $\mathcal{E} = 0$. (Note that we are in the case $\dim V = n$.)*

Proof. Let B (for “bad”) be the (finite) set of non-archimedean places $v \notin S$ of F_0 where $R_{\alpha,v} = R_\alpha \otimes_{O_{F_0}} O_{F_{0,v}}$ is not a product of DVRs.

By [Proposition 13.3](#) and our choice of Φ and Φ' , the function $\mathcal{E}(h) \in \mathcal{A}_{\text{hol}}(\mathbf{H}(\mathbb{A}_0), K, n)$ where the compact open subgroup $K = \prod_{v \nmid \infty} K_v \subset \mathbf{H}(\mathbb{A}_{0,f})$ can be chosen such that K_v is of the form (13.3) for every $v \in B$. This is easy to see if the additive character ψ_v is of level zero. In general, it is known how the Weil representation depends on ψ_v and the desired K_v -invariance holds for any ψ_v at $v \in B$.

By the vanishing criterion [Lemma 13.6](#) below, it suffices to show that for $\xi \in F_0^\times$,

$$W_{\mathcal{E}, \xi}(h_\infty) = 0$$

whenever $(\xi, B) = 1$ (i.e., $v(\xi) = 0$ for all $v \in B$).

Now let $(\xi, B) = 1$. By (12.33), the ξ -th Fourier coefficient of $\mathbb{J}(h_\infty, \Phi')$ is

$$\sum_{(\gamma, u') \in [(S_n(\alpha) \times V_\xi')(F_0)]} \text{Orb}((\gamma, u'), \omega(h_\infty) \Phi').$$

Similarly, by (11.16), the ξ -th Fourier coefficient of $\mathbb{J}^\flat(\tau, \Phi)$ is

$$\sum_{(\delta, u) \in [(G(\alpha) \times V_\xi)(F_0)]} \text{Orb}((\delta, u), \omega(h_\infty) \Phi).$$

By our choices of partial Gaussian test functions, for every $v \mid \infty$,

$$(13.1) \quad \text{Orb}((\gamma, u'), \omega(h_v) \Phi'_v) = \text{Orb}((\delta, u), \omega(h_v) \Phi_v)$$

holds for every (γ, u') matching (δ, u) .

We now *claim* that the equality

$$(13.2) \quad \text{Orb}((\gamma, u'), \Phi'_v) = \text{Orb}((\delta, u), \Phi_v)$$

holds for every non-archimedean place v and every matching pair $(\gamma, u') \in [(S_n(\alpha) \times V_\xi')(F_0)]$ and $(\delta, u) \in [(G(\alpha) \times V_\xi)(F_0)]$ (when $(\xi, B) = 1$). From the claim it follows that $A_\xi = 0$ whenever $(\xi, B) = 1$.

To show the claim, first let $v \in B$. Since $(\xi, B) = 1$, ξ is a unit at v . Therefore by [Proposition 2.7\(ii\)](#), [Conjecture 2.3](#) part (a), which we have assumed to hold, implies (13.2).

If $v \notin S \cup B$, then $R_{\alpha,v}$ is a maximal order, and (13.2) follows from [Proposition 2.6](#) when v is inert. When v is split, the identity is trivial.

If $v \in S$, by our assumption on Φ'_v and Φ_v , they partially match (relative to the fixed α), and hence (13.2) holds. This proves the claim.

Now by (13.1) and (13.2) we conclude that $W_{\mathcal{E},\xi}(h_\infty) = 0$ whenever $(\xi, B) = 1$. This completes the proof. \square

COROLLARY 13.5. *Under the assumption of Theorem 13.4, we have for all $\xi \in F_0^\times$,*

$$\sum_{(\gamma, u') \in [(S_n(\alpha) \times V_\xi')(F_0)]} \text{Orb}((\gamma, u'), \Phi') = \sum_{(\delta, u) \in [(G(\alpha) \times V_\xi)(F_0)]} \text{Orb}((\delta, u), \Phi).$$

Proof. This follows from Theorem 13.4, comparing the ξ -th coefficients of $\mathbb{J}(h, \Phi')$ and $\mathbb{J}(h, \Phi)$. \square

13.3. A lemma on Fourier coefficients of modular forms. Let ϕ be a continuous function on $\mathbf{H}(\mathbb{A}_0)$, left invariant under $\mathbf{H}(F_0)$. Recall its Fourier expansion from (1.12) and (1.13). Let c_v be the level of ψ_v , i.e., the maximal integer such that ψ_v is trivial on $\varpi_v^{-c_v} O_{F_0, v}$, where ϖ_v is a uniformizer of $F_{0,v}$.

LEMMA 13.6. *Let B be a finite set of non-archimedean places of F_0 . Assume that ϕ is right invariant under a compact open $K = \prod_{v \nmid \infty} K_v \subset \mathbf{H}(\mathbb{A}_{0,f})$, where*

$$(13.3) \quad K_v = m(\varpi_v^{c_v})^{-1} \mathbf{H}(O_{F_0, v}) m(\varpi_v^{c_v}), \quad m(\varpi_v^{c_v}) = \begin{pmatrix} \varpi_v^{c_v} & \\ & 1 \end{pmatrix}$$

for all $v \in B$. Suppose that $W_{\phi,\xi}(h_\infty)$ vanishes identically (as a function in $h_\infty \in \mathbf{H}(F_{0,\infty})$) for all $\xi \in F_0^\times$ such that $(\xi, B) = 1$ (i.e., $v(\xi) = 0$ for all $v \in B$). Then ϕ is a constant function. In particular, if ϕ is of (parallel) weight n with $n \neq 0$, then $\phi = 0$.

Proof. We prove the assertion by induction on $\#B$. If B is empty (i.e., $W_{\phi,\xi}(h_\infty) = 0$ for all $\xi \in F_0^\times$), then $\phi(h_\infty) = W_{\phi,\xi=0}(h_\infty)$ is left invariant under $N(F_{0,\infty})$ and left invariant under $\mathbf{H}(F_0) \cap K$. Now note that for every v , $\mathbf{H}(F_{0,v})$ is generated by $N(F_{0,v})$ and any single element in $\mathbf{H}(F_{0,v}) \setminus B(F_{0,v})$. It follows that $h_\infty \in \mathbf{H}(F_{0,\infty}) \mapsto \phi(h_\infty)$ is constant, and hence $h \in \mathbf{H}(\mathbb{A}_0) \mapsto \phi(h)$ is constant since $\mathbf{H}(F_0) \mathbf{H}(F_{0,\infty})$ is dense in $\mathbf{H}(\mathbb{A}_0)$ by the strong approximation theorem for $\mathbf{H} = \text{SL}_{2, F_0}$.

Now assume that B contains at least one element, say $v_0 \in B$. Consider $b_{v_0} \in \varpi_{v_0}^{-c_{v_0}-1} O_{F_0, v_0}$ and the unipotent matrix

$$n(b_{v_0}) := \begin{pmatrix} 1 & b_{v_0} \\ & 1 \end{pmatrix} \in N(F_{0, v_0}).$$

Consider the function

$$\tilde{\phi}(h) := \phi(h n(b_{v_0})) - \phi(h), \quad h \in \mathbf{H}(\mathbb{A}_0).$$

Then we *claim* that $W_{\tilde{\phi}, \xi}(h_\infty) = 0$ for all $\xi \in F_0^\times$ such that $(\xi, B \setminus \{v_0\}) = 1$.

To show the claim, the case $v_0(b_{v_0}) \geq -c_{v_0}$ is obvious, and therefore it suffices to consider the case $v_0(b_{v_0}) = -c_{v_0} - 1$. From the K_{v_0} -invariance it follows that $W_{\phi, \xi}(h_\infty) = 0$ and $W_{\phi, \xi}(h_\infty n(b_{v_0})) = 0$ unless $v_0(\xi) \geq 0$. If $v_0(\xi) \geq 1$, then $v_0(\xi b_{v_0}) \geq -c_{v_0}$ and hence $\psi_{v_0}(\xi b_{v_0}) = 1$. It follows that, unless $v_0(\xi) = 0$, we have

$$\begin{aligned} W_{\tilde{\phi}, \xi}(h_\infty) &= W_{\phi, \xi}(h_\infty n(b_{v_0})) - W_{\phi, \xi}(h_\infty) \\ &= \psi_{v_0}(\xi b_{v_0}) W_{\phi, \xi}(h_\infty) - W_{\phi, \xi}(h_\infty) = 0. \end{aligned}$$

The claim now follows.

By induction, we conclude that $\tilde{\phi}$ is a constant function, i.e., $W_{\tilde{\phi}, \xi}(h) = 0$ for all $\xi \in F_0^\times$. It follows that $W_{\phi, \xi}(h)$ is right invariant under $N(\varpi_{v_0}^{-c_{v_0}-1} O_{F_0, v_0})$. It is well known that the groups $N(\varpi_{v_0}^{-c_{v_0}-1} O_{F_0, v_0})$ and $N_-(\varpi_{v_0}^{c_{v_0}} O_{F_0, v_0}) \subset K_{v_0}$ generate $\mathbf{H}(F_{0, v_0})$. (This is equivalent to the fact that $N(\varpi_{v_0}^{-1} O_{F_0, v_0})$ and $N_-(O_{F_0, v_0})$ generate $\mathbf{H}(F_{0, v_0})$; for a proof, see [27, Prop. 8.1.2].) Here N_- denotes the transpose of N . It follows that, for all $\xi \in F_0^\times$, $W_{\phi, \xi}$ is right invariant under $\mathbf{H}(F_{0, v_0})$, and therefore it must vanish. Finally the assertion follows from the case when B is empty. \square

13.4. *A refinement of Corollary 13.5.* We recall from (2.10) that $\mathcal{B} = \mathcal{B}_n$ is identified with the categorical quotients $(\mathbf{U}(V) \times V)_{/\!/\mathbf{U}(V)}$ and $(S_n \times V'_n)_{/\!/\mathbf{GL}_n}$.

LEMMA 13.7. *Fix $b_0 \in \mathcal{B}(F_0)$. Fix a non-archimedean place v_1 of F_0 , split in F . For every place $v \neq v_1$ of F_0 , we fix a compact subset $\Omega_v \subset \mathcal{B}(F_{0, v})$ containing b_0 , such that for all but finitely many non-archimedean places v , Ω_v is equal to $\mathcal{B}(O_{F_{0, v}})$. Then there exists a neighborhood $\Omega_{v_1} \subset \mathcal{B}(F_{0, v_1})$ of b_0 such that*

$$\mathcal{B}(F_0) \cap \prod_v \Omega_v = \{b_0\},$$

where the intersection is taken inside $\mathcal{B}(\mathbb{A}_0)$.

Proof. We may embed \mathcal{B} as a closed sub-variety of some affine space $Y = \mathbf{A}^m$ over F_0 (e.g., by (2.10)) such that for almost all v , $\mathcal{B}(O_{F_{0, v}}) = Y(O_{F_{0, v}}) \cap \mathcal{B}(F_{0, v})$. For every $v \neq v_1$, by the compactness of Ω_v we may choose a compact subset $\tilde{\Omega}_v \subset Y(F_{0, v})$ such that $\Omega_v = \tilde{\Omega}_v \cap \mathcal{B}(F_{0, v})$, and such that $\tilde{\Omega}_v = Y(O_{F_{0, v}})$ for almost all v . By a standard argument using the product formula (i.e., $\prod_v |x|_v = 1$ for $x \in F_0^\times$), there must be a small neighborhood $\tilde{\Omega}_{v_1} \subset Y(F_{0, v_1})$ such that

$$Y(F_0) \cap \prod_v \tilde{\Omega}_v = \{b_0\}.$$

Set $\Omega_{v_1} = \tilde{\Omega}_{v_1} \cap \mathcal{B}(F_{0, v_1})$ to complete the proof. \square

We are now ready to refine the result of Corollary 13.5.

PROPOSITION 13.8. *Under the assumption of Theorem 13.4, for every $(\delta, u) \in (U(V)(\alpha) \times V)(F_0)_{\text{rs}}$ matching $(\gamma, u') \in (S_n(\alpha) \times V'_n)(F_0)_{\text{rs}}$ such that $\xi = \mathfrak{q}(u) \neq 0$, we have*

$$\text{Orb}((\gamma, u'), \Phi') = \text{Orb}((\delta, u), \Phi).$$

Proof. Let $b_0 \in \mathcal{B}(F_0)$ be the (common) image of (δ, u) and (γ, u') , and let $\xi = \mathfrak{q}(u) = \mathfrak{q}(u')$. Fix a non-archimedean place v_1 of F_0 , split in F . Decompose

$$\text{Orb}((\delta, u), \Phi) = \text{Orb}((\delta, u), \Phi^{v_1}) \text{Orb}((\delta, u), \Phi_{v_1})$$

and

$$\text{Orb}((\gamma, u'), \Phi') = \text{Orb}((\gamma, u'), \Phi'^{v_1}) \text{Orb}((\gamma, u'), \Phi'_{v_1}),$$

where the local orbital integral $\text{Orb}((\delta, u), \Phi_{v_1}) = \text{Orb}((\gamma, u'), \Phi'_{v_1})$. We may assume that the local orbital integrals at v_1 are nonzero (otherwise both sides vanish). It remains to show

$$(13.4) \quad \text{Orb}((\delta, u), \Phi^{v_1}) = \text{Orb}((\gamma, u'), \Phi'^{v_1}).$$

For every non-archimedean $v \neq v_1$, we define a compact set $\Omega_v \subset \mathcal{B}(F_{0,v})$ to be the image of the support of Φ_v for $v \in S$, and $\Omega_v = \mathcal{B}(O_{F_{0,v}})$ for all $v \notin S$.

For $v \mid \infty$, we define Ω_v to be the image of $(U(V)(\alpha) \times V_\xi)(F_{0,v})$. Since the hermitian space $V \otimes_{F_0} F_{0,v}$ is positive definite, the set $(U(V)(\alpha) \times V_\xi)(F_{0,v})$ is compact, and hence so is Ω_v .

Now apply Lemma 13.7 to choose a small neighborhood $\Omega_{v_1} \subset \mathcal{B}(F_{0,v_1})$ of b_0 such that $\mathcal{B}(F_0) \cap \Omega = \{b_0\}$, where $\Omega = \prod_v \Omega_v$. Then we choose a *point-wise non-negative* function $\tilde{\Phi}_{v_1}$ with non-empty support whose image in $\mathcal{B}(F_{0,v_1})$ contains b_0 and is contained in Ω_{v_1} . Choose $\tilde{\Phi}'_{v_1}$ to match $\tilde{\Phi}_{v_1}$ in the elementary way (cf. Remark 13.2). Now apply Corollary 13.5 to this new pair of functions $\tilde{\Phi} = \Phi^{v_1} \otimes \tilde{\Phi}_{v_1}$ and $\tilde{\Phi}' = \Phi'^{v_1} \otimes \tilde{\Phi}'_{v_1}$:

$$\sum_{(\gamma, u') \in [(S_n(\alpha) \times V'_n)(F_0)]} \text{Orb}((\gamma, u'), \tilde{\Phi}') = \sum_{(\delta, u) \in [(G(\alpha) \times V_\xi)(F_0)]} \text{Orb}((\delta, u), \tilde{\Phi}).$$

Now the non-zero terms in each side only involve regular semisimple orbits with invariants in Ω : this is clear for the unitary side by our choice of Ω , and it is true for the left-hand side because $\tilde{\Phi}'_{v_1}$ partially (relative to α) transfers to $\tilde{\Phi}_v$ for all $v \in S$ and $v \mid \infty$. It follows that each side has one term left, namely, the one with invariant $b_0 \in \mathcal{B}(F_0)$:

$$\text{Orb}((\gamma, u'), \tilde{\Phi}') = \text{Orb}((\delta, u), \tilde{\Phi}).$$

By the point-wise positivity of $\tilde{\Phi}_{v_1}$, the local orbital integral at the place v_1 does not vanish. We hence deduce the desired equality (13.4). This completes the proof. \square

13.5. *Proof of the FL conjecture.* Now we return to the set up of [Conjecture 2.3](#) in [Section 2.4](#).

THEOREM 13.9. *Conjecture 2.3 holds for F_0 with $q \geq n$. (Recall that q denotes the cardinality of the residue field of O_{F_0} .)*

Proof. By [Proposition 2.7](#), it suffices to prove [Conjecture 2.3](#) part (b). We will do so by induction on $\dim V_0$.

The case $\dim V_0 = 1$ is trivial. Assume now that [Conjecture 2.3](#) part (b) holds when $\dim V_0 = n - 1$. Then by [Proposition 2.7](#) part (i), [Conjecture 2.3](#) part (a) holds for S_n over F_0 with $q \geq n$.

We now want to apply [Proposition 13.8](#). We now change the notation: we denote by F_0 a totally real field with a place v_0 such that F_{0,v_0} is the local field in [Conjecture 2.3](#). We then choose the following local data:

- an unramified (local) quadratic extension $F_{w_0}/F_{0,v_0}$;
- the split $F_{w_0}/F_{0,v_0}$ -hermitian space V_{v_0} of dimension n ;
- an element $(g_{v_0}, u_{v_0}) \in (U(V_{v_0}) \times V_{v_0})(F_{0,v_0})_{\text{srs}}$ — we further assume that the characteristic polynomial of g_{v_0} has integral coefficients (in $O_{F_{w_0}}$), $\det(1 - g_{v_0})$ is a unit, and $\langle u_{v_0}, u_{v_0} \rangle \neq 0$;
- an element $(\gamma_{v_0}, u'_{v_0}) \in (S_n \times V'_n)(F_{0,v_0})_{\text{srs}}$ matching (g_{v_0}, u_{v_0}) .

To globalize the data, we first use the Cayley transform; cf. [\(4.1\)](#). Let $x_{v_0} = \mathfrak{c}^{-1}(g_{v_0}) = \frac{1+g_{v_0}}{1-g_{v_0}}$, an element in the Lie algebra $\mathfrak{u}(V_{v_0}) \subset \text{End}_{F_{w_0}}(V_{v_0})$.

We now choose a totally negative element $\epsilon \in F_0^\times$ such that v_0 is inert in $F = F_0[\sqrt{\epsilon}]$. Denote by w_0 the place of F above v_0 . Consider $x_{v_0}^\natural = \frac{x_{v_0}}{\sqrt{\epsilon}}$. Then the characteristic polynomial of $x_{v_0}^\natural$ has coefficients in the base field F_{v_0} .

Next we choose a totally real field F'_0 with $[F'_0 : F_0] = n$ and an element $x^\natural \in F'_0$ such that, when setting $F' = F'_0[\sqrt{\epsilon}]$ and

$$x = \sqrt{\epsilon} x^\natural, \quad g = \mathfrak{c}(x) = -\frac{1-x}{1+x},$$

we have $O_{F_{w_0}}[g] = O_{F_{w_0}}[g_{v_0}]$ as subrings of $F' \otimes_F F_{w_0}$. To achieve this, it suffices to approximate the characteristic polynomial of $x_{v_0}^\natural$ by a polynomial with coefficient in F_0 , and we may prescribe its local behavior at finitely many places by weak approximation. (Here the regular semi-simplicity allows us to determine the isomorphism class of the local field $F[g_{v_0}]$ by the characteristic polynomial of g_{v_0} .)

With such a choice, we have $g \in F'^1$. For the CM extension F'/F'_0 , there exists a one-dimensional F'/F'_0 -hermitian space W such that W is totally positive definite, and $V := \text{Res}_{F'_0/F_0} W$, as an n -dimensional F/F_0 -hermitian space, is locally at v_0 isometric to V_{v_0} . Such a hermitian space W exists because we are only imposing local conditions at finitely many places. Then we have

an embedding $\text{Res}_{F'_0/F_0} \text{U}(W) \rightarrow \text{U}(V)$, and we will view g as an element of $\text{U}(V)(F_0)$. Let $\alpha \in \mathcal{A}_n(F_0)$ denote its characteristic polynomial.

Now we choose $u \in V$ (and possibly replacing g by an element v_0 -adically closer to g_{v_0}) such that the pair (g, u) is v_0 -adically close to (g_{v_0}, u_{v_0}) and such that

(13.5)

$$\text{Orb}((g, u), \mathbf{1}_{(\text{U}(V_{v_0}) \times V_{v_0})(O_{F_0, v_0})}) = \text{Orb}((g_{v_0}, u_{v_0}), \mathbf{1}_{(\text{U}(V_{v_0}) \times V_{v_0})(O_{F_0, v_0})}).$$

This is possible due to the local constancy of orbital integrals near a regular semisimple element. Let $(\gamma, u') \in (S_n \times V'_n)(F_0)$ be a regular semisimple element matching (g, u) . Again by local constancy of orbital integrals we may assume that, possibly replacing (g, u) by an element in $(\text{U}(V) \times V)(F_0)_{\text{srs}}$ that is v_0 -adically closer to (g_{v_0}, u_{v_0}) ,

$$(13.6) \quad \text{Orb}((\gamma, u'), \mathbf{1}_{(S_n \times V'_n)(O_{F_0, v_0})}) = \text{Orb}((\gamma_{v_0}, u'_{v_0}), \mathbf{1}_{(S_n \times V'_n)(O_{F_0, v_0})}).$$

Next, we let S be a finite set of non-archimedean places of F_0 , such that

- $v_0 \notin S$;
- S contains all places with residue cardinality less than n ;
- for every non-archimedean $v \notin S \cup \{v_0\}$, the ring R_α is locally maximal at v , and V_v is a split hermitian space.

Choose functions $\Phi = \otimes_v \Phi_v$ and $\Phi' = \otimes_v \Phi'_v$ satisfying the following conditions:

- For every archimedean v , Φ_v and Φ'_v are the (partial) Gaussian test functions (relative to a small neighborhood of γ in $S_n(F_{0,v})$).
- For every non-archimedean $v \in S$, Φ'_v partially (relative to α) transfers to $\Phi_v \in \mathcal{S}((\text{U}(V_v) \times V_v)(F_{0,v}))$ and the local orbital integrals do not vanish at (g, u) and (γ, u') ; cf. [Definition 13.1](#).
- For every non-archimedean $v \notin S$ (in particular at v_0), noting that the hermitian space V_v is split, choose $\Phi_v = \mathbf{1}_{(\text{U}(V) \times V)(O_{F_0, v})}$ (with respect to a self-dual lattice in V_v) and $\Phi'_v = \mathbf{1}_{(S_n \times V')(O_{F_0, v})}$. By enlarging S suitably (while keeping $v_0 \notin S$), we may further assume that, for every $v \notin S$, the image of (g, u) in $\mathcal{B}(F_0)$ lies in the image of the support of $(\text{U}(V) \times V)(O_{F_0, v})$.

By the last condition, for every non-archimedean $v \notin S$, the local orbital integral of Φ_v does not vanish at (g, u) (since the function Φ_v is point-wisely positive on its support). It follows from the special case [Proposition 2.6](#) that the same non-vanishing holds for Φ'_v for every place $v \notin S \cup \{v_0\}$.

Now, by the induction hypothesis, we are ready to apply [Proposition 13.8](#) to conclude

$$\text{Orb}((\gamma, u'), \Phi') = \text{Orb}((g, u), \Phi).$$

By our choices, $\text{Orb}((\gamma, u'), \Phi'_v) = \text{Orb}((g, u), \Phi_v)$ for all $v \neq v_0$, and they do not vanish. It follows that

$$\text{Orb}((\gamma, u'), \Phi'_{v_0}) = \text{Orb}((g, u), \Phi_{v_0}).$$

By (13.5) and (13.6), we have

$$\text{Orb}((\gamma_{v_0}, u'_{v_0}), \mathbf{1}_{(S_n \times V'_n)(O_{F_{v_0}, v_0})}) = \text{Orb}((g_{v_0}, u_{v_0}), \mathbf{1}_{(U(V_{v_0}) \times V_{v_0})(O_{F_{v_0}, v_0})}).$$

We have assumed that $g_{v_0} \in U(V_{v_0})$ is regular semisimple with $\det(1 - g_{v_0}) \in O_{F_{v_0}}^\times$ and $\langle u_{v_0}, u_{v_0} \rangle \neq 0$. We now remove these assumptions. The condition $\det(1 - g_{v_0}) \in O_{F_{v_0}}^\times$ is harmless since we may multiply g_{v_0} by a suitable element in $F_{v_0}^1$ (cf. the proof of [Proposition 4.12](#)). The set of elements $(g_{v_0}, u_{v_0}) \in (U(V_{v_0}) \times V_{v_0})(F_{0, v_0})_{\text{rs}}$ with $\langle u_{v_0}, u_{v_0} \rangle \neq 0$ is dense in $(U(V_{v_0}) \times V_{v_0})(F_{0, v_0})_{\text{rs}}$. By local constancy of orbital integrals at regular semisimple elements, [Conjecture 2.3](#) part (b) holds when $\dim V_0 = n$ (over F_{0, v_0} with $q \geq n$). This completes the induction. \square

14. The comparison for arithmetic intersections

As a preparation for the proof of the AFL conjecture, in this section we compare $\partial\mathbb{J}(h, \Phi')$ with the arithmetic intersection number $\text{Int}(\tau, \Phi)$ (cf. (9.5) in [Section 9](#)).

Let V be the n -dimensional F/F_0 -hermitian space that we use to define the Shimura variety $\text{Sh}_{K_{\widetilde{G}}}(\widetilde{G}, \{h_{\widetilde{G}}\})$ in [Section 6.1](#).

As in [Section 13.1](#), we fix an irreducible $\alpha \in \mathcal{A}_n(F_0) \subset F[T]_{\deg=n}$ and fix $\gamma \in S_n(\alpha)(F_0)$ (cf. [Section 12.6](#)). Let $\Phi = \otimes_{v < \infty} \Phi_v \in \mathcal{S}((U(V) \times V)(\mathbb{A}_{0,f}))$ be a pure tensor. Let $\Phi' = \otimes_v \Phi'_v \in \mathcal{S}((S_n \times V'_n)(\mathbb{A}_0))$ be a pure tensor such that

- for every $v \mid \infty$, Φ'_v is the partial Gaussian test function; and
- for every non-archimedean v , Φ'_v partially (relative to α) transfers to Φ_v .

Remark 14.1. We are now in the “incoherent” case in the following sense. Due to the signature of such V at the archimedean places, there does not exist any *global* F/F_0 -hermitian space \widetilde{V} such that Φ' transfer to a function in $\mathcal{S}((U(\widetilde{V}) \times \widetilde{V})(\mathbb{A}_0))$; cf. [\[46, §3.2\]](#).

We now study $\partial\mathbb{J}(h, \Phi')$, and we recall $\partial\mathbb{J}_v(h, \Phi')$ from (12.36).

LEMMA 14.2. *Let v be a place of F_0 split in F . Then $\partial\mathbb{J}_v(h, \Phi') = 0$.*

Proof. This follows from the same argument in [\[46, Prop. 3.6\]](#) (also cf. [\[40, §7.2\]](#)). \square

If v is non-split (including the archimedean places), then let $V(v)$ be the “nearby” F/F_0 -hermitian space at v ; cf. [Theorem 9.4](#) (resp. [Theorem 10.1](#)) for non-archimedean (resp. archimedean) places. Then, for a non-archimedean v ,

the regular semisimple terms in $\partial\mathbb{J}_v(h, \Phi')$ (cf. (12.36)) is a sum over orbits $(\gamma, u') \in [(S_n(\alpha) \times V')(F_0)]_{\text{rs}}$ matching $(\delta, u) \in [(\text{U}(V(v)) \times V(v))(F_0)]_{\text{rs}}$; we will show that the same holds for archimedean places v ; cf. Lemma 14.4. Moreover, we have a Fourier expansion (cf. (1.12))

$$(14.1) \quad \partial\mathbb{J}_v(h, \Phi') = \sum_{\xi \in F_0} \partial\mathbb{J}_v(\xi, h, \Phi'),$$

where $\partial\mathbb{J}_v(\xi, h, \Phi')$ is the sub-sum,

$$(14.2) \quad \begin{aligned} \partial\mathbb{J}_v(\xi, h, \Phi') &= \sum_{\substack{(\gamma, u') \in [(S_n(\alpha) \times V'_\xi)(F_0)] \\ u' \neq 0}} \partial\text{Orb}((\gamma, u'), \omega(h_v)\Phi'_v) \cdot \text{Orb}((\gamma, u'), \omega(h^v)\Phi'^v). \end{aligned}$$

14.1. *The archimedean places.* Let $v \mid \infty$. Recall that $V' = V_0 \times V_0^*$ for $V_0 = F_0^n$ carries the tautological quadratic form (11.17)

$$\mathfrak{q} : V_0 \times V_0^* \longrightarrow F_0,$$

and an induced quadratic form (12.25)

$$\mathfrak{q}' : V_0 \times V_0^* \longrightarrow F'_0,$$

such that for all $u' \in V'$, $\mathfrak{q}'(u') = \text{tr}_{F'_0/F_0} \mathfrak{q}'(u')$. Set

$$F'_{0,v} := F'_0 \otimes_{F_0, v} \mathbb{R} \simeq \prod_{v' \in \text{Hom}(F'_0, \mathbb{R}), v' \mid v} \mathbb{R}.$$

LEMMA 14.3. *Let $\xi' \in F'_{0,v}$ be an invertible element, and let $u' \in V'(F_{0,v})$ with $\mathfrak{q}'(u') = \xi'$.*

(a) *We have*

$$\text{Orb}((\gamma, u'), \Phi'_v) = \begin{cases} e^{-\pi \text{tr}_{F'_{0,v}/F_{0,v}}(\xi')}, & \text{when } \xi' \in F'_{0,v} \text{ is totally positive,} \\ 0, & \text{otherwise.} \end{cases}$$

(b) *Now assume that ξ' is not totally positive. Then $\partial\text{Orb}((\gamma, u'), \Phi'_v) = 0$, unless ξ' is negative at exactly one archimedean place v' of F'_0 and this place v' is above v , in which case*

$$\partial\text{Orb}((\gamma, u'), \Phi'_v) = \frac{1}{2} e^{-\pi \text{tr}_{F'_{0,v}/F_{0,v}}(\xi')} \text{Ei}(-2\pi|\xi'|_{v'}).$$

Proof. This follows from Lemmas 12.3 and 12.6. \square

LEMMA 14.4. *Let $\xi \in F_0^\times$. Then*

$$\text{Int}_v^\mathbf{K}(\xi, \Phi) = -2\partial\mathbb{J}_v(\xi, \Phi').$$

Proof. It follows from the previous [Lemma 14.3](#) that

$$\partial \mathbb{J}_v(\xi, \Phi') = \frac{1}{2} \sum \text{Ei}(-2\pi|\xi'|_{v'}) \cdot \text{Orb}((\gamma, u'), \Phi'),$$

where the sum runs over $(\gamma, u') \in [(S_n(\alpha) \times V'_\xi)(F_0)]$ such that the refined invariant $q'(u') = \xi' \in F'_0$ is negative at exactly one archimedean place v' of F'_0 and this place is above v .

By [Corollary 10.3](#),

$$\text{Int}_v^K(\xi, \Phi) = - \sum \text{Ei}(-2\pi|\xi'|_{v'}) \cdot \text{Orb}((\delta, u), \Phi),$$

where the sum runs over $(\delta, u) \in [(U(V(v))(\alpha) \times V(v)_\xi)(F_0)]$ such that the refined invariant ξ' is negative at exactly one archimedean place v' of F'_0 and this place is above v .

Therefore, the orbits (δ, u) in the sum in $\text{Int}_v^K(\xi, \Phi)$ are bijective to the orbits (γ, u') in the sum in $\partial \mathbb{J}_v(\xi, \Phi')$. Now the assertion follows from the fact that Φ^∞ and Φ'^∞ are partial transfers of each other. \square

14.2. “Holomorphic projection”. For the rest of this section, we assume that $F_0 = \mathbb{Q}$. Recall that the difference $\text{Int}^{K-B}(h, \Phi)$ between the two Green functions is given by [\(10.9\)](#) and [\(10.10\)](#). (Note that this makes sense for any Schwartz function Φ^∞ .) The following result plays the role of “holomorphic projection” of the modular generating function on the analytic side.

PROPOSITION 14.5. *Let $F_0 = \mathbb{Q}$. The sum*

$$\partial \mathbb{J}_{\text{hol}}(h) := 2\partial \mathbb{J}(h, \Phi') + \text{Int}^{K-B}(h, \Phi), \quad h \in \mathbf{H}(\mathbb{A}_0),$$

lies in $\mathcal{A}_{\text{hol}}(\mathbf{H}(\mathbb{A}_0), K, n)$, where K is the compact open subgroup of $\text{SL}_2(\mathbb{A}_{0,f})$ that acts trivially on both Φ and Φ' .

Proof. First of all, note that the function $h \in \mathbf{H}(\mathbb{A}_0) \mapsto 2\partial \mathbb{J}(h, \Phi')$ belongs to $\mathcal{A}_{\text{exp}}(\mathbf{H}(\mathbb{A}_0), K, n)$. One way to see this is to use the Fourier expansion directly. Another way is to identify it with a linear combination to $\text{SL}_2(\mathbb{A}_0)$ of the restriction of (the first derivative at $s = 0$ of) a degenerate Siegel–Eisenstein series of parallel weight one on $\text{SL}_2(\mathbb{A}_{F'_0})$; cf. [Remark 12.10](#).

By [Corollary 10.4](#), the second summand $\text{Int}^{K-B}(\cdot, \Phi)$ also belongs to $\mathcal{A}_{\text{exp}}(\mathbf{H}(\mathbb{A}_0), K, n)$. Therefore to complete the proof, it suffices to show the holomorphy of the sum $\partial \mathbb{J}_{\text{hol}}(h)$ on the complex upper half plane \mathcal{H} and at all cusps. Equivalently, for any $h_f \in \mathbf{H}(\mathbb{A}_{0,f})$, the function $\partial \mathbb{J}_{\text{hol}, h_f}^\flat$ (associated to $\partial \mathbb{J}_{\text{hol}}$ via [\(1.11\)](#)) is holomorphic, and holomorphic at the cusp $i\infty$. Since we can vary Φ^∞ and Φ'^∞ , and by [Theorem A.1](#) the Weil representation commutes with (partially relative to α) smooth transfer, it suffices to consider the case $h_f = 1$ (but allow all matching Φ^∞ and Φ'^∞).

We *claim* that the Fourier expansion of $\partial\mathbb{J}_{\text{hol}}^{\flat}$ takes the form

$$(14.3) \quad \partial\mathbb{J}_{\text{hol}}^{\flat}(\tau) = \sum_{\xi \in F_0, \xi \geq 0} A_{\xi} q^{\xi}, \quad A_{\xi} \in \mathbb{C},$$

where $A_{\xi} = 0$ unless ξ lies in a (fractional) ideal of F_0 (depending on Φ', Φ). In other words, the non-holomorphic terms all cancel out. The desired holomorphy follows from the claim.

To show the claim, we use the decomposition (12.38) as a sum over places v of F_0 .

First, by Lemma 14.2, $\partial\mathbb{J}_v(h, \Phi') = 0$ if v is a split place.

Next let v be a non-archimedean non-split place. By (14.1) and (14.2), and the fact that Φ'_{∞} is a (partial) Gaussian test function, we have $\partial\mathbb{J}_v(\xi, h, \Phi') = 0$ unless $\xi \geq 0$. We obtain

$$(14.4) \quad \partial\mathbb{J}_v^{\flat}(\tau, \Phi') = \sum_{\xi \in F_0, \xi \geq 0} \partial\mathbb{J}_v(\xi, \Phi') q^{\xi},$$

where, for $\xi \geq 0$,

$$(14.5) \quad \partial\mathbb{J}_v(\xi, \Phi') = \sum_{\substack{(\gamma, u') \in [(S_n(\alpha) \times V_{\xi}^{\flat})(F_0)] \\ u' \neq 0}} \partial\text{Orb}((\gamma, u'), \Phi'_v) \cdot \text{Orb}((\gamma, u'), \Phi'^{v, \infty}).$$

It follows that $2\partial\mathbb{J}_v^{\flat}(\tau, \Phi')$ has the desired form of Fourier expansion as in (14.3).

Finally let $v \mid \infty$. We observe that by Lemma 14.4, modulo the constant terms, the sum $2\partial\mathbb{J}_v(h, \Phi') + \text{Int}_v^{\mathbf{K}-\mathbf{B}}(h, \Phi)$ is equal to $-\text{Int}_v^{\mathbf{B}}(h, \Phi)$, which has the desired form of Fourier expansion. It remains to consider the constant terms. Note that Lemma 14.3 also applies to all $u' \in V'$ with refined invariant $\mathfrak{q}'(u') = \xi' \in F_0'^{\times}$ (possibly $\xi = \text{tr}_{F_0'/F_0} \xi' = 0$). Similarly, Theorem 10.1 also applies to all $u \in V(v)$ with refined invariant $\mathfrak{q}'(u) = \xi' \in F_0'^{\times}$ (i.e., $u \neq 0$). Therefore, by the proof of Lemma 14.4, the contribution from null-norm ($\xi = 0$) non-zero vectors $u \in V(v)$ cancels that from $u' \in V'$ with $\mathfrak{q}'(u') = \xi' \neq 0 \in F_0'$. It follows that the constant term of $2\partial\mathbb{J}_v(h, \Phi') + \text{Int}_v^{\mathbf{K}-\mathbf{B}}(h, \Phi)$ is the sum of the nilpotent term $2\partial\text{Orb}(0_{\pm}, \omega(h)\Phi')$ (from $2\partial\mathbb{J}(h, \Phi')$; cf. (12.37)) and the only term that has not been cancelled in (10.9), which by (8.12) is

$$- \sum_{\delta \in [\text{U}(V(v))(\alpha)(F_0)]} \text{Orb}((\delta, 0), \Phi) \log |a|_v;$$

cf. (11.6) and (11.7). By Lemmas 12.11 and 12.6, the nilpotent term is

$$2\partial\text{Orb}(0_{\pm}, \Phi') = 2\text{Orb}((\gamma, 0_{\pm}), \Phi') \log |a|_v + C$$

for some constant C (depending on γ, Φ'). Since Φ matches Φ' , we claim

$$\text{Orb}((\gamma, 0_+), \Phi') = -\text{Orb}((\gamma, 0_-), \Phi') = \frac{1}{2} \sum_{\delta \in [\text{U}(V(v))(\alpha)(F_0)]} \text{Orb}((\delta, 0), \Phi).$$

In fact, this follows from the argument in [19, (10.4)] (for the quadratic extension F'/F'_0). In loc. cit., Jacquet proves the analogous identity in the “coherent” case. (Here “coherent” is in the sense of Kudla for one-dimensional hermitian spaces.) Since the proof verbatim applies to the current setting, we omit the detail. Therefore the two terms with $\log |a|_v$ cancel, and the sum is a constant independent of α . This shows that $2\partial\mathbb{J}_v(h, \Phi') + \text{Int}_v^{\mathbf{K}-\mathbf{B}}(h, \Phi)$ also has the desired form of Fourier expansion when $v \mid \infty$.

The proof is now complete. \square

14.3. *The comparison.* Now let $\mathcal{M} = \mathcal{M}_{K_{\tilde{G}}}(\tilde{G})$ be the moduli stack introduced in [Definition 6.1](#). Let S be a finite set of non-archimedean places of F_0 such that

- S contains all places $v \mid \mathfrak{d}$ and all places with residue cardinality $< \dim V$; and
- for every non-archimedean $v \notin S$, the hermitian space V_v is split,

$$\Phi_v = \mathbf{1}_{(\text{U}(V) \times V)(O_{F_0, v})}$$

(with respect to a self-dual lattice in V_v), and

$$\Phi'_v = \mathbf{1}_{(S_n \times V'_n)(O_{F_0, v})}.$$

Now we have the FL for all places $v \notin S$ by [Theorem 13.9](#), hence Φ_v and Φ'_v match for every place $v \nmid S$. Then in [Proposition 14.5](#), we can assume that the compact open subgroup $K \subset \text{H}(\mathbb{A}_{0,f})$ is the principal congruence subgroup $K(N)$ of level N , where the prime factors of N are all contained in S .

We have been assuming that the function $\Phi \in \mathcal{S}((\text{U}(V) \times V)(\mathbb{A}_{0,f}))$ is valued in \mathbb{Q} . By [Proposition 14.5](#), $\partial\mathbb{J}_{\text{hol}}^b(\cdot, \Phi')$ lies in $\mathcal{A}_{\text{hol}}(\Gamma(N), n)_{\mathbb{Q}} \otimes_{\mathbb{Q}} \mathbb{R}$ (the Green function takes values in \mathbb{R}). By passing to the quotient $\mathbb{R} \rightarrow \mathbb{R}_S$ (cf. (9.2)) and then extending coefficients $\mathbb{R}_S \rightarrow \mathbb{R}_{S, \overline{\mathbb{Q}}}$, we obtain an element, still denoted by $\partial\mathbb{J}_{\text{hol}}^b(\cdot, \Phi')$, in $\mathcal{A}_{\text{hol}}(\Gamma(N), n)_{\overline{\mathbb{Q}}} \otimes_{\overline{\mathbb{Q}}} \mathbb{R}_{S, \overline{\mathbb{Q}}}$.

By [Theorem 8.6](#) (cf. (9.7)), $\text{Int}(\cdot, \Phi)$ (defined by (9.5)) also belongs to $\mathcal{A}_{\text{hol}}(\Gamma(N), n)_{\overline{\mathbb{Q}}} \otimes_{\overline{\mathbb{Q}}} \mathbb{R}_{S, \overline{\mathbb{Q}}}$, hence so does the sum

$$\mathcal{E}^b(\tau) = 2\partial\mathbb{J}_{\text{hol}}^b(\tau, \Phi') + \text{Int}(\tau, \Phi), \quad \tau \in \mathcal{H}.$$

Write the Fourier expansion (at the cusp $i\infty$) as

$$\mathcal{E}^b(\tau) = \sum_{\xi \in F_0, \xi \geq 0} A_{\xi} q^{\xi}, \quad A_{\xi} \in \mathbb{R}_{S, \overline{\mathbb{Q}}}.$$

THEOREM 14.6. *Assume that Conjecture 3.8 part (a) holds for all p -adic field \mathbb{Q}_p with $p \notin S$ and for S_n . Then*

- (a) $\mathcal{E}^b = 0$;
- (b) *for every non-archimedean place $v \notin S$, and $\xi \in F_0^\times$, we have*

$$-2\partial\mathbb{J}_v(\xi, \Phi') = \text{Int}_v(\xi, \Phi)$$

as an equality in $\mathbb{Q} \log p_v$ where p_v denotes the residue characteristic of $F_{0,v}$; here $\partial\mathbb{J}_v(\xi, \Phi')$ (resp. $\text{Int}_v(\xi, \Phi)$) is defined by (14.5) (resp. (9.13)).

Proof. The proof of part (a) is analogous to that of Theorem 13.4. Recall from (9.15) and (9.16) that we have a decomposition of the generating function $\text{Int}(\tau, \Phi)$ (excluding the constant term $\text{Int}(0, \Phi)$ defined by (9.9)) as a sum of $\text{Int}_v(\tau, \Phi)$ over the places v of F_0 . We have the following equalities, as formal power series in $\mathbb{R}_{S, \overline{\mathbb{Q}}}[q^{1/N}]$ modulo constant terms:

$$\begin{aligned} \mathcal{E}^b(\tau) - \text{Int}(0, \Phi) &= (2\partial\mathbb{J}^b(\tau, \Phi') + \text{Int}^{\mathbf{K}-\mathbf{B}}(\tau, \Phi)) + (\text{Int}(\tau, \Phi) - \text{Int}(0, \Phi)) \\ &= \sum_{v|\infty} (2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v^{\mathbf{K}-\mathbf{B}}(\tau, \Phi) + \text{Int}_v(\tau, \Phi)) \\ &\quad + \sum_{v<\infty} (2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v(\tau, \Phi)) \\ &= \sum_{v|\infty} (2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v^{\mathbf{K}}(\tau, \Phi)) \\ &\quad + \sum_{v<\infty} (2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v(\tau, \Phi)) \\ &= \sum_{v|\infty, v \notin S} (2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v(\tau, \Phi)), \end{aligned}$$

where the last equality (modulo the constant term) follows from Lemma 14.4. Here the sums $2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v^{\mathbf{K}-\mathbf{B}}(\tau, \Phi)$ (resp. $2\partial\mathbb{J}_v^b(\tau, \Phi') + \text{Int}_v^{\mathbf{K}}(\tau, \Phi)$) both belong to $\mathbb{R}_{S, \overline{\mathbb{Q}}}[q^{1/N}]$, even though each summand does not due to the presence of “non-holomorphic” terms.

Recall from (9.16) that we have the expansion of $\text{Int}_v(\tau, \Phi)$ in terms of $\text{Int}_v(\xi, \Phi)$ defined by (9.13). Also recall from (14.4) that we have an expansion of $\partial\mathbb{J}_v^b(\tau, \Phi')$ in terms of $\partial\mathbb{J}_v(\xi, \Phi')$ defined by (14.5).

Let B be the (finite) set of non-archimedean inert places $v \notin S$ of F_0 where R_v is not a maximal order in $F'_v = F' \otimes_{F_0} F_{0,v}$. By the vanishing criterion Lemma 13.6, to show $\mathcal{E}^b = 0$, it remains to show the vanishing of the ξ -th Fourier coefficients when $(\xi, B) = 1$.

If $v \notin S$ is split in F , the intersection number $\text{Int}_v(\tau, \Phi) = 0$ vanishes by Corollary 9.3, and $2\partial\mathbb{J}_v^b(\tau, \Phi') = 0$ by Lemma 14.2.

If $v \notin S$ is inert, then by [Theorem 9.4](#) and [\(9.16\)](#) we obtain the q -expansion of $\text{Int}_v(\tau, \Phi)$. Similarly, [\(14.4\)](#) and [\(14.5\)](#) give the q -expansion of $2\partial\mathbb{J}_v^\flat(\tau, \Phi')$. There are two cases:

- (i) If $v \notin S \cup B$, then R_v is an maximal order and we apply [Proposition 3.9](#) at v to conclude that the v -th summand $2\partial\mathbb{J}_v^\flat(\tau, \Phi') + \text{Int}_v(\tau, \Phi)$ is zero. (Note that now $\Phi^{(v)}$ and $\Phi'^{(v)}$ match.)
- (ii) If $v \in B$, then the v -th term $2\partial\mathbb{J}_v^\flat(\tau, \Phi') + \text{Int}_v(\tau, \Phi)$ is a formal power series in $q^{1/N}$ with coefficients in $\mathbb{Q} \log q_v$ (or its image in $\mathbb{R}_{S, \overline{\mathbb{Q}}}$). By [Proposition 4.12](#), our assumption on [Conjecture 3.8](#) part (a) (for all p -adic field \mathbb{Q}_p with $p \notin S$ and for S_n) implies that for (γ, u') matching (δ, u) ,

$$-\partial\text{Orb}((\gamma, u'), \mathbf{1}_{(S_n \times V'_n)(O_{F_{0,v}})}) = \text{Int}_v(\delta, u) \cdot \log q_v,$$

whenever $\mathbf{q}(u) = \mathbf{q}(u') = \xi$ is a unit at v . In other words, the ξ -th Fourier coefficient of $2\partial\mathbb{J}_v^\flat(\tau, \Phi') + \text{Int}_v(\tau, \Phi)$ vanishes if ξ is a unit v , which holds if $(\xi, B) = 1$.

Therefore, whenever $(\xi, B) = 1$, the ξ -th Fourier coefficient of $\mathcal{E}^\flat - \text{Int}(0, \Phi)$ (equivalently \mathcal{E}^\flat) vanishes. By [Lemma 13.6](#) this completes the proof of part (a).

Now we turn to part (b). By $\mathcal{E}^\flat = 0$, taking the ξ -th Fourier coefficient (for $\xi > 0$) yields

$$\sum_{v \nmid \infty, v \notin S} (2\partial\mathbb{J}_v(\xi, \Phi') + \text{Int}_v(\xi, \Phi)) = 0$$

as an equality in $\mathbb{R}_{S, \overline{\mathbb{Q}}}$. Note that there are finitely many nonzero terms in the sum; in fact, we have proved the v -term vanishes unless $v \in B$. Since both $\partial\mathbb{J}_v(\xi, \Phi')$ and $\text{Int}_v(\xi, \Phi)$ lie in $\mathbb{Q} \log p_v$, and $\{\log p_v \mid v \in B\}$ are $\overline{\mathbb{Q}}$ -linearly independent inside $\mathbb{R}_{S, \overline{\mathbb{Q}}}$, the v -th term for each v must vanish. This completes the proof of part (b). \square

Remark 14.7. A byproduct of [Theorem 14.6](#) is that the constant term of \mathcal{E}^\flat vanishes. This amounts to an equality relating a certain part of the nilpotent term [\(12.37\)](#) to the arithmetic degree of the restriction of the metrized line bundle $\widehat{\omega}$ to the derived CM cycle. This may be of some independent interest.

COROLLARY 14.8. *Let $v \notin S$ be inert, and assume that [Conjecture 3.8](#) part (a) holds for all p -adic field \mathbb{Q}_p with $p \notin S$ and for S_n . Let $(\delta, u) \in (\text{U}(V(v))(\alpha) \times V(v))(F_0)_{\text{srs}}$ be an element matching $(\gamma, u') \in (S_n(\alpha) \times V')(F_0)_{\text{srs}}$. If $\mathbf{q}(u) \neq 0$, then*

$$-\partial\text{Orb}((\gamma, u'), \mathbf{1}_{(S_n \times V'_n)(O_{F_{0,v}})}) = \text{Int}_v(\delta, u) \cdot \log q_v.$$

Proof. We run the same argument as in the proof of [Proposition 13.8](#), where we note that the compactness (modulo $\text{U}(\mathbb{V}_n)(F_0, v)$) of the support of

the function $\text{Int}_v(\cdot, \cdot)$ holds by [Theorem 5.5](#). We then obtain a refinement of the equality in part (ii) of [Theorem 14.6](#):

$$-\partial\text{Orb}((\gamma, u'), \Phi'_v) \cdot \text{Orb}((\gamma, u'), \Phi'^{v, \infty}) = \text{Int}_v(\delta, u) \cdot \text{Orb}((\delta, u), \Phi^v).$$

(We warn the reader that here Φ does not have the archimedean component.) Here we note that $\text{Int}_v(\xi, \Phi)$ in part (ii) of [Theorem 14.6](#) is given by [\(9.18\)](#). Now the away from v factors on the two sides are equal and can be chosen to be non-zero (e.g., the function $\Phi^{(v)}$ can be chosen point-wise non-negative with non-empty support containing (δ, u)). \square

15. The proof of AFL

Now we return to the set up of [Conjecture 3.8](#) in [Section 3](#).

THEOREM 15.1. *[Conjecture 3.8](#) holds when $F_0 = \mathbb{Q}_p$ and $p \geq n$.*

Proof. The proof is parallel to that of [Theorem 13.9](#). We prove [Conjecture 3.8](#) part (b) by induction on $n = \dim \mathbb{V}_n$. The case $n = 1$ is known [\[46\]](#). Assume now that [Conjecture 3.8](#) part (b) holds for \mathbb{V}_{n-1} . Then by [Proposition 4.12](#) part (i), [Conjecture 3.8](#) part (a) holds for S_n . We now want to globalize the situation in order to apply [Corollary 14.8](#).

We start with the following local data:

- a place v_0 of $F_0 = \mathbb{Q}$, and an unramified (local) quadratic extension $F_{w_0}/F_{0, v_0}$;
- the non-split $F_{v_0}/F_{0, v_0}$ -hermitian space \mathbb{V}_n of dimension n ;
- $(g_{v_0}, u_{v_0}) \in (\text{U}(\mathbb{V}_n) \times \mathbb{V}_n)(F_{0, v_0})_{\text{srs}}$ — we further assume that the characteristic polynomial of g_{v_0} has integral coefficients (in $O_{F_{w_0}}$) and $\det(1 - g_{v_0})$ is a unit, and $\langle u_{v_0}, u_{v_0} \rangle \neq 0$;
- $(\gamma_{v_0}, u'_{v_0}) \in (S_n \times V')(F_{0, v_0})(F_{0, v_0})_{\text{srs}}$ matching (g_{v_0}, u_{v_0}) .

By the proof of [Theorem 13.9](#), there exist the following global data:

- an imaginary quadratic field F/F_0 such that $F \otimes_{F_0} F_{0, v_0} \simeq F_{w_0}$;
- a totally real number field F'_0 , and its quadratic extension $F' = F'_0 \otimes_{F_0} F$;
- an element $g \in F'^1$ such that $O_{F_{w_0}}[g] = O_{F_{w_0}}[g_{v_0}]$ as subrings of $F' \otimes_F F_{w_0}$;
- a totally positive definite n -dimensional F/F_0 -hermitian space $V(v_0)$ that is locally at v_0 isometric to \mathbb{V}_n , and an embedding $F'^1 \hookrightarrow \text{U}(V(v_0))(F_0)$;
- $u \in V(v_0)$ such that the pair (g, u) is v_0 -adically close to (g_{v_0}, u_{v_0}) (in particular, $\langle u, u \rangle \neq 0$).

Let $\alpha \in \mathcal{A}_n(F_0)$ denote the characteristic polynomial of g as an element in $\text{U}(V(v_0))(F_0)$.

Now we define the Shimura variety and its integral model \mathcal{M} as in [Definition 6.1](#) for the nearby hermitian space V of $V(v_0)$ at v_0 (that is, non-split at v_0 , with signature $(n-1, 1)$ at $v \mid \infty$, and isomorphic to $V(v_0)$ elsewhere).

Let \mathfrak{d} be a finite set of places as in [Section 6.2.2](#) such that $v_0 \nmid \mathfrak{d}$. Let S the set of non-archimedean places such that

- $v_0 \notin S$;
- S contains all places dividing \mathfrak{d} and all primes less than n ;
- for every non-archimedean $v \notin S \cup \{v_0\}$, the ring R_α is locally maximal at v .

Then we proceed as in the proof of [Theorem 13.9](#) to choose $(\gamma, u') \in (S_n \times V'_n)(F_0)$ to match (g, u) , and choose (partial) Gaussian test functions Φ and Φ' .

Now we apply [Corollary 14.8](#) to obtain

$$-\partial\text{Orb}((\gamma, u'), \Phi'_{v_0}) = \text{Int}_{v_0}(g, u) \log q_{v_0}.$$

Therefore [Conjecture 3.8](#) part (b) holds when $(g, u) \in (\text{U}(\mathbb{V}_n) \times \mathbb{V}_n)_{\text{srs}}$. By the local constancy of the orbital integral, and of the intersection numbers by [Theorem 5.5](#), near a strongly regular semisimple (g, u) , we conclude that [Conjecture 3.8](#) part (b) holds when $(g_{v_0}, u_{v_0}) \in (\text{U}(\mathbb{V}_n) \times \mathbb{V}_n)_{\text{srs}}$. This completes the induction. \square

Appendix A. Weil representation commutes with smooth transfer

We retain the notation from [Section 2](#). Let F/F_0 be a quadratic extension of local fields. (The case $E = F \times F$ could also be allowed but in that case the result below is trivial.) Recall that $V_n = F_0^n \times (F_0^n)^*$. We have a bijection of regular semisimple orbits (cf. [Section 2.3](#)):

$$\coprod_V [(\text{U}(V) \times V)(F_0)]_{\text{rs}} \xrightarrow{\sim} [S_n(F_0) \times V'_n]_{\text{rs}},$$

where the disjoint union runs over the set of isometry classes of F/F_0 -hermitian spaces V of dimension n . The notion of smooth transfer is as in [Definition 2.2](#) (with respect to the transfer factor there). Here let us focus on one hermitian space V at a time.

The Weil representation (for even dimensional quadratic space) is defined in [Section 11](#). Here we apply the formula (11.1) to the second variable in the functions in $\mathcal{S}(S_n \times V'_n)$ and $\mathcal{S}(\text{U}(V) \times V)$ respectively. To fix the set up, we recall that the structure of F_0 -bilinear symmetric pairing on V'_n is the tautological pairing

$$\langle u', u' \rangle = 2u_2(u_1), \quad u' = (u_1, u_2) \in F_0^n \times (F_0^n)^*,$$

and on V the quadratic form is the induced one, i.e.,

$$\langle u, u \rangle_{F_0} = \text{tr}_{F/F_0} \langle u, u \rangle_F, \quad u \in V,$$

where $\langle \cdot, \cdot \rangle_F : V \times V \rightarrow F$ is the hermitian pairing (F -linear on the first factor and conjugate F -linear on the second one).

We now deduce the following result from [\[47\]](#) when F is non-archimedean and [\[43\]](#) when F is archimedean.

THEOREM A.1 (Weil representation commutes with smooth transfer). *If $\Phi' \in \mathcal{S}(S_n \times V'_n)$ matches a function $\Phi \in \mathcal{S}(\mathrm{U}(V) \times V)$, then $\omega(h)\Phi'$ also matches $\omega(h)\Phi$ for any $h \in \mathrm{H}(F)$.*

Remark A.2. Similar results hold for the partial Fourier transforms on the Lie algebra $\mathfrak{s}_n \times V'_n$ and $\mathfrak{u}(V) \times V$. A similar result for the endoscopic transfer can be deduced from a theorem of Waldspurger.

Proof. We need to check the assertion for h of the form $\begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$, $\begin{pmatrix} 1 & b \\ & 1 \end{pmatrix}$ and $\begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$, as in (11.1).

The assertion for $h = \begin{pmatrix} 1 & b \\ & 1 \end{pmatrix}$, $b \in F$ is trivial.

Now let $h_a = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix}$. Then

$$\mathrm{Orb}((g, u), \omega(h_a)\Phi) = \chi_V(a)|a|^n \mathrm{Orb}((g, au), \Phi)$$

for all $(g, u) \in (\mathrm{U}(V) \times V)_{\mathrm{rs}}$. Here

$$\chi_V(a) = (a, (-1)^{\dim_F V} \det(V)),$$

where $\det(V)$ is the discriminant of V as a quadratic space. We claim

$$(A.1) \quad \chi_V(a) = \eta(a)^{\dim_F V}.$$

Since $\det(V_1 \oplus V_2) = \det(V_1) \det(V_2)$ (in $F_0^\times / (F_0^\times)^2$) for orthogonal direct sum $V_1 \oplus V_2$, it suffices to prove the claim when $\dim_F V = 1$. Then there are only two isometry classes, and one can check the claim directly.

On the other hand,

$$\mathrm{Orb}((\gamma, u'), \omega(h_a)\Phi', s) = \chi_{V'_n}(a)|a|^n \mathrm{Orb}((\gamma, au'), \Phi', s).$$

Now $\chi_{V'_n}$ is the trivial character since V'_n is an orthogonal direct sum of n -copies of the hyperbolic 2-space. We now note that the transfer factor (2.17) obeys

$$\omega(\gamma, au') = \eta(a)^n \omega(\gamma, u').$$

This proves the assertion for $\omega(h_a)$, $a \in F^\times$.

Finally, let $h = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$. Then

$$\mathrm{Orb}((g, u), \omega(h)\Phi) = \gamma_V \mathrm{Orb}((g, u), \widehat{\Phi}),$$

where γ_V is the Weil constant. We claim that for our V induced from a hermitian form,

$$\gamma_V = \eta(\det(V)_{F/F_0}) \epsilon(\eta, 1/2, \psi)^{\dim_F V},$$

where $\det(V)_{F/F_0} \in F_0^\times / \mathrm{Nm} F^\times$ is the hermitian discriminant of V (as an F/F_0 -hermitian space). First note that the right-hand side is multiplicative with respect to orthogonal direct sum $V_1 \oplus V_2$,

$$\det(V_1 \oplus V_2)_{F/F_0} = \det(V_1)_{F/F_0} \det(V_2)_{F/F_0}.$$

Note that, by definition, the Weil constant γ_V satisfies

$$\widehat{\psi \circ q} = \gamma_V \psi \circ (-q),$$

where $\psi \circ q : V \rightarrow F_0 \rightarrow \mathbb{C}$ (resp. $\psi \circ (-q)$) is the function precomposing ψ with q (resp. $-q$). Here the Fourier transform is understood as applied to distributions. It follows that it is also multiplicative with respect to orthogonal direct sum $V_1 \oplus V_2$:

$$\gamma_{V_1 \oplus V_2} = \gamma_{V_1} \cdot \gamma_{V_2}.$$

Therefore it suffices to show the claim when $\dim_F V = 1$. Then one can check the claim directly. In fact, it is easy to see that we have $\gamma_{V_a} = \eta(a)^{\dim_F V} \gamma_V$, where V_a denotes the new hermitian space by multiplying the hermitian form by $a \in F_0^\times$. Hence we may just check the case $\det(V)_{F/F_0} \in \mathrm{Nm} F^\times$, which is done in [20, Lemma 1.2] (where the constant $\lambda_{F/F_0}(\psi)$ in loc. cit. is the same as $\epsilon(\eta, 1/2, \psi)$).

On the other hand, the Weil constant $\gamma_{V'_n} = 1$ since V'_n is an orthogonal direct sum of n -copies of the hyperbolic 2-space. Hence

$$\mathrm{Orb}((\gamma, u'), \omega(h)\Phi', s) = \mathrm{Orb}((\gamma, u'), \widehat{\Phi}', s).$$

Now the desired assertion follows from [47, Th. 4.17]¹⁰ when F is non-archimedean, and the proof of [43, Th. 9.1] when F is archimedean. Note that in [43], $\epsilon(\eta, 1/2, \psi) = \sqrt{-1}$ for the choice of the additive character $\psi(x) = e^{2\pi\sqrt{-1}x}$, $x \in \mathbb{R}$. \square

Appendix B. Grothendieck groups for formal schemes

We collect some facts regarding formal schemes and the Grothendieck group of coherent sheaves, largely following the work by Gillet–Soulé [11]. No result here is new.

B.1. Grothendieck groups. Let (X, \mathcal{O}_X) be a noetherian formal scheme [15, §10]. Let Y be a closed formal subscheme of X (i.e., closed subscheme of a formal scheme in the terminology in loc. cit.). Let \mathcal{J} be the sheaf of ideals defining Y . A coherent sheaf \mathcal{F} of \mathcal{O}_X -module is said to be *formally supported* on Y if it is annihilated by \mathcal{J}^n for some $n \geq 1$. We make this explicit when (X, \mathcal{O}_X) is an affine formal scheme, say, the formal completion of $\mathrm{Spec} A$ at $\mathrm{Spec} A/I$ for an ideal I of A , where $A = \varprojlim_n A/I^n$ is I -adically complete. Then we may assume that Y is defined by an ideal J of A (i.e., $\mathcal{J} = J^\Delta$; cf. [15, §10.10]). Then a coherent sheaf \mathcal{F} of an \mathcal{O}_X -module is formally supported on Y if $M = \Gamma(X, \mathcal{F})$ as an A -module (equivalently the sheaf \widetilde{J} of an $\mathcal{O}_{\mathrm{Spec} A}$ -module) has support contained in the closed subset $\mathrm{Spec}(A/J)$ of $\mathrm{Spec} A$.

¹⁰Note that in [47], the factor $\eta(\det(V)_{F/F_0})$ is missing.

Then the definitions in [11, §1] for noetherian schemes carry over to the setting of noetherian formal schemes. Let $K'_0(X)$ denote the Grothendieck group of coherent sheaves of \mathcal{O}_X -modules. Let $K_0^Y(X)$ denote the Grothendieck group of finite complexes of coherent locally free \mathcal{O}_X -modules, acyclic outside Y (i.e., the homology sheaves are supported on Y); cf. [11, §1.2]. Let $K_0(X) = K_0^X(X)$. The tensor product of (complex of) locally free sheaves induces the cup product

$$\cup : K_0^Y(X) \times K_0^Z(X) \longrightarrow K_0^{Y \cap Z}(X)$$

by $[\mathcal{F}] \cup [\mathcal{G}] = [\mathcal{F} \otimes \mathcal{G}]$; cf. [11, §1.4].

There is a descending filtration on $K_0^Y(X)$ by the subgroups

$$(B.1) \quad F^i K_0^Y(X) = \cup_{Z \subset Y, \text{codim}_X Z \geq i} \text{Im}(K_0^Z(X) \longrightarrow K_0^Y(X)).$$

The associated graded groups are

$$(B.2) \quad \text{Gr}^i K_0^Y(X) = F^i K_0^Y(X) / F^{i+1} K_0^Y(X).$$

Similarly, there is an ascending filtration $F_i K'_0(X)$ on $K'_0(X)$,

$$F_i K'_0(X) = \cup_{Z \subset X, \dim Z \leq i} \text{Im}(K'_0(Z) \longrightarrow K'_0(X)).$$

From now on we assume that X is regular of pure dimension d . Then we have natural isomorphisms

$$K_0^Y(X) \xrightarrow{\sim} K'_0(Y)$$

and

$$F^{d-i} K_0^Y(X) \xrightarrow{\sim} F_i K'_0(Y).$$

When X is a scheme, the construction of the Adam operations $\{\psi^k \mid k \in \mathbb{Z}_{\geq 1}\}$ in [11] induce a decomposition

$$K_0^Y(X)_{\mathbb{Q}} = \bigoplus_{i \geq 0} K_0^Y(X)_{\mathbb{Q}}^i,$$

where ψ^k acts on (the “weight- i ” part) $K_0^Y(X)_{\mathbb{Q}}^i$ by the scalar k^i . Moreover, by [11, Prop. 5.3],

$$F^j K_0^Y(X)_{\mathbb{Q}} = \bigoplus_{i \geq j} K_0^Y(X)_{\mathbb{Q}}^i,$$

and for $j_1, j_2 \geq 0$, by [11, Prop. 5.5], the cup product has image

$$(B.3) \quad F^{j_1} K_0^Y(X)_{\mathbb{Q}} \cdot F^{j_2} K_0^Z(X)_{\mathbb{Q}} \subset F^{j_1+j_2} K_0^{Y \cap Z}(X)_{\mathbb{Q}}.$$

This inclusion is used in (7.15). When X is a formal scheme, we expect the same argument to prove (B.3), and we use this case of (B.3) only in the proof of [Proposition 5.2](#). However, due to the lack of reference, we also indicate a proof of [Proposition 5.2](#) without using (B.3); cf. [Remark 5.3](#).

Finally, we relax the noetherian hypothesis. For our purpose, we only consider locally noetherian formal schemes (X, \mathcal{O}_X) . It can be written as an increasing union indexed by a poset I ,

$$(X, \mathcal{O}_X) = \cup_{i \in I} (X_i, \mathcal{O}_{X_i}),$$

of noetherian formal subschemes such that the transition maps $f_{i,i'} : X_i \rightarrow X_{i'}$ are open immersions of formal schemes. We then define

$$K_0(X) = \varprojlim_{i \in I} K_0(X_i), \quad K'_0(X) = \varprojlim_{i \in I} K'_0(X_i).$$

If Y is a closed formal subscheme of X , setting $Y_i = Y \times_X X_i$ to write Y as the union of Y_i 's, we define

$$K_0^Y(X) = \varprojlim_{i \in I} K_0^{Y_i}(X_i).$$

Similarly, we have the filtrations $F^i K_0^Y(X)$ and $F_i K'_0(X)$, and they have the same properties as in the noetherian case. All of these K-groups depend only on X , rather than the choice of such unions $\cup_{i \in I} X_i$.

Now let $\pi : W \rightarrow S = \text{Spf } A$ be a morphism of formal schemes, where A is a complete discrete valuation ring. When π is proper [16, III, 3.4.1] and W is a *scheme* (not only a formal scheme), we have a “degree” map

$$\begin{aligned} K'_0(W) &\longrightarrow \mathbb{Z} \\ [\mathcal{E}] &\longmapsto \sum_{i \in \mathbb{Z}} (-1)^i \text{length}_{\mathcal{O}_S} R^i \pi_* \mathcal{E}. \end{aligned}$$

The assumption on π and W implies that all $R^i \pi_* \mathcal{E}$ are torsion coherent sheaves and hence have finite lengths. It is easy to see that this is independent of the choice of \mathcal{E} in its equivalence class. Now let X be regular with two closed formal subscheme Y and Z . If $\pi : W = Y \cap Z \rightarrow S = \text{Spf } A$ is proper and W is a scheme, we obtain a homomorphism

$$\begin{aligned} K_0^Y(X) \times K_0^Z(X) &\longrightarrow \mathbb{Z} \\ ([\mathcal{F}], [\mathcal{G}]) &\longmapsto \chi(X, \mathcal{F} \overset{\mathbb{L}}{\otimes} \mathcal{G}), \end{aligned}$$

where the Euler–Poincaré characteristic is defined by

$$(B.4) \quad \chi(X, \mathcal{F} \overset{\mathbb{L}}{\otimes} \mathcal{G}) := \sum_{i,j \in \mathbb{Z}} (-1)^{i+j} \text{length}_{\mathcal{O}_S} R^i \pi_* (\text{Tor}_j^{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})).$$

We also denote

$$(B.5) \quad Y \overset{\mathbb{L}}{\cap}_X Z := \mathcal{O}_Y \overset{\mathbb{L}}{\otimes}_{\mathcal{O}_X} \mathcal{O}_Z \in K'_0(Y \cap Z) \simeq K_0^{Y \cap Z}(X),$$

and if the ambient formal scheme X is self-evident, we simply write it as $Y \overset{\mathbb{L}}{\cap} Z$.

B.2. *A few lemmas.* For convenience we record the following results.

LEMMA B.1. *Let X be a locally noetherian formal schemes of the above type. Let $X = X_1 \cup X_2$ be a union of two closed formal subschemes. Then there is a natural isomorphism¹¹*

$$\begin{aligned} \frac{K'_0(X)}{K'_0(X_1 \cap X_2)} &\xrightarrow{\sim} \frac{K'_0(X_1)}{K'_0(X_1 \cap X_2)} \oplus \frac{K'_0(X_2)}{K'_0(X_1 \cap X_2)} \\ [\mathcal{E}] &\longmapsto ([\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}_{X_1}], [\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{O}_{X_2}]). \end{aligned}$$

Proof. We immediately reduce the question to the case when X is noetherian, which we assume now. Let \mathcal{I} and \mathcal{J} be the ideal sheaf of \mathcal{O}_X defining X_1 and X_2 respectively. Consider the exact sequence of \mathcal{O}_X -modules

$$0 \longrightarrow \mathcal{O}_X/(\mathcal{I} \cap \mathcal{J}) \longrightarrow \mathcal{O}_X/\mathcal{I} \oplus \mathcal{O}_X/\mathcal{J} \longrightarrow \mathcal{O}_X/(\mathcal{I} + \mathcal{J}) \longrightarrow 0.$$

Tensoring \mathcal{E} , we obtain an exact sequence

$$\begin{aligned} \mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{E}, \mathcal{O}_{X_1 \cap X_2}) &\longrightarrow \mathcal{E} \otimes \mathcal{O}_X/(\mathcal{I} \cap \mathcal{J}) \longrightarrow \mathcal{E} \otimes \mathcal{O}_{X_1} \oplus \mathcal{E} \otimes \mathcal{O}_{X_2} \\ &\longrightarrow \mathcal{E} \otimes \mathcal{O}_{X_1 \cap X_2} \longrightarrow 0. \end{aligned}$$

Since both $\mathrm{Tor}_1^{\mathcal{O}_X}(\mathcal{E}, \mathcal{O}_{X_1 \cap X_2})$ and $\mathcal{E} \otimes \mathcal{O}_{X_1 \cap X_2}$ lie in $K'_0(X_1 \cap X_2)$, we have

$$[\mathcal{E} \otimes \mathcal{O}_{X_1}] + [\mathcal{E} \otimes \mathcal{O}_{X_2}] = [\mathcal{E} \otimes \mathcal{O}_X/(\mathcal{I} \cap \mathcal{J})] \in \frac{K'_0(X)}{K'_0(X_1 \cap X_2)}.$$

Since $X = X_1 \cup X_2$, we have $\mathcal{I} \cap \mathcal{J} = 0$ and the proof is complete. \square

In the case of “proper intersection,” the derived tensor product can be simplified:

LEMMA B.2. *Let X be a (locally noetherian) pure finite dimensional formal scheme of the above type, and let Z_1, Z_2 be two pure dimensional closed formal subschemes on X . Assume that the closed immersion $Z_1 \rightarrow X$ is a regular immersion (e.g., if both X and Z_1 are regular), and Z_2 is Cohen–Macaulay:*

$$\begin{array}{ccc} Z_1 \cap Z_2 & \longrightarrow & Z_2 \\ \downarrow & \square & \downarrow \\ Z_1 & \longrightarrow & X. \end{array}$$

(i) *If $Z_1 \cap Z_2$ has the expected dimension (i.e., $\mathrm{codim}_X Z_1 \cap Z_2 = \mathrm{codim}_X Z_1 + \mathrm{codim}_X Z_2$ at every point of $Z_1 \cap Z_2$), then the higher Tor sheaves vanish, i.e.,*

$$\mathrm{Tor}_i^{\mathcal{O}_X}(\mathcal{O}_{Z_1}, \mathcal{O}_{Z_2}) = 0, \quad i > 0.$$

¹¹Here $K'_0(X_1 \cap X_2) \rightarrow K'_0(X_1)$ is not necessarily injective, so the quotient simply denotes the cokernel.

In particular, as elements in $K'_0(Z_1 \cap Z_2)$,

$$\mathcal{O}_{Z_1} \overset{\mathbb{L}}{\otimes} \mathcal{O}_{Z_2} = \mathcal{O}_{Z_1} \otimes \mathcal{O}_{Z_2}.$$

(ii) Let $Z_1 \cap Z_2 = Y \cup Y'$ be a union of closed formal subschemes such that Y has the expected dimension. Then

$$\mathrm{Tor}_i^{\mathcal{O}_X}(\mathcal{O}_{Z_1}, \mathcal{O}_{Z_2})|_Y \equiv 0, \quad i > 0,$$

as an element in $K'_0(Y)/K'_0(Y \cap Y')$.

Proof. This follows from the same argument as in the proof of [37, Prop. 8.10] regarding the vanishing of higher Tor terms. We prove the first part; the second part is proved similarly by combining Lemma B.1.

Let x be a point on $Z_1 \cap Z_2$. We need to show that $(\mathcal{O}_{Z_1} \overset{\mathbb{L}}{\otimes} \mathcal{O}_{Z_2})_x$ is represented by $\mathcal{O}_{Z_1 \cap Z_2, x}$. Let R be the local ring of x on X . Since the closed immersion $Z_1 \rightarrow X$ is a regular immersion, by definition Z_1 is defined at x by a regular sequence f_1, \dots, f_m of R . Then the Koszul complex $K(f_1, \dots, f_m)$ is a free resolution of the R -module $\mathcal{O}_{Z_1, x}$. It follows that the complex $K(f_1, \dots, f_m) \otimes_R \mathcal{O}_{Z_2, x}$ represents $(\mathcal{O}_{Z_1} \overset{\mathbb{L}}{\otimes} \mathcal{O}_{Z_2})_x$.

Now, since Z_2 is Cohen–Macaulay, the dimension hypothesis implies that the images $\bar{f}_1, \dots, \bar{f}_m$ of f_1, \dots, f_m in $\mathcal{O}_{Z_2, x}$ again form a regular sequence which generates the ideal defining $Z_1 \cap Z_2$ at x in Z_2 . Hence $K(\bar{f}_1, \dots, \bar{f}_m)$ is a free resolution of the $\mathcal{O}_{Z_2, x}$ -module $\mathcal{O}_{Z_1 \cap Z_2, x}$. On the other hand, we have

$$K(f_1, \dots, f_m) \otimes_R \mathcal{O}_{Z_2, x} = K(\bar{f}_1, \dots, \bar{f}_m).$$

It follows that $(\mathcal{O}_{Z_1} \overset{\mathbb{L}}{\otimes} \mathcal{O}_{Z_2})_x$ is represented by $K(\bar{f}_1, \dots, \bar{f}_m)$, or equivalently by $\mathcal{O}_{Z_1 \cap Z_2, x}$. This completes the proof. \square

References

- [1] R. BEUZART-PLESSIS, Comparison of local spherical characters and the Ichino–Ikeda conjecture for unitary groups, 2016. [arXiv 1602.06538](https://arxiv.org/abs/1602.06538).
- [2] R. BEUZART-PLESSIS, A new proof of Jacquet–Rallis’s fundamental lemma, 2019. [arXiv 1901.02653](https://arxiv.org/abs/1901.02653).
- [3] J.-B. BOST, H. GILLET, and C. SOULÉ, Heights of projective varieties and positive Green forms, *J. Amer. Math. Soc.* **7** no. 4 (1994), 903–1027. [MR 1260106](https://doi.org/10.1090/S0894-0347-1994-01140-1). [Zbl 0973.14013](https://doi.org/10.2307/2152736). <https://doi.org/10.2307/2152736>.
- [4] J. H. BRUINIER, Regularized theta lifts for orthogonal groups over totally real fields, *J. Reine Angew. Math.* **672** (2012), 177–222. [MR 2995436](https://doi.org/10.1515/crelle.2011.163). [Zbl 1268.11058](https://doi.org/10.1515/crelle.2011.163). <https://doi.org/10.1515/crelle.2011.163>.
- [5] J. H. BRUINIER, B. HOWARD, S. KUDLA, M. RAPOPORT, and T. YANG, Modularity of generating series of divisors on unitary Shimura varieties, 2017, to appear in *Astérisque*. [arXiv 1702.07812](https://arxiv.org/abs/1702.07812).

- [6] J. H. BRUINIER, S. S. KUDLA, and T. YANG, Special values of Green functions at big CM points, *Int. Math. Res. Not. IMRN* no. 9 (2012), 1917–1967. [MR 2920820](#). [Zbl 1281.11063](#). <https://doi.org/10.1093/imrn/rnr095>.
- [7] J. I. BURGOS GIL, J. KRAMER, and U. KÜHN, Cohomological arithmetic Chow rings, *J. Inst. Math. Jussieu* 6 no. 1 (2007), 1–172. [MR 2285241](#). [Zbl 1115.14013](#). <https://doi.org/10.1017/S1474748007000011>.
- [8] S. EHLEN and S. SANKARAN, On two arithmetic theta lifts, *Compos. Math.* 154 no. 10 (2018), 2090–2149. [MR 3867297](#). [Zbl 1436.11046](#). <https://doi.org/10.1112/s0010437x18007327>.
- [9] W. T. GAN, B. H. GROSS, and D. PRASAD, Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups, in *Sur les Conjectures de Gross et Prasad. I*, Astérisque 346, Math. Soc. France, Paris, 2012, pp. 1–109. [MR 3202556](#). [Zbl 1280.22019](#).
- [10] H. GILLET, Arithmetic intersection theory on Deligne-Mumford stacks, in *Motives and Algebraic Cycles, Fields Inst. Commun.* 56, Amer. Math. Soc., Providence, RI, 2009, pp. 93–109. [MR 2562454](#). [Zbl 1196.14022](#). <https://doi.org/10.1090/fic/056/04>.
- [11] H. GILLET and C. SOULÉ, Intersection theory using Adams operations, *Invent. Math.* 90 no. 2 (1987), 243–277. [MR 0910201](#). [Zbl 0632.14009](#). <https://doi.org/10.1007/BF01388705>.
- [12] H. GILLET and C. SOULÉ, Arithmetic intersection theory, *Inst. Hautes Études Sci. Publ. Math.* no. 72 (1990), 93–174. [MR 1087394](#). [Zbl 0741.14012](#). Available at http://www.numdam.org/item?id=PMIHES_1990_72_93_0.
- [13] J. GORDON, Transfer to characteristic zero, appendix to [45].
- [14] B. H. GROSS and D. B. ZAGIER, Heegner points and derivatives of L -series, *Invent. Math.* 84 no. 2 (1986), 225–320. [MR 0833192](#). [Zbl 0608.14019](#). <https://doi.org/10.1007/BF01388809>.
- [15] A. GROTHENDIECK, Éléments de géométrie algébrique. I. Le langage des schémas, *Inst. Hautes Études Sci. Publ. Math.* no. 4 (1960), 5–228. [MR 0217083](#). [Zbl 0118.36206](#). Available at http://www.numdam.org/item/PMIHES_1960_4_5_0/.
- [16] A. GROTHENDIECK, Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. première partie, *Inst. Hautes Études Sci. Publ. Math.* no. 11 (1961), 5–167. [MR 0217085](#). Available at http://www.numdam.org/item/PMIHES_1961_11_5_0/.
- [17] X. HE, C. LI, and Y. ZHU, Fine Deligne-Lusztig varieties and arithmetic fundamental lemmas, *Forum Math. Sigma* 7 (2019), e47, 55. [MR 4061970](#). [Zbl 1439.11142](#). <https://doi.org/10.1017/fms.2019.45>.
- [18] B. HOWARD, Complex multiplication cycles and Kudla-Rapoport divisors, *Ann. of Math.* (2) 176 no. 2 (2012), 1097–1171. [MR 2950771](#). [Zbl 1327.14126](#). <https://doi.org/10.4007/annals.2012.176.2.9>.
- [19] H. JACQUET, Sur un résultat de Waldspurger, *Ann. Sci. École Norm. Sup. (4)* 19 no. 2 (1986), 185–229. [MR 0868299](#). [Zbl 0605.10015](#). <https://doi.org/10.24033/asens.1506>.

- [20] H. JACQUET and R. P. LANGLANDS, *Automorphic Forms on $GL(2)$* , *Lecture Notes in Math* 114, Springer-Verlag, New York, 1970. MR 0401654. Zbl 0236.12010. <https://doi.org/10.1007/BFb0058988>.
- [21] H. JACQUET and S. RALLIS, On the Gross-Prasad conjecture for unitary groups, in *On Certain L-Functions*, *Clay Math. Proc.* 13, Amer. Math. Soc., Providence, RI, 2011, pp. 205–264. MR 2767518. Zbl 1222.22018.
- [22] S. S. KUDLA, Algebraic cycles on Shimura varieties of orthogonal type, *Duke Math. J.* 86 no. 1 (1997), 39–78. MR 1427845. Zbl 0879.11026. <https://doi.org/10.1215/S0012-7094-97-08602-6>.
- [23] S. S. KUDLA, Central derivatives of Eisenstein series and height pairings, *Ann. of Math.* (2) 146 no. 3 (1997), 545–646. MR 1491448. Zbl 0990.11032. <https://doi.org/10.2307/2952456>.
- [24] S. KUDLA and M. RAPOPORT, Special cycles on unitary Shimura varieties I. Unramified local theory, *Invent. Math.* 184 no. 3 (2011), 629–682. MR 2800697. Zbl 1229.14020. <https://doi.org/10.1007/s00222-010-0298-z>.
- [25] S. KUDLA and M. RAPOPORT, Special cycles on unitary Shimura varieties II: Global theory, *J. Reine Angew. Math.* 697 (2014), 91–157. MR 3281653. Zbl 1316.11050. <https://doi.org/10.1515/crelle-2012-0121>.
- [26] C. LI and W. ZHANG, Kudla–Rapoport cycles and derivatives of local densities, 2019. [arXiv 1908.01701](https://arxiv.org/abs/1908.01701).
- [27] C. LI and Y. ZHU, Remarks on the arithmetic fundamental lemma, *Algebra Number Theory* 11 no. 10 (2017), 2425–2445. MR 3744362. Zbl 1439.11143. <https://doi.org/10.2140/ant.2017.11.2425>.
- [28] Y. LIU, Arithmetic theta lifting and L -derivatives for unitary groups, I, *Algebra Number Theory* 5 no. 7 (2011), 849–921. MR 2928563. Zbl 1258.11060. <https://doi.org/10.2140/ant.2011.5.849>.
- [29] Y. LIU, Relative trace formulae toward Bessel and Fourier–Jacobi periods on unitary groups, *Manuscripta Math.* 145 no. 1–2 (2014), 1–69. MR 3244725. Zbl 1301.11050. <https://doi.org/10.1007/s00229-014-0666-x>.
- [30] Y. LIU, Fourier–Jacobi cycles and arithmetic relative trace formula (with an appendix by C. Li and Y. Zhu), 2021. [arXiv 2102.11518](https://arxiv.org/abs/2102.11518).
- [31] A. MIHATSCH, Relative unitary RZ-spaces and the arithmetic fundamental lemma, *Journal of the Institute of Mathematics of Jussieu*, 61 pp., published online 14 March 2020. <https://doi.org/10.1017/S1474748020000079>.
- [32] A. MIHATSCH, Local constancy of intersection numbers, 2020. [arXiv 2004.12172](https://arxiv.org/abs/2004.12172).
- [33] A. MIHATSCH, On the arithmetic fundamental lemma through Lie algebras, *Math. Z.* 287 no. 1–2 (2017), 181–197. MR 3694673. Zbl 1410.11071. <https://doi.org/10.1007/s00209-016-1822-7>.
- [34] B. C. NGÔ, Le lemme fondamental pour les algèbres de Lie, *Publ. Math. Inst. Hautes Études Sci.* no. 111 (2010), 1–169. MR 2653248. Zbl 1200.22011. <https://doi.org/10.1007/s10240-010-0026-7>.
- [35] F. OBERHETTINGER, On the derivative of Bessel functions with respect to the order, *J. Math. and Phys.* 37 (1958), 75–78. MR 0092871. Zbl 0111.06607. <https://doi.org/10.1002/sapm195837175>.

- [36] T. ODA and M. TSUZUKI, Automorphic Green functions associated with the secondary spherical functions, *Publ. Res. Inst. Math. Sci.* **39** no. 3 (2003), 451–533. [MR 2001185](#). [Zbl 1044.11033](#). <https://doi.org/10.2977/prims/1145476077>.
- [37] M. RAPOPORT, B. SMITHLING, and W. ZHANG, On the arithmetic transfer conjecture for exotic smooth formal moduli spaces, *Duke Math. J.* **166** no. 12 (2017), 2183–2336. [MR 3694568](#). [Zbl 1411.11056](#). <https://doi.org/10.1215/00127094-2017-0003>.
- [38] M. RAPOPORT, B. SMITHLING, and W. ZHANG, Regular formal moduli spaces and arithmetic transfer conjectures, *Math. Ann.* **370** no. 3-4 (2018), 1079–1175. [MR 3770164](#). [Zbl 1408.14143](#). <https://doi.org/10.1007/s00208-017-1526-2>.
- [39] M. RAPOPORT, B. SMITHLING, and W. ZHANG, On Shimura varieties for unitary groups, 2019, *Pure Appl. Math. Q.* (special issue in honor of D. Mumford), to appear. [arXiv 1906.12346](#).
- [40] M. RAPOPORT, B. SMITHLING, and W. ZHANG, Arithmetic diagonal cycles on unitary Shimura varieties, *Compos. Math.* **156** no. 9 (2020), 1745–1824. [MR 4167594](#). [Zbl 07270080](#). <https://doi.org/10.1112/s0010437x20007289>.
- [41] M. RAPOPORT, U. TERSTIEGE, and W. ZHANG, On the arithmetic fundamental lemma in the minuscule case, *Compos. Math.* **149** no. 10 (2013), 1631–1666. [MR 3123304](#). [Zbl 1300.11069](#). <https://doi.org/10.1112/S0010437X13007239>.
- [42] I. VOLLAARD and T. WEDHORN, The supersingular locus of the Shimura variety of $GU(1, n - 1)$ II, *Invent. Math.* **184** no. 3 (2011), 591–627. [MR 2800696](#). [Zbl 1227.14027](#). <https://doi.org/10.1007/s00222-010-0299-y>.
- [43] H. XUE, On the global Gan–Gross–Prasad conjecture for unitary groups: approximating smooth transfer of Jacquet–Rallis, *J. Reine Angew. Math.* **756** (2019), 65–100. [MR 4026449](#). [Zbl 07126236](#). <https://doi.org/10.1515/crelle-2017-0016>.
- [44] X. YUAN, S.-W. ZHANG, and W. ZHANG, The Gross–Kohnen–Zagier theorem over totally real fields, *Compos. Math.* **145** no. 5 (2009), 1147–1162. [MR 2551992](#). [Zbl 1235.11056](#). <https://doi.org/10.1112/S0010437X08003734>.
- [45] Z. YUN, The fundamental lemma of Jacquet and Rallis, *Duke Math. J.* **156** no. 2 (2011), 167–227, with an appendix by Julia Gordon. [MR 2769216](#). [Zbl 1211.14039](#). <https://doi.org/10.1215/00127094-2010-210>.
- [46] W. ZHANG, On arithmetic fundamental lemmas, *Invent. Math.* **188** no. 1 (2012), 197–252. [MR 2897697](#). [Zbl 1247.14031](#). <https://doi.org/10.1007/s00222-011-0348-1>.
- [47] W. ZHANG, Fourier transform and the global Gan–Gross–Prasad conjecture for unitary groups, *Ann. of Math.* (2) **180** no. 3 (2014), 971–1049. [MR 3245011](#). [Zbl 1322.11048](#). <https://doi.org/10.4007/annals.2014.180.3.4>.

(Received: September 10, 2019)

(Revised: July 6, 2020)

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA

E-mail: weizhang@mit.edu