

SPECTRAL BAND DEGENERACIES OF $\frac{\pi}{2}$ -ROTATIONALLY INVARIANT PERIODIC SCHRÖDINGER OPERATORS*

R. T. KELLER[†], J. L. MARZUOLA[‡], B. OSTING[§], AND M. I. WEINSTEIN[¶]

Abstract. The dynamics of waves in periodic media is determined by the band structure of the underlying periodic Hamiltonian. Symmetries of the Hamiltonian can give rise to novel properties of the band structure. Here we consider a class of periodic Schrödinger operators, $H_V = -\Delta + V$, where V is periodic with respect to the lattice of translates $\Lambda = \mathbb{Z}^2$. The potential is also assumed to be real-valued, sufficiently regular, and such that, with respect to some origin of coordinates, inversion symmetric (even) and invariant under $\pi/2$ rotation. We present general conditions ensuring that the band structure of H_V contains dispersion surfaces which touch at multiplicity two eigenvalues at the vertices (high-symmetry quasi momenta) of the Brillouin zone. Locally, the band structure consists of two intersecting dispersion surfaces described by a normal form which is $\pi/2$ -rotationally invariant, and to leading order homogeneous of degree two. Furthermore, the effective dynamics of wave-packets, which are spectrally concentrated near high-symmetry quasi momenta, is given by a system of coupled Schrödinger equations with indefinite effective mass tensor. For small amplitude potentials, εV with ε small or weak coupling, certain distinguished Fourier coefficients of the potential control which of the low-lying dispersion surfaces (first four) of $H^\varepsilon = H_{\varepsilon V}$ intersect and have the above local behavior. The existence of quadratically touching dispersion surfaces with the above properties persists for all real ε , without restriction on the size of ε , except for ε in a discrete set. Our results apply to periodic superpositions of spatially localized “atomic potentials” centered on the square (\mathbb{Z}^2) and Lieb lattices. We show, in particular, that the well-known conical plus flat-band structure of the three dispersion surfaces of the Lieb lattice tight-binding model does not persist in the corresponding Schrödinger operator with finite depth potential wells. Finally, we corroborate our analytical results with extensive numerical simulations. The present results are the \mathbb{Z}^2 -analogue of results obtained for conical degenerate points (Dirac points) in the band structure for honeycomb structures.

Key words. band structure, periodic potentials, Lieb lattice, Schrödinger operator, spectral band degeneracies, homogenization

AMS subject classifications. 35Q40, 35Q60, 35P99

DOI. 10.1137/18M1171527

1. Introduction. The dynamics of waves in periodic media are determined by the band structure of the underlying Hamiltonian; see, for example, [2, 13, 20, 14, 15]. Symmetries of the underlying Hamiltonian give rise to novel properties of the band structure. An important example is the band structure of the single electron model

*Received by the editors February 20, 2018; accepted for publication (in revised form) August 20, 2018; published electronically October 25, 2018.

<http://www.siam.org/journals/mms/16-4/M117152.html>

Funding: This research of the first author was supported by U.S. National Science Foundation (NSF) grants DGE-1144155 and DGE 16-44869. The work of the second author was supported by NSF grant DMS-1312874 and NSF CAREER grant DMS-1352353. The work of the third author was supported by NSF grant DMS 16-19755. The work of the first and fourth authors was supported by NSF grants DMS-1412560 and DMS-1620418 and RNMS grant 11-07444 (Ki-Net). The fourth author was supported by Simons Foundation Math + X Investigator Award 376319.

[†]Department of Applied Physics and Applied Mathematics, Columbia University, New York, NY 10027 (rachael.keller@columbia.edu).

[‡]Department of Mathematics, University of North Carolina–Chapel Hill, Chapel Hill, NC 27599 (marzuola@math.unc.edu).

[§]Department of Mathematics, University of Utah, Salt Lake City, UT 84112 (osting@math.utah.edu).

[¶]Department of Applied Physics and Applied Mathematics and Department of Mathematics, Columbia University, New York, NY 10027 (miw2103@columbia.edu).




FIG. 1.1. An illustration of the square (left) and Lieb (right) lattices with corresponding fundamental cells, Ω (shaded). The Lieb lattice is the union of three interpenetrating sublattices, labeled A (red), B (black), and C (blue).

of graphene and its artificial analogues. Here, $H_V = -\Delta + V$, with V a real-valued potential with the symmetries of a hexagonal tiling of the plane. It is well-known that the band structure contains Dirac points, conical singularities at the intersections of dispersion surfaces which occur at the vertices (high-symmetry quasi momenta) of the hexagonal Brillouin zone; see, for example, [18, 9, 7, 4]. A consequence is the massless Dirac dynamics of wave-packets (quasi particles) which evolve from initial data which are spectrally localized near Dirac points [18, 10].

In this article, we consider a class of periodic Schrödinger operators on \mathbb{R}^2 , whose underlying period lattice is \mathbb{Z}^2 and such that the potential is real, inversion symmetric, and invariant under $\pi/2$ rotation. We call such potentials *admissible*; see Definition 2.3. The class of potentials to which our results apply includes those which are superpositions of localized potentials (say, potential wells or potential barriers) centered on a discrete structure with the appropriate symmetries. Two such examples are illustrated in Figure 1.1; the square lattice (left) and the Lieb lattice (right) are displayed together with corresponding choices of fundamental cells. We call these two types of potentials *square lattice potentials* and *Lieb lattice potentials*; see Examples 2.5 and 2.6 and the potentials in Figures 7.1–7.5.

Our goal is to study symmetry-induced characteristics in the band structure of such operators and to explore these in the context of the above two examples. The present results are the \mathbb{Z}^2 -analogue of results obtained in honeycomb structures [5, 11, 9, 8, 16, 4]. Our proofs make use of the framework developed in [9, 8].

For the class of potentials we consider, the nature of band degeneracies at high-symmetry quasi momenta is described by two intersecting dispersion surfaces which are locally characterized by a normal form which is $\pi/2$ -rotationally invariant, and to leading order homogeneous of degree two. A consequence is that the dynamics of wave-packets, which are spectrally concentrated near such high-symmetry quasi momenta, is given by an effective system of coupled linear time-dependent Schrödinger equations with indefinite effective mass tensor. This is in contrast to the case of honeycomb structures where the band degeneracies at the high-symmetry points are conical (so-called Dirac points) and the effective dynamics is given by a time-dependent system of Dirac equations.

1.1. A quick review of Floquet–Bloch theory [20, 14, 15, 6, 13]. Consider the periodic Schrödinger operator, $H_V = -\Delta + V$, where V is real-valued and periodic with respect to a lattice $\Lambda = \mathbb{Z}\mathbf{v}_1 \oplus \mathbb{Z}\mathbf{v}_2$; for all $\mathbf{x} \in \mathbb{R}^2$ and $\mathbf{v} \in \Lambda$, we have $V(\mathbf{x} + \mathbf{v}) =$




FIG. 1.2. Brillouin zone, \mathcal{B} , for the square lattice $\Gamma = \mathbb{Z}^2$. The points $\Gamma = (0, 0)$, $\mathbf{M} = (\pi, \pi)$, and $\mathbf{X} = (\pi, 0)$ are labeled, along with successive $\pi/2$ rotations of \mathbf{M} by R ; see (2.5). An irreducible Brillouin zone is shaded in blue. In later figures, the graphs of dispersion relation are plotted along the cyclic path $\Gamma \rightarrow \mathbf{X} \rightarrow \mathbf{M} \rightarrow \Gamma$.

$V(\mathbf{x})$. Introduce the dual lattice, $\Lambda^* = \mathbb{Z}\mathbf{k}_1 \oplus \mathbb{Z}\mathbf{k}_2$, such that $\mathbf{k}_l \cdot \mathbf{v}_m = 2\pi\delta_{lm}$, and spaces of Λ -periodic and \mathbf{k} -pseudoperiodic functions:

$$L^2(\mathbb{R}^2/\Lambda) = \left\{ f \in L^2_{\text{loc}}(\mathbb{R}^2) : f(\mathbf{x} + \mathbf{v}) = f(\mathbf{x}) \text{ almost everywhere in } \mathbf{x} \text{ for all } \mathbf{v} \in \Lambda \right\}$$

and

$$L^2_{\mathbf{k}} = \left\{ f \in L^2_{\text{loc}}(\mathbb{R}^2) : e^{-i\mathbf{k} \cdot \mathbf{x}} f(\mathbf{x}) \in L^2(\mathbb{R}^2/\Lambda) \right\}.$$

Functions $f \in L^2_{\mathbf{k}}$ satisfy $f(\mathbf{x} + \mathbf{v}) = e^{i\mathbf{k} \cdot \mathbf{v}} f(\mathbf{x})$ almost everywhere in \mathbf{x} for all $\mathbf{v} \in \Lambda$. The inner product on $L^2(\mathbb{R}^2/\Lambda)$ is given by $\langle f, g \rangle_{L^2(\mathbb{R}^2/\Lambda)} = \int_{\Omega} \bar{f}g$, where $\Omega \subset \mathbb{R}^2$ is a period cell. Since $f, g \in L^2_{\mathbf{k}}$ implies that $\bar{f}g \in L^1(\mathbb{R}^2/\Lambda)$, the same expression defines an inner product on $L^2_{\mathbf{k}}$.

Floquet–Bloch states are solutions of the self-adjoint \mathbf{k} -pseudoperiodic eigenvalue problem:

$$(1.1) \quad H_V \Phi = \mu \Phi, \quad \Phi \in L^2_{\mathbf{k}}.$$

Since this boundary condition satisfied by $f \in L^2_{\mathbf{k}}$ is invariant under $\mathbf{k} \mapsto \mathbf{k} + \tilde{\mathbf{k}}$, for any $\tilde{\mathbf{k}} \in \Lambda^*$, we may restrict to \mathbf{k} varying over a fundamental period cell in the dual variable. This Brillouin zone, \mathcal{B} , is often taken to be the set of all points in $\mathbf{k} \in \mathbb{R}^2$ which are closer to the origin than to any other point in Λ^* . For the case of the square lattice, $\Lambda = \mathbb{Z}^2$,

$$(1.2) \quad \mathcal{B} = \{\mathbf{k} = (k^{(1)}, k^{(2)}) : -\pi \leq k^{(1)} \leq \pi, -\pi \leq k^{(2)} \leq \pi\};$$

see Figure 1.2.

An alternative formulation is to write $\Phi(\mathbf{x}) = e^{i\mathbf{k} \cdot \mathbf{x}} \phi(\mathbf{x})$ and seek, for all $\mathbf{k} \in \mathcal{B}$, solutions of the self-adjoint periodic eigenvalue problem:

$$(1.3) \quad H_V(\mathbf{k})\phi(\mathbf{x}) \equiv (-\nabla_{\mathbf{x}} + i\mathbf{k})^2 \phi = \mu\phi(\mathbf{x}), \quad \phi \in L^2(\mathbb{R}^2/\Lambda).$$

For each $\mathbf{k} \in \mathcal{B}$, the self-adjoint elliptic eigenvalue problem (1.3) has a discrete set of eigenpairs $(\mu_b(\mathbf{k}), \phi_b(\mathbf{x}; \mathbf{k}))$, $b = 1, 2, 3, \dots$, where the eigenvalues may be listed with multiplicity, in order:

$$\mu_1(\mathbf{k}) \leq \mu_2(\mathbf{k}) \leq \dots \leq \mu_b(\mathbf{k}) \leq \dots,$$

where $\{\phi_b(\mathbf{x}; \mathbf{k})\}_{b \geq 1}$ can be taken to be a complete orthonormal sequence in $L^2(\mathbb{R}^2/\Lambda)$. Moreover, the family of states $\Phi_b(\mathbf{x}; \mathbf{k}) = e^{i\mathbf{k} \cdot \mathbf{x}} \phi_b(\mathbf{x}; \mathbf{k})$ where $b \geq 1$ and $\mathbf{k} \in \mathcal{B}$ are complete in $L^2(\mathbb{R}^2)$. The eigenvalue mappings $\mu_b: \mathcal{B} \rightarrow \mathbb{R}$ are Lipschitz continuous [3, 10] and are called *dispersion relations*, and their graphs are called *dispersion surfaces*. The collection of all dispersion relations is called the *band structure* of the periodic Schrödinger operator, H_V . The spectrum of H_V acting in $L^2(\mathbb{R}^2)$ is the union of the closed real intervals: $\sigma(H_V) = \mu_1(\mathcal{B}) \cup \mu_2(\mathcal{B}) \cup \cdots \cup \mu_b(\mathcal{B}) \cup \cdots$.

1.2. Summary of results.

We summarize our main results.

1. Theorem 4.1:

(A) We present general conditions on the admissible potential, V , for the following scenario: there exists $\mu_S \in \mathbb{R}$ such that for all vertices, \mathbf{M}_* of \mathcal{B} , μ_S is an $L^2_{\mathbf{M}_*}$ -eigenvalue of H_V of geometric multiplicity two, with $L^2_{\mathbf{M}_*}$ -kernel $(H_V - \mu_S I) = \text{span}\{\Phi_1, \Phi_2\}$.

(B) There exist dispersion maps $\mathbf{k} \mapsto \mu_-(\mathbf{k})$ and $\mathbf{k} \mapsto \mu_+(\mathbf{k})$, defined for $\mathbf{k} \in \mathcal{B}$ with $\mu_-(\mathbf{k}) \leq \mu_+(\mathbf{k})$ and such that for all vertices, \mathbf{M}_* , of \mathcal{B} we have $\mu_{\pm}(\mathbf{M}_*) = \mu_S$; the dispersion surfaces associated with μ_- and μ_+ touch at the energy/quasi momentum pairs (\mathbf{M}_*, μ_S) .

(C) The two touching dispersion surfaces are locally described by a normal form which is $\pi/2$ -rotationally invariant and, at leading order in $|\mathbf{k} - \mathbf{M}_*|$, homogeneous of degree two. In particular, for $|\kappa| = |(\kappa_1, \kappa_2)| = \sqrt{\kappa_1^2 + \kappa_2^2}$ small,

(1.4)

$$\mu_{\pm}(\mathbf{M} + \kappa) - \mu_S = (1 - \alpha)|\kappa|^2 + \mathcal{Q}_6(\kappa) \pm \sqrt{\left| \gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2 \right|^2 + \mathcal{Q}_8(\kappa)},$$

where $\alpha \in \mathbb{R}$ and $\beta, \gamma \in \mathbb{C}$ are constants, and $\mathcal{Q}_6(\kappa) = \mathcal{O}(|\kappa|^6)$ and $\mathcal{Q}_8(\kappa) = \mathcal{O}(|\kappa|^8)$ are analytic functions of κ and invariant under $\pi/2$ -rotation: $(\kappa_1, \kappa_2) \mapsto (-\kappa_2, \kappa_1)$.

(D) Corollary 4.2: If V is, in addition, assumed to be *reflection-invariant* with respect to the diagonal of the fundamental cell, Ω (see Figure 1.1), then (1.4) reduces to

$$(1.5) \quad \mu_{\pm}(\mathbf{M} + \kappa) - \mu_S = (1 - \alpha)|\kappa|^2 + \mathcal{Q}_6(\kappa) \pm \sqrt{\left| 2\beta\kappa_1\kappa_2 \right|^2 + \mathcal{Q}_8(\kappa)},$$

with the same notation as in (C).

2. Theorem 5.3: Consider $H^\varepsilon \equiv H_{\varepsilon V} = -\Delta + \varepsilon V$, where V is admissible and ε is real. Let $V_{1,1}$ and $V_{1,0}$ denote the $(1,1)$ and $(1,0)$ indexed Fourier coefficients of V (see (2.4)) and assume the (generically satisfied) nondegeneracy condition: $V_{1,1} \neq V_{1,0}$. Then, for all nonzero and sufficiently small ε , there are two dispersion surfaces of H^ε , among the lowest four, that touch at the vertices of \mathcal{B} . In a neighborhood of each vertex, the local character given in (1.4) or (1.5).

3. Theorem 6: There exists a discrete set $\tilde{\mathcal{C}} \subset \mathbb{R}$ such that if $\varepsilon \notin \tilde{\mathcal{C}}$, then two dispersion surfaces of H^ε touch at the vertices of \mathcal{B} with local behavior described by (1.4). The constants $\alpha^\varepsilon \in \mathbb{R}$, $\beta^\varepsilon \in \mathbb{C}$, and $\gamma^\varepsilon \in \mathbb{C}$ in Theorem 5.3, and α^ε and β^ε in Theorem 6, displayed in (4.27), depend on the degenerate eigenspace, $\text{span}\{\Phi_1^\varepsilon, \Phi_2^\varepsilon\}$, for quasi momentum. Hence, the property of quadratically touching dispersion surfaces with local behavior given by (1.5) holds for generic, even arbitrarily large, values of ε .

4. *Lieb lattice potentials: Tight binding versus strong binding:* The well-known conical plus flat-band structure dispersion surfaces of the three-band tight-binding model for the Lieb lattice *does not* persist in the Schrödinger operator with periodic Lieb lattice potential consisting of potential wells centered on the Lieb lattice; see section 1.3.
5. *Numerical studies:* In section 7 we discuss numerical simulations for various admissible potentials for a full range of coupling parameters, ε small to ε large. These include potentials which are superpositions of spatially localized potentials, centered on vertices of the square lattice or the vertices of a Lieb lattice. Our simulations corroborate our analytical results and are discussed in this context.
6. *Wave-packet dynamics; Appendix B:* A multiscale analysis demonstrates that the envelope of the solution of the time-dependent Schrödinger equation, $i\partial_t\psi(\mathbf{x}, t) = (-\Delta_{\mathbf{x}} + V(\mathbf{x}))\psi(\mathbf{x}, t)$, for wave-packet initial data: $\psi(\mathbf{x}, 0) = C_{10}(\mathbf{X})\Phi_1(\mathbf{x}) + C_{20}(\mathbf{X})\Phi_2(\mathbf{x})$, where $\mathbf{X} \equiv \delta\mathbf{x} = (X_1, X_2)$, and $C_{j0}(\mathbf{X})$, $j = 1, 2$, are in Schwartz class, evolves on large but finite time scales according to a coupled system of Schrödinger equations ($T = \delta^2 t$):

$$(1.6) \quad i\frac{\partial}{\partial T}C_p(\mathbf{X}, T) = -\sum_{q=1}^2\sum_{r,s=1}^2\frac{\partial}{\partial X_r}\Upsilon_{r,s}^{p,q}\frac{\partial}{\partial X_s}C_q(\mathbf{X}, T), \quad p = 1, 2.$$

Here, $\Upsilon_{r,s}^{p,q}$ are the matrix elements of an (indefinite) inverse effective mass tensor; see section 4.1. The branches of the dispersion relation of (1.6) are given by the expression in (1.4).

A derivation of (1.6) is presented in Appendix B. A rigorous proof of the long (finite) time validity would be along the lines of [1] or [10], for example.

1.3. The Lieb lattice, tight-binding versus continuum models, a band structure instability. Our original motivation for the present study was to contrast the band structure of the Lieb lattice tight-binding Hamiltonian (see, for example, Niță, Ostahie, and Aldea [19], Weeks and Franz [22]) with that of the corresponding continuum Schrödinger operator with Lieb lattice potential in the high-contrast (strong binding) regime, as studied via simulation in Guzmán-Silva et al. [12].

Our results demonstrate that the well-known flat-band plus conical structure near the vertices of \mathcal{B} does not persist in the regime of strong binding (finite deep atomic wells at each lattice site) for the continuum Schrödinger operator.

This is in contrast to the case of honeycomb structures, where the Dirac (conical) points of the tight-binding model of Wallace [21, 18] persist in the continuum honeycomb Schrödinger operators [7, 9]. We now explain this in some detail.

The tight-binding model for the Lieb lattice is given by a Hamiltonian acting on wave functions $\psi \in l^2(\mathbb{Z}^2; \mathbb{C}^3)$; for each $(m, n) \in \mathbb{Z}^2$, $\psi_{m,n} \in \mathbb{C}^3$ is a vector of three complex amplitudes assigned to the A, B , and C sites in the (m, n) th cell; see Figure 1.1 (right). This model has $\pi/2$ -rotational invariance about any B site. The detailed setup is presented in Appendix A. The band structure has three band dispersion relations, whose graphs (dispersion surfaces) intersect at the vertices of the Brillouin zone, \mathcal{B} . Figure 1.3 reveals that the three dispersion surfaces which meet at the vertex, \mathbf{M} , of \mathcal{B} consists of a constant energy dispersion surface (a *flat band*, $\mathbf{k} \mapsto E_0^{\text{TB}}(\mathbf{k}) \equiv 0$), and two other dispersion surfaces, $\mathbf{k} \mapsto E_{\pm}^{\text{TB}}(\mathbf{k})$, which meet conically at \mathbf{M} . An analogous picture applies in a neighborhood of each vertex of \mathcal{B} , as seen in the left plot of Figure 1.3. The latter observation follows from symmetry considerations.




FIG. 1.3. Left: Dispersion surfaces of the three-band tight-binding model for the Lieb lattice. The Brillouin zone is $[-\pi, \pi]^2$. Right: A plot of the dispersion surface along the path $\Gamma \rightarrow X \rightarrow M \rightarrow \Gamma$, as described in Figure 1.2. At the vertices of the Brillouin zone, the dispersion relation has the structure of a conical intersection and a flat band. Our analysis shows that this structure does not persist in the continuum Schrödinger operator which limits to the tight-binding model.

It is natural to contrast this behavior with that of the two-band tight-binding model of Wallace (1947), introduced in his pioneering work on graphite [21, 18]. In this model, there are two dispersion surfaces which meet conically at the high-symmetry quasi momenta, located at the vertices of the (hexagonal) Brillouin zone. These conical points are called *Dirac points*.

The tight-binding model gives an approximation to the low-lying spectrum of the continuum Schrödinger operator $-\Delta + \lambda^2 V$, where V is a superposition of identical potential wells, centered at the sites of a honeycomb structure, and λ is sufficiently large. It was proved in [7] for this *strong-binding regime* that Dirac points occur at the vertices of the Brillouin zone and that, after a rescaling, the first two dispersion surfaces converge uniformly to those of Wallace's two-band tight-binding model. It had earlier been proved in [9, 8] that generic Schrödinger operators, for the class of honeycomb lattice potentials, have Dirac points within their band structure and that these Dirac points persist against small perturbations of the potential which break inversion or complex-conjugate symmetries; see Definition 2.3(ii) and (iii).

QUESTION 1.1. Consider a potential $V_L(\mathbf{x})$, formed as a superposition of identical deep potential wells centered at the points of the Lieb lattice. Does the local band structure near the vertices of \mathcal{B} (a flat band plus two conically touching surfaces) of the tight-binding model persist in the band structure of $-\Delta + V_L$, i.e., does this local structure persist into the strong binding regime?

The answer is *no*, and the precise character of the local band structure is a consequence of our analysis of $\pi/2$ -rotationally invariant potentials. Figure 1.4 displays the family of three curves obtained by sampling three dispersion surfaces of $-\Delta + V_L$, two surfaces that touch at the vertices of \mathcal{B} and the nearest to these among all others, for a choice of deep atomic potential wells. The two curves which intersect are locally described by (1.4).

Remark 1.2. In [1, section 6] results on homogenized effective equations were obtained for the dynamics of wave packets, which are spectrally localized near an isolated (quadratic) spectral band edge. The authors also consider the cases of both simple and degenerate eigenvalues occurring at a band edge, derive a system of coupled Schrödinger envelope equations, and remark on the nongenericity of degenerate




FIG. 1.4. Dispersion surfaces for $-\Delta + V_L$ sampled along two different quasi-momentum segments through the high symmetry quasi momentum, \mathbf{M} . Here, V_L is the periodic potential whose restriction to the primitive cell is $\tilde{V}_L(\mathbf{x}) = -V_0(e^{-|\mathbf{x}|^2/\sigma} + e^{-|\mathbf{x}-(1/2,0)|^2/\sigma} + e^{-|\mathbf{x}-(0,1/2)|^2/\sigma})$. Segments are of the form $\mathbf{k} = \mathbf{M} \pm \lambda(\cos \theta, \sin \theta)^t$ for $\lambda \in (-\lambda_0, \lambda_0)$ with $\theta = \frac{\pi}{4}$ (left) and $\theta = \frac{15\pi}{16}$ (right).

eigenvalues at a band edge. Although nongeneric in the space of all potentials, the results of this article show in the presence of symmetry, for example, the symmetries of our $\mathcal{P} \circ \mathcal{C}$ and $\pi/2$ -rotationally invariant (admissible) potentials, such systems are realized in many physical settings of interest.

1.4. Outline of the paper. In section 2, we define the class of *admissible potentials* considered here, characterize the Fourier series of such potentials, and develop tools for Fourier analysis in the spaces, $L^2_{\mathbf{k}}$ for $\mathbf{k} \in \mathcal{B}$. In particular, $L^2_{\mathbf{M}}$ has direct sum (orthogonal) decomposition into eigenspaces, $L^2_{\mathbf{M},\sigma}$ ($\sigma = \pm 1, \pm i$), of the $\pi/2$ -rotation operator, \mathcal{R} , which acts unitarily on $L^2_{\mathbf{M}}$.

In section 3 we study spectral properties of the free Laplacian, $-\Delta$, in $L^2_{\mathbf{M}}$. In particular, we show that $-\Delta$ has fourfold degenerate $L^2_{\mathbf{M}}$ eigenvalues (we consider the least energy such), which correspond to four simple eigenvalues, one in each of the invariant subspaces, $L^2_{\mathbf{M},\sigma}$.

In section 4 we state and prove Theorem 4.1 and Corollary 4.2, which give sufficient conditions for the quadratic touching of two dispersion surfaces at the high-symmetry quasi momenta, situated at the vertices, \mathbf{M}_* , of \mathcal{B} . These results also display the detailed expansion of the dispersion maps in a neighborhood of these quasi momenta.

In section 5 we verify the hypotheses of Theorems 4.1 and Corollary 4.2 for Hamiltonians of the form $H^\varepsilon = -\Delta + \varepsilon V$, where V is admissible, for all ε real, sufficiently

small and nonzero. This then proves the existence of quadratic degeneracies at high-symmetry quasi momenta as described in Theorem 4.1 and Corollary 4.2 for the regime of sufficiently weak coupling (low-contrast).

In section 6, we consider $H^\varepsilon = -\Delta + \varepsilon V$, where ε is real and nonzero, but not restricted by size. Theorem 6.1 states that the small ε -results persist for all nonzero real ε except for a possible discrete set of ε -values. The proof, based on arguments in [9, 8], is sketched.

Section 7 summarizes a range of computational experiments that corroborate our analytical results.

Appendix A summarizes the formulation of the tight binding for the Lieb lattice; see section 1.3.

Appendix B summarizes a multiple scales asymptotic expansion yielding a coupled system of time-dependent Schrödinger equations, which govern the evolution of wave-packets, which are spectrally concentrated in a neighborhood of high-symmetry quasi momenta, $\mathbf{M}_\star \in \mathcal{B}$.

Finally, in Appendix C we provide the details of a calculation of coefficients occurring in the leading order expression for the touching dispersion surfaces in the weak coupling (ε nonzero and small) regime; see also Corollary 5.4.

1.5. Notation. We remark on some frequently used notation and conventions.

1. Repeated indices are understood to be summed unless otherwise specified.
2. For $\mathbf{m} \in \mathbb{Z}^2$, $\mathbf{m}\vec{\mathbf{k}} = m_1\mathbf{k}_1 + m_2\mathbf{k}_2$, where \mathbf{k}_1 and \mathbf{k}_2 are the dual lattice basis vectors defined below in (2.2).
3. $\langle f, g \rangle_{L^2(\mathbb{R}^2/\Lambda)} = \langle f, g \rangle$; if not indicated the inner product is understood to be taken in $L^2(\mathbb{R}^2/\Lambda)$.

2. The lattice Λ and admissible potentials.

2.1. The square lattice. We begin with the lattice $\Lambda = \mathbb{Z}\mathbf{v}_1 \oplus \mathbb{Z}\mathbf{v}_2$, where

$$(2.1) \quad \mathbf{v}_1 = a \begin{pmatrix} 1 \\ 0 \end{pmatrix} \text{ and } \mathbf{v}_2 = a \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

The constant, a , is the lattice constant giving the distance between nearest neighbor sites. The dual lattice, Λ^* , is generated by the vectors

$$(2.2) \quad \mathbf{k}_1 = a^{-1} \begin{pmatrix} 2\pi \\ 0 \end{pmatrix} \text{ and } \mathbf{k}_2 = a^{-1} \begin{pmatrix} 0 \\ 2\pi \end{pmatrix}.$$

For simplicity, we assume $a = 1$. The Brillouin zone, \mathcal{B} , is given in (1.2); see Figure 1.2. We have the relations $\mathbf{k}_l \cdot \mathbf{v}_m = 2\pi\delta_{l,m}$ for $l, m = 1, 2$.

Let R be the $\pi/2$ -clockwise rotation matrix

$$(2.3) \quad R = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

We record the elementary relations: $R^*\mathbf{v}_1 = \mathbf{v}_2$ and $R^*\mathbf{v}_2 = -\mathbf{v}_1$, and $R\mathbf{k}_1 = -\mathbf{k}_2$ and $R\mathbf{k}_2 = \mathbf{k}_1$. It follows that R and R^* map Λ to itself and Λ^* to itself. Denote the vertices of \mathcal{B} by

$$\mathbf{M} \equiv \mathbf{M}_{++} = (\pi, \pi)^T, \quad \mathbf{M}_{+-} = (\pi, -\pi)^T, \quad \mathbf{M}_{--} = (-\pi, -\pi)^T, \quad \mathbf{M}_{-+} = (-\pi, \pi)^T.$$

Then, the set of vertices of \mathcal{B} is mapped by R to itself:

$$\mathbf{M}_{+-} = R\mathbf{M} = \mathbf{M} - \mathbf{k}_2, \quad \mathbf{M}_{--} = R^2\mathbf{M} = \mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2, \quad \mathbf{M}_{-+} = R^3\mathbf{M} = \mathbf{M} - \mathbf{k}_1.$$

Furthermore, R maps the affine sublattice $\mathbf{M} + \Lambda^*$ one to one and onto itself. See Figure 1.2.

Remark 2.1. Note that the pseudoperiodic boundary condition associated with quasi momenta located at the vertices of \mathcal{B} are the same and correspond to \mathbf{M} -pseudoperiodicity, i.e., for any choice of $(a, b) \in \{+, -\}$, we have $\psi(\mathbf{x} + \mathbf{v}; \mathbf{M}_{ab}) = e^{i\mathbf{M} \cdot \mathbf{v}} \psi(\mathbf{x}; \mathbf{M}_{ab})$, $\mathbf{x} \in \mathbb{R}$, $\mathbf{v} \in \Lambda$. Therefore, for any $\mathbf{M}_* \in \mathbf{M} + \Lambda$, the space $L^2_{\mathbf{M}_*}$ can be identified with $L^2_{\mathbf{M}}$. Furthermore, the local character of dispersion surfaces in a neighborhood of any vertex of \mathcal{B} determines the local character in a neighborhood of any other vertex of \mathcal{B} .

2.2. Admissible potentials. For any function f defined on \mathbb{R}^2 , we define the $\pi/2$ -rotational operator

$$\mathcal{R}[f](\mathbf{x}) \equiv f(R^*\mathbf{x}),$$

where R is the $\pi/2$ -clockwise rotation matrix displayed in (2.5).

We consider smooth (say, C^∞) periodic potentials $V(\mathbf{x}) = V(x_1, x_2)$ defined on \mathbb{R}^2 , with fundamental period cell $\Omega = [0, 1] \times [0, 1]$. Any such V can be represented as a Fourier series:

$$(2.4) \quad V(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^2} V_{\mathbf{m}} e^{i\mathbf{m}\vec{\mathbf{k}} \cdot \mathbf{x}} = \sum_{(m_1, m_2) \in \mathbb{Z}^2} V_{m_1 m_2} e^{2\pi i(m_1 x_1 + m_2 x_2)},$$

where $V_{\mathbf{m}} = (2\pi)^{-2} \int_{\Omega} e^{-i\mathbf{m}\vec{\mathbf{k}} \cdot \mathbf{x}} V(\mathbf{x}) d\mathbf{x}$, and $\mathbf{m}\vec{\mathbf{k}} = m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2$.

DEFINITION 2.2 (\mathcal{P} , \mathcal{C} , and \mathcal{R} invariance).

1. Given a point $\mathbf{x} \in \mathbb{R}^2$, its $\pi/2$ -counterclockwise rotation about \mathbf{x}_c , denoted $\widehat{\mathbf{x}}_{\mathcal{R}}$, satisfies $\widehat{\mathbf{x}}_{\mathcal{R}} - \mathbf{x}_c = R^*(\mathbf{x} - \mathbf{x}_c)$. If

$$(2.5) \quad \mathcal{R}[V](\mathbf{x}) \equiv V(\widehat{\mathbf{x}}_{\mathcal{R}}) = V(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{R}^2$ we say that $V(\mathbf{x})$ is \mathcal{R} -invariant, or $\pi/2$ -rotationally invariant, with respect to \mathbf{x}_c .

2. Given a point $\mathbf{x} \in \mathbb{R}^2$, its inversion with respect to \mathbf{x}_c , denoted $\widehat{\mathbf{x}}_{\mathcal{I}}$, satisfies $\widehat{\mathbf{x}}_{\mathcal{I}} - \mathbf{x}_c = -(\mathbf{x} - \mathbf{x}_c)$. If $V(\widehat{\mathbf{x}}_{\mathcal{I}}) = V(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^2$ we say that $V(\mathbf{x})$ is \mathcal{P} -invariant, parity invariant, or inversion symmetric with respect to \mathbf{x}_c .
3. We say that $V(\mathbf{x})$ is \mathcal{C} -invariant or invariant under complex conjugation if $\overline{V(\mathbf{x})} = V(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^2$.

We shall study potentials which are real-valued, smooth, and invariant under $\mathcal{P} \circ \mathcal{C}$ and \mathcal{R} -invariant (invariant under $\pi/2$ rotation). We call such potentials *admissible*.

DEFINITION 2.3 (admissible potentials). An admissible potential is a smooth function, $V(\mathbf{x})$, defined on \mathbb{R}^2 with the following properties:

- (i) Λ -periodicity: $V(\mathbf{x} + \mathbf{v}) = V(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^2$ and $\mathbf{v} \in \Lambda$.

There exists $\mathbf{x}_c \in \mathbb{R}^2$ with respect to which (in the sense of Definition 2.2)

- (ii) V is \mathcal{C} -invariant;
- (iii) V is \mathcal{P} -invariant; and
- (iv) V is $\pi/2$ -rotationally invariant.

Throughout this paper we shall assume, with no loss of generality, $\mathbf{x}_c = 0$.

In some of our results we consider admissible potentials which are also reflection invariant. Such potentials have the full symmetry of the square lattice.

DEFINITION 2.4 (reflection invariance). *An admissible potential, $V(\mathbf{x})$, defined on \mathbb{R}^2 is reflection invariant if $V(x_1, x_2) = V(x_2, x_1)$.*

We introduce two basic examples of admissible potentials, each obtained as a sum of translates of a fixed atomic potential:

Example 2.5 (square lattice potential). Let $V(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^2} V_0(\mathbf{x} + \mathbf{m})$, where $V_0 = V_0(|\mathbf{x}|)$ is a real-valued, radially symmetric, sufficiently rapidly decaying “atomic potential.” We call $V(\mathbf{x})$ a *square lattice potential*. It is easily seen that this class of potentials is admissible with $\mathbf{x}_c = 0$.

Example 2.6 (Lieb lattice potential). We fix the fundamental period cell to be the square with side-length one. Within a fixed cell are three points, labeled \mathbf{A} , \mathbf{B} , and \mathbf{C} ; see Figure 1.1 (right). The Lieb lattice, \mathbb{L} , is the union of three sublattices: $\mathbf{A} + \mathbb{Z}^2$, $\mathbf{B} + \mathbb{Z}^2$, and $\mathbf{C} + \mathbb{Z}^2$. A *Lieb lattice potential* is given by $V(\mathbf{x}) = \sum_{\mathbf{w} \in \mathbb{L}} V_0(\mathbf{x} + \mathbf{w})$, where $V_0 = V_0(|\mathbf{x}|)$ is a real-valued, radially symmetric, and rapidly decaying atomic potential. Lieb lattice potentials are admissible with $\mathbf{x}_c = 0$. An example of an atomic Lieb lattice potential is displayed in Figure 7.4.

The next proposition states that if V is an admissible potential in the sense of Definition 2.3, then the operator H_V acting in $L^2_{\mathbf{k}}$ has an additional symmetry for $\mathbf{k} \in \mathbf{M} + \Lambda^*$.

PROPOSITION 2.7. *Assume V is an admissible potential in the sense of Definition 2.3. Then $H_V = -\Delta + V$ and \mathcal{R} map a dense subspace of $L^2_{\mathbf{M}}$ to itself. Moreover, restricted to this dense subspace of $L^2_{\mathbf{M}}$, the commutator $[H_V, \mathcal{R}] \equiv H_V \mathcal{R} - \mathcal{R} H_V = 0$. In particular, if $\Phi(\mathbf{x})$ is a solution of the \mathbf{M} -pseudoperiodic eigenvalue problem for H_V for energy E , then $\mathcal{R}[\Phi](\mathbf{x})$ is also a solution of the \mathbf{M} -pseudoperiodic eigenvalue problem for H_V with energy E .*

Proof. Note first that $-\Delta$ commutes with rotations and the operator $\Phi \mapsto V\Phi$, where V is an admissible potential, commutes with $\pi/2$ rotations. Furthermore, assume $\Phi(\mathbf{x})$ is \mathbf{M} -pseudoperiodic and define $\Phi_R(\mathbf{x}) \equiv \Phi(R^*\mathbf{x})$. Then, for all $\mathbf{v} \in \Lambda = \mathbb{Z}^2$ and all $\mathbf{x} \in \mathbb{R}^2$: $\Phi_R(\mathbf{x} + \mathbf{v}) = \Phi(R^*\mathbf{x} + R^*\mathbf{v}) = e^{i\mathbf{M} \cdot R^*\mathbf{v}} \Phi(R^*\mathbf{x}) = e^{iR\mathbf{M} \cdot \mathbf{v}} \Phi(R^*\mathbf{x}) = e^{i(\mathbf{M} - \mathbf{k}_2) \cdot \mathbf{v}} \Phi(R^*\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{v}} \Phi_R(\mathbf{x})$. For more detail, see an analogous result in [9]. \square

2.3. Fourier series of admissible potentials. The following proposition implies constraints on the Fourier coefficients of admissible potentials.

PROPOSITION 2.8. *Let $V(\mathbf{x})$ be in $C^\infty(\mathbb{R}^2/\Lambda)$. Then, for all $\mathbf{m} = (m_1, m_2) \in \mathbb{Z}^2$*

- (i) $V(-\mathbf{x}) = V(\mathbf{x}) \implies V_{\mathbf{m}} = V_{-\mathbf{m}}$,
- (ii) $\overline{V(\mathbf{x})} = V(\mathbf{x}) \implies V_{\mathbf{m}} = \overline{V_{-\mathbf{m}}}$,
- (iii) $\mathcal{R}[V](\mathbf{x}) = V(\mathbf{x}) \implies V_{m_1, m_2} = V_{-m_2, m_1}$.

Proof. Parts (i) and (ii) are straightforward. Part (iii) makes use of the action of R on the dual lattice basis $\{\mathbf{k}_1, \mathbf{k}_2\}$ or equivalently R^* on the lattice basis $\{\mathbf{v}_1, \mathbf{v}_2\}$. \square

Note that we may iterate the relation in part (iii) of Proposition 2.8 to obtain

$$(2.6) \quad V_{m_1, m_2} = V_{-m_2, m_1} = V_{-m_1, -m_2} = V_{m_2, -m_1}.$$

Introduce the mapping $\tilde{\mathcal{R}} : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ defined $\tilde{\mathcal{R}}(m_1, m_2) = (-m_2, m_1)$ and therefore $\tilde{\mathcal{R}}^2(m_1, m_2) = (-m_1, -m_2)$, $\tilde{\mathcal{R}}^3(m_1, m_2) = (m_2, -m_1)$, and $\tilde{\mathcal{R}}^4(m_1, m_2) = (m_1, m_2)$. Thus, $\tilde{\mathcal{R}}^4 = \tilde{\mathcal{R}}^0 = Id$, and

$$(2.7) \quad V_{\mathbf{m}} = V_{\tilde{\mathcal{R}}\mathbf{m}} = V_{\tilde{\mathcal{R}}^2\mathbf{m}} = V_{\tilde{\mathcal{R}}^3\mathbf{m}}.$$

Note that $\mathbf{0}$ is the unique element of the kernel (and fixed point) of $\tilde{\mathcal{R}}$ and furthermore every $\mathbf{m} \neq \mathbf{0}$ lies on a nontrivial 4-cycle of $\tilde{\mathcal{R}}$, the set $\{(m_1, m_2), (-m_2, m_1), (-m_1, -m_2), (m_2, -m_1)\}$. Let \mathbf{m} and \mathbf{n} be elements of $\mathbb{Z}^2 \setminus \{\mathbf{0}\}$. We say that $\mathbf{m} \sim \mathbf{n}$ if \mathbf{m} and \mathbf{n} lie on the same 4-cycle of $\tilde{\mathcal{R}}$. The relation \sim is an equivalence relation and partitions $\mathbb{Z}^2 \setminus \{\mathbf{0}\}$ into equivalence classes, $(\mathbb{Z}^2 \setminus \{\mathbf{0}\}) / \sim$. Let $\tilde{\mathcal{S}}$ denote a set consisting of one representative element from each equivalence class.

PROPOSITION 2.9. (a) Let V denote an admissible potential in the sense of Definition 2.3 and let $V_{\mathbf{m}} = V_{m_1, m_2}$, for $\mathbf{m} \in \mathbb{Z}^2$, denote its Fourier coefficients; see (2.4). Then,

$$(2.8) \quad V(x_1, x_2) = V_{0,0} + \sum_{(m_1, m_2) \in \tilde{\mathcal{S}}} 2V_{m_1, m_2} [\cos(2\pi(m_1 x_1 + m_2 x_2)) + \cos(2\pi(m_2 x_1 - m_1 x_2))].$$

(b) If V is also reflection invariant ($V(x_1, x_2) = V(x_2, x_1)$), then

$$(2.9) \quad V(x_1, x_2) = V_{0,0} + \sum_{m \in \mathbb{Z}} 2V_{m,m} [\cos(2\pi m(x_1 + x_2)) + \cos(2\pi m(x_1 - x_2))].$$

Proof. Expanding $V(\mathbf{x})$ in a Fourier series, and using the relations in (2.7), we obtain

$$V(\mathbf{x}) = V_0 + \sum_{\mathbf{m} \in \tilde{\mathcal{S}}} V_{\mathbf{m}} \left(e^{i\mathbf{m}\vec{k} \cdot \mathbf{x}} + e^{i(\tilde{\mathcal{R}}\mathbf{m})\vec{k} \cdot \mathbf{x}} + e^{i(\tilde{\mathcal{R}}^2\mathbf{m})\vec{k} \cdot \mathbf{x}} + e^{i(\tilde{\mathcal{R}}^3\mathbf{m})\vec{k} \cdot \mathbf{x}} \right).$$

Adding this expression to its complex conjugate and dividing by two and using that the coefficients $V_{\mathbf{m}}$ are real (Proposition 2.8(i) and (ii)) implies

$$V(\mathbf{x}) = V_0 + \sum_{\mathbf{m} \in \tilde{\mathcal{S}}} V_{\mathbf{m}} \left(\cos(\mathbf{m}\vec{k} \cdot \mathbf{x}) + \cos((\tilde{\mathcal{R}}\mathbf{m})\vec{k} \cdot \mathbf{x}) + \cos((\tilde{\mathcal{R}}^2\mathbf{m})\vec{k} \cdot \mathbf{x}) + \cos((\tilde{\mathcal{R}}^3\mathbf{m})\vec{k} \cdot \mathbf{x}) \right),$$

which reduces to the expression in (2.8). Thus part (2.9) is proved.

Suppose V additionally is reflection invariant in the sense of Definition 2.4. By (2.8) this is equivalent to

$$\begin{aligned} & \sum_{(m_1, m_2) \in \tilde{\mathcal{S}}} 2V_{m_1, m_2} [\cos(2\pi(m_2 x_1 + m_1 x_2)) + \cos(2\pi(m_2 x_2 - m_1 x_1))] \\ &= \sum_{(m_1, m_2) \in \tilde{\mathcal{S}}} 2V_{m_1, m_2} [\cos(2\pi(m_1 x_1 + m_2 x_2)) + \cos(2\pi(m_2 x_1 - m_1 x_2))]. \end{aligned}$$

It follows that for all x_1, x_2 ,

$$(2.10) \quad \cos[2\pi(m_2 x_1 + m_1 x_2)] + \cos[2\pi(m_2 x_2 - m_1 x_1)] - \cos[2\pi(m_1 x_1 + m_2 x_2)] - \cos[2\pi(m_2 x_1 - m_1 x_2)] = 0.$$

Using trigonometric identities, (2.10) reduces to

$$f_1(x_1, x_2) \equiv \sin(2\pi m_2 x_1) \sin(2\pi m_1 x_2) = \sin(2\pi m_1 x_1) \sin(2\pi m_2 x_2) \equiv f_2(x_1, x_2).$$

This implies that the Fourier coefficients of f_1 and f_2 match and therefore $m_1 = m_2$. \square

2.4. Fourier analysis in $L^2_{\mathbf{M}^*}$. In this subsection, we characterize the Fourier series of functions $\phi \in L^2_{\mathbf{M}}$. Such functions may be expressed in the form $\Phi(\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} \phi(\mathbf{x})$, where $\phi(\mathbf{x})$ is $\Lambda = \mathbb{Z}^2$ -periodic. Thus, Φ has the Fourier representation

$$(2.11) \quad \Phi(\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} \sum_{(m_1, m_2) \in \mathbb{Z}^2} c(m_1, m_2) e^{i(m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2) \cdot \mathbf{x}},$$

which we rewrite as

$$(2.12) \quad \Phi(\mathbf{x}) = \sum_{(m_1, m_2) \in \mathbb{Z}^2} c_{\Phi}(m_1, m_2) e^{i(\mathbf{M} + m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2) \cdot \mathbf{x}} = \sum_{\mathbf{m} \in \mathbb{Z}^2} c_{\Phi}(\mathbf{m}) e^{i\mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}},$$

where $\mathbf{M}^{\mathbf{m}} = \mathbf{M} + \mathbf{m}\vec{\mathbf{k}} = \mathbf{M} + m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2 \in \mathbf{M} + \Lambda^*$. We denote the Fourier coefficients of a specific $\Phi \in L^2_{\mathbf{M}}$ shown in (2.12) as $c_{\Phi}(\mathbf{m})$ or $c(\mathbf{m}; \Phi)$.

Next, observe that the transformation $\mathcal{R}: f \mapsto \mathcal{R}[f](\mathbf{x}) = f(R^* \mathbf{x})$ is unitary on $L^2_{\mathbf{k}}$ and so its eigenvalues lie on the unit circle in \mathbb{C} . Furthermore, if $\mathcal{R}\Phi = \sigma\Phi$ and $\Phi \neq 0$, then since $\mathcal{R}^4 = Id$, we have that $\Phi = \mathcal{R}^4\Phi = \sigma^4\Phi$, so that $\sigma^4 = 1$. Therefore, $\sigma \in \{+1, -1, +i, -i\}$.

This induces a decomposition of $L^2_{\mathbf{M}}$ as an orthogonal sum of eigenspaces of \mathcal{R} :

$$(2.13) \quad L^2_{\mathbf{M}} = L^2_{\mathbf{M},1} \oplus L^2_{\mathbf{M},-1} \oplus L^2_{\mathbf{M},i} \oplus L^2_{\mathbf{M},-i},$$

where

$$(2.14) \quad L^2_{\mathbf{M},\sigma} = \{ f \in L^2_{\mathbf{M}} : \mathcal{R}[f] = \sigma f \}, \quad \sigma \in \{1, -1, i, -i\}.$$

Remark 2.10. The spectral theory of H in $L^2_{\mathbf{M}}$ can be reduced to its independent study in each summand subspace in the orthogonal sum (2.13).

Our next goal is to characterize Fourier series of functions in the orthogonal summands $L^2_{\mathbf{M},\sigma}$ for $\sigma = \pm 1, \pm i$. We first apply \mathcal{R} to Φ , represented as a Fourier series in (2.12). Note that

$$\begin{aligned} R\mathbf{M}^{\mathbf{m}} &= R(\mathbf{M} + \mathbf{m}\vec{\mathbf{k}}) = R\mathbf{M} + R(m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2) \\ &= \mathbf{M} + m_2 \mathbf{k}_1 + (-1 - m_1) \mathbf{k}_2 = \mathbf{M}^{m_2, -1-m_1} \end{aligned}$$

and define (taking some liberty with notation)

$$\mathcal{R}\mathbf{m} \equiv \mathcal{R}(m_1, m_2) = (m_2, -1 - m_1)$$

and hence $\mathcal{R}^{-1}\mathbf{m} = \mathcal{R}^{-1}(m_1, m_2) = (-m_2 - 1, m_1)$. The mapping \mathcal{R} acting on $L^2_{\mathbf{M}}$ induces a decomposition of \mathbb{Z}^2 into orbits of minimal length four,

$$(2.15) \quad \begin{aligned} (m_1, m_2)^{\mathcal{R}} &\mapsto (m_2, -1 - m_1)^{\mathcal{R}} \mapsto (-1 - m_1, -1 - m_2)^{\mathcal{R}} \\ &\mapsto (-1 - m_2, m_1)^{\mathcal{R}} \mapsto (m_1, m_2), \end{aligned}$$

and we write $\mathcal{R}^2\mathbf{m} = (-1 - m_1, -1 - m_2)$, $\mathcal{R}^3\mathbf{m} = (-1 - m_2, m_1)$, and $\mathcal{R}^4\mathbf{m} = (m_1, m_2)$.

For $\mathbf{m}, \mathbf{n} \in \mathbb{Z}^2$ we write $\mathbf{m} \approx \mathbf{n}$ if \mathbf{m} and \mathbf{n} lie on the same orbit under \mathcal{R} . We denote by \mathcal{S} any set containing exactly one representative from each equivalence class in \mathbb{Z}^2 / \approx .

Remark 2.11. One such equivalence class is $\{(0, 0), (0, -1), (-1, -1), (-1, 0)\}$ and we choose $(-1, 0)$ as its representative in \mathcal{S} . In section 5 we shall define $\mathcal{S}^\perp = \mathcal{S} \setminus \{(0, -1)\}$ and write $\mathcal{S} \equiv \{(-1, 0)\} \cup \mathcal{S}^\perp$.

In terms of the above notation we have

(2.16)

$$R\mathbf{M} = \mathbf{M}^{m_2, -1-m_1} = \mathbf{M}^{\mathcal{R}\mathbf{m}},$$

$$\mathcal{R}[\Phi](\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} \sum_{(m_1, m_2) \in \mathbb{Z}^2} c_\Phi(m_1, m_2) e^{i(m_2 \mathbf{k}_1 + (-1-m_1) \mathbf{k}_2) \cdot \mathbf{x}} = \sum_{\mathbf{m} \in \mathbb{Z}^2} c_\Phi(\mathbf{m}) e^{i\mathbf{M}^{\mathcal{R}\mathbf{m}} \cdot \mathbf{x}}.$$

Therefore, $c_{\mathcal{R}\Phi}(\mathcal{R}\mathbf{m}) = c_\Phi(\mathbf{m})$. Note that

$$(2.17) \quad \mathcal{R}^{-1}\mathbf{m} = \mathcal{R}^3\mathbf{m} = (-m_2 - 1, m_1),$$

$$(2.18) \quad \mathcal{R}^{-2}\mathbf{m} = \mathcal{R}^2\mathbf{m} = (-1 - m_1, -1 - m_2), \text{ and}$$

$$(2.19) \quad \mathcal{R}^{-3}\mathbf{m} = \mathcal{R}\mathbf{m} = (m_2, -m_1 - 1).$$

Hence,

$$(2.20) \quad c_{\mathcal{R}^j\Phi}(\mathbf{m}) = c_\Phi(\mathcal{R}^{4-j}\mathbf{m}), \quad j = 0, 1, 2, 3.$$

The Fourier series of $\Phi \in L^2_{\mathbf{k}}$, satisfying the pseudoperiodic boundary conditions, may be expressed as a sum over 4-cycles of \mathcal{R} :

(2.21)

$$\phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c_\Phi(\mathbf{m}) e^{i\mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} + c_\Phi(\mathcal{R}\mathbf{m}) e^{i\mathbf{M}^{\mathcal{R}\mathbf{m}} \cdot \mathbf{x}} + c_\Phi(\mathcal{R}^2\mathbf{m}) e^{i\mathbf{M}^{\mathcal{R}^2\mathbf{m}} \cdot \mathbf{x}} + c_\Phi(\mathcal{R}^3\mathbf{m}) e^{i\mathbf{M}^{\mathcal{R}^3\mathbf{m}} \cdot \mathbf{x}}.$$

We next study the Fourier representation (2.21) in the case where $\Phi \in L^2_{\mathbf{M}, \sigma}$ for $\sigma = \pm 1, \pm i$.

PROPOSITION 2.12. *Let $\Phi \in L^2_{\mathbf{M}}$. Then,*

$$\Phi \in L^2_{\mathbf{M}, \sigma} \iff c_\Phi(\mathcal{R}^j\mathbf{m}) = \sigma^{4-j} c_\Phi(\mathbf{m}), \quad j = 0, 1, 2, 3.$$

In particular,

$$\mathcal{R}\Phi = \Phi \iff c(\mathbf{m}) = c(\mathcal{R}\mathbf{m}) = c(\mathcal{R}^2\mathbf{m}) = c(\mathcal{R}^3\mathbf{m}),$$

$$\mathcal{R}\Phi = -\Phi \iff c(\mathcal{R}\mathbf{m}) = -c(\mathbf{m}), \quad c(\mathcal{R}^2\mathbf{m}) = c(\mathbf{m}), \quad c(\mathcal{R}^3\mathbf{m}) = -c(\mathbf{m}),$$

$$\mathcal{R}\Phi = i\Phi \iff c(\mathcal{R}\mathbf{m}) = -ic(\mathbf{m}), \quad c(\mathcal{R}^2\mathbf{m}) = -c(\mathbf{m}), \quad c(\mathcal{R}^3\mathbf{m}) = +ic(\mathbf{m}),$$

$$\mathcal{R}\Phi = -i\Phi \iff c(\mathcal{R}\mathbf{m}) = +ic(\mathbf{m}), \quad c(\mathcal{R}^2\mathbf{m}) = -c(\mathbf{m}), \quad c(\mathcal{R}^3\mathbf{m}) = -ic(\mathbf{m}).$$

Proof. Suppose $\Phi \in L^2_{\mathbf{M}, \sigma}$ and $\mathcal{R}\Phi = \sigma\Phi$. Then, $\mathcal{R}^2\Phi = \sigma^2\Phi$ and $\mathcal{R}^3\Phi = \sigma^3\Phi$. Correspondingly, $c_{\mathcal{R}^j\Phi}(\mathbf{m}) = \sigma^j c_\Phi(\mathbf{m})$ for $j = 0, 1, 2, 3$. By relations (2.20) we have $c_\Phi(\mathcal{R}^{4-j}\mathbf{m}) = c_{\mathcal{R}^j\Phi}(\mathbf{m}) = \sigma^j c_\Phi(\mathbf{m})$ for $j = 0, 1, 2, 3$. Replacing j by $4-j$ completes the proof. \square

Applying Proposition 2.12 to (2.21) we obtain the following.

PROPOSITION 2.13. *Let $\sigma \in \{+1, -1, +i, -i\}$. Then, $\Phi \in L^2_{\mathbf{M}, \sigma}$ if and only if there exists $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}}$ in $l^2(\mathcal{S})$ such that*

$$(2.22) \quad \Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c(\mathbf{m}) \sum_{j=0}^3 \sigma^{4-j} e^{i\mathbf{R}^j \mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}},$$

In detail,

1. $\Phi \in L^2_{\mathbf{M},i} \iff$ there exists $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}} \in l^2(\mathcal{S})$ such that

$$(2.23) \quad \Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c(\mathbf{m}) \left(e^{i\mathbf{M}^m \cdot \mathbf{x}} - ie^{iR\mathbf{M}^m \cdot \mathbf{x}} - e^{iR^2\mathbf{M}^m \cdot \mathbf{x}} + ie^{iR^3\mathbf{M}^m \cdot \mathbf{x}} \right),$$

2. $\Phi \in L^2_{\mathbf{M},-i} \iff$ there exists $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}} \in l^2(\mathcal{S})$ such that

$$(2.24) \quad \Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c(\mathbf{m}) \left(e^{i\mathbf{M}^m \cdot \mathbf{x}} + ie^{iR\mathbf{M}^m \cdot \mathbf{x}} - e^{iR^2\mathbf{M}^m \cdot \mathbf{x}} - ie^{iR^3\mathbf{M}^m \cdot \mathbf{x}} \right),$$

3. $\Phi \in L^2_{\mathbf{M},1} \iff$ there exists $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}} \in l^2(\mathcal{S})$ such that

$$(2.25) \quad \Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c(\mathbf{m}) \left(e^{i\mathbf{M}^m \cdot \mathbf{x}} + e^{iR\mathbf{M}^m \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^m \cdot \mathbf{x}} + e^{iR^3\mathbf{M}^m \cdot \mathbf{x}} \right),$$

4. $\Phi \in L^2_{\mathbf{M},-1} \iff$ there exists $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}} \in l^2(\mathcal{S})$ such that

$$(2.26) \quad \Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c(\mathbf{m}) \left(e^{i\mathbf{M}^m \cdot \mathbf{x}} - e^{iR\mathbf{M}^m \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^m \cdot \mathbf{x}} - e^{iR^3\mathbf{M}^m \cdot \mathbf{x}} \right).$$

Finally, $\mathcal{P} \circ \mathcal{C}$ is a bijection between $L^2_{\mathbf{M},i}$ and $L^2_{\mathbf{M},-i}$. If $c_\Phi(\mathbf{m}), \mathbf{m} \in \mathcal{S}$ are the Fourier coefficients of $\Phi \in L^2_{\mathbf{M},i}$, then $c_{(\mathcal{P} \circ \mathcal{C})\Phi}(\mathbf{m}) = c_\Phi(\mathbf{m})$, $\mathbf{m} \in \mathcal{S}$ are the Fourier coefficients of $(\mathcal{P} \circ \mathcal{C})\Phi \in L^2_{\mathbf{M},-i}$.

3. $H^{(0)} = -\Delta$ on $L^2_{\mathbf{M}}$: A fourfold degenerate eigenvalue. We consider the eigenvalue problem (1.1) for the case $V \equiv 0$. Let $H^{(0)} = -\Delta$.

$$(3.1) \quad H^{(0)}\Phi^{(0)} = \mu^{(0)}(\mathbf{k})\Phi^{(0)}, \quad \Phi^{(0)} \in L^2_{\mathbf{k}}.$$

Equivalently, take $\Phi^{(0)}(\mathbf{x}; \mathbf{k}) = e^{i\mathbf{k} \cdot \mathbf{x}} \phi^{(0)}(\mathbf{x})$, where $\phi^{(0)}(\mathbf{x}) \in L^2(\mathbb{R}^2/\Lambda)$. We have (see (1.3))

$$(3.2) \quad \begin{aligned} H^{(0)}(\mathbf{k})\phi^{(0)} &= -(\nabla + i\mathbf{k})^2 \phi^{(0)} = \mu^{(0)}(\mathbf{k})\phi^{(0)}, \\ \phi^{(0)}(\mathbf{x} + \mathbf{v}) &= \phi^{(0)}(\mathbf{x}), \quad \mathbf{v} \in \Lambda. \end{aligned}$$

For $m_1, m_2 \in \mathbb{Z}$, the eigenvalue problem (3.2) has solutions of the form

$$\phi^{(0)}_{m_1, m_2}(\mathbf{x}; \mathbf{k}) = e^{i(m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2) \cdot \mathbf{x}}, \quad \mathbf{k} \in \mathcal{B},$$

with corresponding eigenvalues

$$\mu^{(0)}_{m_1, m_2}(\mathbf{k}) = |\mathbf{k} + m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2|^2, \quad \mathbf{k} \in \mathcal{B}.$$

The dispersion relation for the free Hamiltonian, $H^{(0)}$, is plotted in Figure 3.1.

The following result concerns the spectral problem for the high-symmetry quasi momentum $\mathbf{k} = \mathbf{M}$ (and by Remark 2.1 all vertices of \mathcal{B}).

THEOREM 3.1. Let $\mathbf{k} = \mathbf{M}$ and $\sigma = \pm 1, \pm i$:

1. $\mu^{(0)}_S \equiv |\mathbf{M}|^2 = 2\pi^2$ is an $L^2_{\mathbf{M}}$ -eigenvalue of multiplicity four with corresponding four-dimensional eigenspace given by

$$(3.3) \quad L^2_{\mathbf{M}} - \text{Kernel} \left(H^{(0)} - \mu^{(0)}_S \text{Id} \right) = \text{span} \left\{ e^{i\mathbf{M} \cdot \mathbf{x}}, e^{iR\mathbf{M} \cdot \mathbf{x}}, e^{iR^2\mathbf{M} \cdot \mathbf{x}}, e^{iR^3\mathbf{M} \cdot \mathbf{x}} \right\}.$$




FIG. 3.1. Dispersion surfaces of $H^{(0)} = -\Delta$. Left: The first five dispersion surfaces are plotted over the Brillouin zone, $[-\pi, \pi]^2$. Each surface is plotted using a different color. (The first is blue, the second is red, etc.) Some level sets of the dispersion surfaces are indicated with black lines. Right: The same dispersion surfaces are plotted along the circuit $\Gamma \rightarrow \mathbf{X} \rightarrow \mathbf{M} \rightarrow \Gamma$, displayed in Figure 1.2. The colors match the plot on the left. As shown in Theorem 3.1, there is a multiplicity four $L^2_{\mathbf{M}}$ -eigenvalue $\mu^{(0)} = |\mathbf{M}|^2 = 2\pi^2$.

2. $H^{(0)}$ acting in $L^2_{\mathbf{M},\sigma}$ has simple eigenvalue $\mu_S^{(0)} = |\mathbf{M}|^2$ with corresponding eigenspace:

$$L^2_{\mathbf{M},\sigma} - \text{Kernel} \left(H^{(0)} - \mu_S^{(0)} \text{Id} \right) = \text{span} \left\{ \Phi_{\sigma}^{(0)} \right\},$$

where $\Phi_{\sigma}^{(0)}$ are defined as follows:

$$(3.4) \quad \Phi_{+1}^{(0)}(\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} + e^{iR\mathbf{M} \cdot \mathbf{x}} + e^{iR^2\mathbf{M} \cdot \mathbf{x}} + e^{iR^3\mathbf{M} \cdot \mathbf{x}} \in L^2_{\mathbf{M},+1},$$

$$(3.5) \quad \Phi_{-1}^{(0)}(\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} - e^{iR\mathbf{M} \cdot \mathbf{x}} + e^{iR^2\mathbf{M} \cdot \mathbf{x}} - e^{iR^3\mathbf{M} \cdot \mathbf{x}} \in L^2_{\mathbf{M},-1},$$

$$(3.6) \quad \Phi_{+i}^{(0)}(\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} - i e^{iR\mathbf{M} \cdot \mathbf{x}} - e^{iR^2\mathbf{M} \cdot \mathbf{x}} + i e^{iR^3\mathbf{M} \cdot \mathbf{x}} \in L^2_{\mathbf{M},+i},$$

$$(3.7) \quad \Phi_{-i}^{(0)}(\mathbf{x}) = e^{i\mathbf{M} \cdot \mathbf{x}} + i e^{iR\mathbf{M} \cdot \mathbf{x}} - e^{iR^2\mathbf{M} \cdot \mathbf{x}} - i e^{iR^3\mathbf{M} \cdot \mathbf{x}} \in L^2_{\mathbf{M},-i}.$$

Furthermore,

$$\begin{aligned} L^2_{\mathbf{M}} - \text{Kernel} \left(H^{(0)} - \mu_S^{(0)} \text{Id} \right) \\ = \text{span} \left\{ \Phi_{+1}^{(0)}(\mathbf{x}) \right\} \oplus \text{span} \left\{ \Phi_{-1}^{(0)}(\mathbf{x}) \right\} \\ \oplus \text{span} \left\{ \Phi_{+i}^{(0)}(\mathbf{x}) \right\} \oplus \text{span} \left\{ \Phi_{-i}^{(0)}(\mathbf{x}) \right\}. \end{aligned}$$

3. $\mu_S^{(0)} = 2\pi^2$ is the lowest eigenvalue of $H^{(0)}$ in $L^2_{\mathbf{M}}$.

Proof. The function $e^{i\mathbf{k} \cdot \mathbf{x}}$ is an $L^2_{\mathbf{k}}$ -eigenvalue of $-\Delta$ with eigenvalue $|\mathbf{k}|^2$. Since vertices of the Brillouin zone, \mathbf{M} , $R\mathbf{M}$, $R^2\mathbf{M}$, and $R^3\mathbf{M}$, are equidistant from the origin and are all equivalent modulo Λ^* , we have $\mu_S^{(0)} = |\mathbf{M}|^2 = 2\pi^2$ is an $L^2_{\mathbf{M}}$ -eigenvalue of multiplicity at least four with eigenspace contained in the span of the functions $e^{i\mathbf{M} \cdot \mathbf{x}}$, $e^{iR\mathbf{M} \cdot \mathbf{x}}$, $e^{iR^2\mathbf{M} \cdot \mathbf{x}}$, and $e^{iR^3\mathbf{M} \cdot \mathbf{x}}$. To show that $\mu_S^{(0)}$ is of multiplicity exactly four, we seek to find \mathbf{m} for which $|\mathbf{M}^{\mathbf{m}}|^2 = |\mathbf{M}|^2$. Using $\mathbf{M}^{\mathbf{m}} = \mathbf{M} + m_1\mathbf{k}_1 + m_2\mathbf{k}_2$, we have $|\mathbf{M}^{\mathbf{m}}|^2 - |\mathbf{M}|^2 = (2\pi)^2 [m_1^2 + m_2^2 + m_1 + m_2]$, which vanishes only if $\mathbf{m} = (0, 0)$, $(0, -1)$, $(-1, -1)$, or $(-1, 0)$. These four possibilities correspond to the

four vertices of \mathcal{B} . Thus, $\mu_S^{(0)}$ is of multiplicity exactly four. This proves part 1. Part 2 is a consequence of Proposition 2.13 and its proof. Part 3 holds because $m_1^2 + m_2^2 + m_1 + m_2 \geq 1 > 0$ for $\mathbf{m} = (m_1, m_2) \notin \{(0, 0), (0, -1), (-1, -1), (-1, 0)\}$ and therefore $|\mathbf{M}^{\mathbf{m}}|^2 \geq |\mathbf{M}|^2 + (2\pi)^2 > |\mathbf{M}|^2$. \square

4. Twofold degenerate $L_{\mathbf{M}}^2$ eigenvalues imply quadratic touching of dispersion surfaces.

THEOREM 4.1. *Let $H = -\Delta_{\mathbf{x}} + V(\mathbf{x})$, where $V(\mathbf{x})$ is an admissible potential in the sense of Definition 2.3. Assume that μ_* is a twofold degenerate $L_{\mathbf{M}}^2$ eigenvalue of H . More specifically,*

- (H1) *H has a simple $L_{\mathbf{M},+i}^2$ eigenvalue μ_S with corresponding normalized eigenfunction $\Phi_1(\mathbf{x}) = e^{i\mathbf{M}\cdot\mathbf{x}}\phi_1(\mathbf{x})$;*
- (H2) *H has a simple $L_{\mathbf{M},-i}^2$ eigenvalue μ_S with corresponding normalized eigenfunction*

$$\Phi_2 = (\mathcal{P} \circ \mathcal{C})[\Phi_1](\mathbf{x}) = \overline{\Phi_1(-\mathbf{x})} \equiv e^{i\mathbf{M}\cdot\mathbf{x}}\phi_2(\mathbf{x}),$$

and we shall also use the notation $\Phi_1 = \Phi_{(+i)}$ and $\Phi_2 = \Phi_{(-i)}$;

- (H3) *μ_S is neither an $L_{\mathbf{M},+1}^2$ nor an $L_{\mathbf{M},-1}^2$ eigenvalue of H .*

Then, there exist dispersion relations $\mathbf{k} \mapsto \mu_{\pm}(\mathbf{k})$ associated with the $L_{\mathbf{k}}^2$ -eigenvalue problem for H , whose local character in a neighborhood of the high-symmetry quasi momentum, \mathbf{M} (and therefore all vertices of \mathcal{B}), is given by

$$(4.1) \quad \mu_{\pm}(\mathbf{M} + \kappa) - \mu_S = (1 - \alpha)|\kappa|^2 + \mathcal{Q}_6(\kappa) \pm \sqrt{\left| \gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2 \right|^2 + \mathcal{Q}_8(\kappa)}$$

for $|\mathbf{k} - \mathbf{M}| = \sqrt{\kappa_1^2 + \kappa_2^2}$ small. The constants $\alpha \in \mathbb{R}$ and $\beta, \gamma \in \mathbb{C}$ are inner product expressions which are quadratic in the entries of $\nabla_{\mathbf{x}}\Phi_1$ and $\nabla_{\mathbf{x}}\Phi_2$; see (4.27). The functions $\mathcal{Q}_6(\kappa) = \mathcal{O}(|\kappa|^6)$ and $\mathcal{Q}_8(\kappa) = \mathcal{O}(|\kappa|^8)$ are analytic functions of κ and invariant under $\pi/2$ rotation: $(\kappa_1, \kappa_2) \mapsto (-\kappa_2, \kappa_1)$.

The proof of Theorem 4.1 is given in section 4.1.

COROLLARY 4.2. *Assume hypotheses of Theorem 4.1. Assume further that with respect to the origin of coordinates, $\mathbf{x}_c = 0$, we have, in addition, that V is reflection invariant in the sense of Definition 2.4, i.e., $V(x_1, x_2) = V(x_2, x_1)$. Then,*

$$(4.2) \quad \mu_{\pm}(\mathbf{M} + \kappa) - \mu_S = (1 - \alpha)|\kappa|^2 + \mathcal{Q}_6(\kappa) \pm \sqrt{\left| 2\beta\kappa_1\kappa_2 \right|^2 + \mathcal{Q}_8(\kappa)}.$$

Before presenting the proofs of Theorem 4.1 and Corollary 4.2, we state a result on the instability or nonpersistence of the quadratic degeneracies of Theorem 4.1 against a class of real-valued perturbations which preserve \mathbb{Z}^2 -periodicity and inversion symmetry but break $\pi/2$ -rotational invariance.

THEOREM 4.3 (nonpersistence of quadratic degeneracy). *Consider $H^{\eta} = -\Delta + V + \eta W$, where V is admissible. By Theorem 4.1,*

- *H^0 has an $L_{\mathbf{M}}^2$ -eigenvalue μ_S of geometric multiplicity two, and*
- *μ_S has an associated orthonormal basis $\{\Phi_1, \Phi_2\}$ with $\Phi_1 \in L_{\mathbf{M},i}^2$ and $\Phi_2(\mathbf{x}) = \overline{\Phi_1(-\mathbf{x})}$.*

We introduce a class of perturbations, W , consisting of real-valued functions which are \mathbb{Z}^2 -periodic and even but which do not respect $\pi/2$ -rotational invariance, i.e., $\mathcal{R}[W] \neq W$. In particular, we assume that




FIG. 4.1. The admissible potential V in Figure 7.4 with the additional perturbation $W = 2 \cos((\mathbf{k}_1 + \mathbf{k}_2) \cdot \mathbf{x})$.

$$(4.3) \quad \langle \Phi_1, W \Phi_2 \rangle \neq 0.$$

Then, the twofold degenerate eigenvalue splits into two simple eigenvalues, ν_{\pm} , given by

$$(4.4) \quad \nu_{\pm} = \mu_S + \eta \langle \Phi_1, W \Phi_1 \rangle \pm \eta |\langle \Phi_1, W \Phi_2 \rangle| + \mathcal{O}(\eta^2).$$

We omit the proof of Theorem 4.3, which follows from the degenerate perturbation theory argument; see section 9 (particularly Remark 9.2) of [9] and section 5 of [17].

Remark 4.4. It is easy to verify that if W is real-valued, \mathbb{Z}^2 -periodic, even, and $\pi/2$ -rotationally invariant, then $\langle \Phi_1, W \Phi_2 \rangle = 0$.

Remark 4.5. We provide an example of such potential W that is even, is *not* $\pi/2$ -rotationally invariant, and has the property $\langle \Phi_1^\varepsilon, W \Phi_2^\varepsilon \rangle \neq 0$. We set

$$W_0(\mathbf{x}) = 2 \cos((\mathbf{k}_1 + \mathbf{k}_2) \cdot \mathbf{x}).$$

Then, $\mathcal{R}[W_0] = 2 \cos((\mathbf{k}_1 + \mathbf{k}_2) \cdot R^* \mathbf{x}) = 2 \cos(R(\mathbf{k}_1 + \mathbf{k}_2) \cdot \mathbf{x}) = 2 \cos((\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{x}) \neq W_0(\mathbf{x})$. We obtain $\langle \Phi_1^\varepsilon, W_0 \Phi_2^\varepsilon \rangle = -2 + \mathcal{O}(\varepsilon) \neq 0$ for ε small.

Other examples which satisfy the hypotheses of Theorem 4.3 are $W_1(\mathbf{x}) = 2 \cos(\mathbf{k}_1 \cdot \mathbf{x})$, $W_2(\mathbf{x}) = 2 \cos(\mathbf{k}_2 \cdot \mathbf{x})$, and $W_3(\mathbf{x}) = 2 \cos((\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{x})$. We omit the lengthy but elementary verification. A numerical illustration of Theorem 4.3 is presented in Figure 4.1.

4.1. Proof of Theorem 4.1 on conditions for quadratic degeneracy.

4.1.1. Reduction to the study of $\det \mathcal{M}(\mu, \kappa) = 0$ for a 2×2 -matrix-valued analytic function $(\mu, \kappa) \mapsto \mathcal{M}(\mu, \kappa)$ in a neighborhood of $(0, 0)$. The proof follows closely that of Theorem 4.1 of [9]. For $\Phi \in L^2_{\mathbf{k}}$ as $\Phi(\mathbf{x}; \mathbf{k}) = e^{i\mathbf{k} \cdot \mathbf{x}} \phi(\mathbf{x}; \mathbf{k})$, where $\phi(\mathbf{x}; \mathbf{k})$ is Λ -periodic. Let $H(\mathbf{k}) = (-\nabla_{\mathbf{x}} + i\mathbf{k})^2 + V(\mathbf{x})$. We study the eigenvalue problem $H(\mathbf{k})\phi(\mathbf{x}; \mathbf{k}) = \mu\phi(\mathbf{x}; \mathbf{k})$ for $\mathbf{k} = \mathbf{M} + \kappa$ and $|\kappa| \ll 1$. In particular,

$$(4.5) \quad \begin{aligned} & [-(\nabla_{\mathbf{x}} + i(\mathbf{M} + \kappa))^2 + V(\mathbf{x})] \phi = \mu \phi, \\ & \phi(\mathbf{x} + \mathbf{v}) = \phi(\mathbf{x}) \text{ for all } \mathbf{v} \in \Lambda, \quad \mathbf{x} \in \mathbb{R}^2. \end{aligned}$$

We seek a solution of (4.5), $\mu = \mu(\mathbf{M} + \kappa)$ and $\phi = \phi(\mathbf{x}; \mathbf{M} + \kappa)$, in the form

$$(4.6) \quad \mu(\mathbf{M} + \kappa) = \mu_S + \mu^{(1)}, \quad \phi(\mathbf{x}; \mathbf{M} + \kappa) = \phi^{(0)} + \phi^{(1)},$$

where

$$\phi^{(0)} \in \ker(H(\mathbf{M}) - \mu_S I), \quad \phi^{(1)} \perp \ker(H(\mathbf{M}) - \mu_S I),$$

and $\mu^{(1)}$ are to be determined. Substituting (4.6) into (4.5), we obtain

$$(4.7) \quad (H(\mathbf{M}) - \mu_S I)\phi^{(1)} = \left(2i\kappa \cdot (\nabla + i\mathbf{M}) - \kappa \cdot \kappa + \mu^{(1)}\right) (\phi^{(0)} + \phi^{(1)}) \equiv F^{(1)}.$$

The right-hand side of (4.7) depends on $\phi^{(0)} \in \ker(H(\mathbf{M}) - \mu_0 I)$, which, by hypothesis, is expressed for constants α_1, α_2 to be determined,

$$(4.8) \quad \begin{aligned} \phi^{(0)} &= \alpha_1 \phi_1(\mathbf{x}) + \alpha_2 \phi_2(\mathbf{x}), \\ \phi_j(\mathbf{x}) &= e^{-i\mathbf{M} \cdot \mathbf{x}} \Phi_j(\mathbf{x}), j = 1, 2. \end{aligned}$$

We next construct $\phi^{(1)}$. Introduce the orthogonal projections Q_{\parallel} onto the two-dimensional kernel of $H(\mathbf{M}) - \mu_S I$ and $Q_{\perp} = I - Q_{\parallel}$. Note that $Q_{\parallel}\psi^{(1)} = 0$, $Q_{\perp}\psi^{(0)} = 0$, and $Q_{\perp}\psi^{(1)} = \psi^{(1)}$. Equation (4.7) is of the form $(H(\mathbf{M}) - \mu_S I)\phi^{(1)} = F^{(1)}(\alpha_1, \alpha_2, \kappa, \mu^{(1)}, \phi^{(1)})$ and can be expressed as an equivalent system for $(\phi^{(1)}, \mu^{(1)})$:

$$(4.9) \quad (H(\mathbf{M}) - \mu_S I)\phi^{(1)} = Q_{\perp} F^{(1)}(\alpha_1, \alpha_2, \kappa, \mu^{(1)}, \phi^{(1)}),$$

$$(4.10) \quad 0 = Q_{\parallel} F^{(1)}(\alpha_1, \alpha_2, \kappa, \mu^{(1)}, \phi^{(1)}).$$

We proceed as in [9]. We introduce the resolvent $\mathcal{R}_{\mathbf{M}}(\mu_S) = (H(\mathbf{M}) - \mu_S I)^{-1}$, which is a bounded linear map from $Q_{\perp} L^2(\mathbb{R}^2/\Lambda)$ to $Q_{\perp} H^2(\mathbb{R}^2/\Lambda)$. Equivalently, $\mathcal{R}(\mu_S) = (H - \mu_S I)^{-1}$ is a bounded linear map from $\tilde{Q}_{\perp} L^2_{\mathbf{M}}$ to $\tilde{Q}_{\perp} H^2_{\mathbf{M}}$, where \tilde{Q} and \tilde{Q}_{\perp} are the orthogonal projections onto $\text{span}\{\Phi_1, \Phi_2\}$ and its orthogonal complement. For $|\kappa|$ and $|\mu^{(1)}|$ sufficiently small, we have

$$(4.11) \quad \phi^{(1)} = \alpha_1 c^{(1)}[\kappa, \mu^{(1)}](\mathbf{x}) + \alpha_2 c^{(2)}[\kappa, \mu^{(1)}](\mathbf{x}),$$

where

$$(4.12) \quad \begin{aligned} c^{(j)}[\kappa, \mu^{(1)}](\mathbf{x}) &= \left(I + \mathcal{R}_{\mathbf{M}}(\mu_S) Q_{\perp} \left(-2i\kappa \cdot (\nabla + i\mathbf{M}) + \kappa \cdot \kappa - \mu^{(1)} \right) \right)^{-1} \\ &\quad \circ (\mathcal{R}_{\mathbf{M}}(\mu_S) Q_{\perp} (2i\kappa \cdot (\nabla + i\mathbf{M})) \phi_j), \end{aligned}$$

where $(\kappa, \mu^{(1)}) \rightarrow c^{(j)}[\kappa, \mu^{(1)}]$ is a smooth mapping from a neighborhood $(0, 0) \in \mathbb{R}^2 \times \mathbb{C}$ into $H^2(\mathbb{R}^2/\Lambda)$ satisfying the bound $\|c^{(j)}\|_{H^2} \leq C(|\kappa|^2 + |\mu^{(1)}| |\kappa|); j = 1, 2$.

Substituting (4.11) into (4.10), we obtain a homogeneous system

$$\mathcal{M}(\mu^{(1)}, \kappa) \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix} = 0.$$

We therefore have the following characterization of eigenvalues, $\mu = \mu_S + \mu^{(1)}$ for $|\mu^{(1)}|$ small and $\mathbf{k} = \mathbf{M} + \kappa$, with κ near zero.

PROPOSITION 4.6. *Let $\mathbf{k} = \mathbf{M} + \kappa$ with $|\kappa| < \kappa_{\max}$ sufficiently small. Then, for $\mu = \mu_S + \mu^{(1)}$, with $|\mu^{(1)}|$ in a small neighborhood of 0, is an $L^2_{\mathbf{M}}$ -eigenvalue if and only if $\det \mathcal{M}(\mu^{(1)}, \kappa) = 0$.*

The matrix $\mathcal{M}(\mu^{(1)}, \kappa)$ is given by

$$(4.13) \quad \mathcal{M}(\mu^{(1)}, \kappa) \equiv \mathcal{M}^{(0)}(\mu^{(1)}, \kappa) + \mathcal{M}^{(1)}(\mu^{(1)}, \kappa),$$

where

$$(4.14) \quad \mathcal{M}^{(0)}(\mu^{(1)}, \kappa) \equiv \begin{pmatrix} \mu^{(1)} - \kappa \cdot \kappa + \langle \Phi_1, 2i\kappa \cdot \nabla \Phi_1 \rangle & \langle \Phi_1, 2i\kappa \cdot \nabla \Phi_2 \rangle \\ \langle \Phi_2, 2i\kappa \cdot \nabla \Phi_1 \rangle & \mu^{(1)} - \kappa \cdot \kappa + \langle \Phi_2, 2i\kappa \cdot \nabla \Phi_2 \rangle \end{pmatrix}$$

and

$$(4.15) \quad \mathcal{M}^{(1)}(\mu^{(1)}, \kappa) \equiv \begin{pmatrix} \langle \Phi_1, 2i\kappa \cdot \nabla C^{(1)}(\kappa, \mu^{(1)}) \rangle & \langle \Phi_1, 2i\kappa \cdot \nabla C^{(2)}(\kappa, \mu^{(1)}) \rangle \\ \langle \Phi_2, 2i\kappa \cdot \nabla C^{(1)}(\kappa, \mu^{(1)}) \rangle & \langle \Phi_2, 2i\kappa \cdot \nabla C^{(2)}(\kappa, \mu^{(1)}) \rangle \end{pmatrix}.$$

Using the relations $(\nabla + i\mathbf{M})\phi_j = e^{-i\mathbf{M} \cdot \mathbf{x}} \nabla_{\mathbf{x}} e^{i\mathbf{M} \cdot \mathbf{x}} \phi_j = e^{-i\mathbf{M} \cdot \mathbf{x}} \nabla_{\mathbf{x}} \Phi_j$, we have $C^{(j)}[\kappa, \mu^{(1)}](\mathbf{x}) \equiv e^{i\mathbf{M} \cdot \mathbf{x}} c^{(j)}[\kappa, \mu^{(1)}](\mathbf{x})$, where $\langle \Phi_i, C^{(j)} \rangle = 0$ for $i, j = 1, 2$.

Remark 4.7. Given real $\mu^{(1)}$ and κ , the matrices \mathcal{M} , $\mathcal{M}^{(0)}$, and $\mathcal{M}^{(1)}$ are Hermitian matrices.

We conclude this section with an elementary lemma which we use, along with symmetry, to simplify the matrix entries of $\mathcal{M}(\mu^{(1)}, \kappa)$. We denote $\mathcal{M}_{j_1, j_2}^{(1)}(\mu^{(1)}, \kappa) = \langle \Phi_{j_1}, 2i\kappa \cdot \nabla C^{(j_2)}(\kappa, \mu^{(1)}) \rangle$.

LEMMA 4.8. *Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $f \in L^2(\Omega)$ satisfies $f(R^* \mathbf{x}) = f(\mathbf{x})$, where R^* is the counterclockwise rotation matrix by $\pi/2$. Then,*

$$(4.16) \quad \nabla_{\mathbf{x}} \mathcal{R}[f](\mathbf{x}) = R \mathcal{R}[\nabla_{\mathbf{y}} f](\mathbf{x}) \quad \text{or} \quad \partial_{x_\alpha} \mathcal{R}[f](\mathbf{x}) = R_{\alpha r} \mathcal{R}[\partial_{y_r} f](\mathbf{x}).$$

Proof. Let $\mathbf{y} = R^* \mathbf{x}$ or $\mathbf{y}_r = R_{ir} \mathbf{x}_i$. Therefore, $\frac{\partial \mathbf{y}_r}{\partial \mathbf{x}_\alpha} = R_{\alpha r}$. Fix $\alpha \in \{1, 2\}$. Then,

$$(4.17) \quad \partial_{x_\alpha} \mathcal{R}[f](\mathbf{x}) = \partial_{x_\alpha} f(R^* \mathbf{x}) = R_{\alpha r} \mathcal{R}[\partial_{y_r} f](\mathbf{x}) = \{R \mathcal{R}(\nabla f)\}_\alpha. \quad \square$$

4.1.2. Symmetry implies $\det \mathcal{M}(\mu^{(1)}, \kappa)$ has no linear in κ terms for $|\kappa| \ll 1$.

PROPOSITION 4.9.

$$(4.18) \quad \mathcal{M}^{(0)}(\mu^{(1)}, \kappa) = \begin{pmatrix} \mu^{(1)} - \kappa \cdot \kappa \end{pmatrix} \times \mathbf{I}_{2 \times 2}$$

and therefore

$$(4.19) \quad \mathcal{M}(\mu^{(1)}, \kappa) = \begin{pmatrix} \mu^{(1)} - \kappa \cdot \kappa \end{pmatrix} \times \mathbf{I}_{2 \times 2} + \mathcal{M}^{(1)}(\mu^{(1)}, \kappa).$$

Recall from the hypotheses of Theorem 4.1 that $\Phi_1 \in L^2_{\mathbf{M}, +i}$ and $\Phi_2 \in L^2_{\mathbf{M}, -i}$ and therefore $\mathcal{R}[\Phi_q](\mathbf{x}) = i^{2q-1} \Phi_q(\mathbf{x})$, $q = 1, 2$.

PROPOSITION 4.10. *For $j_1, j_2 = 1, 2$,*

$$\langle \Phi_{j_1}, \nabla \Phi_{j_2} \rangle_{L^2(\Omega)} = \mathbf{0}.$$

Proof. Choose $j_1, j_2 \in \{1, 2\}$. Using that \mathcal{R} is unitary and Lemma 4.8, we have

$$\begin{aligned}
\langle \Phi_{j_1}, \nabla_{\mathbf{y}} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{y}})} &= \langle \mathcal{R}[\Phi_{j_1}], \mathcal{R}[\nabla_{\mathbf{y}} \Phi_{j_2}] \rangle_{L^2(\Omega_{\mathbf{x}})} = \langle \mathcal{R}[\Phi_{j_1}], R^* \nabla_{\mathbf{x}} \mathcal{R}[\Phi_{j_2}] \rangle_{L^2(\Omega_{\mathbf{x}})} \\
&= R^* \langle \mathcal{R}[\Phi_{j_1}], \nabla_{\mathbf{x}} \mathcal{R}[\Phi_{j_2}] \rangle_{L^2(\Omega_{\mathbf{x}})} = R^* \langle i^{2j_1-1} \Phi_{j_1}, \nabla_{\mathbf{x}} i^{2j_2-1} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{x}})} \\
&= i^{2(j_2-j_1)} R^* \langle \Phi_{j_1}, \nabla_{\mathbf{x}} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{x}})}.
\end{aligned}$$

It follows that either $i^{2(j_2-j_1)}$ is an eigenvalue of R or $\langle \Phi_{j_1}, \nabla_{\mathbf{x}} \Phi_{j_2} \rangle = \mathbf{0}$. But the eigenvalues of R are $\pm i$, and since $j_2 - j_1$ is an integer, $i^{2(j_2-j_1)}$ is real. We conclude that $\langle \Phi_{j_1}, \nabla_{\mathbf{x}} \Phi_{j_2} \rangle = \mathbf{0}$ for all $j_1, j_2 = 1, 2$. The proof of Proposition 4.10 is complete.

Remark 4.11. As observed in section 4.1.1, $\|C^{(j)}(\kappa, \mu^{(1)})\|_{H^1} \lesssim |\kappa| + |\kappa|^2 + |\mu^{(1)}|$, we have from Proposition 4.9 that $\mathcal{M}(\mu^{(1)}, \kappa) = (\mu^{(1)} - |\kappa|^2) I_{2 \times 2} + \mathcal{O}_{2 \times 2}(|\kappa|^2 + |\kappa|^3 + |\mu^{(1)}| |\kappa|)$. Therefore, $\det \mathcal{M}(\mu^{(1)}, \kappa) = 0$ has solutions $\mu_{\pm}^{(1)} = \mathcal{O}(|\kappa|^2)$. We next obtain the precise quadratic dependence on κ of $\mathcal{M}^{(1)}(\mu^{(1)}, \kappa)$ and then give a more precise expansion of solutions to $\det \mathcal{M}(\mu^{(1)}, \kappa) = 0$.

4.1.3. Quadratic in κ terms of $\det \mathcal{M}(\mu^{(1)}, \kappa)$ for $|\kappa| \ll 1$. We next expand $\mathcal{M}_{j_1, j_2}^{(1)}(\mu^{(1)}, \kappa)$ for $|\kappa|$ and $|\mu^{(1)}|$ small. Recall first (4.12), the relations listed after (4.15). Then, $\tilde{Q}_{\perp} C^{(j)}[\kappa, \mu^{(1)}] = C^{(j)}[\kappa, \mu^{(1)}]$, where

$$\begin{aligned}
C^{(j)}[\kappa, \mu^{(1)}] &= \left(I + \mathcal{R}(\mu_S) \tilde{Q}_{\perp} \left(-2i\kappa \cdot \nabla + \kappa \cdot \kappa - \mu^{(1)} \right) \right)^{-1} \circ \left(\mathcal{R}(\mu_S) \tilde{Q}_{\perp} (2i\kappa \cdot \nabla) \Phi_j \right) \\
(4.20) \quad &= \left(I + \mathcal{O}_{L^2 \rightarrow L^2}(|\kappa| + |\mu^{(1)}|) \right) \circ \left(\mathcal{R}(\mu_S) \tilde{Q}_{\perp} (2i\kappa \cdot \nabla) \Phi_j \right).
\end{aligned}$$

Furthermore, recalling that $\tilde{Q}_{\perp} \partial_{x_l} \Phi_m = \partial_{x_l} \Phi_m$ (Proposition 4.10), we have

$$\mathcal{M}_{j_1, j_2}^{(1)}(\mu^{(1)}, \kappa) = \left\langle \Phi_{j_1}, 2i\kappa \cdot \nabla C^{(j_2)}(\kappa, \mu^{(1)}) \right\rangle = \left\langle \tilde{Q}_{\perp} 2i\kappa \cdot \nabla \Phi_{j_1}, \tilde{Q}_{\perp} C^{(j_2)}(\kappa, \mu^{(1)}) \right\rangle.$$

Therefore, by (4.20) we have for $j_1, j_2 = 1, 2$ and $\kappa \in \mathbb{R}^2$,

$$\begin{aligned}
\mathcal{M}_{j_1, j_2}^{(1)}(\mu^{(1)}, \kappa) &= 4 \sum_{l, m=1}^2 \langle \tilde{Q}_{\perp} \partial_{x_l} \Phi_{j_1}, \mathcal{R}(\mu_S) \tilde{Q}_{\perp} \partial_{x_m} \Phi_{j_2} \rangle_{\kappa_l \kappa_m} + \mathcal{O}(|\kappa|^3 + |\mu^{(1)}| |\kappa|) \\
(4.21) \quad &= 4 \sum_{l, m=1}^2 \langle \partial_{x_l} \Phi_{j_1}, \mathcal{R}(\mu_S) \partial_{x_m} \Phi_{j_2} \rangle_{\kappa_l \kappa_m} + \mathcal{O}(|\kappa|^3 + |\mu^{(1)}| |\kappa|) \\
(4.22) \quad &\equiv 4 \kappa^T A^{j_1, j_2} \kappa + \mathcal{O}(|\kappa|^3 + |\mu^{(1)}| |\kappa|),
\end{aligned}$$

where (4.22) defines the matrix A^{j_1, j_2} with entries

$$(4.23) \quad a_{l, m}^{j_1, j_2} = \langle \partial_{x_l} \Phi_{j_1}, \mathcal{R}(\mu_S) \partial_{x_m} \Phi_{j_2} \rangle,$$

We proceed now to use symmetry to deduce the structure of the matrices A^{j_1, j_2} .

LEMMA 4.12. *For fixed $j_1, j_2 \in \{1, 2\}$, we have the following:*

$$(4.24) \quad R^T A^{j_1, j_2} R = i^{2(j_2-j_1)} A^{j_1, j_2},$$

where R denotes the $\pi/2$ -rotation matrix displayed in (2.5). Therefore,

$$\begin{aligned}
j_1 = j_2 \equiv j &\implies R^T A^{j, j} R = A^{j, j}, \\
j_1 \neq j_2 &\implies R^T A^{j_1, j_2} R = -A^{j_1, j_2}.
\end{aligned}$$

Proof. We will use Lemma 4.8, $\mathcal{R}[\partial_{x_l} f](\mathbf{x}) = R_{nl} \mathcal{R}[\partial_{y_n} f](\mathbf{x})$. Since \mathcal{R} is unitary and commutes with $\mathcal{R}(\mu_S)$,

$$\begin{aligned} \kappa^T A^{j_1, j_2} \kappa &= \langle \partial_{y_l} \Phi_{j_1}, \mathcal{R}(\mu_S) \partial_{y_m} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{y}})} \kappa_l \kappa_m \\ &= \langle \mathcal{R}[\partial_{y_l} \Phi_{j_1}], \mathcal{R}(\mu_S) \mathcal{R}[\partial_{y_m} \Phi_{j_2}] \rangle_{L^2(\Omega_{\mathbf{x}})} \kappa_l \kappa_m \\ &= \langle R_{nl} \partial_{x_n} \mathcal{R}[\Phi_{j_1}], \mathcal{R}(\mu_S) R_{qm} \mathcal{R}[\partial_{x_q} \Phi_{j_2}] \rangle_{L^2(\Omega_{\mathbf{x}})} \kappa_l \kappa_m \\ &= \langle \partial_{x_n} i^{2j_1-1} \Phi_{j_1}, \mathcal{R}(\mu_S) \partial_{x_q} i^{2j_2-1} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{x}})} R_{nl} \kappa_l R_{qm} \kappa_m \\ &= i^{2(j_2-j_1)} \langle \partial_{x_n} \Phi_{j_1} \mathcal{R}(\mu_S) \partial_{x_q} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{x}})} (R\kappa)_n (R\kappa)_q \\ &= i^{2(j_2-j_1)} (R\kappa)^T A^{j_1, j_2} (R\kappa). \end{aligned}$$

Since κ is arbitrary, $A^{j_1, j_2} = i^{2(j_2-j_1)} R^T A^{j_1, j_2} R$. \square

LEMMA 4.13. Assume R is the $\pi/2$ -rotation matrix, (2.5), and $A = (a_{ij})$. Then,

$$\begin{aligned} (j_1 = j_2) \quad R^T A R &= A \quad \implies \quad A = \begin{pmatrix} a_{11} & a_{12} \\ -a_{12} & a_{11} \end{pmatrix}, \\ (j_1 \neq j_2) \quad R^T A R &= -A \quad \implies \quad A = \begin{pmatrix} a_{11} & a_{12} \\ a_{12} & -a_{11} \end{pmatrix}. \end{aligned}$$

CLAIM 4.14. Let A^\dagger be the conjugate-transpose of A . Then,

$$(A^{1,1})^\dagger = A^{1,1}, \quad (A^{2,2})^\dagger = A^{2,2}, \quad \text{and} \quad (A^{2,1})^\dagger = A^{1,2}.$$

Proof. Pick $j_1, j_2 \in \{1, 2\}$ and $l, m \in \{1, 2\}$.

$$\begin{aligned} (A^{j_1, j_2})_{l, m}^\dagger &= \overline{(a_{m, l}^{j_1, j_2})} = \left[\overline{\langle \partial_{x_m} \Phi_{j_1}, \mathcal{R}(\mu_S) \partial_{x_l} \Phi_{j_2} \rangle} \right] \\ &= \left[\langle \partial_{x_l} \Phi_{j_2}, \mathcal{R}(\mu_S) \partial_{x_m} \Phi_{j_1} \rangle \right] = (a_{l, m}^{j_2, j_1}) = (A^{j_2, j_1})_{l, m}, \end{aligned}$$

and therefore $(A^{j_1, j_2})^\dagger = A^{j_2, j_1}$. \square

CLAIM 4.15. $A^{1,1} = (A^{2,2})^T$. In particular, $a_{11}^{1,1} = a_{11}^{2,2}$.

Proof. Recall $\Phi_1(\mathbf{x}) = (\mathcal{P} \circ \mathcal{C})[\Phi_1](\mathbf{x}) = \overline{\Phi_2(-\mathbf{x})}$. For $l, m \in \{1, 2\}$, using that $\partial_{x_l}(\mathcal{P} \circ \mathcal{C}) = -(\mathcal{P} \circ \mathcal{C})\partial_{y_l}$, we have

$$\begin{aligned} a_{l, m}^{1,1} &= \langle \partial_{x_l} \Phi_1, \mathcal{R}(\mu_S) \partial_{x_m} \Phi_1 \rangle \\ &= \langle \partial_{x_l} (\mathcal{P} \circ \mathcal{C})[\Phi_2], \mathcal{R}(\mu_S) \partial_{x_m} (\mathcal{P} \circ \mathcal{C})[\Phi_2] \rangle \\ &= \langle (\mathcal{P} \circ \mathcal{C})[\partial_{y_l} \Phi_2], \mathcal{R}(\mu_S) (\mathcal{P} \circ \mathcal{C})[\partial_{y_m} \Phi_2] \rangle \\ &= \langle \mathcal{C}[\partial_{y_l} \Phi_2], \mathcal{R}(\mu_S) \mathcal{C}[\partial_{y_m} \Phi_2] \rangle \quad (= \overline{a_{lm}^{2,2}}) \\ &= \langle \partial_{y_m} \Phi_2, \mathcal{R}(\mu_S) \partial_{y_l} \Phi_2 \rangle = a_{m, l}^{2,2}. \end{aligned}$$

Therefore, $A^{1,1} = \overline{A^{2,2}} = (A^{2,2})^T$. \square

By (4.22) we have

$$(4.25) \quad \mathcal{M}^{(1)}(\mu^{(1)}, \kappa) = 4 \begin{pmatrix} \kappa^T A^{1,1} \kappa & \kappa^T A^{2,1} \kappa \\ \kappa^T A^{1,2} \kappa & \kappa^T A^{2,2} \kappa \end{pmatrix} + \mathcal{O}_{2 \times 2}(|\kappa|^3 + |\mu^{(1)}| |\kappa|).$$

Simplifying the leading term in (4.25) we observe, by the above claims, that $a_{1,1}^{1,1} \in \mathbb{R}$ and

$$A^{1,1} = \begin{pmatrix} a_{1,1}^{1,1} & a_{1,2}^{1,1} \\ -a_{1,2}^{1,1} & a_{1,1}^{1,1} \end{pmatrix}; \quad A^{1,2} = \overline{A^{2,1}} = \begin{pmatrix} a_{1,1}^{1,2} & a_{1,2}^{1,2} \\ a_{1,2}^{1,2} & -a_{1,1}^{1,2} \end{pmatrix}; \quad A^{2,2} = \begin{pmatrix} a_{1,1}^{1,1} & -a_{1,2}^{1,1} \\ a_{1,2}^{1,1} & a_{1,1}^{1,1} \end{pmatrix}.$$

Hence,

$$(4.26) \quad \mathcal{M}^{(1)}(\mu^{(1)}, \kappa) = \begin{pmatrix} \alpha (\kappa_1^2 + \kappa_2^2) & \gamma (\kappa_1^2 - \kappa_2^2) + 2\beta \kappa_1 \kappa_2 \\ \bar{\gamma} (\kappa_1^2 - \kappa_2^2) + 2\bar{\beta} \kappa_1 \kappa_2 & \alpha (\kappa_1^2 + \kappa_2^2) \end{pmatrix} \\ + \mathcal{O}_{2 \times 2}(|\kappa|^3 + |\mu^{(1)}||\kappa|),$$

where, by (4.23),

$$(4.27) \quad \begin{aligned} \alpha &= 4a_{1,1}^{1,1} = 4 \langle \partial_{x_1} \Phi_1, \mathcal{R}(\mu_S) \partial_{x_1} \Phi_1 \rangle, \\ \beta &= 4a_{1,2}^{1,2} = 4 \langle \partial_{x_1} \Phi_1, \mathcal{R}(\mu_S) \partial_{x_2} \Phi_2 \rangle, \\ \gamma &= 4a_{1,1}^{1,2} = 4 \langle \partial_{x_1} \Phi_1, \mathcal{R}(\mu_S) \partial_{x_1} \Phi_2 \rangle. \end{aligned}$$

Remark 4.16. Observe by Claim (4.14), the terms $\alpha = 4a_{1,1}^{1,1}$ are real.

4.2. Symmetries of dispersion maps $\mathbf{k} \mapsto \mu_{\pm}(\mathbf{k})$ for \mathbf{k} near a vertex of \mathcal{B} . Before deriving a detailed picture of the local character of dispersion surfaces near vertices of \mathcal{B} , we prove a general result on the structure of dispersion surfaces in a neighborhood of a vertex quasi momentum of \mathcal{B} for which the eigenvalue problem has a degenerate eigenvalue.

Assume that the potential V is admissible in the sense of Definition 2.3, where without loss of generality we take the centering $\mathbf{x}_c = 0$. Suppose in addition that V is reflection invariant, i.e., $V(\mathbf{x}) = V(x_1, x_2) = V(x_2, x_1) = V(\rho\mathbf{x})$, where

$$(4.28) \quad \rho\mathbf{x} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1 \end{pmatrix}.$$

To prove Corollary 4.2 we need to show that $\gamma = 4 \langle \partial_{x_1} \Phi_1, \mathcal{R}(\mu_S) \partial_{x_1} \Phi_2 \rangle$ (see (4.27)) vanishes. We first deduce that either $\beta = 4 \langle \partial_{x_1} \Phi_1, \mathcal{R}(\mu_S) \partial_{x_2} \Phi_2 \rangle$ or γ vanishes based on the following symmetry argument; then, we prove that, in fact, $\gamma = 0$ using the properties of ρ . We write

$$(4.29) \quad T_{\rho}[f] = f(\rho^*\mathbf{x}) = f(\rho\mathbf{x}) = f(x_2, x_1).$$

PROPOSITION 4.17. *Let V be an admissible potential in the sense of Definition 2.3; we take $\mathbf{x}_c = 0$ without any loss of generality. In particular, $V(R^*\mathbf{x}) = V(\mathbf{x})$ and $V(-\mathbf{x}) = V(\mathbf{x})$. Assume the hypotheses of Theorem 4.1 which imply that H_V has a degenerate (multiplicity two) $L_{\mathbf{M}}^2$ -eigenvalue, $\mu_S \in \mathbb{R}$.*

Then, for all $\mathbf{k} = \mathbf{M} + \kappa$ with $0 < |\kappa| < \kappa_0$ sufficiently small there exist two eigenvalues given by $\mu_{\pm}(\mathbf{M} + \kappa) = \mu_S \pm \mu^{(1)}(\kappa)$.

1. *For all $0 < |\kappa| < \kappa_0$, we have*

$$(4.30) \quad \{\mu_{-}(\mathbf{M} + \kappa), \mu_{+}(\mathbf{M} + \kappa)\} = \{\mu_{-}(\mathbf{M} + R\kappa), \mu_{+}(\mathbf{M} + R\kappa)\}.$$

2. *Suppose in addition that $V(\rho^*\mathbf{x}) = V(\rho\mathbf{x})$ (recall $\rho = \rho^*$). Then, for $0 < |\kappa| < \kappa_0$, we have*

$$(4.31) \quad \{\mu_{-}(\mathbf{M} + \kappa), \mu_{+}(\mathbf{M} + \kappa)\} = \{\mu_{-}(\mathbf{M} + \rho\kappa), \mu_{+}(\mathbf{M} + \rho\kappa)\}.$$

Proof of Proposition 4.17. Let (μ_{κ}, ψ) denote an $L_{\mathbf{M}+\kappa}^2$ eigenpair of $-\Delta + V$, where μ_{κ} is assumed to be near μ_S . Then, $\mu_{\kappa} = \mu_{-}(\mathbf{M} + \kappa)$ or $\mu_{\kappa} = \mu_{+}(\mathbf{M} + \kappa)$.

Consider now $\tilde{\psi}(\mathbf{x}) \equiv \mathcal{R}[\psi](\mathbf{x}) = \psi(R^*\mathbf{x})$. Note that $(-\Delta + V)\tilde{\psi} = \mu_\kappa \tilde{\psi}$ since \mathcal{R} commutes with $-\Delta + V$. Moreover, for all $\mathbf{v} \in \mathbb{Z}^2$, we have $\tilde{\psi}(\mathbf{x} + \mathbf{v}) = \psi(R^*(\mathbf{x} + \mathbf{v})) = \psi(R^*\mathbf{x} + R^*\mathbf{v}) = e^{i(\mathbf{M} + \kappa) \cdot R^*\mathbf{v}} \tilde{\psi}(\mathbf{x}) = e^{i(R\mathbf{M} + R\kappa) \cdot \mathbf{v}} \tilde{\psi}(\mathbf{x}) = e^{i(\mathbf{M} + R\kappa) \cdot \mathbf{v}} \tilde{\psi}(\mathbf{x})$, where we have used that $R\mathbf{M} \in \mathbf{M} + \Lambda$. Therefore, μ_κ is an $L_{\mathbf{M} + R\kappa}^2$ eigenvalue in a neighborhood of μ_S . Hence, $\{\mu_-(\mathbf{M} + \kappa), \mu_+(\mathbf{M} + \kappa)\} \subset \{\mu_-(\mathbf{M} + R\kappa), \mu_+(\mathbf{M} + R\kappa)\}$. To prove the reverse inclusion, assume $(\hat{\mu}_\kappa, \phi)$ is an $L_{\mathbf{M} + R\kappa}^2$ eigenpair with $\hat{\mu}_\kappa$ near μ_S . Now let $\tilde{\phi} = \mathcal{R}^3[\phi](\mathbf{x})$ and note that $\tilde{\phi} \in L_{\mathbf{M} + \kappa}^2$. Hence $(\hat{\mu}_\kappa, \tilde{\phi})$ is an $L_{\mathbf{M} + \kappa}^2$ -eigenpair of $-\Delta + V$. Therefore, $\{\mu_-(\mathbf{M} + R\kappa), \mu_+(\mathbf{M} + R\kappa)\} \subset \{\mu_-(\mathbf{M} + \kappa), \mu_+(\mathbf{M} + \kappa)\}$ and the proof of part 1 is complete. The proof of part 2 is analogous. This completes the proof of Proposition 4.17. \square

4.3. Local behavior of degenerate dispersion surfaces near the \mathbf{M} -point.

We need to study the solutions of $\det(\mathcal{M}(\mu^{(1)}, \kappa)) = 0$ for κ in a neighborhood of zero. Our strategy is based on a general approach used in [7] (section 13). We extend $\mathcal{M}(\mu^{(1)}, \kappa)$ to be defined as a matrix-valued analytic function of $(\mu^{(1)}, \kappa)$ in a neighborhood of the origin in $\mathbb{C} \times \mathbb{C}^2$ and which agrees with $\mathcal{M}(\mu^{(1)}, \kappa)$ as defined above for $(\mu^{(1)}, \kappa) \in \mathbb{R} \times \mathbb{R}^2$:

$$(4.32) \quad \mathcal{M}(\mu^{(1)}, \kappa) = \begin{pmatrix} m_{11}(\kappa, \mu^{(1)}) & m_{12}(\kappa, \mu^{(1)}) \\ \overline{m_{12}(\bar{\kappa}, \bar{\mu}^{(1)})} & m_{22}(\kappa, \mu^{(1)}) \end{pmatrix}.$$

Here,

$$\begin{aligned} m_{11}(\kappa, \mu^{(1)}) &= (\alpha - 1) |\kappa|^2 + \mu^{(1)} + \mathcal{O}_{2 \times 2}(|\kappa|^3 + |\mu^{(1)}| |\kappa|), \\ m_{12}(\kappa, \mu^{(1)}) &= \gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2 + \mathcal{O}_{2 \times 2}(|\kappa|^3 + |\mu^{(1)}| |\kappa|), \\ m_{21}(\kappa, \mu^{(1)}) &= \overline{m_{12}(\bar{\kappa}, \bar{\mu}^{(1)})} = \bar{\gamma}(\kappa_1^2 - \kappa_2^2) + 2\bar{\beta}\kappa_1\kappa_2 + \mathcal{O}_{2 \times 2}(|\kappa|^3 + |\mu^{(1)}| |\kappa|), \\ m_{22}(\kappa, \mu^{(1)}) &= m_{11}(\kappa, \mu^{(1)}). \end{aligned}$$

Note in particular that $\mathcal{M}(\mu^{(1)}, \kappa)$ given by (4.32) is Hermitian for real $\mu^{(1)}$ and κ .

We first make the simple change of variables

$$(4.33) \quad \nu \equiv (\alpha - 1)(\kappa_1^2 + \kappa_2^2) + \mu^{(1)}.$$

Define

$$(4.34) \quad \widetilde{\mathcal{M}}(\nu, \kappa) = \widetilde{\mathcal{M}}(\nu, \kappa_1, \kappa_2) \equiv \mathcal{M}(\nu - (\alpha - 1)(\kappa_1^2 + \kappa_2^2), \kappa_1, \kappa_2),$$

and study the equivalent equation of $\det(\mathcal{M}(\mu, \kappa)) = 0$ for ν :

$$(4.35) \quad \det(\widetilde{\mathcal{M}}(\nu, \kappa)) = 0.$$

The entries of $\widetilde{\mathcal{M}}(\nu, \kappa)$ are analytic functions of (ν, κ) in a neighborhood of the origin in $\mathbb{C}_\nu \times \mathbb{C}_\kappa^2$. The matrix $\widetilde{\mathcal{M}}(\mu, \kappa)$ has the expansion

$$(4.36) \quad \widetilde{\mathcal{M}}(\nu, \kappa) = \widetilde{\mathcal{M}}_{\text{app}}(\nu, \kappa) + \mathcal{O}_{2 \times 2}(|\kappa|^3 + |\nu| |\kappa|),$$

where

$$(4.37) \quad \widetilde{\mathcal{M}}_{\text{app}}(\nu, \kappa) = \begin{pmatrix} \nu & q(\kappa_1, \kappa_2) \\ q_{\dagger}(\kappa_1, \kappa_2) & \nu \end{pmatrix}$$

and

$$(4.38) \quad q(\kappa_1, \kappa_2) \equiv \gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2,$$

$$(4.39) \quad q_{\sharp}(\kappa_1, \kappa_2) \equiv \overline{q(\overline{\kappa_1}, \overline{\kappa_2})} = \bar{\gamma}(\kappa_1^2 - \kappa_2^2) + 2\bar{\beta}\kappa_1\kappa_2.$$

Calculating the determinant of $\widetilde{\mathcal{M}}(\nu, \kappa)$ we obtain

$$(4.40) \quad D(\nu, \kappa) \equiv \det \left(\widetilde{\mathcal{M}}(\nu, \kappa) \right) = \nu^2 - q(\kappa) q_{\sharp}(\kappa) + g(\nu, \kappa),$$

where $g(\nu, \kappa)$ and hence $D(\nu, \kappa)$ are analytic in a neighborhood of the origin in $\mathbb{C} \times \mathbb{C}^2$. Note also that $g(\nu, \kappa)$ and $\partial_{\nu} g(\nu, \kappa)$ satisfy the following bounds for $|\kappa|$ and $|\nu|$ small:

$$(4.41) \quad |g(\nu, \kappa)| \leq C_g (|\kappa|^5 + |\nu| |\kappa|^3 + |\nu|^2 |\kappa|),$$

$$(4.42) \quad |\partial_{\nu} g(\nu, \kappa)| \leq C'_g (|\kappa|^3 + |\nu| |\kappa|)$$

for some positive constants C_g and C'_g .

The problem of finding eigenvalues $\mu = \mu_S + \nu$ near μ_S for $\mathbf{k} = \mathbf{M} + \kappa$ near \mathbf{M} has been reduced to the study of the solutions to the equation

$$D(\nu, \kappa) = 0$$

for κ near $(0, 0) \in \mathbb{R}_{\kappa}^2$. We shall study the roots of $D(\nu, \kappa)$ using Rouché's theorem.

Consider $\kappa \in \mathbb{R}^2$ such that $|\kappa| < \kappa_{\max}$. We shall eventually take κ_{\max} to be small. For such κ we have

$$\left| \nu^2 - q(\kappa) q_{\sharp}(\kappa) \right| \geq \nu^2 - C_{\gamma, \beta} |\kappa_{\max}|^2,$$

where $C_{\gamma, \beta}$ is a positive constant depending only on γ and β . Note also that for ν constrained to the circle $|\nu| = 2C_{\gamma, \beta} \kappa_{\max}^2$ we have the lower bound:

$$(4.43) \quad \left| \nu^2 - q(\kappa) q_{\sharp}(\kappa) \right| \geq C_{\gamma, \beta} |\kappa_{\max}|^2.$$

Thus, if $|\nu| = 2C_{\gamma, \beta} \kappa_{\max}^2$, then

$$|g(\nu, \kappa)| \leq C_{\gamma, \beta, g} \kappa_{\max}^5.$$

Note also that $C_{\gamma, \beta} \kappa_{\max}^2 > C_{\gamma, \beta, g} \kappa_{\max}^5$ provided $\kappa_{\max}^3 < C_{\gamma, \beta} / C_{\gamma, \beta, g}$. Therefore, if we choose κ_{\max} to be any constant satisfying

$$(4.44) \quad 0 < \kappa_{\max} < \frac{1}{2} (C_{\gamma, \beta} / C_{\gamma, \beta, g})^{\frac{1}{3}},$$

then for $|\kappa| \leq \kappa_{\max}$ we have

$$(4.45) \quad |\nu| = 2C_{\gamma, \beta} \kappa_{\max}^2 \implies |g(\nu, \kappa)| < \left| \nu^2 - q(\kappa) q_{\sharp}(\kappa) \right|$$

for all $|\nu| = 2C_{\gamma, \beta} \kappa_{\max}^2$. Therefore by Rouché's theorem, the functions

$$\nu^2 - q(\kappa) q_{\sharp}(\kappa) \quad \text{and} \quad D(\nu, \kappa) = \nu^2 - q(\kappa) q_{\sharp}(\kappa) + g(\nu, \kappa)$$

have the same number of zeros in the disc: $|\nu| < 2C_{\gamma, \beta} \kappa_{\max}^2$. We denote these zeros: $\nu_+(\kappa)$ and $\nu_-(\kappa)$. For real κ these zeros are real by self-adjointness and we have

$$\nu_+(\kappa), \nu_-(\kappa) \in [-C_{\gamma, \beta} \kappa_{\max}^2, C_{\gamma, \beta} \kappa_{\max}^2], \quad |\kappa| < \kappa_{\max}, \quad \kappa \in \mathbb{R}^2.$$

Next, observe by a residue calculation that for $l = 1, 2$,

$$(4.46) \quad (\nu_+(\kappa))^l + (\nu_-(\kappa))^l = \frac{1}{2\pi i} \int_{|\nu|=2C_{\gamma,\beta}\kappa_{\max}^2} \frac{\nu^l \partial_\nu D(\nu, \kappa)}{D(\nu, \kappa)} d\nu.$$

Since $\partial_\nu D(\nu, \kappa) = 2\nu + \partial_\nu g(\nu, \kappa)$, we have

$$(4.47)$$

$$\begin{aligned} & (\nu_+(\kappa))^l + (\nu_-(\kappa))^l \\ &= \frac{1}{2\pi i} \int_{|\nu|=2C_{\gamma,\beta}\kappa_{\max}^2} \frac{2\nu^{l+1}}{\nu^2 - q(\kappa) q_{\mathfrak{h}}(\kappa) + g(\nu, \kappa)} d\nu \\ & \quad + \frac{1}{2\pi i} \int_{|\nu|=2C_{\gamma,\beta}\kappa_{\max}^2} \frac{\nu^l \partial_\nu g(\nu, \kappa)}{\nu^2 - q(\kappa) q_{\mathfrak{h}}(\kappa) + g(\nu, \kappa)} d\nu \\ &= \frac{1}{2\pi i} \int_{|\nu|=2C_{\gamma,\beta}\kappa_{\max}^2} \frac{2\nu^{l+1}}{\nu^2 - q(\kappa) q_{\mathfrak{h}}(\kappa)} d\nu \\ & \quad - \frac{1}{2\pi i} \int_{|\nu|=2C_{\gamma,\beta}\kappa_{\max}^2} \frac{2\nu^{l+1} g(\nu, \kappa)}{(\nu^2 - q(\kappa) q_{\mathfrak{h}}(\kappa) + g(\nu, \kappa)) \cdot (\nu^2 - q(\kappa) q_{\mathfrak{h}}(\kappa))} d\nu \\ & \quad + \frac{1}{2\pi i} \int_{|\nu|=2C_{\gamma,\beta}\kappa_{\max}^2} \frac{\nu^l \partial_\nu g(\nu, \kappa)}{\nu^2 - q(\kappa) q_{\mathfrak{h}}(\kappa) + g(\nu, \kappa)} d\nu. \end{aligned} \quad (4.48)$$

We can use the identity, for $r > a$,

$$(4.49) \quad \frac{1}{2\pi i} \int_{|z|=r} \frac{z^{l+1}}{z^2 - a^2} dz = \begin{cases} 0 & \text{if } l = 1, \\ a^2 & \text{if } l = 2 \end{cases}$$

to evaluate the first integral and bound the remaining two integrals using (4.41) and (4.42). This gives

$$(4.50) \quad (\nu_+(\kappa))^l + (\nu_-(\kappa))^l = \begin{cases} \mathcal{O}(\kappa_{\max}^4 |\kappa|), & l = 1, \\ 2q(\kappa) q_{\mathfrak{h}}(\kappa) + \mathcal{O}(\kappa_{\max}^6 |\kappa|), & l = 2, \end{cases}$$

for κ_{\max} sufficiently small.

Now note

$$(4.51) \quad \nu_+(\kappa) \cdot \nu_-(\kappa) = \frac{1}{2} (\nu_+(\kappa) + \nu_-(\kappa))^2 - \frac{1}{2} [(\nu_+(\kappa))^2 + (\nu_-(\kappa))^2].$$

Therefore, $\kappa \mapsto \nu_+(\kappa) \cdot \nu_-(\kappa)$ is analytic in a \mathbb{C}^2 neighborhood of $\kappa = 0$. Moreover, we have

$$\begin{aligned} \nu_+(\kappa) \cdot \nu_-(\kappa) &= -\frac{1}{2} (2q(\kappa) q_{\mathfrak{h}}(\kappa) + \mathcal{O}(\kappa_{\max}^6 |\kappa|)) + \mathcal{O}(\kappa_{\max}^8 |\kappa|^2) \\ (4.52) \quad &= -q(\kappa) q_{\mathfrak{h}}(\kappa) + \mathcal{O}(\kappa_{\max}^6 |\kappa|). \end{aligned}$$

Consider now the equation $(\nu - \nu_+(\kappa))(\nu - \nu_-(\kappa)) = 0$ satisfied by $\nu = \nu_{\pm}(\kappa)$.

LEMMA 4.18. *The roots of $D(\nu, \kappa) = 0$ in the disc $|\kappa| < 2C_{\gamma, \beta} \kappa_{\max}^2$ coincide with the roots of the quadratic equation $\nu^2 - t(\kappa)\nu + d(\kappa) = 0$, where $t(\kappa)$ and $d(\kappa)$ are analytic and*

$$\begin{aligned} t(\kappa) &= \nu_+(\kappa) + \nu_-(\kappa) = \mathcal{O}(\kappa_{\max}^4 |\kappa|), \\ d(\kappa) &= \nu_+(\kappa) \cdot \nu_-(\kappa) = -q(\kappa) q_{\mathfrak{h}}(\kappa) + \mathcal{O}(\kappa_{\max}^6 |\kappa|). \end{aligned}$$

Solving for ν we have, for $|\kappa| < 2C_{\gamma, \beta} \kappa_{\max}^2$, two roots:

$$(4.53) \quad \nu_{\pm}(\kappa) = \frac{1}{2}t(\kappa) \pm \sqrt{-d(\kappa) + \frac{1}{4}t^2(\kappa)}$$

$$(4.54) \quad = c_0(\kappa) \pm \sqrt{q(\kappa) q_{\mathfrak{h}}(\kappa) + c_1(\kappa)},$$

where $c_0(\kappa)$ and $c_1(\kappa)$ are analytic in κ in a \mathbb{C}^2 neighborhood of the origin satisfying

$$|c_0(\kappa)| \lesssim \kappa_{\max}^4 |\kappa|, \quad |c_1(\kappa)| \lesssim \kappa_{\max}^6 |\kappa|$$

for all $|\kappa| \leq \kappa_{\max}$ and κ_{\max} is any constant satisfying (4.44).

Since κ can be taken arbitrary and $|\kappa| = \kappa_{\max}$ can be an arbitrarily small positive number, it follows that the analytic functions $c_0(\kappa) = \mathcal{Q}_5(\kappa)$ and $c_1(\kappa) = \mathcal{Q}_7(\kappa)$ which satisfy, for $|\kappa| \rightarrow 0$, $\mathcal{Q}_r(\kappa) = \mathcal{O}(|\kappa|^r)$.

If we now restrict to real $\kappa = (\kappa_1, \kappa_2)$; then from (4.38)–(4.39) we have that

$$(4.55) \quad q(\kappa) q_{\mathfrak{h}}(\kappa) = |q(\kappa)|^2 = \left| \gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2 \right|^2.$$

Therefore, $\nu_{\pm}(\kappa) = \mathcal{Q}_5(\kappa) \pm \sqrt{|\gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2|^2 + \mathcal{Q}_7(\kappa)}$.

We finally return to (4.33), which relates ν to our eigenvalue parameter μ : $\mu(\mathbf{M} + \kappa) = (1 - \alpha)|\kappa|^2 + \nu(\kappa)$. Proposition 4.17 implies constraints, due to symmetry, on the mappings $\mathbf{k} \mapsto \mu_{\pm}(\mathbf{k})$ for \mathbf{k} near \mathbf{M} . Clearly $|q(\kappa)|^2$ is invariant under the $\pi/2$ rotation, $(\kappa_1, \kappa_2) \mapsto (-\kappa_2, \kappa_1)$, and by part 1 of Proposition 4.17 we must have $c_0(\kappa) = \mathcal{Q}_6(\kappa)$ and $c_1(\kappa) = \mathcal{Q}_8(\kappa)$. We therefore have

$$(4.56) \quad \mu_{\pm}(\mathbf{M} + \kappa) = (1 - \alpha)|\kappa|^2 + \mathcal{Q}_6(\kappa) \pm \sqrt{\left| \gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2 \right|^2 + \mathcal{Q}_8(\kappa)}.$$

This completes the proof of Theorem 4.1. \square

4.4. Dispersion surfaces near \mathbf{M} for v admissible and reflection invariant; proof of Corollary 4.2. In addition to V being admissible in the sense of Definition 2.3, we now assume that $V(x_1, x_2)$ is reflection invariant about the line $x_1 = x_2$.

LEMMA 4.19. *Suppose $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $f \in L^2(\Omega)$ satisfies $T_{\rho}[f](\mathbf{x}) \equiv f(\rho^*\mathbf{x}) = f(\mathbf{x})$, where $\rho = \rho^*$ is the reflection (permutation) matrix mapping $(x_1, x_2) \rightarrow (x_2, x_1)$. Then,*

$$(4.57) \quad \partial_{x_n} T_{\rho}[f](\mathbf{x}) = \rho_{nm} T_{\rho}[\partial_{y_m} f](\mathbf{x}) = T_{\rho}[\rho_{nm} \partial_{y_m} f](\mathbf{x}).$$

Proof. Let $\mathbf{y} = \rho^*\mathbf{x}$. Then $\mathbf{y}_m = \rho_{nm}\mathbf{x}_n$, and furthermore $\frac{\partial \mathbf{y}_m}{\partial \mathbf{x}_n} = \rho_{nm}$. Thus,

$$\begin{aligned} \{\nabla \rho[f](\mathbf{x})\}_n &= \frac{\partial}{\partial x_n} f(\rho\mathbf{x}) = \rho_{nm} \frac{\partial}{\partial y_m} f(\mathbf{y}) \Big|_{\mathbf{y}=\rho^*\mathbf{x}} = \left\{ \rho[\nabla_{\mathbf{y}} f] \right\}_n \Big|_{\mathbf{y}=\rho^*\mathbf{x}} \\ &= \left\{ T_{\rho}[\rho \nabla f](\mathbf{x}) \right\}_n. \end{aligned}$$

\square

CLAIM 4.20. Let $\sigma \in \{\pm 1, \pm i\}$. If $\psi(\mathbf{x})$ solves the Floquet–Bloch eigenvalue problem with V admissible and reflection-invariant, and $\psi(\mathbf{x}) \in L^2_{\mathbf{M},\sigma}$, then $\tilde{\psi} = \psi(x_2, x_1)$ is also a solution where $\tilde{\psi} \in L^2_{\mathbf{M},\sigma^3}$. That is, ρ maps $L^2_{\mathbf{M},\sigma} \rightarrow L^2_{\mathbf{M},\sigma^3}$.

Proof. First we'll show if $\psi(\mathbf{x}) \in L^2_{\mathbf{M},\sigma}$, then $\tilde{\psi} = \psi(x_2, x_1) \in L^2_{\mathbf{M},\sigma^3}$. Observe that ρ is self-adjoint, and note

$$(4.58) \quad \rho^* R^* \rho = R$$

since

$$R^* \rho = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \rho R.$$

We seek to show

$$\mathcal{R}[\tilde{\psi}] = \sigma^3 \tilde{\psi}$$

or, equivalently, $\mathcal{R}[\rho[\psi]](\mathbf{x}) = \sigma^3 \rho[\psi](\mathbf{x})$. If that holds, then $\tilde{\psi} \in L^2_{\mathbf{M},\sigma^3}$.

$$\begin{aligned} \mathcal{R}[\tilde{\psi}] &= \mathcal{R}[\rho[\psi]](\mathbf{x}) \\ &= \psi(R^* \rho \mathbf{x}) \stackrel{(4.58)}{=} \psi(\rho R \mathbf{x}) = \psi(\rho(R^*)^3 \mathbf{x}) \\ &= \rho(\psi((R^*)^3 \mathbf{x})) = \rho(\mathcal{R}^3 \psi(\mathbf{x})) \\ &= \sigma^3 \rho[\psi](\mathbf{x}) = \sigma^3 \tilde{\psi}. \end{aligned}$$

Therefore, if $\psi(\mathbf{x}) \in L^2_{\mathbf{M},\sigma}$, then $\tilde{\psi} = \psi(x_2, x_1) \in L^2_{\mathbf{M},\sigma^3}$. \square

CLAIM 4.21. $\gamma = 4a_{1,1}^{1,2} = 0$.

Proof.

$$\begin{aligned} \kappa^T A^{j_1, j_2} \kappa &= \langle \partial_{y_l} \Phi_{j_1}, \mathcal{R}(\mu_*) \partial_{y_m} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{y}})} \kappa_l \kappa_m \\ &= \langle \mathcal{R}(\rho[\partial_{y_l} \Phi_{j_1}]), \mathcal{R}(\mu_*) \mathcal{R}(\rho[\partial_{y_m} \Phi_{j_2}]) \rangle_{L^2(\Omega_{\mathbf{x}})} \kappa_l \kappa_m \\ &= \langle R_{ns} \rho_{ln} \partial_{x_s} \mathcal{R}[\rho \Phi_{j_1}], \mathcal{R}(\mu_*) R_{qt} \rho_{mq} \partial_{x_t} \mathcal{R}[\rho \Phi_{j_2}] \rangle_{L^2(\Omega_{\mathbf{x}})} \kappa_l \kappa_m \\ &= \langle \partial_{x_s} (i^{2j_1-1})^3 \Phi_{j_1}, \mathcal{R}(\mu_*) \partial_{x_q} (i^{2j_2-1})^3 \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{x}})} R_{ns}(\rho_{ln} \kappa_l) R_{qt}(\rho_{mq} \kappa_m) \\ &= \left(i^{2(j_2-j_1)} \right)^3 \langle \partial_{x_s} \Phi_{j_1}, \mathcal{R}(\mu_*) \partial_{x_t} \Phi_{j_2} \rangle_{L^2(\Omega_{\mathbf{x}})} R_{ns}(\rho \kappa)_n R_{qt}(\rho \kappa)_q \\ &= \left(i^{2(j_2-j_1)} \right)^3 (R \rho \kappa)^T \kappa A^{j_1, j_2} R \rho \kappa \end{aligned}$$

for any choice of pairs (j_1, j_2) with $j_1, j_2 \in \{1, 2\}$. Since κ is arbitrary, $A^{j_1, j_2} = (i^{2(j_2-j_1)})^3 \rho R A^{j_1, j_2} R^T \rho$.

Now for any pair $j_1, j_2 \in \{1, 2\}$, letting $A^{j_1, j_2} = A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we have shown

$$(4.59) \quad R A R^T = \begin{pmatrix} d & -c \\ -b & a \end{pmatrix}.$$

Consider $j_1 = 1$ and $j_2 = 2$. From the above analysis, $A^{1,2} = -\rho R A^{1,2} R^T \rho$, so

$$\begin{aligned} -\rho R A^{1,2} R^T \rho &= -\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ &= -\begin{pmatrix} -b & a \\ d & -c \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ &= -\begin{pmatrix} a & -b \\ -c & d \end{pmatrix} = \begin{pmatrix} -a & b \\ c & -d \end{pmatrix}. \end{aligned}$$

Therefore, $A_{1,1}^{1,2} = -A_{1,1}^{1,2}$, and $A_{2,2}^{1,2} = A_{2,2}^{1,2}$, so that both entries must be zero. Using the previous notation,

$$a_{1,1}^{1,2} = a_{2,2}^{1,2} = 0 \implies \gamma = 4a_{1,1}^{1,2} = 0.$$

Corollary 4.2 is now an immediate consequence of Part 2 of Proposition 4.17. \square

5. Spectral band degeneracies for small amplitude potentials. In this section we apply Theorem 4.1 to the case of small amplitude potentials. Our analysis follows that of section 6 in [9] in the case of honeycomb potentials. We consider the Floquet–Bloch eigenvalue problem:

$$(5.1) \quad H^\varepsilon \Phi(\mathbf{x}) \equiv (-\Delta + \varepsilon V(\mathbf{x})) \Phi(\mathbf{x}) = \mu \Phi(\mathbf{x}), \quad \Phi \in L_{\mathbf{M},\sigma}^2,$$

where $\sigma \in \{+1, -1, +i, -i\}$, and ε is small and nonzero.

By Proposition 2.13, we may seek an $L_{\mathbf{M},\sigma}^2$ eigenstate of the form

$$\Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c_\Phi(\mathbf{m}) \left(\sigma^4 e^{i\mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} + \sigma^3 e^{iR\mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} + \sigma^2 e^{iR^2\mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} + \sigma e^{iR^3\mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} \right).$$

We use the notation $c(\mathbf{m}) = c_\Phi(\mathbf{m}) = c(\mathbf{m}; \Phi)$. Applying $(-\Delta - \mu)$ to Φ and using that R is an orthogonal matrix yields

$$(-\Delta - \mu) \Phi = \sum_{\mathbf{m} \in \mathcal{S}} \left(|\mathbf{M}^{\mathbf{m}}|^2 - \mu \right) c(\mathbf{m}; \Phi) \left(\sum_{i=0}^3 \sigma^{4-i} e^{iR^i \mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} \right).$$

Since $V(R^* \mathbf{x}) = V(\mathbf{x})$, we have that $V(\mathbf{x}) \Phi(\mathbf{x}) \in L_{\mathbf{M},\sigma}^2$. Therefore, by Proposition 2.13, $V\Phi$ has an expansion

$$(5.2) \quad V(\mathbf{x}) \Phi(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} \left[\sum_{j=0}^4 \sigma^{4-j} e^{iR^j \mathbf{M}^{\mathbf{m}} \cdot \mathbf{x}} \right] c(\mathbf{m}; V\Phi),$$

where

$$(5.3) \quad c(\mathbf{m}; V\Phi) = \frac{1}{|\Omega|} \int_{\Omega} e^{-i\mathbf{M}^{\mathbf{m}} \cdot \mathbf{y}} V(\mathbf{y}) \Phi(\mathbf{y}) d\mathbf{y}.$$

Recall that $\mathbf{q}\vec{\mathbf{k}} \cdot \mathbf{x} = (q_1 \mathbf{k}_1 + q_2 \mathbf{k}_2) \cdot \mathbf{x}$ and $R\mathbf{M}^{\mathbf{r}} = \mathbf{M}^{\mathcal{R}\mathbf{r}}$; see (2.16). Thus,

$$\begin{aligned}
 c(\mathbf{m}; V\Phi) &= \frac{1}{|\Omega|} \int_{\Omega} e^{-i\mathbf{M}^{\mathbf{m}} \cdot \mathbf{y}} V(\mathbf{y}) \Phi(\mathbf{y}) d\mathbf{y} \\
 &= \frac{1}{|\Omega|} \int_{\Omega} e^{-i\mathbf{M}^{\mathbf{m}} \cdot \mathbf{y}} \left(\sum_{\mathbf{q} \in \mathbb{Z}^2} V_{\mathbf{q}} e^{i\mathbf{q}\vec{\mathbf{k}} \cdot \mathbf{y}} \right) \left(\sum_{\mathbf{r} \in \mathcal{S}} c(\mathbf{r}; \Phi) \left[\sum_{j=0}^3 \sigma^{4-j} e^{iR^j \mathbf{M}^{\mathbf{r}} \cdot \mathbf{y}} \right] \right) d\mathbf{y} \\
 &= \frac{1}{|\Omega|} \sum_{\mathbf{q} \in \mathbb{Z}^2} \sum_{\mathbf{r} \in \mathcal{S}} \int_{\Omega} V_{\mathbf{q}} c(\mathbf{r}; \Phi) \left[\sum_{j=0}^3 \sigma^{4-j} e^{i(\mathbf{M}^{\mathcal{R}^j \mathbf{r}} - \mathbf{M}^{\mathbf{m}} + \mathbf{q}\vec{\mathbf{k}}) \cdot \mathbf{y}} \right] d\mathbf{y} \\
 &= \frac{1}{|\Omega|} \sum_{\mathbf{q} \in \mathbb{Z}^2, \mathbf{r} \in \mathcal{S}} V_{\mathbf{q}} c(\mathbf{r}; \Phi) \int_{\Omega} \sum_{j=0}^3 \left(\sigma^{4-j} e^{i(\mathbf{q} - (\mathbf{m} - \mathcal{R}^j \mathbf{r})) \vec{\mathbf{k}} \cdot \mathbf{y}} \right) d\mathbf{y} \\
 &= \sum_{\mathbf{q} \in \mathbb{Z}^2, \mathbf{r} \in \mathcal{S}} V_{\mathbf{q}} c(\mathbf{r}; \Phi) \sum_{j=0}^3 \left[\sigma^{4-j} \delta(\mathbf{q} - (\mathbf{m} - \mathcal{R}^j \mathbf{r})) \right] \\
 &= \sum_{\mathbf{r} \in \mathcal{S}} \left[\sum_{j=0}^3 \sigma^{4-j} V_{\mathbf{m} - \mathcal{R}^j \mathbf{r}} \right] c(\mathbf{r}; \Phi) \\
 &= \sum_{\mathbf{r} \in \mathcal{S}} \left(V_{\mathbf{m} - \mathbf{r}} + \sigma^3 V_{\mathbf{m} - \mathcal{R}\mathbf{r}} + \sigma^2 V_{\mathbf{m} - \mathcal{R}^2 \mathbf{r}} + \sigma V_{\mathbf{m} - \mathcal{R}^3 \mathbf{r}} \right) c(\mathbf{r}; \Phi).
 \end{aligned}$$

Thus,

$$(5.4) \quad c(\mathbf{m}; V\Phi) = \sum_{\mathbf{r} \in \mathcal{S}} \mathcal{K}_{\sigma}(\mathbf{m}, \mathbf{r}) c(\mathbf{r}; \Phi).$$

$$\begin{aligned}
 (5.5) \quad K_{\sigma}(\mathbf{m}, \mathbf{r}) &= \sum_{j=0}^3 \sigma^{4-j} V_{\mathbf{m} - \mathcal{R}^j \mathbf{r}} \\
 &= V_{\mathbf{m} - \mathbf{r}} + \sigma^3 V_{\mathbf{m} - \mathcal{R}\mathbf{r}} + \sigma^2 V_{\mathbf{m} - \mathcal{R}^2 \mathbf{r}} + \sigma V_{\mathbf{m} - \mathcal{R}^3 \mathbf{r}} \\
 &= V_{m_1 - r_1, m_2 - r_2} + \sigma^3 V_{m_1 - r_2, m_2 + r_1 + 1} \\
 &\quad + \sigma^2 V_{m_1 + r_1 + 1, m_2 + r_2 + 1} + \sigma V_{m_1 + r_2 + 1, m_2 - r_1}.
 \end{aligned}$$

We have the following analogue of Proposition 6.1 of [9].

PROPOSITION 5.1. *Let $\sigma \in \{+i, -i, +1, -1\}$. The $L^2_{\mathbf{M}, \sigma}$ -spectral problem (5.1) is equivalent to the following algebraic eigenvalue problem for $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}}$ and μ :*

$$(5.6) \quad \left(|\mathbf{M}^{\mathbf{m}}|^2 - \mu \right) c(\mathbf{m}) + \varepsilon \sum_{\mathbf{r} \in \mathcal{S}} \mathcal{K}_{\sigma}(\mathbf{m}, \mathbf{r}) c(\mathbf{r}) = 0, \quad \mathbf{m} \in \mathcal{S}.$$

For each σ , we next solve (5.6) for $\varepsilon \mapsto \{c^{\varepsilon}(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}}$, μ^{ε} , perturbatively in ε .

We first set $\varepsilon = 0$ in (5.6). Then, (5.6) reduces to the eigenvalue problem

$$(5.7) \quad \left(|\mathbf{M} + m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2|^2 - \mu^{(0)} \right) c^{(0)}(\mathbf{m}) = 0, \quad \mathbf{m} = (m_1, m_2) \in \mathcal{S},$$

which has a simple eigenpair:

$$(5.8) \quad \mu^{(0)} = |\mathbf{M}|^2 = 2\pi^2 = \mu_S^{(0)}, \quad c^{(0)}(\mathbf{m}) = \delta_{m_1+1, m_2},$$

In arriving at (5.8), recall from Theorem 3.1 that $\mu_S^{(0)} = |\mathbf{M}|^2 = 2\pi^2$ is an eigenvalue of multiplicity four with four-dimensional eigenspace spanned by the eigenvectors $c(m_1, m_2) = \delta_{m_1, m_2}$, δ_{m_1, m_2+1} , δ_{m_1+1, m_2+1} , and δ_{m_1+1, m_2} , corresponding to those

(m_1, m_2) such that $\mathbf{M}^{\mathbf{m}} = \mathbf{M} + m_1 \mathbf{k}_1 + m_2 \mathbf{k}_2$ is a vertex of \mathcal{B} and to the orbit under $\mathcal{R} : \{(0, 0), (0, -1), (-1, -1), (-1, 0)\} \in \mathbb{Z}^2 / \sim$. Recall from Remark 2.11 that $(m_1, m_2) = (-1, 0)$ is the representative from this equivalence class and hence the choice of eigenstate in (5.8).

Note also that the solution (5.8) of the $\varepsilon = 0$ algebraic eigenvalue problem corresponds the simple $L^2_{\mathbf{M}, \sigma}$ -eigenpair of $H^{(\varepsilon=0)}$, $\mu^{(0)} = |\mathbf{M}^{(0, -1)}|^2 = |\mathbf{M}|^2$, with corresponding eigenstate

$$\begin{aligned} \Phi^{(\varepsilon=0)}(\mathbf{x}) &= \sum_{j=0}^3 \sigma^{4-j} e^{i(R^j \mathbf{M}^{0, -1}) \cdot \mathbf{x}} \\ (5.9) \quad &= \sigma e^{i\mathbf{M} \cdot \mathbf{x}} \left(1 + \sigma e^{-i\mathbf{k}_1 \cdot \mathbf{x}} + \sigma^2 e^{-i(\mathbf{k}_1 + \mathbf{k}_2) \cdot \mathbf{x}} + \sigma^3 e^{-i\mathbf{k}_2 \cdot \mathbf{x}} \right). \end{aligned}$$

We next proceed to solve the system (5.6) for a smooth curve of eigenpairs: $\varepsilon \mapsto \mu^\varepsilon$, $\{c^\varepsilon(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}}$ for all ε sufficiently small. We write $\{c(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}} \in l^2(\mathcal{S}) = (c_{\parallel}, c_{\perp}) \in \mathbb{C} \times l^2(\mathcal{S}^{\perp})$, where

$$c_{\parallel} \equiv c(0, -1), \quad \text{and} \quad c_{\perp} = \{c_{\perp}(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}^{\perp}}, \quad \mathcal{S}^{\perp} \equiv \mathcal{S} \setminus \{(-1, 0)\}.$$

Then (5.6) is equivalent to the following coupled system for c_{\parallel} and $c_{\perp} = \{c_{\perp}(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}^{\perp}}$:

$$(5.10) \quad [|\mathbf{M}|^2 - \mu + \varepsilon \mathcal{K}_{\sigma}(0, -1, 0, -1)] c_{\parallel} + \varepsilon \sum_{\mathbf{r} \in \mathcal{S}^{\perp}} \mathcal{K}_{\sigma}(0, -1, \mathbf{r}) c_{\perp}(\mathbf{r}) = 0,$$

$$(5.11) \quad \varepsilon \mathcal{K}_{\sigma}(\mathbf{m}, 0, -1) c_{\parallel} + (|\mathbf{M}^{\mathbf{m}}|^2 - \mu) c_{\perp}(\mathbf{m}) + \varepsilon \sum_{\mathbf{r} \in \mathcal{S}^{\perp}} \mathcal{K}_{\sigma}(\mathbf{m}, \mathbf{r}) c_{\perp}(\mathbf{r}) = 0, \quad \mathbf{m} \in \mathcal{S}^{\perp}.$$

For ε small, we shall solve (5.10)–(5.11) in a neighborhood of the solution to the $\varepsilon = 0$ problem: $c_{\parallel}^{(0)} = 1$, $\mu_S^{(0)} = |\mathbf{M}|^2$, $c_{\perp}^{(0)}(\mathbf{r}) = 0$, $\mathbf{r} \in \mathcal{S}^{\perp}$. We proceed by a Lyapunov–Schmidt reduction strategy in which we first solve (5.11) for the mapping $c_{\parallel} \mapsto c_{\perp}[c_{\parallel}, \mu]$ in a neighborhood of $\mu = \mu_S^{(0)} = |\mathbf{M}|^2$, and then substitute this mapping into (5.10) to obtain a closed equation for c_{\parallel} . We provided a sketch of the argument. For the details, see [9].

Equation (5.11) for c_{\perp} may be expressed in the form

$$(5.12) \quad (I + \varepsilon \mathcal{T}_{\mathcal{K}_{\sigma}}(\mu)) c_{\perp} = \varepsilon c_{\parallel} F_{\sigma}(\mu)$$

with the definitions $F_{\sigma}(\mu) = \{F_{\sigma}(\mathbf{m}, \mu)\}_{\mathbf{m} \in \mathcal{S}^{\perp}}$, $F_{\sigma}(\mathbf{m}, \mu) \equiv \frac{-\mathcal{K}_{\sigma}(\mathbf{m}, 0, -1)}{|\mathbf{M}^{\mathbf{m}}|^2 - \mu}$, and

$$(5.13) \quad [(I + \varepsilon \mathcal{T}_{\mathcal{K}_{\sigma}}(\mu)) c_{\perp}](\mathbf{m}) \equiv \left[\delta_{\mathbf{m}, \mathbf{r}} + \frac{\varepsilon}{|\mathbf{M}^{\mathbf{m}}|^2 - \mu} \sum_{\mathbf{r} \in \mathcal{S}^{\perp}} \mathcal{K}_{\sigma}(\mathbf{m}, \mathbf{r}) \right] c_{\perp}(\mathbf{r}), \quad \mathbf{m} \in \mathcal{S}^{\perp}.$$

For all μ in a fixed neighborhood of $\mu^{(0)} = |\mathbf{M}|^2$, we have for all $\mathbf{m} \in \mathcal{S}^{\perp}$ that $||\mathbf{M}^{\mathbf{m}}|^2 - \mu| \geq \theta > 0$ independent of ε . It follows that for all $|\varepsilon| < \varepsilon^0$ sufficiently small, $(I + \varepsilon \mathcal{T}_{\mathcal{K}_{\sigma}}(\mu))^{-1}$ is well-defined as a bounded operator on $l^2(\mathcal{S}^{\perp})$. Solving (5.12) for $c_{\perp}^{\varepsilon}[c_{\parallel}, \mu]$ and substituting into (5.10) we conclude as follows.

PROPOSITION 5.2. For all $|\varepsilon| < \varepsilon^0$, μ is an $L^2_{\mathbf{M},\sigma}$ eigenvalue if and only if $\mathcal{M}_\sigma(\mu, \varepsilon) = 0$, where

$$(5.14) \quad \begin{aligned} \mathcal{M}_\sigma(\mu, \varepsilon) \equiv & |\mathbf{M}|^2 - \mu + \varepsilon \mathcal{K}_\sigma(0, -1, 0, -1) \\ & + \varepsilon^2 \sum_{\mathbf{r} \in \mathcal{S}^\perp} \mathcal{K}_\sigma(0, -1, \mathbf{r}) [(I + \varepsilon \mathcal{T}_{\mathcal{K}_\sigma}(\mu))^{-1} F_\sigma(\mathbf{m}, \mu)](\mathbf{r}) \end{aligned}$$

is analytic in a neighborhood about $(\varepsilon, \mu) = (0, \mu_S^{(0)}) = (0, |\mathbf{M}|^2)$.

Since $\mathcal{M}_\sigma(\mu_S^{(0)}, 0) = 0$ and $\partial_\mu \mathcal{M}_\sigma(\mu_S^{(0)}, 0) = -1 \neq 0$, the implicit function theorem, implies that there is a function $\varepsilon \rightarrow \mu^\varepsilon$, defined and analytic for $|\varepsilon| < \varepsilon^1 \leq \varepsilon^0$, such that $\mathcal{M}_\sigma(\mu^\varepsilon, \varepsilon) = 0$. Thus taking $c_\parallel^\varepsilon \equiv 1$, the solution of the coupled system (5.10)–(5.11), for $|\varepsilon| < \varepsilon^1$, is

$$(5.15) \quad \mu_S^\varepsilon = |\mathbf{M}|^2 + \varepsilon \mathcal{K}_\sigma(0, -1, 0, -1) + \mathcal{O}(\varepsilon^2),$$

$$c_\parallel^\varepsilon = c(0, -1) \equiv 1,$$

$$(5.16) \quad c_\perp^\varepsilon = \{c^\varepsilon(\mathbf{m})\}_{\mathbf{m} \in \mathcal{S}^\perp} = \varepsilon(I + \varepsilon \mathcal{T}_{\mathcal{K}_\sigma}(\mu))^{-1} F_\sigma(\mathbf{m}, \mu^\varepsilon), \quad \mathbf{m} \in \mathcal{S}^\perp,$$

$$\text{where } F_\sigma(\mathbf{m}, \mu^\varepsilon) = \frac{-\mathcal{K}_\sigma(\mathbf{m}, 0, -1)}{|\mathbf{M}^\mathbf{m}|^2 - \mu^\varepsilon}.$$

From (5.15) we may now make explicit the splitting of the fourfold degenerate $L^2_{\mathbf{M}}$ eigenvalue of $H^\varepsilon = -\Delta + \varepsilon V$ for ε nonzero and small. By (5.5) and the relations among the Fourier coefficients of V displayed in (2.6) we have

$$(5.17) \quad \mathcal{K}_\sigma(0, -1, 0, -1) = V_{00} + (\sigma^3 + \sigma) V_{01} + \sigma^2 V_{11}.$$

Therefore,

$$(5.18) \quad \mathcal{K}_{\pm 1}(0, -1, 0, -1) = V_{0,0} \pm 2V_{0,1} + V_{1,1},$$

$$(5.19) \quad \mathcal{K}_{+i}(0, -1, 0, -1) = \mathcal{K}_{-i}(0, -1, 0, -1) = V_{0,0} - V_{1,1}.$$

This establishes that for typical choices of Fourier coefficients—specifically $V_{0,0} \pm 2V_{0,1} + V_{1,1} \neq V_{0,0} - V_{1,1}$ or equivalently $V_{1,1} \neq V_{0,1}$ —and for ε small and nonzero, the multiplicity four $L^2_{\mathbf{M}}$ eigenvalue, $\mu_S^{(0)} = |\mathbf{M}|^2$, splits, at order ε , distinct $L^2_{\mathbf{M},+1}$ and $L^2_{\mathbf{M},-1}$ eigenvalues and an $L^2_{\mathbf{M}}$ -double eigenvalue in the subspace $L^2_{\mathbf{M},+i} \oplus L^2_{\mathbf{M},-i}$. Note in fact that the latter is an exact (to all orders) double eigenvalue in $L^2_{\mathbf{M},+i} \oplus L^2_{\mathbf{M},-i}$. Indeed, consider the simple $L^2_{\mathbf{M},+i}$ eigenvalue, whose existence is guaranteed by the above proof for $\sigma = +i$. Applying $\mathcal{P} \circ \mathcal{C}$ to the corresponding eigenfunction we obtain an eigenfunction in the space $L^2_{\mathbf{M},+i}$ with the identical eigenvalue. This must coincide with the simple eigenvalue constructed for $\sigma = -i$. Summarizing we have the following.

THEOREM 5.3. Consider $H^\varepsilon = -\Delta + \varepsilon V$, where V is admissible (Definition 2.3) and $0 < |\varepsilon| < \varepsilon_1$ is sufficiently small. Assume the nondegeneracy condition on distinguished Fourier coefficients:

$$(5.20) \quad V_{1,1} \neq \pm V_{0,1},$$

where V_{m_1, m_2} denotes the (m_1, m_2) Fourier coefficient of V . Then the four-dimensional eigenspace of H^0 corresponding to the eigenvalue $\mu_S^{(0)} = |\mathbf{M}|^2$ perturbs to a two-dimensional eigenspace with eigenvalue μ_S^ε and additionally two one-dimensional eigenspaces with eigenvalues $\mu_{(+1)}^\varepsilon$ and $\mu_{(-1)}^\varepsilon$ as follows:

1. μ_S^ε is of geometric multiplicity two eigenvalue, with a two-dimensional eigenspace $\mathbb{X}_i \subset L_{\mathbf{M},i}^2$ and $\mathbb{X}_{-i} \subset L_{\mathbf{M},-i}^2$, and has the expansion

$$(5.21) \quad \mu_S^\varepsilon = |\mathbf{M}|^2 + \varepsilon(V_{0,0} - V_{1,1}) + \mathcal{O}(\varepsilon^2).$$

Associated with it are the eigenstates $\Phi_1^\varepsilon \in L_{\mathbf{M},i}^2$ and $\Phi_2^\varepsilon \in L_{\mathbf{M},-i}^2$, related by the symmetry

$$\Phi_2(\mathbf{x}) = (\mathcal{P} \circ \mathcal{C})[\Phi_1](\mathbf{x}) = \overline{\Phi_1(-\mathbf{x})},$$

which we also denote $\Phi_{(+i)}^\varepsilon$ and $\Phi_{(-i)}^\varepsilon$, respectively. Their Fourier expansions are

$$(5.22) \quad \Phi_{(+i)}^\varepsilon = \Phi_1^\varepsilon(\mathbf{x}) = \sum_{\mathbf{m} \in S} c_{(+i)}^\varepsilon(\mathbf{m}) \left(e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - ie^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + ie^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right)$$

and

$$(5.23) \quad \Phi_{(-i)}^\varepsilon = \Phi_2^\varepsilon(\mathbf{x}) = \sum_{\mathbf{m} \in S} \overline{c_{(+i)}^\varepsilon(\mathbf{m})} \left(e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + ie^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - ie^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right).$$

2. The distinct eigenvalues $\mu_{(+1)}^\varepsilon$ and $\mu_{(-1)}^\varepsilon$ are each of geometric multiplicity one, with corresponding one-dimensional eigenspaces $\mathbb{X}_{\pm 1} \subset L_{\mathbf{M},\pm 1}^2$, and they are given by

$$(5.24) \quad \mu_{(\pm 1)}^\varepsilon = |\mathbf{M}|^2 + \varepsilon(V_{0,0} \pm 2V_{0,1} + V_{1,1}) + \mathcal{O}(\varepsilon^2)$$

with associated eigenstates $\Phi_{(\pm 1)}^\varepsilon$:

$$(5.25) \quad \Phi_{(\pm 1)}^\varepsilon(\mathbf{x}) = \sum_{\mathbf{m} \in S} c_{(\pm 1)}^\varepsilon(\mathbf{m}) \left(e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right).$$

Theorems 5.3 and 4.1 imply the following.

COROLLARY 5.4. *Consider the setup of Theorem 5.3. There exists $\varepsilon_1 > 0$ and $\kappa_1(\varepsilon) > 0$, which tends to zero as $\varepsilon \rightarrow 0$, such that the following holds. Fix $\varepsilon \in (-\varepsilon_1, \varepsilon_1) \setminus \{0\}$. Then the following hold:*

1. For all $|\kappa| = \sqrt{\kappa_1^2 + \kappa_2^2} < \kappa_1(\varepsilon)$, the two dispersion surfaces which touch at \mathbf{M} (and therefore the vertices of \mathcal{B}) are locally described by

$$(5.26) \quad \begin{aligned} \mu_\pm^\varepsilon(\mathbf{M} + \kappa) - \mu_S^\varepsilon \\ = (1 - \alpha^\varepsilon)|\kappa|^2 + \mathcal{Q}_6^\varepsilon(\kappa) \pm \sqrt{\left| \gamma^\varepsilon(\kappa_1^2 - \kappa_2^2) + 2\beta^\varepsilon\kappa_1\kappa_2 \right|^2 + \mathcal{Q}_8^\varepsilon(\kappa)}. \end{aligned}$$

The coefficients α^ε , β^ε , and γ^ε are expressions which depend on $\{\Phi_{(+i)}^\varepsilon, \Phi_{(-i)}^\varepsilon\}$ and have the following expansions for ε positive and small:

$$(5.27) \quad \alpha^\varepsilon = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1);$$

$$(5.28) \quad \beta^\varepsilon = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1);$$

$$(5.29) \quad \gamma^\varepsilon = -\frac{32\pi^2}{\varepsilon} i \left(\frac{V_{01}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1).$$

The functions $\mathcal{Q}_6^\varepsilon(\kappa) = \mathcal{O}(|\kappa|^6)$ and $\mathcal{Q}_8^\varepsilon(\kappa) = \mathcal{O}(|\kappa|^8)$ are analytic in a complex neighborhood of the origin in \mathbb{C}^2 ; see Theorem 4.1 for more discussion.

2. Let V be admissible and assume the following (generically satisfied) conditions on Fourier coefficients:

$$V_{11} \neq \pm V_{01}, \quad V_{11} \neq 0 \quad \text{and} \quad V_{01} \neq 0.$$

Then, for all $\varepsilon \in (-\varepsilon_1, \varepsilon_1) \setminus \{0\}$, the coefficients α^ε , β^ε , and γ^ε are all nonzero.

3. In the case where V is admissible and also reflection invariant, by Corollary 4.2, we have $\gamma^\varepsilon = 0$ for any ε , and in particular $V_{01} = 0$. If $V_{11} \neq 0$, then for all $\varepsilon \in (-\varepsilon_1, \varepsilon_1) \setminus \{0\}$, and the coefficients α^ε and β^ε are both nonzero.

The proof of Corollary 5.4 is completed in Appendix C with the derivation of expansions (5.27), (5.28), and (5.29).

6. $H^\varepsilon = -\Delta + \varepsilon V$ for V admissible and ε generic. Theorem 5.3 considers degeneracies among dispersion surfaces for all ε non-zero, real and small. In this section we extend this result to generic real values of ε , with no constraint on its size. Thus, generic large (*high contrast*) potentials are included.

THEOREM 6.1. *Let V denote an admissible potential (Definition 2.3). Let ε_1 be as in Theorem 5.3 and Corollary 5.4. Consider either of two scenarios:*

- (I) V admissible with $V_{11} \neq \pm V_{01}$, $V_{11} \neq 0$ and $V_{01} \neq 0$, or
- (II) V admissible and reflection invariant with $V_{11} \neq 0$.

Then there exists a discrete set $\tilde{\mathcal{C}} \subset \mathbb{R} \setminus (0, \varepsilon_1)$ such that if $\varepsilon \notin \tilde{\mathcal{C}}$, the conditions of Theorem 4.1 are satisfied and two dispersion surfaces touch at the vertices of \mathcal{B} . Moreover, in scenario (I), α^ε , β^ε , and γ^ε are all nonzero for all $\varepsilon \notin \tilde{\mathcal{C}}$, and α^ε and β^ε are both nonzero for all $\varepsilon \notin \tilde{\mathcal{C}}$.

Remark 6.2. For $\varepsilon \in (-\varepsilon_1, \varepsilon_1) \setminus \{0\}$, Theorem 5.3 ensures that pairs of dispersion surfaces, among the first four, touch at band degeneracies for quasi momenta at the vertices of \mathcal{B} ; for the specific scenarios see section 7. For general $\varepsilon \notin \tilde{\mathcal{C}}$ we make no assertions about which of the infinitely many dispersion surfaces touch at high-symmetry quasi momenta. But in analogy with the honeycomb setting studied in [7], we expect for the case of a potential which is a superposition of potential wells that in the strong binding regime there will exist quadratic degeneracies at the intersection of the first two dispersion surfaces.

We discuss the strategy for the proof of Theorem 6.1 but do not present all details. Arguments of this type, rooted in complex analysis, were developed in [9] and Appendix D of [8]. The strategy is based on a continuation argument in the parameter ε , starting with ε varying in the open interval $(0, \varepsilon_1)$; analogous arguments apply to negative values of ε . Eigenvalues, μ , of the operator $H^\varepsilon = -\Delta + \varepsilon V$, in the spaces $L^2_{\mathbf{M}, \sigma}$, for $\sigma = \pm 1, \pm i$, are realized as zeros of a modified Fredholm determinant $\mathcal{E}_\sigma(\mu, \varepsilon)$. The mapping $(\mu, \varepsilon) \mapsto \mathcal{E}_\sigma(\mu, \varepsilon)$ is analytic. For ε real, μ is an $L^2_{\mathbf{M}, \sigma}$ eigenvalue of geometric multiplicity m if and only if μ is a zero of $\mathcal{E}_\sigma(\mu, \varepsilon)$ of multiplicity m . The strategy of [9] (see also Appendix D of [8]) can be used to establish that there is a discrete set $\tilde{\mathcal{C}} \subset \mathbb{R} \setminus (0, \varepsilon_1)$ such that for all $\varepsilon \notin \tilde{\mathcal{C}}$, there exists $\mu_S^\varepsilon \in \mathbb{R}$ such that

1. $\mu = \mu_S^\varepsilon \in \mathbb{R}$ is a simple zero of $\mathcal{E}_{+i}(\mu, \varepsilon)$ and of $\mathcal{E}_{-i}(\mu, \varepsilon)$,
2. $\mathcal{E}_1(\mu_S^\varepsilon, \varepsilon) \neq 0$ and $\mathcal{E}_{-1}(\mu_S^\varepsilon, \varepsilon) \neq 0$,
3. for scenario (I), $\alpha^\varepsilon, \beta^\varepsilon$, and γ^ε are all nonzero and α^ε , and for scenario (II), α^ε and β^ε are nonzero.

Therefore, by Theorem 4.1, for all $\varepsilon \notin \tilde{\mathcal{C}}$ there exist quadratic degeneracies at the quasi-momentum/energy pairs $(\mu_S^\varepsilon, \mathbf{M}_*)$, where \mathbf{M}_* varies over the four vertices of \mathcal{B} . These are locally described by (4.1) of Theorem 4.1; see also (4.2) of Corollary 4.2.

7. Computational experiments. In this section, we describe numerical computations of the spectrum of periodic Schrödinger operators with admissible potentials in the sense of Definition 2.3. We discuss these numerical results in the context of our analytical results. First, in section 7.1, we consider examples of admissible potentials which exhibit quadratic intersections of dispersion surfaces at \mathbf{M} ; see Theorems 4.1, 5.3, and 6.1. In order to observe clear numerical separation of the bands, we work here with ε generically not so small, meaning many of our results fit more into Theorem 6.1 but display many of the effects described in the small ε setting.

In section 7.2, we consider large amplitude potentials and revisit the question posed in section 1.3 regarding Lieb lattice potentials.

To numerically approximate the dispersion surfaces for the Schrödinger operator, we use the periodic formulation of the self-adjoint eigenvalue problem (1.3). For the numerical experiments of section 7.1 we discretized the fundamental period cell, Ω , and used a finite difference approximation to $H_V(\mathbf{k})$ for a fixed $\mathbf{k} \in \mathcal{B}$. In the numerical experiments of section 7.2, we discretized the mapping $f \mapsto [-(\partial_x + ik_x)^2 - (\partial_y + ik_y)^2]f$ in Fourier space and the operator $f \mapsto Vf$ in physical space. For both approaches, we used the MATLAB function `eigs` using the 'sr' flag. For each fixed \mathbf{k} varying over a discretization of an appropriate subset of \mathcal{B} , i.e., the irreducible Brillouin zone or the circuit $\Gamma \rightarrow \mathbf{X} \rightarrow \mathbf{M} \rightarrow \Gamma$ outlined in Figure 1.2, we compute the five smallest eigenvalues.

7.1. Computations of quadratic degeneracy of dispersion surfaces near \mathbf{M} and comparison with Theorem 5.3. We consider a class of periodic potentials, which are the \mathbb{Z}^2 -periodic extension of the function

$$(7.1) \quad V(\mathbf{x}) = \sum_i s_i f(|\mathbf{x} - \mathbf{x}_i|), \quad \mathbf{x} \in \Omega = [0, 1]^2.$$

Here $f(x) = \frac{1}{2}(1 + \cos(\pi x/r))\chi_{\{x < r\}}$ is a compactly supported, C^1 function and $r = 0.2$. The points $\{\mathbf{x}_i\}$ are lattice points within the primitive cell, $[0, 1]^2$. The binary variables, $s_i \in \{+1, -1\}$, determine the sign of the potential at the lattice points. We choose $\{\mathbf{x}_i\}$ and $\{s_i\}$ so that $V(\mathbf{x})$ is an admissible potential; see Definition 2.3. By varying $\{\mathbf{x}_i\}$ and $\{s_i\}$, we show that the combinations of Fourier coefficients V_{00}, V_{01} , and V_{11} ($V_{m,n} = (2\pi)^{-2} \int_0^1 \int_0^1 V(x, y) e^{-2\pi i(mx+ny)} dx dy$), appearing in Theorem 5.3 can be varied in order to exhibit different local behavior of the first four dispersion surfaces near \mathbf{M} . We remark that while Theorem 5.3 describes the dispersion surfaces for the operator $H_V = -\Delta + \varepsilon V$ near the point $\mathbf{k} = \mathbf{M}$ for sufficiently small and non-zero and real ε , our computations are performed for $\varepsilon = O(1)$.

In each of Figures 7.1 to 7.5 we plot, over one period cell, an admissible potential of the form in (7.1) and the first five dispersion curves for $H_V = -\Delta + \varepsilon V$ and compare the numerically computed results with the assertions of Theorem 5.3. For each potential, the lattice sites, $\{\mathbf{x}_i\}$, and signs, $\{s_i\}$, are easily inferred from the plot of the potential. They are also summarized in Table 7.1.

The potentials in Figures 7.1, 7.2, and 7.4 are Lieb lattice potentials (see Example 2.6), while the potential in Figure 7.3 is a square lattice potential (see Example 2.5); see Figure 1.1. The dispersion curves which pass through $\mu_i^\varepsilon, \mu_{-i}^\varepsilon, \mu_1^\varepsilon$, and μ_{-1}^ε are plotted for quasi momentum \mathbf{K} along the cyclic path $\Gamma \rightarrow \mathbf{X} \rightarrow \mathbf{M} \rightarrow \Gamma$ in \mathcal{B} ; see

TABLE 7.1

Figure	ε	\mathbf{x}_i	s_i
7.1	2	$[(0, 0), (1/2, 0), (0, 1/2)]$	$(1, -1, -1)$
7.2	2	$[(0, 0), (1/2, 0), (0, 1/2)]$	$(-1, 1, 1)$
7.3	2	$[(1/2, 1/4), (1/4, 1/2), (3/4, 1/2), (1/2, 3/4)]$	$(1, 1, 1, 1)$
7.4	4	$[(0, 0), (1/2, 0), (0, 1/2)]$	$(-1, -1, -1)$
7.5	2	$[(1/2, 1/2)]$	(-1)

TABLE 7.2

	Fourier coefficient condition	Dispersion surface ordering
(1a)	$V_{1,1} > 0, V_{0,1} < 0$ and $ V_{0,1} < V_{1,1}$	$\mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{+1}^\varepsilon < \mu_{-1}^\varepsilon$
(1b)	$0 < V_{0,1} < V_{1,1}$	$\mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{-1}^\varepsilon < \mu_{+1}^\varepsilon$
(2a)	$V_{1,1}, V_{0,1} < 0$ and $ V_{0,1} > V_{1,1} $	$\mu_{+1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{-1}^\varepsilon$
(2b)	$V_{1,1} < 0$ and $V_{0,1} > V_{1,1} $	$\mu_{-1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{+1}^\varepsilon$
(3a)	$V_{1,1}, V_{0,1} < 0$ and $ V_{0,1} < V_{1,1} $	$\mu_{+1}^\varepsilon < \mu_{-1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon$
(3b)	$V_{1,1} < 0, V_{0,1} > 0$, and $ V_{0,1} < V_{1,1} $	$\mu_{-1}^\varepsilon < \mu_{+1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon$

Figure 1.2. In all dispersion plots, we observe saddle-like touching of the dispersion surfaces as described in Theorem 4.1.

We note that the ordering of the dispersion surfaces, specified by certain Fourier coefficients of the potential, in Theorem 5.3 (which applies to ε sufficiently small) persists for larger (order 1) values of ε . These dispersion surface orderings are summarized in Table 7.2.

Here, we've enumerated the various cases so that, e.g., the multiplicity occurs in the first eigenvalue for cases (1a) and (1b). For larger values of ε this ordering may be violated. However, by Theorem 6.1, quadratic degeneracies in the band structure of $-\Delta + \varepsilon V$ will still occur for all but a discrete set of ε -values.

The potential in Figure 7.1 satisfies condition (1b) and we observe that the first two surfaces intersect at \mathbf{M} , with the others separated and lying above. The potential in Figure 7.2 satisfies condition (3a) and we observe that the third and fourth surfaces intersect at \mathbf{M} , with the others separated and lying below.

In Figure 7.3 the coefficients satisfy (2a), except that $V_{1,1} = 0$. Here, the first and fourth bands are strongly separated from the second and third bands. The latter not only intersect at \mathbf{M} , but coincide on the interval $\mathbf{M} \rightarrow \Gamma$.

In Figure 7.4, the potential consists of potential wells centered on the Lieb lattice sites and satisfies condition (2b). Consequently, we observe an intersection at \mathbf{M} between the second and third surfaces. The coefficients satisfy $V_{0,1} \approx -V_{1,1}$, so the first dispersion surface is quite close to the second and third surfaces at \mathbf{M} but does not touch. The first three bands are separating from the fourth and we observe the emergence of the tight-binding dispersion relation, as shown in Figure 1.3. This will be further investigated in section 7.2; see Figure 7.6.

Finally, in Figure 7.5, we consider a potential that consists of potential wells centered on the square lattice sites and satisfies condition (2a). We see that the second and third bands remaining touching with the fourth band close but lying above. The first band is well separated and lies below. Interestingly, the linear prediction is that $\mu_{+1}^\varepsilon = 2\pi^2 + 0.0005\varepsilon + o(\varepsilon)$, but for this value of ε , we have $\mu_{+1}^\varepsilon < 2\pi^2$ so we are already in a higher order regime. This will be further investigated in section 7.2; see Figure 7.7.




FIG. 7.1. *Eigenvalue ordering, up to $\mathcal{O}(\varepsilon)$: $\mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{-1}^\varepsilon < \mu_{+1}^\varepsilon$.*




FIG. 7.2. *Eigenvalue ordering, up to $\mathcal{O}(\varepsilon)$: $\mu_{+1}^\varepsilon < \mu_{-1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon$.*




FIG. 7.3. *Eigenvalue ordering, up to $\mathcal{O}(\varepsilon)$: $\mu_{+1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{-1}^\varepsilon$.*

7.2. Periodic arrays of deep potential wells; the strong binding regime.

We will consider here potentials which are a sum over translates of a fixed potential \tilde{V} , which is localized within a unit cell:

$$V(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^2} \tilde{V}(\mathbf{x} - \mathbf{m}).$$




FIG. 7.4. *Eigenvalue ordering, up to $\mathcal{O}(\varepsilon)$: $\mu_{-1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_1^\varepsilon$.*




FIG. 7.5. *Eigenvalue ordering, up to $\mathcal{O}(\varepsilon)$: $\mu_{+1}^\varepsilon < \mu_{+i}^\varepsilon = \mu_{-i}^\varepsilon < \mu_{-1}^\varepsilon$.*

The potential \tilde{V} is taken to be a finite sum of well-localized identical negative Gaussians (potential wells) with centers located within the primitive cell $[0, 1]^2$, i.e., an arrangement of “atoms” within the unit cell. The regime where the depth of the atomic potentials is large is the regime of *strong binding*. The spectral properties associated with the “low-lying” bands are expected to be well-approximated, after appropriate rescaling, by a *tight-binding* limiting model; see, for example, [2]. A rigorous analysis of the low-lying dispersion surfaces and their approximation by those of the tight-binding model was carried out for honeycomb structures in [7].

Figures 7.6 to 7.8 of this section display dispersion surfaces for $-\Delta + V$, for different choices of V , each for increasing atomic well depths. We now discuss these examples.

Example 7.1 (superposition over the Lieb lattice of Gaussian wells). To address the structural stability of Question 1.1 in the introduction, we consider a periodic potential, whose restriction to the primitive cell is

$$\tilde{V}_L(\mathbf{x}) = -V_0(e^{-|\mathbf{x}|^2/\sigma} + e^{-|\mathbf{x}-(1/2,0)|^2/\sigma} + e^{-|\mathbf{x}-(0,1/2)|^2/\sigma}), \quad V_0 > 0.$$

As a typical value of σ we take $\sigma = .001$ and vary the depth of the atomic wells by increasing V_0 . In Figure 7.6, we observe that as V_0 is increased the first three (the low-lying) dispersion surfaces come together in a manner approaching that of




FIG. 7.6. A plot of the dispersion surfaces for the Gaussian Lieb lattice potential with depth $V_0 = 10$ (top left), 500 (top right), 1,000 (bottom left), and 2,000 (bottom right).

the tight-binding limit shown in Figure 1.3; see Appendix A for a discussion of the tight-binding model. Also related is the discussion around Figure 7.4.

However, as indicated in Theorem 4.1 for finite V_0 , there are only two surfaces touching at $\mathbf{M} = (\pi, \pi)$ and the second band has hyperbolic character. The inset in Figure 7.4 displays this character.

Example 7.2 (superposition of \mathbb{Z}^2 -translates of Gaussian wells). We consider the potential, whose restriction to the primitive cell is given by

$$\tilde{V}(\mathbf{x}) = -V_0 e^{-|\mathbf{x}|^2/\sigma}.$$

For V_0 positive and large, we expect that the lowest dispersion surfaces will be governed by a tight-binding model with a single band, namely, the discrete Laplacian on a square lattice. Indeed, Figure 7.7 shows that the first (lowest) dispersion surface separates from the other (higher) dispersion surfaces and takes on a quadratic character in a neighborhood of $\mathbf{k} = 0$. The second and third surfaces intersect for all V_0 as is consistent with the curves in Figure 7.5.

Example 7.3 (a potential which is not invariant under reflection about the line $x_1 = x_2$). In Figure 7.8 we demonstrate how the absence of reflection symmetry about the line $x_1 = x_2$ in the physical domain manifests itself in less symmetry in the dispersion surfaces. In this case, Theorem 4.1 anticipates lesser symmetry, manifested in the nonzero cross-term, β , in the normal form (4.1). To illustrate this we take a simple potential of the form

$$\tilde{V}(\mathbf{x}) = -V_0 (\cos(2\pi(x_1 + 2x_2)) + \cos(2\pi(2x_1 - x_2))).$$




FIG. 7.7. A plot of the dispersion surfaces for the Gaussian square lattice potential with depth $V_0 = 10$ (top left), 500 (top right), 1,000 (bottom left), and a close-up of the bottom surface at depth $V_0 = 1,000$ (bottom right).

Thus, in Figure 7.8 we observe some lack of reflection symmetry about the line $\kappa_1 = \kappa_2$.

Appendix A. Tight-binding model for the Lieb lattice. Denote by $\Psi_A^{(m,n)}$, $\Psi_B^{(m,n)}$, $\Psi_C^{(m,n)}$ the amplitudes associated with the sites of the three sublattices comprising the Lieb lattice; see Figure 1.1 (right). We consider the nearest neighbor tight-binding model for the Lieb lattice is given by $(H^{TB}\Psi)_{mn} = E\Psi_{mn}$ for $m, n \in \mathbb{Z}$:

$$(A.1) \quad \begin{pmatrix} \Psi_B^{(m,n)} + \Psi_B^{(m,n+1)} \\ \Psi_A^{(m,n)} + \Psi_C^{(m,n)} + \Psi_C^{(m-1,n)} + \Psi_A^{(m,n-1)} \\ \Psi_B^{(m,n)} + \Psi_B^{(m+1,n)} \end{pmatrix} = E \begin{pmatrix} \Psi_A^{(m,n)} \\ \Psi_B^{(m,n)} \\ \Psi_C^{(m,n)} \end{pmatrix}, \quad m, n \in \mathbb{Z}.$$

Quasi-periodic (plane-wave) solutions with Bloch momentum $\mathbf{k} = (k_1, k_2) \in \mathcal{B} = [-\pi, \pi]^2$ are of the form $\Psi^{(m,n)} = e^{i(mk_1 + nk_2)}\psi$, where $\psi \in \mathbb{C}^3$ is independent of (m, n) . Substituting into (A.1) we obtain the algebraic eigenvalue problem $(A(\mathbf{k}) - E(\mathbf{k}) I_{3 \times 3})\psi = 0$, where

$$A(\mathbf{k}) = \begin{pmatrix} 0 & 1 + e^{ik_2} & 0 \\ 1 + e^{-ik_2} & 0 & 1 + e^{-ik_1} \\ 0 & 1 + e^{ik_1} & 0 \end{pmatrix}.$$




FIG. 7.8. A plot of the dispersion surfaces for the nonreflection symmetric potential with depth $V_0 = 10$ (top left) and 100 (top right), $V_0 = 500$ (bottom left) showing the lack of reflection symmetry in the surfaces. To highlight the asymmetry, we have included a plot of cross-sections of the first three dispersion surfaces with $V_0 = 100$ along the line from $X = (0, \pi)$ to $\rho(X) = (\pi, 0)$.

The three bands of the tight-binding model are given by the three solution branches of the algebraic equation $\det(A(\mathbf{k}) - E(\mathbf{k})\mathbf{I}) = 0$, which are given explicitly:

$$E_0(\mathbf{k}) = 0, \quad E_{\pm}(\mathbf{k}) = \pm \sqrt{4 + 2 \cos k_1 + 2 \cos k_2}, \quad \mathbf{k} \in \mathcal{B}.$$

These three dispersion surfaces are plotted in Figure 1.3.

Appendix B. Dynamics of wave-packets spectrally localized near \mathbf{M} .

In this section, we study the Schrödinger evolution $i\partial_t \psi = H_V \psi$ for wave-packet initial conditions which are spectrally located near the \mathbf{M} -point. In particular, denote by μ_S a multiplicity two $L^2_{\mathbf{M}}$ eigenvalue with corresponding eigenspace given by $\text{span}\{\Phi_1, \Phi_2\}$; see Theorem 4.1. We consider initial conditions of the form $\psi(\mathbf{x}, t = 0) = C_{10}(\delta \mathbf{x})\Phi_1(\mathbf{x}) + C_{20}(\delta \mathbf{x})\Phi_2(\mathbf{x})$, where $0 < \delta \ll 1$ and C_{10} and C_{20} are smooth and localized functions on \mathbb{R}^2 . The Floquet–Bloch components of such initial conditions are concentrated near \mathbf{M} .

Here we derive a formal multiscale solution. A proof of the validity of this expansion can be derived along the lines of [10] in the context of honeycomb structures; see also [1].

We seek a solution depending on multiple spatial and temporal scales $\psi = \psi^\delta(\mathbf{x}, t; \vec{\mathbf{X}}, \vec{T})$, where $\vec{\mathbf{X}} = (\mathbf{X}_1, \mathbf{X}_2, \dots) = (\delta \mathbf{x}_1, \delta^2 \mathbf{x}_2, \dots)$ and $\vec{T} = (T_1, T_2, \dots) = (\delta t_1, \delta^2 t_2, \dots)$. In terms of this extended set of variables, the time-dependent Schrödinger equation becomes

$$(B.1) \quad i(\partial_t + \delta \partial_{T_1} + \delta^2 \partial_{T_2} + \dots) \psi^\delta = - \left(\nabla_{\mathbf{x}} + \delta \nabla_{\mathbf{x}_1} + \delta^2 \nabla_{\mathbf{x}_2} \right)^2 \psi^\delta + V(\mathbf{x}) \psi^\delta + \dots$$

We seek a solution which is time-harmonic with respect to the fast time variable, t :

$$\psi^\delta = e^{-i\mu_S t} \sum_{j \geq 0} \delta^j \psi_j(\mathbf{x}; \vec{\mathbf{X}}, \vec{T}).$$

Substitution into (B.1) and equating terms of order $\mathcal{O}(\delta^j)$, $j \geq 0$, we obtain the following hierarchy of equations at each order in the small parameter δ . The first several are

(B.2)

$$\mathcal{O}(\delta^0): \quad (\mu_S - H_V) \psi_0 = 0;$$

(B.3)

$$\mathcal{O}(\delta): \quad (\mu_S - H_V) \psi_1 = - (i \partial_{T_1} + 2 \nabla_{\mathbf{x}} \cdot \nabla_{\mathbf{x}_1}) \psi_0;$$

(B.4)

$$\mathcal{O}(\delta^2): \quad (\mu_S - H_V) \psi_2 = - (i \partial_{T_2} + 2 \nabla_{\mathbf{x}} \cdot \nabla_{\mathbf{x}_2} + \Delta_{\mathbf{x}_1}) \psi_0 - (i \partial_{T_1} + 2 \nabla_{\mathbf{x}} \cdot \nabla_{\mathbf{x}_1}) \psi_1.$$

We regard each equation as a linear elliptic equation (nonhomogeneous for $j \geq 1$) in the variable $\mathbf{x} \in \mathbb{R}^2$, depending on slow variables $(\vec{\mathbf{X}}, \vec{T})$, treated as frozen parameters. The precise dependence of ψ_j on these parameters is determined through solvability conditions for the above hierarchy. In fact, to the order in δ that we solve, $\mathcal{O}(\delta^2)$, we find it necessary to modulate only the scales $\mathbf{X}_1, \mathbf{X}_2, T_1$, and T_2 , and so henceforth we assume $\vec{\mathbf{X}} = (\mathbf{X}_1, \mathbf{X}_2)$ and $\vec{T} = (T_1, T_2)$.

Consider the $\mathcal{O}(\delta^0)$ equation (B.2). By assumption, the general solution is

$$(B.5) \quad \psi_0(\mathbf{x}; \vec{\mathbf{X}}, \vec{T}) = C_1(\vec{\mathbf{X}}, \vec{T}) \Phi_1(\mathbf{x}) + C_2(\vec{\mathbf{X}}, \vec{T}) \Phi_2(\mathbf{x}),$$

where $C_1(\vec{\mathbf{X}}, \vec{T})$ and $C_2(\vec{\mathbf{X}}, \vec{T})$ are to be determined. The expression (B.5) satisfies the \mathbf{M} -pseudoperiodic boundary condition $\psi_0(\mathbf{x} + \mathbf{v}; \vec{\mathbf{X}}, \vec{T}) = e^{i\mathbf{M} \cdot \mathbf{v}} \psi_0(\mathbf{x}; \vec{\mathbf{X}}, \vec{T})$ for all $\mathbf{v} \in \mathbb{Z}^2$ and all $\mathbf{x} \in \mathbb{R}^2$ and we shall impose this same pseudoperiodicity at all subsequent orders:

$$\psi_j(\mathbf{x} + \mathbf{v}; \vec{\mathbf{X}}, \vec{T}) = e^{i\mathbf{M} \cdot \mathbf{v}} \psi_j(\mathbf{x}; \vec{\mathbf{X}}, \vec{T}), \quad j \geq 1.$$

Continuing on to $\mathcal{O}(\delta^2)$, we have from (B.3) that

$$(B.6) \quad (\mu_S - H) \psi_1 = \sum_{q=1}^2 [-i \partial_{T_1} C_q \Phi_q(\mathbf{x}) + 2 \nabla_{\mathbf{x}_1} C_q \cdot \nabla_{\mathbf{x}} \Phi_q(\mathbf{x})],$$

$$\psi_1(\mathbf{x} + \mathbf{v}; \vec{\mathbf{X}}, \vec{T}) = e^{i\mathbf{M} \cdot \mathbf{v}} \psi_1(\mathbf{x}; \vec{\mathbf{X}}, \vec{T}).$$

Solvability of (B.6) requires that orthogonality of the right-hand side of (B.6) to the span of $\{\Phi_1, \Phi_2\}$. Using the orthonormality relations $\langle \Phi_p, \Phi_q \rangle = \delta_{p,q}$, $p, q = 1, 2$, and Proposition 4.10, $\langle \Phi_p, \nabla_{\mathbf{x}} \Phi_q \rangle = \mathbf{0}$, $p, q = 1, 2$, we obtain that $\partial_{T_1} C_p = 0$, $p = 1, 2$. Therefore, in terms of the resolvent operator $\mathcal{R}(\mu_S) = (H - \mu_S I)^{-1}$, we have

$$(B.7) \quad \psi_1(\mathbf{x}; \mathbf{X}_1, \mathbf{X}_2, T_2) = 2 \mathcal{R}(\mu_S) \sum_{q=1}^2 \nabla_{\mathbf{x}_1} C_q(\mathbf{X}_1, \mathbf{X}_2, T_2) \cdot \nabla_{\mathbf{x}} \Phi_q(\mathbf{x}).$$

We proceed finally to the equation and boundary conditions for ψ_2 at (B.4):

$$(B.8) \quad (\mu_S - H) \psi_2 = - (i \partial_{T_2} + 2 \nabla_{\mathbf{x}} \cdot \nabla_{\mathbf{x}_2} + \Delta_{\mathbf{x}_1}) \psi_0 - 2 \nabla_{\mathbf{x}} \cdot \nabla_{\mathbf{x}_1} \psi_1,$$

$$\psi_2(\mathbf{x} + \mathbf{v}; \vec{\mathbf{X}}, \vec{T}) = e^{i\mathbf{M} \cdot \mathbf{v}} \psi_2(\mathbf{x}; \vec{\mathbf{X}}, \vec{T}),$$

where the expressions for ψ_0 and ψ_1 are displayed in (B.5) and (B.7).

PROPOSITION B.1. *A sufficient condition for the solvability of (B.8) is that the pair of amplitudes $C_1 = C_1(\mathbf{X}_1, T_2)$ and $C_2 = C_2(\mathbf{X}_1, T_2)$ satisfy the coupled system of constant coefficient homogenized Schrödinger equations:*

$$(B.9) \quad i\partial_{T_2} C_p = -\Delta_{\mathbf{X}_1} C_p + 4 \sum_{q=1}^2 \sum_{r,s=1}^2 a_{r,s}^{p,q} \frac{\partial^2 C_q}{\partial X_{1r} \partial X_{1s}}, \quad p = 1, 2.$$

Here, $a_{s,r}^{p,q}$ are the matrix elements of $(A^{p,q})_{sr}$ displayed in (4.23), $a_{r,s}^{p,q} = \langle \partial_r \Phi_p, \mathcal{R}(\mu_S) \partial_s \Phi_q \rangle$, which were simplified using symmetry arguments in section 4.1.

As a consequence of Proposition B.1, we have item 6 in our summary of results, section 1.2. Namely, solutions of the time-dependent Schrödinger equation with initial conditions of the form

$$\psi(\mathbf{x}, 0) = \psi_0^\delta(\mathbf{x}) = C_1(\mathbf{X}) \Phi_1(\mathbf{x}) + C_2(\mathbf{X}) \Phi_2(\mathbf{x})$$

evolve on large, finite time scales according to

$$i \frac{\partial}{\partial T} C_p = - \sum_{q=1}^2 \sum_{r,s=1}^2 \frac{\partial}{\partial X_r} \Upsilon_{r,s}^{p,q} \frac{\partial}{\partial X_s} C_q, \quad p = 1, 2,$$

where $T = T_2 = \delta^2 t$ and $\Upsilon_{r,s}^{p,q}$ depend on $(A^{p,q})_{sr}$ as stated above.

Proof. Taking the inner product of the right-hand side of (B.8) with $\Phi_p(\mathbf{x})$, $p = 1, 2$, substituting in the expressions for ψ_0 and ψ_1 , recalling that $\partial_{T_1} C_p = 0$, and applying Proposition 4.10, we have

$$(B.10) \quad (i\partial_{T_2} + \Delta_{\mathbf{X}_1}) C_p - 4 \sum_{q=1}^2 \langle \nabla_{\mathbf{x}} \Phi_p \cdot \nabla_{\mathbf{x}_1}, \mathcal{R}(\mu_S) \nabla_{\mathbf{x}} \Phi_q \cdot \nabla_{\mathbf{x}_1} C_q \rangle = 0, \quad p = 1, 2.$$

Expansion of the dot products yields (B.9). \square

Remark B.2. The dispersion relation for system (B.9) is

$$\det \widetilde{\mathcal{M}}_{\text{app}}(\nu, \kappa) = \nu^2 - |q(\kappa)|^2 = \nu^2 - |\gamma(\kappa_1^2 - \kappa_2^2) + 2\beta\kappa_1\kappa_2|^2 = 0$$

for $\kappa = (\kappa_1, \kappa_2) \in \mathbb{R}^2$ yielding two branches given by the leading order expressions in (1.4).

Appendix C. Calculations for $-\Delta + \varepsilon V$, V admissible and ε small. Theorem 4.1 and Corollary 4.2 give a precise description of touching dispersion surfaces at the vertices of \mathcal{B} . The local expansions of these dispersion surfaces are given in terms of coefficients α , β , and γ . In this section we consider $H^\varepsilon = -\Delta + \varepsilon V$, where ε is real, nonzero, and sufficiently small, and we obtain the leading order expressions of α^ε , β^ε , and γ^ε for $|\varepsilon| > 0$ small, required to complete the proof of Theorem 5.4. The expressions for α^ε , β^ε , and γ^ε are displayed in (4.27) and are given by

$$(C.1) \quad a_{\ell,m}^{(r),(s)} = \langle \partial_{x_\ell} \Phi_{(r)}^\varepsilon, \mathcal{R}^\varepsilon(\mu_S^\varepsilon) \partial_{x_m} \Phi_{(s)}^\varepsilon \rangle, \quad r, s \in \{+i, -i\}, \quad \ell, m \in \{1, 2\}.$$

In (C.1) we have used the notation $\Phi_1 = \Phi_{(+i)}$ and $\Phi_2 = \Phi_{(-i)}$ of Theorem 5.3. We now seek the leading order behavior of $a_{\ell,m}^{(r),(s)}$ for ε small and nonzero.

Recall by Theorem 5.3 there are four $L_{\mathbf{M}}^2$ eigenvalues counting multiplicity on an $\mathcal{O}(\varepsilon)$ neighborhood of μ_S^ε . Two correspond to multiplicity one $L_{\mathbf{M},+i}^2$ and $L_{\mathbf{M},-i}^2$ eigenstates $\Phi_{(+i)}^\varepsilon$ and $\Phi_{(-i)}^\varepsilon$ with corresponding eigenvalue which we denote μ_S^ε . The other two eigenpairs consist of two distinct $L_{\mathbf{M},+1}^2$ and $L_{\mathbf{M},-1}^2$ eigenstates $\Phi_{(+1)}^\varepsilon$ and $\Phi_{(-1)}^\varepsilon$ with corresponding eigenvalues we denote $\mu_{(+1)}^\varepsilon = \mu_S + \mathcal{O}(\varepsilon)$ and $\mu_{(-1)}^\varepsilon = \mu_S + \mathcal{O}(\varepsilon)$, respectively; see (5.21) and (5.24). All other eigenvalues of H^ε are at a distance of order 1 from μ_S^ε for ε small, and hence we express $a_{\ell,m}^{(r),(s)}$ as dominant parts coming from these “nearby” eigenvalues, with a remainder which is smaller for $\varepsilon > 0$ and small.

We have

$$(C.2) \quad \begin{aligned} a_{\ell,m}^{(r),(s)} &= \langle \partial_{x_\ell} \Phi_{(r)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_m} \Phi_{(s)}^\varepsilon \rangle \\ &= \sum_{q \in \{+1, -1\}} \frac{\langle \Phi_{(q)}^\varepsilon, \partial_{x_\ell} \Phi_{(r)}^\varepsilon \rangle \langle \Phi_{(q)}^\varepsilon, \partial_{x_m} \Phi_{(s)}^\varepsilon \rangle}{\mu_{(q)}^\varepsilon - \mu_S^\varepsilon} + \sum_{b \geq 5} \frac{\langle \Phi_b^\varepsilon, \partial_{x_\ell} \Phi_{(r)}^\varepsilon \rangle \langle \Phi_b^\varepsilon, \partial_{x_m} \Phi_{(s)}^\varepsilon \rangle}{\mu_b^\varepsilon - \mu_S^\varepsilon}. \end{aligned}$$

Observe the terms of (C.2), for $q \in \{+1, -1\}$, are of order $\mathcal{O}(\frac{1}{\varepsilon})$, while the contributions from the higher order bands $b \geq 5$ are $\mathcal{O}(1)$. From the expansion of $a_{\ell,m}^{(r),(s)}$, we show, in fact, the following.

PROPOSITION C.1. *Given ε sufficiently small and nonzero,*

$$(C.3) \quad \alpha^\varepsilon = 4a_{1,1}^{1,1} = 4a_{1,1}^{(+i),(+i)} = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1);$$

$$(C.4) \quad \beta^\varepsilon = 4a_{1,2}^{1,2} = 4a_{1,2}^{(+i),(-i)} = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1);$$

$$(C.5) \quad \gamma^\varepsilon = 4a_{1,1}^{1,2} = 4a_{1,1}^{(+i),(-i)} = -\frac{32\pi^2}{\varepsilon} i \left(\frac{V_{01}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1).$$

C.1. Derivation of terms in summand. First, note that $\mathbf{k}_1 = (2\pi, 0)^T$ and $\mathbf{k}_2 = (0, 2\pi)^T$ are the dual lattice basis vectors for $\Lambda = \mathbb{Z}^2$.

$$(C.6) \quad \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle = (\mathbf{k}_1 - \mathbf{k}_2)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon)$$

$$(C.7) \quad \langle \Phi_{(-1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle = (\mathbf{k}_2 - \mathbf{k}_1)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon);$$

$$(C.8) \quad \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle = (\mathbf{k}_2 - \mathbf{k}_1)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon);$$

$$(C.9) \quad \langle \Phi_{(-1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle = (\mathbf{k}_1 - \mathbf{k}_2)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon).$$

We have derived the expressions

$$\begin{aligned} \Phi_{(+i)}^\varepsilon(\mathbf{x}) &= \sum_{\mathbf{m} \in \mathcal{S}} c_{(+i)}^\varepsilon(\mathbf{m}) (e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - ie^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + ie^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}}), \\ \partial_{x_\ell} \Phi_{(+i)}^\varepsilon(\mathbf{x}) &= \sum_{\mathbf{m} \in \mathcal{S}} c_{(+i)}^\varepsilon(\mathbf{m}) \left[i(\mathbf{M} + m_1\mathbf{k}_1 + m_2\mathbf{k}_2)_\ell e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right. \\ &\quad - i^2(\mathbf{M} + m_2\mathbf{k}_1 - (m_1 + 1)\mathbf{k}_2)_\ell e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \\ &\quad - i(\mathbf{M} - (m_1 + 1)\mathbf{k}_1 - (m_2 + 1)\mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \\ &\quad \left. + i^2(\mathbf{M} - (1 + m_2)\mathbf{k}_1 + m_1\mathbf{k}_2)_\ell e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right] \end{aligned}$$

$$\Phi_{(\pm 1)}^\varepsilon(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c_{(\pm 1)}^\varepsilon(\mathbf{m}) (e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \pm e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \pm e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}}).$$

Now we proceed to prove the results in (C.6)–(C.9).

$$\begin{aligned} & \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle \\ &= \int_{\Omega} \sum_{\mathbf{n} \in \mathcal{S}} \sum_{\mathbf{m} \in \mathcal{S}} \overline{c_{(+1)}^\varepsilon(\mathbf{n})} c_{(+i)}^\varepsilon(\mathbf{m}) (e^{i\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^3\mathbf{M}^\mathbf{n} \cdot \mathbf{x}}) \\ & \quad * \{i(\mathbf{M}^\mathbf{m})_\ell e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + (R\mathbf{M}^\mathbf{m})_\ell e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - i(R^2\mathbf{M}^\mathbf{m})_\ell e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - (R^3\mathbf{M}^\mathbf{m})_\ell e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}}\} d\mathbf{x} \\ &= \int_{\Omega} \sum_{\mathbf{n} \in \mathcal{S}} \sum_{\mathbf{m} \in \mathcal{S}} \overline{c_{(+1)}^\varepsilon(\mathbf{n})} c_{(+i)}^\varepsilon(\mathbf{m}) (e^{i\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^3\mathbf{M}^\mathbf{n} \cdot \mathbf{x}}) \\ & \quad * \left[i(\mathbf{M} + m_1\mathbf{k}_1 + m_2\mathbf{k}_2)_\ell e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - i^2(\mathbf{M} + m_2\mathbf{k}_1 - (m_1 + 1)\mathbf{k}_2)_\ell e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right. \\ & \quad \left. - i(\mathbf{M} - (m_1 + 1)\mathbf{k}_1 - (m_2 + 1)\mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + i^2(\mathbf{M} - (1 + m_2)\mathbf{k}_1 + m_1\mathbf{k}_2)_\ell e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right]. \end{aligned}$$

The leading order term is associated with $\mathbf{n} = \mathbf{m} = (0, 0) \in \mathcal{S}$, for which $c_{(+1)}^\varepsilon(\mathbf{n}) = c_{(-1)}^\varepsilon(\mathbf{n}) = c_{(+i)}^\varepsilon(\mathbf{m}) = 1$. Then,

$$\begin{aligned} \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle &= \int_{\Omega} (e^{i\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^3\mathbf{M}^\mathbf{n} \cdot \mathbf{x}}) \\ & \quad \cdot (i\mathbf{M}_\ell e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + (\mathbf{M} - \mathbf{k}_2)_\ell e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - i(\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \\ & \quad - (\mathbf{M} - \mathbf{k}_1)_\ell e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}}) d\mathbf{x} + \mathcal{O}(\varepsilon) \\ &= [(\mathbf{M} - \mathbf{k}_2) - (\mathbf{M} - \mathbf{k}_1)]_\ell + i[\mathbf{M} - (\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)]_\ell + \mathcal{O}(\varepsilon) \\ &= (\mathbf{k}_1 - \mathbf{k}_2)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon). \end{aligned}$$

Similarly, we have

$$\begin{aligned} \langle \Phi_{(-1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle &= \int_{\Omega} (e^{i\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} - e^{iR\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^\mathbf{n} \cdot \mathbf{x}} - e^{iR^3\mathbf{M}^\mathbf{n} \cdot \mathbf{x}}) \\ & \quad \cdot (i\mathbf{M}_\ell e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + (\mathbf{M} - \mathbf{k}_2)_\ell e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - i(\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \\ & \quad - (\mathbf{M} - \mathbf{k}_1)_\ell e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}}) d\mathbf{x} + \mathcal{O}(\varepsilon) \\ &= -[(\mathbf{M} - \mathbf{k}_2) - (\mathbf{M} - \mathbf{k}_1)]_\ell + i[\mathbf{M} - (\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)]_\ell + \mathcal{O}(\varepsilon) \\ &= (\mathbf{k}_2 - \mathbf{k}_1)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon). \end{aligned}$$

The inner products of $\Phi_{(+1)}$ and $\Phi_{(-1)}$ with $\partial_{x_\ell} \Phi_{(-i)}$ are calculated similarly. Recall that

$$\Phi_{(-i)}^\varepsilon(\mathbf{x}) = \sum_{\mathbf{m} \in \mathcal{S}} c_{(-i)}^\varepsilon(\mathbf{m}) (e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} + ie^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} - ie^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}}).$$

Then,

$$\begin{aligned} \partial_{x_\ell} \Phi_{(-i)}^\varepsilon(\mathbf{x}) &= \sum_{\mathbf{m} \in \mathcal{S}} c_{(-i)}^\varepsilon(\mathbf{m}) \left[i(\mathbf{M} + m_1\mathbf{k}_1 + m_2\mathbf{k}_2)_\ell e^{i\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right. \\ & \quad + i^2(\mathbf{M} + m_2\mathbf{k}_1 - (m_1 + 1)\mathbf{k}_2)_\ell e^{iR\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \\ & \quad - i(\mathbf{M} - (m_1 + 1)\mathbf{k}_1 - (m_2 + 1)\mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \\ & \quad \left. - i^2(\mathbf{M} - (1 + m_2)\mathbf{k}_1 + m_1\mathbf{k}_2)_\ell e^{iR^3\mathbf{M}^\mathbf{m} \cdot \mathbf{x}} \right]. \end{aligned}$$

Expanding $\langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle$, we obtain

$$\begin{aligned} \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle &= \int_{\Omega} (e^{i\mathbf{M}^n \cdot \mathbf{x}} + e^{iR\mathbf{M}^n \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^n \cdot \mathbf{x}} + e^{iR^3\mathbf{M}^n \cdot \mathbf{x}}) \\ &\quad \cdot (i\mathbf{M}_\ell e^{i\mathbf{M}^m \cdot \mathbf{x}} - (\mathbf{M} - \mathbf{k}_2)_\ell e^{iR\mathbf{M}^m \cdot \mathbf{x}} - i(\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^m \cdot \mathbf{x}} \\ &\quad + (\mathbf{M} - \mathbf{k}_1)_\ell e^{iR^3\mathbf{M}^m \cdot \mathbf{x}}) d\mathbf{x} + \mathcal{O}(\varepsilon) \\ &= -[(\mathbf{M} - \mathbf{k}_2) - (\mathbf{M} - \mathbf{k}_1)]_\ell + i[\mathbf{M} - (\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)]_\ell + \mathcal{O}(\varepsilon) \\ &= (\mathbf{k}_2 - \mathbf{k}_1)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon). \end{aligned}$$

Similarly,

$$\begin{aligned} \langle \Phi_{(-1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle &= \int_{\Omega} (e^{i\mathbf{M}^n \cdot \mathbf{x}} - e^{iR\mathbf{M}^n \cdot \mathbf{x}} + e^{iR^2\mathbf{M}^n \cdot \mathbf{x}} - e^{iR^3\mathbf{M}^n \cdot \mathbf{x}}) \\ &\quad \cdot (i\mathbf{M}_\ell e^{i\mathbf{M}^m \cdot \mathbf{x}} - (\mathbf{M} - \mathbf{k}_2)_\ell e^{iR\mathbf{M}^m \cdot \mathbf{x}} - i(\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)_\ell e^{iR^2\mathbf{M}^m \cdot \mathbf{x}} \\ &\quad + (\mathbf{M} - \mathbf{k}_1)_\ell e^{iR^3\mathbf{M}^m \cdot \mathbf{x}}) d\mathbf{x} + \mathcal{O}(\varepsilon) \\ &= [(\mathbf{M} - \mathbf{k}_2) - (\mathbf{M} - \mathbf{k}_1)]_\ell + i[\mathbf{M} - (\mathbf{M} - \mathbf{k}_1 - \mathbf{k}_2)]_\ell + \mathcal{O}(\varepsilon) \\ &= (\mathbf{k}_1 - \mathbf{k}_2)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon). \end{aligned}$$

Collecting the results, we have (C.6)–(C.9). That is,

$$\begin{aligned} \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle &= (\mathbf{k}_1 - \mathbf{k}_2)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon); \\ \langle \Phi_{(-1)}^\varepsilon, \partial_{x_\ell} \Phi_{(+i)}^\varepsilon \rangle &= (\mathbf{k}_2 - \mathbf{k}_1)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon); \\ \langle \Phi_{(+1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle &= (\mathbf{k}_2 - \mathbf{k}_1)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon); \\ \langle \Phi_{(-1)}^\varepsilon, \partial_{x_\ell} \Phi_{(-i)}^\varepsilon \rangle &= (\mathbf{k}_1 - \mathbf{k}_2)_\ell + i(\mathbf{k}_1 + \mathbf{k}_2)_\ell + \mathcal{O}(\varepsilon). \end{aligned}$$

C.2. Coefficient calculations: α^ε , β^ε , γ^ε . For $\ell, m \in \{1, 2\}$ and $r, s \in \{+i, -i\}$

$$\begin{aligned} a_{\ell, m}^{(r), (s)} &= \langle \partial_{x_\ell} \Phi_{(r)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_m} \Phi_{(s)}^\varepsilon \rangle \\ &= \sum_{q \in \{+1, -1\}} \frac{\langle \Phi_{(q)}^\varepsilon, \partial_{x_\ell} \Phi_{(r)}^\varepsilon \rangle \langle \Phi_{(q)}^\varepsilon, \partial_{x_m} \Phi_{(s)}^\varepsilon \rangle}{\mu_{(q)}^\varepsilon - \mu_S^\varepsilon} + \mathcal{O}(1). \end{aligned}$$

Further, recall $\alpha^\varepsilon = 4a_{1,1}^{(+i), (+i)} = 4\langle \partial_{x_1} \Phi_{(+i)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle$. From the analysis above,

$$\begin{aligned} &\frac{\langle \Phi_{(+1)}^\varepsilon, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle \langle \Phi_{(+1)}^\varepsilon, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle}{\mu_{(+1)}^\varepsilon - \mu_S^\varepsilon} \\ &= \frac{(\mathbf{k}_1 - \mathbf{k}_2)_1 + i(\mathbf{k}_1 + \mathbf{k}_2)_1 [(\mathbf{k}_1 - \mathbf{k}_2)_1 + i(\mathbf{k}_1 + \mathbf{k}_2)_1] + \mathcal{O}(\varepsilon)}{\varepsilon(V_{00} + 2V_{01} + V_{11}) - \varepsilon(V_{00} - V_{11}) + \mathcal{O}(\varepsilon^2)} \\ &= \frac{4\pi^2 (1+i)(1+i) + \mathcal{O}(\varepsilon)}{2\varepsilon(V_{01} + V_{11}) + \mathcal{O}(\varepsilon^2)} \\ &= \frac{2\pi^2}{\varepsilon} \left(\frac{|1+i|^2 + \mathcal{O}(\varepsilon)}{(V_{01} + V_{11}) + \mathcal{O}(\varepsilon)} \right) \\ (C.10) \quad &= \frac{2\pi^2}{\varepsilon} \left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{01} + V_{11} + \mathcal{O}(\varepsilon)} \right). \end{aligned}$$

We calculate the inner products now with $q = -1$:

$$\begin{aligned}
 & \frac{\overline{\langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle} \langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle}{\mu_{(-1)}^\varepsilon - \mu_S^\varepsilon} \\
 &= \frac{(\mathbf{k}_2 - \mathbf{k}_1)_1 + i(\mathbf{k}_1 + \mathbf{k}_2)_1 [(\mathbf{k}_2 - \mathbf{k}_1)_1 + i(\mathbf{k}_1 + \mathbf{k}_2)_1] + \mathcal{O}(\varepsilon)}{\varepsilon(V_{00} - 2V_{01} + V_{11}) - \varepsilon(V_{00} - V_{11}) + \mathcal{O}(\varepsilon^2)} \\
 &= \frac{4\pi^2 (-1+i)(-1+i) + \mathcal{O}(\varepsilon)}{2\varepsilon(V_{11} - V_{01}) + \mathcal{O}(\varepsilon^2)} \\
 &= \frac{2\pi^2}{\varepsilon} \left(\frac{|-1+i|^2 + \mathcal{O}(\varepsilon)}{(V_{11} - V_{01}) + \mathcal{O}(\varepsilon)} \right) \\
 (C.11) \quad &= \frac{2\pi^2}{\varepsilon} \left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{11} - V_{01} + \mathcal{O}(\varepsilon)} \right).
 \end{aligned}$$

Substituting (C.10) and (C.11) into the expression for $a_{1,1}^{(+i),(+i)}(\varepsilon)$,

$$\begin{aligned}
 \alpha^\varepsilon &= 4a_{1,1}^{(+i),(+i)}(\varepsilon) = \frac{8\pi^2}{\varepsilon} \left[\left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{01} + V_{11} + \mathcal{O}(\varepsilon)} \right) + \left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{11} - V_{01} + \mathcal{O}(\varepsilon)} \right) \right] + \mathcal{O}(1) \\
 &= \frac{8\pi^2}{\varepsilon} \left(\frac{2(V_{11} - V_{01}) + 2(V_{01} + V_{11}) + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} \right) + \mathcal{O}(1) \\
 &= \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11} + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} \right) + \mathcal{O}(1).
 \end{aligned}$$

Therefore, we have (C.3):

$$\alpha^\varepsilon = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11} + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} \right) + \mathcal{O}(1) = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1).$$

The results for β^ε and γ^ε follow similarly. First, consider β^ε .

$$\begin{aligned}
 (C.12) \quad \beta^\varepsilon &= 4a_{1,2}^{(+i),(-i)}(\varepsilon) = 4\langle \partial_{x_1} \Phi_1^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_2} \Phi_{(-i)}^\varepsilon \rangle \\
 &= \frac{4\overline{\langle \Phi_{(+1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle} \langle \Phi_{(+1)}, \partial_{x_2} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(+1)}^\varepsilon - \mu_S^\varepsilon} + \frac{4\overline{\langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle} \langle \Phi_{(-1)}, \partial_{x_2} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(-1)}^\varepsilon - \mu_S^\varepsilon} + \mathcal{O}(1).
 \end{aligned}$$

Following the analysis as for α^ε , we find the following:

$$(C.13) \quad \frac{\overline{\langle \Phi_{(+1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle} \langle \Phi_{(+1)}, \partial_{x_2} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(+1)}^\varepsilon - \mu_S^\varepsilon} = \frac{2\pi^2}{\varepsilon} \left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{01} + V_{11} + \mathcal{O}(\varepsilon)} \right);$$

$$(C.14) \quad \frac{\overline{\langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle} \langle \Phi_{(-1)}, \partial_{x_2} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(-1)}^\varepsilon - \mu_S^\varepsilon} = \frac{2\pi^2}{\varepsilon} \left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{11} - V_{01} + \mathcal{O}(\varepsilon)} \right).$$

Combining (C.13) and (C.14),

$$\begin{aligned}\beta^\varepsilon &= 4\langle \partial_{x_1} \Phi_{(+i)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_2} \Phi_{(-i)}^\varepsilon \rangle = \frac{8\pi^2}{\varepsilon} \left[\left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{01} + V_{11} + \mathcal{O}(\varepsilon)} \right) + \left(\frac{2 + \mathcal{O}(\varepsilon)}{V_{11} - V_{01} + \mathcal{O}(\varepsilon)} \right) \right] \\ &= \left(\frac{8\pi^2}{\varepsilon} \right) \frac{2(V_{11} - V_{01}) + 2(V_{01} + V_{11}) + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} \\ &= \left(\frac{8\pi^2}{\varepsilon} \right) \frac{4V_{11} + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)}.\end{aligned}$$

Thus we have (C.4):

$$\beta^\varepsilon = \left(\frac{32\pi^2}{\varepsilon} \right) \frac{V_{11} + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} = \frac{32\pi^2}{\varepsilon} \left(\frac{V_{11}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1).$$

Finally, consider the decomposition for γ^ε .

$$\begin{aligned}\text{(C.15)} \quad \gamma^\varepsilon &= 4a_{1,1}^{(+i),(-i)}(\varepsilon) = 4\langle \partial_{x_1} \Phi_{(+i)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle \\ &= \frac{4\langle \Phi_{(+1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle \langle \Phi_{(+1)}, \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(+1)}^\varepsilon - \mu_S^\varepsilon} + \frac{4\langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle \langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(-1)}^\varepsilon - \mu_S^\varepsilon} + \mathcal{O}(1).\end{aligned}$$

Again, similarly as for α^ε , the first two terms of (C.15) yield the following:

$$\text{(C.16)} \quad \frac{\langle \Phi_{(+1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle \langle \Phi_{(+1)}, \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(+1)}^\varepsilon - \mu_S^\varepsilon} = \frac{2\pi^2}{\varepsilon} \left(\frac{2i + \mathcal{O}(\varepsilon)}{V_{01} + V_{11} + \mathcal{O}(\varepsilon)} \right);$$

$$\text{(C.17)} \quad \frac{\langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(+i)}^\varepsilon \rangle \langle \Phi_{(-1)}, \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle}{\mu_{(-1)}^\varepsilon - \mu_S^\varepsilon} = \frac{2\pi^2}{\varepsilon} \left(\frac{-2i + \mathcal{O}(\varepsilon)}{V_{11} - V_{01} + \mathcal{O}(\varepsilon)} \right).$$

Combining (C.16) and (C.17),

$$\begin{aligned}\langle \partial_{x_1} \Phi_{(+i)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle &= \frac{2\pi^2}{\varepsilon} \left[\left(\frac{2i + \mathcal{O}(\varepsilon)}{V_{01} + V_{11} + \mathcal{O}(\varepsilon)} \right) + \left(\frac{-2i + \mathcal{O}(\varepsilon)}{V_{11} - V_{01} + \mathcal{O}(\varepsilon)} \right) \right] \\ &= \left(\frac{2\pi^2}{\varepsilon} \right) \frac{2i(V_{11} - V_{01}) - 2i(V_{01} + V_{11}) + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} \\ &= \left(\frac{2\pi^2}{\varepsilon} \right) \frac{(-4i)V_{01} + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)}.\end{aligned}$$

Therefore, we have (C.5):

$$\begin{aligned}\gamma^\varepsilon &= 4\langle \partial_{x_1} \Phi_{(+i)}^\varepsilon, \mathcal{R}(\mu_S^\varepsilon) \partial_{x_1} \Phi_{(-i)}^\varepsilon \rangle = - \left(\frac{32\pi^2}{\varepsilon} i \right) \frac{V_{01} + \mathcal{O}(\varepsilon)}{(V_{11}^2 - V_{01}^2) + \mathcal{O}(\varepsilon)} + \mathcal{O}(1) \\ &= - \frac{32\pi^2}{\varepsilon} i \left(\frac{V_{01}}{V_{11}^2 - V_{01}^2} \right) + \mathcal{O}(1).\end{aligned}$$

With the coefficients $\alpha^\varepsilon, \beta^\varepsilon$, and γ^ε , expanded, the description for the expression (5.26),

$$\mu_\pm^\varepsilon(\mathbf{M} + \kappa) - \mu_S^\varepsilon = (1 - \alpha^\varepsilon)|\kappa|^2 + \mathcal{Q}_6^\varepsilon(\kappa) \pm \sqrt{\left| \gamma^\varepsilon(\kappa_1^2 - \kappa_2^2) + 2\beta^\varepsilon\kappa_1\kappa_2 \right|^2} + \mathcal{Q}_8^\varepsilon(\kappa),$$

described in Corollary 5.4 is complete.

Acknowledgments. The authors thank M. Rechtsman and C. L. Fefferman for very stimulating discussions. M. Rechtsman also provided very helpful advice on numerical spectral calculations. The fourth author would like to thank the Department of Mathematics of Stanford University for its excellent hospitality and environment during the Winter 2018 quarter, when part of this work was done as Bergman Visiting Professor.

REFERENCES

- [1] G. ALLAIRE AND A. PIATNITSKI, *Homogenization of the Schrödinger equation and effective mass theorems*, Comm. Math. Phys., 258 (2005), pp. 1–22, <https://doi.org/10.1007/s00220-005-1329-2>.
- [2] N. ASHCROFT AND N. MERMIN, *Solid State Physics*, Saunders College Publishing, Philadelphia, 1976.
- [3] J. AVRON AND B. SIMON, *Analytic properties of band functions*, Ann. Phys., 110 (1978), pp. 85–101, [https://doi.org/10.1016/0003-4916\(78\)90143-4](https://doi.org/10.1016/0003-4916(78)90143-4).
- [4] G. BERKOLAİKO AND A. COMECH, *Symmetry and Dirac points in graphene spectrum*, J. Spectral Theory, 8 (2018), pp. 1099–1147, <https://doi.org/10.4171/jst/223>.
- [5] Y. C. DE VERDIERE, *Sur les singularites de van hove generiques*, Mem. Soc. Math. Fr. Ser. 2, 46 (1991), pp. 99–109, <https://doi.org/10.24033/msmf.356>.
- [6] M. EASTHAM, *The Spectral Theory of Periodic Differential Equations*, Scottish Academic Press, Edinburgh, 1974.
- [7] C. L. FEFFERMAN, J. P. LEE-THORP, AND M. I. WEINSTEIN, *Honeycomb Schrödinger operators in the strong binding regime*, Comm. Pure Appl. Math., 71 (2017), pp. 1178–1270, <https://doi.org/10.1002/cpa.21735>.
- [8] C. L. FEFFERMAN, J. P. LEE-THORP, AND M. I. WEINSTEIN, *Topologically protected states in one-dimensional systems*, Mem. Amer. Math. Soc., 247 (2017), <https://doi.org/10.1090/memo/1173>.
- [9] C. L. FEFFERMAN AND M. I. WEINSTEIN, *Honeycomb lattice potentials and Dirac points*, J. Amer. Math. Soc., 25 (2012), pp. 1169–1220, <https://doi.org/10.1090/s0894-0347-2012-00745-0>.
- [10] C. L. FEFFERMAN AND M. I. WEINSTEIN, *Wave packets in honeycomb structures and two-dimensional Dirac equations*, Comm. Math. Phys., 326 (2014), pp. 251–286, <https://doi.org/10.1007/s00220-013-1847-2>.
- [11] V. V. GRUSHIN, *Multiparameter perturbation theory of Fredholm operators applied to Bloch functions*, Math. Notes, 86 (2009), pp. 767–774, <https://doi.org/10.1134/s0001434609110194>.
- [12] D. GUZMÁN-SILVA, C. MEJÍA-CORTÉS, M. A. BANDRES, M. C. RECHTSMAN, S. WEIMANN, S. NOLTE, M. SEGEV, A. SZAMEIT, AND R. A. VICENCIO, *Experimental observation of bulk and edge transport in photonic Lieb lattices*, New J. Phys., 16 (2014), 063061, <https://doi.org/10.1088/1367-2630/16/6/063061>.
- [13] J. JOANNOPOULOS, S. JOHNSON, J. WINN, AND R. MEADE, *Photonic Crystals: Molding the Flow of Light*, 2nd ed., Princeton University Press, Princeton, NJ, 2008.
- [14] P. KUCHMENT, *Floquet Theory for Partial Differential Equations*, Oper. Theory Adv. Appl. 60, Birkhauser, Basel, 2012.
- [15] P. KUCHMENT, *An overview of periodic elliptic operators*, Bull. Amer. Math. Soc., 53 (2016), pp. 343–414, <https://doi.org/10.1090/bull/1528>.
- [16] M. LEE, *Dirac cones for point scatterers on a honeycomb lattice*, SIAM J. Math. Anal., 48 (2016), pp. 1459–1488, <https://doi.org/10.1137/14095827x>.
- [17] J. LEE-THORP, M. I. WEINSTEIN, AND Y. ZHU, *Elliptic Operators with Honeycomb Symmetry: Dirac Points, Edge States and Applications to Photonic Graphene*, arXiv:1710.03389, 2017.
- [18] A. H. C. NETO, F. GUINEA, N. M. R. PERES, K. S. NOVOSELOV, AND A. K. GEIM, *The electronic properties of graphene*, Rev. Modern Phys., 81 (2009), pp. 109–162, <https://doi.org/10.1103/revmodphys.81.109>.
- [19] M. NIȚĂ, B. OSTAHIE, AND A. ALDEA, *Spectral and transport properties of the two-dimensional Lieb lattice*, Phys. Rev. B, 87 (2013), 125428, <https://doi.org/10.1103/physrevb.87.125428>.
- [20] M. REED AND B. SIMON, *Methods of Modern Mathematical Physics: Analysis of Operators*, Vol. 4, Academic Press, New York, 1978.
- [21] P. R. WALLACE, *The band theory of graphite*, Phys. Rev., 71 (1947), pp. 622–634, <https://doi.org/10.1103/physrev.71.622>.
- [22] C. WEEKS AND M. FRANZ, *Topological insulators on the Lieb and Perovskite lattices*, Phys. Rev. B, 82 (2010), 085310, <https://doi.org/10.1103/physrevb.82.085310>.