

An ultraweak-local discontinuous Galerkin method for PDEs with high order spatial derivatives

Qi Tao*, Yan Xu[†], Chi-Wang Shu[‡]

Abstract: In this paper, we develop a new discontinuous Galerkin method for solving several types of partial differential equations (PDEs) with high order spatial derivatives. We combine the advantages of local discontinuous Galerkin (LDG) method and ultraweak discontinuous Galerkin (UWDG) method. Firstly, we rewrite the PDEs with high order spatial derivatives into a lower order system, then apply the UWDG method to the system. We first consider the fourth order and fifth order nonlinear PDEs in one space dimension, and then extend our method to general high order problems and two space dimensions. The main advantage of our method over the LDG method is that we have introduced fewer auxiliary variables, thereby reducing memory and computational costs. The main advantage of our method over the UWDG method is that no internal penalty terms are necessary in order to ensure stability for both even and odd order PDEs. We prove stability of our method in the general nonlinear case and provide optimal error estimates for linear PDEs for the solution itself as well as for the auxiliary variables approximating its derivatives. A key ingredient in the proof of the error estimates is the construction of the relationship between the derivative and the element interface jump of the numerical solution and the auxiliary variable solution of the solution derivative. With this relationship, we can then use the discrete Sobolev and Poincaré inequalities to obtain the optimal error estimates. The theoretical findings are confirmed by numerical experiments.

*School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, P.R. China. Email: taoq@mail.ustc.edu.cn. Research supported by China Scholarship Council.

[†]Corresponding author. School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, P.R. China. Email: yxu@ustc.edu.cn. Research supported by National Numerical Windtunnel grants NNW2019ZT4-B08, Science Challenge Project TZT2019-A2.3, NSFC grants 11722112.

[‡]Division of Applied Mathematics, Brown University, Providence, RI 02912, USA. Email: shu@dam.brown.edu. Research supported by NSF grant DMS-1719410.

Keywords: Discontinuous Galerkin (DG) method; High order equation; Error estimate; Discrete Sobolev and Poincaré inequalities

MSC (2010): Primary 65M60; Secondary 35G25

1 Introduction

In this paper, we propose a new class of discontinuous Galerkin (DG) methods for solving several types of partial differential equations (PDEs) with high order spatial derivatives. The first two examples we consider are:

- The fourth order equation

$$u_t + (b(u)u_{xx})_{xx} = 0, \quad b(u) \geq 0 \quad (1.1)$$

- The fifth order equation

$$u_t + f(u_{xx})_{xxx} = 0. \quad (1.2)$$

The boundary conditions are assumed to be periodic for simplicity, although most of our discussions can be adapted for other types of boundary conditions. These equations are classical model equations for many very important physical applications. The fourth order problem has wide applications in the modeling of thin beams and plates, strain gradient elasticity, and phase separation in binary mixtures [14]. The fifth order nonlinear evolution equation is known as the critical surface-tension model [15].

Discontinuous Galerkin (DG) methods are a class of finite element methods (FEMs) using completely discontinuous basis functions. The first DG method was introduced in 1973 by Reed and Hill [20] in the framework of neutron transport. It was later developed for time-dependent nonlinear hyperbolic conservation laws, coupled with the Runge-Kutta time discretization, by Cockburn et al [5, 7, 8, 21]. Since then, the DG method has been intensively studied and successfully applied to various problems in a wide range of applications due to its flexibility with meshing, its compactness and its high parallel efficiency. For the equations containing higher order spatial derivatives, there are several different ways to approximate them by discontinuous Galerkin methods. One way is to use the local discontinuous Galerkin (LDG) method [9, 10, 13, 17, 25, 27, 28]. The idea of the LDG methods is to rewrite the equations with higher order spatial derivatives into a first order system, then apply the DG method to this system and

design suitable numerical fluxes to ensure stability. Another way is to use the penalty methods that add penalty terms at cell interfaces in the DG formulation for numerical stability [11, 19]. The third way is to use the ultra-weak DG (UWDG) methods [3]. It is based on repeated integration by parts to move all spatial derivatives to the test function in the weak formulation, and on a careful choice of the numerical fluxes to ensure stability and optimal accuracy. Unlike the traditional LDG method, the UWDG method can be applied without introducing any auxiliary variables or rewriting the original equation into a system. Recently, Liu et al. introduced a mixed DG method [16], by first rewriting the fourth order PDEs into a second order coupled system and then using a direct DG discretization for the second order system. L^2 stability was obtained without internal penalty.

In this paper, we design a new class of DG methods, combining the advantages of LDG and UWDG methodologies, to solve PDEs with high order spatial derivatives. The two PDEs (1.1) and (1.2) are used first as examples to develop our method. The method is then extended to a wider class of PDEs both in one and in two dimensions. Similar to the mixed DG method in [16], we first rewrite the higher order equation into a lower order (but not all first order) system. For example, we rewrite the fourth order problem into a second order system and rewrite the fifth order problem into a system with two second order equations and a first order equation, then we repeat the application of integration parts, and choose suitable numerical fluxes to ensure stability. For the equations with spatial derivative order less than or equal to three, our method will be the same as the LDG methods or ultra-weak DG method, but for higher order PDEs our method combines the advantages of the two type of methods, and is more efficient. It is known that the proof of optimal accuracy for LDG methods solving high order time-dependent wave equations is very difficult. The work in [26] by Xu and Shu might be the first to prove optimal order of accuracy in L^2 for not only the solution but also the auxiliary variables. In their work, the main idea is to derive energy stability for the auxiliary variables in the LDG scheme by using the scheme and its time derivatives. In [12] Fu et al. identified a sub-family of the numerical fluxes by choosing the coefficients in the linear combinations, so that the solution and some auxiliary variables of the proposed DG methods are optimally accurate in the L^2 norm. In [10] Dong and Shu proved the optimal error estimates for the higher even-order equations, including the cases both in one dimension and in multidimensional triangular meshes. In this paper, we prove the optimal error estimates for both the even order equations and the odd order equations. The main idea is to use an important relationship between the derivative and

the element interface jump of the numerical solution and the auxiliary variable numerical solution of the derivative [22, 23]. Then we can obtain suitable estimates to the auxiliary variables, which lead to the optimal error estimates for both the numerical solution and the auxiliary variables. This is a different approach from that in [10, 26], since in this way we do not need to estimate many energy equations, and can get the relationship between the solution and auxiliary variables directly.

The organization of the paper is as follows. In Section 2, we introduce some notations and projections that will be used later. In Section 3, the scheme for the fourth order equation is discussed, including the discussion on the L^2 stability and optimal error estimates. In Section 4, we follow the lines of Section 3 and consider the fifth order equation. In Section 5, we extend the schemes in Sections 3 and 4 to arbitrary even and odd order equations, respectively. We also extend the scheme for the fourth order equations to multidimensional Cartesian meshes as an example of multi-dimensions in Section 6. The theoretical results are confirmed numerically in Section 7. In Section 8, we give some concluding remarks.

2 Notations and projections

In this section, we will introduce some notations, definitions and projections that will be used later for the one-dimensional equations.

Throughout this paper, we adopt standard notations for the Sobolev spaces such as $W^{m,p}(D)$ on the subdomain $D \in \Omega$ equipped with the norm $\|\cdot\|_{m,p,D}$. If $D = \Omega$, we omit the index D ; and if $p = 2$, we set $W^{m,p}(D) = H^m(D)$, $\|\cdot\|_{m,p,D} = \|\cdot\|_{m,D}$; and we use $\|\cdot\|_D$ to denote the L^2 norm in D .

2.1 Basic notations

Let $\Omega = [0, 2\pi]$ and $0 = x_{\frac{1}{2}} < x_{\frac{3}{2}} < \cdots < x_{N+\frac{1}{2}} = 2\pi$ be $N+1$ distinct points on Ω . For each positive integer r , we define $Z_r = (1, 2, \cdots, r)$ and denote by

$$I_j = (x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}}), \quad x_j = \frac{1}{2}(x_{j-\frac{1}{2}} + x_{j+\frac{1}{2}}), \quad j \in Z_N,$$

the cells and cell centers, respectively. Let $h_j = x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}$, and $h = \max_j h_j$. We assume that the mesh is regular. Define

$$V_h = \{v_h : v_h|_{I_j} \in \mathcal{P}^k(I_j), j \in Z_N\}$$

to be the finite element space, where $\mathcal{P}^k$ denotes the space of polynomials of degree at most k . For any $v \in V_h$, $v_{j+\frac{1}{2}}^+$ and $v_{j+\frac{1}{2}}^-$ denote the right and left limit values of v at $j + \frac{1}{2}$, respectively. As usual, the average and the jump of the function v at $j + \frac{1}{2}$ are denoted as

$$\{v\}_{j+\frac{1}{2}} = \frac{1}{2}(v_{j+\frac{1}{2}}^+ + v_{j+\frac{1}{2}}^-), \quad \llbracket v \rrbracket_{j+\frac{1}{2}} = v_{j+\frac{1}{2}}^+ - v_{j+\frac{1}{2}}^-,$$

respectively.

2.2 Projections

Next, we will introduce some projections used in the error estimates. For example, we can choose the Gauss-Radau projections $P_h^\pm$ into V_h , such that for any u we have:

$$\int_{I_j} uv_h dx = \int_{I_j} P_h^\pm uv_h dx, \quad P_h^\pm u \left(x_{j \mp \frac{1}{2}}^\pm \right) = u \left(x_{j \mp \frac{1}{2}} \right), \quad (2.1)$$

$\forall j \in Z_N, v_h \in \mathcal{P}^{k-1}(I_j)$. Furthermore, for $k \geq 1$ we can define the projection $P_{1h}^\pm$ into V_h such that, for any u , the projection $P_{1h}^\pm u$ satisfies: $\forall j \in Z_N$

$$\int_{I_j} uv_h dx = \int_{I_j} P_{1h}^\pm uv_h dx, \quad (2.2)$$

for any $v_h \in \mathcal{P}^{k-2}(I_j)$ and

$$P_{1h}^\pm u \left(x_{j \mp \frac{1}{2}}^\pm \right) = u \left(x_{j \mp \frac{1}{2}} \right), \quad (P_{1h}^\pm u)_x \left(x_{j \mp \frac{1}{2}}^\pm \right) = u_x \left(x_{j \mp \frac{1}{2}} \right). \quad (2.3)$$

Similarly, for $k \geq 2$ we can define the projection $P_{2h}^\pm$ into V_h such that, for any u , it satisfies:

$$\int_{I_j} uv_h dx = \int_{I_j} P_{2h}^\pm uv_h dx, \quad (2.4)$$

and

$$P_{2h}^\pm u \left(x_{j \mp \frac{1}{2}}^\pm \right) = u \left(x_{j \mp \frac{1}{2}} \right), \quad (P_{2h}^\pm u)_x \left(x_{j \mp \frac{1}{2}}^\pm \right) = u_x \left(x_{j \mp \frac{1}{2}} \right), \quad (P_{2h}^\pm u)_{xx} \left(x_{j \mp \frac{1}{2}}^\pm \right) = u_{xx} \left(x_{j \mp \frac{1}{2}} \right), \quad (2.5)$$

for any $j \in Z_N, v_h \in \mathcal{P}^{k-3}(I_j)$. We will use different projections according to the need in each proof. For all these projections, the following inequality holds [4]:

$$\|u^e\| + h\|u^e\|_\infty + h^{\frac{1}{2}}\|u^e\|_{\Gamma_h} \leq Ch^{k+1}\|u\|_{k+1}, \quad (2.6)$$

where $u^e = \pi_h^\pm u - u$, $\pi_h = P_h$, P_{1h} , P_{2h} , and Γ_h denotes the set of boundary points of all elements I_j , and C is a positive constant dependent on k but not on h .

3 The fourth order problem

We start from the fourth order problem. Firstly, we consider the following one-dimensional nonlinear equation

$$u_t + (b(u)u_{xx})_{xx} = 0, \quad b(u) \geq 0, \quad (x, t) \in [0, 2\pi] \times (0, T], \quad (3.1)$$

$$u(x, 0) = u_0(x), \quad x \in \mathbb{R}, \quad (3.2)$$

where $u_0(x)$ is a smooth function. Without loss of generality, we only consider the periodic boundary conditions.

3.1 The numerical scheme

Before we introduce our DG method, we rewrite the fourth order equation (3.1) into a system of second order equations

$$u_t + v_{xx} = 0, \quad (3.3)$$

$$v - b(u)w = 0, \quad (3.4)$$

$$w - u_{xx} = 0. \quad (3.5)$$

Notice that, unlike the LDG method, we stop at second order equations and do not go all the way to a first order system. Our DG method is defined as follows: find $u_h, v_h, w_h \in V_h$ such that for all $p, s, q \in V_h$, we have

$$((u_h)_t, p)_j + (v_h, p_{xx})_j + \tilde{v}_x p^-|_{j+\frac{1}{2}} - \tilde{v}_x p^+|_{j-\frac{1}{2}} - \hat{v} p_x^-|_{j+\frac{1}{2}} + \hat{v} p_x^+|_{j-\frac{1}{2}} = 0, \quad (3.6)$$

$$(v_h, s)_j - (b(u_h)w_h, s)_j = 0, \quad (3.7)$$

$$(w_h, q)_j - (u_h, q_{xx})_j - \tilde{u}_x q^-|_{j+\frac{1}{2}} + \tilde{u}_x q^+|_{j-\frac{1}{2}} + \hat{u} q_x^-|_{j+\frac{1}{2}} - \hat{u} q_x^+|_{j-\frac{1}{2}} = 0. \quad (3.8)$$

Here $(u, v)_j = \int_{I_j} uv dx$ and $\hat{v}, \tilde{v}_x, \hat{u}, \tilde{u}_x$ are the numerical fluxes. The terms involving these fluxes appear from repeated integration by parts, and a suitable choice for these fluxes is the key ingredient for the stability of the DG scheme. We can take either of the following four choices of alternating fluxes for these four fluxes

$$\hat{v} = v_h^-, \quad \tilde{v}_x = (v_h)_x^-, \quad \hat{u} = u_h^+, \quad \tilde{u}_x = (u_h)_x^+; \quad (3.9)$$

$$\hat{v} = v_h^+, \quad \tilde{v}_x = (v_h)_x^+, \quad \hat{u} = u_h^-, \quad \tilde{u}_x = (u_h)_x^-; \quad (3.10)$$

$$\widehat{v} = v_h^-, \quad \widetilde{v}_x = (v_h)_x^+, \quad \widehat{u} = u_h^-, \quad \widetilde{u}_x = (u_h)_x^+; \quad (3.11)$$

$$\widehat{v} = v_h^+, \quad \widetilde{v}_x = (v_h)_x^-, \quad \widehat{u} = u_h^+, \quad \widetilde{u}_x = (u_h)_x^-. \quad (3.12)$$

It is crucial that $\widehat{v}$ and $\widetilde{u}_x$ come from the opposite sides, and $\widetilde{v}_x$ and $\widehat{u}$ come from the opposite sides (alternating fluxes).

Remark 3.1. *For the numerical fluxes, we can also take the following numerical fluxes*

$$\widehat{v} = \theta v_h^- + (1 - \theta) v_h^+, \quad \widetilde{v}_x = \theta (v_h)_x^- + (1 - \theta) (v_h)_x^+, \quad (3.13a)$$

$$\widehat{u} = \theta u_h^+ + (1 - \theta) u_h^-, \quad \widetilde{u}_x = \theta (u_h)_x^+ + (1 - \theta) (u_h)_x^-, \quad (3.13b)$$

where $0 \leq \theta \leq 1$. For $\theta = 1/2$, we would have the central fluxes as in [16] for the linear case. We note that, unlike in the UWDG method [3], here we do not need to add extra internal penalty terms to ensure stability.

3.2 Stability analysis

In this subsection, we will show the stability property of the scheme (3.6)-(3.8) with the choice of fluxes (3.9)-(3.13).

Theorem 3.1. *Our numerical scheme (3.6)-(3.8) with the choice of fluxes (3.9)-(3.13) is L^2 stable, i.e.*

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx + \int_{\Omega} b(u_h) w_h^2(x, t) dx = 0. \quad (3.14)$$

Proof. We integrate by parts in the scheme (3.6) and (3.8) and sum over j to obtain

$$((u_h)_t, p)_{\Omega} - ((v_h)_x, p_x)_{\Omega} + B_1(v_h, p) = 0, \quad (3.15)$$

$$(v_h, s)_{\Omega} - (b(u_h) w_h, s)_{\Omega} = 0, \quad (3.16)$$

$$(w_h, q)_{\Omega} + ((u_h)_x, q_x)_{\Omega} + B_2(u_h, q) = 0, \quad (3.17)$$

where

$$B_1(v_h, p) = \sum_{j=1}^N \left(v_h^- p_x^-|_{j+\frac{1}{2}} - v_h^+ p_x^+|_{j-\frac{1}{2}} + \widetilde{v}_x p^-|_{j+\frac{1}{2}} - \widetilde{v}_x p^+|_{j-\frac{1}{2}} - \widehat{v} p_x^-|_{j+\frac{1}{2}} + \widehat{v} p_x^+|_{j-\frac{1}{2}} \right), \quad (3.18)$$

$$B_2(u_h, q) = \sum_{j=1}^N \left(-u_h^- q_x^-|_{j+\frac{1}{2}} + u_h^+ q_x^+|_{j-\frac{1}{2}} - \widetilde{u}_x q^-|_{j+\frac{1}{2}} + \widetilde{u}_x q^+|_{j-\frac{1}{2}} \right)$$

$$+\widehat{u}q_x^-|_{j+\frac{1}{2}}-\widehat{u}q_x^+|_{j-\frac{1}{2}}\Big). \quad (3.19)$$

Then we take $p = u_h$, $s = -w_h$ and $q = v_h$ and add the three equalities (3.15)-(3.17) to obtain

$$\frac{1}{2}\frac{d}{dt}\int_{\Omega}u_h^2(x,t)dx+\int_{\Omega}b(u_h)w_h^2(x,t)dx+B_1(v_h,u_h)+B_2(u_h,v_h)=0. \quad (3.20)$$

However,

$$\begin{aligned} & B_1(v_h,u_h)+B_2(u_h,v_h) \\ &= \sum_{j=1}^N \left(v_h^-(u_h)_x^- - v_h^+(u_h)_x^+ + \widetilde{v}_x u_h^- - \widetilde{v}_x u_h^+ - \widehat{v}(u_h)_x^- + \widehat{v}(u_h)_x^+ \right. \\ & \quad \left. - u_h^-(v_h)_x^- + u_h^+(v_h)_x^+ - \widetilde{u}_x v_h^- + \widetilde{u}_x v_h^+ + \widehat{u}(v_h)_x^- - \widehat{u}(v_h)_x^+ \right) |_{j-\frac{1}{2}} \\ &= 0, \end{aligned} \quad (3.21)$$

for all of our flux choices (3.9)-(3.13). Then we have (3.14). $\square$

3.3 Error estimates

In this subsection, we state the error estimates of our scheme in the linear case, namely $b(u) = 1$. In this case, (3.7) in the scheme becomes a trivial statement $v_h = w_h$.

Theorem 3.2. *Let u be the exact solution of equation (3.1) with $b(u) = 1$, and $w = u_{xx}$, which are sufficiently smooth with bounded derivatives. Let u_h and w_h be solutions of (3.6), (3.8), with any choice of fluxes (3.9)-(3.12), and let V_h be the space of piecewise polynomials $\mathcal{P}^k$, $k \geq 1$, then we have the following error estimate:*

$$\|u(t) - u_h(t)\| + \int_0^t \|w(t) - w_h(t)\| dt \leq Ch^{k+1}, \quad (3.22)$$

where C is a constant independent of h and dependent on $\|u\|_{k+3}$, and on t .

Proof. Without loss of generality, we choose the flux (3.9). Let

$$e_u = u - u_h, \quad e_w = w - w_h$$

be the errors between the numerical and exact solutions. Since u and w clearly satisfy the scheme (3.6) and (3.8) as well, we can obtain the cell error equations: for all $p, q \in V_h$

$$((e_u)_t, p)_j + (e_w, p_{xx})_j + (e_w)_x^- p^-|_{j+\frac{1}{2}} - (e_w)_x^- p^+|_{j-\frac{1}{2}} - e_w^- p_x^-|_{j+\frac{1}{2}} + e_w^- p_x^+|_{j-\frac{1}{2}} = 0, \quad (3.23)$$

$$(e_w, q)_j - (e_u, q_{xx})_j - (e_u)_x^+ q^-|_{j+\frac{1}{2}} + (e_u)_x^+ q^+|_{j-\frac{1}{2}} + e_u^+ q_x^-|_{j+\frac{1}{2}} - e_u^+ q_x^+|_{j-\frac{1}{2}} = 0. \quad (3.24)$$

Since $k \geq 1$, we can choose a projection $P_{1h}^\pm$ defined in (2.2) and (2.3). Denote

$$\eta_u = u - P_{1h}^+ u, \quad \xi_u = u_h - P_{1h}^+ u, \quad \eta_w = w - P_{1h}^- w, \quad \xi_w = w_h - P_{1h}^- w,$$

and take $p = \xi_w$ and $q = \xi_u$ in (3.23) and (3.24) respectively. By the stability and property of projection $P_{1h}^\pm$ we have

$$((\xi_u)_t, \xi_u)_\Omega + (\xi_w, \xi_w)_\Omega = ((\eta_u)_t, \xi_u)_\Omega + (\eta_w, \xi_w)_\Omega. \quad (3.25)$$

Then

$$\frac{d}{dt} \|\xi_u\|^2 + \|\xi_w\|^2 \leq Ch^{k+1} \|\xi_u\| + Ch^{k+1} \|\xi_w\|.$$

Next we use Gronwall's inequality and choose $u_h(0) = P_{1h}^+ u(0)$ to obtain

$$\|\xi_u\|(t) + \int_0^t \|\xi_w\| dt \leq Ch^{k+1},$$

and

$$\|e_u\|(t) + \int_0^t \|e_w\| dt \leq \|\xi_u\|(t) + \int_0^t \|\xi_w\| dt + \|\eta_u\|(t) + \int_0^t \|\eta_w\| dt \leq Ch^{k+1},$$

where C is a constant independent of h and dependent on $\|u\|_{k+3}$, $\|u_t\|_{k+1}$, k and t . $\square$

4 The fifth order problem

Next we study the DG method for the following one-dimensional nonlinear fifth order equation

$$u_t + f(u_{xx})_{xxx} = 0, \quad (x, t) \in [0, 2\pi] \times (0, T], \quad (4.1)$$

$$u(x, 0) = u_0(x), \quad x \in \mathbb{R}, \quad (4.2)$$

with periodic boundary conditions, where $u_0(x)$ is a smooth function.

4.1 The numerical scheme

Similar to the fourth order problem (3.1), we rewrite (4.1) into a system:

$$u_t + w_{xx} = 0, \quad (4.3)$$

$$w - f(v)_x = 0, \quad (4.4)$$

$$v - u_{xx} = 0. \quad (4.5)$$

Then our DG method is defined as follows: find $u_h, w_h, v_h \in V_h$ such that for all $p, s, q \in V_h$, we have

$$((u_h)_t, p)_j + (w_h, p_{xx})_j + \widetilde{w}_x p^-|_{j+\frac{1}{2}} - \widetilde{w}_x p^+|_{j-\frac{1}{2}} - \widehat{w} p_x^-|_{j+\frac{1}{2}} + \widehat{w} p_x^+|_{j-\frac{1}{2}} = 0, \quad (4.6)$$

$$(w_h, s)_j + (f(v_h), s_x)_j - \widehat{f} s^-|_{j+\frac{1}{2}} + \widehat{f} s^+|_{j-\frac{1}{2}} = 0, \quad (4.7)$$

$$(v_h, q)_j - (u_h, q_{xx})_j - \widetilde{u}_x q^-|_{j+\frac{1}{2}} + \widetilde{u}_x q^+|_{j-\frac{1}{2}} + \widehat{u} q_x^-|_{j+\frac{1}{2}} - \widehat{u} q_x^+|_{j-\frac{1}{2}} = 0. \quad (4.8)$$

Here $\widehat{w}, \widetilde{w}_x, \widehat{f}, \widehat{u}, \widetilde{u}_x$ are numerical fluxes. We can take either of the following two choices for these five fluxes

$$\widehat{w} = w_h^-, \widetilde{w}_x = (w_h)_x^-, \widehat{f} = \widehat{f}(v_h^-, v_h^+), \widehat{u} = u_h^+, \widetilde{u}_x = (u_h)_x^+, \quad (4.9)$$

or

$$\widehat{w} = w_h^+, \widetilde{w}_x = (w_h)_x^+, \widehat{f} = \widehat{f}(v_h^-, v_h^+), \widehat{u} = u_h^-, \widetilde{u}_x = (u_h)_x^-, \quad (4.10)$$

where $\widehat{f}(v^-, v^+)$ is a monotone flux for $f(v)$. Here monotone flux means that the function $\widehat{f}$ is a non-decreasing function of its first argument and a non-increasing function of its second argument. It is also assumed to be at least Lipschitz continuous with respect to each argument and to be consistent with the physical flux $f(v)$ in the sense that $\widehat{f}(v, v) = f(v)$.

Remark 4.1. *It is crucial that $\widehat{w}$ and $\widetilde{u}_x$ come from the opposite sides, $\widetilde{w}_x$ and $\widehat{u}$ come from the opposite sides. We have at least four choices of these alternating fluxes or similar fluxes in (3.13), as in fourth order case. But here we just give the rule of alternating, and list part of them for simplicity.*

4.2 Stability analysis

In this subsection, we will show the stability property of the scheme (4.6)-(4.8) with the choice of fluxes (4.9) or (4.10).

Theorem 4.1. *Our scheme (4.6), (4.7) and (4.8) with the choice of fluxes (4.9) or (4.10) is stable, i.e*

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx \leq 0. \quad (4.11)$$

Proof. Integrate by parts in the scheme (4.6), (4.8) and sum over j , we obtain

$$((u_h)_t, p)_\Omega - ((w_h)_x, p_x)_\Omega + B_1(w_h, p) = 0, \quad (4.12)$$

$$(w_h, s)_\Omega + (f(v_h), s_x)_\Omega + B_3(f, s) = 0, \quad (4.13)$$

$$(v_h, q)_\Omega + ((u_h)_x, q_x)_\Omega + B_2(u_h, q) = 0, \quad (4.14)$$

where B_1 and B_2 have been defined before in (3.18) and (3.19), and

$$B_3(f, s) = \sum_{j=1}^N \left(-\widehat{f}s^-|_{j+\frac{1}{2}} + \widehat{f}s^+|_{j-\frac{1}{2}} \right). \quad (4.15)$$

Then we take $p = u_h$, $s = -v_h$ and $q = w_h$ and add the three equations to obtain

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx - (f(v_h), (v_h)_x)_\Omega + B_1(w_h, u_h) + B_3(f, -v_h) + B_2(u_h, w_h) = 0. \quad (4.16)$$

By (3.21), we have $B_1(w_h, u_h) + B_2(u_h, w_h) = 0$, then

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx + \sum_{j=1}^N (\widehat{G}_{j+\frac{1}{2}} - \widehat{G}_{j-\frac{1}{2}} + \Theta_{j-\frac{1}{2}}) = 0, \quad (4.17)$$

where

$$\widehat{G}_{j+\frac{1}{2}} = (-F(v_h^-) + \widehat{f}v_h^-)|_{j+\frac{1}{2}}, \quad F(v_h) = \int^{v_h} f(\tau) d\tau, \quad (4.18)$$

$$\Theta_{j-\frac{1}{2}} = (F(v_h^+) - F(v_h^-) + \widehat{f}v_h^- - \widehat{f}v_h^+)|_{j-\frac{1}{2}}, \quad (4.19)$$

for both of our flux choices (4.9) and (4.10). By the monotonicity of the fluxes $\widehat{f}$ and periodic boundary condition we obtain

$$\Theta_{j-\frac{1}{2}} \geq 0. \quad (4.20)$$

Then we have (4.11). □

Remark 4.2. We can also choose the central flux for nonlinear term $f(v)$

$$\widehat{f}_{j-\frac{1}{2}} = \frac{F(v_h^+) - F(v_h^-)}{v_h^+ - v_h^-} \Big|_{j-\frac{1}{2}},$$

then our scheme will be conservative, that means $\Theta_{j-\frac{1}{2}} = 0$ in (4.20) and

$$\frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx = 0.$$

4.3 Error estimates

In this subsection we consider the linear case, $f(v) = v$. Then we have the following optimal error estimate:

Theorem 4.2. *Let u be the exact solution of equation (4.1) with $f(v) = v$, and $w = u_{xxx}$, $v = u_{xx}$, which are sufficiently smooth with bounded derivatives. Let u_h, v_h, w_h be the numerical solutions obtained from the scheme (4.6)-(4.8) with the choice of fluxes (4.9) or (4.10) and $\hat{f}(v) = v^-$. If we use the V_h space with piecewise polynomials $\mathcal{P}^k$, $k \geq 1$, then we have the following error estimate:*

$$\|u(t) - u_h(t)\| + \|v(t) - v_h(t)\| + \|w(t) - w_h(t)\| \leq Ch^{k+1}, \quad (4.21)$$

where C is a constant independent of h and dependent on $\|u\|_{k+4}$, $\|u_t\|_{k+1}$, k and t .

To prove Theorem 4.2 we need some lemmas, addressing the relationship between the derivative and the element interface jump of the numerical solution and the auxiliary variable numerical solution of the derivative. This plays an important role in the error estimates analysis. Firstly, we have Lemma 4.1, which was proved in [22] for the LDG method and extended to the multi-dimensional case in [23].

Lemma 4.1. [22] *Suppose $(w_h, v_h) \in V_h \times V_h$ is the solution of the scheme (4.7) with $f(v) = v$, then there exists a positive constant C which is independent of h , such that $\forall j \in Z_N$*

$$\|(v_h)_x\|_{I_j} + h^{-\frac{1}{2}}|\llbracket v_h \rrbracket|_{j-\frac{1}{2}} \leq C\|w_h\|_{I_j}. \quad (4.22)$$

Next, we establish similar results for w_h in the equation (4.6) as in [22].

Lemma 4.2. *Suppose $(u_h, w_h) \in V_h \times V_h$ is the solution of the scheme (4.6), then there exists a positive constant C which is independent of h , such that $\forall j \in Z_N$*

$$\|(w_h)_{xx}\|_{I_j} + h^{-\frac{1}{2}}|\llbracket (w_h)_x \rrbracket|_{j+\frac{1}{2}} + h^{-\frac{3}{2}}|\llbracket w_h \rrbracket|_{j+\frac{1}{2}} \leq C\|(u_h)_t\|_{I_j}. \quad (4.23)$$

Proof. Without loss of generality, we choose the flux (4.10)

$$\hat{w} = w_h^+, \quad \widetilde{w}_x = (w_h)_x^+, \quad \hat{f} = v^-, \quad \hat{u} = u_h^-, \quad \widetilde{u}_x = (u_h)_x^-.$$

Recalling the equation (4.6), after integration by parts we have

$$((u_h)_t, p)_j + ((w_h)_{xx}, p)_j - \llbracket w_h \rrbracket_{j+\frac{1}{2}}(p_x)_{j+\frac{1}{2}}^- + \llbracket (w_h)_x \rrbracket_{j+\frac{1}{2}} p_{j+\frac{1}{2}}^- = 0. \quad (4.24)$$

Let L_k be the standard Legendre polynomial of degree k in $[-1, 1]$, we have $L_k(1) = 1$ and L_k is orthogonal to any polynomials with degree at most $k - 1$. First we take

$$p(x)|_{I_j} = (w_h)_{xx}(x) + AL_k(\xi) + BL_{k-1}(\xi),$$

in (4.6), with $\xi = \frac{2(x - x_j)}{h_j}$

$$A = -\frac{h_j(w_h)_{xxx}(x_{j+\frac{1}{2}}^-)}{2k} + \frac{L'_{k-1}(1)(w_h)_{xx}(x_{j+\frac{1}{2}}^-)}{k},$$

and

$$B = \frac{h_j(w_h)_{xxx}(x_{j+\frac{1}{2}}^-)}{2k} - \frac{L'_{k-1}(1)(w_h)_{xx}(x_{j+\frac{1}{2}}^-)}{k} - (w_h)_{xx}(x_{j+\frac{1}{2}}^-),$$

$p(x) \in V_h$ and is well defined since $k \geq 1$ in our function space. Clearly, there hold $p(x_{j+\frac{1}{2}}^-) = 0$, $p_x(x_{j+\frac{1}{2}}^-) = 0$, and $((w_h)_{xx}, p)_j = ((w_h)_{xx}, (w_h)_{xx})_j$. By (4.24) we have

$$((u_h)_t, p)_j + ((w_h)_{xx}, (w_h)_{xx})_j = 0.$$

Thus

$$\begin{aligned} \|(w_h)_{xx}\|_j^2 &\leq \|(u_h)_t\|_j (\|(w_h)_{xx}\|_j + |A|\|L_k(\xi)\|_j + |B|\|L_{k-1}(\xi)\|_j) \\ &\leq C\|(u_h)_t\|_j \|(w_h)_{xx}\|_j, \end{aligned}$$

where the first inequality is obtained by using the Cauchy-Schwartz inequality and the second is derived by using the inverse inequality and the fact $\|L_k(\xi)\|_j \leq Ch^{\frac{1}{2}}$. Therefore,

$$\|(w_h)_{xx}\|_j \leq C\|(u_h)_t\|_j. \quad (4.25)$$

Next we take $p = 1$ in (4.24) to obtain

$$((u_h)_t, 1)_j + ((w_h)_{xx}, 1)_j + \llbracket (w_h)_x \rrbracket_{j+\frac{1}{2}} = 0,$$

then, by (4.25) and the Cauchy-Schwartz inequality we get

$$|\llbracket (w_h)_x \rrbracket_{j+\frac{1}{2}}| \leq h^{\frac{1}{2}} (\|(u_h)_t\|_j + \|(w_h)_{xx}\|_j) \leq Ch^{\frac{1}{2}} \|(u_h)_t\|_j. \quad (4.26)$$

Our next choice of the test function is $p = \xi$ in (4.24), which gives

$$((u_h)_t, \xi)_j + ((w_h)_{xx}, \xi)_j - \frac{2}{h_j} \llbracket w_h \rrbracket_{j+\frac{1}{2}} + \llbracket (w_h)_x \rrbracket_{j+\frac{1}{2}} = 0.$$

By (4.25), (4.26) and the Cauchy-Schwartz inequality we get

$$|\llbracket w_h \rrbracket_{j+\frac{1}{2}}| \leq Ch^{\frac{3}{2}} (\|(u_h)_t\|_j + \|(w_h)_{xx}\|_j) \leq Ch^{\frac{3}{2}} \|(u_h)_t\|_j. \quad (4.27)$$

Finally, we get the desired result (4.23). $\square$

Based on the relationship constructed in the Lemma 4.1 and Lemma 4.2, we can easily use the discrete Poincaré inequalities [1, 2] to estimate w_h and v_h .

Lemma 4.3. *Let $(u_h, v_h, w_h) \in V_h$ be the solutions of the scheme (4.6)-(4.8), then there exists a positive constant C which are independent of h , such that*

$$\|(w_h)_x\| \leq C\|(u_h)_t\|, \quad (4.28)$$

$$\|w_h\| \leq C\|(u_h)_t\|, \quad (4.29)$$

$$\|v_h\| \leq C\|w_h\|. \quad (4.30)$$

With all these preparations, we can start the proof of Theorem 4.2.

Proof. (**The proof of Theorem 4.2**)

Without loss of generality, we choose the flux (4.10). Let

$$e_u = u - u_h, \quad e_v = v - v_h, \quad e_w = w - w_h$$

be the errors between the numerical and exact solutions. Since u , v and w clearly satisfy (4.6)-(4.8) we can obtain the cell error equations: for all $p, s, q \in V_h$

$$((e_u)_t, p)_j + (e_w, p_{xx})_j + (e_w)_x p^-|_{j+\frac{1}{2}} - (e_w)_x p^+|_{j-\frac{1}{2}} - e_w^+ p_x^-|_{j+\frac{1}{2}} + e_w^+ p_x^+|_{j-\frac{1}{2}} = 0, \quad (4.31)$$

$$(e_w, s)_j + (e_v, s_x)_j - e_v^- s^-|_{j+\frac{1}{2}} + e_v^- s^+|_{j-\frac{1}{2}} = 0, \quad (4.32)$$

$$(e_v, q)_j - (e_u, q_{xx})_j - (e_u)_x q^-|_{j+\frac{1}{2}} + (e_u)_x q^+|_{j-\frac{1}{2}} + e_u^- q_x^-|_{j+\frac{1}{2}} - e_u^- q_x^+|_{j-\frac{1}{2}} = 0. \quad (4.33)$$

Since $k \geq 1$ we choose the projections P_{1h}^+ , and P_h^- , which are defined in (2.1)-(2.3). Denote

$$\begin{aligned} \eta_u &= u - P_{1h}^- u, \quad \xi_u = u_h - P_{1h}^- u, \\ \eta_w &= w - P_{1h}^+ w, \quad \xi_w = w_h - P_{1h}^+ w, \\ \eta_v &= v - P_h^- v, \quad \xi_v = v_h - P_h^- v. \end{aligned}$$

Furthermore by the error equations (4.31)-(4.33) and Lemma 4.1, Lemma 4.2 and Lemma 4.3 we have

$$\|\xi_w\| \leq C\|(e_u)_t\| \leq C\|(\xi_u)_t\| + Ch^{k+1}, \quad (4.34)$$

$$\|\xi_v\| \leq C\|e_w\| \leq C\|\xi_w\| + Ch^{k+1}. \quad (4.35)$$

- **Error estimates for the initial condition.**

We choose the initial condition $u_h(x, 0)$ such that

$$w_h(x, 0) = P_{1h}^+ w(x, 0), \quad w(x, 0) = u_{xxx}(x, 0). \quad (4.36)$$

Then we have

$$\|w(x, 0) - w_h(x, 0)\| \leq Ch^{k+1}.$$

By (4.34) and (4.35) we get

$$\|\xi_v\| \leq \|\xi_w\| + Ch^{k+1} \leq Ch^{k+1},$$

$$\|\xi_u\| \leq \|\xi_v\| + Ch^{k+1} \leq Ch^{k+1},$$

and we have the following estimates:

$$\|u(x, 0) - u_h(x, 0)\| + \|v(x, 0) - v_h(x, 0)\| + \|w(x, 0) - w_h(x, 0)\| \leq Ch^{k+1}. \quad (4.37)$$

Next we choose $t = 0$ in (4.31), due to the choice of $w_h(x, 0)$ we have

$$(u_t(0) - (u_h)_t(0), p)_j = 0.$$

Now, we choose $p = (u_h)_t(0) - P(u_t(0))$, P is the standard L^2 projection, and obtain

$$\|u_t(x, 0) - (u_h)_t(0)\| \leq Ch^{k+1}. \quad (4.38)$$

• **Error estimates for $t > 0$.**

Then we take $p = \xi_u$, $s = -\xi_v$ and $q = \xi_w$, and add the three equations (4.31)-(4.33) and also sum over j . By the stability and the properties of the projections we can obtain

$$((\xi_u)_t, \xi_u)_\Omega + \sum_{j=1}^N \|\xi_v\|_{j-\frac{1}{2}}^2 = ((\eta_u)_t, \xi_u)_\Omega - (\eta_w, \xi_v)_\Omega + (\eta_v, \xi_w)_\Omega.$$

Next, we take the time derivative of the three error equations (4.31)-(4.33), and take $p = (\xi_u)_t$, $s = -(\xi_v)_t$ and $q = (\xi_w)_t$ to obtain

$$((\xi_u)_{tt}, (\xi_u)_t)_\Omega + \sum_{j=1}^N \|(\xi_v)_t\|_{j-\frac{1}{2}}^2 = ((\eta_u)_{tt}, (\xi_u)_t)_\Omega - ((\eta_w)_t, (\xi_v)_t)_\Omega + ((\eta_v)_t, (\xi_w)_t)_\Omega.$$

Now, combining the energy equations we get

$$\frac{1}{2} \frac{d}{dt} (\|\xi_u\|^2 + \|(\xi_u)_t\|^2) + \sum_{j=1}^N (\|\xi_v\|_{j-\frac{1}{2}}^2 + \|(\xi_v)_t\|_{j-\frac{1}{2}}^2) = \Upsilon + \Lambda, \quad (4.39)$$

where

$$\begin{aligned}\Upsilon &= ((\eta_u)_t, \xi_u)_\Omega - (\eta_w, \xi_v)_\Omega + (\eta_v, \xi_w)_\Omega + ((\eta_u)_{tt}, (\xi_u)_t)_\Omega, \\ \Lambda &= -((\eta_w)_t, (\xi_v)_t)_\Omega + ((\eta_v)_t, (\xi_w)_t)_\Omega.\end{aligned}$$

By (4.34), (4.35) we have the estimate

$$\|\xi_v\| \leq C\|\xi_w\| + Ch^{k+1}, \quad \|\xi_w\| \leq C\|(\xi_u)_t\| + Ch^{k+1},$$

then we can easily get

$$\Upsilon \leq Ch^{k+1}\|\xi_u\| + Ch^{k+1}\|(\xi_u)_t\| + Ch^{2k+2}.$$

Next, integrating Λ with respect to time between 0 and t , we can get the following equation after integration by parts:

$$\int_0^t \Lambda dt = -((\eta_w)_t, \xi_v)_\Omega|_0^t + \int_0^t ((\eta_w)_{tt}, \xi_v)_\Omega dt + ((\eta_v)_t, \xi_w)_\Omega|_0^t - \int_0^t ((\eta_v)_{tt}, \xi_w)_\Omega dt.$$

We can easily get the following estimates using the approximation property of the projections and the estimates for the initial condition

$$\begin{aligned}\left| \int_0^t \Lambda dt \right| &\leq Ch^{2k+2} + \|\xi_v\|^2 + \|\xi_w\|^2 + \int_0^t (\|\xi_v\|^2 + \|\xi_w\|^2) dt \\ &\leq Ch^{2k+2} + Ch^{k+1} \int_0^t \|(\xi_u)_t\| dt.\end{aligned}$$

Now we integrate (4.39) with respect to the time between 0 to t , using the Cauchy-Schwartz inequality and (4.37), (4.38) to obtain

$$\frac{1}{2}(\|\xi_u\|^2 + \|(\xi_u)_t\|^2) \leq \frac{1}{4} \int_0^t \|\xi_u\|^2 + \|(\xi_u)_t\|^2 dt + Ch^{2k+2}.$$

After employing the Gronwall's inequality, we get

$$\max_t \|\xi_u\| + \max_t \|(\xi_u)_t\| \leq Ch^{k+1},$$

and also

$$\max_t \|\xi_w\| + \max_t \|\xi_v\| \leq Ch^{k+1}.$$

After using the standard approximation results, we can get (4.21). $\square$

5 Extension to high order equations

The DG method introduced in the previous sections as well as the theoretical analysis for the stability and error estimates can be extended to more general high order PDEs, and to multidimensional cases. Firstly, we consider the extension to the general high order equations,

$$u_t + (-1)^{[\frac{n}{2}]} u_x^n = 0, \quad (5.1)$$

with n being any positive integer. Here u_x^n denotes the n -th derivative of u with respect to x , and $[\frac{n}{2}]$ is the integer part of $\frac{n}{2}$.

In the first two subsections, we will give two specific examples to introduce our scheme to sixth and seventh order equations. Then we will summarize to the general case.

5.1 Extension to sixth order equations

In this subsection, we will consider the sixth order equation:

$$u_t - u_x^{(6)} = 0, \quad (x, t) \in [0, 2\pi] \times (0, T], \quad (5.2)$$

$$u(x, 0) = u_0(x), \quad x \in \mathbb{R}, \quad (5.3)$$

where $u_0(x)$ is a smooth function, as an example of even order diffusive equations. For simplicity of discussion, we will again only consider the periodic boundary conditions. Firstly, we rewrite the sixth order equation into a system of third order equations

$$u_t - w_{xxx} = 0, \quad (5.4)$$

$$w - u_{xxx} = 0. \quad (5.5)$$

Then our DG method is defined as follows: find $u_h, w_h \in V_h$ such that for all $p, q \in V_h$, we have

$$\begin{aligned} & ((u_h)_t, p)_j + (w_h, p_{xxx})_j - \widetilde{w_{xx}p^-}|_{j+\frac{1}{2}} + \widetilde{w_{xx}p^+}|_{j-\frac{1}{2}} + \widetilde{w_xp^-}|_{j+\frac{1}{2}} - \widetilde{w_xp^+}|_{j-\frac{1}{2}} \\ & - \widetilde{w}p_{xx}^-|_{j+\frac{1}{2}} + \widetilde{w}p_{xx}^+|_{j-\frac{1}{2}} = 0, \end{aligned} \quad (5.6)$$

$$\begin{aligned} & (w_h, q)_j + (u_h, q_{xxx})_j - \widetilde{u_{xx}q^-}|_{j+\frac{1}{2}} + \widetilde{u_{xx}q^+}|_{j-\frac{1}{2}} + \widetilde{u_xq^-}|_{j+\frac{1}{2}} - \widetilde{u_xq^+}|_{j-\frac{1}{2}} \\ & - \widetilde{u}q_{xx}^-|_{j+\frac{1}{2}} + \widetilde{u}q_{xx}^+|_{j-\frac{1}{2}} = 0. \end{aligned} \quad (5.7)$$

Here $\widetilde{w}$, $\widetilde{w_x}$, $\widetilde{w_{xx}}$, $\widetilde{u}$, $\widetilde{u_x}$, and $\widetilde{u_{xx}}$ are the numerical fluxes. The terms involving these numerical fluxes appear from repeated integration by parts. We can take either of the

following two choices for these six fluxes

$$\widetilde{w} = w_h^-, \quad \widetilde{w}_x = (w_h)_x^-, \quad \widetilde{w}_{xx} = (w_h)_{xx}^-, \quad \widehat{u} = u_h^+, \quad \widehat{u}_x = (u_h)_x^+, \quad \widehat{u}_{xx} = (u_h)_{xx}^+, \quad (5.8)$$

or

$$\widetilde{w} = w_h^+, \quad \widetilde{w}_x = (w_h)_x^+, \quad \widetilde{w}_{xx} = (w_h)_{xx}^+, \quad \widehat{u} = u_h^-, \quad \widehat{u}_x = (u_h)_x^-, \quad \widehat{u}_{xx} = (u_h)_{xx}^-. \quad (5.9)$$

It is crucial that we take the pair $\widehat{u}$ and $\widetilde{w}_{xx}$ from opposite sides, the pair $\widehat{u}_x$ and $\widetilde{w}_x$ from opposite sides, and the pair $\widehat{u}_{xx}$ and $\widetilde{w}$ from opposite sides.

Theorem 5.1. (Stability) *Our scheme (5.6)-(5.7) with the choice of fluxes (5.8) or (5.9) is L^2 stable, i.e.*

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx + \int_{\Omega} w_h^2(x, t) dx = 0. \quad (5.10)$$

Proof. Integrating by parts in the scheme (5.6)-(5.7) and summing over j , we have

$$((u_h)_t, p)_{\Omega} - ((w_h)_{xxx}, p)_{\Omega} + B_4(w_h, p) = 0, \quad (5.11)$$

$$(w_h, q)_{\Omega} + (u_h, q_{xxx})_{\Omega} + B_5(u_h, q) = 0, \quad (5.12)$$

where

$$\begin{aligned} B_4(w_h, p) = & \sum_{j=1}^N \left(w_h^- p_{xx}^-|_{j+\frac{1}{2}} - w_h^+ p_{xx}^+|_{j-\frac{1}{2}} - (w_h)_x^- p_x^-|_{j+\frac{1}{2}} + (w_h)_x^+ p_x^+|_{j-\frac{1}{2}} \right. \\ & + (w_h)_{xx}^- p^-|_{j+\frac{1}{2}} - (w_h)_{xx}^+ p^+|_{j-\frac{1}{2}} - \widetilde{w}_{xx} p^-|_{j+\frac{1}{2}} + \widetilde{w}_{xx} p^+|_{j-\frac{1}{2}} \\ & \left. + \widetilde{w}_x p_x^-|_{j+\frac{1}{2}} - \widetilde{w}_x p_x^+|_{j-\frac{1}{2}} - \widetilde{w} p_{xx}^-|_{j+\frac{1}{2}} + \widetilde{w} p_{xx}^+|_{j-\frac{1}{2}} \right), \end{aligned} \quad (5.13)$$

$$\begin{aligned} B_5(u_h, q) = & \sum_{j=1}^N \left(-\widehat{u}_{xx} q^-|_{j+\frac{1}{2}} + \widehat{u}_{xx} q^+|_{j-\frac{1}{2}} + \widehat{u}_x q_x^-|_{j+\frac{1}{2}} - \widehat{u}_x q_x^+|_{j-\frac{1}{2}} \right. \\ & \left. - \widehat{u} q_{xx}^-|_{j+\frac{1}{2}} + \widehat{u} q_{xx}^+|_{j-\frac{1}{2}} \right). \end{aligned} \quad (5.14)$$

Then we take $p = u_h$ and $q = w_h$ and add the two equations (5.11)-(5.12) to obtain

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx + \int_{\Omega} w_h^2(x, t) dx + B_4(w_h, u_h) + B_5(u_h, w_h) = 0. \quad (5.15)$$

We can easily check that

$$B_4(w_h, u_h) + B_5(u_h, w_h) = 0,$$

for both of our flux choices (5.8) and (5.9). Then we have (5.10). $\square$

Theorem 5.2. (Error estimates) Let u be the exact solution of the equation (5.2) and $w = u_{xxx}$, which are sufficiently smooth with bounded derivatives. Let u_h and w_h be solutions of the scheme (5.6)-(5.7) with either (5.8) or (5.9) as the numerical fluxes, and let V_h be the space of piecewise polynomials $\mathcal{P}^k$, $k \geq 2$, then we have the following error estimate

$$\|u(t) - u_h(t)\| + \int_0^t \|w(t) - w_h(t)\| dt \leq Ch^{k+1}, \quad (5.16)$$

where C is a constant independent of h and dependent on $\|u\|_{k+4}$, and t .

Proof. The proof is similar to that of Theorem 3.2. By using the projection $P_{2h}^\pm$ defined in (2.4)-(2.5) for $k \geq 2$ and then following the line of proof for Theorem 3.2, we can easily get the result (5.16). $\square$

5.2 Extension to seventh order equations

In this subsection, we will give the formulation of the scheme as well as its theoretical results for the seventh order wave equation

$$u_t - u_x^{(7)} = 0, \quad (x, t) \in [0, 2\pi] \times (0, T], \quad (5.17)$$

$$u(x, 0) = u_0(x), \quad x \in \mathbb{R}, \quad (5.18)$$

where $u_0(x)$ is a smooth function, as an example of general odd order wave equations. As mentioned before, we only consider the periodic boundary conditions. Similar to the sixth order equation, firstly, we rewrite (5.17) into a system:

$$u_t - w_{xxx} = 0, \quad (5.19)$$

$$w - v_x = 0, \quad (5.20)$$

$$v - u_{xxx} = 0. \quad (5.21)$$

Then our DG method defined as follows: find $u_h, v_h, w_h \in V_h$ such that for all $p, s, q \in V_h$, we have

$$\begin{aligned} & ((u_h)_t, p)_j + (w_h, p_{xxx})_j - \widetilde{w_{xx}p^-}|_{j+\frac{1}{2}} + \widetilde{w_{xx}p^+}|_{j-\frac{1}{2}} + \widetilde{w_xp^-}|_{j+\frac{1}{2}} - \widetilde{w_xp^+}|_{j-\frac{1}{2}} \\ & - \widetilde{wp_{xx}^-}|_{j+\frac{1}{2}} + \widetilde{wp_{xx}^+}|_{j-\frac{1}{2}} = 0, \end{aligned} \quad (5.22)$$

$$(w_h, s)_j + (v_h, s_x)_j - \widehat{vs^-}|_{j+\frac{1}{2}} + \widehat{vs^+}|_{j-\frac{1}{2}} = 0, \quad (5.23)$$

$$\begin{aligned} & (v_h, q)_j + (u_h, q_{xxx})_j - \widehat{u_{xx}q^-}|_{j+\frac{1}{2}} + \widehat{u_{xx}q^+}|_{j-\frac{1}{2}} + \widehat{u_xq^-}|_{j+\frac{1}{2}} - \widehat{u_xq^+}|_{j-\frac{1}{2}} \\ & - \widehat{uq_{xx}^-}|_{j+\frac{1}{2}} + \widehat{uq_{xx}^+}|_{j-\frac{1}{2}} = 0. \end{aligned} \quad (5.24)$$

Here $\widetilde{w}$, $\widetilde{w}_x$, $\widetilde{w}_{xx}$, $\widehat{v}$, $\widehat{u}$, $\widehat{u}_x$, $\widehat{u}_{xx}$ are numerical fluxes. For example, we can take either of the following two choices for these fluxes

$$\widetilde{w} = w_h^-, \quad \widetilde{w}_x = (w_h)_x^-, \quad \widetilde{w}_{xx} = (w_h)_{xx}^-, \quad \widehat{v} = v_h^-, \quad \widehat{u} = u_h^+, \quad \widehat{u}_x = (u_h)_x^+, \quad \widehat{u}_{xx} = (u_h)_{xx}^+, \quad (5.25)$$

or

$$\widetilde{w} = w_h^+, \quad \widetilde{w}_x = (w_h)_x^+, \quad \widetilde{w}_{xx} = (w_h)_{xx}^+, \quad \widehat{v} = v_h^-, \quad \widehat{u} = u_h^-, \quad \widehat{u}_x = (u_h)_x^-, \quad \widehat{u}_{xx} = (u_h)_{xx}^-. \quad (5.26)$$

It is crucial that we take $\widehat{v} = v_h^-$ by upwinding, the pair $\widehat{u}$ and $\widetilde{w}_{xx}$ from opposite sides, the pair $\widehat{u}_x$ and $\widetilde{w}_x$ from opposite sides, and the pair $\widehat{u}_{xx}$ and $\widetilde{w}$ from opposite sides.

Theorem 5.3. (*Stability*) *Our scheme (5.22)-(5.24) with the choice of fluxes (5.25) or (5.26) is stable, i.e.*

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx \leq 0. \quad (5.27)$$

Proof. Integrating by parts in the scheme (5.22)-(5.24) and summing over j , we have

$$((u_h)_t, p)_{\Omega} - ((w_h)_{xxx}, p)_{\Omega} + B_4(w_h, p) = 0, \quad (5.28)$$

$$(w_h, s)_{\Omega} + (v_h, s_x)_{\Omega} + B_3(v_h, s) = 0, \quad (5.29)$$

$$(v_h, q)_{\Omega} + (u_h, q_{xxx})_{\Omega} + B_5(u_h, q) = 0, \quad (5.30)$$

where B_3 , B_4 and B_5 are defined in (4.15), (5.13) and (5.14), respectively. Then we take $p = u_h$, $s = -v_h$ and $q = w_h$ in (5.28), (5.29) and (5.30) respectively, add the three equations to obtain

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx + \frac{1}{2} \sum_{j=1}^N (\llbracket v_h \rrbracket)_{j-\frac{1}{2}}^2 = 0, \quad (5.31)$$

for both of our flux choices (5.25) and (5.26). Then we have (5.27). $\square$

Theorem 5.4. (*Error estimates*) *Let u be the exact solution of the equation (5.17), and $w = u_{xxxx}$, $v = u_{xxx}$, which are sufficiently smooth with bounded derivatives. Let u_h , v_h , w_h be the numerical solutions of (5.22)-(5.24). If we use V_h as the space with piecewise polynomials $\mathcal{P}^k$, $k \geq 2$, then we have the following error estimate:*

$$\|u(t) - u_h(t)\| + \|v(t) - v_h(t)\| + \|w(t) - w_h(t)\| \leq Ch^{k+1}, \quad (5.32)$$

where C is a constant independent of h and dependent on $\|u\|_{k+5}$, $\|u_t\|_{k+1}$, k and t .

Proof. The proof is similar to that of Theorem 4.2 and is thus omitted to save space. $\square$

5.3 Extension to general high order cases

We have introduced the numerical schemes for sixth and seventh order cases. More generally, we summarize the scheme for any high order case. The proof of stability and error estimate is similar to the sixth and seventh equations, therefore we just list the results and omit the proof. Again, we only consider the periodic boundary conditions.

5.3.1 General even order case

Let n be a positive even number, and consider the equation

$$u_t + (-1)^{\frac{n}{2}} u_x^n = 0. \quad (5.33)$$

Firstly, we rewrite it into a $\frac{n}{2}$ -th order system,

$$u_t + (-1)^{\frac{n}{2}} w_x^{\frac{n}{2}} = 0, \quad (5.34)$$

$$w - u_x^{\frac{n}{2}} = 0. \quad (5.35)$$

Then our DG method is defined as follows: find $u_h, w_h \in V_h$ such that for all $p, q \in V_h$, we have

$$((u_h)_t, p)_j + (w_h, p_x^{\frac{n}{2}})_j + \sum_{m=0}^{\frac{n}{2}-1} \left((-1)^{\frac{n}{2}+m} \left(\widetilde{w}_x^{\frac{n}{2}-1-m} (p_x^m)^-|_{j+\frac{1}{2}} - \widetilde{w}_x^{\frac{n}{2}-1-m} (p_x^m)^+|_{j-\frac{1}{2}} \right) \right) = 0, \quad (5.36)$$

$$(w_h, q)_j - (-1)^{\frac{n}{2}} (u_h, q_x^{\frac{n}{2}})_j + \sum_{m=0}^{\frac{n}{2}-1} \left((-1)^{m+1} \left(\widehat{u}_x^{\frac{n}{2}-1-m} (q_x^m)^-|_{j+\frac{1}{2}} - \widehat{u}_x^{\frac{n}{2}-1-m} (q_x^m)^+|_{j-\frac{1}{2}} \right) \right) = 0. \quad (5.37)$$

Remark 5.1. We choose alternating fluxes. It is crucial that we take $\widetilde{w}_x^{\frac{n}{2}-1-m}$ and $\widehat{u}_x^m$ from opposite sides, $m = 0, 1, \dots, \frac{n}{2} - 1$.

Theorem 5.5. (Stability) Our scheme (5.36)-(5.37) with the choice of alternating fluxes in Remark 5.1 is L^2 stable, i.e.

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx + \int_{\Omega} w_h^2(x, t) dx = 0. \quad (5.38)$$

Theorem 5.6. (Error estimates) Let u be the exact solution of the equation (5.33), and $w = u_x^{\frac{n}{2}}$, which are sufficiently smooth with bounded derivatives. Let u_h, w_h be the numerical solutions of (5.36)-(5.37) with alternating fluxes in Remark 5.1. If we use V_h

as the space with piecewise polynomials $\mathcal{P}^k$, $k \geq \frac{n}{2} - 1$, then we have the following error estimate:

$$\|u(t) - u_h(t)\| + \int_0^t \|w(t) - w_h(t)\| dt \leq Ch^{k+1}, \quad (5.39)$$

where C is a constant independent of h .

5.3.2 General odd order case

Let n be an odd number, and $n \geq 3$. We consider the following equation:

$$u_t + u_x^n = 0, \quad (5.40)$$

Firstly, we rewrite it into a $(\frac{n-1}{2})$ -th order system,

$$u_t + w_x^{\frac{n-1}{2}} = 0, \quad (5.41)$$

$$w - v_x = 0, \quad (5.42)$$

$$v - u_x^{\frac{n-1}{2}} = 0. \quad (5.43)$$

Then our DG method is defined as follows: find $u_h, v_h, w_h \in V_h$ such that for all $p, s, q \in V_h$, we have

$$((u_h)_t, p)_j + (-1)^{\frac{n-1}{2}} (w_h, p_x^{\frac{n-1}{2}})_j + \sum_{m=0}^{\frac{n-3}{2}} \left((-1)^m \left(\widetilde{w}_x^{\frac{n-3}{2}-m} (p_x^m)^-|_{j+\frac{1}{2}} - \widetilde{w}_x^{\frac{n-3}{2}-m} (p_x^m)^+|_{j-\frac{1}{2}} \right) \right) = 0. \quad (5.44)$$

$$(w_h, s)_j + (v_h, s_x)_j - \widehat{v} s^-|_{j+\frac{1}{2}} + \widehat{v} s^+|_{j-\frac{1}{2}} = 0, \quad (5.45)$$

$$(v_h, q)_j - (-1)^{\frac{n-1}{2}} (u_h, q_x^{\frac{n-1}{2}})_j + \sum_{m=0}^{\frac{n-3}{2}} \left((-1)^{m+1} \left(\widetilde{u}_x^{\frac{n-3}{2}-m} (q_x^m)^-|_{j+\frac{1}{2}} - \widetilde{u}_x^{\frac{n-3}{2}-m} (q_x^m)^+|_{j-\frac{1}{2}} \right) \right) = 0. \quad (5.46)$$

Remark 5.2. It is crucial that we take $\widehat{v}$ by upwinding, the pairs $\widetilde{w}_x^{\frac{n-3}{2}-m}$ and $\widetilde{u}_x^m$ from opposite sides, $m = 0, 1, \dots, \frac{n-3}{2}$.

Theorem 5.7. (Stability) Our scheme (5.44)-(5.46) with the choice of fluxes in Remark 5.2 is stable, i.e.

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} u_h^2(x, t) dx \leq 0. \quad (5.47)$$

Theorem 5.8. (*Error estimates*) *Let u be the exact solution of the equation (5.40), and $v = u_x^{\frac{n-1}{2}}$, $w = v_x$, , which are sufficiently smooth with bounded derivatives. Let u_h , v_h , w_h be the numerical solutions of (5.44)-(5.46) with the choice of fluxes in Remark 5.2. If we use V_h as the space with piecewise polynomials $\mathcal{P}^k$, $k \geq \frac{n-3}{2}$, then we have the following error estimate:*

$$\|u(t) - u_h(t)\| + \|v(t) - v_h(t)\| + \|w(t) - w_h(t)\| \leq Ch^{k+1}, \quad (5.48)$$

where C is a constant independent of h .

6 Extension to the fourth order equation in multi-dimensional Cartesian meshes

In this section, we will extend our DG scheme to multi-dimensional Cartesian meshes for fourth-order equation, as an example of multi-dimensional extension of our schemes. Without loss of generality, we describe our DG method and prove a priori optimal error estimates in two dimensions ($d = 2$), however all the arguments we present in our analysis depend on the tensor product structure of the meshes and can be easily extended to higher dimensions ($d > 2$).

Hence, from now on, we shall restrict ourselves to the following two-dimensional problem:

$$u_t + \Delta^2 u = 0, \quad (\mathbf{x}, t) \in \Omega \times (0, T], \quad (6.1)$$

with the periodic boundary condition and initial condition

$$u(\mathbf{x}, 0) = u_0(\mathbf{x}),$$

where $u_0(\mathbf{x})$ is a smooth function of $\mathbf{x} = (x, y)$, $\Omega \in \mathbb{R}^2$ is a bounded rectangular domain.

6.1 The numerical scheme

Firstly, we rewrite the fourth-order equation (6.1) into a system of second-order equations,

$$u_t + \Delta w = 0, \quad (6.2)$$

$$w - \Delta u = 0. \quad (6.3)$$

In order to define our DG method for the system (6.2)-(6.3), let us introduce some notations. Let Ω_h denote a tessellation of Ω with shape-regular elements K , and the union of the boundary face of element $K \in \Omega_h$, denoted as $\partial\Omega = \bigcup_{K \in \Omega_h} \partial K$. We denote the diameter of K by h_K , and set $h = \max_K h_K$. The finite element spaces with the mesh Ω_h are of the form

$$W_h = \{\eta \in L^2(\Omega) : \eta|_K \in \mathcal{Q}^k(K), \forall K \in \Omega_h\},$$

where $\mathcal{Q}^k(K)$ is the space of tensor product of polynomials of degree at most $k \geq 0$ on $K \in \Omega_h$ in each variable defined on K .

Since the approximation space in discontinuous Galerkin methods consists of piecewise polynomials, we need to have a way of denoting the value of the approximation on the “left” and “right” side of an element boundary e . We give the designation K_L for element to the left side of e , and K_R for element to the right side of e (We refer to [27] for a proper definition of “left” and “right” in our context, for rectangular meshes these are the usual left and bottom directions denoted as “left” and right and top directions denoted as “right”). The normal vector ν_L and ν_R on the edge e point exterior to K_L and K_R respectively. Assuming ψ is a function defined on K_L and K_R , let ψ^- denote $(\psi|_{K_L})|_e$ and ψ^+ denote $(\psi|_{K_R})|_e$, the left and right traces, respectively. The DG method is defined as following: we seek u_h and w_h in the finite element space $W_h \times W_h$, such that for all $p, q \in W_h$ we have

$$((u_h)_t, p)_K + (w_h, \Delta p)_K + \langle \widehat{\nabla w} \cdot \mathbf{n}, p \rangle_{\partial K} - \langle \widetilde{w}, \nabla p \cdot \mathbf{n} \rangle_{\partial K} = 0, \quad (6.4)$$

$$(w_h, q)_K - (u_h, \Delta q)_K - \langle \widehat{\nabla u} \cdot \mathbf{n}, q \rangle_{\partial K} + \langle \widehat{u}, \nabla q \cdot \mathbf{n} \rangle_{\partial K} = 0. \quad (6.5)$$

Here $\mathbf{n}$ denotes the outward unit vector to ∂K , and

$$(p, q)_K := \int_K p(x, y) q(x, y) dx dy, \quad \langle p, \nabla q \cdot \mathbf{n} \rangle = \int_{\partial K} p(x, y) (\nabla q(x, y) \cdot \mathbf{n}) ds, \quad (6.6)$$

for any $p, q \in H_{\Omega_h}^1$. To complete the definition of the DG scheme we need to define the numerical fluxes $\widehat{u}$, $\widehat{\nabla u}$, $\widetilde{w}$, $\widetilde{\nabla w}$. We can choose the alternating fluxes

$$\widehat{u} = u_h^+, \quad \widehat{\nabla u} = (\nabla u_h)^+, \quad \widetilde{w} = w_h^-, \quad \widetilde{\nabla w} = (\nabla w_h)^-, \quad (6.7)$$

or

$$\widehat{u} = u_h^-, \quad \widehat{\nabla u} = (\nabla u_h)^-, \quad \widetilde{w} = w_h^+, \quad \widetilde{\nabla w} = (\nabla w_h)^+. \quad (6.8)$$

6.2 L^2 stability

In this subsection, we will prove the DG method defined in (6.4)-(6.5) for the fourth-order equation satisfies the following L^2 stability.

Theorem 6.1. *The solution given by the DG method defined by (6.4)-(6.5) satisfies*

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega_h} u_h^2(\mathbf{x}, t) d\mathbf{x} + \int_{\Omega_h} w_h^2(\mathbf{x}, t) d\mathbf{x} = 0. \quad (6.9)$$

Proof. We take the test functions $p = u_h$, $q = w_h$ in (6.4) and (6.5) respectively, and integrate by parts to obtain

$$((u_h)_t, u_h)_K + (w_h, w_h)_K + H_{\partial K}(u_h, w_h) = 0,$$

where

$$\begin{aligned} H_{\partial K}(p, q) = & \langle w_h, \nabla u_h \cdot \mathbf{n} \rangle_{\partial K} + \langle \widetilde{\nabla w} \cdot \mathbf{n}, p \rangle_{\partial K} - \langle \widetilde{w}, \nabla p \cdot \mathbf{n} \rangle_{\partial K} - \langle u_h, \nabla w_h \cdot \mathbf{n} \rangle_{\partial K} \\ & - \langle \widehat{\nabla u} \cdot \mathbf{n}, q \rangle_{\partial K} + \langle \widehat{u}, \nabla q \cdot \mathbf{n} \rangle_{\partial K}. \end{aligned}$$

Next we sum over the K . Since

$$H_{\partial K_1 \cap e}(u_h, w_h) + H_{\partial K_2 \cap e}(u_h, w_h) = 0, \quad (6.10)$$

with the numerical flux (6.7) or (6.8), here we suppose e is an inter-element face shared with the elements K_1 and K_2 , we can immediately get the L^2 -stability result (6.9). $\square$

6.3 Error estimates

In this subsection, we obtain a priori error estimates for the approximation (u_h, w_h) given by the DG scheme (6.4)-(6.5). The proof of optimal error estimate in the multi-dimensional case is different from that in the one-dimensional case, in the definition and analysis of suitable projections. Since the projection terms in the error equations do not vanish as in the one-dimensional case, we need to obtain certain superconvergence properties of the projections to deal with these terms.

Theorem 6.2. *Let u be the solution of the equation (6.1) with periodic boundary condition, and $w = \Delta u$. Let u_h and w_h be the numerical solution of the DG scheme (6.4)-(6.5). If we use W_h as the space with piecewise polynomials $\mathcal{Q}^k$, $k \geq 1$. Then for Cartesian meshes, we have*

$$\|u(t) - u_h(t)\| + \int_0^t \|w(t) - w_h(t)\| dt \leq Ch^{k+1}.$$

Here C depends on $\|u\|_{L^\infty((0,T);W^{2k+6,\infty})}$, $\|u_t\|_{L^\infty((0,T);W^{k+1,\infty})}$, and on t , but is independent of h .

6.4 Proof of the error estimates

In this subsection we prove Theorem 6.2 stated in the previous section. To do that, firstly, we define the special projection in Cartesian meshes, similar to the Gauss-Radau projections in Cartesian meshes [6, 18, 26].

On a rectangle $K_{i,j} = I_i \times J_j$, for $u \in W^{1,\infty}(\bar{K})$, we define

$$\Pi^\pm u := P_{1hx}^\pm \otimes P_{1hy}^\pm u, \quad (6.11)$$

with the subscripts indicating the application of the one-dimensional operators $P_{1h}^\pm$ with respect to the corresponding variable. To be more specific, we shall list explicitly the formulations for $\Pi^- u$, on a rectangular element $K_{i,j} = I_i \times J_j := (x_{i-\frac{1}{2}}, x_{i+\frac{1}{2}}) \times (y_{j-\frac{1}{2}}, y_{j+\frac{1}{2}})$. We have

$$\int_{K_{i,j}} \Pi^- u(x, y) v_h(x, y) dx dy = \int_{K_{i,j}} u(x, y) v_h(x, y) dx dy, \quad (6.12a)$$

$$\int_{I_i} \Pi^- u(x, y_{j+\frac{1}{2}}^-) v_h(x, y_{j+\frac{1}{2}}^-) dx = \int_{I_i} u(x, y_{j+\frac{1}{2}}^-) v_h(x, y_{j+\frac{1}{2}}^-) dx, \quad (6.12b)$$

$$\int_{I_i} (\Pi^- u)_y(x, y_{j+\frac{1}{2}}^-) v_h(x, y_{j+\frac{1}{2}}^-) dx = \int_{I_i} u_y(x, y_{j+\frac{1}{2}}^-) v_h(x, y_{j+\frac{1}{2}}^-) dx, \quad (6.12c)$$

$$\int_{J_j} \Pi^- u(x_{i+\frac{1}{2}}^-, y) v_h(x_{i+\frac{1}{2}}^-, y) dy = \int_{J_j} u(x_{i+\frac{1}{2}}^-, y) v_h(x_{i+\frac{1}{2}}^-, y) dy, \quad (6.12d)$$

$$\int_{J_j} (\Pi^- u)_x(x_{i+\frac{1}{2}}^-, y) v_h(x_{i+\frac{1}{2}}^-, y) dy = \int_{J_j} u_x(x_{i+\frac{1}{2}}^-, y) v_h(x_{i+\frac{1}{2}}^-, y) dy, \quad (6.12e)$$

$$\Pi^- u(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-) = u(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-), \quad (6.12f)$$

$$(\Pi^- u)_x(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-) = u_x(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-), \quad (6.12g)$$

$$(\Pi^- u)_y(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-) = u_y(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-), \quad (6.12h)$$

$$(\Pi^- u)_{xy}(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-) = u_{xy}(x_{i+\frac{1}{2}}^-, y_{j+\frac{1}{2}}^-), \quad (6.12i)$$

for all $v_h \in \mathcal{Q}^{k-2}(K)$ and $K \in \Omega_h$. Similarly, we can define the projection Π^+ . Existence and the optimal approximation property of the projection $\Pi^\pm$ are established in the following lemma.

Lemma 6.1. *Assume u is sufficiently smooth, then there exists a unique $\Pi^- u \in W_h$, satisfying (6.12). Moreover, there holds the following approximation property*

$$\|v - \Pi^\pm v\|_{L^2(K)} + h\|v - \Pi^\pm v\|_{H^1(K)} \leq Ch^{k+1}\|u\|_{H^{k+1}(K)}.$$

Proof. Assume that $u \equiv 0$, then by (6.12b), (6.12f) and (6.12g) we have

$$\Pi^- u(x, y_{j+\frac{1}{2}}^-) = 0.$$

Furthermore, by (6.12c), (6.12h) and (6.12i) we get

$$(\Pi^- u)_y(x, y_{j+\frac{1}{2}}^-) = 0.$$

Similarly, we have $\Pi^- u(x_{i+\frac{1}{2}}^-, y) = 0$, and $(\Pi^- u)_x(x_{i+\frac{1}{2}}^-, y) = 0$, then we obtain

$$\Pi^- u = (x - x_{i+\frac{1}{2}}^-)^2 (y - y_{j+\frac{1}{2}}^-)^2 Q(x, y), \quad Q(x, y) \in \mathcal{Q}^{k-2}.$$

Finally, we take $v_h = Q(x, y)$ in (6.12a) to get $Q(x, y) \equiv 0$, therefore $\Pi^- u \equiv 0$, and we have finished the proof of the uniqueness and also existence. Since the one-dimensional operators $P_{1h}^\pm$ satisfy $\|P_{1h}^\pm u\|_{L^\infty(I_j)} \leq C\|u\|_{L^\infty(I_j)}$, similarly in the two-dimensional case we also have $\|\Pi^\pm u\|_{L^\infty(K_{i,j})} \leq C\|u\|_{L^\infty(K_{i,j})}$, here C is a constant independent of h . Again, standard approximation theory implies the optimal approximating estimates. $\square$

To prove Theorem 6.2, firstly we need to write the error equations. Let

$$e_u = u - u_h = \eta_u - \xi_u, \quad e_w = w - w_h = \eta_w - \xi_w$$

with

$$\eta_u = u - \Pi^+ u, \quad \eta_w = w - \Pi^- w, \quad \xi_u = u_h - \Pi^+ u, \quad \xi_w = w_h - \Pi^- w,$$

then

$$((\xi_u)_t, p)_K + B_K^1(\xi_w, p) = ((\eta_u)_t, p)_K + B_K^1(\eta_w, p), \quad (6.13)$$

$$(\xi_w, q)_K - B_K^2(\xi_u, q) = (\eta_w, q)_K - B_K^2(\eta_u, q)_K, \quad (6.14)$$

where

$$B_K^1(w, p) = (w, \Delta p)_K - \langle w^-, (\nabla p \cdot \mathbf{n}) \rangle_{\partial K} + \langle (\nabla w^- \cdot \mathbf{n}), p \rangle_{\partial K}, \quad (6.15)$$

$$B_K^2(u, q) = (u, \Delta q)_K - \langle u^+, (\nabla q \cdot \mathbf{n}) \rangle_{\partial K} + \langle (\nabla u^+ \cdot \mathbf{n}), q \rangle_{\partial K}. \quad (6.16)$$

Besides the standard approximation results, we will also prove superconvergence results for the projections $\Pi^\pm$ in Lemma 6.2 and 6.3. The proof is using similar strategies and skills in [6].

Lemma 6.2. *Let $B_K^1(\eta_w, p)$ and $B_K^2(\eta_u, q)$ be defined by (6.15) and (6.16). Then we have for $k \geq 1$,*

$$B_K^1(\eta_w, p) = 0, \quad B_K^2(\eta_u, q) = 0, \quad \forall u, w \in \mathcal{P}^{k+2}(K), \quad p, q \in \mathcal{Q}^k(K). \quad (6.17)$$

Proof. The proof of the results for B_K^1 and B_K^2 are analogous; therefore we just prove the one for $B_K^2(\eta_u, q)$. Let us consider the rectangular element $K_{ij} = I_i \times J_j = (x_{i-\frac{1}{2}}, x_{i+\frac{1}{2}}) \times (y_{j-\frac{1}{2}}, y_{j+\frac{1}{2}})$. By the definition of $B_K^2(\eta_u, q)$ we have

$$\begin{aligned} B_K^2(\eta_u, q) &= \int_{K_{i,j}} (u - \Pi^+ u)(q_{xx} + q_{yy}) dx dy \\ &\quad - \int_{y_{j-\frac{1}{2}}}^{y_{j+\frac{1}{2}}} (u - \Pi^+ u)(x_{i+\frac{1}{2}}^+, y) q_x(x_{i+\frac{1}{2}}^-, y) - (u - \Pi^+ u)(x_{i-\frac{1}{2}}^+, y) q_x(x_{i-\frac{1}{2}}^-, y) dy \\ &\quad - \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} (u - \Pi^+ u)(x, y_{j+\frac{1}{2}}^+) q_y(x, y_{j+\frac{1}{2}}^-) - (u - \Pi^+ u)(x, y_{j-\frac{1}{2}}^+) q_y(x, y_{j-\frac{1}{2}}^-) dx \\ &\quad + \int_{y_{j-\frac{1}{2}}}^{y_{j+\frac{1}{2}}} (u - \Pi^+ u)_x(x_{i+\frac{1}{2}}^+, y) q(x_{i+\frac{1}{2}}^-, y) - (u - \Pi^+ u)_x(x_{i-\frac{1}{2}}^+, y) q(x_{i-\frac{1}{2}}^-, y) dy \\ &\quad + \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} (u - \Pi^+ u)_y(x, y_{j+\frac{1}{2}}^+) q(x, y_{j+\frac{1}{2}}^-) - (u - \Pi^+ u)_y(x, y_{j-\frac{1}{2}}^+) q(x, y_{j-\frac{1}{2}}^-) dx. \end{aligned}$$

Since Π^+ is polynomial preserving operator, (6.17) holds true for every $u \in \mathcal{Q}^k(K)$. Therefore, we have to consider the cases $u(x, y) = x^{k+1}, y^{k+1}, x^{k+2}, y^{k+2}, x^{k+1}y, y^{k+1}x$.

Let us start with $u(x, y) = x^{k+1}$. We have $(u - \Pi^+ u)_y(x, y) = 0$, by (6.12f) and (6.12g), $u(x_{i+\frac{1}{2}}^+, y) = \Pi^+ u(x_{i+\frac{1}{2}}^+, y)$, $u_x(x_{i+\frac{1}{2}}^+, y) = (\Pi^+ u)_x(x_{i+\frac{1}{2}}^+, y)$. Then

$$\begin{aligned} \int_{y_{j-\frac{1}{2}}}^{y_{j+\frac{1}{2}}} (u - \Pi^+ u)(x_{i+\frac{1}{2}}^+, y) q_x(x_{i+\frac{1}{2}}^-, y) - (u - \Pi^+ u)(x_{i-\frac{1}{2}}^+, y) q_x(x_{i-\frac{1}{2}}^-, y) dy &= 0, \\ \int_{y_{j-\frac{1}{2}}}^{y_{j+\frac{1}{2}}} (u - \Pi^+ u)_x(x_{i+\frac{1}{2}}^+, y) q(x_{i+\frac{1}{2}}^-, y) - (u - \Pi^+ u)_x(x_{i-\frac{1}{2}}^+, y) q(x_{i-\frac{1}{2}}^-, y) dy &= 0, \end{aligned}$$

and $\int_{K_{ij}} (u - \Pi^+ u) q_{xx} dx dy = 0$. Next we integrate by parts

$$\begin{aligned} &\int_{K_{i,j}} (u - \Pi^+ u) q_{yy} dx dy \\ &= \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} (u - \Pi^+ u)(x, y_{j+\frac{1}{2}}^-) q_y(x, y_{j+\frac{1}{2}}^-) - (u - \Pi^+ u)(x, y_{j-\frac{1}{2}}^-) q_y(x, y_{j-\frac{1}{2}}^-) dx. \end{aligned}$$

Therefore, sum all the parts in the definition of $B_K^2(\eta_u, q)$, we have

$$B_K^2(\eta_u, q) = 0.$$

Next, we consider the case $u(x, y) = x^{k+1}y$, in this case $\Pi^+ u = P_{1hx}^+(x^{k+1})y$, and

$$\int_{K_{ij}} (u - \Pi^+ u) q_{xx} dx dy = \int_{K_{ij}} y(x^{k+1} - P_{1hx}^+(x^{k+1})) q_{xx} dx dy = 0,$$

and

$$\begin{aligned}
& \int_{K_{i,j}} (u - \Pi^+ u) q_{yy} dx dy \\
&= \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} y_{j+\frac{1}{2}} (x^{k+1} - P_{1hx}^+(x^{k+1})) q_y(x, y_{j+\frac{1}{2}}^-) - y_{j-\frac{1}{2}}^+ (x^{k+1} - P_{1hx}^+(x^{k+1})) q_y(x, y_{j-\frac{1}{2}}^+) dx \\
&\quad - \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} (x^{k+1} - P_{1hx}^+(x^{k+1})) q(x, y_{j+\frac{1}{2}}^-) - (x^{k+1} - P_{1hx}^+(x^{k+1})) q(x, y_{j-\frac{1}{2}}^+) dx.
\end{aligned}$$

Then summing all the parts in the definition of $B_K^2(\eta_u, q)$, we have

$$B_K^2(\eta_u, q) = 0.$$

The proof of the cases $u(x, y) = y^{k+1}, x^{k+2}, y^{k+2}$ and $u(x, y) = y^{k+1}x$ are analogous. This completes the proof of (6.17). $\square$

Lemma 6.3. *Let $B_K^1(\eta_w, p)$ and $B_K^2(\eta_u, q)$ defined by (6.15) and (6.16). Then we have*

$$|B_K^1(\eta_w, p)| \leq Ch^{k+2} \|w\|_{W^{2k+4,\infty}(\Omega_h)} \|p\|_{L^2(K)}, \quad (6.18)$$

$$|B_K^2(\eta_u, q)| \leq Ch^{k+2} \|u\|_{W^{2k+4,\infty}(\Omega_h)} \|q\|_{L^2(K)}, \quad (6.19)$$

where $p, q \in \mathcal{Q}^k(K)$ and the constant C is independent of h .

Proof. On each element $K = I_i \times J_j$, consider the Taylor expansion of u around (x_i, y_j)

$$u = Tu + R_{k+3},$$

where

$$\begin{aligned}
Tu &= \sum_{l=0}^{k+2} \sum_{m=0}^l \frac{1}{(l-m)!m!} \frac{\partial^l u(x_i, y_j)}{\partial x^{l-m} \partial y^m} (x - x_i)^{l-m} (y - y_j)^m, \\
R_{k+3} &= (k+3) \sum_{m=0}^{k+3} \frac{(x - x_i)^{k+3-m} (y - y_j)^m}{(k+3-m)!m!} \int_0^1 (1-s)^{k+2} \frac{\partial^{k+3} u(x_i^s, y_j^s)}{\partial x^{k+3-m} \partial y^m} ds
\end{aligned}$$

with $x_i^s = x_i + s(x - x_i)$, $y_j^s = y_j + s(y - y_j)$. Clearly, $Tu \in \mathcal{P}^{k+2}$ and by Lemma 6.2 we have

$$B_K^2(Tu - \Pi^+(Tu), q) = 0,$$

then we have

$$B_K^2(\eta_u, q) = T_1 + T_2 + T_3 + T_4 + T_5,$$

where

$$\begin{aligned}
T_1 &= \int_{K_{ij}} (R_{k+3} - \Pi^+ R_{k+3})(p_{xx} + p_{yy}) dx dy, \\
T_2 &= - \int_{y_{j-\frac{1}{2}}}^{y_{j+\frac{1}{2}}} (R_{k+3} - \Pi^+ R_{k+3})(x_{i+\frac{1}{2}}^+, y) p_x(x_{i+\frac{1}{2}}^-, y) - (R_{k+3} - \Pi^+ R_{k+3})(x_{i-\frac{1}{2}}^+, y) p_x(x_{i-\frac{1}{2}}^-, y) dy, \\
T_3 &= - \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} (R_{k+3} - \Pi^+ R_{k+3})(x, y_{j+\frac{1}{2}}^+) p_y(x, y_{j+\frac{1}{2}}^-) - (R_{k+3} - \Pi^+ R_{k+3})(x, y_{j-\frac{1}{2}}^+) p_y(x, y_{j-\frac{1}{2}}^-) dx, \\
T_4 &= \int_{y_{j-\frac{1}{2}}}^{y_{j+\frac{1}{2}}} (R_{k+3} - \Pi^+ R_{k+3})_x(x_{i+\frac{1}{2}}^+, y) p(x_{i+\frac{1}{2}}^-, y) - (R_{k+3} - \Pi^+ R_{k+3})_x(x_{i-\frac{1}{2}}^+, y) p(x_{i-\frac{1}{2}}^-, y) dy, \\
T_5 &= \int_{x_{i-\frac{1}{2}}}^{x_{i+\frac{1}{2}}} (R_{k+3} - \Pi^+ R_{k+3})_y(x, y_{j+\frac{1}{2}}^+) p(x, y_{j+\frac{1}{2}}^-) - (R_{k+3} - \Pi^+ R_{k+3})_y(x, y_{j-\frac{1}{2}}^+) p(x, y_{j-\frac{1}{2}}^-) dx.
\end{aligned}$$

which will be estimated one by one below. From the approximation properties of the projection Π^+ , we have

$$\|R_{k+3} - \Pi^+ R_{k+3}\|_{L^2(K)} \leq Ch^{k+2} \|R_{k+3}\|_{W^{k+1,\infty}(\Omega_h)},$$

and

$$\|R_{k+3}\|_{W^{k+1,\infty}(\Omega_h)} = \max_K \|R_{k+3}\|_{W^{k+1,\infty}(K)} \leq Ch^2 \|u\|_{W^{2k+4,\infty}(\Omega_h)}.$$

Combining the above two estimates, we arrive at

$$\|R_{k+3} - \Pi^+ R_{k+3}\|_{L^2(K)} \leq Ch^{k+4} \|u\|_{W^{2k+4,\infty}(\Omega_h)}. \quad (6.20)$$

Similarly, we have that

$$\|R_{k+3} - \Pi^+ R_{k+3}\|_{H^1(K)} \leq Ch^{k+3} \|u\|_{W^{2k+4,\infty}(\Omega_h)}. \quad (6.21)$$

It follows from the Cauchy-Schwartz inequality, and the inverse inequality that

$$|T_1| \leq \|R_{k+3} - \Pi^+ R_{k+3}\|_{L^2(K)} \|q_{xx}\|_{L^2(K)} \leq Ch^{k+2} \|u\|_{W^{2k+4,\infty}(\Omega_h)} \|q\|_{L^2(K)}.$$

In order to estimate the remaining terms we need to use the trace inequality to get

$$\|R_{k+3} - \Pi^+ R_{k+3}\|_{L^2(\partial K)} \leq Ch^{k+\frac{7}{2}} \|u\|_{W^{2k+4,\infty}(\Omega_h)}$$

and

$$\|R_{k+3} - \Pi^+ R_{k+3}\|_{H^1(\partial K)} \leq Ch^{k+\frac{5}{2}} \|u\|_{W^{2k+4,\infty}(\Omega_h)}$$

Next, by the Cauchy-Schwartz inequality and the inverse inequality, we arrive at

$$|T_2| \leq \|R_{k+3} - \Pi^+ R_{k+3}\|_{L^2(\partial K)} \|q_x\|_{L^2(\partial K)} \leq Ch^{k+2} \|u\|_{W^{2k+4,\infty}(\Omega_h)} \|q\|_{L^2(K)}.$$

Analogously, we have that

$$|T_m| \leq Ch^{k+2} \|u\|_{W^{2k+4,\infty}(\Omega_h)} \|q\|_{L^2(K)}, \quad m = 3, 4, 5.$$

The estimates for $B^1(\eta_u, q)$ now follows by collecting the results for T_m , $m = 1, 2, 3, 4, 5$ obtained above. The proof of Lemma is thus completed. $\square$

Next, we will use these lemmas to prove our final result, Theorem 6.2.

Proof. (The proof of Theorem 6.2). We take $p = \xi_u$ and $q = \xi_w$ in the error equations (6.13)-(6.14), to obtain

$$((\xi_u)_t, \xi_u)_{\Omega_h} + (\xi_w, \xi_w)_{\Omega_h} = ((\eta_u)_t, \xi_u)_{\Omega_h} + (\eta_w, \xi_w)_{\Omega_h} + \sum_K (B_K^1(\eta_w, \xi_u) - B_K^2(\eta_u, \xi_w)).$$

Then by the Cauchy-Schwartz inequality and Lemma 6.3, we have

$$\frac{1}{2} \frac{d}{dt} \|\xi_u\|^2 + \|\xi_w\|^2 \leq Ch^{k+1} \|\xi_u\|^2 + Ch^{k+1} \|\xi_w\|^2.$$

Next, by Gronwall's inequality and choosing $u_h(0) = \Pi_h^+ u(0)$, we have

$$\|\xi_u\|(t) + \int_0^t \|\xi_w\|(t) dt \leq Ch^{k+1},$$

and

$$\|e_u\|(t) + \int_0^t \|e_w\| dt \leq \|\xi_u\|(t) + \int_0^t \|\xi_w\| dt + \|\eta_u\|(t) + \int_0^t \|\eta_w\| dt \leq Ch^{k+1},$$

where C is a constant independent of h and dependent on $\|u\|_{W^{2k+6,\infty}}$, $\|u_t\|_{W^{k+1,\infty}}$ and t . $\square$

7 Numerical results

In this section, we present numerical examples to verify our theoretical convergence properties of the DG method for high order PDEs.

Firstly, we consider the one-dimensional linear fourth and fifth order time-dependent equations with the periodic boundary condition in Examples 7.1 and 7.2, respectively. Time discretization is not our major concern in this paper, hence we use the spectral deferred correction (SDC) [24] time discretization for its simplicity. Our computation is based on the flux choice (3.9) and (4.9), respectively. The errors and numerical orders of accuracy for P^k elements with $1 \leq k \leq 3$ are listed in Table 7.1 and Table 7.2. We observe that our scheme gives the optimal $(k+1)$ -th order of the accuracy when $k \geq 1$.

Example 7.1. (*Accuracy test for a linear fourth-order problem.*) We consider the following fourth-order time-dependent problem

$$\begin{aligned} u_t + u_{xxxx} &= 0, & (x, t) &\in [0, 2\pi] \times (0, 1], \\ u(x, 0) &= \sin(x). \end{aligned}$$

The exact solution is

$$u(x, t) = e^{-t} \sin(x).$$

Table 7.1: Errors and the corresponding convergence rates for Example 7.1 when using $\mathcal{P}^k$ polynomials and SDC time discretization on a uniform mesh of N cells. Final time $t = 1$.

	N	L^1	order	L^2	order	L^∞	order
$\mathcal{P}^1$	10	2.97E-02	—	3.61E-02	—	9.45E-02	—
	20	7.66E-03	1.96	9.31E-03	1.96	2.39E-02	1.98
	40	1.93E-03	1.99	2.35E-03	1.99	6.04E-03	1.99
	80	4.83E-04	2.00	5.88E-04	2.00	1.51E-03	2.00
	160	1.21E-04	2.00	1.47E-04	2.00	3.79E-04	2.00
	320	3.02E-05	2.00	3.68E-05	2.00	9.46E-05	2.00
$\mathcal{P}^2$	10	2.63E-02	—	2.92E-02	—	4.19E-02	—
	20	3.57E-03	2.88	3.97E-03	2.88	5.70E-03	2.88
	40	4.54E-04	2.98	5.04E-04	2.98	7.18E-04	2.99
	80	5.68E-05	3.00	6.31E-05	3.00	8.98E-05	3.00
	160	7.10E-06	3.00	7.88E-06	3.00	1.12E-05	3.00
	320	8.87E-07	3.00	9.85E-07	3.00	1.40E-06	3.00
$\mathcal{P}^3$	10	1.54E-03	—	1.71E-03	—	2.44E-03	—
	20	1.40E-04	3.46	1.55E-04	3.46	2.22E-04	3.46
	40	9.35E-06	3.90	1.04E-05	3.90	1.49E-05	3.90
	80	5.99E-07	3.96	6.66E-07	3.96	9.54E-07	3.96
	160	3.76E-08	3.99	4.18E-08	3.99	5.99E-08	3.99
	320	2.36E-09	4.00	2.62E-09	4.00	3.75E-09	4.00

Example 7.2. (*Accuracy test for a linear fifth-order problem.*) We consider the following linear fifth-order time-dependent problem.

$$u_t + u_{xxxxx} = 0, \quad (x, t) \in [0, 2\pi] \times (0, 1],$$

$$u(x, 0) = \sin(x).$$

The exact solution is

$$u(x, t) = \sin(x - t).$$

Table 7.2: Errors and the corresponding convergence rates for Example 7.2 when using $\mathcal{P}^k$ polynomials and SDC time discretization on a uniform mesh of N cells. Final time $t = 1$.

	N	L^1	order	L^2	order	L^∞	order
$\mathcal{P}^1$	10	8.13E-02	—	9.08E-02	—	1.44E-01	—
	20	2.22E-02	1.87	2.47E-02	1.88	3.97E-02	1.86
	40	5.68E-03	1.97	6.32E-03	1.97	1.08E-02	1.88
	80	1.43E-03	1.99	1.59E-03	1.99	2.81E-03	1.94
	160	3.57E-04	2.00	3.98E-04	2.00	7.15E-04	1.98
	320	8.92E-05	2.00	9.95E-05	2.00	1.80E-04	1.99
$\mathcal{P}^2$	10	7.25E-02	—	8.07E-02	—	1.14E-01	—
	20	9.74E-03	2.90	1.08E-02	2.90	1.53E-02	2.90
	40	1.23E-03	2.98	1.37E-03	2.98	1.94E-03	2.98
	80	1.54E-04	3.00	1.71E-04	3.00	2.42E-04	3.00
	160	1.93E-05	3.00	2.14E-05	3.00	3.03E-05	3.00
	320	2.41E-06	3.00	2.68E-06	3.00	3.79E-06	3.00
$\mathcal{P}^3$	10	5.44E-03	—	6.04E-03	—	8.56E-03	—
	20	4.13E-04	3.72	4.59E-04	3.72	6.49E-04	3.72
	40	2.60E-05	3.99	2.89E-05	3.99	4.08E-05	3.99
	80	1.64E-06	3.99	1.82E-06	3.99	2.58E-06	3.99
	160	1.02E-07	4.00	1.14E-07	4.00	1.61E-07	4.00
	320	6.41E-09	4.00	7.12E-09	4.00	1.01E-08	4.00

Example 7.3. (*Accuracy test for a nonlinear fourth-order problem.*) We consider the following nonlinear fourth-order time-dependent problem.

$$u_t + (u^2 u_{xx})_{xx} = f, \quad x \in [0, 2\pi].$$

The source term f is chosen so that the exact solution is

$$u(x, t) = e^{-t} \sin(x).$$

We test this example by the DG scheme (3.6)-(3.8). Both errors and orders of accuracy are listed in Table 7.3. We again observe that our scheme gives the optimal $(k + 1)$ -th order of the accuracy for this nonlinear problem.

Table 7.3: Errors and the corresponding convergence rates for Example 7.3 when using $\mathcal{P}^k$ polynomials on a uniform mesh of N cells. Final time $t = 0.1$.

	N	L^1	order	L^2	order	L^∞	order
$\mathcal{P}^1$	4	1.47E-01	—	1.93E-01	—	3.97E-01	—
	8	6.74E-02	1.12	8.10E-02	1.25	2.28E-01	0.80
	16	1.94E-02	1.80	2.58E-02	1.65	8.21E-02	1.47
	32	5.05E-03	1.94	6.36E-03	2.02	2.45E-02	1.75
	64	1.19E-03	2.08	1.41E-03	2.17	4.33E-03	2.50
$\mathcal{P}^2$	4	4.85E-02	—	6.72E-02	—	2.63E-01	—
	8	2.63E-03	4.21	3.77E-03	4.16	1.37E-02	4.26
	16	8.22E-04	1.68	1.38E-03	1.45	5.87E-03	1.23
	32	1.19E-04	2.79	2.12E-04	2.71	1.00E-03	2.55
	64	1.55E-05	2.94	2.68E-05	2.99	1.58E-04	2.67
$\mathcal{P}^3$	4	4.86E-03	—	5.91E-03	—	1.81E-02	—
	8	1.07E-03	2.19	1.75E-03	1.75	8.99E-03	1.01
	16	3.54E-05	4.92	6.61E-05	4.73	4.42E-04	4.35
	32	1.16E-06	4.93	2.04E-06	5.02	1.68E-05	4.71
	64	4.65E-08	4.64	6.99E-08	4.87	5.99E-07	4.81

Example 7.4. (*Accuracy test for a nonlinear fifth-order problem.*) We consider the following nonlinear fifth-order time-dependent problem

$$u_t + (u_{xx})_{xxx}^3 = f, \quad x \in [0, 2\pi],$$

where the source term f is chosen such that the exact solution is

$$u(x, t) = \sin(x - t).$$

We test this example by the DG scheme (4.6)-(4.8). Both the errors and the numerical orders of accuracy are listed in Table 7.4. We once again observe the designed $(k + 1)$ -th order of accuracy for this nonlinear problem.

The last example we consider is a two-dimensional fourth-order problem.

Table 7.4: Errors and the corresponding convergence rates for Example 7.4 when using $\mathcal{P}^k$ polynomials on a uniform mesh of N cells. Final time $t = 0.1$.

	N	L^1	order	L^2	order	L^∞	order
$\mathcal{P}^1$	4	2.06E-01	–	2.33E-01	–	5.05E-01	–
	8	5.44E-02	1.92	6.94E-02	1.75	2.09E-01	1.28
	16	1.64E-02	1.73	2.01E-02	1.79	6.13E-02	1.77
	32	3.67E-03	2.16	4.47E-03	2.16	1.42E-02	2.11
	64	1.19E-03	1.62	1.44E-03	1.63	4.17E-03	1.77
$\mathcal{P}^2$	4	3.06E-02	–	4.39E-02	–	1.72E-01	–
	8	4.14E-03	2.88	6.34E-03	2.79	2.80E-02	2.62
	16	4.01E-04	3.37	5.56E-04	3.51	2.44E-03	3.52
	32	4.73E-05	3.08	6.78E-05	3.04	3.29E-04	2.89
	64	5.57E-06	3.09	8.34E-06	3.02	4.07E-05	3.02
$\mathcal{P}^3$	4	4.91E-03	–	6.45E-03	–	2.00E-02	–
	8	1.42E-04	5.12	1.96E-04	5.04	1.03E-03	4.28
	16	8.95E-06	3.98	1.25E-05	3.98	6.73E-05	3.93
	32	5.06E-07	4.15	7.38E-07	4.08	4.21E-06	4.00

Example 7.5. (*Accuracy test for a two-dimensional linear fourth-order problem.*) We consider the following fourth-order time-dependent problem with the periodic boundary condition

$$\begin{aligned} u_t + \Delta^2 u &= 0, & (x, y) &\in [0, 2\pi] \times [0, 2\pi], \\ u(x, 0) &= \sin(x + y). \end{aligned}$$

The exact solution is

$$u(x, t) = e^{-4t} \sin(x + y).$$

Our computation is based on the flux choice (6.7). The errors and numerical orders of accuracy for the $\mathcal{Q}^k$ elements with $1 \leq k \leq 3$ are listed in Table 7.5. We observe that our scheme gives the optimal $(k + 1)$ -th order of the accuracy when $k \geq 1$.

8 Concluding remarks

In this paper, we have constructed a new class of discontinuous Galerkin methods combining the LDG and UWDG methods for solving high order PDEs, namely time-

Table 7.5: Errors and the corresponding convergence rates for Example 7.5 when using $\mathcal{Q}^k$ polynomials on a uniform mesh of $N \times N$ cells. Final time $t = 1$.

	$N \times N$	L^1	order	L^2	order	L^∞	order
$\mathcal{Q}^1$	4×4	1.67E-01	–	2.46E-01	–	1.13E+00	–
	8×8	5.29E-02	1.66	7.93E-02	1.63	4.04E-01	1.49
	16×16	1.25E-02	2.08	2.03E-02	1.97	1.07E-01	1.92
	32×32	3.02E-03	2.05	5.09E-03	2.00	2.70E-02	1.98
	64×64	7.46E-04	2.02	1.27E-03	2.00	6.78E-03	2.00
$\mathcal{Q}^2$	2×2	3.41E-01	–	5.14E-01	–	2.55E+00	–
	4×4	4.49E-02	2.92	7.29E-02	2.82	5.20E-01	2.29
	8×8	5.41E-03	3.05	9.03E-03	3.01	6.73E-02	2.95
	16×16	6.70E-04	3.01	1.12E-03	3.01	8.45E-03	2.99
	32×32	8.35E-05	3.00	1.40E-04	3.00	1.06E-03	3.00
	64×64	1.04E-05	3.00	1.75E-05	3.00	1.32E-04	3.00

dependent PDEs with high order spatial derivatives. The idea is to rewrite the PDE into a lower order system, but not to a system with only first order spatial derivatives as in LDG methods. The ideas in designing numerical fluxes to obtain stable and accurate DG schemes from both the LDG schemes and the UWDG schemes, including the usage of alternating and upwinding numerical fluxes when appropriate, are then used to obtain stable and optimally convergent DG schemes for a wide variety of linear and nonlinear PDEs with high order spatial derivatives in both one and two spatial dimensions. The main advantage of our method over the LDG method is that we have introduced fewer auxiliary variables, thereby reducing memory and computational costs. The main advantage of our method over the UWDG method is that no internal penalty terms are necessary in order to ensure stability for both even and odd order PDEs. Detailed algorithm formulation, stability analysis and optimal L^2 error estimates are given for several examples, including fourth order linear and nonlinear equations in one dimension and a fourth order linear equation in two dimension, and fifth order linear and nonlinear wave equations in one dimension. In our error estimates, a key ingredient is the study of the relationship between the derivative and the element interface jumps of the numerical solution and the auxiliary variable numerical solution of the derivative. With this relationship and by using the discrete Sobolev and Poincaré inequalities, we can obtain optimal error estimates for both even order diffusive PDEs and odd order wave PDEs.

Numerical examples are provided both for linear and nonlinear equations and both in one dimension and in two dimensions, to verify the theoretical results. Extension of the optimal error estimates to the nonlinear equations is highly nontrivial and is left for future work.

References

- [1] S.C. Brenner. Discrete Sobolev and Poincaré inequalities for piecewise polynomial functions. *Electronic Transactions on Numerical Analysis*, v18 (2004), pp.42-48.
- [2] S.C. Brenner. Poincaré–Friedrichs inequalities for piecewise H^1 functions. *SIAM Journal on Numerical Analysis*, v41 (2003), pp.309-324.
- [3] Y. Cheng and C.-W. Shu. A discontinuous Galerkin finite element method for time dependent partial differential equations with higher order derivatives. *Mathematics of Computation*, v77 (2009), pp.699-730.
- [4] P. G. Ciarlet. The Finite Element Method for Elliptic Problems. *Classics in Applied Mathematics, vol. 40, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2002.*
- [5] B. Cockburn, S. Hou, and C.-W. Shu. The Runge-Kutta local projection discontinuous Galerkin finite element method for conservation laws. IV. The multidimensional case. *Mathematics of Computation*, v54 (1990), pp.545-581.
- [6] B. Cockburn, G. Kanschat, L. Perugia and D. Schötzau. Superconvergence of the local discontinuous Galerkin method for elliptic problems on Cartesian grids. *SIAM Journal on Numerical Analysis*, v 39(2001), pp.264-285.
- [7] B. Cockburn and C.-W. Shu. TVB Runge-Kutta local projection discontinuous Galerkin finite element method for conservation laws. II. General framework. *Mathematics of Computation*, v52 (1989), pp.411-435.
- [8] B. Cockburn and C.-W. Shu. The Runge-Kutta discontinuous Galerkin method for conservation laws. V. Multidimensional systems. *Journal of Computational Physics*, v141 (1998), pp.199-224.
- [9] B. Cockburn and C.-W. Shu, The local discontinuous Galerkin method for time-dependent convection-diffusion systems. *SIAM Journal on Numerical Analysis*, v35 (1998), pp.2440-2463.

- [10] B. Dong and C.-W. Shu, Analysis of a local discontinuous Galerkin method for linear time-dependent fourth-order problems. *SIAM Journal on Numerical Analysis*, v47 (2009), pp.3240-3268.
- [11] J. Douglas and T. Dupont. Interior penalty procedures for elliptic and parabolic Galerkin methods. *Computing Methods in Applied Sciences*, pp.207-216. Springer, Berlin, Heidelberg.
- [12] P. Fu, Y. Cheng, F. Li and Y. Xu. Discontinuous Galerkin methods with optimal L^2 accuracy for one dimensional linear PDEs with high order spatial derivatives. *Journal of Scientific Computing*, v78 (2019), pp.816-863.
- [13] L. Ji and Y. Xu. Optimal error estimates of the local discontinuous Galerkin method for Willmore flow of graphs on Cartesian meshes. *International Journal of Numerical Analysis & Modeling*, v8 (2011), pp.252-283.
- [14] S.M. Han, H Benaroya and T. Wei. Dynamics of transversely vibrating beams using four engineering theories. *Journal of Sound and Vibration*, v225 (1999), pp.935-988.
- [15] J.H. Hunter and J.M. Vanden-Broeck. Solitary and periodic gravity capillary waves of finite amplitude. *Journal of Fluid Mechanics*, v134 (1983), pp.205-219.
- [16] H.-L. Liu and P. Yin. A Mixed discontinuous Galerkin method without interior penalty for time-dependent fourth order problems. *Journal of Scientific Computing*, v77 (2018), pp.467-501.
- [17] H.-L. Liu and J. Yan. A local discontinuous Galerkin method for the Korteweg de Vries equation with boundary effect. *Journal of Computational Physics*, v215 (2006), pp.197-218.
- [18] X. Meng, C.-W. Shu and B. Wu. Optimal error estimates for discontinuous Galerkin methods based on upwind-biased fluxes for linear hyperbolic equations. *Mathematics of Computation*, v85 (2016), pp.1225-1261.
- [19] I. Mozolevski, E. Süli, and P.R. Bösing. hp-version a priori error analysis of interior penalty discontinuous Galerkin finite element approximations to the biharmonic equation. *Journal of Scientific Computing*, v30 (2007), pp.465-491.
- [20] W. Reed and T. Hill. Triangular mesh methods for the neutron transport equation. La-ur-73-479, Los Alamos Scientific Laboratory, 1973.

- [21] C.-W. Shu, Discontinuous Galerkin method for time dependent problems: Survey and recent developments, *Recent Developments in Discontinuous Galerkin Finite Element Methods for Partial Differential Equations (2012 John H. Barrett Memorial Lectures)*, X. Feng, O. Karakashian and Y. Xing, editors. *The IMA Volumes in Mathematics and Its Applications*, volume 157, Springer, Switzerland, 2014, pp.25-62.
- [22] H. Wang, C.-W. Shu and Q. Zhang. Stability and error estimates of local discontinuous Galerkin methods with implicit-explicit time-marching for advection-diffusion problems. *SIAM Journal on Numerical Analysis*, v53 (2015), pp.209-227.
- [23] H. Wang, S. Wang, C.-W. Shu and Q. Zhang. Local discontinuous Galerkin methods with implicit-explicit time-marching for multi-dimensional convection-diffusion problems. *ESAIM: Mathematical Modelling and Numerical Analysis (M²AN)*, v50 (2016), pp.1083-1105.
- [24] Y. Xia, Y. Xu and C.-W. Shu. Efficient time discretization for local discontinuous Galerkin methods. *Discrete and Continuous Dynamical Systems - Series B*, v8 (2007), pp.677-693.
- [25] Y. Xu and C.-W. Shu. Local discontinuous Galerkin methods for high-order time-dependent partial differential equations. *Communications in Computational Physics*, v7 (2010), pp.1-46.
- [26] Y. Xu and C.-W. Shu. Optimal error estimates of the semi-discrete local discontinuous Galerkin methods for high order wave equations. *SIAM Journal on Numerical Analysis*, v50 (2012), pp.79-104.
- [27] J. Yan and C.-W. Shu. A local discontinuous Galerkin method for KdV type equations. *SIAM Journal on Numerical Analysis*, v40 (2002), pp.769-791.
- [28] J. Yan and C.-W. Shu. Local discontinuous Galerkin methods for partial differential equations with higher order derivatives. *Journal of Scientific Computing*, v17 (2002), pp.27-47.