

Integrative . Parall SIN A Journal of the Society for Integrative and Comparative Biology

academic.oup.com/icb







RESEARCH ARTICLE

Post-Transcriptional Mechanisms Respond Rapidly to Ecologically Relevant Thermal Fluctuations During Temperature-Dependent Sex Determination

Samantha L. Bock $igodots,^{1,*,\dagger}$ Matthew D. Hale, ‡ Faith M. Leri, § Philip M. Wilkinson, ¶ Thomas R. Rainwater $^{\P,\parallel}$ and Benjamin B. Parrott *,†

*Odum School of Ecology, University of Georgia, Athens, GA 30602, USA; †Savannah River Ecology Laboratory, Aiken, SC 29802, USA; ‡Department of Biology, University of Virginia, Charlottesville, VA 22904, USA; †Department of Biology, University of Oklahoma, Norman, OK 73019, USA; Tom Yawkey Wildlife Center, Georgetown, SC 29440, USA; Belle W. Baruch Institute of Coastal Ecology & Forest Science, Clemson University, Georgetown, SC 29442, USA

¹E-mail: samantha.bock@uga.edu

Synopsis An organism's ability to integrate transient environmental cues experienced during development into molecular and physiological responses forms the basis for adaptive shifts in phenotypic trajectories. During temperaturedependent sex determination (TSD), thermal cues during discrete periods in development coordinate molecular changes that ultimately dictate sexual fate and contribute to patterns of inter- and intra-sexual variation. How these mechanisms interface with dynamic thermal environments in nature remain largely unknown. By deploying thermal loggers in wild nests of the American alligator (Alligator mississippiensis) over two consecutive breeding seasons, we observed that 80% of nests exhibit both male- and female-promoting thermal cues during the thermosensitive period, and of these nests, all exhibited both male- and female-promoting temperatures within the span of a single day. These observations raise a critical question—how are opposing environmental cues integrated into sexually dimorphic transcriptional programs across short temporal scales? To address this question, alligator embryos were exposed to fluctuating temperatures based on nest thermal profiles and sampled over the course of a daily thermal fluctuation. We examined the expression dynamics of upstream genes in the temperature-sensing pathway and find that post-transcriptional alternative splicing and transcript abundance of epigenetic modifier genes JARID2 and KDM6B respond rapidly to thermal fluctuations while transcriptional changes of downstream effector genes, SOX9 and DMRT1, occur on a delayed timescale. Our findings reveal how the basic mechanisms of TSD operate in an ecologically relevant context. We present a hypothetical hierarchical model based on our findings as well as previous studies, in which temperature-sensitive alternative splicing incrementally influences the epigenetic landscape to affect the transcriptional activity of key sex-determining genes.

Introduction

Embryos often experience dynamic environmental conditions during development which can exert lasting influences on organismal phenotypes. This is especially true for species with temperature-dependent sex determination (TSD) in which the sex of offspring is determined by temperature cues experienced during discrete periods in development. Occurring in a diverse range of taxa, including all crocodilians, many turtles, and some fish (Bull 1980; Lang and Andrews 1994; Ospina-Alvarez and Piferrer

2008), TSD provides a unique window through which to examine how transient thermal cues coordinate molecular changes that ultimately establish patterns of inter- and intra-sexual variation. Major advances have recently been made in understanding the proximate developmental mechanisms underlying TSD (Czerwinski et al. 2016; Yatsu et al. 2016; Deveson et al. 2017; Ge et al. 2017, 2018), yet little is known regarding how these mechanisms interface with the dynamic thermal environments experienced by embryos in nature.

For many species with TSD, nest temperatures fluctuate considerably on daily, monthly, and seatimescales (Warner and Shine Escobedo-Galván et al. 2016; Carter et al. 2017; 2018). While the consequences of these temperature fluctuations for offspring phenotype are incompletely understood, in some species including the red eared slider turtle, Trachemys scripta, temperature fluctuations appear to play critical roles in influencing population sex ratios. Mean temperatures of T. scripta nests in the field rarely exceed those necessary to produce females under constant laboratory conditions, yet mark-recapture studies indicate females are frequently recruited into these populations. This apparent discordance is reconciled by the observation that female development can be triggered in T. scripta by as few as 5 days of exposure to female-promoting temperatures (FPTs; Carter et al. 2018). How do embryos integrate brief periods of FPT cues experienced in the field during "heatwaves" into lasting physiological responses that drive sexual fate commitment? And, how do these mechanisms function in other TSD species? These questions remain unanswered and hold important implications for our understanding of TSD as it occurs in ecologically relevant contexts.

Epigenetic mechanisms, including DNA methylation and histone modifications, appear to play upstream roles in integrating transient temperature cues into persistent transcriptional responses during TSD (Navarro-Martín et al. 2011; Matsumoto et al. 2013; Parrott et al. 2014; Ge et al. 2018). Two genes with roles in chromatin regulation, JARID2 and KDM6B (or JMJD3), are among the earliest genes displaying temperature-sensitive differential expression during TSD in both T. scripta and Alligator mississippiensis (Czerwinski et al. 2016; Yatsu et al. 2016). JARID2 is a component of the master Polycomb repressive complex 2 (PRC2) which plays a key role in localizing the PRC2 complex to its target loci for subsequent transcriptional silencing via histone 3 lysine 27 trimethylation (H3K27me3) (Peng et al. 2009; Landeira and Fisher 2011; da Rocha et al. 2014; Kaneko et al. 2014; Sanulli et al. 2015). KDM6B encodes a histone demethylase which removes the repressive H3K27me3 mark to activate transcription of target loci (Agger et al. 2007; Lan et al. 2007). Intriguingly, interruption of KDM6B activity via RNA interference in T. scripta results in the inhibition of male-development at a malepromoting temperature (MPT) and is associated with increased H3K27me3 at the promoter of DMRT1, a transcription factor involved in male development (Ge et al. 2018). Together, these findings

suggest that epigenetic modifications and the conserved genes that regulate them play causal roles in integrating transient temperature cues experienced during development into sexually dimorphic transcriptional programs during TSD.

Interestingly, the role of these upstream epigenetic regulators in mediating TSD is likely more nuanced than previously realized. While much of the focus of TSD research has centered on investigating transcriptional responses to temperature, mounting evidence suggests post-transcriptional mechanisms, particularly alternative splicing, may contribute an added layer of regulatory complexity (Harry et al. 1990; Anand et al. 2008; Agrawal et al. 2009; Rhen et al. 2015; Deveson et al. 2017). Alternative splicing, the process by which pre-mRNA generates variable mature mRNA transcripts via differential exon usage and intron retention, plays a widespread role in contributing to transcriptional diversity across metazoan taxa (Tapial et al. 2017) and serves as the decisive sex-determining signal in some species (Salz 2011). Notably, multiple species possessing different forms of TSD including A. mississippiensis, T. scripta, and Pogona vitticeps, exhibit temperature dependent alternative splicing of the epigenetic modifier genes, JARID2 and KDM6B (Deveson et al. 2017). In particular, a unique intron containing premature stop codons is differentially retained depending on incubation temperature in mature transcripts of each of these genes (Deveson et al. 2017). How environmental variability experienced in the field interfaces with these molecular pathways is currently unknown.

Here, we quantified the level of thermal variation experienced by alligator embryos in wild nests and implemented experimental thermal fluctuations based on empirically derived nest thermal profiles examine how transcription and to transcriptional alternative splicing of genes involved in sex determination and sexual differentiation change over the course of a daily thermal fluctuation during TSD. We found that alternative splicing of the epigenetic modifier JARID2, in both the gonad and brain, responded rapidly to thermal fluctuations. Further, transcript abundance of both JARID2 and KDM6B also fluctuated with temperature in the gonad, albeit with an apparent delay of \sim 7 h. In contrast to these epigenetic regulators, genes with downstream roles in sex determination and differentiation, SOX9 and DMRT1, did not exhibit clear responses to thermal fluctuations. Together our findings suggest fluctuating thermal cues experienced in nature are integrated via a temporal hierarchy of molecular responses during TSD.

Methods

Field-derived nest thermal profiles

Twenty alligator nests were monitored at the Tom Yawkey Wildlife Center (YWC; Georgetown, SC; Bock et al. 2020). At each nest (Fig. 1A), one Onset (UTBI-001) HOBO temperature logger preprogrammed to record temperature at 5-min intervals was deployed in the center of the nest cavity and one temperature logger was attached to vegetation in close proximity to the nest out of direct sunlight to record air temperatures of the nest microclimate. Nests were left undisturbed until after hatchlings emerged (second week of September), after which temperature loggers were retrieved. Raw data from the temperature loggers were processed to only include measurements for the dates encompassing the thermosensitive period (TSP), Ferguson Stages 15-24 (Ferguson 1985; Lang and Andrews 1994; McCoy et al. 2015). Dates of the TSP were estimated based on hatch date, average nest temperature during the first 49 days of the incubation period, and the established relationship between temperature and developmental rate in A. mississippiensis (Kohno and Guillette 2013).

Laboratory incubation experiment

Field collections were performed under permits obtained from the Florida Fish and Wildlife Conservation Commission and the US Fish and Wildlife Service. In 2018, three clutches of alligator eggs were collected shortly after oviposition at Lake Woodruff National Wildlife Refuge (De Leon Springs, FL). Following transport to the Savannah River Ecology Laboratory (Jackson, SC), a representative embryo was examined to determine the developmental stage of each clutch (Ferguson 1985). Eggs were kept in damp sphagnum moss and incubated at 32°C, a temperature promoting the development of both males and females, until the opening of the TSP at Ferguson Stage 15. Thermal sensitivity of sex determination does not begin until Stage 15 and embryos were kept at a single constant temperature prior to this stage to minimize variation resulting from other thermosensitive embryonic traits. Eggs were then shifted to one of the three temperature treatments—a constant 29°C (FPT), constant 33.5°C (MPT), or fluctuating thermal regime based on field-derived nest thermal profiles (average 31.25°C, minimum 29°C, maximum 33.5°C) (González et al. 2019). We derived the fluctuating thermal regime from the eight nest temperature profiles measured during 2015 by determining the average deviation from the mean nest temperature for every 5 min of the day during the TSP, then increasing that deviation by a factor of seven (Fig. 1B). This resulted in a thermal regime that fluctuated on a daily basis between the male-promoting (33.5°C) and female-promoting (29°C) temperatures and exhibited the same periodicity as a wild nest. While daily thermal variation of the experimental fluctuation was greater than average (average daily nest temperature shift = 0.84°C ; experimental daily temperature shift = 4.5°C), it was not outside the scope of measured daily fluctuations (e.g., nests were observed to experience shifts of $\geq 7^{\circ}\text{C}$ within a day).

At Stage 22, the middle of the TSP, alligator embryos was sampled at four time points spanning a daily thermal fluctuation in the fluctuating temperature treatment (FLUX; Fig. 2). The sampling time points corresponded to the minimum temperature (29°C, FPT; T2; n=10), intermediate increasing temperature (31.5°C, T3; n = 10), maximum temperature (33.5°C, MPT, T4; n=9), and intermediate decreasing temperature (31.5°C, T1; n=8) of the FLUX. Embryos exposed to a constant FPT (29°C; n = 11[T1], 10 [T2], 7 [T3], 8 [T4]), and constant MPT (33.5°C; n=9 [T1], 8 [T2], 7 [T3], 11 [T4]) were sampled at the same time points to assess daily variability in gene expression unrelated to temperature and to yield a baseline to which the FLUX group could be compared.

Nucleic acid isolation, cDNA synthesis, and qPCR

Gonadal and brain RNA were extracted using the SV total RNA isolation system (Promega; Madison, WI, USA) and resulting concentrations were quantified via spectrophotometry (NanoDrop One; Thermo Fisher Scientific; Waltham, MA, USA). Synthesis of cDNA was carried out using the iScript cDNA synthesis kit (Bio-Rad; Hercules, CA, USA) according to the manufacturer protocol and using 903.4 and 999 ng of input RNA from the gonad and brain, respectively. Gene expression was assessed via quantitative real-time PCR and reactions were performed in triplicate using an SYBR green reaction mix. Intron-retaining transcripts of JARID2 and KDM6B were selectively targeted for quantification via specially designed primer sets. In particular, we targeted transcripts retaining the same introns reported to be differentially retained in previous studies (11th intron in JARID2 and 19th intron in KDM6B [Deveson et al. 2017]). Information for all associated primers is provided in Fig. 3. All primers were designed to be intron-spanning. Those primers targeting intron-retaining isoforms were designed to span the preceding intron. For each gene and isoform, all expression values were derived from triplicate replicates (for biological and standard samples)




Fig. 1 (a) Image of a representative alligator nest. (B) Design of experimental daily thermal fluctuation using the temperature deviation from the daily mean at 5-min increments in empirically derived nest thermal profiles. Solid black line depicts the average deviation from the mean daily nest temperature across all days in the TSP for eight nests monitored in 2015. Dark gray shading depicts the middle 50th percentile of deviations from the daily mean temperature and light gray shading depicts the middle 90th percentile of deviations from the daily mean temperature. Red and blue lines depict the 95th and 5th percentile of temperature deviations, respectively. Black dotted line depicts the deviation from the daily mean temperature of the experimental thermal fluctuation. Although the amplitude of the experimental trace is greater than the average thermal fluctuation observed in nature, the experimental trace is not outside of ecological relevance as natural nests are observed to fluctuate by 7°C within a single day. (C) Nest thermal profiles from nests monitored during 2015 on which the experimental thermal fluctuation was based. Horizontal dotted lines represent constant temperatures promoting female (blue, maroon) and male (red) development.





Fig. 2 Experimental daily thermal regimes—constant MPT, fluctuating temperature, and constant FPT. Vertical dotted lines depict timepoints at which Stage 22 embryos were sampled.

using a standard curve comprised of target-containing plasmid standards of known concentration (copies/µL) that was run on each qPCR plate.


Expression values for gonad and brain samples were normalized to that of beta-actin (*ACTB*) and ribosomal protein L8 (*RPL8*), respectively, which served as internal controls and were also derived from standard curves. There were no significant effects of temperature, sampling timepoint, or their interaction on the absolute expression levels of either of these housekeeping genes (Supplementary Fig. S2).

Statistical analyses

Statistical analyses were conducted using R statistical software version 3.6.1. Distributions of relative expression values for each gene were tested for normality and homoscedasticity using a Shapiro–Wilk test and Levene's test, respectively. If necessary, a transformation was applied to relative expression values to achieve normality. A log-transformation was applied to values for *JARID2* (all transcripts, intron–



Target	Primer (5'-3')	Annealing temperature (°C)	Product (bp)
(i) All transcripts (non-discriminating)	F:CGAGACCAAAATCACCTTCC R:CACGTCATCAAGTTTGTTCTCC	61.2	102
(ii) Spliced transcripts	F:TATCAGATTGCAACGGCAGA R:TGCCTCATAAAGCTGCTGTC	61.2	120
(iii) Intron-retaining transcripts	F:GCACTTTGCTACCACCCAGT R:GCCCATGGAGCACAAAGTAA	64.5	224



Target	Primer (5'-3')	temperature (°C)	Product (bp)
(i) All transcripts (non-discriminating)	F:ACTTGGCATGAACACAGTGC R:ACAGGGATATTGGCCTGGTA	62.5	243
(ii) Spliced transcripts	F:GCAAGAAGATCGCCTACCAG R:CTCACTTGTCACCAGCAGGA	63.6	101
(iii) Intron-retaining transcripts	F:TGTCCCCCTCCTCCACAG R:GATTCCATCTGCCCAGAGC	59.4	250

Fig. 3 Gene isoforms with primer locations. (A) JARID2 gene model for A. mississippiensis. (B) KDM6B gene model for A. mississippiensis. (i) Entire transcript with locations of primers selecting for all transcripts (not discriminating based on intron–retention status). (ii) Locations of primers selecting for intron-spliced transcripts. (iii) Locations of primers selecting for intron-retaining transcripts (11th intron in JARID2 and 19th intron in KDM6B). Primer information is included in associated tables on right.

retention ratio), KDM6B (all transcripts, intron-retention ratio), and SOX9. A square-root transformaapplied to values Transformations were not sufficient to alleviate non-normality for the IARID2 intron-retention ratio in both the gonad and brain. We examined the influence of temperature treatment, sampling timepoint, and their interaction on relative expression of each gene and isoform using linear mixed effects models with clutch identity included as a random effect. These models were fit using the "lme4" package in R. We obtained P-values from the models using likelihood ratio tests. Due to the persistent non-normality of the JARID2 intron-retention ratio following transformation, we also assessed the effect of temperature and sampling timepoint in the gonad and brain by conducting non-parametric Kruskal-Wallis tests. In both cases, results of the Kruskal-Wallis test conformed to those of the linear mixed effects model. To further examine potential relationships among the responses of genes in the gonad to thermal regime, we assessed pairwise Pearson correlations between each of the isoforms of JARID2 and KDM6B, as well as DMRT1 and SOX9.

Results

Nest thermal profiles

Thermal profiles of 20 alligator nest cavities were monitored between 2015 and 2016 at the YWC (Georgetown, SC, USA). The overall mean

temperature during the TSP for all nests was 32.02°C, a temperature predicted to produce 69.8% males (Lang and Andrews 1994). The coolest nest monitored occurred in 2015 and had a mean nest temperature of 29.00°C, a temperature predicted to produce 100% females. The warmest nest monitored occurred in 2016 and had a mean nest temperature of 33.85°C, a temperature predicted to produce 51.9% males (Lang and Andrews 1994). We observed considerable within-nest variability in temperature (Fig. 1C), with 80% (16 of 20) of nests exhibiting both male- and FPT s within the TSP. All of these nests also exhibited both male- and FPTs within the span of a single day during at least one day during the TSP. On average, nest temperatures varied by 0.84°C within the span of a day. The maximum daily temperature range for an individual nest was 7.61°C and the minimum daily temperature range was 0.08°C. While individual nests varied considerably in their mean temperatures during the TSP, there were distinct commonalities in the shape of their daily thermal cycles (Supplementary Fig. S1). In particular, nests tended to exhibit their coolest temperatures between 06:00 and 18:00 h followed by a temperature maximum occurring between 18:00 and 24:00 h. Nests also tended to warm faster than they cooled (Supplementary Fig. S1). Each of these thermal characteristics was retained in our experimental thermal fluctuation wherein embryos experienced male- and FPTs over the course of a daily thermal cycle.




Fig. 4 Gonadal gene expression of upstream epigenetic modifiers and downstream effectors of sex determination in response to different thermal regimes. (**A**) *JARID2* relative expression ratio of intron-retaining transcripts to all (nondiscriminating) transcripts (IR: ND) for each treatment across sampling time points. (**B**) *JARID2* relative expression of all transcripts. (**C**) *KDM6B* relative expression ratio of intron-retaining transcripts to all transcripts (IR: ND). (**D**) *KDM6B* relative expression of all transcripts. (**E**) *SOX9* relative expression. (**F**) *DMR71* relative expression. All gene expression values normalized to the housekeeping gene *ACTB*. Central dots depict mean, vertical lines depict ±1SD.

Epigenetic modifiers

In the gonad, there was a significant interactive effect of temperature treatment and sampling timepoint on the relative expression ratio of the JARID2 intronretaining transcripts to all transcripts (IR: ND; X²(6) =49.86, P < 0.001; Fig. 4A). A Kruskal–Wallis test of the combined effect of temperature and timepoint on the *JARID2* intron-retention ratio supported this result (H [11] = 81.28, P < 0.001). While embryos exposed to the constant FPT exhibited consistently higher intron-retention in JARID2 compared to embryos exposed to the constant MPT, embryos exposed to the FLUX exhibited variable levels of intron-retention. In particular, intron-retention of JARID2 peaked in FLUX embryos at the timepoint in which these embryos were experiencing a FPT (T2), consistent with a rapid response of intron-retention to temperature (Fig. 4A). We also detected a significant interactive effect of temperature treatment and sampling timepoint on overall transcript abundance of JARID2 (non-discriminating; X² [6] =20.64, P=0.002; Fig. 4B). Similar to the pattern of intron-retention in JARID2, transcript abundance

of *JARID2* was more variable in FLUX embryos. However, in contrast to intron–retention, transcript abundance of *JARID2* peaked at the timepoint after they experienced a FPT (T3; ~7 h after T2; Fig. 4B).

In the brain, intron–retention in *JARID2* exhibited a nearly identical response to temperature as it did in the gonad suggesting a tissue-independent mechanism. There was a significant interactive effect of temperature treatment and sampling timepoint on JARID2 intron-retention in the brain (IR: ND; X²) [6] = 34.105, P < 0.001; Fig. 5A), with intron-retention peaking in FLUX embryos at the FPT timepoint, T2. This result was also supported by a Kruskal-Wallis test of the combined effect of temperature and timepoint on JARID2 intron-retention (H [11] = 83.70, P < 0.001). The response of JARID2 transcript abundance to temperature in the brain was less clear, but there was a significant interactive effect of temperature treatment and sampling time-(non-discriminating; X^2 [6] =point P < 0.001; Fig. 5B).

Levels of intron–retention in *KDM6B* were lower than those of *JARID2* in both the gonad and brain.




Fig. 5 Brain gene expression of upstream epigenetic modifiers in response to different thermal regimes. (**A**) *JARID2* relative expression ratio of intron-retaining transcripts to all transcripts (IR: ND) for each treatment across sampling time points. (**B**) *JARID2* relative expression of all transcripts. (**C**) *KDM6B* relative expression ratio of intron-retaining transcripts to all transcripts (IR: ND). (**D**) *KDM6B* relative expression of all transcripts. All gene expression values normalized to the housekeeping gene *RPL8*. Central dots depict mean, vertical lines depict ±1SD.

There was a significant effect of temperature treatment $(X^2 [2] = 8.6665, P = 0.013)$ but not sampling timepoint or their interaction on the ratio of KDM6B intron-retaining transcripts to all transcripts in the gonad (Fig. 5C). There was also a significant effect of temperature treatment $(X^2 [2] = 84.346,$ P < 0.001) but not sampling timepoint or their interaction on KDM6B transcript abundance in the gonad (Fig. 5D). While gonadal KDM6B transcript abundance appeared highest in the FLUX embryos at the timepoint after embryos received an FPT cue (T3), this variation was not statistically significant. In contrast to JARID2, there was no significant effect of the temperature-by-timepoint interaction on KDM6B transcript abundance (Fig. 5D). In the brain, intron-retention levels in KDM6B were low and there were no effects of temperature, sampling timepoint, or their interaction on the ratio of intronretaining to all KDM6B transcripts. We did, however, detect a significant interactive effect of temperature and timepoint on KDM6Btranscript

abundance in the brain $(X^2 \ [6] = 14.478, P = 0.025)$ (Fig. 5D).

Downstream effector genes of sex determination

There was a significant interactive effect of temperature and timepoint on gonadal SOX9 relative expression (X^2 [6] = 19.2, P = 0.004). Consistent with its role in promoting testicular differentiation, SOX9 relative expression was higher in embryos incubated at MPT compared to those incubated at FPT (Fig. 4E). FLUX embryos exhibited variation in their relative expression of SOX9 with highest relative expression observed at timepoint T3 (one timestep after experiencing an FPT or three timesteps after experiencing an MPT; Fig. 4E). Given the current experimental design, it is not yet possible to distinguish whether this reflects a rapid response of SOX9 relative expression to increasing temperatures following exposure to FPT or a delayed response to MPT.

We observed a similar pattern of relative expression of *DMRT1* compared to that of *SOX9*, though

we did not detect a significant interactive effect of temperature and timepoint on DMRT1 relative expression. There was, however, a significant additive effect of temperature and timepoint (X^2 [3] = 9.3691, P=0.025) on DMRT1 relative expression in the gonad. Similar to SOX9, relative expression of DMRT1 in FLUX embryos was highest at timepoint T3 (one timestep after experiencing an FPT or three timesteps after experiencing an MPT; Fig. 4F), though this did not result in a significant temperature-by-timepoint interactive effect.

The strongest gene expression correlations we observed in the gonad were between the intronretaining and non-discriminating isoforms for *JARID2* and *KDM6B*, the non-discriminating isoform for *JARID2* and the non-discriminating isoform for *KDM6B*, and the intron-retaining isoform for *JARID2* and non-discriminating isoform for *KDM6B* (Supplementary Fig. S3). While we observed a correlation between *DMRT1* and *SOX9* expression, correlations between expression of upstream epigenetic regulators, *KDM6B* and *SOX9*, and downstream effector genes, *DMRT1* and *SOX9*, were comparatively weaker (Supplementary Fig. S3).

Discussion

During TSD, alligator embryos experience rapid fluctuations between FPT and MPT in wild nests. Findings from this study suggest embryos integrate these opposing environmental signals via a temporal hierarchy of responses. Alternative splicing of the epigenetic modifier gene JARID2 responds quickly to thermal fluctuations during sex determination in both the gonad and brain. Transcript abundance of both JARID2 and KDM6B similarly fluctuates with temperature, though with an apparent time delay relative to the splicing response. In contrast to these upstream epigenetic regulators, expression patterns of downstream effector genes of sex determination, DMRT1 and SOX9, are highly variable and do not exhibit as clear a relationship to temperature under fluctuating conditions. These results are consistent with a hypothetical model wherein alternative splicing and transcript abundance of upstream chromatin modifiers fluctuate with temperature resulting in incremental epigenetic changes that influence the transcriptional activity of key sex determining genes (Fig. 6). However, additional experiments testing other key components of this conceptual model, such as how temperature-dependent alternative splicing affects the function of KDM6B and JARID2, are clearly still needed.

The relatively rapid responses of JARID2 and KDM6B to thermal fluctuations suggest these factors are likely proximate targets of a cellular thermosensory mechanism(s) and thus occupy upstream positions in the sex-determining transcriptional cascade. Despite extensive research efforts, the thermosensory mechanism governing TSD has remained elusive. Several candidates have been hypothesized to initially translate incubation temperature into a biological response during TSD including transient receptor potential channels (Yatsu et al. 2015; Lin et al. 2018), heat shock proteins (Kohno et al. 2010; Bentley et al. 2017;), and cold-inducible binding protein (CIRBP) (Rhen and Schroeder 2010; Chojnowski and Braun 2012; Schroeder et al. 2016; Lin et al. 2018) (Fig. 6). All of these factors are generally well-conserved and coordinate cellular responses to temperature across diverse systems (Clapham 2003; Kohno et al. 2010; Zhong and Huang 2017) though evidence for their role in TSD is currently equivocal. CIRBP represents an intriguing candidate as it belongs to a family of RNA recognition motif binding proteins with roles in regulating splicing, translation, and mRNA stability (Dreyfuss et al. 2002; De Leeuw et al. 2007; Xia et al. 2012). Based on data from Chelydra serpentina, temperature-induced transcriptional responses of CIRBP would appear to lag behind the shifts in alternative splicing and transcript abundance of JARID2 and KDM6B reported here (Schroeder et al. 2016). This, however, does not rule out the possibility of rapid post-transcriptional regulation of CIRBP by temperature (Gotic et al. 2016; Haltenhof et al. 2020). A recent report demonstrated role for CDC-like kinases in regulating temperature-sensitive alternative splicing of both CIRBP and JARID2 (Haltenhof et al. 2020), lending support to the possibility that multiple thermosensory mechanisms may act in a coordinated manner during TSD. Further investigations into temperature-induced changes in intracellular signaling cascades (Rottingen and Iversen 2000; Stamm 2008), protein conformation (Haltenhof et al. 2020), and/or conformation of RNA molecules themselves (Johansson et al. 2002; Chowdhury et al. 2006; Loh et al. 2013) during TSD are likely to shed light on the molecular interactions that transduce temperature into a sex-determining signal.

Exposure to FPTs in alligator embryos resulted in both increased intron–retention and increased overall transcript abundance of *JARID2*. Counterintuitively, the intronic sequence retained in mature *JARID2* transcripts contains premature stop codons and intron-retaining transcripts are




Fig. 6 Hypothesized model linking functional and temporal hierarchies of molecular responses to fluctuating thermal cues during TSD. Temperature is initially transduced into a biological signal via one or more thermo-sensory mechanisms. Possible mechanisms include TRP channels which alter intracellular calcium levels, HSPs, CIRBP, RNA thermosensors, or CLK4 which phosphorylate serine arginine-rich proteins with roles in alternative splicing. Upstream epigenetic regulators, JARID2 and KDM6B, are proximate targets of this cellular thermo-sensory mechanism and their transcription and post-transcriptional regulation respond rapidly to thermal fluctuations as a result (1) In particular, transcript abundance of JARID2 and KDM6B increases in the bipotential gonad following exposure to transient female-promoting thermal signals. Intron–retention in JARID2 also responds rapidly to temperature potentially leading to the generation of a truncated protein with altered function or sequestration of "sentinel RNAs" in the nucleus that are later spliced and translated. Once translated, JARID2 and KDM6B influence H3K27me3 marks at the promoters of genes associated with the male and female pathway resulting in incremental changes to the chromatin landscape (2) in response to temperature. Over time, the repressive H3K27me3 mark accumulates at the promoters of genes involved in the male-pathway via the actions of JARID2 in complex with PRC2, while genes involved in the female-pathway progressively lose H3K27me3 at their promoters via actions of KDM6B. As a result, expression of genes involved in the female-pathway increases gradually (3) until reaching a threshold (asterisks) for sexual fate commitment. (References: \(^{1}\)Lin et al. 2018; \(^{2}\)Yatsu et al. 2015; \(^{3}\)Bentley et al. 2017; \(^{4}\)Kohno et al. 2010; \(^{5}\)Rhen and Schroeder 2010; \(^{6}\)Schroeder et al. 2016; \(^{8}\)Johansson et al. 2002; \(^{9}\)Haltenhof et al. 2020).

predicted to produce truncated products with altered or ameliorated function if they avoid nonsensemediated decay and are translated (Deveson et al. 2017). This result is consistent with previous findings in which intron–retention in both *JARID2* and *KDM6B* was more frequent at the temperature promoting expression of these genes in *A. mississippiensis* and *T. scripta* (Deveson et al. 2017; Georges and Holleley 2018). Various explanations have been posited to reconcile these seemingly incongruent observations. For example, intron–retention in *JARID2* and *KDM6B* may influence which genes these factors target for transcriptional regulation or alter how

these factors interact with other chromatin modifier complexes, though these ideas have yet to be tested experimentally (Wong et al. 2016; Georges and Holleley 2018; Marasca et al. 2018). Our finding that intron–retention and transcript abundance respond to fluctuating temperatures on different timescales may provide additional insight into the consequence of this temperature-sensitive alternative splicing in ecologically relevant contexts. It is possible that rapid intron–retention in *JARID2* stabilizes the female-promoting signal during transient decreases in nest temperature during alligator TSD by promoting the storage of intron-retaining

transcripts in the nucleus which are later spliced and transcribed to functional proteins after the FPT cue. An analogous phenomenon is observed in a fern Marsilea vestita, and sea Nematostella vectensis (Moran et al. 2008; Boothby et al. 2013; Wong et al. 2016) in which intronretaining transcripts are stored in the nucleus and serve as "sentinel RNAs" that are later spliced facilitating rapid stage-specific protein translation. Testing whether a similar process plays a role in TSD would necessitate further experiments to determine if intron-retaining transcripts are eventually spliced and translated to functional products (e.g., experiments employing pulse-chase Alternatively, intron-retention serves widespread roles in gene regulation and repression of protein translation (Wong et al. 2013, 2016; Braunschweig et al. 2014), and may serve to buffer against premature sexual fate commitment resulting from temperature-induced transcription of JARID2 and KDM6B, thereby extending the period temperature-sensitivity during TSD. Previous work suggests intron-retention can take on diverse functions across taxa, especially depending on the genomic context of the retained introns (Wong et al. 2016), and thus further experiments to resolve the function of intron-retention in chromatin-modifier genes are likely to yield important insights into TSD.

Many genes with conserved roles in sex determination and differentiation display robust sexually dimorphic expression patterns in TSD species after the TSP (Czerwinski et al. 2016; Yatsu et al. 2016). Yet in this study, we detect highly variable expression patterns of two of these genes, SOX9 and DMRT1, in the middle of the TSP, especially in embryos exposed to a thermal fluctuation. This finding raises the question—how do sexually dimorphic transcriptional patterns arise during TSD and how does this relate to the regulatory actions of JARID2 and KDM6B over time? Ge et al. demonstrated that in T. scripta under constant MPTs, KDM6B promotes commitment to the testicular fate via direct removal of the repressive H3K27me3 mark at the promoter of DMRT1 (Ge et al. 2018). However, under fluctuating conditions in alligator embryos, JARID2 and KDM6B transcript abundance fluctuates with temperature, while downstream effector genes exhibit apparent delays in their responses. This raises the possibility that under fluctuating thermal conditions, chromatin landscapes resulting from the actions of JARID2 and KDM6B may be dynamically remodeled over time to eventually repress or activate target downstream effector genes. However, it remains unknown whether incremental epigenetic changes occur during TSD

under fluctuating conditions, and if so, what ultimately regulates the threshold for sexual fate commitment (Fig. 6). Further, the full battery of genes targeted by *JARID2* and *KDM6B* is yet to be resolved in most TSD species, including *A. mississippiensis*.

While this study revealed intriguing patterns in the responses of upstream epigenetic regulators and downstream effector genes of sex determination to a thermal fluctuation between MPT and FPTs, there are limitations to the current experimental design that leave several important questions unanswered. In particular, while transcript abundance of JARID2 and KDM6B reached a maximum at the timestep after receiving a FPT cue (~7-h delay), this observation is, in part, a result of our sampling scheme and may not reflect the true timescale of this response. In order to better resolve the temporal scale on which alternative splicing, transcription, and translation respond to fluctuating thermal cues during TSD, future experiments must sample embryos across a higher resolution time-series and across multiple days. Furthermore, the thermal regime implemented here represents a simplification of the thermal fluctuations occurring in wild nests. Nests frequently exhibit thermal fluctuations of smaller magnitudes and daily fluctuations often vary over the course of the TSP. It remains unknown how characteristics such as the magnitude and repeatability of fluctuations impact molecular responses to these thermal cues and, ultimately, how these complex thermal profiles shape sex ratio outcomes in species with TSD.

Many questions regarding the epigenetic mechanisms of TSD await further exploration and studies like the one presented here underscore the importance of implementing thermal regimes that accurately reflect the dynamic environment in which TSD evolved and examining these processes across multiple temporal scales in future experimental investigations of this phenomenon.

Acknowledgments

We would like to acknowledge Arnold Brunell and other colleagues from the Florida Fish and Wildlife Conservation Commission for their assistance with permitting and field support during egg collections. Further, we would like to thank Jamie Dozier for his support in permitting and field support at the YWC.

Funding

This work was supported in part by the National Science Foundation [Award # 1754903 to B.B.P.]. Furthermore, this material is based upon work

supported by the Department of Energy Office of Environmental Management [Award Number DE-EM0004391 to the University of Georgia Research Foundation].

Conflict of interest

The authors declare no competing interests.

Disclaimer

This report was prepared as an account of work sponsored by an agency of the US Government. Neither the US Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the US Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the US Government or any agency thereof.

Supplementary data

Supplementary data available at IOB online.

References

- Agger K, Cloos PAC, Christensen J, Pasini D, Rose S, Rappsilber J, Issaeva I, Canaani E, Salcini AE, Helin K. 2007. UTX and JMJD3 are histone H3K27 demethylases involved in HOX gene regulation and development. Nature 449:731–4.
- Agrawal R, Wessely O, Anand A, Singh L, Aggarwal RK. 2009. Male-specific expression of Sox9 during gonad development of crocodile and mouse is mediated by alternative splicing of its proline-glutamine-alanine rich domain. FEBS J 276:4184–96.
- Anand A, Patel M, Lalremruata A, Singh AP, Agrawal R, Singh L, Aggarwal RK. 2008. Multiple alternative splicing of Dmrt1 during gonadogenesis in Indian mugger, a species exhibiting temperature-dependent sex determination. Gene 425:56–63.
- Bentley BP, Haas BJ, Tedeschi JN, Berry O. 2017. Loggerhead sea turtle embryos (*Caretta caretta*) regulate expression of stress response and developmental genes when exposed to a biologically realistic heat stress. Mol Ecol 26:2978–92.
- Bock SL, Lowers RH, Rainwater TR, Stolen E, Drake JM, Wilkinson PM, Weiss S, Back B, Guillette L, Parrott BB. 2020. Spatial and temporal variation in nest temperatures forecasts sex ratio skews in a crocodilian with

- environmental sex determination. Proc Royal Soc B Biol Sci 287:20200210.
- Boothby TC, Zipper RS, Van der Weele CM, Wolniak SM. 2013. Removal of retained introns regulates translation in the rapidly developing gametophyte of *Marsilea vestita*. Dev Cell 24:517–29.
- Braunschweig U, Barbosa-morais NL, Pan Q, Nachman EN, Alipanahi B, Gonatopoulos-Pournatzis T, Frey B, Irimia M, Blencowe BJ. 2014. Widespread intron retention in mammals functionally tunes transcriptomes. Genome Res 24:1774–86.
- Bull JJ. 1980. Sex determination in reptiles. Quart Rev Biol 55:3-21.
- Carter AW, Bowden RM, Paitz RT. 2017. Seasonal shifts in sex ratios are mediated by maternal effects and fluctuating incubation temperatures. Funct Ecol 31:876–84.
- Carter AW, Sadd BM, Tuberville TD, Paitz RT, Bowden RM. 2018. Short heatwaves during fluctuating incubation regimes produce females under temperature-dependent sex determination with implications for sex ratios in nature. Sci Rep 8:1–13.
- Chojnowski JL, Braun EL. 2012. An unbiased approach to identify genes involved in development in a turtle with temperature-dependent sex determination. BMC Genomics 13:308.
- Chowdhury S, Maris C, Allain FHT, Narberhaus F. 2006. Molecular basis for temperature sensing by an RNA thermometer. EMBO J 25:2487–97.
- Clapham DE. 2003. TRP channels as cellular sensors. Nature 426:517–24.
- Czerwinski M, Natarajan A, Barske L, Looger LL, Capel B. 2016. A timecourse analysis of systemic and gonadal effects of temperature on sexual development of the red-eared slider turtle *Trachemys scripta elegans*. Dev Biol 420:166–77.
- da Rocha ST, Boeva V, Escamilla-Del-Arenal M, Ancelin K, Granier C, Matias NR, Sanulli S, Chow J, Schulz E, Picard C, et al. 2014. Jarid2 is implicated in the initial xist-induced targeting of PRC2 to the inactive X chromosome. Mol Cell 53:301–16.
- De Leeuw F, Zhang T, Wauquier C, Huez G, Kruys V, Gueydan C. 2007. The cold-inducible RNA-binding protein migrates from the nucleus to cytoplasmic stress granules by a methylation-dependent mechanism and acts as a translational repressor. Exp Cell Res 313:4130–44.
- Deveson IW, Holleley CE, Blackburn J, Marshall Graves JA, Mattick JS, Waters PD, Georges A. 2017. Differential intron retention in Jumonji chromatin modifier genes is implicated in reptile temperature-dependent sex determination. Sci Adv 3:e1700731.
- Dreyfuss G, Kim VN, Kataoka N. 2002. Messenger-RNA-binding proteins and the messages they carry. Nat Rev Mol Cell Biol 3:195–205.
- Escobedo-Galván AH, López-Luna MA, Cupul-Magaña FG. 2016. Thermal fluctuation within nests and predicted sex ratio of Morelet's Crocodile. J Therm Biol 58:23–8.
- Ferguson MWJ. 1985. Reproductive biology and embryology of the crocodilians. Biology of the reptilia. Vol. 14. New York: John Wiley and Sons.
- Ge C, Ye J, Weber C, Sun W, Zhang H, Zhou Y, Cai C, Qian G, Capel B. 2018. The histone demethylase KDM6B

regulates temperature-dependent sex determination in a turtle species. Science 360:645–8.

- Ge C, Ye J, Zhang H, Zhang Y, Sun W, Sang Y, Capel B, Qian G. 2017. Dmrt1 induces the male pathway in a turtle species with temperature-dependent sex determination. Development 144:2222–33.
- Georges A, Holleley CE. 2018. How does temperature determine sex? Science 360:601–2.
- González EJ, Martínez-López M, Morales-Garduza MA, García-Morales R, Charruau P, Gallardo-Cruz JA. 2019. The sex-determination pattern in crocodilians: a systematic review of three decades of research. J Anim Ecol 88:1417–1.
- Gotic I, Omidi S, Fleury-Olela F, Molina N, Naef F, Schibler U. 2016. Temperature regulates splicing efficiency of the cold-inducible RNA-binding protein gene Cirbp. Genes Dev 30:2005–17.
- Haltenhof T, Kotte A, De Bortoli F, Schiefer S, Meinke S, Emmerichs AK, Petermann KK, Timmermann B, Imhof P, Franz A, et al. 2020. A conserved kinase-based body-temperature sensor globally controls alternative splicing and gene expression. Mol Cell 78:57–13.
- Harry JL, Williams KL, Briscoe DA. 1990. Sex determination in loggerhead turtles: differential expression of two hnRNP proteins. Development 109:305–12.
- Johansson J, Mandin P, Renzoni A, Chiaruttini C, Springer M, Cossart P. 2002. An RNA thermosensor controls expression of virulence genes in Listeria monocytogenes. Cell 110:551–61.
- Kaneko S, Bonasio R, Saldaña-Meyer R, Yoshida T, Son J, Nishino K, Umezawa A, Reinberg D. 2014. Interactions between JARID2 and noncoding RNAs regulate PRC2 recruitment to chromatin. Mol Cell 53:290–300.
- Kohno S, Katsu Y, Urushitani H, Ohta Y, Iguchi T, Guillette LJ. 2010. Potential contributions of heat shock proteins to temperature-dependent sex determination in the American alligator. Sex Dev 4:73–87.
- Kohno S, Guillette LJ. 2013. Endocrine disruption and reptiles: using the unique attributes of temperature-dependent sex determination to assess impacts. In: Matthiessen P, editor. Endocrine disrupters: hazard testing and assessment methods. Hoboken (NJ): John Wiley & Sons. p. 245–71.
- Lan F, Bayliss PE, Rinn JL, Whetstine JR, Wang JK, Chen S, Iwase S, Alpatov R, Issaeva I, Canaani E, et al. 2007. A histone H3 lysine 27 demethylase regulates animal posterior development. Nature 449:689–94.
- Landeira D, Fisher AG. 2011. Inactive yet indispensable: the tale of Jarid2. Trends Cell Biol 21:74–80.
- Lang JW, Andrews HV. 1994. Temperature-dependent sex determination in crocodilians. J Exp Zool 270:28–44.
- Lin JQ, Zhou Q, Yang HQ, Fang LM, Tang KY, Sun L, Wan QH, Fang SG. 2018. Molecular mechanism of temperature-dependent sex determination and differentiation in Chinese alligator revealed by developmental transcriptome profiling. Sci Bull 63:209–12.
- Loh E, Kugelberg E, Tracy A, Zhang Q, Gollan B, Ewles H, Chalmers R, Pelicic V, Tang CM. 2013. Temperature triggers immune evasion by *Neisseria meningitidis*. Nature 502:237–40.
- Marasca F, Bodega B, Orlando V. 2018. How polycombmediated cell memory deals with a changing environment: variations in PcG complexes and proteins assortment

convey plasticity to epigenetic regulation as a response to environment. BioEssays 40:1700137–13.

- Matsumoto Y, Buemio A, Chu R, Vafaee M, Crews D. 2013. Epigenetic control of gonadal aromatase (cyp19a1) in temperature-dependent sex determination of red-eared slider turtles. PLoS One 8:e63599.
- McCoy JA, Parrott BB, Rainwater TR, Wilkinson PM, Guillette LJ. 2015. Incubation history prior to the canonical thermosensitive period determines sex in the American alligator. Reproduction 150:279–87.
- Moran Y, Weinberger H, Reitzel AM, Sullivan JC, Kahn R, Gordon D, Finnerty JR, Gurevitz M. 2008. Intron retention as a posttranscriptional regulatory mechanism of neurotoxin expression at early life stages of the starlet anemone *Nematostella vectensis*. J Mol Biol 380:437–43.
- Navarro-Martín L, Viñas J, Ribas L, Díaz N, Gutiérrez A, Di Croce L, Piferrer F. 2011. DNA methylation of the gonadal aromatase (cyp19a) promoter is involved in temperature-dependent sex ratio shifts in the European sea bass. PLoS Genetics 7:e1002447.
- Ospina-Alvarez N, Piferrer F. 2008. Temperature-dependent sex determination in fish revisited: prevalence, a single sex ratio response pattern, and possible effects of climate change. PLoS One 3:e2837.
- Parrott BB, Kohno S, Cloy-McCoy JA, Guillette LJ. 2014. Differential incubation temperatures result in dimorphic DNA methylation patterning of the SOX9 and aromatase promoters in gonads of alligator (*Alligator mississippiensis*) embryos. Biol Reprod 90:1–11.
- Peng JC, Valouev A, Swigut T, Zhang J, Zhao Y, Sidow A, Wysocka J. 2009. Jarid2/Jumonji coordinates control of PRC2 enzymatic activity and target gene occupancy in pluripotent cells. Cell 139:1290–302.
- Rhen T, Fagerlie R, Schroeder A, Crossley DA, Lang JW. 2015. Molecular and morphological differentiation of testes and ovaries in relation to the thermosensitive period of gonad development in the snapping turtle, *Chelydra serpentina*. Differentiation 89:31–41.
- Rhen T, Schroeder A. 2010. Molecular mechanisms of sex determination in reptiles. Sex Dev 4:16–28.
- Rottingen JA, Iversen JG. 2000. Ruled by waves? Intracellular and intercellular calcium signalling. Acta Physiol Scand 169:203–19.
- Salz HK. 2011. Sex determination in insects: a binary decision based on alternative splicing. Curr Opin Genet Dev 21:395–400.
- Sanulli S, Justin N, Teissandier A, Ancelin K, Portoso M, Caron M, Michaud A, Lombard B, da Rocha ST, Offer J, et al. 2015. Jarid2 methylation via the PRC2 complex regulates H3K27me3 deposition during cell differentiation. Mol Cell 57:769–83.
- Schroeder AL, Metzger KJ, Miller A, Rhen T. 2016. A novel candidate gene for temperature-dependent sex determination in the common snapping turtle. Genetics 203:557–71.
- Stamm S. 2008. Regulation of alternative splicing by reversible protein phosphorylation. J Biol Chem 283:1223–7.
- Tapial J, Ha KCH, Sterne-Weiler T, Gohr A, Braunschweig U, Hermoso-Pulido A, Quesnel-Vallières M, Permanyer J, Sodaei R, Marquez Y, et al. 2017. An atlas of alternative splicing profiles and functional associations reveals new

- regulatory programs and genes that simultaneously express multiple major isoforms. Genome Res 27:1759–68.
- Warner DA, Shine R. 2011. Interactions among thermal parameters determine offspring sex under temperature-dependent sex determination. Proc Royal Soc B Biol Sci 278:256–65.
- Wong JJL, Au AYM, Ritchie W, Rasko JEJ. 2016. Intron retention in mRNA: no longer nonsense: known and putative roles of intron retention in normal and disease biology. BioEssays 38:41–9.
- Wong JJL, Ritchie W, Ebner OA, Selbach M, Wong JWH, Huang Y, Gao D, Pinello N, Gonzalez M, Baidya K, et al. 2013. Orchestrated intron retention regulates normal granulocyte differentiation. Cell 154:583–95.
- Xia Z, Zheng X, Zheng H, Liu X, Yang Z, Wang X. 2012. Cold-inducible RNA-binding protein (CIRP) regulates

- target mRNA stabilization in the mouse testis. FEBS Lett 586:3299–308.
- Yatsu R, Miyagawa S, Kohno S, Parrott BB, Yamaguchi K, Ogino Y, Miyakawa H, Lowers RH, Shigenobu S, Guillette LJ, et al. 2016. RNA-seq analysis of the gonadal transcriptome during *Alligator mississippiensis* temperaturedependent sex determination and differentiation. BMC Genomics 17:1–13.
- Yatsu R, Miyagawa S, Kohno S, Saito S, Lowers RH, Ogino Y, Fukuta N, Katsu Y, Ohta Y, Tominaga M, et al. 2015. TRPV4 associates environmental temperature and sex determination in the American alligator. Sci Rep 5:1–10.
- Zhong P, Huang H. 2017. Recent progress in the research of cold-inducible RNA-binding protein. Fut Sci OA 3:FSO246.