



Complete Mitochondrial Genomes Reveal Population-Level Patterns in the Widespread Red Alga *Gelidiella fanii* (Gelidiales, Rhodophyta)

Ga Hun Boo^{1,2,3*}, Mayalen Zubia⁴, Jeffery R. Hughey⁵, Alison R. Sherwood⁶, Mutue T. Fujii¹, Sung Min Boo² and Kathy Ann Miller³

¹ Nucleus for Research in Phycology, Institute of Botany, São Paulo, Brazil, ² Department of Biology, Chungnam National University, Daejeon, South Korea, ³ University Herbarium, University of California, Berkeley, Berkeley, CA, United States, ⁴ UMR EIO (UPF-IRD-ILM-IFREMER), Université de la Polynésie Française, Labex Corail, Faa'a, French Polynesia, ⁵ Division of Mathematics, Science, and Engineering, Hartnell College, Salinas, CA, United States, ⁶ School of Life Sciences, University of Hawai'i at Mānoa, Honolulu, HI, United States

OPEN ACCESS

Edited by:

Zhijun Dong,
Yantai Institute of Coastal Zone
Research (CAS), China

Reviewed by:

Xiubao Li,
Hainan University, China
Juan E. Uribe,
Smithsonian National Museum
of Natural History (SI), United States

*Correspondence:

Ga Hun Boo
gahunboo@gmail.com

Specialty section:

This article was submitted to
Marine Evolutionary Biology,
Biogeography and Species Diversity,
a section of the journal
Frontiers in Marine Science

Received: 16 July 2020

Accepted: 22 September 2020

Published: 08 October 2020

Citation:

Boo GH, Zubia M, Hughey JR,
Sherwood AR, Fujii MT, Boo SM and
Miller KA (2020) Complete
Mitochondrial Genomes Reveal
Population-Level Patterns
in the Widespread Red Alga *Gelidiella*
fanii (Gelidiales, Rhodophyta).
Front. Mar. Sci. 7:583957.
doi: 10.3389/fmars.2020.583957

Although complete mitogenomic data have been widely applied in human and other animal population studies, they are extremely limited for florideophycean red algal populations. *Gelidiella fanii* is a recently described rhodophyta, previously misidentified as *G. acerosa*, a cosmopolitan agar-yielding species from tropical to subtropical waters. To decipher patterns in genetic diversity and geographic distribution for *G. fanii*, we obtained 10 complete mitogenomes including two outgroups, *G. acerosa* and *G. flabella*. The mitogenomes ranged in size from 25,223 to 25,281 bp and had 48 genes, which are similar in general structure, gene order and content, and presence of a group II intron. Phylogenomic analysis revealed that *G. fanii* was monophyletic and clearly separate from *G. acerosa*. The range of *G. fanii* was extended from Southeast Asia and northern Australia to Eritrea, Juan de Nova Island, and Kenya in the west, and to Hawai'i and Tetiaroa Atoll to the east. Haplotype network analysis of *cox1* revealed seven geographically structured groups: Southeast Asia, Kenya/Juan de Nova Island, Indonesia, northern Australia, the Philippines, Tetiaroa Atoll, and Hawai'i. This regional structure has likely resulted from the separation and isolation of an ancient widespread population during the Pleistocene. This study demonstrates that mitogenome sequencing is a powerful genotyping tool for studies of genetic diversity, biogeography, and conservation of economically valuable marine algal species.

Keywords: agar-yielding algae, biogeography, Gelidiales, haplotype lineages, mitogenomes, widespread species

INTRODUCTION

In red algae, the mitochondrial genome (mitogenome) comprises circular, maternally inherited chromosomes with fast evolving genes (Hughey et al., 2014; Yang et al., 2015). First sequenced from *Chondrus crispus* Stackhouse (Leblanc et al., 1995), mitogenomes have improved our knowledge of evolution, genetics and the taxonomy of red algae. Hughey et al. (2014) was the first to analyze

mitogenomes from archival type specimens of bangiophycidean *Pyropia* species, and proposed a genome-based metric for distinguishing species. Yang et al. (2015) analyzed the mitogenomes of representatives of all five subclasses in the Florideophycidae. They documented the rapid radiation of the class, but concluded that gene synteny in these phylogenetically diverse red algae was highly conserved. In a review of 16 mitogenomes among 30 datasets publicly available at that time, Salomaki and Lane (2016) reported that red algal mitogenomes were more highly conserved than previously reported. Boo et al. (2016a) analyzed 10 mitogenomes from type specimens of species in the order Gelidiales and proposed that, of 23 protein-coding genes (PCGs) present, six PCGs with low non-synonymous (Ka)/synonymous (Ks) ratios were suitable markers for species identification. Mitogenomes have also been used for the purpose of merging morphological species, describing new species and genera, and identifying introduced species (Hughey and Boo, 2016; Suzuki et al., 2016; Song et al., 2017; Gabrielson et al., 2018; Boo and Hughey, 2019; Bustamante et al., 2019). However, red algal mitogenomes have yet to be analyzed from a population perspective.

Gelidiella fanii S.-M.Lin (Gelidiellaceae) is one of 15 species in the widespread genus *Gelidiella* Feldmann and Hamel, found on coral reefs and intertidal rocky shores in tropical to subtropical waters (Abbott, 1999; Costa et al., 2002; Boo et al., 2015; Huisman et al., 2018). *Gelidiella fanii* is morphologically similar to *G. acerosa* (Forsskål) Feldmann and Hamel, a cosmopolitan species yielding high-grade bacteriological and pharmaceutical agarose, as well as agar for food (Rioux and Turgeon, 2015). *Gelidiella fanii* is a recently described species from Taiwan that was previously misidentified as *G. acerosa* (Lin and Freshwater, 2008). It has been shown to be clearly distinct using mitochondrial *cox1* and plastid *rbcL* DNA sequences, as well as by morphology. Thalli are iridescent under water and have downward curved branches with slender unilateral branchlets, numerous surface hairs on the distal end of branches and branchlets, and smaller tetrasporangia (Lin and Freshwater, 2008; Wiriyadamrikul et al., 2010). There have been no reports of sexual reproduction in *G. fanii* (Lin and Freshwater, 2008; Wiriyadamrikul et al., 2010; Boo et al., 2016c; Huisman et al., 2018), a characteristic common in other species of *Gelidiella*. Molecular analyses have confirmed the presence of *G. fanii* in Indonesia, Japan, the Philippines, Thailand, Vietnam, and northern Australia, in addition to Taiwan (Lin and Freshwater, 2008; Wiriyadamrikul et al., 2010; Boo et al., 2016c; Huisman et al., 2018). However, the taxonomy and distribution of *G. fanii* remains understudied outside of Southeast Asia and northern Australia because of its morphological similarity to *G. acerosa*, a species that is recorded globally and is morphologically variable depending on habitat.

To date, 40 individual gene sequences of *G. fanii* are publicly available in GenBank; 23 mitochondrial *cox1*, and 14 plastid *rbcL*, two *psaA*, and one *psbA* from Southeast Asia and northern Australia (Lin and Freshwater, 2008; Wiriyadamrikul et al., 2010; Boo et al., 2016b,c; Huisman et al., 2018). These data have been used for taxonomic studies and analyses of the biogeographical structure of *cox1* haplotype networks in

Southeast Asia. However, limited taxon sampling and individual gene datasets are insufficient for full molecular characterization of the populations of this presumably widespread species. We wished to determine how complete mitogenomes vary among geographically isolated populations of *G. fanii*, and which individual genes have useful resolution at the population level. The aims of this study were i) to obtain complete mitogenomes using High Throughput Sequencing (HTS) techniques, ii) to investigate the utility of intergenic spacer regions as well as PCGs for population studies, and iii) to establish haplotype networks of *cox1* to understand the distribution of *G. fanii* in the Indo-Pacific Ocean.

MATERIALS AND METHODS

Habitat and Collection of Specimens

Like *G. acerosa*, *G. fanii* occurred on subtidal coral reefs and/or intertidal rocky reefs in the Indo-Pacific regions; however, *G. fanii* was very rare compared with *G. acerosa*. Specimens were mostly typical of the species, but they sometimes displayed fewer unilateral branchlets. All specimens collected in this study were vegetative or tetrasporic; and neither spermatangial (male) nor cystocarpic (female) plants were found.

Fresh specimens were collected in Hawai'i, Juan de Nova Island in the Mozambique Channel, Tetiaroa Atoll in French Polynesia, the Philippines, Taiwan and Vietnam, and were placed in individual bags with silica gel until processed. Specimens were identified based on morphological observation as well as analyses using mitochondrial *cox1* and plastid *rbcL* sequences. In addition, herbarium specimens identified as *G. acerosa* were studied on loan from University of California at Berkeley, United States. Voucher specimens are housed in the Natural History Museum, Chungnam National University, Korea (CNUK) and the herbarium of the University of French Polynesia, Tahiti (UPF). Information on all specimens used in this investigation is provided in **Supplementary Table S1**.

DNA Extraction and Genome Sequencing

Eight *Gelidiella fanii* individuals were selected to represent the species' broad geographic distribution in the Indo-Pacific Ocean and the topology of the *cox1* phylogeny (below). DNA was extracted from ~5 mg of dried tissue using NucleoSpin Plant II Kit (Macherey-Nagel, Düren, Germany) according to the manufacturer's protocol or DNeasy Blood and Tissue Kit (Qiagen, Valencia, California, USA) following Boo et al. (2016a). Two genomic DNAs extracted in the previous studies (Wiriyadamrikul et al., 2010; Huisman et al., 2018), one from the Philippines (CNU026931) and the other from northern Australia (CNU066493), were also used for genome sequencing. Two outgroup species, *G. acerosa* and *G. flabella* G.H.Boo and L.Le Gall, were included for HTS. Library preparation and HTS were performed by Genotech Co. (Daejeon, Korea) with Illumina platforms, HiSeq2500 or HiSeqX, using 100 bp or 150 bp paired-end library constructions, respectively.

Genome Assembly and Annotation

The raw reads were assembled using a combination of approaches, NOVOPlasty 3.5 (Dierckxsens et al., 2016), SPAdes assembler 3.13.0 (Nurk et al., 2013) and MEGAHIT (Li et al., 2015). Assembled contigs were sorted and reassembled using Geneious Prime 2019.2.3¹ to construct consensus mitochondrial genome sequences. To confirm the accuracy of the assembly, raw reads were mapped to the draft mitochondrial genomes using Bowtie2 in Geneious Prime.

Mitochondrial genomes were annotated using BlastX and NCBI ORFfinder. Transfer RNA (tRNA) genes were predicted by tRNAscan-SE v. 2.0 (Lowe and Chan, 2016) with default settings and the “Mold/Protozoan Mito” model, RNAweasel server², and Aragorn v.1.2.38 (Laslett and Canback, 2004). Only tRNA positions supported by at least two programs were retained for further analysis. Ribosomal RNA (rDNA) genes were predicted by RNAmmer 1.2 server (Lagesen et al., 2007).

Comparison of Mitogenome Structure and Phylogenetic Analysis

The physical map of the mitogenome was prepared for visualization using OrganellarGenomeDRAW (OGDraw) (Lohse et al., 2013). Locally collinear blocks (LCBs) alignments were generated using ProgressiveMauve (Darling et al., 2004) with a seed of 21 for the mitochondrial alignments and the ‘Use seed families’ option selected. CREx (Bernt et al., 2007) was performed to compare the gene order and their rearrangement events (e.g., inversion, transpositions, reverse transpositions, and tandem-duplication-random-loss [TDRL]) using heuristic pairwise comparisons with the common interval measurement in the genus *Gelidiella*. Twenty-three PCGs were translated into amino acid sequences and aligned using MAFFT 7.450 (Katoh and Standley, 2013) under the default setting in Geneious Prime. The alignment was manually adjusted and the ambiguous sites were deleted by GBlocks v.0.19b (Castresana, 2000). The phylogenetic tree of the concatenated dataset of 23 PCGs was reconstructed using Maximum Likelihood (ML) and Bayesian inference (BI). For the ML analysis, the best-fitting partitioning schemes and models of molecular evolution were inferred by using ModelFinder (Kalyanamoorthy et al., 2017). Based on the Bayesian information criterion (BIC), ModelFinder identified two partitions: mtVer + F + I + G4 for *atp4* + *atp8* + *rpl16* + *rps3* + *rps11* + *sdh3* + *sdhD* + *tatC* and mtVer + F + I + G4 for *atp6* + *atp9* + *cob* + *cox1* + *cox2* + *cox3* + *nad1* + *nad2* + *nad3* + *nad4* + *nad4L* + *nad5* + *nad6* + *rps12* + *sdh2*. The ML analyses were performed using the W-IQ-tree webserver (Trifinopoulos et al., 2016) with 1,000 ultrafast bootstrap (BS) replicates. For the BI analysis, PartitionFinder v.2.1.1 (Lanfear et al., 2016) was used to select the best-fitting partitioning schemes and models of molecular evolution using the greedy algorithm with unlinked branch lengths. The PartitionFinder identified three partitions: CPREV + I + G for *nad2* + *nad5* + *nad3* + *nad4* + *sdh2* + *rps12* + *nad6* + *rpl16* +

atp8 + *rps11* + *rps3* + *atp4*, MTMAM + I + G for *cox1* + *atp9* + *cob* + *cox2* + *cox3* + *atp6* + *nad1* + *nad4L*, and MTMAM + I + G for *tatC* + *sdh3* + *sdhD*. The BI was performed with MrBayes v.3.2.1 (Ronquist et al., 2012) using the Metropolis-coupled Markov Chain Monte Carlo (MC3) with the models selected by PartitionFinder. For each matrix, four million generations of two independent runs were performed with four chains and sampling trees every 100 generations. The burn-in period was identified graphically by tracking the likelihoods at each generation to determine when they reached a plateau. Twenty-five percent of saved trees were removed, and the remaining trees used to calculate the Bayesian posterior probabilities (BPP).

Genetic diversities, in mitogenomes, PCGs, and intergenic spacer levels, were calculated by DnaSP v.5.10.1 (Librado and Rozas, 2009). To test the selection pressure of mitochondrial PCGs, ratios of non-synonymous (*Ka*) versus synonymous substitutions (*Ks*) were measured using DnaSP. Pairwise genetic distances were computed to assess the divergences of *G. fanii*, *G. acerosa* and *G. flabella*. Haplotype networks were constructed from PCGs and intergenic spacer regions with PopART v.1.7 (Leigh and Bryant, 2015) using the median-joining networks (MJN) (Bandelt et al., 1999).

Molecular Analyses of Mitochondrial *cox1* and Plastid *rbcL*

DNA extractions, polymerase chain reaction (PCR) amplification, and sequencing followed Boo et al. (2016b). The primers used for amplifying and sequencing were F7, F645, R753, and RrbcS start for *rbcL* (Freshwater and Rueness, 1994; Lin et al., 2001; Gavio and Fredericq, 2002), and COXI43F and COXI1549R for *cox1* (Geraldino et al., 2006). When large fragments of the analyzed loci could not be amplified in one herbarium specimen (UC1461694, collected in 1962), we were able to amplify and sequence 237 bp of *rbcL* using primers F577, F993, and R753 (Freshwater and Rueness, 1994). Sequences of the forward and reverse strands were determined for all taxa, and the electropherograms were edited using MEGA7 (Kumar et al., 2016) and checked manually. Newly generated sequences were deposited in GenBank. Sequences were aligned using the MUSCLE algorithm in MEGA7 with default parameters and the alignment was manually adjusted.

Phylogenies of individual datasets were reconstructed with Maximum Likelihood (ML) analysis using the W-IQ-tree webserver. The best-fitting substitution model was determined with the model test option (auto), followed by the ML tree search, and 1,000 ultrafast bootstrap replicates. Haplotype network of mitochondrial *cox1* was constructed with PopART v.1.7 using the median-joining networks (MJN). Haplotype diversity (*h*) and nucleotide diversity (π) were calculated for each population and at the species level using DnaSP v.5.1. Non-hierarchical and hierarchical analyses of molecular variance (AMOVA) was performed using Arlequin v.3.5 (Excoffier and Lischer, 2010) with Φ -statistics to quantify the proportion of total genetic variance, with significance of fixation indices tested using 10,000 permutations. The hierarchical partition was set to seven

¹<https://www.geneious.com>

²<https://megasun.bch.umontreal.ca/cgi-bin/RNAweasel/RNAweaselInterface.pl>

groups (I–VII) based on the phylogeny and haplotype network of *cox1*.

RESULTS

Gelidiella Mitogenomes

Ten complete mitogenomes were sequenced using HTS; eight from *Gelidiella fanii* and one of each from *G. acerosa* and *G. flabella* (Table 1). The sequences of *Gelidiella* mitogenomes showed 91.5% identity in most regions relative to *G. fanii*. Pairwise divergences of mitogenomes within *G. fanii* were in a range of $2.4 \pm 0.8\%$ (between Hawai'i and northern Australia). Interspecific pairwise divergences were $14.4 \pm 0.1\%$ between *G. fanii* and *G. acerosa* and $18.0 \pm 0.08\%$ between *G. fanii* with *G. flabella*.

The mitogenomes of *G. fanii* ranged in size from 25,223 bp (JN071, Juan de Nova Island) to 25,263 bp (CNU026931, the Philippines), with highly conserved gene synteny (Table 1 and Supplementary Figure S1). The GC content was in a range of $30.3 \pm 0.1\%$. The mitogenomes contained 48 genes, consisting of 23 PCGs, 23 tRNAs, and 2 rRNA subunits, a result similar to the publicly available mitogenomes of *Gelidium* J.V.Lamoureux and *Pterocladia* B.Santelices and Hommersand (Boo et al., 2016a), which were included in our phylogenomic analysis. All eight *G. fanii* mitogenomes had a group II intron between the *nad5* and *nad4* genes. Three types of tRNAs were found between the *trnA* and *trnN* regions (Figure 1A). TYPE1 (*trnY-trnR-trnS* insertion) was found in most *G. fanii* and *G. acerosa* samples. TYPE2 (*trnS-trnR-trnY* inversion) was present in *G. fanii* from northern Australia (CNU066493). This tRNA inversion rearrangement was inferred from the putative ancestral state of *G. acerosa* (*trnY-trnR-trnS*). TYPE3 (lacking *trnY-trnR-trnS*) was most common in the *Gelidiales* and *Gelidiella flabella* (Figure 1A and Supplementary Table S2).

Phylogenomics and Gene Characteristics

The concatenated dataset of 23 PCGs (5,555 amino acid positions) from 23 mitogenomes, including 13 previously

published mitogenomes (nine *Gelidium* and four *Pterocladia*), was used for phylogenetic analysis. Because the topologies of ML and BI were identical, we show the ML tree with branch supports of MLBS and BPP. Mitochondrial phylogenomics revealed that the three species of *Gelidiella* formed a fully supported clade. *Gelidiella fanii* was monophyletic (BS: 100, BPP: 1.0) and clearly segregated from *G. acerosa* and *G. flabella* (Figure 1B). Two *G. fanii* mitogenomes from the Philippines clustered with full support with those from Tetiaroa Atoll (BS: 100, BPP: 1.0) and those formed a moderately supported clade with Hawai'i (BS: 79, BPP: 0.97). *Gelidiella fanii* from Taiwan formed a clade (BS: 100, BPP: 1.0) with Vietnam, and was sister to Juan de Nova Island.

The characteristics of the 23 PCGs are provided in Table 2. The mean value of the ratio of non-synonymous (Ka) versus synonymous substitutions (Ks) for 23 PCGs was in a range of 0.0000–0.2827 (Supplementary Figure S2). The value was low in *atp9* (0.0000), *cox1* (0.0189), *cob* (0.0258), *cox2* (0.0416), *cox3* (0.0519), and *nad4L* (0.0538) compared to the other PCGs. The number of haplotypes for 23 PCGs is provided in Table 2. The haplotype networks of six genes with a low Ka/Ks ratio are shown in Supplementary Figure S3.

Intergenic spacers, mostly less than 100 bp in size, were detected in 46 regions (Table 3). Six spacer regions were > 100 bp in size: 144 bp in *trnA-trnY*, 145 bp in *rps11-nad3*, 158 bp in *cox2-cox3*, 247 bp in *trnM-rns*, 409 bp in *trnW-trnA*, and 526 bp in *nad4-nad5*. Of these, five regions contained six to seven haplotypes (Figure 2 and Table 3), except the *trnA-trnY* region in northern Australian sequences, which was excluded due to an inversion rearrangement (Table 3).

Phylogeny of Mitochondrial *cox1* and Plastid *rbcl*

Both markers newly confirmed the occurrence of *G. fanii* in Kenya, Juan de Nova Island, and Tetiaroa Atoll; *rbcl* sequences confirmed its presence in Eritrea (Supplementary Figure S4). The unrooted phylogeny of *G. fanii* based on 47 *cox1* sequences, including 21 newly generated sequences, consisted of seven groups (I–VII) (Figure 3); I from Southeast Asia (Taiwan, Thailand, the Philippines, Vietnam, Japan), II from Eastern Africa (Juan de Nova Island, Kenya), III

TABLE 1 | Characteristics of 10 complete mitogenomes of *Gelidiella fanii*, *G. acerosa*, and *G. flabella*.

Species	Voucher	Length (bp)	G + C (%)	PCGs	tRNAs	rRNAs	Group II intron	Total genes
<i>Gelidiella fanii</i>	CNU026931	25,263	30.4	23	23	2	17123–17078 (c)	48
<i>Gelidiella fanii</i>	CNU040585	25,257	30.4	23	23	2	17123–17078 (c)	48
<i>Gelidiella fanii</i>	CNU066493	25,260	30.4	23	23	2	17127–17082 (c)	48
<i>Gelidiella fanii</i>	CNU070344	25,239	30.1	23	23	2	17108–17063 (c)	48
<i>Gelidiella fanii</i>	CNU070353	25,229	30.3	23	23	2	17124–17079 (c)	48
<i>Gelidiella fanii</i>	CNU080505	25,228	30.4	23	23	2	17123–17078 (c)	48
<i>Gelidiella fanii</i>	JN071	25,223	30.3	23	23	2	17120–17075 (c)	48
<i>Gelidiella fanii</i>	PF447	25,230	30.4	23	23	2	17112–17067 (c)	48
<i>Gelidiella acerosa</i>	CNU026920	25,281	29.3	23	23	2	17123–17078 (c)	48
<i>Gelidiella flabella</i>	PC0166624a	24,948	31.2	23	20	2	17149–17104 (c)	45

PCGs, protein-coding genes; (c), complement.




FIGURE 1 | (A) The tRNA insertion and inversion rearrangement of *trnY-trnR-trnS* region in Gelidiales. **(B)** Maximum likelihood phylogeny of *Gelidiella* using 23 protein-coding genes generated in this study and previously published data of the Gelidiales. Both ML bootstrap values ($\geq 50\%$) and Bayesian posterior probabilities (≥ 0.9) are indicated near branches. Bold letter indicates newly generated sequences in this study. Red bar indicates tRNA inversion rearrangement inferred by CREX analysis.

from Indonesia, IV from northern Australia, V from the Philippines, VI from Tetiaroa Atoll, and VII from Hawai'i. Groups III and IV formed a strongly supported clade (BS: 94); groups V and VI were clustered with moderate support (BS: 84).

Of a total of 23 *rbcL* sequences from *G. fanii*, 10 were generated in this study. The short *rbcL* sequences (237 bp) were successfully amplified from the herbarium specimen from Eritrea (UC1461694, as *G. acerosa*) that we identified as *G. fanii*. It differed by 0 to 1 bp (0–0.4%) from all other *G. fanii* specimens (Supplementary Table S3), but was 6 bp (2.5%) different from

G. acerosa and 16 bp (6.8%) from *G. flabella*. The unrooted phylogeny of *rbcL* was similar to that of *cox1* (Supplementary Figure S4), except that the Hawaiian sequences grouped with Kenya and Juan de Nova Island.

Genetic Diversity, Haplotype Network, and Population Structure of Mitochondrial *cox1*

The *cox1* dataset revealed high estimates of haplotype diversity ($h = 0.931$) and nucleotide diversity ($\pi = 0.00951$), indicating

TABLE 2 | A comparison of 23 protein-coding and two ribosomal genes in *Gelidiella fanii*.

Gene	Length ^a	V (%)	S	PI	$\pi \pm SD$	π_s	π_{ns}	<i>h</i>	Indel sites	Ka/Ks ratio ^b
<i>atp4</i>	543	50 (9.2)	36	14	0.0291 $\pm$ 0.0049	0.0819	0.0152	7	NA	0.2095
<i>atp6</i>	762	53 (7.0)	33	20	0.0238 $\pm$ 0.0026	0.0838	0.0055	7	NA	0.0614
<i>atp8</i>	402	39 (9.7)	25	14	0.0336 $\pm$ 0.0052	0.0884	0.0193	7	15	0.2369
<i>atp9</i>	231	14 (6.1)	9	5	0.0206 $\pm$ 0.0028	0.0848	0.0000	7	NA	0.0000
<i>cob</i>	1146	78 (6.8)	47	31	0.0236 $\pm$ 0.0037	0.0939	0.0027	7	NA	0.0258
COI-5P [†]	664	35 (5.3)	23	12	0.0175 $\pm$ 0.0026	0.0728	0.0000	7	NA	0.0000
<i>cox1</i>	1602	85 (5.3)	58	27	0.0170 $\pm$ 0.0026	0.0677	0.0014	8	NA	0.0189
<i>cox2</i>	783	49 (6.3)	31	18	0.0208 $\pm$ 0.0025	0.0835	0.0034	6	NA	0.0416
<i>cox3</i>	819	55 (6.7)	36	19	0.0225 $\pm$ 0.0029	0.0854	0.0039	7	NA	0.0519
<i>nad1</i>	981	54 (5.5)	30	24	0.0196 $\pm$ 0.0021	0.0695	0.0039	7	NA	0.0635
<i>nad2</i>	1473	117 (7.9)	58	59	0.0294 $\pm$ 0.0029	0.0831	0.0135	7	NA	0.1553
<i>nad3</i>	366	27 (7.4)	21	6	0.0220 $\pm$ 0.0035	0.0769	0.0051	7	NA	0.0627
<i>nad4</i>	1473	108 (7.3)	68	40	0.0246 $\pm$ 0.0030	0.0835	0.0069	7	NA	0.0833
<i>nad4L</i>	306	16 (5.2)	8	8	0.0186 $\pm$ 0.0025	0.0718	0.0029	7	NA	0.0538
<i>nad5</i>	1986	151 (7.6)	99	52	0.0253 $\pm$ 0.0035	0.0811	0.0081	8	NA	0.0950
<i>nad6</i>	609	56 (9.2)	28	28	0.0344 $\pm$ 0.0037	0.1174	0.0094	8	NA	0.0740
<i>rpl16</i>	405	36 (8.9)	20	16	0.0323 $\pm$ 0.0044	0.1106	0.0127	8	NA	0.1173
<i>rps3</i>	723	80 (11.1)	45	35	0.0399 $\pm$ 0.0050	0.0909	0.0260	7	NA	0.2827
<i>rps11</i>	363	37 (10.2)	23	14	0.0345 $\pm$ 0.0046	0.0961	0.0184	7	NA	0.1940
<i>rps12</i>	366	24 (6.6)	11	13	0.0249 $\pm$ 0.0024	0.0833	0.0066	6	NA	0.0836
<i>sdh2</i>	753	57 (7.6)	40	17	0.0247 $\pm$ 0.0033	0.0950	0.0084	8	NA	0.0724
<i>sdh3</i>	381	42 (11.0)	19	23	0.0436 $\pm$ 0.0046	0.1094	0.0251	7	NA	0.2236
<i>sdhD</i>	243	14 (5.8)	10	4	0.0190 $\pm$ 0.0035	0.0695	0.0062	6	NA	0.0979
<i>tatC</i>	750	76 (10.1)	42	34	0.0361 $\pm$ 0.0039	0.1033	0.0179	8	NA	0.1825
LSU	2538	78 (3.1)	56	22	0.0098 $\pm$ 0.0018	NA	NA	8	6	NA
SSU	1354	36 (2.7)	29	7	0.0080 $\pm$ 0.0013	NA	NA	7	4	NA

Length, variable sites, nucleotide diversity, and Ka/Ks ratio. V, variable site; S, singleton; PI, parsimony informative site, π , nucleotide diversity; SD, standard deviation; π_s , synonymous diversity; π_{ns} , non-synonymous diversity; *h*, number of haplotype; ^aaligned length; ^bmean value; [†]DNA barcoding region; NA, not applicable.

genetic heterogeneity within the species. The median-joining network consisted of 19 haplotypes in seven groups, connected by many missing haplotypes related to the geographical distances between sampling locations (**Figure 4**). Haplotypes from Indonesia (C13) and northern Australia (C14) were distantly related (32 missing haplotypes) while Southeast Asian haplotypes (C1–C10) were connected by 1–3 missing haplotypes. Two haplotypes from Taiwan shared sequences with Thailand (C2) and the Philippines (C6), respectively. Haplotypes from the Philippines were separated into two groups: one was restricted to Cebu (C15–C17), while the other (C6), shared with Taiwan, clustered with Southeast Asian haplotypes. The Indonesian haplotype (C13) was distantly connected to those from Southeast Asia. Haplotypes from northern Australia (C14), Tetiaroa Atoll (C18), and Hawai'i (C19) were also isolated.

Non-hierarchical AMOVA showed that most of the *cox1* variation within species was found among populations (85.36%; **Table 4**), while a smaller amount of genetic variation was found within populations (14.64%). Genetic subdivision was highly significant among populations ($\Phi_{ST} = 0.85$, $P < 0.001$). The hierarchical AMOVA indicated the total genetic variance was mainly explained by the variance among groups (88.16%, $\Phi_{CT} = 0.88$, $P < 0.001$); the variances among populations within groups (6.23%, $\Phi_{SC} = 0.53$, $P < 0.001$) and within

populations (5.62%, $\Phi_{ST} = 0.94$, $P < 0.001$) were much lower (**Table 4**).

DISCUSSION

We sequenced eight complete mitogenomes of *Gelidiella fanii* and two outgroup species, *G. acerosa* and *G. flabella*. Gene content and organization in the newly determined *Gelidiella* mitogenomes are similar to published reports for *Gelidium* and *Pterocladia*. However, the *Gelidiella* mitogenome is slightly larger (45–48 genes and 25,223–25,281 bp) than those (43–44 genes and 24,901–24,970 bp) of *Gelidium* and *Pterocladia* (Yang et al., 2015; Boo et al., 2016a; Boo and Hughey, 2019). The differences are due to the addition of three tRNA genes centrally positioned in the tRNA track between *atp6-tatC* of all three species of *Gelidiella*. The three tRNAs are tyrosine, arginine, and serine, which are located between arginine and asparagine. Interestingly, an insertion of tRNA track, *trnY-trnR-trnS* (TYPE1) was characteristic of *G. fanii* and *G. acerosa*, supporting their close relationship compared to *G. flabella* and other species in the Gelidiales. The Australian *G. fanii* differs, however, by having an inversion, *trnS-trnR-trnY* (TYPE2).

The pairwise divergences in mitogenome data ranged from 0.07 to 3.4% within *G. fanii*, which is higher than

TABLE 3 | A comparison of intergenic spacers in *Gelidiella fanii*.


Intergenic region	Length ^a	V (%)	S	PI	$\pi \pm SD$	<i>h</i>	Indel site	Indel event
nad4L-rnl	16	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
LSU-rps3	18	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
rpl16-trnD	7	1 (14.3)	1	0	0.0833 $\pm$ 0.0601	2	4	1
trnD-cox1	41	4 (9.8)	3	1	0.0288 $\pm$ 0.0091	4	0	0
cox1-cox2	4	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
cox2-cox3	158	10 (6.3)	7	3	0.0204 $\pm$ 0.0036	6	9	1
cox3-atp4	5	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
atp4-trnG	7	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
trnG-trnQ	3	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
trnQ-trnL	10	3 (30.0)	2	1	0.1295 $\pm$ 0.0527	3	2	2
trnL-cob	45	3 (6.7)	3	0	0.0174 $\pm$ 0.0126	2	2	2
cob-trnL	31	3 (9.7)	3	0	0.0268 $\pm$ 0.0136	3	3	2
trnL-nad6	2	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
nad6-trnG	23	5 (21.7)	1	4	0.1007 $\pm$ 0.0157	5	1	1
trnG-trnH	8	4 (50.0)	4	0	0.1250 $\pm$ 0.0593	3	0	0
trnH-sdh2	31	5 (16.1)	3	2	0.0744 $\pm$ 0.0190	4	7	4
sdh2-sdh3	5	0 (0)	0	0	0.0000 $\pm$ 0.0000	1	0	0
sdh3-trnF	19	2 (10.5)	2	0	0.0263 $\pm$ 0.0125	3	0	0
trnF-trnS	8	1 (12.5)	1	0	0.0313 $\pm$ 0.0225	2	0	0
trnS-trnP	6	1 (16.7)	1	0	0.0417 $\pm$ 0.0300	2	0	0
trnP-atp9	5	0	0	0	0.0000 $\pm$ 0.0000	1	1	1
atp9-trnC	49	4 (8.2)	1	3	0.0350 $\pm$ 0.0074	6	2	2
trnC-trnM	5	2 (40.0)	2	0	0.1000 $\pm$ 0.0474	3	0	0
trnM-tps11	1	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
rps11-nad3	145	8 (5.5)	3	5	0.0235 $\pm$ 0.0037	6	2	2
nad3-nad1	10	1 (10.0)	1	0	0.0250 $\pm$ 0.0180	2	0	0
nad1-nad2	11	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
nad2-sdhD	9	2 (22.2)	1	1	0.0873 $\pm$ 0.0221	3	0	0
sdhD-nad4	9	0	0	0	0.0000 $\pm$ 0.0000	1	1	1
nad4-nad5	526	20 (3.8)	11	9	0.0131 $\pm$ 0.0024	7	7	3
nad5-atp8	23	2 (8.7)	1	1	0.0295 $\pm$ 0.0137	3	0	0
atp8-atp6	3	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
atp6-trnW	19	1 (5.3)	1	0	0.0132 $\pm$ 0.0095	2	0	0
trnW-trnA	409	48 (11.7)	28	20	0.0422 $\pm$ 0.0050	7	3	3
trnA-trnY*	144	17 (11.8)	6	11	0.0670 $\pm$ 0.0115	6	6	2
trnY-trnR*	4	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
trnR-trnS*	9	3 (33.3)	1	2	0.1556 $\pm$ 0.0409	3	0	0
trnS-trnN	68	5 (7.4)	0	5	0.0315 $\pm$ 0.0096	3	0	0
trnN-trnV	12	1 (8.3)	1	0	0.0227 $\pm$ 0.0164	2	1	1
trnV-trnR	2	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
trnR-trnK	4	1 (25.0)	0	1	0.1071 $\pm$ 0.0422	2	0	0
trnK-tatC	14	2 (14.3)	1	1	0.0522 $\pm$ 0.0171	3	1	1
tatC-rps12	1	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
trnE-trnM	3	0	0	0	0.0000 $\pm$ 0.0000	1	0	0
trnM-rns	247	31 (12.6)	13	18	0.0483 $\pm$ 0.0046	7	3	3
SSU-nad4L	25	2 (8.0)	1	1	0.0327 $\pm$ 0.0083	3	1	1

Length, variable site, nucleotide diversity, and indel. V, variable site; S, singleton; PI, parsimony informative site, π nucleotide diversity; SD, standard deviation; *h*, number of haplotype; ^aaligned length; *excluded CNU066493 due to the rearrangement of tRNAs.

that in *Pterocladia capillacea* (S.G.Gmelin) Santelices and Hommersand from the Galápagos Islands and Pacific Mexico (Boo et al., 2016a). However, pairwise divergence in *rbcL* was 0–0.5%, a value lower than interspecific divergences (0.8–0.9%) in the Gelidiales, with the exception

of a closely related sibling species (Boo et al., 2014). The pairwise divergences in *rbcL* fell within the large range of reported rhodomelacean divergences, e.g., 0.4–1% in *Symphyocladia dendroidea* (Montagne) D.Bustamante, B.Y.Won, S.C.Lindstrom and T.O.Cho, and 0.3–0.7% in





previous studies have recently distinguished *G. fanii* from its very similar and widespread congener, *G. acerosa*, with which it has been long confused.

The phylogeny and haplotype network of *Gelidiella fanii* detected strong signals of genetic differentiation into seven geographical groups; i) Southeast Asia, ii) Kenya/Juan de Nova Island, iii) Indonesia, iv) northern Australia, v) the Philippines, vi) Tetiaroa Atoll, and vii) Hawai'i. The origin of these groups may be interpreted via two scenarios. One hypothesis is that ancient populations of *G. fanii* were widely distributed in the Tethys Ocean and the local populations accumulated genetic changes at the geographical periphery. Subsequently, local populations have likely been geographically isolated by changes in sea level and other oceanographic conditions during the Pleistocene epoch (2.6 Ma – 11.7 ka), which experienced many glacial-interglacial cycles (Herzschuh et al., 2016). A similar hypothesis has been proposed for *Gelidiophycus* G.H.Boo, J.K.Park and S.M.Boo and *Mazzaella* G.De Toni (Montecinos et al., 2012; Boo et al., 2019). This hypothesis is supported by low dispersal capacity of *G. fanii*, a small species lacking flotation and with asexual reproduction. Our second hypothesis is that Southeast Asia, rich in haplotypes, was the center of origin of *G. fanii*. Populations may have subsequently migrated to marginal regions of the Indo-Pacific Ocean. This hypothesis has been proposed for *Portieria* Zanardini species (Gigartinales) that

originated in the Central Indo-Pacific and then dispersed in all directions, although the origins of species in remote islands were uncertain (Leliaert et al., 2018). Additional sampling in more regions is necessary to differentiate the two hypotheses, but the logistics of sampling are currently difficult. It is unlikely that *G. fanii* has recently spread via introductions because of the paucity of shared haplotypes among the seven groups. The many missing haplotypes also indicate deep history.

The mitochondrial genome is a significant source of molecular markers at different levels of resolution. The *cox1* gene, with the lowest Ka/Ks ratio (0.0189), is considered a good marker at the population level and has a high degree of resolution at the species level (Boo et al., 2016b). The large size (1,602 bp) of *cox1* also provides evolutionarily important information with many singletons (58) and haplotypes (8). Its utility as a population marker has been demonstrated for *Gelidiophycus* species (Boo et al., 2019). COI-5P (~664 bp), a conserved part of *cox1* gene, is the well-known DNA barcoding marker for identifying red algal species and populations (e.g., Saunders, 2005; Sherwood et al., 2010; Hu et al., 2015; Freshwater et al., 2017; Leliaert et al., 2018). The haplotype number (7) and nucleotide diversity (0.0175 ± 0.0026) of COI-5P are similar to those of *cox1* gene, supporting its utility in population studies. Both *cox2* and *cox3* can also be useful markers; *cox2* (783 bp) contained 31 singletons and six haplotypes, and *cox3* (819 bp) contained 36 singletons and




TABLE 4 | Non-hierarchical and hierarchical analyses of molecular variance (AMOVA) of mitochondrial *cox1* from *Gelidiella fanii*.

Dataset	Source of variation	d.f.	Sum of squares	Variance component	Percentage of variance	Fixation index
Non-hierarchical	Among populations	10	220.351	6.409	85.36	$\Phi_{ST} = 0.85^{***}$
	Within populations	31	34.078	1.099	14.64	
Hierarchical	Among groups	6	217.595	10.284	88.16	$\Phi_{CT} = 0.88^{***}$
	Among populations within groups	5	17.172	0.726	6.23	$\Phi_{SC} = 0.53^{***}$
	Within populations	30	19.661	0.655	5.62	$\Phi_{ST} = 0.94^{***}$

The hierarchical level "group" indicates 7 groups based on the phylogeny and haplotype network of mitochondrial *cox1*. d.f. = degrees of freedom, *** $P < 0.001$.

seven haplotypes. The cytochrome b gene (*cob*, 1,146 bp), with 47 singletons and seven haplotypes, has been used as a DNA barcode at the species and population levels (Saunders and Moore, 2013; Yoon et al., 2014).

Five intergenic spacers described in this study are candidates for markers to distinguish populations. The *cox2-cox3* (158 bp) contained six haplotypes with a nucleotide diversity of 0.0204 ± 0.0036 . This spacer region (~350–400 bp including flanking parts) has been used to identify red algal species and populations (e.g., Zuccarello et al., 1999; Lim et al., 2013; Kamiya and West, 2014; Pezzolesi et al., 2019). *Rps11-nad3* (145 bp) may be a suitable marker with its six haplotypes and nucleotide

diversity of 0.0235 ± 0.0037 . *Nad4-nad5*, *trnW-trnA*, and *trnM-rns* contained seven haplotypes; however, nucleotide diversity was higher in *trnW-trnA* (0.0422 ± 0.0050) and *trnM-rns* (0.0483 ± 0.0046) than *nad4-nad5* (0.0131 ± 0.0024). Suitable primers for amplifying the above spacer regions have yet to be designed and tested for their practical use in population studies.

CONSERVATION AND CONCLUSION

This study, a detailed characterization of the phylogeography of *G. fanii* using data from the mitogenomes of eight populations,

reveals significant genetic structure correlated with geographic distribution. The continued persistence of *G. fanii* is unclear, considering its rarity and close association with *G. acerosa*, with which it likely competes for substrate, nutrients and light. There is a critical shortage of gelidioid biomass for the production of high-grade bacteriological and pharmaceutical agar (Callaway, 2015; Santos and Melo, 2018). The harvest of *Gelidiella* is possible due to the higher productivity of populations in tropical waters compared to those of temperate species of *Gelidium* (Ganzon-Forbes, 1994; Santos and Melo, 2018). Commercial harvest of *G. acerosa* for agar impacts *G. fanii* and, because it is rare, population reduction over time is probable. Minimizing the loss of genetic diversity is a key factor for conserving economically important marine species. Populations with distinct haplotypes can be recognized and managed as independent conservation units. Populations could also be protected within marine reserves or marine protected areas. Systematic management of natural populations can improve biomass yields for the agar industry. Ecological and phenological studies will be needed in the future to detect changes in population sizes and shifts in geographical ranges due to overharvesting and anticipated oceanographic climate change.

Mitogenome data are currently unavailable for population studies of florideophycean red algae, a large and diverse group comprising 7,000 species. Our results represent a new path toward a mitogenomic picture of the evolution and biogeography of widespread red algal species. This study is the first to describe and analyze complete mitogenomes from the family Gelidiellaceae and to test their utility for population-level studies. We have extended the range of *G. fanii* from Southeast Asia and northern Australia to the eastern and western margins of the Indo-Pacific Ocean, and have suggested that the observed patterns of genetic groups may have resulted from sea level fluctuations or other environmental changes that occurred during the Pleistocene. On the basis of our mitogenome sequences, we propose that intergenic spacer regions as well as PCGs can be explored as suitable markers at the population level for red algae. Markers derived from mitogenomic analyses are important tools for tracking species ranges and developing future conservation strategies for ecologically and economically important red algal species.

DATA AVAILABILITY STATEMENT

The datasets generated for this study can be found in GenBank under the accession numbers, MT742595–MT742604

REFERENCES

- ## DISCUSSION
- reveals significant genetic structure correlated with geographic distribution. The continued persistence of *G. fanii* is unclear, considering its rarity and close association with *G. acerosa*, with which it likely competes for substrate, nutrients and light. There is a critical shortage of gelidioid biomass for the production of high-grade bacteriological and pharmaceutical agar (Callaway, 2015; Santos and Melo, 2018). The harvest of *Gelidiella* is possible due to the higher productivity of populations in tropical waters compared to those of temperate species of *Gelidium* (Ganzon-Forbes, 1994; Santos and Melo, 2018). Commercial harvest of *G. acerosa* for agar impacts *G. fanii* and, because it is rare, population reduction over time is probable. Minimizing the loss of genetic diversity is a key factor for conserving economically important marine species. Populations with distinct haplotypes can be recognized and managed as independent conservation units. Populations could also be protected within marine reserves or marine protected areas. Systematic management of natural populations can improve biomass yields for the agar industry. Ecological and phenological studies will be needed in the future to detect changes in population sizes and shifts in geographical ranges due to overharvesting and anticipated oceanographic climate change.
- Mitogenome data are currently unavailable for population studies of florideophycean red algae, a large and diverse group comprising 7,000 species. Our results represent a new path toward a mitogenomic picture of the evolution and biogeography of widespread red algal species. This study is the first to describe and analyze complete mitogenomes from the family Gelidiellaceae and to test their utility for population-level studies. We have extended the range of *G. fanii* from Southeast Asia and northern Australia to the eastern and western margins of the Indo-Pacific Ocean, and have suggested that the observed patterns of genetic groups may have resulted from sea level fluctuations or other environmental changes that occurred during the Pleistocene. On the basis of our mitogenome sequences, we propose that intergenic spacer regions as well as PCGs can be explored as suitable markers at the population level for red algae. Markers derived from mitogenomic analyses are important tools for tracking species ranges and developing future conservation strategies for ecologically and economically important red algal species.
- ## DATA AVAILABILITY STATEMENT
- The datasets generated for this study can be found in GenBank under the accession numbers, MT742595–MT742604
- ## REFERENCES
- Abbott, I. A. (1999). *Marine Red Algae of the Hawaiian Islands*. Honolulu, HI: Bishop Museum Press.
- Bandelt, H. J., Forster, P., and Röhl, A. (1999). Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16, 37–48. doi: 10.1093/oxfordjournals.molbev.a026036
- Bernt, M., Merkle, D., Ramsch, K., Fritsch, G., Perseke, M., Bernhard, D., et al. (2007). CREx: inferring genomic rearrangements based on common intervals. *Bioinformatics* 23, 2957–2958. doi: 10.1093/bioinformatics/btm468
- Boo, G. H., and Hughey, J. R. (2019). Phylogenomics and multigene phylogenies decipher two new cryptic marine algae from California, *Gelidium gabrielsonii* and *Gelidium kathyanniae* (Gelidiales, Rhodophyta). *J. Phycol.* 55, 160–172. doi: 10.1111/jpy.12802
- Callaway, J. (2015). *Gelidium* spp. and *Gelidium* spp. for agar production. *Mar. Biotechnol.* 17, 1–10. doi: 10.1007/s10291-015-0400-0
- Ganzon-Forbes, S. (1994). *Gelidium* spp. and *Gelidium* spp. for agar production. *Mar. Biotechnol.* 17, 1–10. doi: 10.1007/s10291-015-0400-0
- Santos, M. A., and Melo, F. (2018). *Gelidium* spp. and *Gelidium* spp. for agar production. *Mar. Biotechnol.* 17, 1–10. doi: 10.1007/s10291-015-0400-0
- Santos, M. A., and Melo, F. (2018). *Gelidium* spp. and *Gelidium* spp. for agar production. *Mar. Biotechnol.* 17, 1–10. doi: 10.1007/s10291-015-0400-0
- ## ACKNOWLEDGMENTS
- The authors thank Olivier de Clerck, Eric Coppejans, John Huisman, Paul Geraldino, Line Le Gall, Shao-Lun Liu, and Tu van Nguyen for providing materials or help with collection trips. The authors also thank Tetiaroa Society for the logistical support in Tetiaroa atoll sampling.
- ## AUTHOR CONTRIBUTIONS
- GHB, MTF, SMB, and KAM conceived and designed the study. GHB, MZ, ARS, SMB, and KAM performed the collections and provided samples. GHB and JRH performed laboratory work and the data analyses. All authors participated in the interpretation of the data and writing the manuscript.
- ## FUNDING
- This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) by the Ministry of Education (2018R1A6A3A03012648), the São Paulo Research Foundation (FAPESP) for Bolsas Concedidas como Itens Orçamentários em Auxílios/BCO – Pós-Doutorado (2019/16956-2), and postdoctoral funding from the Silva Center for Psychological Documentation, University Herbarium, University of California at Berkeley to GHB; BIORECIE program with the financial support of the Institut National des Sciences de l'Univers (INSU), the Institut de Recherche pour le Développement (IRD), the Agence des Aires Marines Protégées (AAMP), the Fondation pour la Recherche sur la Biodiversité (FRB), and the TAAF (“Terres Australes et Antarctiques Françaises”) to MZ; the U.S. National Science Foundation (DEB-1754117) to ARS; and the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2016/50370-7) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq for the Productivity Fellowship (304899/2017-8) to MTF.
- ## SUPPLEMENTARY MATERIAL
- The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2020.583957/full#supplementary-material>

AUTHOR CONTRIBUTIONS

GHB, MTF, SMB, and KAM conceived and designed the study. GHB, MZ, ARS, SMB, and KAM performed the collections and provided samples. GHB and JRH performed laboratory work and the data analyses. All authors participated in the interpretation of the data and writing the manuscript.

FUNDING

This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) by the Ministry of Education (2018R1A6A3A03012648), the São Paulo Research Foundation (FAPESP) for Bolsas Concedidas como Itens Orçamentários em Auxílios/BCO – Pós-Doutorado (2019/16956-2), and postdoctoral funding from the Silva Center for Psychological Documentation, University Herbarium, University of California at Berkeley to GHB; BIORECIE program with the financial support of the Institut National des Sciences de l'Univers (INSU), the Institut de Recherche pour le Développement (IRD), the Agence des Aires Marines Protégées (AAMP), the Fondation pour la Recherche sur la Biodiversité (FRB), and the TAAF (“Terres Australes et Antarctiques Françaises”) to MZ; the U.S. National Science Foundation (DEB-1754117) to ARS; and the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2016/50370-7) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq for the Productivity Fellowship (304899/2017-8) to MTF.

ACKNOWLEDGMENTS

The authors thank Olivier de Clerck, Eric Coppejans, John Huisman, Paul Geraldino, Line Le Gall, Shao-Lun Liu, and Tu van Nguyen for providing materials or help with collection trips. The authors also thank Tetiaroa Society for the logistical support in Tetiaroa atoll sampling.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2020.583957/full#supplementary-material>

- Boo, G. H., Hughey, J. R., Miller, K. A., and Boo, S. M. (2016a). Mitogenomes from type specimens, a genotyping tool for morphologically simple species: ten genomes of agar-producing red algae. *Sci. Rep.* 6:e35337. doi: 10.1038/srep35337
- Boo, G. H., Le Gall, L., Miller, K. A., Freshwater, D. W., Wernberg, T., Terada, R., et al. (2016b). A novel phylogeny of the Gelidiales (Rhodophyta) based on five genes including nuclear *CesA*, with descriptions of *Orthogonaccladia* gen. nov. and *Orthogonaccladiaceae* fam. nov. *Mol. Phylogenet. Evol.* 101, 359–372. doi: 10.1016/j.ympev.2016.05.018
- Boo, G. H., Nguyen, T. V., Kim, J. Y., Le Gall, L., Rico, J. M., Bottalico, A., et al. (2016c). A revised classification of the Gelidiellaceae (Rhodophyta) with description of three new genera: *Huismaniella*, *Millerella*, and *Perronella*. *Taxon* 65, 965–979. doi: 10.12705/655.2
- Boo, G. H., Kim, K. M., Nelson, W. A., Riosmena-Rodríguez, R., Yoon, K. J., and Boo, S. M. (2014). Taxonomy and distribution of selected species of the agarophyte genus *Gelidium* (Gelidiales, Rhodophyta). *J. Appl. Phycol.* 26, 1243–1251. doi: 10.1007/s10811-013-0111-7
- Boo, G. H., Le Gall, L., Rousseau, F., de Reviers, B., Coppejans, E., Anderson, R., et al. (2015). Phylogenetic relationships of *Gelidiella* (Gelidiales, Rhodophyta) from Madagascar with a description of *Gelidiella incrassata* sp. nov. *Cryptogam. Algal.* 36, 219–237. doi: 10.7872/crya.v36.iss2.2015.219
- Boo, G. H., Qui, Y.-X., Kim, J. Y., Ang, P. O., Bosch, S., De Clerck, O., et al. (2019). Contrasting patterns of genetic structure and phylogeography in the marine agarophytes *Gelidiophycus divaricatus* and *G. freshwateri* (Gelidiales, Rhodophyta) from East Asia. *J. Phycol.* 55, 1319–1334. doi: 10.1111/jpy.12910
- Boo, G. H., Robledo, D., Andrade-Sorcia, G., and Boo, S. M. (2018). Genetic discontinuity of *Digenea* (Rhodomelaceae, Rhodophyta) from Mexico supports recognition of two new species, *D. mexicana* and *D. rafaellii*. *Algae* 33, 231–241. doi: 10.4490/algae.2018.33.8.20
- Bustamante, D. E., Calderon, M. S., and Hughey, J. R. (2019). Conspecificity of the Peruvian *Corallina ferreyrae* with *C. caespitosa* (Corallinales, Rhodophyta) inferred from genomic analysis of the type specimen. *Mitochondrial DNA B Resour.* 4, 1285–1286. doi: 10.1080/23802359.2019.1591203
- Callaway, E. (2015). Lab staple agar hit by seaweed shortage. *Nature* 528, 171–172. doi: 10.1038/528171a
- Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol. Biol. Evol.* 17, 540–552. doi: 10.1093/oxfordjournals.molbev.a026334
- Costa, O. S., Attrilla, M. J., Pedrinib, A. G., and De-Paulab, J. C. (2002). Spatial and seasonal distribution of seaweeds on coral reefs from Southern Bahia, Brazil. *Bot. Mar.* 45, 346–355. doi: 10.1515/BOT.2002.035
- Darling, A. C. E., Mau, B., Blattner, F. R., and Perna, N. T. (2004). Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res.* 14, 1394–1403. doi: 10.1101/gr.2289704
- Díaz-Tapia, P., Maggs, C. A., Macaya, E. C., and Verbruggen, H. (2018). Widely distributed red algae often represent hidden introductions, complex of cryptic species or species with strong phylogeographic structure. *J. Phycol.* 54, 829–839. doi: 10.1111/jpy.12778
- Dierckxsens, N., Mardulyn, P., and Smits, G. (2016). NOVOPlasty: *de novo* assembly of organelle genomes from whole genome data. *Nucleic Acids Res.* 45:e18. doi: 10.1093/nar/gkw955
- Excoffier, L., and Lischer, H. E. L. (2010). Arlequin suite ver. 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Res.* 10, 564–567. doi: 10.1111/j.1755-0998.2010.02847.x
- Freshwater, D. W., Idol, J. N., Parham, S. L., Fernández-García, C., Léon, N., and Gabrielson, P. W. et al. (2017). Molecular assisted identification reveals hidden red algae diversity from the Burica Peninsula, Pacific Panama. *Diversity* 9:e19. doi: 10.3390/d9020019
- Freshwater, D. W., and Rueness, J. (1994). Phylogenetic relationships of some European *Gelidium* (Gelidiales, Rhodophyta) species, based on *rbcL* nucleotide sequence analysis. *Phycologia* 33, 187–194. doi: 10.2216/i0031-8884-33-3-187.1
- Gabrielson, P. W., Hughey, J. R., and Díaz-Pulido, G. (2018). Genomics reveals abundant speciation in the coral reef building alga *Porolithon onkodes* (Corallinales, Rhodophyta). *J. Phycol.* 54, 429–434. doi: 10.1111/jpy.12761
- Ganzon-Fortes, E. T. (1994). “Gelidiella,” in *Biology of Economic Algae*, ed. I. Akatsuka (The Hague: SPB Academic Publishing), 149–184.
- Gavio, B., and Fredericq, S. (2002). *Grateloupia turuturu* (Halymeniaceae, Rhodophyta) is the correct name of the nonnative species in the Atlantic known as *Grateloupia doryphora*. *Eur. J. Phycol.* 37, 349–359. doi: 10.1017/S0967026202003839
- Geraldino, P. J. L., Yang, E. C., and Boo, S. M. (2006). Morphology and molecular phylogeny of *Hypnea flexicaulis* (Gigartinales, Rhodophyta) from Korea. *Algae* 21, 417–423. doi: 10.4490/algae.2006.21.4.417
- Herzschuh, U., Birks, H. J. B., Laepple, T., Andreev, A., Melles, M., and Brigham-Grette, J. (2016). Glacial legacies on interglacial vegetation at the Pliocene-Pleistocene transition in NE Asia. *Nat. Commun.* 7:11967. doi: 10.1038/ncomms11967
- Hu, Z. M., Li, J. J., Sun, Z. M., Oak, J. H., Zhang, J., Fresia, P., et al. (2015). Phylogeographic structure and deep lineage diversification of the red alga *Chondrus ocellatus* Holmes in the northwest Pacific. *Mol. Ecol.* 24, 5020–5033. doi: 10.1111/eva.12455
- Hughey, J. R., and Boo, G. H. (2016). Genomic and phylogenetic analysis of *Ceramium cimbricum* (Ceramiales, Rhodophyta) from the Atlantic and Pacific oceans supports the naming of a new invasive Pacific entity *Ceramium sungminbooi* sp. nov. *Bot. Mar.* 59, 211–222. doi: 10.1515/bot-2016-0036
- Hughey, J. R., Gabrielson, P. W., Rohmer, L., Tortolani, J., Silva, M., Miller, K. A., et al. (2014). Minimally destructive sampling of type specimens of *Pyropia* (Bangiales, Rhodophyta) recovers complete plastid and mitochondrial genomes. *Sci. Rep.* 4:5113. doi: 10.1038/srep05113
- Huisman, J. M., Boo, G. H., and Boo, S. M. (2018). “Gelidiales,” in *Algae of Australia: Marine Benthic Algae of North-Western Australia*, 2. *Red Algae*, ed. J. M. Huisman (Canberra: ABRIS & CSIRO), 245–264.
- Kalyanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., and Jermini, L. S. (2017). ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Meth.* 14, 587–589. doi: 10.1038/nmeth.4285
- Kamiya, M., and West, J. A. (2014). Cryptic diversity in the euryhaline red alga *Caloglossa ogasawaraensis* (Delesseriaceae, Ceramiales). *Phycologia* 53, 374–382. doi: 10.2216/13-242.1
- Katoh, K., and Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780. doi: 10.1093/molbev/mst010
- Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* 33, 1870–1874. doi: 10.1093/molbev/msw054
- Lagesen, K., Hallin, P., Rødland, E. A., Staerfeldt, H.-H., Rognes, T., and Ussery, D. W. (2007). RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35, 3100–3108. doi: 10.1093/nar/gkm160
- Lanfear, R., Frandsen, P. B., Wright, A. M., Senfeld, T., and Calcott, B. (2016). PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Mol. Biol. Evol.* 34, 772–773. doi: 10.1093/molbev/msw260
- Laslett, D., and Canback, B. (2004). ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res.* 32, 11–16. doi: 10.1093/nar/gkh152
- Leblanc, C., Boyen, C., Richard, O., Bonnard, G., Grienemberger, J. M., and Kloareg, B. (1995). Complete sequence of the mitochondrial DNA of the rhodophyte *Chondrus crispus* (Gigartinales). Gene content and genome organization. *J. Mol. Biol.* 250, 484–495. doi: 10.1006/jmbi.1995.0392
- Leigh, J. W., and Bryant, D. (2015). PopART: full-feature software for haplotype network construction. *Methods Ecol. Evol.* 6, 1110–1116. doi: 10.1111/2041-210X.12410
- Leliaert, F., Payo, D. A., Gurgel, C. F. D., Schils, T., Draisma, S. G. A., Saunders, G. W., et al. (2018). Patterns and drivers of species diversity in the Indo-Pacific red seaweed *Portieria*. *J. Biogeogr.* 45, 2299–2313. doi: 10.1111/jbi.13410
- Li, D., Liu, C. M., Luo, R., Sadakane, K., and Lam, T. W. (2015). MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31, 1674–1676. doi: 10.1093/bioinformatics/btv033
- Librado, P., and Rozas, J. (2009). DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452. doi: 10.1093/bioinformatics/btp187
- Lim, P.-E., Tan, J., Phang, S. M., Nikmatullah, A., Hong, D. D., Sunarpi, H., et al. (2013). Genetic diversity of *Kappaphycus* Doty and *Eucheuma* J. Agardh

- (Solieriaceae, Rhodophyta) in Southeast Asia. *J. Appl. Phycol.* 26, 1253–1272. doi: 10.1007/s10811-013-0197-y
- Lin, S.-M., Fredericq, S., and Hommersand, M. H. (2001). Systematics of the Delesseriaceae (Ceramiales, Rhodophyta) based on large subunit rDNA and *rbcL* sequences, including the Phycodryoidae subfam. nov. *J. Phycol.* 37, 881–899. doi: 10.1046/j.1529-8817.2001.01012.x
- Lin, S.-M., and Freshwater, D. W. (2008). The red algal genus *Gelidiella* (Gelidiales, Rhodophyta) from Taiwan, including *Gelidiella fanii* sp. nov. *Phycologia* 47, 168–176. doi: 10.2216/07-30.1
- Lohse, M., Drechsel, O., Kahlau, S., and Bock, R. (2013). OrganellarGenomeDRAW – a suite of tools for generating physical maps of plastid and mitochondrial genomes and visualizing expression data sets. *Nucleic Acids Res.* 41, W575–W581. doi: 10.1093/nar/gkt289
- Lowe, T. M., and Chan, P. P. (2016). tRNAscan-SE On-line: search and contextual analysis of transfer RNA genes. *Nucleic Acids Res.* 44, W54–W57. doi: 10.1093/nar/gkw413
- Montecinos, A., Broitman, B. R., Faugeron, F., Haye, P. A., Tellier, F., and Guillemain, M. L. (2012). Species replacement along a linear coastal habitat: phylogeography and speciation in the red alga *Mazzaella laminarioides* along the south east Pacific. *BMC Evol. Biol.* 12:97. doi: 10.1186/1471-2148-12-97
- Nurk, S., Bankevich, A., Antipov, D., Gurevich, A. A., Korobeynikov, A., Lapidus, A., et al. (2013). Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J. Comput. Biol.* 20, 714–737. doi: 10.1089/cmb.2013.0084
- Pezzolesi, L., Peña, V., Le Gall, L., Gabrielson, P. W., Kaleb, S., Hughey, J. R., et al. (2019). Mediterranean *Lithophyllum strictiforme* (Corallinales, Rhodophyta) is a genetically diverse species complex: implications for species circumscription, biogeography and conservation of coralligenous habitats. *J. Phycol.* 55, 473–492. doi: 10.1111/jpy.12837
- Rioux, L., and Turgeon, S. L. (2015). “Seaweed carbohydrates,” in *Seaweed Sustainability: Food and Non-Food Applications*, eds B. J. Tiwari, and D. J. Troy (London: CRC Press), 141–192. doi: 10.1016/b978-0-12-418697-2.00007-6
- Ronquist, F., Teslenko, M., van Der Mark, P., Ayres, D., Darling, A., Höhna, S., et al. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–542. doi: 10.1093/sysbio/sys029
- Salomaki, E. D., and Lane, C. E. (2016). Red algal mitochondrial genomes are more complete than previously reported. *Genome Biol. Evol.* 9, 48–63. doi: 10.1093/gbe/evw267
- Santos, R., and Melo, R. A. (2018). Global shortage of technical agars: back to basics (resource management). *J. Appl. Phycol.* 30, 2463–2473. doi: 10.1007/s10811-018-1425-2
- Saunders, G. W. (2005). Applying DNA barcoding to red macroalgae: a preliminary appraisal holds promise for future applications. *Philos. Trans. R. Soc. B* 360, 1879–1888. doi: 10.1098/rstb.2005.1719
- Saunders, G. W., and Moore, T. E. (2013). Refinements for the amplification and sequencing of red algal DNA barcode and RedToL phylogenetic markers: a summary of current primers, profiles and strategies. *Algae* 28, 31–43. doi: 10.4490/algae.2013.28.1.031
- Sherwood, A. R., Sauvage, T., Kurihara, A., Conklin, K. Y., and Presting, G. G. (2010). The Hawaiian Rhodophyta biodiversity survey (2006–2010): a summary of principal findings. *BMC Plant Biol.* 10:258. doi: 10.1186/1471-2229-10-258
- Song, S. L., Yong, H. S., Lim, P. E., and Phang, S. M. (2017). Complete mitochondrial genome of *Gracilaria changii* (Rhodophyta: Gracilariaceae). *J. Appl. Phycol.* 29, 2129–2134. doi: 10.1007/s10811-017-1100-z
- Suzuki, M., Segawa, T., Mori, H., Akiyoshi, A., Ootsuki, R., Kurihara, A., et al. (2016). Next-generation sequencing of an 88-year-old specimen of the poorly known species *Liagora japonica* (Nemaliales, Rhodophyta) supports the recognition of *Otohimella* gen. nov. *PLoS One* 11:e0158944. doi: 10.1371/journal.pone.0158944
- Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., and Minh, B. Q. (2016). W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 44, W232–W235. doi: 10.1093/nar/gkw256
- Wiriyadamrikul, J., Park, J. K., Khan, W., and Boo, S. M. (2010). Additional records of *Gelidiella fanii* (Gelidiales, Rhodophyta) from the southeast Pacific based on morphology, *rbcL* and *cox1* analyses. *Bot. Mar.* 53, 343–350. doi: 10.1515/BOT.2010.037
- Won, B. Y., Cho, T. O., and Fredericq, S. (2009). Morphological and molecular characterization of species of the genus *Centroceras* (Ceramiales, Rhodophyta), including two new species. *J. Phycol.* 45, 227–250. doi: 10.1111/j.1529-8817.2008.00620.x
- Yang, E. C., Kim, K. M., Kim, S. Y., Lee, J. M., Boo, G. H., Lee, J.-H., et al. (2015). Highly conserved mitochondrial genomes among multicellular red algae of the Florideophyceae. *Genome Biol. Evol.* 7, 2394–2406. doi: 10.1093/gbe/evv147
- Yoon, K. J., Kim, K. M., Boo, G. H., Miller, K. A., and Boo, S. M. (2014). Mitochondrial *cox1* and *cob* sequence diversities in *Gelidium vagum* (Gelidiales, Rhodophyta) in Korea. *Algae* 29, 15–25. doi: 10.4490/algae.2014.29.1.015
- Zuccarello, G. C., Burger, G., West, J. A., and King, R. J. (1999). A mitochondrial marker for red algal intraspecific relationships. *Mol. Ecol.* 8, 1443–1447. doi: 10.1046/j.1365-294x.1999.00710.x

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Boo, Zubia, Hughey, Sherwood, Fujii, Boo and Miller. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.