



Available online at www.sciencedirect.com

ScienceDirect

Stochastic Processes and their Applications 130 (2020) 3394–3407

stochastic
processes
and their
applications

www.elsevier.com/locate/spa

When is it best to follow the leader?

Philip A. Ernst^{a,*}, L.C.G. Rogers^b, Quan Zhou^c

^a Department of Statistics, Rice University, Houston, Texas 77005, USA

^b Statistical Laboratory, Center for Mathematical Sciences, Cambridge CB5 0WB, UK

^c Department of Statistics, Texas A&M University, College Station, Texas 77843, USA

Received 30 March 2018; received in revised form 7 September 2019; accepted 27 September 2019

Available online 5 October 2019

Abstract

An object is hidden in one of N boxes. Initially, the probability that it is in box i is $\pi_i(0)$. You then search in continuous time, observing box J_t at time t , and receiving a signal as you observe: if the box you are observing does not contain the object, your signal is a Brownian motion, but if it does contain the object your signal is a Brownian motion with positive drift μ . It is straightforward to derive the evolution of the posterior distribution $\pi(t)$ for the location of the object. If T denotes the first time that one of the $\pi_j(t)$ reaches a desired threshold $1 - \varepsilon$, then the goal is to find a search policy $(J_t)_{t \geq 0}$ which minimizes the mean of T . This problem was studied by Posner and Rumsey (1966) and by Zigangirov (1966), who derive an expression for the mean time of a conjectured optimal policy, which we call *follow the leader* (FTL); at all times, observe the box with the highest posterior probability. Posner and Rumsey assert without proof that this is optimal, and Zigangirov offers a proof that if the prior distribution is uniform then FTL is optimal. In this paper, we show that if the prior is not uniform, then FTL is *not* always optimal; for uniform prior, the question remains open.

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

MSC: primary 60H30; secondary 60G35

Keywords: Follow the leader; Optimal scanning; Quickest search; Tanaka's stochastic differential equation; First exit time

1. Introduction

This paper studies a classical search problem first considered independently by Posner and Rumsey [12] and Zigangirov [17]. An object is hidden in one of N boxes; we denote by j^*

* Corresponding author.

E-mail addresses: philip.ernst@rice.edu (P.A. Ernst), chris@statslab.cam.ac.uk (L.C.G. Rogers), quan@stat.tamu.edu (Q. Zhou).

the index of the true box. Initially,

$$P(j^* = i) = \pi_i(0).$$

We then observe in continuous time, choosing to search box J_t at time t . We see a signal process Y whose dynamics are

$$dY(t) = dW(t) + \mu I_{\{J_t=j^*\}} dt, \quad (1)$$

where $\mu > 0$ is a known constant, and W is a Brownian motion. It is straightforward to derive the evolution of $\pi(t)$, the posterior distribution at time t . The objective proposed by Posner and Rumsey [12] is to choose $(J_t)_{t \geq 0}$ to minimize ET , where

$$T = \inf\{t : \max_j \pi_j(t) \geq 1 - \varepsilon\}, \quad (2)$$

where $\varepsilon \in (\frac{1}{2}, 1)$ is some desired error bound.

To the best of our knowledge, the solution to this specific problem was studied by three papers: Posner and Rumsey [12], Zigangirov [17] and Klimko and Yackel [8]. The earliest of these references [12] motivates the problem in the context of search for a space probe by radar. In this framework, the sky is assumed divided into n equal regions (equivalently, “boxes”). One box emits a signal of Brownian motion with positive drift μ and the remaining $n - 1$ boxes emit a signal of Brownian motion. Posner and Rumsey [12] asserted, without proof, that the optimal strategy is to always search the box with the largest posterior probability. We call this policy the *follow the leader* (FTL) policy. They formulated the FTL strategy as the limit of a sequence of discrete-time approximations, which was later shown by Klimko and Yackel [7] not to be tight. Zigangirov [17] considered only the case of uniform prior distribution, that is, $\pi_i(0) = 1/N$ for $i = 1, \dots, N$, and offered a proof for the optimality of FTL. However, this proof lacks clarity on a number of points, and we were not able to verify the arguments given. Klimko and Yackel [8] provided a proof for the optimality of FTL for arbitrary prior distribution, but, as we will explain later, their proof is in error. The main result of our paper is to give counterexamples that clearly show FTL is not optimal for some specific values of $(\pi_1(0), \dots, \pi_N(0))$. An additional contribution is the characterization of the solution to a class of stochastic differential equations, which plays a key role in our calculations, and can be considered to be generalizations of Tanaka’s SDE.

1.1. Literature review

Optimal scanning problems apparently date back to Shiryaev [14]. Our specific problem of interest was considered in the works of Posner and Rumsey [12], Zigangirov [17] and Klimko and Yackel [8]. We now briefly review other variants of the problem that are closely related to this work.

Liptser and Shiryaev [10] considered a setup with two boxes ($N = 2$) and allow for the possibility that the object may not be in either box. The task is to determine if the missing object is in one of the two boxes. Dragalin [3] considered general stochastic processes other than Brownian motion and proposed a scanning rule based on the sequential probability ratio test of Wald [16].

Another class of problems similar to optimal scanning is problems of “quickest search”. These problems are often formulated under the setting $N \rightarrow \infty$ with an unknown number of boxes containing the hidden objects. We refer readers to Lai et al. [9] for a discrete-time solution and Bayraktar and Kravitz [1] for a continuous-time solution to these quickest search

problems. We also note that optimal scanning problems and quickest search problems are known collectively as “screening problems”. We refer readers to the references given in Heydari et al. [5] for other variants of screening problems. More generally, such problems can be viewed as sequential decision problems. References on this topic include Dvoretzky et al. [4], Shiryaev [15, Ch. 4], and Peskir and Shiryaev [11, Ch. VI].

2. The evolution of the posterior

As we noted at (1), the signal process Y evolves as¹

$$dY_t = dW_t + \mu I_{\{J_t=j^*\}} dt.$$

If $(\mathcal{Y}_t)_{t \geq 0}$ is the filtration of the observation process, and we choose to search box J_t at time t , then the posterior likelihood (relative to Wiener measure) that the true box is j , given \mathcal{Y}_t , is

$$z_j(t) = \pi_j(0) \exp \left(\int_0^t \mu I_{\{J_s=j\}} dY_s - \frac{1}{2} \mu^2 \int_0^t I_{\{J_s=j\}} ds \right). \quad (3)$$

The posterior probabilities are obtained by normalizing the z_j :

$$\pi_j(t) = z_j(t)/\bar{z}(t), \quad (4)$$

where of course $\bar{z}(t) = \sum_{j=1}^N z_j(t)$. Now

$$dz_j(t) = z_j(t) \mu I_{\{J_t=j\}} dY_t, \quad (5)$$

so if we write $X_j(t) = \mu^{-1} \log(z_j(t))$ then we have

$$dX_j(t) = I_{\{J_t=j\}} (dY_t - \frac{1}{2} \mu dt). \quad (6)$$

The evolution (1) of Y is expressed in the filtration of W , but familiar results of filtering theory (see Kallianpur et al. [6]) establish that we can rewrite the evolution in the filtration \mathcal{Y} as

$$dY_t = d\hat{W}_t + \mu \pi_{J_t}(t) dt \quad (7)$$

where \hat{W} is the innovation process, a \mathcal{Y} -Brownian motion, and $\mu \pi_{J_t}(t)$ is the \mathcal{Y} -optional projection of the drift $\mu I_{\{J_t=j^*\}}$ of (1).

Formulating the dynamics slightly more generally, as

$$dz_j(t) = z_j(t) \theta_j(t) dY_t \quad (8)$$

where θ is a bounded previsible N -vector process, we can consider the evolution of $\pi(t)$ defined in terms of $z(t)$ by (4). Some routine calculations with Itô’s formula give us

$$d\pi_j(t) = \pi_j(t) \{ \theta_j(t) - \theta(t) \cdot \pi(t) \} \{ dY_t - \theta(t) \cdot \pi(t) dt \}, \quad (9)$$

where $\theta(t) \cdot \pi(t) = \sum_{i=1}^N \theta_i(t) \pi_i(t)$. In the case of special interest to us, where $\theta_j(t) = \mu I_{\{J_t=j\}}$, the representation (7) combines with (9) to show that

$$d\pi_j(t) = \pi_j(t) \{ \theta_j(t) - \theta(t) \cdot \pi(t) \} d\hat{W}_t. \quad (10)$$

In particular, $\pi(t)$ is a \mathcal{Y} -local martingale; but we know this already, because $\pi_j(t) = P(j^* = j | \mathcal{Y}_t)$, which is even a martingale.

¹ The notation X_t and $X(t)$ will be used interchangeably according to context.

3. The FTL policy

For the FTL policy, the dynamics (8) has the special form

$$\theta_j(t) = \mu I_j(X(t)), \quad (11)$$

where we define for $x \in \mathbb{R}^N$

$$\begin{aligned} I_j(x) &= 1 && \text{if } x_j = \max\{x_i : i = 1, \dots, j\} > \max\{x_i : i > j\} \\ &= 0 && \text{else.} \end{aligned} \quad (12)$$

Thus $I_j(x)$ is the index where the N -vector x is maximal, taking care to avoid ambiguities when there are ties, and to ensure that $\sum_{j=1}^N I_j(x) = 1$.

In these terms, the evolution of $X_j(t)$ can be expressed as²

$$\begin{aligned} dX_j(t) &= I_j(X(t))(dY_t - \frac{1}{2}\mu dt) \\ &= I_j(X(t))(d\hat{W}_t + \mu(\pi_j(t) - \frac{1}{2})dt) \end{aligned} \quad (13)$$

$$\equiv I_j(X(t))dZ_t, \quad (14)$$

say. Since

$$\pi_j(t) = \frac{e^{\mu X_j(t)}}{\sum_{i=1}^N e^{\mu X_i(t)}} \quad (15)$$

is a function of $X(t)$, the SDE (13) is an autonomous SDE, but the coefficients are not Lipschitz, or even continuous, so the sense in which the SDE has a solution needs to be clarified.

We shall address this by firstly studying the SDE (13) without the drift term:

$$dX_j(t) = I_j(X(t))dW_t, \quad (16)$$

where W is a standard Brownian motion. To appreciate the issues involved, let us first consider the case $N = 2$, when the SDE is

$$dX_1(t) = I_{\{X_1(t) > X_2(t)\}}dW_t = I_{\{X_1(t) - X_2(t) > 0\}}dW_t \quad (17)$$

$$dX_2(t) = I_{\{X_2(t) \geq X_1(t)\}}dW_t = I_{\{X_1(t) - X_2(t) \leq 0\}}dW_t. \quad (18)$$

So if $y_t \equiv X_1(t) - X_2(t)$ we have the celebrated Tanaka SDE

$$dy_t = \text{sgn}(y_t)dW_t \quad (19)$$

for y , where the definition of sgn is the correct one for the definition of semimartingale local time — see Theorem IV.43.3 of Rogers and Williams [13]. There is no strong solution to this SDE, but there is a weak solution, represented by taking Y to be a Brownian motion started at $y_0 = X_1(0) - X_2(0)$, which we may as well suppose is positive, and then defining

$$dW_t = \text{sgn}(y_t)dy_t = d|y_t| - dL_t \quad (20)$$

where L is the local time of y at zero. Then we have

$$dX_1(t) = I_{\{y_t > 0\}}dy_t = d(y_t^+) - \frac{1}{2}dL_t. \quad (21)$$

² In Posner and Rumsey [12], the signal has a constant volatility σ , as well as the drift μ . We could always scale the signal to turn the volatility to 1, and indeed we could also replace μ by any desired positive value; this is equivalent to a constant rescaling of time, which will not affect optimality of a search policy.

Thus

$$X_1(t) = X_1(0) + y_t^+ - \frac{1}{2} L_t \quad (22)$$

$$X_2(t) = X_2(0) + y_t^- - \frac{1}{2} L_t. \quad (23)$$

In view of the above, we realize:

- We cannot hope for (16) to have a strong solution;
- We might obtain uniqueness in law for all initial values;
- If all the X_j start from 0, the sum $\sum_{j=1}^N X_j(t)$ is a Brownian motion when $N = 2$;
- If all the X_j start from 0, the running minimum $\underline{X}_j(t) \equiv \inf\{X_j(s) : s \leq t\}$ is the same for all j when $N = 2$.

For general N , we have the analogous conclusions.

Theorem 1. *For all starting values $X(0)$, the SDE (16) has a weak solution which is unique in law. If $X(0) = 0$, then*

(1) $\sum_{j=1}^N X_j(t) \equiv W_t$ is a Brownian motion;

(2) the running minimum processes coincide:

$$\underline{X}_j(t) \equiv \inf\{X_j(s) : s \leq t\} = N^{-1} \underline{W}_t \equiv \underline{X}(t) \quad (24)$$

(3) for all $t \geq 0$,

$$X_j(t) = \underline{X}(t) \quad \text{for all but at most one index } j; \quad (25)$$

(4) $X_j(t) - \underline{X}(t) = W_t - \underline{W}_t$ if $X_j(t) > \underline{X}(t)$;

(5) The process

$$X_k(t) - (N-1)^{-1} \sum_{1 \leq j \neq k \leq N} X_j(t) = \frac{NI_k(X(t)) - 1}{N-1} (W_t - \underline{W}_t) \quad (26)$$

is a martingale.

Theorem 1 is proved in the appendix. It deals immediately with the question of existence and uniqueness of solutions of (13), because any weak solution to (16) can be transformed by change of measure into a weak solution to (13), and vice versa.

4. The value of FTL

Our aim in this section is to discover the value function $V(x_1, \dots, x_N)$ of the FTL policy, where x_j denotes the initial value of the process $X_j(t)$. Formally,

$$V(x_1, \dots, x_N) = E(T \mid X_1(0) = x_1, \dots, X_N(0) = x_N, \text{policy} = \text{FTL}),$$

where T is as in (2). It does not appear possible to express this in closed form, but we can find a recursive algorithm for computing the value numerically. It is obvious that V is a symmetric function of its arguments, so we will make the convention in what follows that the arguments of V have been arranged in decreasing order:

$$x_1 \geq x_2 \geq \dots \geq x_N. \quad (27)$$

It is also obvious that the value will not be changed if we add the same constant to all the arguments.

Now suppose that all the inequalities in (27) are strict, and we apply the FTL rule. What happens is that initially we observe the most likely box, box 1, up until the time τ_2 when X_1 first falls³ to x_2 . At that time, X_2 begins to move, and in accordance with Theorem 1 we find that

$$dX_1(t) = dX_2(t) = \frac{1}{2} dZ_t, \quad \max\{X_1(t), X_2(t)\} - X_1(t) = Z_t - Z_t. \quad (28)$$

This continues until the first time τ_3 that one of X_1, X_2 (and hence both) falls to x_3 . Thereafter we observe the boxes 1, 2, 3 with

$$d\underline{X}_1(t) = d\underline{X}_2(t) = d\underline{X}_3(t) = \frac{1}{3} dZ_t, \quad \max\{X_1(t), X_2(t), X_3(t)\} - \underline{X}_1(t) = Z_t - Z_t.$$

This continues sequentially, with the $X_j(t)$ starting to move one after another, until all the $X_j(t)$ achieve a common minimum x_N at time τ_N , or, of course, the termination criterion is achieved. So we see that calculation of the value comes down to solving a sequence of first-exit problems, which we formalize in the following result.

Theorem 2. *Suppose that $x_1 > x_2 > \dots > x_N$, and let $V_n(x, x_{n+1}, \dots, x_N)$ denote the value if we start with $x_1 = x_2 = \dots = x_n = x > x_{n+1} > \dots > x_N$. Then the values $V_n(x_n, x_{n+1}, \dots, x_N)$ can be calculated recursively as*

$$V_n(x_n, x_{n+1}, \dots, x_N) = A_n(x_n) + \frac{B_n(x_n)}{1 + K_n(x_n)}, \quad (29)$$

where

$$K_n(y) \equiv n - 1 + b_n e^{-\mu y}, \quad b_n \equiv e^{\mu x_{n+1}} + \dots + e^{\mu x_N}, \quad (30)$$

where B_n is the solution to the ODE ($p_0 \equiv 1 - \varepsilon$)

$$(1 - p_0(1 + K_n(y)))B'(y) = (n - 1)\mu B(y) + \frac{2n(K_n(y) - 1)}{\mu} - \frac{2(1 - 2p_0)}{\mu K_n(y)}(K_n(y) - n + 1)(K_n(y) + 1), \quad (31)$$

with boundary condition

$$B(x_{n+1}) = \frac{(1 + K_n(x_{n+1}))\{\mu V_{n+1}(x_{n+1}, \dots, x_N) - 2q_n(x_{n+1})(1 - 2\varepsilon)\}}{\mu\{1 - p_0(1 + K_n(x_{n+1}))\}}, \quad (32)$$

and where

$$0 = A_n(y) + (1 - \varepsilon)B_n(y) - 2q_n(y)(1 - 2\varepsilon)/\mu, \quad (33)$$

$$q_n(y) \equiv \mu^{-1} \log\{(1 - \varepsilon)(n - 1 + b_n e^{-\mu y})/\varepsilon\}. \quad (34)$$

The recursion begins at $n = N$ with the Posner-Rumsey value:

$$V_N(x_N) = M_{PR} \equiv \frac{2}{\mu^2} \left[\frac{N-2}{N-1} (Np_0 - 1) + (2p_0 - 1) \log\left(\frac{1-\varepsilon}{\varepsilon/(N-1)}\right) \right]. \quad (35)$$

Proof. Holding x_{n+1}, \dots, x_N fixed, we let $f(s, y)$ denote the value when the running minimum of X_1, \dots, X_n is $y \in [x_{n+1}, x_n]$, and the unique leading particle is at $y + s \geq y$. The success criterion will be satisfied if

$$\frac{e^{\mu(y+s)}}{e^{\mu(y+s)} + (n-1)e^{\mu y} + b_n} = 1 - \varepsilon, \quad (36)$$

³ Of course, the termination condition may have been achieved before that time.

where b_n is given by (30). Hence successful termination occurs when

$$s = \mu^{-1} \log\{(1 - \varepsilon)(n - 1 + b_n e^{-\mu y})/\varepsilon\} \equiv q_n(y). \quad (37)$$

Now up until τ_{n+1} , in terms of Z we have that

$$f(Z_t - Z_t, n^{-1}Z_t) + t \quad \text{is a martingale.}$$

The probability that we are viewing the correct box is

$$\begin{aligned} p(s, y) &= \frac{e^{\mu(y+s)}}{e^{\mu(y+s)} + (n-1)e^{\mu y} + b_n} \\ &= \frac{e^{\mu s}}{e^{\mu s} + (n-1) + b_n e^{-\mu y}} \\ &\equiv \frac{e^{\mu s}}{e^{\mu s} + K_n(y)}, \end{aligned} \quad (38)$$

where K_n is defined at (30). Using Itô's formula, we arrive at the equations

$$0 = \frac{1}{2} f_{ss} + (p(s, y) - \frac{1}{2})\mu f_s + 1, \quad (39)$$

$$0 = -nf_s + f_y \quad (s = 0) \quad (40)$$

with boundary conditions

$$f(q_n(y), y) = 0 \quad \forall y \in [x_{n+1}, x_n], \quad (41)$$

$$f(0, x_{n+1}) = V_{n+1}(x_{n+1}, \dots, x_N). \quad (42)$$

We see that (39) is a second-order linear ODE in the variable s , whose general solution can be shown by routine calculations to be

$$f(s, y) = \tilde{A}(y) + \frac{\tilde{B}(y)}{e^{\mu s} + K_n(y)} + \frac{2s}{\mu} (1 - 2p(s, y)) \quad (43)$$

for some functions \tilde{A} , \tilde{B} . Equivalently, we may express the solution as

$$f(s, y) = A(y) + B(y)p(s, y) + \frac{2s}{\mu} (1 - 2p(s, y)) \quad (44)$$

for some functions A , B . The boundary condition at reflection (40) leads to the equation

$$\frac{(n-1)\mu B(y)}{1 + K_n(y)} + \frac{2n(K_n(y) - 1)}{\mu(K_n(y) + 1)} = A'(y) + \frac{B'(y)}{1 + K_n(y)} \quad (45)$$

and the boundary condition (41) gives us

$$0 = A(y) + (1 - \varepsilon)B(y) - 2q_n(y)(1 - 2\varepsilon)/\mu. \quad (46)$$

This allows us to express $A(y)$ as a function of y and $B(y)$, reducing the ODE (45) to a first-order linear ODE for B . From (46) we find that

$$A'(y) = -p_0 B'(y) + \frac{2(1 - 2p_0)}{\mu K_n(y)} (K_n(y) - n + 1). \quad (47)$$

Returning this to (45) leads to the first-order ODE (31) for B in $y \geq x_{n+1}$. The boundary condition (42) together with (46) becomes the boundary condition (32).

At the final stage, V_N is a function of just one variable, and $p(s, y)$ is independent of y ; the form (44) collapses to

$$f(s) = A_N + B_N p(s, 0) + \frac{2s}{\mu} (1 - 2p(s, 0)) \quad (48)$$

with the boundary conditions

$$f(q_N(0)) = 0, \quad f'(0) = 0. \quad (49)$$

Solving this for A_N , B_N leads to the Posner-Rumsey solution (35). Finally, the expression (29) is obtained by letting $s = 0$. \square

5. Counterexamples for the optimality of FTL

In this section, we introduce an alternative strategy that can beat FTL in some circumstances. We make use of a classical result for the exit time of a Brownian motion at two boundaries. It was apparently first derived by Darling and Siegert [2].

Lemma 1 (Darling and Siegert [2]). *Let $W(t)$ be a Brownian motion with drift λt and variance $\sigma^2 t$, and started at x . Let $\rho = \lambda/\sigma^2$. Consider the boundaries a and b such that $a > x > b$. Then exit at one of the boundaries occurs with probability 1, and the probability of exit at a is given by*

$$P(x, a, b, \lambda, \sigma^2) = \frac{e^{-2\rho b} - e^{-2\rho x}}{e^{-2\rho b} - e^{-2\rho a}}.$$

Conditional on exiting at a , the expected time is given by

$$F_a(x, a, b, \lambda, \sigma^2) = \frac{1}{\lambda} \left[(a - x) + \frac{2(a - b)e^{-2\rho a}}{e^{-2\rho b} - e^{-2\rho a}} - \frac{2(x - b)e^{-2\rho x}}{e^{-2\rho b} - e^{-2\rho x}} \right].$$

Conditional on exiting at b , the expected time is given by

$$F_b(x, a, b, \lambda, \sigma^2) = \frac{1}{\lambda} \left[(x - b) + \frac{2(a - b)e^{-2\rho a}}{e^{-2\rho b} - e^{-2\rho a}} - \frac{2(a - x)e^{-2\rho a}}{e^{-2\rho b} - e^{-2\rho a}} \right].$$

Remark. Observe that $F_a(x, a, b, \lambda, \sigma^2) = F_a(x, a, b, -\lambda, \sigma^2)$. The same equality holds true for F_b . This is useful since, by (6), the drift of $X_{J_t}(t)$ is either $\mu/2$ or $-\mu/2$.

Now we introduce an *alternative strategy*, which we call “Strategy B”. For simplicity, let us consider three boxes with initial values $x_1 > x_2 > x_3$ (and thus prior probabilities $\pi_1(0) > \pi_2(0) > \pi_3(0)$). We shall suppose that

$$\pi_1(0) \equiv \frac{e^{\mu x_1}}{e^{\mu x_1} + e^{\mu x_2} + e^{\mu x_3}} < 1 - \varepsilon < \frac{e^{\mu x_1}}{e^{\mu x_1} + 2e^{\mu x_3}}, \quad (50)$$

so that there exists a unique $a \in (x_3, x_2)$ such that

$$\frac{e^{\mu x_1}}{e^{\mu x_1} + e^{\mu a} + e^{\mu x_3}} = 1 - \varepsilon. \quad (51)$$

Strategy B observes $X_2(t)$ until it reaches a or x_1 . If X_2 reaches a before x_1 , then the objective is achieved, in view of (51). Because of Lemma 1, we know the mean of this stopping time, and the probability that exit happens at x_1 . Otherwise, if X_2 reaches x_1 before a , we now continue with the FTL policy, whose mean remaining time to finish will be $V(x_1, x_1, x_3)$, which can be calculated according to Theorem 2.

Table 1

Counterexamples for the optimality of FTL with $N = 3$, $\mu = 1$ and $x_1 > x_2 > x_3 = 0$. $E_A(T)$ denotes the expected search time of the FTL strategy, which can be computed using [Theorem 2](#). $E_B(T)$ denotes the expected search time of Strategy B, which can be computed as explained above.

ε	x_1	x_2	$E_A(T) \times 10^2$	$E_B(T) \times 10^2$
0.4	2	1.4	3.633	3.464
0.3	2.7	1.7	3.053	2.936
0.2	4.05	2.6	1.832	1.797
0.1	6.2	4.0	3.749	3.738
0.05	10.3	7.4	10.6482	10.6476

By fixing $\mu = 1$, $x_3 = 0$ and searching over the corresponding domain of (x_1, x_2) , we obtain a few counterexamples for different values of ε . These are presented in [Table 1](#).

6. Further discussion

6.1. Discussion of the work of Klimko and Yackel [8]

Klimko and Yackel [8] gave a proof for the optimality of FTL for arbitrary prior distribution; however, according to our counterexamples, this conclusion cannot be correct. Here we explain why their proof is incorrect.

Consider that we start the search by choosing one box to observe until $\pi_1(t)$ reaches either $\pi_1(0) + \alpha$ or $\pi_1(0) - \beta$, assuming $\pi_1(0) > \dots > \pi_N(0)$. Denote this stopping time by τ and assume that there is no switch of the observed box before τ . Thus for $t \in [0, \tau]$, there are N possible search rules. In both the proofs of Lemma 3.4 and Theorem 3.5 in Klimko and Yackel [8], the authors assume that the posterior distribution at τ is *independent of the search rule*, which is incorrect. For example, their proof for case (i) of Theorem 3.5 relies on this incorrect assumption (the authors write that “furthermore, the posterior distribution at exit time are also independent of the rule used”.)

For a concrete example, consider $N = 3$ and $\pi(0) = (0.5, 0.25, 0.25)$. We wish to use this example to further explain why the reasoning of Klimko and Yackel [8] is incorrect. Let τ_1 be the exit time of π_1 at either $1 - \varepsilon$ or 0.4 , and assume no switch of observed box before τ_1 . Lemma 3.1 of Klimko and Yackel [8] correctly states that in order to minimize $E(\tau_1)$ we should choose to observe box 1. However, if box 1 is observed and π_1 exits at 0.4 , at τ_1 we have the posterior probability $\pi(\tau_1) = (0.4, 0.3, 0.3) \equiv \pi_A$; if box 2 is observed and π_1 exits at 0.4 , we have $\pi(\tau_1) = (0.4, 0.4, 0.2) \equiv \pi_B$. The inductive argument used in the proof of Lemma 3.4 of Klimko and Yackel [8] now fails because it is no longer clear whether π_A or π_B would lead to a smaller expected search time after τ_1 . In fact, according to our numerics, π_B would give a smaller expected search time if FTL is applied.

6.2. Open problems

Our work gives rise to several open problems. First, what is the optimal strategy for this optimal scanning problem for any prior distribution? For decades, it has been (incorrectly) assumed that FTL is optimal. Indeed, as we have shown, FTL is sub-optimal at least for some values of $(\pi_1(0), \dots, \pi_N(0))$.

Another open problem concerns whether FTL is optimal for the case of uniform prior distribution. There is already a proof given by Zigangirov [17], but as already mentioned, we

found the argument presented to lack clarity in various places, so we cannot be confident that the result is established.

So the answer to the question in the title is, ‘We don’t know, but we know that it is not *always* best to follow the leader!’

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

Philip Ernst gratefully acknowledges support from United States Army Research Office (ARO) Grant ARO-YIP-71636-MA, National Science Foundation (NSF), USA Grant DMS-1811936, and United States Office of Naval Research (ONR) Grant N00014-18-1-2192.

Appendix. The SDE $dX = I(X) dW$

The aim in this appendix is to prove [Theorem 1](#). Until further notice, we shall focus on the case $X_j(0) = 0$ for all j . To begin with, we present two results about any solution of the SDE [\(16\)](#)

$$dX_j(t) = I_j(X(t)) dW_t, \quad X_j(0) = 0 \quad \forall j. \quad (\text{A.1})$$

Proposition 1. *For all i , $\underline{X}_i(t) = \underline{X}_1(t) \equiv \underline{X}(t)$.*

Proof. For any $a < 0$, we let $H_j(a) = \inf\{t : X_j(t) \leq a\}$. If it were the case that for some $i \neq j$ we have $X_i(H_j(a)) > X_j(H_j(a)) = a$, then there has to be some time interval (s, u) containing $H_j(a)$ throughout which $X_i(t) > a$. This means that throughout (s, u) the process X_j is not the leader, so it does not move. This contradicts the definition of $H_j(a)$. Therefore $X_i(H_j(a)) \leq a$ for all $i \neq j$, and hence $H_i(a) \leq H_j(a)$ for all $i \neq j$. Since we can interchange the rôles of i and j , it must be that $H_i(a) = H_j(a)$ for all $i \neq j$, and the result follows. \square

Proposition 2. *For all $t \geq 0$, $\underline{W}_t = N\underline{X}(t)$.*

Proof. Observe that $\sum_{j=1}^N X_j(t) = W_t$, since $\sum_{j=1}^N I_j(x) \equiv 1$, which proves statement (1) of [Theorem 1](#). With the notation of the proof of [Proposition 1](#), we have that for any $a < 0$

$$X_j(H(a)) = \underline{X}(H(a)) = a, \quad \forall j,$$

where $H(a)$ denotes the common value $H_j(a)$. Therefore $W(H(a)) = Na$. Further, for any $t < H(a)$ we have $X_j(t) > a$, and thus $W_t > Na$. So it must be that $H(a) = \inf\{t : W_t \leq Na\}$, and the result follows. \square

Remark. Statement (2) of [Theorem 1](#) is now proved.

Proposition 3. *On a suitable probability space, a solution to [\(A.1\)](#) may be constructed.*

Proof. Consider the Itô excursion point process description of Brownian motion, using notation and terminology from Rogers and Williams [13, Ch. VI]. According to Proposition VI.51.2, the rate of excursions which get at least $x > 0$ away from zero is

$$n(\{f \in U : \sup_t |f(t)| > x\}) = 1/x, \quad (\text{A.2})$$

and the full excursion law is specified in various ways. We write U_+ for the space of non-negative excursions:

$$U_+ = \{f : \mathbb{R}^+ \rightarrow \mathbb{R}^+ | f^{-1}((0, \infty)) = (0, \zeta) \text{ for some } \zeta > 0\}. \quad (\text{A.3})$$

We let n_+ be the law of excursions away from zero of $|W|$, so that

$$n_+(\{f \in U : \sup_t f(t) > x\}) = 1/x. \quad (\text{A.4})$$

Now let Π be a Poisson random measure on $(0, \infty) \times U_+ \times \{1, \dots, k\}$ with mean measure

$$M_N \equiv N^{-1} dt \times n_+(df) \times \mu_N, \quad (\text{A.5})$$

where $\mu_N(\{j\}) = 1$ for each $j = 1, \dots, N$. Now we define the clock

$$T(\ell) = \iint_{(0, \ell] \times U_+ \times S_N} \zeta(f) \Pi(ds, df, dj), \quad (\text{A.6})$$

with inverse

$$L_t = \inf\{\ell : T(\ell) > t\}. \quad (\text{A.7})$$

The local time L remains constant through all excursion intervals; let

$$g_t \equiv \sup\{s : L_s < L_t\} \quad (\text{A.8})$$

denote the left end of the excursion including time t . Finally, we may define

$$X_j(t) = -N^{-1} L_t + f(t - g_t) \quad \text{if } (L_t, f, j) \text{ is a point of } \Pi; \quad (\text{A.9})$$

$$= -N^{-1} L_t \quad \text{else. } \square \quad (\text{A.10})$$

Proposition 4. *Uniqueness in law holds for (A.1).*

Proof. Firstly, let us deal with the case $N = 2$. We saw in Section 3 that any solution to the SDE (16) can be represented in terms of the Brownian motion Y defined at (19) by Eqs. (22), (23), so the law of the solution (X_1, X_2) is uniquely determined.

The case $N \geq 3$ requires a little more subtlety. Take any $j \neq k$ in $\{1, \dots, N\}$, and define

$$A_t = \int_0^t (I_j(X(s)) + I_k(X(s))) ds, \quad (\text{A.11})$$

$$\tau_t = \inf\{u : A_u > t\}. \quad (\text{A.12})$$

Notice that

$$(1 - I_j(X(t)) - I_k(X(t))) dX_j(t) = I_j(X(t))(1 - I_j(X(t)) - I_k(X(t))) dW_t = 0 \quad (\text{A.13})$$

so X_j, X_k do not change when the clock A is not growing. Therefore if we define

$$\tilde{X}_j(t) = X_j(\tau_t), \quad \tilde{X}_k(t) = X_k(\tau_t), \quad (\text{A.14})$$

we have

$$\inf\{\tilde{X}_j(s) : s \leq t\} = \underline{X}_j(\tau_t) = \underline{X}(\tau_t). \quad (\text{A.15})$$

With a slight overloading of notation, we have

$$\begin{aligned} d\tilde{X}_j(t) &= I_j(\tilde{X}(t)) d\tilde{W}_t, \\ d\tilde{X}_k(t) &= I_k(\tilde{X}(t)) d\tilde{W}_t, \end{aligned}$$

so the pair $(\tilde{X}_j, \tilde{X}_k)$ is a solution of the SDE for the case $d = 2$. But we know that uniqueness in law holds for this situation, so in particular we know that at any time t such that $\tilde{X}_j(t) > \tilde{X}_k(t)$ we must have

$$\tilde{X}_k(t) = \inf\{\tilde{X}_k(s) : s \leq t\} = \inf\{\tilde{X}_j(s) : s \leq t\} = \underline{X}(\tau_t). \quad (\text{A.16})$$

But the choice of the pair j, k was arbitrary, so we deduce that

whenever $X_k(t) > \underline{X}(t)$, it must be that $X_j(t) = \underline{X}(t)$ for all $j \neq k$.

This is statement (3) of [Theorem 1](#). In view of the facts that $\sum_{j=1}^N X_j = W$ and $\underline{X} = N^{-1}W$, if at time t we have $X_k(t) > \underline{X}(t)$ then

$$X_k(t) - \underline{X}(t) = \sum_j \{X_j(t) - \underline{X}(t)\} = W_t - \underline{W}_t, \quad (\text{A.17})$$

proving statement (4) of [Theorem 1](#).

The picture of the solution of [\(A.1\)](#) is now clearer: *each excursion away from 0 of $W - \underline{W}$ is assigned to exactly one of the X_j .* We shall now prove that the probabilistic structure of any solution to [\(A.1\)](#) coincides with the probabilistic structure of the solution constructed in [Proposition 3](#).

For this, define

$$A_j(t) \equiv \int_0^t I_j(X(s)) ds, \quad \tau_j(t) \equiv \inf\{s : A_j(s) > t\}, \quad X^j(t) = X_j(\tau_j(t)). \quad (\text{A.18})$$

Thus each X^j is a standard Brownian motion. In fact, the X^j are *independent* Brownian motions, as we see by the following argument. Fix any $t_1, \dots, t_N > 0$, and any $\theta_1, \dots, \theta_N \in \mathbb{R}$. Then

$$M_t \equiv \exp \left[\sum_{j=1}^N \{i\theta_j X_j(t \wedge \tau_j(t_j)) - \frac{1}{2} \theta_j^2 A_j(t \wedge \tau_j(t_j))\} \right]$$

is a bounded martingale,⁴ so

$$1 = EM_0 = EM_\infty = E \left(\exp \left[\sum_{j=1}^N \{i\theta_j X^j(t_j) - \frac{1}{2} \theta_j^2 t_j\} \right] \right).$$

Hence the $X^j(t_j)$ are independent zero-mean Gaussian, and the independence of the X^j follows. Thus if we decompose each X^j into its Poisson process Π^j of excursions away from the minimum as in [Proposition 3](#), then the Π^j are independent. Therefore if we define a random measure $\bar{\Pi}$ on $(0, \infty) \times U_+ \times \{1, \dots, k\}$ by

$$\bar{\Pi}(B \times \{j\}) = \Pi^j(B)$$

⁴ This is proved using Itô's formula, and the fact that $d\langle X_j, X_k \rangle = 0$ for $j \neq k$.

for any Borel $B \subseteq (0, \infty) \times U_+$, it can be seen that \bar{H} is a Poisson random measure, with mean measure $dt \times n_+(df) \times \mu_N$, where as before $\mu_N(\{j\}) = 1$ for $j = 1, \dots, N$. This is the measure M_N defined at (A.5), but scaled up by a factor of N . A point in $(0, \infty) \times U_+ \times \{1, \dots, k\}$ is a triple, where the final component in $\{1, \dots, N\}$ we refer to as the *label*. If we take all points in \bar{H} with label j , we see the Poisson point process of excursions of a Brownian motion (in fact, X^j); if we take all points in \bar{H} , we see the Poisson point process of a Brownian motion (in fact, W) but scaled up by a factor of N . This means that the local time of the totality of all points in \bar{H} is growing N times as fast as the local time of the corresponding Brownian motion, explaining the factor N^{-1} in expressions (A.9), (A.10). \square

Proposition 5. *For any k , the process*

$$M_t \equiv X_k(t) - (N-1)^{-1} \sum_{j \neq k} X_j(t) = \frac{NI_k(X(t)) - 1}{N-1} (W_t - \underline{W}_t) \quad (\text{A.19})$$

is a martingale.

Proof. Firstly we verify the algebraic equivalence of the two sides of (A.19). If X_k is the lead process at time t (that is, $I_k(X(t)) = 1$), then from (A.17) we have $X_k(t) = W_t - \underline{W}_t + \underline{X}(t)$, and $X_j(t) = \underline{X}(t)$ for $j \neq k$, so the two sides of (A.19) agree in this case. If X_k is not the lead process at time t , then similarly the left-hand side of (A.19) is equal to $-(N-1)^{-1}(W_t - \underline{W}_t)$, as required.

Now take $0 < s < t$, any $A \in \mathcal{F}_s$, and let $\tau = \inf\{u > s : W_u = \underline{W}_u\}$. Notice that

$$\begin{aligned} (N-1)E[M_t - M_\tau : A, \tau < t] &= E[(NI_k(X(t)) - 1)(W_t - \underline{W}_t) : A, \tau < t] \\ &= E[NI_k(X(t)) - 1 : A, \tau < t] E[W_t - \underline{W}_t : A, \tau < t] \\ &= 0 \end{aligned}$$

since the label of any excursion is independent of the path of that excursion, and each label has equal probability $1/N$. Therefore

$$\begin{aligned} E[M_t - M_s : A] &= E[M_t - M_s : A, \tau \leq t] + E[M_t - M_s : A, \tau > t] \\ &= E[M_\tau - M_s : A, \tau \leq t] + E[M_t - M_s : A, \tau > t] \\ &= E[M_{\tau \wedge t} - M_s : A] \\ &= E[(N-1)^{-1}(I_k(X(s)) - 1)(W_{\tau \wedge t} - W_s) : A] \\ &= E[E(W_{\tau \wedge t} - W_s | \mathcal{F}_s) (N-1)^{-1}(I_k(X(s)) - 1) I_A] \\ &= 0, \end{aligned}$$

because the label of the lead process does not change during $[s, \tau]$, nor does \underline{W} . \square

This completes the proof of Theorem 1 in the case where $X(0) = 0$. The general case follows by concatenation. So if we have $X_1(0) > X_2(0) > \dots > X_N(0)$, then up until the time $T_j \equiv \inf\{t : X_1(t) = X_j(t)\}$ none of the processes X_i , $i \geq j$ has moved. Up until T_2 , only X_1 is moving, so this behaves like Brownian motion. Between T_2 and T_3 , both X_1 and X_2 are moving with a common minimum, so we may apply Theorem 1 with two processes, both starting at the same place; then between T_3 and T_4 we have three moving processes, and so on.

References

- [1] E. Bayraktar, R. Kravitz, Quickest search over Brownian channels, *Stochastics* 86 (3) (2014) 473–490.
- [2] D.A. Darling, A.J.F. Siegert, The first passage problem for a continuous Markov process, *Ann. Math. Stat.* 24 (4) (1953) 624–639.
- [3] V. Dragalin, A simple and effective scanning rule for a multi-channel system, *Metrika* 43 (1) (1996) 165–182.
- [4] A. Dvoretzky, J. Kiefer, J. Wolfowitz, Sequential decision problems for processes with continuous time parameter. Testing hypotheses, *Ann. Math. Stat.* 24 (2) (1953) 254–264.
- [5] J. Heydari, A. Tajer, H.V. Poor, Quickest linear search over correlated sequences, *IEEE Trans. Inform. Theory* 62 (10) (2016) 5786–5808.
- [6] G. Kallianpur, M. Fujisaki, H. Kunita, Stochastic differential equations for the non linear filtering problem, *Osaka J. Math.* 9 (1) (1972) 19–40.
- [7] E.M. Klimko, J. Yackel, Some aspects of search strategies for Wiener processes. Statistical decision theory and related topics (Proc. Sympos., Purdue Univ., Lafayette, Ind., 1970), 1971, pp. 225–237.
- [8] E.M. Klimko, J. Yackel, Optimal search strategies for Wiener processes, *Stochastic Process. Appl.* 3 (1) (1975) 19–33.
- [9] L. Lai, H.V. Poor, Y. Xin, G. Georgiadis, Quickest search over multiple sequences, *IEEE Trans. Inform. Theory* 57 (8) (2011) 5375–5386.
- [10] R. Liptser, A.N. Shiryaev, A Bayesian problem of sequential search in diffusion approximation, *Theory Probab. Appl.* 10 (1) (1965) 178–186.
- [11] G. Peskir, A.N. Shiryaev, *Optimal Stopping and Free-boundary Problems*, Springer, 2006.
- [12] E. Posner, H. Rumsey, Continuous sequential decision in the presence of a finite number of hypotheses, *IEEE Trans. Inform. Theory* 12 (2) (1966) 248–255.
- [13] L.C.G. Rogers, D. Williams, *Diffusions, Markov Processes and Martingales: Volume 2, Itô calculus*, Vol. 2, Cambridge University Press, 2000.
- [14] A.N. Shiryaev, On the theory of decision functions and control by an observation process with incomplete data, in: *Trans. Third Prague Conf. Information Theory, Statist. Decision Functions, Random Processes* (Liblice, 1962), Publ. House Czech. Acad. Sci. Prague, 1964, p. 657.
- [15] A.N. Shiryaev, *Optimal Stopping Rules*, Vol. 8, Springer Science & Business Media, 2007.
- [16] A. Wald, Sequential tests of statistical hypotheses, *Ann. Math. Stat.* 16 (2) (1945) 117–186.
- [17] K.Sh. Zigangirov, On a problem in optimal scanning, *Theory Probab. Appl.* 11 (2) (1966) 294–298.