

1 **Oxygen isotope systematics of chondrules in the Paris CM2 chondrite: 2**
3 **indication for a single large formation region across snow line 3**

Noël Chaumard^{a,b}, Céline Defouilloy^{a,c}, Andreas T. Hertwig^{a,d}, Noriko T. Kita^{a,*} 4 5

6 ^aWiscSIMS, Department of Geoscience, University of Wisconsin-Madison, 1215 W. Dayton 7
Street, Madison, WI 53706-1692, USA.

^b 8 Fi Group, Direction scientifique, 14 terrasse Bellini, 92800 Puteaux, France. ^c

9 CAMECA, 29 quai des Grésillons, 92622 Gennevilliers Cedex, France.

^d 10 Institut für Geowissenschaften, Universität Heidelberg, Im Neuenheimer Feld 234-236, 69120
11 Heidelberg, Germany.

12

13 *Revised version submitted to Geochimica et Cosmochimica Acta* 14

15 February 6, 2020

16

17

18

* 19 Corresponding author. E-mail: noriko@geology.wisc.edu

20

21 Keywords: Carbonaceous chondrites, chondrules, oxygen three-isotope measurements, SIMS 22
analyses

23

25 In-situ oxygen three-isotope analyses of chondrules and isolated olivine grains in the Paris 26 (CM)
 chondrite were conducted by secondary ion mass spectrometry (SIMS). Multiple analyses of olivine
 and/or pyroxene in each chondrule show indistinguishable $\Delta 27^{17}\text{O}$ values, except for minor
 occurrences of relict olivine grains (and one low-Ca pyroxene). A mean $\Delta 28^{17}\text{O}$ value of these 29
 homogeneous multiple analyses was obtained for each chondrule, which represent oxygen isotope
 ratios of the chondrule melt. The $\Delta 30^{17}\text{O}$ values of individual chondrules range from -7‰ to -2‰ 31
 and generally increase with decreasing Mg# of olivine and pyroxene in individual chondrules. Most
 type I (FeO-poor) chondrules have high Mg# (~ 99) and variable $\Delta 32^{17}\text{O}$ values from -7.0‰ to 33
 -3.3‰ . Other type I chondrules (Mg# ≤ 97), type II (FeO-rich) chondrules, and two isolated FeO rich
 olivine grains have host $\Delta 34^{17}\text{O}$ values from -3‰ to -2‰ . Eight chondrules contain relict grains that
 are either ^{16}O -rich or ^{16}O -poor relative to their host chondrule and show a wide range of $\Delta 35^{17}\text{O}$ 36
 values from -13‰ to 0‰ .

37 The results from chondrules in the Paris meteorite are similar to those in Murchison (CM).
 Collectively, the $\Delta 38^{17}\text{O}$ values of chondrules in CM chondrites continuously increase from -7‰ to
 -2‰ with decreasing Mg# from 99 to 37. The majority of type I chondrules (Mg# > 98) show $\Delta 39^{17}\text{O}$
 values from -6‰ to -4‰ , while the majority of and type II chondrules (Mg# 60-70) show $\Delta 40^{17}\text{O}$
 values of -2.5‰ . The covariation of $\Delta 41^{17}\text{O}$ versus Mg# observed among chondrules in CM 42
 chondrites may suggest that most chondrules in carbonaceous chondrites formed in a single large
 region across the snow line where the contribution of ^{16}O -poor ice to chondrule precursors and 44
 dust enrichment factors varied significantly.

46 1. INTRODUCTION

47

48 Among carbonaceous chondrites (CCs), CM (Mighei-like) chondrites are the most 49 abundant
group (Weisberg et al., 2006). They are recognized as xenoliths in numerous other groups 50 of CCs
and meteorite classes and thus may correspond to the most abundant and/or widely 51 dispersed
material in the main belt (e.g., Zolensky et al., 1996; Gounelle et al., 2003; Bischoff et 52 al., 2006; and
references therein; Briani et al., 2012), hence their importance to deciphering the 53 formation and
evolution of the early Solar System. In addition to low to mild thermal 54 metamorphism (e.g.,
Nakamura, 2006; Kimura et al., 2011; Tonui et al., 2014), CM chondrites 55 experienced intense parent
body aqueous alteration (e.g., Sears and Dodd, 1988; Brearley, 2003; 56 Busemann et al., 2007;
Schrader and Davidson, 2017). Most CM chondrites are of petrologic type 57 2 (e.g., McSween 1979;
Kallemeyn and Wasson, 1981; Zolensky et al., 1993) and display various 58 degrees of aqueous
alteration and have been divided into petrologic subtypes from 3.0 to 2.0, where 59 numbers decreases
with increasing alteration (e.g., Zolensky et al., 1997; Rubin et al., 2007; Rubin, 60 2015; Kimura et
al., 2020). Some CM chondrites are almost completely altered and assigned to be 61 subtype 2.0, which
was previously classified as CM1 (Zolensky et al., 1997; Rubin et al., 2007).

62 Paris is one of the least altered CM chondrites and has been used to investigate the early 63 stages of
the parent body aqueous alteration of CM chondrites (Hewins et al., 2014; Marrocchi et 64 al., 2014;
Rubin, 2015; Leroux et al., 2015; Pignatelli et al., 2016; Vacher et al., 2016, 2017; 65 Verdier-Paoletti
et al., 2017). Based on a detailed petrographic and mineralogical survey, 66 Marrocchi et al. (2014)
classified Paris as a CM2.7. However, Paris contains both highly and less 67 altered lithologies

(Hewins et al., 2014). Based on the PCP (Poorly Characterized Phases) index 68 defined by Rubin et al. (2007), the less altered lithology of Paris is of petrologic subtype 2.9

69 (Hewins et al., 2014), which is consistent with the observation of a significant amount of Fe-Ni 70 metal blebs and the presence of a pristine matrix (Leroux et al., 2015; Rubin, 2015). The oxygen isotope ratios of the less altered lithologies are as low as $\delta^{17}\text{O} = -2.1\text{‰}$ and $\delta^{18}\text{O} = 2.4\text{‰}$, which 72 is at the lower end of a linear trend defined by Paris subsamples and bulk CM2 chondrites (Hewins 73 et al., 2014). The variation of oxygen isotope ratios among bulk CM chondrites is ascribed to 74 heterogeneity in the extent of secondary aqueous alteration processes (Clayton and Mayeda, 1999; 75 Hewins et al., 2014). The less altered lithology of Paris thus offers a unique opportunity to 76 investigate the origin and petrogenesis of CM chondrules.

77 In situ SIMS (secondary ion mass spectrometry) oxygen 3-isotope analysis of individual 78 chondrules is a powerful tool to constrain the conditions of their formation (e.g., Kita et al., 2010; 79 Ushikubo et al., 2012; Schrader et al., 2013; Marrocchi et al. 2018; 2019). Many chondrules in 80 CCs contain olivine grains with heterogeneous oxygen isotope ratios (e.g., Kunihiro et al., 2004, 81 2005; Jones et al., 2004; Wasson et al., 2004; Connolly and Huss, 2010; Rudraswami et al., 2011; 82 Ushikubo et al., 2012; Schrader et al., 2013; 2017; Tenner et al., 2013; Marrocchi et al., 2018; 83 2019). They are considered as “relict” and interpreted as unmelted material that survived the final 84 high-temperature event of chondrule formation. Because of the slow diffusivity of oxygen isotopes 85 in olivine (e.g., Chakraborty, 2010), these relict grains preserved their initial oxygen isotope ratios 86 and would provide important knowledge about precursor solids that formed chondrules. However, 87 multiple high precision SIMS analyses of olivine and/or pyroxene in each chondrules are mostly 88 indistinguishable (e.g., Rudraswami et al., 2011; Ushikubo et al., 2012; Tenner et al., 2013; 2015; 89

2017; Hertwig et al., 2018; 2019a; Chaumard et al., 2018). Ushikubo et al. (2012) showed ⁹⁰ plagioclase and glass in chondrule mesostasis from Acfer 094 are in agreement with those of ⁹¹ olivine and pyroxene phenocrysts of the same chondrules. Internally homogeneous oxygen isotope

4

⁹² ratios within individual chondrules represent those of the final chondrule melt from which the ⁹³ “non-relict” olivine and other minerals crystallized.

⁹⁴ The degree of mass independent isotope fractionation of oxygen 3-isotopes, commonly expressed as $\Delta^{17}\text{O}$ ($= \delta^{17}\text{O} - 0.52 \times \delta^{18}\text{O}$), determined for individual chondrules in CCs, ⁹⁶ systematically vary against the Mg# ($= \text{MgO}/[\text{MgO}+\text{FeO}]$ in mol.%) of olivine and pyroxene ⁹⁷ (Ushikubo et al., 2012; Tenner et al., 2013; 2015; 2017; Hertwig et al., 2018; 2019a). Variation of

⁹⁸ Mg#s among chondrules indicates that the redox state of the environments they formed in were ⁹⁹ variable. Such variations were probably due to metal-silicate equilibria (Zanda et al., 1994) which ¹⁰⁰ were influenced by varying proportions of anhydrous dust, H₂O ice, and organic matters relative ¹⁰¹ to the solar-composition nebula gas in the outer disk regions where CC chondrules formed (e.g., ¹⁰² Wood and Hashimoto, 1993; Grossman et al., 2008). Heterogeneous oxygen isotope ratios among chondrule precursor phases, such as ¹⁶O-rich anhydrous dust and ¹⁰³ ¹⁶O-poor H₂O-ice (Krot et al., 2006; Sakamoto et al., 2007), could explain the negative correlation between Δ ¹⁰⁴ ¹⁷O values and Mg# ¹⁰⁵ among chondrules in CCs (e.g., Connolly and Huss, 2010; Schrader et al., 2013; Tenner et al., ¹⁰⁶ 2015; Hertwig et al., 2018).

In contrast, Δ ¹⁰⁷ ¹⁷O values among chondrules in non-carbonaceous chondrites, such as ¹⁰⁸ ordinary and enstatite chondrites that likely derived from inner disk material (Kruijer et al., 2017), show narrower ranges and are ¹⁰⁹ ¹⁶O-depleted relative to those in CCs (e.g., Kita et al., 2010; ¹¹⁰ Weisberg et al., 2011; Libourel and Chaussidon, 2011; Schneider et al., 2020). Kita et al. (2010) argued that solid

precursors of ordinary chondrite chondrules were fractionated in $\delta^{18}\text{O}$ as a result of condensation of solids from ^{16}O -poor nebula gas at high temperatures. It is likely that the oxygen isotope ratios of precursor phases were homogenized in the inner disk region prior to the formation of chondrules.

5

Chaumard et al. (2018) studied 29 chondrules in the Murchison CM2 chondrite and found that the distribution of Mg#s and $\Delta^{17}\text{O}$ of individual chondrules are similar to those in Acfer 094 and CO3 chondrites. Co-existing olivine and pyroxene in individual Murchison chondrules show indistinguishable oxygen isotope ratios excluding the minor occurrence of relict grains, which further indicate that chondrules solidified from numerous melt droplets with homogeneous oxygen isotope ratios. Chondrule oxygen isotope systematics in Murchison further indicated that the CO and CM chondrite parent bodies collected similar populations of chondrules, while other studies suggest that the timing of the CM parent body accretion may have been delayed relative to the CO parent body accretion (e.g., Sugiura and Fujiya, 2014).

In Murchison, chondrules have an extremely limited range of Mg#s ranging from 99.6 to 99.0 for type I chondrules (with one exception with Mg#~96) and a slightly larger range, from ~65 to ~70, for type II chondrules. The $\Delta^{17}\text{O}$ values among Murchison chondrules show a hint of systematic increase with decreasing Mg#s. Here we report *in situ* high precision SIMS oxygen 3-

isotope measurements of olivine and pyroxene in chondrules from the less altered lithology of the Paris CM2 chondrite. The Paris meteorite provides an opportunity to study a diverse range of less altered chondrules in CM chondrites in petrologic context, including chondrule textures, mineralogy, and mineral chemistry (Hewins et al., 2014; Rubin, 2015; Stephant et al., 2017). This will enable us to further constrain the nature of pristine chondrules from carbonaceous chondrites in general.

134

135 **2. ANALYTICAL PROCEDURES**

136

137 **2.1. Sample and chondrule selection**

6

138

139 Within the less altered lithology of the Paris CM chondrite (type 2.9; Hewins et al., 2014), 140 we
selected and analyzed 29 chondrules from one polished section allocated by the Muséum 141 national
d'Histoire naturelle of Paris (MNHN 4029-11). We intended to obtain the most diverse 142 selection of
chondrules based on their size, texture, and mineralogy. Selected chondrules include 143 25 type I
(Mg# $\geq$ 90) chondrules (from $\sim$ 200 μ m to $\sim$ 2 mm in diameter), three type II (Mg# $<$ 90) 144 chondrules
($\sim$ 250–600 μ m in diameter), and one fragment of a type I chondrule ($\sim$ 600 μ m long). 145 We also
analyzed two isolated grains of FeO-rich olivine ($\sim$ 250–400 μ m in size), one amoeboid 146 olivine
aggregate (AOA) ($\sim$ 500 μ m long), and one FeO-poor olivine-bearing object ($\sim$ 400 μ m 147 long).

148 Nine type I chondrules are porphyritic olivine-pyroxene (POP; 20–80% modal olivine), 149 three
are porphyritic olivine (PO; $>$ 80% modal olivine), and eight are porphyritic pyroxene (PP; 150 $<$ 20%
modal olivine). One type I chondrule is barred olivine (BO) and two are composed of a BO 151 core
surrounded by a POP rim. We also analyzed one type I granular olivine (GO) chondrule and 152 one
granular olivine-pyroxene (GOP) chondrule. The fragment of the type I chondrule displays a 153 BO
texture. The three type II chondrules have a PO texture.

154 In all groups of CCs, chondrules are predominantly type I (e.g., Jones, 2012). It has been 155
reported that $\sim$ 95% of chondrules in CM chondrites are porphyritic, 10–40% of these porphyritic 156
chondrules being type II (Jones, 2012; and references therein). Approximately 80% of the 157

chondrules analyzed are porphyritic and ~13% of them are type II. This selected population is thus roughly representative of chondrules in CM chondrites. Moreover, PO, POP, and PP chondrules in our selection represent approximately 10%, 31%, and 28%, respectively, of the entire population of chondrules analyzed in this study.

7

161

162 **2.2. Scanning electron microscopy and electron microprobe analysis**

163

164 Backscattered electron (BSE), secondary electron (SE) imaging and energy dispersive X ray
spectrometry (EDS) analyses of chondrules were performed using a Hitachi S-3400N scanning
electron microscope (SEM) at the University of Wisconsin-Madison. The accelerating voltage was
set to 15 kV. The locations of SIMS analyses were selected for olivine and pyroxene grains in each
chondrule that are free of cracks and other phases as identified in BSE and SE images. We used
a Cameca SXFive FE electron microprobe at the University of Wisconsin Madison to obtain
quantitative chemical analyses of olivine and pyroxene grains with an accelerating voltage of 15
keV and a beam current of 20 nA. Counting times for the peak and background were 10 and 5 s,
respectively. Several standards were analyzed for matrix correction of individual elements: natural
olivine and synthetic forsterite and enstatite (Mg, Si), jadeite (Na, Al), microcline (K), chromian
augite (Ca), TiO₂ (Ti), synthetic Cr₂O₃ (Cr), fayalite (Fe), and synthetic Mn₂SiO₄ (Mn). We analyzed
Mg, Al, Si, and Na with a LTAP crystal, Ca and K with a LPET crystal, Cr and Ti with a PET
crystal, and Mn and Fe with a LLIF crystal. We used the Probe for EPMATM (PFE) software
(Donovan, 2015) for data reduction and matrix corrections (ZAF and $\phi_{\rho(z)}$). For each analysis, we
calculated the Mg# based on EPMA measurements.

180 **2.3. SIMS oxygen three-isotope analysis**

182 We performed *in situ* oxygen three-isotope analyses of olivine and pyroxene using the 183 Cameca
 IMS 1280 at the WiscSIMS laboratory, University of Wisconsin-Madison. Analytical

8

184 conditions and data reduction methods were generally similar to those of Kita et al. (2010) and
 Tenner et al. (2013, 2015) using multi-collector Faraday cups; ^{16}O and 185 ^{18}O on multi-collector array
 and 186 ^{17}O on a fixed mono-collector (axial detector FC2). The analyses were performed in two
 sessions with different primary Cs^+ 187 beam conditions; a 15 μm (session #1) and 10 μm (session #2)
 188 diameters with intensities of $\sim 3\text{ nA}$ and $\sim 1\text{ nA}$, respectively. During the analysis session #1 (15 μm
 diameter spot), secondary ion intensities of ^{16}O , ^{17}O , and ^{18}O were $\sim 3.5 \times 10^9$, $\sim 1.5 \times 10^6$ 189 , and
 $\sim 7.5 \times 10^6$ counts per second (cps), respectively. FC amplifier resistors were $10^{11}\ \Omega$ for ^{17}O and 190 ^{18}O ,
 and $10^{10}\ \Omega$ for 191 ^{16}O as in the previous studies (e.g., Kita et al., 2010), though a newer version of the
 FC amplifier board was used for the detection of 192 ^{17}O on the axial FC2 detector. This newer FC 193
 amplifier board from Cameca (for IMS 1280-HR) shows improved thermal noise that is close to 194 the
 theoretical limit (1SD $\sim 1,300$ cps for 4 s integrations) compared to the original FC amplifier 195 board
 installed to IMS 1280 (1SD $\sim 2,000$ cps for 4 s integrations). Taking advantage of lower noise level on
 196 ^{17}O , the acquisition time for a single analysis was reduced from 200 s to 100 s and 197 the total
 analysis time (including pre-sputtering and secondary beam centering time) was reduced 198 from 7 min
 to 4 min compared to previous oxygen 3-isotope analyses at the WiscSIMS laboratory. 199 Analyses
 session #2 (10 μm) was performed at the same time that some of chondrules from the Murchison were
 analyzed (Chaumard et al., 2018). Secondary ion intensities of ^{16}O , ^{17}O , and 200 ^{18}O decreased to
 $\sim 1.5 \times 10^9$, $\sim 5.5 \times 10^5$, and $\sim 2.9 \times 10^6$ 201 cps, respectively. In order to reduce the noise level of the FC

amplifier for ^{17}O analyses below 10^6 cps intensity, we replaced the resistor and capacitor pair on the original IMS 1280 FC amplifier board by a $10^{23} \Omega$ resistor and 1 pF capacitor from a Finnigan MAT 251 stable mass spectrometer (noise level was reduced to $1\text{SD} \sim 1,200$ cps for 4 s integrations; Chaumard et al., 2018). The mass resolving power (MRP at 10% peak height) for both sessions was set to ~ 2200 for ^{16}O and ^{18}O , and 5000 for ^{17}O . We measured $^{16}\text{OH}^-$ at the end

9
of each analysis to determine its contribution to the $^{17}\text{O}^-$ signal following the methods described by Heck et al. (2010). The correction of the $^{17}\text{O}^-$ signal from $^{16}\text{OH}^-$ was negligible ($<0.08\text{‰}$) for both standards and unknowns.

We normalized the measured $^{18}\text{O}/^{16}\text{O}$ and $^{17}\text{O}/^{16}\text{O}$ ratios to the VSMOW scale ($\delta^{210} ^{18}\text{O}$ and $\delta^{211} ^{17}\text{O}$ expressed as a deviation from standard mean ocean water in the unit of 1/1000; VSMOW scale, Baertschi 1976). The external reproducibility has been determined by intermittent measurements of a San Carlos olivine (SC-Ol) standard ($\delta^{213} ^{18}\text{O} = 5.32\text{‰}$; Kita et al., 2010). We bracketed 4 to 19 unknown chondrule analyses with 8 SC-Ol analyses, 4 before and 4 after (Kita et al., 2009). External reproducibility is calculated as the 2SD of the SC-Ol brackets, with average values during session #1 (15 μm) of 0.2‰, 0.3‰, and 0.3‰ for $\delta^{18}\text{O}$, $\delta^{17}\text{O}$, and $\Delta^{16} ^{17}\text{O}$, respectively. For session #2 (10 μm), the 2SD average values are 0.4‰, 0.5‰, and 0.4‰ for $\delta^{18}\text{O}$, $\delta^{17}\text{O}$, and $\Delta^{16} ^{17}\text{O}$, respectively. These values represent the spot-to-spot reproducibility and were thus assigned as the uncertainties of each individual spot analysis (Kita et al., 2009). Measurements of four olivine ($\text{Fo}_{0.6-100}$), three low-Ca pyroxene (En_{70-97}), and one diopside (Wo_{50}) standards with known oxygen isotope ratios (Eiler et al., 1997; Kita et al., 2010) were used to estimate corrections for instrumental biases of unknown olivine and pyroxene analyses (EA1). The compositional ranges of standards

cover those of unknowns measured.

224 We obtained multiple SIMS analyses per chondrule ($n=4$ to 12, typically 8) to examine the 225
homogeneity of the isotope ratios. As in the previous studies, a specific analysis in a single chondrule is
identified as a relict when its Δ 226 ^{17}O value deviates more than 0.5‰ and 0.7‰ (3SD 227 limits of
bracket standard analyses in each analysis session) from the chondrule mean (Ushikubo 228 et al., 2012;
Tenner et al., 2013, 2015; 2017; Hertwig et al., 2018; 2019a; Chaumard et al., 2018).

229 To identify all chondrules containing rare relict grains, a multitude of analyses per chondrules

10

230 (e.g., ~50; Marrocchi et al., 2018; 2019) is required; however, a total of ~8 analyses per chondrule is
sufficient to identify the Δ 231 ^{17}O value of most phenocrysts that in turn represents the value of the 232
chondrule melt, if the chondrule consists largely of minerals with homogeneous oxygen isotope 233
ratios. Thus, we can calculate mean oxygen isotope ratios (referred to as “host chondrule” oxygen 234
isotope ratios) from multiple analyses within each chondrule excluding relict grains. The

uncertainties of host δ ^{18}O and δ 235 ^{17}O values is the propagation of (i) the 2 standard error of the mean
236 of multiple analyses that constitute the host chondrule value ($=2\text{SD}/\sqrt{\text{number of analyses}}$; 2SD of
237 the bracket standard analyses or 2SD of multiple analyses within a chondrule, whichever is larger),
238 (ii) the 2SE of associated SC-OI bracketing analyses for instrumental bias correction, and (iii) the
239 uncertainty due to the sample topography and/or sample positioning on the SIMS stage as well as
uncertainty of instrumental bias corrections, estimated to be 0.3‰ for δ ^{18}O and 0.15‰ for δ 240 ^{17}O
(Kita et al., 2009). Because (iii) is mass-dependent and does not affect Δ 241 ^{17}O , we only use (i) and
(ii) for the propagated uncertainty of Δ ^{17}O . The uncertainties of Δ 242 ^{17}O in the relict grains are the
243 spot-to-spot reproducibility (2SD) as determined by bracketing analyses of San Carlos olivine.

244 We inspected each SIMS spot by obtaining BSE and SE images using SEM after each 245 SIMS
session. Ten of 252 pits were rejected from our final dataset because they either overlapped cracks,
imperfections, or display a contribution of ^{16}OH to the $^{246} \text{}^{17}\text{O}$ signal of $\sim 0.1\%$ or higher. 247

248 **3. RESULTS**

249

250 **3.1. Texture, petrography, and mineralogy**

251

11

252 Chondrules display a wide diversity of sizes and textures, as previously observed by 253 Hewins et
al. (2014) for the Paris CM chondrite. In the following summary, textural features and 254 chemistry of
chondrule minerals are described. BSE images of individual chondrules are shown in 255 Figs. 1-4 and
EA2, in which locations of EPMA and SIMS analyses are annotated. EPMA major 256 element data for
olivine and pyroxene are shown in EA3. Additional petrologic descriptions of 257 individual chondrules
can be found in EA4. Within 19 of the 29 chondrules analyzed, coexisting 258 olivine and low-Ca
pyroxene have similar Mg#s. For the other 10 chondrules, we were only able 259 to obtain quantitative
analyses either for olivine or low-Ca pyroxene. In type I chondrules where 260 both olivine and
pyroxene were analyzed, Mg#s of olivine and low-Ca pyroxene are 261 indistinguishable, similar to
type I chondrules in other pristine chondrites (e.g., Jones, 1994; 262 Tachibana et al., 2003; Ushikubo et
al., 2012; Tenner et al., 2013, 2015; Schrader and Davidson, 263 2017; Schrader et al., 2017; Chaumard
et al., 2018; Hertwig et al., 2018; 2019a).

264 *3.1.1 Type I porphyritic chondrules*

265 Olivine in type I porphyritic chondrules is present as large anhedral phenocrysts and 266 euhedral

grains (up to $\sim 300\ \mu\text{m}$) (Fig. 1, 2). In several chondrules (e.g., C4, C8, C7; Fig. 2a for 267 C7), olivine is located in the cores while the pyroxene is more abundant at the periphery. In type I 268 POP chondrules, low-Ca pyroxene often displays a poikilitic texture with numerous cracks and 269 pores (e.g., C7; Fig. 2a). Low-Ca pyroxene was also observed as large euhedral grains, i.e., up to 270 $\sim 150\ \mu\text{m}$ length in chondrule C7 (Fig. 2a). In type I PP chondrules, low-Ca pyroxene display less 271 cracks and pores (e.g., C21; Fig. 1c). High-Ca pyroxene is also present in many type I porphyritic 272 chondrules as small grains (from a few μm to $\sim 70\ \mu\text{m}$ length), either in association with Low-Ca 273 pyroxene or olivine (e.g., C30; Fig. 1b). Olivine grains in type I porphyritic chondrules are 274 chemically homogeneous, with an average Mg# of 99.0 ± 0.8 (2SD) (97.6–99.8). They contain up

12

275 to 0.44 wt% Al_2O_3 , 0.84 wt% CaO, 0.67 wt% Cr_2O_3 , and 0.46 wt% MnO. Low-Ca pyroxene grains 276 have Mg#s ranging from 92.5 to 99.2 and compositions of $\text{En}_{89-98}\text{Fs}_{1-7}\text{Wo}_{0-4}$. We measured 0.17– 277 2.13 and 0.30–1.07 wt% Al_2O_3 and Cr_2O_3 , respectively.

278 The two POP chondrules C5 (Fig. 2b) and C14 (Fig. 3) contain fragments of BO chondrules 279 in their cores with sizes of $\sim 200\ \mu\text{m}$ and $\sim 600\ \mu\text{m}$, respectively. Olivine and pyroxene 280 compositions do not show significant differences between BO core and POP rim (see analysis 281 locations on BSE images in EA2 and EPMA data in EA3). There are two type I granular chondrules 282 (C1, GOP; C9, GO) that are composed of evenly sized grains. Olivine grains in the chondrule C9 283 display linear trails of micron-sized inclusions of metal (Fig. 2e). Mg# of olivine and low-Ca 284 pyroxene in these type I chondrules are all close to ~ 99 .

285 3.1.2 Type II PO chondrules

286 The three type II chondrules analyzed display a porphyritic texture and are mainly 287 composed of olivine phenocrysts. Most of the olivine grains in chondrule C15 have euhedral 288 shapes (Fig. 4a).

The chondrules C15 and C17 contain olivine grains with forsteritic cores ($\sim 20\text{--}289\text{--}80\text{ }\mu\text{m}$, $\text{Fo}_{99.91}$). The ranges of olivine Mg#s excluding forsteritic core in chondrule C15 and C17 are 58.9–77.3 and 69.9–79.7, respectively. In chondrule C22, olivine displays Mg#s ranging from 60.3–79.0. Olivine grains in these three type II chondrules contain 0.20–0.60 wt% Cr_2O_3 and 0.02–0.40 wt% MnO.

3.1.3 Isolated olivine grains

The isolated FeO-rich olivine grains G32 and G33 (Fig. 4b-c) are 200–300 μm in sizes. The Fo contents of olivine grains show ranges of 67–75 and 30–49 for G32 and G33, respectively, which becomes more FeO-rich towards the margin of grains. Another isolated object G24 is FeO-poor olivine-bearing object (Fig. 4d) with Mg# of $\sim 99.4 \pm 0.1$. The interior of the object G24 is composed

of olivine that contains numerous μm -sized Fe-metal inclusions. The texture of olivine is similar to that of dusty olivine grains, in which FeO was reduced to form Fe-rich metal (Nagahara, 1981; Rambaldi, 1981).

3.1.4 Amoeboid olivine aggregate (AOA)

The AOA I3 texturally resembles the amoeboid olivine inclusions (AOI) previously observed by Rubin (2015) in Paris, as well as AOAs in CV and CK chondrites (e.g., Grossman and Steele, 1976; Rubin, 2013). Small grains of diopside ($< 5\text{ }\mu\text{m}$) are enclosed within large, porous, forsteritic olivine grains. The AOA I3 mostly contains chemically homogeneous olivine with Mg# 99.5 ± 0.1 and 0.35 ± 0.11 and 0.28 ± 0.03 wt% of Cr_2O_3 and MnO, respectively.

3.2. Oxygen isotope ratios

310 A total of 242 oxygen 3-isotope analyses were obtained in 2 SIMS sessions from 33 objects 311
 (including 29 chondrules) in the CM2 Paris, which are listed in EA5. Typically, 8 analyses were 312
 performed in each object at the same location as the EPMA analyses (EA2 and EA3), including 313
 multiple phases (olivine, low-Ca pyroxene, and high-Ca pyroxene) where available. Only four 314 good
 analyses were obtained from BO chondrule C13 because several analyses were rejected due 315 to
 significant surface roughness (EA2). Only four analyses each were taken from two FeO-rich 316
 isolated olivine grains that were relatively small (200-300 μm). We performed four analyses in 317
 AOA I13, though two were rejected after inspections of the SIMS pits. Our new data plot between 318
 the CCAM (carbonaceous chondrite anhydrous mineral; Clayton et al., 1977) and the Y&R (Young 319
 and Russell, 1998) lines, close to the PCM (primitive chondrule minerals; Ushikubo et al., 2012) line.
 The $\delta^{18}\text{O}$ and $\delta^{17}\text{O}$ data from chondrules range from -23‰ to $+4\text{‰}$ and from -25‰ to $+2\text{‰}$,

14

321 respectively, for olivine, and from -11‰ to $+2\text{‰}$ and from -14‰ to -1‰ , respectively, for 322
 pyroxene (Fig. 5). These ranges are very similar to those observed in chondrules from Murchison 323
 (Chaumard et al., 2018).

324

325 3.2.1. Chondrule oxygen isotope analyses

326 Table 1 lists the host oxygen isotope ratio calculated for all chondrules except for two (C9, 327
 C15), as well as their texture, Mg#, the number of measurements per mineral, and the individual 328
 analyses that were not included in host chondrule calculations. In 18 chondrules, multiple analyses
 within a single chondrule display indistinguishable $\Delta^{17}\text{O}$ values within the 3SD external 330
 reproducibility (0.5‰ and 0.7‰ , for 15 μm and 10 μm spot analyses, respectively). These 331

chondrules are considered to be internally homogeneous in oxygen 3-isotope ratios. Eight chondrules contain olivine grains showing $\Delta^{17}\text{O}$ values that differ by more than the 3SD external reproducibility from the average values calculated using the remaining multiple (≥ 5) analyses. As described in section 2.3, these olivine grains with distinct oxygen isotope ratios are considered to be relict and were not used to calculate the individual host chondrule values. In the case of chondrule C1, one each of olivine and low-Ca pyroxene show $\Delta^{17}\text{O}$ values of -9.5‰ and -7.0‰ , respectively, which are significantly lower than remaining five analyses (two olivine and three low-Ca pyroxene grains) with the mean $\Delta^{17}\text{O} = -5.0 \pm 0.6\text{‰}$ (2SD) (Fig. 6). We considered the ^{16}O rich olivine and low-Ca pyroxene grains as relict grains. By excluding relict grain data, we calculate the host chondrule oxygen ratios from multiple (4–11) analyses for all but two chondrules. We consider the host chondrule $\Delta^{17}\text{O}$ values to represent oxygen isotope ratios of the chondrule melt during their formation (e.g., Ushikubo et al., 2012; Tenner et al., 2013; 2015).

15

In chondrule C18, five olivine grains display variable $\Delta^{17}\text{O}$ value of $-1.3 \pm 0.6\text{‰}$ (2SD), which differ significantly from three pyroxene analyses ($-3.3 \pm 0.3\text{‰}$; Fig. 1a). We consider pyroxene data to represent the host chondrule value and olivine grains to be relict grains, as was discussed in previous studies (chondrule Y22 in Tenner et al., 2013; chondrule A6 in Chaumard et al., 2018). Similarly, chondrule C8 contains dusty olivine grains (SIMS spot #252, EA2 and EA5) with a $\Delta^{17}\text{O}$ value of $0.1 \pm 0.4\text{‰}$, which are significantly different from the host chondrule $\Delta^{17}\text{O}$ value of $-5.4 \pm 0.2\text{‰}$ that are calculated from five olivine and pyroxene analyses (Table 1).

In chondrule C9, seven olivine grains display homogeneous isotope ratios with the mean $\Delta^{17}\text{O}$

value of $-0.15 \pm 0.43\text{‰}$ (2SD), which differ significantly from a single analysis of pyroxene ($-352.7.8\text{‰}$; Fig. 2e). All olivine grains in C9 contain numerous micron-sized inclusions of metal, which suggests they are dusty olivine grains and thus are likely to be relict. The pyroxene data are more likely to represent the host value. However, since there is only a single pyroxene analysis, we do not assign it to be host $\Delta 355^{17}\text{O}$ value.

We also did not assign a host $\Delta 356^{17}\text{O}$ value to the type II chondrule C15 (Fig. 4a) that show a large range of $\delta^{18}\text{O}$ and $\delta 357^{17}\text{O}$ values between PCM and CCAM lines from -6.6‰ to $+2.5\text{‰}$ and from -9.8‰ to -1.0‰ , respectively, among 8 olivine analyses. Chondrule C15 contains forsteritic olivine grains in its core (Fo_{91-99}), indicating that these grains represent relict grains (based on FeO contents, Fe/Mn ratios; Berlin et al., 2011; Frank et al., 2014; Schrader and Davidson, 2017). This is consistent with the lower $\Delta 361^{17}\text{O}$ values (-4.1‰ to -6.3‰) of forsteritic olivine grains compared to those of the more FeO-rich olivine grains with variable $\Delta 362^{17}\text{O}$ values (-3.3‰ to -2.3‰).

Host $\Delta 363^{17}\text{O}$ values from most of the type I chondrules vary continuously from $-7.0 \pm 0.2\text{‰}$ to $-3.3 \pm 0.3\text{‰}$ (Fig. 6). The two other type I chondrules (C11; Fig. 1d, and C12) and the two type II chondrules (C17 and C22) display a narrow range of host $\Delta 365^{17}\text{O}$ values ranging from $-2.5 \pm 0.2\text{‰}$

16

to $-2.1 \pm 0.2\text{‰}$ (Fig. 6). Most of the relict olivine have $\Delta 366^{17}\text{O}$ values between $\sim -2\text{‰}$ and -8‰ (Fig. 6) overlapping those of the host values of other chondrules. Similar observation have been made in other carbonaceous chondrites such as Acfer 094, Murchison, Y-82094, CV, CO, and CR chondrites (Rudraswami et al., 2011; Ushikubo et al., 2012; Schrader et al., 2013, 2014, 2017; Tenner et al., 2013, 2015, 2017; Hertwig et al., 2018; 2019a; Chaumard et al., 2018). Only two relict grains have $\Delta 371^{17}\text{O}$ values lower than -8‰ ; the relict grains in chondrule C1 (-9.5‰) and C14

(−13.0‰) (Fig. 6). Among the 29 relict grains analyzed, 13 (only olivine) display Δ 372 ^{17}O values 373 between −1.6‰ and 0.2‰ (Fig. 6). Most of these were found in the chondrule C9 that contains 7 relict olivine grains with Δ 374 ^{17}O values ranging from −0.4‰ to 0.2‰ (Fig. 2e), and in chondrule C18 that contains 5 relict olivine grains with Δ 375 ^{17}O values between −1.6‰ to −0.8‰ (Fig. 1a).

376

377 3.2.2. Oxygen isotope ratios of isolated grains and AOA

378 Table 2 shows the oxygen isotope ratios of three isolated grains and AOA. Four analyses of FeO-rich olivine grains G32 and G33 are homogeneous with Δ 379 ^{17}O values of $-2.7 \pm 0.5\text{‰}$ and $-1.9 \pm 0.3\text{‰}$, respectively (Fig. 4b-c). These analyses are in agreement with type II chondrules C17 381 and C22. In FeO-poor olivine G24, 8 analysis of olivine are widely distributed close to PCM line with $\delta^{18}\text{O}$ and δ 382 ^{17}O values from -3.2‰ to $+3.3\text{‰}$ and from -5.5‰ to $+1.7\text{‰}$, respectively. The range of $\Delta^{17}\text{O}$ values is from -4‰ to 0‰ . As shown in Fig. 4d, the core of G24 is ^{16}O -poor (Δ 383 ^{17}O : 384 -0.8‰ to 0.0‰) compared to the coarse rim (-2‰ to -4‰).

385 We analyzed four spots in olivine from AOA I3. While we rejected two analyses that show numerous large cavities in their SIMS spots (EA2), the mean of $\delta^{18}\text{O}$, $\delta^{17}\text{O}$, and Δ 386 ^{17}O values of the 387 two remaining analyses of olivine in AOA I3 are $-45.5 \pm 0.5\text{‰}$, $-47.3 \pm 0.4\text{‰}$, and $-23.7 \pm 0.2\text{‰}$, 388 respectively.

17

389

390 3.3. Chondrule Mg#

391

392 Following the method of Ushikubo et al. (2012) and Tenner et al. (2013, 2015, 2017), we 393 calculated the “host chondrule Mg#” in taking a mean value of olivine and/or low-Ca pyroxene 394

excluding relict grains of each chondrule investigated. These values are shown in Table 1. 395

Uncertainties of host chondrule Mg# are defined so that they represent the range of Mg#s in each 396
chondrule, by taking differences between maxima, or minima, and the mean Mg# of olivine and 397
low-Ca pyroxene, respectively. The host chondrules Mg#s of type I chondrules show a narrow 398 range
between 98.6 and 99.6, except for the three type I PP chondrules C31, C11 and C12, with 399 Mg# of
97.3, 93.3 and 93.6, respectively. The host chondrule Mg#s for the type II chondrules are 400 calculated
to be ~70 by excluding obvious relict forsteritic olivine.

401

402

403 4. DISCUSSION

404

405 4.1. A single or two separate isotope reservoir(s)?

406

The Δ 407 ^{17}O values of host chondrules and isolated olivine grains ($n=30$) are shown in Fig. 6 in the
sequence of decreasing Δ 408 ^{17}O values from $-1.9 \pm 0.3\text{‰}$ to $-7.0 \pm 0.2\text{‰}$. The majority of type I
chondrules show a narrow range of Δ 409 ^{17}O between -6‰ and -4‰ , while all FeO-rich chondrules
and isolated olivine grains have Δ 410 ^{17}O between -3‰ and -2‰ . This is very similar to the results
from Murchison, where the Δ 411 ^{17}O values of high Mg# (>98) chondrules range from -6‰ to -4‰

18

412 and those of lower Mg# (96-65) chondrules are between -3‰ and -2‰ (Chaumard et al., 2018).

413 Compared to Murchison data, Paris chondrule data are more continuous, without a gap between $-$
 4‰ and -3‰ and extend to lower Δ 414 $^{17}\text{O} \sim -7\text{‰}$.

In Fig. 7, the host chondrule $\Delta 415$ ^{17}O values in Paris are shown against their Mg#s along with 416 data from Murchison (Chaumard et al., 2018). The majority of type I chondrules in Paris plot at highest Mg#s ~ 99 with $\Delta 417$ ^{17}O 418 values increase with decreasing Mg#s. In Murchison, chondrules with high Mg#s (>98.5) display host $\Delta 419$ ^{17}O values ranging from -6.0‰ to -4.1‰ while chondrules with lower Mg# ($\sim 96\text{--}65$) have host $\Delta 420$ ^{17}O values of $\sim -2.5\text{‰}$. Thus, the chondrule Mg#- $\Delta 420$ ^{17}O relationship in Paris is nearly the 421 same as that observed for Murchison. Indeed, the two isolated FeO-rich olivine grains analyzed in 422 Paris plot along and extend the trend defined by type II chondrules (in both Paris and Murchison), 423 towards lower Mg#s (~ 37 for the grain G33) than was previously measured in Murchison. One PP (C31) and one POP (C18) chondrule with host $\Delta 424$ ^{17}O values of -3.3‰ also plot between Mg# $\sim 96\text{--}99$ and two other PP chondrules (C11, C12) with $\Delta 425$ ^{17}O values of $\sim -2.5\text{‰}$ have Mg#s $\sim 93\text{--}94$, a 426 range of Mg#s not found in Murchison (Chaumard et al., 2018). Thus, by combining data from two CM chondrites, CM chondrule data define a single continuous trend of increasing $\Delta 427$ ^{17}O values with decreasing Mg#s. Marrocchi et al. (2018) also reported similar host chondrule $\Delta 428$ ^{17}O values 429 and Mg# data for three chondrules in NWA 5958, which is a CM-related ungrouped carbonaceous 430 chondrite.

The Mg#- $\Delta 431$ ^{17}O relationship among chondrules observed in CM chondrites is similar to 432 those obtained from Acfer 094 (Ushikubo et al., 2012) and the Y-81020 CO chondrite (Tenner et al., 433 al., 2013), as shown in Fig. 8, and to a lesser extent for chondrules in the Y-82094 ungrouped 434 carbonaceous chondrite and CV chondrites (Tenner et al., 2017; Hertwig et al., 2018; 2019a).

^{17}O at $\sim -5\text{‰}$ and -2.5‰ (Fig. 8). Ushikubo et al. (2012) and Tenner et al. (2013) argued that these chondrules formed in two distinct isotope reservoirs; (1) a reducing environment at lower dust enrichments ($100 \times$ solar composition gas) in which chondrules with $\text{Mg\#} > 97$ and $\Delta^{17}\text{O} \sim -5\text{‰}$ formed and (2) an oxidizing environment showing higher dust enrichments ($1000 \times$ solar) in which chondrules with $\text{Mg\#} < 97$ chondrules and $\Delta^{17}\text{O} \sim -2.5\text{‰}$ formed. They interpreted the difference in oxygen isotope ratios between the two reservoirs as a result of varying amounts of ^{16}O -poor H_2O ice mixed in with other chondrule precursor solids.

Chondrule data from Acfer 094 and Y-810202 overlap almost exactly with the CM chondrule trend (Fig. 8), so that the two separate isotope reservoirs indicated from earlier studies could represent subsets of a single chondrule-forming region, which could have been shared by many CC chondrules. Chondrules in the ungrouped carbonaceous chondrite Y-82094 also show a similar $\text{Mg\#}$ - $\Delta^{17}\text{O}$ relationship that overlaps with that of CM chondrites. In CV chondrites, chondrules generally have high $\text{Mg\#}$ s (> 98) with $\Delta^{17}\text{O}$ values predominantly ranging from -6‰ to -4‰ (Hertwig et al., 2018; 2019a; Marrocchi et al., 2019). These values overlap exactly with the CM chondrule data for $\text{Mg\#} > 98$. In all these CCs, type I chondrules with $\text{Mg\#} \leq 97$ and $\Delta^{17}\text{O}$ values between -4‰ and -3‰ are scarce. The Paris chondrule C31 (PP) with $\text{Mg\#} \sim 97.3$ ($+0.5/-$

1.6) shows intermediate $\Delta^{17}\text{O} = -3.3 \pm 0.2\text{‰}$ and is located in the gap between the two groups of chondrules ($\Delta^{17}\text{O} \sim -5\text{‰}$ and $\sim -2.5\text{‰}$ on the $\Delta^{17}\text{O}$ versus $\text{Mg\#}$ plot, Fig. 8), so is one of the three chondrules studied by Marrocchi et al. (2018) in NWA 5958 with $\text{Mg\#} \sim 95$ and $\Delta^{17}\text{O}$ of -3‰ .

Identifying more chondrules with intermediate $\text{Mg\#}$ s and $\Delta^{17}\text{O}$ values would further test if CC chondrules derived from a single continuous region, or if they were derived from two separate

reservoirs in terms of time and locations, as argued earlier from Acfer 094 and CO chondrite chondrule data (Ushikubo et al., 2012; Tenner et al., 2013).

459

4.2. Nature of the oxygen isotope reservoir(s)

461

Tenner et al. (2015) presented a mass balance model to describe the observed negative correlation between $\Delta^{17}\text{O}$ values and Mg#s among CR chondrules. As discussed by Tenner et al. (2015), Mg#s of olivine and pyroxene in chondrules mainly depend on the redox state of iron that in turn was controlled by the oxygen fugacity of the chondrule-forming environment. Equilibrium thermodynamic calculations (e.g., Ebel and Grossman 2000) indicate that type I chondrules with high Mg#s formed under reducing conditions, e.g., $f\text{O}_2 \sim 3.5$ to ~ 2.5 log units below the IW buffer for Mg# 99 and 96, respectively (Tenner et al., 2015). In the protoplanetary disk, H_2O ice would reside in fine-grained dust particles enriched in organic matter and amorphous silicates, similar to chondritic porous interplanetary dust particles and the matrix in primitive meteorites (e.g., Bradley and Brownlee, 1986; Abreu et al., 2010). Therefore, the model supposes that the chondrule forming region consisted of varying relative proportions of these components. In the scope of the model, H_2O ice, anhydrous dust, organics, and solar-composition gas possessed $\Delta^{17}\text{O}$ values of 5.1‰, -5.9‰, 11.3‰, and -28.4‰, respectively (Tenner et al., 2015).

The model assumes that chondrules formed in an environment whose effective oxygen isotopic composition and redox state is mostly controlled by the mixture of these four components. During transient heating, this environment comprised the solar-composition gas enriched in the evaporated

portions of the precursors (= ambient gas), a melt component, and unmelted solids.

479 Being a mixing model, no predictions are made about the actual physical processes operating

21
480 during chondrule formation. It is implied by the model, however, that after evaporation, 481
recondensation into the melt (e.g., Libourel et al., 2006; Nagahara et al., 2008; Libourel and Portail, 482
2018) lead to an effective oxygen exchange between melt and ambient gas (e.g., Kita et al., 2010; 483
Ushikubo et al., 2012; Tenner et al., 2015; Marrocchi and Chaussidon, 2015; Marrocchi et al., 2019), so
that melt and ambient gas display the same effective Δ 484 ^{17}O value at the time of olivine 485
crystallization. The Mg#s of olivine and pyroxene in chondrules would be determined by the
486 oxygen fugacity of the ambient gas, which can be estimated from the relative abundances of H, C,
487 and O of the mixture of ambient gas and the melt component. Further details of the mass balance
488 model are described in Tenner et al. (2015; 2018).

489 Using the model calculation, Tenner et al. (2015) suggested that the majority of type I 490
chondrules in CR chondrites formed at a dust enrichment factor of 100-200 relative to solar 491
composition gas and with a H_2O -ice enhancement of 0-0.8 times relative to CI-composition dust.
Subsequently, Hertwig et al. (2018) modified the model parameters by applying Δ 492 ^{17}O values of 493
the anhydrous dust and H_2O ice of -8‰ and $+2\text{‰}$, respectively, in order to explain the distribution of
Mg#s and Δ 494 ^{17}O values of chondrules in CV chondrites. These parameters were estimated based on
the assumptions that (1) the lowest host-chondrule Δ 495 ^{17}O value is representative of the anhydrous
silicate dust component and (2) the average $\Delta^{17}\text{O}$ value of type II chondrules constrain the Δ 496 ^{17}O
497 values of H_2O ice based on the oxygen isotope mass balance at high dust enrichments (Tenner et al.,
2015). In Paris, the lowest host Δ 498 ^{17}O value measured is -7.0‰ , while the type II chondrules and
FeO-rich olivine G33 reach $\Delta^{17}\text{O}$ values as high as $\sim -2\text{‰}$. Consequently, the Δ 499 ^{17}O values for 500

anhydrous silicate dust and H₂O ice used by Hertwig et al. (2018) are useful to estimate both the 501 dust and H₂O enrichments of the precursors during the formation of CM chondrules. Applying the 502 model of Tenner et al. (2015) and the calculations of Hertwig et al. (2018), Mg# ~99 chondrules

22

503 in Paris correspond to dust enrichment factors of 50-100× relative to solar-composition gas and to 504 amounts of H₂O in the dust ranging from 0 (anhydrous dust) to 0.8× relative to the nominal 505 abundance of H₂O ice in the CI dust (Fig. 9). For chondrules with Mg#s lower than 98 and down 506 to mostly 60-70, the ice enrichment factor is roughly constant (between 0.8 and 1× the nominal 507 abundance of H₂O ice, relative to the CI dust) while the dust enrichment factor increases from 508 ~100× to ≥1,000× (Fig. 9). We note that Mg# should be controlled not only by oxygen fugacity, 509 but also by iron abundance of the system relative to Mg and Si, which is assumed to be CI 510 chondritic in the model. Chondrules with lower Mg# (<60 down to ~35) might have been formed 511 from precursors that were significantly enriched in iron and may not represent chondrule formation 512 under extremely high dust-enrichments (>3,000×).

The observed relationship between Δ 513 ¹⁷O values and Mg#s from chondrules in CM 514 chondrites might have resulted from their formation in a single large region in the protoplanetary disk that was radially zoned with respect to the effective Δ 515 ¹⁷O values, possibly due to variable abundances of 516 ¹⁶O-poor H₂O ice. The region would likely have existed near to the snow line, the condensation front of H₂O ice. Inside the snow line, the relative abundance of 517 ¹⁶O-poor H₂O ice 518 among the solid chondrule precursors was low in contrast to outside of the snow line (e.g., 519 Morbidelli et al., 2016). The most reduced chondrules (Mg#s >99) formed inside the snow line, 520 within this large region, where dust enrichment factors were up to ~100×. More oxidized chondrules

with higher $\Delta 521$ ^{17}O values and lower Mg#s would have formed towards the external part of this single chondrule-forming region where there was addition of ^{16}O -poor H_2O ice to the nearly anhydrous chondrule precursors and/or an increase of the dust enrichment factor from $\sim 100\times$ to $\sim 1,000\times$, as proposed for CR chondrites by Tenner et al. (2015).

23

Alternatively, systematic changes in Mg#- $\Delta 525$ ^{17}O could have developed over time. Most type I chondrules with Mg#>97 and lower $\Delta 526$ ^{17}O values $\sim -5\text{‰}$ formed in regions mainly inside of the snow line with dust enrichments lower than $\sim 200\times$. The local disk temperatures would decrease with time and, consequently, the snow line could migrate through the chondrule forming regions. At the same time, the dust layer became thicker prior to the formation of asteroidal bodies (e.g., Alexander et al., 2008). Later-forming chondrules would form under oxidizing environments and with higher $\Delta^{17}\text{O}$ values $\sim -2\text{‰}$ due to addition of ^{16}O -poor H_2O ice to chondrule precursors. Hartlep and Cuzzi (2020) estimated dust enrichment factors of the turbulent disk midplane to be typically $\sim 100\times$ where cm-sized pebbles formed by streaming instability. They further discussed that chondrules and their precursors existed in the form of cm-sized pebbles and argued that the estimated particle concentrations are consistent with those of high Mg# chondrules, which are abundant in carbonaceous chondrites. Their model also predicts that particle concentrations may reach $\sim 1,000\times$, but probability is low. Therefore, in principle, formation of type II chondrules under higher dust-enrichment factors of $\sim 1,000\times$ might have occurred in the protoplanetary disk. Many type I POP chondrules show a mineralogical zoning with olivine grains being located in the core and pyroxene in periphery of individual chondrules (e.g., Krot et al., 2004; Hezel et al., 2006; Friend et al., 2016). Further, Villeneuve et al. (2020) observed a large mass-dependent Si isotope

fractionation in type I chondrules, especially in those with high Mg#s. Both observations 543 provide evidence for SiO molecules condensing from the ambient gas to the melt. Based on 544 detailed CL-mapping of chondrule minerals and correlated oxygen isotope systematics, Marrocchi et al. (2019) proposed that SiO (along with Mg atoms) condensing from an 545 ^{16}O -poor nebula gas ($\Delta^{17}\text{O} \sim 0\text{‰}$) interacted with ^{16}O -rich AOA-like chondrule precursors ($\Delta^{17}\text{O} \sim -20\text{‰}$) to form type I chondrules with $\Delta^{17}\text{O}$ values ranging from -6‰ to -3‰ . However, 2-3 Ma after CAIs, ambient

24

548 temperatures of the protoplanetary disk should have been significantly lower than the condensation 549 temperature of Si (e.g., Desch et al., 2018). Hence, the partial pressure of SiO in the nebula gas is 550 expected to be low, prior to the heating events causing chondrule formation. Instead, heating and 551 evaporation of solid precursors, such as silicates, during chondrule formation would produce the 552 majority of gaseous SiO, residing in the ambient gas (e.g., Nagahara et al., 2008) and being 553 available for exchange and interaction with the chondrule melt and solids. Thus, the observed correlation of $\Delta^{17}\text{O}$ values and Mg#s is better explained by the contribution of 554 ^{16}O -poor H_2O ice (or icy fine-grained dust) rather than addition of 555 ^{16}O -poor SiO gas from nebula gas.

556

557 4.3. Origin of relict grains in CM chondrules

558

559 Some chondrules contain grains predating the host minerals that crystallized from the final 560 chondrule melts. These grains are defined as relict grains and can be identified chemically and/or 561 isotopically (e.g., Jones et al., 2004; Kunihiro et al., 2004; Krot et al., 2006; Rudraswami et al., 562 2011; Ushikubo et al., 2012; Schrader et al., 2013; Tenner et al., 2013; Schrader and Davidson, 563 2017; Hertwig et al., 2018; 2019a; Chaumard et al., 2018; Marrocchi et al., 2018; 2019). Ten of 564 the

27 chondrules in this study contain relict olivine and/or low-Ca pyroxene grains (Table 1, Fig. 565 6). The abundance of relict-grain bearing chondrules (~37%) in Paris is similar to Murchison 566 (CM2) (~38%; Chaumard et al., 2018), Acfer 094 (~ 45%; Ushikubo et al., 2012), and the Y-81020 567 CO chondrite (~ 42%; Tenner et al., 2013). We note that the abundance of relict-grain bearing 568 chondrules in these chondrites is probably underestimated due to the limited number of analyses 569 per chondrules (≤ 8), in comparison to the 30-50 analyses per chondrules performed in other studies 570 (Marrocchi et al., 2018; 2019). While eight out of the nine relict grains in Murchison chondrules

25

were 571 ^{16}O -rich compared to their host chondrules (Chaumard et al., 2018), four out of 10 chondrules in Paris contain relict olivine grains with Δ 572 ^{17}O values higher than their host chondrules, which 573 reach as high as 0‰. This difference might be due to selection bias for chondrules in Paris in this work (or in Murchison in previous work) and may not be statistically significant. Similar 574 ^{16}O -poor 575 relict olivine grains have also been reported from other CCs but seem to be not as common 576 (Ushikubo et al., 2012; Tenner et al., 2013; Hertwig et al., 2018; 2019a; Marrocchi et al., 2018; 577 Schrader et al., 2020).

578 In type II chondrules, forsteritic cores in FeO-rich olivine phenocrysts are easily recognized 579 as relict grains (e.g., Jones, 1990; Wasson and Rubin, 2003; Ruzicka et al., 2007, 2008). These 580 observations indicate that at least a part of the type II chondrule precursors formed in more 581 reducing conditions, similar to those during formation of type I chondrules (e.g., Ruzicka et al., 582 2008). Based on re-heating experiments of type I precursor materials at 1450–1500°C under 583 oxidizing conditions (between the IW and NNO buffers), Villeneuve et al. (2015) proposed that 584 type I chondrules (or fragments) could have been the main precursor material of type II chondrules. These results and the recognition of ^{16}O -rich relict grains in the type II chondrule C17 with Δ 585 ^{17}O 586 values similar to

those of host type I chondrules (Fig. 6) support this precursor origin of type II 587 chondrules. There are many other examples where type II chondrules enclose FeO-poor olivine 588 grains that are similar to those observed in type I chondrules (Kunihiro et al. 2004; 2005; Ushikubo 589 et al., 2012; Tenner et al. 2013; 2017; Krot and Nagashima, 2017; Krot et al., 2018). Villeneuve et al. (2020) reported negative δ^{30} 590 Si in relict olivine from type II chondrules in carbonaceous 591 chondrites, indicating they could have originated from forsteritic olivine of type I chondrules.

Thirteen relict grains from 7 chondrules have Δ 592 ^{17}O values within the range of the host values calculated for chondrules (from -7.0‰ to -2.1‰ ; Fig. 6). Including the 593 ^{16}O -rich relict

26

594 olivine in type II chondrules, it had been suggested that most relict grains were result of mixing 595 between two distinct major isotope reservoirs (e.g., Jones et al., 2005; Hewins and Zanda, 2012; 596 Ushikubo et al., 2012; Tenner et al., 2013, 2015). The new dataset for Paris chondrules instead 597 may point to mixing within a single chondrule-forming region. Within this region, the radial 598 transport of solids would have occurred (Cuzzi et al., 2010) between an inner area, enriched in reduced and ^{16}O -rich chondrules, and an outer area, enriched in more oxidized and 599 ^{16}O -poor 600 chondrules.

The remaining relict grains are significantly ^{16}O -rich ($\Delta^{17}\text{O}$: -13‰ and -9.5‰) or 601 ^{16}O depleted ($\Delta^{17}\text{O}$: from -1.9‰ to 0.2‰ ; Fig. 6). Similar 602 ^{16}O -rich relict grains were also reported in Murchison chondrules, reaching values of down to -18‰ (Chaumard et al., 2018). The 603 ^{16}O -rich 604 relict grains might be related to CAI and AOA-like refractory precursors (Ushikubo et al., 2012; 605 Marrocchi et al., 2019), which formed during the earliest stage of the Solar System evolution (≤ 0.2 606 Ma; Kita et al., 2013). However, relict olivine grains in some chondrules (C1 and C14) do neither 607 show any special textural nor major-element compositional difference compared to other 608 coexisting

(non-relict) olivine in the same chondrules. Identification of small differences in minor elements would require detailed elemental mapping of olivine at a high electron-beam intensity, as demonstrated by Marrocchi et al. (2018; 2019) who found that relict grains tend to have lower Ca, Ti, and Al concentrations. In contrast, chondrules C8 and C9 and the isolated olivine grain G24 all show dusty olivine textures and $\Delta^{17}\text{O}$ values $\sim 0\text{‰}$, which are significantly higher than other olivine analyses in the same objects (Table 1). Schrader et al. (2020) reported ^{16}O -poor dusty olivine with $\Delta^{17}\text{O} \sim 0\text{‰}$ in four type I chondrules from Murchison and Murray CM chondrites and suggested that precursors of dusty olivine in CM include those unrelated to type I and II chondrules, but originated instead from an unequilibrated ordinary chondrite source. Dusty olivine

27

grains are thought to form by the reduction during partial melting of more FeO-rich olivine, derived from a previous generation of chondrules (Nagahara, 1981; Rambaldi, 1981; Jones and Danielson, 1997; Leroux et al., 2003). Consequently, their oxygen isotope ratios likely reflect those of the FeO-rich precursors. Our results from C8, C9 and G24 are consistent with those of Schrader et al. (2020) and imply an origin of these relict grains from a different reservoir than the one in which type I ($\Delta^{17}\text{O}: \sim 5\text{‰}$) and type II ($\Delta^{17}\text{O}: \sim 2.5\text{‰}$) chondrules in CM and other groups of CCs formed. The ^{16}O -poor relict grains may be related to OC-like chondrules in CCs that show homogeneous oxygen isotope ratios with $\Delta^{17}\text{O} \sim 0\text{‰}$ (e.g., Tenner et al., 2017). Tenner et al. (2017) argued that three type II chondrules in Y-82094 (ungroup C) are characterized by intermediate Mg#s (80-90), higher MnO contents in olivine, and oxygen isotope ratios on the terrestrial fractionation line but to the left side of the PCM line, which are similar to chondrules in ordinary chondrites. Recently, Williams et al. (2020) and Schneider et al. (2020) obtained coordinated $\epsilon^{54}\text{Cr}$, $\epsilon^{50}\text{Ti}$, and SIMS oxygen isotope analyses of individual chondrules from multiple chondrite groups and found

that chondrules generally show isotope signatures similar to those of their host bulk meteorites. However, Williams et al. (2020) also found three BO chondrules in Allende (CV) with Mg#s of 80-90 that show negative $\epsilon^{54}\text{Cr}$ and $\epsilon^{50}\text{Ti}$ values and $\Delta^{17}\text{O} = 0\text{‰}$, which are very different from those of the bulk Allende meteorite. These chondrules do not exactly match ordinary chondrites, but show similar isotope ratios to achondrites.

The Al-Mg ages of two OC-like chondrules in Acfer 094 are older by 0.4-0.8 Ma than other chondrules in the same meteorite (Ushikubo et al., 2013; Hertwig et al., 2019b). The age difference could represent the transit time for these chondrules to travel from the outside of the CC chondrule forming regions to the Acfer 094 accretion region. Schrader et al. (2020) and Williams et al. (2020)

28

suggested that dusty olivine in CM chondrite chondrules and the FeO-rich BO chondrules in Allende, respectively, had migrated outward from inner to outer disk across the Jupiter's gap. Collectively, both refractory ^{16}O -rich precursors and OC-like ^{16}O -poor precursors represent inner disk solids that would have been added to the solid precursors of CM chondrules.

644

645 4.4. Oxygen isotope ratios of AOA in CM

646 Here we compare oxygen isotope ratios of AOA I3 in Paris obtained during this work to AOAs in other CCs and refractory inclusions in CM chondrites, which were previously obtained in the WiscSIMS laboratory. The mean of 2 analyses of AOA I3 gives $\delta^{18}\text{O} = -45.5 \pm 0.5\text{‰}$, $\delta^{17}\text{O} = -47.3 \pm 0.4\text{‰}$, and $\Delta^{17}\text{O} = -23.7 \pm 0.2\text{‰}$ (Table 1). This is in agreement with AOA analyses in Acfer 094 (mean of four AOA and 2SD; $\delta^{18}\text{O} = -45.8 \pm 0.7\text{‰}$, $\delta^{17}\text{O} = -47.7 \pm 0.7\text{‰}$, and $\Delta^{17}\text{O} = -23.8 \pm 0.6\text{‰}$; Ushikubo et al., 2017) and those in DOM 08006 (CO3.0; $\delta^{18}\text{O} = -45.8 \pm 0.7\text{‰}$ and $\Delta^{17}\text{O} = -24.0 \pm 0.4\text{‰}$; Fukuda et al., 2021). Data from I3 are also consistent with spinel-hibonite

inclusions (SHIBs) in Murchison studied by Kööp et al. (2016) with $\Delta 653 \text{ }^{17}\text{O} = -23.4 \pm 1.1\%$. Other studies reporting oxygen isotope ratios of AOA from multiple CCs (Fagan et al., 2004; Krot et al., 2005; 2014; Komatsu et al., 2017), Kakangari (Nagashima et al., 2015), ordinary chondrites (Itoh et al., 2007; Ebert et al., 2020), and enstatite chondrite (Guan et al., 2000) also show similar ranges. Thus, AOA I3 formed in a homogeneous ^{16}O -rich isotope reservoir that formed a wide variety of CAIs (e.g., Ushikubo et al., 2017).

659

660 4.5. Implications for the chondrule formation region

661 The large chondrule formation region discussed in 4.2. would likely be located at around 3 AU, where the snow line would be located at the time of chondrule formation (a few Ma after

29

663 CAIs; Morbidelli et al., 2016). This region would have been outside of the OC chondrite accretion region, so that a minor fraction of inner disk materials might have been transported from the inner disk to the outer disk (Hertwig et al., 2018; Schrader et al., 2020; Williams et al., 2020). Several recent studies suggest that CM and other CCs form outside of proto-Jupiter in order to explain the isotope dichotomy between carbonaceous meteorites and non-carbonaceous meteorites (e.g., Kruijer et al., 2017; Desch et al., 2018; Nanne et al., 2019; Van Kooten et al., 2020), a hypothesis first advocated based on bulk meteorite nucleosynthetic anomalies of ^{54}Cr and ^{50}Ti by Warren (2011). Kruijer et al. (2017) further suggested that the isolation of carbonaceous and non carbonaceous isotope reservoirs occurred early, within 1 Ma after CAI formation. This predates the events forming CC chondrules which are estimated to have occurred 2-3 Ma after CAI formation (e.g., Ushikubo et al., 2013; Nagashima et al., 2017; Hertwig et al., 2019b). Desch et al. (2018) proposed a comprehensive evolutionary model involving the proto-Jupiter's gap that predicts a

location and formation time for each meteorite parent body in order to explain the 676 chemical diversity among meteorites. In their model, Jupiter originally formed ≥ 4 AU and 677 migrated inward to 3 AU to open a gap that separated the two isotope reservoirs. Within this 678 framework, chondrules in CCs should form beyond the snow line, which is in contrast to the dry environments expected for the abundant CM chondrules with high Mg# (>98) and Δ 679 $^{17}\text{O} \sim -5\text{‰}$. 680 Furthermore, complete isolation of carbonaceous and non-carbonaceous meteorite forming regions 681 can not explain chondrules and relict grains with OC-like oxygen isotope ratios in CM chondrites. 682 Schrader et al. (2020) argued that small fragments of UOC chondrules ($<300\text{ }\mu\text{m}$) could migrate 683 outward beyond the Jupiter's gap and so were incorporated to chondrule precursors for CM chondrites. A few chondrules in Paris contain dusty olivine grains with Δ 684 $^{17}\text{O} \sim 0\text{‰}$ (C8, C9, C24), 685 which are also $\sim 300\text{ }\mu\text{m}$ or smaller and could have migrated from UOC regions across the Jupiter's

30

gap. Chondrules with Δ 686 $^{17}\text{O} \sim 0\text{‰}$ in Acfer 094 and Y-82094 (ungrouped C) are also $\leq 300\text{ }\mu\text{m}$ in 687 size, though those in CV are from $\sim 500\text{ }\mu\text{m}$ (Hertwig et al., 2018; 2019) to $\geq 2\text{ mm}$ (Williams et 688 al., 2020).

689 Alternatively, proto-Jupiter could have been located close to the current Jupiter orbit that 690 is outside of the major CC chondrule-forming regions. Recent numerical simulation by Tanaka et 691 al. (2020) indicated that Jupiter-sized giant planets would not experience a significant radial 692 migration as opposed to those indicated in previous studies (e.g., Lin and Papaloizou 1986). If 693 there were radial transport of solids in the protoplanetary disk (e.g., Cuzzi et al., 2010), a small 694 amount of OC-like chondrules and their fragments could have migrated outward to CM and other 695 major CC forming regions. The main drawback in this scenario is that inward drift of significant 696 amounts of dust from the CC forming regions to the OC forming regions might have occurred.

697 This would be in disagreement with the observed distinct isotope signatures between bulk 698
carbonaceous and non-carbonaceous meteorites (e.g., Budde et al., 2016; Kruijer et al., 2017; 699 Kleine
et al. 2020).

700

701 CONCLUSIONS

702

703 In situ SIMS oxygen 3-isotope analyses of 29 chondrules and 3 isolated olivine grains in 704 the
least-altered CM chondrite Paris were performed. The results were used to estimate host chondrule Δ
705 ^{17}O values. By combining this data set with that from Murchison chondrules (Chaumard et al.,
2018), the Mg#- Δ 706 ^{17}O relationship of CM chondrite chondrules was evaluated.

707

(1) Host chondrule Δ 708 ^{17}O values in CM chondrites increase continuously from -7‰ to -2‰ 31
709
with decreasing Mg# from 99 to 37. The majority of types I and II chondrules in CM show high Mg#s
~99 with $\Delta^{17}\text{O}$ from -6‰ to -4‰ and lower Mg#s of 60-70 with Δ 710 ^{17}O of -2.5‰ , respectively,
while a few type I chondrules with lower Mg# (97-93) show Δ 711 ^{17}O 712 between -3.3‰ and -2.5‰ ,
We suggest that bimodal distribution of chondrule Mg# and Δ 713 ^{17}O from Acfer 094 and Y-81020
(CO3) previously reported by Ushikubo et al. (2012)

714 and Tenner et al. (2013), respectively, are parts of the CM chondrule trend. (2) Relict grains in CM
chondrite chondrules are either ^{16}O -rich or 715 ^{16}O -poor relative to their host chondrules, with the total
range from -18‰ to 0‰ . The majority of relict grain Δ 716 ^{17}O values overlap the range of host

chondrule $\Delta 717^{17}\text{O}$ values, suggesting they were derived from a precursor in the same CM chondrite chondrule forming region. Relict grains with $\Delta 718^{17}\text{O}$ values $<-8\text{‰}$ and $\sim 0\text{‰}$ are likely from refractory precursors and OC-like precursors, respectively.

(3) By adapting the oxygen isotope mass balance model of Tenner et al. (2015) and Hertwig et al. (2018), we argue that the majority of type I chondrules formed under relatively low dust enrichment factors ($50\text{--}100\times$ Solar) but with a range of ^{16}O -poor H_2O ice ($0\text{--}0.8\times$ CI) in the precursors. Other chondrules formed under oxidizing environments with higher dust enrichments ($100\text{--}1,000\times$ Solar) and abundant ^{16}O -poor H_2O ice ($0.8\text{--}1\times$ CI).

(4) CM chondrite chondrule formation could have occurred in a large single region that spanned both sides of the snow line, facilitating a wide range of dust-enrichments and ^{16}O poor H_2O -ice enhancements. It is possible that the conditions in this region evolved with time, from reducing to more oxidizing, as the local disk became denser and colder. This

32

chondrule forming region would have likely exist inside the proto-Jupiter orbit if Jupiter was large enough to produce a gap in the accretion disk.

(5) In addition to chondrules, we also analyzed one AOA in Paris, and the data from this AOA are in good agreement with AOAs and other refractory inclusions from Acfer 094, DOM 08006 (CO_3), and Murchison (CM2).

735

Acknowledgements We gratefully thank Brigitte Zanda for allocating us Paris thin sections for the SIMS work. We thank John Fournelle and Jim Kern for assistance with electron probe micro analysis and SIMS instrument, respectively. We thank Michael Spicuzza for his reading of earlier

version of the paper to improving the quality of the manuscript. We thank Kazu Nagashima and 740
Yves Marrocchi for their constructive reviews and Sasha Krot for editorial handling, which 741
improved clarity of the discussions. This work is supported by the NASA Cosmochemistry 742 program
(NNX14AG29G). WiscSIMS is partly supported by NSF (EAR13-55590, EAR 743 1658823).

744

745 REFERENCES

746

747 Abreu N. M. and Brearley A. J. (2010) Early solar system processes recorded in the matrices of 748
two highly pristine CR3 carbonaceous chondrites, MET 00426 and QUE 99177. *Geochim.* 749
Cosmochim. Acta **74**, 1146-1171.

750 Alexander C. M. O'D., Grossman J. N., Ebel D. S., and Ciesla F. J. (2008) The formation 751
conditions of chondrules and chondrites. *Science* **320**, 1617-1619.

Baertschi P. (1976) Absolute 752 ^{18}O content of standard mean ocean water. *Earth Planet. Sci. Lett.*

33

753 **31**, 341-344.

754 Berlin J., Jones R. H., and Brearley A. J. (2011) Fe-Mn systematics of type IIA chondrules in 755
unequilibrated CO, CR, and ordinary chondrites. *Meteor. Planet. Sci.* **45**, 513-533. 756 Bischoff A., Scott
E. R. D., Metzler K., and Goodrich C. A. (2006) Nature and origins of meteoritic 757 breccias. In
Meteorites and the early solar system II, edited by Lauretta D. S. and McSween H. 758 Y. Jr. Tucson,
Arizona: The University of Arizona Press. pp. 679-712. 759 Brearley A. J. (2003) Nebular versus
parent-body processing. In *Meteorites, comets, and planets*, 760 (ed. A. M. Davis). Vol. 1. *Treatise on*
Geochemistry (eds. H. D. Holland, and K. K. Turekian) 761 Elsevier-Pergamon, Oxford. pp. 247-268.

762 Bradley J. P. and Brownlee D. E. (1986) Cometary Particles: Thin Sectioning and Electron Beam

763 Analysis. *Science* **231**, 1542-1544.

764 Briani G., Gounelle M., Bourot-Denise M., and Zolensky M. (2012) Xenoliths and microxenoliths
765 in H chondrites: Sampling of the zodiacal cloud in the asteroid Main Belt. *Meteor. Planet. Sci.* 766
47, 880-902.

767 Budde G., Burkhardt C., Brennecke G. A., Fischer-Gödde M., Kruijer T. S., Kleine T. (2016) 768
Molybdenum isotopic evidence for the origin of chondrules and a distinct genetic heritage of 769
carbonaceous and non-carbonaceous meteorites. *Earth Planet. Sci. Lett.* **454**, 293–303. 770 Busemann
H., Alexander C. M. O'D, and Nittler L. R. (2007) Characterization of insoluble organic 771 matter in
primitive meteorites by microRaman spectroscopy. *Meteor. Planet. Sci.* **42**, 1387- 772 1416.

773 Chakraborty S. (2010) Diffusion coefficients in olivine, wadsleyite, and ringwoodite. In *Reviews*
774 in *Mineralogy and Geochemistry*, vol. 72 (eds. Y. Zhang and D. J. Cherniak). Mineralogical 775
Society of America, Washington, DC. pp. 603-639.

34

776 Chaumard N., Defouilloy C., and Kita N. T. (2018) Oxygen isotope systematics of chondrules in
777 the Murchison CM2 chondrite and implications for the CO-CM relationship. *Geochim.* 778
Cosmochim. Acta **228**, 220-242.

779 Clayton R. N. and Mayeda T. K. (1999) Oxygen isotope studies of carbonaceous chondrites. 780
Geochim. Cosmochim. Acta **63**, 2089-2104.

781 Clayton R. N., Onuma N., Grossman L., and Mayeda T. K. (1977) Distribution of pre-solar 782
component in Allende and other carbonaceous chondrites. *Earth Planet. Sci. Lett.* **34**, 209-224. 783

Connolly, Jr., H. C. and Huss G. R. (2010) Compositional evolution of the protoplanetary disk: 784
oxygen isotopes of type-II chondrules from CR2 chondrites. *Geochim. Cosmochim. Acta* **74**, 785
2473-2483.

786 Cuzzi J. N., Hogan R. C., and Bottke W. F. (2010) Toward initial mass functions for asteroids and
787 Kuiper Belt objects. *Icarus* **208**, 518-538.

788 Desch S. J., Kalyaan A. and Alexander C. M. O'D. (2018) The effect of Jupiter's formation on the
789 distribution of refractory elements and inclusions in meteorites. *Astrophys. J. Suppl. Ser.* **238**, 790
11.

791 Donovan J. J. (2015) Probe for EPMA v. 11.1.5. User's Guide and References. Probe Software, 792
Inc., Eugene, OR.

793 Ebel S. D. and Grossman L. (2000) Condensation in dust-enriched systems. *Geochim. Cosmochim.*
794 *Acta* **64**, 339-366.

795 Ebert S., Nagashima K., Krot A. N., Bischoff A. (2020) Oxygen-isotope heterogeneity in the 796
Northwest Africa 3358 (H3.1) refractory inclusions - Fluid-assisted isotopic exchange on the 797
H-chondrite parent body. *Geochim. Cosmochim. Acta* **282**, 98-112.

798 Eiler J. M., Graham C., Valley J. W. (1997) SIMS analysis of oxygen isotopes: matrix effects in

35

799 complex minerals and glasses. *Chem. Geol.* **138**, 221-244.

800 Fagan T. J., Krot A. N., Keil K. and Yurimoto H. (2004) Oxygen isotopic evolution of amoeboid
801 olivine aggregates in the reduced CV3 chondrites Efremovka, Vigarano, and Leoville. 802
Geochim. Cosmochim. Acta **68**, 2591-2611.

803 Frank D. R., Zolensky M. E., and Le L. (2014) Olivine in terminal particles of Stardust aerogel 804
tracks and analogous grains in chondrite matrix. *Geochim. Cosmochim. Acta* **142**, 240-259. 805 Friend
P., Hezel D. C. and Mucerschi D. (2016) The conditions of chondrule formation, Part II: 806 Open
system. *Geochim. Cosmochim. Acta* **173**, 198-209.

807 Fukuda K., Brownlee D. E., Joswiak D. J., Tenner T. J., Kimura M., and Kita N. T. (2021) 808

Correlated isotopic and chemical evidence for condensation origins of olivine in comet 809 81P/Wild 2
and in AOAs from CV and CO chondrites. *Geochim. Cosmochim. Acta* **293**, 544- 810 574.

811 Gounelle M., Zolensky M. E., Liou J.-C., Bland P. A., and Alard O. (2003) Mineralogy of 812
carbonaceous chondritic microclasts in howardites: Identification of C2 fossil micrometeorites. 813
Geochim. Cosmochim. Acta **67**, 507-527.

814 Grossman L. and Steele I. M. (1976) Amoeboid olivine aggregates in the Allende meteorite. 815
Geochim. Cosmochim. Acta **40**, 149-155.

816 Grossman L., Beckett J. R., Fedkin A. V., Simon S. B., and Ciesla F. J. (2008) Redox conditions
817 in the solar nebula: Observational, experimental, and theoretical constraints. In *Reviews in* 818
Mineralogy and Geochemistry, vol. 68 (ed. G. J. MacPherson). Mineralogical Society of 819 America,
Washington, D.C. pp. 93-140.

820 Guan Y., McKeegan K. D. and MacPherson G. J. (2000) Oxygen isotopes in calcium-aluminum 821
rich inclusions from enstatite chondrites: new evidence for a single CAI source in the solar

36

822 nebula. *Earth Planet. Sci. Lett.* **181**, 271-277.

823 Hartlep T. and Cuzzi J. N. (2020) Cascade Model for Planetesimal Formation by Turbulent 824
Clustering. *Astrophys. J.* **892**, 120.

825 Heck P. R., Ushikubo T., Schmitz B., Kita N. T., Spicuzza M. J. and Valley J. W. (2010) A single
826 asteroidal source for extraterrestrial Ordovician chromite grains from Sweden and China: high 827
precision oxygen three-isotope SIMS analysis. *Geochim. Cosmochim. Acta* **74**, 497-509. 828 Hertwig A,
Defouilloy C., and Kita N. T. (2018) Formation of chondrules in a moderately high 829 dust enriched
disk: Evidence from oxygen isotopes of chondrules from the Kaba CV3 830 chondrite. *Geochim.*
Cosmochim. Acta **224**, 116-131.

831 Hertwig A. T., Kimura M., Defouilloy C. and Kita N. T. (2019a) Oxygen isotope systematics of 832
chondrule olivine, pyroxene, and plagioclase in one of the most pristine CV_{3Red} chondrites 833
(Northwest Africa 8613). *Meteorit. Planet. Sci.* **54**, 2666-2685.

Hertwig A. T., Kimura M., Ushikubo T., Defouilloy C. and Kita N. T. (2019b) The ²⁶Al- 834 ²⁶Mg 835
systematics of FeO-rich chondrules from Acfer 094: Two chondrule generations distinct in age 836 and
oxygen isotope ratios. *Geochim. Cosmochim. Acta* **253**, 111-126. 837 Hewins R. H. and Zanda B. (2012)
Chondrules: Precursors and interactions with the nebular gas. 838 *Meteorit. Planet. Sci.* **47**, 1120-1138.

839 Hewins R. H., Bourot-Denise M., Zanda B., Leroux H., Barrat J.-A., Humayun M., Göpel C., 840
Greenwood R. C., Franchi I. A., Pont S., Lorand J.-P., Cournède C., Gattacceca J., Rochette 841 P.,
Kuga M., Marrocchi Y., and Marty B. (2014) The Paris meteorite, the least altered CM 842 chondrite so
far. *Geochim. Cosmochim. Acta* **124**, 190-222.

843 Hezel D. C., Palme H., Nasdala L., Brenker F. E., (2006) Origin of SiO₂-rich components in 844
ordinary chondrites. *Geochim. Cosmochim. Acta* **70**, 1548-1564.

845 Itoh S., Russell S. S., Yurimoto H. (2007) Oxygen and magnesium isotopic compositions of 846
amoeboid olivine aggregates from the Semarkona LL3.0 chondrite. *Meteorit. Planet. Sci.* **42**, 847
1241-1247.

848 Jones R. H. (1990) Petrology and mineralogy of type-II, FeO-rich chondrules in Semarkona 849
(LL3.0) – origin by closed-system fractional crystallization, with evidence for supercooling. 850
Geochim. Cosmochim. Acta **54**, 1785-1802.

851 Jones R. H. (1994) Petrology of FeO-poor, porphyritic pyroxene chondrules in the Semarkona 852
chondrite. *Geochim. Cosmochim. Acta* **58**, 5325-5340.

853 Jones R. H. (2012) Petrographic constraints on the diversity of chondrule reservoirs in the 854

protoplanetary disk. *Meteorit. Planet. Sci.* **47**, 1176-1190.

855 Jones R. H. and Danielson L. R. (1997) A chondrule origin for dusty relict olivine in

856 unequilibrated chondrites. *Meteorit. Planet. Sci.* **32**, 753-760.

857 Jones R. H., Leshin L. A., Guan Y., Sharp Z. D. Durakiewicz, T., and Schilk A. J. (2004) 858

Oxygen isotope heterogeneity in chondrules from the Mokoia CV3 carbonaceous chondrite. 859

Geochim. Cosmochim. Acta **68**, 3423-3438.

860 Jones R. H., Grossman J. N., and Rubin A. E. (2005) Chemical, mineralogical and isotopic 861

properties of chondrules: clues to their origin. In *Chondrules and the Protoplanetary Disk, ASP* 862

Conference Series (eds. A. N. Krot, E. R. D. Scott and B. Reipurth). Sheridan Books, Ann 863 Arbor,

Michigan, pp. 251-285.

864 Kallemeyn G. W. and Wasson J. T. (1981) The compositional classification of chondrites: I. The

865 carbonaceous chondrite groups. *Geochim. Cosmochim. Acta* **45**, 1217-1230. 866 Kimura M.,

Grossman J. N., and Weisberg M. K. (2011) Fe-Ni metal and sulfide minerals in CM 867 chondrites: An

indicator of thermal history. *Meteorit. Planet. Sci.* **46**, 431-442.

38

868 Kimura M., Imae N., Komatsu M., Barrat J.A., Greenwood R.C., Yamaguchi A., Noguchi T. 869

(2020) The most primitive CM chondrites, Asuka 12085, 12169, and 12236, of subtypes 3.0– 870 2.8:

Their characteristic features and classification, *Polar Science* **26**, 100565. Kita N. K. and Ushikubo T.

(2012) Evolution of protoplanetary disk inferred from 871 ²⁶Al chronology 872 of individual

chondrules. *Meteorit. Planet. Sci.* **47**, 1108-1119.

873 Kita N. T., Ushikubo T., Fu B. and Valley J. W. (2009) High precision SIMS oxygen isotope 874

analysis and the effect of sample topography. *Chem. Geol.* **264**, 43-57. 875 Kita N. T., Nagahara H.,

Tachibana S., Tomomura S., Spicuzza M. J., Fournelle J. H. and Valley 876 J. W. (2010) High precision

SIMS oxygen three isotope study of chondrules in LL3 chondrites: 877 role of ambient gas during
chondrule formation. *Geochim. Cosmochim. Acta* **74**, 6610-6635. 878 Kita N. T., Yin Q.-Z., MacPherson
G. J., Ushikubo T., Jacobsen B., Nagashima K., Kurahashi E., Krot A. N. and Jacobsen S. B. (2013)
 ^{26}Al – ^{26}Mg isotope systematics of the first solids in the 880 early solar system. *Meteorit. Planet.*
Sci. **48**, 1383-1400.

881 Kleine T., Budde G., Burkhardt C., Kruijer T. S., Worsham E. A., Morbidelli A., Nimmo F. (2020)
882 the Non-carbonaceous–Carbonaceous Meteorite Dichotomy. *Space Sci. Rev.* **216**, 55. 883 Komatsu
M., Fagan T. J., Krot A. N., Nagashima K., Petaev M. I., Kimura M. and Yamaguchi A. 884 (2018) First
evidence for silica condensation within the solar protoplanetary disk. *Proc. Natl. Acad. Sci. USA*
115, 7497-7502.

886 Kööp L., Nakashima D., Heck P. R., Kita N. T., Tenner T. J., Krot A. N., Nagashima K., Park C.,
and Davis A. M. (2016) New constraints on the relationship between ^{26}Al and oxygen, calcium, 888
and titanium isotopic variation in the early Solar System from a multielement isotopic study 889 of
spinel-hibonite inclusions. *Geochim. Cosmochim. Acta* **184**, 151-172.

890 Krot A. N. and Nagashima K. (2017) Constraints on mechanisms of chondrule formation from

891 chondrule precursors and chronology of transient heating events in the protoplanetary disk. 892
Geochim. J. **51**, 45-68.

893 Krot A. N., Libourel G., Goodrich C., Petaev M. I. (2004) Silica-igneous rims around magnesian
894 chondrules in CR carbonaceous chondrites: evidence for fractional condensation during 895
chondrule formation. *Meteorit. Planet. Sci.* **39**, 1931-1955.

896 Krot A. N., Fagan T. J., Nagashima K., Petaev M. I. and Yurimoto H. (2005) Origin of low-Ca 897
pyroxene in amoeboid olivine aggregates: Evidence from oxygen isotopic compositions. 898 *Geochim.*

Cosmochim. Acta **69**, 1873-1881.

899 Krot A. N., Yurimoto H., McKeegan K. D., Leshin L., Chaussidon M., Libourel G., Yoshitake M.,
900 Huss G. R., Guan Y., and Zanda B. (2006) Oxygen isotopic compositions of chondrules: 901
implications for evolution of oxygen isotopic reservoirs in the inner Solar nebula. *Chemie der* 902 *Erde –*
Geochem. **66**, 249-276.

903 Krot A. N., Park C., Nagashima K. (2014) Amoeboid olivine aggregates from CH carbonaceous
904 chondrites. *Geochim. Cosmochim. Acta* **139**, 131-153.

905 Krot A. N., Nagashima K., Libourel G., and Miller K. E. (2018) Multiple mechanisms of transient
906 heating events in the protoplanetary disk: Evidence from precursors of chondrules and igneous 907
Ca, Al-rich inclusions. In *Chondrules: Records of the Protoplanetary Disk Processes*. (eds. S. 908 S.
Russell, H. C. Connolly Jr., and A. N. Krot) Cambridge University Press, U.K., pp. 11-57.

909 Kruijer T. S., Burkhardt C., Budde G., Kleine T. (2017) Age of Jupiter inferred from the distinct 910
genetics and formation times of meteorites. *Proc. Natl. Acad. Sci. USA* **114**, 6712-6716. 911 Kunihiro T.,
Rubin A. E., McKeegan K. D. and Wasson J. T. (2004) Oxygen-isotopic compositions 912 of relict and
host grains in chondrules in the Yamato 81020 CO3.0 chondrite. *Geochim.* 913 *Cosmochim. Acta* **68**,
3599-3606.

40

914 Kunihiro T., Rubin A. E. and Wasson J. T. (2005) Oxygen-isotopic compositions of low-FeO 915
relicts in high FeO-host chondrules in Acfer 094, a type 3.0 carbonaceous chondrite closely 916 related
to CM. *Geochim. Cosmochim. Acta* **69**, 3831-3840.

917 Leroux H., Libourel G., Lemelle L., Guyot F. (2003) Experimental study and TEM 918
characterization of dusty olivines in chondrites: Evidence for formation by in-situ reduction. 919
Meteorit. Planet. Sci. **38**, 81-94.

920 Leroux H., Cuvillier P., Zanda B., and Hewins, R. H. (2015) GEMS-like material in the matrix of
921 the Paris meteorite and the early stages of alteration of CM chondrites. *Geochim. Cosmochim. Acta* **170**, 247-265.

923 Libourel G. and Chaussidon M. (2011) Oxygen isotopic constraints on the origin of Mg-rich
olivines from chondritic meteorites. *Earth Planet. Sci. Lett.* **301**, 9-21. 925 Libourel G. and Portail M.
(2018) Chondrules as direct thermo- chemical sensors of solar 926 protoplanetary disk gas. *Sci. Adv.* **4**,
eaar3321.

927 Libourel G., Krot A. and Tissandier L. (2006) Role of gas-melt interaction during chondrule 928
formation. *Earth Planet. Sci. Lett.* **251**, 232-240.

929 Lin D. N. C. and Papaloizou J. (1986) On the tidal interaction between protoplanets and the 930
protoplanetary disk. III—Orbital migration of protoplanets. *Astrophys. J* **309**, 846-857. 931 Marrocchi Y.
and Chaussidon M. (2015) A systematic for oxygen isotopic variation in meteoritic 932 chondrules.
Earth Planet. Sci. Lett. **430**, 308-315.

933 Marrocchi Y., Gounelle M., Blanchard I., Caste F., and Kearsley A.T. (2014), The Paris CM 934
chondrite: Secondary minerals and asteroidal processing. *Meteorit Planet Sci* **49**, 1232-1249. 935

Marrocchi Y., Villeneuve J., Batanova V., Piani L., Jacquet E. (2018) Oxygen isotopic diversity 936 of
chondrule precursors and the nebular origin of chondrules. *Earth Planet. Sci. Lett.* **496**, 132-

41

937 141.

938 Marrocchi Y., Euverte R., Villeneuve J., Batanova V., Welsch B., Ferrière L., Jacquet E. (2019) 939
Formation of CV chondrules by recycling of amoeboid olivine aggregate-like precursors. 940 *Geochim.*
Cosmochim. Acta **247**, 121-141.

941 McSween H. Y. (1979) Alteration in CM carbonaceous chondrites inferred from modal and 942

chemical variations in matrix. *Geochim. Cosmochim. Acta* **43**, 1761-1770. 943 Morbidelli A., Bitsch B., Crida A., Gounelle M., Guillot T., Jacobson S., Johansen A., Lambrechts 944 M., Lega E. (2016) Fossilized condensation lines in the Solar System protoplanetary disk. 945 *Icarus* **267**, 368-376.

946 Nanne J. A. M., Nimmo F., Cuzzi J. N., Kleine T., (2019) Origin of the non- carbonaceous 947 carbonaceous meteorite dichotomy. *Earth Planet. Sci. Lett.* **511**, 44–54. 948 Nagahara H. (1981) Evidence for secondary origin of chondrules. *Nature* **292**, 135–136. 949 Nagahara H., Kita N. T., Ozawa K. and Morishita Y. (2008) Condensation of major elements 950 during chondrule formation and its implication to the origin of chondrules. *Geochim. 951 Cosmochim. Acta* **72**, 1442-1465.

952 Nagashima K., Krot A. N. and Huss G. R. (2015) Oxygen-isotope compositions of chondrule 953 phenocrysts and matrix grains in Kakangari K-grouplet chondrite: Implication to a chondrule 954 matrix genetic relationship. *Geochim. Cosmochim. Acta* **151**, 49-67.

955 Nakamura T. (2005) Post-hydration thermal metamorphism of carbonaceous chondrites. *J. 956 Mineral. Petrol. Sci.* **100**, 260-274.

957 Pignatelli I., Marrocchi Y., Vacher L. G., Delon R., and Gounelle M. (2016) Multiple precursors 958 of secondary mineralogical assemblages in CM chondrites. *Meteorit. Planet. Sci.* **51**, 785–805. 959 Rambaldi E. R. (1981) Relict grains in chondrules. *Nature* **292**, 558-561.

42

960 Rubin A. E. (2013) An amoeboid olivine inclusion (AOI) in CK3 NWA 1559, comparison to AOIs 961 in CV3 Allende, and the origin of AOIs in CK and CV chondrites. *Meteorit. Planet. Sci.* **48**, 962 432-441.

963 Rubin A. E. (2015) An American in Paris: Extent of aqueous alteration of a CM chondrite and the 964 petrography of its refractory and amoeboid olivine inclusions. *Meteorit. Planet. Sci.* **50**, 1595- 965 1612.

966 Rubin A. E., Trigo-Rodríguez J. M., Huber H., and Wasson J. T. (2007) Progressive aqueous 967 alteration of CM carbonaceous chondrites. *Geochim. Cosmochim. Acta* **71**, 2361–382. 968 Rudraswami N. G., Ushikubo T., Nakashima D. and Kita N. T. (2011) Oxygen isotope systematics 969 of chondrules in the Allende CV3 chondrite: high precision ion microprobe studies. *Geochim. 970 Cosmochim. Acta* **75**, 7596-7611.

971 Ruzicka A., Hiyagon H., Hutson M., and Floss C. (2007) Relict olivine, chondrule recycling, and 972 the evolution of nebular oxygen reservoirs. *Earth Planet. Sci. Lett.* **257**, 274-289. 973 Ruzicka A., Floss C., and Hutson M. (2008) Relict olivine grains, chondrules recycling, and 974 implications for the chemical, thermal, and mechanical processing of nebular materials. 975 *Geochim. Cosmochim. Acta* **72**, 5530-5557.

976 Sakamoto N., Seto Y., Itoh S., Kuramoto K., Fujino K., Nagashima K., Krot A. N. and Yurimoto 977 H. (2007) Remnants of the early solar system water enriched in heavy oxygen isotopes. *Science* 978 **317**, 231-233.

979 Schneider J. M., Burkhardt C., Marrocchi Y., Brennecka G., Kleine T. (2020) Early evolution of 980 the solar accretion disk inferred from Cr-Ti-O isotopes in individual chondrules. *Earth Planet. 981 Sci. Lett.* **551**, 116585.

982 Schrader D. L. and Davidson J. (2017) CM and CO chondrites: A common parent body or

983 asteroidal neighbors? Insights from chondrule silicates. *Geochim. Cosmochim. Acta* **214**, 157- 43 984 171.

985 Schrader D. L., Connolly, Jr., H. C., Lauretta D. S., Nagashima K., Huss G. R., Davidson J. and 986 Domanik K. J. (2013) The formation and alteration of the Renazzo-like carbonaceous 987 chondrites II: linking O-isotope composition and oxidation state of chondrule olivine. 988 *Geochim. Cosmochim. Acta*

101, 302-327.

989 Schrader D. L., Nagashima K., Krot A. N., Ogliore R. C., and Hellebrand E. (2014) Variations in
990 the O-isotope composition of gas during the formation of chondrules from the CR chondrites. 991
Geochim. Cosmochim. Acta **132**, 50-74.

992 Schrader D. L., Nagashima K., Krot A. N., Ogliore R. C., Yin Q.-Z., Amelin Y., Stirling C. H.,
Kaltenbach A. (2017) Distribution of 993 ^{26}Al in the CR chondrite chondrule-forming region of 994 the
protoplanetary disk. *Geochim. Cosmochim. Acta* **201**, 275-302.

995 Schrader D. L., Nagashima K., Fu R. R., Davidson J., and Ogliore R. C. (2020) Outward migration
996 of chondrule fragments in the early Solar System: O-isotopic evidence for rocky material 997
crossing the Jupiter Gap? *Geochim. Cosmochim. Acta* **282**, 133-155.

998 Sears D. W. G. and Dodd R. T. (1988) Overview and classification of meteorites. In *Meteorites* 999
and the early solar system. (eds. J. F. Kerridge and M. S. Matthews) University of Arizona 1000 Press,
Tucson, Arizona, pp 3–31.

1001 Stephant A., Remusat L., and Robert F. (2017) Water in type I chondrules of Paris CM chondrite.
1002 *Geochim. Cosmochim. Acta* **199**, 75-90.

Sugiura N. and Fujiya W. (2014) Correlated accretion ages and $\epsilon^{54}\text{Cr}$ of meteorite parent bodies
1004 and the evolution of the solar nebula. *Meteorit. Planet. Sci.* **49**, 772-787. 1005 Tachibana S.,
Nagahara H., Mostefaoui S., and Kita N. T. (2003) Correlation between relative ages

44

inferred from 1006 ^{26}Al and bulk compositions of ferromagnesian chondrules in least equilibrated 1007
ordinary chondrites. *Meteorit. Planet. Sci.* **38**, 939-962.

1008 Tanaka H., Murase K., and Tanigawa T. (2020) Final masses of giant planets. III. Effect of 1009

photoevaporation and a new planetary migration model. *Astrophys. J.* **891**, 143. 1010 Tenner T. J., Ushikubo T., Kurahashi E., Kita N. T. and Nagahara H. (2013) Oxygen isotope 1011 systematics of chondrule phenocrysts from the CO3.0 chondrite Yamato 81020: evidence for 1012 two distinct oxygen isotope reservoirs. *Geochim. Cosmochim. Acta* **102**, 226-245. 1013 Tenner T. J., Nakashima D., Ushikubo T., Kita N. T. and Weisberg M. K. (2015) Oxygen isotope 1014 ratios of Fe-poor chondrules in CR3 chondrites: Influence of dust enrichment and H₂O during 1015 chondrule formation. *Geochim. Cosmochim. Acta* **148**, 228-250.

1016 Tenner T. J., Kimura M., and Kita N. T. (2017) Oxygen isotope characteristics of chondrules from 1017 the Yamato-82094 ungrouped carbonaceous chondrite: Further evidence for common O 1018 isotope environments sampled among carbonaceous chondrites. *Meteorit. Planet. Sci.* **52**, 268- 1019 294.

1020 Tenner T. J., Ushikubo T., Nakashima D., Schrader D. L., Weisberg M. K., Kimura M. and Kita 1021 N. T. (2018) Oxygen isotope characteristics of chondrules from recent studies by secondary 1022 ion mass spectrometry. In *Chondrules: Records of the Protoplanetary Disk Processes*. (eds. 1023 S.S. Russell, H. C. Connolly Jr., A. N. Krot) Cambridge University Press, U.K., pp. 196-246.

1024 Tonui E., Zolensky M. E., Hiroi T., Nakamura T., Lipshutz M. E., Wang M.-S., and Okudaira K. 1025 (2014) Petrographic, chemical and spectroscopic evidence for thermal metamorphism in 1026 carbonaceous chondrites I: CI and CM chondrites. *Geochim. Cosmochim. Acta* **126**, 284-306. 1027

Ushikubo T., Kimura M., Kita N. T. and Valley J. W. (2012) Primordial oxygen isotope reservoirs 1028 of the solar nebula recorded in chondrules in Acfer 094 carbonaceous chondrite. *Geochim.*

1029 *Cosmochim. Acta* **90**, 242-264.

1030 Ushikubo T., Nakashima D., Kimura M., Tenner T. J. and Kita N. T. (2013) Contemporaneous 1031 formation of chondrules in distinct oxygen isotope reservoirs. *Geochim. Cosmochim. Acta* **109**, 1032

280-295.

Ushikubo T., Tenner T. J., Hiyagon H., and Kita N. T. (2017) A long duration of the ^{16}O -rich reservoir in the solar nebula, as recorded in fine-grained refractory inclusions from the least metamorphosed carbonaceous chondrites. *Geochim. Cosmochim. Acta* **201**, 103-122.

Vacher L. G., Marrocchi Y., Verdier-Paoletti M. J., Villeneuve J., and Gounelle M. (2016) Inward radial mixing of interstellar water ices in the solar protoplanetary disk. *Astrophys. J. Lett.* **827**, L1.

Vacher L. G., Marrocchi Y., Verdier-Paoletti M. J., Villeneuve J., and Gounelle M. (2017) Petrographic and C & O isotopic characteristics of the earliest stages of aqueous alteration of CM chondrites. *Geochim. Cosmochim. Acta* **213**, 271-290.

Van Kooten E., Cavalcante L., Wielandt D., and Bizzarro M. (2020) The role of Bells in the continuous accretion between the CM and CR chondrite reservoirs. *Meteorit. Planet. Sci.* **55**, 575-590.

Verdier-Paoletti M. J., Marrocchi Y., Avice G., Roskosz M., Gurenko A., and Gounelle M. (2017) Oxygen isotope constraints on the alteration temperatures of CM chondrites. *Earth Planet. Sci. Lett.* **458**, 273-281.

Villeneuve J., Libourel G., Soulié C. (2015) Relationships between type I and type II chondrules: implications on chondrule formation processes. *Geochim. Cosmochim. Acta* **160**, 277–305.

Villeneuve J., Marrocchi Y., Jacquet E. (2020) Silicon isotopic compositions of chondrule silicates in carbonaceous chondrites and the formation of primordial solids in the accretion disk. *Earth*

Wasson J. T. and Rubin A. E. (2003) Ubiquitous low-FeO relict grains in type II chondrules and limited overgrowths on phenocrysts following the final melting event. *Geochim. Cosmochim.*

Acta **67**, 2239–2250.

1056 Wasson J. T., Rubin A. E. and Yurimoto H. (2004) Evidence in CO3.0 chondrules for a drift in
1057 the O isotopic composition of the solar nebula. *Meteorit. Planet. Sci.* **39**, 1591-1598. 1058 Warren P.
H. (2011) Stable-isotopic anomalies and the accretionary assemblage of the Earth and 1059 Mars: A
subordinate role for carbonaceous chondrites. *Earth Planet. Sci. Lett.* **311**, 93-100. 1060 Weisberg M. K.,
McCoy T. J., and Krot A. N. (2006) Systematics and evaluation of meteorite 1061 classification. In
Meteorites and the Early Solar System II (eds. D. S. Lauretta and H. Y. 1062 McSween Jr.) University of
Arizona Press, Tucson, Arizona, pp. 19-52. 1063 Weisberg M. K., Ebel D. S., Connolly H. C., Kita N. T.
and Ushikubo T. (2011) Petrology and 1064 oxygen isotope compositions of chondrules in E3 chondrites.
Geochim. Cosmochim. Acta **75**, 1065 6556–6569.

1066 Williams C. D., Sanborn M. E., Defouilloy D., Yin Q.-Z., Kita N. T., Ebel D. S., Yamakawa A.
1067 and Yamashita K. (2020) Chondrules reveal large-scale outward transport of inner Solar 1068
System materials in the protoplanetary disk. *Proc. Natl. Acad. Sci. USA* **117**, 23426-23435 1069 Wood J.
A. and Hashimoto A. (1993) Mineral equilibrium in fractionated nebular systems. 1070 *Geochim.*
Cosmochim. Acta **57**, 2377-2388.

1071 Young E. D. and Russell S. S. (1998) Oxygen reservoirs in the early solar nebula inferred from an
1072 Allende CAI. *Science* **282**, 2377-2388.

1073 Zanda B., Bourot-Denise M., Perron C. and Hewins R. H. (1994) Origin and metamorphic 1074
redistribution of silicon, chromium, and phosphorous in the metal of chondrites. *Science* **265**,

1075 1846-1849.

1076 Zolensky M. E., Barrett R., and Browning L. (1993) Mineralogy and composition of matrix and
1077 chondrule rims in carbonaceous chondrites. *Geochim. Cosmochim. Acta* **57**, 3123–3148. 1078

Zolensky M. E., Weisberg M. K., Buchanan P. C., Mittlefehldt D. W. (1996) Mineralogy of 1079
carbonaceous chondrite clasts in HED achondrites and the Moon. Meteoritics and Planetary 1080 *Science*
31, 518–537.

1081 Zolensky M. E., Mittlefehldt D. W., Lipshutz M. E., Wang M.-S., Clayton R. N., Mayeda T. K.,
1082 Grady M. M., Pillinger C. T., and Barber D. (1997) CM chondrites exhibit the complete 1083
petrologic range from type 2 to 1. *Geochim. Cosmochim. Acta* **61**, 5099–5115. 1084

terms of oxygen isotope compositions (C18: POP, C30: PO, C21 and C11: PP). SIMS oxygen 1089 three-isotope analysis points are shown by the vertex of the triangles, color-coded for mineral phases (olivine: blue, relict olivine: white, low-Ca pyroxene: green). Δ 1090 ^{17}O values of individual 1091 analyses are indicated. (For interpretation of the references to color in this figure legend, the reader 1092 is referred to the web version of this article).

1093 Fig 2. BSE images of type I chondrules analyzed in Paris displaying various textures (C7: POP, 1094 C5: POP + BO core, C16: BO fragment, C25: PP, and C9: GO). SIMS oxygen three-isotope 1095 analysis points are shown by the vertex of the triangles, color-coded for mineral phases (olivine: blue, relict olivine: white, low-Ca pyroxene: green, high-Ca pyroxene: orange). Δ 1096 ^{17}O values of 1097 individual analyses are indicated. (For interpretation of the references to color in this figure legend, 1098 the reader is referred to the web version of this article).

1099 Fig 3. BSE images of the type I chondrule C14 analyzed in Paris. SIMS oxygen three-isotope 1100 analysis points are shown by the vertex of the triangles, color-coded for mineral phases (olivine: blue, relict olivine: white, low-Ca pyroxene: green). Δ 1101 ^{17}O values of individual analyses are 1102 indicated. (For interpretation of the reference to color in this figure legend, the reader is referred 1103 to the web version of this article).

1104 Fig. 4. BSE images of a type II chondrule (C15: PO), two isolated Fe-rich olivine grains (G32 and 1105 G33), and an isolated Fe-poor olivine grain (G24) analyzed in Paris. SIMS oxygen three-isotope analysis points are shown by the vertex of the triangles, color-coded for olivine (blue). Δ 1106 ^{17}O values

1107 of individual analyses are indicated. (For interpretation of the reference to color in this figure 1108 legend, the reader is referred to the web version of this article).

Fig. 5. Oxygen 3-isotope diagram of individual spot analyses of olivine, Low-Ca pyroxene (Lpx), high-Ca pyroxene (Hpx), and relict grains in chondrules in Paris. Error bars, corresponding to the spot-to-spot reproducibility (2SD), are smaller than the symbol sizes. The CCAM (carbonaceous chondrite anhydrous mineral; Clayton et al., 1977), Y&R (Young and Russell, 1998), and PCM (primitive chondrule minerals; Ushikubo et al., 2012) lines are shown for reference. The terrestrial fractionation line (TFL) is also shown.

1115

Fig. 6. $\Delta^{17}\text{O}$ values of individual host chondrules and isolated olivine grains in Paris. Relict olivine (ol.) and low-Ca pyroxene (Lpx) grains are shown together. Data are sorted according to the host $\Delta^{17}\text{O}$ values. Chondrule C9 (GO) contains homogeneously ^{16}O -poor olivine, which is likely relict. A single analysis of pyroxene would represent melt oxygen isotope ratios that show lower $\Delta^{17}\text{O}$ compared to other host chondrule values. For the heterogeneous chondrule C15 (II PO) and isolated Fe-poor olivine grain G24, individual analyses are shown. Error bars represent the propagated uncertainties for host chondrules and olivine in C9 and external reproducibility of individual analyses for relict grains, pyroxene in C9, and all data in C15 and G24.

Fig. 7. $\Delta^{17}\text{O}$ values of individual host chondrules in Paris vs. Mg#s (open diamonds). Data from the Murchison CM2 chondrite (black and white squares; Chaumard et al., 2018) are shown for comparison. Chondrule Mg# uncertainties correspond to the range of measured values, while uncertainties in $\Delta^{17}\text{O}$ are the propagated 2SE.

Fig. 8. $\Delta^{17}\text{O}$ values vs. Mg#s of individual host chondrules in CM (open diamonds; Chaumard et

1129 al., 2018; this work), CO (grey circles; Tenner et al., 2013), and Acfer 094 (black circles; Ushikubo
1130 et al., 2012) chondrites. Chondrule Mg# uncertainties correspond to the range of measured values,
while uncertainties in $\Delta 1131^{17}\text{O}$ are the propagated 2SE.

Fig. 9. $\Delta 1132^{17}\text{O}$ values of individual host chondrules in the CM chondrites Paris (this work) and 1133
Murchison (Chaumard et al., 2018) and Mg#s superimposed by oxygen isotope mixing curves of 1134
constant dust enrichment and ice enhancement from Tenner et al. (2015) and Hertwig et al. (2018). 1135
In this model, the anhydrous silicate dust, Solar gas, water ice, and organics in the dust are considered to
have $\Delta 1136^{17}\text{O}$ values of -8.0‰ , -28.4‰ , $+2.0\text{‰}$, and $+11.3\text{‰}$, respectively (Hertwig 1137 et al.,
2018). Chondrule Mg# uncertainties correspond to the range of measured values, while uncertainties in Δ
1138 ^{17}O are the propagated 2SE.

1139

1140

Table 1. Mg#’s and O-isotope ratios of host chondrules and relict grains.

(ol, lpx, hpx)^c $\delta^{18}\text{O}$ unc. $\delta^{17}\text{O}$ unc. $\Delta^{17}\text{O}$ unc. $\Delta^{17}\text{O}$ Chondrule Type, texture Mg#^a +/- ^b n

Beam

2SD

(μm)

Homogeneous

C21 I, PP 99.0 0.2/0.3 2,5,0 -7.7 0.3 -10.9 0.3 -6.93 0.22 0.6 15 C6 I, PP 98.8 0.3/0.8 2,4,0 -6.5 0.4 -9.5 0.4 -6.10 0.34 0.3 15 C28 I, PO 99.6 0.1/0.2 8,0,0 -6.3 0.3 -9.3 0.3 -5.99 0.18 0.4 15 C23 I, POP 98.7 0.3/0.3 3,4,0 -5.7 0.6 -8.8 0.3 -5.84 0.17 0.4 15 C10 I, PO 99.2 0.1/0.2 5,1,0 -6.4 0.5 -9.1 0.3 -5.72 0.16 0.3 15

relict ol 99.3 n.a. 1,0,0 -5.8 0.2 -9.5 0.3 -6.5 0.3 n.a. 15 relict ol 99.0 n.a. 1,0,0 -7.3 0.2 -10.3 0.3 -6.6 0.3 n.a. 15 C27 I, PP 99.3 0.5/0.9 6,2,0 -4.5 0.4 -7.9 0.3 -5.57 0.13 0.2 15 C4 I, POP 99.0 0.2/0.3 6,2,0 -5.9 0.4 -8.7 0.4 -5.63 0.31 0.2 15 C26 I, POP 99.0 0.1/0.1 2,3,0 -4.7 0.4 -7.9 0.3 -5.51 0.25 0.4 15

relict ol 99.3 n.a. 1,0,0 -6.8 0.2 -9.8 0.5 -6.3 0.4 n.a. 15 relict ol 99.3 n.a. 1,0,0 -6.8 0.2 -10.1 0.5 -6.6 0.4 n.a. 15 relict ol 99.2 n.a. 1,0,0 -6.3 0.2 -9.9 0.5 -6.6 0.4 n.a. 15 C29 I, PP 99.0 0.3/0.9 2,4,0 -4.8 0.4 -7.9 0.3 -5.43 0.19 0.4 15 C20 I, POP 98.9 0.4/0.2 4,5,0 -5.0 0.3 -7.9 0.2 -5.32 0.18 0.4 15 C13 I, BO 99.1 0.2/0.1 4,0,0 -4.2 0.4 -7.4 0.3 -5.18 0.19 0.3 15 C2 I, POP 98.9 0.3/0.4 4,4,0 -4.8 0.4 -7.7 0.2 -5.19 0.17 0.4 15 C19 I, POP 98.6 0.2/0.2 5,3,0 -4.8 0.3 -7.5 0.2 -4.98 0.14 0.3 15 C30 I, PO 98.8 0.1/0.1 6,0,0 -3.2 0.3 -5.6 0.3 -3.90 0.26 0.2 15 relict ol 98.7 n.a. 1,0,0 1.0 0.3 -1.3 0.4 -1.9 0.5 n.a. 15 relict ol 98.4 n.a. 1,0,0 -4.3 0.3 -6.3 0.4 -4.1 0.5 n.a. 15 C18 I, POP 98.7 0.1/0.3 0,3,0 -1.2 0.5 -4.0 0.4 -3.31 0.29 0.4 15 relict ol 99.3 n.a. 1,0,0 2.8 0.2 0.7 0.5 -0.8 0.4 n.a. 15 relict ol 99.1 n.a. 1,0,0 2.4 0.2 0.0 0.5

–1.2 0.4 n.a. 15 relict ol 99.1 n.a. 1,0,0 2.3 0.2 0.0 0.5 –1.2 0.4 n.a. 15 relict ol 99.1 n.a. 1,0,0 2.0 0.2 –0.5 0.5 –1.5 0.4 n.a. 15 relict ol 99.1 n.a. 1,0,0 2.4 0.2 –0.4 0.5 –1.6 0.4 n.a. 15 C11 I, PP 93.2 0.2/0.7 0,8,0 1.5 0.3 –1.7 0.3 –2.51 0.18 0.5 15 C12 I, PP 93.6 0.8/0.6 0,7,0 1.3 0.4 –1.7 0.2 –2.37 0.13 0.3 15 C16 I, BO, frag. 99.4 0.2/0.2 8,0,0 –7.9 0.4 –11.1 0.3 –6.99 0.24 0.4 10 C7 I, POP 99.0 0.1/0.2 6,2,0 –6.7 0.5 –9.4 0.4 –5.90 0.24 0.4 10
C8 I, POP 98.9 0.0/0.1 3,2,0 –5.3 0.4 –8.2 0.3 –5.42 0.22 0.4 10 relict ol 98.8 n.a. 1,0,0 2.2 0.4 1.2 0.4 0.1 0.4 n.a. 10 relict ol 98.7 n.a. 1,0,0 –2.1 0.4 –5.7 0.4 –4.6 0.4 n.a. 10 relict ol 98.7 n.a. 1,0,0 –0.9 0.4 –3.8 0.4 –3.3 0.4 n.a. 10 C14 I, POP+BO

core 99.0 0.2/0.3 7,4,0 –5.3 0.5 –8.0 0.3 –5.26 0.23 0.6 10 relict ol 99.1 n.a. 1,0,0 –22.7 0.3 –24.8 0.4 –13.0 0.4 n.a. 10 C1 I, GOP 98.6 0.3/0.3 2,3,0 –3.7 0.5 –6.9 0.4 –4.99 0.33 0.6 10 relict ol 99.0 n.a. 1,0,0 –14.6 0.6 –17.2 0.6 –9.5 0.5 n.a. 10 relict lpx 98.8 n.a. 0,1,0 –8.4 0.6 –11.4 0.6 –7.0 0.5 n.a. 10 C25 I, PP 98.8 0.4/0.8 3,4,1 –3.2 0.4 –6.3 0.4 –4.66 0.26 0.6 10 C5 I, POP+BO

core 98.6 0.2/0.3 4,4,0 –4.8 0.8 –7.3 0.4 –4.77 0.23 0.4 10 C31 I, PP 97.3 0.5/1.6 1,5,0 0.0 0.4 –3.3 0.4 –3.28 0.23 0.4 10 relict ol 97.8 n.a. 1,0,0 –3.5 0.5 –7.1 0.6 –5.3 0.4 n.a. 10 relict ol 97.6 n.a. 1,0,0 –5.2 0.5 –8.9 0.6 –6.2 0.4 n.a. 10 C22 II, PO 69.7 9.3/9.4 7,0,0 2.3 0.4 –1.0 0.3 –2.21 0.15 0.3 15 C17 II, PO 74.4 5.2/4.5 5,0,0 2.8 0.3 –0.6 0.3 –2.09 0.25 0.5 15 relict ol 99 n.a. 1,0,0 –7.1 0.1 –10.2 0.3 –6.5 0.3 n.a. 15 relict ol 76.6 n.a. 1,0,0 –3.7 0.1 –7.0 0.3 –5.1 0.3 n.a. 15 relict ol 71.4 n.a. 1,0,0 0.5 0.1 –2.6 0.3 –2.9 0.3 n.a. 15

Heterogeneous

C9 I, GO 99.4 0.1/0.1 7,0,0 3.5 0.4 1.7 0.3 –0.15 0.20 0.4 10 99.3 n.a. 0,1,0 –11.3 0.4 –13.7 0.4 –7.8 0.4 n.a. 10 C15 II, PO 98.5 n.a. 1,0,0 –6.6 0.1 –9.8 0.2 –6.3 0.2 n.a. 15 98.3 n.a. 1,0,0 –3.4 0.1 –6.6 0.2 –4.8 0.2 n.a. 15 97.7 n.a. 1,0,0 –2.6 0.1 –5.9 0.2 –4.5 0.2 n.a. 15 91.3 n.a. 1,0,0 –2.4 0.1 –5.4 0.2 –4.1 0.2 n.a. 15 77.3 n.a. 1,0,0 1.2 0.1 –2.7 0.2 –3.3 0.2 n.a. 15 66.9 n.a. 1,0,0 1.6 0.1 –2.4 0.2 –3.3 0.2 n.a. 15 63.4 n.a. 1,0,0 2.5 0.1 –1.0 0.2 –2.3 0.2 n.a. 15 58.9 n.a. 1,0,0 1.9 0.1 –2.2 0.2 –3.2 0.2 n.a. 15^a Mg# = Mg/(Fe+Mg) molar % of olivine and/or pyroxene in each chondrule.

^bUncertainties represent the range in measured Mg#'s of olivine and/or pyroxenes.

^cNumbers of mineral phases analyzed (olivine, low-Ca pyroxene, and high-Ca pyroxene).

Table 2. Mg#'s and O-isotope ratios of isolated olivine grains and aggregate olivine inclusion.

Object Type, texture Mg# +/- n (ol) $\delta^{18}\text{O}$ unc. $\delta^{17}\text{O}$ unc. $\Delta^{17}\text{O}$ unc. $\Delta^{17}\text{O}$

Beam

2SD

(μm)

G32 Isolated FeO-rich olivine 72.2 2.3/5.7 4 2.2 0.7 –1.6 0.5 –2.7 0.5 0.8 15 G33 Isolated FeO-rich olivine 37.3 11.4/7.7 4 1.8 0.9 –0.9 0.5 –1.9 0.3 0.3 15 G24 Isolated FeO-poor olivine 99.4 n.c. 1 3.3 0.3 1.7 0.5 0.0 0.5 n.a. 10 99.4 n.c. 1 2.0 0.3 0.4 0.5 –0.6 0.5 n.a. 10 99.4 n.c. 1 –1.8 0.3 –4.0 0.5 –3.1 0.5 n.a. 10 99.4 n.c. 1 –3.2 0.3 –5.5 0.5 –3.8 0.5 n.a. 10 99.3 n.c. 1 2.6 0.3 0.6 0.5 –0.8 0.5 n.a. 10 99.3 n.c. 1 0.2 0.3 –2.2 0.5 –2.3 0.5 n.a. 10 99.3 n.c. 1 –3.1 0.3 –5.4 0.5 –3.9 0.5 n.a. 10 I3 AOI 99.5 0.1/0.0 2 –45.5 0.5 –47.3 0.4 –23.7 0.2 0.4 10