



Initialized Earth System prediction from subseasonal to decadal timescales

Gerald A. Meehl¹✉, Jadwiga H. Richter¹, Haiyan Teng², Antonietta Capotondi^{3,4}, Kim Cobb⁵, Francisco Doblas-Reyes^{6,7}, Markus G. Donat⁶, Matthew H. England⁸, John C. Fyfe⁹, Weiqing Han¹⁰, Hyemi Kim¹¹, Ben P. Kirtman¹², Yochanan Kushnir¹³, Nicole S. Lovenduski^{10,14}, Michael E. Mann^{15,16}, William J. Merryfield⁹, Veronica Nieves¹⁷, Kathy Pegion¹⁸, Nan Rosenbloom¹, Sara C. Sanchez¹⁹, Adam A. Scaife^{20,21}, Doug Smith²⁰, Aneesh C. Subramanian¹⁰, Lantao Sun²², Diane Thompson²³, Caroline C. Ummenhofer^{24,25} and Shang-Ping Xie²⁶

Abstract | Initialized Earth System predictions are made by starting a numerical prediction model in a state as consistent as possible to observations and running it forward in time for up to 10 years. Skilful predictions at time slices from subseasonal to seasonal (S2S), seasonal to interannual (S2I) and seasonal to decadal (S2D) offer information useful for various stakeholders, ranging from agriculture to water resource management to human and infrastructure safety. In this Review, we examine the processes influencing predictability, and discuss estimates of skill across S2S, S2I and S2D timescales. There are encouraging signs that skilful predictions can be made: on S2S timescales, there has been some skill in predicting the Madden–Julian Oscillation and North Atlantic Oscillation; on S2I, in predicting the El Niño–Southern Oscillation; and on S2D, in predicting ocean and atmosphere variability in the North Atlantic region. However, challenges remain, and future work must prioritize reducing model error, more effectively communicating forecasts to users, and increasing process and mechanistic understanding that could enhance predictive skill and, in turn, confidence. As numerical models progress towards Earth System models, initialized predictions are expanding to include prediction of sea ice, air pollution, and terrestrial and ocean biochemistry that can bring clear benefit to society and various stakeholders.

There has been an increasing desire for climatic information on timescales from weeks to months, seasons and years. Such information offers clear benefits to society and various stakeholders alike. For instance, prediction of the hydroclimate could allow for better water resource management and improved agricultural maintenance, whereas temperature and wind predictions could provide critical information for infrastructure planning and expected energy consumption. To obtain this climatic information, initialized predictions on various near-term timescales must be used.

Initialized Earth System prediction describes a suite of climate model simulations wherein the starting conditions are set as close to observations as possible and the model is run forward for up to 10 years¹. Internally generated, naturally occurring variability is therefore

considered a key aspect of these time-evolving climate predictions². They differ from uninitialized simulations — or climate change projections — where internal variability is removed through ensemble averaging, and focus is instead given to quantifying the effects of external forcing such as anthropogenic greenhouse gases^{3,4}.

Given the duration of simulations, initialized predictions span various timescales (FIG. 1a): subseasonal to seasonal (S2S; ~2 weeks–2 months)^{5,6}, seasonal to interannual (S2I; 2–12 months)⁷ and seasonal to decadal (S2D; 3 months–10 years)^{1,2}. In each case, efforts have focused on climate phenomena that also operate on similar timescales. For example, S2S research has concentrated on the Madden–Julian Oscillation (MJO) and sudden stratospheric warmings (SSWs); S2I on the

✉e-mail: meehl@ucar.edu
<https://doi.org/10.1038/s43017-021-00155-x>

Key points

- Initialization methods vary greatly across different prediction timescales, creating difficulties for seamless prediction.
- Model error and drift limit predictability across all timescales. Although higher resolution models show promise in reducing these errors, improvements in physical parameterizations are needed to improve predictability.
- The effects of land processes, interactions across various ocean basins and the role of stratospheric processes in predictability are not well understood.
- Predictability on seasonal to decadal timescales is largely associated with predictability of the major modes of variability in the atmosphere and the ocean.
- Evolution of Earth System models will lead to predictability of more societal-relevant variables spanning multiple parts of the Earth System.

El Niño–Southern Oscillation (ENSO), North Atlantic Oscillation (NAO), Indian Ocean Dipole (IOD), Southern Annular Mode (SAM) and Quasi-Biennial Oscillation (QBO); and S2D on slowly evolving oceanic processes such as Pacific decadal variability (PDV) and Atlantic multi-decadal variability (AMV).

Distinct communities have therefore formed to coordinate research and perform initialized predictions on

each timescale. Efforts such as the S2S Prediction Project and Database⁵ and the Subseasonal Experiment (SubX⁶) emerged for S2S; the [North American Multi-Model Ensemble](#)⁷, the [Asia-Pacific Economic Cooperation \(APEC\) Climate Center \(APCC\)](#), and the [Copernicus Climate Change Service](#) for S2I; and sets of hindcasts and predictions as part of the Coupled Model Intercomparison Project phase 5 (CMIP5)^{1,2} and CMIP6 (REF.⁸) for S2D.

Although these communities are often separate, however, they all rely on similar methodologies (TABLE 1; see Supplementary Tables 1–3). Thus, there is potential for ‘seamless prediction’⁹, whereby one framework can be used to address prediction across all timescales, with skill increasingly associated with external forcing as simulations progress¹⁰ (FIG. 1b). Yet, in practice, community differences with regards to initialization frequency, for example, make seamless prediction challenging^{1,2}.

In this Review, we bring together research on initialized predictions on timescales of weeks to years. We begin by outlining current methodologies for initialized predictions, incorporating discussion of the process, ensemble size, verification and prediction skill. We subsequently outline prediction on S2S, S2I and S2D timescales, before discussing priorities for future research that will increase the feasibility for seamless prediction.

Making predictions

S2I research using initialized prediction has been taking place since the late 1980s (REF.¹¹). In contrast, it was not until 20 years later that initialized S2D climate predictions began, in turn, initiating a rapid acceleration of research from which operational systems are now routinely produced¹². We begin by describing the process of initialized prediction, focusing on the methodological aspects involving forecast verification and measures of prediction skill (the level of agreement between an initialized prediction and the observed state it is meant to predict).

Process of initialized prediction. Predictions for S2S, S2I and S2D timescales, ranging from weeks to years, use numerical models with components of (at least) atmosphere, ocean, land and sea ice that are started from a particular observed state. The process of bringing the model components into close correspondence with that observed state is termed initialization, and predictions that are started from such observed states are referred to as initialized predictions. There are currently many activities taking place in the S2S, S2I and S2D communities with regards to initialized prediction, with key differences amongst centres regarding how models are used (TABLE 1; see Supplementary Tables 1–3).

One key difference between the subseasonal and longer timescale systems is the origin of the model. Many S2S (and some S2I) prediction systems originate in the numerical weather prediction community. As such, they tend to have the highest horizontal resolution in the atmosphere, largely ~0.25–0.5° (TABLE 1). Atmospheric initialization in these numerical weather prediction-derived models uses data assimilation¹³, such as 3D variational assimilation (as in the CMA model). Moreover, to produce the initial perturbations for ensemble generation,

Author addresses

¹Climate and Global Dynamics Laboratory, National Center for Atmospheric Research, Boulder, CO, USA.

²Pacific Northwest National Laboratory, Richland, WA, USA.

³Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA.

⁴NOAA Physical Sciences Laboratory, Boulder, CO, USA.

⁵School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA.

⁶Department of Earth Sciences, Barcelona Supercomputing Center-Centro Nacional de Supercomputación, Barcelona, Spain.

⁷Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain.

⁸Climate Change Research Centre, University of New South Wales, Sydney, New South Wales, Australia.

⁹Canadian Centre for Climate Modeling and Analysis, Environment and Climate Change Canada, Victoria, British Columbia, Canada.

¹⁰Department of Atmospheric and Oceanic Sciences, University of Colorado, Boulder, CO, USA.

¹¹School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY, USA.

¹²Rosenstiel School for Marine and Atmospheric Science, University of Miami, Miami, FL, USA.

¹³Lamont Doherty Earth Observatory, Columbia University, Palisades, NY, USA.

¹⁴Institute of Arctic and Alpine Research, University of Colorado, Boulder, CO, USA.

¹⁵Department of Meteorology & Atmospheric Science, Pennsylvania State University, State College, PA, USA.

¹⁶Earth & Environmental Systems Institute, Pennsylvania State University, State College, PA, USA.

¹⁷Image Processing Laboratory, University of Valencia, Valencia, Spain.

¹⁸Department of Atmospheric Oceanic and Earth Sciences, George Mason University, Fairfax, VA, USA.

¹⁹Joint Institute for the Study of the Atmosphere and Ocean, University of Washington, Seattle, WA, USA.

²⁰Hadley Centre, Exeter, UK.

²¹College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK.

²²Department of Atmospheric Science, Colorado State University, Ft. Collins, CO, USA.


²³Department of Geosciences, University of Arizona, Tucson, AZ, USA.

²⁴Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, MA, USA.

²⁵ARC Centre of Excellence for Climate Extremes, Sydney, New South Wales, Australia.

²⁶Scripps Institution of Oceanography, La Jolla, CA, USA.

a Predictability sources and timescales



b Predictability skill from initial state versus external forcing




Fig. 1 | Timescales and processes involved with initialized predictions. a | Timescales and sources of predictability for subseasonal to seasonal (S2S), seasonal to interannual (S2I) and seasonal to decadal (S2D) timescales. Lighter green shading indicates larger uncertainty. **b** | Skill in predicting the upper 300 m of the Atlantic Ocean temperature, as measured by relative entropy, in initialized models (blue) and those forced by RCP4.5 (red). Skill is high for initialized predictions on S2S and S2I timescales (<2 years), but decreases towards S2D (years 3–9), after which time the skill from external forcing increases. AMV, Atlantic multi-decadal variability; ENSO, El Niño–Southern Oscillation; GHG, greenhouse gas; GMST, global mean surface temperature; MJO, Madden–Julian Oscillation; NAO, North Atlantic Oscillation; PDV, Pacific decadal variability; QBO, Quasi-Biennial Oscillation; SSW, sudden stratospheric warming. Panel **b** adapted with permission from REF.¹⁰, Wiley ©2012. American Geophysical Union. All Rights Reserved.

they sometimes use data assimilation with an ensemble Kalman filter¹⁴ (as in the ECCO model) or singular vectors¹⁵ (as in the JMA model). In comparison, most S2I, and all but one S2D, prediction systems are based on climate or Earth System models (ESMs) previously used for IPCC climate projections. In these cases, the majority of models have a horizontal resolution of $\sim 0.5\text{--}1^\circ$ (TABLE 1).

In addition to differences in the models and their resolution across prediction timescales, contrasts are also evident in the components that are initialized and

the degree of coupling between Earth System components. In S2S predictions, for example, coupling between the atmosphere, ocean, land and sea ice is not considered crucial (FIG. 1a). As such, only a small number of models initialize the ocean and employ atmosphere–ocean coupling, but the majority initialize land surface conditions (Supplementary Table 1). For S2D predictions, however, oceanic processes are vital and, as a result, all models initialize the ocean and have at least partial coupling with the atmosphere and sea ice; only a fraction initialize the

Table 1 | General characteristics of models used for S2S, S2I and S2D initialized predictions^a

Timescale	Number of models	Atmospheric resolution and levels	Ocean resolution and levels	Components initialized	Initialization	Number of ensembles	Prediction length
S2S	18	25–200 km 17–91 levels	8–200 km 25–75 levels	Most initialize atmosphere, ocean, land and sea ice	Full field	4–51	31–62 days
S2I	13	36–200 km 24–95 levels	25–200 km 24–74 levels	All initialize atmosphere, ocean, land and sea ice	Full field	10–51	6–12 months
S2D	14	50–200 km 26–95 levels	25–100 km 30–75 levels	Models range from initializing only ocean to initializing atmosphere, ocean, land and sea ice	Full field, anomaly	10–40	5–10 years

S2D, seasonal to decadal; S2I, seasonal to interannual; S2S, subseasonal to seasonal. ^aA full and more complete account of model features is given in Supplementary Tables 1, 2 and 3 for S2S, S2I and S2D models, respectively.

atmosphere and land surface (Supplementary Table 3). As S2I falls in the time window where predictability comes from all Earth System components (FIG. 1a), care is typically taken to initialize each of them.

Atmospheric initialization is often achieved by interpolating an existing analysis to the model grid and generating an ensemble spread using the random field perturbation method¹⁶ (as in CESM1 for S2S), the lagged ensemble method^{17,18} (as in CCSM3) or nudging to reanalyses in coupled mode¹⁹ (as in the CCCma model). Various approaches have also been used to initialize the ocean state, including a hindcast spin-up in an ocean forced by observed atmospheric conditions²⁰, nudging the ocean model to some observed ocean state²¹ or using full ocean data assimilation²². Land variables are initialized either by assimilation of land observations²³ or by running an offline land-only model that is forced with observed atmospheric conditions²⁴. The initialization strategy also differs between the shorter and longer-term prediction models. All S2S and S2I prediction models use full fields (such as sea surface temperature (SST)). By contrast, about half of the S2D modes use anomaly initialization, meaning an initial condition is constructed by adding observed (or reanalysis) anomalies to the model's climatology in order to minimize initialization shock and model drift^{25–27}.

As individual model components are often initialized in different ways, there is frequently no coupling between initial conditions for various parts of the Earth System, thereby creating an imbalance in the initial state of the model. New methodologies, such as weakly coupled and strongly coupled data assimilation, offer promising approaches to reduce initialization shock and imbalance in the model²⁸. In the weakly coupled approach, the assimilation is applied to each of the components of the coupled model independently, whereas interaction between the components is provided by the coupled forecasting system²⁸. In the strongly coupled method, however, assimilation is applied to the full Earth System state simultaneously, treating the coupled system as a single integrated system²⁸.

There are currently very few modelling centres that have been able to apply seamless prediction owing to numerous practical aspects (including the initialization

method, initialization frequency, number of ensemble members, among others). The most seamless system is currently operated by the UK Met Office, which is providing S2S, S2I and S2D forecasts operationally using almost identical configurations of the model for all prediction systems²⁹. NCAR, although not an operational centre, is also using the same models, CESM1 and CESM2, to generate S2S, S2I and S2D hindcasts (and predictions for research purposes) using the same modelling framework, although at this time initialization details vary among the three prediction systems.

Ensemble size. Ensemble size is an important aspect determining predictive skill and reliability. In most prediction systems, ensemble sizes typically range between 10 and 50 (TABLE 1). There is potential to increase the number of ensembles by combining those from multiple systems³⁰ or time-lagged ensembles³¹, or using other techniques such as subsampling^{32,33} to improve the ensemble properties. Typically, the more ensemble members, the higher the anomaly correlation coefficient (ACC), a measure of prediction skill. For example, on S2S timescales, the ACC of global surface air temperature over land is ~0.29 when using only 4 CESM1 hindcast ensemble members³⁴, increasing to ~0.33 for 8 members and ~0.36 for 16 members (FIG. 2a).

Large ensembles are also advantageous for improving seasonal prediction skill of the NAO³⁵, including on S2D timescales^{33,36}. For example, ACC values are ~0.6 for an average of years 2–8 when using 40 ensemble members³⁷ (FIG. 2b). Further increases in multi-year NAO skill with an ACC of 0.8 are possible with a lagged ensemble of several hundred members³³ as a result of the modelled signal to noise ratio being too small.

There are consequences and trade-offs in terms of computing costs when using more ensemble members. For instance, an S2S reforecast could run 16 years (SubX) × 4 members × 2 months long × weekly start dates for ~600 model years; an S2I example could run 30 years × 9 members × 1 year long × 4 start dates per year for ~1,000 model years; and an S2D example (DCPP) could run 60 years × 10 members × 10 years long for ~6,000 model years.

Verification using observations. A key element of initialized prediction is having a solid understanding of the climate phenomena that are being predicted. Analyses of observations in comparison with the model simulations are thus required. On S2S and S2I timescales, the observational record provides a good source of data to verify initialized hindcasts. For example, observations cover roughly 30 ENSO events and as many as 300 MJO cycles. However, these data have their limitations. For instance, 3D observations of the atmosphere and ocean are desired for prediction verification, for understanding of processes and mechanisms, and for initialization of the predictions in the first place³⁸. Yet such 3D gridded

data are limited to the period of the satellite record (dating from the late 1970s) and to reanalyses that assimilate all available observations. Moreover, although several ENSO (and similar timescale) events have been observed, these can exhibit different expressions³⁹ and undergo large decadal to millennial variations^{40–42}, requiring a long observational record to perform robust analyses.

Researchers in the field of initialized Earth System prediction on S2D timescales often cite the short observational record as a factor inhibiting understanding. For example, with reliable observations limited to the latter half of the twentieth century⁴³, only approximately three PDV or AMV transitions have occurred by which to compare predictions. Although some observations are available earlier in the twentieth century, these are sparse and reanalyses are highly uncertain, making consistent comparisons of prediction skill between the pre and post-satellite eras difficult. Added to that, subsurface ocean observations and critical state atmospheric variables (such as surface winds) are crucial to understanding slow variations in the climate system⁴⁴, but such observations also have a very short duration. Moreover, it is also difficult to objectively separate forced (natural and anthropogenic) and internal decadal to multi-decadal climate variability, adding further challenges for S2D prediction verification and triggering debate on best practices for signal separation^{45–48}.

Nevertheless, efforts are underway to improve methodological approaches and data provisions for prediction verification. The crucial need for better observations of the full depth of the ocean have started to be addressed by Argo floats, first for the upper 2,000 m (REF.⁴⁹) but with plans to be expanded to the full ocean depth⁵⁰.

Proxy-based reconstructions are also increasingly available, shedding light on processes associated with interannual and decadal timescales of variability⁵¹ beyond that possible by instrumental observations. Indeed, the particular limitations of instrumental data length and coverage for verification of S2D predictions have pointed to palaeoclimate reconstructions — using trees, corals and speleothems — to extend observations and provide further realizations of decadal variability^{40,42,52–56} (FIG. 3). Additionally, such records can provide insights into the physical mechanisms associated with this variability, including westerly wind anomalies⁵¹, upwelling, gyre circulation⁵⁷ and links among major modes of variability⁵⁸. Together with further advances in palaeoclimate research — including palaeoclimate synthesis^{59–62}, palaeo data assimilation techniques^{63–65} and development and expansion of proxy system models and toolboxes^{66,67} — palaeoclimate data will not only help with the verification of climate model simulations, particularly on the S2D timescale, but also provide context for initialized predictions by providing insights into the timescales of variability beyond the instrumental record.

Bias correction and prediction skill. To account for model drifts and biases, the skill of initialized predictions is typically evaluated in terms of forecast time-dependent anomalies that are departures from some




Fig. 2 | Influence of ensemble size and lead year ranges on predictive skill. **a** | Skill (as measured by the anomaly correlation coefficient (ACC)) in predicting subseasonal to seasonal (S2S) globally averaged NDJFM surface air temperature (SAT) (excluding the Antarctic) from CESM initialized hindcasts of various ensemble size (grey line). Shading denotes the 5% and 95% significance levels. Blue and red whiskers illustrate predictive skill for NCEP Climate Forecast System version 2 (CFSv2) and ECMWF subseasonal hindcasts, respectively. **b** | Skill (as measured by the anomaly correlation coefficient) in predicting seasonal to decadal (S2D) wintertime North Atlantic Oscillation (NAO) using ensembles of different sizes from the Decadal Prediction Large Ensemble (DPLE)²⁰. Each line depicts a different lead year range, with those that are coloured corresponding to statistically significant correlations; the darker the shading, the greater the statistical significance. Dashed-dotted line shows the skill of the sub-ensemble mean against a single member of the ensemble (averaged for all possible combinations). Both panels illustrate that the more ensemble members, the higher the skill for longer lead year ranges. Panel **b** is adapted from REF.³⁷, CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>).




Fig. 3 | Extending proxy observations of S2D variability back in time derived from corals. **a** | Global mean surface temperature anomalies. **b** | 30-year running mean standard deviations of the coral-based Indian Ocean Dipole (IOD) (blue) and El Niño–Southern Oscillation (ENSO) (red). **c** | Scatter plot of the 30-year running mean standard deviations of coral-based IOD and ENSO. **d** | Equatorial Pacific west–east (W–E) sea surface temperature (SST) gradient (shading represents uncertainty). **e** | Central and eastern Pacific (CP/EP) El Niño derived from teleconnected climate patterns. **f** | An indication of reconstructions considered robust in panel **e**. Collectively, the panels illustrate a strengthening of IOD–ENSO decadal variability after ~1590. S2D, seasonal to decadal. Adapted with permission from REF.⁵⁸, Springer Nature Limited.

measure of mean climate. However, a prediction will drift rapidly from the initial observed state towards its own climatology owing to model error. These drifts start almost immediately in a prediction, and by lead year 1 are already considerable (FIG. 4).

The calculation of anomalies and correction of model biases are addressed together, typically by calculating and removing the model climatology. For S2S predictions, the common methodology is to calculate a lead time-dependent model climatology from a set of hindcasts and to compute anomalies from this climatology. However, such a procedure is complicated owing to the inhomogeneous nature of current subseasonal prediction systems⁶. The climatology for S2I predictions is similarly accomplished by averaging over all years of the hindcast for a particular start time and lead or target time⁶⁸, thereby assuming stationarity of biases and drifts in the predictions.

For S2D predictions, model drift is acute and is addressed by multiple approaches for computing anomalies (FIG. 4). One method is to calculate the model

climatology of drifts from hindcasts over a prediction period of interest (for example, the average of lead years 3–7) and, then, subtract that climatology from each prediction for years 3–7 (REF.⁶⁹); this approach works well for short timescale predictions where externally forced trends are less of a factor, but can be problematic for longer timescales. An alternative method is to compute a mean time-evolving drift from a set of hindcasts, subtract that mean drift from a prediction and compute anomalies as differences from the drift-adjusted prediction and time period (such as the previous 15-year average) immediately prior to the prediction⁷⁰. This alternative approach better reduces the effects of an externally forced trend, but raises the issue of how great a role the recent observed period should play in prediction verification. When long-term trends in the hindcasts differ from observations, a further method is to correct biases in the trends in addition to those in the mean model climatology over the hindcast period⁷¹, although such an approach can yield an overestimation of the skill of the system.

Models can also underestimate the magnitude of predictable signals relative to unpredictable internal variability, especially at seasonal and longer timescales in the extratropical North Atlantic sector³³. This underestimation leads to the counter-intuitive implication that models are better at predicting the real climate variability than they are at predicting themselves, a phenomenon termed the ‘signal to noise paradox’, when observed signal to noise ratios are larger than those in models⁷². Given that such features also occur in uninitialized climate simulations of the historical period^{73,74}, and potentially in modelled responses to volcanoes and solar variations⁷², they are not believed to arise from initialization itself. As a result of the signal to noise paradox, it is necessary to take the mean of a very large ensemble to extract the predictable signal and then adjust its variance³³.

Although discrepancies between signal to noise measures in models and observations highlight an important model deficiency, they also imply an optimistic potential to use adjusted climate model outputs to predict the observed system^{33,36}. Additionally, there has been growing interest in the influence of decadal variability on the predictability and skill of seasonal forecasts⁷⁵. Sometimes, the impact of this variability can obscure the gradual skill improvements that are found from advancing the science and modelling⁷⁶.

Clearly, a major challenge for initialized prediction at any timescale is the mean drift of the model away from its initialized state to its preferred systematic error state (FIG. 4). All of the efforts at bias adjustment and drift

correction arise from this fundamental characteristic of model error, but improvements in initialized prediction require increased understanding of the processes and mechanisms at work in the climate system in order to reduce model error.

S2S initialized predictions

All initialized predictions start with a particular observed state that could contribute to some combination of externally forced and internally generated variability. However, owing to the relatively short timescales, sub-seasonal (S2S) predictability is largely an initial value problem in which the atmosphere, ocean, land and sea ice contribute to prediction skill through their memory of the initial state, and not external forcing (FIG. 1). Considerable resources are therefore allocated to initialization of atmosphere and land, including generation of ensemble spread. Ocean initialization and coupling are additionally important, especially in tropical regions, where sources of predictability can come from modes of variability such as the MJO^{6,77}, as well as the stratosphere, both of which are now discussed.

Modes of variability. The MJO is recognized as one of the leading sources of S2S predictability⁷⁸ owing to the strong interaction between the tropics and extratropics on subseasonal timescales⁷⁹. For example, forecast models involved in the SubX and the S2S Prediction Project can predict the MJO skilfully up to 4 weeks^{5,80,81}. Furthermore, skill has been shown in predicting the MJO in a multi-model framework consisting of six SubX models for




Fig. 4 | **Impact of model drift on initialized predictions.** Globally averaged surface air temperature (SAT) predictions from the Decadal Prediction Large Ensemble (DPLE)²⁰ as a function of simulation year. Initial state predictions (blue dots) compare well with observations (black line), but drift (progression of blue dots to red dots) towards the model's systematic error state represented by the uninitialized state (dark grey line; grey shading is range of uninitialized projections).




Fig. 5 | Initialized S2S predictions of the MJO. **a** | Observed outgoing long-wave radiation (OLR) anomalies averaged over 5° S–5° N as a function of the stage of the Madden–Julian Oscillation (MJO). **b–g** | As panel **a** but for various initialized predictions, with OLR anomalies taken as the average of simulations days 15–21. MJO events are identified based on Real-time Multivariate MJO (RMM) index amplitude ≥ 1 . Eastward propagation of MJO-related OLR anomalies is well captured by all six models. S2S, subseasonal to seasonal. Adapted from REF.⁶, © American Meteorological Society. Used with permission.

week 3 predictions averaged over days 15–21 (REF.⁶) (FIG. 5), whereby most reproduce the eastward propagation of outgoing long-wave radiation anomalies. Some models, however, have difficulty in simulating the propagation of the MJO across the Maritime Continent (eastward of 120° E), the so-called Maritime Continent ‘barrier’⁷⁸. MJO-related Rossby wave propagation into the extratropics also provides predictability for extreme events such as storm tracks⁸², atmospheric rivers⁸³ and tornadoes⁸⁴.

S2S predictability is also influenced by the NAO (itself influenced by ENSO⁸⁵), sea ice and the stratosphere⁸⁶, which has a bearing on extremes in large regions of Europe and North America. Using the NCEP Climate Forecast System version 2 (CFSv2) and the Met Office Global Seasonal forecast System 5 (GloSea5), it has been suggested that the NAO exhibits predictability to at least several months ahead^{35,87,88}. Indeed, all SubX models

demonstrate significant NAO skill at week 3, specifically an ACC of ~ 0.27 – 0.5 (REF.⁶).

Similarly, the SAM is a source of predictability and prediction skill of rainfall, temperature and heat extremes over Australia^{89,90}. Although SAM predictability is typically low beyond ~ 2 weeks, there is the potential to make seasonal predictions⁹¹ because of its association with ENSO⁹² and the influence of the stratosphere^{81,93}.

Consideration of these modes offers ‘windows of opportunity’ in S2S prediction, where in certain situations there could be better predictability owing to active periods of the MJO or certain large-scale atmospheric regimes, for example⁹⁴.

Initial state. Given that the land surface varies more slowly than the atmosphere, it also provides a source of predictability for temperature and precipitation on S2S

timescales, the greatest contribution coming from soil moisture⁹⁵. This predictability is most pronounced during boreal spring and summer when synoptic systems have a smaller influence on soil moisture variability. The contribution of soil moisture anomalies to sub-seasonal predictability also varies regionally, with the largest contribution in areas of strong land–atmosphere interactions⁹⁶. As such, the land surface is initialized in most current operational subseasonal prediction systems and all research subseasonal systems (Supplementary Tables 1 and 2). In doing so, improved skill for S2S predictions of temperature and precipitation have been observed, although model errors impact the full realization of this skill^{95,97,98}.

The coupling of the atmosphere to the ocean and sea ice is further thought to be important for predictability at lead times longer than 2 weeks, and, accordingly, ocean–sea ice–atmosphere coupled models are routinely used in operational S2S initialized predictions. For Arctic sea ice, there is rising demand for reliable projections up to months ahead owing to increased human activities. Currently, the best subseasonal models show skilful forecasts of more than 1.5 months ahead⁹⁹. Yet many current operational forecast models lack skill even on timescales of a week¹⁰⁰. Hence, there is more work to be done to improve the S2S forecast skill of Arctic sea ice variables, although many systems are capable of predicting the sea ice extent at seasonal timescales, at least in some regions and seasons^{101–104}.

Sea ice conditions (such as the location of the sea ice edge) can have significant feedback with the atmosphere and, thus, impact the forecast of the coupled system in initialized predictions¹⁰⁵. For example, the largest mid-latitude forecast skill improvements have occurred owing to improved Arctic predictions over eastern Europe, northern Asia and North America relating to sea ice reductions and anomalous anticyclonic circulation¹⁰⁶.

The stratosphere. The largest recognized influence of the stratosphere on the troposphere comes from extreme states of the stratospheric polar vortex, particularly SSWs. SSWs are followed by tropospheric circulation anomalies that can last up to 60 days and resemble the negative phase of the NAO^{107,108}. S2S forecasts initialized near the onset of an SSW thus show increased skill for mid-latitude to high-latitude surface climate¹⁰⁹, and seasonal predictability of the NAO is dependent on the presence of SSWs in ensemble predictions¹¹⁰. Although SSWs are not as common in the southern hemisphere, weakening and warming of the stratospheric polar vortex is predictable a season in advance and, through connections with a negative SAM, can offer some predictability of hot and dry extremes over Australia^{81,93}.

The QBO can further influence the troposphere on S2S timescales. Specifically, phase changes in the QBO modify the strength of the stratospheric polar vortex¹¹¹, in turn affecting the subtropical jet and storm tracks and, hence, surface climate^{112,113}, and the strength of the MJO^{114,115}. For example, the phase of the QBO in the initial state influences the prediction skill of the MJO, with higher skill during easterly QBO boreal winters

compared with westerly QBO winters and improved skill for lead times of 1–10 days¹¹⁶. The prediction skill of the QBO itself is very high on the S2S timescales, with an ACC of 0.85–1.0 at a 1-month timescale⁹³.

S2I initialized predictions

S2I initialized predictions are relatively mature compared with S2S and S2D, as evidenced by the number of national operational meteorological services that maintain state-of-the-art initialized S2I prediction systems^{7,117}. Primary sources and mechanisms of S2I predictability consist of slowly evolving boundary conditions of SST, land surface conditions (moisture, snow cover), sea ice variations¹¹⁸ and stratospheric state. Additional predictability might be gained from the atmospheric composition, not typically represented in S2I models. Each of these factors are now discussed.

ENSO. The largest source of S2I predictability is associated with ENSO. ENSO provides skill in predicting rainfall across the tropics¹¹⁹ and surface climate across the globe given their teleconnections¹²⁰. This predictability skill is primarily derived from subsurface ocean processes¹²¹. Specifically, given that winds and SSTs in the deep tropical Pacific are largely in equilibrium, and the subsurface temperature or thermocline variations are in disequilibrium, capturing the latter in the initial state of ESMs offers predictability¹²¹.

However, ENSO events exhibit a large diversity in spatial patterns, with the location of maximum SST anomalies ranging from the central Pacific to the far-eastern Pacific^{39,122}. ENSO diversity raises predictability issues in terms of precursor mechanisms such as Pacific Meridional Modes^{123–127}, forecast skill^{128,129}, teleconnections¹³⁰, multi-year events¹³¹ and interpretation in the palaeo record¹³² — many of which remain unresolved.

Overall, current state-of-the-art prediction systems are able to predict SSTs in the eastern Pacific up to 6–9 months in advance with modest skill, especially for forecasts initialized in June and verified in the following boreal winter. Yet current prediction systems consistently struggle to predict through the boreal spring season, that is, the so-called spring prediction barrier. The rapid onset or initiation of canonical, eastern Pacific ENSO events also remains a challenge to predict, largely because onset often requires stochastic triggers such as westerly wind bursts^{133,134}. Indeed, inclusion of westerly wind bursts (or other triggers) as stochastic parameterizations has been found to improve model simulations of ENSO¹³⁵ and forecast skill¹³⁶. Prediction of different ENSO types appears to be limited to about 1 month¹³⁷ and, owing to the models' systematic tendency to produce more warming in the east, strong eastern Pacific events are generally better predicted (that is, exhibit better forecast skill) than central Pacific events⁷.

Other modes of variability. Tropical Atlantic SST anomalies are also predictable on S2I timescales. SST anomaly variability in this region is broadly categorized into two spatial patterns. The first is often referred to as the 'Atlantic Niño' and involves many of the feedback

mechanisms noted for ENSO¹³⁸, but is shorter lived and weaker. In comparison with ENSO, however, the Atlantic Niño is less studied and also less predictable^{139,140}. The second pattern of variability is referred to as the Atlantic Meridional Mode⁸⁷. It is estimated that the Atlantic Meridional Mode is predictable one to two seasons in advance, with the mechanisms for predictability largely stemming from near-surface air–sea interactions (thermocline variability is of secondary importance). However, even with some indications of successful predictions in certain circumstances including interactions with the tropical Pacific¹³⁸, as with all timescales of initialized predictions, persistent regional systematic errors with current initialized Earth prediction systems continue to be a factor in limiting the predictive abilities of tropical Atlantic S2I variability^{141,142}.

Much like the Atlantic, Indian Ocean SST anomaly variability is weaker and may be less predictable than the Pacific, but is important for regional teleconnections and impacts. Indian Ocean SST variability has three distinct patterns of interest: the IOD, which can be triggered by ENSO but can also emerge independently^{88,143}; a basin-wide pattern that is an ENSO teleconnection¹⁴⁴; and a meridional mode pattern that depends on near-surface air–sea interactions similar to that in the Atlantic¹⁴⁵. Earth System prediction models typically struggle to predict the connection between ENSO and the IOD, the northward propagation of the meridional mode and the persistence of the IOD, except in large-amplitude cases¹⁴⁶. The IOD also can affect processes on the S2S timescale¹⁴⁷, including the MJO, and even the extratropics. There are also other possible sources of S2I predictive skill involving the NAO¹⁴⁸ and the Atlantic Ocean state that appear to drive aspects of summer European rainfall¹⁴⁹.

Land surface processes. Slowly varying S2I soil moisture anomalies influence the prediction skill for precipitation and temperature¹⁵⁰. Currently, the memory resulting from large soil moisture anomalies in the initial conditions is believed to last ~2–3 months¹⁵¹, but there are case by case examples where predictability can be considerably longer under conditions where soil moisture anomalies persist for more than one season, particularly for surface temperature. Indeed, some seasonal temperature predictability has been confirmed to arise from soil moisture, but the realization of skill is severely hampered by model biases^{152,153}. Thus, reducing model error in the land surface components could considerably improve forecast skill, as seen in a large sample of initialized Earth System prediction experiments¹⁷.

The stratosphere. Improved surface prediction resulting from stratosphere-related processes has been demonstrated on the seasonal timescale: having a higher vertical resolution in the stratosphere in a GCM captures SSWs earlier compared with the standard model configuration and has a positive influence on the simulations of European surface climate¹⁵⁴. Southern hemisphere SSWs also affect predictions of Australian extremes^{81,93}. The QBO, discussed earlier with respect to S2S predictability, has also been shown to lead to enhanced predictability

on seasonal timescales^{155,156}, is predictable up to several years ahead¹⁵⁷ and can also involve the MJO¹¹⁶.

Other sources of predictive skill. There are additional sources and mechanisms for S2I predictability that are not particularly well modelled in S2I prediction. For example, slowly evolving greenhouse gases such as carbon dioxide and methane are known to be a source of forecast skill owing to their role as external forcing agents¹⁵⁸. However, an approximate time history of carbon dioxide, methane and chlorofluorocarbons is typically specified and not predicted, thus limiting the potential to capture S2I variability or regional effects. Moreover, dust and aerosol concentrations are known to affect human health, but these changes in atmospheric composition are usually not included in prediction systems.

S2D initialized predictions

There is a high level of interest in, and expectations of, initialized Earth System predictions on timescales beyond S2S and S2I. For example, even with their limitations, there is evidence of skill in predicting surface temperature over and above that of simple persistence (FIG. 6a,b), and also precipitation and sea level pressure when using large multi-model ensembles, albeit with less skill³⁶. These skilful multi-year predictions of precipitation over land indicate potential benefit to communities, as demonstrated with summer drought indicators in major European agricultural regions being predictable on multi-year timescales¹⁵⁹. Here, we review the evidence for processes and mechanisms acting on the S2D timescale that could contribute to the skill of initialized predictions^{12,36}.

Modes of decadal SST variability. Processes and mechanisms have been identified that could provide skill for fundamental quantities such as SST in initialized predictions. Attention has been focused on AMV¹⁶⁰, but predictions of PDV^{160,161} — which are often described in terms of the Interdecadal Pacific Oscillation (IPO)¹⁶² over the Pacific basin and the Pacific Decadal Oscillation^{163,164} over the north Pacific — are also of interest. Other modes of variability associated with decadal timescales include the Meridional Modes¹⁶⁵ and the North Pacific Gyre Oscillation¹⁶⁶.

Basin-wide warming and cooling patterns of SSTs and upper ocean heat content (averaged temperature for 0–400 m) have also been shown to characterize decadal timescale variability in the Indian Ocean^{167–169}, as have decadal variations of the IOD^{56,170}. Decadal variability in the Indian Ocean could influence warming events near the Australian west coast^{171,172}. Furthermore, a rapid rise in Indian Ocean subsurface heat content in the 2000s in observations and model simulations is associated with a redistribution of heat from the Pacific to the Indian Ocean and has been suggested to account for a large portion of the global ocean heat gain during that period^{173,174}. IPO variability could thus be affecting Indian Ocean variability, transmitted through both the atmospheric and oceanic bridges¹⁷⁵. These low-frequency connections have been implicated in




Fig. 6 | S2D predictions and aspects of time-evolving globally averaged temperature. **a** | Prediction skill, measured as the anomaly correlation coefficient, of sea surface temperature (SST) averaged over lead years 5–9 from the Decadal Prediction Large Ensemble (DPLE)²⁰; darker red indicates higher skill. **b** | Improvement in prediction skill from initialized predictions in panel **a** over and above a persistence prediction; darker red indicates better skill in the initialized predictions, thus showing the value-added of initialized predictions. **c** | Schematic of the ‘rising staircase’, illustrating how natural decadal-scale temperature fluctuations (blue) are tilted upwards owing to anthropogenic greenhouse gas emissions (red), producing accelerated warming in some decades and reduced warming in others. **d** | Time series of observed global mean surface air temperature (SAT) anomalies (land surface air temperature (LSAT) and sea surface temperature (SST)) showing characteristics of the rising staircase: accelerated warming over 1980–2000 and 2014–present, and a slowdown in the rate of warming over 2000–2014. S2D, seasonal to decadal. Panels **a** and **b** adapted from REF.²⁰ © American Meteorological Society. Used with permission. Panel **c** adapted with permission from National Academies Press, from REF.²⁶⁷; permission conveyed through Copyright Clearance Center Inc. Panel **d** adapted, with permission, from REFS^{270,271}, NASA.

modulating interannual variability associated with the IOD on decadal timescales^{172,176}.

One issue that remains to be resolved for S2D related to prediction skill is whether there are well-defined timescales of variability that are distinct from the background of climatic noise; that is, whether there are modes of large-scale variability that might display a statistically significant spectral peak in the decadal to multi-decadal range and that could be predicted. Such signals could offer the best prospect for long-term predictability, but on this timescale there is more of a broadband spectral peak. For example, CMIP5 control simulations showed patterns and multi-decadal timescales of variability in the Pacific associated with the IPO that resemble observations but with lower amplitude¹⁷⁷. Moreover, analysis of three generations of climate models (CMIP3, CMIP5 and CMIP6) shows progressive improvement of climate models’ simulations of PDV¹⁷⁸. However, there was no convincing evidence across these state-of-the-art coupled models for distinct oscillatory signals, other than on the

interannual (years 3–7) ENSO timescales¹⁷⁹. These observations suggest, as noted previously, that low frequency variability on interdecadal timescales is characterized by broadband rather than oscillatory behaviour.

Global temperatures. The idealized ‘rising staircase’ (FIG. 6c) of global mean surface temperature (GMST) trends represents actual epochs of larger or smaller amplitude-positive GMST trends (FIG. 6d) in a world with steadily increasing positive radiative forcing from increasing greenhouse gases¹⁸⁰. This increase in radiative forcing means that the entire Earth System warms continuously, but the manifestation of that warming at the Earth’s surface on decadal timescales depends on how heat is redistributed in the climate system: if more heat remains near the ocean surface, the GMST rate of warming will be larger, but if more heat is distributed into the deeper ocean, then the GMST trend will be reduced^{44,181}.

It is recognized that the slowdown in the rate of GMST warming in the early 2000s was likely a combination

of internal variability from the negative phase of the IPO^{182–186} and/or variations in the strength of the Atlantic meridional overturning circulation¹⁸⁷, both of which acted to redistribute heat into the subsurface ocean. However, there is disagreement on whether the heat is primarily stored in the tropics¹⁷⁴ or at high latitudes¹⁸¹. External forcing from a collection of moderate-sized volcanic eruptions¹⁸⁸ and from anthropogenic aerosols¹⁸⁹ might have also played a role in the slowdown, although their contribution is not entirely settled¹⁹⁰.

Initialized predictions have been shown to successfully predict the onset of the GMST warming slowdown, linked to increased ocean heat uptake in the tropical Pacific and Atlantic Oceans^{183,191}. Spatial patterns of predicted 20-year surface air temperature trends have been shown to depend on the initial state of the Pacific Ocean¹⁹², with initialized model predictions exhibiting a large spread in projected multi-decadal global warming unless the initial state of the Pacific Ocean is known and well represented in the model. Apart from its connection to the recent global warming slowdown, the negative phase of the IPO has also been linked to regional climate changes at higher latitudes, including the rate of Arctic sea ice decrease in the early 2000s (REF.¹⁹³) and Antarctic sea ice expansion during that same period^{194,195}.

Statistical methods⁴⁷ and initialized predictions^{70,196} foretold a transition of the IPO in the tropical Pacific from negative to positive in the 2014–2015 time frame, with a resumption of more rapid rates of global warming thereafter. There is observational evidence that this IPO transition also contributed to initiating rapid Antarctic sea ice retreat¹⁹⁷.

There is a chronic shortage of observed data in the ocean to document heat redistribution. In models, this redistribution has been shown to involve the subtropical cells in the Pacific, Antarctic Bottom Water formation and the AMOC in the Atlantic^{2,44}, as well as changes in the zonal slope of the equatorial thermocline^{182,198} associated with changes in tropical winds. However, deciphering decadal timescale variability in the observed climate system, and interpreting such variability in the context of initialized predictions, is complicated by the presence of external forcings (such as anthropogenic and volcanic aerosols and solar forcing) that can produce decadal variability in the Pacific¹⁸⁹ or Atlantic^{199,200} with similar patterns to presumptive internally generated decadal climate variability^{180,201,202}.

Interactions between ocean basins. Interactions between various ocean basins are one of the most compelling science questions that have arisen regarding the origins and nature of decadal climate variability, with implications for initialized prediction skill^{160,203,204}. For instance, if a skilful prediction of climate in one basin is achieved, then skilful simulations in the other basins could follow (if the models capture these connections realistically), thus improving the skill of initialized S2D predictions.

SST variability in one ocean basin can affect the others through the tropical large-scale east–west atmospheric Walker Circulation, although the direction of those influences differs^{204,205}. For example, model simulations

have indicated that decadal timescale variability in the Atlantic could produce decadal timescale variability in the Pacific^{61,206–208}. PDV can also affect the Atlantic^{194,209,210} and control a large fraction of decadal variability in the Indian Ocean^{58,172,211–213}. Similarly, the Indian Ocean could influence decadal variability in the Pacific^{168,203,214}. There also could be staggered responses based on decadal timescales, with the tropical Pacific driving the tropical Atlantic on interannual timescales, with the Atlantic then affecting the Indian Ocean and, subsequently, the Pacific on decadal timescales^{215,216}. It has further been postulated that the tropical Atlantic and Pacific Oceans are mutually interactive on decadal timescales, with each alternately affecting the other²⁰⁵, and that the tropical Pacific could be driving the extratropical Pacific²¹⁷.

External forcing, particularly from time-evolving anthropogenic aerosols, is another factor that could produce decadal climate variability and inter-basin connections^{189,199,218}. Such fundamental interactions all currently fall under the heading of a compelling research frontier that, with increased understanding, will certainly advance the science of initialized prediction.

Summary and future perspectives

Numerical models initialized with observations for specific time periods and integrated forward in time provide a continuum of predictions on different timescales from S2S to S2I and S2D. Results so far demonstrate initialized prediction skill for variables such as surface temperature and key modes of atmospheric and ocean variability. Such skill has been demonstrated, for example, for the MJO on S2S timescales, for ENSO on S2I timescales and for surface temperatures in most ocean regions on S2D timescales. Yet, despite progress in predictions and processes, there are still many challenges and priorities for future research.

Model error. Almost every science-related aspect of subseasonal to decadal climate variability has considerable uncertainty associated with it. Therefore, apart from fundamental scientific understanding, perhaps the key obstacle to progress is model error, particularly with regards to biases and drifts. Progress thus requires model improvement, developments of which are difficult but not impossible. In recent years, for instance, model development work has been undertaken in the coupled space, improving simulation of atmosphere–ocean phenomena that give rise to predictability (such as the MJO and ENSO), and therefore minimizing the exacerbation of drift when developed in isolation. Model improvements depend critically on our understanding of processes and mechanisms and how they work in the climate system, as it is difficult to model what is not understood. Therefore, enhanced observational and analysis projects must continue to provide the knowledge base from which to make improvements to the model simulations.

Model error remains a significant obstacle against which future progress will be measured, with profound implications for possible applications to stakeholder communities. Such applications could include

energy supply (wind, solar) and demand²¹⁹, agriculture (drought, freezing), transport²²⁰ and numerous others spanning a range of timescales. Notably, S2S prediction could inform preparedness for specific large-scale extreme events weeks ahead⁵, and S2I and S2D initialized predictions are beginning to inform planning at ranges between the seasonal and multi-decadal climate change timescales²²¹.

In addition to coupled model development, increased model resolution has also shown the ability to improve model bias and the signal to noise ratio. Consequently, the benefit of increased model resolution is one of the research frontiers of initialized prediction. However, such increased resolution must also be accompanied by comparable increases in the quality of the physical parameterizations such as cloud feedback and cloud–aerosol interactions. Although we are still very likely decades away from having global coupled models (and suitable machines) capable of explicitly resolving processes that would improve model bias (such as atmospheric convection and ocean eddies), approaches have been developed to reduce computational cost and bias. These approaches include flux correction techniques²²², parameter estimation²²³, reducing the precision of some variables²²⁴ and stochastic modelling²²⁵. Additionally, machine learning techniques are providing indications of improving predictive skill. For example, a deep-learning approach using a statistical forecast model has been shown to produce skilful ENSO forecasts for lead times of up to 1.5 years²²⁶. Utilization of GPU-based computer architectures could become useful and open the way to better parametrizations that depend on intensive calculations that can be addressed with GPU architectures.

Initialization. Integrating the vast amount of observed information into an ESM is central to the S2D prediction. Traditionally, the most advanced data assimilation techniques were implemented in the atmospheric component. In the last decade, however, there has been growing interest in how to fully utilize relevant satellite and in situ observations to improve S2S and S2I predictions. Coupled ocean–atmosphere data assimilation^{28,227,228} shows promising evidence that coupling can reduce ‘initialization shock’ and improve forecast performance on timescales of weeks to decades²²⁹. The advancement has led to coupled reanalysis products for both the ocean and the atmosphere (CFSR by NCEP²³⁰ and CERA by ECMWF²³¹) and is expected to substantially improve S2S and S2I predictions.

Compared with S2S and S2I predictions, there remain critical obstacles to how to initialize decadal predictions. First, there is a lack of observations. S2D models need to be initialized in the 1960s and 1970s in order to calibrate the decadal prediction systems and achieve the potential to capture the evolution of low-frequency modes of variability (such as PDV and AMV). Reconstruction of the global ocean subsurface temperature and salinity prior to the advent of Argo floats remains a large problem. Currently, most modelling centres performing decadal predictions do not carry out their own assimilation exercise; rather, they simply nudge some reanalysis products in the ocean and atmosphere (Supplementary Table 3).

How to best initialize the ocean without reliable subsurface observations, and how the inhomogeneity of the observations can impact model performance, have not been carefully investigated.

Building ensembles is another key obstacle to decadal prediction, as common practice in the community is to use an ensemble of ten members following the CMIP5 and CMIP6 experimental designs. A large ensemble consisting of 40 members can provide better opportunities for skilful predictions of low-frequency climate variability over land in selected regions²⁰. However, compared with the atmosphere, there is very limited understanding of the mechanisms and uncertainty associated with the low-frequency internal variability in the ocean owing to the lack of long-term observations of the subsurface ocean, and thus lack of guidance as to how to build the ensemble. Machine learning methods could help address this problem, although the lack of long-term subsurface ocean observations will always be a factor for the S2D timescale. Finally, a major constraint is computational capability, both for initialization and for running adequate numbers of ensembles to improve skill³³. The future of initialized prediction will depend on computational resources balanced with factors involving increased resolution, machine learning, use of new high-performance computing architectures and developments in exascale computing.

Predictability of internal variability. There are considerable future challenges for understanding internal variability in the context of initialized prediction. These include the need to have a better understanding and better estimates of predictability. Additionally, research is needed regarding why models appear to underestimate the magnitude of predictable signals compared with unpredictable variability, and this involves the response to external forcing as well²³².

One issue that remains to be resolved for S2D initialized predictions is whether there are well-defined processes and mechanisms that, if initialized properly, could provide predictable signals distinct from the background of climatic noise. Signals from PDV and AMV offer the best prospect for long-term predictability. Strong low-frequency variability in palaeoclimate ‘proxy’ records, which is not captured by most climate models, suggests either that models do indeed underestimate low-frequency modes of variability or that proxy observations contain significant residual non-climatic sources of variation, or some combination thereof^{233–236}. Even if there is no distinct low-frequency (oscillating) phenomenon, predictability on decadal timescales could also come from memory and slowly varying components of the Earth System, such as the slow propagation of oceanic planetary waves^{237,238} or natural volcanic forcing⁴⁷, and initialization could be expected to contribute to skill in such cases.

Expanding predicted variables. There is interest in, and corresponding applications for, expanding beyond the prediction of surface temperature, precipitation and SST. Predictions of the frequency of extreme events such as tropical storms and hurricanes have great potential as

climate services. There have been efforts at predicting soil moisture with implications for drought prediction²³⁹ and ecosystem respiration²⁴⁰, as well as snowpack with ramifications for water resources^{241,242} and marine heatwaves²⁴³. There is also a great societal need for prediction of sea ice on S2I and S2D timescales. Some S2I models show some skill in predicting the sea ice edge in the Arctic²⁴⁴, whereas S2S models show a very wide range of skill in predicting the sea ice edge in the Arctic, with the most skilful models producing useful forecasts up to 45 days⁹⁹. Although the potential for skilful initialized predictions of Arctic sea ice on S2S timescales has improved in the last decade, there is still a lot more to be explored and improved¹⁰¹. We still need to understand what are the key processes driving subseasonal variations of sea ice and to improve the representation of these processes in the S2S models. Improved coupled data assimilation of the ocean, sea ice and atmospheric coupled system can help improve initial conditions for coupled forecasts and, concomitantly, the forecast skill of features that are sensitive to the initial state^{14,245,246}.

Other important aspects of the cryosphere relevant to initialized prediction on S2D timescales are ice sheets. As new interactive ice sheet simulations and spin-up procedures come increasingly online²⁴⁷, this will provide an additional opportunity for initialized S2D predictions.

Air pollution and air quality are other very society-relevant applications that have been largely unexplored owing to the lack of inclusion of interactive tropospheric chemistry in most S2S, S2I and S2D models. However, new comprehensive ESMs, such as the Community Earth System Model with the Whole Atmosphere Community Climate Model as its atmospheric component (CESM2-WACCM²⁴⁸), will be able to explore this research area.

In the broader Earth System, there is growing interest in predicting the biosphere and biogeochemical state variables and fluxes that could inform management decisions. Skilful initialized predictions of SST on S2S timescales can engender predictability of fish yields in the California Current System²⁴⁹ and other large marine ecosystems²⁵⁰. S2S initialized predictions of heat stress and coral bleaching risk have also demonstrated considerable skill and have provided critical advanced warning for coral reef scientists, managers and stakeholders²⁵¹. SST anomalies in the western tropical Pacific and northern subtropics, often associated with ENSO events, appear to be skilful precursors for variations in temperature and related biological productivity along the US West Coast on S2I timescales²⁵².

Emerging literature on S2D predictions of biogeochemistry in the terrestrial biosphere and ocean suggests that slowly evolving state variables could enable prediction of biogeochemically relevant quantities with

greater skill than physical state variables such as temperature and precipitation. For example, predictions of marine net primary production by photosynthesizing phytoplankton (including algae, eukaryotes and cyanobacteria) might foretell future potential fisheries catches, predict harmful algal blooms²⁵³ and aid with fisheries management strategies^{253–256}, as would skilful predictions of ocean oxygen content or acidity^{257,258}. Reliable forecasts of the changing global carbon budget, including the rate of ocean carbon absorption^{216,259–261} or the rate of terrestrial biosphere–atmosphere net ecosystem exchange^{240,259}, could help to generate forecasts of atmospheric CO₂ growth rate and contribute to CO₂ emission management strategies. Additionally, there has been demonstrated S2I skill at predicting net primary production related to fire risk²⁶².

Recently reported skilful predictions of chlorophyll concentrations over the global oceans at seasonal to multi-annual timescales have been related to the successful simulation of the chlorophyll response to ENSO, and to the winter re-emergence of subsurface nutrient anomalies in the extratropics²⁵⁵. Chlorophyll not only responds to ENSO, but can also constitute a potentially useful ENSO precursor²⁶³.

In the ocean biogeochemical system, variables of interest for prediction are rarely directly observed at the spatial and temporal scales needed for forecast verification, regardless of the timescale of the prediction^{264,265}. Thus, most of the literature is focused on the potential to make predictions of these quantities, rather than on skill as measured by historical observations^{254,256,259,260}, with exceptions^{216,257,258}. On the global scale, verification is limited to variables measured or derived from satellite observations, such as ocean chlorophyll²⁵⁵, marine primary productivity²⁰ or interpolated estimates of the surface ocean partial pressure of CO₂ (REF²⁶¹). Nevertheless, there is promising potential to make ocean biogeochemical initialized predictions across multiple timescales.

For S2S, S2I and S2D initialized predictions to be useful, they must be shown to be not only skilful but reliable²⁶⁶, and this is a considerable challenge that the community is only starting to attempt to address^{5,21}. The ultimate challenge in this emerging area of research, and one that is igniting excitement and interest in the scientific community, is to provide predictions with maximum skill that take into account all relevant processes across subseasonal to decadal timescales^{267–269}. Towards that end, initialized prediction is already put to task and being applied in various sectors even as improvements in understanding and prediction capability are being improved, thus driving rapid advances in this burgeoning field.

Published online: 13 April 2021

1. Meehl, G. A. et al. Decadal prediction. *Bull. Am. Meteorol. Soc.* **90**, 1467–1486 (2009).
2. Meehl, G. A., Hu, A., Arblaster, J. M., Fasullo, J. & Trenberth, K. E. Externally forced and internally generated decadal climate variability associated with the Interdecadal Pacific Oscillation. *J. Clim.* **26**, 7298–7310 (2013).
3. Hawkins, E. & Sutton, R. (2009). The potential to narrow uncertainty in regional climate predictions.

- Bull. Am. Meteorol. Soc.* **90**, 1095–1108 (2009).
4. Lehner, F. et al. Partitioning climate projection uncertainty with multiple large ensembles and CMIP5/6. *Earth Syst. Dynam.* **11**, 491–508 (2020).
5. Vitart, F. & Robertson, A. W. The Sub-Seasonal to Seasonal Prediction Project (S2S) and the prediction of extreme events. *npj Clim. Atmos. Sci.* **1**, 3 (2018).

6. Pegion, K. et al. The Subseasonal Experiment (SubX). *Bull. Amer. Meteorol. Soc.* **100**, 2043–2060 (2019).
7. Kirtman, B. P. et al. The North American multimodel ensemble: phase-1 seasonal to interannual prediction; phase-2 toward developing intraseasonal prediction. *Bull. Amer. Meteorol. Soc.* <https://doi.org/10.1175/BAMS-D-12-00050.1> (2014).

8. Boer, G. J. et al. The Decadal Climate Prediction Project (DCPP) contribution to CMIP6. *Geosci. Model Dev.* **9**, 3751–3777 (2016).
9. Palmer, T. N., Doblas-Reyes, F. J., Weisheimer, A. & Rodwell, J. J. Toward seamless prediction: calibration of climate change projections using seasonal forecasts. *Bull. Amer. Meteorol. Soc.* **89**, 459–470 (2008).
10. Branstator, G. & Teng, H. Potential impact of initialization on decadal predictions as assessed for CMIP5 models. *Geophys. Res. Lett.* <https://doi.org/10.1029/2012GL051974> (2012).
11. Barnett, T. et al. On the prediction of the El Niño of 1986–1987. *Science* **241**, 192–196 (1988).
12. Kushnir, Y. et al. Towards operational predictions of the near-term climate. *Nat. Clim. Chang.* <https://doi.org/10.1038/s41558-018-0359-7> (2019).
13. Lean, P. et al. Continuous data assimilation for global numerical weather prediction. *Q J R Meteorol. Soc.* <https://doi.org/10.1002/qj.3917> (2020).
14. Sandery, P. A., O’Kane, T. J., Kitsios, V. & Sakov, P. Climate model state estimation using variants of EnKF coupled data assimilation. *Mon. Weather Rev.* **148**, 2411–2431 (2020).
15. Johnson, C., Hoskins, B. J. & Nichols, N. K. A singular vector perspective of 4D-Var: filtering and interpolation. *Q J R Meteorol. Soc.* **131**, 1–19 (2005).
16. Magnusson, L., Nycander, J. & Kallen, E. Flow-dependent versus flow-independent initial perturbations for ensemble prediction. *Tellus A* **61**, 194–209 (2009).
17. Infanti, J. M. & Kirtman, B. P. Prediction and predictability of land and atmosphere initialized CCSM4 climate forecasts over North America. *J. Geophys. Res. Atmos.* **121**, 12,690–12,701 (2016b).
18. Trenary, L., DelSole, T., Tippet, M. K. & Pegion, K. A new method for determining the optimal lagged ensemble. *J. Adv. Model Earth Syst.* **9**, 291–306 (2017).
19. Kirtman, B. P. & Min, D. Multi-model ensemble ENSO prediction with CCSM and CFS. *Mon. Weather Rev.* <https://doi.org/10.1175/2009MWR2672.1> (2009).
20. Yeager, S. G. et al. Predicting near-term changes in the earth system: a large ensemble of initialized decadal prediction simulations using the community earth system model. *Bull. Am. Meteorol. Soc.* **99**, 1867–1886 (2018).
21. Smith, D. M. et al. Real-time multi-model decadal climate predictions. *Clim. Dyn.* **41**, 2875–2888 (2013a).
22. MacLachlan, C. et al. Global Seasonal Forecast System version 5 (GloSea5): a high resolution seasonal forecast system. *Q J R Meteorol Soc.* <https://doi.org/10.1002/qj.2396> (2014).
23. Muñoz-Sabater et al. Assimilation of SMOS brightness temperatures in the ECMWF integrated forecasting system. *Q J R Meteorol Soc.* <https://doi.org/10.1002/qj.3577> (2019).
24. Drewitt, G., Berg, A. A., Merryfield, W. J. & Lee, W.-S. Effect of realistic soil moisture initialization on the Canadian CanCM3 seasonal forecast model. *Atmos. Ocean* **50**, 466–474 (2012).
25. Polkova, I., Köhl, A. & Stammer Climate-mode initialization for decadal climate predictions. *Clim. Dyn.* **53**, 7097–7111 (2019).
26. Smith, D. M., Eade, R. & Pohlmann, H. A comparison of full-field and anomaly initialization for seasonal to decadal climate prediction. *Clim. Dyn.* **41**, 3325–3338 (2013).
27. Volpi, D., Guemas, V. & Doblas-Reyes, F. J. Comparison of full field and anomaly initialization for decadal climate prediction: towards an optimal consistency between the ocean and sea-ice anomaly initialization state. *Clim. Dyn.* **49**, 1181–1195 (2017).
28. Penny, S. G., et al. Coupled data assimilation for integrated earth system analysis and prediction: goals, challenges and recommendations. Technical report (World Meteorological Organisation, 2017).
29. Williams, K. D. et al. The Met Office Global Coupled Model 2.0 (GC2) configuration. *Geosci. Model Dev.* **8**, 1509–1524 (2015).
30. Becker, E. & Van Den Dool, H. Probabilistic seasonal forecasts in the North American multimodel ensemble: a baseline skill assessment. *J. Clim.* **29**, 3015–3026 (2016).
31. Kadow, C. et al. Decadal climate predictions improved by ocean ensemble dispersion filtering. *J. Adv. Model. Earth Syst.* **9.2**, 1138–1149 (2017).
32. Dobrynin, M. et al. Improved teleconnection-based dynamical seasonal predictions of boreal winter. *Geophys. Res. Lett.* **45**, 3605–3614 (2018).
33. Smith, D. M. et al. North Atlantic climate far more predictable than models imply. *Nature* **583**, 796–800 (2020).
34. Richter, J. H. et al. Subseasonal prediction with and without a well-represented stratosphere in CESM1. Weather and Forecasting, <https://journals.ametsoc.org/view/journals/efw/aop/WAF-D-20-0029.1/WAF-D-20-0029.1.xml> (2020).
35. Scaife, A. A. et al. Skillful long-range prediction of European and North American winters. *Geophys. Res. Lett.* **41**, 2514–2519 (2014).
36. Smith, D. M. et al. Robust skill of decadal climate predictions. *npj Clim. Atmos. Sci.* **2**, 13 (2019).
37. Athanasiadis, P. J. et al. Decadal predictability of North Atlantic blocking and the NAO. *NPJ Clim. Atmos. Sci.* **3**, 20 (2020).
38. Nie, Y. et al. Stratospheric initial conditions provide seasonal predictability of the North Atlantic and Arctic oscillations. *Env. Res. Lett.* **14**, 3 (2019).
39. Capotondi, A. et al. Understanding ENSO diversity. *Bull. Am. Meteorol. Soc.* **96**, 921–938 (2015).
40. Cobb, K. M. et al. Highly variable El Niño–Southern Oscillation throughout the Holocene. *Science* **339**, 67–70 (2013).
41. Capotondi, A. & Sardeshmukh, P. D. Is El Niño really changing? *Geophys. Res. Lett.* <https://doi.org/10.1002/2017GL074515> (2017).
42. Grothe, P. R. et al. Enhanced El Niño–Southern Oscillation variability in recent decades. *Geophys. Res. Lett.* <https://doi.org/10.1029/2019GL083906> (2019).
43. Deser, C., Phillips, A. S. & Alexander, M. A. Twentieth century tropical sea surface temperature trends revisited. *Geophys. Res. Lett.* **37**, L10701 (2010).
44. Meehl, G. A., Arblaster, J. M., Fasullo, J., Hu, A. & Trenberth, K. E. Model-based evidence of deep ocean heat uptake during surface temperature hiatus periods. *Nat. Clim. Change* **1**, 360–364 (2011).
45. Mann, M. E. & Emanuel, K. A. Atlantic hurricane trends linked to climate change. *Eos* **87**, 233–241 (2006).
46. Mann, M. E., Steinman, B. A. & Miller, S. K. On forced temperature changes, internal variability and the AMO (‘Frontier’ article). *Geophys. Res. Lett.* **41**, 3211–3219 (2014).
47. Mann, M. E. et al. Predictability of the recent slowdown and subsequent recovery of large-scale surface warming using statistical methods. *Geophys. Res. Lett.* **43**, 3459–3467 (2016).
48. Steinman, B. A., Frankcombe, L. M., Mann, M. E., Miller, S. K. & England, M. H. Response to comment on “Atlantic and Pacific multidecadal oscillations and Northern Hemisphere temperatures”. *Science* **350**, 1326 (2015).
49. Roemmich, D. & Gilson, J. The 2004–2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. *Prog. Oceanogr.* **82**, 81–100 (2009).
50. Roemmich, D. et al. On the future of Argo: a global, full-depth, multi-disciplinary array. *Front. Mar. Sci.* <https://doi.org/10.3389/fmars.2019.00439> (2019).
51. Thompson, D. M., Cole, J. E., Shen, G. T., Tudhope, A. W. & Meehl, G. A. Early twentieth-century warming linked to tropical Pacific wind strength. *Nat. Geosci.* **8**, 117–121 (2015).
52. Cook, E. R. et al. Megadroughts in North America: placing IPCC projections of hydroclimatic change in a long-term paleoclimatic context. *J. Quat. Sci.* **25**, 48–61 (2010).
53. Emile-Geay, J., Cobb, K. M., Mann, M. E. & Wittenberg, A. T. Estimating central equatorial Pacific SST variability over the past millennium. Part II: reconstructions and implications. *J. Clim.* **26**, 2329–2352 (2013).
54. Linsley, B. K., Wu, H. C., Dassié, E. P. & Schrag, D. P. Decadal changes in South Pacific sea surface temperatures and the relationship to the Pacific decadal oscillation and upper ocean heat content. *Geophys. Res. Lett.* **42**, 2358–2366 (2015).
55. Buckley, B. M. et al. Interdecadal Pacific Oscillation reconstructed from trans-Pacific tree rings: 1350–2004 CE. *Clim. Dyn.* **53**, 3181–3196 (2019).
56. Abram, N. J. et al. Paleoclimatic perspectives on the Indian Ocean dipole. *Quat. Sci. Rev.* <https://doi.org/10.1016/j.quascirev.2020.106302> (2020).
57. Sanchez, S. C., Charles, C. D., Carriguiry, J. D. & Villaseca, J. A. Two centuries of coherent decadal climate variability across the Pacific North American region. *Geophys. Res. Lett.* **43**, 9208–9216 (2016).
58. Abram, N. J. et al. Coupling of Indo-Pacific climate variability over the last millennium. *Nature* **579**, 385–392 (2020).
59. Konecky, B., Dee, S. G. & Noone, D. WaxPSM: a forward model of leaf wax hydrogen isotope ratios to bridge proxy and model estimates of past climate. *J. Geophys. Res. Biogeosci.* <https://doi.org/10.1029/2018JG004708> (2019).
60. Neukom, R. et al. Consistent multi-decadal variability in global temperature reconstructions and simulations over the common era. *Nat. Geosci.* **12**, 643 (2019).
61. McGregor, H. V. et al. Robust global ocean cooling trend for the pre-industrial common era. *Nat. Geosci.* **8**, 671–677 (2015).
62. Tierney, J. E. et al. Tropical sea surface temperatures for the past four centuries reconstructed from coral archives. *Paleoceanography* **30**, 226–252 (2015).
63. Goosse, H. et al. Reconstructing surface temperature changes over the past 600 years using climate model simulations with data assimilation. *J. Geophys. Res.* **115**, D09108 (2010).
64. Hakim, G. J. et al. The last millennium climate reanalysis project: framework and first results. *J. Geophys. Res. Atmos.* **121**, 6745–6764 (2016).
65. Steiger, N. J., Jason, E. S., Cook, E. R. & Cook, B. I. A reconstruction of global hydroclimatic and dynamical variables over the common era. *Sci. Data* **5**, 180086 (2018).
66. Evans, M. N., Tolwinski-Ward, S. E., Thompson, D. M. & Anchukaitis, K. J. Applications of proxy system modeling in high resolution paleoclimatology. *Quat. Sci. Rev.* **76**, 16–28 (2013).
67. Dee, S. et al. PRISM: an open-source framework for PROXY system modeling, with applications to oxygen-isotope systems. *J. Adv. Model. Earth Syst.* **7**, 1220–1247 (2015).
68. Becker, E., Dool, den, H. V. & Zhang, Q. Predictability and forecast skill in NMME. *J. Clim.* **27**, 5891–5906 (2014).
69. Doblas-Reyes, F. J. et al. Initialized near-term regional climate change prediction. *Nat. Commun.* **4**, 1715 (2013).
70. Meehl, G. A., Hu, A. & Teng, H. Initialized decadal prediction for transition to positive phase of the Interdecadal Pacific Oscillation. *Nat. Commun.* <https://doi.org/10.1038/NCOMMS11718> (2016).
71. Kharin, V. V., Boer, G. J., Merryfield, W. J., Scinocca, J. F. & Lee, W.-S. Statistical adjustment of decadal predictions in a changing climate. *Geophys. Res. Lett.* **39**, L19705 (2012).
72. Scaife, A. A. & Smith, D. A signal-to-noise paradox in climate science. *npj Clim. Atmos. Sci.* **1**, 28 (2018).
73. Sévellec, F. & Drijfhout, S. S. The signal-to-noise paradox for interannual surface atmospheric temperature predictions. *Geophys. Res. Lett.* **46**, 9031–9041 (2019).
74. Zhang, W. & Kirtman, B. Estimates of decadal climate predictability from an interactive ensemble model. *Geophys. Res. Lett.* **46**, 3387–3397 (2019).
75. Weisheimer, A. et al. How confident are predictability estimates of the winter North Atlantic oscillation? *Quart. J. R. Meteorol. Soc.* <https://doi.org/10.1002/qj.3446> (2019).
76. Barnston, A. G., Tippet, M. K., L’Heureux, M. L., Li, S. & DeWitt, D. G. Skill of real-time seasonal ENSO model predictions during 2002–11: is our capability increasing? *Bull. Am. Meteorol. Soc.* **93**, 631–651 (2012).
77. Robertson, A. W. & Vitart, F. (eds) *Sub-seasonal to Seasonal Prediction* (Elsevier, 2018).
78. Kim, H., Vitart, F. & Waliser, D. E. Prediction of the Madden–Julian Oscillation: a review. *J. Clim.* **31**, 9425–9443 (2018).
79. Stan, C. et al. Review of tropical–extratropical teleconnections on intraseasonal time scales. *Rev. Geophys.* **55**, 902–937 (2017).
80. Kim, H., Richter, J. H. & Martin, Z. Insignificant QBO–MJO prediction skill relationship in the SubX and S2S subseasonal reforecasts. *J. Geophys. Res. Atmos.* <https://doi.org/10.1029/2019JD031416> (2019).
81. Lim, E.-P., Hendon, H. H. & Thompson, D. W. J. Seasonal evolution of stratosphere–troposphere coupling in the southern hemisphere and implications for the predictability of surface climate. *J. Geophys. Res. Atmos.* **123**, 1–15 (2018).
82. Zheng, C., Chang, E. K. M., Kim, H., Zhang, M. & Wang, W. Subseasonal to seasonal prediction of wintertime northern hemisphere extratropical cyclone activity by S2S and NMME models. *J. Geophys. Res. Atmos.* <https://doi.org/10.1029/2019JD031252> (2019).
83. DeFlorio, M. J. et al. Global evaluation of atmospheric river subseasonal prediction skill. *Clim. Dyn.* **52**, 3039–3060 (2019).

84. Baggett, C. et al. Skillful subseasonal forecasts of weekly tornado and hail activity using the Madden-Julian Oscillation. *J. Geophys. Res. Atmos.* **123**, 12,661–12,675 (2018).
85. Broennimann, S. Impact of El Niño–Southern Oscillation on European climate. *Rev. Geophys.* **45**, RG3003 (2007).
86. Ambaum, P. & Hoskins, B. J. The NAO troposphere–stratosphere connection. *J. Clim.* **15**, 1969–1978 (2002).
87. Kushnir, Y., Robinson, W. A., Chang, P. & Robertson, A. W. The physical basis for predicting Atlantic sector seasonal-to-interannual climate variability. *J. Clim.* **19**, 5949–5970 (2006).
88. Riddle, E. E., Butler, A. H., Furtado, J. C., Cohen, J. L. & Kumar, A. CFSv2 ensemble prediction of the wintertime Arctic oscillation. *Clim. Dyn.* **41**, 1099–1116 (2013).
89. Hendon, H. H., Thompson, D. W. J. & Wheeler, M. C. Australian rainfall and surface temperature variations associated with the southern hemisphere annular mode. *J. Clim.* **20**, 2452–2467 (2007).
90. Marshall, A. G. et al. Intra-seasonal drivers of extreme heat over Australia in observations and POAMA-2. *Clim. Dyn.* **43**, 1915–1937 (2014).
91. Seviour, W. J. M. et al. Skillful seasonal prediction of the southern annular mode and Antarctic ozone. *J. Clim.* **27**, 7462–7474 (2014).
92. Lim, E.-P., Hendon, H. H. & Rashid, H. A. Seasonal predictability of the southern annular mode due to its association with ENSO. *J. Clim.* **26**, 8037–8054 (2013).
93. Lim, E. et al. Australian hot and dry extremes induced by weakenings of the stratospheric polar vortex. *Nat. Geosci.* **12**, 896–901 (2019).
94. Mariotti, A. et al. Windows of opportunity for skillful forecasts subseasonal to seasonal and beyond. *Bull. Am. Meteorol. Soc.* <https://doi.org/10.1175/BAMS-D-18-0326.1> (2020).
95. Dirmeyer, P. A., Halder, S. & Bombardi, R. On the harvest of predictability from land states in a global forecast model. *J. Geophys. Res. Atmos.* **123**, 13111–13127 (2018).
96. Koster, R. D. et al. Regions of strong coupling between soil moisture and precipitation. *Science* **305**, 1138–1140 (2004).
97. Koster, R. D. et al. The second phase of the global land–atmosphere coupling experiment: soil moisture contributions to subseasonal forecast skill. *J. Hydrometeorol.* **12**, 805–822 (2011).
98. Seo, E. et al. Impact of soil moisture initialization on boreal summer subseasonal forecasts: mid-latitude surface air temperature and heat wave events. *Clim. Dyn.* <https://doi.org/10.1007/s00382-018-4221-4> (2018).
99. Zampieri, L., Goessling, H. F. & Jung, T. Bright prospects for Arctic sea ice prediction on subseasonal time scales. *Geophys. Res. Lett.* **45**, 9731–9738 (2018).
100. Zampieri, L., Goessling, H. F. & Jung, T. Predictability of Antarctic sea ice edge on subseasonal time scales. *Geophys. Res. Lett.* **46**, 9719–9727 (2019).
101. Bushuk, M. et al. A mechanism for the Arctic sea ice spring predictability barrier. *Geophys. Res. Lett.* <https://doi.org/10.1029/2020GL088335> (2020).
102. Kimmritz, M. et al. Impact of ocean and sea ice initialisation on seasonal prediction skill in the Arctic. *JAMES* <https://doi.org/10.1029/2019MS001825> (2019).
103. Ono, J., Komuro, Y. & Tatebe, H. Impact of sea-ice thickness initialized in April on Arctic sea-ice extent predictability with the MIROC climate model. *Ann. Glaciol.* **61**, 97–105 (2020).
104. Liu, J. et al. Towards reliable Arctic sea ice prediction using multivariate data assimilation. *Sci. Bull.* **64**, 63–72 (2019).
105. Jung, T. et al. Advancing polar prediction capabilities on daily to seasonal time scales. *Bull. Am. Meteorol. Soc.* **97**, 1631–1647 (2016).
106. Jung, T., Kasper, M. A., Semmler, T. & Serraz, S. Arctic influence on subseasonal midlatitude prediction. *Geophys. Res. Lett.* **41**, 3676–3680 (2014).
107. Baldwin, M. P. et al. Stratospheric memory and skill of extended-range weather forecasts. *Science* **301**, 636–640 (2003).
108. Butler, A. H., Polvani, L. M. & Deser, C. Separating the stratospheric and tropospheric pathways of El Niño–Southern Oscillation teleconnections. *Environ. Res. Lett.* <https://doi.org/10.1088/1748-9326/9/2/024014> (2014).
109. Sigmund, M. et al. Enhanced seasonal forecast skill following stratospheric sudden warmings. *Nat. Geosci.* **6**, 98–102 (2013).
110. Scaife, A. A. et al. Seasonal winter forecasts and the stratosphere. *Atmos. Sci. Lett.* <https://doi.org/10.1002/asl.598> (2016).
111. Anstey, J. A. & Shepherd, T. G. High-latitude influence of the quasi-biennial oscillation (Review article). *Quart. J. Roy. Meteorol. Soc.* **140**, 1–21 (2014).
112. Garinkel, C. I. & Hartmann, D. L. Influence of the quasi-biennial oscillation on the North Pacific and El Niño teleconnections. *J. Geophys. Res.* **115**, D20116 (2010).
113. Wang, J., Kim, H.-M. & Chang, E. K. M. Interannual modulation of northern hemisphere winter storm tracks by the QBO. *Geophys. Res. Lett.* **45**, 2786–2794 (2018).
114. Yoo, C. & Son, S.-W. Modulation of the boreal wintertime Madden-Julian Oscillation by the stratospheric quasi-biennial oscillation. *Geophys. Res. Lett.* **43**, 1392–1398 (2016).
115. Son, S.-W., Lim, Y., Yoo, C., Hendon, H. H. & Kim, J. Stratospheric control of Madden-Julian Oscillation. *J. Clim.* **30**, 1909–1922 (2017).
116. Lim, Y. et al. Influence of the QBO on MJO prediction skill in the subseasonal-to-seasonal prediction models. *Clim. Dyn.* <https://doi.org/10.1007/s00382-019-04719-y> (2019).
117. Tompkins, A. M. et al. The climate-system historical forecast project: providing open access to seasonal forecast ensembles from centers around the globe. *Bull. Am. Meteorol. Soc.* **98**(11), 2293–2301 (2017).
118. Acosta Navarro, J. C. et al. Link between autumnal Arctic sea ice and northern hemisphere winter forecast skill. *Geophys. Res. Lett.* **47**, e2019GL086753 (2020).
119. Scaife, A. A. et al. Skill of tropical rainfall predictions in multiple seasonal forecast systems. *Int. J. Climatol.* <https://doi.org/10.1002/joc.5855> (2018).
120. Hu, Z. et al. How much of monthly mean precipitation variability over global land is associated with SST anomalies? *Clim. Dyn.* **54**, 701–712 (2020).
121. Kirtman, B. P. et al. in *Climate Science for Serving Society: Research, Modelling and Prediction Priorities* (eds Asrar, G. R. & Hurrell, J. W.) 205–235 (Springer, 2013).
122. Capotondi, A., Wittenberg, A. T., Kug, J.-S., Takahashi, K. & McPhaden, M. J., in *El Niño Southern Oscillation in a Changing Climate* (eds McPhaden, M., Santoso, A. & Cai, W.) 65–86 (AGU, 2020).
123. Vimont, D. J., Alexander, M. A. & Newman, M. Optimal growth of central and east Pacific ENSO events. *Geophys. Res. Lett.* **41**, 4027–4034 (2014).
124. Zhang, H., Clement, A. & DiNezio, L. The south Pacific meridional mode: a mechanism for ENSO-like variability. *J. Clim.* **27**, 769–783 (2014).
125. Larson, S. & Kirtman, B. P. The Pacific meridional mode as a trigger for ENSO in a high-resolution coupled model. *Geophys. Res. Lett.* <https://doi.org/10.1002/grl.50571> (2013).
126. Capotondi, A. & Sardeshmukh, P. D. Optimal precursors of different types of ENSO events. *Geophys. Res. Lett.* **42**, 9952–9960 (2015).
127. Amaya, D. The Pacific meridional mode and ENSO: a review. *Curr. Clim. Change Rep.* <https://doi.org/10.1007/s40641-019-00142-x> (2019).
128. Larson, S. M. & Kirtman, B. P. Assessing Pacific Meridional Mode forecasts and its role as an ENSO precursor and predictor in the North American multi-model ensemble. *J. Clim.* **27**, 7018–7032 (2014).
129. Ren, H. F.-F., Jin, B. & Tian, A. A. Scaife distinct persistence barriers in two types of ENSO. *Geophys. Res. Lett.* **43**, 10,973–10,979 (2016).
130. Infanti, J. M. & Kirtman, B. P. North American rainfall and temperature prediction response to the diversity of ENSO. *Clim. Dyn.* <https://doi.org/10.1007/s00382-015-2749-0> (2016).
131. DiNezio, P. et al. A two-year forecast for a 60–80% chance of La Niña in 2017–2018. *Geophys. Res. Lett.* <https://doi.org/10.1002/2017GL074904> (2017).
132. Freund, M. B. et al. Higher frequency of central Pacific El Niño events in recent decades relative to past centuries. *Nat. Geosci.* **12**, 450–455 (2019).
133. McPhaden, M. J. Genesis and evolution of the 1997–98 El Niño. *Science* **283**, 950–954 (1999).
134. Capotondi, A., Sardeshmukh, P. D. & Ricciardulli, L. The nature of the stochastic wind forcing of ENSO. *J. Clim.* **31**, 8081–8099 (2018).
135. Tan, X. et al. A study of the effects of westerly wind bursts on ENSO based on CESM. *Clim. Dyn.* **54**, 885–899 (2020).
136. Lopez, H. & WWBs, B. P. K. ENSO predictability, the spring barrier and extreme events. *J. Geophys. Res. Atmos.* **119**, 10,114–10,138 (2014).
137. Ren, H. L. et al. Seasonal predictability of winter ENSO types in operational dynamical model predictions. *Clim. Dyn.* **52**, 3869–3890 (2019).
138. Chang, P. et al. Climate fluctuations of tropical coupled systems: the role of ocean dynamics. *J. Clim.* **19**, 5122–5174 (2006).
139. Lübbecke, J. F. & McPhaden, M. J. Symmetry of the Atlantic Niño mode. *Geophys. Res. Lett.* **44**, 965–973 (2017).
140. Richter, I. et al. On the link between mean state biases and prediction skill in the tropics: an atmospheric perspective. *Clim. Dyn.* **50**, 3355–3374 (2018).
141. Stockdale, T. N., Balmaseda, M. A. & Vidard, A. Tropical Atlantic SST prediction with coupled ocean–atmosphere GCMs. *J. Clim.* **19**, 6047–6061 (2006).
142. Ding, H. et al. The impact of mean state errors on equatorial Atlantic interannual variability in a climate model. *J. Geophys. Res. Oceans* **120**, 1133–1151 (2015).
143. Saji, N. H., Goswami, B. N., Vinayachandran, P. N. & Yamagata, T. A dipole mode in the tropical Indian Ocean. *Nature* **401**, 360–363 (1999).
144. Krishnamurthy, V. & Kirtman, B. P. Variability of the Indian Ocean: relation to monsoon and ENSO. *Q. J. R. Meteorol. Soc.* **129**, 1623–1646 (2003).
145. Wu, R., Kirtman, B. P. & Krishnamurthy, V. An asymmetric mode of tropical Indian Ocean rainfall variability in boreal spring. *J. Geophys. Res. Atmos.* <https://doi.org/10.1029/2007JD009316> (2008).
146. Lu, B. et al. An extreme negative Indian Ocean dipole event in 2016: dynamics and predictability. *Clim. Dyn.* <https://doi.org/10.1007/s00382-017-3908-2> (2017).
147. Shinoda, T. & Han, W. Influence of Indian Ocean dipole on atmospheric subseasonal variability. *J. Clim.* **18**, 3891–3909 (2005).
148. Dunstone, N. et al. Skillful predictions of the winter North Atlantic Oscillation one year ahead. *Nat. Geosci.* **9**, 809–814 (2016).
149. Dunstone, N. et al. Skillful seasonal predictions of summer European rainfall. *Geophys. Res. Lett.* **45**, 3246–3254 (2018).
150. Paolino, D. A., Kinter, J. L., Kirtman, B. P., Min, D. & Straus, D. M. The impact of land surface and atmospheric initialization on seasonal forecasts with CCSM. *J. Clim.* **25**, 1007–1021 (2011).
151. Dirmeyer, P. A. The role of the land surface background state in climate predictability. *J. Hydrometeorol.* **4**, 599–610 (2003).
152. Prodhomme, C., Doblas-Reyes, F., Bellprat, O. & Dutra, E. Impact of land-surface initialization on sub-seasonal to seasonal forecasts over Europe. *Clim. Dyn.* **47**, 919–935 (2016).
153. Ardilouze, C., Batté, L., Decharme, B. & Déqué, M. On the link between summer dry bias over the US Great Plains and seasonal temperature prediction skill in a dynamical forecast system. *Weather Forecast.* **34**, 1161–1172 (2019).
154. Marshall, A. G. & Scaife, A. A. Improved predictability of stratospheric sudden warming events in an atmospheric general circulation model with enhanced stratospheric resolution. *J. Geophys. Res.* **115**, D16114 (2010).
155. Boer, G. J. & Hamilton, K. QBO influence on extratropical predictive skill. *Clim. Dyn.* **31**, 987–1000 (2008).
156. Marshall, A. G. & Scaife, A. A. Impact of the QBO on surface winter climate. *J. Geophys. Res.* **114**, D18110 (2009).
157. Scaife, A. A. et al. Predictability of the Quasi-Biennial Oscillation and its northern winter teleconnection on seasonal to decadal timescales. *Geophys. Res. Lett.* **41**, 1752–1758 (2014).
158. Doblas-Reyes, F. J., Hagedorn, R., Palmer, T. N. & Morcrette, J.-J. Impact of increasing greenhouse gas concentrations in seasonal ensemble forecasts. *Geophys. Res. Lett.* **33**, L07708 (2006).
159. Solaraju-Murali, B., Caron, L.-P., González-Reviriego, N. & Doblas-Reyes, F. J. Multi-year prediction of European summer drought conditions for the agricultural sector. *Environ. Res. Lett.* <https://doi.org/10.1088/1748-9326/ab5043> (2019).
160. Cassou, C. et al. Decadal climate variability and predictability: challenges and opportunities. *Bull. Am. Meteorol. Soc.* **99**, 479–490 (2018).
161. Liu, Z. & Di Lorenzo, E. Mechanisms and predictability of Pacific decadal variability. *Curr. Clim. Change Rep.* **4**, 128–144 (2018).
162. Power, S., Casey, T., Folland, C., Colman, A. & Mehta, V. Inter-decadal modulation of the impact of ENSO on Australia. *Clim. Dyn.* **15**, 319–324 (1999).

163. Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M. & Francis, R. C. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bull. Am. Meteorol. Soc.* **78**, 1069–1079 (1997).
164. Newman, M. et al. The Pacific Decadal Oscillation, revisited. *J. Clim.* **29**, 4399–4427 (2016).
165. Chiang, J. C. H. & Vimont, D. J. Analogous Pacific and Atlantic Meridional Modes of tropical atmosphere–ocean variability. *J. Clim.* **17**, 4143–4158 (2004).
166. Di Lorenzo, E. et al. North Pacific Gyre Oscillation links ocean climate and ecosystem change. *GRL* **35**, L08607 (2008).
167. Han, W. et al. Indian Ocean decadal variability: a review. *Bull. Am. Meteor. Soc.* **95**, 1679–1703 (2014).
168. Han, W. et al. Intensification of decadal and multi-decadal sea level variability in the western tropical Pacific during recent decades. *Clim. Dyn.* **43**, 1357–1379 (2014).
169. Li, Y., Han, W., Wang, F., Zhang, L. & Duan, J. Vertical structure of the upper-Indian Ocean thermal variability. *J. Clim.* **33**, 7233–7253 (2020).
170. Tozuka, T., Luo, J., Masson, S. & Yamagata, T. Decadal modulations of the Indian Ocean dipole in the SINTEX-F1 coupled GCM. *J. Clim.* **20**, 2881–2894 (2007).
171. Feng, M. H. H. et al. Decadal increase in Ningaloo Niño since the late 1990s. *Geophys. Res. Lett.* **42**, 104–112 (2015).
172. Ummenhofer, C. C., Biastoch, A. & Böning, C. W. Multi-decadal Indian Ocean variability linked to the Pacific and implications for preconditioning Indian Ocean Dipole events. *J. Clim.* **30**, 1739–1751 (2017).
173. Lee, S.-K. et al. Pacific origin of the abrupt increase in Indian Ocean heat content during the warming hiatus. *Nat. Geosci.* **8**, 445–450 (2015).
174. Nieves, V., Willis, J. K. & Patzert, W. C. Recent hiatus caused by decadal shift in Indo-Pacific heating. *Science* **349**, 532–535 (2015).
175. Jin, X. et al. Distinct mechanisms of decadal subsurface heat content variations in the eastern and western Indian Ocean modulated by tropical Pacific SST. *J. Clim.* **31**, 7751–7769 (2018).
176. Annamalai, H., Potemra, J., Murtugudde, R. & McCreary, J. P. Effect of preconditioning on the extreme climate events in the tropical Indian Ocean. *J. Clim.* **18**, 3450–3469 (2005).
177. Henley, B. J. et al. Spatial and temporal agreement in climate model simulations of the Interdecadal Pacific Oscillation. *Environ. Res. Lett.* **12**, 044011 (2017).
178. Fasullo, J. T., Phillips, A. S. & Deser, C. Evaluation of leading modes of climate variability in the CMIP archives. *J. Clim.* <https://doi.org/10.1175/JCLI-D-19-1024.1> (2020).
179. Mann, M. E., Steinman, B. A. & Miller, S. K. Absence of internal multidecadal and interdecadal oscillations in climate model simulations. *Nat. Commun.* <https://doi.org/10.1038/s41467-019-13823-w> (2020).
180. Kosaka, Y. & Xie, S.-P. The tropical Pacific as a key pacemaker of the variable rates of global warming. *Nat. Geosci.* <https://doi.org/10.1038/NGEO2770> (2016).
181. Tung, K.-K. & Chen, X. Understanding the recent global surface warming slowdown: a review. *Climate* **6**, 82 (2018).
182. England, M. H. et al. Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus. *Nat. Clim. Change* **4**, 222–227 (2014).
183. Meehl, G. A., Teng, H. & Arblaster, J. M. Climate model simulations of the observed early-2000s hiatus of global warming. *Nat. Clim. Change* **4**, 898–902 (2014).
184. Fyfe, J. C. et al. Making sense of the early-2000s warming slowdown. *Nat. Clim. Change* **6**, 224–228 (2016).
185. Xie, S.-P. & Kosaka, Y. What caused the global surface warming hiatus of 1998–2013? *Curr. Clim. Change Rep.* **3**, 128–140 (2017).
186. Seager, R. et al. Strengthening tropical Pacific zonal sea surface temperature gradient consistent with rising greenhouse gases. *Nat. Clim. Change* **9**, 517–522 (2019).
187. Chen, X. & Tung, K.-K. Varying planetary heat sink led to global-warming slowdown and acceleration. *Science* **345**, 897–903 (2014).
188. Santer, B. D. et al. Observed multivariable signals of late 20th and early 21st century volcanic activity. *Geophys. Res. Lett.* **42**, 500–509 (2015).
189. Smith, D. M. et al. Role of volcanic and anthropogenic aerosols in the recent global surface warming slowdown. *Nat. Clim. Chang.* **6**, 936 (2016).
190. Oudar, T., Kushner, P. J., Fyfe, J. & Sigmond, M. No impact of anthropogenic aerosols on early 21st century global temperature trends in a large initial-condition ensemble. *Geophys. Res. Lett.* **45**, 9245–9252 (2018).
191. Guemas, V., Doblas-Reyes, F. J., Andreu-Burillo, I. & Asif, M. Retrospective prediction of the global warming slowdown in the past decade. *Nat. Clim. Change* **3**, 649–653 (2013).
192. Bordbar, M. H. et al. Uncertainty in near-term global surface warming linked to tropical Pacific climate variability. *Nat. Commun.* **10**, 1990 (2019).
193. Meehl, G. A., Chung, C. T. Y., Arblaster, J. M., Holland, M. M. & Bitz, C. M. Tropical decadal variability and the rate of Arctic sea ice retreat. *Geophys. Res. Lett.* <https://doi.org/10.1029/2018GL079989> (2018).
194. Meehl, G. A., Arblaster, J. M., Bitz, C., Chung, C. T. Y. & Teng, H. Antarctic sea ice expansion between 2000–2014 driven by tropical Pacific decadal climate variability. *Nat. Geosci.* <https://doi.org/10.1038/NGEO2751> (2016).
195. Purich, A. et al. Tropical Pacific SST drivers of recent Antarctic sea ice trends. *J. Clim.* **29**, 8931–8948 (2016).
196. Thoma, M., Greatbatch, R. J., Kadow, C. & Gerdes, R. Decadal hindcasts initialized using observed surface wind stress: evaluation and prediction out to 2024. *Geophys. Res. Lett.* **42**, 6454–6461 (2015).
197. Meehl, G. A. et al. Recent sudden Antarctic sea ice retreat caused by connections to the tropics and sustained ocean changes around Antarctica. *Nat. Commun.* **10**, 14 (2019).
198. Yin, J., Overpeck, J., Peyser, C. & Stouffer, R. Big jump of record warm global mean surface temperature in 2014–2016 related to unusually large oceanic heat releases. *Geophys. Res. Lett.* **45**, 1069–1078 (2018).
199. Booth, B. B. B., Dunstone, N. J., Halloran, P. R., Andrews, T. & Bellouin, N. Aerosols implicated as a prime driver of twentieth-century North Atlantic climate variability. *Nature* **484**, 228 (2012).
200. Watanabe, M. & Tabebe, H. Reconciling roles of sulphate aerosol forcing and internal variability in Atlantic multidecadal climate changes. *Clim. Dyn.* **53**, 4651–4665 (2019).
201. Hermanson, L. et al. Robust multiyear climate impacts of volcanic eruptions in decadal prediction systems. *J. Geophys. Res. Atmos.* <https://doi.org/10.1029/2019JD031739> (2020).
202. Menary, M. B. & Scaife, A. A. Naturally forced multidecadal variability of the Atlantic meridional overturning circulation. *Clim. Dyn.* **42**, 1347–1362 (2014).
203. Cai, W. et al. Pan-tropical climate interactions. *Science* **363**, eaav4236 (2019).
204. Mechoso, R. (ed.) *Interacting Climates of Ocean Basins: Observations, Mechanisms, Predictability, and Impacts* (Cambridge Univ. Press, 2020).
205. Meehl, G. A. et al. Atlantic and Pacific tropics connected by mutually interactive decadal-timescale processes. *Nat. Geosci.* <https://doi.org/10.1038/s41561-020-00669-x> (2020).
206. Chikamoto, Y. et al. Skillful multi-year predictions of tropical trans-basin climate variability. *Nat. Commun.* **6**, 6869 (2015).
207. Ruprich-Robert, Y. et al. Assessing the climate impacts of the observed Atlantic multidecadal variability using the GFDL CM2.1 and NCAR CESM1 global coupled models. *J. Clim.* **30**, 2785–2810 (2017).
208. Levine, A. F. Z., McPhaden, M. J. & Frierson, D. M. W. The impact of the AMV on multidecadal ENSO variability. *Geophys. Res. Lett.* **44**, 3877–3886 (2017).
209. Kumar, A., Bhaskar, J. & Wang, H. Attribution of SST variability in global oceans and the role of ENSO. *Clim. Dyn.* **43**, 209–220 (2014).
210. Taschetto, A. S., Rodrigues, R. R., Meehl, G. A., McGregor, S. & England, M. H. How sensitive are the Pacific-North Atlantic teleconnections to the position and intensity of El Niño-related warming. *Clim. Dyn.* <https://doi.org/10.1007/s00382-015-2679-x> (2015).
211. Han, W. et al. Decadal variability of Indian and Pacific Walker Cells: do they co-vary on decadal timescales? *J. Clim.* **30**, 8447–8468 (2017).
212. Han, W. et al. Multi-decadal trend and decadal variability of the regional sea level over the Indian Ocean since the 1960s: roles of climate modes and external forcing. *Climate* **6**, 51 (2018).
213. Deepa, J. S. et al. The tropical Indian Ocean decadal sea level response to the Pacific decadal oscillation forcing. *Clim. Dyn.* **52**, 5045 (2019).
214. Zhang, R. et al. A review of the role of the Atlantic meridional overturning circulation in Atlantic multidecadal variability and associated climate impacts. *Rev. Geophys.* **57**, 316–375 (2019).
215. Li, X., Xie, S.-P., Gille, S. T. & Yoo, C. Atlantic-induced pan-tropical climate change over the past three decades. *Nat. Clim. Change* <https://doi.org/10.1038/NCLIMATE2840> (2015).
216. Li, H., Ilyina, T., Müller, W. A. & Seitz, F. Decadal prediction of the North Atlantic CO₂ uptake. *Nat. Commun.* **7**, 11076 (2016).
217. Jin, D. & Kirtman, B. P. How the annual cycle affects the extratropical response to ENSO. *J. Geophys. Res.* **115**, D06102 (2010).
218. Zhang, L., Han, W. & Sieng, F. Unraveling causes for the changing behavior of tropical Indian Ocean in the past few decades. *J. Clim.* **31**, 2377–2388 (2018).
219. Thornton, H. et al. Skillful seasonal prediction of winter gas demand. *Env. Res. Lett.* **14**, 024009 (2019).
220. Palin, E. J. et al. Skillful seasonal forecasts of winter disruption to the U.K. transport system. *J. Appl. Meteor. Climatol.* **55**, 325–344 (2016).
221. Towler, E., Paimazumder, D. & Done, J. Toward application of decadal climate predictions. *J. Appl. Meteorol. Climatol.* **57**, 555–568 (2018).
222. Vecchi, G. A. et al. On the seasonal forecasting of regional tropical cyclone activity. *J. Clim.* **27**, 7994–8016 (2014).
223. Annan, J. D. et al. Parameter estimation in an atmospheric GCM using the ensemble Kalman filter. *Nonlinear Processes Geophys.* <https://doi.org/10.5194/npg-12-363-2005> (2005).
224. Düben, P. D., Hugh McNamee, H. & Palmer, T. N. The use of imprecise processing to improve accuracy in weather & climate prediction. *J. Comput. Phys.* **271**, 2–18 (2014).
225. Palmer, T. N., Peter Düben, P. & McNamee, H. Stochastic modelling and energy-efficient computing for weather and climate prediction. *Phil. Trans. Roy. Soc. A* <https://doi.org/10.1098/rsta.2014.0118> (2014).
226. Ham, Y.-G., Kim, J.-H. & Luo, J.-J. Deep learning for multi-year ENSO forecasts. *Nature* <https://doi.org/10.1038/s41586-019-1559-7> (2019).
227. Zhang, S. et al. Coupled data assimilation and parameter estimation in coupled ocean–atmosphere models: a review. *Clim. Dyn.* **54**, 5127–5144 (2020).
228. Karspeck, A. R. et al. A global coupled ensemble data assimilation system using the community earth system model and the data assimilation research testbed. *Q. J. R. Meteorol. Soc.* **144**, 2404–2430 (2018).
229. Mulholland, D., Lalouaux, P., Haines, K. & Balmaseda, M. Origin and impact of initialization shocks in coupled atmosphere–ocean forecasts. *Monthly Weather. Rev.* **143**, 4631–4644 (2015).
230. Saha, S. et al. The NCEP climate forecast system reanalysis. *Bull. Am. Meteorol. Soc.* **91**, 1015–1057 (2010).
231. Lalouaux, P. et al. A coupled data assimilation system for climate reanalysis. *Q. J. R. Meteorol. Soc.* **142**, 65–78 (2016).
232. Herman, R. J. et al. The effects of anthropogenic and volcanic aerosols and greenhouse gases on twentieth century Sahel precipitation. *Sci. Rep.* **10**, 12203 (2020).
233. Schurer, A., Hegerl, G., Mann, M. E. & Tett, S. F. B. Separating forcing from chaotic climate variability over the past millennium. *J. Clim.* **26**, 6954–6973 (2013).
234. Ault, T. R. et al. The continuum of hydroclimate variability in western North America during the last millennium. *J. Clim.* **26**, 5863–5878 (2013).
235. Laepple, T. & Huybers, P. Global and regional variability in marine surface temperatures. *Geophys. Res. Lett.* **41**, 2528–2534 (2014).
236. Loope, G., Thompson, D. M., Cole, J. E. & Overpeck, J. Is there a low-frequency bias in multiproxy reconstructions of Pacific SST variability? *Quat. Sci. Rev.* **246**, 106530 (2020).
237. Frankignoul, C., Muller, P. & Zorita, E. A simple model of the decadal response of the ocean to stochastic wind forcing. *J. Phys. Oceanogr.* **27**, 1533–1546 (1997).
238. Capotondi, A., Alexander, M. A. & Deser, C. Why are there Rossby wave maxima in the Pacific at 10S and 13N? *J. Phys. Oceanogr.* **33**, 1549–1563 (2003).
239. Chikamoto, Y., Timmermann, A., Widlansky, M. J., M. A., & L. Multi-year predictability of climate, drought, and wildfire in southwestern North America. *Sci. Rep.* <https://www.ncbi.nlm.nih.gov/pubmed/28747719> (2017).
240. Lovenduski, N. S., Bonan, G. B., Yeager, S. G., Lindsay, K. & Lombardozzi, D. L. High predictability of terrestrial carbon fluxes from an initialized decadal prediction system. *Environ. Res. Lett.* **14**, 124074 (2019).
241. Sospedra-Alfonso, R., Merryfield, W. J. & Kharin, V. V. Representation of snow in the Canadian seasonal to

- interannual prediction system: part II. Potential predictability and hindcast skill. *J. Hydrometeorol.* **17**, 2511–2535 (2016).
242. Kapnick, S. B. et al. Potential for western US seasonal snowpack prediction. *Proc. Natl Acad. Sci. USA* **115**, 1180–1185 (2018).
 243. Holbrook, N. J. et al. Keeping pace with marine heatwaves. *Nat. Rev. Earth Environ.* **1**, 482–493 (2020).
 244. Batté, L. et al. Summer predictions of Arctic sea ice edge in multi-model seasonal re-forecasts. *Clim. Dyn.* **54**, 5013–5029 (2020).
 245. Subramanian, A., Juricke, S., Dueben, P. & Palmer, T. A stochastic representation of subgrid uncertainty for dynamical core development. *Bull. Am. Meteorol. Soc.* **100**, 1091–1101 (2019).
 246. Penny, S. G. et al. Observational needs for improving ocean and coupled reanalysis, S2S prediction, and decadal prediction. *Front. Mar. Sci.* <https://doi.org/10.3389/fmars.2019.00391> (2019).
 247. Lofverstrom et al. An efficient ice-sheet/Earth System model spin-up procedure for CESM2.1 and CISM2.1: description, evaluation, and broader applicability. *JAMES* <https://doi.org/10.1029/2019MS001984> (2020).
 248. Gettelman, A. et al. The Whole Atmosphere Community Climate Model version 6 (WACCM6). *J. Geophys. Res. Atmos.* <https://doi.org/10.1029/2019JD030943> (2019).
 249. Tommasi, D. C. et al. Managing living marine resources in a dynamic environment: the role of seasonal to decadal climate forecasts. *Prog. Oceanogr.* **152**, 15–49 (2017).
 250. Stock, C. A. et al. Seasonal sea surface temperature anomaly prediction for coastal ecosystems. *Prog. Oceanogr.* **137**, 219–236 (2015).
 251. Liu, G. et al. Predicting heat stress to inform reef management: NOAA Coral Reef Watch's 4-month coral bleaching outlook. *Front. Mar. Sci.* <https://doi.org/10.3389/fmars.2018.00057> (2018).
 252. Capotondi, A., Sardeshmukh, P. D., Di Lorenzo, E., Subramanian, A. & Miller, A. J. Predictability of US West Coast ocean temperatures is not solely due to ENSO. *Sci. Rep.* **9**, 10993 (2019).
 253. Wells, M. L. et al. Harmful algal blooms and climate change: learning from the past and present to forecast the future. *Harmful Algae* **49**, 68–93 (2015).
 254. Séférian, R. et al. Multiyear predictability of tropical marine productivity. *Proc. Natl Acad. Sci. USA* **111**, 11646–11651 (2014).
 255. Park, J.-Y., Stock, C. A., Dunne, J. P., Yang, X. & Rosati, A. Seasonal to multiannual marine ecosystem prediction with a global Earth System model. *Science* **365**, 284–288 (2019).
 256. Krumhardt, K. M. et al. Potential predictability of net primary production in the ocean. *Glob. Biogeochem. Cycles* **34**, e2020GB006531 (2020).
 257. Siedlecki, S. A. et al. Experiments with seasonal forecasts of ocean conditions for the northern region of the California Current upwelling system. *Sci. Rep.* **6**, 1–18 (2016).
 258. Brady, R. X., Lovenduski, N. S., Yeager, S. G., Long, M. C. & Lindsay, K. Skillful multiyear predictions of ocean acidification in the California Current System. *Nat. Commun.* **11**, 2166 (2020).
 259. Séférian, R., Berthet, S. & Chevallier, M. Assessing the decadal predictability of land and ocean carbon uptake. *Geophys. Res. Lett.* **45**, 2455–2466 (2018).
 260. Lovenduski, N. S., Yeager, S. G., Lindsay, K. & Long, M. C. Predicting near-term variability in ocean carbon uptake. *Earth Syst. Dyn.* **10**, 45–57 (2019).
 261. Li, H., Ilyina, T., Müller, W. A. & Landschützer, P. Predicting the variable ocean carbon sink. *Sci. Adv.* <https://doi.org/10.1126/sciadv.aav6471> (2019).
 262. Bett, P. E. et al. Skillful seasonal prediction of key carbon cycle components: NPP and fire risk. *Environ. Res. Commun.* **2**, 055002 (2020).
 263. Park, J.-Y., Dunne, J. P. & Stock, C. A. Ocean chlorophyll as a precursor of ENSO: an earth system modeling study. *Geophys. Res. Lett.* <https://doi.org/10.1002/2017GL076077> (2018).
 264. Capotondi, A. et al. Observational needs supporting marine ecosystem modeling and forecasting: from the global ocean to regional and coastal systems. *Front. Mar. Sci.* **6**, 623 (2019).
 265. Fennel, K. et al. Advancing marine biogeochemical and ecosystem reanalyses and forecasts as tools for monitoring and managing ecosystem health. *Front. Mar. Sci.* **6**, 89 (2019).
 266. Weisheimer, A. & Palmer, T. N. On the reliability of seasonal climate forecasts. *J. R. Soc. Interface* <https://doi.org/10.1098/rsif.2013.1162> (2014).
 267. National Academies of Sciences, Engineering and Medicine. *Next Generation Earth System Prediction: Strategies for Subseasonal to Seasonal Forecasts* 1–351 (National Academies Press, 2017).
 268. National Research Council. *Assessment of Intraseasonal to Interannual Climate Prediction and Predictability* 1–193 (National Academies Press, 2010).
 269. Mehta, V. Natural Decadal Climate Variability: Phenomena, Mechanisms, and Predictability 1–374 (CRC Press, 2020).
 270. GISTEMP Team, 2020: GISS Surface Temperature Analysis (GISTEMP), version 4. *NASA Goddard Institute for Space Studies Dataset* accessed 2021-02-25 at <https://data.giss.nasa.gov/gistemp/> (2020).
 271. Lenssen, N. J. L. et al. Improvements in the GISTEMP Uncertainty Model. *J. Geophys. Res. Atmos.* **124**, 6307–6326 (2019).

Acknowledgements

The foundations of this Review emerged from a workshop held by the National Academies of Sciences, Engineering and Medicine in 2015 at Woods Hole, MA, USA, and the authors gratefully acknowledge support from A. Purcell and N. Huddleston. G.A.M., J.H.R. and N.R. were supported in part by the Regional and Global Model Analysis (RGMA) component of the Earth and Environmental System Modeling Program of the US Department of Energy's Office of Biological & Environmental Research (BER) via National Science Foundation IA 1844590, and by the National Center for Atmospheric Research (NCAR), which is a major facility

sponsored by the National Science Foundation (NSF) under Cooperative Agreement No. 1852977. M.H.E. acknowledges support from the Australian Research Council (Grant CE170100023). H.T. was partly supported by DOE/BER RGMA HiLAT-RASM. M.E.M. was supported by a grant from the NSF Paleoclimate Program #1748097. F.D.-R. and M.G.D. were supported by the H2020 EUCP project under Grant agreement no. 776613, and M.G.D. also by the Ramón y Cajal 2017 grant reference RYC-2017-22964. A.C. acknowledges support from the National Oceanic and Atmospheric Administration (NOAA) Climate Program Office's Modeling Analysis, Prediction and Projections (MAPP) Program and from the NOAA Climate Program Office's Climate Variability and Predictability (CVP) Program. A.C.S. acknowledges support from the NOAA Climate Variability and Predictability Program (Award NA18OAR4310405) and NOAA-MAPP (NA17OAR4310106). N.S.L. is grateful for support from the NSF (OCE-1752724). D.T. acknowledges support from the NCAR Advanced Study Program and NSF (OCE-1931242). S.C.S. was supported by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) Postdoctoral Fellowship. A.A.S. and D.S. were supported by the Met Office Hadley Centre Climate Programme funded by the Department for Business, Energy & Industrial Strategy (BEIS) and Department for Environment, Food and Rural Affairs (Defra) and by the European Commission Horizon 2020 EUCP project (GA 776613).

Author contributions

H.T. suggested the original concept. G.A.M. led the overall conceptual design and coordinated the writing. J.H.R. and H.T. made major contributions to the conceptual design and organization. J.H.R. generated Fig. 1a. H.T. generated Fig. 4. All authors discussed the concepts presented and contributed to the writing.

Competing interests

The authors declare no competing interests.

Peer review information

Nature Reviews Earth & Environment thanks Debra Hudson, Constantin Ardilouze, Terrence O'Kane and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1038/s43017-021-00155-x>.

RELATED LINKS

APCC climate predictions: <https://www.apcc21.org/>
Copernicus Climate Change Service: <https://www.copernicus.eu/en>
North American Multi-model Ensemble: <https://www.cpc.ncep.noaa.gov/products/NMME/>

© Springer Nature Limited 2021