

Received 29 May 2020

Accepted 16 June 2020

Edited by M. Zeller, Purdue University, USA

Keywords: lanthanide; caesium; cyclopentadienide; oligomer; crystal structure.

CCDC reference: 2010185

Supporting information: this article has supporting information at journals.iucr.org/e

Crystal structure of the $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}]_n$ oligomer



Daniel N. Huh, Joseph W. Ziller and William J. Evans*

Department of Chemistry, University of California, Irvine, California, 92697, USA. *Correspondence e-mail: wevans@uci.edu

The green compound poly[(tetrahydrofuran)tris[$\mu\text{-}\eta^5\text{:}\eta^5\text{-1-(trimethylsilyl)cyclopentadienyl}$]caesium(I)ytterbium(II)], $[\text{CsYb}(\text{C}_8\text{H}_{13}\text{Si})_3(\text{C}_4\text{H}_8\text{O})]_n$ or $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}^{\text{II}}]_n$ was synthesized by reduction of a red THF solution of $(\text{C}_5\text{H}_4\text{SiMe}_3)_3\text{Yb}^{\text{III}}$ with excess Cs metal and identified by X-ray diffraction. The compound crystallizes as a two-dimensional array of hexagons with alternating Cs^I and Yb^{II} ions at the vertices and cyclopentadienyl groups bridging each edge. This, based off the six-electron cyclopentadienyl rings occupying three coordination positions, gives a formally nine-coordinate tris(cyclopentadienyl) coordination environment to Yb and the Cs is ten-coordinate due to the three cyclopentadienyl rings and a coordinated molecule of THF. The complex comprises layers of Cs₃Yb₃ hexagons with THF ligands and Me₃Si groups in between the layers. The Yb–C metrical parameters are consistent with a $4f^{14}$ Yb^{II} electron configuration.

1. Chemical context

The new +2 oxidation states for the rare-earth metals Y, La, Ce, Pr, Gd, Tb, Ho, Er, and Lu were recently discovered by reduction of Cp_3^xLn ($\text{Cp}^x = \text{C}_5\text{H}_4\text{SiMe}_3$, $\text{C}_5\text{H}_3(\text{SiMe}_3)_2$; Ln = rare-earth metal) using alkali metal reductants Li, Na, K, and KC_8 (Fig. 1) (Hitchcock *et al.*, 2008; MacDonald *et al.*, 2013; Fieser *et al.*, 2015; Evans, 2016; Palumbo *et al.*, 2018). In each of these cases, 2.2.2-cryptand was added in these reactions to encapsulate the alkali metal. It was thought that chelating agents were necessary to sequester the alkali metal to prevent interactions with cyclopentadienide ligands and subsequent ligand dissociation leading to product decomposition. This idea was challenged by examining reduction reactions of Cp_3^xM ($\text{Cp}^x = \text{C}_5\text{H}_3(\text{SiMe}_3)_2$; M = La, Ce, U) with Li and Cs in the absence of chelating agents (Huh *et al.*, 2018). The reaction resulted in the isolation of the first chelate-free synthesis of La^{II}, Ce^{II}, and U^{II} complexes. The $[\text{Li}(\text{THF})_4]^{1+}$ cation of the




R = H, Ln = Y, La, Ce, Pr, Nd, Gd, Tb, Dy, Ho, Er, and Lu
 R = SiMe₃, Ln = La, Ce, Pr, and Nd
 M = Li, Na, K, KC₈

Figure 1

 Synthesis of $(\text{Cp}_3\text{Ln}^{\text{II}})^{1-}$ complexes by alkali metal reduction of $\text{Cp}_3\text{Ln}^{\text{III}}$ precursors; $\text{Cp}^x = \text{C}_5\text{H}_4\text{SiMe}_3$, $\text{C}_5\text{H}_3(\text{SiMe}_3)_2$.

Li salts in these chelate-free M^{II} complexes were well-separated from the $(\text{Cp}''M)^{1-}$ anion. However, the Cs reductions yielded polymeric complexes of general formula $[\text{Cp}''M(\mu\text{-Cp}'')_2\text{Cs}(\text{THF})_2]_n$ where the Cs cation has coordinated THF and cyclopentadienide ligands. Attempts to extend this chemistry to smaller rare-earth metals by reduction of $\text{Cp}'_3\text{Ln}$ ($\text{Cp}' = \text{C}_5\text{H}_4\text{SiMe}_3$; $\text{Ln} = \text{Y}, \text{Tb}, \text{Dy}$) showed evidence of Ln^{II} in solution; however, the reduction products were highly unstable and decomposed even at 238 K.



In this study, we were interested in examining the reduction of $\text{Cp}'_3\text{Yb}^{\text{III}}$ with Cs metal. Unlike Y^{II} , Tb^{II} , and Dy^{II} ions, Yb^{II} complexes are more easily obtainable, as reflected by their less negative reduction potentials (Morss, 1976). A crystal containing the oligomeric compound; $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}]_n$, **1** ($\text{Cp}' = \text{C}_5\text{H}_4\text{SiMe}_3$) was isolated by reduction of the $\text{Cp}'_3\text{Yb}^{\text{III}}$ complex (Fieser *et al.*, 2015) in THF using Cs metal (Figs. 2 and 3).

2. Structural commentary

All three Cp' rings remain coordinated to the Yb metal center after reduction and are coordinated in a trigonal-planar fashion. The Yb atom is within 0.107 Å of the plane of the three ring centroids. Each ring bridges Yb to Cs, which also is surrounded by three cyclopentadienyl ligands as well as a coordinated molecule of THF. The three ring centroids and the oxygen of THF are arranged in a pseudo-tetrahedral geometry around Cs with a calculated four-coordinate Cs τ_4 value of 0.76 ($\tau_4 = 1$ for tetrahedral; $\tau_4 = 0$ for square planar; Rosiak *et al.*, 2018). The Cs metal center has a pseudo-tetrahedral geometry with $\text{Cp}'(\text{centroid})\cdots\text{Cs}\cdots\text{Cp}'(\text{centroid})$ angles of 109.0, 114.3, and 121.4° and $\text{Cp}'(\text{centroid})\cdots\text{Cs}\cdots\text{O}(\text{THF})$ angles of 88.8, 94.1, and 127.8°.




Figure 2
Synthesis of $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}^{\text{II}}]_n$, **1**, by caesium metal reduction of the $\text{Cp}'_3\text{Yb}^{\text{III}}$ precursor.

Table 1
Selected bond distances and angles for $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}]_n$, **1**.

Centroid1, centroid2, and centroid3 are the centroids of the Cp rings connected to Si1, Si2, and Si3, respectively.

Yb1···centroid1	2.510 (1)
Yb1···centroid2	2.513 (2)
Yb1···centroid3	2.504 (1)
Cs1···centroid1	3.197 (1)
Cs1···centroid2	3.268 (2)
Cs1···centroid3	3.159 (1)
Cs1—O1	3.095 (3)
centroid1—Yb1···centroid2	120.1
centroid1—Yb1···centroid3	116.6
centroid2—Yb1···centroid3	122.8
centroid1—Cs1···centroid2	121.4
centroid1—Cs1···centroid3	109.0
centroid2—Cs1···centroid3	114.3
Yb1···centroid1···Cs1	175.3
Yb1···centroid2···Cs1	172.3
Yb1···centroid3···Cs1	176.7
centroid1···Cs1···O1	88.8
centroid2···Cs1···O1	94.1
centroid3···Cs1···O1	127.8

Table 2

Bond distance (Å) ranges for $\text{Yb}\cdots\text{Cp}'(\text{centroid})$ and bond angle (°) ranges for $\text{Cp}'(\text{centroid})\cdots\text{Yb}\cdots\text{Cp}'(\text{centroid})$ in $\text{Cp}'_3\text{Yb}$ (Fieser *et al.*, 2015), $[\text{K}(\text{crypt})][\text{Cp}'_3\text{Yb}]$ (Fieser *et al.*, 2015), and $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}]_n$.

	$\text{Cp}'_3\text{Yb}$	$[\text{K}(\text{crypt})][\text{Cp}'_3\text{Yb}]$	1
$\text{Yb}\cdots\text{Cp}'(\text{centroid})$	2.363–2.368	2.503–2.513	2.504 (1)–2.513 (2)
$\text{Cs}\cdots\text{Cp}'(\text{centroid})$			3.159 (1)–3.268 (2)
$\text{Cp}'\cdots\text{Yb}\cdots\text{Cp}'$	118.85–120.55	118.10–122.93	116.64–122.76
$\text{Cp}'\cdots\text{Cs}\cdots\text{Cp}'$			109.0–121.4

The bond distances and angles in **1** are summarized in Table 1. The range of 2.504 (1)–2.513 (2) Å $\text{Cp}'(\text{centroid})\cdots\text{Yb}$ bond distances in **1** is the same as that in the complex $[\text{K}(\text{crypt})][\text{Cp}'_3\text{Yb}^{\text{II}}]$ (crypt = 2.2.2-cryptand), which was




Figure 3
ORTEP representation of an asymmetric unit of $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}]_n$, **1**, with probability ellipsoids drawn at the 50% probability level. Hydrogen atoms were omitted for clarity.




Figure 4
CHEMDRAW (Mills, 2006) representation of [K(2.2.2-cryptand)][Cp₃Yb^{II}] (left) and [(THF)Cs(μ-η⁵:η⁵-Cp')₃Yb^{II}]_n, **1**, (right).

fully characterized as a 4f¹⁴ Yb^{II} complex, Table 2 and Fig. 4. In Cp₃Ln reduction chemistry, the difference in Ln···Cp(centroid) distances between the Ln^{III} and Ln^{II} complexes provides important information on the electronic configuration of the lanthanide ion (Evans, 2016). Differences in Ln···Cp(centroid) distances for reduction of 4fⁿ Ln^{III} ions to 4fⁿ⁺¹ Ln^{II} ions range from 0.1 to 0.2 Å (Fieser *et al.*, 2015). In this study, the difference of 0.14 Å in the Ln···Cp(centroid) distance is characteristic of a 4f¹³ Yb^{III} reduction to a 4f¹⁴ Yb^{II} ion. In contrast, Ln^{II} ions with 4fⁿ5d¹ configurations where the additional electron populates a d-orbital instead of the an f-orbital have differences of only 0.02–0.05 Å (Evans, 2016).

3. Supramolecular features

In **1**, all of the cyclopentadienyl ligands are bridging. The threefold symmetry of three bridging Cp' ligands on each metal generates a hexagonal pattern as shown in Fig. 5. The




Figure 5
Top view of the extended structure of [(THF)Cs(μ-η⁵:η⁵-Cp')₃Yb^{II}]_n, **1**, with the SiMe₃ substituent of the C₅H₄SiMe₃ group and the THF attached to Cs removed for clarity.

Yb···Cp'(centroid)···Cs angles are 172.5–176.7° such that each side of the hexagon is nearly linear. The 112.4–117.3° Yb···Cs···Yb angles are smaller than the 120.8–125.6° Cs···Yb···Cs angles, which makes the hexagon slightly irregular. This could be of interest to quantum scientists trying to make thin-film layers of magnetic materials since the hexagonal pattern could lead to spin frustration with a paramagnetic lanthanide.

The side view of these layers in Fig. 6 shows how the space in between them is filled with THF and Me₃Si substituent groups. The 116.6–122.8° Cp'(centroid)···Yb···Cp'(centroid) and 109.0–121.4° Cp'(centroid)···Cs···Cp'(centroid) angles generate the undulation of the hexagons shown in Fig. 6.

4. Database survey

The 3.159 (1), 3.197 (1), and 3.268 (2) Å Cs···Cp'(centroid) distances in **1** are shorter than the 3.278 and 3.435 Å Cs···Cp''(centroid) distances in [(THF)₂Cs][(μ-η⁵:η⁵-Cp'')₂U^{II}(η⁵-Cp'')]_n, (Huh *et al.*, 2018), the 3.396 Å Cs···C₅H₅(centroid) distances in {[(Me₃Si)₂NCs]₂[(C₅H₅)₂Fe] 0.5·(C₆H₅Me)}_n, (Morris *et al.*, 2007) and the 3.337 Å Cs···C₅Me₅(centroid) distances in [(THF)₂Cs(μ₃-O)₃{[Ti(C₅Me₅)₃(μ₃-CCH₂)]}] (González-del Moral *et al.*, 2005). The 3.095 (3) Å Cs—O(THF) bond distance is consistent with the Cs—O(THF) distances of 3.081 (7) to 3.119 (8) Å in [(THF)₂Cs][(μ-η⁵:η⁵-Cp'')₂U^{II}(η⁵-Cp'')]_n (Huh *et al.*, 2018) and 3.034 (9)–3.06 (1) Å in [(THF)₂Cs(μ₃-O)₃{[Ti(C₅Me₅)₃(μ₃-CCH₂)]}] (González-del Moral *et al.*, 2005).

The extended structure of **1** differs from that of the [(THF)₂Cs][(μ-η⁵:η⁵-Cp'')₂M^{II}(η⁵-Cp'')]_n, complexes (M = La, U), which comprise zigzag chains of –M–(μ-Cp'')–Cs–(μ-Cp'')– repeat units with a terminal Cp'' attached to M and two terminal THF ligands attached to Cs (Huh *et al.*, 2018). These were obtained by reduction of Cp''₃M^{III} compounds with Cs in THF. In those structures, La and U have a trigonal-planar tris(cyclopentadienyl) coordination like Yb in **1**, but the Cs is




Figure 6
Side view of the extended structure of [(THF)Cs(μ-η⁵:η⁵-Cp')₃Yb^{II}]_n, **1**. Magenta, Yb; brown, Cs; green, Si; red, O.

coordinated by only two cyclopentadienyl ligands to give a bent metallocene $\text{Cp}'_2\text{Cs}(\text{THF})_2$ sub-structure with these larger rings.

A survey of the Cambridge Structural Database (CSD, version 5.41, March 2020; Groom *et al.*, 2016) also revealed four oligomeric complexes containing Yb–Cp^x moieties with various types of cyclopentadienyl rings (Cp^x): $[\text{Na}(\mu\text{-}\eta^5\text{:}\eta^5\text{-C}_5\text{H}_5)_3\text{Yb}^{\text{II}}]_n$ (Apostolidis *et al.*, 1997), $[\text{Na}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}'')_2\text{Yb}^{\text{II}}]_n$ (Voskoboynikov *et al.*, 1997), $[(\text{C}_5\text{Me}_5)\text{Yb}(\mu\text{-I})(\mu\text{-}\eta^5\text{:}\eta^5\text{-C}_5\text{Me}_5)\text{Yb}(\text{C}_5\text{Me}_5)]_n$ (Evans *et al.*, 2006) and $[\text{Yb}(\mu\text{-}\eta^5\text{:}\eta^5\text{-C}_5\text{H}_5)(\text{Ph}_2\text{Pz})(\text{THF})]_n$ (Ph₂Pz = 3,5-diphenylpyrazolate) (Ali *et al.*, 2018). The $[\text{Na}(\mu\text{-}\eta^5\text{:}\eta^5\text{-C}_5\text{H}_5)_3\text{Yb}^{\text{II}}]_n$ (Apostolidis *et al.*, 1997) complex adopts a hexagonal net extended structure similar to that in **1** except the alkali metal does not have a coordinated solvent. The structure of $[\text{Na}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}'^{\text{Bu}})_3\text{Sm}^{\text{II}}]$ is similar (Bel'sky *et al.*, 1990). Three oligomeric complexes containing Cs–cyclopentadienyl moieties have previously been reported: $[(\text{THF})_2\text{Cs}][(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}'')_2\text{U}^{\text{IV}}(\eta^5\text{-Cp}'')]_n$ (Huh *et al.*, 2018), $\{[(\text{Me}_3\text{Si})_2\text{NCs}]_2[(\text{C}_5\text{H}_5)_2\text{Fe}]\cdot 0.5(\text{C}_6\text{H}_5\text{Me})\}_n$ (Morris *et al.*, 2007) and $[(\text{THF})_2\text{Cs}(\mu_3\text{-O})_3\{\text{Ti}(\text{C}_5\text{Me}_5)_3(\mu_3\text{-CCH}_2)\}]_n$ (González-del Moral *et al.*, 2005). An oligomeric, base-free Li–Cp' compound was also previously reported in the literature, $[(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')\text{Li}]_n$ (Evans *et al.*, 1992).

5. Synthesis and crystallization

In an argon-filled glovebox, addition of a red solution of Cp'₃Yb (50 mg, 0.085 mmol) in THF (2 mL) to excess Cs as a smear produced a green solution. This was stirred for 15 min at room temperature and then layered at the bottom of a vial below an Et₂O (10 mL) layer for crystallization at –35°C. After 1 d, X-ray quality dark-green crystals of $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}^{\text{II}}]_n$ were isolated. A small number of crystals were obtained and used for crystallographic analysis. Too little sample was available for other characterization.

6. Refinement

Crystal data and structure refinement for $[(\text{THF})\text{Cs}(\mu\text{-}\eta^5\text{:}\eta^5\text{-Cp}')_3\text{Yb}^{\text{II}}]_n$, **1** are summarized in Table 3. Hydrogen atoms were included using a riding model with $U_{\text{iso}}(\text{H})$ values of $1.2U_{\text{eq}}(\text{C})$ for CH₂ and aromatic hydrogens and $1.5U_{\text{eq}}(\text{C})$ for CH₃ hydrogens with C–H distances of 0.99 (CH₂), 0.95 (aromatic), and 0.98 Å (CH₃).

Acknowledgements

WJE would like to acknowledge the National Science Foundation for continued support.

Funding information

Funding for this research was provided by: National Science Foundation, Directorate for Mathematical and Physical Sciences (grant No. CHE-1855328 to William J. Evans).

Table 3
Experimental details.

Crystal data	
Chemical formula	$[\text{CsYb}(\text{C}_8\text{H}_{13}\text{Si})_3(\text{C}_4\text{H}_8\text{O})]$
M_r	789.87
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	88
a, b, c (Å)	9.4401 (4), 16.8718 (8), 21.0246 (10)
β (°)	92.0668 (6)
V (Å ³)	3346.4 (3)
Z	4
Radiation type	Mo $K\alpha$
μ (mm ^{–1})	3.99
Crystal size (mm)	0.15 × 0.09 × 0.08
Data collection	
Diffractometer	Bruker SMART APEXII CCD
Absorption correction	Multi-scan (SADABS; Bruker, 2014)
$T_{\text{min}}, T_{\text{max}}$	0.374, 0.432
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	40586, 8223, 6580
R_{int}	0.055
$(\sin \theta/\lambda)_{\text{max}}$ (Å ^{–1})	0.667
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.032, 0.056, 1.02
No. of reflections	8223
No. of parameters	316
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ (e Å ^{–3})	1.14, –0.62

Computer programs: APEX2 (Bruker, 2014), SAINT (Bruker, 2013), SHELXT2014/4 (Sheldrick, 2015a), SHELXL2014/7 (Sheldrick, 2015b) and SHELXTL (Sheldrick, 2008).

References

- Ali, S. H., Deacon, G. B., Junk, P. C., Hamidi, S., Wiecko, M. & Wang, J. (2018). *Chem. Eur. J.* **24**, 230–242.
- Apostolidis, C. B., Deacon, G., Dornberger, E. T., Edelmann, F., Kanellakopoulos, B., MacKinnon, P. & Stalke, D. (1997). *Chem. Commun.* pp. 1047–1048.
- Bel'sky, V. K., Gunko, Y. K., Bulychev, B. M., Sizov, A. I. & Soloveichik, G. L. (1990). *J. Organomet. Chem.* **390**, 35–44.
- Bruker (2013). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2014). APEX2 and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Evans, W. J. (2016). *Organometallics*, **35**, 3088–3100.
- Evans, W. J., Boyle, T. J. & Ziller, J. W. (1992). *Organometallics*, **11**, 3903–3907.
- Evans, W. J., Champagne, T. M., Davis, B. L., Allen, N. T., Nyce, G. W., Johnston, M. A., Lin, Y.-C., Khvostov, A. & Ziller, J. W. (2006). *J. Coord. Chem.* **59**, 1069–1087.
- Fieser, M. E., MacDonald, M. R., Krull, B. T., Bates, J. E., Ziller, J. W., Furche, F. & Evans, W. J. (2015). *J. Am. Chem. Soc.* **137**, 369–382.
- González-del Moral, O., Martín, A., Mena, M., Morales-Varela, M. del C. & Santamaría, C. (2005). *Chem. Commun.* pp. 3682–3684.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Hitchcock, P. B., Lappert, M. F., Maron, L. & Protchenko, A. V. (2008). *Angew. Chem. Int. Ed.* **47**, 1488–1491.
- Huh, D. N., Ziller, J. W. & Evans, W. J. (2018). *Inorg. Chem.* **57**, 11809–11814.
- MacDonald, M. R., Bates, J. E., Ziller, J. W., Furche, F. & Evans, W. J. (2013). *J. Am. Chem. Soc.* **135**, 9857–9868.
- Mills, N. (2006). *J. Am. Chem. Soc.* **128**, 13649–13650.

- Morris, J. J., Noll, B. C., Honeyman, G. W., O'Hara, C. T., Kennedy, A. R., Mulvey, R. E. & Henderson, K. W. (2007). *Chem. Eur. J.* **13**, 4418–4432.
- Morss, L. R. (1976). *Chem. Rev.* **76**, 827–841.
- Palumbo, C. T., Darago, L. E., Windorff, C. J., Ziller, J. W. & Evans, W. J. (2018). *Organometallics*, **37**, 900–905.
- Rosiak, D., Okuniewski, A. & Chojnacki, J. (2018). *Polyhedron*, **146**, 35–41.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015a). *Acta Cryst.* **A71**, 3–8.
- Sheldrick, G. M. (2015b). *Acta Cryst.* **C71**, 3–8.
- Voskoboinikov, A. Z., Argarkov, A. Y., Shestakova, A. K., Beletskaya, I. P., Kuz'mina, L. O. & Howard, J. A. K. (1997). *J. Organomet. Chem.* **544**, 65–68.

supporting information

Acta Cryst. (2020). E76, 1131-1135 [https://doi.org/10.1107/S2056989020008051]

Crystal structure of the [(THF)Cs(μ - η^5 : η^5 -Cp')₃Yb]_n oligomer

Daniel N. Huh, Joseph W. Ziller and William J. Evans

Computing details

Data collection: *APEX2* (Bruker, 2014); cell refinement: *SAINT* (Bruker, 2013); data reduction: *SAINT* (Bruker, 2013); program(s) used to solve structure: *SHELXT2014/4* (Sheldrick, 2015a); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015b); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Poly[(tetrahydrofuran)tris[μ - η^5 : η^5 -1-(trimethylsilyl)cyclopentadienyl]caesium(I)ytterbium(II)]

Crystal data

[CsYb(C₈H₁₃Si)₃(C₄H₈O)]

$M_r = 789.87$

Monoclinic, $P2_1/n$

$a = 9.4401$ (4) Å

$b = 16.8718$ (8) Å

$c = 21.0246$ (10) Å

$\beta = 92.0668$ (6)°

$V = 3346.4$ (3) Å³

$Z = 4$

$F(000) = 1560$

$D_x = 1.568$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9869 reflections

$\theta = 2.3$ – 28.5 °

$\mu = 3.99$ mm⁻¹

$T = 88$ K

Prism, green

$0.15 \times 0.09 \times 0.08$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2014)

$T_{\min} = 0.374$, $T_{\max} = 0.432$

40586 measured reflections

8223 independent reflections

6580 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.055$

$\theta_{\max} = 28.3$ °, $\theta_{\min} = 1.6$ °

$h = -12 \rightarrow 12$

$k = -22 \rightarrow 22$

$l = -27 \rightarrow 27$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.056$

$S = 1.02$

8223 reflections

316 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0229P)^2 + 0.1327P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 1.14$ e Å⁻³

$\Delta\rho_{\min} = -0.62$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. A green crystal of approximate dimensions 0.079 x 0.086 x 0.148 mm was mounted in a cryoloop and transferred to a Bruker SMART APEX II diffractometer. The APEX2 program package was used to determine the unit-cell parameters and for data collection (90 sec/frame scan time for a sphere of diffraction data). The raw frame data was processed using SAINT and SADABS to yield the reflection data file. Subsequent calculations were carried out using the SHELXTL program. The diffraction symmetry was 2/m and the systematic absences were consistent with the monoclinic space group P21/n that was later determined to be correct.

The structure was solved by dual space methods and refined on F2 by full-matrix least-squares techniques. The analytical scattering factors for neutral atoms were used throughout the analysis. Hydrogen atoms were included using a riding model. The structure is polymeric.

Least-squares analysis yielded $wR2 = 0.0562$ and $Goof = 1.017$ for 316 variables refined against 8223 data (0.75 Å), $R1 = 0.0315$ for those 6580 data with $I > 2.0\sigma(I)$.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}
Yb1	0.48657 (2)	0.44122 (2)	0.25290 (2)	0.01341 (4)
Cs1	0.99437 (2)	0.27157 (2)	0.31696 (2)	0.01493 (5)
Si1	0.69834 (10)	0.29048 (6)	0.15029 (5)	0.0184 (2)
Si2	0.23952 (11)	0.57037 (6)	0.11663 (5)	0.0204 (2)
Si3	0.25791 (10)	0.45182 (6)	0.40988 (5)	0.0187 (2)
O1	0.8795 (3)	0.24593 (16)	0.45145 (12)	0.0285 (6)
C1	0.7108 (3)	0.33895 (19)	0.22942 (16)	0.0161 (7)
C2	0.6551 (3)	0.3115 (2)	0.28743 (17)	0.0174 (8)
H2A	0.6070	0.2626	0.2929	0.021*
C3	0.6826 (3)	0.3680 (2)	0.33517 (17)	0.0196 (8)
H3A	0.6554	0.3643	0.3781	0.024*
C4	0.7575 (4)	0.4310 (2)	0.30831 (17)	0.0194 (8)
H4A	0.7902	0.4774	0.3298	0.023*
C5	0.7751 (3)	0.4129 (2)	0.24430 (17)	0.0184 (8)
H5A	0.8231	0.4453	0.2150	0.022*
C6	0.6270 (4)	0.1878 (2)	0.15741 (18)	0.0257 (9)
H6A	0.6848	0.1582	0.1890	0.039*
H6B	0.6299	0.1611	0.1161	0.039*
H6C	0.5288	0.1902	0.1709	0.039*
C7	0.5764 (4)	0.3487 (2)	0.09635 (17)	0.0276 (9)
H7A	0.6159	0.4017	0.0900	0.041*
H7B	0.4836	0.3533	0.1155	0.041*
H7C	0.5656	0.3216	0.0552	0.041*
C8	0.8767 (4)	0.2878 (2)	0.11484 (18)	0.0263 (9)
H8A	0.9435	0.2601	0.1438	0.039*
H8B	0.9099	0.3421	0.1081	0.039*
H8C	0.8702	0.2599	0.0740	0.039*
C9	0.3846 (3)	0.57009 (19)	0.17806 (16)	0.0162 (7)
C10	0.5320 (4)	0.56244 (19)	0.16482 (17)	0.0173 (7)

H10A	0.5679	0.5462	0.1253	0.021*
C11	0.6148 (4)	0.58254 (19)	0.21886 (17)	0.0171 (8)
H11A	0.7154	0.5833	0.2220	0.021*
C12	0.5224 (4)	0.60152 (19)	0.26788 (17)	0.0190 (8)
H12A	0.5497	0.6163	0.3102	0.023*
C13	0.3823 (4)	0.59465 (19)	0.24280 (16)	0.0180 (8)
H13A	0.2992	0.6048	0.2656	0.022*
C14	0.2553 (5)	0.4858 (2)	0.0598 (2)	0.0380 (11)
H14A	0.2394	0.4358	0.0822	0.057*
H14B	0.3503	0.4856	0.0425	0.057*
H14C	0.1843	0.4916	0.0249	0.057*
C15	0.0612 (4)	0.5700 (3)	0.1518 (2)	0.0379 (11)
H15A	0.0433	0.5178	0.1705	0.057*
H15B	-0.0113	0.5811	0.1185	0.057*
H15C	0.0577	0.6107	0.1849	0.057*
C16	0.2543 (4)	0.6620 (2)	0.06760 (17)	0.0230 (8)
H16A	0.3509	0.6665	0.0524	0.034*
H16B	0.2329	0.7085	0.0936	0.034*
H16C	0.1869	0.6593	0.0311	0.034*
C17	0.2651 (3)	0.4025 (2)	0.33118 (16)	0.0151 (7)
C18	0.1959 (3)	0.42821 (19)	0.27347 (17)	0.0169 (7)
H18A	0.1408	0.4751	0.2686	0.020*
C19	0.2216 (3)	0.3741 (2)	0.22535 (17)	0.0187 (8)
H19A	0.1879	0.3780	0.1823	0.022*
C20	0.3061 (4)	0.3127 (2)	0.25131 (17)	0.0206 (8)
H20A	0.3395	0.2678	0.2291	0.025*
C21	0.3320 (3)	0.3300 (2)	0.31617 (17)	0.0173 (8)
H21A	0.3858	0.2981	0.3453	0.021*
C22	0.2748 (5)	0.3775 (2)	0.47533 (18)	0.0346 (10)
H22A	0.1956	0.3400	0.4718	0.052*
H22B	0.2732	0.4047	0.5165	0.052*
H22C	0.3644	0.3487	0.4721	0.052*
C23	0.4026 (4)	0.5264 (2)	0.42253 (18)	0.0280 (9)
H23A	0.3884	0.5702	0.3923	0.042*
H23B	0.4944	0.5012	0.4158	0.042*
H23C	0.4008	0.5469	0.4661	0.042*
C24	0.0843 (4)	0.5039 (3)	0.4147 (2)	0.0399 (11)
H24A	0.0071	0.4652	0.4106	0.060*
H24B	0.0748	0.5429	0.3803	0.060*
H24C	0.0798	0.5309	0.4559	0.060*
C25	0.8427 (4)	0.3003 (2)	0.50059 (19)	0.0306 (9)
H25A	0.8093	0.3512	0.4820	0.037*
H25B	0.9255	0.3108	0.5296	0.037*
C26	0.7248 (4)	0.2603 (2)	0.53629 (19)	0.0338 (10)
H26A	0.7634	0.2246	0.5700	0.041*
H26B	0.6615	0.2998	0.5555	0.041*
C27	0.6489 (4)	0.2145 (2)	0.4835 (2)	0.0315 (10)
H27A	0.5959	0.1690	0.5005	0.038*

H27B	0.5827	0.2489	0.4585	0.038*
C28	0.7700 (4)	0.1873 (2)	0.44422 (18)	0.0278 (9)
H28A	0.8052	0.1351	0.4593	0.033*
H28B	0.7388	0.1823	0.3989	0.033*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Yb1	0.01020 (7)	0.01233 (7)	0.01766 (8)	-0.00136 (6)	-0.00014 (6)	0.00251 (6)
Cs1	0.01034 (10)	0.01249 (10)	0.02208 (12)	-0.00018 (8)	0.00232 (8)	0.00028 (9)
Si1	0.0150 (5)	0.0213 (5)	0.0190 (5)	0.0031 (4)	0.0009 (4)	-0.0011 (4)
Si2	0.0214 (5)	0.0157 (5)	0.0236 (6)	-0.0009 (4)	-0.0052 (4)	0.0028 (4)
Si3	0.0177 (5)	0.0192 (5)	0.0191 (5)	-0.0009 (4)	0.0012 (4)	-0.0017 (4)
O1	0.0207 (14)	0.0355 (16)	0.0295 (16)	-0.0014 (12)	0.0062 (12)	-0.0009 (13)
C1	0.0112 (16)	0.0161 (17)	0.0209 (19)	0.0018 (14)	0.0018 (14)	0.0027 (15)
C2	0.0130 (17)	0.0140 (17)	0.025 (2)	0.0030 (14)	0.0021 (15)	0.0029 (15)
C3	0.0184 (18)	0.024 (2)	0.0162 (19)	0.0082 (15)	0.0003 (15)	0.0016 (15)
C4	0.0151 (17)	0.0177 (19)	0.025 (2)	0.0022 (14)	-0.0071 (15)	-0.0016 (16)
C5	0.0083 (16)	0.0175 (18)	0.030 (2)	-0.0002 (13)	0.0033 (15)	0.0049 (16)
C6	0.027 (2)	0.023 (2)	0.027 (2)	0.0011 (16)	-0.0006 (17)	-0.0044 (17)
C7	0.029 (2)	0.033 (2)	0.021 (2)	0.0055 (18)	-0.0059 (17)	0.0013 (17)
C8	0.023 (2)	0.031 (2)	0.024 (2)	0.0038 (17)	0.0045 (17)	-0.0018 (17)
C9	0.0162 (17)	0.0111 (17)	0.0212 (19)	0.0006 (13)	0.0003 (14)	0.0012 (14)
C10	0.0208 (18)	0.0103 (16)	0.0210 (19)	0.0006 (14)	0.0037 (15)	0.0040 (15)
C11	0.0133 (17)	0.0139 (17)	0.024 (2)	-0.0047 (13)	-0.0014 (15)	0.0047 (15)
C12	0.0241 (19)	0.0118 (17)	0.021 (2)	-0.0067 (15)	-0.0054 (15)	0.0020 (15)
C13	0.0208 (18)	0.0118 (17)	0.022 (2)	-0.0011 (14)	0.0012 (15)	0.0009 (15)
C14	0.056 (3)	0.020 (2)	0.037 (3)	0.004 (2)	-0.023 (2)	-0.0014 (19)
C15	0.022 (2)	0.049 (3)	0.043 (3)	0.0003 (19)	-0.0048 (19)	0.024 (2)
C16	0.028 (2)	0.0204 (19)	0.021 (2)	0.0025 (16)	-0.0007 (16)	0.0022 (16)
C17	0.0097 (16)	0.0196 (18)	0.0162 (19)	-0.0037 (14)	0.0024 (14)	0.0003 (15)
C18	0.0099 (16)	0.0156 (18)	0.025 (2)	-0.0029 (13)	0.0026 (14)	0.0024 (15)
C19	0.0129 (17)	0.028 (2)	0.0149 (19)	-0.0086 (15)	-0.0025 (14)	0.0029 (15)
C20	0.0165 (18)	0.0192 (19)	0.027 (2)	-0.0062 (15)	0.0096 (16)	-0.0061 (16)
C21	0.0124 (17)	0.0176 (18)	0.022 (2)	-0.0025 (14)	0.0044 (14)	0.0051 (15)
C22	0.056 (3)	0.026 (2)	0.023 (2)	-0.009 (2)	0.006 (2)	-0.0032 (18)
C23	0.031 (2)	0.029 (2)	0.024 (2)	-0.0056 (18)	0.0015 (18)	-0.0038 (17)
C24	0.028 (2)	0.044 (3)	0.048 (3)	0.008 (2)	0.002 (2)	-0.020 (2)
C25	0.034 (2)	0.030 (2)	0.028 (2)	-0.0032 (18)	-0.0060 (19)	0.0018 (18)
C26	0.042 (3)	0.032 (2)	0.028 (2)	0.006 (2)	0.012 (2)	-0.0031 (19)
C27	0.024 (2)	0.026 (2)	0.044 (3)	0.0008 (17)	0.0085 (19)	0.0089 (19)
C28	0.033 (2)	0.025 (2)	0.026 (2)	-0.0010 (17)	0.0071 (18)	0.0006 (17)

Geometric parameters (Å, °)

Yb1—Cnt1	2.510	C7—H7B	0.9800
Yb1—Cnt2	2.513	C7—H7C	0.9800
Yb1—Cnt3	2.504	C8—H8A	0.9800

Cs1—Cnt1	3.197	C8—H8B	0.9800
Cs1—Cnt2	3.268	C8—H8C	0.9800
Cs1—Cnt3	3.159	C9—C13	1.424 (5)
Yb1—C12	2.742 (3)	C9—C10	1.434 (5)
Yb1—C21	2.750 (3)	C9—Cs1 ⁱⁱⁱ	3.587 (3)
Yb1—C20	2.757 (3)	C10—C11	1.398 (5)
Yb1—C13	2.775 (3)	C10—Cs1 ⁱⁱⁱ	3.559 (3)
Yb1—C3	2.777 (3)	C10—H10A	0.9500
Yb1—C4	2.778 (3)	C11—C12	1.410 (5)
Yb1—C5	2.778 (3)	C11—Cs1 ⁱⁱⁱ	3.427 (3)
Yb1—C11	2.779 (3)	C11—H11A	0.9500
Yb1—C17	2.785 (3)	C12—C13	1.411 (5)
Yb1—C2	2.787 (3)	C12—Cs1 ⁱⁱⁱ	3.379 (3)
Yb1—C19	2.787 (3)	C12—H12A	0.9500
Yb1—C1	2.788 (3)	C13—Cs1 ⁱⁱⁱ	3.457 (3)
Yb1—C18	2.802 (3)	C13—H13A	0.9500
Yb1—C10	2.802 (3)	C14—H14A	0.9800
Yb1—C9	2.832 (3)	C14—H14B	0.9800
Cs1—O1	3.095 (3)	C14—H14C	0.9800
Cs1—C2	3.309 (3)	C15—H15A	0.9800
Cs1—C21 ⁱ	3.337 (3)	C15—H15B	0.9800
Cs1—C20 ⁱ	3.367 (3)	C15—H15C	0.9800
Cs1—C12 ⁱⁱ	3.379 (3)	C16—Cs1 ⁱⁱⁱ	3.810 (4)
Cs1—C17 ⁱ	3.383 (3)	C16—H16A	0.9800
Cs1—C1	3.390 (3)	C16—H16B	0.9800
Cs1—C3	3.396 (3)	C16—H16C	0.9800
Cs1—C18 ⁱ	3.401 (3)	C17—C21	1.417 (5)
Cs1—C19 ⁱ	3.407 (3)	C17—C18	1.425 (5)
Cs1—C11 ⁱⁱ	3.427 (3)	C17—Cs1 ^{iv}	3.383 (3)
Cs1—C13 ⁱⁱ	3.457 (3)	C18—C19	1.390 (5)
Cs1—C5	3.475 (3)	C18—Cs1 ^{iv}	3.401 (3)
Cs1—C4	3.499 (3)	C18—H18A	0.9500
Cs1—C10 ⁱⁱ	3.559 (3)	C19—C20	1.406 (5)
Cs1—C9 ⁱⁱ	3.587 (3)	C19—Cs1 ^{iv}	3.407 (3)
Si1—C1	1.854 (4)	C19—H19A	0.9500
Si1—C8	1.866 (4)	C20—C21	1.407 (5)
Si1—C7	1.866 (4)	C20—Cs1 ^{iv}	3.367 (3)
Si1—C6	1.867 (4)	C20—H20A	0.9500
Si2—C9	1.848 (4)	C21—Cs1 ^{iv}	3.337 (3)
Si2—C15	1.863 (4)	C21—H21A	0.9500
Si2—C16	1.867 (3)	C22—H22A	0.9800
Si2—C14	1.871 (4)	C22—H22B	0.9800
Si3—C17	1.856 (3)	C22—H22C	0.9800
Si3—C22	1.864 (4)	C23—H23A	0.9800
Si3—C24	1.865 (4)	C23—H23B	0.9800
Si3—C23	1.869 (4)	C23—H23C	0.9800
O1—C25	1.434 (5)	C24—H24A	0.9800
O1—C28	1.435 (4)	C24—H24B	0.9800

C1—C5	1.418 (5)	C24—H24C	0.9800
C1—C2	1.423 (5)	C25—C26	1.522 (5)
C2—C3	1.402 (5)	C25—H25A	0.9900
C2—H2A	0.9500	C25—H25B	0.9900
C3—C4	1.407 (5)	C26—C27	1.511 (6)
C3—H3A	0.9500	C26—H26A	0.9900
C4—C5	1.396 (5)	C26—H26B	0.9900
C4—H4A	0.9500	C27—C28	1.507 (5)
C5—H5A	0.9500	C27—H27A	0.9900
C6—H6A	0.9800	C27—H27B	0.9900
C6—H6B	0.9800	C28—H28A	0.9900
C6—H6C	0.9800	C28—H28B	0.9900
C7—H7A	0.9800		
Cnt1—Yb1—Cnt2	120.1	C15—Si2—C14	110.1 (2)
Cnt1—Yb1—Cnt3	116.6	C16—Si2—C14	105.67 (18)
Cnt2—Yb1—Cnt3	122.8	C17—Si3—C22	110.60 (16)
Cnt1—Cs1—O1	88.8	C17—Si3—C24	108.68 (17)
Cnt2—Cs1—O1	94.1	C22—Si3—C24	109.2 (2)
Cnt3—Cs1—O1	127.8	C17—Si3—C23	112.26 (16)
Cnt1—Cs1—Cnt3	109.0	C22—Si3—C23	107.77 (19)
Cnt1—Cs1—Cnt2	121.4	C24—Si3—C23	108.32 (19)
Cnt2—Cs1—Cnt3	114.3	C25—O1—C28	109.0 (3)
Yb1—Cnt1—Cs1	175.3	C25—O1—Cs1	132.2 (2)
Yb1—Cnt2—Cs1	172.5	C28—O1—Cs1	105.9 (2)
Yb1—Cnt3—Cs1	176.7	C5—C1—C2	105.4 (3)
C12—Yb1—C21	133.07 (10)	C5—C1—Si1	126.8 (3)
C12—Yb1—C20	148.28 (10)	C2—C1—Si1	127.7 (3)
C21—Yb1—C20	29.62 (10)	C5—C1—Yb1	74.82 (18)
C12—Yb1—C13	29.63 (10)	C2—C1—Yb1	75.16 (18)
C21—Yb1—C13	118.73 (10)	Si1—C1—Yb1	114.08 (15)
C20—Yb1—C13	121.02 (10)	C5—C1—Cs1	81.47 (19)
C12—Yb1—C3	106.89 (10)	C2—C1—Cs1	74.58 (18)
C21—Yb1—C3	75.43 (10)	Si1—C1—Cs1	111.25 (13)
C20—Yb1—C3	93.17 (11)	Yb1—C1—Cs1	134.51 (12)
C13—Yb1—C3	133.57 (10)	C3—C2—C1	109.1 (3)
C12—Yb1—C4	84.54 (10)	C3—C2—Yb1	75.02 (19)
C21—Yb1—C4	104.58 (10)	C1—C2—Yb1	75.27 (19)
C20—Yb1—C4	121.05 (10)	C3—C2—Cs1	81.44 (19)
C13—Yb1—C4	114.11 (10)	C1—C2—Cs1	80.94 (19)
C3—Yb1—C4	29.34 (10)	Yb1—C2—Cs1	138.44 (12)
C12—Yb1—C5	93.42 (10)	C3—C2—H2A	125.4
C21—Yb1—C5	116.94 (10)	C1—C2—H2A	125.4
C20—Yb1—C5	118.11 (10)	Yb1—C2—H2A	116.2
C13—Yb1—C5	120.08 (10)	Cs1—C2—H2A	105.3
C3—Yb1—C5	48.02 (10)	C2—C3—C4	108.0 (3)
C4—Yb1—C5	29.10 (10)	C2—C3—Yb1	75.79 (19)
C12—Yb1—C11	29.58 (10)	C4—C3—Yb1	75.34 (19)

C21—Yb1—C11	162.49 (10)	C2—C3—Cs1	74.48 (18)
C20—Yb1—C11	161.12 (11)	C4—C3—Cs1	82.34 (19)
C13—Yb1—C11	48.43 (10)	Yb1—C3—Cs1	134.71 (12)
C3—Yb1—C11	104.80 (10)	C2—C3—H3A	126.0
C4—Yb1—C11	76.00 (10)	C4—C3—H3A	126.0
C5—Yb1—C11	72.15 (10)	Yb1—C3—H3A	115.1
C12—Yb1—C17	104.81 (10)	Cs1—C3—H3A	110.0
C21—Yb1—C17	29.67 (9)	C5—C4—C3	107.5 (3)
C20—Yb1—C17	49.15 (10)	C5—C4—Yb1	75.45 (19)
C13—Yb1—C17	89.55 (10)	C3—C4—Yb1	75.31 (19)
C3—Yb1—C17	91.45 (10)	C5—C4—Cs1	77.52 (19)
C4—Yb1—C17	115.93 (10)	C3—C4—Cs1	74.17 (18)
C5—Yb1—C17	139.21 (10)	Yb1—C4—Cs1	130.28 (12)
C11—Yb1—C17	134.15 (10)	C5—C4—H4A	126.3
C12—Yb1—C2	132.66 (10)	C3—C4—H4A	126.3
C21—Yb1—C2	69.27 (10)	Yb1—C4—H4A	115.3
C20—Yb1—C2	74.47 (10)	Cs1—C4—H4A	114.4
C13—Yb1—C2	161.85 (10)	C4—C5—C1	109.9 (3)
C3—Yb1—C2	29.18 (10)	C4—C5—Yb1	75.44 (19)
C4—Yb1—C2	48.21 (10)	C1—C5—Yb1	75.66 (18)
C5—Yb1—C2	47.93 (10)	C4—C5—Cs1	79.4 (2)
C11—Yb1—C2	119.41 (10)	C1—C5—Cs1	74.73 (19)
C17—Yb1—C2	95.49 (10)	Yb1—C5—Cs1	131.24 (12)
C12—Yb1—C19	122.04 (10)	C4—C5—H5A	125.0
C21—Yb1—C19	48.35 (10)	C1—C5—H5A	125.0
C20—Yb1—C19	29.37 (10)	Yb1—C5—H5A	115.8
C13—Yb1—C19	92.85 (10)	Cs1—C5—H5A	112.9
C3—Yb1—C19	121.44 (10)	Si1—C6—H6A	109.5
C4—Yb1—C19	150.23 (10)	Si1—C6—H6B	109.5
C5—Yb1—C19	142.33 (10)	H6A—C6—H6B	109.5
C11—Yb1—C19	133.69 (10)	Si1—C6—H6C	109.5
C17—Yb1—C19	48.72 (10)	H6A—C6—H6C	109.5
C2—Yb1—C19	103.58 (10)	H6B—C6—H6C	109.5
C12—Yb1—C1	122.63 (10)	Si1—C7—H7A	109.5
C21—Yb1—C1	94.75 (10)	Si1—C7—H7B	109.5
C20—Yb1—C1	89.08 (10)	H7A—C7—H7B	109.5
C13—Yb1—C1	146.41 (10)	Si1—C7—H7C	109.5
C3—Yb1—C1	48.85 (10)	H7A—C7—H7C	109.5
C4—Yb1—C1	48.90 (10)	H7B—C7—H7C	109.5
C5—Yb1—C1	29.52 (9)	Si1—C8—H8A	109.5
C11—Yb1—C1	98.33 (10)	Si1—C8—H8B	109.5
C17—Yb1—C1	123.39 (10)	H8A—C8—H8B	109.5
C2—Yb1—C1	29.57 (9)	Si1—C8—H8C	109.5
C19—Yb1—C1	113.15 (10)	H8A—C8—H8C	109.5
C12—Yb1—C18	100.20 (10)	H8B—C8—H8C	109.5
C21—Yb1—C18	48.09 (10)	C13—C9—C10	105.1 (3)
C20—Yb1—C18	48.09 (10)	C13—C9—Si2	129.1 (3)
C13—Yb1—C18	74.83 (10)	C10—C9—Si2	124.3 (3)

C3—Yb1—C18	120.22 (10)	C13—C9—Yb1	73.08 (19)
C4—Yb1—C18	145.35 (10)	C10—C9—Yb1	74.11 (18)
C5—Yb1—C18	164.74 (10)	Si2—C9—Yb1	128.22 (15)
C11—Yb1—C18	123.02 (10)	C13—C9—Cs1 ⁱⁱⁱ	73.26 (18)
C17—Yb1—C18	29.55 (9)	C10—C9—Cs1 ⁱⁱⁱ	77.33 (18)
C2—Yb1—C18	116.83 (10)	Si2—C9—Cs1 ⁱⁱⁱ	104.19 (12)
C19—Yb1—C18	28.80 (10)	Yb1—C9—Cs1 ⁱⁱⁱ	127.59 (11)
C1—Yb1—C18	137.11 (10)	C11—C10—C9	109.8 (3)
C12—Yb1—C10	48.31 (10)	C11—C10—Yb1	74.58 (19)
C21—Yb1—C10	156.10 (10)	C9—C10—Yb1	76.40 (19)
C20—Yb1—C10	132.46 (11)	C11—C10—Cs1 ⁱⁱⁱ	73.21 (18)
C13—Yb1—C10	48.00 (10)	C9—C10—Cs1 ⁱⁱⁱ	79.51 (18)
C3—Yb1—C10	128.46 (10)	Yb1—C10—Cs1 ⁱⁱⁱ	129.76 (12)
C4—Yb1—C10	99.32 (10)	C11—C10—H10A	125.1
C5—Yb1—C10	84.76 (10)	C9—C10—H10A	125.1
C11—Yb1—C10	29.00 (10)	Yb1—C10—H10A	115.8
C17—Yb1—C10	134.24 (10)	Cs1 ⁱⁱⁱ —C10—H10A	114.2
C2—Yb1—C10	130.23 (10)	C10—C11—C12	107.9 (3)
C19—Yb1—C10	108.45 (10)	C10—C11—Yb1	76.42 (18)
C1—Yb1—C10	101.51 (10)	C12—C11—Yb1	73.76 (18)
C18—Yb1—C10	109.46 (10)	C10—C11—Cs1 ⁱⁱⁱ	83.81 (19)
C12—Yb1—C9	49.06 (10)	C12—C11—Cs1 ⁱⁱⁱ	76.14 (19)
C21—Yb1—C9	128.10 (10)	Yb1—C11—Cs1 ⁱⁱⁱ	136.45 (12)
C20—Yb1—C9	113.55 (10)	C10—C11—H11A	126.1
C13—Yb1—C9	29.39 (9)	C12—C11—H11A	126.1
C3—Yb1—C9	153.27 (10)	Yb1—C11—H11A	115.9
C4—Yb1—C9	124.75 (10)	Cs1 ⁱⁱⁱ —C11—H11A	107.2
C5—Yb1—C9	114.22 (10)	C11—C12—C13	107.7 (3)
C11—Yb1—C9	48.76 (10)	C11—C12—Yb1	76.66 (19)
C17—Yb1—C9	105.13 (10)	C13—C12—Yb1	76.46 (19)
C2—Yb1—C9	158.00 (10)	C11—C12—Cs1 ⁱⁱⁱ	79.95 (19)
C19—Yb1—C9	84.94 (10)	C13—C12—Cs1 ⁱⁱⁱ	81.20 (19)
C1—Yb1—C9	128.44 (10)	Yb1—C12—Cs1 ⁱⁱⁱ	140.67 (13)
C18—Yb1—C9	80.25 (10)	C11—C12—H12A	126.1
C10—Yb1—C9	29.49 (9)	C13—C12—H12A	126.1
O1—Cs1—C2	80.28 (8)	Yb1—C12—H12A	113.2
O1—Cs1—C21 ⁱ	114.33 (8)	Cs1 ⁱⁱⁱ —C12—H12A	106.1
C2—Cs1—C21 ⁱ	148.98 (8)	C12—C13—C9	109.6 (3)
O1—Cs1—C20 ⁱ	138.03 (8)	C12—C13—Yb1	73.92 (19)
C2—Cs1—C20 ⁱ	137.32 (9)	C9—C13—Yb1	77.53 (19)
C21 ⁱ —Cs1—C20 ⁱ	24.23 (8)	C12—C13—Cs1 ⁱⁱⁱ	75.02 (18)
O1—Cs1—C12 ⁱⁱ	110.63 (8)	C9—C13—Cs1 ⁱⁱⁱ	83.51 (19)
C2—Cs1—C12 ⁱⁱ	92.69 (8)	Yb1—C13—Cs1 ⁱⁱⁱ	135.26 (12)
C21 ⁱ —Cs1—C12 ⁱⁱ	105.96 (8)	C12—C13—H13A	125.2
C20 ⁱ —Cs1—C12 ⁱⁱ	89.07 (8)	C9—C13—H13A	125.2
O1—Cs1—C17 ⁱ	107.40 (7)	Yb1—C13—H13A	115.3
C2—Cs1—C17 ⁱ	127.34 (8)	Cs1 ⁱⁱⁱ —C13—H13A	108.8
C21 ⁱ —Cs1—C17 ⁱ	24.34 (8)	Si2—C14—H14A	109.5

C20 ⁱ —Cs1—C17 ⁱ	39.93 (8)	Si2—C14—H14B	109.5
C12 ⁱⁱ —Cs1—C17 ⁱ	128.47 (8)	H14A—C14—H14B	109.5
O1—Cs1—C1	104.30 (7)	Si2—C14—H14C	109.5
C2—Cs1—C1	24.49 (8)	H14A—C14—H14C	109.5
C21 ⁱ —Cs1—C1	129.35 (8)	H14B—C14—H14C	109.5
C20 ⁱ —Cs1—C1	113.14 (8)	Si2—C15—H15A	109.5
C12 ⁱⁱ —Cs1—C1	88.68 (8)	Si2—C15—H15B	109.5
C17 ⁱ —Cs1—C1	114.09 (8)	H15A—C15—H15B	109.5
O1—Cs1—C3	68.30 (8)	Si2—C15—H15C	109.5
C2—Cs1—C3	24.09 (8)	H15A—C15—H15C	109.5
C21 ⁱ —Cs1—C3	133.69 (8)	H15B—C15—H15C	109.5
C20 ⁱ —Cs1—C3	136.14 (9)	Si2—C16—Cs1 ⁱⁱⁱ	96.16 (13)
C12 ⁱⁱ —Cs1—C3	116.22 (8)	Si2—C16—H16A	109.5
C17 ⁱ —Cs1—C3	109.40 (8)	Cs1 ⁱⁱⁱ —C16—H16A	66.8
C1—Cs1—C3	39.65 (8)	Si2—C16—H16B	109.5
O1—Cs1—C18 ⁱ	124.90 (8)	Cs1 ⁱⁱⁱ —C16—H16B	52.7
C2—Cs1—C18 ⁱ	109.78 (8)	H16A—C16—H16B	109.5
C21 ⁱ —Cs1—C18 ⁱ	39.23 (8)	Si2—C16—H16C	109.5
C20 ⁱ —Cs1—C18 ⁱ	39.10 (8)	Cs1 ⁱⁱⁱ —C16—H16C	153.3
C12 ⁱⁱ —Cs1—C18 ⁱ	122.27 (8)	H16A—C16—H16C	109.5
C17 ⁱ —Cs1—C18 ⁱ	24.24 (8)	H16B—C16—H16C	109.5
C1—Cs1—C18 ⁱ	91.83 (8)	C21—C17—C18	105.5 (3)
C3—Cs1—C18 ⁱ	98.84 (8)	C21—C17—Si3	128.0 (3)
O1—Cs1—C19 ⁱ	146.73 (8)	C18—C17—Si3	126.4 (3)
C2—Cs1—C19 ⁱ	114.55 (8)	C21—C17—Yb1	73.79 (18)
C21 ⁱ —Cs1—C19 ⁱ	39.29 (8)	C18—C17—Yb1	75.89 (18)
C20 ⁱ —Cs1—C19 ⁱ	23.95 (8)	Si3—C17—Yb1	118.31 (14)
C12 ⁱⁱ —Cs1—C19 ⁱ	98.71 (9)	C21—C17—Cs1 ^{iv}	75.99 (18)
C17 ⁱ —Cs1—C19 ⁱ	39.57 (8)	C18—C17—Cs1 ^{iv}	78.58 (18)
C1—Cs1—C19 ⁱ	91.36 (8)	Si3—C17—Cs1 ^{iv}	108.65 (13)
C3—Cs1—C19 ⁱ	112.73 (8)	Yb1—C17—Cs1 ^{iv}	132.98 (11)
C18 ⁱ —Cs1—C19 ⁱ	23.56 (8)	C19—C18—C17	109.5 (3)
O1—Cs1—C11 ⁱⁱ	87.61 (8)	C19—C18—Yb1	75.02 (19)
C2—Cs1—C11 ⁱⁱ	82.35 (8)	C17—C18—Yb1	74.57 (18)
C21 ⁱ —Cs1—C11 ⁱⁱ	123.61 (8)	C19—C18—Cs1 ^{iv}	78.45 (19)
C20 ⁱ —Cs1—C11 ⁱⁱ	111.22 (8)	C17—C18—Cs1 ^{iv}	77.18 (18)
C12 ⁱⁱ —Cs1—C11 ⁱⁱ	23.90 (8)	Yb1—C18—Cs1 ^{iv}	131.52 (11)
C17 ⁱ —Cs1—C11 ⁱⁱ	147.92 (8)	C19—C18—H18A	125.3
C1—Cs1—C11 ⁱⁱ	88.14 (8)	C17—C18—H18A	125.3
C3—Cs1—C11 ⁱⁱ	102.45 (8)	Yb1—C18—H18A	117.0
C18 ⁱ —Cs1—C11 ⁱⁱ	146.18 (8)	Cs1 ^{iv} —C18—H18A	111.4
C19 ⁱ —Cs1—C11 ⁱⁱ	122.61 (8)	C18—C19—C20	108.2 (3)
O1—Cs1—C13 ⁱⁱ	110.19 (8)	C18—C19—Yb1	76.18 (19)
C2—Cs1—C13 ⁱⁱ	116.34 (8)	C20—C19—Yb1	74.12 (19)
C21 ⁱ —Cs1—C13 ⁱⁱ	85.36 (8)	C18—C19—Cs1 ^{iv}	77.99 (19)
C20 ⁱ —Cs1—C13 ⁱⁱ	73.64 (8)	C20—C19—Cs1 ^{iv}	76.46 (19)
C12 ⁱⁱ —Cs1—C13 ⁱⁱ	23.79 (8)	Yb1—C19—Cs1 ^{iv}	131.87 (12)
C17 ⁱ —Cs1—C13 ⁱⁱ	109.44 (8)	C18—C19—H19A	125.9

C1—Cs1—C13 ⁱⁱ	111.22 (8)	C20—C19—H19A	125.9
C3—Cs1—C13 ⁱⁱ	139.38 (8)	Yb1—C19—H19A	115.9
C18 ⁱ —Cs1—C13 ⁱⁱ	111.98 (8)	Cs1 ^{iv} —C19—H19A	112.2
C19 ⁱ —Cs1—C13 ⁱⁱ	90.34 (8)	C19—C20—C21	107.4 (3)
C11 ⁱⁱ —Cs1—C13 ⁱⁱ	38.65 (8)	C19—C20—Yb1	76.50 (19)
O1—Cs1—C5	105.94 (7)	C21—C20—Yb1	74.91 (19)
C2—Cs1—C5	38.83 (8)	C19—C20—Cs1 ^{iv}	79.59 (19)
C21 ⁱ —Cs1—C5	110.42 (8)	C21—C20—Cs1 ^{iv}	76.66 (18)
C20 ⁱ —Cs1—C5	101.17 (8)	Yb1—C20—Cs1 ^{iv}	134.91 (12)
C12 ⁱⁱ —Cs1—C5	109.56 (8)	C19—C20—H20A	126.3
C17 ⁱ —Cs1—C5	91.53 (8)	C21—C20—H20A	126.3
C1—Cs1—C5	23.80 (8)	Yb1—C20—H20A	114.7
C3—Cs1—C5	38.38 (8)	Cs1 ^{iv} —C20—H20A	110.3
C18 ⁱ —Cs1—C5	71.25 (8)	C20—C21—C17	109.3 (3)
C19 ⁱ —Cs1—C5	77.29 (8)	C20—C21—Yb1	75.48 (19)
C11 ⁱⁱ —Cs1—C5	111.87 (8)	C17—C21—Yb1	76.54 (18)
C13 ⁱⁱ —Cs1—C5	129.48 (8)	C20—C21—Cs1 ^{iv}	79.11 (19)
O1—Cs1—C4	84.87 (8)	C17—C21—Cs1 ^{iv}	79.67 (18)
C2—Cs1—C4	38.88 (8)	Yb1—C21—Cs1 ^{iv}	136.70 (12)
C21 ⁱ —Cs1—C4	112.41 (8)	C20—C21—H21A	125.3
C20 ⁱ —Cs1—C4	112.73 (8)	C17—C21—H21A	125.3
C12 ⁱⁱ —Cs1—C4	127.43 (8)	Yb1—C21—H21A	114.7
C17 ⁱ —Cs1—C4	88.95 (8)	Cs1 ^{iv} —C21—H21A	108.6
C1—Cs1—C4	39.03 (8)	Si3—C22—H22A	109.5
C3—Cs1—C4	23.49 (8)	Si3—C22—H22B	109.5
C18 ⁱ —Cs1—C4	75.59 (8)	H22A—C22—H22B	109.5
C19 ⁱ —Cs1—C4	89.67 (8)	Si3—C22—H22C	109.5
C11 ⁱⁱ —Cs1—C4	121.18 (8)	H22A—C22—H22C	109.5
C13 ⁱⁱ —Cs1—C4	150.24 (8)	H22B—C22—H22C	109.5
C5—Cs1—C4	23.09 (8)	Si3—C23—H23A	109.5
O1—Cs1—C10 ⁱⁱ	74.63 (7)	Si3—C23—H23B	109.5
C2—Cs1—C10 ⁱⁱ	98.69 (8)	H23A—C23—H23B	109.5
C21 ⁱ —Cs1—C10 ⁱⁱ	111.34 (8)	Si3—C23—H23C	109.5
C20 ⁱ —Cs1—C10 ⁱⁱ	108.30 (8)	H23A—C23—H23C	109.5
C12 ⁱⁱ —Cs1—C10 ⁱⁱ	38.08 (8)	H23B—C23—H23C	109.5
C17 ⁱ —Cs1—C10 ⁱⁱ	133.90 (8)	Si3—C24—H24A	109.5
C1—Cs1—C10 ⁱⁱ	109.52 (8)	Si3—C24—H24B	109.5
C3—Cs1—C10 ⁱⁱ	113.49 (8)	H24A—C24—H24B	109.5
C18 ⁱ —Cs1—C10 ⁱⁱ	147.28 (8)	Si3—C24—H24C	109.5
C19 ⁱ —Cs1—C10 ⁱⁱ	127.76 (8)	H24A—C24—H24C	109.5
C11 ⁱⁱ —Cs1—C10 ⁱⁱ	22.98 (8)	H24B—C24—H24C	109.5
C13 ⁱⁱ —Cs1—C10 ⁱⁱ	37.70 (8)	O1—C25—C26	105.8 (3)
C5—Cs1—C10 ⁱⁱ	133.23 (8)	O1—C25—H25A	110.6
C4—Cs1—C10 ⁱⁱ	136.14 (8)	C26—C25—H25A	110.6
O1—Cs1—C9 ⁱⁱ	87.86 (7)	O1—C25—H25B	110.6
C2—Cs1—C9 ⁱⁱ	120.23 (8)	C26—C25—H25B	110.6
C21 ⁱ —Cs1—C9 ⁱⁱ	88.67 (8)	H25A—C25—H25B	108.7
C20 ⁱ —Cs1—C9 ⁱⁱ	85.71 (8)	C27—C26—C25	101.6 (3)

C12 ⁱⁱ —Cs1—C9 ⁱⁱ	38.69 (8)	C27—C26—H26A	111.5
C17 ⁱ —Cs1—C9 ⁱⁱ	112.18 (8)	C25—C26—H26A	111.5
C1—Cs1—C9 ⁱⁱ	125.32 (8)	C27—C26—H26B	111.5
C3—Cs1—C9 ⁱⁱ	136.64 (8)	C25—C26—H26B	111.5
C18 ⁱ —Cs1—C9 ⁱⁱ	124.31 (8)	H26A—C26—H26B	109.3
C19 ⁱ —Cs1—C9 ⁱⁱ	106.95 (8)	C28—C27—C26	102.1 (3)
C11 ⁱⁱ —Cs1—C9 ⁱⁱ	38.49 (8)	C28—C27—H27A	111.3
C13 ⁱⁱ —Cs1—C9 ⁱⁱ	23.23 (8)	C26—C27—H27A	111.3
C5—Cs1—C9 ⁱⁱ	147.92 (8)	C28—C27—H27B	111.3
C4—Cs1—C9 ⁱⁱ	158.85 (8)	C26—C27—H27B	111.3
C10 ⁱⁱ —Cs1—C9 ⁱⁱ	23.16 (7)	H27A—C27—H27B	109.2
C1—Si1—C8	109.90 (16)	O1—C28—C27	106.8 (3)
C1—Si1—C7	109.33 (16)	O1—C28—Cs1	52.50 (16)
C8—Si1—C7	108.40 (17)	C27—C28—Cs1	137.3 (2)
C1—Si1—C6	110.42 (16)	O1—C28—H28A	110.4
C8—Si1—C6	110.06 (17)	C27—C28—H28A	110.4
C7—Si1—C6	108.70 (18)	Cs1—C28—H28A	112.0
C9—Si2—C15	112.34 (18)	O1—C28—H28B	110.4
C9—Si2—C16	108.64 (16)	C27—C28—H28B	110.4
C15—Si2—C16	107.99 (17)	Cs1—C28—H28B	60.1
C9—Si2—C14	111.75 (17)	H28A—C28—H28B	108.6

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+3/2, y-1/2, -z+1/2$; (iii) $-x+3/2, y+1/2, -z+1/2$; (iv) $x-1, y, z$.