

Eclogite-driven subsidence of the Columbia Basin (Washington State, USA) caused by deposition of Columbia River Basalt

Jonathan Perry-Houts and Eugene Humphreys

Department of Earth Sciences, University of Oregon, Eugene, Oregon 97403-1272, USA

ABSTRACT

The south-central portion of the Columbia Basin in Washington State (USA), although not the primary eruptive center for the Columbia River Basalt (CRB), contains the thickest package of CRB lavas, at over 4 km. Prior to CRB deposition the region was a sediment-filled rift basin dating back to the early Eocene. Previous studies interpret a lower-crust, high-velocity body below the thick basalt deposits as a syn-extensional magmatic underplate. CRB layers thicken toward the center of the basin, revealing a pulse of subsidence during deposition that ceased coincident with cessation of CRB eruptions. We propose a subsidence mechanism based on metamorphic transition of the underplate from basalt to eclogite facies, driven by the increased pressure of CRB loading. We demonstrate the plausibility of our interpretation using numerical models of the mechanical and chemical components of the system. Further, we constrain the effective elastic thickness of the Miocene Columbia Basin lithosphere, finding it to be much thinner (5–10 km) than previous estimates.

INTRODUCTION

The Columbia River Flood Basalt (CRB) erupted at ca. 17–6 Ma from northeast Oregon and adjacent portions of Idaho and Washington, USA (Fig. 1). These flows covered the Columbia Basin as nearly flat, self-leveling inflation flows. Basalt accumulation was greatest not near the primary source of eruption, but in the center of the Columbia Basin (Fig. 1), where >4 km of CRB ponded (Reidel et al., 1989). As evidenced by the CRB flows thickening toward the center of the basin (Fig. 2), progressive basin subsidence occurred during CRB activity, deepening a depression that CRB flows repeatedly filled and over-topped. Subsidence ended with the cessation of CRB deposition, and the region has not experienced significant subsidence since (Reidel et al., 2013). We use the name central Columbia Basin downwarp (CCBD) to refer to the region of subsidence roughly outlined by the 2 km CRB isopach in Figure 1.

GEOLOGIC CONTEXT

Basic aspects of crustal structure in the greater Columbia Basin region are represented in Figures 1 and 2. A major phase of regional extension and volcanism occurred during the ca. 53 Ma accretion of Siletzia, accompanied by an outward stepping of subduction and establishment of the present Cascadia subduction zone (Christiansen and Yeats, 1992). Regional extension includes the creation of large core complexes in northeastern Washington and trans-tensional basins now exposed along the eastern flank of the north Cascades uplift. The Swauk and Chiwaukum basins (Fig. 1) preserve the most complete record of sedimentation. They are filled with Eocene sediments and volcanic rocks ranging in age from ca. 55 to 40 Ma (Taylor et al., 1988). These basins trend southeast into the CCBD, where they are covered by CRB flows. Drill holes through the CCBD penetrate these same Eocene sedimentary units (Campbell, 1989). The formation ages of the early Swauk, Chiwaukum, and (where resolved) Eocene rift basin in the CCBD are similar to the unroofing of nearby core complexes (54–47 Ma; Kruckenberg et al., 2008), and vigorous regional volcanism (Christiansen and Yeats, 1992; Gaschnig et al., 2010).




Figure 1. Map showing features that correlate with the central Columbia Basin downwarp. Accumulated thickness of the Columbia River Basalt (CRB) flows is indicated with black dashed lines (thickness in meters). CRB ponded in the downwarp, which subsided during the CRB eruptions. Subsidence centered on an Eocene rift basin associated with coeval core complex and trans-tensional basin activity in northern Washington State, USA. The Swauk (S), Chiwaukum (C), Republic (R), and Methow (M) grabens are labeled. The rift basin is imaged with active-source seismology along the cross section (indicated with triangles) (Catchings and Mooney, 1988; Fig. 2) and in three dimensions using surface waves (Gao et al., 2011). The anomalous crustal structure imaged in the immediate vicinity of the downwarp includes 5–6 km of sediment beneath ~4 km of CRB, and ~16 km of high-velocity lowermost crust interpreted by Catchings and Mooney (1988) to be rift-related magmatic underplate. Red and blue lines outline areas of low- and high-velocity crust (Gao et al., 2011). Topography adapted from Global Multi-Resolution Topography (GMRT) synthesis (Ryan et al., 2009).




Figure 2. The upper crust along the cross section indicated in Figure 1 (after Catchings and Mooney, 1988). Numbers on plot give the P-wave velocity in km/s. Tan color is post-Columbia River Basalt (CRB) sediment; blue is CRB flows; green is Eocene sediment; white is basement. Note the deep sediment-filled rift basin, and CRB flows thickening toward the basin center, indicating that the basin subsided during deposition, and not after.

The active source seismic study of Catchings and Mooney (1988) and the surface wave and receiver function study of Gao et al. (2011) provide images of the deep structure in the CCB and surrounding area. Eocene sediments in the CCB locally extend to ~11 km beneath the surface, defining a deep, local (~110 × 80 km) rift structure. Immediately below the area of deep sediment accumulation, and of similar horizontal dimensions, is an anomalous region of seismically high-velocity ($V_p = 7.5$ km/s) lower crust extending from ~23 km to ~40 km depth (Fig. 1). We assume this to be the basaltic underplate portion of the rift structure, as proposed by Catchings and Mooney (1988), with the 40 km seismic Moho representing local basalt-eclogite coexistence pressure, and underlying root seismically indistinguishable from surrounding mantle. The crustal section from 11 km to 23 km depth has been interpreted as either crystalline North America (Catchings and Mooney, 1988) or Farallon slab connected at depth to Siletzia (Gao et al., 2011). Either interpretation is consistent with our modeling.

ECLOGITE-DRIVEN SUBSIDENCE

The rate and magnitude of Miocene subsidence in south-central Washington is far greater than any other CRB depocenter, at up to 10 mm/yr throughout Grande Ronde deposition (Reidel et al., 2013). Because this time period does not coincide with any major change in regional tectonics (Christiansen and Yeats, 1992), we infer subsidence was driven by local vertical forcing rather than far-field horizontal tectonics.

Isostasy and Flexure

The core logic of our model is a simple isostatic argument. Loading from CRB deposition increases lithostatic pressure throughout the crust, pushing lower-crustal basaltic rocks across the phase transition to eclogite facies. Rocks densify as they metamorphose from basalt to eclogite, contracting the crust and creating a topographic low relative to a column that did not eclogitize. The resulting depression traps subsequent lava flows, compounding the effect over time.

Conceptually, we model the mechanical aspects of this system in two distinct steps. First, when a new basalt flow is deposited, we assume that its weight is transmitted hydrostatically to the lower crust. This implies that an equal volume of lower crust is eclogitized directly below the new deposit.

Mechanical contraction is filtered by the elastic strength of the plate, producing a smoothed topographic response. We model this effect as an equivalent load applied to an isostatically supported elastic plate. We use a thin-plate assumption (Turcotte and Schubert, 2014), for which the dominant forces are buoyancy and elastic bending stress, as expressed by the equation

$$\nabla^2 D \nabla^2 w(x, y) + [\rho_m - \rho_b] g w(x, y) = L(x, y), \quad (1)$$

where w is displacement as a function of position (x, y) ; ρ_b and ρ_m are the density of basalt and mantle, respectively; g is gravity; L is vertical loading stress; and D is flexural rigidity, defined as

$$D = \frac{ET_e^3}{12(1-\nu^2)}, \quad (2)$$

where E is Young's Modulus, T_e is effective elastic thickness, and ν is Poisson's ratio.

We numerically solved Equation 1 with the finite difference method. We initialize our model with a small topographic depression, representing the Eocene basin just prior to CRB onset, which we take to be in isostatic equilibrium. Assuming that an equal volume of lower-crustal basalt eclogitizes upon deposition, we expect a proportional reduction in crustal volume.

We use reference densities of 3000, 3300, and 3370 kg m⁻³ for basalt, mantle, and eclogite, respectively, to calculate the effective load of

eclogitization. We assume an elastic plate with Young's Modulus of 7×10^{10} Pa, and Poisson's ratio of 0.25. Each time step represents a lava flow filling the existing depression, and the crustal downwarping caused by the newly formed eclogite load. Each event adds a proportional volume of eclogite to the root structure, for a cumulative root growth proportional to cumulative deposition.

Our calculations assume the system reaches isostatic equilibrium between lava flows, and omits dynamic effects such as mantle convection and lateral crustal flow, which may affect the long-term evolution of the system (Wang and Currie, 2017).

The nature of Equation 1 results in a smooth subsidence response, making the model relatively insensitive to the detailed shape of the initial basin. For the models presented in Figure 3, our initial geometry is a radial Gaussian function with root mean square (RMS) width of 15 km and depth of 50 m. Any short-wavelength load excites the natural bending wavelength of the crust. If the initial basin is much wider than the natural bending wavelength, the basin localizes to the natural bending wavelength with repeated lava flows. A deeper initial basin results in faster basin development. The long-term resulting structure is qualitatively the same in all cases. Model results are shown in Figure 3, and further details of our methods can be found in the GSA Data Repository¹.

Based on our modeling, the observed basin geometry is possible only if the crust has an effective elastic thickness between 5 and 10 km. This differs from the 25–35 km estimate of Lowry and Pérez-Gussinyé (2011), which is based on the ratio of topography to Bouguer gravity. We think the discrepancy in estimated effective elastic thickness is a consequence of topographic flattening created by the CRB deposits.

Reaction Kinetics

Although we describe our flexural model as an iteration over individual lava flows, it makes no predictions of the actual time elapsed between steps. Rather, each model step represents time taken for the slowest physical process in the system to act. In the following paragraphs we show that neither reaction processes, nor latent heat release, hinders the basalt to eclogite transition on scales relevant to eclogite-driven subsidence. Therefore, we conclude that the duration of the CRB deposition is the limiting mechanism for this system. That is, by the time each new flow begins, metamorphism owing to the previous flow has come to equilibrium.

Previous studies on the basalt to eclogite transition have found that eclogitization takes place on geologically relevant time scales at temperatures above 400–600 °C in dry basalt (Ito and Kennedy, 1971; Ahrens and Schubert, 1975), although dry gabbroic facies may remain metastable for substantial time, even at high temperatures (Hacker, 2013). However, the same reactions progress rapidly to stable equilibrium in the presence of small amounts of water (Ahrens and Schubert, 1975; Hacker, 2013), as we expect to exist in abundance in the Cascadia backarc. Hence, we assume the lower crust of the Columbia Basin is capable of reacting quickly and completely from basalt to eclogite upon local changes in pressure and temperature.

The latent heat of reaction inhibits eclogitization. As this heat diffuses away, the reaction can continue. The following analysis shows the release of latent heat has a negligible effect on the reactions we consider. Following O'Connell and Wasserburg (1972), we consider a vertical thermochemical profile through the crust. The deposition of lava both advects the geotherm down with the sinking crust, and causes the release of latent heat at depth. This modified geotherm then conductively cools back toward its original equilibrium slope.

We model the thermo-chemical system with a heat conservation equation and an Arrhenius reaction equation, respectively:

¹GSA Data Repository item 2018221, physical and thermo-chemical models of eclogite-induced subsidence, is available online at <http://www.geosociety.org/datarepository/2018/> or on request from editing@geosociety.org.




Figure 3. Models of time-evolving eclogite loading on an isostatically supported elastic plate. A: Beginning with a small depocenter, a larger basin develops by repeatedly filling the existing topography and modeling new subsidence owing to lower-crustal eclogite formation. B: Depositional interfaces over time, illustrating the compounding effect of the eclogitization subsidence mechanism. Each flow is deposited as a flat layer filling the previous depression, and subsequently downwarps as a result of newly generated eclogite. Color represents the sequence of deposition. Line thickness is proportional to the thickness of the basalt layer. Inset shows zoom-in of initial topography. C: Basin geometry characteristics over time, for varying elastic plate thicknesses. Other parameters are held constant across models. Basin depth increases slowly at first, and accelerates as the basin size, and hence load, increases. Basin radius rapidly adjusts to reflect the plate's natural flexural wavelength. Aspect ratio (radius/depth) reveals that, in this adjustment stage, radial growth briefly out-paces deepening. Long-term basin growth is dominated by deepening. D: The effects demonstrated in C trade off differently for different elastic plate thicknesses. Contours denote basin depth and radius over time for different plate thicknesses. Models whose geometries are similar to the Columbia Basin are plotted in filled circles, setting narrow limits on possible Miocene plate thickness.

$$\rho C_p \left(\frac{\partial T}{\partial t} + u \cdot \nabla T \right) - \nabla \cdot (\rho C_p \alpha \nabla T) = \rho T \Delta S \left(\frac{\partial X}{\partial t} + u \cdot \nabla X \right), \quad (3)$$

and

$$\frac{\partial X}{\partial t} = A \exp \left(\frac{-E_a}{RT} \right) \left\{ \frac{1}{2} \left[1 + \tanh \left(\frac{P - P_c}{\delta_p} \right) \right] - X \right\}, \quad (4)$$

where ρ is density, C_p is specific heat capacity, T is temperature, t is time, u is deposition rate, α is thermal diffusivity, ΔS is specific entropy change of eclogitization, X is fraction eclogite facies, A is an Arrhenius coefficient, E_a is activation energy of reaction, R is the universal gas constant, P is pressure, P_c is facies coexistence pressure as a function of temperature, and δ_p is a smoothing factor for the coexistence boundary as a function of pressure.

We non-dimensionalized Equations 3 and 4 using a modified Péclet number for chemically reactive continuum,

$$Pe_r = \frac{U^3}{SG\alpha}, \quad (5)$$

where G is a reference geothermal gradient, and other boldface variables are representative values of variables in Equation 3. Finite-element solutions to Equations 3 and 4, and instructions for reproduction, are provided in the Data Repository. However, one can easily verify that, for any geological system, $Pe_r \ll 1$. Therefore, the system tends toward the limiting case of instantaneous thermal diffusion, implying that latent heat has negligible effect on this system.

DISCUSSION

Eclogitization has been discussed before as a driver of localized dynamic topography. A model similar to the one presented here has been proposed for sediment-loaded intracratonic basins (Naimark and Ismail-Zadeh, 1995; Hamdani et al., 1994), where eclogitization is driven primarily by a cooling geotherm, and surface loading is of secondary importance. Compressional tectonics may also drive eclogitization, forming basins like the East Barents Sea (Gac et al., 2013), and affecting the buoyancy structure of aging mountain belts (Fischer, 2002).

The minimum requirement for eclogite-driven subsidence is the presence of basaltic lower crust. Rift structures are common localities for

lower-crustal magmatic underplates. The CCB, and intracratonic basins mentioned above, all have rift histories. However, lower-crustal gabbro may also form in significant quantities in other settings, such as at arcs and hot spots.

Additional lower-crustal density owing to eclogitization could be a precursor to lithospheric delamination, and associated lithospheric recycling, uplift, and volcanism (Krystopowicz and Currie, 2013). It is noteworthy that the mechanisms discussed here are examples of so-called vertical tectonics, driven not by far-field horizontal forces, but by localized changes in lithospheric buoyancy. This class of tectonic processes appears to be a significant driver of tectonic evolution from the local and concentrated scale of the CCB, to uplift of the entire North American Cordillera.

CONCLUSION

The occurrence of a local downwarp only during the short duration of CRB deposition, and at the site of an apparent earlier magmatic underplate, strongly suggests that the event was driven by creation of a local load on the crust. With simple numerical models of the physical and chemical components of the system, and inferring from the pre-Miocene geology a basaltic lower crust straddling the basalt-eclogite transition, we show that formation of lower-crustal eclogite during CRB loading would drive localized subsidence in a manner consistent with the modern Columbia Basin geometry and structure. We can think of no other satisfactory explanations for the anomalous, localized Miocene subsidence of the central Columbia Basin. Further, our models constrain the effective elastic thickness of the Miocene Columbia Basin crust to <10 km.

Our study supports the hypothesis, proposed by others, that similar mechanisms may explain other localities of anomalous subsidence. In general, lower-crustal eclogitization is capable of producing deep basins without far-field tectonic driving, and may be responsible for anomalous topographic features globally.

ACKNOWLEDGMENTS

We thank Claire Currie, Vic Camp, and an anonymous reviewer for constructive reviews that improved the quality and clarity of this paper. This work was supported by National Science Foundation grant EAR-1727451.

REFERENCES CITED

- Ahrens, T.J., and Schubert, G., 1975, Gabbro-eclogite reaction rate and its geophysical significance: *Reviews of Geophysics*, v. 13, p. 383–400, <https://doi.org/10.1029/RG013i002p00383>.
- Campbell, N., 1989, Structural and stratigraphic interpretation of rocks under the Yakima fold belt, Columbia Basin, based on recent surface mapping and well data, in Reidel, S.P., and Hooper, P.R., eds., *Volcanism and Tectonism in the Columbia River Flood-basalt Province*: Geological Society of America Special Papers, v. 239, p. 209–222, <https://doi.org/10.1130/SPE239-p209>.
- Catchings, R.D., and Mooney, W.D., 1988, Crustal structure of the Columbia Plateau: Evidence for continental rifting: *Journal of Geophysical Research: Solid Earth*, v. 93, p. 459–474, <https://doi.org/10.1029/JB093iB01p00459>.
- Christiansen, R.L., and Yeats, R.S., 1992, Post-Laramide geology of the U.S. Cordilleran region, in Burchfiel, B.C., et al., eds., *The Cordilleran Orogen: Conterminous U.S.*: Boulder, Colorado, Geological Society of America, *Geology of North America*, v. G-3, p. 261–406, <https://doi.org/10.1130/DNAG-GNA-G3.261>.
- Fischer, K.M., 2002, Waning buoyancy in the crustal roots of old mountains: *Nature*, v. 417, p. 933–936 <https://doi.org/10.1038/nature00855>.
- Gac, S., Huisman, R.S., Simon, N.S.C., Podladchikov, Y.Y., and Faleide, J.I., 2013, Formation of intracratonic basins by lithospheric shortening and phase changes: A case study from the ultra-deep East Barents Sea basin: *Terra Nova*, v. 25, p. 459–464, <https://doi.org/10.1111/ter.12057>.
- Gao, H., Humphreys, E.D., Yao, H., and van der Hilst, R.D., 2011, Crust and lithosphere structure of the northwestern U.S. with ambient noise tomography: Terrane accretion and Cascade arc development: *Earth and Planetary Science Letters*, v. 304, p. 202–211, <https://doi.org/10.1016/j.epsl.2011.01.033>.
- Gaschnig, R.M., Vervoort, J.D., Lewis, R.S., and McClelland, W.C., 2010, Migrating magmatism in the northern US Cordillera: In situ U-Pb geochronology of the Idaho batholith: *Contributions to Mineralogy and Petrology*, v. 159, p. 863–883, <https://doi.org/10.1007/s00410-009-0459-5>.
- Hacker, B.R., 2013, Eclogite formation and the rheology, buoyancy, seismicity, and H₂O content of oceanic crust, in Bebout, G.E., et al., eds., *Subduction Top to Bottom: American Geophysical Union Geophysical Monograph Series*, v. 96, p. 337–346, <https://doi.org/10.1029/GM096p0337>.
- Hamdani, Y., Mareschal, J.-C., and Arkani-Hamed, J., 1994, Phase change and thermal subsidence of the Williston basin: *Geophysical Journal International*, v. 116, p. 585–597, <https://doi.org/10.1111/j.1365-246X.1994.tb03282.x>.
- Ito, K., and Kennedy, G.C., 1971, An Experimental Study of the Basalt-Garnet Granulite-Eclogite Transition, in Heacock, J.G., ed., *The Structure and Physical Properties of the Earth's Crust: American Geophysical Union, Geophysical Monograph Series*, v. 14, p. 303–314, <https://doi.org/10.1029/GM014p0303>.
- Kruckenberger, S.C., Whitney, D.L., Teyssier, C., Fanning, C.M., and Dunlap, W.J., 2008, Paleocene-Eocene migmatite crystallization, extension, and exhumation in the hinterland of the northern Cordillera: Okanogan dome, Washington, USA: *Geological Society of America Bulletin*, v. 120, p. 912–929, <https://doi.org/10.1130/B26153.1>.
- Krystopowicz, N.J., and Currie, C.A., 2013, Crustal eclogitization and lithosphere delamination in orogens: *Earth and Planetary Science Letters*, v. 361, p. 195–207, <https://doi.org/10.1016/j.epsl.2012.09.056>.
- Lowry, A.R., and Pérez-Gussinyé, M., 2011, The role of crustal quartz in controlling Cordilleran deformation: *Nature*, v. 471, p. 353, <https://doi.org/10.1038/nature09912>.
- Naimark, B.M., and Ismail-Zadeh, A.T., 1995, Numerical models of a subsidence mechanism in intracratonic basins: applications to North American basins: *Geophysical Journal International*, v. 123, p. 149–160, <https://doi.org/10.1111/j.1365-246X.1995.tb06667.x>.
- O'Connell, R.J., and Wasserburg, G.J., 1972, Dynamics of submergence and uplift of a sedimentary basin underlain by a phase-change boundary: *Reviews of Geophysics*, v. 10, p. 335–368, <https://doi.org/10.1029/RG010i001p00335>.
- Reidel, S.P., Tolan, T.L., Hooper, P.R., Beeson, M.H., Focht, K.R., Bentley, R.D., and Anderson, J.L., 1989, The Grande Ronde Basalt, Columbia River Basalt Group: Stratigraphic descriptions and correlations in Washington, Oregon, and Idaho, in Reidel, S.P., and Hooper, P.R., eds., *Volcanism and Tectonism in the Columbia River Flood-basalt Province*: Geological Society of America Special Papers, v. 239, p. 21–54, <https://doi.org/10.1130/SPE239-p21>.
- Reidel, S.P., Camp, V.E., Tolan, T.L., and Martin, B.S., 2013, The Columbia River flood basalt province: Stratigraphy, areal extent, volume, and physical volcanology, in Reidel, S.P., et al., eds., *The Columbia River Flood Basalt Province*: Geological Society of America Special Papers, v. 497, p. 1–43, doi: 10.1130/2013.2497(01).
- Ryan, W.B., Carbotte, S.M., Coplan, J.O., O'Hara, S., Melkonian, A., Arko, R., Weissel, R.A., Ferrini, V., Goodwillie, A., Nitsche, F., et al., 2009, Global multi-resolution topography synthesis: *Geochemistry Geophysics Geosystems*, v. 10, Q03014, <https://doi.org/10.1029/2008GC002332>.
- Taylor, S.B., Johnson, S.Y., Fraser, G.T., and Roberts, J.W., 1988, Sedimentation and tectonics of the lower and middle Eocene Swauk Formation in eastern Swauk Basin, central Cascades, central Washington: *Canadian Journal of Earth Sciences*, v. 25, p. 1020–1036, <https://doi.org/10.1139/e88-100>.
- Turcotte, D., and Schubert, G., 2014, *Geodynamics*: Cambridge, UK, Cambridge University Press, 636 p., <https://doi.org/10.1017/CBO9780511843877>.
- Wang, H., and Currie, C.A., 2017, Crustal deformation induced by mantle dynamics: Insights from models of gravitational lithosphere removal: *Geophysical Journal International*, v. 210, p. 1070–1091, <https://doi.org/10.1093/gji/ggx209>.

Manuscript received 28 March 2018

Revised manuscript received 31 May 2018

Manuscript accepted 4 June 2018

Printed in USA