

May 13, 2020

Holographic Line Defects in $D = 4$, $N = 2$ Gauged Supergravity

Kevin Chen, Michael Gutperle, and Matteo Vicino

*Mani L. Bhaumik Institute for Theoretical Physics
Department of Physics and Astronomy
University of California, Los Angeles, CA 90095, USA*

Abstract

We construct solutions of four-dimensional $N = 2$ gauged supergravity coupled to vector multiplets which are holographically dual to superconformal line defects. For the gauged STU and the $SU(1, n)/U(1) \times SU(n)$ coset models, we use the solutions to calculate holographic observables such as the expectation value of the defect and one-point functions in the presence of the defect.

1 Introduction

Extended objects such as defects, line operators, and interfaces are important tools in the study of quantum field theories. For conformal field theories (CFTs) with holographic duals, in many cases the extended objects have a realization on the gravity side. One example of such a duality is the fundamental string in $\text{AdS}_5 \times S^5$ describing a Wilson line in the fundamental representation [1, 2], and its generalization to D3 probe branes with an $\text{AdS}_2 \times S^2$ worldvolume and D5 probe branes with an $\text{AdS}_2 \times S^4$ worldvolume [3, 4, 5, 6, 7]. When the number of probe branes becomes large, the backreaction can not be neglected and a fully backreacted type IIB supergravity solution replaces the probe description. For the case of the half-BPS Wilson loop, this solution was found in [8].¹ In general, the fully backreacted solutions in type II and M-theory are warped product geometries and the solutions are complicated and difficult to obtain.² A simpler setting is to consider lower-dimensional gauged supergravities, where the ansatz is simpler and the number of fields is smaller. If the gauged supergravity is a consistent truncation of a higher-dimensional theory, the resulting lower-dimensional solution can be uplifted to produce solutions in the higher-dimensional theory.

In this paper, we consider four-dimensional $N = 2$ gauged supergravity, which has been used in the past to describe condensed matter systems in three dimensions in order to find holographic models for superfluids and superconductors, see e.g. [12, 13, 14]. We construct half-BPS supergravity solutions which are dual to line defects in three-dimensional $\mathcal{N} = 2$ superconformal field theories. The metric ansatz is given by $\text{AdS}_2 \times S^1$ warped over an interval. We generalize the analysis of [15], which considered pure gauged supergravity, to the case of matter couplings.

The structure of the paper is as follows. In section 2, we review our conventions for four-dimensional $N = 2$ gauged supergravity coupled to vector multiplets. In section 3, we give a general solution describing a half-BPS line defect, obtained by a double analytic continuation of the black hole solutions first found by Sabra [16]. Since the behavior of the vector multiplet scalars can only be determined implicitly, we consider three examples, namely a single scalar model, the gauged STU model, and the $\text{SU}(1, n)$ coset model to obtain explicit solutions. In section 4, we use the machinery of holographic renormalization to calculate holographic observables for the solutions, namely the on-shell action and the expectation values of operators dual to the supergravity fields. In section 5, we explore the conditions for a regular geometry and calculate their consequences. In section 6, we discuss our results and possible directions for future research. Our conventions and some details of the calculations presented in the main body of the paper are relegated to several appendices.

¹See [9] for earlier work on IIB supergravity solutions for Wilson loops.

²See [10, 11] for gravity solutions dual to conformal defects in both type IIB and D=11 supergravity.

2 $D = 4$, $N = 2$ Gauged Supergravity

In this section, we review four-dimensional $N = 2$ gauged supergravity coupled to n vector multiplets. We use the conventions and notations of [17, 18, 19].

The field content of the gauged supergravity theory is as follows. The supergravity contains one graviton e_μ^a , two gravitinos ψ_μ^i , and one graviphoton A_μ^0 . The gravity multiplet can be coupled to $N = 2$ matter, and in particular we consider n vector multiplets, which are labeled by an index $\alpha = 1, 2, \dots, n$. Each vector multiplet contains one vector field A_μ^α , two gauginos λ_i^α , and one complex scalar τ^α . In this paper we do not consider adding $N = 2$ hypermultiplets.

It is convenient to introduce a new index $I = 0, 1, \dots, n$ and include the graviphoton with the other vector fields as A_μ^I . The complex scalars τ^α parametrize a special Kähler manifold equipped with a holomorphic symplectic vector

$$v(\tau) = \begin{pmatrix} Z^I(\tau) \\ \mathcal{F}_I(\tau) \end{pmatrix}, \quad (2.1)$$

where the Kähler potential $\mathcal{K}(\tau, \bar{\tau})$ is determined by

$$e^{-\mathcal{K}(\tau, \bar{\tau})} = -i \langle v, \bar{v} \rangle \equiv -i(Z^I \bar{\mathcal{F}}_I - \mathcal{F}_I \bar{Z}^I). \quad (2.2)$$

In the models we will consider, there exists a holomorphic function $\mathcal{F}(Z)$, called the prepotential, that is homogeneous of second order in Z such that

$$\mathcal{F}_I(\tau) = \frac{\partial}{\partial Z^I} \mathcal{F}(Z(\tau)). \quad (2.3)$$

The supergravity theory is fully specified by the prepotential $\mathcal{F}(Z)$ and the choice of gauging of the $SU(2)$ R -symmetry. We will choose the $U(1)$ Fayet-Iliopoulos (FI) gauging. The only charged fields of the theory are the gravitinos, which couple to the gauge fields through the linear combination $\xi_I A^I$, for some real constants ξ_I . The two gravitinos have opposite charges $\pm g \xi_I$ for each $U(1)$ gauge factor, where g is the gauge coupling.

The bosonic action is³

$$\begin{aligned} e^{-1} \mathcal{L}_{\text{bos}} = & \frac{1}{2} R - g_{\alpha\bar{\beta}} \partial^\mu \tau^\alpha \partial_\mu \bar{\tau}^{\bar{\beta}} - V(\tau, \bar{\tau}) \\ & + \frac{1}{4} (\text{Im} \mathcal{N})_{IJ} F^{I\mu\nu} F_{\mu\nu}^J - \frac{1}{8} (\text{Re} \mathcal{N})_{IJ} e^{-1} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^I F_{\rho\sigma}^J, \end{aligned} \quad (2.4)$$

where $F_{\mu\nu}^I = \partial_\mu A_\nu^I - \partial_\nu A_\mu^I$ are the field strengths and $g_{\alpha\bar{\beta}} = \partial_\alpha \partial_{\bar{\beta}} \mathcal{K}$ is the Kähler metric of the scalar manifold. We use $G_{\mu\nu}$ to denote the four-dimensional metric, so $e = \sqrt{-\det G}$. The scalar potential is

$$V(\tau, \bar{\tau}) = -2g^2 \xi_I \xi_J ((\text{Im} \mathcal{N})^{-1})^{IJ} + 8e^{\mathcal{K}} \bar{Z}^I Z^J, \quad (2.5)$$

³We set $8\pi G_N = 1$.

where the kinetic matrix $\mathcal{N}_{IJ}$ is given by

$$\mathcal{N}_{IJ}(\tau, \bar{\tau}) = \bar{\mathcal{F}}_{IJ} + 2i \frac{(\text{Im } \mathcal{F}_{IL})(\text{Im } \mathcal{F}_{JK})Z^L Z^K}{(\text{Im } \mathcal{F}_{MN})Z^M Z^N}, \quad \mathcal{F}_{IJ} \equiv \frac{\partial}{\partial Z^I} \frac{\partial}{\partial Z^J} \mathcal{F}(Z). \quad (2.6)$$

This is equivalently defined as the matrix which solves the equations

$$\mathcal{F}_I = \mathcal{N}_{IJ} Z^J, \quad \mathcal{D}_{\bar{\alpha}} \bar{\mathcal{F}}_I = \mathcal{N}_{IJ} \mathcal{D}_{\bar{\alpha}} \bar{Z}^J, \quad (2.7)$$

where $\mathcal{D}$ is the Kähler covariant derivative,

$$\begin{aligned} \mathcal{D}_{\alpha} v &= (\partial_{\alpha} + \partial_{\alpha} \mathcal{K}) v, \\ \mathcal{D}_{\bar{\alpha}} \bar{v} &= (\partial_{\bar{\alpha}} + \partial_{\bar{\alpha}} \mathcal{K}) \bar{v}, \\ \mathcal{D}_{\alpha} \bar{v} &= \partial_{\alpha} \bar{v} = 0, \\ \mathcal{D}_{\bar{\alpha}} v &= \partial_{\bar{\alpha}} v = 0. \end{aligned} \quad (2.8)$$

The equations of motion are obtained by varying the Lagrangian (2.4)

$$\begin{aligned} R_{\mu\nu} &= 2g_{\alpha\bar{\beta}} \partial_{\mu} \tau^{\alpha} \partial_{\nu} \bar{\tau}^{\bar{\beta}} + V G_{\mu\nu} + (\text{Im } \mathcal{N})_{IJ} \left(-F_{\mu}^I{}^{\rho} F_{\nu\rho}^J + \frac{1}{4} F^{I\rho\sigma} F_{\rho\sigma}^J G_{\mu\nu} \right), \\ \partial_{\mu} \left(e g_{\alpha\bar{\beta}} \partial^{\mu} \bar{\tau}^{\bar{\beta}} \right) &= e \left((\partial_{\alpha} g_{\beta\bar{\gamma}}) \partial^{\mu} \tau^{\beta} \partial_{\mu} \bar{\tau}^{\bar{\gamma}} - \frac{1}{4} \partial_{\alpha} (\text{Im } \mathcal{N})_{IJ} F^{I\mu\nu} F_{\mu\nu}^J + \partial_{\alpha} V \right) \\ &\quad + \frac{1}{8} \partial_{\alpha} (\text{Re } \mathcal{N})_{IJ} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^I F_{\rho\sigma}^J, \\ 0 &= \partial_{\mu} \left(e (\text{Im } \mathcal{N})_{IJ} F^{J\mu\nu} - \frac{1}{2} (\text{Re } \mathcal{N})_{IJ} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}^J \right). \end{aligned} \quad (2.9)$$

The supersymmetry transformations are given in appendix A.

3 Line Defect Solution

In this section, we give a general solution describing a half-BPS line defect in four-dimensional $N = 2$ gauged supergravity, and then construct the solution for three specific choices of the prepotential.

3.1 Holographic Line Defects

A conformal line defect in three dimensions is a codimension-two defect which breaks the three-dimensional conformal group $\text{SO}(3, 2)$ down to an $\text{SO}(2, 1) \times \text{SO}(2)$ subgroup. The subgroup factors represent the unbroken conformal symmetry along the defect and transverse rotations about the defect, respectively. Minkowski space $\mathbb{R}^{1,2}$ is related by a Weyl transformation to $\text{AdS}_2 \times S^1$, namely

$$-dt^2 + dr^2 + r^2 d\phi^2 = \Omega(r) \left(\frac{-dt^2 + dr^2}{r^2} + d\phi^2 \right). \quad (3.1)$$

Hence in the holographic dual, the $\text{SO}(2, 1) \times \text{SO}(2)$ symmetry can be realized as the isometries of $\text{AdS}_2 \times S^1$, which we choose as the boundary of the four-dimensional asymptotically anti-de Sitter space. Therefore we consider a metric ansatz with $\text{AdS}_2 \times S^1$ warped over a radial coordinate. We note that the location of the defect at $r = 0$ in Minkowski space gets mapped to the boundary of AdS_2 in the $\text{AdS}_2 \times S^1$ geometry. Secondly, the absence of a conical singularity on the boundary fixes the periodicity of the angle ϕ to be 2π .

The superconformal algebras in three dimensions are $\text{OSp}(\mathcal{N}|4)$, where $\mathcal{N} = 1, 2, \dots, 6, 8$. For the CFT dual of four-dimensional $N = 2$ gauged supergravity, the relevant superalgebra is $\text{OSp}(2|4)$ which has four Poincaré and four conformal supercharges. A conformal line defect is called superconformal if it preserves some supersymmetry. In the present paper, we will consider half-BPS defects which preserve an $\text{OSp}(2|2)$ superalgebra and hence four of the eight supersymmetries.

3.2 General Solution

Four-dimensional $N = 2$, $\text{U}(1)$ FI gauged supergravity admits half-BPS black hole solutions first found in [16]. The line defect solutions with $\text{AdS}_2 \times S^1$ geometry are constructed by a double analytic continuation of the black hole solution. The metric and gauge fields are given by

$$\begin{aligned} ds^2 &= r^2 \sqrt{H(r)} ds_{\text{AdS}_2}^2 + \frac{f(r)}{\sqrt{H(r)}} ds_{S^1}^2 + \frac{\sqrt{H(r)}}{f(r)} dr^2, \\ f(r) &= -1 + 8g^2 r^2 H(r), \\ H(r)^{1/4} &= \frac{1}{\sqrt{2}} e^{\mathcal{K}/2} Z^I H_I(r), \\ H_I(r) &= \xi_I + \frac{q_I}{r}, \quad I = 0, 1, \dots, n, \\ A^I &= (-2H(r)^{-1/4} e^{\mathcal{K}/2} Z^I + \mu^I) d\theta, \quad I = 0, 1, \dots, n, \end{aligned} \quad (3.2)$$

for some real constants q_I and μ_I , where $Z^I = \bar{Z}^I$. Given a prepotential $\mathcal{F}(Z)$ and choice of parametrization of the symplectic sections $Z^I(\tau)$, the scalars τ^α are given implicitly by the equation

$$iH^{1/4} e^{\mathcal{K}/2} (\mathcal{F}_I - \bar{\mathcal{F}}_I) = \frac{1}{\sqrt{2}} H_I. \quad (3.3)$$

At the conformal boundary where $r \rightarrow \infty$, in order to have asymptotic AdS_4 we need $2\sqrt{2}g\theta$ to be 2π -periodic, i.e. $\theta \sim \theta + \pi/\sqrt{2}g$. The AdS_4 length scale is then given by

$$L^{-2} = 8g^2 H(r = \infty)^{1/2}. \quad (3.4)$$

We will set $8g^2 = 1$ to obtain the usual S^1 periodicity $\theta \sim \theta + 2\pi$.

The center of the space⁴ $r = r_+$ corresponds to the largest value of r where $f(r) = 0$. We consider radii taking values in the range $r \in [r_+, \infty)$. Demanding a regular geometry also requires $r_+ > 0$ and the absence of a conical singularity at the center of the space, both of which can be done by tuning the q_I and ξ_I parameters. This is explored in further detail in section 5.

3.3 Examples

For a general prepotential, the equation (3.3) is very complicated and can only be solved numerically. Consequently, we will explicitly work out the line defect solution for three specific prepotentials, for which we can find explicit expressions for the scalars. An important requirement is the existence of an AdS_4 vacuum, which not all prepotentials admit, see e.g. [19, 20].

3.3.1 Single Scalar Model

Consider a single ($n = 1$) vector multiplet with the prepotential $\mathcal{F}(Z) = -iZ^0Z^1$. This theory has a single complex scalar τ and the scalar manifold is $\frac{\text{SU}(1,1)}{\text{U}(1)}$. Using the parametrization $(Z^0, Z^1) = (1, \tau)$, we can calculate the Kähler potential, kinetic matrix, and scalar potential,

$$\begin{aligned} e^{\mathcal{K}(\tau, \bar{\tau})} &= \frac{1}{2(\tau + \bar{\tau})} , \\ \mathcal{N}(\tau, \bar{\tau}) &= -i \begin{pmatrix} \tau & 0 \\ 0 & 1/\tau \end{pmatrix} , \\ V(\tau, \bar{\tau}) &= -\frac{1}{2(\tau + \bar{\tau})} (\xi_0^2 + 2\xi_0\xi_1(\tau + \bar{\tau}) + \xi_1^2\tau\bar{\tau}) . \end{aligned} \quad (3.5)$$

The potential has extrema at $\tau = \pm\xi_0/\xi_1$, but only $\tau = \xi_0/\xi_1$ maintains $e^{\mathcal{K}} > 0$ for $\xi_I > 0$. The cosmological constant at this extremum gives the AdS_4 length scale

$$L^{-2} = \frac{1}{2}\xi_0\xi_1. \quad (3.6)$$

⁴For the black hole geometry this is the location of the horizon.

We choose $\xi_1 = 2/\xi_0$ to set the AdS_4 length scale to unity. The line defect solution (3.2) has the explicit form,

$$\begin{aligned}
ds^2 &= r^2 \sqrt{H} ds_{\text{AdS}_2}^2 + \frac{f}{\sqrt{H}} ds_{S^1}^2 + \frac{\sqrt{H}}{f} dr^2 , \\
f(r) &= -1 + r^2 H(r) , \\
\sqrt{H(r)} &= \frac{1}{2} H_0 H_1 , \\
H_I(r) &= \xi_I + \frac{q_I}{r} , \quad I = 0, 1 , \\
A^I &= \left(-\frac{\sqrt{2}}{H_I} + \mu^I \right) d\theta , \quad I = 0, 1 .
\end{aligned} \tag{3.7}$$

The scalar is given by

$$\tau = \frac{H_0}{H_1} . \tag{3.8}$$

We have verified that the above fields obey the equations of motion (2.9).

3.3.2 Gauged STU Model

The STU model is given by considering $n = 3$ vector multiplets with the prepotential

$$\mathcal{F}(Z) = -2i\sqrt{Z^0 Z^1 Z^2 Z^3} . \tag{3.9}$$

This theory has three complex scalars τ^1, τ^2, τ^3 and the scalar manifold is three copies of $\frac{\text{SU}(1,1)}{\text{U}(1)}$. When all $\xi_I = \xi > 0$ are equal, this theory is a consistent truncation of $\mathcal{N} = 8$ gauged supergravity [21, 22]. For reference on this model, see [23]. Using the parametrization $(Z^0, Z^1, Z^2, Z^3) = (1, \tau^2 \tau^3, \tau^1 \tau^3, \tau^1 \tau^2)$, the Kähler potential is

$$e^{\mathcal{K}(\tau, \bar{\tau})} = \frac{1}{(\tau^1 + \bar{\tau}^1)(\tau^2 + \bar{\tau}^2)(\tau^3 + \bar{\tau}^3)} . \tag{3.10}$$

The expressions for the kinetic matrix and scalar potential are complicated, but simplify for real scalars $\tau^\alpha = \bar{\tau}^\alpha$, which will be the case for the line defect solution.

$$\begin{aligned}
\mathcal{N}(\tau, \bar{\tau} = \tau) &= -i \text{diag} \left(\tau^1 \tau^2 \tau^3, \frac{\tau^1}{\tau^2 \tau^3}, \frac{\tau^2}{\tau^1 \tau^3}, \frac{\tau^3}{\tau^1 \tau^2} \right) , \\
V(\tau, \bar{\tau} = \tau) &= -\frac{1}{2} \left(\xi_0 \left(\frac{\xi_1}{\tau^1} + \frac{\xi_2}{\tau^2} + \frac{\xi_3}{\tau^3} \right) + (\tau^1 \xi_2 \xi_3 + \xi_1 \tau^2 \xi_3 + \xi_1 \xi_2 \tau^3) \right) .
\end{aligned} \tag{3.11}$$

The potential has extrema at

$$\tau^1 = \pm \sqrt{\frac{\xi_0 \xi_1}{\xi_2 \xi_3}} , \quad \tau^2 = \pm \sqrt{\frac{\xi_0 \xi_2}{\xi_1 \xi_3}} , \quad \tau^3 = \pm \sqrt{\frac{\xi_0 \xi_3}{\xi_1 \xi_2}} . \tag{3.12}$$

Positivity of $e^{\mathcal{K}}$ requires us to choose the positive root. The cosmological constant at this extremum gives the AdS₄ length scale

$$L^{-2} = \sqrt{\xi_0 \xi_1 \xi_2 \xi_3} . \quad (3.13)$$

We pick the non-zero constants ξ_I in a way that sets the AdS₄ length scale to unity. The line defect solution (3.2) has the explicit form,

$$\begin{aligned} ds^2 &= r^2 \sqrt{H} ds_{\text{AdS}_2}^2 + \frac{f}{\sqrt{H}} ds_{S^1}^2 + \frac{\sqrt{H}}{f} dr^2 , \\ f(r) &= -1 + r^2 H(r) , \\ H(r) &= H_0 H_1 H_2 H_3 , \\ H_I(r) &= \xi_I + \frac{q_I}{r} , \quad I = 0, 1, 2, 3 , \\ A^I &= \left(-\frac{1}{\sqrt{2} H_I} + \mu^I \right) d\theta , \quad I = 0, 1, 2, 3 . \end{aligned} \quad (3.14)$$

The scalars are

$$\tau^1 = \sqrt{\frac{H_0 H_1}{H_2 H_3}} , \quad \tau^2 = \sqrt{\frac{H_0 H_2}{H_1 H_3}} , \quad \tau^3 = \sqrt{\frac{H_0 H_3}{H_1 H_2}} . \quad (3.15)$$

This solution is also the double analytic continuation of the hyperbolic black hole solution in [24]. As consistency checks, we have verified that the above solution obeys the equations of motion (2.9) and is half-BPS. The latter was done by a direct calculation, independent of [16], which can be found in appendix A.

3.3.3 SU(1, n) Coset Model

Another model which admits an AdS₄ vacuum has the prepotential $\mathcal{F}(Z) = \frac{i}{4} Z^I \eta_{IJ} Z^J$, and can be formulated with any number of vector multiplets. η_{IJ} is a Minkowski metric, which we will take to be $\eta = \text{diag}(-1, +1, \dots, +1)$. The scalar manifold of this theory is $\frac{\text{SU}(1, n)}{\text{U}(1) \times \text{SU}(n)}$. Using the parametrization $(Z^0, Z^\alpha) = (1, \tau^\alpha)$, the Kähler potential is

$$e^{\mathcal{K}(\tau, \bar{\tau})} = \frac{1}{1 - \sum_\alpha \tau^\alpha \bar{\tau}^\alpha} . \quad (3.16)$$

Once again, the kinetic matrix and scalar potential have simpler forms for real scalars $\tau^\alpha = \bar{\tau}^\alpha$. The matrix η_{IJ} is used to lower indices, e.g. $Z_I = \eta_{IJ} Z^J$.

$$\begin{aligned} \mathcal{N}_{IJ}(\tau, \bar{\tau} = \tau) &= -\frac{i}{2} \eta_{IJ} - i e^{\mathcal{K}(\tau, \tau)} Z_I Z_J , \\ V(\tau, \bar{\tau} = \tau) &= \frac{1}{2} \xi_I \eta^{IJ} \xi_J - \frac{(\xi_0 + \sum_\alpha \xi_\alpha \tau^\alpha)^2}{1 - \sum_\alpha (\tau^\alpha)^2} . \end{aligned} \quad (3.17)$$

This potential has an extremum at $\tau^\alpha = -\xi_\alpha/\xi_0$.⁵ The cosmological constant at this extremum gives us the AdS₄ length scale

$$L^{-2} = -\xi^2/2 , \quad (3.18)$$

where $\xi^2 = \xi_I \eta^{IJ} \xi_J$. We pick a time-like ξ_I with $\xi^2 = -2$ that will set the AdS₄ length scale to unity. The line defect solution (3.2) has the explicit form,

$$\begin{aligned} ds^2 &= r^2 \sqrt{H} ds_{\text{AdS}_2}^2 + \frac{f}{\sqrt{H}} ds_{S^1}^2 + \frac{\sqrt{H}}{f} dr^2 , \\ f(r) &= -1 + r^2 H(r) , \\ \sqrt{H(r)} &= -\frac{1}{2} H_I \eta^{IJ} H_J , \\ H_I(r) &= \xi_I + \frac{q_I}{r} , \quad I = 0, 1, \dots, n , \\ A^I &= \left(\frac{\sqrt{2} \eta^{IJ} H_J}{\sqrt{H}} + \mu^I \right) d\theta , \quad I = 0, 1, \dots, n . \end{aligned} \quad (3.19)$$

The scalars are

$$\tau^\alpha = -\frac{H_\alpha}{H_0} . \quad (3.20)$$

We have verified that the above fields obey the equations of motion (2.9).

4 Holographic Calculations

In this section, we use the machinery of holographic renormalization [25, 26] to calculate the on-shell action and the one-point functions of dual operators of the boundary CFT in the presence of the defect, namely the stress tensor, scalar, and currents. This is done explicitly for the three examples in section 3.3.

4.1 General Procedure

First, we put the metric into the Fefferman-Graham (FG) form,

$$ds^2 = \frac{1}{z^2} (dz^2 + g_{ij}(x, z) dx^i dx^j) , \quad (4.1)$$

where $i, j = 1, 2, 3$ run over the AdS₂ and S^1 indices and $z \rightarrow 0$ is the conformal boundary. This is done by taking $z = z(r)$ so that the appropriate coordinate change is obtained by the solution to the ordinary differential equation

$$-\frac{H(r)^{1/4}}{f(r)^{1/2}} dr = \frac{dz}{z} , \quad (4.2)$$

⁵The other extrema at $\xi_0 + \sum_\alpha \xi_\alpha \tau^\alpha = 0$ do not admit AdS₄ vacua while maintaining $e^{\mathcal{K}}$ positive.

which can be integrated perturbatively in $1/r$. This coordinate change gives the FG expansions of the fields, which we assume will take the form

$$\begin{aligned} g_{ij} &= g_{0ij} + z^2 g_{2ij} + z^3 g_{3ij} + \mathcal{O}(z^4) , \\ A^I &= A_0^I + z A_1^I + \mathcal{O}(z^2) , \\ \tau^\alpha &= \tau_0^\alpha + z \tau_1^\alpha + z^2 \tau_2^\alpha + \mathcal{O}(z^3) , \\ \bar{\tau}^{\bar{\alpha}} &= \tau_0^\alpha + z \tau_1^\alpha + z^2 \tau_2^\alpha + \mathcal{O}(z^3) , \end{aligned} \tag{4.3}$$

where A_0^I and A_1^I are 1-forms on the x^1, x^2, x^3 coordinates. The constants τ_0^α are the AdS₄ vacuum values of the scalars, which depend on the model. There is no gravitational conformal anomaly (i.e. a term proportional to $z^3 \log z$ in the expansion of g_{ij}) since $d = 3$ is odd.

In the three-dimensional boundary CFT, the conformal dimensions of the dual operators corresponding to the scalars τ^α and vector fields A^I are determined by the linearized bulk equations of motion near the AdS boundary. For instance, using the expansion $\tau^\alpha \sim \tau_0^\alpha + z^{\Delta_\tau}$ in the linearized equation of motion for the scalar, we find that the scaling dimension of the dual operator is related to the mass-squared of the field by the equation

$$\Delta_\tau(\Delta_\tau - 3) = -2 . \tag{4.4}$$

The mass-squared is -2 for all scalars of the three examples considered in this paper. This mass-squared is within the window where both standard and alternative quantization are possible [27], which implies that the scaling dimension of the dual operator can be either $\Delta_\tau = 1$ or $\Delta_\tau = 2$. Similarly, using the expansion $A^I \sim z^{\Delta_A - 1} d\theta$ in the linearized equation of motion for the vector field gives us

$$(\Delta_A - 1)(\Delta_A - 2) = 0 . \tag{4.5}$$

We must have $\Delta_A = 2$ as the vector field sources a conserved current of the boundary CFT.

These scaling dimensions naturally fit into the flavor current multiplet $A_2 \bar{A}_2 [0]_1^{(0)}$ of the $d = 3$, $\mathcal{N} = 2$ boundary CFT, using the notation of [28]. This short multiplet contains, in addition to the spin-1 operator $[2]_2^{(0)}$ with scaling dimension $\Delta = 2$, two scalar operators $[0]_1^{(0)}$ and $[0]_2^{(0)}$ as bottom and top components with scaling dimensions $\Delta = 1$ and 2 respectively. The stress tensor multiplet $A_1 \bar{A}_1 [2]_2^{(0)}$ is also present, as usual.

In the four-dimensional gauged supergravity, for a well-defined variational principle of the metric we need to add to the bulk action given by the Lagrangian (2.4) the Gibbons-Hawking boundary term,

$$\begin{aligned} I_{\text{bulk}} &= \int_M d^4x \mathcal{L}_{\text{bos}} , \\ I_{\text{GH}} &= \int_{\partial M} d^3x \sqrt{-h} \text{Tr}(h^{-1} K) , \end{aligned} \tag{4.6}$$

where h_{ij} is the induced metric on the boundary and K_{ij} is the extrinsic curvature. In FG coordinates, these take the form,

$$h_{ij} = \frac{1}{z^2} g_{ij} , \quad K_{ij} = -\frac{z}{2} \partial_z h_{ij} . \quad (4.7)$$

The action $I_{\text{bulk}} + I_{\text{GH}}$ diverges due to the infinite volume of integration. To regulate the theory, we restrict the bulk integral to the region $z \geq \epsilon$ and evaluate the boundary term at $z = \epsilon$. Divergences in the action then appear as $1/\epsilon^k$ poles.⁶ Counterterms I_{ct} are added on the boundary which subtract these divergent terms. The counterterms have been constructed in [23] and are compatible with supersymmetry. They are

$$I_{\text{ct}} = \int_{\partial M} d^3x \sqrt{-h} \left(\mathcal{W} - \frac{1}{2} R[h] \right) , \quad \mathcal{W} \equiv -\sqrt{2} e^{\mathcal{K}/2} |\xi_I Z^I| , \quad (4.8)$$

where $R[h]$ is the Ricci scalar of the boundary metric and $\mathcal{W}$ is the superpotential. In all, the renormalized action,

$$I_{\text{ren}} = I_{\text{bulk}} + I_{\text{GH}} + I_{\text{ct}} , \quad (4.9)$$

is finite. We can then take functional derivatives to obtain finite expectation values of the dual CFT operators. Let T_{ij} be the boundary stress tensor, $\mathcal{O}_\alpha$ be the operators dual to τ^α , and J_{I_i} be the current operators dual to A_μ^I .

4.1.1 Stress Tensor Expectation Value

The expectation value of the boundary stress tensor is defined to be [29]

$$\langle T_{ij} \rangle \equiv \frac{-2}{\sqrt{-g_0}} \frac{\delta I_{\text{ren}}}{\delta g_0^{ij}} . \quad (4.10)$$

The variation decomposes into two contributions: one coming from the regularized action and one coming from the counterterms. As usual [30], the former is given by

$$T_{ij}^{\text{reg}}[h] \equiv \frac{-2}{\sqrt{-h}} \frac{\delta(I_{\text{bulk}} + I_{\text{GH}})}{\delta h^{ij}} = -K_{ij} + h_{ij} \text{Tr}(h^{-1} K) . \quad (4.11)$$

The latter is straightforward to compute, and is given by

$$T_{ij}^{\text{ct}}[h] \equiv \frac{-2}{\sqrt{-h}} \frac{\delta I_{\text{ct}}}{\delta h^{ij}} = h_{ij} \left(\mathcal{W} - \frac{1}{2} R[h] \right) + R_{ij}[h] . \quad (4.12)$$

Therefore,

$$\langle T_{ij} \rangle = \lim_{\epsilon \rightarrow 0} \left[\epsilon^{-1} \left(T_{ij}^{\text{reg}}[h] + T_{ij}^{\text{ct}}[h] \right) \right]_{z=\epsilon} . \quad (4.13)$$

By construction of the counterterms, this limit exists.

⁶In even boundary dimensions, a term proportional to $\log \epsilon$ may also appear.

4.1.2 Scalar Expectation Values

The expectation value of the operator $\mathcal{O}_\alpha$ is similarly defined by

$$\langle \mathcal{O}_\alpha \rangle \equiv \frac{1}{\sqrt{-g_0}} \frac{\delta I_{\text{ren}}}{\delta \tau_1^\alpha} = \lim_{\epsilon \rightarrow 0} \left[\epsilon^{-2} \frac{1}{\sqrt{-h}} \frac{\delta I_{\text{ren}}}{\delta \tau^\alpha} \Big|_{z=\epsilon} \right]. \quad (4.14)$$

The variation has contributions from the bulk action and the counterterms, and is

$$\frac{1}{\sqrt{-h}} \frac{\delta I_{\text{ren}}}{\delta \tau^\alpha} = g_{\alpha\bar{\beta}} z \partial_z \bar{\tau}^{\bar{\beta}} + \partial_\alpha \mathcal{W}. \quad (4.15)$$

For real scalars, supersymmetry implies $\langle \mathcal{O}_\alpha \rangle = 0$. A proof of this statement can be found in appendix B.

4.1.3 Current Expectation Values

The expectation value of the current operator J_I is defined by

$$\langle J_I^i \rangle \equiv \frac{1}{\sqrt{-g_0}} \frac{\delta I_{\text{ren}}}{\delta A_{0i}^I} = \lim_{\epsilon \rightarrow 0} \left[\epsilon^{-3} \frac{1}{\sqrt{-h}} \frac{\delta I_{\text{ren}}}{\delta A_i^I} \Big|_{z=\epsilon} \right]. \quad (4.16)$$

The only contribution to the variation comes from the bulk action, and is

$$\frac{1}{\sqrt{-h}} \frac{\delta I_{\text{ren}}}{\delta A_i^I} = -(\text{Im } \mathcal{N})_{IJ} h^{ij} z \partial_z A_j^J. \quad (4.17)$$

4.1.4 On-Shell Action

We can evaluate the on-shell action for the line defect solution by further simplifying the bulk action to a total derivative [31],

$$I_{\text{bulk}} \Big|_{\text{on-shell}} = \text{Vol}(\text{AdS}_2) \text{Vol}(S^1) \left[-\frac{H'(r)}{4H(r)} r^2 f(r) - r(f(r) + 1) \right] \Big|_{r_+}^{\infty}, \quad (4.18)$$

where $\text{Vol}(S^1) = 2\pi$ and $\text{Vol}(\text{AdS}_2) = -2\pi$ is the regularized volume of AdS_2 .

4.2 Examples

In this section, we use the general expressions derived in section 4.1 to compute observables for the three examples considered in this paper.

4.2.1 Single Scalar Model

Let us consider the defect solution (3.7, 3.8) for the single scalar model. The FG expansion of the radial coordinate r from solving the ordinary differential equation (4.2) is

$$\begin{aligned} \frac{1}{r} = z + \frac{1}{2} \left(\sum_{I=0}^1 \frac{q_I}{\xi_I} \right) z^2 + \frac{-16 + (3q_1\xi_0 + q_0\xi_1)(3q_0\xi_1 + q_1\xi_0)}{64} z^3 \\ + \frac{(q_1\xi_0 + q_0\xi_1)(-16 + 12q_0q_1\xi_0\xi_1 + 3(q_0\xi_1 + q_1\xi_0)^2)}{384} z^4 + \mathcal{O}(z^5). \end{aligned} \quad (4.19)$$

Using this coordinate change, the metric, gauge fields, and scalar can be expanded in FG coordinates. The one-point functions in the presence of the line defect can then be evaluated by computing the limits (4.13, 4.14, 4.16) directly. For the renormalized on-shell action (4.9), the finite terms at the conformal boundary cancel, leaving just the term obtained by evaluating (4.18) at $r = r_+$. In the end, we obtain the following expectation values:

$$\begin{aligned}
I_{\text{ren}} &= \text{Vol}(\text{AdS}_2)\text{Vol}(S^1)r_+ , \\
\langle T_{ij} \rangle &= \frac{1}{2} \left(\sum_{I=0}^1 \frac{q_I}{\xi_I} \right) \begin{pmatrix} -g_{\text{AdS}_2} & 0 \\ 0 & 2g_{S^1} \end{pmatrix}_{ij} , \\
\langle T_i^i \rangle &= 0 , \\
\langle \mathcal{O} \rangle &= 0 , \\
\langle J_{Ii} \rangle &= \frac{q_I}{\sqrt{2}} \delta_{i\theta} .
\end{aligned} \tag{4.20}$$

4.2.2 Gauged STU Model

Let us consider the defect solution (3.14, 3.15) for the gauged STU model. Some of the calculations for this model are identical to those found in [24]. The FG expansion of the radial coordinate r from solving the ODE (4.2) is

$$\frac{1}{r} = z + \frac{A}{4} z^2 + \frac{-16 + B_1 + 10B_2}{64} z^3 + \frac{-16A + C_1 + 11C_2 + 62C_3}{384} z^4 + \mathcal{O}(z^5) , \tag{4.21}$$

where we have defined the constants

$$\begin{aligned}
A &= \sum_{I=0}^3 \frac{q_I}{\xi_I} , & B_1 &= \sum_{I=0}^3 \left(\frac{q_I}{\xi_I} \right)^2 , & B_2 &= \sum_{I<J} \frac{q_I q_J}{\xi_I \xi_J} , \\
C_1 &= \sum_{I=0}^3 \left(\frac{q_I}{\xi_I} \right)^3 , & C_2 &= \sum_{I \neq J} \left(\frac{q_I}{\xi_I} \right)^2 \frac{q_J}{\xi_J} , & C_3 &= \sum_{I<J<K} \frac{q_I q_J q_K}{\xi_I \xi_J \xi_K} .
\end{aligned} \tag{4.22}$$

Using this coordinate change, the fields of the defect solution can be expanded in FG coordinates. We obtain the following on-shell action and one-point functions,

$$\begin{aligned}
I_{\text{ren}} &= \text{Vol}(\text{AdS}_2)\text{Vol}(S^1)r_+ , \\
\langle T_{ij} \rangle &= \frac{1}{4} \left(\sum_{I=0}^3 \frac{q_I}{\xi_I} \right) \begin{pmatrix} -g_{\text{AdS}_2} & 0 \\ 0 & 2g_{S^1} \end{pmatrix}_{ij} , \\
\langle T_i^i \rangle &= 0 , \\
\langle \mathcal{O}_1 \rangle = \langle \mathcal{O}_2 \rangle = \langle \mathcal{O}_3 \rangle &= 0 , \\
\langle J_{Ii} \rangle &= \frac{q_I}{\sqrt{2}} \delta_{i\theta} .
\end{aligned} \tag{4.23}$$

Note that the expression for I_{ren} is identical to that of the single scalar model, but the radius $r_+ = r_+(\xi_I, q_I)$ will be different.

4.2.3 SU(1, n) Coset Model

For the defect solution (3.19, 3.20) of the SU(1, n) coset model, the FG expansion of the radial coordinate r is

$$\begin{aligned} \frac{1}{r} = & z - \frac{1}{2} q_I \xi^I z^2 - \frac{1}{4} \left[1 + \frac{1}{2} q_I q^I - \frac{3}{4} (q_I \xi^I)^2 \right] z^3 \\ & + \frac{1}{12} q_I \xi^I \left[1 + \frac{3}{2} q_I q^I - \frac{3}{4} (q_I \xi^I)^2 \right] z^4 + \mathcal{O}(z^5) , \end{aligned} \quad (4.24)$$

where η^{IJ} is used to raise the indices of ξ_I and q_I . Using this coordinate change and expanding the fields in FG coordinates, the on-shell action and one-point functions are

$$\begin{aligned} I_{\text{ren}} &= \text{Vol}(\text{AdS}_2) \text{Vol}(S^1) r_+ , \\ \langle T_{ij} \rangle &= -\frac{q_I \xi^I}{2} \begin{pmatrix} -g_{\text{AdS}_2} & 0 \\ 0 & 2g_{S^1} \end{pmatrix}_{ij} , \\ \langle T_i^i \rangle &= 0 , \\ \langle \mathcal{O}_\alpha \rangle &= 0 , \\ \langle J_{Ii} \rangle &= \frac{q_I}{\sqrt{2}} \delta_{i\theta} . \end{aligned} \quad (4.25)$$

5 Regularity

In this section, we impose two regularity conditions on the solutions. First, we demand that the geometry smoothly closes off at the largest positive zero of $f(r)$ without a conical singularity in the bulk spacetime. This condition is analogous to the regularity condition imposed on Euclidean black hole solutions. Second, we fix the periodicity of the S^1 at the conformal boundary such that when the $\text{AdS}_2 \times S^1$ boundary is conformally mapped to $\mathbb{R}^{1,2}$ there is no conical deficit on the boundary. This condition is different from the one imposed in the holographic calculation of supersymmetric Rényi entropies [32, 33, 34, 35], which use solutions that are related by double analytic continuation. For these solutions, the periodicity is related to the Rényi index n .

The regularity conditions will impose constraints on the parameters of the solutions. Since the general solution is only implicit, a detailed analysis is performed for the examples presented in this paper. We will show that for the single scalar and coset models, these conditions imply a bound on the expectation value of the boundary stress tensor.

5.1 General Statements

Given the metric

$$ds^2 = r^2 \sqrt{H(r)} ds_{\text{AdS}_2}^2 + \frac{f(r)}{\sqrt{H(r)}} ds_{S^1}^2 + \frac{\sqrt{H(r)}}{f(r)} dr^2 , \quad (5.1)$$

the center of the space $r = r_+$ is defined to be the largest zero of $f(r) = -1 + r^2 H(r)$. We can identify four criteria a regular geometry should satisfy:

- (a) positivity of the zero, $r_+ > 0$,
- (b) $0 < H(r) < \infty$ on $r \in [r_+, \infty)$,
- (c) $0 < f(r) < \infty$ on $r \in (r_+, \infty)$, and
- (d) no conical singularity at $r = r_+$.

Criteria (b) and (c) are satisfied if $H(r)$ is continuous: the AdS length scale (3.4) is well-defined if and only if the limit $H(r = \infty)$ is positive and finite. Since a zero of $H(r)$ occurs at $f(r) < 0$, positivity of $H(r)$ at large r and continuity imply that the spacetime closes off before a zero of $H(r)$ is ever encountered.

By expanding the metric around the center of the space, criterion (d) is satisfied when

$$f'(r_+)^2 = 4H(r_+) . \quad (5.2)$$

This can be simplified to

$$H'(r_+)(r_+^2 f'(r_+) + 2r_+) = 0 . \quad (5.3)$$

As the second factor is the sum of two positive quantities, a conical singularity can be avoided if we satisfy the condition $H'(r_+) = 0$. As r_+ is determined implicitly in terms of the q_I, ξ_I constants through the equation $f(r_+) = 0$, this condition can be viewed as a constraint on the possible values q_I, ξ_I can take. Additionally, we will see that criterion (a) manifests as an inequality on q_I, ξ_I that we must satisfy.

5.2 Single Scalar Model

The single scalar model is simple enough that the conditions for a regular geometry can be solved exactly. Let us define $x_I \equiv q_I/\xi_I$, but still pick the AdS length scale to be unity, i.e. keep $\xi_0 \xi_1 = 2$. The metric functions become

$$\begin{aligned} H(r) &= \left(1 + \frac{x_0}{r}\right)^2 \left(1 + \frac{x_1}{r}\right)^2 , \\ f(r) &= -1 + \frac{1}{r^2}(r + x_0)^2(r + x_1)^2 . \end{aligned} \quad (5.4)$$

Let us first satisfy the criterion $r_+ > 0$. Solving $f(r) = 0$,

$$0 = (r^2 + r(x_0 + x_1 - 1) + x_0 x_1)(r^2 + r(x_0 + x_1 + 1) + x_0 x_1) . \quad (5.5)$$

When the first factor is zero, we have a solution

$$r_1 = \frac{1}{2} \left(-(x_0 + x_1 - 1) + \sqrt{(x_0 + x_1 - 1)^2 - 4x_0 x_1} \right) , \quad (5.6)$$

where we took the $+$ sign to get the largest root. This solution exists when $(x_0 + x_1 - 1)^2 - 4x_0x_1 \geq 0$, which is a region on the x_0x_1 -plane bounded by a parabola, shown in figure 1a. The red shaded region indicates where r_1 does not exist and the blue shaded region indicates where $r_1 > 0$. When the second factor of (5.5) is zero, we have another solution

$$r_2 = \frac{1}{2} \left(-(x_0 + x_1 + 1) + \sqrt{(x_0 + x_1 + 1)^2 - 4x_0x_1} \right), \quad (5.7)$$

where we also took the $+$ sign. We have also marked regions where this solution exists and is positive in figure 1b. In regions where r_1 and r_2 both exist and $r_1 > 0$, we have $r_1 > r_2$. Therefore, we can take $r_+ = r_1$ and restrict the (x_0, x_1) parameter space to the blue shaded region of figure 1a.




Figure 1: Candidate r_+ for the single scalar model.

Let us now avoid the conical singularity by satisfying $H'(r_+) = 0$. Calculating the derivative of $H(r)$ in (5.4) and plugging in $r_+ = r_1$ from (5.6), we get the condition

$$0 = (x_0 - x_1)^2 - 2(x_0 + x_1). \quad (5.8)$$

This is a parabola, marked by the black curve in figure 1a in the region where $r_+ > 0$. For the single scalar model to admit a regular geometry, the parameters $x_I = q_I/\xi_I$ must satisfy this condition. As a corollary, we can note that

$$0 \leq x_0 + x_1 < 2. \quad (5.9)$$

This implies that the components of the boundary stress tensor (4.20) have bounded expectation value. Additionally, the pure AdS_4 vacuum ($x_0 = x_1 = 0$) is the only solution with regular geometry and $\langle T_{ij} \rangle = 0$.

5.3 $SU(1, n)$ Coset Model

The coset model is also simple enough that the conditions for a regular geometry can be solved exactly. We can note that

$$H(r) = \left(1 - \frac{q_I \xi^I}{r} - \frac{q_I q^I}{2r^2}\right)^2, \quad (5.10)$$

actually has the same form as (5.4), where

$$x_0 = \frac{-q_I \xi^I - \sqrt{(q_I \xi^I)^2 + 2q_I q^I}}{2}, \quad x_1 = \frac{-q_I \xi^I + \sqrt{(q_I \xi^I)^2 + 2q_I q^I}}{2}. \quad (5.11)$$

This map is always well-defined as $(q_I \xi^I)^2 + 2q_I q^I \geq 0$, which can be checked by rotating to the frame where $\xi_I = (\sqrt{2}, 0, 0, \dots)$. Thus all our results for the single scalar model can be carried over. The bound (5.9) for the single scalar model translates to the same bound on $\langle T_{ij} \rangle$ for the coset model:

$$0 \leq -q_I \xi^I < 2. \quad (5.12)$$

The condition (5.8) for a regular geometry translates to

$$0 = (q_I \xi^I)^2 + 2q_I q^I + 2q_I \xi^I. \quad (5.13)$$

We can show that the only regular geometry with vanishing $\langle T_{ij} \rangle$ is the AdS_4 vacuum. If we rotate to the frame where $\xi_I = (\sqrt{2}, 0, 0, \dots)$, the only q which satisfies $q_I \xi^I = 0$ and $q_I q^I = 0$ is $q_I = 0$. A general ξ then has a q in the orbit of $q_I = 0$, which is still the zero vector.

5.4 Gauged STU Model

For the gauged STU model, it is not practical to solve $f(r) = 0$ to find r_+ as f is a quartic polynomial. However, we still expect the criterion $r_+ > 0$ to impose an inequality on the four-dimensional parameter space (x_0, x_1, x_2, x_3) and the condition of avoiding a conical singularity to reduce this to a three-dimensional hypersurface. However, note that unlike the single scalar and coset models, the expectation value $\langle T_{ij} \rangle$ is not bounded. In appendix C we give special cases of the STU model with regular geometry which can have arbitrarily large $x_0 + x_1 + x_2 + x_3$.

6 Discussion

In this paper, we constructed solutions of four-dimensional $N = 2$ gauged supergravity by a double analytic continuation of the half-BPS black hole solutions first found by Sabra [16]. While the black hole solutions exist for arbitrary prepotentials, explicit expressions for the

scalars fields involve algebraic equations which in general can only be solved numerically. We considered three explicit examples of matter-coupled gauged supergravities, namely the single scalar model, the gauged STU model, and the $SU(1, n)/U(1) \times SU(n)$ coset model to find solutions and calculate holographic observables.

The solutions we find are holographic duals to line defects in three-dimensional SCFTs. The defect is characterized by a non-trivial expectation value of the R -symmetry and flavor currents along the S^1 factor in the $AdS_2 \times S^1$ description of the defect. After conformally mapping to Minkowski space, this corresponds to a holonomy when encircling the line defect. The expectation values of the real scalar operators vanish for general models as a consequence of supersymmetry.

For a conformal defect on $AdS_2 \times S^1$, the expectation value of the stress tensor can be parameterized by a single coefficient h ,

$$\langle T_{ab} \rangle = h g_{ab}^{AdS_2} , \quad \langle T_{\theta\theta} \rangle = -2h g_{\theta\theta} , \quad (6.1)$$

in analogy to the scaling dimension of local operators [36, 37]. However, there are in general no unitarity bounds on h which follow from the superconformal algebra. For line operators in $N = 4$ SYM and ABJM theories, h can be related to the so-called Bremsstrahlung function B [38, 39, 40, 41, 42] which has been used in the application of conformal bootstrap techniques to the study of defects [43, 44, 45, 46]. For the single scalar and coset models studied in this paper we find that $-2 < h \leq 0$, where the upper bound is saturated only by the AdS_4 vacuum. However, such a bound does not seem to generally hold, since for the gauged STU model, h can become arbitrarily negative. Based on numerical searches we conjecture that only the AdS_4 vacuum has vanishing h . Note that recently the relation of h and B , as well as the negativity of h has been established on the SCFT side for various defect theories [47, 48, 49, 50] and the arguments should carry over to the defects dual to the solutions studied in the paper.⁷

The solutions we find are related to supergravity solutions [24, 33, 34, 35] which are holographic duals for a supersymmetric version of Rényi entropy first formulated in [32]. We note two differences. First, the solutions we find in Minkowski time signature have real gauge fields, unlike the duals cited above.⁸ Second, we impose the condition that the periodicity of the circle in $AdS_2 \times S^1$ boundary is such that after a conformal map we obtain flat space without a conical singularity. On the other hand, in the holographic duals to the Super-Rényi entropy, the conical singularity is related to the Rényi index n . We note that in [24, 33, 34, 35] the holographic calculation of the Rényi entropy was compared to a localization calculation and agreement was found, and it would be interesting to see whether such a calculation can be performed for the holonomy defects described in this paper.

Another interesting question is whether more general solutions going beyond the examples discussed in this paper can be found. First, it would be interesting to study (numerical) solutions for more complicated superpotentials. Second, it would be interesting to see whether

⁷We thank Marco Meineri and Lorenzo Bianchi for a useful correspondence regarding these matters.

⁸After analytic continuation to Euclidean signature, the gauge fields in both cases are real.

one can go beyond the gauged supergravity approximation and find solutions dual to holonomy defects in ten- or eleven-dimensional duals of $N = 2$ SCFTs. Uplifting the solutions found in this paper might prove to be a useful guide in this direction [22].

Acknowledgements

The work of M. G. was supported, in part, by the National Science Foundation under grant PHY-19-14412. All the authors are grateful to the Mani L. Bhaumik Institute for Theoretical Physics for support.

A Supersymmetry

We use the metric conventions $\eta = (-+++)$ and $\epsilon_{0123} = -\epsilon^{0123} = 1$. The gamma matrices are defined as usual, e.g.

$$\{\gamma_a, \gamma_b\} = 2\eta_{ab} , \quad \gamma_{ab} = \frac{1}{2}[\gamma_a, \gamma_b] , \quad \gamma_5 = i\gamma_0\gamma_1\gamma_2\gamma_3 . \quad (\text{A.1})$$

The two chiral gravitinos can be written in terms of a single complex (Dirac) spinor ψ_μ , and likewise for the gauginos λ^α . The supersymmetry transformations of the four-dimensional gauged supergravity are [18]

$$\begin{aligned} \delta\psi_\mu &= \left(\partial_\mu + \frac{1}{4}\omega_\mu^{ab}\gamma_{ab} + \frac{i}{2}Q_\mu\gamma_5 + ig\xi_I A_\mu^I + ge^{\mathcal{K}/2}\gamma_\mu\xi_I(\text{Im } Z^I + i\gamma_5 \text{Re } Z^I) \right. \\ &\quad \left. + \frac{i}{4}e^{\mathcal{K}/2}\gamma^{ab}(\text{Im } \mathcal{N})_{IJ}(\text{Im}(F_{ab}^{-I}Z^J) - i\gamma_5 \text{Re}(F_{ab}^{-I}Z^J))\gamma_\mu \right) \epsilon , \\ \delta\lambda^\alpha &= \left(\gamma^\mu\partial_\mu(\text{Re } z^\alpha - i\gamma_5 \text{Im } z^\alpha) + 2ge^{\mathcal{K}/2}\xi_I \left(\text{Im}(\mathcal{D}_{\bar{\beta}}\bar{Z}^I g^{\alpha\bar{\beta}}) - i\gamma_5 \text{Re}(\mathcal{D}_{\bar{\beta}}\bar{Z}^I g^{\alpha\bar{\beta}}) \right) \right. \\ &\quad \left. + \frac{i}{2}e^{\mathcal{K}/2}\gamma^{ab}(\text{Im } \mathcal{N})_{IJ} \left(\text{Im}(F_{ab}^{-I}\mathcal{D}_{\bar{\beta}}\bar{Z}^J g^{\alpha\bar{\beta}}) - i\gamma_5 \text{Re}(F_{ab}^{-I}\mathcal{D}_{\bar{\beta}}\bar{Z}^J g^{\alpha\bar{\beta}}) \right) \right) \epsilon , \end{aligned} \quad (\text{A.2})$$

where ϵ is a complex spinor, and we have defined

$$F_{ab}^{\pm I} \equiv \frac{1}{2}(F_{ab}^I \pm \tilde{F}_{ab}^I) , \quad \tilde{F}_{ab}^I \equiv -\frac{i}{2}\epsilon_{abcd}F^{cd} . \quad (\text{A.3})$$

The Kähler connection Q_μ is

$$Q_\mu = -\frac{i}{2}(\partial_\mu\tau^\alpha\partial_\alpha\mathcal{K} - \partial_\mu\bar{\tau}^{\bar{\alpha}}\partial_{\bar{\alpha}}\mathcal{K}) . \quad (\text{A.4})$$

For the gauged STU model defect solution (3.14), we can work with the explicit coordinates $(x^0, x^1, x^2, x^3) = (t, \eta, \theta, r)$ and the metric

$$ds^2 = r^2\sqrt{H}\left(\frac{-dt^2 + d\eta^2}{\eta^2}\right) + \frac{f}{\sqrt{H}}d\theta^2 + \frac{\sqrt{H}}{f}dr^2 . \quad (\text{A.5})$$

The non-vanishing spin connection 1-forms of the metric are

$$\begin{aligned} \omega^{01} &= -\frac{dt}{\eta} , & \omega^{03} &= \frac{f^{1/2}}{H^{1/4}}\frac{d}{dr}(rH^{1/4})\frac{dt}{\eta} , \\ \omega^{13} &= \frac{f^{1/2}}{H^{1/4}}\frac{d}{dr}(rH^{1/4})\frac{d\eta}{\eta} , & \omega^{23} &= \frac{f^{1/2}}{H^{1/4}}\frac{d}{dr}\left(\frac{f^{1/2}}{H^{1/4}}\right)d\theta . \end{aligned} \quad (\text{A.6})$$

For the following calculations, we use the parametrization $(Z^0, Z^1, Z^2, Z^3) = (i, iz^2z^3, iz^1z^3, iz^1z^2)$. The BPS equations (A.2) simplify to

$$\begin{aligned} 0 &= \delta\psi_\mu = \left(\partial_\mu + \frac{1}{4}\omega_\mu^{ab}\gamma_{ab} + ig\xi_I A_\mu^I + \sqrt{2}g\gamma_\mu\frac{d}{dr}(rH^{1/4}) - \frac{i}{2}\gamma_{23}\gamma_\mu\frac{d}{dr}(H^{-1/4}) \right) , \\ 0 &= \delta\lambda^\alpha = \frac{dz^\alpha}{dr}\left(\frac{f^{1/2}}{H^{1/4}}\gamma_3 + 2\sqrt{2}grH^{1/4} + \frac{i}{H^{1/4}}\gamma_{23}\right)\epsilon . \end{aligned} \quad (\text{A.7})$$

The gaugino equation implies the projector

$$0 = \left(1 + \frac{2\sqrt{2}gr\sqrt{H}}{\sqrt{f}}\gamma_3 - \frac{i}{\sqrt{f}}\gamma_2 \right) \epsilon . \quad (\text{A.8})$$

The $\mu = t, \eta, \theta$ components of the gravitino equation then simplify to

$$\begin{aligned} 0 &= \left(\partial_t - \frac{1}{2\eta}\gamma_{01} - \frac{i}{2\eta}\gamma_{023} \right) \epsilon , \\ 0 &= \left(\partial_\eta - \frac{i}{2\eta}\gamma_{123} \right) \epsilon , \\ 0 &= \left(\partial_\theta + i\sqrt{2}g \left(-1 + \frac{1}{\sqrt{2}}\xi_I\mu^I \right) \right) \epsilon . \end{aligned} \quad (\text{A.9})$$

These can be integrated to

$$\epsilon = \exp \left(-i\sqrt{2}g\theta \left(-1 + \frac{1}{\sqrt{2}}\xi_I\mu^I \right) \right) \exp \left(\frac{i}{2}\gamma_{123} \ln \eta \right) \exp \left(\frac{t}{2}(\gamma_{01} + i\gamma_{023}) \right) \tilde{\epsilon}(r) . \quad (\text{A.10})$$

We can see that we need $\xi_I\mu^I \in 2\sqrt{2}\mathbb{Z}$ in order for ϵ to be anti-periodic under the identification $\theta \sim \theta + \pi/\sqrt{2}g$. The $\mu = r$ component of the gravitino equation simplifies to

$$\left(\partial_r + \frac{1}{8}\frac{H'}{H} + \frac{f'}{8\sqrt{2}gr\sqrt{H}\sqrt{f}}\gamma_3 \right) \epsilon . \quad (\text{A.11})$$

The gaugino projector (A.8) and the radial equation (A.11) take the form of the equation solved in the appendix of [51], by identifying

$$\begin{aligned} x &\equiv \frac{2\sqrt{2}gr\sqrt{H}}{\sqrt{f}} , & y &\equiv \frac{-i}{\sqrt{f}} , \\ \Gamma_1 &\equiv \gamma_3 , & \Gamma_2 &\equiv \gamma_2 . \end{aligned} \quad (\text{A.12})$$

The solution is

$$\tilde{\epsilon}(r) = \frac{1}{H^{1/8}} \left(\sqrt{\sqrt{f} + 2\sqrt{2}gr\sqrt{H}} - \gamma_2 \sqrt{\sqrt{f} - 2\sqrt{2}gr\sqrt{H}} \right) (1 - \gamma_3) \epsilon_0 , \quad (\text{A.13})$$

where ϵ_0 is a constant spinor.

B Vanishing of Scalar One-Point Functions from Supersymmetry

The scalar one-point function is given by

$$\langle \bar{\mathcal{O}}_{\bar{\alpha}} \rangle = \lim_{\epsilon \rightarrow 0} \left[\frac{1}{\epsilon^2} (z g_{\beta\bar{\alpha}} \partial_z \tau^\beta + \partial_{\bar{\alpha}} \mathcal{W}) \right] . \quad (\text{B.1})$$

The derivative of the superpotential $\mathcal{W}$ simplifies to

$$\begin{aligned}\partial_{\bar{\alpha}}\mathcal{W} &= \partial_{\bar{\alpha}}\left(-\sqrt{2}e^{\mathcal{K}/2}\xi_I|Z^I|\right) \\ &= -\frac{1}{\sqrt{2}}e^{\mathcal{K}/2}\xi_I\left(\sqrt{\frac{Z^I}{\bar{Z}^I}}\partial_{\bar{\alpha}}\bar{Z}^I + (\partial_{\bar{\alpha}}\mathcal{K})|Z^I|\right),\end{aligned}\tag{B.2}$$

where $|Z^I|^2 = Z^I(\tau)\bar{Z}^I(\bar{\tau})$. For real scalars, we can choose a parameterization such that $\bar{Z}^I = Z^I$. This implies

$$\partial_{\bar{\alpha}}\mathcal{W} = -\frac{1}{\sqrt{2}}e^{\mathcal{K}/2}\xi_I(\partial_{\bar{\alpha}}\bar{Z}^I + (\partial_{\bar{\alpha}}\mathcal{K})\bar{Z}^I) = -\frac{1}{\sqrt{2}}e^{\mathcal{K}/2}\xi_I\mathcal{D}_{\bar{\alpha}}\bar{Z}^I, \tag{B.3}$$

so that

$$\langle\bar{\mathcal{O}}_{\bar{\alpha}}\rangle = \lim_{\epsilon\rightarrow 0}\left[\frac{1}{\epsilon^2}\left(zg_{\beta\bar{\alpha}}\partial_z\tau^{\beta} - \frac{1}{\sqrt{2}}e^{\mathcal{K}/2}\xi_I\mathcal{D}_{\bar{\alpha}}\bar{Z}^I\right)\right]_{z=\epsilon}.\tag{B.4}$$

The gaugino BPS variation in FG coordinates is

$$(z\gamma_3\partial_z\tau^{\beta} - 2ige^{\mathcal{K}/2}\xi_Ig^{\beta\bar{\alpha}}\mathcal{D}_{\bar{\alpha}}\bar{Z}^I\gamma_5)\epsilon + \mathcal{O}(z^3)\epsilon = 0, \tag{B.5}$$

since $F_{ab} \sim 1/r^2 \sim \mathcal{O}(z^2)$. At $\mathcal{O}(z^2)$, the BPS equations imply

$$z\partial_z\tau^{\beta} = \pm 2ige^{\mathcal{K}/2}\xi_Ig^{\beta\bar{\alpha}}\mathcal{D}_{\bar{\alpha}}\bar{Z}^I. \tag{B.6}$$

Without loss of generality, we can choose the upper sign by sending $g \rightarrow -g$ if necessary. After setting $g^2 = 1/8$ we have

$$\langle\bar{\mathcal{O}}_{\bar{\alpha}}\rangle = \langle\mathcal{O}_{\alpha}\rangle = 0. \tag{B.7}$$

C STU Model Special Cases

Here we give a construction for STU models with regular geometry and arbitrarily large $x_0 + x_1 + x_2 + x_3$. The approach we took to find these models was different than that of section 5.2. Instead of solving the condition $f = 0$ and then $H' = 0$, we first solved $H' = 0$ and then $f = 0$. The benefit is that H' is a lower-degree polynomial and is technically simpler to solve. The downside is that this generates spurious solutions: it is possible that the r we obtain is not the largest root r_+ , and r_+ does not satisfy the equation $H' = 0$. These spurious solutions then need to be removed by hand.

To summarize our findings, consider the following construction:

1. Let x_0 be any positive number.
2. Numerically solve the equation

$$27x_1(x_0 - x_1)^4 = -16x_0(x_0 + 3x_1)^2. \tag{C.1}$$

Let x_1 be the unique solution satisfying $-x_0/3 < x_1 < 0$.

3. Consider an STU model with unit AdS_4 length scale where

$$x_0 = \frac{q_0}{\xi_0} , \quad x_1 = \frac{q_1}{\xi_1} = \frac{q_2}{\xi_2} = \frac{q_3}{\xi_3} . \quad (\text{C.2})$$

Numerically solve the equation $f(r) = 0$ for r ,

$$(r + x_0)(r + x_1)^3 = r^2 . \quad (\text{C.3})$$

There exist exactly two solutions: a positive solution greater than $-x_1$, and a negative solution less than $-x_0$. Let r_+ be the positive solution.

4. Check that $H'(r_+) = 0$. This is guaranteed by the following argument. Consider $r^* = -4x_0x_1/(x_0 + 3x_1) > 0$ which satisfies $H'(r^*) = 0$. This also satisfies $f(r^*) = 0$, as plugging $r = r^*$ into (C.3) simplifies to (C.1), which is satisfied by construction of x_1 . But as the positive solution to $f = 0$ is unique, we must have $r_+ = r^*$.

The steps above give a STU model with regular geometry. To prove that $x_0 + 3x_1$ is arbitrarily large, we need a better bound than $-x_0/3 < x_1 < 0$. To satisfy (C.1) for large x_0 , we have

$$x_1 \sim -\frac{16}{27x_0} . \quad (\text{C.4})$$

Therefore $x_0 + 3x_1 \approx x_0$ for large x_0 , and can be arbitrarily large.

References

- [1] J. M. Maldacena, *Wilson loops in large N field theories*, *Phys. Rev. Lett.* **80** (1998) 4859–4862, [[hep-th/9803002](#)].
- [2] S.-J. Rey and J.-T. Yee, *Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity*, *Eur. Phys. J.* **C22** (2001) 379–394, [[hep-th/9803001](#)].
- [3] N. Drukker and B. Fiol, *All-genus calculation of Wilson loops using D -branes*, *JHEP* **02** (2005) 010, [[hep-th/0501109](#)].
- [4] S. A. Hartnoll and S. Kumar, *Higher rank Wilson loops from a matrix model*, *JHEP* **08** (2006) 026, [[hep-th/0605027](#)].
- [5] S. Yamaguchi, *Wilson loops of anti-symmetric representation and $D5$ -branes*, *JHEP* **05** (2006) 037, [[hep-th/0603208](#)].
- [6] J. Gomis and F. Passerini, *Holographic Wilson Loops*, *JHEP* **08** (2006) 074, [[hep-th/0604007](#)].
- [7] J. Gomis and F. Passerini, *Wilson Loops as $D3$ -Branes*, *JHEP* **01** (2007) 097, [[hep-th/0612022](#)].
- [8] E. D’Hoker, J. Estes and M. Gutperle, *Gravity duals of half-BPS Wilson loops*, *JHEP* **06** (2007) 063, [[0705.1004](#)].
- [9] O. Lunin, *On gravitational description of Wilson lines*, *JHEP* **06** (2006) 026, [[hep-th/0604133](#)].
- [10] J. P. Gauntlett, O. A. Mac Conamhna, T. Mateos and D. Waldram, *New supersymmetric $AdS(3)$ solutions*, *Phys. Rev. D* **74** (2006) 106007, [[hep-th/0608055](#)].
- [11] J. P. Gauntlett, N. Kim and D. Waldram, *Supersymmetric $AdS(3)$, $AdS(2)$ and Bubble Solutions*, *JHEP* **04** (2007) 005, [[hep-th/0612253](#)].
- [12] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, *Building a Holographic Superconductor*, *Phys. Rev. Lett.* **101** (2008) 031601, [[0803.3295](#)].
- [13] K. Balasubramanian and J. McGreevy, *Gravity duals for non-relativistic CFTs*, *Phys. Rev. Lett.* **101** (2008) 061601, [[0804.4053](#)].
- [14] S. S. Gubser and S. S. Pufu, *The Gravity dual of a p -wave superconductor*, *JHEP* **11** (2008) 033, [[0805.2960](#)].
- [15] M. Gutperle and M. Vicino, *Conformal defect solutions in $N = 2$, $D = 4$ gauged supergravity*, *Nucl. Phys.* **B942** (2019) 149–163, [[1811.04166](#)].

- [16] W. A. Sabra, *Anti-de Sitter BPS black holes in $N=2$ gauged supergravity*, *Phys. Lett. B* **458** (1999) 36–42, [[hep-th/9903143](#)].
- [17] E. Lauria and A. Van Proeyen, *$N = 2$ Supergravity in $D = 4, 5, 6$ Dimensions*, vol. 966. Springer, 2020, [10.1007/978-3-030-33757-5](#).
- [18] S. L. Cacciatori, D. Klemm, D. S. Mansi and E. Zorzan, *All timelike supersymmetric solutions of $N=2$, $D=4$ gauged supergravity coupled to abelian vector multiplets*, *JHEP* **05** (2008) 097, [[0804.0009](#)].
- [19] S. L. Cacciatori and D. Klemm, *Supersymmetric $AdS(4)$ black holes and attractors*, *JHEP* **01** (2010) 085, [[0911.4926](#)].
- [20] K. Hristov and S. Vandoren, *Static supersymmetric black holes in AdS_4 with spherical symmetry*, *JHEP* **04** (2011) 047, [[1012.4314](#)].
- [21] M. J. Duff and J. T. Liu, *Anti-de Sitter black holes in gauged $N = 8$ supergravity*, *Nucl. Phys. B* **554** (1999) 237–253, [[hep-th/9901149](#)].
- [22] M. Cvetič, M. J. Duff, P. Hoxha, J. T. Liu, H. Lu, J. X. Lu et al., *Embedding AdS black holes in ten-dimensions and eleven-dimensions*, *Nucl. Phys. B* **558** (1999) 96–126, [[hep-th/9903214](#)].
- [23] A. Cabo-Bizet, U. Kol, L. A. Pando Zayas, I. Papadimitriou and V. Rathee, *Entropy functional and the holographic attractor mechanism*, *JHEP* **05** (2018) 155, [[1712.01849](#)].
- [24] S. M. Hosseini, C. Toldo and I. Yaakov, *Supersymmetric Rényi entropy and charged hyperbolic black holes*, [1912.04868](#).
- [25] S. de Haro, S. N. Solodukhin and K. Skenderis, *Holographic reconstruction of space-time and renormalization in the AdS / CFT correspondence*, *Commun. Math. Phys.* **217** (2001) 595–622, [[hep-th/0002230](#)].
- [26] K. Skenderis, *Lecture notes on holographic renormalization*, *Class. Quant. Grav.* **19** (2002) 5849–5876, [[hep-th/0209067](#)].
- [27] I. R. Klebanov and E. Witten, *AdS / CFT correspondence and symmetry breaking*, *Nucl. Phys. B* **556** (1999) 89–114, [[hep-th/9905104](#)].
- [28] C. Cordova, T. T. Dumitrescu and K. Intriligator, *Multiplets of Superconformal Symmetry in Diverse Dimensions*, *JHEP* **03** (2019) 163, [[1612.00809](#)].
- [29] V. Balasubramanian and P. Kraus, *A Stress tensor for Anti-de Sitter gravity*, *Commun. Math. Phys.* **208** (1999) 413–428, [[hep-th/9902121](#)].

- [30] P. Kraus, *Lectures on black holes and the $AdS(3) / CFT(2)$ correspondence*, *Lect. Notes Phys.* **755** (2008) 193–247, [[hep-th/0609074](#)].
- [31] A. Batrachenko, J. T. Liu, R. McNees, W. A. Sabra and W. Y. Wen, *Black hole mass and Hamilton-Jacobi counterterms*, *JHEP* **05** (2005) 034, [[hep-th/0408205](#)].
- [32] T. Nishioka and I. Yaakov, *Supersymmetric Renyi Entropy*, *JHEP* **10** (2013) 155, [[1306.2958](#)].
- [33] T. Nishioka, *The Gravity Dual of Supersymmetric Renyi Entropy*, *JHEP* **07** (2014) 061, [[1401.6764](#)].
- [34] X. Huang and Y. Zhou, *$\mathcal{N} = 4$ Super-Yang-Mills on conic space as hologram of STU topological black hole*, *JHEP* **02** (2015) 068, [[1408.3393](#)].
- [35] M. Crossley, E. Dyer and J. Sonner, *Super-Renyi entropy & Wilson loops for $\mathcal{N} = 4$ SYM and their gravity duals*, *JHEP* **12** (2014) 001, [[1409.0542](#)].
- [36] A. Kapustin, *Wilson-'t Hooft operators in four-dimensional gauge theories and S-duality*, *Phys. Rev. D* **74** (2006) 025005, [[hep-th/0501015](#)].
- [37] N. Drukker, J. Gomis and D. Young, *Vortex Loop Operators, M2-branes and Holography*, *JHEP* **03** (2009) 004, [[0810.4344](#)].
- [38] A. Lewkowycz and J. Maldacena, *Exact results for the entanglement entropy and the energy radiated by a quark*, *JHEP* **05** (2014) 025, [[1312.5682](#)].
- [39] B. Fiol, E. Gerchkovitz and Z. Komargodski, *Exact Bremsstrahlung Function in $N = 2$ Superconformal Field Theories*, *Phys. Rev. Lett.* **116** (2016) 081601, [[1510.01332](#)].
- [40] P. Liendo and C. Meneghelli, *Bootstrap equations for $\mathcal{N} = 4$ SYM with defects*, *JHEP* **01** (2017) 122, [[1608.05126](#)].
- [41] L. Bianchi, L. Griguolo, M. Preti and D. Seminara, *Wilson lines as superconformal defects in ABJM theory: a formula for the emitted radiation*, *JHEP* **10** (2017) 050, [[1706.06590](#)].
- [42] L. Bianchi, M. Preti and E. Vescovi, *Exact Bremsstrahlung functions in ABJM theory*, *JHEP* **07** (2018) 060, [[1802.07726](#)].
- [43] A. Gadde, *Conformal constraints on defects*, *JHEP* **01** (2020) 038, [[1602.06354](#)].
- [44] M. Fukuda, N. Kobayashi and T. Nishioka, *Operator product expansion for conformal defects*, *JHEP* **01** (2018) 013, [[1710.11165](#)].
- [45] M. Billo, V. Goncalves, E. Lauria and M. Meineri, *Defects in conformal field theory*, *JHEP* **04** (2016) 091, [[1601.02883](#)].

- [46] E. Lauria, M. Meineri and E. Trevisani, *Radial coordinates for defect CFTs*, *JHEP* **11** (2018) 148, [[1712.07668](#)].
- [47] L. Bianchi, M. Lemos and M. Meineri, *Line Defects and Radiation in $\mathcal{N} = 2$ Conformal Theories*, *Phys. Rev. Lett.* **121** (2018) 141601, [[1805.04111](#)].
- [48] M. Lemos, P. Liendo, M. Meineri and S. Sarkar, *Universality at large transverse spin in defect CFT*, *JHEP* **09** (2018) 091, [[1712.08185](#)].
- [49] K. Jensen, A. O’Bannon, B. Robinson and R. Rodgers, *From the Weyl Anomaly to Entropy of Two-Dimensional Boundaries and Defects*, *Phys. Rev. Lett.* **122** (2019) 241602, [[1812.08745](#)].
- [50] L. Bianchi and M. Lemos, *Superconformal surfaces in four dimensions*, [1911.05082](#).
- [51] L. J. Romans, *Supersymmetric, cold and lukewarm black holes in cosmological Einstein-Maxwell theory*, *Nucl. Phys.* **B383** (1992) 395–415, [[hep-th/9203018](#)].