

Kinetochore assembly throughout the cell cycle

Alexandra P. Navarro^{1, 2} and Iain M. Cheeseman^{1, 2 *}

¹ Whitehead Institute for Biomedical Research, 455 Main Street, Cambridge, MA 02142

² Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02142

*Corresponding author:
E-mail: icheese@wi.mit.edu
Phone: (617) 324-2503
Fax: (617) 258-5578

Abstract

The kinetochore plays an essential role in facilitating chromosome segregation during cell division. This massive protein complex assembles onto the centromere of chromosomes and enables their attachment to spindle microtubules during mitosis. The kinetochore also functions as a signaling hub to regulate cell cycle progression, and is crucial to ensuring the fidelity of chromosome segregation. Despite the fact that kinetochores are large and robust molecular assemblies, they are also highly dynamic structures that undergo structural and organizational changes throughout the cell cycle. This review will highlight our current understanding of kinetochore structure and function, focusing on the dynamic processes that underlie kinetochore assembly.

Introduction

Cell division is the fundamental process by which one cell gives rise to two genetically identical daughter cells. In eukaryotes, each cell division cycle requires that the genetic material be properly duplicated and segregated across the dividing cell. It is critical that the DNA is equally distributed between the two cells, as chromosome gains or losses can be highly detrimental to an organism. Accurate chromosome segregation depends upon the macromolecular structure called the kinetochore [1], which is assembled upon a specific region of each chromosome known as the centromere. The kinetochore is comprised of multiple individual proteins that assemble together to form a network of interdependent sub-complexes. This large protein assembly can be divided into two distinct regions: the inner kinetochore, which associates with centromere DNA and is assembled throughout the cell cycle; and the outer kinetochore, which binds directly to spindle microtubules and is assembled only in mitosis.

The kinetochore is a massive protein assembly with more than 110 different components and around 400 molecules of each protein at a single human kinetochore, that together form a structure that is roughly 5-10-fold larger than a ribosome [2, 3]. However, despite its large size, the kinetochore is a remarkably dynamic structure, exhibiting changes in its assembly, dynamics, organization, and function throughout the cell cycle. Our understanding of kinetochore assembly often focuses on the recruitment of the outer kinetochore in mitosis. However, a subset of kinetochore components are

present at centromeres constitutively, and the manner in which these protein complexes adapt to the events of the cell cycle is also important to fully understand the principles of kinetochore assembly.

During interphase, three critical dynamic events must occur that result in substantial changes to the centromere. First, during G1, new molecules of the centromere-specific histone CENP-A are deposited into centromeric chromatin, resulting in a substantial increase of CENP-A molecules at centromeres (Figure 1A). During S phase, centromeric DNA undergoes a dramatic structural change, similar to other chromosomal regions, as chromosomes are unwound and replicated, resulting in at least the temporary eviction of chromatin-associated factors (Figure 1B). Finally, during the G2/M transition, chromosomal DNA undergoes further structural changes as chromosomes become condensed with sister chromatids remaining bound together at centromeres (Figure 1C,D). In addition to navigating DNA compaction during the G2/M transition, the kinetochore must assemble its outer structure to prepare for the formation of microtubule attachments in mitosis. The constitutive centromere components that comprise the inner kinetochore must remain assembled despite these vast changes and disruptions that occur to centromeric chromatin over the course of the cell cycle. Once in mitosis, the kinetochore recruits additional dynamically associated proteins that are required to enable the diverse mitotic kinetochore functions. Here, we review kinetochore assembly in the context of the constantly changing centromere chromatin status and discuss how kinetochore organization and function are altered to facilitate these changes and confer the diverse functions of the kinetochore throughout the cell cycle.

The centromere locus

An understanding of kinetochore assembly and organization first requires a consideration of the centromere, the chromosomal locus upon which the kinetochore is built. Despite providing the functional platform for kinetochore assembly, centromere structure, position, and sequence vary widely across organisms [4]. For example, in the budding yeast *S. cerevisiae*, centromeric DNA is 125 base pairs, with a defined sequence that is both necessary and sufficient for centromere function and kinetochore

assembly, referred to as a point centromere. Conversely, other commonly studied organisms, including mammals, *Drosophila*, and fission yeast, have centromeres that consist of long arrays of repetitive DNA sequences that can span ten's of thousands to millions of base pairs in length, termed regional centromeres. Finally, the centromeres found in some plants, nematodes, and insect species are not restricted to a single position, but instead span the complete chromosome, a centromere type called holocentric. Holocentricity makes it such that kinetochores assemble along the entire length of the chromosome [5].

Regional centromeres are often composed of repetitive DNA elements including retro-transposon elements and small units of repetitive sequences called satellite DNA that can span kilobases to megabases of DNA [4, 6]. The organization of these centromeres can be divided into two regions: 1) the core centromere region, which is comprised of homogenous ordered repeats (largely satellite DNA) upon which the kinetochore is assembled, and 2) the pericentromere, which flanks the core centromere and is made of less ordered repetitive sequences that are heterochromatic [7]. In humans, the repeating unit of the core centromere region is known as α -satellite DNA. The α -satellite sequence is 171-bp in length, where each unit is arranged in a head-to-tail manner forming higher-order repeats throughout the core centromere [7-11]. Subsets of α -satellite repeats additionally contain a 17 bp motif known as the CENP-B box [12, 13], which serves to recruit the centromere protein (CENP), CENP-B. Regional centromeres promote the assembly of larger kinetochore structures that bind to multiple microtubules during mitosis [2]. In fact, kinetochore size has been shown to scale with centromere length, which can ultimately impact kinetochore function as chromosomes with larger kinetochores are prone to mis-segregate in mitosis [14].

Although regional centromeres can be highly structured, the underlying DNA sequence is not sufficient to generate a functional centromere [15]. Indeed, although the presence of repetitive sequences at regional centromeres is common, there are exceptions to this, such as the non-repetitive centromeres found in horse, orangutan, and chicken chromosomes [16-20]. Instead, regional centromeres are considered to be epigenetically defined based on the presence of the centromere-specific histone CENP-A, a variant of the canonical histone protein H3 [21]. The presence of CENP-A at

centromeres is not unique to organisms with regional centromeres. CENP-A can be found at the centromeres of many organisms, including those with holocentric and point centromeres. In fact, CENP-A is present at the centromere of most model organisms including *S. cerevisiae*, *S. pombe*, *D. melanogaster*, *C. elegans*, and humans [22-25]. However, there are notable exceptions, including kinetoplastids and some insects such as Lepidoptera, which lack centromere-specific histones [26-28]. The precise mechanism by which centromeres are specified in these organisms is an active area of research. Recent work in Lepidoptera suggests that kinetochores are specifically assembled at regions of the genome with low transcriptional activity and have adapted to rely on the function of other centromere proteins [29, 30].

In addition to CENP-A, multiple factors act to collectively specify the centromere region. Among these, are the proteins that localize to centromeres throughout the cell cycle and form the inner kinetochore, collectively referred to as the Constitutive Centromere Associated Network (CCAN) [31, 32]. The CCAN is composed of 16 proteins that can be further organized into 5 distinct groups based on their physical associations and dependencies for centromere localization: CENP-C, CENP/L/N, CENP-H/I/K/M, CENP-O/P/Q/U/R, and CENP-T/W/S/X [33]. Together, these sub-complexes function to bind specifically to CENP-A and recruit the components of the outer kinetochore. Importantly, although the features that define and characterize the centromere can vary dramatically between species, the requirement for constitutively bound centromere proteins to maintain centromere identity and function throughout the cell cycle remains constant.

Kinetochores assembly during G1 – refreshing centromere chromatin

The first dynamic changes to centromere chromatin and kinetochore assembly state occur in G1, when new CENP-A histones are deposited at the centromere. CENP-A is both necessary and sufficient for the recruitment of all known kinetochore proteins [34-42], and thus this deposition process is a critical event for the specification and maintenance of centromere identity. The deposition of new CENP-A molecules involves the coordination of multiple factors. Importantly, the cell must distinguish between CENP-A nucleosomes and canonical H3 nucleosomes, which share ~62% sequence

identity in their histone fold domain [21]. To achieve this, CENP-A deposition depends on a dedicated chaperone called HJURP (Holliday Junction Recognizing Protein) in vertebrates [43-45]. Functional orthologs of HJURP also exist in *S. cerevisiae* and *S. pombe*, called Scm3, and in *D. melanogaster*, called Cal1 [46-56]. The HJURP chaperone binds specifically to the CENP-A targeting domain (CATD), a region that distinguishes CENP-A from canonical histone H3 [57-60]. In humans, this domain of CENP-A is both necessary and sufficient for targeting CENP-A to centromeres although not sufficient to promote kinetochore assembly [34, 61-64]. HJURP binds to the CATD of soluble CENP-A through its N-terminal CENP-A binding domain [43, 44, 57]. Once bound to CENP-A, HJURP is targeted to centromere regions through its interaction with the Mis18 complex (Mis18 α , Mis18 β , and Mis18BP1) [65-69], with Mis18BP1 binding directly to HJURP via its central domain [39, 70, 71]. In humans, Mis18BP1 also binds to the inner kinetochore protein CENP-C and through this interaction is targeted to the centromere [72-74]. Given that CENP-C localizes to centromeres through its direct interaction with CENP-A, this is one mechanism to ensure that CENP-A is exclusively deposited at centromeres in a self-propagating manner. Although the general paradigms of CENP-A propagation are conserved across vertebrates, the precise interactions vary between organisms. For example, in chicken and Xenopus, Mis18BP1 centromere localization does not depend on CENP-C, and instead involves a unique motif that is similar to the CENP-A binding domain in CENP-C to directly interact with CENP-A [75, 76]. This Mis18BP1-CENP-A interaction provides an alternative strategy for CENP-A to promote centromere specification and its self-propagation.

The timing for deposition of new CENP-A molecules is tightly coordinated with cell cycle progression in a way that is distinct from canonical H3 nucleosomes. Whereas H3 nucleosomes are deposited during S phase, the deposition of new CENP-A in vertebrates occurs in early G1 [61, 77-79]. To enact this temporal restriction, two kinases function as the gatekeepers of CENP-A deposition: Cyclin Dependent Kinase (CDK) and Polo-like Kinase 1 (PLK1). In human cells, CDK phosphorylation of Mis18BP1 results in decreased centromere localization and prevents its interaction with Mis18 α and Mis18 β [68, 74, 80, 81]. CDK phosphorylation of HJURP has also been shown to negatively impact its ability to localize to centromeres [81, 82]. Therefore,

deposition of new CENP-A molecules can only occur following chromosome segregation, when CDK activity is at its lowest. Conversely, the kinase activity of PLK1 acts to promote CENP-A deposition through phosphorylation of upstream targets, including the Mis18 complex [74]. Disrupting this temporal control of CENP-A deposition has negative effects on mitotic progression, highlighting the importance of this regulatory network on centromere function [74].

Once incorporated into centromere chromatin, CENP-A remains remarkably stable in rapidly dividing cells [83-85], with individual CENP-A molecules persisting over multiple divisions [78, 86]. It is currently not well understood how centromeric chromatin adapts to the changes in CENP-A occupancy throughout the cell cycle, although the presence of chromatin remodeling factors and transcription at the centromere may play a role [87]. Additionally, CENP-A has been shown to acquire post-translational modifications throughout the cell cycle that help coordinate different aspects of CENP-A function [88].

Given that maintenance of centromere CENP-A relies on passage through the cell cycle, it is interesting to consider how it is that the centromere is maintained in cells that are not actively dividing. Non-transformed cells and many cells in an organism can enter a state of proliferative hibernation termed quiescence or G0 [89]. Quiescent cells can persist in a non-dividing state for weeks, months, even decades and maintain their proliferative potential. Therefore, it is critical to retain the presence of centromere components to mark this chromosomal locus. In particular, as the centromere is specified primarily by the presence of CENP-A, if CENP-A is evicted from chromosomes, it would eliminate centromere identity and prevent future chromosome segregation. Thus, quiescent cells must maintain the protein components at centromeres over extended periods. One possibility to explain this is that CENP-A and other CCAN components are exceptionally stable. Indeed, studies in mouse oocytes blocking new CENP-A synthesis have indicated that CENP-A protein remains stable over extended periods [84]. Alternatively, quiescent cells may alter CCAN dynamics to promote the deposition of CENP-A and refresh centromere proteins. By studying this in quiescent cultured cells and starfish oocytes, our lab has found that CENP-A undergoes a gradual, but consistent turnover at centromeres [90]. This dynamic behavior requires

the canonical CENP-A deposition machinery and may be promoted by ongoing transcription at centromeres to evict CENP-A molecules providing “holes” to enable new CENP-A deposition. The failure to deposit new CENP-A molecules in quiescent cells results in defective chromosome segregation when quiescent cells return to proliferation. In contrast to CENP-A, some other CCAN components are lost from centromeres in quiescent cells [90] suggesting the potential for additional changes in centromere organization in non-dividing cells. Interestingly, CENP-A is differentially maintained in diverse cell types in a manner that correlates with replicative potential [91]. For example, in work by Swartz et. al. [90], terminally differentiated cells, such as adult mouse cardiomyocytes, lose CENP-A at their centromeres. In contrast, adult mouse hepatocytes, which maintain the capacity to divide and regenerate, maintain centromeric CENP-A. Together, this supports the importance of CENP-A maintenance at the centromere to facilitate cell division.

Together, the dynamic localization and maintenance of CENP-A molecules in the early cell cycle stages and in non-dividing cells provides the foundation onto which the remaining kinetochore architecture is assembled in the subsequent cell cycle phases. Importantly, the large number of CENP-A nucleosomes deposited during G1 represents a substantial change to centromere chromatin, and likely also requires the subsequent recruitment of CENP-A binding factors.

Retaining centromere assembly during Replication and Transcription

As the cell cycle proceeds, the next challenge that centromeres face is the need to replicate the underlying DNA during S phase (Figure 1B). DNA replication requires the displacement of existing nucleosomes to allow for the passage of the replication fork and DNA synthesis [92]. Therefore, the centromere must accommodate the passage of the replication fork while maintaining its identity and function. CENP-A histones are stably inherited across multiple divisions, suggesting that centromere nucleosomes are retained through DNA replication. Given that no new CENP-A molecules are deposited during S phase, CENP-A nucleosomes must be recycled and conservatively distributed between the newly replicated sister centromeres [78]. It is thought that the gaps left due to the 2-fold reduction in the number of CENP-A nucleosomes are instead filled with

histone H3.3 [93]. Interestingly, the position and distribution of CENP-A nucleosomes within the centromere are maintained, implicating a positional memory of CENP-A nucleosomes within the centromere [93, 94], although the CENP-A domain can migrate slightly across divisions [95]. This raises the question as to how CENP-A nucleosomes are distributed during DNA replication. Recent work has suggested that the distribution of CENP-A nucleosomes involves HJURP, which transiently associates with CENP-A during S phase, in conjunction with Mcm2, a helicase involved in DNA replication [96]. This parallels the mechanism by which canonical H3 nucleosomes are deposited in non-centromere chromatin, which also involves Mcm2 and distinct H3/H4 nucleosome chaperones [92].

The passage of a replication fork through centromere DNA has the potential to be highly destabilizing. In fact, DNA replication has been shown to evict CENP-A nucleosomes erroneously incorporated at chromosome arms [94]. So how is it that CENP-A nucleosomes remain enriched at the centromere following DNA replication? A key distinction of CENP-A at centromeres is the presence of the associated CCAN components, which remain assembled on CENP-A nucleosomes during S phase/G2 [94]. Recent work has suggested that the CCAN contributes to centromere specific recycling/retention of CENP-A nucleosomes specifically at centromeres. Specifically, the rapid depletion of CENP-C in early S phase results in a significant loss of CENP-A nucleosomes at centromeres following a round of DNA replication [94]. Together, this supports the idea that the CCAN plays a critical role in supporting the reincorporation of CENP-A nucleosomes at centromeres during DNA replication. How the CCAN remains assembled on CENP-A nucleosomes and interacts with the replication machinery remains an important open question.

Importantly, DNA polymerase is not the only enzyme that acts at centromeres and has the ability to disrupt the underlying chromatin. Despite the fact that centromeres lack coding genes, centromeres are actively transcribed. RNA polymerase II has been found at centromeres in mitosis [97] and centromeric chromatin contains histone modifications, such as H3K4me2, that are permissive for transcription [98, 99]. Importantly, centromere transcription occurs at low levels throughout the cell cycle, including during mitosis [97, 100]. The presence of ongoing centromere transcription

during mitosis may play an active role in centromere function [87, 101]. For example, centromere transcription by RNA polymerase could destabilize nucleosomes and CCAN components. In fact, centromere transcription has been implicated in promoting the turnover of CENP-A [90]. Alternatively, the presence of RNA polymerase II at mitotic centromeres may be due to the persistence of cohesin at centromere regions [102], but may not reflect an active role [103].

Together, the CCAN and CENP-A nucleosomes maintain a reciprocal relationship to ensure centromere identity is maintained despite drastic changes to the DNA structure in S phase. Whereas the CCAN components require the presence of CENP-A within the centromere for their localization, these proteins also help maintain CENP-A localization, including during DNA replication. Additionally, although replication and transcription represent a challenge to centromere identity, it also performs a beneficial role in centromere maintenance by preventing the accumulation of CENP-A at ectopic chromosome regions.

The Inner Kinetochore

The inner kinetochore, or CCAN, is assembled specifically at the centromere due to its interactions with CENP-A. A distinguishing characteristic of the CCAN is that it remains present at the centromere throughout the cell cycle. Therefore, once assembled the interactions between CCAN components must be robust enough so that the entire assembly is maintained at every cell cycle state. Indeed, each of the CCAN sub-complexes (CENP-C, CENP-L/N, CENP-H/I/K/M, CENP-T/W/S/X, and CENP-O/P/Q/U/R) [104-106] has a distinct role in CCAN organization, and together form a hierarchy of interactions that promote the stability of this protein complex (Figure 2A).

CENP-C plays a central role in kinetochore organization as it interacts with multiple components of the inner kinetochore, including CENP-A, and can recruit components of the outer kinetochore directly [107-111]. In fact, for some organisms, such as *D. melanogaster* and *C. elegans*, CENP-C is the only identified CCAN component [26, 31, 112]. Based on its network of interactions, CENP-C is considered to be an organizing scaffold within the CCAN [107, 108, 113-115]. In addition to CENP-C, CENP-N is the only other protein within the kinetochore that can interact directly with

CENP-A [114, 116]. CENP-C and CENP-N interact with CENP-A in a distinct manner [114], providing the potential for both proteins to interact with CENP-A simultaneously. Within the CCAN, CENP-N forms an obligate complex with CENP-L where together they interact directly with CENP-HIKM and CENP-C [108, 115, 117]. The CENP-HIKM complex also sits at the core of the CCAN as it has been shown to directly interact with CENP-C, CENP-LN, and CENP-TWSX [104, 115, 118]. The CENP-TWSX complex contains histone fold domains that promote centromere recruitment by conferring DNA binding abilities to this complex in addition to facilitating protein interactions [32, 107, 108, 119, 120]. Like CENP-C, the CENP-TWSX complex interacts directly with proteins of the outer kinetochore and plays an important role in their regulated recruitment in mitosis [32, 121-123]. Finally, the CENP-OPQUR complex is recruited to the kinetochore through its interaction with CENP-C/HIKM/LN [118]. The CENP-OPQUR complex is the most peripheral CCAN component, as depletion of this complex does not affect the localization of the other CCAN components [106, 108, 118]. The CENP-OPQUR complex facilitates functions that are important for mitotic progression, including recruitment of CENP-E and PLK1, and has been suggested to interact directly with microtubules [118, 124-130]. The interdependencies that exist between the CCAN components enable this complex of proteins to adapt to changes that occur in the centromere so that they remain assembled throughout the cell cycle.

The Dynamic Nature of the Inner Kinetochore

The naming of the CCAN as “constitutive” centromere factors can imply that they are unchanged throughout the cell cycle. However, despite their constitutive localization, the association of individual components of the CCAN to the centromere is instead quite dynamic. This includes cell cycle changes in the abundance of individual CCAN components [131-133], the timing by which new proteins are incorporated at centromeres [85, 132, 134, 135], and changes in the physical associations and functional relationships between the CCAN sub-complexes (Figure 2B).

Of the CCAN proteins, CENP-C and CENP-N provide important examples of these distinct cell cycle behaviors. Although CENP-C is always present at centromeres, it displays dynamic and changing properties. First, CENP-C levels at kinetochores

remain relatively constant throughout the cell cycle but increase upon entry into mitosis [133]. The molecular requirements for CENP-C localization to centromeres also differ between interphase and mitosis. In interphase, CENP-C localization depends on its interaction with CENP-HIKM and CENP-L/N as depletion of either complex result in the loss of CENP-C at the centromere [108, 136]. In contrast, during mitosis CENP-C can localize to kinetochores in the absence of CENP-LN and CENP-HIKM. Given the important role that CENP-C plays in CCAN organization, this change in its localization requirements suggests that there is a cell cycle-dependent reorganization of the CCAN. Recent work in chicken cells found that CDK-dependent phosphorylation of CENP-C in mitosis promotes its interaction with CENP-A [137, 138]. Thus, the requirements for CENP-C localization change throughout the cell cycle, such that its interaction with other CCAN subunits is important in interphase, but its interaction with CENP-A becomes more critical in mitosis. This change may be important in adapting the CCAN for the recruitment of outer kinetochore components, or rewiring CCAN organization to accommodate interphase events at centromeres.

The CENP-L/N complex also plays a critical role in the organization of the CCAN and displays distinct cell cycle dependent behaviors. CENP-N interacts directly with the CATD of CENP-A [116, 139]. This interaction is mediated by the N-terminus of CENP-N, which is both necessary and sufficient for CENP-N localization to the kinetochore during interphase, but not mitosis [108, 116]. The C-terminus of CENP-N, which mediates its direct interaction with CENP-L and indirectly with the CCAN, is critical for the mitotic localization of CENP-N [108, 140]. Therefore, the localization of CENP-N relies more on its interactions with the CCAN in mitosis, a behavior that directly contrasts with that of CENP-C. Additionally, CENP-N displays distinct localization behaviors throughout the cell cycle. Unlike CENP-C, CENP-N is enriched at kinetochores specifically during S-phase and then dissociates during the G2/M transition such that its levels are low in mitosis [141, 142]. Finally, CENP-N has been shown to associate dynamically with centromeres during G1 and early S-phase [141]. However, the basis and the significance of this change in CENP-N kinetochore levels remain unclear. One aspect of CENP-N behavior that may contribute to this difference in CENP-N association is the fact that CENP-N loading appears to be dependent on

chromatin state. In vitro assays suggest that CENP-N more readily interacts with CENP-A nucleosomes present in open chromatin where the RG loop is exposed, as compared to a more compact chromatin state where the RG loop is not accessible [143]. Therefore, the underlying changes in chromatin state that occur throughout the cell cycle may affect the ability of CENP-N to remain stably bound at kinetochores.

The fact that CENP-N and CENP-C, two components essential for CCAN recruitment and CENP-A recognition, have distinct cell cycle dependent behaviors suggests the CCAN complex as a whole undergoes substantial changes. However, it remains unclear how these behaviors and differences in centromere associations affect CCAN organization and function. The underlying mechanisms that contribute to the varying association behaviors within the CCAN have not been elucidated, but post-translational modifications, such as phosphorylation, may play a role [137, 138, 144].

Role of CENP-B in Centromere Function

Given the critical requirement to maintain the assembly of inner kinetochore components at centromeres, it is important to highlight the potential contributions that CENP-B plays in maintaining centromere function. CENP-B is the only protein within the inner kinetochore that binds to a specific DNA sequence - the 17 bp CENP-B box present in the alpha-satellite sequences [12]. The functional importance of CENP-B at centromeres has been debated given that it is not essential for viability in either cell culture or mouse models and that it is absent from both neocentromeres and the Y chromosome [145]. However, chromosomes that lack CENP-B have been shown to mis-segregate more frequently [146] suggesting that CENP-B plays important roles in centromere function. Within the centromere, CENP-B has been shown to interact directly with CENP-A, and through this interaction increases the stability of CENP-A in reconstituted nucleosomes [147-149]. CENP-B and CENP-C also interact directly. This interaction has been proposed to serve as an additional recruitment mechanism for CENP-C independently from CENP-A, as depletion of CENP-B results in a decrease of CENP-C at kinetochores [147]. In fact, in the absence of CENP-A, CENP-B is sufficient to maintain CENP-C at the kinetochore and facilitate proper chromosome segregation [150], at least for a short period of time. Recent work has supported a model in which

CENP-B becomes more important at centromeres in the absence of CENP-A [146]. Therefore, the presence of CENP-B at the centromere may help to ensure centromere function in conjunction with CENP-A by providing an additional mechanism to tether the inner kinetochore to the centromere.

Higher order centromere structure and changes to chromatin state

A major consideration for understanding the assembly and dynamics of the kinetochore is the higher order organization of the underlying centromere DNA and chromatin (Figure 1C). The centromere does not exist as a linear segment of DNA. Instead, centromere DNA displays a complex three-dimensional organization, similar to other chromosomal loci. Within the centromere, CENP-A nucleosomes are interspersed with H3 nucleosomes [93, 151, 152]. At human centromeres, quantification of CENP-A molecules indicate that ~100 CENP-A nucleosomes are dispersed throughout the centromere (likely over ~100 kb) at a ratio of ~1:25 relative to H3 [153]. Therefore, to enable kinetochore assembly and proper microtubule attachment in mitosis, as chromosomes are compacted and condensed at mitotic entry, the centromere must be organized to position CENP-A nucleosomes together facing away from the chromosome to facilitate kinetochore assembly. How this occurs remains unclear, but multiple models have been proposed [15, 154, 155]. Although the mechanism and precise structure of centromere chromatin upon entry into mitosis is not known, CENP-A and the CCAN confer properties to centromere chromatin that are important for enacting its mitotic function. For example, in *C. elegans*, loss of CENP-A causes defects in chromosome structure and affects the dynamics of chromosome condensation [156]. A similar behavior has also been shown in human cells, where CENP-A chromatin withstands unfolding to a higher degree than the surrounding heterochromatin and mitotic CENP-A enriched chromatin can withstand unfolding to a higher degree than interphase chromatin [151]. This suggests that the presence of CENP-A distinctly impacts chromatin architecture at the centromere, and that this may confer an added rigidity that allows the centromere to withstand mitotic forces.

The presence and association of the CCAN at the centromere also contributes to centromere rigidity [151]. Reciprocally, the changes that happen to centromere

chromatin upon mitotic entry can affect the organization of the CCAN as demonstrated by changes in CENP-C interdependencies and the effect that CENP-A chromatin state has on CENP-N binding *in vitro* [143]. Efforts to reconstitute the kinetochore suggest that two CCAN complexes (containing CENP-C/HIKM/LN) are capable of binding a single CENP-A nucleosome, forming a symmetric structure [115]. Recent structural work reconstituting the core centromere nucleosome complex (CCNC) consisting of CENP-A, CENP-N, and CENP-C from human and chickens has provided insights into the potential changes in CCAN associations at centromeres between interphase and mitosis [138, 142]. In work reconstituting the human CCNC, Allu and colleagues [142] observed two CCNC populations - one in which two CENP-C and CENP-N molecules bind to CENP-A, and one in which only a single molecule of CENP-N and two molecules of CENP-C are bound to the CENP-A nucleosome. Given that this corresponds to the differing ratios of CENP-C and CENP-N present at centromeres in interphase and mitosis, it is possible that these two structures represent the cell cycle differences in CCAN centromere associations. Alternatively, studies analyzing centromere proteins from chicken cells support an alternative model for the changes in CCNC assembly throughout the cell cycle [138]. In this work, the authors argue that CENP-C and CENP-N do not bind CENP-A simultaneously and instead do so in a mutually exclusive and cell cycle-dependent manner. In this model, CENP-N binds directly to CENP-A nucleosomes during interphase, whereas CENP-C does so in mitosis. A key distinction between these two studies is the CENP-C fragments used to reconstitute the CCNC. Ariyoshi and colleagues utilized a longer CENP-C C-terminal fragment, which resulted in structures where CENP-C obscured the RG loop in CENP-A preventing CENP-N binding. Importantly, CENP-C was phosphorylated in these experiments, modeling mitotic CENP-C [137]. Together, these studies provide molecular insights into the mechanisms that may underlie CCAN reorganization during mitosis. However, future studies will be required to determine the physiological relevance of these two proposed mechanisms, including structural studies with full length CENP-C and CENP-N. Overall, progression into mitosis requires that the centromere adapt a structure that will withstand the forces transmitted during chromosome segregation and provide a platform for outer kinetochore assembly.

The Outer Kinetochore

The next critical phase in kinetochore assembly occurs during entry into mitosis when the CCAN directs recruitment of the outer kinetochore, which in turn interacts directly with spindle microtubules. Outer kinetochore assembly is a highly controlled process that occurs through multiple regulatory layers (Figure 3). First, the core components of the outer kinetochore are recruited upon mitotic entry in a manner that depends upon the phospho-regulation of inner kinetochore components to create a platform for outer kinetochore assembly. Second, kinetochores alter their functional properties at different stages within mitosis by altering outer kinetochore composition, adding or removing specific components to enable distinct activities. Finally, under circumstances where kinetochores fail to interact with microtubules, the outer kinetochore seeds formation of an extended fibrous corona structure.

The core of the outer kinetochore is made up of 10 proteins that assemble into the three sub-complexes Knl1, Mis12, and Ndc80 - also referred to as the KMN network [157-167]. In particular, the four-subunit Ndc80 complex acts as the primary kinetochore receptor for microtubule binding [157, 162, 168]. The KMN network is recruited in a stepwise manner throughout the cell cycle via two distinct pathways mediated by CENP-C and CENP-T. Each of these pathways is sufficient to recruit the outer kinetochore in vertebrates [41, 133, 169]. The coordinated assembly of the outer kinetochore relies on the activity of cell cycle-associated kinases and cell cycle-specific cellular events [170]. The Mis12 complex and Knl1 are the first to associate to the kinetochore starting in late S phase [133, 171] (Figure 3). This recruitment is mediated through the direct interaction between Mis12 and CENP-C [109-111, 121, 172]. Following their initial association, Mis12 and Knl1 levels gradually increase and are maximally enriched at metaphase [133]. The recruitment of Mis12 is facilitated by phosphorylation of Mis12 by the mitotic kinase Aurora B, which results in a conformational change that promotes the interaction between Mis12 and CENP-C [109, 173]. Upon entry into mitosis and onset of nuclear envelope breakdown, Ndc80 is recruited to the kinetochore by two distinct mechanisms [133, 171] (Figure 3). First, the Mis12 complex, via its interaction with CENP-C, recruits a single molecule of Ndc80

upon mitotic entry [174, 175]. Second, CENP-T can directly bind to two molecules of Ndc80 [123, 176, 177], following phosphorylation of two distinct sites of CENP-T by mitotic CDK1 [121, 123, 169, 176]. Additionally, CENP-T has been shown to independently recruit Mis12 following CDK phosphorylation, therefore indirectly recruiting an additional molecule of Ndc80 [169, 176]. Although both CENP-C and CENP-T recruit outer kinetochore proteins, work in chicken cells has suggested that CENP-T plays a more critical role in the recruitment of the KMN network. In this work, researchers found that expressing a CENP-C mutant that lacks its Mis12 complex binding domain did not affect mitotic progression in chicken cells. In contrast, CENP-T mutants that lack the ability to bind the Ndc80 or Mis12 complexes in the absence of endogenous CENP-T resulted in cell death [172]. Further work is needed to determine whether this is true in mammalian cells.

Although Ndc80 is essential for mediating kinetochore/microtubule interactions, additional microtubule-binding proteins are recruited to the outer kinetochore throughout mitosis to strengthen and stabilize this interaction. The first of these proteins is the Ska1 complex, which functions to enhance the microtubule binding activity of Ndc80 [178-180]. Recruitment of the Ska1 to kinetochores requires Ndc80 and the presence of kinetochore/microtubule interactions [181, 182]. The Ska1 complex is first recruited during prometaphase and accumulates at kinetochores throughout chromosome congression [133, 183, 184]. Once kinetochores have established stable attachments to microtubules and are properly bi-oriented, the Astrin-SKAP protein complex is recruited to kinetochores [185-188]. The Astrin-SKAP localizes to bi-oriented kinetochores at metaphase and persists into anaphase, such that this complex is thought to play an important role in stabilizing proper kinetochore/microtubule attachments [186, 188-190]. Additional proteins also associate with kinetochores to regulate spindle microtubule dynamics and facilitate microtubule organization (see [191, 192]).

Finally, under conditions where kinetochores fail to generate microtubule interactions, kinetochores undergo a dramatic additional assembly step and physical transformation where they can expand to form a crescent-like structure on its surface called the fibrous corona [193-196]. This fibrous structure forms at kinetochores that lack attached microtubules and compacts once kinetochore-microtubule interactions are

established [193, 197]. This process involves a complex network of proteins, including the motor proteins CENP-E [198], the dynein/dynactin motor complex [199], the microtubule binding protein CENP-F [200], and the RZZ (Rod-ZW10-Zwilch) complex [201, 202]. This structure is thought to function to increase the likelihood of microtubule capture and promote Spindle Assembly Checkpoint (SAC) signaling in the presence of unattached kinetochores [195, 197]. The formation of the fibrous corona also precedes the formation of end-on attachments in unperturbed cells [195]. The molecular mechanisms underlying kinetochore expansion have implicated the RZZ complex and Spindly protein, which together have been shown to oligomerize *in vitro* [203-205]. Depletion of Spindly or RZZ complex components prevents kinetochore expansion in the absence of microtubules [203-205]. It remains unclear how the RZZ complex is tethered to kinetochores, although Knl1 and Bub1 have been proposed to contribute to this process [203, 206]. Once end-on attachments are established, the majority of the fibrous corona components are stripped from the kinetochore in a dynein-dependent manner, resulting in compaction of the kinetochore structure [207-211]. However, CENP-E and CENP-F perform additional roles in chromosome segregation and remain at the kinetochore until anaphase [212, 213]. This general process of kinetochore compaction is important for proper mitotic progression as disruption of this process results in erroneous microtubule attachments as well as chromosome mis-segregation [14, 195, 205, 211]. Therefore, this regulated morphological change in the outer kinetochore is critical to ensure faithful chromosome segregation.

Ensuring the fidelity of mitosis

In mitosis, the kinetochore functions as a scaffold to recruit signaling proteins and monitor kinetochore-microtubule attachments. For faithful chromosome segregation to occur, sister kinetochores must attach to microtubules from opposing spindle poles to establish bi-orientation. Through a process called error correction, kinetochore-microtubule attachments are monitored by the activity of kinetochore-associated kinases and phosphatases [214]. Together, the dynamic association of these factors modulates kinetochore function in mitosis to not only form interactions with microtubules but also to ensure that kinetochores undergo proper chromosome segregation.

Although changes in kinetochore composition can dynamically alter kinetochore function, some kinetochore activities are modulated without changes to kinetochore assembly state. For example, Aurora B kinase regulates kinetochore-microtubule interactions by destabilizing the interaction between components of the KMN and microtubules. Ndc80 is a critical substrate of Aurora B within the KMN, where Aurora B phosphorylates the regulatory N-terminal tail of Ndc80, decreasing its affinity for microtubules and therefore disrupting erroneous interactions with microtubules [157, 168, 215]. Similarly, Aurora B phosphorylates the Ska1 complex reducing its binding affinity for microtubules [179, 216]. Once proper kinetochore-microtubule attachments have been achieved, dephosphorylation of these substrates is mediated by the activity of two mitotic phosphatases: PP1 and PP2A-B56 [217-219]. Once dephosphorylated, Ndc80 and Ska1 are able to establish stable interactions with microtubules [157, 215, 220-222].

Accurate chromosome segregation is also monitored by an additional set of proteins that collectively make up the Spindle Assembly Checkpoint (SAC), which associate preferentially with unattached kinetochores. The key function of the SAC is to prevent anaphase onset until all chromosomes are stably attached to spindle microtubules [223]. This checkpoint is regulated by the activity of two complexes called the mitotic checkpoint complex (MCC), which prevents anaphase onset, and the Anaphase Promoting Complex/Cyclosome (APC/C), an E3 ubiquitin ligase that induces anaphase onset and mitotic exit through the targeted degradation of important mitotic proteins [224-226]. In addition to its role in mediating kinetochore-microtubule interactions, the KMN acts as a scaffold for the recruitment of spindle assembly checkpoint proteins. For example, the MCC is recruited to the kinetochore by phosphorylation of Knl1 by Mps1, a kinase that is critical for initiating the SAC response, as Mps1 phosphorylation activity is required for the recruitment of downstream SAC components [227, 228]. Once stable microtubule attachments have been achieved and sister kinetochores are bi-oriented, the SAC is inactivated to allow for mitotic progression [229]. The inactivation of the SAC involves the disassembly of the MCC, removal of SAC proteins by dynein, and dephosphorylation of kinetochore and SAC

components by the PP1, localized to the kinetochore through its interaction with Knl1 and PP2A-B56 [223, 230].

In summary, the kinetochore continues to be a dynamic structure even once it is fully assembled in mitosis. Through the coordinated recruitment of regulatory proteins, the kinetochore is able to perform the many functions required to facilitate proper chromosome segregation.

Conclusions

The kinetochore is a dynamic molecular protein complex that changes in composition and structure throughout the cell cycle to enable the kinetochore to perform distinct functions at each cell cycle stage. During interphase, the minimal kinetochore structure present at the centromere, the CCAN, performs the important function of maintaining centromere identity through its role in maintaining CENP-A at the centromere. As the cell progresses into mitosis, the CCAN reorganizes to enable the recruitment of the outer kinetochore. Once in mitosis, the kinetochore functions to recruit proteins required to facilitate the formation of kinetochore-microtubule interactions, promote checkpoint signaling to ensure proper attachments are made, and expand to enable kinetochore-microtubule interactions to be formed. Given the importance of the regulation of these protein-protein interactions, future work in understanding the proteins and regulatory mechanisms that enable kinetochore function and disassembly will provide a more comprehensive understanding of the kinetochore's role in chromosome segregation and centromere maintenance.

Acknowledgements

The authors thank Océane Marescal and Ally Nguyen for critical reading of the manuscript, and B. Knowles and members of the Cheeseman lab for their support. The work in the Cheeseman lab is supported by grants from the NIH/National Institute of General Medical Sciences (R35GM126930), the Moore Foundation, the National Science Foundation (2029868), and a Pilot award from the Global Consortium for Reproductive Longevity and Equity (GCRLE), and an NSF graduate research fellowship to APN.

Declarations of interest: none

Figure Legends

Figure 1. Dynamic events throughout the cell cycle affect centromere structure.

A. In G1, new CENP-A molecules are deposited at centromeres. This process is tightly regulated to ensure this only happens once each cell division cycle. **B.** During S phase, centromere DNA must be replicated. Because no new molecules of CENP-A are deposited, this involves the distribution of previously deposited CENP-A molecules such that CENP-A molecules are diluted and H3.3 histones are incorporated as placeholders to fill in gaps. Throughout S-phase, the CCAN remains assembled upon centromeric CENP-A and helps to retain CENP-A centromere following passage of the replication fork. **C.** In late G2, chromatin is condensed to form the distinct chromosome structures observed in mitosis. In addition, centromere DNA must reorganize to orient the centromere proteins to recruit of outer kinetochore components and binding to spindle microtubules. **D.** In mitosis, following nuclear envelope breakdown and phosphorylation by CDK1, outer kinetochore proteins are recruited to the centromere completing the assembly of the kinetochore and enable kinetochores to interact directly with spindle microtubules and facilitate chromosome segregation.

Figure 2. Organization of the Inner Kinetochore. **A.** The Inner kinetochore is made up of 5 distinct groups: CENP-C, CENP-N/L, CENP-H/I/K/M, CENP-T/W/S/X, and CENP-O/P/Q/U/R. These sub complexes are largely interdependent for their localization to the centromere. Direct interactions between inner kinetochore sub-complexes, as defined by biochemical reconstitutions and functional dependencies, are represented by arrows. **B.** Although all components of the CCAN are present at centromeres constitutively, these proteins undergo reorganization and changes throughout the cell cycle. This model represents snapshots of inner kinetochore organization at distinct cell cycle phases.

Figure 3. Organization and assembly of the outer kinetochore. Recruitment of outer kinetochore components occurs in a step-wise manner. In late G2, Mis12/Knl1 are recruited through their interaction with CENP-C. This interaction is enabled by Aurora B

phosphorylation of the Mis12 complex. Ndc80 is then recruited to the kinetochore following entry into mitosis. Once in mitosis, additional proteins are transiently recruited to the kinetochore to enable the multiple kinetochore functions in mitosis. In the absence of a microtubule interaction, proteins involved in the spindle assembly checkpoint (MCC) and the fibrous corona (RZZ, spindly, CENP-E, CENP-F, dynein/dynactin) are recruited to kinetochore's. As kinetochore-microtubule interactions are established, the Ska1 complex and Astrin/SKAP are recruited to promote Ndc80 binding to microtubules. Once stable kinetochore-microtubule interactions are established and are properly bi-oriented, the spindle assembly checkpoint is satisfied and chromosome segregation occurs. CENP-E and CENP-F perform additional roles in chromosome segregation and remain at the kinetochore until anaphase.

References

- [1] I.M. Cheeseman, The kinetochore, *Cold Spring Harb Perspect Biol* 6(7) (2014) a015826.
- [2] A. Musacchio, A. Desai, A Molecular View of Kinetochore Assembly and Function, *Biology (Basel)* 6(1) (2017).
- [3] J. de la Cruz, K. Karbstein, J.L. Woolford, Jr., Functions of ribosomal proteins in assembly of eukaryotic ribosomes *in vivo*, *Annu Rev Biochem* 84 (2015) 93-129.
- [4] K.L. McKinley, I.M. Cheeseman, The molecular basis for centromere identity and function, *Nat Rev Mol Cell Biol* 17(1) (2016) 16-29.
- [5] P.S. Maddox, K. Oegema, A. Desai, I.M. Cheeseman, "Holo"er than thou: Chromosome segregation and kinetochore function in *C. elegans*, *Chromosome Res* 12(6) (2004) 641-53.
- [6] S. Kit, Equilibrium sedimentation in density gradients of DNA preparations from animal tissues, *J Mol Biol* 3 (1961) 711-6.
- [7] M.E. Aldrup-Macdonald, B.A. Sullivan, The past, present, and future of human centromere genomics, *Genes (Basel)* 5(1) (2014) 33-50.
- [8] B. Vissel, K.H. Choo, Human alpha satellite DNA--consensus sequence and conserved regions, *Nucleic Acids Res* 15(16) (1987) 6751-2.
- [9] L. Manuelidis, Chromosomal localization of complex and simple repeated human DNAs, *Chromosoma* 66(1) (1978) 23-32.
- [10] L. Manuelidis, Complex and simple sequences in human repeated DNAs, *Chromosoma* 66(1) (1978) 1-21.
- [11] J.S. Waye, H.F. Willard, Human beta satellite DNA: genomic organization and sequence definition of a class of highly repetitive tandem DNA, *Proc Natl Acad Sci U S A* 86(16) (1989) 6250-4.
- [12] H. Masumoto, H. Masukata, Y. Muro, N. Nozaki, T. Okazaki, A human centromere antigen (CENP-B) interacts with a short specific sequence in alphoid DNA, a human centromeric satellite, *J Cell Biol* 109(5) (1989) 1963-73.
- [13] Y. Muro, H. Masumoto, K. Yoda, N. Nozaki, M. Ohashi, T. Okazaki, Centromere protein B assembles human centromeric alpha-satellite DNA at the 17-bp sequence, CENP-B box, *J Cell Biol* 116(3) (1992) 585-96.
- [14] D. Drpic, A.C. Almeida, P. Aguiar, F. Renda, J. Damas, H.A. Lewin, D.M. Larkin, A. Khodjakov, H. Maiato, Chromosome Segregation Is Biased by Kinetochore Size, *Curr Biol* 28(9) (2018) 1344-1356 e5.
- [15] T. Fukagawa, W.C. Earnshaw, The centromere: chromatin foundation for the kinetochore machinery, *Dev Cell* 30(5) (2014) 496-508.
- [16] W.H. Shang, T. Hori, A. Toyoda, J. Kato, K. Popendorf, Y. Sakakibara, A. Fujiyama, T. Fukagawa, Chickens possess centromeres with both extended tandem repeats and short non-tandem-repetitive sequences, *Genome Res* 20(9) (2010) 1219-28.
- [17] D.P. Locke, L.W. Hillier, W.C. Warren, K.C. Worley, L.V. Nazareth, D.M. Muzny, S.P. Yang, Z. Wang, A.T. Chinwalla, P. Minx, M. Mitreva, L. Cook, K.D. Delehaunty, C. Fronick, H. Schmidt, L.A. Fulton, R.S. Fulton, J.O. Nelson, V. Magrini, C. Pohl, T.A. Graves, C. Markovic, A. Cree, H.H. Dinh, J. Hume, C.L. Kovar, G.R. Fowler, G. Lunter, S. Meader, A. Heger, C.P. Ponting, T. Marques-Bonet, C. Alkan, L. Chen, Z. Cheng, J.M. Kidd, E.E. Eichler, S. White, S. Searle, A.J. Vilella, Y. Chen, P. Flicek, J. Ma, B. Raney, B. Suh, R. Burhans, J. Herrero, D. Haussler, R. Faria, O.

Fernando, F. Darre, D. Farre, E. Gazave, M. Oliva, A. Navarro, R. Roberto, O. Capozzi, N. Archidiacono, G. Della Valle, S. Purgato, M. Rocchi, M.K. Konkel, J.A. Walker, B. Ullmer, M.A. Batzer, A.F. Smit, R. Hubley, C. Casola, D.R. Schrider, M.W. Hahn, V. Quesada, X.S. Puente, G.R. Ordóñez, C. López-Otín, T. Vinar, B. Brejova, A. Ratan, R.S. Harris, W. Miller, C. Kosiol, H.A. Lawson, V. Taliwal, A.L. Martins, A. Siepel, A. Roychoudhury, X. Ma, J. Degenhardt, C.D. Bustamante, R.N. Gutenkunst, T. Mailund, J.Y. Dutheil, A. Hobolth, M.H. Schierup, O.A. Ryder, Y. Yoshinaga, P.J. de Jong, G.M. Weinstock, J. Rogers, E.R. Mardis, R.A. Gibbs, R.K. Wilson, Comparative and demographic analysis of orang-utan genomes, *Nature* 469(7331) (2011) 529-33.

[18] F.M. Piras, S.G. Nergadze, E. Magnani, L. Bertoni, C. Attolini, L. Khoriauli, E. Raimondi, E. Giulotto, Uncoupling of satellite DNA and centromeric function in the genus *Equus*, *PLoS Genet* 6(2) (2010) e1000845.

[19] C.M. Wade, E. Giulotto, S. Sigurdsson, M. Zoli, S. Gnerre, F. Imsland, T.L. Lear, D.L. Adelson, E. Bailey, R.R. Bellone, H. Blocker, O. Distl, R.C. Edgar, M. Garber, T. Leeb, E. Mauceli, J.N. MacLeod, M.C. Penedo, J.M. Raison, T. Sharpe, J. Vogel, L. Andersson, D.F. Antczak, T. Biagi, M.M. Binns, B.P. Chowdhary, S.J. Coleman, G. Della Valle, S. Fryc, G. Guerin, T. Hasegawa, E.W. Hill, J. Jurka, A. Kialainen, G. Lindgren, J. Liu, E. Magnani, J.R. Mickelson, J. Murray, S.G. Nergadze, R. Onofrio, S. Pedroni, M.F. Piras, T. Raudsepp, M. Rocchi, K.H. Roed, O.A. Ryder, S. Searle, L. Skow, J.E. Swinburne, A.C. Syvanen, T. Tozaki, S.J. Valberg, M. Vaudin, J.R. White, M.C. Zody, P. Broad Institute Genome Sequencing, T. Broad Institute Whole Genome Assembly, E.S. Lander, K. Lindblad-Toh, Genome sequence, comparative analysis, and population genetics of the domestic horse, *Science* 326(5954) (2009) 865-7.

[20] K. Sanyal, M. Baum, J. Carbon, Centromeric DNA sequences in the pathogenic yeast *Candida albicans* are all different and unique, *Proc Natl Acad Sci U S A* 101(31) (2004) 11374-9.

[21] K.F. Sullivan, M. Hechenberger, K. Masri, Human CENP-A contains a histone H3 related histone fold domain that is required for targeting to the centromere, *J Cell Biol* 127(3) (1994) 581-92.

[22] B.J. Buchwitz, K. Ahmad, L.L. Moore, M.B. Roth, S. Henikoff, A histone-H3-like protein in *C. elegans*, *Nature* 401(6753) (1999) 547-8.

[23] K. Takahashi, E.S. Chen, M. Yanagida, Requirement of Mis6 centromere connector for localizing a CENP-A-like protein in fission yeast, *Science* 288(5474) (2000) 2215-9.

[24] S. Henikoff, K. Ahmad, J.S. Platero, B. van Steensel, Heterochromatic deposition of centromeric histone H3-like proteins, *Proc Natl Acad Sci U S A* 97(2) (2000) 716-21.

[25] S. Stoler, K.C. Keith, K.E. Curnick, M. Fitzgerald-Hayes, A mutation in CSE4, an essential gene encoding a novel chromatin-associated protein in yeast, causes chromosome nondisjunction and cell cycle arrest at mitosis, *Genes Dev* 9(5) (1995) 573-86.

[26] I.A. Drinnenberg, D. deYoung, S. Henikoff, H.S. Malik, Recurrent loss of CenH3 is associated with independent transitions to holocentricity in insects, *eLife* 3 (2014).

[27] A.P. Navarro, I.M. Cheeseman, Chromosome Segregation: Evolving a Plastic Chromosome-Microtubule Interface, *Curr Biol* 30(4) (2020) R174-R177.

[28] B. Akiyoshi, K. Gull, Discovery of unconventional kinetochores in kinetoplastids, *Cell* 156(6) (2014) 1247-1258.

[29] N. Cortes-Silva, J. Ulmer, T. Kiuchi, E. Hsieh, G. Cornilleau, I. Ladid, F. Dingli, D. Loew, S. Katsuma, I.A. Drinnenberg, CenH3-Independent Kinetochore Assembly in Lepidoptera Requires CCAN, Including CENP-T, *Curr Biol* 30(4) (2020) 561-572 e10.

[30] A.P. Senaratne, H. Muller, K.A. Fryer, M. Kawamoto, S. Katsuma, I.A. Drinnenberg, Formation of the CenH3-Deficient Holocentromere in Lepidoptera Avoids Active Chromatin, *Curr Biol* 31(1) (2021) 173-181 e7.

[31] I.M. Cheeseman, A. Desai, Molecular architecture of the kinetochore-microtubule interface, *Nat Rev Mol Cell Biol* 9(1) (2008) 33-46.

[32] T. Hori, M. Amano, A. Suzuki, C.B. Backer, J.P. Welburn, Y. Dong, B.F. McEwen, W.H. Shang, E. Suzuki, K. Okawa, I.M. Cheeseman, T. Fukagawa, CCAN makes multiple contacts with centromeric DNA to provide distinct pathways to the outer kinetochore, *Cell* 135(6) (2008) 1039-52.

[33] M. Hara, T. Fukagawa, Critical Foundation of the Kinetochore: The Constitutive Centromere-Associated Network (CCAN), *Prog Mol Subcell Biol* 56 (2017) 29-57.

[34] D. Fachinetti, H.D. Folco, Y. Nechemia-Arbely, L.P. Valente, K. Nguyen, A.J. Wong, Q. Zhu, A.J. Holland, A. Desai, L.E. Jansen, D.W. Cleveland, A two-step mechanism for epigenetic specification of centromere identity and function, *Nat Cell Biol* 15(9) (2013) 1056-66.

[35] S.T. Liu, J.B. Rattner, S.A. Jablonski, T.J. Yen, Mapping the assembly pathways that specify formation of the trilaminar kinetochore plates in human cells, *J Cell Biol* 175(1) (2006) 41-53.

[36] G.A. Logsdon, E.J. Barrey, E.A. Bassett, J.E. DeNizio, L.Y. Guo, T. Panchenko, J.M. Dawicki-McKenna, P. Heun, B.E. Black, Both tails and the centromere targeting domain of CENP-A are required for centromere establishment, *J Cell Biol* 208(5) (2015) 521-31.

[37] P. Heun, S. Erhardt, M.D. Blower, S. Weiss, A.D. Skora, G.H. Karpen, Mislocalization of the *Drosophila* centromere-specific histone CID promotes formation of functional ectopic kinetochores, *Dev Cell* 10(3) (2006) 303-15.

[38] M.J. Mendiburo, J. Padeken, S. Fulop, A. Schepers, P. Heun, *Drosophila* CENH3 is sufficient for centromere formation, *Science* 334(6056) (2011) 686-90.

[39] M.C. Barnhart, P.H. Kuich, M.E. Stellfox, J.A. Ward, E.A. Bassett, B.E. Black, D.R. Foltz, HJURP is a CENP-A chromatin assembly factor sufficient to form a functional de novo kinetochore, *J Cell Biol* 194(2) (2011) 229-43.

[40] A.A. Van Hooser, Ouspenski, II, H.C. Gregson, D.A. Starr, T.J. Yen, M.L. Goldberg, K. Yokomori, W.C. Earnshaw, K.F. Sullivan, B.R. Brinkley, Specification of kinetochore-forming chromatin by the histone H3 variant CENP-A, *J Cell Sci* 114(Pt 19) (2001) 3529-42.

[41] T. Hori, W.H. Shang, K. Takeuchi, T. Fukagawa, The CCAN recruits CENP-A to the centromere and forms the structural core for kinetochore assembly, *J Cell Biol* 200(1) (2013) 45-60.

[42] A. Guse, C.W. Carroll, B. Moree, C.J. Fuller, A.F. Straight, In vitro centromere and kinetochore assembly on defined chromatin templates, *Nature* 477(7364) (2011) 354-8.

[43] E.M. Dunleavy, D. Roche, H. Tagami, N. Lacoste, D. Ray-Gallet, Y. Nakamura, Y. Daigo, Y. Nakatani, G. Almouzni-Pettinotti, HJURP is a cell-cycle-dependent maintenance and deposition factor of CENP-A at centromeres, *Cell* 137(3) (2009) 485-97.

[44] D.R. Foltz, L.E. Jansen, A.O. Bailey, J.R. Yates, 3rd, E.A. Bassett, S. Wood, B.E. Black, D.W. Cleveland, Centromere-specific assembly of CENP-a nucleosomes is mediated by HJURP, *Cell* 137(3) (2009) 472-84.

[45] R. Bernad, P. Sanchez, T. Rivera, M. Rodriguez-Corsino, E. Boyarchuk, I. Vassias, D. Ray-Gallet, A. Arnaoutov, M. Dasso, G. Almouzni, A. Losada, Xenopus HJURP and condensin II are required for CENP-A assembly, *J Cell Biol* 192(4) (2011) 569-82.

[46] L. Aravind, L.M. Iyer, C. Wu, Domain architectures of the Scm3p protein provide insights into centromere function and evolution, *Cell Cycle* 6(20) (2007) 2511-5.

[47] S. Stoler, K. Rogers, S. Weitze, L. Morey, M. Fitzgerald-Hayes, R.E. Baker, Scm3, an essential *Saccharomyces cerevisiae* centromere protein required for G2/M progression and Cse4 localization, *Proc Natl Acad Sci U S A* 104(25) (2007) 10571-6.

[48] R. Camahort, B. Li, L. Florens, S.K. Swanson, M.P. Washburn, J.L. Gerton, Scm3 is essential to recruit the histone h3 variant cse4 to centromeres and to maintain a functional kinetochore, *Mol Cell* 26(6) (2007) 853-65.

[49] A.L. Pidoux, E.S. Choi, J.K. Abbott, X. Liu, A. Kagansky, A.G. Castillo, G.L. Hamilton, W. Richardson, J. Rappaport, X. He, R.C. Allshire, Fission yeast Scm3: A CENP-A receptor required for integrity of subkinetochore chromatin, *Mol Cell* 33(3) (2009) 299-311.

[50] G. Mizuguchi, H. Xiao, J. Wisniewski, M.M. Smith, C. Wu, Nonhistone Scm3 and histones CenH3-H4 assemble the core of centromere-specific nucleosomes, *Cell* 129(6) (2007) 1153-64.

[51] C.C. Chen, M.L. Dechassa, E. Bettini, M.B. Ledoux, C. Belisario, P. Heun, K. Luger, B.G. Mellone, CAL1 is the *Drosophila* CENP-A assembly factor, *J Cell Biol* 204(3) (2014) 313-29.

[52] J.S. Williams, T. Hayashi, M. Yanagida, P. Russell, Fission yeast Scm3 mediates stable assembly of Cnp1/CENP-A into centromeric chromatin, *Mol Cell* 33(3) (2009) 287-98.

[53] L. Sanchez-Pulido, A.L. Pidoux, C.P. Ponting, R.C. Allshire, Common ancestry of the CENP-A chaperones Scm3 and HJURP, *Cell* 137(7) (2009) 1173-4.

[54] S. Erhardt, B.G. Mellone, C.M. Betts, W. Zhang, G.H. Karpen, A.F. Straight, Genome-wide analysis reveals a cell cycle-dependent mechanism controlling centromere propagation, *J Cell Biol* 183(5) (2008) 805-18.

[55] R.B. Schittenhelm, F. Althoff, S. Heidmann, C.F. Lehner, Detrimental incorporation of excess Cnp1/Cid and Cnp2/C into *Drosophila* centromeres is prevented by limiting amounts of the bridging factor Cal1, *J Cell Sci* 123(Pt 21) (2010) 3768-79.

[56] L. Rosin, B.G. Mellone, Co-evolving CENP-A and CAL1 Domains Mediate Centromeric CENP-A Deposition across *Drosophila* Species, *Dev Cell* 37(2) (2016) 136-47.

[57] M. Shuaib, K. Ouararhni, S. Dimitrov, A. Hamiche, HJURP binds CENP-A via a highly conserved N-terminal domain and mediates its deposition at centromeres, *Proc Natl Acad Sci U S A* 107(4) (2010) 1349-54.

[58] H. Hu, Y. Liu, M. Wang, J. Fang, H. Huang, N. Yang, Y. Li, J. Wang, X. Yao, Y. Shi, G. Li, R.M. Xu, Structure of a CENP-A-histone H4 heterodimer in complex with chaperone HJURP, *Genes Dev* 25(9) (2011) 901-6.

[59] E.A. Bassett, J. DeNizio, M.C. Barnhart-Dailey, T. Panchenko, N. Sekulic, D.J. Rogers, D.R. Foltz, B.E. Black, HJURP uses distinct CENP-A surfaces to recognize and to stabilize CENP-A/histone H4 for centromere assembly, *Dev Cell* 22(4) (2012) 749-62.

[60] Z. Zhou, H. Feng, B.R. Zhou, R. Ghirlando, K. Hu, A. Zwolak, L.M. Miller Jenkins, H. Xiao, N. Tjandra, C. Wu, Y. Bai, Structural basis for recognition of centromere histone variant CenH3 by the chaperone Scm3, *Nature* 472(7342) (2011) 234-7.

[61] B.E. Black, M.A. Brock, S. Bedard, V.L. Woods, Jr., D.W. Cleveland, An epigenetic mark generated by the incorporation of CENP-A into centromeric nucleosomes, *Proc Natl Acad Sci U S A* 104(12) (2007) 5008-13.

[62] B.E. Black, D.R. Foltz, S. Chakravarthy, K. Luger, V.L. Woods, Jr., D.W. Cleveland, Structural determinants for generating centromeric chromatin, *Nature* 430(6999) (2004) 578-82.

[63] B.E. Black, L.E. Jansen, P.S. Maddox, D.R. Foltz, A.B. Desai, J.V. Shah, D.W. Cleveland, Centromere identity maintained by nucleosomes assembled with histone H3 containing the CENP-A targeting domain, *Mol Cell* 25(2) (2007) 309-22.

[64] D. Goutte-Gattat, M. Shuaib, K. Ouararhni, T. Gautier, D.A. Skoufias, A. Hamiche, S. Dimitrov, Phosphorylation of the CENP-A amino-terminus in mitotic centromeric chromatin is required for kinetochore function, *Proc Natl Acad Sci U S A* 110(21) (2013) 8579-84.

[65] Y. Fujita, T. Hayashi, T. Kiyomitsu, Y. Toyoda, A. Kokubu, C. Obuse, M. Yanagida, Priming of centromere for CENP-A recruitment by human hMis18alpha, hMis18beta, and M18BP1, *Dev Cell* 12(1) (2007) 17-30.

[66] P.S. Maddox, F. Hyndman, J. Monen, K. Oegema, A. Desai, Functional genomics identifies a Myb domain-containing protein family required for assembly of CENP-A chromatin, *J Cell Biol* 176(6) (2007) 757-63.

[67] F. Spiller, B. Medina-Pritchard, M.A. Abad, M.A. Wear, O. Molina, W.C. Earnshaw, A.A. Jeyaprakash, Molecular basis for Cdk1-regulated timing of Mis18 complex assembly and CENP-A deposition, *EMBO Rep* 18(6) (2017) 894-905.

[68] D. Pan, K. Klare, A. Petrovic, A. Take, K. Walstein, P. Singh, A. Rondelet, A.W. Bird, A. Musacchio, CDK-regulated dimerization of M18BP1 on a Mis18 hexamer is necessary for CENP-A loading, *Elife* 6 (2017).

[69] M. Korntner-Vetter, S. Lefevre, X.W. Hu, R. George, M.R. Singleton, Subunit interactions and arrangements in the fission yeast Mis16-Mis18-Mis19 complex, *Life Sci Alliance* 2(4) (2019).

[70] M. Perpelescu, T. Hori, A. Toyoda, S. Misu, N. Monma, K. Ikeo, C. Obuse, A. Fujiyama, T. Fukagawa, HJURP is involved in the expansion of centromeric chromatin, *Mol Biol Cell* 26(15) (2015) 2742-54.

[71] J. Wang, X. Liu, Z. Dou, L. Chen, H. Jiang, C. Fu, G. Fu, D. Liu, J. Zhang, T. Zhu, J. Fang, J. Zang, J. Cheng, M. Teng, X. Ding, X. Yao, Mitotic regulator Mis18beta

interacts with and specifies the centromeric assembly of molecular chaperone holliday junction recognition protein (HJURP), *J Biol Chem* 289(12) (2014) 8326-36.

[72] S. Dambacher, W. Deng, M. Hahn, D. Sadic, J. Frohlich, A. Nuber, C. Hoischen, S. Diekmann, H. Leonhardt, G. Schotta, CENP-C facilitates the recruitment of M18BP1 to centromeric chromatin, *Nucleus* 3(1) (2012) 101-10.

[73] B. Moree, C.B. Meyer, C.J. Fuller, A.F. Straight, CENP-C recruits M18BP1 to centromeres to promote CENP-A chromatin assembly, *J Cell Biol* 194(6) (2011) 855-71.

[74] K.L. McKinley, I.M. Cheeseman, Polo-like kinase 1 licenses CENP-A deposition at centromeres, *Cell* 158(2) (2014) 397-411.

[75] B.T. French, F.G. Westhorpe, C. Limouse, A.F. Straight, *Xenopus laevis* M18BP1 Directly Binds Existing CENP-A Nucleosomes to Promote Centromeric Chromatin Assembly, *Dev Cell* 42(2) (2017) 190-199 e10.

[76] T. Hori, W.H. Shang, M. Hara, M. Ariyoshi, Y. Arimura, R. Fujita, H. Kurumizaka, T. Fukagawa, Association of M18BP1/KNL2 with CENP-A Nucleosome Is Essential for Centromere Formation in Non-mammalian Vertebrates, *Dev Cell* 42(2) (2017) 181-189 e3.

[77] R.D. Shelby, O. Vafa, K.F. Sullivan, Assembly of CENP-A into centromeric chromatin requires a cooperative array of nucleosomal DNA contact sites, *J Cell Biol* 136(3) (1997) 501-13.

[78] L.E. Jansen, B.E. Black, D.R. Foltz, D.W. Cleveland, Propagation of centromeric chromatin requires exit from mitosis, *J Cell Biol* 176(6) (2007) 795-805.

[79] M. Schuh, C.F. Lehner, S. Heidmann, Incorporation of *Drosophila* CID/CENP-A and CENP-C into centromeres during early embryonic anaphase, *Curr Biol* 17(3) (2007) 237-43.

[80] M.C. Silva, D.L. Bodor, M.E. Stellfox, N.M. Martins, H. Hochegger, D.R. Foltz, L.E. Jansen, Cdk activity couples epigenetic centromere inheritance to cell cycle progression, *Dev Cell* 22(1) (2012) 52-63.

[81] A. Stankovic, L.Y. Guo, J.F. Mata, D.L. Bodor, X.J. Cao, A.O. Bailey, J. Shabanowitz, D.F. Hunt, B.A. Garcia, B.E. Black, L.E.T. Jansen, A Dual Inhibitory Mechanism Sufficient to Maintain Cell-Cycle-Restricted CENP-A Assembly, *Mol Cell* 65(2) (2017) 231-246.

[82] S. Muller, R. Montes de Oca, N. Lacoste, F. Dingli, D. Loew, G. Almouzni, Phosphorylation and DNA binding of HJURP determine its centromeric recruitment and function in CenH3(CENP-A) loading, *Cell Rep* 8(1) (2014) 190-203.

[83] S.J. Falk, L.Y. Guo, N. Sekulic, E.M. Smoak, T. Mani, G.A. Logsdon, K. Gupta, L.E. Jansen, G.D. Van Duyne, S.A. Vinogradov, M.A. Lampson, B.E. Black, Chromosomes. CENP-C reshapes and stabilizes CENP-A nucleosomes at the centromere, *Science* 348(6235) (2015) 699-703.

[84] E.M. Smoak, P. Stein, R.M. Schultz, M.A. Lampson, B.E. Black, Long-Term Retention of CENP-A Nucleosomes in Mammalian Oocytes Underpins Transgenerational Inheritance of Centromere Identity, *Curr Biol* 26(8) (2016) 1110-6.

[85] P. Hemmerich, S. Weidtkamp-Peters, C. Hoischen, L. Schmiedeberg, I. Erliandri, S. Diekmann, Dynamics of inner kinetochore assembly and maintenance in living cells, *J Cell Biol* 180(6) (2008) 1101-14.

[86] D.L. Bodor, L.P. Valente, J.F. Mata, B.E. Black, L.E. Jansen, Assembly in G1 phase and long-term stability are unique intrinsic features of CENP-A nucleosomes, *Mol Biol Cell* 24(7) (2013) 923-32.

[87] S. Mitra, B. Srinivasan, L.E.T. Jansen, Stable inheritance of CENP-A chromatin: Inner strength versus dynamic control, *J Cell Biol* 219(10) (2020).

[88] S. Srivastava, D.R. Foltz, Posttranslational modifications of CENP-A: marks of distinction, *Chromosoma* 127(3) (2018) 279-290.

[89] O. Marescal, I.M. Cheeseman, Cellular Mechanisms and Regulation of Quiescence, *Dev Cell* 55(3) (2020) 259-271.

[90] S.Z. Swartz, L.S. McKay, K.C. Su, L. Bury, A. Padeganeh, P.S. Maddox, K.A. Knouse, I.M. Cheeseman, Quiescent Cells Actively Replenish CENP-A Nucleosomes to Maintain Centromere Identity and Proliferative Potential, *Dev Cell* 51(1) (2019) 35-48 e7.

[91] S.H. Lee, P. Itkin-Ansari, F. Levine, CENP-A, a protein required for chromosome segregation in mitosis, declines with age in islet but not exocrine cells, *Aging (Albany NY)* 2(11) (2010) 785-90.

[92] W. Zhang, J. Feng, Q. Li, The replisome guides nucleosome assembly during DNA replication, *Cell Biosci* 10 (2020) 37.

[93] E.M. Dunleavy, G. Almouzni, G.H. Karpen, H3.3 is deposited at centromeres in S phase as a placeholder for newly assembled CENP-A in G(1) phase, *Nucleus* 2(2) (2011) 146-57.

[94] Y. Nechemia-Arbely, K.H. Miga, O. Shoshani, A. Aslanian, M.A. McMahon, A.Y. Lee, D. Fachinetti, J.R. Yates, 3rd, B. Ren, D.W. Cleveland, DNA replication acts as an error correction mechanism to maintain centromere identity by restricting CENP-A to centromeres, *Nat Cell Biol* 21(6) (2019) 743-754.

[95] T. Hori, N. Kagawa, A. Toyoda, A. Fujiyama, S. Misu, N. Monma, F. Makino, K. Ikeo, T. Fukagawa, Constitutive centromere-associated network controls centromere drift in vertebrate cells, *J Cell Biol* 216(1) (2017) 101-113.

[96] E. Zasadzinska, J. Huang, A.O. Bailey, L.Y. Guo, N.S. Lee, S. Srivastava, K.A. Wong, B.T. French, B.E. Black, D.R. Foltz, Inheritance of CENP-A Nucleosomes during DNA Replication Requires HJURP, *Dev Cell* 47(3) (2018) 348-362 e7.

[97] F.L. Chan, O.J. Marshall, R. Saffery, B.W. Kim, E. Earle, K.H. Choo, L.H. Wong, Active transcription and essential role of RNA polymerase II at the centromere during mitosis, *Proc Natl Acad Sci U S A* 109(6) (2012) 1979-84.

[98] J.H. Bergmann, J.N. Jakubsche, N.M. Martins, A. Kagansky, M. Nakano, H. Kimura, D.A. Kelly, B.M. Turner, H. Masumoto, V. Larionov, W.C. Earnshaw, Epigenetic engineering: histone H3K9 acetylation is compatible with kinetochore structure and function, *J Cell Sci* 125(Pt 2) (2012) 411-21.

[99] J.H. Bergmann, M.G. Rodriguez, N.M. Martins, H. Kimura, D.A. Kelly, H. Masumoto, V. Larionov, L.E. Jansen, W.C. Earnshaw, Epigenetic engineering shows H3K4me2 is required for HJURP targeting and CENP-A assembly on a synthetic human kinetochore, *EMBO J* 30(2) (2011) 328-40.

[100] L. Bury, B. Moodie, J. Ly, L.S. McKay, K.H. Miga, I.M. Cheeseman, Alpha-satellite RNA transcripts are repressed by centromere-nucleolus associations, *Elife* 9 (2020).

[101] C. Perea-Resa, M.D. Blower, Centromere Biology: Transcription Goes on Stage, *Mol Cell Biol* 38(18) (2018).

[102] C. Perea-Resa, L. Bury, I.M. Cheeseman, M.D. Blower, Cohesin Removal Reprograms Gene Expression upon Mitotic Entry, *Mol Cell* 78(1) (2020) 127-140 e7.

[103] M. Novais-Cruz, M. Alba Abad, I.W.F. van, N. Galjart, A.A. Jeyaprakash, H. Maiato, C. Ferras, Mitotic progression, arrest, exit or death relies on centromere structural integrity, rather than *de novo* transcription, *Elife* 7 (2018).

[104] M. Okada, I.M. Cheeseman, T. Hori, K. Okawa, I.X. McLeod, J.R. Yates, 3rd, A. Desai, T. Fukagawa, The CENP-H-I complex is required for the efficient incorporation of newly synthesized CENP-A into centromeres, *Nat Cell Biol* 8(5) (2006) 446-57.

[105] T. Hori, M. Okada, K. Maenaka, T. Fukagawa, CENP-O class proteins form a stable complex and are required for proper kinetochore function, *Mol Biol Cell* 19(3) (2008) 843-54.

[106] D.R. Foltz, L.E. Jansen, B.E. Black, A.O. Bailey, J.R. Yates, 3rd, D.W. Cleveland, The human CENP-A centromeric nucleosome-associated complex, *Nat Cell Biol* 8(5) (2006) 458-69.

[107] K. Klare, J.R. Weir, F. Basilico, T. Zimniak, L. Massimiliano, N. Ludwigs, F. Herzog, A. Musacchio, CENP-C is a blueprint for constitutive centromere-associated network assembly within human kinetochores, *J Cell Biol* 210(1) (2015) 11-22.

[108] K.L. McKinley, N. Sekulic, L.Y. Guo, T. Tsinman, B.E. Black, I.M. Cheeseman, The CENP-L-N Complex Forms a Critical Node in an Integrated Meshwork of Interactions at the Centromere-Kinetochore Interface, *Mol Cell* 60(6) (2015) 886-98.

[109] A. Petrovic, J. Keller, Y. Liu, K. Overlack, J. John, Y.N. Dimitrova, S. Jenni, S. van Gerwen, P. Stege, S. Wohlgemuth, P. Rombaut, F. Herzog, S.C. Harrison, I.R. Vetter, A. Musacchio, Structure of the MIS12 Complex and Molecular Basis of Its Interaction with CENP-C at Human Kinetochores, *Cell* 167(4) (2016) 1028-1040 e15.

[110] E. Scrpanti, A. De Antoni, G.M. Alushin, A. Petrovic, T. Melis, E. Nogales, A. Musacchio, Direct binding of Cenp-C to the Mis12 complex joins the inner and outer kinetochore, *Curr Biol* 21(5) (2011) 391-8.

[111] M.R. Przewloka, Z. Venkei, V.M. Bolanos-Garcia, J. Debski, M. Dadlez, D.M. Glover, CENP-C is a structural platform for kinetochore assembly, *Curr Biol* 21(5) (2011) 399-405.

[112] J.J. van Hooff, E. Tromer, L.M. van Wijk, B. Snel, G.J. Kops, Evolutionary dynamics of the kinetochore network in eukaryotes as revealed by comparative genomics, *EMBO Rep* 18(9) (2017) 1559-1571.

[113] K. Tanaka, H.L. Chang, A. Kagami, Y. Watanabe, CENP-C functions as a scaffold for effectors with essential kinetochore functions in mitosis and meiosis, *Dev Cell* 17(3) (2009) 334-43.

[114] C.W. Carroll, K.J. Milks, A.F. Straight, Dual recognition of CENP-A nucleosomes is required for centromere assembly, *J Cell Biol* 189(7) (2010) 1143-55.

[115] J.R. Weir, A.C. Faesen, K. Klare, A. Petrovic, F. Basilico, J. Fischbock, S. Pentakota, J. Keller, M.E. Pesenti, D. Pan, D. Vogt, S. Wohlgemuth, F. Herzog, A. Musacchio, Insights from biochemical reconstitution into the architecture of human kinetochores, *Nature* 537(7619) (2016) 249-253.

[116] C.W. Carroll, M.C. Silva, K.M. Godek, L.E. Jansen, A.F. Straight, Centromere assembly requires the direct recognition of CENP-A nucleosomes by CENP-N, *Nat Cell Biol* 11(7) (2009) 896-902.

[117] F. Basilico, S. Maffini, J.R. Weir, D. Prumbaum, A.M. Rojas, T. Zimniak, A. De Antoni, S. Jeganathan, B. Voss, S. van Gerwen, V. Krenn, L. Massimiliano, A. Valencia, I.R. Vetter, F. Herzog, S. Raunser, S. Pasqualato, A. Musacchio, The pseudo GTPase CENP-M drives human kinetochore assembly, *Elife* 3 (2014) e02978.

[118] M.E. Pesenti, D. Prumbaum, P. Auckland, C.M. Smith, A.C. Faesen, A. Petrovic, M. Erent, S. Maffini, S. Pentakota, J.R. Weir, Y.C. Lin, S. Raunser, A.D. McAinsh, A. Musacchio, Reconstitution of a 26-Subunit Human Kinetochore Reveals Cooperative Microtubule Binding by CENP-OPQUR and NDC80, *Mol Cell* 71(6) (2018) 923-939 e10.

[119] T. Nishino, K. Takeuchi, K.E. Gascoigne, A. Suzuki, T. Hori, T. Oyama, K. Morikawa, I.M. Cheeseman, T. Fukagawa, CENP-T-W-S-X forms a unique centromeric chromatin structure with a histone-like fold, *Cell* 148(3) (2012) 487-501.

[120] K. Takeuchi, T. Nishino, K. Mayanagi, N. Horikoshi, A. Osakabe, H. Tachiwana, T. Hori, H. Kurumizaka, T. Fukagawa, The centromeric nucleosome-like CENP-T-W-S-X complex induces positive supercoils into DNA, *Nucleic Acids Res* 42(3) (2014) 1644-55.

[121] K.E. Gascoigne, K. Takeuchi, A. Suzuki, T. Hori, T. Fukagawa, I.M. Cheeseman, Induced ectopic kinetochore assembly bypasses the requirement for CENP-A nucleosomes, *Cell* 145(3) (2011) 410-22.

[122] A. Schleiffer, M. Maier, G. Litos, F. Lampert, P. Hornung, K. Mechtler, S. Westermann, CENP-T proteins are conserved centromere receptors of the Ndc80 complex, *Nat Cell Biol* 14(6) (2012) 604-13.

[123] T. Nishino, F. Rago, T. Hori, K. Tomii, I.M. Cheeseman, T. Fukagawa, CENP-T provides a structural platform for outer kinetochore assembly, *EMBO J* 32(3) (2013) 424-36.

[124] J. Bancroft, P. Auckland, C.P. Samora, A.D. McAinsh, Chromosome congression is promoted by CENP-Q- and CENP-E-dependent pathways, *J Cell Sci* 128(1) (2015) 171-84.

[125] S. Hua, Z. Wang, K. Jiang, Y. Huang, T. Ward, L. Zhao, Z. Dou, X. Yao, CENP-U cooperates with Hec1 to orchestrate kinetochore-microtubule attachment, *J Biol Chem* 286(2) (2011) 1627-38.

[126] A.C. Amaro, C.P. Samora, R. Holtackers, E. Wang, I.J. Kingston, M. Alonso, M. Lampson, A.D. McAinsh, P. Meraldi, Molecular control of kinetochore-microtubule dynamics and chromosome oscillations, *Nat Cell Biol* 12(4) (2010) 319-29.

[127] Y.H. Kang, C.H. Park, T.S. Kim, N.K. Soung, J.K. Bang, B.Y. Kim, J.E. Park, K.S. Lee, Mammalian polo-like kinase 1-dependent regulation of the PBIP1-CENP-Q complex at kinetochores, *J Biol Chem* 286(22) (2011) 19744-57.

[128] Y.H. Kang, J.E. Park, L.R. Yu, N.K. Soung, S.M. Yun, J.K. Bang, Y.S. Seong, H. Yu, S. Garfield, T.D. Veenstra, K.S. Lee, Self-regulated Plk1 recruitment to kinetochores by the Plk1-PBIP1 interaction is critical for proper chromosome segregation, *Mol Cell* 24(3) (2006) 409-22.

[129] P. Singh, M.E. Pesenti, S. Maffini, S. Carmignani, M. Hedtfeld, A. Petrovic, A. Srinivasamani, T. Bange, A. Musacchio, BUB1 and CENP-U, Primed by CDK1, Are the Main PLK1 Kinetochore Receptors in Mitosis, *Mol Cell* 81(1) (2021) 67-87 e9.

[130] A.L. Nguyen, M.D. Fadel, I.M. Cheeseman, Differential Requirements for the CENP-O Complex Reveal Parallel PLK1 Kinetochore Recruitment Pathways, *Mol Biol Cell* (2021) mbcE20110751.

[131] S.E. McClelland, S. Borusu, A.C. Amaro, J.R. Winter, M. Belwal, A.D. McAinsh, P. Meraldi, The CENP-A NAC/CAD kinetochore complex controls chromosome congression and spindle bipolarity, *EMBO J* 26(24) (2007) 5033-47.

[132] A. Eskat, W. Deng, A. Hofmeister, S. Rudolphi, S. Emmerth, D. Hellwig, T. Ulbricht, V. Doring, J.M. Bancroft, A.D. McAinsh, M.C. Cardoso, P. Meraldi, C. Hoischen, H. Leonhardt, S. Diekmann, Step-wise assembly, maturation and dynamic behavior of the human CENP-P/O/R/Q/U kinetochore sub-complex, *PLoS One* 7(9) (2012) e44717.

[133] K.E. Gascoigne, I.M. Cheeseman, CDK-dependent phosphorylation and nuclear exclusion coordinately control kinetochore assembly state, *J Cell Biol* 201(1) (2013) 23-32.

[134] D. Hellwig, S. Munch, S. Orthaus, C. Hoischen, P. Hemmerich, S. Diekmann, Live-cell imaging reveals sustained centromere binding of CENP-T via CENP-A and CENP-B, *J Biophotonics* 1(3) (2008) 245-54.

[135] L. Prendergast, C. van Vuuren, A. Kaczmarczyk, V. Doering, D. Hellwig, N. Quinn, C. Hoischen, S. Diekmann, K.F. Sullivan, Premitotic assembly of human CENPs -T and -W switches centromeric chromatin to a mitotic state, *PLoS Biol* 9(6) (2011) e1001082.

[136] T. Fukagawa, Y. Mikami, A. Nishihashi, V. Regnier, T. Haraguchi, Y. Hirakawa, N. Sugata, K. Todokoro, W. Brown, T. Ikemura, CENP-H, a constitutive centromere component, is required for centromere targeting of CENP-C in vertebrate cells, *EMBO J* 20(16) (2001) 4603-17.

[137] R. Watanabe, M. Hara, E.I. Okumura, S. Herve, D. Fachinetti, M. Ariyoshi, T. Fukagawa, CDK1-mediated CENP-C phosphorylation modulates CENP-A binding and mitotic kinetochore localization, *J Cell Biol* 218(12) (2019) 4042-4062.

[138] M. Ariyoshi, F. Makino, R. Watanabe, R. Nakagawa, T. Kato, K. Namba, Y. Arimura, R. Fujita, H. Kurumizaka, E.I. Okumura, M. Hara, T. Fukagawa, Cryo-EM structure of the CENP-A nucleosome in complex with phosphorylated CENP-C, *EMBO J* (2021) e105671.

[139] L.Y. Guo, P.K. Allu, L. Zandarashvili, K.L. McKinley, N. Sekulic, J.M. Dawicki-McKenna, D. Fachinetti, G.A. Logsdon, R.M. Jamiolkowski, D.W. Cleveland, I.M. Cheeseman, B.E. Black, Centromeres are maintained by fastening CENP-A to DNA and directing an arginine anchor-dependent nucleosome transition, *Nat Commun* 8 (2017) 15775.

[140] S. Pentakota, K. Zhou, C. Smith, S. Maffini, A. Petrovic, G.P. Morgan, J.R. Weir, I.R. Vetter, A. Musacchio, K. Luger, Decoding the centromeric nucleosome through CENP-N, *Elife* 6 (2017).

[141] D. Hellwig, S. Emmerth, T. Ulbricht, V. Doring, C. Hoischen, R. Martin, C.P. Samora, A.D. McAinsh, C.W. Carroll, A.F. Straight, P. Meraldi, S. Diekmann, Dynamics of CENP-N kinetochore binding during the cell cycle, *J Cell Sci* 124(Pt 22) (2011) 3871-83.

[142] P.K. Allu, J.M. Dawicki-McKenna, T. Van Eeuwen, M. Slavin, M. Braitbard, C. Xu, N. Kalisman, K. Murakami, B.E. Black, Structure of the Human Core Centromeric Nucleosome Complex, *Curr Biol* 29(16) (2019) 2625-2639 e5.

[143] J. Fang, Y. Liu, Y. Wei, W. Deng, Z. Yu, L. Huang, Y. Teng, T. Yao, Q. You, H. Ruan, P. Chen, R.M. Xu, G. Li, Structural transitions of centromeric chromatin regulate the cell cycle-dependent recruitment of CENP-N, *Genes Dev* 29(10) (2015) 1058-73.

[144] S. Ohta, M. Kimura, S. Takagi, I. Toramoto, Y. Ishihama, Identification of Mitosis-Specific Phosphorylation in Mitotic Chromosome-Associated Proteins, *J Proteome Res* 15(9) (2016) 3331-41.

[145] R. Gamba, D. Fachinetti, From evolution to function: Two sides of the same CENP-B coin?, *Exp Cell Res* 390(2) (2020) 111959.

[146] M. Dumont, R. Gamba, P. Gestraud, S. Klaasen, J.T. Worrall, S.G. De Vries, V. Boudreau, C. Salinas-Luypaert, P.S. Maddox, S.M. Lens, G.J. Kops, S.E. McClelland, K.H. Miga, D. Fachinetti, Human chromosome-specific aneuploidy is influenced by DNA-dependent centromeric features, *EMBO J* 39(2) (2020) e102924.

[147] D. Fachinetti, J.S. Han, M.A. McMahon, P. Ly, A. Abdullah, A.J. Wong, D.W. Cleveland, DNA Sequence-Specific Binding of CENP-B Enhances the Fidelity of Human Centromere Function, *Dev Cell* 33(3) (2015) 314-27.

[148] R. Fujita, K. Otake, Y. Arimura, N. Horikoshi, Y. Miya, T. Shiga, A. Osakabe, H. Tachiwana, J. Ohzeki, V. Larionov, H. Masumoto, H. Kurumizaka, Stable complex formation of CENP-B with the CENP-A nucleosome, *Nucleic Acids Res* 43(10) (2015) 4909-22.

[149] D. Hasson, T. Panchenko, K.J. Salimian, M.U. Salman, N. Sekulic, A. Alonso, P.E. Warburton, B.E. Black, The octamer is the major form of CENP-A nucleosomes at human centromeres, *Nat Struct Mol Biol* 20(6) (2013) 687-95.

[150] S. Hoffmann, M. Dumont, V. Barra, P. Ly, Y. Nechemia-Arbely, M.A. McMahon, S. Herve, D.W. Cleveland, D. Fachinetti, CENP-A Is Dispensable for Mitotic Centromere Function after Initial Centromere/Kinetochore Assembly, *Cell Rep* 17(9) (2016) 2394-2404.

[151] S.A. Ribeiro, P. Vagnarelli, Y. Dong, T. Hori, B.F. McEwen, T. Fukagawa, C. Flors, W.C. Earnshaw, A super-resolution map of the vertebrate kinetochore, *Proc Natl Acad Sci U S A* 107(23) (2010) 10484-9.

[152] L.L. Sullivan, C.D. Boivin, B. Mravinac, I.Y. Song, B.A. Sullivan, Genomic size of CENP-A domain is proportional to total alpha satellite array size at human centromeres and expands in cancer cells, *Chromosome Res* 19(4) (2011) 457-70.

[153] D.L. Bodor, J.F. Mata, M. Sergeev, A.F. David, K.J. Salimian, T. Panchenko, D.W. Cleveland, B.E. Black, J.V. Shah, L.E. Jansen, The quantitative architecture of centromeric chromatin, *Elife* 3 (2014) e02137.

[154] G. Vargiu, A.A. Makarov, J. Allan, T. Fukagawa, D.G. Booth, W.C. Earnshaw, Stepwise unfolding supports a subunit model for vertebrate kinetochores, *Proc Natl Acad Sci U S A* 114(12) (2017) 3133-3138.

[155] T. Schalch, F.A. Steiner, Structure of centromere chromatin: from nucleosome to chromosomal architecture, *Chromosoma* 126(4) (2017) 443-455.

[156] P.S. Maddox, N. Portier, A. Desai, K. Oegema, Molecular analysis of mitotic chromosome condensation using a quantitative time-resolved fluorescence microscopy assay, *Proc Natl Acad Sci U S A* 103(41) (2006) 15097-102.

[157] I.M. Cheeseman, J.S. Chappie, E.M. Wilson-Kubalek, A. Desai, The conserved KMN network constitutes the core microtubule-binding site of the kinetochore, *Cell* 127(5) (2006) 983-97.

[158] I.M. Cheeseman, S. Niessen, S. Anderson, F. Hyndman, J.R. Yates, 3rd, K. Oegema, A. Desai, A conserved protein network controls assembly of the outer kinetochore and its ability to sustain tension, *Genes Dev* 18(18) (2004) 2255-68.

[159] A. Desai, S. Rybina, T. Muller-Reichert, A. Shevchenko, A. Shevchenko, A. Hyman, K. Oegema, KNL-1 directs assembly of the microtubule-binding interface of the kinetochore in *C. elegans*, *Genes Dev* 17(19) (2003) 2421-35.

[160] S.L. Kline, I.M. Cheeseman, T. Hori, T. Fukagawa, A. Desai, The human Mis12 complex is required for kinetochore assembly and proper chromosome segregation, *J Cell Biol* 173(1) (2006) 9-17.

[161] P.A. Wigge, J.V. Kilmartin, The Ndc80p complex from *Saccharomyces cerevisiae* contains conserved centromere components and has a function in chromosome segregation, *J Cell Biol* 152(2) (2001) 349-60.

[162] R. Bharadwaj, W. Qi, H. Yu, Identification of two novel components of the human NDC80 kinetochore complex, *J Biol Chem* 279(13) (2004) 13076-85.

[163] M.L. McCleland, R.D. Gardner, M.J. Kallio, J.R. Daum, G.J. Gorbsky, D.J. Burke, P.T. Stukenberg, The highly conserved Ndc80 complex is required for kinetochore assembly, chromosome congression, and spindle checkpoint activity, *Genes Dev* 17(1) (2003) 101-14.

[164] C. Obuse, O. Iwasaki, T. Kiyomitsu, G. Goshima, Y. Toyoda, M. Yanagida, A conserved Mis12 centromere complex is linked to heterochromatic HP1 and outer kinetochore protein Zwint-1, *Nat Cell Biol* 6(11) (2004) 1135-41.

[165] S. Westermann, I.M. Cheeseman, S. Anderson, J.R. Yates, 3rd, D.G. Drubin, G. Barnes, Architecture of the budding yeast kinetochore reveals a conserved molecular core, *J Cell Biol* 163(2) (2003) 215-22.

[166] V.S. Nekrasov, M.A. Smith, S. Peak-Chew, J.V. Kilmartin, Interactions between centromere complexes in *Saccharomyces cerevisiae*, *Mol Biol Cell* 14(12) (2003) 4931-46.

[167] B.A. Pinsky, S.Y. Tatsutani, K.A. Collins, S. Biggins, An Mtw1 complex promotes kinetochore biorientation that is monitored by the Ipl1/Aurora protein kinase, *Dev Cell* 5(5) (2003) 735-45.

[168] J.G. DeLuca, W.E. Gall, C. Ciferri, D. Cimini, A. Musacchio, E.D. Salmon, Kinetochore microtubule dynamics and attachment stability are regulated by Hec1, *Cell* 127(5) (2006) 969-82.

[169] F. Rago, K.E. Gascoigne, I.M. Cheeseman, Distinct organization and regulation of the outer kinetochore KMN network downstream of CENP-C and CENP-T, *Curr Biol* 25(5) (2015) 671-7.

[170] K.E. Gascoigne, I.M. Cheeseman, Kinetochore assembly: if you build it, they will come, *Curr Opin Cell Biol* 23(1) (2011) 102-8.

[171] I.M. Cheeseman, T. Hori, T. Fukagawa, A. Desai, KNL1 and the CENP-H/I/K complex coordinately direct kinetochore assembly in vertebrates, *Mol Biol Cell* 19(2) (2008) 587-94.

[172] M. Hara, M. Ariyoshi, E.I. Okumura, T. Hori, T. Fukagawa, Multiple phosphorylations control recruitment of the KMN network onto kinetochores, *Nat Cell Biol* 20(12) (2018) 1378-1388.

[173] Y.N. Dimitrova, S. Jenni, R. Valverde, Y. Khin, S.C. Harrison, Structure of the MIND Complex Defines a Regulatory Focus for Yeast Kinetochore Assembly, *Cell* 167(4) (2016) 1014-1027 e12.

[174] F. Malvezzi, G. Litos, A. Schleiffer, A. Heuck, K. Mechtler, T. Clausen, S. Westermann, A structural basis for kinetochore recruitment of the Ndc80 complex via two distinct centromere receptors, *EMBO J* 32(3) (2013) 409-23.

[175] A. Petrovic, S. Pasqualato, P. Dube, V. Krenn, S. Santaguida, D. Cittaro, S. Monzani, L. Massimiliano, J. Keller, A. Tarricone, A. Maiolica, H. Stark, A. Musacchio, The MIS12 complex is a protein interaction hub for outer kinetochore assembly, *J Cell Biol* 190(5) (2010) 835-52.

[176] P.J. Huis In 't Veld, S. Jeganathan, A. Petrovic, P. Singh, J. John, V. Krenn, F. Weissmann, T. Bange, A. Musacchio, Molecular basis of outer kinetochore assembly on CENP-T, *Elife* 5 (2016).

[177] G. Pekgoz Altunkaya, F. Malvezzi, Z. Demianova, T. Zimniak, G. Litos, F. Weissmann, K. Mechtler, F. Herzog, S. Westermann, CCAN Assembly Configures Composite Binding Interfaces to Promote Cross-Linking of Ndc80 Complexes at the Kinetochore, *Curr Biol* 26(17) (2016) 2370-8.

[178] T.N. Gaitanos, A. Santamaria, A.A. Jeyaprakash, B. Wang, E. Conti, E.A. Nigg, Stable kinetochore-microtubule interactions depend on the Ska complex and its new component Ska3/C13Orf3, *EMBO J* 28(10) (2009) 1442-52.

[179] J.C. Schmidt, H. Arthanari, A. Boeszoermenyi, N.M. Dashkevich, E.M. Wilson-Kubalek, N. Monnier, M. Markus, M. Oberer, R.A. Milligan, M. Bathe, G. Wagner, E.L. Grishchuk, I.M. Cheeseman, The kinetochore-bound Ska1 complex tracks depolymerizing microtubules and binds to curved protofilaments, *Dev Cell* 23(5) (2012) 968-80.

[180] A.A. Jeyaprakash, A. Santamaria, U. Jayachandran, Y.W. Chan, C. Benda, E.A. Nigg, E. Conti, Structural and functional organization of the Ska complex, a key component of the kinetochore-microtubule interface, *Mol Cell* 46(3) (2012) 274-86.

[181] D.K. Cheerambathur, B. Prevo, N. Hattersley, L. Lewellyn, K.D. Corbett, K. Oegema, A. Desai, Dephosphorylation of the Ndc80 Tail Stabilizes Kinetochore-Microtubule Attachments via the Ska Complex, *Dev Cell* 41(4) (2017) 424-437 e4.

[182] P.L. Janczyk, K.A. Skorupka, J.G. Tooley, D.R. Matson, C.A. Kestner, T. West, O. Pornillos, P.T. Stukenberg, Mechanism of Ska Recruitment by Ndc80 Complexes to Kinetochores, *Dev Cell* 41(4) (2017) 438-449 e4.

[183] J.P. Welburn, E.L. Grishchuk, C.B. Backer, E.M. Wilson-Kubalek, J.R. Yates, 3rd, I.M. Cheeseman, The human kinetochore Ska1 complex facilitates microtubule depolymerization-coupled motility, *Dev Cell* 16(3) (2009) 374-85.

[184] P. Auckland, N.I. Clarke, S.J. Royle, A.D. McAinsh, Congressing kinetochores progressively load Ska complexes to prevent force-dependent detachment, *J Cell Biol* 216(6) (2017) 1623-1639.

[185] J. Gruber, J. Harborth, J. Schnabel, K. Weber, M. Hatzfeld, The mitotic-spindle-associated protein astrin is essential for progression through mitosis, *J Cell Sci* 115(Pt 21) (2002) 4053-9.

[186] A.L. Manning, S.F. Bakhour, S. Maffini, C. Correia-Melo, H. Maiato, D.A. Compton, CLASP1, astrin and Kif2b form a molecular switch that regulates kinetochore-microtubule dynamics to promote mitotic progression and fidelity, *EMBO J* 29(20) (2010) 3531-43.

[187] K.H. Thein, J. Kleylein-Sohn, E.A. Nigg, U. Gruneberg, Astrin is required for the maintenance of sister chromatid cohesion and centrosome integrity, *J Cell Biol* 178(3) (2007) 345-54.

[188] J.C. Schmidt, T. Kiyomitsu, T. Hori, C.B. Backer, T. Fukagawa, I.M. Cheeseman, Aurora B kinase controls the targeting of the Astrin-SKAP complex to bioriented kinetochores, *J Cell Biol* 191(2) (2010) 269-80.

[189] D.M. Kern, J.K. Monda, K.C. Su, E.M. Wilson-Kubalek, I.M. Cheeseman, Astrin-SKAP complex reconstitution reveals its kinetochore interaction with microtubule-bound Ndc80, *Elife* 6 (2017).

[190] G.J. Mack, D.A. Compton, Analysis of mitotic microtubule-associated proteins using mass spectrometry identifies astrin, a spindle-associated protein, *Proc Natl Acad Sci U S A* 98(25) (2001) 14434-9.

[191] J.K. Monda, I.M. Cheeseman, The kinetochore-microtubule interface at a glance, *J Cell Sci* 131(16) (2018).

[192] E.J. Lawrence, M. Zanic, L.M. Rice, CLASPs at a glance, *J Cell Sci* 133(8) (2020).

[193] C.L. Rieder, The formation, structure, and composition of the mammalian kinetochore and kinetochore fiber, *Int Rev Cytol* 79 (1982) 1-58.

[194] B.F. McEwen, J.T. Arena, J. Frank, C.L. Rieder, Structure of the colcemid-treated PtK1 kinetochore outer plate as determined by high voltage electron microscopic tomography, *J Cell Biol* 120(2) (1993) 301-12.

[195] V. Magidson, R. Paul, N. Yang, J.G. Ault, C.B. O'Connell, I. Tikhonenko, B.F. McEwen, A. Mogilner, A. Khodjakov, Adaptive changes in the kinetochore architecture facilitate proper spindle assembly, *Nat Cell Biol* 17(9) (2015) 1134-44.

[196] P.T. Jokelainen, The ultrastructure and spatial organization of the metaphase kinetochore in mitotic rat cells, *J Ultrastruct Res* 19(1) (1967) 19-44.

[197] V. Magidson, J. He, J.G. Ault, C.B. O'Connell, N. Yang, I. Tikhonenko, B.F. McEwen, H. Sui, A. Khodjakov, Unattached kinetochores rather than intrakinetochore tension arrest mitosis in taxol-treated cells, *J Cell Biol* 212(3) (2016) 307-19.

[198] C.A. Cooke, B. Schaar, T.J. Yen, W.C. Earnshaw, Localization of CENP-E in the fibrous corona and outer plate of mammalian kinetochores from prometaphase through anaphase, *Chromosoma* 106(7) (1997) 446-55.

[199] L. Wordeman, E.R. Steuer, M.P. Sheetz, T. Mitchison, Chemical subdomains within the kinetochore domain of isolated CHO mitotic chromosomes, *J Cell Biol* 114(2) (1991) 285-94.

[200] J.B. Rattner, A. Rao, M.J. Fritzler, D.W. Valencia, T.J. Yen, CENP-F is a .ca 400 kDa kinetochore protein that exhibits a cell-cycle dependent localization, *Cell Motil Cytoskeleton* 26(3) (1993) 214-26.

[201] R. Basto, F. Scaerou, S. Mische, E. Wojcik, C. Lefebvre, R. Gomes, T. Hays, R. Karess, In vivo dynamics of the rough deal checkpoint protein during Drosophila mitosis, *Curr Biol* 14(1) (2004) 56-61.

[202] D.A. Starr, B.C. Williams, T.S. Hays, M.L. Goldberg, ZW10 helps recruit dynein and dynein to the kinetochore, *J Cell Biol* 142(3) (1998) 763-74.

[203] J.A. Rodriguez-Rodriguez, C. Lewis, K.L. McKinley, V. Sikirzhytski, J. Corona, J. Maciejowski, A. Khodjakov, I.M. Cheeseman, P.V. Jallepalli, Distinct Roles of RZZ and Bub1-KNL1 in Mitotic Checkpoint Signaling and Kinetochore Expansion, *Curr Biol* 28(21) (2018) 3422-3429 e5.

[204] C. Pereira, R.M. Reis, J.B. Gama, R. Celestino, D.K. Cheerambathur, A.X. Carvalho, R. Gassmann, Self-Assembly of the RZZ Complex into Filaments Drives Kinetochore Expansion in the Absence of Microtubule Attachment, *Curr Biol* 28(21) (2018) 3408-3421 e8.

[205] C. Sacristan, M.U.D. Ahmad, J. Keller, J. Fermie, V. Groenewold, E. Tromer, A. Fish, R. Melero, J.M. Carazo, J. Klumperman, A. Musacchio, A. Perrakis, G.J. Kops, Dynamic kinetochore size regulation promotes microtubule capture and chromosome biorientation in mitosis, *Nat Cell Biol* 20(7) (2018) 800-810.

[206] V. Silio, A.D. McAinsh, J.B. Millar, KNL1-Bubs and RZZ Provide Two Separable Pathways for Checkpoint Activation at Human Kinetochores, *Dev Cell* 35(5) (2015) 600-613.

[207] B.J. Howell, B.F. McEwen, J.C. Canman, D.B. Hoffman, E.M. Farrar, C.L. Rieder, E.D. Salmon, Cytoplasmic dynein/dynactin drives kinetochore protein transport to the spindle poles and has a role in mitotic spindle checkpoint inactivation, *J Cell Biol* 155(7) (2001) 1159-72.

[208] E. Wojcik, R. Basto, M. Serr, F. Scaerou, R. Karess, T. Hays, Kinetochore dynein: its dynamics and role in the transport of the Rough deal checkpoint protein, *Nat Cell Biol* 3(11) (2001) 1001-7.

[209] S. Mische, Y. He, L. Ma, M. Li, M. Serr, T.S. Hays, Dynein light intermediate chain: an essential subunit that contributes to spindle checkpoint inactivation, *Mol Biol Cell* 19(11) (2008) 4918-29.

[210] M.V. Sivaram, T.L. Wadzinski, S.D. Redick, T. Manna, S.J. Doxsey, Dynein light intermediate chain 1 is required for progress through the spindle assembly checkpoint, *EMBO J* 28(7) (2009) 902-14.

[211] R. Gassmann, A.J. Holland, D. Varma, X. Wan, F. Civril, D.W. Cleveland, K. Oegema, E.D. Salmon, A. Desai, Removal of Spindly from microtubule-attached kinetochores controls spindle checkpoint silencing in human cells, *Genes Dev* 24(9) (2010) 957-71.

[212] K.D. Brown, K.W. Wood, D.W. Cleveland, The kinesin-like protein CENP-E is kinetochore-associated throughout poleward chromosome segregation during anaphase-A, *J Cell Sci* 109 (Pt 5) (1996) 961-9.

[213] H. Liao, R.J. Winkfein, G. Mack, J.B. Rattner, T.J. Yen, CENP-F is a protein of the nuclear matrix that assembles onto kinetochores at late G2 and is rapidly degraded after mitosis, *J Cell Biol* 130(3) (1995) 507-18.

[214] M.A. Lampson, E.L. Grishchuk, Mechanisms to Avoid and Correct Erroneous Kinetochore-Microtubule Attachments, *Biology (Basel)* 6(1) (2017).

[215] J.P. Welburn, M. Vleugel, D. Liu, J.R. Yates, 3rd, M.A. Lampson, T. Fukagawa, I.M. Cheeseman, Aurora B phosphorylates spatially distinct targets to differentially regulate the kinetochore-microtubule interface, *Mol Cell* 38(3) (2010) 383-92.

[216] J. Maciejowski, H. Drechsler, K. Grundner-Culemann, E.R. Ballister, J.A. Rodriguez-Rodriguez, V. Rodriguez-Bravo, M.J.K. Jones, E. Foley, M.A. Lampson, H. Daub, A.D. McAinsh, P.V. Jallepalli, Mps1 Regulates Kinetochore-Microtubule Attachment Stability via the Ska Complex to Ensure Error-Free Chromosome Segregation, *Dev Cell* 41(2) (2017) 143-156 e6.

[217] D. Liu, M. Vleugel, C.B. Backer, T. Hori, T. Fukagawa, I.M. Cheeseman, M.A. Lampson, Regulated targeting of protein phosphatase 1 to the outer kinetochore by KNL1 opposes Aurora B kinase, *J Cell Biol* 188(6) (2010) 809-20.

[218] M. Posch, G.A. Khoudoli, S. Swift, E.M. King, J.G. Deluca, J.R. Swedlow, Sds22 regulates aurora B activity and microtubule-kinetochore interactions at mitosis, *J Cell Biol* 191(1) (2010) 61-74.

[219] K. Schleicher, M. Porter, S. Ten Have, R. Sundaramoorthy, I.M. Porter, J.R. Swedlow, The Ndc80 complex targets Bod1 to human mitotic kinetochores, *Open Biol* 7(11) (2017).

[220] I.M. Cheeseman, S. Anderson, M. Jwa, E.M. Green, J. Kang, J.R. Yates, 3rd, C.S. Chan, D.G. Drubin, G. Barnes, Phospho-regulation of kinetochore-microtubule attachments by the Aurora kinase Ipl1p, *Cell* 111(2) (2002) 163-72.

[221] J.G. DeLuca, Y. Dong, P. Hergert, J. Strauss, J.M. Hickey, E.D. Salmon, B.F. McEwen, Hec1 and nuf2 are core components of the kinetochore outer plate essential for organizing microtubule attachment sites, *Mol Biol Cell* 16(2) (2005) 519-31.

[222] K.F. DeLuca, S.M. Lens, J.G. DeLuca, Temporal changes in Hec1 phosphorylation control kinetochore-microtubule attachment stability during mitosis, *J Cell Sci* 124(Pt 4) (2011) 622-34.

[223] A. Musacchio, The Molecular Biology of Spindle Assembly Checkpoint Signaling Dynamics, *Curr Biol* 25(20) (2015) R1002-18.

[224] N. London, S. Biggins, Mad1 kinetochore recruitment by Mps1-mediated phosphorylation of Bub1 signals the spindle checkpoint, *Genes Dev* 28(2) (2014) 140-52.

[225] A. Musacchio, E.D. Salmon, The spindle-assembly checkpoint in space and time, *Nat Rev Mol Cell Biol* 8(5) (2007) 379-93.

[226] N. London, S. Biggins, Signalling dynamics in the spindle checkpoint response, *Nat Rev Mol Cell Biol* 15(11) (2014) 736-47.

[227] S.T. Pachis, G. Kops, Leader of the SAC: molecular mechanisms of Mps1/TTK regulation in mitosis, *Open Biol* 8(8) (2018).

[228] G.V. Caldas, J.G. DeLuca, KNL1: bringing order to the kinetochore, *Chromosoma* 123(3) (2014) 169-81.

[229] Z. Dou, D.K. Prifti, P. Gui, X. Liu, S. Elowe, X. Yao, Recent Progress on the Localization of the Spindle Assembly Checkpoint Machinery to Kinetochores, *Cells* 8(3) (2019).

[230] C. Wurzenberger, D.W. Gerlich, Phosphatases: providing safe passage through mitotic exit, *Nat Rev Mol Cell Biol* 12(8) (2011) 469-82.

Figure 1. Cell cycle specific events result in changes to centromere.





Figure 2. Network of interactions within the inner kinetochore

A.



B.




Figure 3. Outer kinetochore assembly throughout mitosis

