

Simplifying 4D $\mathcal{N}=3$ harmonic superspace

Dharmesh Jain^{*}

Department of Theoretical Sciences, S. N. Bose National Centre for Basic Sciences,
Block JD, Sector III, Salt Lake City, Kolkata 700106, India

Chia-Yi Ju[†]

Department of Physics, National Chung Hsing University,
145 Xingda Road, South District, Taichung City 40227, Taiwan

Warren Siegel[‡]

C. N. Yang Institute for Theoretical Physics, Stony Brook University,
100 Nicolls Road, Stony Brook, New York 11794-3840, USA



(Received 5 August 2020; accepted 27 August 2020; published 24 September 2020)

We quantize super Yang-Mills action in $\mathcal{N} = 3$ harmonic superspace using “Fermi-Feynman” gauge and also develop the background field formalism. This leads to simpler propagators and Feynman rules that are useful in performing explicit calculations. The superspace rules are used to show that divergences do not appear at one loop and beyond. We also compute a finite contribution to the effective action from a four-point diagram at one loop, which matches the expected covariant result.

DOI: 10.1103/PhysRevD.102.066007

I. INTRODUCTION

$\mathcal{N} = 3$ harmonic superspace in four dimensions was developed by GIKOS around three and a half decades ago [1,2], and it provided the first successful off-shell formulation of four-dimensional (4D) $\mathcal{N} = 3$ super Yang-Mills (SYM) theory. This theory was quantized in “Landau” gauge a few years later by Delduc and McCabe [3]; however, the propagators obtained did not lend themselves to easier calculations. It is well known that the field content of a $\mathcal{N} = 3$ vector multiplet is the same as that of a $\mathcal{N} = 4$ one, and Zupnik explicitly showed this hidden supersymmetry of the $\mathcal{N} = 3$ SYM in [4]. The $\mathcal{N} = 3$ superspace also manifests the full superconformal symmetry, and using such symmetry arguments, low-energy effective action for $\mathcal{N} = 3$ and $\mathcal{N} = 4$ were considered by Zupnik and collaborators in [5–8]. A “twistorial” perspective on the $\mathcal{N} = 3$ SYM action was presented in [9] a few years ago, but no concrete progress has yet been made “to bring the quantization scheme into a form suitable for computations” [2].

We present evidence of some progress in the direction of simplifying computations here. We choose “Fermi-Feynman” gauge to drastically reduce the number ($9 \rightarrow 1$) and simplify the form ($\{\text{chiral, antichiral, linear}\}$ -analytic \rightarrow just analytic) of propagators when compared to [3]. This simplifies the proof of the nonrenormalization theorem as one might expect. Moreover, we also introduce the background field formalism in the $\mathcal{N} = 3$ harmonic superspace to simplify computations further.

In Sec. II, we review the basic ingredients of the $\mathcal{N} = 3$ harmonic superspace and the $\mathcal{N} = 3$ SYM action. In Sec. III, we introduce the Fermi-Feynman gauge to gauge fix this SYM action and derive the propagators. As an application, we prove the nonrenormalization theorem. In Sec. IV, we introduce the background field gauge to simplify the diagrammatic computations and present a sample calculation. Finally, we conclude with some discussion in Sec. V.

II. REVIEW

Our notation will closely follow [3], and we review it here for orientation purposes. The full 4D $\mathcal{N} = 3$ superspace has the usual set of ordinary bosonic ($x^{\alpha\dot{\alpha}}$) and fermionic ($\theta_i^\alpha, \bar{\theta}^{i\dot{\alpha}}$) coordinates with $i = 1, 2, 3$. The harmonic superspace augments these with six internal bosonic coordinates of the R -symmetry coset $SU(3)/U(1) \times U(1)$, denoted collectively as u . Using these internal coordinates, an “analytic” subspace with eight out of the twelve θ ’s of the full superspace is identified, which allows one to

^{*}dharmesh.jain@bose.res.in

[†]cju@nchu.edu.tw

[‡]siegel@insti.physics.sunysb.edu

<http://insti.physics.sunysb.edu/~siegel/plan.html>

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP³.

construct an off-shell action for the $\mathcal{N} = 3$ SYM and proceed with its quantization. In this section, we discuss the internal coordinates in some detail first, then the fermionic ones, and finally the superspace action of $\mathcal{N} = 3$ SYM. The quantization is dealt with in subsequent sections.

A. Internal coordinates

A $SU(3)/U(1) \times U(1)$ coset element can be parametrized in matrix form as follows:

$$\mathcal{U} = (u_i^1, u_i^2, u_i^3) \equiv (u_i^{(1,1)}, u_i^{(-1,1)}, u_i^{(0,-2)}),$$

$$\mathcal{U}^\dagger = \begin{pmatrix} \bar{u}_1^i \\ \bar{u}_2^i \\ \bar{u}_3^i \end{pmatrix} \equiv \begin{pmatrix} \bar{u}^{i(-1,-1)} \\ \bar{u}^{i(1,-1)} \\ \bar{u}^{i(0,2)} \end{pmatrix}, \quad (2.1)$$

Constraints: $\mathcal{U}^\dagger \mathcal{U} = \mathcal{U} \mathcal{U}^\dagger = 1$, $\det \mathcal{U} = 1$,

$$\Rightarrow \bar{u}_a \cdot u^b = \bar{u}_a^i u_i^b = \delta_a^b, \quad u_a^a \bar{u}_a^j = \delta_i^j, \quad (2.2)$$

$$\epsilon^{ijk} u_i^1 u_j^2 u_k^3 = 1.$$

The notation (q_1, q_2) denotes the charges corresponding to the two Cartan $U(1)$ generators Q_1, Q_2 of $SU(3)$. Given the constraints in Eq. (2.2), we have eight independent coordinates in the \mathcal{U} matrix, as expected for $SU(3)$. However, we also require the two $U(1)$ charges to be fixed (i.e., $Q_i u = q_i u$), which effectively implements the $U(1)^2$ quotient, and we are left with six independent coordinates.

Furthermore, the six harmonic covariant derivatives acting on these coordinates are

$$D_b^a = u_i^a \frac{\partial}{\partial u_i^b} - \bar{u}_b^i \frac{\partial}{\partial \bar{u}_a^i} \quad (a \neq b). \quad (2.3)$$

These derivatives satisfy the $SU(3)$ Lie algebra given by

$$[D_b^a, D_d^c] = \delta_b^c D_d^a - \delta_d^a D_b^c. \quad (2.4)$$

We note here that the two Cartan generators are given in terms of D_a^a (no sum over a) as follows:

$$\begin{aligned} Q_1 &= D_1^1 - D_2^2 & \Rightarrow & D_1^1 - D_3^3 = \frac{1}{2}(Q_1 + Q_2) \\ Q_2 &= D_1^1 + D_2^2 - 2D_3^3 & \Rightarrow & D_2^2 - D_3^3 = \frac{1}{2}(Q_2 - Q_1). \end{aligned} \quad (2.5)$$

Their commutators with the harmonic derivatives are $[Q_i, D_b^a] = q_i D_b^a$ with the charges given by

$$\begin{array}{c|ccccccc} U(1)^2 & D_3^1 & D_2^3 & D_2^1 & D_1^3 & D_3^2 & D_1^2 \\ \hline q_1 & 1 & 1 & 2 & -1 & -1 & -2 \\ q_2 & 3 & -3 & 0 & -3 & 3 & 0 \end{array}$$

In what follows, we will mostly be dealing with functions defined at two different points in this coset space, labeled as u and v . We denote their products by the notation $U^a{}_b = u^a \cdot \bar{v}_b$ and $\bar{U}_a{}^b = \bar{u}_a \cdot v^b$ such that the covariant derivatives in this basis simply read

$$D_b^a = U^a{}_c \frac{\partial}{\partial U^b{}_c} - \bar{U}_b{}^c \frac{\partial}{\partial \bar{U}_a{}^c} \quad (a \neq b). \quad (2.6)$$

Finally, the integration over this coset space is defined such that only a $SU(3)$ singlet integrand gives a non-vanishing result, i.e.,

$$\int du 1 = 1, \quad \int du D_b^a f(u) = 0. \quad (2.7)$$

The latter integral allows one to integrate by parts in the u space.

B. Fermionic coordinates

We make a coordinate transformation of the usual θ 's with $SU(3)$ indices to θ 's having definite $U(1)$ charges as follows:

$$\theta_a^\alpha = \bar{u}_a^i \theta_i^\alpha, \quad \bar{\theta}^{a\dot{\alpha}} = u_i^a \bar{\theta}^{i\dot{\alpha}}. \quad (2.8)$$

The index a identifies the $U(1)$ charges straightforwardly via Eq. (2.1). Then the corresponding spinorial covariant derivatives satisfy the following commutators:

$$\{D_\alpha^a, D_\beta^b\} = 0, \quad \{\bar{D}_{a\dot{\alpha}}, \bar{D}_{b\dot{\beta}}\} = 0, \quad \{D_\alpha^a, \bar{D}_{b\dot{\beta}}\} = i\delta_\alpha^a \partial_{a\dot{\beta}}, \quad (2.9)$$

$$[D_b^a, D_\alpha^c] = \delta_b^c D_\alpha^a, \quad [D_b^a, \bar{D}_{c\dot{\alpha}}] = -\delta_c^a \bar{D}_{b\dot{\alpha}}. \quad (2.10)$$

Explicitly, the $U(1)^2$ charges of the spinorial derivatives are

$$\begin{array}{c|cccccc} U(1)^2 & D_1^1 & D_2^2 & D_3^3 & \bar{D}_{1\dot{\alpha}} & \bar{D}_{2\dot{\alpha}} & \bar{D}_{3\dot{\alpha}} \\ \hline q_1 & 1 & -1 & 0 & -1 & 1 & 0 \\ q_2 & 1 & 1 & -2 & -1 & -1 & 2 \end{array}$$

The harmonic superspace is an analytic subspace of the full superspace, where the coordinates θ_1^α and $\bar{\theta}^{2\dot{\alpha}}$ do not appear explicitly in a given harmonic superfield $\Phi^{(q_1, q_2)}(x, \theta, u)$, i.e.,

$$D_\alpha^1 \Phi^{(q_1, q_2)} = \bar{D}_{2\dot{\alpha}} \Phi^{(q_1, q_2)} = 0. \quad (2.11)$$

Note that these analytic constraints are preserved by the three harmonic derivatives D_2^1, D_3^1, D_2^3 .

Finally, we can define an analytic measure $\int dud\zeta$ on harmonic superspace via the full superspace as follows:

$$\begin{aligned} \int d^4 x d^{12} \theta du &\equiv \int dud\zeta_{11}^{22} (D^1)^2 (\bar{D}_2)^2 \\ &\Rightarrow \int d\zeta_{11}^{22} = \int d^4 x_A (D^2)^2 (D^3)^2 (\bar{D}_1)^2 (\bar{D}_3)^2. \end{aligned} \quad (2.12)$$

We frequently use the notation $[D_\theta^4]_{22}^{11} \equiv (D^1)^2 (\bar{D}_2)^2$ to denote the four θ 's that are not part of the harmonic

superspace. The $[D_{\vartheta}^4]_{22}^{11}$ has $U(1)^2$ charge $(4,0)$, the negative of that for the measure $d\zeta_{11}^{22}$.

C. SYM action

We do not review here the procedure for finding prepotentials of the $\mathcal{N} = 3$ SYM in harmonic superspace but instead simply state the results. The $\mathcal{N} = 3$ prepotentials are the gauge connections of the analyticity-preserving harmonic derivatives, i.e., we have three connections defined by $\nabla = D + iA$. The gauge transformations read as usual: $\delta A = -\nabla\lambda$. The field strengths are introduced via the “flat” commutation relations as follows:

$$[\nabla_2^1, \nabla_3^1] = F_{23}^{11}, \quad [\nabla_2^3, \nabla_2^1] = F_{22}^{31}, \quad [\nabla_3^1, \nabla_2^3] = \nabla_2^1 + F_2^1. \quad (2.13)$$

The equations of motion are, of course, all $F = 0$. These are generated simply by the following Chern-Simons-like action:

$$\mathcal{S} = \text{tr} \int dud\zeta_{11}^{22} (A_2^1 F_2^1 + A_2^3 F_{23}^{11} + A_3^1 F_{22}^{31} - iA_2^1 [A_3^1, A_2^3]). \quad (2.14)$$

Also, notice that one of the three prepotentials is related algebraically to the other two on shell,

$$D_3^1 A_2^3 - D_2^3 A_3^1 + i[A_3^1, A_2^3] = A_2^1, \quad (2.15)$$

from which we start the quantization procedure in the next section.

III. QUANTIZING SYM IN FERMI-FEYNMAN GAUGE

The $\mathcal{N} = 3$ SYM action (2.14), after substituting the algebraic equation defining A_2^1 (2.15), depends only on two harmonic connections and reads

$$\begin{aligned} \mathcal{S} = \text{tr} \int du d\zeta_{11}^{22} & \{ (D_3^1 A_2^3)^2 + (D_2^3 A_3^1)^2 + 2A_3^1 (D_2^3 D_3^1 A_2^3) \\ & - 2A_3^1 D_2^1 A_2^3 + 2i[A_3^1, A_2^3] (D_3^1 A_2^3 - D_2^3 A_3^1) \\ & - [A_3^1, A_2^3]^2 \}. \end{aligned} \quad (3.1)$$

We choose the following gauge-fixing function:

$$\begin{aligned} \mathcal{S}_{gf} &= -\text{tr} \int dud\zeta_{11}^{22} (D_3^1 A_2^3 + D_2^3 A_3^1)^2 \\ &= -\text{tr} \int dud\zeta_{11}^{22} \{ (D_3^1 A_2^3)^2 + (D_2^3 A_3^1)^2 \\ & \quad - 2A_3^1 (D_2^3 D_3^1 A_2^3) \}, \end{aligned} \quad (3.2)$$

such that the gauge-fixed action for SYM in Fermi-Feynman gauge becomes

$$\begin{aligned} \mathcal{S} + \mathcal{S}_{gf} &= \text{tr} \int dud\zeta_{11}^{22} \{ 2A_3^1 (D_3^1 D_2^3 + D_2^3 D_3^1 - 2D_2^1) A_2^3 \\ & \quad + 2i[A_3^1, A_2^3] (D_3^1 A_2^3 - D_2^3 A_3^1) - [A_3^1, A_2^3]^2 \}, \end{aligned} \quad (3.3)$$

where we use $[D_3^1, D_2^3] = D_2^1$ once. The ghost action follows from the Becchi-Rouet-Stora-Tyutin formalism (BRST) straightforwardly by using $\delta A = -D\lambda - i[A, \lambda]$:

$$\begin{aligned} \mathcal{S}_{gh} &= -\text{tr} \int dud\zeta_{11}^{22} \{ b_2^1 (D_3^1 D_2^3 + D_2^3 D_3^1) c \\ & \quad + i(D_3^1 b_2^1 [A_2^3, c] + D_2^3 b_2^1 [A_3^1, c]) \}. \end{aligned} \quad (3.4)$$

Having just introduced new fields, let us recap the $U(1)^2$ charges of all the fields here (which are straightforwardly deduced from the covariant derivatives):

$U(1)^2$	A_3^1	A_2^3	A_2^1	λ	c	b_2^1
q_1	1	1	2	0	0	2
q_2	3	-3	0	0	0	0

A. Propagators

From the gauge-fixed SYM and ghost actions given above, we can derive the equations to solve for the Green’s functions for vector and ghost superfields:

$$\begin{aligned} (K_2^1)_0 G_{0|2}^{0|1}(1, 2) &= [\delta_A]_{2|2}^{1|1}(1, 2), \\ (K_2^1)_0 G_{2|0}^{1|0}(1, 2) &= [\delta_A]_{22|0}^{11|0}(1, 2), \\ (K_2^1)_{+1} G_{3|2}^{1|3}(1, 2) &= [\delta_A]_{23|2}^{11|3}(1, 2), \\ (K_2^1)_{-1} G_{2|3}^{3|1}(1, 2) &= [\delta_A]_{22|3}^{13|1}(1, 2), \end{aligned} \quad (3.5)$$

where $(K_2^1)_a = (1 + |a|)(\{D_3^1, D_2^3\} + 2aD_2^1)$ and the analytic delta functions explicitly read

$$\begin{aligned} [\delta_A]_{2|2}^{1|1}(1, 2) &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta_{2|1}^{1|2}(u, v) \\ &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) (U^1{}_1 \bar{U}_2{}^2) \delta^6(u, v), \\ [\delta_A]_{22|0}^{11|0}(1, 2) &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta_{22|11}^{11|22}(u, v) \\ &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) (U^1{}_1 \bar{U}_2{}^2)^2 \delta^6(u, v), \\ [\delta_A]_{23|2}^{11|3}(1, 2) &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta_{23|11}^{11|23}(u, v) \\ &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) (U^1{}_1)^3 \delta^6(u, v), \\ [\delta_A]_{22|3}^{13|1}(1, 2) &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta_{22|13}^{13|22}(u, v) \\ &= \delta(x_{12}) [D_{v\vartheta}^4]_{22}^{11} \delta^{12}(\theta_{12}) (\bar{U}_2{}^2)^3 \delta^6(u, v). \end{aligned} \quad (3.6)$$

The general form of G ’s which satisfy the Green’s function equations then looks like

$$\begin{aligned} \langle c(1)b_2^1(2) \rangle &\equiv G_{0|2}^{0|1}(1,2) \\ &= \mathcal{F}_{11|1}^{22|2}(u,v) \frac{1}{\square} [D_{u\theta}^4]_{22}^{11} [D_{v\theta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta(x_{12}), \end{aligned} \quad (3.7)$$

$$\begin{aligned} \langle b_2^1(1)c(2) \rangle &\equiv G_{2|0}^{1|0}(1,2) \\ &= \mathcal{F}_{1|11}^{2|22}(u,v) \frac{1}{\square} [D_{u\theta}^4]_{22}^{11} [D_{v\theta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta(x_{12}), \end{aligned} \quad (3.8)$$

$$\begin{aligned} \langle A_3^1(1)A_2^3(2) \rangle &\equiv G_{3|2}^{1|3}(1,2) \\ &= \mathcal{F}_{13|11}^{22|23}(u,v) \frac{1}{\square} [D_{u\theta}^4]_{22}^{11} [D_{v\theta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta(x_{12}), \end{aligned} \quad (3.9)$$

$$\begin{aligned} \langle A_2^3(1)A_3^1(2) \rangle &\equiv G_{2|3}^{3|1}(1,2) \\ &= \mathcal{F}_{11|13}^{23|22}(u,v) \frac{1}{\square} [D_{u\theta}^4]_{22}^{11} [D_{v\theta}^4]_{22}^{11} \delta^{12}(\theta_{12}) \delta(x_{12}), \end{aligned} \quad (3.10)$$

such that

$$(K_2^1)_a \mathcal{F}_{\cdot|}^{|\cdot}(u,v) = \frac{1}{2} (D_1^2)^2 \delta_{\cdot|}^{|\cdot}(u,v). \quad (3.11)$$

The $\delta_{\cdot|}^{|\cdot}(u,v)$ functions are the same δ -functions appearing in the corresponding $[\delta_A]_{\cdot|}^{|\cdot}(1,2)$ defined in Eq. (3.6). Equation (3.11) is motivated by the identity¹

$$(D_1^2)^2 D_{u\theta}^4 D_{v\theta}^4 \delta^6(u,v) = 2 \square D_{v\theta}^4 \delta^6(u,v), \quad (3.12)$$

which can be used to prove that Eqs. (3.7)–(3.10) indeed satisfy Eq. (3.5).

In order to make the above equations simpler and more tractable, we choose the following “independent” internal coordinates in the $\{U^a{}_b, \bar{U}_a{}^b\}$ basis:

$$U^1{}_1, U^1{}_3, U^2{}_1, U^3{}_1, \bar{U}_2{}^1, \bar{U}_2{}^2, \bar{U}_2{}^3, \bar{U}_3{}^2, \quad (3.13)$$

and the “zero charge” δ -function in these coordinates reads

$$\begin{aligned} \delta^6(u,v) &= \pi U^1{}_1 \bar{U}_2{}^2 \delta(U^1{}_3) \delta(U^3{}_1) \delta(\bar{U}_2{}^3) \\ &\quad \times \delta(\bar{U}_3{}^2) \delta(U^2{}_1) \delta(\bar{U}_2{}^1). \end{aligned} \quad (3.14)$$

The rest of the U coordinates can be written in terms of the chosen ones as follows:

$$\begin{aligned} U^1{}_2 &= -\frac{U^1{}_1 \bar{U}_2{}^1 + U^1{}_3 \bar{U}_2{}^3}{\bar{U}_2{}^2}, \\ U^2{}_2 &= -\frac{U^1{}_1 U^2{}_1 \bar{U}_2{}^1 + U^1{}_3 U^2{}_1 \bar{U}_2{}^3 - \bar{U}_2{}^3 \bar{U}_3{}^2 - U^1{}_1}{U^1{}_1 \bar{U}_2{}^2}, \\ U^2{}_3 &= \frac{U^1{}_3 U^2{}_1 - \bar{U}_3{}^2}{U^1{}_1}, \\ U^3{}_2 &= -\frac{U^1{}_1 U^3{}_1 \bar{U}_2{}^1 + U^1{}_3 U^3{}_1 \bar{U}_2{}^3 + \bar{U}_2{}^2 \bar{U}_2{}^3}{U^1{}_1 \bar{U}_2{}^2}, \\ U^3{}_3 &= \frac{U^1{}_3 U^3{}_1 + \bar{U}_2{}^2}{U^1{}_1}, \\ \bar{U}_1{}^1 &= \frac{-U^2{}_1 \bar{U}_2{}^1 \bar{U}_2{}^2 - U^3{}_1 \bar{U}_2{}^1 \bar{U}_3{}^2 + U^1{}_3 U^3{}_1 + \bar{U}_2{}^2}{U^1{}_1 \bar{U}_2{}^2}, \\ \bar{U}_1{}^2 &= -\frac{U^2{}_1 \bar{U}_2{}^2 + U^3{}_1 \bar{U}_3{}^2}{U^1{}_1}, \\ \bar{U}_1{}^3 &= -\frac{U^2{}_1 \bar{U}_2{}^2 \bar{U}_2{}^3 + U^3{}_1 \bar{U}_2{}^3 \bar{U}_3{}^2 + U^1{}_1 U^3{}_1}{U^1{}_1 \bar{U}_2{}^2}, \\ \bar{U}_3{}^1 &= \frac{\bar{U}_2{}^1 \bar{U}_3{}^2 - U^1{}_3}{\bar{U}_2{}^2}, \quad \bar{U}_3{}^3 = \frac{\bar{U}_2{}^3 \bar{U}_3{}^2 + U^1{}_1}{\bar{U}_2{}^2}. \end{aligned} \quad (3.15)$$

The differential operators get modified as well, leading to the following expressions:

$$\begin{aligned} D^1{}_2 &= U^1{}_1 \frac{\partial}{\partial U^2{}_1}, \quad D^1{}_3 = U^1{}_1 \frac{\partial}{\partial U^3{}_1}, \\ D^3{}_2 &= U^3{}_1 \frac{\partial}{\partial U^2{}_1} - \bar{U}_2{}^2 \frac{\partial}{\partial \bar{U}_3{}^2}, \\ D^2{}_1 &= U^2{}_1 \frac{\partial}{\partial U^1{}_1} + U^2{}_3 \frac{\partial}{\partial U^1{}_3} - \bar{U}_1{}^2 \frac{\partial}{\partial \bar{U}_2{}^1} \\ &\quad - \bar{U}_1{}^2 \frac{\partial}{\partial \bar{U}_2{}^2} - \bar{U}_1{}^3 \frac{\partial}{\partial \bar{U}_2{}^3} \\ &= U^2{}_1 \frac{\partial}{\partial U^1{}_1} + \left(\frac{U^1{}_3 U^2{}_1 - \bar{U}_3{}^2}{U^1{}_1} \right) \frac{\partial}{\partial U^1{}_3} \\ &\quad - \left(\frac{-U^2{}_1 \bar{U}_2{}^1 \bar{U}_2{}^2 - U^3{}_1 \bar{U}_2{}^1 \bar{U}_3{}^2 + U^1{}_3 U^3{}_1 + \bar{U}_2{}^2}{U^1{}_1 \bar{U}_2{}^2} \right) \\ &\quad \times \frac{\partial}{\partial \bar{U}_2{}^1} + \left(\frac{U^2{}_1 \bar{U}_2{}^2 + U^3{}_1 \bar{U}_3{}^2}{U^1{}_1} \right) \frac{\partial}{\partial \bar{U}_2{}^2} \\ &\quad + \left(\frac{U^2{}_1 \bar{U}_2{}^2 \bar{U}_2{}^3 + U^3{}_1 \bar{U}_2{}^3 \bar{U}_3{}^2 + U^1{}_1 U^3{}_1}{U^1{}_1 \bar{U}_2{}^2} \right) \frac{\partial}{\partial \bar{U}_2{}^3}, \end{aligned} \quad (3.16)$$

$$\begin{aligned} D^3{}_1 &= U^3{}_1 \frac{\partial}{\partial U^1{}_1} + U^3{}_3 \frac{\partial}{\partial U^1{}_3} - \bar{U}_1{}^2 \frac{\partial}{\partial \bar{U}_3{}^2} \\ &= U^3{}_1 \frac{\partial}{\partial U^1{}_1} + \left(\frac{U^1{}_3 U^3{}_1 + \bar{U}_2{}^2}{U^1{}_1} \right) \frac{\partial}{\partial U^1{}_3} \\ &\quad + \left(\frac{U^2{}_1 \bar{U}_2{}^2 + U^3{}_1 \bar{U}_3{}^2}{U^1{}_1} \right) \frac{\partial}{\partial \bar{U}_3{}^2}, \end{aligned} \quad (3.17)$$

¹We will suppress the $SU(3)$ “indices” on $[D_\theta^4]_{22}^{11}$ from now on.

$$\begin{aligned} D^2_3 &= U^2_1 \frac{\partial}{\partial U^3_1} - \bar{U}_3^1 \frac{\partial}{\partial \bar{U}_2^1} - \bar{U}_3^2 \frac{\partial}{\partial \bar{U}_2^2} - \bar{U}_3^3 \frac{\partial}{\partial \bar{U}_2^3} \\ &= U^2_1 \frac{\partial}{\partial U^3_1} + \left(\frac{U^1_3 - \bar{U}_2^1 \bar{U}_3^2}{\bar{U}_2^2} \right) \frac{\partial}{\partial \bar{U}_2^1} - \bar{U}_3^2 \frac{\partial}{\partial \bar{U}_2^2} \\ &\quad - \left(\frac{U^1_1 + \bar{U}_2^3 \bar{U}_3^2}{\bar{U}_2^2} \right) \frac{\partial}{\partial \bar{U}_2^3}. \end{aligned} \quad (3.19)$$

Using all these expressions, we can write Eq. (3.11) explicitly in the following form:

$$(1+|a|) \left[U^3_1 \frac{\partial^2}{\partial U^2_1 \partial U^3_1} - \bar{U}_2^2 \frac{\partial^2}{\partial \bar{U}_3^2 \partial \bar{U}_1^3} \right. \\ \left. + \left(a + \frac{1}{2} \right) \frac{\partial}{\partial U^2_1} \right] \mathcal{F}_{\cdot\cdot}^{\cdot\cdot} = \Delta_{\cdot\cdot}^{\cdot\cdot} \delta(U^3_1) \delta(U^2_1) \delta(\bar{U}_3^2), \quad (3.20)$$

where

$$\Delta_{\cdot\cdot}^{\cdot\cdot} = \frac{\pi}{2} (U^1_1)^{n_1} (\bar{U}_2^2)^{n_2} \delta(U^1_3) \delta(\bar{U}_2^3) \delta''(\bar{U}_2^1) \quad \text{with} \\ (n_1, n_2) = \begin{cases} (-1, 2) & \text{for } \langle c(1) b_2^1(2) \rangle, \\ (0, 3) & \text{for } \langle b_2^1(1) c(2) \rangle, \\ (1, 1) & \text{for } \langle A_3^1(1) A_2^3(2) \rangle, \\ (-2, 4) & \text{for } \langle A_2^3(1) A_3^1(2) \rangle. \end{cases}$$

Relabeling $U^3_1 = x$, $U^1_3 = \hat{x}$, $U^2_1 = y$, $\bar{U}_2^1 = \bar{y}$, $\bar{U}_3^2 = z$, $\bar{U}_2^3 = \hat{z}$, $\bar{U}_2^2 = A$, $U^1_1 = B$, and $a + \frac{1}{2} = b$, we get a simple looking partial differential equation

$$(x \partial_x \partial_y - A \partial_x \partial_z + b \partial_y) \mathcal{F}_{\cdot\cdot}^{\cdot\cdot} \\ = \frac{\pi}{2(1+|a|)} B^{n_1} A^{n_2} \delta(x) \delta(\hat{x}) \delta(y) \delta''(\bar{y}) \delta(z) \delta(\hat{z}). \quad (3.21)$$

We solve it by choosing an ansatz of the form

$$\mathcal{F}_{\cdot\cdot}^{\cdot\cdot} = C_a A^p B^q \frac{y^r \bar{y}^s}{(\bar{y} + \frac{\epsilon^2}{y})^3} \left(\frac{Ay + xz}{B} \right)^t \left(\frac{B\bar{y} + \hat{x}\hat{z}}{A} \right)^{-1}, \quad (3.22)$$

where ϵ is an infinitesimal parameter and the exponents p , q , r , s , t along with the normalization factor C_a are to be determined. Plugging Eq. (3.22) into the lhs of Eq. (3.21), we find that the values $t = -b$, $r = b - 1$, $s = 0$ simplify the expression to a single term as follows:

$$\begin{aligned} \text{lhs of Eq. (3.21)} &= \frac{3C_a b A^{p+2} B^{q+b} y^{b+2} (Ay + xz)^{-b} \epsilon^2}{(Ay + xz)(B\bar{y} + \hat{x}\hat{z})(y\bar{y} + \epsilon^2)^4} \\ &= \frac{C_a b A^{p+2} B^{q+b} (A + \frac{xz}{y})^{-b}}{2(Ay + xz)(B\bar{y} + \hat{x}\hat{z})} \pi \delta(y) \delta''(\bar{y}) \\ &= \frac{\pi}{2} C_a b A^{p+2} B^{q+b} \left(A + \frac{xz}{y} \right)^{-b} \delta(x) \delta(z) \delta(\hat{x}) \delta(\hat{z}) \delta(y) \delta''(\bar{y}) \\ &= \frac{\pi}{2} C_a b B^{q+b} A^{p+2-b} \delta(x) \delta(\hat{x}) \delta(y) \delta''(\bar{y}) \delta(z) \delta(\hat{z}), \end{aligned} \quad (3.23)$$

where we use the fact that δ 's for y , \bar{y} are produced in the limit of $\epsilon \rightarrow 0$ and $y \rightarrow 0$ via the following identity:

$$\frac{(m+1)!(-y)^m \epsilon^2}{(y\bar{y} + \epsilon^2)^{m+2}} \rightarrow \pi \delta(y) \delta^{(m)}(\bar{y}), \quad (3.24)$$

and $\delta(z) = \frac{1}{z}$, etc., for rest of the four nonconjugate complex variables. Now, comparing Eq. (3.23) to the rhs of Eq. (3.21), we can deduce that $p = n_2 + b - 2$, $q = n_1 - b$, and $C_a = \frac{1}{b(1+|a|)}$. Thus, the final form of $\mathcal{F}_{\cdot\cdot}^{\cdot\cdot}$ that solves Eq. (3.21) reads

$$\begin{aligned} \mathcal{F}_{\cdot\cdot}^{\cdot\cdot} &= \frac{1}{b(1+|a|)} A^{n_2+b-2} B^{n_1-b} \\ &\quad \times \frac{y^{b-1}}{(\bar{y} + \frac{\epsilon^2}{y})^3} \left(\frac{Ay + xz}{B} \right)^{-b} \left(\frac{B\bar{y} + \hat{x}\hat{z}}{A} \right)^{-1}. \end{aligned} \quad (3.25)$$

Finally, the complete propagators in terms of the U variables read as follows:

$$\begin{aligned} G_{\cdot\cdot}^{\cdot\cdot}(1, 2) &= \frac{(\bar{U}_2^2)^{n_2+b-2} (U_1^1)^{n_1-b}}{b(1+|a|)} \frac{(U^2_1)^{b-1}}{(\bar{U}_2^1 + \frac{\epsilon^2}{U_1^2})^3} \\ &\quad \times \frac{(-\bar{U}_1^2)^{-b}}{(-U_1^2)^{-1}} \frac{1}{\square} D_{u\theta}^4 D_{v\theta}^4 \delta^{12}(\theta_{12}) \delta(x_{12}). \end{aligned} \quad (3.26)$$

B. Feynman rules

The Feynman rules are now derived as usual. The vector and ghost propagators are given in Eq. (3.26), but we reproduce them here individually with explicit harmonic factors in momentum space (replace $\frac{1}{\square} \delta(x_{12}) \rightarrow \frac{1}{-k^2}$):

$$\langle c(1)b_2^1(2) \rangle = \frac{2(\bar{U}_2^2)^{\frac{1}{2}}}{(U^1_1)^{\frac{3}{2}}} \frac{(U^2_1)^{-\frac{1}{2}}}{(\bar{U}_2^1 + \frac{\epsilon^2}{U^2_1})^3} \frac{(-\bar{U}_1^2)^{-\frac{1}{2}}}{U^1_2} \times \frac{1}{k^2} D_{u\theta}^4 D_{v\theta}^4 \delta^{12}(\theta_{12}), \quad (3.27)$$

$$\langle b_2^1(1)c(2) \rangle = \frac{2(\bar{U}_2^2)^{\frac{3}{2}}}{(U^1_1)^{\frac{1}{2}}} \frac{(U^2_1)^{-\frac{1}{2}}}{(\bar{U}_2^1 + \frac{\epsilon^2}{U^2_1})^3} \frac{(-\bar{U}_1^2)^{-\frac{1}{2}}}{U^1_2} \times \frac{1}{k^2} D_{u\theta}^4 D_{v\theta}^4 \delta^{12}(\theta_{12}), \quad (3.28)$$

$$\langle A_3^1(1)A_2^3(2) \rangle = \frac{(\bar{U}_2^2)^{\frac{1}{2}}}{3(U^1_1)^{\frac{1}{2}}} \frac{(U^2_1)^{\frac{1}{2}}}{(\bar{U}_2^1 + \frac{\epsilon^2}{U^2_1})^3} \frac{(-\bar{U}_1^2)^{-\frac{3}{2}}}{U^1_2} \times \frac{1}{k^2} D_{u\theta}^4 D_{v\theta}^4 \delta^{12}(\theta_{12}), \quad (3.29)$$

$$\langle A_2^3(1)A_3^1(2) \rangle = -\frac{(\bar{U}_2^2)^{\frac{3}{2}}}{(U^1_1)^{\frac{3}{2}}} \frac{(U^2_1)^{-\frac{3}{2}}}{(\bar{U}_2^1 + \frac{\epsilon^2}{U^2_1})^3} \frac{(-\bar{U}_1^2)^{\frac{1}{2}}}{U^1_2} \times \frac{1}{k^2} D_{u\theta}^4 D_{v\theta}^4 \delta^{12}(\theta_{12}), \quad (3.30)$$

where we have kept the ϵ prescription explicit. The vertices can be read from Eqs. (3.3) and (3.4):

$$\begin{aligned} & \langle (A_3^1)^a (A_2^3)^b (A_2^3)^c \rangle / \langle (A_2^3)^a (A_3^1)^b (A_3^1)^c \rangle \\ & \rightarrow 2 \int du d^8 \theta f^{abc} [D_3^1 / D_2^3], \\ & \langle (A_3^1)^a (A_2^3)^b (A_3^1)^c (A_2^3)^d \rangle \rightarrow i \int du d^8 \theta f^{abec} f^{ecd}, \\ & \langle (A_2^3)^a c^b (b_2^1)^c \rangle / \langle (A_3^1)^a c^b (b_2^1)^c \rangle \rightarrow - \int du d^8 \theta f^{abc} [D_3^1 / D_2^3], \end{aligned} \quad (3.31)$$

where $\int d^8 \theta \equiv (D^2)^2 (D^3)^2 (\bar{D}_1)^2 (\bar{D}_3)^2$, f^{abc} are structure constants of the gauge group and the harmonic derivatives act on the leg corresponding to the group index ‘ c ’.

Evaluating loop graphs

Let us first focus on one-loop graphs. We can generate $\int d^{12} \theta$ from the analytic measure at vertices by taking off one factor of D_θ^4 from the propagators. After this, we are left with $(v_3 + v_4) \int d^{12} \theta$ integrals from three- and four-point vertices, $p D_\theta^4 \delta^{12}(\theta)$ ’s from propagators. As usual, we need to saturate all but one $\int d^{12} \theta$, which means that we need to kill one of the $\delta^{12}(\theta)$. This is achieved by using three D_θ^4 ’s and the identity

$$\begin{aligned} D_{w\theta}^4 D_{v\theta}^4 D_{u\theta}^4 = & [((W^1_3)^2 (\bar{W}_2^3)^2 (V^1_2)^2 (\bar{V}_2^1)^2 + (W^1_3)^2 (\bar{W}_2^1)^2 (V^1_2)^2 (\bar{V}_2^3)^2 + (W^1_2)^2 (\bar{W}_2^3)^2 (V^1_3)^2 (\bar{V}_2^1)^2 \\ & + (W^1_2)^2 (\bar{W}_2^1)^2 (V^1_3)^2 (\bar{V}_2^3)^2) D_{u\theta}^8 - i (W^1_2)^2 (\bar{W}_2^3)^2 (V^1_3)^2 (\bar{V}_2^1)^2 D^3 \cdot \partial \cdot \bar{D}_3 (D^2)^2 (\bar{D}_1)^2 \\ & - \frac{1}{2} \square ((W^1_2)^2 (\bar{W}_2^3)^2 (V^1_3)^2 (\bar{V}_2^1)^2 (D^2)^2 (\bar{D}_1)^2 + (W^1_3)^2 (\bar{W}_2^3)^2 (V^1_3)^2 (\bar{V}_2^1)^2 (D^3)^2 (\bar{D}_1)^2 \\ & + (W^1_2)^2 (\bar{W}_2^3)^2 (V^1_3)^2 (\bar{V}_2^3)^2 (D^2)^2 (\bar{D}_3)^2 + (W^1_3)^2 (\bar{W}_2^3)^2 (V^1_3)^2 (\bar{V}_2^3)^2 (D^3)^2 (\bar{D}_3)^2)] D_{u\theta}^4, \end{aligned} \quad (3.32)$$

where $D_\theta^8 \equiv (D^2)^2 (D^3)^2 (\bar{D}_1)^2 (\bar{D}_3)^2$, $V^a_b \equiv v^a \bar{u}_b$, $\bar{W}_a^b \equiv \bar{w}_a u^b$, etc. The first term which contains $D_\theta^8 D_\theta^4$ [recall that $D_\theta^4 \equiv (D^1)^2 (\bar{D}_2)^2$] can be used to kill one $\delta^{12}(\theta)$. This means that a two-point function trivially vanishes, as it does not have enough D_θ^4 ’s, and a three-point function cannot have any divergent piece due to the presence of three \square ’s in the denominator, i.e., $\int \frac{d^4 k}{(k^2)^3}$ is finite. In fact, no higher-point function can have any divergent piece at one loop because the numerator can generate at most $(p-3)$ \square ’s (together with a D_θ^8) compared to p \square ’s in the denominator, so the difference is always $n \geq 3$, i.e., $\int \frac{d^4 k}{(k^2)^n}$ is finite.

This power counting readily generalizes to multiloop graphs because each one-loop subgraph needs to follow this procedure of D algebra, and hence the whole graph is

rendered finite. This proves the nonrenormalization theorem at all loops for $\mathcal{N} = 3$ SYM or, equivalently, $\mathcal{N} = 4$ SYM.

IV. QUANTIZING SYM IN BACKGROUND FIELD GAUGE

The computation of finite terms for loop graphs is still cumbersome with the Feynman rules discussed in the previous section because manifestly covariant expressions are not obtained for individual graphs. For that purpose, we develop the background field formalism in this section.

Let us gauge covariantize all the differential operators ($D \rightarrow \nabla = D + iA$). Then we do a background splitting of the connections in a straightforward manner, $A \rightarrow A_{bg} + a_q$, where the subscripts are suppressed in favor

of self-explanatory fonts. Next, we choose different representations for these connections: the “real” rep for background \mathcal{A} ’s, meaning that the three harmonic connections vanish ($\mathcal{A}_2^1 = \mathcal{A}_3^1 = \mathcal{A}_2^3 = 0$), and the “analytic” rep for the quantum \mathbf{a} ’s, meaning that the four fermionic connections vanish ($\mathbf{a}_\alpha^1 = \bar{\mathbf{a}}_{2\dot{\alpha}} = 0 \Rightarrow D_g \rightarrow \nabla_g \equiv \mathcal{D}_g$).² Let us write down the consequences of these choices on connections and field strengths from various commutators:

$$\begin{aligned} \{\nabla_\alpha^a, \bar{\nabla}_{b\dot{\beta}}\} &= i\delta_b^a \nabla_{\dot{\alpha}\dot{\beta}} \text{ (“unchanged”),} \\ \{\nabla_\alpha^a, \nabla_\beta^b\} &= \epsilon_{a\beta} \bar{W}^{ab}, \\ \{\bar{\nabla}_{a\dot{\alpha}}, \bar{\nabla}_{b\dot{\beta}}\} &= \epsilon_{\dot{\alpha}\dot{\beta}} W_{ab}, \\ [\nabla_b^a, \nabla_\alpha^c] &= \delta_b^c \nabla_\alpha^a, \\ [\nabla_b^a, \bar{\nabla}_{c\dot{\alpha}}] &= -\delta_c^a \bar{\nabla}_{b\dot{\alpha}}, \\ [\nabla_b^a, \nabla_d^c] &= \delta_b^c \nabla_d^a - \delta_d^a \nabla_b^c. \end{aligned} \quad (4.1)$$

Of course, the W ’s are antisymmetric in the two indices, and they satisfy a few Bianchi identities along with some analytic plus harmonic constraints [5,8]. The most relevant identity for us is

$$D_\alpha^a W_{bc} = \frac{1}{2} (\delta_b^a D_\alpha^k W_{kc} - \delta_c^a D_\alpha^k W_{kb}). \quad (4.2)$$

Another thing to note is that the spinorial background-covariant derivatives \mathcal{D}_α^1 and $\bar{\mathcal{D}}_{2\dot{\alpha}}$ still possess the structure of Eq. (2.9), so that the “background analytic” superfields can be defined: $\mathcal{D}_\alpha^1 \Phi = \bar{\mathcal{D}}_{2\dot{\alpha}} \Phi = 0$. Moreover, the fourth and fifth equations of Eq. (4.1) tell us that the harmonic connections we are most interested in are now background analytic. We will also need the following identity defining a generalized d’Alembertian:

$$\begin{aligned} \mathcal{D}_g^4 (\nabla_1^2)^2 \mathcal{D}_g^4 &= 2 \hat{\square} \mathcal{D}_g^4 \\ &= 2 [(\square - 2\bar{W}^{12} W_{21}) - \bar{\nabla}_1 \cdot \bar{\nabla}_a \bar{W}^{1a} \\ &\quad + 2\nabla^2 \cdot \mathcal{D}^1 W_{21} + \nabla_1^2 (\mathcal{D}^1)^2 W_{21}] \mathcal{D}_g^4. \end{aligned} \quad (4.3)$$

A. Feynman rules

Since the harmonic connections appearing in the action as well as the ghosts are purely quantum superfields but background analytic, only \square changes to $\hat{\square}$ in the propagators were derived in the previous section, whereas the vertices in the Lagrangian remain the same. However, we can expand the $\hat{\square}$ to get vertices with explicit field strengths, which leads to covariant results in the loop calculations directly. With this structure, we can write down the background Feynman rules as follows:

$$\begin{aligned} \text{Propagators: } & \begin{cases} \mathcal{F}_{\cdot\cdot\cdot}^1(u, v) \frac{1}{-k^2} \mathcal{D}_{u\theta}^4 \mathcal{D}_{v\theta}^4 \delta^{12}(\theta_{12}), & \text{ignore background vertices,} \\ \frac{1}{-k^2} \delta^8(\theta_{12}) \delta^6(u, v), & \text{consider all such vertices but one,} \end{cases} \\ \text{All but one background vertex: } & \int du d^8\theta (\hat{\square} - \square_0), \\ \text{One background vertex: } & \int du dv d^8\theta (K_2^1)_a \delta^6(u, v), \\ \text{All quantum vertices: } & \text{same as Eq. (3.31).} \end{aligned}$$

Note that the background d’Alembertian is expanded as $\square \equiv \mathcal{D}^{a\dot{\alpha}} \mathcal{D}_{a\dot{\alpha}} = \square_0 + \dots$, where \square_0 corresponds to $-k^2$ in the momentum space.

One-loop graph computation

Let us focus on the computation of a four-point function here. It is finite and, from the symmetry arguments of

Sec. 5.5 of [8], it is known to look like $(\bar{W}^{13} W_{23})^2$ in $\mathcal{N} = 3$ harmonic superspace.

We can obtain such a contribution by evaluating a bubble graph and expanding the $(\hat{\square} - \square_0)$ -vertex factors to get the relevant W ’s using Eqs. (4.3) and (4.2). The above Feynman rules give the following expression for the graph with vector loop shown in Fig. 1 (after performing θ and u integrals at two of the vertices):

$$\begin{aligned} \Gamma_4^{(A)} \sim \hat{\mathcal{A}}_4 \int du dv \int d^8\theta_{1,2} (\bar{\nabla}_1 \cdot \bar{\nabla}_3 \bar{W}^{13}(p_1)) \\ \times (\bar{\nabla}_1 \cdot \bar{\nabla}_3 \bar{W}^{13}(p_2)) ((U^2{}_1 U^1{}_2) \nabla^2 \cdot \nabla^3 W_{23}(p_3)) \\ \times \delta^8(\theta_{12}) (K_2^1)_{-1} \mathcal{F}_{13|11}^{22|23}(u, v) \mathcal{D}_{u\theta}^4(p_4) \mathcal{D}_{v\theta}^4(p_4) \\ \times \delta^{12}(\theta_{12}) \delta^6(u, v), \end{aligned} \quad (4.4)$$

²Such a choice was used in the case of $\mathcal{N} = 2$ projective superspace to construct the background field formalism [10]. It ensures that the effective action is independent of background fields with dimension 0 (like the harmonic connections), which is required for the nonrenormalization theorems to hold [11]. However, the simple splitting of $\mathcal{N} = 3$ prepotentials is reminiscent of the background field formalism for the $\mathcal{N} = 2$ harmonic superspace developed in [12], with further refinements and explicit calculations appearing in [13,14].




FIG. 1. One-loop four-point graph with external background field strengths.

where $\hat{\mathcal{A}}_4 \sim \int d^4k \frac{1}{(k_1^2)(k_2^2)(k_3^2)(k_4^2)}$ is the scalar box integral with the subscripts on loop momentum k denoting the external momenta (p_i) dependence. Note that we had to partially integrate $(K_2^1)_{+1}$ so that it changed to $(K_2^1)_{-1}$ acting on $\mathcal{F}_{\cdot\cdot}^{\cdot\cdot}$, which gives, using Eq. (3.20), $(K_2^1)_{-1}\mathcal{F}_{13|11}^{22|23}(u, v) = \frac{1}{2}(D_1^2)^2\delta^6(u, v) - 4\frac{\partial}{\partial U_1^2}\mathcal{F}_{13|11}^{22|23}$. The first term leads to a harmonic singularity with two

$\delta^6(u, v)$ functions, but this singularity will cancel with the analogous contribution $\Gamma_4^{(bc)}$ from the ghost loop in Fig. 1. So we focus only on the second term, which has no analog from the ghost loop graph (as $a = 0$) and thus gives the complete four-point function. To get rid of the $\delta^{12}(\theta_{12})$ function, eight spinorial derivatives should be gathered off of W 's in addition to \mathcal{D}_θ^4 as follows³:

$$\begin{aligned} \Gamma_4 &\sim \hat{\mathcal{A}}_4 \int dudv \int d^8\theta_{1,2} \bar{W}^{13}(p_1) \bar{W}^{13}(p_2) W_{23}(p_3) \delta^8(\theta_{12}) D_2^1 \mathcal{F}_{13|11}^{22|23}(u, v) \\ &\quad \times (\bar{\nabla}_1)^2 (\bar{\nabla}_3)^2 ((U^2{}_1 U^1{}_2) \nabla^2 \cdot \nabla^3) (\bar{U}_2^1 \bar{U}_2^1 U^2{}_1 U^1{}_2 \nabla^2 \cdot \nabla^3) W_{23}(p_4) \mathcal{D}_{v\theta}^4(p_4) \delta^{12}(\theta_{12}) \delta^6(u, v) \\ &\sim \hat{\mathcal{A}}_4 \int du \int d^8\theta \bar{W}^{13}(p_1) \bar{W}^{13}(p_2) W_{23}(p_3) W_{23}(p_4) \left[\left(\frac{U^1{}_1}{(\bar{U}_2^1)^4 (U^2{}_1)^2} \right) (U^2{}_1 U^1{}_2)^2 (\bar{U}_2^1)^2 \right]_{u \rightarrow v} \\ &\sim \hat{\mathcal{A}}_4 \int du \int d^8\theta \bar{W}^{13}(p_1) \bar{W}^{13}(p_2) W_{23}(p_3) W_{23}(p_4) \left[\frac{(U^1{}_2)^2}{(\bar{U}_2^1)^2} \right]_{u \rightarrow v}, \end{aligned} \quad (4.5)$$

where we use $\nabla_\theta^8 \mathcal{D}_\theta^4 \delta^{12}(\theta_{12}) = 1$ in the second step. In the last step, the apparent harmonic singularity cancels as we take the limit $u \rightarrow v$, leading to the expected result for the one-loop effective action,

$$\Gamma_4 \sim \int dud\zeta_{11}^{22} \hat{\mathcal{A}}_4 (\bar{W}^{13} W_{23})^2. \quad (4.6)$$

V. DISCUSSION

We have introduced a new gauge-fixing action to quantize $\mathcal{N} = 3$ SYM in $\mathcal{N} = 3$ harmonic superspace. This leads to simpler (and fewer) propagators for vector and ghost superfields in a Fermi-Feynman gauge. These are sufficient to prove the nonrenormalization theorem for $\mathcal{N} = 3, 4$ SYM at all loops. However, computation of loop graphs beyond the divergent terms can be simplified more with the background field formalism. With the background Feynman rules in

hand, we have computed the one-loop four-point contribution to the effective action, which gets a finite contribution purely from the vector loop diagram.

The way that we derived the harmonic propagators here is reminiscent of how $\mathcal{N} = 2$ projective superspace [15,16] is derived from $\mathcal{N} = 2$ harmonic superspace [2,17] in [18]: by choosing a special parametrization of the harmonic R -symmetry coordinates on $SU(2)/U(1) \simeq S^2$ to obtain a single (complex) coordinate on CP^1 that forms the internal coordinate for the projective case.⁴ Even though we did not take this route to its full conclusion, it should be possible to derive a $\mathcal{N} = 3$ projective superspace in such a way that it simplifies the $\int du$ integrals to something more tractable. For example, viewing the R -symmetry coset $SU(3)/U(1)^2 \simeq [SU(3)/(SU(2) \times U(1))] \times [SU(2)/U(1)] \simeq CP^2 \times CP^1$, one can expect to reduce the six $(x, \hat{x}, y, \bar{y}, z, \hat{z})$ R -symmetry coordinates of $\mathcal{N} = 3$ harmonic superspace to only three (x, y, z) for a possible $\mathcal{N} = 3$ projective superspace. This should then be followed by a projection of the harmonic

³The requirement of collecting eight spinorial derivatives to get ∇_θ^8 [which is analogous to extracting D_θ^8 from Eq. (3.32) while evaluating graphs in Fermi-Feynman gauge] is sufficient to make $\Gamma_2 = \Gamma_3 = 0$ identically at one loop.

⁴The relation between these two hyperspaces was explored from different points of view in [19,20].

gauge condition and equations of motion for the gauge and ghost fields to the projective superspace, which we leave for future work.

ACKNOWLEDGMENTS

D. J. thanks Daniel Butter and Yu-tin Huang for the encouraging discussions. D. J. would also like to thank the

Simons Center for Geometry and Physics for its hospitality during the “2016 Simons Summer Workshop,” where a part of this work was done. C.-Y. J. is supported in part by the National Center for Theoretical Sciences and the Ministry of Science and Technology, Taiwan, through MOST Grant No. 109-2811-M-005-509. W. S. is supported by NSF Grant No. PHY-1915093.

- [1] A. Galperin, E. Ivanov, S. Kalitsyn, V. Ogievetsky, and E. Sokatchev, Unconstrained off-shell $\mathcal{N} = 3$ supersymmetric Yang-Mills theory, *Classical Quantum Gravity* **2**, 155 (1985).
- [2] A. Galperin, E. Ivanov, V. Ogievetsky, and E. Sokatchev, *Harmonic Superspace*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2007).
- [3] F. Delduc and J. McCabe, The quantization of $\mathcal{N} = 3$ super Yang-Mills off-shell in harmonic superspace, *Classical Quantum Gravity* **6**, 233 (1989).
- [4] B. Zupnik, $\mathcal{N} = 4$ multiplets in $\mathcal{N} = 3$ harmonic superspace, *Theor. Math. Phys.* **140**, 1121 (2004).
- [5] E. Ivanov and B. Zupnik, $\mathcal{N} = 3$ supersymmetric Born-Infeld theory, *Nucl. Phys.* **B618**, 3 (2001).
- [6] I. Buchbinder, E. Ivanov, I. Samsonov, and B. Zupnik, Scale invariant low-energy effective action in $\mathcal{N} = 3$ SYM theory, *Nucl. Phys.* **B689**, 91 (2004).
- [7] I. Buchbinder, E. Ivanov, I. Samsonov, and B. Zupnik, Superconformal $\mathcal{N} = 3$ SYM low-energy effective action, *J. High Energy Phys.* **10** (2012) 001.
- [8] I. Buchbinder, E. Ivanov, and I. Samsonov, The low-energy $\mathcal{N} = 4$ SYM effective action in diverse harmonic superspaces, *Phys. Part. Nucl.* **48**, 333 (2017).
- [9] B. U. Schwab and C. Vergu, Twistors, harmonics and holomorphic Chern-Simons, *J. High Energy Phys.* **03** (2013) 046.
- [10] D. Jain and W. Siegel, Improved methods for hypergraphs, *Phys. Rev. D* **88**, 025018 (2013).
- [11] M. T. Grisaru and W. Siegel, Supergraphity: (II). Manifestly covariant rules and higher loop finiteness, *Nucl. Phys.* **B201**, 292 (1982); Erratum, *Nucl. Phys.* **206**, 496 (1982).
- [12] I. Buchbinder, E. Buchbinder, S. Kuzenko, and B. A. Ovrut, The background field method for $\mathcal{N} = 2$ super Yang-Mills theories in harmonic superspace, *Phys. Lett. B* **417**, 61 (1998).
- [13] I. L. Buchbinder, S. M. Kuzenko, and B. A. Ovrut, On the $D = 4$, $\mathcal{N} = 2$ nonrenormalization theorem, *Phys. Lett. B* **433**, 335 (1998).
- [14] I. L. Buchbinder and S. M. Kuzenko, Comments on the background field method in harmonic superspace: Non-holomorphic corrections in $\mathcal{N} = 4$ SYM, *Mod. Phys. Lett. A* **13**, 1623 (1998).
- [15] U. Lindström and M. Roček, New hyperkähler metrics and new supermultiplets, *Commun. Math. Phys.* **115**, 21 (1988).
- [16] U. Lindström and M. Roček, $\mathcal{N} = 2$ super Yang-Mills theory in projective superspace, *Commun. Math. Phys.* **128**, 191 (1990).
- [17] A. Galperin, E. Ivanov, S. Kalitsyn, V. Ogievetsky, and E. Sokatchev, Unconstrained $\mathcal{N} = 2$ matter, Yang-Mills and supergravity theories in harmonic superspace, *Classical Quantum Gravity* **1**, 469 (1984); Erratum, *Classical Quantum Gravity* **2**, 127 (1985).
- [18] D. Jain and W. Siegel, Deriving projective hyperspace from harmonic, *Phys. Rev. D* **80**, 045024 (2009).
- [19] S. M. Kuzenko, Projective superspace as a double punctured harmonic superspace, *Int. J. Mod. Phys. A* **14**, 1737 (1999).
- [20] D. Butter, Relating harmonic and projective descriptions of $\mathcal{N} = 2$ nonlinear sigma models, *J. High Energy Phys.* **11** (2012) 120.