

POINTWISE ERROR ESTIMATES FOR C^0 INTERIOR PENALTY APPROXIMATION OF BIHARMONIC PROBLEMS

D. LEYKEKHMAN

ABSTRACT. The aim of this paper is to derive pointwise global and local best approximation type error estimates for biharmonic problems using the C^0 interior penalty method. The analysis uses the technique of dyadic decompositions of the domain, which is assumed to be a convex polygon. The proofs require local energy estimates and new pointwise Green's function estimates for the continuous problem which has independent interest.

1. INTRODUCTION

We consider the fourth order problem:

$$(1) \quad \begin{aligned} \Delta^2 u &= f && \text{in } \Omega, \\ u &= \frac{\partial u}{\partial n} = 0 && \text{on } \partial\Omega, \end{aligned}$$

where $\Omega \subset \mathbb{R}^2$ is a convex polygonal domain, $f \in L^2(\Omega)$, and n denotes the outward unit normal of $\partial\Omega$. Finite element discretization of the above problem is not straightforward and various approaches to approximate the above problem were proposed over the years. However, they all have some drawbacks. For instance, conformal C^1 elements are rather complicated even in two dimensions [2, 7], the classical non-conformal elements [13, 20] must be altered in the presence of low order terms (cf. [14]). Furthermore, the construction of higher order nonconforming finite elements for fourth order problems is also not easy. The C^0 interior penalty method is a sound alternative. This method is attractive since the finite elements consist of usual Lagrange elements of arbitrary order and straightforward to implement. A detailed description of the method with energy based error estimates on convex and nonconvex domains can be found in [4].

Pointwise error estimates is well developed area for the second order problems. However, there are few such results for fourth order problems. Many such pointwise error estimates are obtained via Sobolev embedding. This is not satisfactory since such results are usually not optimal and often the discrepancy between norms makes them hard to use for applications, for example in optimal control problems. In addition, it is hard to localize them. The only exceptions we are aware of are the papers by Rolf Rannacher [16] and Ming Wang [23], where pointwise error estimates were established for some nonconforming and mixed elements. Both

Received by the editor July 21, 2019, and, in revised form, August 16, 2020, and August 29, 2020.

2020 *Mathematics Subject Classification*. Primary 65N30, 65N15.

Key words and phrases. Maximum norm, finite element method, pointwise error estimates, Green's function, biharmonic equation, interior penalty, local error estimates.

The author was partially supported by NSF grant DMS-1913133.

papers used a weighted technique, and their $W^{1,\infty}$ error estimates are in the form of quasi-optimal order error estimates with $H^3(\Omega)$ norm on the right hand side, which is consistent with regularity of the biharmonic problem on convex domains. However, the error estimate for the second derivatives in [16], which is important for computing bending moments, requires $W^{3,\infty}$ and H^4 regularity for the solution, that can not be guaranteed on convex polygons. In addition, it is not clear how to localize those results.

In this paper we take a different approach and show the following best approximation type results.

Theorem 1.1. *Let u satisfy (1) and u_h be its C^0 interior penalty finite element approximation, and x_0 be an arbitrary point of $\bar{\Omega}$. Then there exists a constant C independent of h such that*

$$|D^2(u - u_h)(x_0)| \leq C |\ln h|^{\frac{3}{2}} \min_{\chi \in S_h^r} \|\|u - \chi\|\|_{W_h^{2,\infty}(\Omega)},$$

where D^2 denotes a general second order differential operator and $W_h^{2,\infty}$ is a mesh dependent norm defined in (17c).

There is some ambiguity in the meaning of $D^2u_h(x_0)$ in the case of x_0 be on the edge of two elements. In this situation the value of $D^2u_h(x_0)$ can be taken by restricting u_h to either element. We also want to point out that from the Theorem 3.4.4 from [8], for example, it follows that on convex polygonal domains the solution u to (1) is in $H^{3+\varepsilon}(\Omega)$ for some $\varepsilon > 0$ and by the Sobolev embedding in two dimensions $u \in W^{2,\infty}(\Omega)$. As a result the theorem makes sense without any additional smoothness assumptions on the solution or domain. In general, we can not expect even a first order convergence since the solution is not in $W^{3,\infty}(\Omega)$. However, similarly to the second order elliptic equations [17, 18], we can establish the following local result using which we can obtain first order convergence in the interior of the convex domains without any additional assumptions on the regularity.

Theorem 1.2 (Local error estimate). *Let and x_0 be an arbitrary point of $\bar{\Omega}$ and define $B_d = B_d(x_0) \cap \Omega$, where $B_d(x_0)$ denotes a ball of radius d centered at x_0 . Then there exists a constant C independent of h and d such that*

$$|D^2(u - u_h)(x_0)| \leq C |\ln h|^{\frac{3}{2}} \left(\|\|u - \chi\|\|_{W_h^{2,\infty}(B_d)} + d^{-1} \|\|u - \chi\|\|_{2,h,\Omega} \right),$$

where $\|\cdot\|_{W_h^{2,\infty}(B_d)}$ and $\|\cdot\|_{2,h,\Omega}$ are mesh dependent norms defined in the next section.

The analysis of the method uses a combination of well-established technique of dyadic decomposition together with local energy estimates for second order elliptic problem [22] and pointwise Green's function estimates. In this paper we only consider two-dimensional convex polygonal domains. However, even on such simple domains we had to overcome several technical difficulties in order to obtain the results. Our error analysis framework has a lot in common with pointwise error analysis for discontinuous Galerkin methods for the second order elliptic problem [5, 9]. The details are of course rather different and the main difficulty lies in the regularity results. In contrast to the second order elliptic problems on convex domains, where H^2 norm of the solution is bounded by the L^2 of the right hand side, for the biharmonic problem we can control H^3 norm of the solution by H^{-1} norm of the right hand side, which causes many difficulties due to the global nature of the H^{-1}

norm. Although pointwise error analysis for the second order problem is available in three dimensions too, the pointwise Green's function estimates and the duality argument are the main obstacles for extending our main results to non-convex case or to three dimensions.

The rest of the paper is organized as follows. In the next chapter we introduce notation and some basic results for continuous problem. In chapter 3, we define the C^0 interior penalty discretization, and establish key lemmas. The central result is the Local Energy Estimates, Lemma 3.9. In chapter 4 we provide a proof of the global pointwise error estimate, namely Theorem 1.1 and in the chapter 5, we establish a localized version, Theorem 1.2.

2. NOTATION AND PRELIMINARY RESULTS

In the paper we use the usual notation for Sobolev and Lebesgue spaces. We denote by (\cdot, \cdot) the inner product in $L^2(\Omega)$, and by $(\cdot, \cdot)_{\Omega_0}$ the L^2 inner product in over a subdomain $\Omega_0 \subset \Omega$.

We let D denote a general first order differential operator, D^2 - second order differential operator, D^3 - third and etc. The partial derivatives we will denote by ∂ . We will also use the multi-index notation when it is important.

The weak solution u to (1) given by

$$(2) \quad B(u, \varphi) := (D^2 u, D^2 \varphi) = \sum_{i,j=1}^2 (\partial_{ij}^2 u, \partial_{ij}^2 \varphi) = (f, \varphi), \quad \forall \varphi \in H_0^2(\Omega),$$

is naturally in $H_0^2(\Omega)$ with the following estimate

$$(3) \quad \|u\|_{H^2(\Omega)} \leq C \|f\|_{H^{-2}(\Omega)}.$$

On convex domains we also have that $u \in H_0^2(\Omega) \cap H^3(\Omega)$ with the following estimate (cf. [10] sec. 5.9)

$$(4) \quad \|u\|_{H^3(\Omega)} \leq C \|f\|_{H^{-1}(\Omega)}.$$

2.1. Green's function estimates. In the proof of our main results we will make a heavy use of pointwise estimates for the Green's function and its derivatives. The following form of the estimate seems to be new for convex polygons, but follows rather easily from available results.

Lemma 2.1. *Let Ω be a bounded convex polygonal domain in \mathbb{R}^2 and $G(x, y)$ be the corresponding Green's function for the problem (1). Then for all multi-indices $\alpha = (\alpha_1, \alpha_2)$ and $\beta = (\beta_1, \beta_2)$ with the range $1 \leq |\alpha|, |\beta| \leq 2$ and $3 \leq |\alpha| + |\beta| \leq 4$ the following estimates hold,*

$$(5) \quad |D_x^\alpha D_y^\beta G(x, y)| \leq C |x - y|^{2-|\alpha|-|\beta|}.$$

Proof. The estimate (5) essentially follows from the estimates from [12] (see also [11, p. 286]). Let $\omega_1, \omega_2, \dots, \omega_M$ be the interior angles of the convex polygon Ω . Let ω be an arbitrary angle, and let $\rho(x)$ denotes a distance from x to the vertex of ω .

Theorem 2.1 in [12] states that for Green's function the following estimates hold

(6a)

$$|D_x^\alpha D_y^\beta G(x, y)| \leq C|x - y|^{2-|\alpha|-|\beta|}, \quad \text{for } \rho(x)/2 \leq \rho(y) \leq 2\rho(x), \quad 3 \leq |\alpha| + |\beta|,$$

(6b)

$$|D_x^\alpha D_y^\beta G(x, y)| \leq C\rho(x)^{\gamma-|\alpha|}\rho(y)^{-\gamma+2-|\beta|}, \quad \text{for } 2\rho(x) \leq \rho(y), \quad 0 \leq |\alpha| + |\beta|,$$

(6c)

$$|D_x^\alpha D_y^\beta G(x, y)| \leq C\rho(x)^{-\gamma+2-|\alpha|}\rho(y)^{\gamma-|\beta|}, \quad \text{for } 2\rho(y) \leq \rho(x), \quad 0 \leq |\alpha| + |\beta|,$$

where γ is any real number that satisfies inequality $|\gamma - 1| < C_\omega$. Since $\omega < \pi$, according to Lemma 1.2 in [12] (see also [3]), $C_\omega > 1$ and as a result we may take $\gamma = 2 + \varepsilon$ for some $\varepsilon > 0$.

With this choice of γ from (6b) it follows that

$$|D_x^\alpha D_y^\beta G(x, y)| \leq C\rho(x)^{2+\varepsilon-|\alpha|}\rho(y)^{-2-\varepsilon+2-|\beta|} \leq C\rho(y)^{2-|\alpha|-|\beta|},$$

where we used that this is the case when $2\rho(x) \leq \rho(y)$ and $2 + \varepsilon - |\alpha| > 0$. Using the triangle inequality

$$|x - y| \leq \rho(y) + \rho(x) \leq \frac{3}{2}\rho(y)$$

and that by the assumptions on $2 - |\alpha| - |\beta| < 0$ we have that in this case

$$(7) \quad |D_x^\alpha D_y^\beta G(x, y)| \leq C\rho(y)^{2-|\alpha|-|\beta|} \leq C|x - y|^{2-|\alpha|-|\beta|}.$$

The case $2\rho(y) \leq \rho(x)$ is very similar. From (6c) with $\gamma = 2 + \varepsilon$ we have

$$|D_x^\alpha D_y^\beta G(x, y)| \leq C\rho(x)^{-2-\varepsilon+2-|\alpha|}\rho(y)^{2+\varepsilon-|\beta|} \leq C\rho(x)^{2-|\alpha|-|\beta|}.$$

Again using the triangle inequality

$$|x - y| \leq \rho(x) + \rho(y) \leq \frac{3}{2}\rho(x)$$

together with the assumption $2 - |\alpha| - |\beta| < 0$ we obtain

$$(8) \quad |D_x^\alpha D_y^\beta G(x, y)| \leq C\rho(x)^{2-|\alpha|-|\beta|} \leq C|x - y|^{2-|\alpha|-|\beta|}.$$

Combining (7), (8), and (6a), we obtain the lemma. \square

3. DISCRETIZATION

To define finite element approximation of the solution to (1), for $h \in (0, h_0]$; $h_0 > 0$, we let \mathcal{T}_h denote a quasi-uniform triangulation of Ω with mesh size h , i.e., $\mathcal{T}_h = \{T\}$ is a partition of Ω into triangles T of diameter h_T such that for $h = \max_T h_T$,

$$\text{diam}(T) \leq h \leq C|T|^{\frac{1}{2}}, \quad \text{for all } T \in \mathcal{T}_h.$$

Let $S_h^r \subset H_0^1(\Omega)$ denote the Lagrange finite element space consisting of polynomials of degree $r \geq 2$; that is,

$$S_h^r = \{v_h \in H_0^1(\Omega); v_h|_T \in \mathbb{P}^r(T) \ \forall T \in \mathcal{T}_h\}.$$

By $I_h : C^0(\Omega) \rightarrow S_h^r$ we denote the usual nodal interpolant which satisfies

$$(9) \quad \|v - I_h v\|_{W^{j,p}(T)} \leq Ch^{s-j}\|v\|_{W^{s,p}(T)}, \quad \forall v \in W^{s,p}(T),$$

for $1 \leq p \leq \infty$, $j \leq s \leq r + 1$, and $s > \frac{2}{p}$.

3.1. Trace and inverse inequalities. We will frequently use the following trace and inverse inequalities. For $T \in \mathcal{T}_h$ and $v \in H_0^2(\Omega)$, $v_h \in S_h^r$ there exist positive constants C_{tr} and C_{inv} independent of τ and v, v_h such that

$$(10a) \quad \|v\|_{L^2(\partial T)}^2 \leq C_{tr}^2 \|v\|_{L^2(T)} \|\nabla v\|_{L^2(T)} \leq C_{tr}^2 (h^{-1} \|v\|_{L^2(T)}^2 + h \|\nabla v\|_{L^2(T)}^2),$$

$$(10b) \quad \|\nabla v_h\|_{L^2(T)} \leq C_{inv} h^{-1} \|v_h\|_{L^2(T)},$$

$$(10c) \quad \|v_h\|_{L^2(\partial T)} \leq C_{tr} (1 + C_{inv}) h^{-1/2} \|v_h\|_{L^2(T)}.$$

3.2. The C^0 interior penalty method. To define the method, we need some additional notation. Let \mathcal{E}_h be the set of edges in \mathcal{T}_h . For $e \in \mathcal{E}_h$ and $v \in H^2(\Omega, \mathcal{T}_h)$, where

$$H^2(\Omega, \mathcal{T}_h) = \{v \in L^2(\Omega) : v|_T \in H^2(T) \quad \forall T \in \mathcal{T}_h\},$$

we define the jump $\llbracket \frac{\partial v}{\partial n} \rrbracket$ of the normal derivative of v across an edge e and the average of the second normal derivative $\left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\}$ of v on an edge e as follows. If $e \subset \Omega$, we take n_e to be one of the two unit vectors normal to e . Then e is the common side of two triangles $T_+ \in \mathcal{T}_h$ and $T_- \in \mathcal{T}_h$, where n_e is pointing from T_- to T_+ . Thus, on such e we define

$$\llbracket \frac{\partial v}{\partial n} \rrbracket = \frac{\partial v_{T_+}}{\partial n}|_e - \frac{\partial v_{T_-}}{\partial n}|_e \quad \text{and} \quad \left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} = \frac{\partial^2 v_{T_+}}{\partial n^2}|_e + \frac{\partial^2 v_{T_-}}{\partial n^2}|_e.$$

We note that the above definitions do not depend on the choice of n_e . If $e \subset \partial\Omega$, we take n_e to be the unit normal pointing outside Ω and

$$\llbracket \frac{\partial v}{\partial n} \rrbracket = -\frac{\partial v}{\partial n_e} \quad \text{and} \quad \left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} = -\frac{\partial^2 v}{\partial n_e^2}.$$

Next, we define the bilinear form $B_h(\cdot, \cdot)$ by

$$\begin{aligned} B_h(v, w) = & \sum_{T \in \mathcal{T}_h} \int_T D^2 v : D^2 w \, dx \\ & + \sum_{e \in \mathcal{E}_h} \int_e \left(\left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} \llbracket \frac{\partial w}{\partial n} \rrbracket + \llbracket \frac{\partial v}{\partial n} \rrbracket \left\{ \left\{ \frac{\partial^2 w}{\partial n^2} \right\} \right\} + \frac{\eta}{|e|} \llbracket \frac{\partial v}{\partial n} \rrbracket \llbracket \frac{\partial w}{\partial n} \rrbracket \right) ds, \end{aligned}$$

where

$$D^2 v : D^2 w = \sum_{i,j=1}^2 \partial_{ij}^2 v \partial_{ij}^2 w.$$

The discrete problem is to find $u_h \in S_h^r$ s.t.

$$(11) \quad B_h(u_h, v_h) = (f, v_h), \quad \forall v_h \in S_h^r.$$

Similarly to [4] we define the following mesh-dependent norm

$$\|v\|_{2,h}^2 = \sum_{T \in \mathcal{T}_h} \|D^2 v\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} \left(|e| \left\| \left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 + |e|^{-1} \left\| \llbracket \frac{\partial v}{\partial n} \rrbracket \right\|_{L^2(e)}^2 \right).$$

Easy to see that the bilinear form $B_h(\cdot, \cdot)$ is bounded

$$(12) \quad B_h(v, w) \leq C \|v\|_{2,h} \|w\|_{2,h},$$

and for η sufficiently large, the bilinear form $B_h(\cdot, \cdot)$ is coercive (cf. [4]) on S_h^r ,

$$(13) \quad B_h(v_h, v_h) \geq C \|v_h\|_{2,h}^2, \quad \forall v_h \in S_h^r.$$

As a result by the Lax-Milgram lemma, u_h is well-defined. From [4, eq. (4.9)] we also have that the exact solution u satisfies

$$B_h(u, \chi) = (f, \chi), \quad \forall \chi \in S_h^r,$$

and as a result u and u_h satisfy the usual Galerkin orthogonality

$$(14) \quad B_h(u - u_h, \chi) = 0 \quad \forall \chi \in S_h^r.$$

In [4, Theorem 3], on convex domains the following error estimate was established

$$(15) \quad \|u - u_h\|_{2,h} \leq Ch\|f\|_{H^{-1}(\Omega)}.$$

Note that for $v_h \in S_h^r$ using the inequality (10c), we have

$$\sum_{e \in \mathcal{E}_h} |e| \left\| \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \leq C \sum_{T \in \mathcal{T}_h} \|D^2 v_h\|_{L^2(T)}^2,$$

and thus on S_h^r the $\|\cdot\|_{2,h}$ norm is equivalent to the following norm without the terms involving averages, namely

$$\|v\|_{2,h}^2 := \sum_{T \in \mathcal{T}_h} \|D^2 v\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \left[\left[\frac{\partial v}{\partial n} \right] \right] \right\|_{L^2(e)}^2.$$

We will also require the following norms on subsets $D \subset \Omega$. First we define

$$(16) \quad \mathcal{T}_h \cap D = \{T \in \mathcal{T}_h : T \cap D \neq \emptyset\} \quad \text{and} \quad \mathcal{E}_h \cap D = \{e \in \mathcal{E}_h : e \cap D \neq \emptyset\}.$$

Then,

$$(17a) \quad \|v\|_{W_h^{2,1}(D)} = \sum_{T \in \mathcal{T}_h \cap D} |v|_{W^{2,1}(T)} + \sum_{e \in \mathcal{E}_h \cap D} \int_e \left(|e| \left| \left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} \right| + \left| \left[\left[\frac{\partial v}{\partial n} \right] \right] \right) ds,$$

(17b)

$$\|v\|_{2,h,D}^2 = \sum_{T \in \mathcal{T}_h \cap D} |v|_{H^2(T)}^2 + \sum_{e \in \mathcal{E}_h \cap D} \left(|e| \left\| \left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 + |e|^{-1} \left\| \left[\left[\frac{\partial v}{\partial n} \right] \right] \right\|_{L^2(e)}^2 \right),$$

and

(17c)

$$\begin{aligned} \|v\|_{W_h^{2,\infty}(D)} &= \max_{T \in \mathcal{T}_h \cap D} |v|_{W^{2,\infty}(T)} + \max_{e \in \mathcal{E}_h \cap D} \left\| \left\{ \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right\} \right\|_{L^\infty(e)} + \max_{e \in \mathcal{E}_h \cap D} |e|^{-1} \left\| \left[\left[\frac{\partial v}{\partial n} \right] \right] \right\|_{L^\infty(e)}. \end{aligned}$$

Using the above norms we also have

$$(18) \quad B_h(v, w) \leq C \|v\|_{W_h^{2,1}(\Omega)} \|w\|_{W_h^{2,\infty}(\Omega)}.$$

3.3. Superapproximation. Superapproximation is an essential tool in pointwise finite element error estimates [15]. To describe them we denote by $B_r(x_0)$ a ball of radius $r > 0$ centered at x_0 and introduce a smooth cut-off function $\omega \in C^\infty(\Omega)$ with the properties:

$$(19a) \quad \omega(x) \equiv 1, \quad x \in B_{d/2}(x_0)$$

$$(19b) \quad \omega(x) \equiv 0, \quad x \in B_{3d/4}^c(x_0)$$

$$(19c) \quad |\omega|_{W_\infty^j} \leq C d^{-j}, \quad j = 0, 1, 2.$$

In [6] an improved superapproximation result was obtained

$$\|\omega^2\chi - I_h(\omega^2\chi)\|_{H^1(T)} \leq C \frac{h_T}{d} (\|\nabla(\omega\chi)\|_{L^2(T)} + d^{-1}\|\chi\|_{L^2(T)}), \quad \chi \in S_h^r.$$

We will follow the ideas from that paper. Let $P_1(\omega)$ and $P_0(\omega)$ be linear and constant order approximations to ω on element T , respectively, such that

$$(20) \quad \|\omega - P_0(\omega)\|_{L^\infty(T)} \leq Ch_T \|\omega\|_{W_\infty^1(T)} \leq Ch_T d^{-1}$$

$$(21) \quad \|\omega - P_1(\omega)\|_{W_\infty^s(T)} \leq Ch_T^{2-s} \|\omega\|_{W_\infty^2(T)} \leq Ch_T^{2-s} d^{-2}, \quad s = 0, 1.$$

We take $P_0(\omega) = \widehat{\omega} = |T|^{-1} \int_T \omega$ and $P_1(\omega)$ a linear interpolant of ω .

Lemma 3.1 (Superapproximation). *Let ω be as above and $\chi \in S_h^r$ with $r \geq 2$. Then there exists a constant C independent of ω , d and h such that for $s = 1, 2, 3$*

$$|\omega^4\chi - I_h(\omega^4\chi)|_{H^s(T)} \leq C \frac{h_T^{3-s}}{d} (|\omega^2\chi|_{H^2(T)} + d^{-1}|\omega\chi|_{H^1(T)} + d^{-2}\|\chi\|_{L^2(T)}).$$

Proof. We give a detailed proof for piecewise quadratic elements $r = 2$ only. The proof for general $r \geq 2$ is very similar. By the standard approximation theory we have

$$(22) \quad |\omega^4\chi - I_h(\omega^4\chi)|_{H^s(T)} \leq Ch_T^{3-s} |\omega^4\chi|_{H^3(T)}, \quad \text{for } s = 1, 2, 3.$$

Then

$$(23) \quad D^3(\omega^4\chi) = D^3(\omega^4)\chi + 3D^2(\omega^4)D\chi + 3D(\omega^4)D^2\chi + \omega^4 D^3\chi := I_1 + I_2 + I_3 + I_4.$$

Since we are dealing with quadratic elements, $I_4 = 0$. Using (19c), I_1 can easily be estimated as

$$(24) \quad I_1 = \|D^3(\omega^4)\chi\|_{L^2(T)} \leq Cd^{-3}\|\chi\|_{L^2(T)}.$$

Using

$$(25) \quad D(\omega^4) = 4\omega^3 D\omega \quad \text{and} \quad D^2(\omega^4) = 4D(\omega^3 D\omega) = 4(3\omega^2 |D\omega|^2 + \omega^3 D^2\omega),$$

we have

$$(26) \quad I_2 = \|3D(\omega^4)D^2\chi\|_{L^2(T)} \leq \|12\omega^2 |D\omega|^2 D\chi\|_{L^2(T)} + \|\omega^3 D^2\omega D\chi\|_{L^2(T)} = I_{21} + I_{22}.$$

Using (19c) and adding and subtracting $\widehat{\omega}$, where $\widehat{\omega} = |T|^{-1} \int_T \omega$, we have

$$(27) \quad \begin{aligned} I_{21} &\leq Cd^{-2}\|\omega D\chi\|_{L^2(T)} \\ &\leq Cd^{-2}\|(\omega - \widehat{\omega})D\chi\|_{L^2(T)} + Cd^{-2}\|\widehat{\omega}D\chi\|_{L^2(T)} \\ &= I_{211} + I_{212}. \end{aligned}$$

Using the approximation and the inverse estimate

$$I_{211} \leq Cd^{-2}h_T d^{-1}\|D\chi\|_{L^2(T)} \leq Cd^{-3}\|\chi\|_{L^2(T)}.$$

By using that $\widehat{\omega}$ is constant on T and using the triangle inequality, properties of ω and the inverse estimates, we have

$$\begin{aligned}
I_{212} &= Cd^{-2} \|D(\widehat{\omega}\chi)\|_{L^2(T)} \\
&\leq Cd^{-2} (\|D((\widehat{\omega} - \omega)\chi)\|_{L^2(T)} + \|D(\omega\chi)\|_{L^2(T)}) \\
&\leq Cd^{-2} (\|(D\omega)\chi\|_{L^2(T)} + \|(\widehat{\omega} - \omega)D\chi\|_{L^2(T)} + |\omega\chi|_{H^1(T)}) \\
&\leq Cd^{-3} \|\chi\|_{L^2(T)} + Cd^{-2} h_T d^{-1} \|D\chi\|_{L^2(T)} + Cd^{-2} |\omega\chi|_{H^1(T)} \\
&\leq Cd^{-3} \|\chi\|_{L^2(T)} + Cd^{-2} |\omega\chi|_{H^1(T)}.
\end{aligned}$$

Similarly to the analysis above, by the properties of ω

$$\begin{aligned}
I_{22} &\leq Cd^{-2} \|\omega^3 D\chi\|_{L^2(T)} \leq Cd^{-2} \|\omega D\chi\|_{L^2(T)} \\
&\leq Cd^{-2} \|(\widehat{\omega} - \omega)D\chi\|_{L^2(T)} + Cd^{-2} \|D(\widehat{\omega}\chi)\|_{L^2(T)} \\
&\leq Cd^{-3} \|\chi\|_{L^2(T)} + Cd^{-2} |\omega\chi|_{H^1(T)}.
\end{aligned}$$

The next term,

$$(28) \quad I_3 = \|3D(\omega^4)D^2\chi\|_{L^2(T)} = \|12\omega^3 D\omega D^2\chi\|_{L^2(T)}.$$

Thus adding and subtracting $P_1(\omega^2)$, we have

$$\begin{aligned}
I_3 &\leq Cd^{-1} \|\omega^2 D^2\chi\|_{L^2(T)} \\
&\leq Cd^{-1} (\|(\omega^2 - P_1(\omega^2))D^2\chi\|_{L^2(T)} + \|P_1(\omega^2)D^2\chi\|_{L^2(T)}) \\
&= I_{31} + I_{32}.
\end{aligned}$$

By the approximation and the inverse estimates

$$I_{31} \leq Cd^{-1} h_T^2 \|D^2(\omega^2)\|_{L^\infty(T)} \|D^2\chi\|_{L^2(T)} \leq Cd^{-3} \|\chi\|_{L^2(T)}.$$

Since $D^2(P_1(\omega^2)) = 0$, by the triangle inequality,

$$I_{32} \leq Cd^{-1} (\|D^2(P_1(\omega^2)\chi)\|_{L^2(T)} + \|D(P_1(\omega^2))D\chi\|_{L^2(T)}) = I_{321} + I_{322}.$$

Using $D^2(P_1(\omega^2)) = 0$, the approximation and the inverse inequality,

$$\begin{aligned}
I_{321} &\leq Cd^{-1} (\|D^2((P_1(\omega^2) - \omega^2)\chi)\|_{L^2(T)} + \|D^2(\omega^2\chi)\|_{L^2(T)}) \\
&\leq Cd^{-1} (\|D^2(\omega^2)\chi\|_{L^2(T)} + \|D(P_1(\omega^2) - \omega^2)D\chi\|_{L^2(T)}) \\
&\quad + Cd^{-1} (\|(P_1(\omega^2) - \omega^2)D^2\chi\|_{L^2(T)} + |\omega^2\chi|_{H^2(T)}) \\
&\leq Cd^{-1} (d^{-2} \|\chi\|_{L^2(T)} + h_T d^{-2} \|D\chi\|_{L^2(T)} + h_T^2 d^{-2} \|D^2\chi\|_{L^2(T)} + |\omega^2\chi|_{H^2(T)}) \\
&\leq Cd^{-3} \|\chi\|_{L^2(T)} + Cd^{-1} |\omega^2\chi|_{H^2(T)}.
\end{aligned}$$

Adding and subtracting ω^2 , and using the estimate for I_{21} (27), we have

$$\begin{aligned}
I_{322} &\leq Cd^{-1} (\|D(P_1(\omega^2) - \omega^2)D\chi\|_{L^2(T)} + \|D(\omega^2)D\chi\|_{L^2(T)}) \\
&\leq Cd^{-1} (h_T d^{-2} \|D\chi\|_{L^2(T)} + 2 \|\omega D\omega D\chi\|_{L^2(T)}) \\
&\leq Cd^{-3} \|\chi\|_{L^2(T)} + Cd^{-2} \|\omega D\chi\|_{L^2(T)} \\
&\leq Cd^{-3} \|\chi\|_{L^2(T)} + Cd^{-2} |\omega\chi|_{H^1(T)}.
\end{aligned}$$

Combining all the estimates we complete the proof. \square

Remark 3.2. The straightforward approach [15] gives the following superapproximation result

$$|\omega^4 \chi - I_h(\omega^4 \chi)|_{H^s(T)} \leq C \frac{h_T^{3-s}}{d} (|\chi|_{H^2(T)} + d^{-1} |\chi|_{H^1(T)} + d^{-2} \|\chi\|_{L^2(T)}), \quad s=1, 2, 3.$$

Although, such a result would be sufficient for our goals in this paper, it would make the analysis of local energy error estimates in Lemma 3.9 more cumbersome.

Corollary 3.3. *Let ω be as above and $\chi \in S_h^r$ with $r \geq 2$. Then there exists a constant C independent of ω , d and h such that*

(29a)

$$\| |\omega^4 \chi - I_h(\omega^4 \chi)| \|_{2,h} \leq C \frac{h_T}{d} (|\omega^2 \chi|_{H^2(\Omega)} + d^{-1} |\omega \chi|_{H^1(\Omega)} + d^{-2} \|\chi\|_{L^2(B_d)})$$

(29b)

$$\| |\omega^4 \chi - I_h(\omega^4 \chi)| \|_{2,h} \leq C \frac{h_T}{d} (\|\omega^2 D^2 \chi\|_{L^2(\Omega)} + d^{-1} \|\omega \nabla \chi\|_{L^2(\Omega)} + d^{-2} \|\chi\|_{L^2(B_d)}).$$

Proof. The proof follows from Lemma 3.1, the trace and inverse inequalities and the product rule. \square

3.4. Preliminary weighted results. First we show the following supplementary result.

Lemma 3.4. *Let ω be as at the beginning of section 3.3 with the properties (19a)-(19c). Then for any $\varepsilon > 0$ and any $v_h \in S_h^r$*

$$\begin{aligned} \|\omega \nabla v_h\|_{L^2(\Omega)}^2 &\leq \varepsilon d^2 \left(\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 \right) \\ &\quad + C_\varepsilon d^{-2} \|v_h\|_{L^2(B_d)}^2. \end{aligned}$$

Proof. Integrating by parts on an element T we have

$$\begin{aligned} \|\omega \nabla v_h\|_{L^2(T)}^2 &= (\omega^2 \nabla v_h, \nabla v_h)_T = -(\nabla \cdot (\omega^2 \nabla v_h), v_h)_T + \left(\omega^2 \frac{\partial v_h}{\partial n}, v_h \right)_{\partial T} \\ &= -(\nabla(\omega^2) \cdot \nabla v_h, v_h)_T - (\omega^2 \Delta v_h, v_h)_T + \left(\omega^2 \frac{\partial v_h}{\partial n}, v_h \right)_{\partial T}. \end{aligned}$$

Summing over elements we obtain

$$\begin{aligned} \|\omega \nabla v_h\|_{L^2(\Omega)}^2 &= \sum_{T \in \mathcal{T}_h} -(\nabla(\omega^2) \cdot \nabla v_h, v_h)_T - (\omega^2 \Delta v_h, v_h)_T + \left(\omega^2 \frac{\partial v_h}{\partial n}, v_h \right)_{\partial T} \\ &:= J_1 + J_2 + J_3. \end{aligned}$$

Using the Cauchy-Schwarz and Young's inequalities, and using the definition of $\mathcal{T}_h \cap B_{3d/4}$ in (16), we have

$$\begin{aligned}
J_1 &= - \sum_{T \in \mathcal{T}_h} 2(\omega \nabla \omega \cdot \nabla v_h, v_h)_T \\
&\leq \sum_{T \in \mathcal{T}_h \cap B_{3d/4}} C d^{-1} \|\omega \nabla v_h\|_{L^2(T)} \|v_h\|_{L^2(T)} \\
&\leq \frac{1}{4} \sum_{T \in \mathcal{T}_h} \|\omega \nabla v_h\|_{L^2(T)}^2 + C d^{-2} \sum_{T \in \mathcal{T}_h \cap B_{3d/4}} \|v_h\|_{L^2(T)}^2 \\
&\leq \frac{1}{4} \|\omega \nabla v_h\|_{L^2(\Omega)}^2 + C d^{-2} \|v_h\|_{L^2(B_d)}^2.
\end{aligned}$$

Similarly using the Cauchy-Schwarz and Young's inequalities

$$\begin{aligned}
J_2 &= - \sum_{T \in \mathcal{T}_h} (\omega^2 \Delta v_h, v_h)_T \\
&\leq \sum_{T \in \mathcal{T}_h \cap B_{3d/4}} \|\omega^2 \Delta v_h\|_{L^2(T)} \|v_h\|_{L^2(T)} \\
&\leq \frac{\varepsilon d^2}{2} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + C_\varepsilon d^{-2} \sum_{T \in \mathcal{T}_h \cap B_{3d/4}} \|v_h\|_{L^2(T)}^2 \\
&\leq \frac{\varepsilon d^2}{2} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + C_\varepsilon d^{-2} \|v_h\|_{L^2(B_d)}^2.
\end{aligned}$$

Finally, summing over the elements and using the trace and inverse inequalities and using the definition of $\mathcal{T}_h \cap B_{3d/4}$ in (16), we obtain

$$\begin{aligned}
J_3 &= \sum_{T \in \mathcal{T}_h} \left(\omega^2 \frac{\partial v_h}{\partial n}, v_h \right)_{\partial T} = \sum_{e \in \mathcal{E}_h} \left(\omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket, v_h \right)_e \\
&\leq \frac{\varepsilon d^2}{2} \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + C_\varepsilon d^{-2} \sum_{e \in \mathcal{E}_h \cap B_{3d/4}} |e| \|v_h\|_{L^2(e)}^2 \\
&\leq \frac{\varepsilon d^2}{2} \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + C_\varepsilon C_{inv} C_{tr} d^{-2} \sum_{T \in \mathcal{T}_h \cap B_{3d/4}} \|v_h\|_{L^2(T)}^2 \\
&\leq \frac{\varepsilon d^2}{2} \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + C_\varepsilon d^{-2} \|v_h\|_{L^2(B_d)}^2.
\end{aligned}$$

Combining the above estimates we obtain the lemma. \square

We also need the following results.

Lemma 3.5. *Let ω be as above. There exists a constant C independent of d and h such that for any $v_h \in S_h^r$ and any $\varepsilon > 0$*

$$\begin{aligned}
&\sum_{e \in \mathcal{E}_h} |e| \left\| \omega \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 \\
&\leq C_\varepsilon d^2 \left(\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 \right) + C_\varepsilon d^{-2} \|v_h\|_{L^2(B_d)}^2.
\end{aligned}$$

Proof. Adding and subtracting $P_0(\omega)$ we have

$$\left\| \omega \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 \leq \left\| P_0(\omega) \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 + \left\| (\omega - P_0(\omega)) \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2.$$

By the trace and inverse inequalities

$$\begin{aligned} \sum_{e \in \mathcal{E}_h} |e| \left\| P_0(\omega) \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 &\leq C \sum_{T \in \mathcal{T}_h} \|P_0(\omega) \nabla v_h\|_{L^2(T)}^2 \\ &\leq C \sum_{T \in \mathcal{T}_h} \|\omega \nabla v_h\|_{L^2(T)}^2 + \|(P_0(\omega) - \omega) \nabla v_h\|_{L^2(T)}^2 \\ &= C \left(\|\omega \nabla v_h\|_{L^2(\Omega)}^2 + \|(P_0(\omega) - \omega) \nabla v_h\|_{L^2(\Omega)}^2 \right). \end{aligned}$$

By the approximation and the inverse inequality, we have

$$\begin{aligned} \|(P_0(\omega) - \omega) \nabla v_h\|_{L^2(\Omega)} &\leq \|P_0(\omega) - \omega\|_{L^\infty(B_d)} \|\nabla v_h\|_{L^2(B_d)} \\ &\leq Ch \|\nabla \omega\|_{L^\infty(B_d)} \|\nabla v_h\|_{L^2(B_d)} \leq Cd^{-1} \|v_h\|_{L^2(B_d)}. \end{aligned}$$

Combining the above estimate with Lemma 3.4, we obtain

$$\begin{aligned} \sum_{e \in \mathcal{E}_h} |e| \left\| P_0(\omega) \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 &\leq \\ C\varepsilon d^2 \left(\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \left[\left[\frac{\partial v_h}{\partial n} \right] \right] \right\|_{L^2(e)}^2 \right) \\ + C\varepsilon d^{-2} \|v_h\|_{L^2(B_d)}^2. \end{aligned}$$

Similarly, by the approximation, properties of ω , trace and inverse inequalities,

$$\sum_{e \in \mathcal{E}_h} |e| \left\| (P_0(\omega) - \omega) \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 \leq Cd^{-2} \|v_h\|_{L^2(B_d)}^2.$$

Combining, we obtain the lemma. \square

Lemma 3.6. *Let ω be as above. There exists a constant C independent of d and h such that for any $v_h \in S_h^r$*

$$\sum_{e \in \mathcal{E}_h} |e| \left\| \omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \leq C \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + Cd^{-4} \|v_h\|_{L^2(B_d)}^2.$$

Proof. Adding and subtracting $P_1(\omega^2)$ we have

$$\left\| \omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \leq \left\| P_1(\omega^2) \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 + \left\| (\omega^2 - P_1(\omega^2)) \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2.$$

By the trace and inverse inequalities

$$\begin{aligned} \sum_{e \in \mathcal{E}_h} |e| \left\| P_1(\omega^2) \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 &\leq \\ C \sum_{T \in \mathcal{T}_h} \|P_1(\omega^2) D^2 v_h\|_{L^2(T)}^2 &\leq \\ C \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \|(P_1(\omega^2) - \omega^2) D^2 v_h\|_{L^2(T)}^2. \end{aligned}$$

By the approximation and the inverse inequality, we have

$$\begin{aligned} \|(P_1(\omega^2) - \omega^2)D^2v_h\|_{L^2(T)} &\leq \|P_1(\omega^2) - \omega^2\|_{L^\infty(T)} \|D^2v_h\|_{L^2(T)} \\ &\leq Ch^2 \|D^2(\omega^2)\|_{L^\infty(T)} \|D^2v_h\|_{L^2(T)} \leq Cd^{-2} \|v_h\|_{L^2(T)} \end{aligned}$$

and as a result

$$\sum_{e \in \mathcal{E}_h} |e| \left\| P_1(\omega^2) \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \leq C \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + Cd^{-4} \|v_h\|_{L^2(B_d)}^2.$$

Similarly, by the approximation, properties of ω , trace and inverse inequalities,

$$\sum_{e \in \mathcal{E}_h} |e| \left\| (P_1(\omega^2) - \omega^2) \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \leq C \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + Cd^{-4} \|v_h\|_{L^2(B_d)}^2.$$

□

Next we show the following coersivity type result.

Lemma 3.7. *Let ω be defined as above. Then, for any $v_h \in S_h^r$*

$$\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \left[\frac{\partial v_h}{\partial n} \right] \right\|_{L^2(e)}^2 \leq CB_h(v_h, \omega^4 v_h) + Cd^{-4} \|v_h\|_{L^2(B_d)}^2.$$

Proof. From the definition of the bilinear form

$$\begin{aligned} B_h(v_h, \omega^4 v_h) &= \sum_{T \in \mathcal{T}_h} \int_T D^2 v_h : D^2(\omega^4 v_h) dx + \\ &\quad + \sum_{e \in \mathcal{E}_h} \int_e \left(\left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \left[\frac{\partial(\omega^4 v_h)}{\partial n} \right] + \left[\frac{\partial v_h}{\partial n} \right] \left\{ \left\{ \frac{\partial^2(\omega^4 v_h)}{\partial n^2} \right\} \right\} \right) ds \\ &\quad + \sum_{e \in \mathcal{E}_h} \frac{\eta}{|e|} \int_e \left[\frac{\partial v_h}{\partial n} \right] \left[\frac{\partial(\omega^4 v_h)}{\partial n} \right] ds \\ &:= J_1 + J_2 + J_3 + J_4. \end{aligned}$$

First we will address

$$J_1 = \sum_{T \in \mathcal{T}_h} \int_T D^2 v_h : D^2(\omega^4 v_h) dx := \sum_{T \in \mathcal{T}_h} \sum_{i,j=1}^2 (\partial_{ij}^2 v_h, \partial_{ij}^2(\omega^4 v_h))_T.$$

Since

$$\begin{aligned} \partial_{ij}^2(\omega^4 v_h) &= \partial_i(4\omega^3 \partial_j \omega v_h + \omega^4 \partial_j v_h) \\ &= 12\omega^2 \partial_i \omega \partial_j \omega v_h + 4\omega^3 \partial_{ij}^2 \omega v_h + 4\omega^3 \partial_j \omega \partial_i v_h + 4\omega^3 \partial_i \omega \partial_j v_h + \omega^4 \partial_{ij}^2 v_h, \end{aligned}$$

we have

$$J_1 = \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + J_{12} + J_{13},$$

where

$$J_{12} = \sum_{T \in \mathcal{T}_h} \sum_{i,j=1}^2 (\omega^2 \partial_{ij}^2 v_h, (12\partial_i \omega \partial_j \omega + 4\omega \partial_{ij}^2 \omega) v_h)_T$$

and

$$J_{13} = \sum_{T \in \mathcal{T}_h} \sum_{i,j=1}^2 (\omega^2 \partial_{ij}^2 v_h, 4\omega \partial_j \omega \partial_i v_h + 4\omega \partial_i \omega \partial_j v_h)_T.$$

Since $v_h \in C^0(\Omega)$ and $\omega \in C^\infty(\Omega)$ we have

$$\llbracket \frac{\partial(\omega^4 v_h)}{\partial n} \rrbracket = \omega^4 \llbracket \frac{\partial v_h}{\partial n} \rrbracket$$

and

$$\left\{ \left\{ \frac{\partial^2(\omega^4 v_h)}{\partial n^2} \right\} \right\} = \omega^4 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} + 2 \frac{\partial(\omega^4)}{\partial n} \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} + \frac{\partial^2(\omega^4)}{\partial n^2} v_h.$$

Thus,

$$\begin{aligned} J_2 &= \sum_{e \in \mathcal{E}_h} \left(\omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\}, \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e, \\ J_4 &= \sum_{e \in \mathcal{E}_h} \frac{\eta}{|e|} \int_e \llbracket \frac{\partial v_h}{\partial n} \rrbracket \llbracket \frac{\partial(\omega^4 v_h)}{\partial n} \rrbracket ds = \sum_{e \in \mathcal{E}_h} \frac{\eta}{|e|} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 \end{aligned}$$

and

$$\begin{aligned} J_3 &= \sum_{e \in \mathcal{E}_h} \left(\omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\}, \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e \\ &\quad + 2 \left(\frac{\partial \omega^4}{\partial n} \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\}, \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e + \left(\frac{\partial^2 \omega^4}{\partial n^2} v_h, \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e \\ &= \sum_{e \in \mathcal{E}_h} \left(\omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\}, \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e + J_{31} + J_{32}. \end{aligned}$$

Combining the above estimates we obtain

$$\begin{aligned} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 &= B(v_h, \omega^4 v_h) \\ &\quad - J_{12} - J_{13} - 2 \sum_{e \in \mathcal{E}_h} \left(\omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\}, \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e - J_{31} - J_{32} \\ &\quad - (\eta - 1) \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2. \end{aligned}$$

Using the Cauchy-Schwarz and Young's inequalities

$$\begin{aligned} (\omega^2 \partial_{ij}^2 v_h, (12\partial_i \omega \partial_j \omega + 4\omega \partial_{ij}^2 \omega) v_h)_T &\leq C d^{-2} \|\omega^2 D^2 v_h\|_{L^2(T)} \|v_h\|_{L^2(T \cap B_{3d/4})} \\ &\leq \frac{1}{8} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + C d^{-4} \|v_h\|_{L^2(T \cap B_{3d/4})}^2 \end{aligned}$$

and summing up we have

$$-J_{12} \leq \frac{1}{8} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + C d^{-4} \|v_h\|_{L^2(B_d)}^2.$$

Again using the Cauchy-Schwarz and Young's inequalities

$$\begin{aligned} 4(\omega^2 \partial_{ij}^2 v_h, \omega \partial_j \omega \partial_i v_h + \omega \partial_i \omega \partial_j v_h)_T &\leq C d^{-1} \|\omega^2 D^2 v_h\|_{L^2(T)} \|\omega \nabla v_h\|_{L^2(T)} \\ &\leq \frac{1}{8} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + C d^{-2} \|\omega \nabla v_h\|_{L^2(T)}^2. \end{aligned}$$

Summing and using Lemma 3.4, we have

$$\begin{aligned} -J_{13} &\leq \frac{1}{8} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + Cd^{-2} \sum_{T \in \mathcal{T}_h} \|\omega \nabla v_h\|_{L^2(T)}^2 \\ &\leq \left(\frac{1}{8} + \varepsilon C \right) \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \varepsilon C \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 \\ &\quad + C_\varepsilon d^{-4} \|v_h\|_{L^2(B_d)}^2. \end{aligned}$$

Using the Cauchy-Schwarz and Young's inequalities and Lemma 3.6

$$\begin{aligned} 2 \sum_{e \in \mathcal{E}_h} \left(\omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\}, \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right)_e &\leq \frac{1}{8} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + Cd^{-4} \|v_h\|_{L^2(B_d)}^2 \\ &\quad + C \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2. \end{aligned}$$

Using properties of ω and Lemma 3.5,

$$\begin{aligned} -J_{31} &\leq Cd^{-1} \sum_{e \in \mathcal{E}_h} \left\| \omega \left\{ \left\{ \frac{\partial v_h}{\partial n} \right\} \right\} \right\|_e \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_e \\ &\leq C\varepsilon \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + C \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + C_\varepsilon d^{-4} \|v_h\|_{L^2(B_d)}^2. \end{aligned}$$

Similarly, using properties of ω and the trace inequality,

$$\begin{aligned} -J_{32} &\leq Cd^{-2} \sum_{e \in \mathcal{E}_h \cap B_{3d/4}} \|v_h\|_e \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_e \\ &\leq C \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + Cd^{-4} \|v_h\|_{L^2(B_d)}^2. \end{aligned}$$

Choosing ε such that $C\varepsilon \leq \frac{1}{8}$ and providing η is sufficiently large we obtain the lemma. \square

In view of Lemma 3.6, we also have

Corollary 3.8. *Under the assumption of Lemma 3.7, we have*

$$\begin{aligned} \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 v_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial v_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + \sum_{e \in \mathcal{E}_h} |e| \left\| \omega^2 \left\{ \left\{ \frac{\partial^2 v_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \\ \leq CB(v_h, \omega^4 v_h) + Cd^{-4} \|v_h\|_{L^2(B_d)}^2. \end{aligned}$$

3.5. Local energy estimates. Next we establish the following key result.

Lemma 3.9 (Local energy estimate). *Let u and u_h satisfy $B_h(u - u_h, \chi) = 0$ for any $\chi \in S_h^r$. Given $D \subset \Omega$, $d \geq \kappa h > 0$ for sufficiently large κ , and $D \subset D_d$ with $\text{dist}(D_d, \partial D \setminus \partial \Omega) \geq d$, there exists a constant C independent of d and h such that*

$$\|u - u_h\|_{2,h,D} \leq C \inf_{\chi \in S_h^r} \left(\|u - \chi\|_{2,h,D_d} + d^{-2} \|u - \chi\|_{L^2(D_d)} \right) + Cd^{-2} \|u - u_h\|_{L^2(D_d)}.$$

Proof. To show the lemma, it is sufficient to establish the following estimate

$$(30) \quad \|\|u_h\|\|_{2,h,B} \leq C\|u\|_{2,h,B_d} + Cd^{-2}\|u_h\|_{L^2(B_d)}$$

for a ball $B := B_{d/2}$. The lemma then would follow by standard covering argument and by replacing u with $u - \chi$ and $u_h - \chi$ and the triangle inequality.

Let ω be the cut-off function as in (19). Then by Corollary 3.8,

$$\begin{aligned} \|\|u_h\|\|_{2,h,B}^2 &\leq \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 u_h\|_{L^2(T)}^2 \\ &\quad + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial u_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + \sum_{e \in \mathcal{E}_h} |e| \left\| \omega^2 \left\{ \left\{ \frac{\partial^2 u_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \\ &\leq CB_h(u_h, \omega^4 u_h) + Cd^{-4}\|u_h\|_{L^2(B_d)}^2. \end{aligned}$$

We use the identity

$$\begin{aligned} B_h(u_h, \omega^4 u_h) &= B_h(u_h, \omega^4 u_h - I_h(\omega^4 u_h)) + B_h(u, \omega^4 u_h) - B_h(u, \omega^4 u_h - I_h(\omega^4 u_h)) \\ &:= J_1 + J_2 + J_3. \end{aligned}$$

Now we treat all three terms separately. Using the boundedness of the bilinear from and Young's inequality

$$J_2 \leq C\|u\|_{2,h,B_d} \|\omega^4 u_h\|_{2,h}.$$

By the definition of the $\|\cdot\|_{2,h}$ norm,

$$\begin{aligned} \|\omega^4 u_h\|_{2,h}^2 &= \sum_{T \in \mathcal{T}_h} \|D^2(\omega^4 u_h)\|_{L^2(T)}^2 \\ &\quad + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \llbracket \frac{\partial(\omega^4 u_h)}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + \sum_{e \in \mathcal{E}_h} |e| \left\| \left\{ \left\{ \frac{\partial^2(\omega^4 u_h)}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 \\ &:= I_{21} + I_{22} + I_{23}. \end{aligned}$$

Using the product rule, the properties of ω and Lemma 3.4, we have

$$\begin{aligned} I_{21} &\leq \sum_{T \in \mathcal{T}_h} \left(\|\omega^4 D^2 u_h\|_{L^2(T)}^2 + 4\|D(\omega^4) D u_h\|_{L^2(T)}^2 + \|D^2(\omega^4) u_h\|_{L^2(T)}^2 \right) \\ &\leq C \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 u_h\|_{L^2(T)}^2 + C \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial u_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 + Cd^{-4}\|u_h\|_{L^2(B_d)}^2. \end{aligned}$$

Since

$$\llbracket \frac{\partial(\omega^4 u_h)}{\partial n} \rrbracket = \omega^4 \llbracket \frac{\partial u_h}{\partial n} \rrbracket,$$

we have

$$I_{22} = \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^4 \llbracket \frac{\partial u_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2 \leq \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \llbracket \frac{\partial u_h}{\partial n} \rrbracket \right\|_{L^2(e)}^2.$$

Using

$$\left\{ \left\{ \frac{\partial^2(\omega^4 u_h)}{\partial n^2} \right\} \right\} = \omega^4 \left\{ \left\{ \frac{\partial^2 u_h}{\partial n^2} \right\} \right\} + 2 \frac{\partial(\omega^4)}{\partial n} \left\{ \left\{ \frac{\partial u_h}{\partial n} \right\} \right\} + \frac{\partial^2(\omega^4)}{\partial n^2} u_h,$$

Lemma 3.6 and Lemma 3.5, the properties of ω , and inverse inequality, we also obtain

$$\begin{aligned} I_{23} &\leq \sum_{e \in \mathcal{E}_h} |e| \left(\left\| \omega^2 \left\{ \left\{ \frac{\partial^2 u_h}{\partial n^2} \right\} \right\} \right\|_{L^2(e)}^2 + Cd^{-2} \left\| \omega \left\{ \left\{ \frac{\partial u_h}{\partial n} \right\} \right\} \right\|_{L^2(e)}^2 + Cd^{-4} \|u_h\|_{L^2(B_d)}^2 \right) \\ &\leq C \sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 u_h\|_{L^2(T)}^2 + C \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \left[\frac{\partial u_h}{\partial n} \right] \right\|_{L^2(e)}^2 + Cd^{-4} \|u_h\|_{L^2(B_d)}^2. \end{aligned}$$

Hence, by the Young's inequality

$$\begin{aligned} J_2 &\leq \frac{1}{4} \left(\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 u_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \left[\frac{\partial u_h}{\partial n} \right] \right\|_{L^2(e)}^2 \right) \\ &\quad + Cd^{-4} \|u_h\|_{L^2(B_d)}^2 + C \|u\|_{2,h,B_d}^2. \end{aligned}$$

Using the superapproximation result Corollary 3.3 and Lemma 3.4 and similarly to the above

$$\begin{aligned} J_3 &\leq C \|u\|_{2,h,B_d} \| \omega^4 u_h - I_h(\omega^4 u_h) \|_{2,h} \\ &\leq C \frac{h}{d} \|u\|_{2,h,B_d} (\|\omega^2 D^2 u_h\|_{L^2(\Omega)} + d^{-1} \|\omega \nabla u_h\|_{L^2(\Omega)} + d^{-2} \|u_h\|_{L^2(B_d)}) \\ &\leq \frac{1}{4} \left(\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 u_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \left[\frac{\partial u_h}{\partial n} \right] \right\|_{L^2(e)}^2 \right) \\ &\quad + C \|u\|_{2,h,B_d}^2 + Cd^{-4} \|u_h\|_{L^2(B_d)}^2. \end{aligned}$$

To estimate J_1 we use again the superapproximation result Corollary 3.3 and Lemma 3.4 to obtain

$$\begin{aligned} J_1 &\leq C \|u_h\|_{2,h,B_d} \| \omega^4 u_h - I_h(\omega^4 u_h) \|_{2,h} \\ &\leq C \frac{h}{d} \|u_h\|_{2,h,B_d} (\|\omega^2 D^2 u_h\|_{L^2(\Omega)} + d^{-1} \|\omega \nabla u_h\|_{L^2(\Omega)} + d^{-2} \|u_h\|_{L^2(B_d)}) \\ &\leq \frac{1}{4} \left(\sum_{T \in \mathcal{T}_h} \|\omega^2 D^2 u_h\|_{L^2(T)}^2 + \sum_{e \in \mathcal{E}_h} |e|^{-1} \left\| \omega^2 \left[\frac{\partial u_h}{\partial n} \right] \right\|_{L^2(e)}^2 \right) \\ &\quad + C \frac{h^2}{d^2} \|u_h\|_{2,h,B_d}^2 + Cd^{-4} \|u_h\|_{L^2(B_d)}^2. \end{aligned}$$

Combining the estimates and kicking back the terms, we obtain

$$\|u_h\|_{2,h,B}^2 \leq C \|u\|_{2,h,B_d}^2 + C \frac{h^2}{d^2} \|u_h\|_{2,h,B_d}^2 + Cd^{-4} \|u_h\|_{L^2(B_d)}^2.$$

Iterating the argument once again and using the inverse inequality we obtain

$$\begin{aligned} \|u_h\|_{2,h,B}^2 &\leq C \|u\|_{2,h,B_{2d}}^2 + C \frac{h^4}{d^4} \|u_h\|_{2,h,B_{2d}}^2 + Cd^{-4} \|u_h\|_{L^2(B_{2d})}^2 \\ &\leq C \|u\|_{2,h,B_{2d}}^2 + Cd^{-4} \|u_h\|_{L^2(B_{2d})}^2, \end{aligned}$$

which establishes (30) with insignificant difference of having subset B_{2d} instead of B_d on the right hand side. \square

4. POINTWISE ERROR ESTIMATES

Let $x_0 \in \overline{\Omega}$ be a fixed (but arbitrary) point. Let $T_0 \in \mathcal{T}_h$ be an element such that $x_0 \in \overline{T}_0$. Associated to this point we introduce a smooth Delta function [21, Lemma 2.2], which we will denote by $\tilde{\delta} = \tilde{\delta}_{x_0}$, cf. also [19]. This function is supported in T_0 and satisfies

$$(\chi, \tilde{\delta})_{T_0} = \chi(x_0), \quad \forall \chi \in \mathbb{P}^r(T_0).$$

In addition from [21, Lemma 2.2] we also have

$$(31) \quad \|\tilde{\delta}\|_{W^{s,p}(\Omega)} \leq Ch^{-s-2(1-\frac{1}{p})}, \quad 1 \leq p \leq \infty, \quad s = 0, 1, 2.$$

In particular $\|\tilde{\delta}\|_{L^1(\Omega)} \leq C$, $\|\tilde{\delta}\|_{L^2(\Omega)} \leq Ch^{-1}$, and $\|\tilde{\delta}\|_{L^\infty(\Omega)} \leq Ch^{-2}$.

Now we define a function $g \in H_0^2(\Omega)$, by the equation

$$(32) \quad B(g, \varphi) = (D^2\tilde{\delta}, \varphi), \quad \forall \varphi \in H_0^2(\Omega),$$

which also satisfy

$$B_h(g, \chi) = (D^2\tilde{\delta}, \chi) = D^2\chi(x_0), \quad \forall \chi \in S_h^r,$$

and correspondingly we define $g_h \in S_h^r$ by

$$(33) \quad B_h(g - g_h, \chi) = 0, \quad \forall \chi \in S_h^r.$$

Thus, for any $\chi \in S_h^r$ using (33), (14), (18) and that $\tilde{\delta}$ is supported on T_0 , we have

$$\begin{aligned} D^2(u_h - \chi)(x_0) &= B_h(u_h - \chi, g) \\ &= B_h(u_h - \chi, g_h) \\ &= B_h(u - \chi, g_h) \\ &= B_h(u - \chi, g) - B_h(u - \chi, g - g_h) \\ &= (u - \chi, D^2\tilde{\delta})_{T_0} - B_h(u - \chi, g - g_h) \\ &= (D^2(u - \chi), \tilde{\delta})_{T_0} - B_h(u - \chi, g - g_h) \\ &\leq \|D^2(u - \chi)\|_{L^\infty(T_0)} \|\tilde{\delta}\|_{L^1(T_0)} + C \|u - \chi\|_{W_h^{2,\infty}(\Omega)} \|g - g_h\|_{W_h^{2,1}(\Omega)} \\ &\leq C \|u - \chi\|_{W_h^{2,\infty}(\Omega)} (1 + \|g - g_h\|_{W_h^{2,1}(\Omega)}). \end{aligned}$$

The main result would follow once we establish the following result.

Lemma 4.1. *There exists a constant C independent of h such that*

$$\|g - g_h\|_{W_h^{2,1}(\Omega)} \leq C |\ln h|^{3/2}.$$

4.1. Proof of Lemma 4.1. In the proof we will use a dyadic decomposition of Ω . Let $j_0 \in \mathbb{Z}$ be the largest integer such that $d_{j_0} := 2^{-j_0} \geq \text{diam}(\Omega)$. We have

$$(34) \quad \Omega = \Omega_* \cup \bigcup_{j=j_0}^J \Omega_j,$$

with

$$\Omega_* = \{x \in \Omega : |x - x_0| \leq d_*,\} \quad d_* = C_* h$$

$$\Omega_j = \{x \in \Omega : d_{j+1} \leq |x - x_0| \leq d_j\}, \quad d_j = 2^{-j},$$

where C_* is a sufficiently large constant and J is the integer such that $\frac{C_*}{2}h \leq d_J \leq C_*h$. Note that $J \approx |\ln h|$. We will also use $\Omega'_j = \Omega_{j-1} \cup \Omega_j \cup \Omega_{j+1}$. Thus by the triangle inequality

$$\|g - g_h\|_{W_h^{2,1}(\Omega)} \leq \|g - g_h\|_{W_h^{2,1}(\Omega_*)} + \sum_{j=j_0}^J \|g - g_h\|_{W_h^{2,1}(\Omega_j)}.$$

By the Cauchy-Schwarz inequality, (15) and using the estimate (31), we obtain (35)

$$\|g - g_h\|_{W_h^{2,1}(\Omega_*)} \leq C_*h\|g - g_h\|_{2,h} \leq Ch^2\|D^2\tilde{\delta}\|_{H^{-1}(\Omega)} \leq Ch^2\|\nabla\tilde{\delta}\|_{L^2(\Omega)} \leq C.$$

Similarly by the Cauchy-Schwarz inequality

$$\sum_{j=j_0}^J \|g - g_h\|_{W_h^{2,1}(\Omega_j)} \leq C \sum_{j=j_0}^J d_j \|g - g_h\|_{2,h,\Omega_j}.$$

Thus we have

$$(36) \quad \|g - g_h\|_{W_h^{2,1}(\Omega)} \leq C + C \sum_{j=j_0}^J d_j \|g - g_h\|_{2,h,\Omega_j}.$$

By the local energy estimate in Lemma 3.9

$$\|g - g_h\|_{2,h,\Omega_j} \leq C \left(\|g - \chi\|_{2,h,\Omega'_j} + d_j^{-2} \|g - \chi\|_{L^2(\Omega'_j)} \right) + Cd_j^{-2} \|g - g_h\|_{L^2(\Omega'_j)}.$$

Taking $\chi = I_h g$, where I_h is the usual nodal interpolant with properties (9), we obtain

$$\begin{aligned} \|g - I_h g\|_{2,h,\Omega'_j}^2 &\leq \sum_{T \in \mathcal{T}_h \cap \Omega'_j} \|D^2(g - I_h g)\|_{L^2(T)}^2 \\ &\quad + \sum_{e \in \mathcal{E}_h \cap \Omega'_j} \int_e \left(|e| \left| \left\{ \frac{\partial^2(g - I_h g)}{\partial n^2} \right\} \right|^2 + |e|^{-1} \left| \left[\frac{\partial(g - I_h g)}{\partial n} \right] \right|^2 \right) ds \\ &= E_1 + E_2 + E_3. \end{aligned}$$

Using the Hölder inequality and (9), we have

$$\begin{aligned} E_1 &\leq \sum_{T \in \mathcal{T}_h \cap \Omega'_j} Ch^2 \|D^2(g - I_h g)\|_{L^\infty(T)}^2 \\ &\leq Ch^2 \|D^2 g\|_{L^\infty(\Omega'_j)}^2 \sum_{T \in \mathcal{T}_h \cap \Omega'_j} 1 \leq Cd_j^2 \|D^2 g\|_{L^\infty(\Omega'_j)}^2, \end{aligned}$$

where in the last step we used that for the quasi-uniform triangulation, the number of triangle in Ω'_j is of order $(d_j/h)^2$. Similarly, using the Hölder inequality and stability of the interpolant, we have

$$\begin{aligned} E_2 &\leq C \sum_{e \in \mathcal{E}_h \cap \Omega_j} \int_e |e|^2 \|D^2(g - I_h g)\|_{L^\infty(e)}^2 \\ &\leq Ch^2 \|D^2 g\|_{L^\infty(\Omega'_j)}^2 \sum_{e \in \mathcal{E}_h \cap \Omega'_j} 1 \leq Cd_j^2 \|D^2 g\|_{L^\infty(\Omega'_j)}^2. \end{aligned}$$

To estimate E_3 we use the trace inequality together with the approximation property and the stability of the interpolant (9). Thus, we obtain

$$\begin{aligned} E_3 &\leq C \sum_{T \in \mathcal{E}_h \cap \Omega'_j} \left(h^{-2} \|D(g - I_h g)\|_{L^2(T)}^2 + \|D^2(g - I_h g)\|_{L^2(T)}^2 \right) \\ &\leq C \sum_{T \in \mathcal{E}_h \cap \Omega'_j} \left(\|D(g - I_h g)\|_{L^\infty(T)}^2 + h^2 \|D^2(g - I_h g)\|_{L^\infty(T)}^2 \right) \\ &\leq Ch^2 \|D^2 g\|_{L^\infty(T)}^2 \sum_{T \in \mathcal{E}_h \cap \Omega'_j} 1 \leq Cd_j^2 \|D^2 g\|_{L^\infty(\Omega'_j)}^2. \end{aligned}$$

Using the Green's function representation and the integration by part, we obtain

$$D^2 g(x) = \int_{T_0} D_x^2 G(x, y) D^2 \tilde{\delta}(y) dy = \int_{T_0} D_x^2 D_y^2 G(x, y) \tilde{\delta}(y) dy.$$

Using the pointwise Green's function estimate from Lemma 2.1 and that $\text{dist}(\Omega'_j, T_0) \approx d_j$, for any $x \in \Omega'_j$ we have

$$|D^2 g(x)| \leq \int_{T_0} |D_x^2 D_y^2 G(x, y)| |\tilde{\delta}(y)| dy \leq \int_{T_0} \frac{|\tilde{\delta}(y)|}{|x - y|^2} dy \leq Cd_j^{-2}$$

and as a result

$$\sum_{j=j_0}^J d_j \|g - I_h g\|_{2,h,\Omega'_j} \leq C \sum_{j=j_0}^J d_j^2 \|g\|_{W^{2,\infty}(\Omega'_j)} \leq C \sum_{j=j_0}^J 1 \leq CJ \leq C|\ln h|.$$

Similarly, by the approximation, the Hölder inequality, and the estimates for $|D^2 g(x)|$ above, we also have

$$\|g - I_h g\|_{L^2(\Omega'_j)} \leq Ch^2 \|D^2 g\|_{L^2(\Omega'_j)} \leq Ch^2 d_j \|D^2 g\|_{L^\infty(\Omega'_j)} \leq Ch^2 d_j^{-1},$$

and as a result

$$C \sum_{j=j_0}^J d_j^{-1} \|g - I_h g\|_{L^2(\Omega'_j)} \leq C \sum_{j=j_0}^J h^2 d_j^{-2} \leq C.$$

Thus so far we have established

$$(37) \quad \|g - g_h\|_{W_h^{2,1}(\Omega)} \leq C|\ln h| + \sum_{j=j_0}^J d_j^{-1} \|g - g_h\|_{L^2(\Omega'_j)}.$$

To complete the proof of Lemma 4.1, we need to establish

$$(38) \quad \sum_{j=j_0}^J d_j^{-1} \|g - g_h\|_{L^2(\Omega'_j)} \leq C|\ln h|^{3/2},$$

which we will accomplish in the next section via a duality argument.

4.2. Duality argument. To estimate $\sum_{j=j_0}^J d_j^{-1} \|g - g_h\|_{L^2(\Omega'_j)}$ we use a duality argument. However, in contrast to the second order elliptic problems, on convex polygonal domains the regularity shift goes from H^3 to H^{-1} and estimating the L^2 norm of the error is problematic. To avoid this difficulty, we first transform $\sum_{j=j_0}^J d_j^{-1} \|g - g_h\|_{L^2(\Omega'_j)}$ to a gradient in L^2 norm.

Lemma 4.2. *There exists a constant C independent of h such that*

$$\sum_{j=j_0}^J d_j^{-1} \|g - g_h\|_{L^2(\Omega'_j)} \leq C |\ln h|^{3/2} \|\nabla(g - g_h)\|_{L^2(\Omega)}.$$

Proof. Define a weight function

$$\sigma(x) = \sqrt{|x - x_0|^2 + h^2}.$$

By the definition of σ it is easy to see that $\sigma \approx d_j$ on Ω_j . Thus, by the Cauchy-Schwarz inequality and using that $J \leq C |\ln h|$, we have

$$\begin{aligned} \sum_{j=j_0}^J d_j^{-1} \|g - g_h\|_{L^2(\Omega'_j)} &\leq C \left(\sum_{j=j_0}^J 1 \right)^{1/2} \left(\sum_{j=j_0}^J d_j^{-2} \|g - g_h\|_{L^2(\Omega'_j)}^2 \right)^{1/2} \\ &\leq C |\ln h|^{1/2} \left(\sum_{j=j_0}^J \|\sigma^{-1}(g - g_h)\|_{L^2(\Omega'_j)}^2 \right)^{1/2} \\ &\leq C |\ln h|^{1/2} \|\sigma^{-1}(g - g_h)\|_{L^2(\Omega)}. \end{aligned}$$

Since $g - g_h$ is zero on $\partial\Omega$, by the argument in Lemma 3.4 in [1], we have

$$\|\sigma^{-1}(g - g_h)\|_{L^2(\Omega)} \leq C |\ln h| \|\nabla(g - g_h)\|_{L^2(\Omega)},$$

which gives us the lemma. \square

Now we proceed with a duality argument.

Lemma 4.3. *There exists a constant C such that*

$$\|\nabla(g - g_h)\|_{L^2(\Omega)} \leq C.$$

Proof. We proceed similarly to [16, section 4], and define a bounded linear functional on $H_0^1(\Omega)$ by

$$(39) \quad F(\varphi) = (\nabla(g - g_h), \nabla\varphi)$$

and the corresponding solution $w \in H_0^2(\Omega)$ by

$$(40) \quad B(w, \varphi) = (D^2w, D^2\varphi) = F(\varphi), \quad \forall \varphi \in H_0^2(\Omega).$$

Since Ω is convex, $w \in H_0^2(\Omega) \cap H^3(\Omega)$ and by the regularity estimate (4), we have

$$(41) \quad \|w\|_{H^3(\Omega)} \leq C \|F\|_{H^{-1}(\Omega)} = C \sup_{\varphi \in H_0^1(\Omega)} \frac{(\nabla(g - g_h), \nabla\varphi)}{\|\nabla\varphi\|_{L^2(\Omega)}} \leq C \|\nabla(g - g_h)\|_{L^2(\Omega)}.$$

From Lemma 5 in [4], w also satisfies

$$B_h(w, \chi) = F(\chi), \quad \forall \chi \in S_h^r.$$

From the definition of w above, a priori error estimate (15), using (41) and (4), and in addition using the estimate (31) for $\|\nabla\tilde{\delta}\|_{L^2(\Omega)}$, we have

$$\begin{aligned}
\|\nabla(g - g_h)\|_{L^2(\Omega)}^2 &= F(g - g_h) = B_h(w, g - g_h) = B_h(w - I_h w, g - g_h) \\
&\leq \|w - I_h w\|_{2,h} \|g - g_h\|_{2,h} \\
&\leq Ch\|w\|_{H^3(\Omega)} Ch\|g\|_{H^3(\Omega)} \\
&\leq Ch^2\|\nabla(g - g_h)\|_{L^2(\Omega)} \|D^2\tilde{\delta}\|_{H^{-1}(\Omega)} \\
&\leq Ch^2\|\nabla(g - g_h)\|_{L^2(\Omega)} \|\nabla\tilde{\delta}\|_{L^2(\Omega)} \\
&\leq C\|\nabla(g - g_h)\|_{L^2(\Omega)}.
\end{aligned}$$

Cancelling, we obtain the result. \square

Combining Lemma 4.2 and Lemma 4.3, which establishes (38) and as a result Lemma 4.1.

5. LOCAL ERROR ESTIMATES

To show local error estimates we use again the weight function

$$(42) \quad \sigma(x) = \sqrt{|x - x_0|^2 + h^2} > 0.$$

One can easily check that σ satisfies the following properties,

$$(43a) \quad \|\sigma^{-1}\|_{L^2(\Omega)} \leq C|\ln h|^{\frac{1}{2}},$$

$$(43b) \quad |\nabla\sigma| \leq C,$$

$$(43c) \quad |\nabla^2\sigma| \leq C|\sigma^{-1}|$$

$$(43d) \quad \max_T \sigma \leq C \min_T \sigma, \quad \forall T \in \mathcal{T}_h.$$

Using this σ , we define the following weighted norms

$$\|v\|_{2,h,\sigma}^2 = \sum_{T \in \mathcal{T}_h} \|\sigma D^2 v\|_T^2 + \sum_{e \in \mathcal{E}_h} \int_e \left(\sigma^2 |e| \left| \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right|^2 + \sigma^2 |e|^{-1} \left| \left[\frac{\partial v}{\partial n} \right] \right|^2 \right) ds$$

and

$$\|v\|_{2,h,\sigma^{-1}}^2 = \sum_{T \in \mathcal{T}_h} \|\sigma^{-1} D^2 v\|_T^2 + \sum_{e \in \mathcal{E}_h} \int_e \left(\sigma^{-2} |e| \left| \left\{ \frac{\partial^2 v}{\partial n^2} \right\} \right|^2 + \sigma^{-2} |e|^{-1} \left| \left[\frac{\partial v}{\partial n} \right] \right|^2 \right) ds.$$

Lemma 5.1. *There exists a constant C independent of h such that*

$$\|g - g_h\|_{2,h,\sigma} \leq C|\ln h|^{\frac{3}{2}}.$$

Proof. Using dyadic decomposition (34) and the property that $\sigma \approx d_j$ on each Ω_j we have

$$\|g - g_h\|_{2,h,\sigma} \leq C_* h \|g - g_h\|_{2,h} + \sum_{j=j_0}^J d_j \|g - g_h\|_{2,h,\Omega_j}.$$

The rest of the proof is identical to the proof of Lemma 4.1. \square

5.1. Proof of Theorem 1.2.

Proof. Similarly to the proof of Theorem 1.1, for any $\chi \in S_h^r$ and using that $\tilde{\delta}$ is supported on T_0

$$\begin{aligned}
D^2(u_h - \chi)(x_0) &= B_h(u_h - \chi, g) \\
&= B_h(u_h - \chi, g_h) \\
&= B_h(u - \chi, g_h) \\
&= B_h(u - \chi, g) - B_h(u - \chi, g - g_h) \\
&= (u - \chi, D^2\tilde{\delta})_{T_0} - B_h(u - \chi, g - g_h)_{B_d} - B_h(u - \chi, g - g_h)_{\Omega \setminus B_d} \\
&= J_1 + J_2 + J_3,
\end{aligned}$$

where by $B(\cdot, \cdot)_D$ we denote the part of the bilinear form restricted to a set $D \subset \Omega$. Exactly as in Theorem 1.1

$$J_1 \leq \|D^2(u - \chi)\|_{L^\infty(T_0)} \|\tilde{\delta}\|_{L^1(T_0)} \leq C \|D^2(u - \chi)\|_{L^\infty(B_d)}.$$

Using the Hölder inequality and Lemma 4.1

$$J_2 \leq C \|u - \chi\|_{W_h^{2,\infty}(B_d)} \|g - g_h\|_{W_h^{2,1}(\Omega)} \leq C |\ln h|^{\frac{3}{2}} \|u - \chi\|_{W_h^{2,\infty}(B_d)}.$$

Using the Cauchy-Schwarz inequality, Lemma 5.1, and using that $\sigma^{-1} \leq Cd^{-1}$ on $\Omega \setminus B_d$, we have

$$J_3 \leq C \|u - \chi\|_{2,h,\Omega \setminus B_d, \sigma^{-1}} \|g - g_h\|_{2,h,\Omega, \sigma} \leq C |\ln h|^{\frac{3}{2}} d^{-1} \|u - \chi\|_{2,h,\Omega}.$$

□

ACKNOWLEDGMENTS

I would like to thank Boris Vexler for valuable discussions and an anonymous referee for carefully reading the manuscript and providing valuable comments.

REFERENCES

- [1] T. Apel, O. Benedix, D. Sirch, and B. Vexler, *A priori mesh grading for an elliptic problem with Dirac right-hand side*, SIAM J. Numer. Anal. **49** (2011), no. 3, 992–1005, DOI 10.1137/090778018. MR2812554
- [2] G. A. Baker, *Finite element methods for elliptic equations using nonconforming elements*, Math. Comp. **31** (1977), no. 137, 45–59, DOI 10.2307/2005779. MR431742
- [3] H. Blum and R. Rannacher, *On the boundary value problem of the biharmonic operator on domains with angular corners*, Math. Methods Appl. Sci. **2** (1980), no. 4, 556–581, DOI 10.1002/mma.1670020416. MR595625
- [4] S. C. Brenner and L.-Y. Sung, *C^0 interior penalty methods for fourth order elliptic boundary value problems on polygonal domains*, J. Sci. Comput. **22/23** (2005), 83–118, DOI 10.1007/s10915-004-4135-7. MR2142191
- [5] Z. Chen and H. Chen, *Pointwise error estimates of discontinuous Galerkin methods with penalty for second-order elliptic problems*, SIAM J. Numer. Anal. **42** (2004), no. 3, 1146–1166, DOI 10.1137/S0036142903421527. MR2113680
- [6] A. Demlow, J. Guzmán, and A. H. Schatz, *Local energy estimates for the finite element method on sharply varying grids*, Math. Comp. **80** (2011), no. 273, 1–9, DOI 10.1090/S0025-5718-2010-02353-1. MR2728969
- [7] J. Douglas Jr., T. Dupont, P. Percell, and R. Scott, *A family of C^1 finite elements with optimal approximation properties for various Galerkin methods for 2nd and 4th order problems* (English, with French summary), RAIRO Anal. Numér. **13** (1979), no. 3, 227–255, DOI 10.1051/m2an/1979130302271. MR543934

- [8] P. Grisvard, *Singularities in boundary value problems*, Recherches en Mathématiques Appliquées [Research in Applied Mathematics], vol. 22, Masson, Paris; Springer-Verlag, Berlin, 1992. MR1173209
- [9] J. Guzmán, *Pointwise error estimates for discontinuous Galerkin methods with lifting operators for elliptic problems*, Math. Comp. **75** (2006), no. 255, 1067–1085, DOI 10.1090/S0025-5718-06-01823-0. MR2219019
- [10] V. A. Kondratév, *Boundary value problems for elliptic equations in domains with conical or angular points*, Trudy Moskov. Mat. Obšč. **16** (1967), 209–292. MR0226187
- [11] V. G. Maz'ya and J. Rossmann, *On the Agmon-Miranda maximum principle for solutions of elliptic equations in polyhedral and polygonal domains*, Ann. Global Anal. Geom. **9** (1991), no. 3, 253–303, DOI 10.1007/BF00136815. MR1143406
- [12] V. G. Maz'ya and B. A. Plamenevskii, *On the maximum principle for the biharmonic equation in a domain with conical points* (Russian), Izv. Vyssh. Uchebn. Zaved. Mat. **2** (1981), 52–59. MR614817
- [13] L. S. D. Morley, *The triangular equilibrium element in the solution of plate bending problems*, Aeronautical Quarterly **19** (1968), no. 2, 149–169.
- [14] T. K. Nilssen, X.-C. Tai, and R. Winther, *A robust nonconforming H^2 -element*, Math. Comp. **70** (2001), no. 234, 489–505, DOI 10.1090/S0025-5718-00-01230-8. MR1709156
- [15] J. A. Nitsche and A. H. Schatz, *Interior estimates for Ritz-Galerkin methods*, Math. Comp. **28** (1974), 937–958, DOI 10.2307/2005356. MR373325
- [16] R. Rannacher, *On nonconforming and mixed finite element method for plate bending problems. The linear case* (English, with French summary), RAIRO Anal. Numér. **13** (1979), no. 4, 369–387, DOI 10.1051/m2an/1979130403691. MR555385
- [17] A. H. Schatz and L. B. Wahlbin, *Interior maximum norm estimates for finite element methods*, Math. Comp. **31** (1977), no. 138, 414–442, DOI 10.2307/2006424. MR431753
- [18] A. H. Schatz and L. B. Wahlbin, *Interior maximum-norm estimates for finite element methods. II*, Math. Comp. **64** (1995), no. 211, 907–928, DOI 10.2307/2153476. MR1297478
- [19] R. Scott, *Finite element convergence for singular data*, Numer. Math. **21** (1973/74), 317–327, DOI 10.1007/BF01436386. MR337032
- [20] Z. C. Shi, *On the convergence of the incomplete biquadratic nonconforming plate element* (Chinese, with English summary), Math. Numer. Sinica **8** (1986), no. 1, 53–62. MR864031
- [21] V. Thomée and L. B. Wahlbin, *Stability and analyticity in maximum-norm for simplicial Lagrange finite element semidiscretizations of parabolic equations with Dirichlet boundary conditions*, Numer. Math. **87** (2000), no. 2, 373–389, DOI 10.1007/s002110000184. MR1804662
- [22] L. B. Wahlbin, *Local behavior in finite element methods*, Handbook of numerical analysis, Vol. II, Handb. Numer. Anal., II, North-Holland, Amsterdam, 1991, pp. 353–522. MR1115238
- [23] M. Wang, *L^∞ error estimates of nonconforming finite elements for the biharmonic equation*, J. Comput. Math. **11** (1993), no. 3, 276–288. MR1229633

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT 06269
Email address: dmitriy.leykekhman@uconn.edu