



Complete Genome Sequence of *Rhodococcus qingshengii* Strain CL-05, Isolated from Concrete

 E. Anders Kiledal,^a Shannon G. McDermott,^b Olga Shevchenko,^c  Jean Ross,^d Stacy Bediako,^b  Julia A. Maresca^b

^aDepartment of Biological Sciences, University of Delaware, Newark, Delaware, USA

^bDepartment of Civil and Environmental Engineering, University of Delaware, Newark, Delaware, USA

^cSequencing and Genotyping Center, University of Delaware, Newark, Delaware, USA

^dBioImaging Center, University of Delaware, Newark, Delaware, USA

ABSTRACT Here, we report the complete genome sequence of *Rhodococcus qingshengii* strain CL-05, which was isolated from pavement concrete in Newark, Delaware. The genome consists of a 6.29-Mbp chromosome and one plasmid (123,183 bp), encodes a total of 5,859 predicted proteins, and has a GC content of 62.5%.

We previously isolated several bacterial strains from concrete, an alkaline, high-salt environment (1), and then sequenced the genome of *Rhodococcus* sp. strain CL-05 to compare it with those of alkaliphilic and halophilic *Rhodococcus* species (2–4).

Pieces of concrete (~1 g) were vortexed in TE buffer (10 mM Tris, 1 mM EDTA), and then 25 μ l of solution was spread onto concrete medium solidified with agar (CM-A) and incubated at room temperature for ~2 weeks (1). Individual colonies were restreaked onto CM-A until axenic, as determined by microscopy. CL-05 cells are ~2.5- to 4- μ m rods (Fig. 1), and this isolate was initially identified as *Rhodococcus erythropolis* by Sanger sequencing of its 16S gene (primers 8F and 1492R [1, 5]).

Strain CL-05 was revived on nutrient agar (NA) (catalog number OXCM0003; Fisher) from a –80°C stock (in 7% dimethyl sulfoxide). One colony was restreaked onto NA and grown overnight at 28°C, and then one colony was transferred to 50 ml nutrient broth and grown overnight. DNA was extracted using a phenol-chloroform extraction protocol optimized for Gram-positive bacteria (6). A single-molecule real-time (SMRT) library was barcoded and prepared using the PacBio SMRTbell Express template preparation kit version 2.0 (7). DNA fragments larger than 6 kb were size selected using BluePippin (Sage Science). The average library fragment size was 15 kb, as measured by a fragment analyzer (Advanced Analytical Technologies, Inc.). Sequencing was completed on a PacBio Sequel single-molecule sequencer in one 1M version 3 LR SMRT Cell with a 20-h movie. Samples were demultiplexed using PacBio SMRT Link version 9.0.0.92188.

We obtained 67,539 barcoded reads (size range, 51 to 91,079 nucleotides [nt]; N_{50} , 12,320 nt). Demultiplexed raw subreads were downloaded from PacBio SMRT Link, converted to .fastq files with bam2fastx version 1.3.1, and then chimera checked, quality controlled, assembled, and circularized using Flye version 2.8.2 (8). The genome was rotated to start at *dnaA* using Circlator version 1.5.5 (fixstart method) (9). The assembly was polished using raw reads aligned with the PacBio minimap2 version 1.3.0 wrapper (10) and the Arrow polishing algorithm in pbcpp version 1.0.0 (11). The assembled genome consists of a 6,290,587-bp circular chromosome (GC content of 62.5%, with 90 \times coverage) and one circular plasmid (123,183 bp; GC content of 62.4%, with 119 \times coverage). Default parameters were used for all software unless otherwise noted; the pipeline is available at github.com/MarescaLab/genome_pipeline.

FastANI in GTDB-Tk version 1.4.0 (12, 13) identified the closest relative of CL-05 as *Rhodococcus qingshengii* strain JCM 15477 (GenBank accession number [GCF_001646745.1](https://www.ncbi.nlm.nih.gov/nuccore/GCF_001646745.1)), with

Citation Kiledal EA, McDermott SG, Shevchenko O, Ross J, Bediako S, Maresca JA. 2021. Complete genome sequence of *Rhodococcus qingshengii* strain CL-05, isolated from concrete. *Microbiol Resour Announc* 10: e00376-21. <https://doi.org/10.1128/MRA.00376-21>.

Editor David A. Baltrus, University of Arizona

Copyright © 2021 Kiledal et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Julia A. Maresca, jmaresca@udel.edu.

Received 12 April 2021

Accepted 21 April 2021

Published 13 May 2021




FIG 1 Scanning electron microscopy (SEM) of CL-05 cells. Cells were grown overnight in nutrient broth for SEM. After centrifugation, cells were resuspended in fresh 4% paraformaldehyde in phosphate-buffered saline. Suspended cells in fixative were placed onto poly-L-lysine-coated silicon wafers. After cells were allowed to attach, the sample wafers were washed with 0.1 M sodium cacodylate buffer and then incubated with 1% osmium tetroxide in 0.1 M sodium cacodylate buffer. The samples were dehydrated in a graded series of ethanol concentrations from 50% to 100%, transferred to hexamethyldisilazane, and air dried. Wafers were mounted onto SEM stubs, sputter coated with 3-nm platinum, and imaged with a Thermo Fisher Scientific Apreo SEM.

an average nucleotide identity (ANI) of 98.78% over 93% of the input sequence. The plasmid is >99% identical to a plasmid from *R. qingshengii* strain IGTS8 (GenBank accession number [GCA_006384225.1](#)).

PGAP build 5132 (14) was used to predict open reading frames and to annotate genes, using default parameters. The chromosome contains 5 rRNA operons, 59 tRNAs, 1 transfer-messenger RNA, and 5,726 predicted protein-coding genes; the plasmid contains 133 protein-coding genes.

Using BLASTp, we identified two putative Na^+/H^+ antiporters (*nhaA* and *mrpABCDEFG* [15, 16]) that are useful in halotolerance and alkalitolerance (17) and the putative betaine transporter *betP*, which is involved in osmoprotection (18), in the CL-05 genome.

Data availability. The raw reads have been submitted to the SRA and have the accession number [SRR13722032](#). The assembled, annotated genome is available in the NCBI database under BioProject number [PRJNA702129](#) (chromosome, accession number [CP072108](#); plasmid, accession number [CP072109](#)).

ACKNOWLEDGMENTS

This publication was made possible by National Science Foundation EPSCoR grant 1757353 and support from the State of Delaware, the University of Delaware early-career research apprenticeship program (UDRAW), and the Delaware Department of Transportation, with funding from the Federal Highway Administration (task 27-1891).

REFERENCES

1. Maresca JA, Moser P, Schumacher T. 2017. Analysis of bacterial communities in and on concrete. *Mater Struct* 50:25. <https://doi.org/10.1617/s11527-016-0929-y>.
2. Mayilraj S, Krishnamurthi S, Saha P, Saini HS. 2006. *Rhodococcus kroppenstedtii* sp. nov., a novel actinobacterium isolated from a cold desert of the Himalayas, India. *Int J Syst Evol Microbiol* 56:979–982. <https://doi.org/10.1099/ijss.0.63831-0>.
3. Goordial J, Raymond-Bouchard I, Ronholm J, Shapiro N, Woyke T, Whyte L, Bakermans C. 2015. Improved-high-quality draft genome sequence of *Rhodococcus* sp. JG-3, a eurypsychrophilic actinobacteria from Antarctic Dry Valley permafrost. *Stand Genomic Sci* 10:61. <https://doi.org/10.1186/s40793-015-0043-8>.
4. Hjerde E, Pierechod MM, Williamson AK, Bjerga GEK, Willassen NP, Smalås AO, Altermark B. 2013. Draft genome sequence of the actinomycete

Rhodococcus sp. strain AW25M09, isolated from the Hadsel Fjord. Genome Announc 1:e0005513. <https://doi.org/10.1128/genomeA.00055-13>.

- Turner S, Prysor KM, Miao VP, Palmer JD. 1999. Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis. *J Eukaryot Microbiol* 46:327–338. <https://doi.org/10.1111/j.1550-7408.1999.tb04612.x>.
- Kiledal EA, Maresca JA. 2021. Chromosomal DNA extraction from Gram-positive bacteria. protocolsio <https://doi.org/10.17504/protocols.io.bs7knhkw>.
- Pacific Biosciences of California, Inc. 2020. Preparing multiplexed microbial libraries using SMRTbell express template prep kit 2.0. Pacific Biosciences of California, Inc., Menlo Park, CA.
- Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol* 37:540–546. <https://doi.org/10.1038/s41587-019-0072-8>.
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Li H. 2018. minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. <https://doi.org/10.1093/bioinformatics/bty191>.
- Pacific Biosciences. GCpp: generate highly accurate reference contigs. github.com/PacificBiosciences/gcpp.
- Jain C, Rodriguez-R LM, Phillippe AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 9:5114. <https://doi.org/10.1038/s41467-018-07641-9>.
- Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 36:1925–1927. <https://doi.org/10.1093/bioinformatics/btz848>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Huang Y, Chen W, Dotson DL, Beckstein O, Shen J. 2016. Mechanism of pH-dependent activation of the sodium-proton antiporter NhaA. *Nat Commun* 7:12940. <https://doi.org/10.1038/ncomms12940>.
- Fang H, Qin X-Y, Zhang K-D, Nie Y, Wu X-L. 2018. Role of the group 2 Mrp sodium/proton antiporter in rapid response to high alkaline shock in the alkaline- and salt-tolerant *Dietzia* sp. DQ12-45-1b. *Appl Microbiol Biotechnol* 102:3765–3777. <https://doi.org/10.1007/s00253-018-8846-3>.
- Krulwich TA, Sachs G, Padan E. 2011. Molecular aspects of bacterial pH sensing and homeostasis. *Nat Rev Microbiol* 9:330–343. <https://doi.org/10.1038/nrmicro2549>.
- Wood JM. 2015. Bacterial responses to osmotic challenges. *J Gen Physiol* 145:381–388. <https://doi.org/10.1085/jgp.201411296>.