

Quantification of Ni-N-O bond angles and NO activation by X-ray emission spectroscopy.

Phan N. Phu,^{1†} Carlos E. Gutierrez,¹ Subrata Kundu,^{2,3} Dimosthenis Sokaras,⁴ Thomas Kroll,⁴ Timothy H. Warren,^{2*} S. Chantal E. Stieber^{1*}

¹Department of Chemistry & Biochemistry, California State Polytechnic University, Pomona, California 91768, United States.

²Department of Chemistry, Georgetown University, Box 571227, Washington, D. C. 20057, United States.

³School of Chemistry, Indian Institute of Science Education and Research Thiruvananthapuram, Kerala 695551, India.

⁴Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, Menlo Park, California 94025, United States.

Supporting Information Placeholder

ABSTRACT: A series of β -diketiminate Ni-NO complexes with a range of NO binding modes and oxidation states were studied by X-ray emission spectroscopy (XES). The results demonstrate that XES can directly probe and distinguish end-on versus side-on NO coordination modes, as well as 1-electron NO reduction. Density functional theory (DFT) calculations show that the transition from the NO 2s2s σ^* orbital has higher intensity for end-on NO coordination than for side-on NO coordination, whereas the 2s2s σ orbital has lower intensity. XES calculations in which the Ni-N-O bond angle was fixed over the range from 80-176° suggest that differences in NO coordination angles of approximately 10 degrees could be experimentally distinguished. Calculations of Cu nitrite reductase (NiR) demonstrate the utility of XES for characterizing NO intermediates in metalloenzymes. This work shows the capability of XES to distinguish NO coordination modes and oxidation states at Ni and highlights applications in quantifying small molecule activation in enzymes.

INTRODUCTION

Metalloenzymes play key roles in biological signaling of nitric oxide (NO) and reduction of NO_x species, with metal nitrosyls (M-NO) oftentimes invoked as key intermediates. M-NO species are known or proposed as intermediates in enzymatic reduction of NO₂⁻,¹ reduction of NO,^{2,3,4} and conversion of hydroxylamine to N₂O,⁵ although full mechanisms are generally unknown for these systems.^{4,6} An understanding of the mechanisms of enzymatic NO or NO_x reduction would greatly impact

development of catalysts for NO and NO_x reduction or activation.

A particular challenge in metal-NO complexes is the ability for both the metal and the ligand to participate in redox events and the difficulty in distinguishing these, especially for first row transition metals.⁷ NO is regularly found as NO¹⁺, NO, NO¹⁻ and NO²⁻, but commonly its interactions with metal centers are highly covalent and difficult to discern. Therefore, the Enemark-Feltham notation is generally used to account for the total valence electrons in a system by using the sum of metal *d*-electrons and NO π^* electrons.⁸ However, a more detailed understanding of the metal and NO oxidation states would add clarity to ligand activation and enzyme mechanisms in NO redox chemistry.

Infrared spectroscopy (IR) is generally used to characterize the presence of a metal nitrosyl, and the IR stretch is sensitive to NO activation/NO bond length and the metal-N-O bond angle.^{9,10,11} The contributions from activation and bond angle, however are difficult to deconvolute and not well-quantified.^{9,12} A further complication is the limited ability to correlate calculated electronic structures and M-N-O bond angles with IR data.¹³ As such, electronic structure characterization of M-NO species generally involves significant spectroscopic and computational efforts. Determination of the M-N-O bond angle is especially important in enzymes such as the reduced forms of nitrite reductase (NiR) that exhibit side-on NO bonding. Crystallography allowed for a side-on nitrosyl to be observed for copper, however X-ray structures are often difficult to obtain for enzymes.¹

X-ray absorption spectroscopy (XAS) is often used to characterize enzyme metal centers and intermediates when crystal structures are not obtainable. Bond distanc-

es from the metal center can be determined to reasonable levels of accuracy (0.02 Å) with XAS.¹⁴ A significant limitation for NO chemistry is that XAS cannot distinguish differences in coordination number of ± 1 , similar light atoms such as C, N, and O, and protonation states. This implies that variations in M-N-O angles including end-on, bent, and side-on coordination modes of NO could not be distinguished or quantified by XAS alone.

X-ray emission spectroscopy (XES) overcomes some of the limitations of XAS,^{15,16} and is perhaps best known for being able to distinguish C and N from O, as demonstrated in Cr compounds,¹⁷ Mn complexes,¹⁸ a cluster of iron centers, and the enzyme nitrogenase.^{19,20} These significant findings highlighted the value of XES to bio-inorganic chemistry the understanding of enzyme active sites. More recent enzyme studies included two-color VtC XES probing both Mn and Fe centers of intermediates in ribonucleotide reductase,^{21,22} peptidylglycine monooxygenase,²³ and galactose oxidase.²⁴

Valence-to-core (VtC) XES probes occupied valence orbitals by detecting relaxation of electrons from ligand-centered $2p$ and $2s$ orbitals to the metal-centered $1s$ orbital (Figure 1.A.).²⁵ Gradual improvements in instrumentation and detectors have allowed the very low intensity VtC to be observed with sufficient resolution to become a useful spectral tool for chemists. However, there is still much development needed using model complexes to fully understand effects of varying coordination environments on spectral features and render the technique more versatile. Recent work highlighted the ability for VtC XES to distinguish a) 1 versus 2 N_2 or NO molecules coordinated at Fe (Figure 1.B.2.),^{26,27} b) N_2 reduction (Figure 1.B.3.),²⁸ c) bis μ -oxo versus μ -oxo versus μ -hydroxo,²⁹ d) O_2 activation (Figure 1.B.3.),³⁰ and nickel hydrides.³¹ In combining these results, it can be generally understood that portions of the VtC $K\beta_{2,5}$ region increase in intensity upon additional ligand coordination (Figure 1.B.2.).^{26,27} Additionally, some $K\beta_{2,5}$ features shift to lower energy and the energy separation between the $2s$ σ and σ^* decreases with increased bond activation (Figure 1.B.3.).^{28,30} The VtC $K\beta''$ is more difficult to experimentally detect, but decreases in energy with increased electronegativity of the ligand (such as C to N to O, Figure 1.B.1.).

XES characterization of four-coordinate Fe-NO complexes with SPh and OPh ancillary ligands determined that one versus two NO ligands bound could be distinguished by XES.^{27,32} Computationally, they reported that XES should be able to distinguish NO oxidation states based on differences in energies of the σ_{2s}^* and σ_{2p} orbitals. However, the complexes in the study were sufficiently similar that differences in NO oxidation states could not be fully experimentally distinguished. In the current work, two new areas for experimentally applying XES are reported, including heteronuclear ligand activation and ligand coordination mode (Figure 1).





Figure 1. Schematic of (A) X-ray emission spectroscopy (XES) and (B) significant applications for determining ligand and atom identity, coordination number, and ligand activation at metal centers.

RESULTS AND DISCUSSION

XES Spectra of Nickel Nitrosyl Complexes. Three previously reported β -diketiminate nickel(II) nitrosyl complexes were studied with end-on versus side-on coordination geometries and NO oxidation states best described as NO^1- versus NO^{2-} .³³ The complexes $[Pr_2NNF_6]NiNO$ (end-on Ni- (NO^1-)), $[Pr_2NNF_6]Ni(\mu-\eta^1:\eta^1-NO)K[2.2.2-cryptand]$ (end-on Ni- (NO^{2-})), and $[Pr_2NNF_6]Ni(\mu-\eta^2:\eta^2-NO)K(18-crown-6)(THF)$ (side-on Ni- (NO^{2-})) were synthesized as previously described (Scheme 1).³³ These nickel(II) complexes were extensively characterized by X-ray crystallography, Ni K-edge XAS, and DFT analysis, making them ideal for probing capabilities of XES to distinguish NO oxidation states and coordination modes. It should be noted that end-on Ni- (NO^{2-}) was crystallographically characterized and modelled as a disordered NO with 77% end-on and 23% side-on NO^{2-} .

Scheme 1. Reduction of end-on Ni- (NO^1-) to form end-on and side-on Ni- (NO^{2-}) .



The K β mainline XES spectra for the three complexes overlay (Figure S1), providing additional experimental support for these complexes having the same oxidation and spin states at nickel. Although spin states and covalency can also affect the K β mainline,³⁴ this suggests a similar electronic environment at the nickel center in each complex. The VtC for the three complexes are distinct (Figure 2a), with differences in both the K β '' (~8300-8320 eV) and K β _{2,5} (~8320-8336 eV) regions.




Figure 2. VtC XES (a) of end-on Ni-(NO¹⁻) (green), end-on Ni-(NO²⁻) (blue), and side-on Ni-(NO²⁻) (red), and calculated VtC XES (b). Left inset shows end-on versus side-on Ni-(NO²⁻) from 8300-8320 eV. A shift of 217.0 eV and a broadening of 2.5 eV were applied to the computed data.

The VtC XES spectrum of end-on Ni-(NO¹⁻) has a distinct shoulder at 8318.2 eV (Figure 2a, peak C), which has lower intensity for end-on Ni-(NO²⁻), and no intensity for side-on Ni-(NO²⁻). The K β _{2,5} feature around 8326 eV (Figure 2a, peak D) has the highest intensity for side-on Ni-(NO²⁻), and decreases in intensity for end-on Ni-(NO²⁻) followed by end-on Ni-(NO¹⁻), with a corresponding shift to lower energy. This suggests that the features at 8318.2 eV and 8326 eV are sensitive to both the NO oxidation state and coordination mode. The feature around 8309 eV (Figure 2a, peak B) was observed for all complexes with nacnac ligands. In the K β '' re-

gion, the side-on Ni-(NO²⁻) has a feature present around 8306 eV (Figure 2a, peak A), which is notably absent in the spectra of the two end-on complexes. This supports the ability for XES to distinguish end-on versus side-on coordination modes for NO²⁻.

DFT Calculations: Simulating XES Spectra. DFT calculations were conducted using the ORCA program³⁵ to offer insight into XES spectral features in the VtC. A simple one-electron, ground state model has been well-established to offer good agreement with experimental VtC XES spectra for Cr,³⁶ Mn,^{29,37} Fe,^{19,20,26,28,38,39} Co,⁴⁰ and Cu.^{30,23} The calculated VtC XES spectra using BP86 with the CP(PPP) basis set⁴¹ on Ni and TZVP⁴² on all other atoms on geometry optimized structures for the three complexes (Figure 2) are distinct, and in good agreement with experimental data with regards to both energies and intensities. The same general trends of a decrease in intensity for the feature at 8322 eV (Figure 2b, peak C) and an increase in intensity for the feature around 8311 eV (Figure 2b, peak A) were observed from end-on Ni-(NO¹⁻) to end-on Ni-(NO²⁻) to side-on Ni-(NO²⁻).

Spectral contributions were analyzed through visualization of molecular orbitals, since the transitions were calculated from valence orbitals to 1s orbitals in a ground-state model. Previous studies reported that higher sigma overlap of ligand orbitals with metal orbitals result in more intense spectral features.²⁵ The calculated VtC XES for end-on Ni-(NO¹⁻), end-on Ni-(NO²⁻), and side-on Ni-(NO²⁻) spectra along with orbital contributions highlight the lowest energy feature as resulting from NO-based orbitals for all three complexes (Figure 3). For all three complexes, the feature at (a) 8308-8311 eV is the 2s(N)-2s(O) σ bonding combination, (b) 8314-8316 is the nacnac ligand σ and b₁ σ^* , and (c) 8322 eV is the 2s(N)-2s(O) σ^* orbital. The higher energy features >8323 eV correlate to orbitals that are combinations of 2p σ , π and π^* interactions of the NO and nacnac ligands, with the feature around 8335 having significant NO π^* contributions.

A closer examination of the orbitals reveals a shift to higher energy for the 2s(N)-2s(O) σ bonding orbital upon NO reduction from end-on NO¹⁻ at 8308.5 eV to end-on NO²⁻ at 8310.1 eV. This is consistent with previous XES observations of increased energy of the 2s σ bonding orbital upon reduction of N₂,²⁸ NO,²⁷ and O₂.³⁰ In the current work, the 2s(N)-2s(O) σ^* orbital shifts to slightly higher energy upon NO reduction from end-on NO¹⁻ at 8322.1 eV to end-on NO²⁻ at 8322.5 eV; however, the shift is much smaller. The energy difference between the 2s(N)-2s(O) σ and σ^* decreases from 13.6 eV for end-on Ni-(NO¹⁻) to 12.4 eV for end-on Ni-(NO²⁻) and 11.5 eV for side-on Ni-(NO²⁻). These are consistent with differences of 8.4 – 12.4 eV as calculated for free NO¹⁻,




Figure 3. Calculated VtC XES spectrum of end-on Ni-(NO¹⁻), (left), end-on Ni-(NO²⁻) (middle), and side-on Ni-(NO²⁻), (right) with the molecular orbitals that strongly contribute to the observed transitions. A shift of 217.0 eV and a broadening of 2.5 eV were applied.

and NO²⁻ (Figure S2). The magnitude is generally larger than reported 2s σ and σ^* energy differences of 5-6 eV in Fe-NO,²⁷ 6-8 eV in Cu-O₂²⁻,^{30,43} and 12 eV in Fe-N₂,²⁸ which is consistent with more activate NO. The highest energy NO π^* signal shifts to lower energy from 8334.0 eV to 8333.6 eV upon reduction. This suggests that XES is sensitive to NO reduction, although the intensity of 2s(N)-2s(O) σ bonding orbital is too low to be experimentally resolved within current detection limits for nickel. Therefore, the shift in higher energy of the 2s(N)-2s(O) σ^* orbital and the shift to lower energy of the NO π^* can be used to assess NO reduction from NO¹⁻ to NO²⁻ using XES.

Connecting XES Spectra with Ni-NO Molecular Orbitals. The observed effects on the XES spectrum of NO coordination geometry as end-on or side-on are clarified by examining the molecular orbitals (Figure 3). The 2s(N)-2s(O) σ bonding feature at 8308-8311 eV has a greater intensity for side-on Ni-(NO²⁻) than end-on Ni-(NO²⁻). This is a result of increased orbital overlap between Ni and the oxygen atom of NO. Strong overlap of the side-on NO²⁻ π orbital with Ni results in the feature at ~8328 eV having higher intensity for side-on NO²⁻ than end-on NO²⁻. For all three complexes, transitions from the NO π^* orbitals are at energies >8325 eV.

From previously reported geometry optimization calculations,³³ strong orbital overlap of the Ni d_{xy} and d_{yz} orbitals with the NO π^* orbitals contribute to complex formation for end-on Ni-(NO¹⁻) and NO activation. Upon one-electron reduction to Ni-(NO²⁻), the Ni-N bond distance lengthens, while the Ni-nacnac bond distances shorten. Since the Ni d_{yz} orbital also has the correct symmetry to interact with nacnac π^* orbitals, the Ni d_{yz} orbital can be stabilized through backbonding to nacnac

π^* orbitals, allowing the NO π^* orbitals to accept the electron and be further activated. These effects are manifest in the VtC XES spectra by the slight shift to lower energy of the nacnac peak b observed upon reduction from Ni-(NO¹⁻) to Ni-(NO²⁻). Similarly, the highest energy peak, which has NO π^* contributions, shifts to lower energy upon reduction from Ni-(NO¹⁻) to Ni-(NO²⁻). Since the d_{yz} orbital has weaker backbonding to NO²⁻, both end-on and side-on coordination are possible with minimal effects on the orbital energies.

Probing the Effect of Ni-N-O Bond Angle. To probe capabilities for XES to distinguish NO coordination modes, XES calculations of Ni-NO complexes with hypothetical Ni-N-O angles ranging from 102-176° for Ni-(NO¹⁻) and 80-160° for Ni-(NO²⁻) were commenced (Figures 4, S3-6). A geometry optimization was conducted at each probed angle, while keeping the Ni-N-O angle fixed. It was not possible to find solutions for Ni-(NO¹⁻) complexes with angles smaller than 102.3°. For both Ni-(NO¹⁻) and Ni-(NO²⁻), the feature to lowest energy (around 8306-8311 eV) increases in intensity for decreasing Ni-N-O angles, while the feature around 8322 eV decreases. The lowest energy feature around 8310.7 eV corresponds to the 2s(N)-2s(O) σ bonding orbital and increases in intensity for smaller Ni-N-O angles due to increased overlap with metal orbitals (Figure S5). The Ni p -character of the 2s(N)-2s(O) σ bonding orbital increases the most strongly with increased intensity by approximately 2%, however both Ni s -character and d -character also increase with intensity over the calculated range of angles for Ni-(NO²⁻). Nitrogen s -character (~25%) and p -character (~13%) as well as oxygen s -character (50%) and p -character (~8%) of the 2s(N)-2s(O) σ bonding orbital remain relatively consistent over the calculated range of angles for Ni-(NO²⁻).

The feature around 8322 eV corresponds to the 2s(N)-2s(O) σ^* orbital and decreases in intensity with decreasing bond angle. The Ni p -character of the 2s(N)-2s(O) σ^* orbital increases by 2.6% with increasing signal intensity over the 70° range of angles probed for Ni-(NO²⁻), while both Ni s -character and d -character increase by approximately 2% (Figure S6). With increasing intensity, the nitrogen s -character and p -character generally increased by 8-10%, while oxygen s -character and p -character generally decreased by 8-10% in the 2s(N)-2s(O) σ^* orbital over the calculated range of angles for Ni-(NO²⁻). One complication in the geometry optimizations is that the Ni-N bond distance lengthened by approximately 0.1 Å with decreasing Ni-N-O angle (Table S1), so the observed effects will be slightly affected by this (see following). The N-O distance only changed by 0.05 Å across the series, so there is no significant activation. These results suggest that XES is potentially sensitive to changes in Ni-N-O bond angle of as small as 10°.




Figure 4. Calculated VtC XES spectrum of Ni-(NO²⁻) with varying the Ni-N-O angle from 80-160°. A shift of 217.0 eV and a broadening of 2.5 eV were applied.

Probing the Effects of Ni-N and N-O Distances. The effect of NO reduction on the XES spectral features was further examined through calculations of Ni-(NO¹⁻) where the N-O bond distance was fixed from 1.459 – 1.859 Å in a series of single point calculations without additional geometry optimization (Figure 5). These calculations reveal that both the lowest energy nacnac feature around 8316 eV and the feature around 8330 eV shift to higher energy upon lengthening of the N-O bond. The lowest energy nacnac feature shifts from 8315.9 eV to 8316.5 eV along with an increase in N-O bond length from 1.459 Å to 1.859 Å. The larger effect is observed in the 2p(N)-2p(O) feature at 8329.3 eV, which shifts to higher energy and significantly decreases in intensity along with an increase in N-O bond length from 1.459 Å to 1.859 Å. Notably, the 2s(N)-2s(O) σ^* feature around 8322 eV is not affected by the N-O bond

distance, supporting that it is not sensitive to NO reduction in this nacnac ligand system. Combined, these calculations support that XES is sensitive to changes in nitrosyl N-O bond distances and therefore sensitive to NO reduction.

Effects of Ni-NO bond lengths were examined through XES calculations of Ni-(NO¹⁻) where the Ni-N bond distance was fixed from 1.426 – 2.126 Å without geometry optimization (Figure 5). The resulting XES spectra reveal that features above 8318 eV sequentially increase in intensity with decreasing Ni-N distance, and that features below 8318 eV are not significantly altered. The increasing intensity of the NO 2s(N)-2s(O) σ^* feature around 8322 eV correlates with the increasing Ni 4s character in the molecular orbital contributing to the observed transition (Figure S7). A correlation of XES VtC intensity, as determined by the total oscillator strength, and Ni p -character was determined to have an exponential correlation (Figure S7).




Figure 5. Calculated VtC XES spectrum of end-on Ni-(NO¹⁻) with fixed N-O distances (top) and fixed Ni-N distances (bottom). A shift of 217.0 eV and a broadening of 2.5 eV were applied.

The correlation of XES VtC intensity with metal np character has been well-established and is consistent with previous reports.^{39,44,45} The XES VtC intensity was also found to correlate exponentially to Ni s -character (Figure S7), which is in contrast to previous reports where no correlation of XES VtC intensity to metal s -character was observed.⁴⁶ It is likely that this is due to the low symmetry of the current system being studied, as compared to previous reports with higher symmetry molecules.

Application of XES to Cu-NO in CuNiR. The applicability of these XES findings to metalloenzymes was probed through XES calculations of the active site in copper nitrite reductase (NiR). The crystal structures of a side-on Cu-NO and Cu-NO₂ moiety within NiR were previously reported (Figure 6), and are a rare example of structural characterization of this motif within an enzyme.¹ Structural coordinates for the side-on [(his)₃Cu-NO]⁺ and [(his)₃Cu-NO₂]⁺ centers and surrounding histidine ligands were used directly for XES calculations without further optimization (Figure 6), and a hypothetical end-on Cu-NO was also calculated.¹ The reported crystal structures were consistent with either side-on Cu(I)-(NO⁺) or side-on Cu(II)-(NO⁻),¹ so both possibilities were input for XES calculations (see Figure S8).




Figure 6. Calculated VtC XES spectrum of NiR for side-on Cu(I)-(NO⁺), end-on Cu(I)-(NO⁺), and Cu(II)-NO₂ along with structural models (orange = copper, red = oxygen, purple = nitrogen, gray = carbon, white = hydrogen). Calculated energies are not calibrated, and a broadening of 2.5 eV was applied to calculated spectra.

The calculated XES spectra for Cu NiR reveal that side-on Cu-NO, end-on Cu-NO, and Cu-NO₂ centers should be experimentally distinguishable by XES. This is significant for potential mechanistic studies using time-resolved XES methods.^{47–50} A closer look at the spectra reveals similar trends and molecular orbital contributions to what was observed in the Ni-NO system (Figures S9–S11). The lowest energy 2s(N)-2s(O) σ bonding feature at 8278.3 eV for side-on Cu(I)-(NO⁺) is

shifted 0.8 eV to lower energy than for end-on Cu(I)-(NO⁺) and has higher intensity. The 2s(N)-2s(O) σ^* feature at 8736.6 eV has significantly lower intensity for side-on Cu(I)-(NO⁺) and shifts 0.7 eV to lower energy for end-on Cu(I)-(NO⁺). There is no equivalent feature around this energy for Cu-NO₂. The 2p(N)-2p(O) feature around 8743 eV significantly decreases in intensity for Cu-NO₂, supporting considerable nitrosyl contributions to this feature as was observed for Ni-NO.

CONCLUSIONS

The Ni K β VtC XES spectra of end-on Ni-(NO¹⁻), end-on Ni-(NO²⁻), and side-on Ni-(NO²⁻) indicate that XES is sensitive to the oxidation states of NO¹⁻ versus NO²⁻, and end-on versus side-on NO coordination, and is among the first Ni K β VtC XES reports.³¹ This is a new example of the sensitivity of XES to the coordination mode of a small molecule, as demonstrated by sensitivity to Ni-N-O bond angle changes of as small as 10°. Calculations of Cu NiR suggest that these results are applicable for characterizing side-on versus end-on NO in Cu metalloenzymes and that these intermediates can be distinguished from Cu-NO₂. This expands capabilities for XES to characterize bond activation of small molecules and demonstrates the potential for XES to be used to characterize enzyme intermediates in NO and NO_x reduction. Specifically, future time-resolved XES experiments^{47–50} have the potential to probe previously uncharacterized intermediates in nitrite reductases.

EXPERIMENTAL SECTION

General Methods. Samples of [¹Pr₂NNF₆]NiNO (end-on Ni-(NO¹⁻)), [¹Pr₂NNF₆]Ni(μ - η^1 : η^1 -NO)K[2.2.2-cryptand] (end-on Ni-(NO²⁻)), and [¹Pr₂NNF₆]Ni(μ - η^2 : η^2 -NO)K(18-crown-6)(THF) (side-on Ni-(NO²⁻)) were synthesized as previously reported.³³ Structural coordinates for Cu NiR active sites including side-on Cu-NO and Cu-NO₂ were taken from reported crystal structures of the full enzyme with PDB codes 1SNR and 1SJM.¹

XES Experiments. The X-ray emission spectroscopy experiments were conducted on beamline 6-2 (3 GeV, 500 mA) at the Stanford Synchrotron Radiation Lightsource (SSRL).⁵¹ Beamline 6-2 is equipped with an LN₂ cooled Si[111] monochromator, which was calibrated to a nickel foil at 8332.8 eV (first inflection point). The multi-crystal Johann spectrometer used three spherically bent Si[551] crystals (100 mm diameter, 1 m radius of curvature) on a Rowland geometry; a Si drift detector (SDD) with a 1.5 mm vertical slit was used. The spectrometer energy was calibrated across its overall operational energy range using a sequence of elastic scattering measurements from the incident tunable monochromatic x-ray beam. Ni samples were sealed in ampules for shipment to the synchrotron facility. The samples were manipulated in an Ar glovebox at the synchro-

tron facility. Samples were finely ground and pressed into a 1 mm Al holder without further dilution and sealed with Kapton tape. The samples were transferred to the beamline under liquid N₂ and attached to the sample rod under liquid N₂. A liquid helium Oxford CFI208 cryostat was attached to maintain a sample temperature of ~10 K. To reduce signal attenuation, a He-filled flight path was employed. To prevent sample damage, aluminum filters were utilized before the sample to attenuate the incident X-ray. Radiation damage scans were conducted for each sample prior to measurements. In addition, two scans were collected per sample spot to assess radiation damage, and only scans that showed no evidence for radiation damage were used for data averages. The incident flux was measured with a He-filled ion chamber and used to normalize the measured emission signal. Data were processed and normalized using PyMCA⁵² and visualized with Kaleidagraph.

DFT Calculations. DFT calculations were conducted using the ORCA program as previously described,^{26,35} and using the XSEDE computational resources.⁵³ Avogadro was used to center coordinates and modify structural inputs.⁵⁴ Geometry optimizations were carried out at the B3LYP level of theory.^{55–57} B3LYP is a hybrid-functional that has been shown to offer a more accurate description for metal-ligand bond lengths and interactions in 3d transition metal compounds.⁵⁸ The all-electron Ahlrichs basis sets were applied,^{59–61} utilizing triple- ξ def2-TZVP basis sets on the metal center and all atoms directly bound to the metal center. All other atoms were described using double- ξ def2-SV(P) basis sets. Auxiliary basis sets were selected to match the orbital basis.^{62,63} The RIJCOSX approximation was used for computational expediency. For broken symmetry calculations, the notation BS(x,y) signifies x spin up electrons and y spin down electrons.^{64,65} The resulting coupled orbital pair was visualized by plotting the corresponding orbitals (uco). Orbitals and spin density plots were visualized using the program Chimera.⁶⁶

The XES spectra were calculated using the simple one-electron approximation as previously described,²⁶ using the BP86 functional, the CP(PPP) basis set on nickel and copper,⁴¹ and the TZVP basis set on all other atoms.⁴² A dense integration grid with an integration accuracy of 7 was applied to increase radial integration accuracy. Calculated XES spectra (XESQ) were plotted using ORCA_MAPSPC over a range of 7500–8500 eV for Ni and 8000–9000 eV for Cu, with a weighting of 2.5 eV, and 10,000 points. Calculated spectra for Ni were shifted by 271.0 eV, and Cu XES spectra were not energy calibrated. All orbitals and spin density distributions were plotted using Chimera.⁶⁶ XAS and XES spectra were visualized using Kaleidagraph.

Supporting Information. The Supporting Information is available free of charge on the ACS Publications website and includes a PDF file containing XES and computational details. Computational xyz input files may be obtained free of charge at <https://zenodo.org> under DOI: 10.5281/zenodo.3911324.

AUTHOR INFORMATION

Corresponding Author

*Email for S.C.E.S.: sestieber@cpp.edu; orcid.org/0000-0003-3778-207X

*Email for T.H.W.: thw@georgetown.edu; orcid.org/0000-0001-9217-8890

Present Addresses

†Department of Chemistry, University of California, Irvine, California 92697, United States.

Funding Sources

No competing financial interests have been declared.

ACKNOWLEDGMENT

S.C.E.S. acknowledges NSF CAREER (1847926), a CSUPERB New Investigator Grant, and NSF XSEDE (CHE160059, ACI-1548562). T.H.W. acknowledges funding from the NIH (R01GM126205) and the Georgetown Environment Initiative. P.N.P. was supported by a Louis Stokes Alliance for Minority Participation Research Fellowship. P.N.P. and C.E.G. were supported by NSF XSEDE EMPOWER. Use of Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, supported by DOE, Office of Science, BES (DE-AC02-76SF00515). The SSRL Structural Molecular Biology Program is supported by the DOE Office of Biological and Environmental Research, and by the National Institutes of Health, National Institute of General Medical Sciences (including P41GM103393). The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of NIGMS or NIH.

REFERENCES

- (1) Tocheva, E. I. Side-On Copper-Nitrosyl Coordination by Nitrite Reductase. *Science* **2004**, *304* (5672), 867–870. <https://doi.org/10.1126/science.1095109>.
- (2) Chakraborty, S.; Reed, J.; Sage, J. T.; Branagan, N. C.; Petrik, I. D.; Miner, K. D.; Hu, M. Y.; Zhao, J.; Alp, E. E.; Lu, Y. Recent Advances in Biosynthetic Modeling of Nitric Oxide Reductases and Insights Gained from Nuclear Resonance Vibrational and Other Spectroscopic Studies. *Inorg. Chem.* **2015**, *54* (19), 9317–9329. <https://doi.org/10.1021/acs.inorgchem.5b01105>.
- (3) Matsumoto, Y.; Tosha, T.; Pislakov, A. V.; Hino, T.; Sugimoto, H.; Nagano, S.; Sugita, Y.; Shiro, Y. Crystal Structure of Quinol-Dependent Nitric Oxide Reductase from *Geobacillus Stearothermophilus*. *Nat. Struct. Mol. Biol.* **2012**, *19* (2), 238–245. <https://doi.org/10.1038/nsmb.2213>.
- (4) Blomberg, M. R. A.; Siegbahn, P. E. M. Mechanism for N₂ O Generation in Bacterial Nitric Oxide Reductase: A Quantum Chemical Study. *Biochemistry* **2012**, *51* (25), 5173–5186. <https://doi.org/10.1021/bi300496e>.

ASSOCIATED CONTENT

- (5) Caranto, J. D.; Vilbert, A. C.; Lancaster, K. M. *Nitrosomonas Europaea* Cytochrome P460 Is a Direct Link between Nitrification and Nitrous Oxide Emission. *Proc. Natl. Acad. Sci.* **2016**, *113* (51), 14704–14709. <https://doi.org/10.1073/pnas.1611051113>.
- (6) Solomon, E. I.; Heppner, D. E.; Johnston, E. M.; Ginsbach, J. W.; Cirera, J.; Qayyum, M.; Kieber-Emmons, M. T.; Kjaergaard, C. H.; Hadt, R. G.; Tian, L. Copper Active Sites in Biology. *Chem. Rev.* **2014**, *114* (7), 3659–3853. <https://doi.org/10.1021/cr400327t>.
- (7) Tomson, N. C.; Williams, K. D.; Dai, X.; Sproules, S.; DeBeer, S.; Warren, T. H.; Wieghardt, K. Re-Evaluating the Cu K Pre-Edge XAS Transition in Complexes with Covalent Metal–Ligand Interactions. *Chem. Sci.* **2015**, *6* (4), 2474–2487. <https://doi.org/10.1039/C4SC03294B>.
- (8) Enemark, J. H.; Feltham, R. D. Principles of Structure, Bonding, and Reactivity for Metal Nitrosyl Complexes. *Coord. Chem. Rev.* **1974**, *13* (4), 339–406. [https://doi.org/10.1016/S0010-8545\(00\)80259-3](https://doi.org/10.1016/S0010-8545(00)80259-3).
- (9) Ampfeler, T.; Monsch, G.; Popp, J.; Riggemann, T.; Salvador, P.; Schröder, D.; Klüfers, P. Not Guilty on Every Count: The “Non-Innocent” Nitrosyl Ligand in the Framework of IUPAC’s Oxidation-State Formalism. *Angew. Chem. Int. Ed.* **2020**, anie.202003122. <https://doi.org/10.1002/anie.202003122>.
- (10) McCleverty, J. A. Chemistry of Nitric Oxide Relevant to Biology. *Chem. Rev.* **2004**, *104* (2), 403–418. <https://doi.org/10.1021/cr020623q>.
- (11) Teillet-Billy, D.; Fiquet-Fayard, F. The NO-3 $\uparrow$ Sigma. **1977**, *10* (4), L111–L117. <https://doi.org/10.1088/0022-3700/10/4/006>.
- (12) *Nitrosyl Complexes in Inorganic Chemistry, Biochemistry and Medicine II*; Mingos, D. M. P., Ed.; Structure and Bonding; Springer-Verlag: Berlin Heidelberg, 2014. <https://doi.org/10.1007/978-3-642-41160-1>.
- (13) Balcells, D.; Carbo, J. J.; Maseras, F.; Eisenstein, O. Self-Consistency versus “Best-Fit” Approaches in Understanding the Structure of Metal Nitrosyl Complexes. *Organometallics* **2004**, *23*, 6008–6014.
- (14) Kowalska, J. K.; Lima, F. A.; Pollock, C. J.; Rees, J. A.; DeBeer, S. A Practical Guide to High-Resolution X-Ray Spectroscopic Measurements and Their Applications in Bioinorganic Chemistry. *Isr. J. Chem.* **2016**, *56* (9–10), 803–815. <https://doi.org/10.1002/ijch.201600037>.
- (15) Bergmann, U.; Glatzel, P. X-Ray Emission Spectroscopy. *Photosynth. Res.* **2009**, *102* (2–3), 255–266. <https://doi.org/10.1007/s1120-009-9483-6>.
- (16) Glatzel, P.; Bergmann, U. High Resolution 1s Core Hole X-Ray Spectroscopy in 3d Transition Metal Complexes—Electronic and Structural Information. *Coord. Chem. Rev.* **2005**, *249* (1), 65–95. <https://doi.org/10.1016/j.ccr.2004.04.011>.
- (17) Eeckhout, S. G.; Safonova, O. V.; Smolentsev, G.; Biasioli, M.; Safonov, V. A.; Vykhodtseva, L. N.; Sikora, M.; Glatzel, P. Cr Local Environment by Valence-to-Core X-Ray Emission Spectroscopy. *J. Anal. At. Spectrom.* **2009**, *24* (2), 215–223. <https://doi.org/10.1039/B808345M>.
- (18) Smolentsev, G.; Soldatov, A. V.; Messinger, J.; Merz, K.; Weyhermüller, T.; Bergmann, U.; Pushkar, Y.; Yano, J.; Yachandra, V. K.; Glatzel, P. X-Ray Emission Spectroscopy To Study Ligand Valence Orbitals in Mn Coordination Complexes. *J. Am. Chem. Soc.* **2009**, *131* (36), 13161–13167. <https://doi.org/10.1021/ja808526m>.
- (19) Delgado-Jaime, M. U.; Dible, B. R.; Chiang, K. P.; Brennessel, W. W.; Bergmann, U.; Holland, P. L.; DeBeer, S. Identification of a Single Light Atom within a Multinuclear Metal Cluster Using Valence-to-Core X-Ray Emission Spectroscopy. *Inorg. Chem.* **2011**, *50* (21), 10709–10717. <https://doi.org/10.1021/ic201173j>.
- (20) Lancaster, K. M.; Roemelt, M.; Ettenhuber, P.; Hu, Y.; Ribbe, M. W.; Neese, F.; Bergmann, U.; DeBeer, S. X-Ray Emission Spectroscopy Evidences a Central Carbon in the Nitrogenase Iron-Molybdenum Cofactor. *Science* **2011**, *334* (6058), 974–977. <https://doi.org/10.1126/science.1206445>.
- (21) Martinie, R. J.; Blaesi, E. J.; Krebs, C.; Bollinger, J. M.; Silakov, A.; Pollock, C. J. Evidence for a Di- μ -Oxo Diamond Core in the Mn(IV)/Fe(IV) Activation Intermediate of Ribonucleotide Reductase from *Chlamydia Trachomatis*. *J. Am. Chem. Soc.* **2017**, *139* (5), 1950–1957. <https://doi.org/10.1021/jacs.6b11563>.
- (22) Martinie, R. J.; Blaesi, E. J.; Bollinger, J. M.; Krebs, C.; Finkelstein, K. D.; Pollock, C. J. Two-Color Valence-to-Core X-Ray Emission Spectroscopy Tracks Cofactor Protonation State in a Class I Ribonucleotide Reductase. *Angew. Chem. Int. Ed.* **2018**, *57* (39), 12754–12758. <https://doi.org/10.1002/anie.201807366>.
- (23) Martin-Diaconescu, V.; Chacón, K. N.; Delgado-Jaime, M. U.; Sokaras, D.; Weng, T.-C.; DeBeer, S.; Blackburn, N. J. K β Valence to Core X-Ray Emission Studies of Cu(I) Binding Proteins with Mixed Methionine – Histidine Coordination. Relevance to the Reactivity of the M- and H-Sites of Peptidylglycine Monooxygenase. *Inorg. Chem.* **2016**, *55* (7), 3431–3439. <https://doi.org/10.1021/acs.inorgchem.5b02842>.
- (24) Lim, H.; Baker, M. L.; Cowley, R. E.; Kim, S.; Bhadra, M.; Siegler, M. A.; Kroll, T.; Sokaras, D.; Weng, T.-C.; Biswas, D. R.; Dooley, D. M.; Karlin, K. D.; Hedman, B.; Hodgson, K. O.; Solomon, E. I. K β X-Ray Emission Spectroscopy as a Probe of Cu(I) Sites: Application to the Cu(I) Site in Preprocessed Galactose Oxidase. *Inorg. Chem.* **2020**, *59* (22), 16567–16581. <https://doi.org/10.1021/acs.inorgchem.0c02495>.
- (25) Pollock, C. J.; DeBeer, S. Insights into the Geometric and Electronic Structure of Transition Metal Centers from Valence-to-Core X-Ray Emission Spectroscopy. *Acc. Chem. Res.* **2015**, *48* (11), 2967–2975. <https://doi.org/10.1021/acs.accounts.5b00309>.
- (26) Stieber, S. C. E.; Milschmann, C.; Hoyt, J. M.; Turner, Z. R.; Finkelstein, K. D.; Wieghardt, K.; DeBeer, S.; Chirik, P. J. Bis(Imino)Pyridine Iron Dinitrogen Compounds Revisited: Differences in Electronic Structure Between Four- and Five-Coordinate Derivatives. *Inorg. Chem.* **2012**, *51* (6), 3770–3785. <https://doi.org/10.1021/ic202750n>.
- (27) Lu, T.-T.; Weng, T.-C.; Liaw, W.-F. X-Ray Emission Spectroscopy: A Spectroscopic Measure for the Determination of NO Oxidation States in Fe-NO Complexes. *Angew. Chem. Int. Ed.* **2014**, *53* (43), 11562–11566. <https://doi.org/10.1002/anie.201407603>.
- (28) Pollock, C. J.; Grubel, K.; Holland, P. L.; DeBeer, S. Experimentally Quantifying Small-Molecule Bond Activation Using Valence-to-Core X-Ray Emission Spectroscopy. *J. Am. Chem. Soc.* **2013**, *135* (32), 11803–11808. <https://doi.org/10.1021/ja3116247>.
- (29) Lassalle-Kaiser, B.; Boron, T. T.; Krewald, V.; Kern, J.; Beckwith, M. A.; Delgado-Jaime, M. U.; Schroeder, H.; Alonso-Mori, R.; Nordlund, D.; Weng, T.-C.; Sokaras, D.; Neese, F.; Bergmann, U.; Yachandra, V. K.; DeBeer, S.; Pecoraro, V. L.; Yano, J. Experimental and Computational X-Ray Emission Spectroscopy as a Direct Probe of Protonation States in Oxo-Bridged Mn^{IV} Dimers Relevant to Redox-Active Metalloproteins. *Inorg. Chem.* **2013**, *52* (22), 12915–12922. <https://doi.org/10.1021/ic400821g>.
- (30) Cutsail, G. E.; Gagnon, N. L.; Spaeth, A. D.; Tolman, W. B.; DeBeer, S. Valence-to-Core X-Ray Emission Spectroscopy as a Probe of O–O Bond Activation in Cu₂O₂ Complexes. *Angew. Chem. Int. Ed.* **2019**, *58* (27), 9114–9119. <https://doi.org/10.1002/anie.201903749>.
- (31) Hugenbruch, S.; Shafaat, H. S.; Krämer, T.; Delgado-Jaime, M. U.; Weber, K.; Neese, F.; Lubitz, W.; DeBeer, S. In Search of Metal Hydrides: An X-Ray Absorption and Emission Study of [NiFe] Hydrogenase Model Complexes. *Phys. Chem. Chem. Phys.* **2016**, *18* (16), 10688–10699. <https://doi.org/10.1039/C5CP07293J>.
- (32) Burkhardt, L.; Holzwarth, M.; Plietker, B.; Bauer, M. Detection and Characterization of Hydride Ligands in Iron Complexes by High-Resolution Hard X-Ray Spectroscopy and Implications for Catalytic Processes. *Inorg. Chem.* **2017**, *56* (21), 13300–13310. <https://doi.org/10.1021/acs.inorgchem.7b02063>.
- (33) Kundu, S.; Phu, P. N.; Ghosh, P.; Kozimor, S. A.; Bertke, J. A.; Stieber, S. C. E.; Warren, T. H. Nitrosyl Linkage Isomers: NO Coupling to N₂O at a Mononuclear Site. *J. Am. Chem. Soc.* **2019**, *141* (4), 1415–1419. <https://doi.org/10.1021/jacs.8b09769>.
- (34) Kowalska, J. K.; Hahn, A. W.; Albers, A.; Schiewer, C. E.; Björnsson, R.; Lima, F. A.; Meyer, F.; DeBeer, S. X-Ray Absorption and Emission Spectroscopic Studies of [L₂FeS₂]ⁿ Model Complexes: Implications for the Experimental Evaluation of Redox States

in Iron–Sulfur Clusters. *Inorg. Chem.* **2016**, *55* (9), 4485–4497. <https://doi.org/10.1021/acs.inorgchem.6b00295>.

(35) Neese, F. *Orca: An Ab Initio, DFT and Semiempirical Electronic Structure Package*; Max Planck Institute for Chemical Energy Conversion: Mülheim an der Ruhr, Germany.

(36) MacMillan, S. N.; Walroth, R. C.; Perry, D. M.; Morsing, T. J.; Lancaster, K. M. Ligand-Sensitive But Not Ligand-Diagnostic: Evaluating Cr Valence-to-Core X-Ray Emission Spectroscopy as a Probe of Inner-Sphere Coordination. *Inorg. Chem.* **2015**, *54* (1), 205–214. <https://doi.org/10.1021/ic502152r>.

(37) Krewald, V.; Lassalle-Kaiser, B.; Boron, T. T.; Pollock, C. J.; Kern, J.; Beckwith, M. A.; Yachandra, V. K.; Pecoraro, V. L.; Yano, J.; Neese, F.; DeBeer, S. The Protonation States of Oxo-Bridged Mn^{IV} Dimers Resolved by Experimental and Computational Mn K-Pre-Edge X-Ray Absorption Spectroscopy. *Inorg. Chem.* **2013**, *52* (22), 12904–12914. <https://doi.org/10.1021/ic4008203>.

(38) Pollock, C. J.; Lancaster, K. M.; Finkelstein, K. D.; DeBeer, S. Study of Iron Dimers Reveals Angular Dependence of Valence-to-Core X-Ray Emission Spectra. *Inorg. Chem.* **2014**, *53* (19), 10378–10385. <https://doi.org/10.1021/ic501462y>.

(39) Lee, N.; Petrenko, T.; Bergmann, U.; Neese, F.; DeBeer, S. Probing Valence Orbital Composition with Iron K β X-Ray Emission Spectroscopy. *J. Am. Chem. Soc.* **2010**, *132* (28), 9715–9727. <https://doi.org/10.1021/ja101281e>.

(40) Schwalenstocker, K.; Paudel, J.; Kohn, A. W.; Dong, C.; Van, K. M.; Farquhar, E. R.; Li, F. Cobalt K β Valence-to-Core X-Ray Emission Spectroscopy: A Study of Low-Spin Octahedral Cobalt(III) Complexes. *Dalton Trans.* **14**.

(41) Kirchner, B.; Wennmohs, F.; Ye, S.; Neese, F. CP(PPP). *Curr Opin Chem Biol* **2007**, *11* (134).

(42) Schaefer, A.; Horn, H.; Ahlrichs, R. TZVP. *J Chem Phys* **1992**, *97*, 2571.

(43) Rees, J. A.; Martin-Diaconescu, V.; Kovacs, J. A.; DeBeer, S. X-Ray Absorption and Emission Study of Dioxxygen Activation by a Small-Molecule Manganese Complex. *Inorg. Chem.* **2015**, *54* (13), 6410–6422. <https://doi.org/10.1021/acs.inorgchem.5b00699>.

(44) Delgado-Jaime, M. U.; DeBeer, S.; Bauer, M. Valence-to-Core X-Ray Emission Spectroscopy of Iron-Carbonyl Complexes: Implications for the Examination of Catalytic Intermediates. *Chem. - Eur. J.* **2013**, *19* (47), 15888–15897. <https://doi.org/10.1002/chem.201301913>.

(45) Bergmann, U.; Horne, C. R.; Collins, T. J.; Workman, J. M.; Cramer, S. P. Chemical Dependence of Interatomic X-Ray Transition Energies and Intensities – a Study of Mn K β '' and K β 2,5 Spectra. *Chem. Phys. Lett.* **1999**, *302* (1), 119–124. [https://doi.org/10.1016/S0009-2614\(99\)00095-0](https://doi.org/10.1016/S0009-2614(99)00095-0).

(46) Pollock, C. J.; DeBeer, S. Valence-to-Core X-Ray Emission Spectroscopy: A Sensitive Probe of the Nature of a Bound Ligand. *J. Am. Chem. Soc.* **2011**, *133* (14), 5594–5601. <https://doi.org/10.1021/ja200560z>.

(47) Szlachetko, J.; Nachtegaal, M.; de Boni, E.; Willmann, M.; Safonova, O.; Sa, J.; Smolentsev, G.; Szlachetko, M.; van Bokhoven, J. A.; Dousse, J.-Cl.; Hoszowska, J.; Kayser, Y.; Jagodzinski, P.; Bergamaschi, A.; Schmitt, B.; David, C.; Lücke, A. A von Hamos X-Ray Spectrometer Based on a Segmented-Type Diffraction Crystal for Single-Shot x-Ray Emission Spectroscopy and Time-Resolved Resonant Inelastic x-Ray Scattering Studies. *Rev. Sci. Instrum.* **2012**, *83* (10), 103105. <https://doi.org/10.1063/1.4756691>.

(48) Alonso-Mori, R.; Kern, J.; Gildea, R. J.; Sokaras, D.; Weng, T.-C.; Lassalle-Kaiser, B.; Tran, R.; Hattne, J.; Laksmono, H.; Hellmich, J.; Glöckner, C.; Echols, N.; Sierra, R. G.; Schafer, D. W.; Sellberg, J.; Kenney, C.; Herbst, R.; Pines, J.; Hart, P.; Herrmann, S.; Grosse-Kunstleve, R. W.; Latimer, M. J.; Fry, A. R.; Messerschmidt, M. M.; Miahnahri, A.; Seibert, M. M.; Zwart, P. H.; White, W. E.; Adams, P. D.; Bogan, M. J.; Boutet, S.; Williams, G. J.; Zouni, A.; Messinger, J.; Glatzel, P.; Sauter, N. K.; Yachandra, V. K.; Yano, J.; Bergmann, U. Energy-Dispersive X-Ray Emission Spectroscopy Using an X-Ray Free-Electron Laser in a Shot-by-Shot Mode. *Proc. Natl. Acad. Sci.* **2012**, *109* (47), 19103–19107. <https://doi.org/10.1073/pnas.1211384109>.

(49) Davis, K. M.; Mattern, B. A.; Pacold, J. I.; Zakharova, T.; Brewé, D.; Kosheleva, I.; Henning, R. W.; Graber, T. J.; Heald, S. M.; Seidler, G. T.; Pushkar, Y. Fast Detection Allowing Analysis of Metalloprotein Electronic Structure by X-Ray Emission Spectroscopy at Room Temperature. *J. Phys. Chem. Lett.* **2012**, *3* (14), 1858–1864. <https://doi.org/10.1021/jz3006223>.

(50) March, A. M.; Assefa, T. A.; Boemer, C.; Bressler, C.; Britz, A.; Diez, M.; Doumy, G.; Galler, A.; Harder, M.; Khakhulin, D.; Németh, Z.; Pápai, M.; Schulz, S.; Southworth, S. H.; Yavaş, H.; Young, L.; Gawelda, W.; Vankó, G. Probing Transient Valence Orbital Changes with Picosecond Valence-to-Core X-Ray Emission Spectroscopy. *J. Phys. Chem. C* **2017**, *121* (5), 2620–2626. <https://doi.org/10.1021/acs.jpcc.6b12940>.

(51) Sokaras, D.; Weng, T.-C.; Nordlund, D.; Alonso-Mori, R.; Velikov, P.; Wenger, D.; Garachtchenko, A.; George, M.; Borzenets, V.; Johnson, B.; Rabedeau, T.; Bergmann, U. A Seven-Crystal Johann-Type Hard x-Ray Spectrometer at the Stanford Synchrotron Radiation Lightsource. *Rev. Sci. Instrum.* **2013**, *84* (5), 053102. <https://doi.org/10.1063/1.4803669>.

(52) Solé, V. A.; Papillon, E.; Cotte, M.; Walter, Ph.; Susini, J. A Multiparameter Code for the Analysis of Energy-Dispersive X-Ray Fluorescence Spectra. *Spectrochim. Acta Part B At. Spectrosc.* **2007**, *62* (1), 63–68. <https://doi.org/10.1016/j.sab.2006.12.002>.

(53) Towns, J.; Cockerill, T.; Dahan, M.; Foster, I.; Gaither, K.; Grimshaw, A.; Hazlewood, V.; Lathrop, S.; Lifka, D.; Peterson, G. D.; Roskies, R.; Scott, J. R.; Wilkins-Diehr, N. XSEDE: Accelerating Scientific Discovery. *Comput. Sci. Eng.* **2014**, *16* (5), 62–74. <https://doi.org/10.1109/MCSE.2014.80>.

(54) Hanwell, M. D.; Curtis, D. E.; Lonie, D. C.; Vandermeersch, T.; Zurek, E.; Hutchison, G. R. Avogadro: An Advanced Semantic Chemical Editor, Visualization, and Analysis Platform. *J. Cheminformatics* **2012**, *4* (1), 17. <https://doi.org/10.1186/1758-2946-4-17>.

(55) Perdew, J. P. Density-Functional Approximation for the Correlation Energy of the Inhomogeneous Electron Gas. *Phys. Rev. B* **1986**, *33* (12), 8822–8824. <https://doi.org/10.1103/PhysRevB.33.8822>.

(56) Perdew, J. P. Erratum: Density-Functional Approximation for the Correlation Energy of the Inhomogeneous Electron Gas. *Phys. Rev. B* **1986**, *34* (10), 7406–7406. <https://doi.org/10.1103/PhysRevB.34.7406>.

(57) Lee, C.; Yang, W.; Parr, R. G. Development of the Colle-Salvetti Correlation-Energy Formula into a Functional of the Electron Density. *Phys. Rev. B* **1988**, *37* (2), 785–789. <https://doi.org/10.1103/PhysRevB.37.785>.

(58) Neese, F.; Solomon, E. I. Interpretation and Calculation of Spin-Hamiltonian Parameters in Transition Metal Complexes. In *Magnetism: Molecules to Materials IV*; John Wiley & Sons, Ltd, 2003; pp 345–466. <https://doi.org/10.1002/3527600698.ch9>.

(59) Schäfer, A.; Horn, H.; Ahlrichs, R. Fully Optimized Contracted Gaussian Basis Sets for Atoms Li to Kr. *J. Chem. Phys.* **1992**, *97* (4), 2571–2577. <https://doi.org/10.1063/1.463096>.

(60) Schäfer, A.; Huber, C.; Ahlrichs, R. Fully Optimized Contracted Gaussian Basis Sets of Triple Zeta Valence Quality for Atoms Li to Kr. *J. Chem. Phys.* **1994**, *100* (8), 5829–5835. <https://doi.org/10.1063/1.467146>.

(61) Weigend, F.; Ahlrichs, R. Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy. *Phys. Chem. Chem. Phys.* **2005**, *7* (18), 3297–3305. <https://doi.org/10.1039/B508541A>.

(62) Eichkorn, K.; Treutler, O.; Öhm, H.; Häser, M.; Ahlrichs, R. Auxiliary Basis Sets to Approximate Coulomb Potentials. *Chem. Phys. Lett.* **1995**, *240* (4), 283–290. [https://doi.org/10.1016/0009-2614\(95\)00621-A](https://doi.org/10.1016/0009-2614(95)00621-A).


(63) Eichkorn, K.; Weigend, F.; Treutler, O.; Ahlrichs, R. Auxiliary Basis Sets for Main Row Atoms and Transition Metals and Their Use to Approximate Coulomb Potentials. *Theor. Chem. Acc.* **1997**, *97* (1), 119–124. <https://doi.org/10.1007/s002140050244>.

(64) Ginsberg, A. P. Magnetic Exchange in Transition Metal Complexes. 12. Calculation of Cluster Exchange Coupling Constants

with the X.Alpha.-Scattered Wave Method. *J. Am. Chem. Soc.* **1980**, *102* (1), 111–117. <https://doi.org/10.1021/ja00521a020>.

(65) Noodleman, L.; Peng, C. Y.; Case, D. A.; Mouesca, J.-M. Orbital Interactions, Electron Delocalization and Spin Coupling in Iron-Sulfur Clusters. *Coord. Chem. Rev.* **1995**, *144*, 199–244. [https://doi.org/10.1016/0010-8545\(95\)07011-L](https://doi.org/10.1016/0010-8545(95)07011-L).

(66) Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E. C.; Ferrin, T. E. Chimera. *J Comput Chem* **2004**, *13*, 1605.



Synopsis: β -diketiminate Ni-NO complexes studied by X-ray emission spectroscopy (XES) demonstrate that XES can directly probe and distinguish end-on versus side-on NO coordination modes, as well as 1-electron NO reduction. This work shows the capability of XES to distinguish NO coordination modes and oxidation states at Ni and highlights applications in quantifying small molecule activation in enzymes.
