

Astrophysical and Theoretical Physics Implications from Multimessenger Neutron Star Observations

Hector O. Silva^{1,2}, A. Miguel Holgado^{3,4,5}, Alejandro Cárdenas-Avendaño^{2,6}, and Nicolás Yunes²

¹Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, D-14476 Potsdam, Germany

²Department of Physics and Illinois Center for Advanced Studies of the Universe, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA

³Department of Physics and McWilliams Center for Cosmology, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

⁴Department of Astronomy, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA

⁵National Center for Supercomputing Applications, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA

⁶Programa de Matemática, Fundación Universitaria Konrad Lorenz, 110231 Bogotá, Colombia



(Received 24 December 2020; accepted 18 March 2021; published 3 May 2021)

The Neutron Star Interior Composition Explorer (NICER) recently measured the mass and equatorial radius of the isolated neutron star PSR J0030+0451. We use these measurements to infer the moment of inertia, the quadrupole moment, and the surface eccentricity of an isolated neutron star for the first time, using relations between these quantities that are insensitive to the unknown equation of state of supranuclear matter. We also use these results to forecast the moment of inertia of neutron star A in the double pulsar binary J0737-3039, a quantity anticipated to be directly measured in the coming decade with radio observations. Combining this information with the measurement of the tidal Love number with LIGO/Virgo observations, we propose and implement the first theory-agnostic and equation-of-state-insensitive test of general relativity. Specializing these constraints to a particular modified theory, we find that consistency with general relativity places the most stringent constraint on gravitational parity violation to date, surpassing all other previously reported bounds by 7 orders of magnitude and opens the path for a future test of general relativity with multimessenger neutron star observations.

DOI: [10.1103/PhysRevLett.126.181101](https://doi.org/10.1103/PhysRevLett.126.181101)

Introduction.—Neutron stars are some of the most extreme objects in nature. Their mass (typically around $1.4 M_{\odot}$) combined with their small radius (between 10–14 km) result in interior densities that can exceed nuclear saturation density ($\rho \geq 2.8 \times 10^{14}$ g/cm³), above which exotic states of matter can arise [1]. Neutron stars are, next to black holes, the strongest gravitational field sources known, with typical gravitational potentials that are 5 orders of magnitude larger than that of the Sun. These properties make neutron stars outstanding laboratories to study both matter and gravity in situations out of reach in terrestrial and Solar System experiments.

Our current poor understanding of the supranuclear equation of state translates, via the equations of stellar equilibrium, to a large variability on observable properties of neutron stars, such as their masses and radii [2]. This variability increases if one lifts the assumption that Einstein's theory of general relativity is valid in the strong-gravity environment of neutron star interiors [3]. Modifications to general relativity generically predict new equations of stellar equilibrium, which, when combined with uncertainties on the nuclear equation of state, jeopardize attempts to test Einstein's theory with isolated, neutron star observations.

One possibility to circumvent this issue is to explore whether relations between neutron-star observables that are

insensitive to either (or both) the equation of state and the gravitational theory exist. Fortunately, they do. For instance, when properly nondimensionalized, the moment of inertia (I), the rotational quadrupole moment (Q), and the tidal Love number (λ) of neutron stars show a remarkable degree of equation-of-state insensitivity, at a level below 1% [4,5]. These “I-Love-Q” relations also exist in some modified theories of gravity, although they are different from their general relativity counterparts [6].

We here combine the first measurements [7,8] by NICER [9] of *both* the mass (M) and equatorial radius (R_e) of the isolated pulsar PSR J0030+0451 [10,11] with known equation-of-state insensitive relations involving the compactness $C = GM/(R_e c^2)$ (see, for instance, Refs. [12–15]) to infer a number of astrophysical and theoretical physics consequences. Before doing so, let us explain how these relations are obtained.

Quasiuniversal relations.—Neutron stars can have short rotation periods of the order of milliseconds, so their surfaces are oblate instead of spherical. The inclusion of this effect is of critical importance to accurately model the thermal x-ray waveform that NICER observes, since the x rays are produced by hotspots at the star's surface [16,17]. The canonical approach to model relativistic rotating stars was developed in the 1970s [18,19]. In this approach, the

star's rotation is treated as a small perturbation $\varepsilon = f/f_0 \ll 1$, involving the star's rotation frequency f and its characteristic mass-shedding frequency $f_0 = (GM/R_e^3)^{1/2}/(2\pi)$. Rotating stars are then found by perturbing in ε an otherwise nonrotating star, which can be obtained by solving the Tolman-Oppenheimer-Volkoff equations [20]. This slow-rotation approximation is well justified for most neutron stars with astrophysically relevant spins. Even for a prototypical millisecond pulsar with $f = 700$ Hz, $M = 1.4 M_\odot$, and $R_e = 11$ km, one has $\varepsilon = 0.37$. In the case of PSR J0030+0451, its rotation frequency is known to be $f_\star = 205.53$ Hz [10,11], so $\varepsilon_\star = 0.14$, when one uses the best-fit M and R_e values obtained by NICER [7,8]. Henceforth, a “ $\star$ ” indicates observables associated with PSR J0030+0451.

Using this technique, we numerically calculated over a thousand neutron star solutions to order ε^2 in this perturbative scheme, using a broad set of 46 different equations of state [21,22], as detailed in the Supplemental Material [23]. From these solutions, we then numerically computed the moment of inertia I , the rotational quadrupole moment Q , the surface eccentricity e , and the electric-type, $\ell = 2$, tidal Love number λ , which is the dominant parameter in the description of tidal effects in the late inspiral of neutron star binaries [41–43]. We nondimensionalized these quantities through division by the appropriate factors of M and dimensionless spin $\chi = (2\pi f_0) \tilde{G} \tilde{M} / c^3$, namely: $\tilde{I} = c^4 I / (G^2 M^3)$, $\tilde{Q} = -c^4 Q / (G^2 M^3 \chi^2)$ and $\tilde{\lambda} = c^{10} \lambda / (GM)^5$. The surface eccentricity e is dimensionless by definition, given in terms of the star's equatorial R_e and polar R_p radii as $e = [(R_e/R_p)^2 - 1]^{1/2}$ [14]. The relations between these nondimensional quantities are strongly insensitive to the equation of state. Because of the small value of $\varepsilon_\star$, we can neglect higher order in spin corrections in this expression.

The first step in using the approximately universal relations on NICER's first observation is to derive equation-of-state-insensitive relations between the observables $\{\tilde{I}, \tilde{Q}, \tilde{\lambda}, e\}$, with respect to the compactness $\mathcal{C}$. Details of these “ $\mathcal{C}$ relations” are given in the Supplemental Material [23]. Our plan of attack is then clear: use the publicly available Markov chain Monte Carlo (MCMC) M - R_e samples [44,45] for the best-fit model inferred by two independent analysis [7,8] of the NICER data [46]. Although each group modeled the surface hotspots differently and used different sampling methods, their results are consistent with each other. Here we use the results for the three-hotspot model inferred by Miller *et al.* [8] and the favored single temperature, two-hotspot ST + PST model from Riley *et al.* [7] to obtain a posterior distribution for the compactness, and then use the approximately universal relations to infer other astrophysical quantities. We detail this procedure next.

Astrophysical implications.—We begin by constructing a posterior distribution $P(\mathcal{C}|\text{NICER})$ for the compactness $\mathcal{C}$

of PSR J0030+0451, using the MCMC chains [44,45]. With this posterior in hand, we then use the $\mathcal{C}$ relations to infer posterior distributions for $\{\tilde{I}, \tilde{\lambda}, \tilde{Q}, e\}$.

The implementation of such an inference procedure requires a particular scheme, and we here follow a proposal that accounts for the approximately universal nature of the relations [22]. In this scheme, the maximum relative error of each fitting function defines the half width of the 90% credible interval of a Gaussian distribution centered at each fitted value. The posterior distribution for each dimensionless quantity is then calculated using the corresponding $\mathcal{C}$ relation and the posterior distribution of the compactness, after marginalizing over the latter. From these posteriors and using the same procedure described above, we can also construct posteriors for the dimensional versions of these quantities by a change of variables, marginalize over the nuisance variables mass M and radius R_e , and then do a final rescaling of the posterior by ε ($= 0.14$) for the surface eccentricity e and by ε^2 for the rotational quadrupole moment Q . We refer to the Supplemental Material for details [23].

The resulting mean and 1σ intervals of these parameters (both the nondimensionalized and the dimensional versions) are shown in Table I; see the Supplemental Material [23] for plots of the inferred posteriors. The reported confidence intervals in all of these quantities account for both the approximate nature of the universal relations and the uncertainties in NICER's observation. These results are the first inferences on the moment of inertia, the surface eccentricity, the Love number, and the quadrupole moment of an isolated neutron star.

We can also use NICER's observation combined with the I-C relation to estimate the moment of inertia of PSR J0737-3039A ($I_{1.3381}$), where the subscript refers to this pulsar's measured mass of $M = (1.3381 \pm 0.0007) M_\odot$ [47]. The double pulsar J0737-3039 is expected to provide

TABLE I. Inferred properties of PSR J0030+0451 using equation-of-state-insensitive relations combined with the MCMC samples by Miller *et al.* [44] and Riley *et al.* [45]. We report the values within 1 standard deviation from the mean, representing approximately 68% confidence intervals. These values are the first inferences of the moment of inertia, the eccentricity, the Love number, and the quadrupole moment of an isolated neutron star.

Parameter	Miller <i>et al.</i>	Riley <i>et al.</i>
$\tilde{I}_\star$ (10)	$1.31^{+0.13}_{-0.11}$	$1.42^{+0.26}_{-0.19}$
$\tilde{\lambda}_\star$ (10^2)	$4.97^{+1.92}_{-1.28}$	$6.75^{+5.52}_{-2.69}$
$\tilde{Q}_\star$	$5.92^{+0.73}_{-0.61}$	$6.50^{+1.38}_{-1.03}$
$I_\star$ (10^{45} g cm ²)	$1.71^{+0.64}_{-0.48}$	$1.42^{+0.81}_{-0.53}$
$Q_\star$ (10^{43} g cm ²)	$1.49^{+0.63}_{-0.45}$	$1.27^{+0.74}_{-0.49}$
$e_\star$ (10^{-1})	$1.56^{+0.25}_{-0.21}$	$1.58^{+0.29}_{-0.28}$

the first direct neutron star measurement of the moment of inertia [48]. This system is the only double-pulsar observed to date, which makes it an unique laboratory for binary stellar astrophysics [49,50]. Moreover, an accurate measurement of $I_{1.3381}$ in combination with its known mass is expected to strongly constrain the nuclear equation of state around once and twice nuclear saturation density [51].

To predict the moment of inertia of PSR J0737-3039A from NICER's observation of PSR J0030+0451, we first need to obtain an estimate for the compactness $\mathcal{C}_{1.3381}$ of PSR J0737-3039A. This can be approximated by the substitution $\{M, R_e\} \mapsto \{M_0 = 1.3381 M_\odot, R_e\}$ at each MCMC sample [44] and then computing $\mathcal{C}_{M_0}$. This yields an approximation to the distribution of compactness for a system with mass M_0 , which is assumed known and identical to PSR J0030+0451. This procedure is only justified as long as M_0 is very close to $M_\star$, as in the case of PSR J0737-3039A, whose inferred mass ($M_0 = 1.3381^{+0.0007}_{-0.0007} M_\odot$) [47] is within the 1σ credible interval of both NICER's mass inference ($M_\star = 1.34^{+0.15}_{-0.16} M_\odot$ [7] and $M_\star = 1.44^{+0.15}_{-0.14} M_\odot$ [8]).

With an estimate of the compactness of PSR J0737-3039A, we can now obtain a prediction for PSR J0030+0451's moment of inertia repeating the procedure applied to PSR J0030+0451. Figure 1 shows our result using both NICER MCMC samples; $I_{1.3381}^{\text{Miller et al.}} = 1.64^{+0.52}_{-0.37} \times 10^{45} \text{ g cm}^2$, and $I_{1.3381}^{\text{Riley et al.}} = 1.68^{+0.53}_{-0.48} \times 10^{45} \text{ g cm}^2$, together with two other independent predictions [52,53]. All predictions are consistent with one another. The anticipated future independent measurement of $I_{1.3381}$ from continued radio timing of




FIG. 1. Predictions for the moment of inertia of PSR J0737-3039A. We compare our predicted $I_{1.3381}$ using both the MCMC samples from Miller *et al.* [44] and Riley *et al.* [45] against: (i) Landry and Kumar [52] (LK18), which used binary Love [54] and I-Love relations with the tidal-deformability constraints from binary neutron-star merger GW170817 [55], and (ii) Lim *et al.* [53] (LHS19) which carried out Bayesian modeling of a number of equations of state. The larger moment of inertia that we predict is due to the larger radii favored by an $M \approx 1.4 M_\odot$ neutron star by NICER's observation relative what is inferred by the two other methods, as $I \sim MR_e^2$.

PSR J0737-3039A will provide another test for nuclear theory and enable an I-Love test of gravity, the latter of which we define next.

Theoretical physics implications.—The combination of the inference of I with NICER data described above, and the independent measurement of λ [55] by the LIGO/Virgo Collaboration from the binary neutron-star merger GW170817 [56], allows for the first implementation of an I-Love test [4]. This test would be the first multimessenger test of general relativity with neutron star observables.

The idea of an I-Love test is as follows [4,5] (see Fig. 2). Consider two independent inferences of $\bar{I}_{1.4}$ and $\bar{\lambda}_{1.4}$ for a $1.4 M_\odot$ neutron star. In the $(\bar{I}, \bar{\lambda})$ plane, these measurements yield a 90% confidence error box. If the I-Love relation in general relativity, including its small equation-of-state variability, *does not* pass through this error box, then there is evidence for a violation of Einstein's theory, regardless of the underlying equation of state. Moreover, if any theory of gravity predicts an I-Love curve that also does not pass through this error box for a given value of its coupling constants, then the I-Love test places a constraint on the couplings of this theory, which is also independent of the equation of state.

Such a test, however, requires the inference of the tidal deformability and the moment of inertia of a neutron star of the *same* mass. The LIGO/Virgo Collaboration used gravitational wave data to infer the tidal deformability of a $1.4 M_\odot$ neutron star to be $\bar{\lambda}_{1.4} = 190^{+390}_{-120}$ at 90%




FIG. 2. Multimessenger test of general relativity using the parametrized I-Love relation. The vertical (horizontal) lines delimit the 90% confidence region (shaded) for $\bar{\lambda}_{1.4}$ [57] ($\bar{I}_{1.4}$, this work), while the circle marks the median (190, 14.6). The solid black line corresponds to the I-Love relation in general relativity [Eq. (1)] and is consistent with the inferred values of $\bar{I}_{1.4}$, $\bar{\lambda}_{1.4}$ at 90% confidence. Starting from $b = -2$ and moving clockwise, we show the parametrized I-Love curves (b, β_{crit}), where $b \in [-2, 5]$ and β_{crit} is the critical value of β above which the parametrized I-Love relation [Eq. (2)] fails to pass by the 90% confidence region in the plane. Here we used the value of $\bar{I}_{1.4}$ inferred using the results by Miller *et al.* [8,44]. We found similar results using the results by Riley *et al.* [7,45] (see Supplemental Material [23]).

confidence [57], obtained under the assumptions that the binary components were described by the same equation of state and were slowly spinning. We can use NICER's data to infer the moment of inertia of a $1.4 M_\odot$ neutron star with the same techniques we used to predict the moment of inertia of PSR J0737-3039A. For concreteness, we use the results from Miller *et al.* [8,44], but we verified (see the Supplemental Material [23] for detail) that our conclusions are essentially the same had we used the results from Riley *et al.* [7,45]. We find that $C_{1.4} = 0.159^{+0.025}_{-0.022}$ and $\bar{I}_{1.4} = 14.6^{+4.5}_{-3.3}$ at 90% confidence. An important underlying assumption behind both inferences is that general relativity is the correct theory of gravity. The rationale behind this test is detailed in the Supplemental Material [23].

Since carrying out such a test on a theory-by-theory basis would, in general, be complicated and time consuming, we here develop and implement a useful *parametrization* of the I-Love test. From Newtonian gravity, we know that $\bar{I}$ scales with $\mathcal{C}^{-2}$, whereas $\bar{\lambda}$ scales with $\mathcal{C}^{-5}$. Therefore, $\bar{I} = C_{\bar{I}\bar{\lambda}} \bar{\lambda}^{2/5}$, with $C_{\bar{I}\bar{\lambda}} \approx 0.52$ a constant that depends on the equation of state very weakly [5]. This calculation can be extended, systematically, in a post-Minkowskian expansion, i.e., an expansion in powers of $\mathcal{C} \ll 1$ [58]. The outcome is that both $\bar{I}$ and $\bar{\lambda}$ can be written as a power series in $\mathcal{C}$ and then be combined (as just described in the Newtonian limit) to obtain $\bar{I} = \bar{I}(\bar{\lambda})$. The resulting I-Love relation has the same degree of equation-of-state independence as the original I-Love relation [4]. For our neutron star catalog, a parametrization in general relativity of the form

$$\bar{I}_{\text{GR}} = \bar{\lambda}^{2/5} (c_0 + c_1 \bar{\lambda}^{-1/5} + c_2 \bar{\lambda}^{-2/5}), \quad (1)$$

with $c_0 = 0.584$, $c_1 = 0.980$, $c_2 = 2.695$, is sufficient to reproduce our numerical data with mean relative error $\langle \epsilon^{\bar{I}} \rangle \leq 2 \times 10^{-3}$. The prefactor $\bar{\lambda}^{2/5}$ is the Newtonian result, while the powers of $\bar{\lambda}^{-1/5}$ inside parenthesis are relativistic (post-Minkowskian) corrections because $\bar{\lambda}^{-1/5} \propto \mathcal{C} \lesssim 0.2$. Given this, we then propose a minimal deformation of the Einsteinian parametrization in Eq. (1) of the form

$$\bar{I}_p = \bar{I}_{\text{GR}} + \beta \bar{\lambda}^{-b/5}, \quad \beta \in \mathbb{R}_+, \quad b \in \mathbb{Z}, \quad (2)$$

where β and b are deformation parameters that control the magnitude and type of the deviations from general relativity in the I-Love relation, respectively. Such a parametrization is similar to that successfully used in gravitational-wave tests of general relativity by the LIGO/Virgo Collaboration, the parametrized post-Einsteinian framework [59].

We performed such a test of general relativity through the procedure described earlier. First, we see that the I-Love relation in general relativity does indeed pass this null-test and it is consistent with the error box. Second, we

considered $b \in [-2, 5]$, where the lower limit is set by requiring no deviations at the Newtonian level and the upper limit is set for simplicity. We then fixed b and calculated what the corresponding value of $\beta = \beta_{\text{crit}}$ is, above which the parametrized I-Love relation (2) would be in tension with the inferred $(\bar{I}_{1.4}, \bar{\lambda}_{1.4})$ region at 90% confidence. Our results are summarized in Fig. 2, where the numbers in parenthesis correspond to (b, β_{crit}) . We stress that our results for $b \leq 0$ are of course dependent on the posterior used for $\bar{\lambda}_{1.4}$. If one treated the tidal deformabilities as independent free parameters in the waveform model [56], then the $\bar{\lambda}_{1.4}$ posterior would not have a lower limit, allowing all curves with $b \leq 0$ to be consistent with both observations.

With these theory-agnostic constraints in hand, we can now map them to specific theories and place constraints on their coupling parameters. As an example, let us consider dynamical Chern-Simons gravity, a theory that modifies general relativity by introducing gravitational parity violation [60]. This theory has found applications to several open problems in cosmology, such as the matter-antimatter asymmetry and leptogenesis [61–64]. It also arises in several approaches to quantum gravity, such as string theory [65] and loop quantum gravity [66–68]. Mathematical well-posedness requires the theory to be treated as an effective field theory [69]. In this formalism, one works in a small-coupling approximation $\zeta \equiv 16\pi\alpha^2 \mathcal{R}^{-4} \ll 1$, where $\mathcal{R} = [c^2 R_e^3 / (GM)]^{1/2}$ is the curvature length scale associated with a neutron star (in our case), and where α is a coupling constant with units of length squared, such that ζ is dimensionless. This theory modifies Einstein's only when gravity is strong, and thus, it passes all Solar System constraints, being only extremely weakly constrained by Gravity Probe B and the LAGEOS satellites, and table-top experiments, to $\alpha^{1/2} \leq 10^8$ km [70–72]. This theory has also evaded gravitational-wave tests [73], making it a key target to test the constraining power of our new I-Love test.

Let us now map the theory-agnostic deformation of the I-Love relations in Eq. (2) to dynamical Chern-Simons gravity, though this methodology could be applied to other theories as well. As we discuss in the Supplemental Material [23], the I-Love relation in this theory can be described by Eq. (2) with $b_{\text{CS}} = 4$ and $\beta_{\text{CS}} = 6.15 \times 10^{-2} \bar{\xi}$, where $\bar{\xi} = 16\pi\alpha^2 / M^4$. We can now use our theory-agnostic constraints on β to place a constraint on α , the coupling constant of dynamical Chern-Simons gravity. Using the constraint on β when $b = 4$, namely, $\beta_{\text{crit}} \leq 8.84 \times 10^2$, and applying the mapping, yields $\beta_{\text{CS}} = 6.15 \times 10^{-2} \bar{\xi} \leq 8.84 \times 10^2$, or simply

$$\alpha^{1/2} \leq 8.5 \text{ km}, \quad (3)$$

at 90% credibility, if the theory is to be consistent with the observational bounds on $\bar{I}_{1.4}$ and $\bar{\lambda}_{1.4}$. Using the mean value

$C_{1,4} = 0.159$, which implies the mean equatorial radius $R_{1,4} = 13.0$ km, we also find that $\zeta \leq 0.23$ when using Eq. (3), implying that the small-coupling approximation is indeed satisfied. This bound is 7 orders of magnitude stronger than any previous constraints and it is unlikely to be improved upon with foreseeable gravitational-wave observations [74].

Conclusions and outlook.—The NICER’s observation of PSR J0030+0451 allows the extraction of new astrophysical and theoretical physics inferences when one uses equation-of-state-insensitive relations. We have here shown the first inferences of the moment of inertia, the quadrupole moment, the surface eccentricity, and the Love number of an isolated neutron star. We have also been able to perform the first theory-agnostic and equation-of-state independent test of general relativity by combining NICER’s and LIGO/Virgo’s observations. This test, in turn, was leveraged to produce the most stringent constraint on gravitational parity violation, improving previous bounds by 7 orders of magnitude. This robust methodology can be applied to future multimessenger observations of neutron stars with NICER and gravitational wave observatories, with important implications to nuclear astrophysics and theoretical physics.

We thank Toral Gupta, Fred Lamb, Philippe Landry, and Helvi Witek for various discussions. We also thank Cole Miller, Sharon Morsink, and Kent Yagi for suggestions that improved this work. We thank the NICER Collaboration for making [44,45] publicly available. H. O. S., A. C.-A., and N. Y. are supported by NASA Grants No. NNX16AB98G, No. 80NSSC17M0041, No. 80NSSC18K1352, and NSF Grant No. 1759615. A. C.-A. also acknowledges funding from the Fundación Universitaria Konrad Lorenz (Project No. 5INV1). A. M. H. was supported by the DOE NNSA Stewardship Science Graduate Fellowship under Grant No. DE-NA0003864.

[1] G. Baym, T. Hatsuda, T. Kojo, P. D. Powell, Y. Song, and T. Takatsuka, From hadrons to quarks in neutron stars: A review, *Rep. Prog. Phys.* **81**, 056902 (2018).
 [2] F. Özel and P. Freire, Masses, radii, and the equation of state of neutron stars, *Annu. Rev. Astron. Astrophys.* **54**, 401 (2016).
 [3] E. Berti *et al.*, Testing general relativity with present and future astrophysical observations, *Classical Quantum Gravity* **32**, 243001 (2015).
 [4] K. Yagi and N. Yunes, I-Love-Q, *Science* **341**, 365 (2013).
 [5] K. Yagi and N. Yunes, I-Love-Q relations in neutron stars and their applications to astrophysics, gravitational waves and fundamental physics, *Phys. Rev. D* **88**, 023009 (2013).
 [6] D. D. Doneva and G. Pappas, Universal relations and alternative gravity theories, *Astrophysics and Space Science Library* **457**, 737 (2018).

[7] T. E. Riley *et al.*, A NICER view of PSR J0030+0451: Millisecond pulsar parameter estimation, *Astrophys. J. Lett.* **887**, L21 (2019).
 [8] M. Miller *et al.*, PSR J0030+0451 mass and radius from NICER data and implications for the properties of neutron star matter, *Astrophys. J. Lett.* **887**, L24 (2019).
 [9] K. C. Gendreau *et al.*, The Neutron star Interior Composition Explorer (NICER): Design and development, in *Proc. SPIE Int. Soc. Opt. Eng.*, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9905 (2016), p. 99051H, <https://doi.org/10.1117/12.2231304>.
 [10] A. N. Lommen, A. Zepka, D. C. Backer, M. McLaughlin, J. C. Cordes, Z. Arzoumanian, and K. Xilouris, New pulsars from an Arecibo drift scan search, *Astrophys. J.* **545**, 1007 (2000).
 [11] Z. Arzoumanian *et al.* (NANOGrav Collaboration), The NANOGrav 11-year Data Set: High-precision timing of 45 Millisecond pulsars, *Astrophys. J. Suppl. Ser.* **235**, 37 (2018).
 [12] J. M. Lattimer and M. Prakash, The equation of state of hot, dense matter and neutron stars, *Phys. Rep.* **621**, 127 (2016).
 [13] A. Maselli, V. Cardoso, V. Ferrari, L. Gualtieri, and P. Pani, Equation-of-state-independent relations in neutron stars, *Phys. Rev. D* **88**, 023007 (2013).
 [14] M. Bauböck, E. Berti, D. Psaltis, and F. Özel, Relations between neutron-star parameters in the Hartle-Thorne approximation, *Astrophys. J.* **777**, 68 (2013).
 [15] C. Breu and L. Rezzolla, Maximum mass, moment of inertia and compactness of relativistic stars, *Mon. Not. R. Astron. Soc.* **459**, 646 (2016).
 [16] S. M. Morsink, D. A. Leahy, C. Cadeau, and J. Braga, The oblate schwarzschild approximation for light curves of rapidly rotating neutron stars, *Astrophys. J.* **663**, 1244 (2007).
 [17] S. Bogdanov *et al.*, Constraining the neutron star mass–radius relation and dense matter equation of state with NICER. II. Emission from hot spots on a rapidly rotating neutron star, *Astrophys. J.* **887**, L26 (2019).
 [18] J. B. Hartle, Slowly rotating relativistic stars. I. Equations of structure, *Astrophys. J.* **150**, 1005 (1967).
 [19] J. B. Hartle and K. S. Thorne, Slowly rotating relativistic stars. II. models for neutron stars and supermassive stars, *Astrophys. J.* **153**, 807 (1968).
 [20] S. L. Shapiro and S. A. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects* (Wiley, New York, 1983).
 [21] J. S. Read, B. D. Lackey, B. J. Owen, and J. L. Friedman, Constraints on a phenomenologically parameterized neutron-star equation of state, *Phys. Rev. D* **79**, 124032 (2009).
 [22] B. Kumar and P. Landry, Inferring neutron star properties from GW170817 with universal relations, *Phys. Rev. D* **99**, 123026 (2019).
 [23] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.126.181101> for further details on the equation of state and neutron star catalogs, the C relations, our inference scheme, the I-Love relation in dynamical Chern-Simons gravity, and the parametrized I-Love test, which includes Refs. [24–40].

- [24] H. T. Cromartie *et al.*, Relativistic Shapiro delay measurements of an extremely massive millisecond pulsar, *Nat. Astron.* **4**, 72 (2020).
- [25] M. C. Miller, C. Chirenti, and F. K. Lamb, Constraining the equation of state of high-density cold matter using nuclear and astronomical measurements, *Astrophys. J.* **888**, 12 (2020).
- [26] K. Yagi, L. C. Stein, N. Yunes, and T. Tanaka, Isolated and binary neutron stars in dynamical Chern-Simons gravity, *Phys. Rev. D* **87**, 084058 (2013); Erratum, *Phys. Rev. D* **93**, 089909 (2016).
- [27] K. Yagi and N. Yunes, Approximate universal relations for neutron stars and quark stars, *Phys. Rep.* **681**, 1 (2017).
- [28] Y. Ali-Haïmoud and Y. Chen, Slowly-rotating stars and black holes in dynamical Chern-Simons gravity, *Phys. Rev. D* **84**, 124033 (2011).
- [29] T. Gupta, B. Majumder, K. Yagi, and N. Yunes, I-Love-Q relations for neutron stars in dynamical Chern Simons gravity, *Classical Quantum Gravity* **35**, 025009 (2018).
- [30] T. Damour and J. H. Taylor, Strong field tests of relativistic gravity and binary pulsars, *Phys. Rev. D* **45**, 1840 (1992).
- [31] T. Damour and G. Esposito-Farèse, Tensor multiscalar theories of gravitation, *Classical Quantum Gravity* **9**, 2093 (1992).
- [32] T. Damour and G. Esposito-Farèse, Nonperturbative Strong Field Effects in Tensor—Scalar Theories of Gravitation, *Phys. Rev. Lett.* **70**, 2220 (1993).
- [33] T. Damour and G. Esposito-Farèse, Tensor—scalar gravity and binary pulsar experiments, *Phys. Rev. D* **54**, 1474 (1996).
- [34] D. Anderson, P. Freire, and N. Yunes, Binary pulsar constraints on massless scalar-tensor theories using Bayesian statistics, *Classical Quantum Gravity* **36**, 225009 (2019).
- [35] P. Wagle, N. Yunes, D. Garfinkle, and L. Bieri, Hair loss in parity violating gravity, *Classical Quantum Gravity* **36**, 115004 (2019).
- [36] N. Yunes, D. Psaltis, F. Ozel, and A. Loeb, Constraining parity violation in gravity with measurements of neutron-star moments of inertia, *Phys. Rev. D* **81**, 064020 (2010).
- [37] H. Sotani, Pulse profiles from a pulsar in scalar-tensor gravity, *Phys. Rev. D* **96**, 104010 (2017).
- [38] H. O. Silva and N. Yunes, Neutron star pulse profiles in scalar-tensor theories of gravity, *Phys. Rev. D* **99**, 044034 (2019).
- [39] R. Xu, Y. Gao, and L. Shao, Strong-field effects in massive scalar-tensor gravity for slowly spinning neutron stars and application to X-ray pulsar pulse profiles, *Phys. Rev. D* **102**, 064057 (2020).
- [40] H. O. Silva and N. Yunes, Neutron star pulse profile observations as extreme gravity probes, *Classical Quantum Gravity* **36**, 17LT01 (2019).
- [41] T. Mora and C. M. Will, Numerically generated quasiequilibrium orbits of black holes: Circular or eccentric?, *Phys. Rev. D* **66**, 101501(R) (2002).
- [42] E. E. Flanagan and T. Hinderer, Constraining neutron star tidal Love numbers with gravitational wave detectors, *Phys. Rev. D* **77**, 021502(R) (2008).
- [43] T. Hinderer, Tidal Love numbers of neutron stars, *Astrophys. J.* **677**, 1216 (2008).
- [44] M. C. Miller *et al.*, NICER PSR J0030+0451 Illinois-Maryland MCMC Samples, <https://doi.org/10.5281/zenodo.3473466> (2019).
- [45] T. E. Riley, A. L. Watts, S. Bogdanov, P. S. Ray, R. M. Ludlam, S. Guillot, Z. Arzoumanian, C. L. Baker, A. V. Bilous, D. Chakrabarty, K. C. Gendreau, A. K. Harding, W. C. G. Ho, J. M. Lattimer, S. M. Morsink, and T. E. Strohmayer, A NICER View of PSR J0030+0451: Nested Samples for Millisecond Pulsar Parameter Estimation, https://zenodo.org/record/3386449#YG1szT_hXIU (2019).
- [46] S. Bogdanov *et al.*, Constraining the neutron star mass–radius relation and dense matter equation of state with NICER. I. The millisecond pulsar X-ray data set, *Astrophys. J. Lett.* **887**, L25 (2019).
- [47] M. Kramer *et al.*, Tests of general relativity from timing the double pulsar, *Science* **314**, 97 (2006).
- [48] M. Kramer and N. Wex, The double pulsar system: A unique laboratory for gravity, *Classical Quantum Gravity* **26**, 073001 (2009).
- [49] I. H. Stairs, S. E. Thorsett, R. J. Dewey, M. Kramer, and C. A. McPhee, The formation of the double pulsar PSR J0737-3039A/B, *Mon. Not. R. Astron. Soc.* **373**, L50 (2006).
- [50] R. D. Ferdman *et al.*, The double pulsar: Evidence for neutron star formation without an iron core-collapse supernova, *Astrophys. J.* **767**, 85 (2013).
- [51] J. M. Lattimer and B. F. Schutz, Constraining the equation of state with moment of inertia measurements, *Astrophys. J.* **629**, 979 (2005).
- [52] P. Landry and B. Kumar, Constraints on the moment of inertia of PSR J0737-3039A from GW170817, *Astrophys. J.* **868**, L22 (2018).
- [53] Y. Lim, J. W. Holt, and R. J. Stahulak, Predicting the moment of inertia of pulsar J0737-3039A from Bayesian modeling of the nuclear equation of state, *Phys. Rev. C* **100**, 035802 (2019).
- [54] K. Yagi and N. Yunes, Binary Love relations, *Classical Quantum Gravity* **33**, 13LT01 (2016).
- [55] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), Properties of the binary neutron star merger GW170817, *Phys. Rev. X* **9**, 011001 (2019).
- [56] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.* **119**, 161101 (2017).
- [57] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), GW170817: Measurements of Neutron Star Radii and Equation of State, *Phys. Rev. Lett.* **121**, 161101 (2018).
- [58] T. K. Chan, A. P. O. Chan, and P. T. Leung, I-Love relations for incompressible stars and realistic stars, *Phys. Rev. D* **91**, 044017 (2015).
- [59] N. Yunes and F. Pretorius, Fundamental theoretical bias in gravitational wave astrophysics and the parameterized post-Einsteinian framework, *Phys. Rev. D* **80**, 122003 (2009).
- [60] R. Jackiw and S. Y. Pi, Chern-Simons modification of general relativity, *Phys. Rev. D* **68**, 104012 (2003).
- [61] S. Weinberg, A tree theorem for inflation, *Phys. Rev. D* **78**, 063534 (2008).

- [62] J. Garcia-Bellido, M. Garcia-Perez, and A. Gonzalez-Arroyo, Chern-Simons production during preheating in hybrid inflation models, *Phys. Rev. D* **69**, 023504 (2004).
- [63] S. H. Alexander and S. J. Gates, Jr., Can the string scale be related to the cosmic baryon asymmetry?, *J. Cosmol. Astropart. Phys.* **06** (2006) 018.
- [64] S. H.-S. Alexander, M. E. Peskin, and M. M. Sheikh-Jabbari, Leptogenesis from Gravity Waves in Models of Inflation, *Phys. Rev. Lett.* **96**, 081301 (2006).
- [65] M. Adak and T. Dereli, String-inspired Chern-Simons modified gravity in 4-dimensions, *Eur. Phys. J. C* **72**, 1979 (2012).
- [66] A. Ashtekar, A. Balachandran, and S. Jo, The CP problem in quantum gravity, *Int. J. Mod. Phys. A* **04**, 1493 (1989).
- [67] S. Mercuri and V. Taveras, Interaction of the Barbero-Immirzi field with matter and pseudo-scalar perturbations, *Phys. Rev. D* **80**, 104007 (2009).
- [68] L. Smolin and C. Soo, The Chern-Simons invariant as the natural time variable for classical and quantum cosmology, *Nucl. Phys.* **B449**, 289 (1995).
- [69] T. Delsate, D. Hilditch, and H. Witek, Initial value formulation of dynamical Chern-Simons gravity, *Phys. Rev. D* **91**, 024027 (2015).
- [70] S. Alexander and N. Yunes, Chern-Simons modified general relativity, *Phys. Rep.* **480**, 1 (2009).
- [71] K. Yagi, N. Yunes, and T. Tanaka, Slowly rotating black holes in dynamical Chern-Simons gravity: Deformation quadratic in the spin, *Phys. Rev. D* **86**, 044037 (2012); Erratum, *Phys. Rev. D* **89**, 049902 (2014).
- [72] Y. Nakamura, D. Kikuchi, K. Yamada, H. Asada, and N. Yunes, Weakly-gravitating objects in dynamical Chern-Simons gravity and constraints with gravity probe B, *Classical Quantum Gravity* **36**, 105006 (2019).
- [73] R. Nair, S. Perkins, H. O. Silva, and N. Yunes, Fundamental Physics Implications for Higher-Curvature Theories from Binary Black Hole Signals in the LIGO-Virgo Catalog GWTC-1, *Phys. Rev. Lett.* **123**, 191101 (2019); Erratum, *Phys. Rev. Lett.* **124**, 169904 (2020).
- [74] S. H. Alexander and N. Yunes, Gravitational wave probes of parity violation in compact binary coalescences, *Phys. Rev. D* **97**, 064033 (2018).