

ASYMPTOTICS OF THE PRINCIPAL EIGENVALUE FOR A LINEAR TIME-PERIODIC PARABOLIC OPERATOR II: SMALL DIFFUSION

SHUANG LIU, YUAN LOU, RUI PENG, AND MAOLIN ZHOU

ABSTRACT. We investigate the effect of small diffusion on the principal eigenvalues of linear time-periodic parabolic operators with zero Neumann boundary conditions in one dimensional space. The asymptotic behaviors of the principal eigenvalues, as the diffusion coefficients tend to zero, are established for non-degenerate and degenerate spatial-temporally varying environments. A new finding is the dependence of these asymptotic behaviors on the periodic solutions of a specific ordinary differential equation induced by the drift. The proofs are based upon delicate constructions of super/sub-solutions and the applications of comparison principles.

1. INTRODUCTION

In this paper, we consider the following linear time-periodic parabolic eigenvalue problem in one dimensional space:

$$(1.1) \quad \begin{cases} \partial_t \varphi - D \partial_{xx} \varphi - \partial_x m \partial_x \varphi + V \varphi = \lambda(D) \varphi & \text{in } (0, 1) \times (0, T), \\ \partial_x \varphi(t, 0) = \partial_x \varphi(1, t) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } (0, 1), \end{cases}$$

where $D > 0$ represents the diffusion rate, and the functions $m \in C^{2,1}([0, 1] \times [0, T])$ and $V \in C([0, 1] \times [0, T])$ are assumed to be periodic in t with a common period T .

By the Krein-Rutman Theorem, (1.1) admits a simple and real eigenvalue (called *principal eigenvalue*), denoted by $\lambda(D)$, which corresponds to a positive eigenfunction (called *principal eigenfunction*) and satisfies $\text{Re}\lambda > \lambda(D)$ for any other eigenvalue λ of (1.1); see Proposition 7.2 of [12]. The principal eigenvalue $\lambda(D)$ plays a fundamental role in the study of reaction-diffusion equations and systems in spatio-temporal media, e.g. in the stability analysis for equilibria [3, 4, 12, 14]. Of particular interest is to understand the dependence of $\lambda(D)$ on the parameters [15, 16, 19, 20]. The present paper continues our previous studies in [17, 18] on the principal eigenvalues for time-periodic parabolic operators, where the dependence of $\lambda(D)$ on frequency and advection rate were investigated. Our main goal here is to establish the asymptotic behavior of $\lambda(D)$ as the diffusion rate D tends to zero.

Received by the editors February 2, 2020, and, in revised form, October 18, 2020.

2020 *Mathematics Subject Classification*. Primary 35P15, 35P20; Secondary 35K10, 35B10.

The first author was partially supported by the Outstanding Innovative Talents Cultivation Funded Programs 2018 of Renmin Univerty of China and the NSFC grant No. 11571364.

The second author was partially supported by the NSF grant DMS-1853561.

The third author was partially supported by the NSFC grant No. 11671175.

The fourth author was partially supported by National Key R&D Program of China (2020YFA0713300) and Nankai ZhiDe Foundation.

For notational convenience, given any T -periodic function $p(x, t)$, we define

$$\hat{p}(x) := \frac{1}{T} \int_0^T p(x, s) \, ds \quad \text{and} \quad p_+(x, t) := \max \{p(x, t), 0\},$$

and redefine $\partial_{xx}\hat{m}(0)$ and $\partial_{xx}\hat{m}(1)$ via

$$(1.2) \quad \partial_{xx}\hat{m}(0) = \begin{cases} -\infty & \text{if } \partial_x\hat{m}(0^+) < 0, \\ \infty & \text{if } \partial_x\hat{m}(0^+) > 0, \end{cases} \quad \text{and} \quad \partial_{xx}\hat{m}(1) = \begin{cases} \infty & \text{if } \partial_x\hat{m}(1^-) < 0, \\ -\infty & \text{if } \partial_x\hat{m}(1^-) > 0. \end{cases}$$

For the case when V and $\partial_x m$ depend upon the space variable alone, i.e. $V(x, t) = V(x)$ and $\partial_x m(x, t) = m'(x)$, problem (1.1) reduces to the following elliptic eigenvalue problem:

$$(1.3) \quad \begin{cases} -D\varphi'' - m'(x)\varphi' + V(x)\varphi = \lambda(D)\varphi & \text{in } (0, 1), \\ \varphi'(0) = \varphi'(1) = 0. \end{cases}$$

This sort of advection-diffusion operator in (1.3) with small diffusion can be regarded as a singular perturbation of the corresponding first order operator [24], and was studied in [11] by the large deviation approach. Therein, the limit of the principal eigenvalue $\lambda(D)$ as $D \rightarrow 0$ plays a pivotal role in studying the large time behavior of the trajectories of stochastic systems; see also [7, 10]. Recently the asymptotic behavior of $\lambda(D)$ for problem (1.3) has been considered in [6] for general bounded domains, and their result in particular implies

Theorem 1.1 ([6]). *Assume $V(x, t) = V(x)$ and $\partial_x m(x, t) = m'(x)$. Suppose that $m'(0) \neq 0$, $m'(1) \neq 0$, and all critical points of m are non-degenerate. Then*

$$\lim_{D \rightarrow 0} \lambda(D) = \min_{x \in \Sigma \cup \{0, 1\}} \{V(x) + [m'']_+(x)\},$$

where $\Sigma := \{x \in (0, 1) : m'(x) = 0\}$ and $m''(0), m''(1)$ are defined by (1.2).

We refer to [21] for recent progress on problem (1.3) under general boundary conditions.

Theorem 1.1 indicates that the limit of $\lambda(D)$ relies upon the set of critical points of function m in the elliptic scenario. Turning to the time-periodic parabolic case where m depends on both spatial and temporal variables, it seems reasonable to anticipate that the limit of $\lambda(D)$ will be associated to the curves $x(t)$ satisfying $\partial_x m(x(t), t) = 0$. This is indeed the case for the limit of the principal eigenvalue with large advection, and we refer to Theorem 1.1 in [18] for further details. However, it turns out that this is generally not true while considering the limit of $\lambda(D)$ as D tends to zero. Instead, the asymptotic behavior of $\lambda(D)$ depends heavily on the periodic solutions of the following ordinary differential equation:

$$(1.4) \quad \begin{cases} \dot{P}(t) = -\partial_x m(P(t), t), \\ P(t) = P(t + T). \end{cases}$$

More specifically, our main result can be stated as follows.

Theorem 1.2. *Assume that $\partial_x m(0, t) \neq 0$ and $\partial_x m(1, t) \neq 0$ for all $t \in [0, T]$. Let $\partial_{xx}\hat{m}(0)$ and $\partial_{xx}\hat{m}(1)$ be defined by (1.2).*

(i) *If (1.4) has at least one but finitely many T -periodic solutions, denoted by $\{P_i(t)\}_{i=1}^N$, satisfying $0 < P_1(t) < \dots < P_N(t) < 1$, and $\partial_{xx}m(P_i(t), t) \neq 0$ for*

$1 \leq i \leq N$ and $t \in [0, T]$, then

$$\lim_{D \rightarrow 0} \lambda(D) = \min_{0 \leq i \leq N+1} \left\{ \frac{1}{T} \int_0^T \left[V(P_i(s), s) + [\partial_{xx}m]_+(P_i(s), s) \right] ds \right\},$$

where $P_0(t) \equiv 0$ and $P_{N+1}(t) \equiv 1$;

(ii) If (1.4) has no periodic solutions, then

$$\lim_{D \rightarrow 0} \lambda(D) = \min \left\{ \hat{V}(0) + [\partial_{xx}\hat{m}]_+(0), \hat{V}(1) + [\partial_{xx}\hat{m}]_+(1) \right\}.$$

If V and m are independent of time, all solutions of (1.4) are constants which correspond to the critical points of function m , and part (i) of Theorem 1.2 is reduced to Theorem 1.1. When $m(x, t)$ is monotone in x , part (ii) of Theorem 1.2 was first established in [22].

One potential application of Theorem 1.2 is the study of large-time behaviours of solutions to the Cauchy problem for singularly perturbed parabolic equations in spatio-temporal media [1, 8, 12], in which the growth or decay rate of the solutions can be described in terms of $\lambda(D)$. In a very recent work [9], the asymptotics of $\lambda(D)$ for small D was considered in a case of underlying advection $\partial_x m$ being a constant, when analyzing the effect of small mutations on phenotypically-structured populations in a shifting and fluctuating environment.

The restriction $\partial_{xx}m(P(t), t) \neq 0$ in Theorem 1.2, in fact guarantees the non-degeneracy of advection $\partial_x m$ along periodic solution P of (1.4). See [5, 18] for the definitions of degeneracy and non-degeneracy. To complement Theorem 1.2, we consider a type of degenerate advection in the following result:

Theorem 1.3. Suppose that for each $1 \leq i \leq N$, $\partial_x m(\kappa_i, t) \equiv 0$ for all $t \in [0, T]$, and $0 < \kappa_1 < \dots < \kappa_N < 1$. Furthermore, assume that $\{i : 0 \leq i \leq N\} = \mathbf{A} \cup \mathbf{B}$, where

$$\begin{aligned} \mathbf{A} &= \{i : 0 \leq i \leq N, \partial_x m(x, t) \neq 0, (x, t) \in (\kappa_i, \kappa_{i+1}) \times [0, T]\}; \\ \mathbf{B} &= \{i : 0 \leq i \leq N, \partial_x m(x, t) \equiv 0, (x, t) \in [\kappa_i, \kappa_{i+1}] \times [0, T]\}, \end{aligned}$$

with $\kappa_0 = 0$ and $\kappa_{N+1} = 1$. Then we have

(1.5)

$$\lim_{D \rightarrow 0} \lambda(D) = \min \left\{ \min_{0 \leq i \leq N+1} \left\{ \hat{V}(\kappa_i) + [\partial_{xx}\hat{m}]_+(\kappa_i) \right\}, \min_{i \in \mathbf{B}} \left\{ \min_{x \in [\kappa_i, \kappa_{i+1}]} \hat{V}(x) \right\} \right\},$$

where $\partial_{xx}\hat{m}(0)$ and $\partial_{xx}\hat{m}(1)$ are defined by (1.2).

The main contribution of Theorem 1.3 is to allow $\mathbf{B} \neq \emptyset$, i.e. the spatial-temporal degeneracy of function $\partial_x m$. When $\mathbf{B} = \emptyset$, which means $\partial_x m(x, t) \neq 0$ for all $x \neq \kappa_i, 0 \leq i \leq N+1$, all solutions of (1.4) are nothing but constant solutions $P \equiv \kappa_i, 1 \leq i \leq N$, and consequently, Theorem 1.3 becomes a special case of Theorem 1.2 when $\mathbf{B} = \emptyset$.

The assumption $i \in \mathbf{A}$ implies there are no periodic solutions of (1.4) in $[\kappa_i, \kappa_{i+1}] \times [0, T]$ except for constant solutions $P \equiv \kappa_i$ and $P \equiv \kappa_{i+1}$. Without this assumption, the situation becomes even more complicated. To illustrate the complexity, we consider the special case $m(x, t) = ab(t)x$ as in [18], where $\alpha > 0$ denotes the advection rate, and the T -periodic function b is Lipschitz continuous. In this case,

problem (1.1) becomes

$$(1.6) \quad \begin{cases} \partial_t \varphi - D \partial_{xx} \varphi - \alpha b(t) \partial_x \varphi + V \varphi = \lambda(D) \varphi & \text{in } (0, 1) \times [0, T], \\ \partial_x \varphi(0, t) = \partial_x \varphi(1, t) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } (0, 1). \end{cases}$$

For different α and b , we have the following result:

Theorem 1.4. *Let $\lambda(D)$ denote the principal eigenvalue of (1.6).*

(i) *If $\hat{b} \neq 0$, then for all $\alpha > 0$,*

$$\lim_{D \rightarrow 0} \lambda(D) = \begin{cases} \hat{V}(1) & \text{for } \hat{b} > 0, \\ \hat{V}(0) & \text{for } \hat{b} < 0. \end{cases}$$

(ii) *If $\hat{b} = 0$, set $P(t) = -\int_0^t b(s) ds$, $\bar{P} = \max_{[0, T]} P$, and $\underline{P} = \min_{[0, T]} P$. Then*

$$\lim_{D \rightarrow 0} \lambda(D) = \begin{cases} \min_{y \in [-\alpha \underline{P}, 1 - \alpha \bar{P}]} \left\{ \frac{1}{T} \int_0^T V(\alpha P(s) + y, s) ds \right\}, & 0 < \alpha \leq \frac{1}{\bar{P} - \underline{P}}, \\ \frac{1}{T} \int_0^T V(\tilde{P}_\alpha(s), s) ds, & \alpha > \frac{1}{\bar{P} - \underline{P}}, \end{cases}$$

where $\tilde{P}_\alpha \in C([0, T]; [0, 1])$ is the unique T -periodic solution of $\dot{\tilde{P}}(t) = -\alpha F(\tilde{P}(t), t)$ in $[0, 1]$, and F is given by

$$(1.7) \quad F(x, t) = \begin{cases} b(t) & 0 < x < 1, t \in [0, T], \\ \min\{b(t), 0\}, & x = 0, t \in [0, T], \\ \max\{b(t), 0\}, & x = 1, t \in [0, T]. \end{cases}$$

Remark 1.5. When $\hat{b} = 0$ and $\alpha = \frac{1}{\bar{P} - \underline{P}}$, part (ii) of Theorem 1.4 implies that $\lambda(D) \rightarrow \frac{1}{T} \int_0^T V \left(\frac{P(s) - \underline{P}}{\bar{P} - \underline{P}}, s \right) ds$ as $D \rightarrow 0$. Direct calculation yields that $\frac{P(t) - \underline{P}}{\bar{P} - \underline{P}}$ is in fact a periodic solution of $\dot{P}(t) = -\frac{1}{\bar{P} - \underline{P}} F(\tilde{P}(t), t)$, so that the uniqueness part in Lemma 4.1 implies $\tilde{P}_\alpha(t) \rightarrow \frac{P(t) - \underline{P}}{\bar{P} - \underline{P}}$ as $\alpha \rightarrow \frac{1}{\bar{P} - \underline{P}}$. This means that the limit of $\lambda(D)$ as $D \rightarrow 0$ is continuous at $\alpha = \frac{1}{\bar{P} - \underline{P}}$.

For $m(x, t) = \alpha b(t) x$, Theorem 1.4 gives a complete description of the behaviors of $\lambda(D)$ as $D \rightarrow 0$, and it provides a type of complicated spatial-temporal degeneracy not covered by Theorem 1.3. To further illustrate Theorem 1.4, consider the case $b(t) = -\frac{\pi}{T} \sin \left(\frac{2\pi t}{T} \right)$, in which

$$P(t) = \frac{1}{2} \cos \left(\frac{2\pi t}{T} \right) - \frac{1}{2}, \quad \bar{P} = 0, \quad \underline{P} = -1.$$

More precisely, (i) when $0 < \alpha < 1$, we could find some $y_\alpha \in [\alpha, 1]$ such that $\lambda(D) \rightarrow \frac{1}{T} \int_0^T V(\alpha P(s) + y_\alpha, s) ds$ as $D \rightarrow 0$, and the trajectory $\{\alpha P(t) + y_\alpha : t \in [0, T]\}$ in x - t plane is illustrated by the red solid curve in Fig. 1(a), where the two red dotted curves represent $\{\alpha P(t) + \alpha : t \in [0, T]\}$ and $\{\alpha P(t) + 1 : t \in [0, T]\}$, respectively; (ii) When $\alpha = 1$, we have $\lambda(D) \rightarrow \frac{1}{T} \int_0^T V(P(s) + 1, s) ds$ as $D \rightarrow 0$, and the trajectory $\{P(t) + 1 : t \in [0, T]\}$ is shown in Fig. 1(b); (iii) When $\alpha > 1$, it follows that $\lambda(D) \rightarrow \frac{1}{T} \int_0^T V(\tilde{P}_\alpha(s), s) ds$, and the corresponding trajectory $\{\tilde{P}_\alpha(t) : t \in [0, T]\}$ is given in Fig. 1(c)–(d).

As the proofs of Theorems 1.2, 1.3, and 1.4 are fairly technical, in the following we briefly outline the main strategies in proving Theorems 1.2 and 1.3:




FIGURE 1. Each rectangle corresponds to the region $[0, 1] \times [0, T]$ in x - t plane. The limit of $\lambda(D)$ as $D \rightarrow 0$ is determined by the average of V over the red solid curves, illustrated for various ranges of α and $m(x, t) = -\frac{\alpha \pi x}{T} \sin\left(\frac{2\pi t}{T}\right)$.

- (i) We note that $\lambda(D)$ for (1.3) in the elliptic situation can be characterized by variational formulation [5, 6, 21, 23]. In contrast, the time-periodic parabolic problem (1.1) has no variational formulations. Our general strategy is to construct super/sub-solutions and apply generalized comparison principle developed in [18, Theorem A.1]. This technique was first introduced by Berestycki and Lions [2] to the elliptic scenario, whereas its adaptation to our context is more subtle because of the presence of temporal variable; see [22] for further discussions.
- (ii) We first establish Theorem 1.3 which assumes that $\partial_x m$ is strictly positive, negative, or identically zero in each sub-interval (κ_i, κ_{i+1}) . The main difficulty is to establish the lower bound of the principal eigenvalue in (1.5). The construction of super-solutions near the curves $\{(\kappa_i, t), t \in [0, T]\}$ is rather subtle, due to the fact that the spatio-temporal derivatives of the principal eigenfunction of (1.1) restricted to the curves may be unbounded as D tends to zero. Our strategy is to construct the super-solution almost coinciding with the principal eigenfunction of (1.1) near these curves, and then use an iterated argument to extend the super-solution to the whole domain.
- (iii) A key ingredient in the proof of Theorem 1.2 is to recognize the critical role of the solutions of (1.4). Our idea is to reduce the proof of Theorem 1.2 to that of Theorem 1.3 with $\mathbf{B} = \emptyset$. As Theorem 1.3 assumes that $\partial_x m$ is either strictly positive or negative in each sub-interval (κ_i, κ_{i+1}) , there are two difficulty in doing so: First, the solutions $P_i(t)$ of (1.4) are not constant ones as specified in Theorem 1.3. This difficulty can be overcome by introducing a proper transformation so that $P_i(t)$ become constant after the transformation. The second difficulty is that *a priori* we do not know the sign of the term $\partial_x m$ in each (κ_i, κ_{i+1}) . Our idea is to introduce another transformation, which is associated with the trajectories of (1.4). We prove that after the second transformation, $\partial_x m$ is indeed either strictly positive

or negative in each (κ_i, κ_{i+1}) , so that the proof of Theorem 1.3 is directly applicable to complete the proof of Theorem 1.2.

This paper is organized as follows: In Section 2 we present some results associated with the case when all of periodic solutions of (1.4) are constants and establish Theorem 1.3. These results are used in Section 3 to give the proof of Theorem 1.2, by combining with an idea of “straightening periodic solutions”. Section 4 is devoted to the proof of Theorem 1.4. A generalized comparison result will be presented in the Appendix.

2. PROOF OF THEOREM 1.3

This section is devoted to the proof of Theorem 1.3. Hereafter, we use \mathcal{L}_D to denote the time-periodic parabolic operator

$$\mathcal{L}_D := \partial_t - D\partial_{xx} - \partial_x m\partial_x + V.$$

For any $x \in [0, 1]$, we define a T -periodic function $f_x : [0, T] \rightarrow (0, \infty)$ by

$$(2.1) \quad f_x(t) = \exp \left[- \int_0^t V(x, s) ds + \hat{V}(x)t \right],$$

which solves, for fixed $x \in [0, 1]$, that $(\log f_x)' = \hat{V}(x) - V(x, t)$.

Proposition 2.1. *For any constant $\kappa \in (0, 1)$, suppose that*

$$\begin{cases} \partial_x m(x, t) > 0, & (x, t) \in [0, \kappa) \times [0, T], \\ \partial_x m(x, t) < 0, & (x, t) \in (\kappa, 1] \times [0, T]. \end{cases}$$

Then we have

$$\lim_{D \rightarrow 0} \lambda(D) = \hat{V}(\kappa).$$

Proof. We first prove the upper bound

$$(2.2) \quad \limsup_{D \rightarrow 0} \lambda(D) \leq \hat{V}(\kappa).$$

Fix any $\epsilon > 0$. For sufficiently small D , we construct a strict non-negative sub-solution $\underline{\varphi}$ in the sense of Definition A.1 (see Appendix A) such that

$$(2.3) \quad \begin{cases} \mathcal{L}_D \underline{\varphi} \leq [\hat{V}(\kappa) + \epsilon] \underline{\varphi} & \text{in } ((0, 1) \setminus \mathbb{X}) \times (0, T), \\ \partial_x \underline{\varphi}(0, t) = \partial_x \underline{\varphi}(1, t) = 0 & \text{on } [0, T], \\ \underline{\varphi}(x, 0) = \underline{\varphi}(x, T) & \text{on } (0, 1), \end{cases}$$

for some point set \mathbb{X} determined later.

To this end, by continuity of V , we choose small $\delta \in (0, 1)$ such that

$$(2.4) \quad |V(x, t) - V(\kappa, t)| < \epsilon/2 \quad \text{on } [\kappa - \delta, \kappa + \delta] \times [0, T].$$

Then we define $\underline{\varphi}$ by

$$\underline{\varphi}(x, t) := f_\kappa(t) \cdot \underline{z}(x),$$

where $f_\kappa(t)$ is defined in (2.1) with $x = \kappa$, and $\underline{z} \in C([0, 1])$ is given by

$$(2.5) \quad \underline{z}(x) := \begin{cases} -(x - \kappa)^2 + \delta^2 & \text{on } [\kappa - \delta, \kappa + \delta], \\ 0 & \text{on } [0, \kappa - \delta] \cup (\kappa + \delta, 1]. \end{cases}$$

Observe that $\partial_x \underline{\varphi}((\kappa \pm \delta)^+, \cdot) > \partial_x \underline{\varphi}((\kappa \pm \delta)^-, \cdot)$. We identify \mathbb{X} in (2.3) as

$$\mathbb{X} = \{\kappa \pm \delta\}.$$

To verify (2.3), note from definition (2.1) that $f'_\kappa = (\hat{V}(\kappa) - V(\kappa, t))f_\kappa$, direct calculations on $[\kappa - \delta, \kappa + \delta] \times [0, T]$ yield that for small D ,

$$\begin{aligned} \mathcal{L}_D \underline{\varphi} &= f'_\kappa(t)\underline{\varphi} - D\partial_{xx}\underline{\varphi} - \partial_x m\partial_x \underline{\varphi} + V(x, t)\underline{\varphi} \\ &= [\hat{V}(\kappa) - V(\kappa, t) + V(x, t)]\underline{\varphi} - \partial_x m\partial_x \underline{\varphi} - D\partial_{xx}\underline{\varphi} \\ &\leq [\hat{V}(\kappa) + \epsilon/2]\underline{\varphi} - \partial_x m\partial_x \underline{\varphi} - D\partial_{xx}\underline{\varphi} \\ &\leq [\hat{V}(\kappa) + \epsilon]\underline{\varphi}, \end{aligned}$$

where $-\partial_x m\partial_x \underline{\varphi} - D\partial_{xx}\underline{\varphi} \leq \frac{\epsilon}{2}\underline{\varphi}$ in the last inequality is due to the fact that $-\partial_x m\partial_x \underline{\varphi} < 0 \leq \epsilon \underline{\varphi}$ in the neighborhoods of $\{\kappa \pm \delta\} \times [0, T]$. Hence (2.3) holds, and (2.2) follows from (2.3) and Proposition A.2 by letting $\epsilon \rightarrow 0^+$.

Next, we show that

$$(2.6) \quad \liminf_{D \rightarrow 0} \lambda(D) \geq \hat{V}(\kappa).$$

Define $\bar{\varphi} \in C^{2,1}([0, 1] \times [0, T])$ by

$$\bar{\varphi}(x, t) := f_\kappa(t) \cdot e^{M_1(x-\kappa)^2}$$

with $M_1 > 0$ to be specified later. For any given $\epsilon > 0$, we shall choose M_1 large so that for sufficiently small D , $\bar{\varphi}$ satisfies

$$(2.7) \quad \begin{cases} \mathcal{L}_D \bar{\varphi} \geq [\hat{V}(\kappa) - \epsilon]\bar{\varphi} & \text{in } (0, 1) \times (0, T), \\ \partial_x \bar{\varphi}(0, t) < 0 < \partial_x \bar{\varphi}(1, t) & \text{on } [0, T], \\ \bar{\varphi}(x, 0) = \bar{\varphi}(x, T) & \text{on } (0, 1). \end{cases}$$

To establish (2.7), we first recall that δ is chosen as in (2.4). For $x \in (0, \kappa - \delta] \cup [\kappa + \delta, 1)$, there exists some $\epsilon_0 > 0$ such that $|\partial_x m| \geq \epsilon_0$, and thus

$$-\partial_x m\partial_x(\log \bar{\varphi}) = 2M_1\partial_x m \cdot (x - \kappa) \geq 2M_1\delta\epsilon_0,$$

from which direct calculation leads to

$$(2.8) \quad \mathcal{L}_D \bar{\varphi} \geq \left\{ \hat{V}(\kappa) + V(x, t) - V(\kappa, t) - D[2M_1 + 4M_1^2(x - \kappa)^2] + 2M_1\delta\epsilon_0 \right\} \bar{\varphi}.$$

We choose M_1 large such that $2M_1\delta\epsilon_0 > 2\|V\|_{L^\infty}$. Letting D be small enough in (2.8), we deduce $\mathcal{L}_D \bar{\varphi} \geq \hat{V}(\kappa)\bar{\varphi}$ as desired.

For $x \in [\kappa - \delta, \kappa + \delta]$, by $-\partial_x m\partial_x \bar{\varphi} \geq 0$ and the definition of δ we have

$$\mathcal{L}_D \bar{\varphi} \geq \left\{ \hat{V}(\kappa) + V(x, t) - V(\kappa, t) - D[2M_1 + 4M_1^2(x - \kappa)^2] \right\} \bar{\varphi} \geq [\hat{V}(\kappa) - \epsilon] \bar{\varphi}$$

for sufficiently small D .

Therefore, (2.7) holds and (2.6) follows from Proposition A.2 with $\mathbb{X} = \emptyset$. \square

To proceed further, we will need the following result:

Lemma 2.2. *Let $\rho(t) \geq 0$ ($\not\equiv 0$) be any T -periodic function. For each $R > 0$, denote by μ_R the principal eigenvalue of the following problem:*

$$(2.9) \quad \begin{cases} \partial_t \varphi - \partial_{xx} \varphi - x \rho(t) \partial_x \varphi = \mu_R \varphi & \text{in } (-R, R) \times (0, T), \\ \varphi(-R, t) = \varphi(R, t) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } [-R, R]. \end{cases}$$

Then we have

$$\lim_{R \rightarrow \infty} \mu_R = \hat{\rho}.$$

Proof. For each $\delta \geq 0$, in view of $\rho(t) \geq 0$ ($\not\equiv 0$) in $[0, T]$, we choose $\beta_\delta(t)$ as the unique positive solution of the problem

$$(2.10) \quad \begin{cases} \frac{\dot{\beta}(t)}{2} = \beta(t) [\rho(t) + \delta - \beta(t)] & \text{in } [0, T], \\ \beta(0) = \beta(T). \end{cases}$$

Denote by $(\alpha_\delta(t), \mu_\delta)$ an eigenpair, with $\alpha_\delta(t) > 0$, of the eigenvalue problem

$$(2.11) \quad \begin{cases} \dot{\alpha}(t) + \beta_\delta(t) \alpha(t) = \mu \alpha(t) & \text{in } [0, T], \\ \alpha(0) = \alpha(T). \end{cases}$$

Dividing both sides of (2.10) by β_δ , and integrating the resulting equation over $[0, T]$, by periodicity of β_δ we have $\hat{\beta}_\delta = \hat{\rho} + \delta$. Similarly, (2.11) implies $\mu_\delta = \hat{\beta}_\delta$. Therefore,

$$(2.12) \quad \mu_\delta = \hat{\beta}_\delta = \hat{\rho} + \delta.$$

For any $\delta \geq 0$, we define T -periodic function $\psi_\delta \in C^{2,1}(\mathbb{R} \times [0, T])$ by

$$(2.13) \quad \psi_\delta(x, t) := \alpha_\delta(t) e^{-\frac{\beta_\delta(t)}{2} x^2},$$

which, by definitions (2.10) and (2.11), solves

$$(2.14) \quad \mathcal{L}_\delta \psi_\delta := \partial_t \psi_\delta - \partial_{xx} \psi_\delta - x(\rho(t) + \delta) \partial_x \psi_\delta = \mu_\delta \psi_\delta \quad \text{in } \mathbb{R} \times [0, T].$$

We first show $\liminf_{R \rightarrow \infty} \mu_R \geq \hat{\rho}$. For $\delta = 0$, ψ_0 defined by (2.13) is a super-solution to (2.9) in the sense of Definition A.1 for any $R > 0$. By Proposition A.2, we have $\mu_R \geq \mu_0 = \hat{\rho}$ for any $R > 0$, and thus $\liminf_{R \rightarrow \infty} \mu_R \geq \hat{\rho}$.

Next, we show $\limsup_{R \rightarrow \infty} \mu_R \leq \hat{\rho}$. Fix any $\delta > 0$. Choose $R_\delta > 0$ large such that $\delta x^2 \beta_\delta(t) \geq \hat{\rho} + \delta$ for all $|x| \geq R_\delta$ and $t \in [0, T]$. Then let $\epsilon = \epsilon(\delta) > 0$ be small so that $\delta \psi_\delta(R_\delta, t) \geq \epsilon(\hat{\rho} + 2\delta)$ for all $t \in [0, T]$. Set $\tilde{\psi}_\delta = \max\{\psi_\delta - \epsilon, 0\}$. Note that we can choose ϵ smaller if necessary such that $\tilde{\psi}_\delta(x, t) > 0$ holds for all $|x| \leq R_\delta$ and $t \in [0, T]$.

On $\{(x, t) : \tilde{\psi}_\delta(x, t) > 0, |x| \geq R_\delta\}$, by (2.12) and (2.14) we calculate that

$$(2.15) \quad \begin{aligned} & \partial_t \tilde{\psi}_\delta - \partial_{xx} \tilde{\psi}_\delta - x \rho(t) \partial_x \tilde{\psi}_\delta - (\hat{\rho} + 2\delta) \tilde{\psi}_\delta \\ &= (\hat{\rho} + \delta) \psi_\delta - \delta x^2 \beta_\delta(t) \psi_\delta - (\hat{\rho} + 2\delta)(\psi_\delta - \epsilon) \\ &\leq (\hat{\rho} + \delta) \psi_\delta - \delta x^2 \beta_\delta(t) \psi_\delta \\ &\leq 0, \end{aligned}$$

where the last inequality follows from the choice of R_δ .

On $\{(x, t) : \tilde{\psi}_\delta(x, t) > 0, |x| \leq R_\delta\}$, we have

$$\begin{aligned}
 & \partial_t \tilde{\psi}_\delta - \partial_{xx} \tilde{\psi}_\delta - x\rho(t) \partial_x \tilde{\psi}_\delta - (\hat{\rho} + 2\delta) \tilde{\psi}_\delta \\
 &= (\hat{\rho} + \delta) \psi_\delta - \delta x^2 \beta_\delta(t) \psi_\delta - (\hat{\rho} + 2\delta)(\psi_\delta - \epsilon) \\
 (2.16) \quad & \leq (\hat{\rho} + \delta) \psi_\delta - (\hat{\rho} + 2\delta)(\psi_\delta - \epsilon) \\
 & \leq \epsilon(\hat{\rho} + 2\delta) - \delta \psi_\delta(R_\delta, t) \\
 & \leq 0,
 \end{aligned}$$

where the last inequality is due to the choice of ϵ .

Choose some large $\tilde{R}_\delta > R_\delta$ such that $\tilde{\psi}_\delta(\tilde{R}_\delta, t) = 0$ for all $t \in [0, T]$. By (2.15) and (2.16), the constructed $\tilde{\psi}_\delta$ is a sub-solution to (2.9) in the sense of Definition A.1 for any $R \geq \tilde{R}_\delta$. A direct application of Proposition A.2 yields $\mu_R \leq \hat{\rho} + 2\delta$ for all $R \geq \tilde{R}_\delta$, and thus $\limsup_{R \rightarrow \infty} \mu_R \leq \hat{\rho} + 2\delta$. Letting $\delta \rightarrow 0$ completes the proof. \square

Proposition 2.3. *For any $\kappa \in (0, 1)$, suppose that*

$$\begin{cases} \partial_x m(x, t) < 0, & (x, t) \in (0, \kappa) \times [0, T], \\ \partial_x m(x, t) > 0, & (x, t) \in (\kappa, 1) \times [0, T], \\ \partial_{xx} m(\kappa, t) \geq (\not\equiv) 0, & t \in [0, T], \\ m(x, 0) = m(x, T), & x \in [0, 1]. \end{cases}$$

Then we have

$$\lim_{D \rightarrow 0} \lambda(D) = \min \left\{ \hat{V}(0), \hat{V}(\kappa) + \partial_{xx} \hat{m}(\kappa), \hat{V}(1) \right\}.$$

Proof. For any given $\epsilon > 0$, we choose some small $\delta > 0$ such that

$$\begin{aligned}
 (2.17) \quad & |V(x, t) - V(0, t)| < \epsilon/2, \quad (x, t) \in [0, \delta] \times [0, T], \\
 & |V(x, t) - V(\kappa, t)| < \epsilon/2, \quad (x, t) \in [\kappa - \delta, \kappa + \delta] \times [0, T], \\
 & |V(x, t) - V(1, t)| < \epsilon/2, \quad (x, t) \in [1 - \delta, 1] \times [0, T].
 \end{aligned}$$

Part I. In this part, we establish the upper bound

$$\limsup_{D \rightarrow 0} \lambda(D) \leq \lambda_{\min} := \min \left\{ \hat{V}(0), \hat{V}(\kappa) + \partial_{xx} \hat{m}(\kappa), \hat{V}(1) \right\}.$$

By a similar argument as in Proposition 2.1, it is straightforward to show that

$$\limsup_{D \rightarrow 0} \lambda(D) \leq \min \left\{ \hat{V}(0), \hat{V}(1) \right\}.$$

It remains to prove

$$(2.18) \quad \limsup_{D \rightarrow 0} \lambda(D) \leq \hat{V}(\kappa) + \partial_{xx} \hat{m}(\kappa).$$

Fix any $\epsilon > 0$. For sufficiently small D , we construct a sub-solution $\underline{\varphi}$ such that

$$\begin{cases} \mathcal{L}_D \underline{\varphi} \leq [\hat{V}(\kappa) + \partial_{xx} \hat{m}(\kappa) + 2\epsilon] \underline{\varphi} & \text{in } ((0, 1) \setminus \mathbb{X}) \times [0, T], \\ \partial_x \underline{\varphi}(0, t) = \partial_x \underline{\varphi}(1, t) = 0 & \text{on } [0, T], \\ \underline{\varphi}(x, 0) = \underline{\varphi}(x, T) & \text{on } (0, 1), \end{cases}$$

where the set \mathbb{X} will be determined later.

To this end, we define

$$\bar{m}(x, t) := [\partial_{xx} m(\kappa, t) + \epsilon] \cdot \frac{(x - \kappa)^2}{2},$$

and further choose δ smaller if necessary such that

$$(2.20) \quad |\partial_x \bar{m}| \geq |\partial_x m| \quad \text{in } [\kappa - \delta, \kappa + \delta] \times [0, T].$$

Let $\bar{\lambda}_D$ denote the principal eigenvalue of the problem

$$(2.21) \quad \begin{cases} \partial_t \psi - D \partial_{xx} \psi - \partial_x \bar{m} \partial_x \psi = \bar{\lambda}_D \psi & \text{in } (\kappa - \delta, \kappa + \delta) \times [0, T], \\ \psi(\kappa - \delta, t) = \psi(\kappa + \delta, t) = 0 & \text{on } [0, T], \\ \psi(x, 0) = \psi(x, T) & \text{on } [\kappa - \delta, \kappa + \delta], \end{cases}$$

and the corresponding eigenfunction $\underline{\psi}_D$ is chosen to be positive in $(\kappa - \delta, \kappa + \delta) \times [0, T]$. Under the scaling $y = \frac{x - \kappa}{\sqrt{D}}$, we set $\underline{\varphi}_D(y, t) := \underline{\psi}_D(\sqrt{D}y + \kappa, t)$, which is the principal eigenfunction (associated to $\bar{\lambda}_D$) of the problem

$$\begin{cases} \partial_t \varphi - \partial_{yy} \varphi - y[\partial_{xx} m(\kappa, t) + \epsilon] \partial_y \varphi = \bar{\lambda}_D \varphi & \text{in } \left(-\frac{\delta}{\sqrt{D}}, \frac{\delta}{\sqrt{D}}\right) \times [0, T], \\ \varphi\left(-\frac{\delta}{\sqrt{D}}, t\right) = \varphi\left(\frac{\delta}{\sqrt{D}}, t\right) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } \left[-\frac{\delta}{\sqrt{D}}, \frac{\delta}{\sqrt{D}}\right]. \end{cases}$$

By Lemma 2.2, we deduce that

$$(2.22) \quad \lim_{D \rightarrow 0} \bar{\lambda}_D = \partial_{xx} \hat{m}(\kappa) + \epsilon.$$

We extend $\underline{\psi}_D$, the principal eigenfunction of (2.21), to $[0, 1] \times [0, T]$ by setting

$$\underline{\psi}_D \equiv 0 \quad \text{on } ([0, \kappa - \delta] \cup [\kappa + \delta, 1]) \times [0, T].$$

Applying the Hopf boundary lemma to (2.21), we have

$$\begin{aligned} \partial_x \underline{\psi}_D((\kappa - \delta)^+, \cdot) &> 0 = \partial_x \underline{\psi}_D((\kappa - \delta)^-, \cdot), \\ \partial_x \underline{\psi}_D((\kappa + \delta)^+, \cdot) &= 0 > \partial_x \underline{\psi}_D((\kappa + \delta)^-, \cdot), \end{aligned}$$

so that we choose \mathbb{X} by $\mathbb{X} = \{\kappa \pm \delta\}$.

Define

$$\underline{\varphi}(x, t) = f_\kappa(t) \cdot \underline{\psi}_D(x, t) \quad \text{in } [0, 1] \times [0, T],$$

where $f_\kappa(t)$ is given by (2.1) with $x = \kappa$. We verify that $\underline{\varphi}$ satisfies (2.19). By properties of $\underline{\psi}_D$ and (2.20) we can derive that

$$-\partial_x m \partial_x \underline{\psi}_D \leq -\partial_x \bar{m} \partial_x \underline{\psi}_D \quad \text{in } [0, 1] \times [0, T].$$

Hence, direct calculations on $((0, 1) \setminus \mathbb{X}) \times [0, T]$ give

$$\begin{aligned} \mathcal{L}_D \underline{\varphi} &= \left[-V(\kappa, t) + \hat{V}(\kappa) + V(x, t) \right] \underline{\varphi} + \left[\partial_t \underline{\psi}_D - D \partial_{xx} \underline{\psi}_D - \partial_x m \partial_x \underline{\psi}_D \right] f_\kappa(t) \\ &\leq \left[\hat{V}(\kappa) + \epsilon/2 \right] \underline{\varphi} + \left[\partial_t \underline{\psi}_D - D \partial_{xx} \underline{\psi}_D - \partial_x \bar{m} \partial_x \underline{\psi}_D \right] f_\kappa(t) \\ &= \left[\hat{V}(\kappa) + \bar{\lambda}_D + \epsilon/2 \right] \underline{\varphi} \\ &\leq \left[\hat{V}(\kappa) + \partial_{xx} \hat{m}(\kappa) + 2\epsilon \right] \underline{\varphi}, \end{aligned}$$

provided that D is small enough, where the last inequality is a consequence of (2.22). Therefore, $\underline{\varphi}$ defines a sub-solution satisfying (2.19), which together with Proposition A.2 implies (2.18).

Part II. We shall establish the lower bound

$$(2.23) \quad \liminf_{D \rightarrow 0} \lambda(D) \geq \lambda_{\min} := \min \left\{ \hat{V}(0), \hat{V}(\kappa) + \partial_{xx}\hat{m}(\kappa), \hat{V}(1) \right\}.$$

For each small $\epsilon > 0$, the main ingredient in the proof is to construct a positive continuous super-solution $\bar{\varphi}$ in the sense of Definition A.1, i.e. for sufficiently small D ,

$$(2.24) \quad \begin{cases} \mathcal{L}_D \bar{\varphi} \geq (1 - \epsilon)(\lambda_{\min} - \epsilon)\bar{\varphi} & \text{in } ((0, 1) \setminus \mathbb{X}) \times [0, T], \\ \partial_x \bar{\varphi}(0, t) = \partial_x \bar{\varphi}(1, t) = 0 & \text{on } [0, T], \\ \bar{\varphi}(x, 0) = \bar{\varphi}(x, T) & \text{on } (0, 1), \end{cases}$$

where the point set \mathbb{X} will be determined in Step 3. Then (2.23) follows from Proposition A.2 and arbitrariness of ϵ .

Step 1. We prepare some notations. First, we choose suitable T -periodic function $\underline{\rho}(t) \geq 0$ and small $\delta > 0$ such that

$$(2.25) \quad \begin{cases} \max\{\partial_{xx}m(\kappa, t) - \epsilon, 0\} \leq \underline{\rho}(t) \leq \partial_{xx}m(\kappa, t), & t \in [0, T], \\ \underline{\rho}(t)|x - \kappa| \leq |\partial_x m(x, t)|, & (x, t) \in [\kappa - \delta, \kappa + \delta] \times [0, T]. \end{cases}$$

Due to $\hat{\rho} > 0$, define $r(t)$ as the unique positive T -periodic solution of

$$(2.26) \quad \frac{\dot{r}(t)}{2 - \ell} = r(t) \left[\underline{\rho}(t) - \left(\frac{4}{(2 - \ell)^2} + \frac{\epsilon}{2} \right) r(t) \right],$$

where the small parameter $\ell \in (0, \epsilon/2]$ can be specified as follows: Note that there exist $0 < \underline{r} < \bar{r}$ independent of $\ell \in (0, \epsilon/2]$ such that

$$0 < \underline{r} < r(t) < \bar{r} \quad \text{for all } t \in [0, T] \text{ and } \ell \in [0, \epsilon/2].$$

We fix $\ell \in (0, \epsilon/2]$ small such that

$$(2.27) \quad \begin{cases} \frac{\frac{2}{2-\ell}}{\frac{4}{(2-\ell)^2} + \frac{\epsilon}{2}} \geq 1 - \epsilon, \\ \nu := \frac{\ell \bar{r}}{2 - \ell} < \left[\sqrt{\frac{4}{(2 - \ell)^2} + \frac{\epsilon}{2}} - 1 \right] \underline{r}. \end{cases}$$

Without loss of generality, we assume there is some $n_* \in \mathbb{N}$ ($n_* > 3$) such that

$$(2.28) \quad 1/\ell = 2^{n_* - 2},$$

and further choose δ smaller if necessary such that

$$\delta < \kappa - (n_* + 1)\delta < \kappa + (n_* + 1)\delta < 1 - \delta.$$

For fixed $r(t)$ and ℓ , we define $(\alpha_1(t), \lambda_\ell)$ as the eigenpair of

$$(2.29) \quad \begin{cases} \dot{\alpha}_1(t) + \frac{2}{2-\ell} \alpha_1(t) r(t) = \lambda_\ell \alpha_1(t) & \text{in } [0, T], \\ \alpha_1(0) = \alpha_1(T). \end{cases}$$

Similar to (2.12), we deduce from (2.26) and (2.29) that

$$\lambda_\ell = \frac{2}{2 - \ell} \hat{r} = \frac{\frac{2}{2-\ell}}{\frac{4}{(2-\ell)^2} + \frac{\epsilon}{2}} \hat{\rho},$$

which, together with (2.27), leads to

$$(2.30) \quad \lambda_\ell \geq (1 - \epsilon) \hat{\rho}.$$

Step 2. We construct a positive super-solution $\bar{\psi} \in C(\mathbb{R} \times [0, T])$ for the auxiliary problem

$$(2.31) \quad \begin{cases} \partial_t \psi - \partial_{yy} \psi - y \underline{\rho}(t) \partial_y \psi = (1 - \epsilon) \hat{\rho} \psi & \text{in } \mathbb{R} \times [0, T], \\ \psi(x, 0) = \psi(x, T) & \text{on } \mathbb{R}. \end{cases}$$

Using the notations introduced in Step 1, we define

$$(2.32) \quad \bar{\psi}(y, t) := \begin{cases} \alpha_1(t) e^{-\frac{r(t)}{2-\ell} y^2} & \text{on } [-y_0, y_0] \times [0, T], \\ \eta_1(t) e^{-\frac{(r(t)+\nu)y_0^\ell}{2-\ell} y^{2-\ell}} & \text{on } (y_0, \infty) \times [0, T], \\ \eta_1(t) e^{-\frac{(r(t)+\nu)y_0^\ell}{2-\ell} (-y)^{2-\ell}} & \text{on } (-\infty, -y_0) \times [0, T], \end{cases}$$

where y_0 is a constant to be determined later, and $\eta_1(t) = \alpha_1(t) e^{\frac{\nu y_0^2}{2-\ell}}$, so that $\bar{\psi} \in C(\mathbb{R} \times [0, T])$ and $(\log \eta_1)' = (\log \alpha_1)'$ independent of y_0 .

By the definition of ν in (2.27), we may assert that for any $y_0 > 0$,

$$(2.33) \quad \partial_y (\log \bar{\psi})(y_0^-, \cdot) = \left[-\frac{2r(\cdot)}{2-\ell} \right] y_0 > [-(r(\cdot) + \nu)] y_0 = \partial_y (\log \bar{\psi})(y_0^+, \cdot),$$

and similarly, $\partial_y \bar{\psi}((-y_0)^-, \cdot) > \partial_y \bar{\psi}((-y_0)^+, \cdot)$. Therefore, in view of (2.30), to verify that $\bar{\psi}$ defined by (2.32) is a super-solution of (2.31), it remains to choose large y_0 such that

$$(2.34) \quad \partial_t \bar{\psi} - \partial_{yy} \bar{\psi} - y \underline{\rho}(t) \partial_y \bar{\psi} \geq \lambda_\ell \bar{\psi} \quad \text{in } (\mathbb{R} \setminus \{\pm y_0\}) \times [0, T],$$

which can be verified by the following computations:

(i) For $y \in (-y_0, y_0)$, by (2.26) and (2.32), direct calculations yield

$$\begin{aligned} & \partial_t \bar{\psi} - \partial_{yy} \bar{\psi} - y \underline{\rho}(t) \partial_y \bar{\psi} \\ &= \left[(\log \alpha_1)' - \frac{\dot{r}(t)}{2-\ell} y^2 + \frac{2r(t)}{2-\ell} - \frac{4r^2(t)}{(2-\ell)^2} y^2 + \frac{2r(t)\underline{\rho}(t)}{2-\ell} y^2 \right] \bar{\psi} \\ &\geq \left[(\log \alpha_1)' + \frac{2r(t)}{2-\ell} \right] \bar{\psi} + \left[-\frac{\dot{r}(t)}{2-\ell} - \frac{4r^2(t)}{(2-\ell)^2} + r(t)\underline{\rho}(t) \right] y^2 \bar{\psi} \\ &\geq \left[(\log \alpha_1)' + \frac{2r(t)}{2-\ell} \right] \bar{\psi} \\ &= \lambda_\ell \bar{\psi}. \end{aligned}$$

(ii) For $y \in (y_0, \infty)$, again by (2.26) and (2.32), we calculate that

$$\begin{aligned} & \partial_t \bar{\psi} - \partial_{yy} \bar{\psi} - y \underline{\rho}(t) \partial_y \bar{\psi} - \lambda_\ell \bar{\psi} \\ &= [(\log \eta_1)' - \lambda_\ell] \bar{\psi} + y_0^\ell y^{2-\ell} \left[-\frac{\dot{r}(t)}{2-\ell} + (1-\ell)(r(t) + \nu) y^{-2} \right] \bar{\psi} \\ &\quad + y_0^\ell y^{2-\ell} [-(r(t) + \nu)^2 y_0^\ell y^{-\ell} + (r(t) + \nu) \underline{\rho}(t)] \bar{\psi} \\ &\geq [(\log \alpha_1)' - \lambda_\ell] \bar{\psi} + y_0^\ell y^{2-\ell} \left[-\frac{\dot{r}(t)}{2-\ell} - (r(t) + \nu)^2 + r(t)\underline{\rho}(t) \right] \bar{\psi} \\ &= [(\log \alpha_1)' - \lambda_\ell] \bar{\psi} + y_0^\ell y^{2-\ell} \left[\left(\frac{4}{(2-\ell)^2} + \frac{\epsilon}{2} \right) r^2(t) - (r(t) + \nu)^2 \right] \bar{\psi}. \end{aligned}$$

In light of $\left(\frac{4}{(2-\ell)^2} + \frac{\epsilon}{2} \right) r^2(t) > (r(t) + \nu)^2$ (due to (2.27)), we may pick y_0 large enough to ensure (2.34) on $(y_0, \infty) \times [0, T]$.

(iii) For $y \in (-\infty, -y_0)$, we can verify (2.34) by the same argument as in (ii).

Consequently, (2.34) holds true, and $\bar{\psi}$ constructed by (2.32) is a super-solution of (2.31) in the sense of Definition A.1.

In what follows, we divide the construction of super-solution $\bar{\varphi}$ which satisfies (2.24) into the following several steps via separating different regions; see Fig. 2 for the profile of $\bar{\varphi}$ to be constructed.




FIGURE 2. The profile of $\bar{\varphi}$ for fixed $t \in [0, T]$

Step 3. We construct super-solution $\bar{\varphi}$ on $[\kappa - \delta, \kappa + \delta] \times [0, T]$ satisfying (2.24). Let $\bar{\psi}$ be given by (2.32) with fixed y_0 chosen in Step 2. We assume $\sqrt{D}y_0 < \delta$, and define \mathbb{X} by

$$(2.35) \quad \mathbb{X} = \bigcup_{n=1}^{n_*} \{\kappa \pm n\delta\} \cup \{\delta, 1 - \delta\} \cup \left\{ \kappa \pm \sqrt{D}y_0 \right\},$$

where n_* is chosen in (2.28). Set

$$(2.36) \quad \bar{\varphi}(x, t) := f_\kappa(t) \cdot \bar{\psi} \left(\frac{x - \kappa}{\sqrt{D}}, t \right) \quad \text{on } [\kappa - \delta, \kappa + \delta] \times [0, T],$$

where $f_\kappa(t)$ is defined by (2.1) with $x = \kappa$. Note that $\bar{\varphi}$ is symmetric in x with respect to $x = \kappa$, and is decreasing in x for $x \geq \kappa$ and $t \in [0, T]$. Thus by (2.25) and (2.36) we arrive at

$$(2.37) \quad \begin{aligned} -\partial_x m \partial_x \bar{\varphi} &= |\partial_x m| \cdot |\partial_x \bar{\varphi}| \geq f_\kappa(t) \underline{\rho}(t) \frac{|x - \kappa|}{\sqrt{D}} \left| \partial_y \bar{\psi} \left(\frac{x - \kappa}{\sqrt{D}}, t \right) \right| \\ &= -f_\kappa(t) \underline{\rho}(t) \cdot y \partial_y \bar{\psi}(y, t), \end{aligned}$$

where $y = \frac{x-\kappa}{\sqrt{D}}$. This implies that on $([\kappa - \delta, \kappa + \delta] \setminus \{\kappa \pm \sqrt{D}y_0\}) \times [0, T]$,

$$\begin{aligned} \mathcal{L}_D \bar{\varphi} &\geq \left[-V(\kappa, t) + \hat{V}(\kappa) + V(x, t) \right] \bar{\varphi} + [\partial_t \bar{\psi} - \partial_{yy} \bar{\psi} - \underline{\rho}(t)y \partial_y \bar{\psi}] f_\kappa(t) \\ &\geq \left[\hat{V}(\kappa) - \epsilon/2 + (1-\epsilon)\hat{\rho} \right] \bar{\varphi} \\ &\geq \left[\hat{V}(\kappa) + (1-\epsilon)\partial_{xx} \hat{m}(\kappa) - \epsilon(1-\epsilon) \right] \bar{\varphi} \\ &\geq (1-\epsilon)(\lambda_{\min} - \epsilon) \bar{\varphi}, \end{aligned}$$

where the first inequality is due to (2.37), the second inequality follows from (2.17) and the fact that $\bar{\psi}$ is a super-solution of (2.31) (see Step 2), and the third inequality follows from (2.25).

On the other hand, by (2.33), we have

$$\partial_x(\log \bar{\varphi})((\kappa + \sqrt{D}y_0)^+, \cdot) < \partial_x(\log \bar{\varphi})((\kappa + \sqrt{D}y_0)^-, \cdot) \quad (\text{as } \kappa + \sqrt{D}y_0 \in \mathbb{X}).$$

Therefore, $\bar{\varphi}$ defined by (2.36) satisfies (2.24) on $[\kappa - \delta, \kappa + \delta] \times [0, T]$.

Step 4. We construct $\bar{\varphi}$ which satisfies (2.24) on $(\kappa + \delta, \kappa + 2\delta] \times [0, T]$. Since $\sqrt{D}y_0 < \delta$, by (2.36) in Step 3 and (2.32) in Step 2, we have

$$(2.38) \quad \begin{cases} \log \bar{\varphi}(\kappa + \delta, t) = \log f_\kappa(t) + \log \eta_1(t) - \frac{(r(t) + \nu)y_0^\ell \delta^{2-\ell}}{(2-\ell)D^{1-\ell/2}}, \\ \partial_x(\log \bar{\varphi})((\kappa + \delta)^-, \cdot) = -(r(\cdot) + \nu) \frac{y_0^\ell \delta^{1-\ell}}{D^{1-\ell/2}}, \end{cases}$$

whence there is some constant $K_0 > 0$ such that

$$(2.39) \quad |\partial_t(\log \bar{\varphi})(\kappa + \delta, \cdot)| = \left| (\log f_\kappa)' + (\log \eta_1)' - \frac{\dot{r}(t)y_0^\ell \delta^{2-\ell}}{(2-\ell)D^{1-\ell/2}} \right| < \frac{K_0}{D^{1-\ell/2}}.$$

We introduce a small parameter $\epsilon_0 > 0$ such that

$$|\partial_x m| \geq \epsilon_0 \quad \text{on } ([\delta, \kappa - \delta] \cup [\kappa + \delta, 1 - \delta]) \times [0, T],$$

and fix constant K_1 so that

$$K_1 > (\bar{r} + \nu)y_0^\ell \delta^{1-\ell} + 2K_0/\epsilon_0.$$

Then we define

$$(2.40) \quad \bar{\varphi}(x, t) := \zeta_1(x, t) \cdot e^{-\frac{K_1(x-\kappa)}{D^{1-\ell/2}}} \quad \text{on } (\kappa + \delta, \kappa + 2\delta] \times [0, T].$$

Here $\zeta_1 \in C^{2,1}((\kappa + \delta, \kappa + 2\delta) \times [0, T])$ is determined by

(2.41)

$$\log \zeta_1(x, t) = \left[\frac{(\kappa+2\delta)-x}{\delta} \right] \cdot \left[\frac{K_1 \delta}{D^{1-\ell/2}} + \log \bar{\varphi}(\kappa + \delta, t) \right] + \left[\frac{x-(\kappa+\delta)}{\delta} \right] \log f_1(t),$$

with T -periodic function $f_1(t)$ defined in (2.1) with $x = 1$, so that

$$\zeta_1(\kappa + \delta, t) = e^{\frac{K_1 \delta}{D^{1-\ell/2}}} \cdot \bar{\varphi}(\kappa + \delta, t).$$

This implies immediately that $\bar{\varphi}$ defined by (2.40) is continuous at $\{\kappa + \delta\} \times [0, T]$. In light of $\partial_x \zeta_1 < 0$ (for small D), using (2.38) and (2.40), by choice of K_1 we can verify that

$$\partial_x(\log \bar{\varphi})((\kappa + \delta)^+, \cdot) < -K_1/D^{1-\ell/2} < \partial_x(\log \bar{\varphi})((\kappa + \delta)^-, \cdot) \quad (\text{as } \kappa + \delta \in \mathbb{X}).$$

On the other hand, combined with (2.38), (2.39), and (2.41), we see that

$$|\partial_t(\log \zeta_1)| < \frac{2K_0}{D^{1-\ell/2}}, \quad -\frac{3K_1}{D^{1-\ell/2}} < \partial_x(\log \zeta_1) < 0, \quad \text{and} \quad \partial_{xx}(\log \zeta_1) = 0$$

for small D , and thus

$$\left| \frac{\partial_{xx}\zeta_1}{\zeta_1} \right| = \left| \partial_{xx}(\log \zeta_1) + [\partial_x(\log \zeta_1)]^2 \right| < \frac{9K_1^2}{D^{2-\ell}},$$

from which, using (2.40) and $-\partial_x m \cdot \partial_x(\log \zeta_1) \geq 0$, we may calculate that

$$\begin{aligned} \mathcal{L}_D \bar{\varphi} &= \left\{ \partial_t(\log \zeta_1) - D \left[\frac{\partial_{xx}\zeta_1}{\zeta_1} - \frac{2K_1}{D^{1-\ell/2}} \partial_x(\log \zeta_1) + \frac{K_1^2}{D^{2-\ell}} \right] \right\} \bar{\varphi} \\ &\quad + \left\{ -\partial_x m \cdot \left[\partial_x(\log \zeta_1) - \frac{K_1}{D^{1-\ell/2}} \right] + V \right\} \bar{\varphi} \\ &\geq \left[-\frac{2K_0}{D^{1-\ell/2}} - \frac{16K_1^2}{D^{1-\ell}} + \frac{\epsilon_0 K_1}{D^{1-\ell/2}} + V \right] \bar{\varphi} \\ &= \frac{1}{D^{1-\ell/2}} \left[-2K_0 + \epsilon_0 K_1 - 16K_1^2 D^{\ell/2} + D^{1-\ell/2} V \right] \bar{\varphi}. \end{aligned}$$

Since $\epsilon_0 K_1 > 2K_0$ (by the definition of K_1), we may choose D small such that (2.24) holds. Step 4 is thereby completed.

Step 5. We construct $\bar{\varphi}$ on $(\kappa + 2\delta, \kappa + 3\delta] \times [0, T]$. By (2.40) and (2.41) in Step 4, we have

$$(2.42) \quad \log \bar{\varphi}(\kappa + 2\delta, t) = \log f_1(t) - \frac{2K_1\delta}{D^{1-\ell/2}} \quad \text{and} \quad \partial_x(\log \bar{\varphi})((\kappa + 2\delta)^-, \cdot) > -\frac{4K_1}{D^{1-\ell/2}}.$$

Fix a constant K_2 such that $K_2 > 16K_1^2/\epsilon_0$, where ϵ_0 is given in Step 4 such that $\partial_x m \geq \epsilon_0$ on $[\kappa + \delta, 1 - \delta] \times [0, T]$. Define

$$(2.43) \quad \bar{\varphi}(x, t) := f_1(t) \cdot \bar{\phi}_2(x) \quad \text{on } (\kappa + 2\delta, \kappa + 3\delta] \times [0, T],$$

where $\bar{\phi}_2$ solves

$$(2.44) \quad \begin{cases} (\log \bar{\phi}_2)'(x) = -\frac{4K_1}{D^{1-\ell/2}} \left[\frac{\kappa+3\delta-x}{\delta} \right] - \frac{K_2}{D^{1-\ell}} \left[\frac{x-(\kappa+2\delta)}{\delta} \right] & \text{in } (\kappa + 2\delta, \kappa + 3\delta], \\ \log \bar{\phi}_2(\kappa + 2\delta) = -\frac{2K_1\delta}{D^{1-\ell/2}}. \end{cases}$$

Together (2.43) with (2.42) and (2.44), we discover that $\bar{\varphi}$ is continuous at $\{\kappa + 2\delta\} \times [0, T]$, and

$$\partial_x(\log \bar{\varphi})((\kappa + 2\delta)^+, \cdot) = -4K_1/D^{1-\ell/2} < \partial_x(\log \bar{\varphi})((\kappa + 2\delta)^-, \cdot) \quad (\text{as } \kappa + 2\delta \in \mathbb{X}).$$

For all $x \in (\kappa + 2\delta, \kappa + 3\delta]$, by (2.44) we have

$$\left| \frac{\bar{\phi}_2''}{\bar{\phi}_2} \right| = \left| (\log \bar{\phi}_2)'' + [(\log \bar{\phi}_2)']^2 \right| \leq \frac{4K_1}{\delta D^{1-\ell/2}} + \frac{16K_1^2}{D^{2-\ell}},$$

from which we arrive at

$$\begin{aligned} \mathcal{L}_D \bar{\varphi} &= \left[(\log f_1)' - D \bar{\phi}_2''/\bar{\phi}_2 - \partial_x m \cdot (\log \bar{\phi}_2)' + V \right] \bar{\varphi} \\ &\geq \left[(\log f_1)' - \frac{4K_1}{\delta} D^{\ell/2} - \frac{16K_1^2}{D^{1-\ell}} + \frac{\epsilon_0 K_2}{D^{1-\ell}} + V \right] \bar{\varphi}. \end{aligned}$$

In view of $\epsilon_0 K_2 > 16K_1^2$, we once more would select D small enough such that (2.24) holds.

Step 6. We construct $\bar{\varphi}$ on $(\kappa + 3\delta, \kappa + (n_* + 1)\delta] \times [0, T]$, where n_* is determined by (2.28) in Step 1. By the definition of $\bar{\phi}_2$ in (2.44), we have

$$(2.45) \quad \partial_x(\log \bar{\varphi})((\kappa + 3\delta)^-, \cdot) = (\log \bar{\phi}_2)'(\kappa + 3\delta) = -K_2/D^{1-\ell}.$$

We introduce a sequence $\{K_n\}_{n=3}^{n_*}$ independent of D such that $K_n > K_{n-1}^2/\epsilon_0$. With $\bar{\phi}_2$ in hand, by induction we define $\bar{\phi}_n \in C^{2,1}([\kappa + n\delta, \kappa + (n+1)\delta])$ ($n = 3, \dots, n_*$) to solve

$$(2.46) \quad \begin{cases} (\log \bar{\phi}_n)'(x) = -\left[\frac{K_{n-1}}{D^{1-2^{n-3}\ell}} + \epsilon\right] \cdot \left[\frac{\kappa + n\delta - x}{\delta}\right] - \frac{K_n}{D^{1-2^{n-2}\ell}} \left[\frac{x - (\kappa + n\delta)}{\delta}\right] & \text{in } (\kappa + n\delta, \kappa + (n+1)\delta], \\ \log \bar{\phi}_n(\kappa + n\delta) = \log \bar{\phi}_{n-1}(\kappa + n\delta). \end{cases}$$

Then we define

$$(2.47) \quad \bar{\varphi}(x, t) := f_1(t) \cdot \bar{\phi}_n(x) \quad \text{on } (\kappa + n\delta, \kappa + (n+1)\delta] \times [0, T].$$

By (2.45) and (2.46), it can be verified that

$$\begin{aligned} \partial_x(\log \bar{\varphi})((\kappa + 3\delta)^+, \cdot) &= \partial_x(\log \bar{\phi}_3)(\kappa + 3\delta) \\ &= -\left[\frac{K_2}{D^{1-\ell}} + \epsilon\right] < \partial_x(\log \bar{\varphi})((\kappa + 3\delta)^-, \cdot), \end{aligned}$$

and similarly for $4 \leq n \leq n_*$,

$$\partial_x(\log \bar{\varphi})((\kappa + n\delta)^+, \cdot) < \partial_x(\log \bar{\varphi})((\kappa + n\delta)^-, \cdot) \quad (\text{as } \kappa + n\delta \in \mathbb{X}).$$

For each $3 \leq n \leq n_*$, it follows from (2.46) that for $x \in (\kappa + n\delta, \kappa + (n+1)\delta]$

$$(2.48) \quad -\left[\frac{K_{n-1}}{D^{1-2^{n-3}\ell}} + \epsilon\right] \leq (\log \bar{\phi}_n)' \leq -\frac{K_n}{D^{1-2^{n-2}\ell}},$$

and then as in Step 5, we derive that

$$(2.49) \quad \left|\frac{\bar{\phi}_n''}{\bar{\phi}_n}\right| = \left|(\log \bar{\phi}_n)'' + [(\log \bar{\phi}_n)']^2\right| \leq \frac{2K_{n-1}}{\delta D^{1-2^{n-3}\ell}} + \frac{K_{n-1}^2}{D^{2-2^{n-2}\ell}}.$$

By (2.48) and (2.49), on $(\kappa + n\delta, \kappa + (n+1)\delta] \times [0, T]$, we calculate that

$$\begin{aligned} \mathcal{L}_D \bar{\varphi} &= \left[(\log f_1)' - D\bar{\phi}_n''/\bar{\phi}_n - \partial_x m \cdot (\log \bar{\phi}_n)' + V\right] \bar{\varphi} \\ &\geq \left[(\log f_1)' - \frac{2K_{n-1}}{\delta} D^{2^{n-3}\ell} - \frac{K_{n-1}^2}{D^{1-2^{n-2}\ell}} + \frac{\epsilon_0 K_n}{D^{1-2^{n-2}\ell}} + V\right] \bar{\varphi}. \end{aligned}$$

Since $\epsilon_0 K_n > K_{n-1}^2$, we choose D to be small so that $\bar{\varphi}$ satisfies (2.24).

Step 7. We construct $\bar{\varphi}$ on $(\kappa + (n_* + 1)\delta, 1] \times [0, T]$. Set $\kappa^* = \kappa + (n_* + 1)\delta$. Observe from Step 6 and the definition of n_* in (2.28) that

$$\partial_x(\log \bar{\varphi})((\kappa^*)^-, \cdot) = -K_{n_*}/D^{1-2^{n_*-2}\ell} = -K_{n_*}.$$

We define

$$(2.50) \quad \bar{\varphi}(x, t) := f_1(t) \bar{\phi}_{n_*}(\kappa^*) \cdot \begin{cases} e^{-K_*(x-\kappa^*)} & \text{on } (\kappa^*, 1-\delta] \times [0, T], \\ e^{\frac{\kappa_*+\epsilon}{2\delta}(1-x)^2+\theta_1} & \text{on } (1-\delta, 1] \times [0, T], \end{cases}$$

where $K_* > K_{n_*}$ will be determined later, and the parameter θ_1 is chosen such that $\bar{\varphi}$ is continuous at $\{1 - \delta\} \times [0, T]$. It follows that

$$\partial_x(\log \bar{\varphi})(x^+, \cdot) < \partial_x(\log \bar{\varphi})(x^-, \cdot) \quad \text{for } x \in \{\kappa^*, 1 - \delta\} \subset \mathbb{X}.$$

It remains to verify that $\bar{\varphi}$ defined by (2.50) satisfies (2.24). For $x \in (\kappa^*, 1 - \delta]$, since $\partial_x m \geq \epsilon_0$, using (2.50) we deduce that

$$\mathcal{L}_D \bar{\varphi} \geq [(\log f_1)' - DK_*^2 + \epsilon_0 K_* + V] \bar{\varphi}.$$

By choosing K_* large and then choosing D small, we see that $\bar{\varphi}$ satisfies (2.24).

For $x \in (1 - \delta, 1]$, since $-\partial_x m \partial_x \bar{\varphi} \geq 0$, by (2.50), letting D be so small that

$$\begin{aligned} \mathcal{L}_D \bar{\varphi} &\geq \left\{ (\log f_1)' - D(K_* + \epsilon)/\delta \cdot \left[(K_* + \epsilon)(x - 1)^2/\delta + 1 \right] + V \right\} \bar{\varphi} \\ &\geq \left[\hat{V}(1) - V(1, t) + V(x, t) - \epsilon/2 \right] \bar{\varphi} \\ &\geq (\lambda_{\min} - \epsilon) \bar{\varphi}, \end{aligned}$$

where the last inequality is due to (2.17).

By Steps 3-7, we have already constructed the strict super-solution $\bar{\varphi}$ satisfying (2.24) on $[\kappa - \delta, 1] \times [0, T]$ with the set \mathbb{X} given by (2.35), which is summarized in the following table for the convenience of readers; see also Fig. 2.

Construction of $\bar{\varphi}$ on $[\kappa - \delta, 1] \times [0, T]$		
$\bar{\varphi}(x, t)$	Region	Defined in
$f_\kappa(t) \cdot \bar{\psi}\left(\frac{x - \kappa}{\sqrt{D}}, t\right)$	$[\kappa - \delta, \kappa + \delta] \times [0, T]$	(2.36) in Step 3
$\zeta_1(x, t) \cdot e^{-\frac{K_1(x - \kappa)}{D^{1-\ell/2}}}$	$[\kappa + \delta, \kappa + 2\delta] \times [0, T]$	(2.40) and (2.41) in Step 4
$f_1(t) \cdot \bar{\phi}_n(x)$	$(\kappa + n\delta, \kappa + (n + 1)\delta] \times [0, T]$ ($n = 2, \dots, n_*$)	(2.46) and (2.47) in Steps 5 and 6
$f_1(t) \bar{\phi}_{n_*}(\kappa^*) \cdot e^{-K_*(x - \kappa^*)}$	$(\kappa^*, 1 - \delta] \times [0, T]$	(2.50) in Step 7
$f_1(t) \bar{\phi}_{n_*}(\kappa^*) \cdot e^{\frac{K_* + \epsilon}{2\delta}(1-x)^2 + \theta_1}$	$(1 - \delta, 1] \times [0, T]$	(2.50) in Step 7

Finally, we construct $\bar{\varphi}$ on $[0, \kappa - \delta) \times [0, T]$ symmetrically; and precisely, define (2.51)

$$\bar{\varphi}(x, t) = \begin{cases} \zeta_2(x, t) \cdot e^{\frac{K_1(\kappa - x)}{D^{1-\ell/2}}} & \text{on } [\kappa - 2\delta, \kappa - \delta) \times [0, T], \\ f_0(t) \cdot \bar{\phi}_n(2\kappa - x) & \text{on } \begin{cases} [\kappa - (n + 1)\delta, \kappa - n\delta) \times [0, T], \\ (n = 2, \dots, n_*), \end{cases} \\ f_0(t) \bar{\phi}_{n_*}(\kappa_*) \cdot e^{K_*(\kappa_* - x)} & \text{on } [\delta, \kappa_*) \times [0, T], \\ f_0(t) \bar{\phi}_{n_*}(\kappa_*) \cdot e^{\frac{K_* + \epsilon}{2\delta}x^2 + \theta_2} & \text{on } [0, \delta) \times [0, T], \end{cases}$$

where $\kappa_* = \kappa - (n_* + 1)\delta$, and similar to (2.41), ζ_2 solves

$$\log \zeta_2(x, t) = \left[\frac{x - (\kappa - 2\delta)}{\delta} \right] \cdot \left[\frac{K_1 \delta}{D^{1-\ell/2}} + \log \bar{\varphi}(\kappa - \delta, t) \right] + \left[\frac{(\kappa - \delta) - x}{\delta} \right] \log f_0(t),$$

with f_0 defined in (2.1) with $x = 0$, and θ_2 is chosen such that $\bar{\varphi}$ is continuous at $\{\delta\} \times [0, T]$. Using the same arguments as in Steps 4-7, we may conclude that $\bar{\varphi}$ defined by (2.51) verifies (2.24), and thus $\bar{\varphi}$ constructed above defines a super-solution on the entire region $[0, 1] \times [0, T]$ with \mathbb{X} given by (2.35). Therefore, (2.23) follows from Proposition A.2. \square

By assuming $\partial_x m(0, t) > 0$ and $\partial_x m(1, t) < 0$ for each $t \in [0, T]$, it is shown in Proposition 2.1 that the limit of $\lambda(D)$ as $D \rightarrow 0$ does not depend upon the value of V on boundary points $\{0, 1\} \times [0, T]$. However, without the positivity assumption of $\partial_x m(0, t)$, one can prove

Lemma 2.4. *Suppose that $\partial_x m(x, t) > 0$ for all $(x, t) \in (0, 1) \times [0, T]$. Then*

$$\lim_{D \rightarrow 0} \lambda(D) = \min \left\{ \hat{V}(0) + [\partial_{xx} \hat{m}]_+(0), \hat{V}(1) \right\},$$

where $\partial_{xx} \hat{m}(0)$ is defined by (1.2).

Proof. If $\partial_x m(0, t) = 0$ for all $t \in [0, T]$, Lemma 2.4 can be proved directly by constructing the same super/sub-solutions as those in Proposition 2.3 defined on $[\kappa, 1] \times [0, T]$. It suffices to consider the remaining case $\partial_x \hat{m}(0) > 0$ and in view of $\partial_{xx} \hat{m}_+(0) = \infty$ in this case, i.e. to show

$$\lim_{D \rightarrow 0} \lambda(D) = \hat{V}(1).$$

First, similarly as in the proof of Proposition 2.1, we may construct a sub-solution to prove $\limsup_{D \rightarrow 0} \lambda(D) \leq \hat{V}(1)$. In the sequel, we show

$$(2.52) \quad \liminf_{D \rightarrow 0} \lambda(D) \geq \hat{V}(1).$$

For any given $\epsilon > 0$, we fix some small $\delta > 0$ such that

$$|V(x, t) - V(1, t)| < \epsilon/2 \quad \text{on } [1 - \delta, 1] \times [0, T].$$

The strategy is to construct a positive super-solution $\bar{\varphi} \in C^{2,1}([0, 1] \times [0, T])$, which satisfies

$$(2.53) \quad \begin{cases} \mathcal{L}_D \bar{\varphi} \geq [\hat{V}(1) - \epsilon] \bar{\varphi} & \text{in } (0, 1) \times [0, T], \\ \partial_x \bar{\varphi}(0, t) < 0, \quad \partial_x \bar{\varphi}(1, t) = 0 & \text{on } [0, T], \\ \bar{\varphi}(x, 0) = \bar{\varphi}(x, T) & \text{on } (0, 1) \end{cases}$$

for sufficiently small D . To this end, we proceed as follows:

On $[1 - \delta, 1] \times [0, T]$, we define

$$\bar{\varphi}(x, t) := f_1(t) \cdot e^{\frac{M_2}{2\delta}(1-x)^2} \quad \text{on } [1 - \delta, 1] \times [0, T],$$

where $M_2 > 0$ will be determined later, and $f_1(t)$ is given by (2.1) with $x = 1$. Similar to Step 2 in Proposition 2.1, one can verify that (2.53) holds on $[1 - \delta, 1] \times [0, T]$.

On $[0, \delta] \times [0, T]$, since $\partial_x m(0, t) \geq (\not\equiv) 0$ for $t \in [0, T]$ (due to $\partial_x \hat{m}(0) > 0$), one can find some $t_0 \in (0, T)$ and positive constants ϵ_0, δ_0 such that

$$\partial_x m(x, t) > \epsilon_0 \quad \text{for any } (x, t) \in [0, \delta] \times [t_0 - \delta_0, t_0 + \delta_0].$$

Fix $\eta_2 \in C^\infty([0, T])$ to be a positive T -periodic function such that

$$(2.54) \quad (\log \eta_2(t))' > \|V(\cdot, t)\|_{L^\infty} + |\hat{V}(1)| \quad \text{for } t \in [0, t_0 - \delta] \cup [t_0 + \delta, T].$$

We then define, for $(x, t) \in [0, \delta] \times [0, T]$,

$$\bar{\varphi}(x, t) := \eta_2(t) \cdot e^{-M_2 x}.$$

On $[0, \delta] \times [t_0 - \delta_0, t_0 + \delta_0]$, since $\partial_x m(x, t) > \epsilon_0$, by straightforward computations we deduce

$$\mathcal{L}_D \bar{\varphi} \geq \left[(\log \eta_2)' - DM_2^2 + M_2 \epsilon_0 - V \right] \bar{\varphi},$$

whence by choosing M_2 large and then choosing D small, we have $\mathcal{L}_D \bar{\varphi} \geq \hat{V}(1) \bar{\varphi}$.

On the other hand, on $[0, \delta] \times ([0, t_0 - \delta] \cup [t_0 + \delta, T])$, in view of (2.54) and $-\partial_x m \partial_x \bar{\varphi} \geq 0$, by letting D be small, we arrive at

$$\mathcal{L}_D \bar{\varphi} \geq \left[(\log \eta_2)' - DM_2^2 - V \right] \bar{\varphi} \geq \hat{V}(1) \bar{\varphi},$$

whence (2.53) is verified on $[0, \delta] \times [0, T]$.

On $(\delta, 1 - \delta) \times [0, T]$, notice from the definition of $\bar{\varphi}$ above that

$$\partial_x (\log \bar{\varphi})(\delta) = \partial_x (\log \bar{\varphi})(1 - \delta) = -M_2.$$

We can always find $\bar{\varphi} \in C^{2,1}([\delta, 1 - \delta] \times [0, T])$ such that $\bar{\varphi}(\cdot, 0) = \bar{\varphi}(\cdot, T)$ and

$$\partial_x \log \bar{\varphi} \leq -M_2 \quad \text{and} \quad |\partial_t \log \bar{\varphi}| \leq 2 \|\|(\log f_1)'| + |(\log \eta_2)'|\|_{L^\infty},$$

and then (2.53) can be verified directly by further choosing M_2 large and D small.

Therefore, such a super-solution $\bar{\varphi}$ defined above satisfies (2.53), and Proposition A.2 concludes the proof. \square

Corollary 2.5. *Assume $V(x, t) = V(x)$ and $\partial_x m(x, t) = m'(x)$. Suppose that $m'(x) > 0$ for all $x \in (0, 1)$. Then we have*

$$\lim_{D \rightarrow 0} \lambda(D) = \min \{V(0) + [m'']_+(0), V(1)\}.$$

Remark 2.6. Corollary 2.5 cannot be covered by Theorem 1.1. It also provides an example such that Theorem 1.2 in [6] fails without the assumption $|\nabla m| \neq 0$ on $\partial\Omega$ therein.

To establish Theorem 1.3, we prepare the following

Lemma 2.7. *Given any $0 \leq \underline{\kappa} < \bar{\kappa} \leq 1$, let $\lambda(D)$ be the principal eigenvalue of the problem*

$$(2.55) \quad \begin{cases} \partial_t \varphi - D \partial_{xx} \varphi + V \varphi = \lambda(D) \varphi & \text{in } (\underline{\kappa}, \bar{\kappa}) \times [0, T], \\ c_1 \partial_x \varphi(\underline{\kappa}, t) - (1 - c_1) \varphi(\underline{\kappa}, t) = 0 & \text{on } [0, T], \\ c_2 \partial_x \varphi(\bar{\kappa}, t) + (1 - c_2) \varphi(\bar{\kappa}, t) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } [\underline{\kappa}, \bar{\kappa}], \end{cases}$$

where $c_1, c_2 \in [0, 1]$. Then we have

$$\lim_{D \rightarrow 0} \lambda(D) = \min_{x \in [\underline{\kappa}, \bar{\kappa}]} \hat{V}(x).$$

Remark 2.8. Lemma 2.7 is proved in Lemma 2.4(c) of [14] for the case $c_1 = c_2 = 1$.

Proof of Lemma 2.7. For the upper bound, it suffices to claim that

$$\limsup_{D \rightarrow 0} \lambda(D) \leq V(\tilde{x}) \quad \text{for any } \tilde{x} \in (\underline{\kappa}, \bar{\kappa}).$$

Indeed, we follow the ideas as in Proposition 2.1 and define a sub-solution

$$(2.56) \quad \underline{\varphi}(x, t) := f_{\tilde{x}}(t) \cdot \tilde{z}(x)$$

with $f_{\tilde{x}}(t)$ defined in (2.1) with $x = \tilde{x}$ and

$$\tilde{z}(x) = \begin{cases} \cos\left(\frac{\pi}{2\tilde{\delta}}(x - \tilde{x})\right) & \text{on } [\tilde{x} - \tilde{\delta}, \tilde{x} + \tilde{\delta}], \\ 0 & \text{on } [0, \tilde{x} - \tilde{\delta}) \cup (\tilde{x} + \tilde{\delta}, 1]. \end{cases}$$

Here $\tilde{\delta}$ is chosen such that $|V(x, t) - V(\tilde{x}, t)| < \epsilon/2$ in $[\tilde{x} - \tilde{\delta}, \tilde{x} + \tilde{\delta}] \times [0, T]$ for any given $\epsilon > 0$. One may verify readily that

$$\mathcal{L}_D \underline{\varphi} \leq [\hat{V}(\tilde{x}) + \epsilon] \underline{\varphi},$$

so that the upper bound follows from Proposition A.2.

It remains to prove

$$(2.57) \quad \liminf_{D \rightarrow 0} \lambda(D) \geq \min_{x \in [\underline{\kappa}, \bar{\kappa}]} \hat{V}(x).$$

For any $\epsilon > 0$, choose some T -periodic function $V_\epsilon \in C^{2,1}([\underline{\kappa}, \bar{\kappa}] \times [0, T])$ such that

$$\|V_\epsilon - V\|_{L^\infty([0, 1] \times [0, T])} \leq \epsilon.$$

Then we define T -periodic function φ_ϵ by

$$(2.58) \quad \varphi_\epsilon(x, t) := \exp \left[- \int_0^t V_\epsilon(x, s) ds + t \hat{V}_\epsilon(x) \right] \beta_\epsilon(x),$$

where $\beta_\epsilon \in C^2([\underline{\kappa}, \bar{\kappa}])$ is a positive function and is chosen such that

$$(2.59) \quad c_1 \partial_x \varphi_\epsilon(\underline{\kappa}, t) - (1 - c_1) \varphi_\epsilon(\underline{\kappa}, t) \leq 0 \quad \text{and} \quad c_2 \partial_x \varphi_\epsilon(\bar{\kappa}, t) + (1 - c_2) \varphi_\epsilon(\bar{\kappa}, t) \geq 0.$$

By (2.58) and the definition of V_ϵ , we may choose D small to derive that

$$\begin{aligned} \partial_t \varphi_\epsilon - D \partial_{xx} \varphi_\epsilon + V \varphi_\epsilon &= \left[\hat{V}_\epsilon(x) - V_\epsilon(x, t) + V(x, t) \right] \varphi_\epsilon - D \partial_{xx} \varphi_\epsilon \\ &\geq \left[\min_{x \in [\underline{\kappa}, \bar{\kappa}]} \hat{V}(x) - 3\epsilon \right] \varphi_\epsilon, \end{aligned}$$

which together with (2.59) implies that φ_ϵ defined by (2.58) is a super-solution in the sense of Definition A.1 with $\mathbb{X} = \emptyset$. Thus (2.57) follows from Proposition A.2, and the proof of Lemma 2.7 is now complete. \square

We are now in a position to prove Theorem 1.3.

Proof of Theorem 1.3. The proof can be carried out by the same ideas as in Propositions 2.1 and 2.3 with the help of Lemmas 2.4 and 2.7. Here we just outline it for completeness.

Step 1. We establish the upper bound of $\limsup_{D \rightarrow 0} \lambda(D)$. First, using a similar argument as in Lemma 2.7, one can establish

$$\limsup_{D \rightarrow 0} \lambda(D) \leq \min_{i \in \mathbf{B}} \left\{ \min_{x \in [\kappa_i, \kappa_{i+1}]} \hat{V}(x) \right\}$$

by constructing a suitable sub-solution like (2.56). Similarly, the estimate

$$\limsup_{D \rightarrow 0} \lambda(D) \leq \min \left\{ \hat{V}(0) + [\partial_{xx} \hat{m}]_+(0), \hat{V}(1) + [\partial_{xx} \hat{m}]_+(1) \right\}$$

can also be proved; the details are omitted here. It remains to show

$$(2.60) \quad \limsup_{D \rightarrow 0} \lambda(D) \leq \hat{V}(\kappa_i) + [\partial_{xx} \hat{m}]_+(\kappa_i) \quad \text{for all } 1 \leq i \leq N.$$

Fix any $\epsilon > 0$. Choose some small $\delta > 0$ such that $|V(x, t) - V(\kappa_i, t)| < \epsilon/2$ on $[\kappa_i - \delta, \kappa_i + \delta] \times [0, T]$ for all $1 \leq i \leq N$. To prove (2.60), we define

$$\underline{\varphi}_i(x, t) := \begin{cases} f_{\kappa_i}(t) \cdot \underline{z}(x) & \text{if } \partial_{xx}\hat{m}(\kappa_i) \leq 0, \\ f_{\kappa_i}(t) \cdot \underline{\psi}_D(x, t) & \text{if } \partial_{xx}\hat{m}(\kappa_i) > 0, \end{cases}$$

where f_{κ_i} and \underline{z} are defined respectively by (2.1) and (2.5), and $\underline{\psi}_D$ denotes the principal eigenfunction of (2.21) with $\kappa = \kappa_i$. The same arguments as in Step 1 of Propositions 2.1 and 2.3 allow us to verify that such a function $\underline{\varphi}_i$ defines a sub-solution in the sense of Definition A.1 such that for sufficiently small D ,

$$\begin{cases} \mathcal{L}_D \underline{\varphi}_i \leq [\hat{V}(\kappa_i) + [\partial_{xx}\hat{m}]_+(\kappa_i) + 2\epsilon] \underline{\varphi}_i & \text{in } ((0, 1) \setminus \{\kappa_i \pm \delta\}) \times [0, T], \\ \partial_x \underline{\varphi}_i(0, t) = \partial_x \underline{\varphi}_i(1, t) = 0 & \text{on } [0, T], \\ \underline{\varphi}_i(x, 0) = \underline{\varphi}_i(x, T) & \text{on } (0, 1). \end{cases}$$

Then (2.60) is a direct consequence of Proposition A.2.

Step 2. We establish the lower bound of $\liminf_{D \rightarrow 0} \lambda(D)$. It suffices to find a super-solution $\bar{\varphi} \in C([0, 1] \times [0, T])$ satisfying (2.24) with λ_{\min} being replaced by the right hand side of (1.5) and \mathbb{X} will be determined later. Recall the sets \mathbf{A} and \mathbf{B} defined in the statement of Theorem 1.3. The construction of $\bar{\varphi}$ can be given as follows; see Fig. 3 for an illustrated example.




FIGURE 3. The black solid curve corresponds to an example of m for fixed t . The super-solution $\bar{\varphi}$ is constructed respectively on different regions (i)–(v).

(i) On $([\kappa_i - \delta, \kappa_{i+1} + \delta] \cap [0, 1]) \times [0, T]$ for $0 \leq i \leq N$ and $i \in \mathbf{B}$ with the small constant $\delta > 0$ to be determined later, we define $\bar{\varphi}$ as in the form of (2.58) in Lemma 2.7 with

$$\underline{\kappa} = \kappa_i - \delta, \quad \bar{\kappa} = \kappa_{i+1} + \delta, \quad \text{and} \quad c_1 = c_2 = \frac{1}{2}.$$

(ii) On $([0, \frac{\kappa_1}{3}] \cup [\frac{2+\kappa_N}{3}, 1]) \times [0, T]$, if $0 \notin \mathbf{B}$ or $N \notin \mathbf{B}$, then such a super-solution $\bar{\varphi}$ can be constructed by adapting the same arguments as in the proof of Lemma 2.4; Otherwise, it has been constructed in (i).

(iii) On $[\frac{\kappa_{i-1}+2\kappa_i}{3}, \frac{2\kappa_i+\kappa_{i+1}}{3}] \times [0, T]$ for $i \in \mathbf{A}$ and $i-1 \in \mathbf{A}$, one constructs $\bar{\varphi}$ by Step 2 of Proposition 2.1 (with $\kappa = \kappa_i$) for the case $\partial_{xx}\hat{m}(\kappa_i) \leq 0$, and by Part II of Proposition 2.3 (with $\kappa = \kappa_i$) for the case $\partial_{xx}\hat{m}(\kappa_i) > 0$.

(iv) On the remaining region $\Omega \times [0, T]$, where

$$\Omega = \begin{cases} \left(\frac{2\kappa_i+\kappa_{i+1}}{3}, \frac{\kappa_i+2\kappa_{i+1}}{3}\right) & \text{for } i \in \mathbf{A} \text{ and } i-1 \in \mathbf{A}, \\ \left(\kappa_{i-1} + \delta, \frac{2\kappa_i+\kappa_{i+1}}{3}\right) & \text{for } i \in \mathbf{A} \text{ and } i-1 \in \mathbf{B}, \\ \left(\frac{\kappa_{i-1}+2\kappa_i}{3}, \kappa_i - \delta\right) & \text{for } i \in \mathbf{B}, \end{cases}$$

we construct $\bar{\varphi}$ by monotonically connecting the endpoints on $\partial\Omega$, such that

- (a) $\bar{\varphi}$ is continuous at $\partial\Omega \times [0, T]$;
- (b) $|\partial_x(\log \bar{\varphi})| > M_3$ for some large M_3 ;
- (c) $\partial_x(\log \bar{\varphi})(x^+, \cdot) < \partial_x(\log \bar{\varphi})(x^-, \cdot)$ for $x \in \partial\Omega$.

Define $\mathbb{X} = \partial\Omega$. By Lemmas 2.4 and 2.7, explicit calculations as in Propositions 2.1 and 2.3 imply that we may choose δ smaller if necessary such that the super-solution $\bar{\varphi}$ defined above satisfies (2.24) with λ_{\min} being replaced by the right hand side of (1.5). Then the lower bound of $\liminf_{D \rightarrow 0} \lambda(D)$ can be established by Proposition A.2. The proof is now complete. \square

3. PROOF OF THEOREM 1.2

In this section, we study the case when the ODE (1.4) possesses finitely many periodic solutions and establish Theorem 1.2 with the help of Theorem 1.3.

Proof of Theorem 1.2. We first prove part (i) of Theorem 1.2. Let $\{\kappa_i\}_{0 \leq i \leq N+1}$ be any strictly increasing sequence such that

$$0 = \kappa_0 < \kappa_1 < \dots < \kappa_N < \kappa_{N+1} = 1.$$

Fix small δ such that $0 < \delta < \min_{0 \leq i \leq N} (\kappa_{i+1} - \kappa_i)/3$ and

$$(3.1) \quad \partial_{xx}m(x, t) \neq 0 \quad \text{for all } x \in [P_i(t) - \delta, P_i(t) + \delta], t \in [0, T], 1 \leq i \leq N.$$

To “straighten the periodic solution $P_i(t)$ ”, we first define a $C^{2,1}$ -diffeomorphism $\Psi : [0, 1] \times [0, T] \rightarrow [0, 1]$ such that $\partial_y \Psi(y, t) \neq 0$ and

$$(3.2) \quad \Psi(y, t) = \begin{cases} y - \kappa_i + P_i(t) & \text{for } y \in [\kappa_i - \delta, \kappa_i + \delta], t \in [0, T], 1 \leq i \leq N, \\ y & \text{for } y \in [0, \delta] \cup [1 - \delta, 1], t \in [0, T]. \end{cases}$$

Define $\tilde{V}(y, t) = V(\Psi(y, t), t)$. By direct calculations, $\lambda(D)$ is also the principal eigenvalue of

$$(3.3) \quad \begin{cases} \partial_t \tilde{\varphi} - D \frac{\partial_{yy} \tilde{\varphi}}{(\partial_y \Psi)^2} - \left[\partial_y \tilde{m} - D \frac{\partial_{yy} \Psi}{(\partial_y \Psi)^3} \right] \partial_y \tilde{\varphi} + \tilde{V}(y, t) \tilde{\varphi} = \lambda(D) \tilde{\varphi} & \text{in } (0, 1) \times [0, T], \\ \partial_y \tilde{\varphi}(0, t) = \partial_y \tilde{\varphi}(1, t) = 0 & \text{on } [0, T], \\ \tilde{\varphi}(y, 0) = \tilde{\varphi}(y, T) & \text{on } (0, 1), \end{cases}$$

for which the principal eigenfunction becomes $\tilde{\varphi}(y, t) = \varphi(\Psi(y, t), t)$. Here φ denotes the principal eigenfunction of problem (1.1), and \tilde{m} is given by

$$(3.4) \quad \partial_y \tilde{m}(y, t) = \frac{\partial_x m(\Psi(y, t), t)}{\partial_y \Psi} + \frac{\partial_t \Psi}{\partial_y \Psi}.$$

Next, we focus on problem (3.3), and divide the proof into several steps.

Step 1. We show that the ODE problem

$$(3.5) \quad \begin{cases} \dot{\tilde{P}}(t) = -\partial_y \tilde{m}(\tilde{P}(t), t), \\ \tilde{P}(t) = \tilde{P}(t + T) \end{cases}$$

has only N periodic solutions $\tilde{P}_i(t) \equiv \kappa_i$, and $\partial_{yy} \tilde{m}(y, t) \neq 0$ for all $(y, t) \in [\kappa_i - \delta, \kappa_i + \delta] \times [0, T]$ and $i = 1, \dots, N$.

First, we claim that $\tilde{P}_i(t) \equiv \kappa_i$ is a solution of (3.5). This is due to the following calculations:

$$\begin{aligned} \partial_y \tilde{m}(\kappa_i, t) &= \frac{\partial_x m(\Psi(\kappa_i, t), t)}{\partial_y \Psi(\kappa_i, t)} + \frac{\partial_t \Psi(\kappa_i, t)}{\partial_y \Psi(\kappa_i, t)} \\ &= \partial_x m(P_i(t), t) + \dot{P}_i(t) = 0, \end{aligned}$$

where the first equality follows from (3.4), and the second equality is due to (3.2).

Suppose on the contrary that there exists a periodic solution $\tilde{P}(t)$ such that $\tilde{P}(t) \not\equiv \kappa_i$ for any $1 \leq i \leq N$. Then by (3.2) and (3.4), one can verify that $\Psi(\tilde{P}(t), t) \not\equiv P_i(t)$ is a periodic solution to (1.4) by the following calculations:

$$\begin{aligned} \dot{\Psi}(\tilde{P}(t), t) &= \dot{\tilde{P}}(t) \partial_y \Psi + \partial_t \Psi = -\partial_y \tilde{m}(\tilde{P}(t), t) \partial_y \Psi + \partial_t \Psi \\ &= -\partial_x m(\Psi(\tilde{P}(t), t), t) - \partial_t \Psi + \partial_t \Psi \\ &= -\partial_x m(\Psi(\tilde{P}(t), t), t), \end{aligned}$$

which is a contradiction. Therefore, (3.5) has only N periodic solutions $\tilde{P}_i(t) \equiv \kappa_i$ ($i = 1, \dots, N$). Furthermore, from (3.1) and (3.2), it is easily seen that $\partial_{yy} \tilde{m}(y, t) \neq 0$ on $[\kappa_i - \delta, \kappa_i + \delta] \times [0, T]$, which completes Step 1.

In the sequel, we aim to find a proper $C^{2,1}$ -transformation $\Phi : [0, 1] \times \mathbb{R} \rightarrow [0, 1]$ such that $\partial_z \Phi > 0$, and if for some $\bar{m} \in C^{2,1}([0, 1] \times [0, T])$ satisfying

$$(3.6) \quad \partial_z \bar{m}(z, r) = \frac{\partial_y \tilde{m}(\Phi(z, r), r)}{\partial_z \Phi} + \frac{\partial_r \Phi}{\partial_z \Phi},$$

then $\partial_z \bar{m} > 0$ or $\partial_z \bar{m} < 0$ holds on $(\kappa_i, \kappa_{i+1}) \times [0, T]$ for each $0 \leq i \leq N$. Then we may apply Theorem 1.3 to complete the proof.

Fix any $0 \leq i \leq N$. We assume without loss of generality that $\partial_{yy} \tilde{m}(\kappa_i, t) < 0$, so that $\partial_y \tilde{m}(\kappa_i + \delta/2, t) < 0$. For any $s \in \mathbb{R}$, denote by $q_s(t)$ the unique solution of

$$(3.7) \quad \begin{cases} \dot{q}(t) = -\partial_y \tilde{m}(q(t), t + s), \\ q(0) = \kappa_i + \delta/2, \end{cases}$$

where \tilde{m} is given by (3.4). Obviously, $q_s(t) = q_{s+T}(t)$ for all $s, t \in \mathbb{R}$. We define


$$(3.8) \quad Q(t) := \{(q_{r-t}(t), r) : r \in \mathbb{R}\},$$

which is a continuous curve and is referred as the isochron of (3.7).

Step 2. Fix any $0 < t_1 < t_2$. We show that $Q(t_1) \prec Q(t_2)$ (see Fig. 4) in the sense that

$$(3.9) \quad q_{r-t_1}(t_1) < q_{r-t_2}(t_2) \quad \text{for any } r \in \mathbb{R}.$$

We argue by contradiction by assuming $Q(t_1) \cap Q(t_2) \neq \emptyset$ or $Q(t_2) \prec Q(t_1)$.

FIGURE 4. The isochrons $Q(t_1)$ and $Q(t_2)$ of (3.7)

(i) If $Q(t_1) \cap Q(t_2) \neq \emptyset$, then by definition (3.8), there exists some $r_0 \in \mathbb{R}$ such that

$$(3.10) \quad q_{r_0-t_1}(t_1) = q_{r_0-t_2}(t_2).$$

Then we define

$$\bar{q}(t) := q_{r_0-t_1}(t - r_0 + t_1) \quad \text{and} \quad \underline{q}(t) := q_{r_0-t_2}(t - r_0 + t_2),$$

both of which satisfy $\dot{q}(t) = -\partial_y \tilde{m}(q(t), t)$, and

$$(3.11) \quad \bar{q}(r_0 - t_1) = \underline{q}(r_0 - t_2) = \kappa_i + \delta/2 \quad \text{and} \quad \bar{q}(r_0) = \underline{q}(r_0),$$

where $\bar{q}(r_0) = \underline{q}(r_0)$ follows from (3.10). In view of $t_1 < t_2$, we have $r_0 - t_1 > r_0 - t_2$. Thanks to the uniqueness of solutions to $\dot{q}(t) = -\partial_y \tilde{m}(q(t), t)$, we conclude from (3.11) that $\bar{q}(t) = \underline{q}(t)$ for any $t \in [r_0 - t_1, r_0]$, and particularly, $\underline{q}(r_0 - t_1) = \bar{q}(r_0 - t_1) = \kappa_i + \delta/2 = \underline{q}(r_0 - t_2)$, i.e. $q_{r_0-t_2}(t_2 - t_1) = \kappa_i + \delta/2 = q_{r_0-t_2}(0)$, which contradicts $\partial_y \tilde{m}(\kappa_i + \delta/2, t) < 0$.

(ii) If $Q(t_2) \prec Q(t_1)$, then given any $(q_{r_1-t_1}(t_1), r_1) \in Q(t_1)$, there is some $t_0 \in (0, t_1)$ such that $(q_{r_1-t_1}(t_0), r_2) \in Q(t_2)$, where $r_2 = r_1 - (t_1 - t_0)$. By definition (3.8), we also have $(q_{r_2-t_2}(t_2), r_2) \in Q(t_2)$, so that $q_{r_1-t_1}(t_0) = q_{r_2-t_2}(t_2)$. This, together with $r_2 - t_0 = r_1 - t_1$, leads to $q_{r_2-t_0}(t_0) = q_{r_2-t_2}(t_2)$, whence $(q_{r_2-t_2}(t_2), r_2) \in Q(t_0) \cap Q(t_2)$, i.e. $Q(t_0) \cap Q(t_2) \neq \emptyset$. Since $t_0 < t_2$, we can apply (i) to reach a contradiction.

Step 3. We show

$$\lim_{t \rightarrow \infty} Q(t) = \{(\kappa_{i+1}, r) : r \in \mathbb{R}\},$$

in the sense that for any $r \in \mathbb{R}$, $q_{r-t}(t) \rightarrow \kappa_{i+1}$ as $t \rightarrow \infty$.

By \mathbb{M} we denote the set of all continuous curves in $[\kappa_i + \delta/2, \kappa_{i+1}] \times [0, T]$. By Step 2, there is some curve $Q_\infty := \{(q_\infty(s), s) : s \in \mathbb{R}\} \in \mathbb{M}$ such that $Q(t) \rightarrow Q_\infty$ as $t \rightarrow \infty$. It suffices to show $q_\infty \equiv \kappa_{i+1}$. To this end, we claim that q_∞ is a periodic solution of (3.5), and then $q_\infty \equiv \kappa_{i+1}$ is a direct consequence of Step 1.

Indeed, the periodicity of q_∞ is due to the fact that $q_s(t) = q_{s+T}(t)$ for all $s, t \in \mathbb{R}$. We show that q_∞ is a solution to (3.5). Suppose not, then for given $s_0 \in \mathbb{R}$, there exists some $t_0 > s_0$ such that the unique solution $p_{s_0}(t)$ of

$$\begin{cases} \dot{p}(t) = -\partial_y \tilde{m}(p(t), t + s_0), \\ p(0) = q_\infty(s_0), \end{cases}$$

satisfies $p_{s_0}(t_0 - s_0) \neq q_\infty(t_0)$. Let $t_* = t_0 - s_0$. For any $\Sigma_q := \{(q(s), s) : s \in \mathbb{R}\} \in \mathbb{M}$, we denote by p_s the unique solution of

$$\begin{cases} \dot{p}(t) = -\partial_y \tilde{m}(p(t), t + s), \\ p(0) = q(s), \end{cases}$$

and define a continuous operator $F : \mathbb{M} \rightarrow \mathbb{M}$ by

$$F(\Sigma_q) := \{(p_s(t_*), t_* + s) : s \in \mathbb{R}\} = \{(p_{r-t_*}(t_*), r) : r \in \mathbb{R}\}.$$

It is straightforward to verify that $F(Q(t)) = Q(t + t_*)$, and thus

$$F(Q_\infty) = Q_\infty,$$

from which we deduce in particular that $p_{t_0-t_*}(t_*) = q_\infty(t_0)$, that is $p_{s_0}(t_0 - s_0) = q_\infty(t_0)$, a contradiction. Therefore, q_∞ is a periodic solution of (3.5). Step 3 is thereby completed.

Step 4. We define the transformation Φ satisfying $\partial_z \Phi > 0$, and for \bar{m} given by (3.6), we show that $\partial_z \bar{m} > 0$ or $\partial_z \bar{m} < 0$ holds in $(\kappa_i, \kappa_{i+1}) \times [0, T]$ for each $0 \leq i \leq N$.

For any $0 \leq i \leq N$, we define $\Phi_i : [\kappa_i + \delta/2, \kappa_{i+1} - \delta/2] \times \mathbb{R} \rightarrow [\kappa_i, \kappa_{i+1}]$ such that for any $(z, r) \in [\kappa_i + \delta, \kappa_{i+1} - \delta] \times \mathbb{R}$,

$$(3.12) \quad \Phi_i(z, r) = q_{r-\tau_i(z)}(\tau_i(z)),$$

where $q_{r-\tau_i(z)}$ is the solution of (3.7) with $s = r - \tau_i(z)$ and $\tau_i(z)$ is determined by

$$(3.13) \quad q_{-\tau_i(z)}(\tau_i(z)) = z.$$

Obviously, $\{(\Phi_i(z, r), r) : r \in \mathbb{R}\} = Q(\tau_i(z))$. It is easily seen that $z \rightarrow \tau_i(z)$ is a bijection (where the surjection follows from Step 3), is of class C^2 and is increasing (by Step 2), so that $\Phi_i \in C^{2,1}([\kappa_i + \delta, \kappa_{i+1} - \delta] \times \mathbb{R})$ and $\partial_z \Phi_i \geq 0$ by (3.9).

We claim that for $(z, r) \in (\kappa_i + \delta/2, \kappa_{i+1} - \delta/2) \times \mathbb{R}$,

$$(3.14) \quad \tau'_i(z) > 0 \quad \text{and} \quad \partial_y \tilde{m}(\Phi_i(z, r), r) + \partial_r \Phi_i = -\frac{\partial_z \Phi_i}{\tau'_i(z)}.$$

For the sake of clarification, write $q_s(t) = q(t; s)$, where q_s is defined by (3.7). Differentiating both sides of (3.13) by z , we derive that

$$\left[\partial_t q_{-\tau_i(z)}(\tau_i(z)) - \partial_s q_{r-\tau_i(z)}(\tau_i(z)) \right] \tau'_i(z) = 1,$$

which implies $\tau'_i(z) \neq 0$, and thus $\tau'_i(z) > 0$ since $\tau_i(z)$ is increasing. Similarly, by (3.12), we deduce that $\partial_r \Phi_i(z, r) = \partial_s q_{r-\tau_i(z)}(\tau_i(z))$, and thus

$$(3.15) \quad \begin{aligned} \partial_z \Phi_i(z, r) &= \left[\partial_t q_{r-\tau_i(z)}(\tau_i(z)) - \partial_s q_{r-\tau_i(z)}(\tau_i(z)) \right] \tau'_i(z) \\ &= \left[\partial_t q_{r-\tau_i(z)}(\tau_i(z)) - \partial_r \Phi_i(z, r) \right] \tau'_i(z). \end{aligned}$$

By the definition of $q_{r-\tau_i(z)}(\tau_i(z))$ in (3.7) with $s = r - \tau_i(z)$ and $t = \tau_i(z)$, we note that

$$\partial_t q_{r-\tau_i(z)}(\tau_i(z)) = -\partial_y \tilde{m}(\Phi_i(z, r), r),$$

which, together with (3.15), implies (3.14).

We then claim that

$$(3.16) \quad \partial_z \Phi_i(z, r) > 0 \quad \text{for any } (z, r) \in (\kappa_i + \delta/2, \kappa_{i+1} - \delta/2) \times \mathbb{R}.$$

To this end, denote by $\tilde{p}(t; s)$ the unique solution of the problem

$$\begin{cases} \dot{\tilde{p}}(t) = -\partial_y \tilde{m}(\tilde{p}(t), t), \\ \tilde{p}(s) = \kappa_i + \delta/2, \end{cases}$$

whence by (3.7), we observe that $q_s(t) = \tilde{p}(t + s; s)$. For any $\tau \in \mathbb{R}$, we have

$$\tilde{p}(\tau) = - \int_s^\tau \partial_y \tilde{m}(\tilde{p}(t), t) dt + \kappa_i + \delta/2,$$

so that

$$\partial_s \tilde{p}(\tau) = \partial_y \tilde{m}(\kappa_i + \delta/2, s) - \int_s^\tau \partial_{yy} \tilde{m}(\tilde{p}(t), t) \partial_s \tilde{p}(t) dt,$$

and thus $\partial_s \tilde{p}(s) = \partial_y \tilde{m}(\kappa_i + \delta/2, s) < 0$. We further calculate that

$$\partial_\tau (\partial_s \tilde{p}(\tau)) = -\partial_{yy} \tilde{m}(\tilde{p}(\tau), \tau) \partial_s \tilde{p}(\tau),$$

which implies immediately that for any $r \in \mathbb{R}$,

$$(3.17) \quad \partial_s \tilde{p}(r) = \partial_s \tilde{p}(s) \exp \left[- \int_s^r \partial_{yy} \tilde{m}(\tilde{p}(\tau), \tau) d\tau \right] < 0.$$

By (3.12) and the fact that $q_s(t) = \tilde{p}(t + s; s)$, we can see $\Phi_i(z, r) = \tilde{p}(r; r - \tau_i(z))$, so that

$$\partial_z \Phi_i(z, r) = -\partial_s \tilde{p}(r) \cdot \tau'_i(z) > 0$$

by noting that $\tau'_i(z) > 0$ in (3.14) and $\partial_s \tilde{p}(r) < 0$ in (3.17).

Then we define a $C^{2,1}$ -transformation $\Phi : [0, 1] \times \mathbb{R} \rightarrow [0, 1]$ such that $\partial_z \Phi > 0$ and for any $0 \leq i \leq N$,

$$(3.18) \quad \Phi(z, r) := \begin{cases} \Phi_i(z, r) & \text{on } [\kappa_i + \delta_1, \kappa_{i+1} - \delta_1] \times \mathbb{R}, \\ z & \text{on } ([\kappa_i, \kappa_i + \delta/2] \cup [\kappa_{i+1} - \delta/2, \kappa_{i+1}]) \times \mathbb{R}, \end{cases}$$

where $\delta_1 \in (\delta/2, \delta]$ is chosen to be close to $\delta/2$ such that

$$(3.19) \quad \partial_y \tilde{m} + \partial_r \Phi < 0 \quad \text{on } ([\kappa_i, \kappa_i + \delta_1] \cup [\kappa_{i+1} - \delta_1, \kappa_{i+1}]) \times \mathbb{R}.$$

This is possible since by (3.18) and Step 1, it follows that

$$\partial_y \tilde{m} + \partial_r \Phi = \partial_y \tilde{m} < 0 \quad \text{on } ([\kappa_i, \kappa_i + \delta/2] \cup [\kappa_{i+1} - \delta/2, \kappa_{i+1}]) \times \mathbb{R}.$$

Let \overline{m} satisfy (3.6) with Φ defined by (3.18). For any $z \in [\kappa_i, \kappa_i + \delta_1] \cup [\kappa_{i+1} - \delta_1, \kappa_{i+1}]$, it follows from (3.6), (3.18), and (3.19) that $\partial_z \overline{m}(z, r) < 0$; For $z \in [\kappa_i + \delta_1, \kappa_{i+1} - \delta_1]$, by (3.18) we have $\Phi(z, r) = \Phi_i(z, r)$, whence comparing (3.6) with (3.14) gives $\partial_z \overline{m}(z, r) = -\frac{1}{\tau'_i(z)} < 0$. This completes Step 4.

Step 5. We apply Theorem 1.3 to complete the proof. Let the $C^{2,1}$ -transformation Φ be defined by (3.18) in Step 4. Denote

$$\bar{V}(z, r) := \tilde{V}(\Phi(z, r), r) \quad \text{and} \quad \bar{\varphi}(z, r) := \tilde{\varphi}(\Phi(z, r), r),$$

where \tilde{V} and $\tilde{\varphi}$ are defined in (3.3). Using the definition of \bar{m} in (3.6), direct calculation enables us to transform (3.3) into the following equation:

$$(3.20) \quad \begin{cases} \partial_r \bar{\varphi} - \frac{D \partial_{zz} \bar{\varphi}}{(\partial_y \Psi)^2 (\partial_z \Phi)^2} - [\partial_z \bar{m} + D \eta_3] \partial_z \bar{\varphi} + \bar{V} \bar{\varphi} = \lambda(D) \bar{\varphi} & \text{in } (0, 1) \times [0, T], \\ \partial_z \bar{\varphi}(0, r) = \partial_z \bar{\varphi}(1, r) = 0 & \text{on } [0, T], \\ \bar{\varphi}(z, 0) = \bar{\varphi}(z, T) & \text{on } (0, 1), \end{cases}$$

where η_3 is given by

$$\eta_3(z, r) := \frac{\partial_{yy} \Psi}{(\partial_y \Psi)^3 \partial_z \Phi} + \frac{\partial_{zz} \Phi}{(\partial_z \Phi)^3 (\partial_y \Psi)^2}.$$

For each $0 \leq i \leq N$, by Step 4, $\partial_z \bar{m} > 0$ or $\partial_z \bar{m} < 0$ holds for all $z \in (\kappa_i, \kappa_{i+1})$; by the definitions of Ψ and Φ in (3.2) and (3.18), we find that for any $z \in [\kappa_i, \kappa_i + \delta/2] \cup [\kappa_{i+1} - \delta/2, \kappa_{i+1}]$, $\partial_{yy} \Psi = \partial_{zz} \Phi = 0$, so that $\eta_3(z, r) = 0$. Therefore, we conclude that for any small $\vartheta > 0$, there exists some $\epsilon_0 = \epsilon_0(\vartheta) > 0$, independent of small D , such that

$$(3.21) \quad \partial_z \bar{m} + D \eta_3 \geq \epsilon_0 \quad \text{or} \quad \partial_z \bar{m} + D \eta_3 \leq -\epsilon_0 \quad \text{on } [\kappa_i + \vartheta, \kappa_{i+1} - \vartheta] \times [0, T].$$

Moreover, from (3.6) and (3.18), we observe that for any $1 \leq i \leq N$,

$$\partial_z \bar{m}(\kappa_i, r) = \partial_y \tilde{m}(\Phi(\kappa_i, r), r) = \partial_y \tilde{m}(\kappa_i, r) = 0,$$

which implies that $\partial_z \bar{m}(\kappa_i, r) + D \eta_3(\kappa_i, r) = 0$ since $\eta_3(\kappa_i, r) = 0$. Together with (3.21), noting that $\partial_y \Psi = \partial_z \Phi = 1$ on $[\kappa_i, \kappa_i + \tilde{\vartheta}] \cup [\kappa_{i+1} - \tilde{\vartheta}, \kappa_{i+1}]$ with some $0 < \tilde{\vartheta} \ll 1$ for any $0 \leq i \leq N$, we can follow directly the same proof of Theorem 1.3 with $\mathbf{B} = \emptyset$ to (3.20) and deduce that

$$(3.22) \quad \lim_{D \rightarrow 0} \lambda(D) = \min_{0 \leq i \leq N+1} \left\{ \hat{\bar{V}}(\kappa_i) + [\partial_{xx} \hat{m}]_+(\kappa_i) \right\}.$$

Noting that $\hat{\bar{V}}(\kappa_i) = \frac{1}{T} \int_0^T V(P_i(s), s) ds$ and

$$\partial_{zz} \bar{m}(\kappa_i, r) = \partial_{yy} \tilde{m}(\kappa_i, r) = \partial_{xx} m(P_i(r), r),$$

part (i) of Theorem 1.2 follows from (3.22).

Finally, part (ii) of Theorem 1.2 can be established by Steps 2–5 with $N = 0$. The proof is now complete. \square

4. PROOF OF THEOREM 1.4

This section is devoted to the case $m(x, t) = ab(t)x$ and the proof of Theorem 1.4. We start with the existence and uniqueness of \tilde{P}_α defined in Theorem 1.4.

Lemma 4.1. *Let F be defined by (1.7) and \bar{P}, \underline{P} be given in Theorem 1.4. If $\alpha \geq \frac{1}{\bar{P} - \underline{P}}$, then*

$$(4.1) \quad \begin{cases} \dot{P}(t) = -\alpha F(P(t), t), \\ P(t) = P(t + T) \end{cases}$$

has a unique T -periodic solution in $W^{1,\infty}(\mathbb{R})$.

Proof. Recalling the definition of F in (1.7), we observe that $F(0, t) = \min\{b(t), 0\} \leq 0$ and $F(1, t) = \max\{b(t), 0\} \geq 0$, so that $P_* \equiv 0$ and $P^* \equiv 1$ are a pair of sub- and super-solutions to (4.1). Hence, as F is bounded, there exists at least one T -periodic solution in $W^{1,\infty}(\mathbb{R})$.

For the uniqueness, given any two T -periodic solutions \tilde{P} and \tilde{P}_α of (4.1), we show $\tilde{P} = \tilde{P}_\alpha$. Suppose not, without loss of generality we may assume $\tilde{P}(0) < \tilde{P}_\alpha(0)$. We consider two cases:

(i) If $\tilde{P}(t_1) = \tilde{P}_\alpha(t_1)$ for some $t_1 \in (0, T)$, in view of $\tilde{P}(T) = \tilde{P}(0) < \tilde{P}_\alpha(0) = \tilde{P}_\alpha(T)$, by continuity there is some $\tilde{t}_1 \in [t_1, T)$ such that $\tilde{P}(\tilde{t}_1) = \tilde{P}_\alpha(\tilde{t}_1)$ and $\tilde{P}(t) < \tilde{P}_\alpha(t)$ for any $t \in (\tilde{t}_1, T]$. Then by the definition of F , it can be verified that for any $t \in [\tilde{t}_1, T]$,

$$(4.2) \quad [\tilde{P}_\alpha(t) - \tilde{P}(t)] - [\tilde{P}_\alpha(\tilde{t}_1) - \tilde{P}(\tilde{t}_1)] = \alpha \int_{\tilde{t}_1}^t [F(\tilde{P}(s), s) - F(\tilde{P}_\alpha(s), s)] \, ds \leq 0,$$

which implies $\tilde{P}_\alpha(T) - \tilde{P}(T) \leq \tilde{P}_\alpha(\tilde{t}_1) - \tilde{P}(\tilde{t}_1) = 0$, a contradiction.

(ii) If $\tilde{P}(t) < \tilde{P}_\alpha(t)$ for all $t \in [0, T]$, then (4.2) holds for all $t \in [0, T]$ and $\tilde{t}_1 = 0$. In view of $\tilde{P}_\alpha(T) - \tilde{P}(T) = \tilde{P}_\alpha(0) - \tilde{P}(0)$, we deduce that

$$\tilde{P}_\alpha(t) - \tilde{P}(t) \equiv \tilde{P}_\alpha(0) - \tilde{P}(0) \quad \text{for all } t \in [0, T].$$

In such a case, again by the definition of F , we infer that

$$\tilde{P}_+ := \tilde{P} + (\tilde{P}_\alpha(0) - \tilde{P}(0))/2 \in (0, 1)$$

defines a T -periodic solution of (4.1), and thus $\dot{\tilde{P}}_+ = -\alpha b(t)$, where $\tilde{P}_+ \in (0, 1)$ is due to $0 \leq \tilde{P} < \tilde{P}_+ < \tilde{P}_\alpha \leq 1$. By recalling $P(t) = -\int_0^t b(s) \, ds$, this implies that $\tilde{P}_+ = \alpha P(t) + c \in (0, 1)$ for some constant $c \in \mathbb{R}$, so that

$$1 > \max_{[0, T]} \tilde{P}_+ - \min_{[0, T]} \tilde{P}_+ = \alpha(\bar{P} - \underline{P}),$$

which contradicts $\alpha \geq \frac{1}{\bar{P} - \underline{P}}$. Lemma 4.1 thus follows. \square

We are now ready to prove Theorem 1.4.

Proof of Theorem 1.4. The proof is divided into three steps.

Step 1. Assume $\hat{b} \neq 0$ and show part (i) of Theorem 1.4. Let $\Psi_1 : [0, 1] \times [0, T] \rightarrow \mathbb{R}$ denote a T -periodic diffeomorphism given by

$$\Psi_1(y, t) = \alpha \left[\hat{b}t - \int_0^t b(s) \, ds \right] + y.$$

Under the transformation $x = \Psi_1(y, t)$, as in (3.3), direct calculation from (1.6) yields that $\lambda(D)$ defines the principal eigenvalue of the problem

$$\begin{cases} \partial_t \varphi - D \partial_{yy} \varphi - \alpha \hat{b} \cdot \partial_y \varphi + V_1 \varphi = \lambda(D) \varphi, & y \in (-\Psi_1(0, t), 1 - \Psi_1(0, t)), t \in [0, T], \\ \partial_y \varphi(-\Psi_1(0, t), t) = \partial_y \varphi(1 - \Psi_1(0, t), t) = 0, & t \in [0, T], \\ \varphi(y, 0) = \varphi(y, T), & y \in [-\Psi_1(0, 0), 1 - \Psi_1(0, 0)], \end{cases}$$

where $V_1(y, t) = V(\Psi_1(y, t), t)$. Then we can conclude that part (i) of Theorem 1.4 is a direct consequence of Theorem 1.2. Indeed, if $\hat{b} > 0$ for example, then ODE (1.4) with $\partial_x m = \alpha\hat{b} > 0$ has no periodic solutions, so that by part (ii) of Theorem 1.2 we deduce that

$$\lim_{D \rightarrow 0} \lambda(D) = \frac{1}{T} \int_0^T V_1(1 - \Psi_1(0, s), s) ds = \hat{V}(1).$$

The same argument can be adapted to the case $\hat{b} < 0$, which completes Step 1.

Step 2. Assume $\hat{b} = 0$ and $0 < \alpha \leq \frac{1}{\underline{P} - \bar{P}}$. We prove the first part of (ii) in Theorem 1.4. Recall $P(t) = -\int_0^t b(s) ds$ defined in Theorem 1.4. Taking the transformation $x = y + \alpha P(t)$ in (1.6), we derive that $\lambda(D)$ is also the principal eigenvalue of the problem

$$\begin{cases} \partial_t \varphi - D \partial_{yy} \varphi + V_2 \varphi = \lambda(D) \varphi, & y \in (-\alpha P(t), 1 - \alpha P(t)), t \in [0, T], \\ \partial_y \varphi(-\alpha P(t), t) = \partial_y \varphi(1 - \alpha P(t), t) = 0, & t \in [0, T], \\ \varphi(y, 0) = \varphi(y, T), & y \in (-\alpha P(0), 1 - \alpha P(0)), \end{cases}$$

where $V_2(y, t) = V(\alpha P(t) + y, t)$. Under the transformation $x = y + \alpha P(t)$, all periodic solutions of (1.4) are constants in the interval $[-\alpha \underline{P}, 1 - \alpha \bar{P}]$. This includes the special case $\alpha = \frac{1}{\underline{P} - \bar{P}}$, for which the interval reduces to a single point. It is desired to show that

$$\lim_{D \rightarrow 0} \lambda(D) = \min_{y \in [-\alpha \underline{P}, 1 - \alpha \bar{P}]} \hat{V}_2(y).$$

First, the upper bound $\limsup_{D \rightarrow 0} \lambda(D) \leq \hat{V}_2(y)$, for any $y \in [-\alpha \underline{P}, 1 - \alpha \bar{P}]$, can be established by the same arguments as in Step 1 of Lemma 2.7 by constructing the sub-solution locally. We thus omit the details here.

It remains to show the lower bound of $\liminf_{D \rightarrow 0} \lambda(D)$. For any $\epsilon > 0$, we define T -periodic function $V_{2\epsilon} \in C^{2,1}(\mathbb{R} \times [0, T])$ satisfying $\|V_{2\epsilon} - V_2\|_{L^\infty} \leq \epsilon$, and choose small $\delta > 0$ such that

$$(4.3) \quad \tilde{\lambda}_{\min} := \min_{y \in [-\alpha \underline{P} - 2\delta, 1 - \alpha \bar{P} + 2\delta]} \hat{V}_{2\epsilon}(y) \geq \min_{y \in [-\alpha \underline{P}, 1 - \alpha \bar{P}]} \hat{V}_2(y) - 2\epsilon.$$

We define $\bar{\phi} \in C^{2,1}([-\alpha \underline{P} - 2\delta, 1 - \alpha \bar{P} + 2\delta] \times [0, T])$ by

$$(4.4) \quad \bar{\phi}(y, t) := \exp \left[- \int_0^t V_{2\epsilon}(y, s) ds + t \hat{V}_{2\epsilon}(y) \right] \beta_\epsilon(y),$$

where $\beta_\epsilon \in C^2([-\alpha \underline{P} - 2\delta, 1 - \alpha \bar{P} + 2\delta])$ is a positive function chosen such that

$$(4.5) \quad \begin{aligned} \partial_y \bar{\phi} &< 0 \quad \text{on } [-\alpha \underline{P} - 2\delta, -\alpha \underline{P}] \times [0, T] \\ \text{and} \quad \partial_y \bar{\phi} &> 0 \quad \text{on } [1 - \alpha \bar{P}, 1 - \alpha \bar{P} + 2\delta] \times [0, T]. \end{aligned}$$

Next, we aim to find a super-solution $\bar{\varphi} \in C([0, 1] \times [0, T])$ which satisfies

$$(4.6) \quad \begin{cases} \partial_t \bar{\varphi} - D \partial_{yy} \bar{\varphi} + V_2 \bar{\varphi} \geq [\tilde{\lambda}_{\min} - 3\epsilon] \bar{\varphi}, & y \in (-\alpha P(t), 1 - \alpha P(t)) \setminus \mathbb{X}, t \in [0, T], \\ \partial_y \bar{\varphi}(-\alpha P(t), t) \leq 0 \leq \partial_y \bar{\varphi}(1 - \alpha P(t), t), & t \in [0, T], \\ \bar{\varphi}(y, 0) = \bar{\varphi}(y, T), & y \in (-\alpha P(0), 1 - \alpha P(0)), \end{cases}$$

where $\mathbb{X} = \{-\alpha\underline{P} - 2\delta, 1 - \alpha\overline{P} + 2\delta\}$. Then it follows from Proposition A.2 and (4.3) that

$$\liminf_{D \rightarrow 0} \lambda(D) \geq \min_{y \in [-\alpha\underline{P}, 1 - \alpha\overline{P}]} \hat{V}_2(y);$$

see also Remark A.3.

We only construct $\bar{\varphi}$ for $y \in (-\alpha P(t), 1 - \alpha\overline{P} + \delta)$ and $t \in [0, T]$. The constructions of the remaining regions are similar. To this end, by the definition of \underline{P} , there exist $t_3 > t_2$ such that

$$[t_2, t_3] \subset \{t \in [0, T] : -\alpha P(t) > -\alpha\underline{P} - \delta\}.$$

We then choose $\eta_4 \in C^{2,1}((-\infty, 1 - \alpha\overline{P} + \delta] \times [0, T])$ to be a positive T -periodic function, and satisfy that $\partial_y \eta_4 \leq 0$ and

$$(4.7) \quad \begin{cases} \eta_4 \equiv 1 & \text{on } [-\alpha\underline{P} - \delta, 1 - \alpha\overline{P} + \delta] \times [0, T], \\ \partial_t(\log \eta_4) > 0 & \text{on } [-\alpha\underline{P} - 2\delta, -\alpha\underline{P} - \delta] \times ([0, T] \setminus [t_2, t_3]), \\ \partial_t(\log \eta_4) \geq M_4 & \text{on } (-\infty, -\alpha\underline{P} - 2\delta] \times ([0, T] \setminus [t_2, t_3]). \end{cases}$$

Here M_4 is chosen such that

$$M_4 > \|V_2\|_{L^\infty} + \tilde{\lambda}_{\min} + \|\partial_t \log \bar{\phi}\|_{L^\infty},$$

where $\bar{\phi}$ is defined by (4.4). Moreover, we extend $\bar{\phi}$ to $(-\infty, 1 - \alpha\underline{P} + 2\delta] \times [0, T]$ by setting $\bar{\phi}(\cdot, t) \equiv \bar{\phi}(-\alpha\underline{P} - 2\delta, t)$ on $(-\infty, -\alpha\underline{P} - 2\delta) \times [0, T]$, so that by (4.5) we have

$$(4.8) \quad \partial_y \bar{\phi}((-\alpha\underline{P} - 2\delta)^+, \cdot) < 0 = \partial_y \bar{\phi}((-\alpha\underline{P} - 2\delta)^-, \cdot).$$

Let $\bar{\phi}$ and η_4 be given by (4.4) and (4.7), then we define

$$(4.9) \quad \bar{\varphi}(y, t) := \eta_4(y, t) \cdot \bar{\phi}(y, t).$$

By (4.8), as η_4 is smooth, one can infer that

$$\partial_y \log \bar{\varphi}((-\alpha\underline{P} - 2\delta)^+, \cdot) < \partial_y \log \bar{\varphi}((-\alpha\underline{P} - 2\delta)^-, \cdot) \quad \text{as } -\alpha\underline{P} - 2\delta \in \mathbb{X}.$$

It remains to check that $\bar{\varphi}$ defined above satisfies (4.6).

(i) For $y \in (-\alpha P(t), 1 - \alpha P(t)) \cap [-\alpha\underline{P} - \delta, 1 - \alpha\overline{P} + \delta]$ and $t \in [0, T]$, since $\eta_4 \equiv 1$ in (4.7), we have $\bar{\varphi}(y, t) = \bar{\phi}(y, t)$. By the definition of $\bar{\phi}$ in (4.4), direct calculations yield that

$$\partial_t \bar{\varphi} - D \partial_{yy} \bar{\varphi} + V_2 \bar{\varphi} = \left[\hat{V}_{2\epsilon}(y) - V_{2\epsilon}(y, t) + V_2(y, t) \right] \bar{\phi} - D \partial_{yy} \bar{\phi}.$$

By the definition of $V_{2\epsilon}$, we can argue as in Lemma 2.7 to choose D small such that the first inequality in (4.6) holds. Then the part of boundary conditions on $\{-\alpha P(t), 1 - \alpha P(t)\} \cap [-\alpha\underline{P} - \delta, 1 - \alpha\overline{P} + \delta]$ and $t \in [0, T]$ can be verified by (4.5).

(ii) For $y \in (-\alpha P(t), 1 - \alpha P(t)) \cap [-\alpha\underline{P} - 2\delta, -\alpha\underline{P} - \delta]$ and $t \in [0, T]$, since $t \in [0, T] \setminus [t_2, t_3]$ in this case, we use (4.7) and (4.9) to deduce that

$$\begin{aligned} \partial_t \bar{\varphi} - D \partial_{yy} \bar{\varphi} + V_2 \bar{\varphi} &= \left[\hat{V}_{2\epsilon}(y) - V_{2\epsilon}(y, t) + V_2(y, t) \right] \bar{\varphi} + [\partial_t(\log \eta_4) - D \partial_{yy} \bar{\varphi}] \bar{\varphi} \\ &\geq \left[\tilde{\lambda}_{\min} - 2\epsilon + \partial_t(\log \eta_4) + O(D) \right] \bar{\varphi}. \end{aligned}$$

Since $\partial_t(\log \eta_4) > 0$ in this case, again we choose D small such that (4.6) holds. And the boundary conditions in this case can be verified by $\partial_y \bar{\phi} \leq 0$ and $\partial_y \eta_4 \leq 0$.

(iii) For $y \in (-\alpha P(t), 1 - \alpha P(t)) \cap (-\infty, -\underline{P} - 2\delta)$ and $t \in [0, T]$, since $\bar{\phi}$ is independent of y , by (4.7) and (4.9) direct calculation yields that

$$\partial_t \bar{\varphi} - D \partial_{yy} \bar{\varphi} + V_2 \bar{\varphi} \geq \left[(\log \bar{\phi})' + M_4 - D \partial_{yy} \eta_4 / \eta_4 + V_2 \right] \bar{\varphi}.$$

Thus the first inequality in (4.6) is verified by the definition of M_4 , and the boundary condition follows from $\partial_y \eta_4 \leq 0$. Step 2 is now completed.

Step 3. Assume $\hat{b} = 0$ and $\alpha > \frac{1}{\underline{P} - \underline{P}}$. We establish the second part of (ii) in Theorem 1.4. Let \tilde{P}_α denote the unique solution of (4.1). We apply the transformation $x = y + \tilde{P}_\alpha(t)$ to rewrite problem (1.6) as

$$\begin{cases} \partial_t \varphi - D \partial_{yy} \varphi - \alpha \tilde{b}(t) \partial_y \varphi + V_3 \varphi = \lambda(D) \varphi, & (y, t) \in \tilde{\Omega}, \\ \partial_y \varphi(-\tilde{P}_\alpha(t), t) = \partial_y \varphi(1 - \tilde{P}_\alpha(t), t) = 0, & t \in [0, T], \\ \varphi(y, 0) = \varphi(y, T), & y \in (-\tilde{P}_\alpha(0), 1 - \tilde{P}_\alpha(0)), \end{cases}$$

where $\tilde{b}(t) := b(t) - F(\tilde{P}_\alpha(t), t)$, $V_3(y, t) = V(\tilde{P}_\alpha(t) + y, t)$, and

$$\tilde{\Omega} = \left\{ (y, t) : y \in (-\tilde{P}_\alpha(t), 1 - \tilde{P}_\alpha(t)), t \in [0, T] \right\}.$$

See Fig. 5 for an example of this transformation.




FIGURE 5. The diagram of $\tilde{\Omega}$ under transformation $x = y + \tilde{P}_\alpha(t)$. The red colored curve in the left side picture corresponds to $\tilde{P}_\alpha(t)$, whereas the red colored line in the right side picture is the image of $\tilde{P}_\alpha(t)$ after the transformation.

It remains to prove

$$\lim_{D \rightarrow 0} \lambda(D) = \hat{V}_3(0).$$

The upper bound $\limsup_{D \rightarrow 0} \lambda(D) \leq \hat{V}_3(0)$ can be established by using the arguments in Step 1 of Proposition 2.1. We next prove $\liminf_{D \rightarrow 0} \lambda(D) \geq \hat{V}_3(0)$.

We claim that if $\alpha > \frac{1}{\underline{P} - \underline{P}}$, then

$$\text{mes} \left\{ t \in [0, T] : \tilde{P}_\alpha(t) \in \{0, 1\} \text{ and } b \neq 0 \right\} > 0,$$

i.e. there exist $0 \leq t_4 < t_5 \leq T$ such that $b \neq 0$, and $\tilde{P}_\alpha(t) \equiv 0$ or $\tilde{P}_\alpha(t) \equiv 1$ on $[t_4, t_5]$. Suppose not, then \tilde{P}_α is also a periodic solution of $\dot{P}(t) = -\alpha b(t)$, so that $\tilde{P}_\alpha(t) = P(t) + c$ for $c \in \mathbb{R}$, where $P(t) = -\int_0^t b(s) ds$ as defined in part (ii) of

Theorem 1.4. Since $\tilde{P}_\alpha \in [0, 1]$, we have

$$1 \geq \max_{[0, T]} \tilde{P}_\alpha - \min_{[0, T]} \tilde{P}_\alpha = \alpha(\bar{P} - \underline{P}),$$

which contradicts $\alpha > \frac{1}{\bar{P} - \underline{P}}$.

In what follows, we assume $\tilde{P}_\alpha(t) \equiv 1$ on $[t_4, t_5]$, and the proof is similar for the other case. To proceed further, we introduce positive functions $\bar{z}_5 \in C^2(\mathbb{R})$ and $\eta_5 \in C^1([0, T])$ as follows: For any $\epsilon > 0$, we choose some small $\delta > 0$ such that

$$(4.10) \quad |V_3(y, t) - V_3(0, t)| < \epsilon/2 \quad \text{on } [-2\delta, 2\delta] \times [0, T].$$

We first choose η_5 to be T -periodic and

$$(4.11) \quad (\log \eta_5)' > 2\|V\|_{L^\infty} + \|(\log f_1)'\|_{L^\infty} \quad \text{on } [0, t_4 + \delta] \cup [t_5 - \delta, T].$$

Then we choose \bar{z}_5 such that

$$(4.12) \quad \begin{cases} \bar{z}_5'(y) < 0 & \text{in } (-\infty, 0), \quad \bar{z}_5'(y) > 0 \quad \text{in } (0, \infty), \\ (\log \bar{z}_5)' \leq -M_5 & \text{in } (-\infty, -\delta), \end{cases}$$

where M_5 is some large constant to be determined later.

We define

$$(4.13) \quad \bar{\varphi}(y, t) := \bar{z}_5(y) \cdot \begin{cases} f_1(t) & \text{for } |y| \leq \delta, \\ \zeta_5(y, t) & \text{for } -2\delta < y < -\delta, \\ \zeta_5(-y, t) & \text{for } \delta < y < 2\delta, \\ \eta_5(t) & \text{for } |y| \geq 2\delta, \end{cases}$$

where f_1 is defined by (2.1) with $x = 1$. Due to the choice of η_5 in (4.11), ζ_5 can be chosen such that $\bar{\varphi} \in C^{2,1}(\mathbb{R} \times [0, T])$ and

$$(4.14) \quad \partial_t(\log \zeta_5) \geq (\log f_1)' \quad \text{on } [0, t_4 + \delta] \cup [t_5 - \delta, T].$$

We shall verify that $\bar{\varphi}$ defined by (4.13) satisfies

$$(4.15) \quad L_D \bar{\varphi} := \partial_t \bar{\varphi} - D \partial_{yy} \bar{\varphi} - \alpha \tilde{b}(t) \partial_y \bar{\varphi} + V_3 \bar{\varphi} \geq (\hat{V}_3(0) - \epsilon) \bar{\varphi} \quad \text{for } (y, t) \in \tilde{\Omega},$$

provided that D is small enough. The verification is divided into the following cases:

(i) For $(y, t) \in ((-\delta, \delta] \times [0, T]) \cap \tilde{\Omega}$, we note that (see Fig. 5)

$$\tilde{b}(t) \geq 0 \quad \text{in } ((-\delta, 0] \times [0, T]) \cap \tilde{\Omega} \quad \text{and} \quad \tilde{b}(t) \leq 0 \quad \text{in } ([0, \delta] \times [0, T]) \cap \tilde{\Omega}.$$

One can check (4.15) by the same arguments as in Step 2 of Proposition 2.1.

(ii) For $(y, t) \in ((-\infty, -\delta] \times [t_4 + \delta, t_5 - \delta]) \cap \tilde{\Omega} = (-1, -\delta) \times [t_4 + \delta, t_5 - \delta]$ (since $\tilde{P}_\alpha(t) \equiv 1$ on $[t_4, t_5]$), there exists some $\epsilon_0 > 0$ such that $\tilde{b}(t) > \epsilon_0$. By the choice of \bar{z}_5 in (4.12) and construction (4.13), direct calculation gives

$$L_D \bar{\varphi} \geq \left[-|(\log \eta_5)' + \partial_t(\log \zeta_5)'| - D \partial_{yy} \bar{\varphi} + \alpha \epsilon_0 M_5 - \alpha \tilde{b} |\zeta_5| + V_3 \right] \bar{\varphi}.$$

By choosing M_5 large and D small, we can verify that (4.15) holds.

(iii) For $(y, t) \in ((-2\delta, -\delta] \times ([0, t_4 + \delta] \cup [t_5 - \delta, T])) \cap \tilde{\Omega}$, by construction, $\bar{\varphi}(y, t) = \bar{z}_5(y) \zeta_5(y, t)$. Observe that $\tilde{b} \geq 0$ in this case. Using (4.12), we choose M_5 large such that

$$-\tilde{b}(t) \partial_y \bar{\varphi} \geq \tilde{b}(t) [M_5 - \partial_y(\log \zeta_5)] \bar{\varphi} \geq 0.$$

Hence, by (4.10) and (4.14), for small D we arrive at

$$\begin{aligned} L_D \bar{\varphi} &\geq \left[\partial_t(\log \zeta_5) + V_3 - \epsilon/2 \right] \bar{\varphi} \\ &\geq \left[(\log f_1)' + V_3 - \epsilon/2 \right] \bar{\varphi} \\ &= \left[\hat{V}_3(0) - V_3(0, t) + V_3(y, t) - \epsilon/2 \right] \bar{\varphi} \\ &\geq \left[\hat{V}_3(0) - \epsilon \right] \bar{\varphi}. \end{aligned}$$

(iv) For $(y, t) \in ((-\infty, -2\delta) \times [0, t_4 + \delta] \cup [t_5 - \delta, T]) \cap \tilde{\Omega}$, by (4.13) we have $\bar{\varphi}(y, t) = \bar{z}_5(y)\eta_5(t)$. Also since $\tilde{b} \geq 0$, the choice of \bar{z}_5 in (4.12) implies $-\tilde{b}(t)\partial_y \bar{\varphi} \geq 0$. Choosing D smaller if necessary, we use (4.11) to deduce that

$$L_D \bar{\varphi} \geq \left[(\log \eta_5)' - D\bar{z}_5''/\bar{z}_5 - V_3 \right] \bar{\varphi} \geq \hat{V}_3(0) \bar{\varphi}.$$

(v) For $(y, t) \in ((\delta, \infty) \times [0, T]) \cap \tilde{\Omega}$, the verification of (4.15) is rather similar to that in cases **(ii)**–**(iv)**, and thus is omitted.

Finally, we verify the boundary conditions

$$(4.16) \quad \partial_y \bar{\varphi}(-\tilde{P}_\alpha(t), t) \leq 0 \quad \text{and} \quad \partial_y \bar{\varphi}(1 - \tilde{P}_\alpha(t), t) \geq 0 \quad \text{for } t \in [0, T].$$

For the set $\{t \in [0, T] : -\alpha \tilde{P}_\alpha(t) \in [-2\delta, -\delta] \text{ or } 1 - \alpha \tilde{P}_\alpha(t) \in [\delta, 2\delta]\}$, we can choose M_5 large such that $M_5 > \|\partial_y(\log \zeta_5)\|_{L^\infty}$ to verify (4.16) as in case **(iii)**. The verification of (4.16) for the remaining cases is straightforward.

By (4.15) and (4.16), we apply Proposition A.2 and Remark A.3 to conclude $\liminf_{D \rightarrow 0} \lambda(D) \geq \hat{V}_3(0)$. The proof of Theorem 1.4 is thereby completed. \square

APPENDIX A. GENERALIZED SUPER/SUB-SOLUTION FOR A PERIODIC PARABOLIC OPERATOR

In this section, we introduce a generalized definition of super/sub-solution for a time-periodic parabolic operator and then present a comparison result. This result is a mortification of Proposition A.1 in [18], and it plays a vital role in this paper.

Let \mathcal{L} denote the following linear parabolic operator over $(0, 1) \times [0, T]$:

$$\mathcal{L} = \partial_t \varphi - a_1(x, t) \partial_{xx} - a_2(x, t) \partial_x + a_0(x, t).$$

In the sequel, we always assume $a_1(x, t) > 0$ so that \mathcal{L} is uniformly elliptic for each $t \in [0, T]$, and assume $a_0, a_1, a_2 \in C([0, 1] \times [0, T])$ are T -periodic in t .

Consider the linear parabolic problem

$$(A.1) \quad \begin{cases} \mathcal{L}\varphi = 0 & \text{in } (0, 1) \times [0, T], \\ c_1 \partial_x \varphi(0, t) - (1 - c_1)\varphi(0, t) = 0 & \text{on } [0, T], \\ c_2 \partial_x \varphi(0, t) + (1 - c_2)\varphi(1, t) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } (0, 1), \end{cases}$$

where $c_1, c_2 \in [0, 1]$. We now define the super/sub-solution corresponding to (A.1) as follows.

Definition A.1. The function $\bar{\varphi}$ in $[0, 1] \times [0, T]$ is called a super-solution of (A.1) if there exists a set \mathbb{X} consisting of at most finitely many points:

$$\mathbb{X} = \emptyset \quad \text{or} \quad \mathbb{X} = \{\kappa_i \in (0, 1) : i = 1, \dots, N\}$$

for some integer $N \geq 1$, such that

- (i) $\bar{\varphi} \in C((0, 1) \times [0, T]) \cap C^2(((0, 1) \setminus \mathbb{X}) \times [0, T])$;
- (ii) $\partial_x \bar{\varphi}(x^+, t) < \partial_x \bar{\varphi}(x^-, t)$ for every $x \in \mathbb{X}$ and $t \in [0, T]$;
- (iii) $\bar{\varphi}$ satisfies

$$\begin{cases} \mathcal{L}\bar{\varphi} \geq 0 & \text{in } ((0, 1) \setminus \mathbb{X}) \times (0, T), \\ c_1 \partial_x \varphi(0, t) - (1 - c_1) \varphi(0, t) \leq 0 & \text{on } [0, T], \\ c_2 \partial_x \varphi(1, t) + (1 - c_2) \varphi(1, t) \geq 0 & \text{on } [0, T], \\ \bar{\varphi}(x, 0) \geq \bar{\varphi}(x, T) & \text{on } (0, 1). \end{cases}$$

A super-solution $\bar{\varphi}$ is called to be strict if it is not a solution of (A.1). Moreover, a function $\underline{\varphi}$ is called a (strict) sub-solution of (A.1) if $-\underline{\varphi}$ is a (strict) super-solution.

Let $\lambda(\mathcal{L})$ denote the principal eigenvalue of the problem

$$(A.2) \quad \begin{cases} \mathcal{L}\varphi = \lambda(\mathcal{L})\varphi & \text{in } (0, 1) \times [0, T], \\ c_1 \partial_x \varphi(0, t) - (1 - c_1) \varphi(0, t) = 0 & \text{on } [0, T], \\ c_2 \partial_x \varphi(1, t) + (1 - c_2) \varphi(1, t) = 0 & \text{on } [0, T], \\ \varphi(x, 0) = \varphi(x, T) & \text{on } (0, 1). \end{cases}$$

The following result was proved in [18, Proposition A.1] for the case $c_1 = c_2 = 1$, and it can be extended to the general case $c_1, c_2 \in [0, 1]$.

Proposition A.2. *Let $\lambda(\mathcal{L})$ denote the principal eigenvalue of (A.2). If (A.1) admits some strict positive super-solution defined in Definition A.1, then $\lambda(\mathcal{L}) \geq 0$. Moreover, if (A.1) admits some strict nonnegative sub-solution defined in Definition A.1, then $\lambda(\mathcal{L}) \leq 0$.*

Remark A.3. Instead of $[0, 1] \times [0, T]$, Proposition A.2 also holds for the general domain given by $\{(x, t) : \beta_1(t) < x < \beta_2(t), t \in [0, T]\}$, where $\beta_1, \beta_2 \in C([0, T])$ satisfy $\beta_1 < \beta_2$. This fact is applied in Section 4 to prove Theorem 1.4.

ACKNOWLEDGMENTS

We sincerely thank the referees for their suggestions which help improve the manuscript.

REFERENCES

- [1] Alex Beltramo and Peter Hess, *On the principal eigenvalue of a periodic-parabolic operator*, Comm. Partial Differential Equations **9** (1984), no. 9, 919–941, DOI 10.1080/03605308408820351. MR749652
- [2] H. Berestycki and P.-L. Lions, *Some applications of the method of super and subsolutions*, Bifurcation and nonlinear eigenvalue problems (Proc., Session, Univ. Paris XIII, Villetaneuse, 1978), Lecture Notes in Math., vol. 782, Springer, Berlin, 1980, pp. 16–41. MR572249
- [3] Robert Stephen Cantrell and Chris Cosner, *Spatial ecology via reaction-diffusion equations*, Wiley Series in Mathematical and Computational Biology, John Wiley & Sons, Ltd., Chichester, 2003. MR2191264
- [4] Cécile Carrère and Grégoire Nadin, *Influence of mutations in phenotypically-structured populations in time periodic environment*, Discrete Contin. Dyn. Syst. Ser. B **25** (2020), no. 9, 3609–3630, DOI 10.3934/dcdsb.2020075. MR4128352
- [5] Xinfu Chen and Yuan Lou, *Principal eigenvalue and eigenfunctions of an elliptic operator with large advection and its application to a competition model*, Indiana Univ. Math. J. **57** (2008), no. 2, 627–658, DOI 10.1512/iumj.2008.57.3204. MR2414330

- [6] Xinfu Chen and Yuan Lou, *Effects of diffusion and advection on the smallest eigenvalue of an elliptic operator and their applications*, Indiana Univ. Math. J. **61** (2012), no. 1, 45–80, DOI 10.1512/iumj.2012.61.4518. MR3029392
- [7] Allen Devinatz, Richard Ellis, and Avner Friedman, *The asymptotic behavior of the first real eigenvalue of second order elliptic operators with a small parameter in the highest derivatives. II*, Indiana Univ. Math. J. **23** (1973/74), 991–1011, DOI 10.1512/iumj.1974.23.23081. MR344709
- [8] Susely Figueroa Iglesias and Sepideh Mirrahimi, *Long time evolutionary dynamics of phenotypically structured populations in time-periodic environments*, SIAM J. Math. Anal. **50** (2018), no. 5, 5537–5568, DOI 10.1137/18M1175185. MR3864685
- [9] S. Figueroa Iglesias and S. Mirrahimi, *Selection and mutation in a shifting and fluctuating environment*, preprint, 2019, <https://hal.archives-ouvertes.fr/hal-02320525>.
- [10] Avner Friedman, *The asymptotic behavior of the first real eigenvalue of a second order elliptic operator with a small parameter in the highest derivatives*, Indiana Univ. Math. J. **22** (1972/73), 1005–1015, DOI 10.1512/iumj.1973.22.22084. MR320551
- [11] M. I. Freidlin and A. D. Wentzell, *Random perturbations of dynamical systems*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 260, Springer-Verlag, New York, 1984. Translated from the Russian by Joseph Szücs. MR722136
- [12] Peter Hess, *Periodic-parabolic boundary value problems and positivity*, Pitman Research Notes in Mathematics Series, vol. 247, Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1991. MR1100011
- [13] V. Hutson, W. Shen, and G. T. Vickers, *Estimates for the principal spectrum point for certain time-dependent parabolic operators*, Proc. Amer. Math. Soc. **129** (2001), no. 6, 1669–1679, DOI 10.1090/S0002-9939-00-05808-1. MR1814096
- [14] V. Hutson, K. Mischaikow, and P. Poláčik, *The evolution of dispersal rates in a heterogeneous time-periodic environment*, J. Math. Biol. **43** (2001), no. 6, 501–533, DOI 10.1007/s002850100106. MR1874400
- [15] King-Yeung Lam and Yuan Lou, *Asymptotic behavior of the principal eigenvalue for cooperative elliptic systems and applications*, J. Dynam. Differential Equations **28** (2016), no. 1, 29–48, DOI 10.1007/s10884-015-9504-4. MR3455183
- [16] S. Liu and Y. Lou, *A functional approach towards eigenvalue problems associated with incompressible flow*, Discrete Contin. Dyn. Syst.-A, **40** (2020) 3715–3736.
- [17] Shuang Liu, Yuan Lou, Rui Peng, and Maolin Zhou, *Monotonicity of the principal eigenvalue for a linear time-periodic parabolic operator*, Proc. Amer. Math. Soc. **147** (2019), no. 12, 5291–5302, DOI 10.1090/proc/14653. MR4021088
- [18] S. Liu, Y. Lou, R. Peng, M. Zhou, *Asymptotics of the principal eigenvalue for a linear time-periodic parabolic operator I: Large advection*, submitted.
- [19] Grégoire Nadin, *The principal eigenvalue of a space-time periodic parabolic operator*, Ann. Mat. Pura Appl. (4) **188** (2009), no. 2, 269–295, DOI 10.1007/s10231-008-0075-4. MR2491804
- [20] Grégoire Nadin, *Some dependence results between the spreading speed and the coefficients of the space-time periodic Fisher-KPP equation*, European J. Appl. Math. **22** (2011), no. 2, 169–185, DOI 10.1017/S0956792511000027. MR2774781
- [21] Rui Peng, Guanghui Zhang, and Maolin Zhou, *Asymptotic behavior of the principal eigenvalue of a linear second order elliptic operator with small/large diffusion coefficient*, SIAM J. Math. Anal. **51** (2019), no. 6, 4724–4753, DOI 10.1137/18M1217577. MR4033689
- [22] Rui Peng and Xiao-Qiang Zhao, *Effects of diffusion and advection on the principal eigenvalue of a periodic-parabolic problem with applications*, Calc. Var. Partial Differential Equations **54** (2015), no. 2, 1611–1642, DOI 10.1007/s00526-015-0838-x. MR3396426
- [23] Rui Peng and Maolin Zhou, *Effects of large degenerate advection and boundary conditions on the principal eigenvalue and its eigenfunction of a linear second-order elliptic operator*, Indiana Univ. Math. J. **67** (2018), no. 6, 2523–2568, DOI 10.1512/iumj.2018.67.7547. MR3900377
- [24] A. D. Wentzell, *On the asymptotic behavior of the first eigenvalue of a second order differential operator with small parameter in higher derivatives*, Theory Probab. Appl., **20** (1975) 599–602.

INSTITUTE FOR MATHEMATICAL SCIENCES, RENMIN UNIVERSITY OF CHINA, BEIJING 100872,
PEOPLE'S REPUBLIC OF CHINA

Email address: liushuangnqkg@ruc.edu.cn

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OHIO 43210
Email address: lou@math.ohio-state.edu

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGSU NORMAL UNIVERSITY, XUZHOU, JIANGSU
221116, PEOPLE'S REPUBLIC OF CHINA

Email address: pengrui_seu@163.com

CHERN INSTITUTE OF MATHEMATICS AND LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEO-
PLE'S REPUBLIC OF CHINA

Email address: zhouml123@nankai.edu.cn