

Notes on *Trochila* (Ascomycota, Leotiomyces), with new species and combinations

Paula Andrea Gómez-Zapata¹, Danny Haelewaters^{1,2,3,4}, Luis Quijada^{2,3},
Donald H. Pfister^{2,3}, M. Catherine Aime¹

1 Department of Botany and Plant Pathology, Purdue University, West Lafayette, Indiana, USA **2** Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, USA **3** Farlow Herbarium and Reference Library of Cryptogamic Botany, Harvard University Herbaria, Harvard University, 22 Divinity Avenue, Cambridge, MA 02138, USA **4** Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic

Corresponding author: Paula Andrea Gómez-Zapata (pgomezza@purdue.edu),
Danny Haelewaters (danny.haelewaters@gmail.com),
M. Catherine Aime (maime@purdue.edu)

Academic editor: N. Wijayawardene | Received 15 December 2020 | Accepted 7 January 2021 | Published 11 February 2021

Citation: Gómez-Zapata PA, Haelewaters D, Quijada L, Pfister DH, Aime MC (2021) Notes on *Trochila* (Ascomycota, Leotiomyces), with new species and combinations. MycoKeys 78: 21–47. <https://doi.org/10.3897/mycokeys.78.62046>

Abstract

Studies of *Trochila* (Leotiomyces, Helotiales, Cenangiaceae) are scarce. Here, we describe two new species based on molecular phylogenetic data and morphology. *Trochila bostonensis* was collected at the Boston Harbor Islands National Recreation Area, Massachusetts. It was found on the stem of *Asclepias syriaca*, representing the first report of any *Trochila* species from a plant host in the family Apocynaceae. *Trochila urediniophila* is associated with the uredinia of the rust fungus *Cerotelium fici*. It was discovered during a survey for rust hyperparasites conducted at the Arthur Fungarium, in a single sample from 1912 collected in Trinidad. Macro- and micromorphological descriptions, illustrations, and molecular phylogenetic analyses are presented. The two new species are placed in *Trochila* with high support in both our six-locus (SSU, ITS, LSU, *rpb1*, *rpb2*, *tef1*) and two-locus (ITS, LSU) phylogenetic reconstructions. In addition, two species are combined in *Trochila*: *Trochila colensoi* (formerly placed in *Pseudopeziza*) and *T. xishuangbanna* (originally described as the only species in *Calycellinopsis*). This study reveals new host plant families, a new ecological strategy, and a new country record for the genus *Trochila*. Finally, our work emphasizes the importance of specimens deposited in biological collections such as fungaria.

Keywords

4 new taxa, biological collections, Boston Harbor Islands, fungarium specimens, fungicolous fungi, South America, taxonomy, Trinidad

Introduction

The genus *Trochila* Fr. (Ascomycota, Leotiomyces) was erected by Fries (1849) to accommodate four species previously placed in *Phacidium* Fr., *Sphaeria* Haller, and *Xyloma* Pers. *Trochila craterium* (DC) Fr. was the first species listed by Fries, based on *Sphaeria craterium* DC., which was later selected by Clements and Shear (1931) as the type species of *Trochila*. The other three species included by Fries (1849) were: *T. ilicis* (Fr.) Fr. [= *Sphaeria ilicis* Fr.], *T. laurocerasi* (Desm.) Fr. [= *Phacidium laurocerasi* Desm.], and *T. taxi* (Fr.) Fr. [= *Xyloma taxi* Fr.]. Only the genus and one species (*T. laurocerasi*) were briefly described by Fries (1849). However, the type species, *T. craterium*, was well described macromorphologically by Lamarck and de Candolle (1805). The description can be translated loosely from French as “a fungus growing on the lower surface of ivy leaves, initially forming a flat white disc, then turning blackish and concave opening by a split along radial lines, the disc usually surrounded by a whitish membrane” (Lamarck and de Candolle 1805). Later, the generic concept was expanded to include other types of apothecial opening. Rehm (1896) remarked that the covering layer of the apothecia could also open completely like a lid depending on host characters such as cuticle thickness. After the inclusion of this new character describing the genus, *Stegia ilicis* (Chevall.) Gillet was transferred as *Trochila ilicina* (Nees ex Fr.) Courtec (Crouan and Crouan 1867; Rehm 1896).

In our current circumscription of the genus *Trochila*, apothecia are sunken in the host tissues and hymenia are exposed either by splitting along radial lines or by splitting into a number of lobes that roll outward exposing the hymenium. The excipulum is composed of dark, globose-angular cells; asci contain eight ellipsoid, hyaline ascospores with oil guttules (except *T. substrictica* Rehm and *T. tetraspora* E. Müll. & Gamundí, which both have asci containing four ascospores); and paraphyses possess yellowish guttules (Dennis 1978; Baral and Marson 2005). Thirty-three names have been applied in the genus (Index Fungorum 2021). Jaklitsch et al. (2016) suggest that only ca. 10 names should be accepted.

Fries (1849) included *Trochila* in “Patellariacei” (= Patellariaceae). Later, it was transferred to Dermateaceae, Helotiales (Fuckel 1869; Karsten 1869; Saccardo 1884; Lambotte 1888). *Trochila* remained in this family (Korf 1973; Dennis 1978) into the molecular era (Lumbsch and Huhndorf 2010). Jaklitsch et al. (2016) placed *Trochila* in the resurrected family Cenangiaceae based on morphological and molecular data. Later, the relationships among genera in this family were supported in another, 5–15-locus phylogeny of Leotiomyces (Johnston et al. 2019).

Most species of *Trochila* have been described from their sexual morph. The asexual morph has the characteristics of the form-genus *Cryptocline* Petr. (Morgan-Jones 1973; Kiffer and Morelet 2000; Hyde et al. 2011). Two species of *Trochila* have been linked to their asexual morphs: *T. craterium* to *C. paradoxa* (De Not.) Arx and *T. laurocerasi* to *C. phacidiella* (Grove) Arx (von Arx 1957). The paucity of culture and molecular data of both *Cryptocline* and *Trochila* species has hindered the linkage of sexual and asexual morphs for most species. *Trochila viburnicola* Crous & Denman was the first species

of the genus to be described based on the combination of morphology and molecular data, but only its asexual morph is known (Crous et al. 2018). The species was named referring to its host, *Viburnum* sp. (Dipsacales, Adoxaceae). In addition to *T. viburnicola*, two other species have been reported on this host genus, but only from their sexual morph, *T. ramulorum* Feltgen and *T. tini* (Duby) Quél. [currently *Pyrenopeziza tini* (Duby) Nannf.]. Due to the lack of sequences or cultures of these two species, a comparison with *T. viburnicola* is impossible (Feltgen 1903; Crous et al. 2018).

Most *Trochila* members have a restricted record of geographical distribution and ecological strategy. *Trochila* records typically originate from the Northern Hemisphere limited to temperate regions in Europe and North America (Ziolo et al. 2005; Stoykov and Assyov 2009; Crous et al. 2018; Stoykov 2019; Global Biodiversity Information Facility 2020). Nonetheless, a number of putative *Trochila* reports are known from the Southern hemisphere (Spegazzini 1888, 1910, 1921; Rehm 1909; Gamundí et al. 1978). In addition, species of *Trochila* are typically recorded as saprotrophs on dead leaves and branches of both herbaceous plants and trees. However, a few species have been found infecting living plant tissues. *Trochila ilicina* is reported as both a weak parasite and a saprotroph because of its presence on living, decaying, and fallen leaves of *Ilex aquifolium* (Aquifoliales, Aquifoliaceae) (Ziolo et al. 2005), *T. laurocerasi* as a parasite of living leaves of *Prunus laurocerasus* (Rosales, Rosaceae) (Gregor 1936), and *T. symploci* as a pathogen of living leaves of *Symplocos japonica* (Ericales, Symplocaceae) (Hennings 1900; Stevenson 1926).

Here, we describe two new species, *T. bostonensis* and *T. urediniophila*, collected at the Boston Harbor Islands National Recreation Area, Massachusetts and at Port of Spain, Trinidad, respectively. We also make two new combinations in *Trochila* based on morphological studies and phylogenetic analyses. We reveal two new host plant families (Apocynaceae and Asparagaceae) and a new ecological strategy (fungicolous symbiont) for the genus. Finally, we provide a comparative table of characters, based on literature review, for all currently accepted species of *Trochila* (*sensu* Index Fungorum 2021).

Material and methods

Collected samples

Samples were collected in the field and from fungaria. One collection of *Trochila* was discovered during the Boston Harbor Islands (BHI) National Recreation Area fungal ATBI (Haelewaters et al. 2018a). In this project, above-ground, ephemeral fruiting bodies of non-lichenized fungi were collected. In the field, specimens were placed in plastic containers or brown paper bags. BHI-F collection numbers were assigned. Date, specific locality when applicable, GPS coordinates, substrate, and habitat notes were recorded. Specimens were dried using a Presto Dehydro food dehydrator (National Presto Industries, Eau Claire, Wisconsin) set at 35 °C for 7–9 hours. Collections were packaged, labeled, and deposited at FH. A second *Trochila* collection came to our attention during

a survey for hyperparasites of rust fungi at PUR. The specimen was found on the uredinia of the rust fungus *Cerotelium fici* on the underside of *Ficus maxima* leaves. Fungarium acronyms follow Thiers (continuously updated).

Morphological studies

Methods to study the morphological characteristics of the *Trochila* specimens followed the process given in Baral (1992). Macro- and micromorphological features were examined on both fresh and dried apothecia for the specimen collected at the BHI and on dried apothecia for the specimen found at PUR. Apothecia from the BHI specimen were observed under an EZ4 stereomicroscope (Leica, Wetzlar, Germany) and studied under a B1 compound microscope (Motic, Barcelona, Spain). Apothecia from the PUR specimen were examined on an SZ2-ILTS dissecting microscope (Olympus, Center Valley, Pennsylvania) and studied using a BH2-RFCA compound microscope (Olympus). Sections of apothecia were cut free-hand and mounted in water or pre-treated in 5% KOH. Sections were also mounted in Melzer's reagent with and without KOH-pretreatment to determine dextrinoid or amyloid reactions. At least 10 measurements were made for each structure at 400–1000× magnification. Measurements for each character are given as $(a-)b-c(-d)$, with $b-c$ indicating the 95% confidence interval and a and d representing the smallest and large single measurement, respectively. Macro- and microphotographs were taken with a USB Moticam 2500 camera (Motic) (BHI specimen) or an Olympus SC30 camera (PUR specimen). Measurements were made using the following software suites: Motic Images Plus 2.0 and cellSens Standard 1.18 Imaging Software (Olympus). Color coding refers to Kelly (1965). Abbreviations were adopted from Baral (1992) and Baral and Marson (2005) as follows:

*	living state;	LBs	lipid bodies;
†	dead state;	MLZ	Melzer's reagent;
IKI	Lugol's solution;	OCI	oil content index;
KOH	potassium hydroxide;	VBs	refractive vacuolar bodies.

DNA isolation, PCR amplifications, sequencing

Genomic DNA was isolated from 1–3 apothecia per specimen using the E.Z.N.A. HP Fungal DNA Kit (Omega Bio-Tek, Norcross, Georgia), QIAamp DNA Micro Kit (Qiagen, Valencia, California), following the manufacturer's instructions, and the Extract-N-Amp Plant PCR Kit (Sigma-Aldrich, St. Louis, Missouri), following Haelewaters et al. (2018a). We amplified the following loci: nuclear small and large ribosomal subunits (SSU and LSU), internal transcribed spacer region of the ribosomal DNA (ITS), RNA polymerase II second largest subunit (*rpb2*), and translation elongation factor 1- α (*tef1*). Primer combinations were as follows: NS1/NS2 and NS1/NS4 for SSU (White et al. 1990); LR0R/LR5 for LSU (Vilgalys and Hester 1990; Hopple 1994); ITS1F/ITS4, ITS9mun/ITS4A, and ITS5/ITS2 for ITS (White et al.

1990; Gardes and Bruns 1993; Egger 1995); RPB2-5F2/fRPB2-7cR for *rpb2* (Liu et al. 1999; Sung et al. 2007); and EF1-983F/EF1-1567R and EF1-983F/EF1-2218R for *tef1* (Rehner and Buckley 2005). All 25- μ l PCR reactions were conducted on a Mastercycler ep gradient Thermal Cycler (Eppendorf model #5341, Hauppauge, New York) and consisted of 12.5 μ l of 2 $\times$ MyTaq Mix (Bioline, Swedesboro, New Jersey), 1 μ l of each 10 μ M primer, and 10.5 μ l of 1/10 diluted DNA extract. Amplifications of rDNA and *rpb2* loci were run under the following conditions: initial denaturation at 95 °C for 5 min (94 °C for LSU); followed by 40 cycles of denaturation at 95 °C for 30 sec (94 °C for LSU), annealing at 45 °C (ITS) / 50 °C (LSU) / 55 °C (SSU, *rpb2*) for 45 sec, and elongation at 72 °C for 45 sec (1 min for LSU); and final extension at 72 °C for 7 min (1 min for SSU). Amplification of *tef1* was done with a touchdown PCR as follows: initial denaturation at 95 °C for 10 min; followed by 30 cycles of 95 °C for 1 min, 62 °C for 1 min (decreasing 1 °C every 3 cycles), 72 °C for 90 sec; then 30 cycles of 95 °C for 30 sec, 55 °C for 30 sec, and 72 °C for 1 min; and final extension at 72 °C for 7 min (Don et al. 1991; Haelewaters et al. 2018b). PCR products were visualized by gel electrophoresis. Purification of successful PCR products and subsequent sequencing in both directions were outsourced to Genewiz (South Plainfield, New Jersey). Raw sequence reads were assembled and edited using Sequencher version 5.2.3 (Gene Codes Co., Ann Arbor, Michigan).

Sequence alignment and phylogenetic analysis

Edited sequences were blasted against the NCBI GenBank nucleotide database (<http://ncbi.nlm.nih.gov/blast/Blast.cgi>) to search for closest relatives. For phylogenetic placement of our isolates, we downloaded SSU, ITS, LSU, *rpb1*, *rpb2*, and *tef1* sequences of *Trochila* from GenBank. We also downloaded sequence data of selected clades of Helotiales, mainly from Pärtel et al. (2017) but also other sources (details in Table 1), as a basis for our six-locus phylogenetic analysis. We selected representative taxa of Cenangiaceae, Cordieritidaceae, Rutstroemiaceae, and Sclerotiniaceae, with taxa in the family Chlorociboriaceae serving as outgroups (Johnston et al. 2019). Alignment of DNA sequences was done for each locus separately using MUSCLE version 3.7 (Edgar 2004), available on the Cipres Science Gateway 3.3 (Miller et al. 2010). The aligned sequences for each locus were concatenated in MEGA7 (Kumar et al. 2016). Maximum likelihood (ML) inference was performed using IQ-TREE from the command line (Nguyen et al. 2015) under partitioned models (Chernomor et al. 2016). Nucleotide substitution models were selected under Akaike's information criterion corrected for small sample size (AICc) with the help of the built-in program ModelFinder (Kalyaanamoorthy et al. 2017). Ultrafast bootstrap analysis was implemented with 1000 replicates (Hoang et al. 2017).

For the purpose of species delimitation, we constructed a second dataset of ITS–LSU consisting of isolates of *Trochila* and closely related taxa in the family Cenangiaceae. We included *Trochila* spp., *Calycellinopsis xishuangbanna*, and *Pseudopeziza colensoi*, with *Cenangiopsis* spp. serving as outgroup. In this analysis, we included *T. ilicina*, for

Table 1. Sequences used in phylogenetic analyses. Accession numbers in boldface indicate sequences that were generated during the course of this study.

Isolate	Species	Family	SSU	ITS	LSU	<i>rpb1</i>	<i>rpb2</i>	<i>tef1</i>	Reference
KL391	<i>Areghiniella australis</i>	Cordieritiaceae	KX090893		KX090841	KX090787		KX090690	Parrel et al. (2017)
AD283531 [†]	<i>Annabellia australensis</i>	Cordieritiaceae		MK328475	MK328476				Fryar et al. (2019)
AFTOL-ID 59	<i>Boryotinia suckeliana</i>	Sclerotiniaceae	AY544695		AY544651	DQ471116	DQ247786	DQ471045	Spatafora et al. (2006)
HMAS:187063	<i>Calyndrinopsis xizhuangbanua</i>	Cenangiaceae	GU936124		KR094163	MH729338	MH729345		W.Y. Zhuang et al. (unpubl.)
KL375	<i>Cenangiopis alpestris</i>	Cenangiaceae			KX090837	KX090784	KX090736		Parrel et al. (2017)
KL378	<i>Cenangiopis alpestris</i>	Cenangiaceae	KX090891	LT158470	KX090839	KX090786	KX090738		Parrel et al. (2017)
KL157	<i>Cenangiopis alpestris</i>	Cenangiaceae	KX090858	LT158421	KX090806		KX090709	KX090663	Parrel et al. (2017)
KL174	<i>Cenangiopis quercicola</i>	Cenangiaceae	KX090862	LT158425	KX090811	KX090760	KX090713		Parrel et al. (2017)
KL377	<i>Cenangiopis</i> sp.	Cenangiaceae	KX090890	KX090900	KX090838	KX090785	KX090737		Parrel et al. (2017)
KL276	" <i>Cenangium</i> " <i>actum</i>	<i>Piccomphale</i> clade	KX090879	LT158445	KX090828	KX090727	KX090680		Parrel et al. (2017)
KL243	" <i>Cenangium</i> " <i>actum</i>	<i>Piccomphale</i> clade	KX090873	LT158439	KX090822	KX090767	KX090720	KX090674	Parrel et al. (2017)
KL390	<i>Cenangium ferruginosum</i>	Cenangiaceae	KX090892	LT158471	KX090840		KX090739		Parrel et al. (2017)
KL167	<i>Chloenocelia torta</i>	Cenangiaceae		LT158424	KX090810	KX090759			Parrel et al. (2017)
KP606	<i>Chloenocelia versiformis</i>	Cenangiaceae	KX090894			KX090788	KX090740	KX090692	Parrel et al. (2017)
KL21	<i>Chloenocelia versiformis</i>	Cenangiaceae	KX090846	LT158427	KX090795				Parrel et al. (2017)
KL152	<i>Chlorociboria aeruginascens</i>	Chlorociboriaceae		LT158419		KX090752	KX090706	KX090657	Parrel et al. (2017)
KL247	<i>Chlorociboria aeruginella</i>	Chlorociboriaceae				KX090769	KX090722	KX090676	Parrel et al. (2017)
KL238	<i>Chlorociboria glauca</i>	Chlorociboriaceae	KX090875	LT158438	KX090821	KX090766		KX090673	Parrel et al. (2017)
KL212	<i>Ciboria nitidifusa</i>	Sclerotiniaceae	KX090863	LT158429	KX090812				Parrel et al. (2017)
KL254	<i>Crumenulopsis sororia</i>	Cenangiaceae		LT158442	KX090826		KX090725		Parrel et al. (2017)
KL317	<i>Diplacarpa bloxamii</i>	Cordieritiaceae	KX090885		KX090834	KX090778	KX090745	KX090688	Parrel et al. (2017)
SK80	<i>Diplacarpia nanula</i>	Cordieritiaceae	KX090896	KP984782	KX090790				Parrel et al. (2017)
TU:109263	<i>Dumontinia tuberosa</i>	Sclerotiniaceae	KX090897	LT158412	KX090843	KX090792			Etayo et al. (2015), Parrel et al. (2017)
KL111	<i>Enocelia fimbriata</i>	Cenangiaceae	KX090852		KX090800		KX090703	KX090697	Parrel et al. (2017)
KL108	<i>Enocelia fufrinacea</i>	Cenangiaceae	KX090851		KX090799		KX090702	KX090654	Parrel et al. (2017)
KL107	<i>Enocelia fufrinacea</i>	Cenangiaceae	KX090850	LT158416	KX090798	KX090749	KX090701	KX090653	Parrel et al. (2017)
KL106	<i>Enocelia fufrinacea</i>	Cenangiaceae	KX090849	LT158415		KX090748		KX090652	Parrel et al. (2017)
KL92	<i>Enocelia fufrinacea</i>	Cenangiaceae	KX090847	LT158482	KX090796			KX090651	Parrel et al. (2017)
KL164	<i>Enocelia heteromera</i>	Cenangiaceae	KX090861		KX090809	KX090758	KX090712	KX090662	Parrel et al. (2017)
KL304	<i>Enocelia heteromera</i>	Cenangiaceae	KX138404			KX138400			Parrel et al. (2017)
KL244	Helotiales sp.	Cenangiaceae		LT158440	KX090823	KX090768	KX090721	KX090675	Parrel et al. (2017)
KL20	<i>Heyderia abietis</i>	Cenangiaceae	KX090874	LT158426		KX090747	KX090699	KX090650	Parrel et al. (2017)
HMAS:71954	<i>Heyderia abietis</i>	Cenangiaceae	AY789295	AY789297					Wang et al. (2005)
KL216	<i>Heyderia pusilla</i>	Cenangiaceae	KX090865	LT158430		KX090762	KX090715	KX090665	Parrel et al. (2017)
KL299	<i>Ionomidotis frondosa</i>	Cordieritiaceae	KX090882			KX090775		KX090685	Parrel et al. (2017)
KL231	<i>Ionomidotis fulvotinctus</i>	Cordieritiaceae	KX090870		KX090819	KX090765	KX090719	KX090671	Parrel et al. (2017)

Isolate	Species	Family	SSU	ITS	LSU	<i>rpb1</i>	<i>rpb2</i>	<i>ref1</i>	Reference
KL239	<i>Ionanidopsis filiostringens</i>	Cordieritiaceae	KX138403		KX138407	KX138399	KX138401		Parrel et al. (2017)
KL154	<i>Ionanidopsis irregularis</i>	Cordieritiaceae	KX090856		KX090804	KX090754		KX090658	Parrel et al. (2017)
KL301	<i>Ionanidopsis olivaceus</i>	Cordieritiaceae	KX090883		KX090833	KX090776	KX090732	KX090686	Parrel et al. (2017)
CBS:811.85	<i>Lambertella subrepenspora</i>	Rustroemiaceae	KF545416	AB956097	MH873604				Zhao et al. (2016), Parrel et al. (2017), Vu et al. (2019)
LL95	<i>Limonietta terricola</i>	Cordieritiaceae	KX090895		KX090842	KX090789	KX090741	KX090693	Parrel et al. (2017)
AFTOL-ID 169	<i>Monilinia laxa</i>	Sclerotiniaceae	AY544714		AY544670	FJ238425	DQ470889	DQ471057	Spatafora et al. (2006)
KL374	<i>Piccomphale bulgaroides</i>	Piccomphale clade	KX090889	LT158469	KX090836	KX090783			Parrel et al. (2017)
KL98	<i>Piccomphale bulgaroides</i>	Piccomphale clade	KX090848	LT158483	KX090797		KX090700		Parrel et al. (2017)
PDD:112240	<i>Pseudopeziza olenoi</i>	Cenangiaceae		MH921874	MH985297	MH986706	MH986705		P.R. Johnston and D. Park (unpubl.)
KL267	<i>Pycnopiezia sejanetii</i>	Sclerotiniaceae	KX090878	LT158443	KX090827	KX090772	KX090726	KX090679	Parrel et al. (2017)
AFTOL-ID 907	<i>Rhodocline laticis</i>	Cenangiaceae	DQ471002		DQ470954	DQ471146	DQ470904	DQ471073	Spatafora et al. (2006)
KL292	<i>Rustroemia firma</i>	Rustroemiaceae	KX090881	LT158450	KX090832	KX090774	KX090731	KX090684	Parrel et al. (2017)
KL291	<i>Rustroemia firma</i>	Rustroemiaceae		LT158449	KX090831		KX090730	KX090683	Parrel et al. (2017)
KL290	<i>Rustroemia firma</i>	Rustroemiaceae			KX090830		KX090729	KX090682	Parrel et al. (2017)
KL222	<i>Rustroemia firma</i>	Rustroemiaceae	KX138402		KX138406			KX138397	Parrel et al. (2017)
KL310	<i>Rustroemia johnstonii</i>	Rustroemiaceae	KX090884	LT158454		KX090777	KX090733	KX090687	Parrel et al. (2017)
KL234	<i>Rustroemia juniperi</i>	Rustroemiaceae	KX090871		KX090820			KX090672	Parrel et al. (2017)
KL217	<i>Rustroemia lateritrescens</i>	Rustroemiaceae		LT158431	KX090814	KX090763	KX090716	KX090666	Parrel et al. (2017)
KL160	<i>Rustroemia tiliacea</i>	Rustroemiaceae	KX090860	LT158423	KX090808	KX090757	KX090711	KX090661	Parrel et al. (2017)
KL393	Rustroemiaceae sp.	Rustroemiaceae	KX138405	LT158472	KX138408		KX138398	KX090691	Parrel et al. (2017)
KL288	Rustroemiaceae sp.	Rustroemiaceae	KX090880	LT158446	KX090829	KX090773	KX090728	KX090681	Parrel et al. (2017)
CBS:273.74 ^T	<i>Sarastrochila longispora</i>	Cenangiaceae		KJ663836	KJ663877		KJ663918		Crous et al. (2014)
KL347	<i>Sclerochaeta fascicularis</i>	Sclerotiniaceae			KX090782				Parrel et al. (2017)
KL156	<i>Sclerochaeta fascicularis</i>	Sclerotiniaceae	KX090857		KX090805	KX090755	KX090708	KX090659	Parrel et al. (2017)
KL344	<i>Sclerochaeta pruinosula</i>	Sclerotiniaceae	KX090888			KX090781	KX090735		Parrel et al. (2017)
CBS:499.50	<i>Sclerotinia sclerotiorum</i>	Sclerotiniaceae	DQ471013		DQ470965		DQ470916		Spatafora et al. (2006)
NY:01231276	<i>Skeptea radiatilis</i>	Cordieritiaceae		KJ559538	KJ559560	KX090791	KX090742	KX090694	Sujia et al. (2015), Parrel et al. (2017)
TH90	<i>Thamnogella crombiei</i>	Cordieritiaceae	KJ559583	KJ559535	KJ559557		KX090743	KX090695	Parrel et al. (2017)
BH1-P974a ^T	<i>Trochila bostonensis</i>	Cenangiaceae	MT873949	MT873947	MT873952		MT861181	MT861183	This study
BH1-P974b ^T	<i>Trochila bostonensis</i>	Cenangiaceae	MT873950	MT873948	MT873948		MT861182	MT861184	This study
KL332	<i>Trochila craterium</i>	Cenangiaceae	KX090886			KX090779			Parrel et al. (2017)
KL336	<i>Trochila lauracensis</i>	Cenangiaceae	KX090887	LT158460	KX090835	KX090780	KX090734	KX090689	Parrel et al. (2017)
F18316 ^T	<i>Trochila undinophila</i>	Cenangiaceae		MT873946	MT873951				This study
CBS:144206 ^T	<i>Trochila viburnicola</i>	Cenangiaceae		MH107921	MH107967		MH108011	MH108031	Crous et al. (2018)
KL253	<i>Velutaria rufopoliacea</i>	Cenangiaceae	KX090877		KX090825	KX090771	KX090724	KX090678	Parrel et al. (2017)

which only a single ITS sequence is available. The same methods as above were applied: alignment using MUSCLE (Edgar 2004), selection of nucleotide substitution models with the help of ModelFinder (Kalyaanamoorthy et al. 2017), ML using IQ-TREE (Nguyen et al. 2015; Chernomor et al. 2016; Hoang et al. 2017). Phylogenetic reconstructions with bootstrap values (BS) were visualized in FigTree version 1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Results

Nucleotide alignment dataset and phylogenetic inferences

The concatenated six-locus dataset consisted of 11343 characters, of which 2655 were parsimony-informative. The percentage of parsimony-informative characters per locus was 9.3% for SSU, 48.1% for ITS, 21.4% for LSU, 48.9% for *rpb1*, 30.0% for *rpb2*, and 19.2% for *tef1*. A total of 71 isolates were included, of which *Chlorociboria aeruginascens* (Nyl.) Kanouse ex C.S. Ramamurthi, Korf & L.R. Batra, *C. aeruginella* (P. Karst.) Dennis, and *C. glauca* (Dennis) Baral & Pärtel (Helotiales, Chlorociboriaceae) served as outgroup taxa. The following models were selected by ModelFinder (AICc): TNe+R3 (SSU, $-\ln L = 23478.796$); GTR+F+I+G4 (ITS, $-\ln L = 18385.043$); TN+F+R4 (LSU, $-\ln L = 28398.591$); SYM+I+G4 (*rpb1*, $-\ln L = 41387.214$); GTR+F+R10 (*rpb2*, $-\ln L = 57025.083$); and GTR+F+R8 (*tef1*, $-\ln L = 35467.940$). Our ML analysis reveals five high to maximum-supported clades (Fig. 1): Cenangiaceae, Cordieritidaceae, Rutstroemiaceae, Sclerotiniaceae, and a clade with *Piceomphale bulgarioides* (P. Karst.) Svrček and “*Cenangium*” *acuum* Cooke & Peck (*Piceomphale* clade *sensu* Pärtel et al. 2017). As previously reported (e.g., Pärtel et al. 2017; Johnston et al. 2019), several genera in their current circumscription are polyphyletic: *Encoelia* (Fr.) P. Karst. in Cenangiaceae and Rutstroemiaceae, *Ionomidotis* E.J. Durand ex Thaxt. in Cordieritidaceae, *Rutstroemia* P. Karst. in Rutstroemiaceae, and *Trochila* in Cenangiaceae. *Trochila laurocerasi* is placed as a sister taxon to *Calycellinopsis xishuangbanna* W.Y. Zhuang and *Pseudopeziza colensoi* (Berk.) Massee. The other species of *Trochila*, including the type species *T. craterium* and the here described species, form a monophyletic clade (BS = 81).

The second two-locus dataset consisted of 2284 characters (ITS: 924, LSU: 1360), of which 2040 were parsimony-informative (ITS: 782, LSU: 1258). A total of 13 isolates were included, of which *Cenangiosis alpestris* (Baral & B. Perić) Baral, B. Perić & Pärtel, *C. quercicola* (Romell) Rehm, and *Cenangiosis* sp. served as outgroup taxa. The following models were selected by ModelFinder (AICc): GTR+F+I+G4 (ITS, $-\ln L = 5810.483$) and TIM+F+R2 (LSU, $-\ln L = 5595.374$). *Calycellinopsis xishuangbanna*, *Pseudopeziza colensoi*, and all *Trochila* species form a monophyletic clade with high support (BS = 96) (Fig. 2). Both new species of *Trochila* are distinct from previously described species. The undescribed *Trochila* species found on uredinia of *Cerotelium fici* is retrieved as sister to *T. viburnicola* (BS = 90).




Figure 1. The best-scoring ML tree ($-\ln L = 87544.854$) of Cenangiaceae, Cordieritidaceae, Rutstroemiaceae, Sclerotiniaceae, and the *Piceomphale* clade, reconstructed from a concatenated six-locus dataset (SSU, ITS, LSU, *rpb1*, *rpb2*, and *tef1*). For each node, the ML bootstrap value (if ≥ 70) is presented above or in front of the branch leading to that node. The arrow denotes the genus *Trochila*. Species with an asterisk (*) are treated in the Taxonomy section.




Figure 2. The best-scoring ML tree ($-\ln L = 5225.551$) of Cenangiaceae, reconstructed from a concatenated ITS–LSU dataset. For each node, the ML bootstrap value (if ≥ 70) is presented above the branch leading to that node. Species treated in the Taxonomy section are highlighted with gray shading.

Taxonomy

Leotiomyces O.E. Erikss. & Winka

Helotiales Nannf. ex Korf & Lizoň

Cenangiaceae Rehm

***Trochila bostonensis* Quijada & Haelew, sp. nov.**

Mycobank No: 836582

Fig. 3

Diagnosis. Differs from *Trochila craterium* and *T. laurocerasi* in its host (Apocynaceae), sizes of asci ($57\text{--}65.5 \times 5\text{--}6 \mu\text{m}$) and ascospores ($6.2\text{--}7.2 \times 2.6\text{--}2.8 \mu\text{m}$), and the inamyloidity of its ascus apex.

Type. Holotype: USA, Massachusetts, Boston Harbor Islands National Recreation Area, Plymouth County, Great Brewster Island, 42.3310722°N , 70.8977667°W , alt. 10 m a.s.l., 16 Oct 2017, leg. D. Haelewaters, J.K. Mitchell & L. Quijada, on hollow dead stem of *Asclepias syriaca* (Gentianales, Apocynaceae), FH:BHI-F0974. Ex-holotype sequences: isolates BHI-F0974a (1 apothecium, SSU: MT873949,




Figure 3. Morphological features of *Trochila bostonensis* (holotype collection FH:BHI-F0974) **a1–3, a5** fresh apothecia **a4** dried apothecia **b1** excipular tissues in median section **b2** cells at the base **b3** cells at the upper and lower flank **c1, c2** paraphyses **d1, d2** asci **d3** ascus pore with inamyloid reaction **d4** crozier at ascus base **e1–e6** ascospores. Mounted in: Congo Red (**c2, d2, d4, e3, e5**), H₂O (**b1–b3, c1, d1, e1, e2**), KOH (**e4**), MLZ (**d3, e6**). Scale bars: 500 μ m (**a1–a5**); 50 μ m (**b1**); 10 μ m (**b1, b2, c1, c2, d1–d4, e1–e6**).

ITS: MT873947, LSU: MT873952, *rpb2*: MT861181, *tefl*: MT861183) and BHI-F0974b (1 apothecium, SSU: MT873950, ITS: MT873948, LSU: MT873953, *rpb2*: MT861182, *tefl*: MT861184).

Etymology. *bostonensis* – referring to Boston, Massachusetts, the locality of the type collection.

Description. *Apothecia* erumpent singly or in groups of 2–3, protruding from the bark by lifting and rolling outward the host periderm, sessile on a broad base, closed and barely visible when dry, rehydrated 0.4–1.1 mm diam., 0.1–0.2 mm thick; mature flat to slightly cupulate, dark grayish red brown (47.D.gy.r.Br) to black (267.Black). Margin toothed and lighter than the disc, apothecia star-shaped, with 3–6 teeth of 0.1–0.3 mm in length, each tooth deep yellowish brown (75.deepyBr). *Asci* $^*(46.5\text{--}55.5\text{--}66.5(-73) \times (5.5\text{--}6.0\text{--}6.5(-7.0) \mu\text{m})$, $^\dagger(50.5\text{--}57\text{--}65.5(-66) \times (4.5\text{--}5.0\text{--}6.0 \mu\text{m})$, 8-spored, cylindrical, pars sporifera $^*30\text{--}52 \mu\text{m}$; apex rounded to subconical, inamyloid (IKI, KOH-pretreated or not), slightly thick-walled at apex, lateral walls thin; base slightly tapered and arising from croziers. *Ascospores* $^*(6.3\text{--}6.7\text{--}7.7(-8.6) \times 2.7\text{--}3.4 \mu\text{m}$, $^\dagger(5.8\text{--}6.2\text{--}7.2 \times 2.6\text{--}2.8 \mu\text{m})$, ellipsoid-cuneate, inequilateral, ends rounded or subacute, aseptate, hyaline, smooth, thick-walled, oligoguttulate, containing 2–5 grayish yellow (90.gy.Y) oil drops (LBs), $1\text{--}2.4 \mu\text{m}$ diam., OCI = $(45\text{--}60\text{--}75(-90)\%$. *Paraphyses* slightly to medium clavate, terminal cell $^*(17.5\text{--}18\text{--}23(-29.5) \times 3\text{--}4 \mu\text{m}$, secondary cells $^*(8\text{--}9\text{--}10(-11) \times 2.5\text{--}3 \mu\text{m}$, lower cells $^*(7.5\text{--}8.5\text{--}10.5(-11.5) \times 2.5\text{--}3 \mu\text{m}$, unbranched, thin-walled, smooth, with one or several cylindric to globose refractive drops (VBs, not present after KOH-pretreated), $^*3.5\text{--}14 \times 2\text{--}3.5 \mu\text{m}$. *Medullary excipulum* $17.5\text{--}54 \mu\text{m}$ thick, grey yellowish brown (80.gy.yBr), upper part of *textura porrecta*, lower part dense *textura intricata*, cells with tiny globose deep yellow (85.deepyY) refractive drops (VBs). *Ectal excipulum* of thin-walled *textura globulosa-angularis* at base and lower flanks, dark yellowish brown (78.d.yBr) to dark brown (59.d.Br), $(40\text{--})55\text{--}78 \mu\text{m}$ thick, cells $^*(7.0\text{--}9.5\text{--}13(-15.5) \times (3.0\text{--})5.0\text{--}8.5(-10) \mu\text{m}$; at upper flanks and margin of *textura prismatica*, $30\text{--}40 \mu\text{m}$ thick, cells $^*(5.5\text{--})6.5\text{--}7.5(-8.5) \times 2.5\text{--}3.5 \mu\text{m}$, entirely without drops and slightly gelatinized, cells slightly thick-walled with irregular patches of dark brown exudates in areas of mutual contact, cortical cells in flanks covered by amorphous refractive deep yellow (88.d.Y) granular exudates, at margin some cells protruding like short hairs ($^*6.5\text{--}14 \times 2.5\text{--}3.5 \mu\text{m}$). *Asexual state* unknown.

Notes. *Trochila bostonensis* is the only species of the genus found on a member of Apocynaceae (Table 2). It was growing in the outer layer of a dead stem of *Asclepias syriaca*, which had fallen on the ground. The host was close to the shore in a shrubby thicket of *Rhus*. There are two similar species. *Trochila laurocerasi* has wider asci ($6.0\text{--}8.0 \mu\text{m}$ vs. $4.5\text{--}6.0 \mu\text{m}$) and larger ascospores ($6.3\text{--}10 \times 2.5\text{--}4.6 \mu\text{m}$ vs. $5.8\text{--}7.2 \times 2.6\text{--}2.8 \mu\text{m}$) compared to *T. bostonensis*. Ascus and ascospore length are similar in *T. bostonensis* and *T. craterium*, although ascospores are slightly larger in *T. craterium*. The two species mostly differ in the width of their asci ($7\text{--}12 \mu\text{m}$ in *T. craterium* vs. $4.5\text{--}6.0 \mu\text{m}$ in *T. bostonensis*). We used the measurements in dead state to compare *T. bostonensis* with other species in the genus (see Table 2).

Table 2. Comparative table of currently accepted species of *Trochila* (except *T. viburnicola*). For each species, the following characters are presented: host plant, host family, measurements of asci and ascospores (dead state). The asterisk (*) indicates a fungal host.

Species	Host Plant	Host Family	Asci (μm)		Ascospores (μm)		Reference
			Length	Width	Length	Width	
<i>T. andromedae</i>	<i>Andromeda polifolia</i>	Ericaceae	80	12	15–18	4–5	Karsten (1871)
<i>T. astragali</i>	<i>Astragalus glycyphyllos</i>	Fabaceae	50–60	6–7	8	4	Rehm (1896)
<i>T. atrosanguinea</i>	<i>Carex rigida</i>	Cyperaceae	45–68	7–8	7–8	2–3	Rostrup (1885)
	<i>Carex vulgaris</i>	Cyperaceae					
<i>T. bostonensis</i>	<i>Asclepias syriaca</i>	Apocynaceae	(50.5)57–65.5(66)	(4.5)5–6	(5.8)6.2–7.2	2.6–2.8	This study
<i>T. chilensis</i>	<i>Lardizabala biternata</i>	Lardizabaleae	70–80	8–9	14–15	4	Spegazzini (1910)
<i>T. cinerea</i>	<i>Pyrola</i> sp.	Ericaceae	no data	no data	6–7	1.5	Patouillard (1886)
<i>T. colensoi</i>	<i>Cordyline</i> sp.	Asparagaceae	60–70	8–10	9–12.5	3.5–5	Dennis (1961)
<i>T. conioselinii</i>	<i>Conioselinum</i> sp.	Apiaceae	38–40	6–7	10–13	3	Rostrup (1886)
	<i>Gmelina</i> sp.	Apiaceae					
<i>T. craterium</i>	<i>Cassiope tetragona</i>	Araliaceae	50–60	8–12	6–8	4–5	Rehm (1896)
	<i>Hedera algeriensis</i>	Araliaceae	no data	7	6–8.2	3–4.5	Greenhalgh and Morgan-Jones (1964)
	<i>Hedera helix</i>	Araliaceae					
<i>T. epilobii</i>	<i>Epilobium angustifolium</i>	Onagraceae	75–95	17–20	15–17	8	Karsten (1871)
<i>T. exigua</i>	<i>Nardus stricta</i>	Poaceae	32	6	8–10	0.8	Rostrup (1888)
<i>T. fallens</i>	<i>Salix</i> sp.	Salicaceae	50–60	7–9	9–14	3.5–4.5	Karsten (1871)
<i>T. ilicina</i>	<i>Ilex aquifolia</i>	Aquifoliaceae	75–80	9–10	9–11	3.5–4.5	Rehm (1896)
	<i>Ilex aquifolium</i>	Aquifoliaceae	60–76	8.5–10	10–12.5	3.5–4.5	Greenhalgh and Morgan-Jones (1964)
	<i>Ilex colchica</i>	Aquifoliaceae					
	<i>Ilex platyphylla</i>	Aquifoliaceae	57.6–93.4	6.6–9.6	9.8–15.9	2.7–5.1	Ziolo et al. (2005)
<i>T. jaffuelii</i>	<i>Lapageria rosea</i>	Philesiaceae	50–70	25	13–14	6–7	Spegazzini (1921)
<i>T. juncicola</i>	<i>Juncus compressus</i>	Juncaceae	40–45	5–6	8–9	1–1.5	Rostrup (1886)
<i>T. laurocerasi</i>	<i>Laurocerasus officinalis</i>	Rosaceae	45–60	8–9	7–10	3.5–4	Rehm (1896)
	<i>Photinia serrulata</i>	Rosaceae					Greenhalgh and Morgan-Jones (1964)
	<i>Prunus laurocerasus</i>	Rosaceae	50–65	6–9	7.5–10	3–3.75	
	<i>Prunus lusitanica</i>	Rosaceae					
<i>T. leopoldina</i>	<i>Nectandra rigida</i>	Lauraceae	45–50	7	8–9	3	Rehm (1909)
<i>T. majalis</i>	<i>Fagus sylvatica</i>	Fagaceae	38–45	7–8	7–9	3–3.5	Kirschstein (1944)
<i>T. molluginea</i>	<i>Galium molluginis</i>	Rubiaceae	55–60	7	10–12	2.5	Mouton (1900)
<i>T. oleae</i>	<i>Olea europaea</i>	Oleaceae	no data	no data	no data	no data	Fries (1849)
<i>T. oxycoccus</i>	<i>Vaccinium oxycoccus</i>	Ericaceae	60–70	11–14	14–18	5	Karsten (1871)
<i>T. perexigua</i>	<i>Hippophae rhamnoides</i>	Elaeagnaceae	80	15	14	7	Spegazzini (1881)
<i>T. perseae</i>	<i>Persea lingue</i>	Lauraceae	50–60	10	9–10	3	Spegazzini (1910)
<i>T. plantaginea</i>	<i>Plantago major</i>	Plantaginaceae	42–50	12–16	18–25	4–4.5	Karsten (1871)
<i>T. prominula</i>	<i>Juniperus sabina</i>	Cupressaceae	65–70	10–12	18–20	6	Saccardo (1878)
<i>T. puccinioidea</i>	<i>Carex</i> sp.	Cyperaceae	no data	no data	no data	no data	De Notaris (1863)
<i>T. ramulorum</i>	<i>Viburnum opulus</i>	Viburnaceae	40–55	5.5–7	5–7	1.5–2	Felgen (1903)
<i>T. rhodiolae</i>	<i>Rhodiola</i> sp.	Crassulaceae	40	5–6	10	1–1.5	Rostrup (1891)
<i>T. staritziana</i>	<i>Ailanthus glandulosa</i>	Simaroubaceae	no data	no data	no data	no data	Kirschstein (1941)
	<i>Rhus glabra</i>	Anacardiaceae					
<i>T. substricta</i>	<i>Solidago virgaurea</i>	Asteraceae	60	9	12–14	6	Rehm (1884)
<i>T. symploci</i>	<i>Symplocos japonica</i>	Symplocaceae	65–85	5–7	8–11	4–5	Hennings (1900)
<i>T. tami</i>	<i>Tamus communis</i>	Dioscoreaceae	40–55	6–7	5–8	2.5–4	Grelet and de Crozals (1928)
<i>T. tetraspora</i>	<i>Nothofagus dombeyi</i>	Nothofagaceae	58–72	7.7–9.6	12–15	3.4–4.8	Gamundi et al. (1978)
<i>T. urediniophila</i>	<i>Cerotelium fici</i> *	Phakopsoraceae	(86.4)102.4–111.2(121.8)	(9.1)10.5–11.6(13.1)	(7.6)9.0–9.7(10.9)	(5.1)6.3–7.1(8.1)	This study
<i>T. xishuangbanna</i>	no data	no data	55–60	3.5–4	8–11	1.2–1.7	Zhuang et al. (1990)
<i>T. winteri</i>	<i>Drymis Winteri</i>	Winteraceae	40–50	10–12	12–13	5	Spegazzini (1888)

***Trochila urediniophila* Gomez-Zap., Haelew. & Aime, sp. nov.**

Mycobank No: 836583

Fig. 4

Diagnosis. Differs from *Trochila ilicina* in ecological strategy (fungicolous symbiont); sizes of asci (102.4–111.2 × 10.5–11.6 µm), ascospores (9.0–9.7 × 6.3–7.1 µm), paraphyses (3.2–3.6 µm wide); and the inamyloidity of its ascus apex.

Type. Holotype: Reliquiae Farlowiana No. 723; Trinidad and Tobago, Port of Spain, Trinidad, Maraval Valley, ca. 10.5°N, 61.25°W, alt. ±301 m a.s.l., 1 Apr 1912, leg. R. Thaxter, on uredinia of *Cerotelium fici* [as *Phakopsora nishidana*] (Pucciniales, Phakopsoraceae) on the underside of *Ficus maxima* (Rosales, Moraceae) leaves, PUL F27668 (ex-PUR F18316). Ex-holotype sequences: isolate F18316 (3 apothecia, ITS: MT873946, LSU: MT873951).

Etymology. Referring to the intimate association of the fungus with the uredinia of *Cerotelium fici*.

Description. *Apothecia* protruding from uredinia of *Cerotelium fici*, gregarious in small groups or rarely solitary, discoid to irregular-ellipsoid when crowded, 0.4–1.0 mm diam., sessile on a broad base, flat to slightly concave at maturity, dark grayish yellow brown (81.d.gy.yBr) to dark grayish brown (62.d.gy.Br), margin marked and lighter than hymenium, light grayish yellow brown (79.l.gr.yBr) to medium yellow brown (77.m.yBr), receptacle concolor with margin and surface slightly pruinose. *Asci* †(86.4–)102.4–111.2(–121.8) × (9.1–)10.5–11.6(–13.1) µm, 8-spored, cylindrical, †uniseriate; apex rounded to subconical, inamyloid (IKI, KOH-pretreated or not), base arising from croziers. *Ascospores* †(7.6–)9.0–9.7(–10.9) × (5.1–)6.3–7.1(–8.1) µm, ovoid to ellipsoid, aseptate, hyaline, smooth-walled, guttulate, containing †one to two pale yellow (89.p.Y) to yellow gray (93.y Gray) oil drops (LBs), 2–5 µm diam., OCI = (40–)55.1–66.9(–81)%. *Paraphyses* cylindrical to slightly or medium clavate-spathulate, unbranched, smooth, septate, hyaline, †(2.3–)3.2–3.6(–4.1) µm wide, apex up to 6.8 µm wide. *Medullary excipulum* †17.4–79.4 µm thick, *textura intricata* strong brown (55.s.Br) to deep brown (56.deepBr). *Ectal excipulum* of *textura globulosa-angularis* at base and lower flanks, strong yellow brown (74.s.yBr) to dark brown (59.d.Br), †32.8–93.5 µm thick, cells †(7.3–)9.0–10.8(–15.3) × (6.0–)7.5–8.7(–11.5) µm; at upper flanks and margin cells vertically oriented of *textura prismatica*, 17–34 µm thick, at margin and upper flank cells protruding like short hairs, hyaline, aseptate, cylindrical, †(9.5–)16–20.6(–29.1) × (3.0–)3.9–4.5(–5.8) µm. *Asexual state* unknown.

Notes. *Trochila urediniophila* is the first known fungicolous member of the genus. The specimen described here was discovered during a survey of hyperparasites of rust fungi at PUR. Apothecia of *T. urediniophila* were never observed in direct contact with the plant tissue; instead, they grew directly on the uredinia of *Cerotelium fici* on the underside of *Ficus maxima* leaves. *Trochila ilicina* is most similar to *T. urediniophila*, but *T. urediniophila* differs from *T. ilicina* in its distinctly wider ascospores, larger




Figure 4. Morphological features of *Trochila urediniophila*, holotype collection (PUL F27668) **a1–a4** dried apothecia growing on uredinia of *Cerotelium fici* **a2, a3** substrate (uredinia) on which the apothecia grow (arrows) **b1** transverse section of apothecia; arrow pointing out the substrate **b2, b3** details of excipulum at margin and upper flanks **b4** cells at base **c1–c3** asci **d1** paraphyses **e1–e3** ascospores **e2, e3** oil drops (LBs) inside ascospores. Mounted in: Congo Red (**c1, e2**), H₂O (**b2, c3, d1, e1, e3**), KOH (**b1, b3, b4, c2**). Scale bars: 1 mm (**a1–a3**); 500 μm (**a4**); 200 μm (**b1**); 50 μm (**b2**); 20 μm (**b3, b4, c2, c3, d1**); 2 μm (**c1, e1–e3**).

asci, inamyloid ascus apex, and wider apex of the paraphyses. The uredinia of the host fungus, *C. fici*, become a solidified mass that changes in color from dark orange yellow (72.d.OY) without apothecia of *Trochila* to brownish black (65.brBlack) where apothecia are present.

A second duplicate of the Reliquiae Farlowiana No. 723 is also deposited at PUR (accession PUR F1098). However, no apothecia were present on this specimen, nor could additional specimens of *T. urediniophila* be found on any of the other specimens of *C. fici* housed at PUR. At least eight other duplicates are housed at BPI, CINC, CUP, F, ISC, MICH, and UC (MyCoPortal 2020). It is unknown whether any of them may host *T. urediniophila*.

New combinations

Trochila colensoi (Berk.) Quijada, comb. nov.

Mycobank No: 836591

≡ *Cenangium colensoi* Berk., Hooker, Bot. Antarct. Voy. Erebus Terror 1839–1843, II, Fl. Nov.-Zeal.: 201 (1855). [Basionym]

= *Pseudopeziza colensoi* (Berk.) Masee, J. Linn. Soc., Bot. 31: 468 (1896)

Notes. *Cenangium colensoi* is described from dead leaves of *Cordyline* sp. (Asparagales, Asparagaceae) in New Zealand (Hooker 1855). The host had been mistakenly reported as *Phormium* (Asparagales, Asphodelaceae) by Berkeley in Hooker (1855) and only recently corrected after re-study of the type collection (Landcare Research 2020). *Cenangium colensoi* was later combined in *Pseudopeziza* and described in more detail by Masee (1896). Both authors commented on the watery-grey disc and brownish receptacle of the apothecia. The apothecia develop among the rigid vascular bundles of the epidermis, first covered by the cuticle, then erumpent and opening by a narrow slit, becoming discoid when mature (Hooker 1855; Masee 1896). The habit of this fungus fits well with typical macromorphological features of the genus *Trochila* – a dark brown to black receptacle, which develops beneath the host tissues and eventually becomes erumpent to expose the hymenium by splitting along radial lines or by its splitting into lobes (von Höhnelt 1917; Greenhalgh and Morgan-Jones 1964; Dennis 1978; Baral and Marson 2005). Microscopically, *P. colensoi* was described with a parenchymatous excipulum (angular-globose or isodiametric cells), hyaline under the hymenium and dark brown at the cortex (Berkeley in Hooker 1855; Masee 1896), which is also in agreement with the excipular features of *Trochila* species. Finally, the hymenium of *P. colensoi* was described as composed of inamyloid, 8-spored asci with elliptical hyaline ascospores and slender paraphyses (*op. cit.*).

In 2018, P.R. Johnston collected two specimens (PDD:112240, PDD:112242, Landcare Research 2020) on leaves of *Cordyline australis* (Asparagaceae). The

morphology, ecology (host), and locality of these new collections agree with *P. colensoi*. The photographs of both specimens reveal features such as guttules in ascospores and paraphyses, protruding hyaline cells in the cortical layer of the upper flank and margin, and hyaline gelatinized hyphae covering the dark globose-angular cells of the ectal excipulum at the base and lower flanks. The latter excipular feature of the receptacle is reminiscent of Zhuang's (1990) description of *Calycellinopsis xishuangbanna*. An ITS sequence of this species was generated from the recent material (PDD:112240) and included in the Leotiomycetes-wide ITS phylogeny of Johnston et al. (2019). Their results and those in this study (Figs 1, 2) show that *P. colensoi* is placed among species of *Trochila*.

***Trochila xishuangbanna* (W.Y. Zhuang) Quijada, comb. nov.**

Mycobank No: 836592

≡ *Calycellinopsis xishuangbanna* W.Y. Zhuang, Mycotaxon 38: 121 (1990). [Basionym]

Notes. The genus *Calycellinopsis* was proposed with a single species, *C. xishuangbanna*, which is a petiole-inhabiting fungus (Zhuang 1990). The genus was placed in Dermateaceae because of its isodiametric dark brownish excipular cells (Zhuang 1990). In 2002, a second collection of the same species was sampled (HMAS:187063), which was sequenced (Zhuang et al. 2010). Additional morphological details were provided, and the genus was placed in Helotiaceae (Zhuang et al. 2010). *Trochila* was treated in Dermateaceae until recently because of its excipular features (Fuckel 1869; Karsten 1869; Saccardo 1884; Lambotte 1888; Lumbsch and Huhndorf 2010). Collections of *Calycellinopsis* have a well-developed excipulum, with an outer layer of angular to isodiametric cells with brownish walls and cortical cells at flanks and margin with protruding hyaline cells. The medullary excipulum is subhyaline and composed of *textura angularis* to *textura intricata* (Zhuang 1990; Zhuang et al. 2010).

Species in *Trochila* usually have a poorly developed excipulum. For example, *T. bostonensis* and *T. craterium* produce only a thin layer of globose to angular dark excipular cells (von Höhnelt 1917; Greenhalgh and Morgan-Jones 1964; Baral and Marson 2005). However, other species, such as *T. laurocerasi* and *T. urediniophila*, have a well-developed excipulum (*op. cit.*). The excipulum of *Calycellinopsis* is very similar to those species of *Trochila* with a well-developed excipulum, composed of an outer layer of dark *textura globulosa-angularis* and an inner layer of hyaline medulla made of *textura angularis-porrecta-intricata*. At the flanks and margin of the excipulum, *Calycellinopsis* has protruding hyaline cells similar to *Trochila* species with a well-developed excipulum (Fig. 4). Although limited details about the living features can be obtained from the original description of *Calycellinopsis*, its hymenial features are consistent with *Trochila*. The ascospores of *Calycellinopsis* are described with several guttules, a feature that is also observed in species of *Trochila*.

Discussion

Taxonomy of *Trochila*

This study represents the first attempt to investigate the systematics of *Trochila* using both morphological features and DNA sequences. We have added four species to *Trochila*, bringing the total number of species described in the genus to 37. Most *Trochila* species have been delimited based on the size of asci and ascospores, but we find that amyloidity of ascus apex, excipular features, details of the paraphyses, and presence vs. absence of guttules are also diagnostic (Table 2). For this study, we also applied a two-dataset approach for phylogenetic analyses (e.g., Aime and Phillips-Mora 2005; Haelewaters et al. 2019). Our phylogenetic reconstruction of a six-locus dataset resolved *Trochila* as polyphyletic with respect to *C. xishuangbanna* and *P. colensoi* (Fig. 1). Because morphological data of these two taxa agree with *Trochila*, we recombined them in this genus. The second, two-locus dataset was used for species delimitation, which showed *T. bostonensis* and *T. urediniophila* as distinct from the other *Trochila* species. Our molecular phylogenetic results (Figs 1, 2) and morphological comparisons of *Trochila* species (Table 2) will facilitate future taxonomic studies in the genus.

Host associations

Thus far, members of *Trochila* have been reported from 31 families of both monocots and dicots (Table 2). In this study, we add two plant family hosts, Apocynaceae (for *T. bostonensis*) and Asparagaceae (for *T. colensoi*). In addition, we reveal a new ecological niche (for *T. urediniophila*) – a species that associates with uredinia of the rust species *Cerotelium fici*. This sample was collected in 1912 as a rust specimen and deposited in the Arthur Fungarium (PUR) at Purdue University. More than a century later, the exsiccatae sample was scanned for the presence of hyperparasites of rust fungi from South America. Apothecia of *T. urediniophila* were found exclusively on uredinia without any direct contact with the host plant. Due to the age and limited available material, ultrastructural examinations of the interaction between these two fungi could not be made. However, *T. urediniophila* is the first species in the genus that fruits exclusively from another fungus, hinting at more complex associations among *Trochila* species and other fungi on which they might act as mycoparasites.

Trochila in the Neotropics

South America is known to be one of the most biodiverse continents in the world (Dourojeanni 1990; Hawksworth 2001). However, its fungal communities are thought to be severely understudied (Mueller and Schmit 2007). Members of *Trochila* are no exception to this. Six species of *Trochila* have been described from South America. These are *T. chilensis* Speg., *T. jaffuelii* Speg., and *T. perseae* Speg. from Chile; *T. leopoldina* Rehm from Brazil; and *T. tetraspora*, and *T. winteri* Speg. from Argentina (Spegazzini 1888, 1910, 1921; Rehm 1909; Gamundí et al. 1978). Their type collections need to be

re-examined to determine if these species are in fact members of *Trochila*. One of our new species, *T. urediniophila*, was collected in Port of Spain, Trinidad. Little data are available regarding the Funga (*sensu* Kuhar et al. 2018) of Trinidad and Tobago (Baker and Dale 1951; Dennis 1954a, b). The most recent work on the fungal diversity from this country was published online (Jodhan and Minter 2006) derived from reference collections and data from scientific literature. Based on the available literature, no records of *Trochila* are known in Trinidad. As a result, *T. urediniophila* represents the first published report of the genus from Trinidad, and by extension from the Caribbean (Minter et al. 2001).

Trochila species are likely more broadly distributed than generally thought, and certainly not limited to the Northern Hemisphere. This is often the case for many fungi that are based on limited regional collecting and thus may not represent the full extent of their distributional ranges due to, for example, the lack of studies in sub-tropical and tropical ecosystems (Groombridge 1992; Hawksworth and Mueller 2005; Mueller and Schmit 2007; Aime and Brearley 2012; Cheek et al. 2020).

The importance of biological collections

Our work emphasizes the importance of specimens preserved in biological collections – such as fungaria and herbaria – for studies of biodiversity and applied biological sciences, and for climate change research (Hawksworth and Lücking 2017; Andrew et al. 2019; Lang et al. 2019; Ristaino 2020; Wijayawardene et al. 2020). Because of the well-preserved specimens deposited at PUR, the genus *Trochila* is now known to be present in Trinidad and to form fungicolous associations. Another interesting example of the use of collections is *Trochila colensoi*. Known only from the type specimen for more than 100 years, additional specimens were only reported following the correction of the host substrate (as *Cordyline* rather than *Phormium*), which was based on re-examination of the type specimen preserved at K. Biological collections are not only important for morphological studies, but also as sources of genetic and genomic information (Bruns et al. 1990; Brock et al. 2009; Redchenko et al. 2012; Dentinger et al. 2016; this study). The single-oldest fungal specimen used for DNA extraction and sequencing was the type of *Hygrophorus cossus* (Sowerby) Fr. (Agaricales, Hygrophoraceae), collected in 1794 and deposited at K (Larsson and Jacobsson 2004). Our material of *T. urediniophila* gathered by Roland Thaxter in 1912 proves again that old samples can be used successfully for modern molecular phylogenetic analyses.

Acknowledgements

The National Park Service at the Boston Harbor Islands (BHI) National Recreation Area and the University of Massachusetts – Boston School for the Environment are acknowledged for facilitating the fungal ATBI. The National Park Service issued the scientific research and collecting permits (#BOHA-2012-SCI-0009, PI B.D. Farrell; #BOHA-2018-SCI-0002, PI D. Haelewaters). Thanks are due to: Marc Albert (Boston Harbor Islands Stewardship Program) for immense support with everything that is

Boston Harbor Islands-related; Russ Bowles and his staff (Division of Marine Operations, University of Massachusetts Boston) for expert navigation and transportation to Great Brewster Island; Peter R. Johnston (Manaaki Whenua Landcare Research) for providing important information about *Pseudopeziza colensoi* and for improvements to the manuscript. D. Haelewaters acknowledges support for fieldwork at the BHI and molecular work from Boston Harbor Now (2017–2018) and the New England Botanical Club (2017 Les Mehrhoff Botanical Research Award). L. Quijada thanks the support of the Farlow Fellowship, the Department of Organismic and Evolutionary Biology at Harvard University, and the Harvard University Herbaria. This work was supported in part by the U.S. National Science Foundation (DEB-2018098 to D. Haelewaters; DEB-1458290 to M.C. Aime) and the U.S. Department of Agriculture (National Institute of Food and Agriculture Hatch project 1010662 to M.C. Aime).

References

- Aime MC, Brearley FQ (2012) Tropical fungal diversity: closing the gap between species estimates and species discovery. *Biodiversity and Conservation* 21: 2177–2180. <https://doi.org/10.1007/s10531-012-0338-7>
- Aime MC, Phillips-Mora W (2005) The causal agents of witches' broom and frosty pod rot of cacao (chocolate, *Theobroma cacao*) form a new lineage of Marasmiaceae. *Mycologia* 97: 1012–1022. <https://doi.org/10.3852/mycologia.97.5.1012>
- Andrew C, Diez J, James TY, Kauserud H (2019) Fungarium specimens: a largely untapped source in global change biology and beyond. *Philosophical Transactions of the Royal Society B* 374(1763): 20170392. <https://doi.org/10.1098/rstb.2017.0392>
- Baker RED, Dale WT (1951) Fungi of Trinidad and Tobago. *Mycological Papers* 33: 1–123.
- Baral H-O (1992) Vital versus herbarium taxonomy: morphological differences between living and dead cells of ascomycetes, and their taxonomic implications. *Mycotaxon* 44: 333–390.
- Baral H-O, Marson G (2005) In vivo veritas. Over 10000 images of fungi and plants (microscopical drawings, water colour plates, photo macro- & microphotographs), with materials on vital taxonomy and xerotolerance. Ed. 3. Privately distributed DVD-ROM.
- Brock PM, Döring H, Bidartondo MI (2009) How to know unknown fungi: The role of a herbarium. *New Phytologist* 181: 719–724. <https://doi.org/10.1111/j.1469-8137.2008.02703.x>
- Bruns TD, Fogel R, Taylor JW (1990) Amplification and sequencing of DNA from fungal herbarium specimens. *Mycologia* 82: 175–184. <https://doi.org/10.2307/3759846>
- Cheek M, Lughadha EN, Kirk P, Lindon H, Carretero J, Looney B, Douglas B, Haelewaters D, Gaya E, Llewellyn T, Ainsworth M, Gafforov Y, Hyde K, Crous P, Hughes M, Walker BE, Forzza RC, Meng WK, Niskanen T (2020) New scientific discoveries: Plants and fungi. *Plants, People, Planet* 2(5): 371–388. <https://doi.org/10.1002/ppp3.10148>
- Chernomor O, Von Haeseler A, Minh BQ (2016) Terrace aware data structure for phylogenomic inference from supermatrices. *Systematic Biology* 65: 997–1008. <https://doi.org/10.1093/sysbio/syw037>
- Clements FE, Shear CL (1931) *The genera of fungi*. H.W. Wilson Company, Bronx, New York, 496 pp.

- Crouan PL, Crouan MH (1867) Florule du Finistère. Contenant les descriptions de 360 espèces nouvelles de sporogames, de nombreuses observations et une synonymie des plantes cellulaires et vasculaires qui croissant spontanément dans ce département. Friedrich Klincksieck and J.B. et A. Lefournier, Paris & Brest, 262 pp. <https://doi.org/10.5962/bhl.title.11601>
- Crous PW, Quaedvlieg W, Hansen K, Hawksworth DL, Groenewald JZ (2014) *Phacidium* and *Ceuthospora* (Phaciaceae) are congeneric: taxonomic and nomenclatural implications. *IMA Fungus* 5: 173–193. <https://doi.org/10.5598/imafungus.2014.05.02.02>
- Crous PW, Schumacher RK, Wingfield MJ, Akulov A, Denman S, Roux J, Braun U, Burgess TI, Carnegie AJ, Váczy KZ, Guatimosim E, Schwartsburd PB, Barreto RW, Hernández-Restrepo M, Lombard L, Groenewald JZ (2018) New and Interesting Fungi. 1. *Fungal Systematics and Evolution* 1: 169–215. <https://doi.org/10.3114/fuse.2018.01.08>
- De Notaris G (1863) [printed 1864] Proposte di alcune rettificazioni al profilo dei Discomiceti. *Commentario della Società Crittogamologica Italiana* 1(5): 357–388.
- Dennis RWG (1954a) Operculate Discomycetes from Trinidad and Jamaica. *Kew Bulletin* 9(3): 417–421. <https://doi.org/10.2307/4108810>
- Dennis RWG (1954b) Some Inoperculate Discomycetes of Tropical America. *Kew Bulletin* 9(2): 289–348. <https://doi.org/10.2307/4114399>
- Dennis RWG (1961) Some Inoperculate Discomycetes from New Zealand. *Kew Bulletin* 15(2): 293–320. <https://doi.org/10.2307/4109373>
- Dennis RWG (1978) *British Ascomycetes*. J. Cramer, Vaduz, Liechtenstein, 585 pp.
- Dentinger BTM, Gaya E, O'Brien H, Suz LM, Lachlan R, Diaz-Valderrama JR, Koch RA, Aime MC (2016) Tales from the crypt: genome mining from fungarium specimens improves resolution of the mushroom tree of life. *Biological Journal of the Linnean Society* 117: 11–32. <https://doi.org/10.1111/bij.12553>
- Don RH, Cox PT, Wainwright BJ, Baker K, Mattick JS (1991) 'Touchdown' PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Research* 19(14): 4008–4008. <https://doi.org/10.1093/nar/19.14.4008>
- Dourojeanni MJ (1990) Entomology and biodiversity conservation in Latin America. *American Entomologist* 36: 88–93. <https://doi.org/10.1093/ae/36.2.88>
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Egger KN (1995) Molecular analysis of ectomycorrhizal fungal communities. *Canadian Journal of Botany* 73: S1415–S1422. <https://doi.org/10.1139/b95-405>
- Etayo J, Flakus A, Suija A, Kukwa M (2015) *Macroskyttea parmotrematis* gen. et sp. nov. (Helotiales, Leotiomycetes, Ascomycota), a new lichenicolous fungus from Bolivia. *Phytotaxa* 224: 247–257. <https://doi.org/10.11646/phytotaxa.224.3.3>
- Feltgen J (1903) Vorstudien zu einer Pilz-flora des Grossherzogthums Luxemburg. I. Theil – Ascomycetes. *Nachträge III. Recueil des mémoires et des travaux publiés par la Société de botanique du grand-duché de Luxembourg* 16: 3–328.
- Fries E (1849) *Summa Vegetabilium Scandinaviae. Sectio Posterior. Cl. XX. Fungi*. A. Bonnier, Stockholm & Leipzig, 261–572.
- Fryar SC, Haelewaters D, Catcheside DE (2019) *Annabella australiensis* gen. & sp. nov. (Helotiales, Cordieritidaceae) from South Australian mangroves. *Mycological Progress* 18: 973–981. <https://doi.org/10.1007/s11557-019-01499-x>

- Fuckel L (1869) *Symbolae mycologicae. Beiträge zur Kenntniss der rheinischen Pilze. Mit VI lithographirten und colorirten Tafeln. Jahrbücher des Nassauischen Vereins für Naturkunde* 23–24: 1–459. <https://doi.org/10.5962/bhl.title.47117>
- Gamundí, IJ, Arambarri, AM, Gaiotti A (1978) Micoflora de la hojarasca de *Nothofagus dombeyi*. *Darwiniana* 21: 81–114.
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2: 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Global Biodiversity Information Facility (2020) *Trochila* Fr. GBIF Backbone Taxonomy. <https://www.gbif.org/species/2575690> [accessed 17 August 2020]
- Greenhalgh G, Morgan-Jones G (1964). Some species of *Trochila* and an undescribed discomycete on leaves of *Prunus laurocerasus*. *Transactions of the British Mycological Society* 47: 311–320. [https://doi.org/10.1016/S0007-1536\(64\)80002-4](https://doi.org/10.1016/S0007-1536(64)80002-4)
- Gregor MJF (1936) A disease of cherry laurel caused by *Trochila laurocerasi* (Desm.) Fr. *Annals of Applied Biology* 23: 700–704. <https://doi.org/10.1111/j.1744-7348.1936.tb06121.x>
- Grelet L-J, de Crozals A (1928) Discomycètes nouveaux (3ième série). *Bulletin trimestriel de la Société mycologique de France* 44: 336–340.
- Haelewaters D, Dirks AC, Kappler LA, Mitchell JK, Quijada L, Vandegrift R, Buyck B, Pfister DH (2018a) A preliminary checklist of fungi at the Boston Harbor islands. *Northeastern Naturalist* 25(sp9): 45–77. <https://doi.org/10.1656/045.025.s904>
- Haelewaters D, De Kesel A, Pfister DH (2018b) Integrative taxonomy reveals hidden species within a common fungal parasite of ladybirds. *Scientific Reports* 8: e15966. <https://doi.org/10.1038/s41598-018-34319-5>
- Haelewaters D, Pfliegler WP, Gorczak M, Pfister DH (2019) Birth of an order: comprehensive molecular phylogenetic study reveals that *Herpomyces* (Fungi, Laboulbeniomycetes) is not part of Laboulbeniales. *Molecular Phylogenetics and Evolution* 133: 286–301. <https://doi.org/10.1016/j.ympev.2019.01.007>
- Hammond PM (1992) Species inventory. In: Groombridge B (Ed.) *Global biodiversity, status of the earth's living resources*. Chapman & Hall, London, 17–39. https://doi.org/10.1007/978-94-011-2282-5_4
- Hawksworth DL (2001) The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycological Research* 105: 1422–1432. <https://doi.org/10.1017/S0953756201004725>
- Hawksworth DL, Lücking R (2017) Fungal diversity revisited: 2.2 to 3.8 million species. In: Heitman J, Howlett B, Crous P, Stukenbrock E, James T, Gow N (Eds) *The Fungal Kingdom*. ASM Press, Washington, 79–95. <https://doi.org/10.1128/microbiolspec.FUNK-0052-2016>
- Hawksworth DL, Mueller GM (2005) Fungal communities: their diversity and distribution. In: Digthton J, White JF, Oudemans P (Eds) *The fungal community: its organisation and role in the ecosystem*. CRC Press, Boca Raton, 27–37. <https://doi.org/10.1201/9781420027891.ch2>
- Hennings P (1900) *Fungi japonici*. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 28: 273–280.
- Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS (2017) UFBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35: 518–522. <https://doi.org/10.1093/molbev/msx281>
- Höhnelt F von (1917) Über die Gattung *Trochila* Fries. *Annales Mycologici* 15: 330–334.

- Hooker JD (1855) The botany of the Antarctic Voyage of H.M. discovery ships Erebus and Terror, in the years 1839–1843. II. Flora Novae-Zelandiae. Part II. Flowerless plants. Lovell Reeve, London, 378 pp.
- Hopple JS (1994) Phylogenetic investigations in the genus *Coprinus* based on morphological and molecular characters. PhD Dissertation, Duke University, Durham.
- Hyde KD, McKenzie EHC, KoKo TW (2011) Towards incorporating anamorphic fungi in a natural classification – checklist and notes for 2010. *Mycosphere* 2: 1–88.
- Index Fungorum (2021) Index Fungorum. <http://www.indexfungorum.org/names/Names.asp> [accessed 22 June 2020]
- Jaklitsch W, Baral H-O, Lücking R, Lumbsch HT (2016) Syllabus of plant families. In: Frey W (Ed.) *Adolf Engler's Syllabus der Pflanzenfamilien*. Part 1/2 Ascomycota (13th edn). Borntraeger Science Publishers, Stuttgart, 322 pp.
- Jodhan D, Minter DW (2006) Fungi of Trinidad & Tobago. <http://www.cybertruffle.org.uk/trinfung> [accessed 14 November 2019]
- Johnston PR, Quijada L, Smith CA, Baral H-O, Hosoya T, Baschien C, Pärtel K, Zhuang K-Y, Haelewaters D, Park D, Carl S, López-Giráldez F, Wang Z, Townsend JP (2019) A multigene phylogeny toward a new phylogenetic classification of Leotiomycetes. *IMA Fungus* 10: 1–22. <https://doi.org/10.1186/s43008-019-0002-x>
- Kalyanamoorthy K, Minh BQ, Wong TKE, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14: 587–589. <https://doi.org/10.1038/nmeth.4285>
- Karsten PA (1869) *Monographia Pezizarum fennicarum*. Notiser ur Sällskapet pro Fauna et Flora fennica förhandlingar X: 99–206.
- Karsten PA (1871) *Mycologia Fennica I. Discomycetes*. Bidrag Kännedom Finland Natur Folk 19: 1–263.
- Kelly KL (1965) ISCC-NBS Colour-name charts illustrated with centroid colors. Inter-Society Colors Council. National Bureau of Standards, Circular 553 (Supplement). US Government Printing Office, Washington, 44 pp.
- Kiffer E, Morelet M (2000) *The Deuteromycetes, mitosporic fungi: classification and generic key*. Science Publishers, Enfield, 273 pp.
- Kirschstein W (1941) De plerisque novis ascomycetibus et paucis novis fungis imperfectis. *Hedwigia* 80: 119–137.
- Kirschstein W (1944) Über neue, seltene und kritische Kleinpilze. *Hedwigia* 81: 193–224.
- Korf RP (1973) Discomycetes and Tuberales. In: Ainsworth GC, Sparrow FK, Sussman AS (Eds) *The Fungi: An Advanced Treatise*. Vol. 4a. Academic Press, London, 249–319.
- Kuhar F, Furci G, Drechsler-Santos ER, Pfister DH (2018) Delimitation of Funga as a valid term for the diversity of fungal communities: the Fauna, Flora & Funga proposal (FF&F). *IMA Fungus* 9: 71–74. <https://doi.org/10.1007/BF03449441>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33: 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Lamarck JBPA, de Candolle AP (1805) *Flore française*, ii. Desray, Paris, 600 pp.
- Lambotte E (1888) *La flore mycologique de la Belgique*. Mémoires de la Société royale des sciences de Liège, sér. 2, 14: 1–350.

- Landcare Research (2020) Collection details. *Pseudopeziza colensoi* (Berk.) Massee (1896) [1895–97]. <https://nzfungi2.landcareresearch.co.nz/default.aspx?selected=NameDetails&TabNum=0&NameId=1CB1B798-36B9-11D5-9548-00D0592D548C> [accessed 16 August 2020]
- Lang PL, Willems FM, Scheepens JF, Burbano HA, Bossdorf O (2019) Using herbaria to study global environmental change. *New Phytologist* 221: 110–122. <https://doi.org/10.1111/nph.15401>
- Larsson E, Jacobsson S (2004) Controversy over *Hygrophorus cossus* settled using ITS sequence data from 200 year-old type material. *Mycological Research* 108: 781–786. <https://doi.org/10.1017/S0953756204000310>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16: 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Lumbsch HT, Huhndorf SM (2010) Myconet Volume 14. Part One. Outline of Ascomycota – 2009. Part Two. Notes on Ascomycete Systematics. Nos. 4751–5113. *Fieldiana Life and Earth Sciences* 2010: 1–64. <https://doi.org/10.3158/1557.1>
- Massee GE (1896) Redescriptions of Berkeley's types of fungi. *Journal of the Linnean Society* 31: 463–524. <https://doi.org/10.1111/j.1095-8339.1896.tb00812.x>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop*, 14 Nov. 2010, New Orleans, LA, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Minter DW, Rodríguez-Hernández M, Mena-Portales J (2001) *Fungi of the Caribbean. An annotated checklist*. PDMS Publishing, Isleworth, 950 pp.
- Morgan-Jones G (1973) *Genera coelomycetarum*. VII. *Cryptocline* Petrak. *Canadian Journal of Botany* 51: 309–325. <https://doi.org/10.1139/b73-039>
- Mouton V (1900) Quatrième notice sur des ascomycètes nouveaux ou peu connus. *Bulletin de la Société Royale de Botanique de Belgique* 39: 37–53.
- Mueller GM, Schmit JP (2007) Fungal biodiversity: what do we know? What can we predict? *Biodiversity and Conservation* 16: 1–5. <https://doi.org/10.1007/s10531-006-9117-7>
- MyCoPortal (2020) Mycology Collections data Portal. <http://mycoportal.org/portal/index.php> [accessed 7 December 2020]
- Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. *Molecular Biology and Evolution* 32: 268–274. <https://doi.org/10.1093/molbev/msu300>
- Pärtel K, Baral H-O, Tamm H, Pöldmaa K (2017) Evidence for the polyphyly of *Encoelia* and *Encoelioideae* with reconsideration of respective families in *Leotiomycetes*. *Fungal Diversity* 82: 183–219. <https://doi.org/10.1007/s13225-016-0370-0>
- Patouillard MN (1886) Quelques champignons de la Chine, récoltés par M. l'abbé Delavay dans la province du Yunnan. *Revue Mycologique* 8: 179–194.
- Redchenko O, Vondrák J, Košnar J (2012) The oldest sequenced fungal herbarium sample. *The Lichenologist* 44: 715–718. <https://doi.org/10.1017/S002428291200031X>
- Rehm H (1896) Abt. 3. Ascomyceten: hysteriaceen und discomyceten. In: Rabenhorst L (Ed.) *Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz*. Verlag von Eduard Kummer, Leipzig.
- Rehm H-J (1909) *Ascomycetes exs. Fasc. 45. Annales Mycologici* 7: 524–530.

- Rehner SA, Buckley E (2005) A *Beauveria* phylogeny inferred from nuclear ITS and EF1- α sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* 97: 84–98. <https://doi.org/10.3852/mycologia.97.1.84>
- Ristaino JB (2020) The importance of mycological and plant herbaria in tracking plant killers. *Frontiers in Ecology and Evolution* 7: e521. <https://doi.org/10.3389/fevo.2019.00521>
- Rostrup E (1885) Islands svampe. *Botanisk Tidsskrifte* 14: 218–229.
- Rostrup E (1886) Svampen fra Finmarken. *Botanisk Tidsskrifte* 15: 229–236.
- Rostrup E (1888) Oversigt over Grönlands svampe. *Meddelelser om Grønland* 3: 516–590.
- Rostrup E (1891) Tillaeg til “Grönlands svampe (1888)”. *Meddelelser om Grønland* 3: 591–643.
- Saccardo PA (1878) *Fungi Veneti novi vel critici*. Series IX. *Michelia* 4: 361–434.
- Saccardo PA (1884) *Conspectus generum discomycetum hucusque cognitorum*. *Botanisches Centralblatt* 18: 247–256.
- Spatafora JW, Sung GH, Johnson D, Hesse C, O'Rourke B, Serdani M, Spotts R, Lutzoni F, Hofstetter V, Miadlikowska J, Reeb V, Gueidan C, Fraker E, Lumbsch T, Lücking R, Schmitt I, Hosaka K, Aptroot A, Roux C, Miller AN, Geiser DM, Hafellner J, Hestmark G, Arnold AE, Büdel B, Rauhut A, Hewitt D, Untereiner WA, Cole MS, Scheidegger C, Schultz M, Sipman H, Schoch CL (2006) A five-gene phylogeny of Pezizomycotina. *Mycologia* 98: 1018–1028. <https://doi.org/10.1080/15572536.2006.11832630>
- Spegazzini C (1888) *Fungi Fuegiani*. *Boletín de la Academia Nacional de Ciencias en Córdoba* 11: 135–311. <https://doi.org/10.5962/bhl.title.4055>
- Spegazzini C (1910) Hongos chilenos. *Revista de la Facultad de Agronomía y Veterinaria*. 6: 130–132.
- Spegazzini C (1921) *Mycetes chilenses*. *Boletín de la Academia Nacional de Ciencias* 25: 1–124.
- Stevenson JA (1926) Foreign plant diseases. A manual of economic plant diseases which are new or not widely distributed in the United States. United States Department of Agriculture Federal Horticulture Board, Washington DC. <https://doi.org/10.5962/bhl.title.114939>
- Stoykov DY, Assyov B (2009) The genus *Trochila* in Bulgaria. *Mycotaxon* 109: 351–359. <https://doi.org/10.5248/109.351>
- Stoykov DY (2019) New records of *Trochila* (Cenangiaceae, Helotiales) from the Balkans. *Phytologia Balcanica* 25: 245–248.
- Suija A, Ertz D, Lawrey JD, Diederich P (2015) Multiple origin of the lichenicolous life habit in Helotiales, based on nuclear ribosomal sequences. *Fungal Diversity* 70: 55–72. <https://doi.org/10.1007/s13225-014-0287-4>
- Sung GH, Sung JM, Hywel-Jones NL, Spatafora JW (2007) A multi-gene phylogeny of Clavicipitaceae (Ascomycota, Fungi): Identification of localized incongruence using a combinational bootstrap approach. *Molecular Phylogenetics and Evolution* 44: 1204–1223. <https://doi.org/10.1016/j.ympev.2007.03.011>
- Thiers B (2020) [continuously updated] Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanic Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [accessed 14 July 2020]
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246. <https://doi.org/10.1128/JB.172.8.4238-4246.1990>

- von Arx JA (1957) Revision der zu *Gloeosporium* gestellten Pilze. Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen, Afd. Natuurkunde, Tweede Reeks 51: 1–153.
- Vu D, Groenewald M, de Vries M, Gehrman T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* 92: 135–154. <https://doi.org/10.1016/j.simyco.2018.05.001>
- Wang Z, Binder M, Hibbett DS (2005) Life history and systematics of the aquatic discomycete *Mitrula* (Helotiales, Ascomycota) based on cultural, morphological, and molecular studies. *American Journal of Botany* 92: 1565–1574. <https://doi.org/10.3732/ajb.92.9.1565>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand H, Sninsky JS, White TJ (Eds) *PCR protocols: A guide to methods and applications*. Academic Press, San Diego, 315–322. <https://doi.org/10.3181/00379727-150-39089>
- Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, Tokarev YS, Dai DQ, Letcher PM, Stephenson SL, Ertz D, Lumbsch HT, Kukwa M, Issi IV, Madrid H, Phillips AJL, Selbmann L, Pfliegler WP, Horváth E, Bensch K, Kirk PM, Kolaříková K, Raja HA, Radek R, Papp V, Dima B, Ma J, Malosso E, Takamatsu S, Rambold G, Gannibal PB, Triebel D, Gautam AK, Avasthi S, Suetrong S, Timdal E, Fryar SC, Delgado G, Réblová M, Doilom M, Dolatabadi S, Pawłowska J, Humber RA, Kodsueb R, Sánchez-Castro I, Goto BT, Silva DKA, de Souza FA, Oehl F, da Silva GA, Silva IR, Błaszczowski J, Jobim K, Maia LC, Barbosa FR, Fiuza PO, Divakar PK, Shenoy BD, Castañeda-Ruiz RF, Somrithipol S, Lateef AA, Karunarathna SC, Tibpromma S, Mortimer PE, Wanasinghe DN, Phookamsak R, Xu J, Wang Y, Tian F, Alvarado P, Li DW, Kušan I, Matočec N, Maharachchikumbura SSN, Papizadeh M, Heredia G, Wartchow F, Bakhshi M, Boehm E, Youssef N, Hustad VP, Lawrey JD, Santiago ALCMA, Bezerra JDP, Souza-Motta CM, Firmino AL, Tian Q, Houbraken J, Hongsanan S, Tanaka K, Dissanayake AJ, Monteiro JS, Grossart HP, Suija A, Weerakoon G, Etayo J, Tsurykau A, Vázquez V, Mungai P, Damm U, Li QR, Zhang H, Boonmee S, Lu YZ, Becerra AG, Kendrick B, Brearley FQ, Motiejūnaitė J, Sharma B, Khare R, Gaikwad S, Wijesundara DSA, Tang LZ, He MQ, Flakus A, Rodriguez-Flakus P, Zhurbenko MP, McKenzie EHC, Stadler M, Bhat DJ, Liu JK, Raza M, Jeewon R, Nassonova ES, Prieto M, Jayalal RGU, Erdoğdu M, Yurkov A, Schnittler M, Shchepin ON, Novozhilov YK, Silva-Filho AGS, Liu P, Cavender JC, Kang Y, Mohammad S, Zhang LF, Xu RF, Li YM, Dayarathne MC, Ekanayaka AH, Wen TC, Deng CY, Pereira OL, Navathe S, Hawksworth DL, Fan XL, Dissanayake LS, Kuhnert E, Grossart HP, Thines M (2020) Outline of Fungi and fungus-like taxa. *Mycosphere* 11: 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>
- Zhao YJ, Hosaka K, Hosoya T (2016) Taxonomic re-evaluation of the genus *Lambertella* (Rutstroemiaceae, Helotiales) and allied stroma-forming fungi. *Mycological Progress* 15: 1215–1228. <https://doi.org/10.1007/s11557-016-1225-5>
- Zhuang WY (1990) *Calycellinopsis Xishuangbanna* gen. et sp. nov. (Dermateaceae), a petiole-inhabiting fungus from China. *Mycotaxon* 38: 121–124.

- Zhuang WY, Luo J, Zhao P (2010) The fungal genus *Calycellinopsis* belongs in Helotiaceae not Dermateaceae. Phytotaxa 3: 54–58. <https://doi.org/10.11646/phytotaxa.3.1.8>
- Ziolo E, Madej T, Blaszkowski J (2005) *Trochila ilicina* (Helotiales, Ascomycota), a fungus newly found in Poland. Acta Mycologica 40: 181–184. <https://doi.org/10.5586/am.2005.016>