

C. elegans MAGU-2/Mpp5 homolog regulates epidermal phagocytosis and synapse density

Salvatore J. Cherra III, Alexandre Goncharov, Daniela Boassa, Mark Ellisman & Yishi Jin

To cite this article: Salvatore J. Cherra III, Alexandre Goncharov, Daniela Boassa, Mark Ellisman & Yishi Jin (2020) *C. elegans* MAGU-2/Mpp5 homolog regulates epidermal phagocytosis and synapse density, *Journal of Neurogenetics*, 34:3-4, 298-306, DOI: [10.1080/01677063.2020.1726915](https://doi.org/10.1080/01677063.2020.1726915)

To link to this article: <https://doi.org/10.1080/01677063.2020.1726915>



[View supplementary material](#)



Published online: 04 May 2020.



[Submit your article to this journal](#)



Article views: 382



[View related articles](#)



[View Crossmark data](#)



Citing articles: 2 [View citing articles](#)

C. elegans MAGU-2/Mpp5 homolog regulates epidermal phagocytosis and synapse density

Salvatore J. Cherra III^{a,b} , Alexandr Goncharov^a, Daniela Boassa^c, Mark Ellisman^{c,d} and Yishi Jin^{a,d,e} 

^aSection of Neurobiology, Division of Biological Sciences, University of California San Diego, La Jolla, CA, USA; ^bDepartment of Neuroscience, University of Kentucky College of Medicine, Lexington, KY, USA; ^cNational Center for Microscopy and Imaging Research, University of California San Diego, La Jolla, CA, USA; ^dDepartment of Neurosciences, School of Medicine, University of California San Diego, La Jolla, CA, USA; ^eDepartment of Cellular and Molecular Medicine, School of Medicine, University of California San Diego, La Jolla, CA, USA

ABSTRACT

Synapses are dynamic connections that underlie essential functions of the nervous system. The addition, removal, and maintenance of synapses govern the flow of information in neural circuits throughout the lifetime of an animal. While extensive studies have elucidated many intrinsic mechanisms that neurons employ to modulate their connections, increasing evidence supports the roles of non-neuronal cells, such as glia, in synapse maintenance and circuit function. We previously showed that *C. elegans* epidermis regulates synapses through ZIG-10, a cell-adhesion protein of the immunoglobulin domain superfamily. Here we identified a member of the Pals1/MPP5 family, MAGU-2, that functions in the epidermis to modulate phagocytosis and the number of synapses by regulating ZIG-10 localization. Furthermore, we used light and electron microscopy to show that this epidermal mechanism removes neuronal membranes from the neuromuscular junction, dependent on the conserved phagocytic receptor CED-1. Together, our study shows that *C. elegans* epidermis constrains synaptic connectivity, in a manner similar to astrocytes and microglia in mammals, allowing optimized output of neural circuits.

ARTICLE HISTORY

Received 5 November 2019
Accepted 4 February 2020

KEYWORDS

MAGUK; synapse elimination; glia; miniSOG; neuromuscular junction; ACR-2

Introduction

Synapses enable the transmission and integration of information within the nervous system. Proper synaptic connectivity is essential to govern nervous system functions such as sensory perception, learning, and coordinated movement. Aberrant synaptic connections have been associated with a variety of neurological disorders. Synapse formation, elimination, and maintenance work together to ensure precision and plasticity of neuronal circuits throughout the lifetime of an animal. Many neuronal-intrinsic mechanisms involve various classes of cell surface proteins and intracellular signaling pathways that establish synaptic connections (Cherra & Jin, 2015; de Wit & Ghosh, 2016; Sudhof, 2018). Additional work has shown how extrinsic mechanisms involving non-neuronal cells, such as astrocytes and microglia, cooperate with neurons to modulate neuronal circuits (Allen & Eroglu, 2017; Chung, Allen, & Eroglu, 2015). It is now well established that astrocytes and microglia play an active role in the pruning of synaptic connections during development of the visual system in mice (Chung *et al.*, 2013; Stevens *et al.*, 2007). In *Drosophila*, glial cells also remove synapses and axonal components as a means to eliminate neuronal connections (Awasaki *et al.*, 2006; Fuentes-Medel *et al.*, 2009).

One common mechanism for non-neuronal cells to assist in the wiring and rewiring of circuits is through the phagocytosis pathway. The phagocytosis pathway is highly conserved throughout evolution and plays essential roles in the removal of cell corpses and cellular debris (Bangs, Franc, & White, 2000; Mangahas & Zhou, 2005). While the core machinery that orchestrates engulfment has been widely studied, additional mechanisms that modulate the initiation or target specificity of this pathway still remain largely unknown. The *C. elegans* motor circuit provides a simple model to understand how non-neuronal cells modulate synapse elimination via the phagocytosis pathway. Our previous work uncovered a cell surface protein, ZIG-10, which mediates an interaction between the epidermis and neurons. The cell-cell interaction mediated by ZIG-10 enables the elimination of synapses through the activation of the phagocytosis pathway (Cherra & Jin, 2016).

In this study we have further investigated how the phagocytosis pathway maintains optimal synaptic connectivity in the motor circuit. We have discovered that a member of the MPP5/Pals1 membrane-associated guanylate kinase (MAGUK) family, MAGU-2, regulates synapse density. MAGU-2 functions in the epidermis and regulates ZIG-10 localization. Furthermore, we show that motor neuron membrane transfer to the epidermis is dependent on the phagocytosis receptor,

CONTACT Salvatore J. Cherra III  scherra@uky.edu  Section of Neurobiology, Division of Biological Sciences, University of Kentucky College of Medicine, 800 Rose Street, Medical Science Building, Room MN208, Lexington, KY 40536, USA; Yishi Jin  yijin@ucsd.edu  Department of Neurosciences, School of Medicine, University of California San Diego, 9500 Gilman Drive, Bonner Hall, Room 2418, La Jolla, CA 92093, USA.

 Supplemental data for this article can be accessed [here](#).

© 2020 Informa UK Limited, trading as Taylor & Francis Group

CED-1. Our study highlights a new role for MAGUKs in regulating synaptic connectivity.

Results

Identification of MAGU-2 in ZIG-10 pathway

We have previously reported that maintenance of synaptic connectivity in the *C. elegans* motor circuit requires components of the phagocytosis pathway acting in the epidermis (Cherra & Jin, 2016). To further understand the regulation of phagocytosis-mediated synapse elimination, we screened for additional genes using a function-based assay. Ectopic expression of ZIG-10 in the ventral cord GABAergic motor neurons causes a reduction in GABAergic synapse number (Cherra & Jin, 2016) and enhances locomotor deficit caused by an acetylcholine receptor mutation, *acr-2(n2420)* (Jospin *et al.*, 2009) (Supplemental Figure 1(a–c)). We employed this ectopic ZIG-10 expression induced phenotype to identify intracellular signaling molecules using RNAi against genes that contained SH3 domains (Cherra & Jin, 2016). One candidate corresponded to the *magu-2* gene, knockdown of which decreased the locomotor defects in *zig-10(tm6127)*; *acr-2(n2420)* animals that also ectopically expressed ZIG-10 in GABAergic neurons. The observed *magu-2(RNAi)* effect on locomotor defects was further verified using genetic null mutations in *magu-2* (Supplementary Figure 1(a)). Additionally, the *magu-2(gk218)* null mutation restored the number of GABAergic synapses to wild type levels in animals ectopically expressing ZIG-10 (Supplementary Figure 1(b–c)). Since loss of function in *magu-2* alone does not cause any defects in GABA synapse morphology (Supplementary Figure 1(b–c)), this suggests that MAGU-2 acts in a cellular context dependent on the ZIG-10 pathway.

MAGU-2 belongs to the membrane-associated guanylate kinase (MAGUK) family that contains a PDZ domain, an SH3 domain, and a guanylate kinase domain in its carboxy-terminus (Supplementary Figure 2(a)) (Hobert, 2013; Zhu, Shang, & Zhang, 2016). The family of MAGUKs modulate various forms of intercellular junctions by regulating protein localization (Zhu *et al.*, 2016). In mammals, CASK and PSD95 modulate intercellular junctions in the nervous system through their regulation of synaptic transmission and neurotransmitter receptor clustering. Based on homology, MAGU-2 belongs to the MAGUK subfamily of membrane palmitoylated proteins (MPP) and displays highest percent of total identity (~32%) with Mpp5/Pals1 (Supplementary Figure 2(b)). In the developing mouse nervous system, Mpp5/Pals1 is expressed in Schwann cells and Muller glia and is essential for establishing cell polarity (Ozcelik *et al.*, 2010; van Rossum *et al.*, 2006). Additionally, Mpp5/Pals1 regulates the maintenance of cerebellar progenitors (Park *et al.*, 2016) and axon sorting in the peripheral nervous system (Zollinger, Chang, Baalman, Kim, & Rasband, 2015). Retinal ganglion cell-specific Mpp5/Pals1 knockout mice display neuronal degeneration early during development, mimicking the clinical degeneration observed in Leber congenital amaurosis patients (Cho *et al.*, 2012; Park *et al.*, 2011). The

Drosophila homolog, Stardust, acts as a scaffolding protein at epithelial intercellular junctions to establish cell polarity (Tepass, 2012). However, it is unclear whether the MPP subfamily also regulates synaptic connectivity.

MAGU-2 acts in the epidermis to regulate the synapse number of cholinergic motor neurons

To determine where MAGU-2 functions to modulate synaptic connectivity, we first investigated the *magu-2* gene structure to identify its promoter based on the information from AceView and Wormbase (Lee *et al.*, 2018; Thierry-Mieg & Thierry-Mieg, 2006). Both sources predicted *magu-2* to span more than 10 kb, producing two potential isoforms (Figure 1(a–b)), with the long isoform a being supported by a recent dataset based on single-cell RNA-seq (Cao *et al.*, 2017). We isolated RNA from a mixed population of wild type N2 animals. We then used isoform-specific forward primers and a common reverse primer to amplify each putative isoform. Using genomic DNA as a control, we found that only the shorter isoform b was detected in our assay as a 270 bp-band whereas 748 bp-band indicative of the longer isoform was not detected (Figure 1(d)). This analysis does not exclude the possibility that isoform a may be expressed at lower levels or in a very limited subset of tissues.

Based on our transcript analysis, we created a MAGU-2::GFP fusion protein driven by the 4 kb DNA sequence upstream of *magu-2* isoform b to determine where MAGU-2 was expressed (Figure 1(c)). We detected MAGU-2::GFP expression from embryo to adulthood (Figure 1(e–f)). MAGU-2::GFP was visible as a diffuse signal throughout the epidermis and in various cells in the head, pharyngeal muscle, and posterior intestine, but was not observed in neurons or muscle (Figure 1(g–h)). Since ZIG-10 acts in both epidermis and cholinergic neurons to regulate cholinergic synapse density (Cherra & Jin, 2016), we investigated whether *magu-2* affects cholinergic synapses. We analyzed two *magu-2* deletion mutants, *ok1059* and *gk218*, which both remove most of the gene, and therefore likely are null alleles (Figure 2(a–d); Supplementary Figure 2(a)). Both alleles displayed an increase in cholinergic synapses, similar to *zig-10(tm6127)* animals (Figure 2(d)). We observed that *zig-10(tm6127)*; *magu-2(gk218)* double mutants showed no further increase in cholinergic synapses (Figure 2(d)). To assess whether the additional cholinergic synapses in *magu-2* mutants formed functional postsynaptic compartments, we assayed the animals' sensitivity to the cholinergic agonist, levamisole. We have previously shown that the excessive synapses observed in the *zig-10(tm6127)* animals caused hypersensitivity to levamisole, leading to more rapid paralysis than wild type animals (Cherra & Jin, 2016). The *magu-2(gk218)* mutants showed an increased sensitivity to levamisole similar to *zig-10(tm6127)* mutants (Figure 2(e)). Overall, these observations are consistent with MAGU-2 and ZIG-10 functioning in the same pathway.

Expression of MAGU-2::GFP in *magu-2(gk218)* animals was sufficient to restore cholinergic synapse density to wild type levels (Figure 2(h,m)), demonstrating that isoform b




Figure 1. MAGU-2 is expressed in the epidermis. (a) Diagram of the *magu-2* genetic locus on chromosome V. The *magu-2* gene is flanked by the neighboring genes, *sup-37* and *rpm-1*. (b) Diagram of *magu-2* gene structure representing the predicted a and b isoforms. Gray arrowheads indicate primers used for isoform analysis. (c) Diagram of *MAGU-2::GFP* transgene. (d) Images of PCR products using *magu-2a* and *magu-2b* specific primers after gel electrophoresis. Genomic DNA (gDNA) from N2 was used as a PCR reaction control. Messenger RNAs from N2 were reverse transcribed into cDNA. PCR of *magu-2a* is expected to produce a 5.7 kb band from gDNA and 748 bp band from cDNA; *magu-2b* is expected to produce a 323 bp band from gDNA and a 270 bp band from cDNA. (e) MAGU-2::GFP expression in an embryo, between 150 and 300 min after fertilization. Arrows indicate cluster of cells expressing MAGU-2::GFP. (f) MAGU-2::GFP expression in a z-plane through the lateral epidermis of an adult animal. (g–h) Representative fluorescence images of L1 larva expressing MAGU-2::GFP. Scale bars are 20 μ m. (g) Maximum projection through superficial layers of lateral aspect of animal showing MAGU-2 expression in epidermis, but not in neuronal commissures. (h) Maximum projection through medial layers around nerve cord showing MAGU-2 expression in epidermal ridge. Arrows indicate epidermal ridge.

was sufficient to replace MAGU-2 function in the *gk218* mutant. We further addressed in which cells MAGU-2 was functioning to regulate synapse density using tissue-specific promoters to express MAGU-2 isoform b cDNA. Expression of MAGU-2 cDNA in the epidermis of *magu-2(gk218)* animals restored the density of cholinergic synapses to wild type levels (Figure 2(j,m)). However, expression of MAGU-2 in the nervous system or muscles was not sufficient to restore cholinergic synapse density (Figure 2(i,k,m)). Moreover, expressing mouse Mpp5 cDNA in the epidermis sufficiently restored synapse number to wild type levels

(Figure 2(l-m)), indicating that mpp5 can function similarly to MAGU-2. Together, these data indicate that MAGU-2 is expressed and functions solely in the epidermis to modulate synaptic connectivity.

MAGU-2 affects phagocytosis in epidermis

To determine if MAGU-2 regulates epidermal phagocytosis, we investigated whether *magu-2(gk218)* animals displayed changes in the epidermal structures marked by GFP::FYVE.




Figure 2. MAGU-2 functions in the epidermis to regulate cholinergic synapse density. Representative images of cholinergic synapses labeled by ELKS-1::Cerulean in wild type (a), *magu-2(gk218)* (b), and *magu-2(ok1059)* (c) animals. Scale bars are 5 μ m. (d) Quantification of synapses from the indicated genotypes of animals expressing ELKS-1::Cerulean in cholinergic neurons. Gray dots represent individual animals; the black lines indicate the means. * p < 0.05; ** p < 0.01; ns: not significant. (e) Quantification of paralysis over time in the presence of 1 mM levamisole; * p < 0.05. Representative images of cholinergic synapses labeled by ELKS-1::Cerulean in wild type (f), *magu-2(gk218)* (g), and *magu-2(gk218); MAGU-2::GFP* (h), *magu-2(gk218); Neuronal::MAGU-2* (i), *magu-2(gk218); Epidermal::MAGU-2* (j), *magu-2(gk218); Muscle::MAGU-2* (k), and *magu-2(gk218); Epidermal::mpp5* (l) animals. Scale bars are 5 μ m. (m) Quantification of cholinergic synapses labeled by ELKS-1::Cerulean in wild type or *magu-2(0)* animals. Transgenic expression of MAGU-2 was achieved using its endogenous promoter, *rgef-1* promoter for neurons, *col-10* promoter for epidermis, or *myo-3* promoter for muscle. Mouse *mpp5* cDNA was expressed using the *col-10* promoter. Gray dots represent individual animals; the black lines indicate the means; * p < 0.05; *** p < 0.001.

FYVE domains associate with the membranes of endosomes and phagosomes in developing embryos of *C. elegans* (Yu, Lu, & Zhou, 2008). We have reported that both fluorescently-labeled FYVE domain and CED-1 expressed in the epidermis show colocalization with puncta labeled for presynaptic proteins (Cherra & Jin, 2016). Here, we analyzed the colocalization between GFP::FYVE structures in the epidermis and cholinergic synapses labeled by the presynaptic marker, mCherry::RAB-3. We observed that wild type animals displayed ~4 colocalization events per 100 micrometers where GFP::FYVE-labeled structures were associated with mCherry::RAB-3 expressed by cholinergic motor neurons (Figure 3(a–b)). In *magu-2(gk218)* animals, we found a significant decrease in the number of phagosomes near cholinergic synapses, similar to that of *ced-1(e1735)* animals (Figure 3(a–b)). These results indicate that MAGU-2 regulates epidermal phagocytosis.

To further evaluate the phagocytosis of neuronal materials by the epidermis, we performed electron microscopy analysis using the mini singlet oxygen generator (miniSOG) to label specific tissues. When stimulated by blue light, miniSOG produces singlet oxygen, and in the presence of diaminobenzidine (DAB), miniSOG oxidizes DAB to generate electron-dense osmiophilic polymers within nanometers

of miniSOG localization (Shu *et al.*, 2011). We expressed miniSOG fused to ZIG-10 in cholinergic neurons to determine if cholinergic membranes were engulfed by the epidermis. On electron micrographs of the nerve cord, miniSOG::ZIG-10 signals were detected in a subset of neuronal processes, whose positions were consistent with being cholinergic axons and dendrites (Figure 3(c–d)). Importantly, in addition to the miniSOG signals found in the nerve cord, we also observed miniSOG-containing vesicles within the epidermis (Figure 3(c)). This suggested that the epidermis engulfs portions of cholinergic neuronal membranes. As further support for this idea, we analyzed cholinergic-driven miniSOG::ZIG-10 in the phagocytic receptor *ced-1(e1735)* animals, which are deficient in executing phagocytosis (Hedgecock, Sulston, & Thomson, 1983; Zhou, Hartwieg, & Horvitz, 2001). In over 300 EM sections from two wild type animals, we found more than twenty miniSOG-labeled vesicles derived from cholinergic neurons in the epidermis. In the same number of sections from three *ced-1(e1735)* animals, we only observed miniSOG-labeled cholinergic axons or dendrites (Figure 3(d)). Thus, these data from miniSOG-mediated correlated-light-electron-microscopy analyses show that neuronal membranes are transferred to epidermis through the phagocytosis pathway.




Figure 3. MAGU-2 modulates epidermal phagocytosis of neuronal membranes. (a) Representative single, 0.5 μm-plane fluorescence images of wild type, *magu-2(gk218)*, and *ced-1(e1735)* animals expressing GFP::FYVE to label epidermal phagosomes and mCherry::RAB-3 to label cholinergic synapses. Arrows indicate GFP::FYVE spots that colocalize with RAB-3::mCherry. Scale bars are 5 μm. (b) Quantification of colocalization between epidermal phagosomes and cholinergic synapses. Gray dots represent individual animals; the black lines indicate the means; *** <0.001 . (c-d) Representative EM images of the ventral nerve cord in animals expressing miniSOG::ZIG-10 in cholinergic neurons. Scale bars are 500 nm. Cholinergic miniSOG-labeled structures are indicated by white arrows. Green shading indicates the muscle, blue indicates the epidermis, and yellow indicates the nerve cord. (c) Representative image of the ventral nerve cord flanked by muscle and epidermis. Cholinergic membranous material labeled by miniSOG::ZIG-10 is visible in the nerve cord and inside the epidermis in a wild type animal. (d) Representative image of the ventral cord in *ced-1(e1735)* animals, which show no miniSOG labeled structures inside the epidermis.

ZIG-10 localization depends on *magu-2*

MAGUKs regulate localization and surface expression of ion channels (El-Husseini *et al.*, 2000), glutamate receptors (Mi *et al.*, 2004), and N-cadherins (Wang *et al.*, 2014). Mutations in the extracellular Ig domain of ZIG-10 or RNAi-mediated knockdown displayed a similar phenotype to the *zig-10(tm6127)* mutation, suggesting that extracellular interactions and cell surface expression levels of ZIG-10 were essential to its function (Cherra & Jin, 2016). We hypothesized that MAGU-2 may regulate ZIG-10 expression or localization, similar to the interactions between cell adhesion proteins and other MAGUKs. To test this possibility, we generated a single-copy insertion of a functional GFP::ZIG-10 (Supplemental Figure 3). We then co-stained for GFP::ZIG-10 and the presynaptic active zone protein, UNC-10 in wild type and *magu-2(gk218)* animals harboring the GFP::ZIG-10 transgene. In *magu-2(gk218)* animals, there was a significant decrease in GFP::ZIG-10 near UNC-10-

labeled synapses (Figure 4(a,b)). However, there was no change in GFP::ZIG-10 intensity in neuronal cell bodies (Figure 4(c,d)). These data suggest that MAGU-2 regulates ZIG-10 localization near synapses but does not alter ZIG-10 expression levels.

Discussion

The regulation of synaptic connections plays an essential role in promoting circuit function and robust signaling. Cell-cell signaling is a common mechanism to modulate synapse formation and maintenance. In mammals, *Drosophila*, and *C. elegans* non-neuronal cells play essential roles in controlling synapse number through multiple mechanisms (Allen, 2013; Cherra & Jin, 2015; Corty & Freeman, 2013). Phagocytosis-mediated synapse pruning requires cell-cell interactions that enable glia or other non-neuronal cells to remove synapses (Cherra & Jin, 2016; Chung *et al.*, 2013; MacDonald *et al.*, 2006). Here, we have presented a new approach for investigating cellular interactions by electron microscopy, using genetically encoded miniSOG enzyme that enables the labeling of specific proteins or cellular compartments for electron microscopic analysis (Shu *et al.*, 2011). miniSOG can be expressed in a tissue-specific or temporal manner to enable the analysis of discrete interactions, such as between the epidermis and neurons. This approach provides a complementary method for immuno-EM analysis of protein localization or cell-cell interactions, including the analysis of phagocytosis.

Within the nervous system, phagocytosis by glia or epidermal cells plays multiple roles in removing dead cells or debris, degrading axons after injury, and pruning axons or synapses to modulate neural circuits (Awasaki *et al.*, 2006; Cherra & Jin, 2016; MacDonald *et al.*, 2006; Rasmussen, Sack, Martin, & Sagasti, 2015; Stevens *et al.*, 2007). While the removal of apoptotic cells by phagocytes has been widely studied, the recognition and engulfment of synapses is not well understood. In addition to phagocyte-corpse interactions mediated by a phagocytosis receptor, CED-1/Draper/Megf, other cell surface proteins further modulate the phagocytosis process, such as integrin and the immunoglobulin domain superfamily member, ZIG-10 (Albert, Kim, & Birge, 2000; Cherra & Jin, 2016). Here, we uncovered a role for the Pals1/Mpp5 homolog, MAGU-2, in regulating non-neuronal phagocytosis and synapse density at the neuromuscular junction. MPP members establish cell polarity, participate in axon sorting, and reduce or prevent neuronal degeneration through the maintenance of cell-cell interactions (Cho *et al.*, 2012; van Rossum *et al.*, 2006; Zollinger *et al.*, 2015). Our data suggest that cell polarity and MPPs may also modulate synaptic connectivity.

We propose that MAGU-2 functions in the epidermis to maintain cellular interactions between the epidermis and neurons by regulating the localization of ZIG-10. In the presence of MAGU-2, ZIG-10 localizes near synapses and enables the execution of phagocytosis to reduce the number of cholinergic synapses at the neuromuscular junction. Since MPPs are expressed in brain regions where glial-mediated




Figure 4. MAGU-2 regulates ZIG-10 localization. (a) Representative single, 0.5 μ m-plane fluorescence images of the dorsal nerve cord in *zig-10(tm6127)* and *zig-10(tm6127); magu-2(gk218)* animals expressing a single-copy transgene of GFP::ZIG-10. Animals were stained for UNC-10 to label synapses and GFP to identify ZIG-10 localization. Scale bars are 5 μ m. (b) Quantification of fluorescence intensity of GFP::ZIG-10 near synapses. Gray dots represent individual animals; the black lines indicate the means; *** p < 0.001. (c) Representative single, 0.5 μ m-plane fluorescence images of neuronal cell bodies from *zig-10(tm6127)* and *zig-10(tm6127); magu-2(gk218)* animals stained for GFP::ZIG-10. Scale bars are 5 μ m. White arrowheads indicate cell bodies. (d) Quantification of fluorescence intensity of GFP::ZIG-10 in neuronal cell bodies. Gray dots represent individual animals; the black lines indicate the means; ns: not significant.

synaptic pruning occurs (Clarke *et al.*, 2018; Zhang *et al.*, 2014), this raises the possibility that a conserved mechanism involving MPPs and phagocytosis may underlie synaptic pruning in mammals. Together these results provide new insights into how the phagocytosis pathway modulates synaptic connectivity.

Materials and methods

Strains and transgenes

All strains were maintained at 20 °C as previously described (Brenner, 1974). Transgenic animals were created by micro-injection as previously described (Mello, Kramer, Stinchcomb, & Ambros, 1991). Single-copy insertion of GFP::ZIG-10 at *cxTi10882* on chromosome IV was generated using CRISPR/Cas9 and modified plasmids (Frokjaer-Jensen, Davis, Ailion, & Jorgensen, 2012; Takayanagi-Kiya, Zhou, & Jin, 2016). For further strain and transgene information, see Table S1 and Supplementary Figure 3.

mRNA Analysis for *magu-2*

To determine if *magu-2* produced both predicted isoforms, we isolated RNA from mixed stage worms using Trizol according to manufacturer's specifications. SuperScript First Strand Synthesis kit (Invitrogen) was used to generate cDNA. We performed RT-PCR using primers: 5'-TCGATACCAACAGGCACTGT CC-3' as *magu-2a* forward, 5'-CATTGCTGCAACATCTGG ACC-3' as *magu-2b* forward, and 5'-TTTCTTCAACTT CAGCGAGTGG-3' as the common *magu-2* reverse primer.

DNA plasmid construction

MAGU-2::GFP was constructed by PCR amplifying the 4 kb sequence upstream of *magu-2 b* as its promoter and cloning it into pCR8 (Invitrogen) to generate pCZGY3355. The genomic *magu-2* coding sequence was amplified by PCR from

wild type worms and Gibson assembly was used to fuse GFP to the C-terminus of *magu-2* to generate MAGU-2::GFP (pCZGY3358). For tissue-specific rescue constructs, *magu-2* cDNA was cloned from wild type animals using SuperScript III First-Strand Synthesis System (Invitrogen). The splice leader 1 sequence was used with gene-specific primers that recognized the longest predicted mRNA to generate *magu-2* cDNA, which then was cloned into pCR8 (Invitrogen) to produce pCZGY3350. Mouse mpp5 cDNA was amplified from a mouse brain cDNA library and cloned into pCR8 to produce pCZGY3359. Generation of tissue-specific rescue constructs was performed using LR reactions (Invitrogen) with destination vectors containing tissue-specific promoters: rgef-1, col-10, unc-17b, myo-3 (Altun-Gultekin *et al.*, 2001; Cherra, & Jin, 2016; Okkema *et al.*, 1993; Rand, 2007). For single-copy insertion, Gibson assembly was used to clone *Pzig-10-gfp::zig-10*, *Prps-0-hygromycin resistance gene*, and homology arms around the *cxTi10882* locus on chromosome IV (pCZGY3354). See Table S2 for more information.

Whole-mount immunocytochemistry

To visualize GFP::ZIG-10 at near endogenous levels, CRISPR technology was used to insert into chromosome IV a single-copy of GFP::ZIG-10, which encodes the genomic locus of ZIG-10 with GFP fused directly after the signal peptide. The localization of GFP::ZIG-10 (*juSi333*) in the *zig-10(tm6127)* background in the presence or absence of the *magu-2(gk218)* allele was analyzed. Adult animals were washed 3 times in M9, followed by one 30-min wash in water. Animals were then placed on poly-L-lysine coated slides for freeze-crack and fixation with methanol and acetone as previously described (Duerr *et al.*, 1999). Following fixation, slides were washed in PBST and blocked with 5% goat serum. Samples were then incubated overnight with antibodies against GFP (1:500, A1112, RRID:AB_10073917, Invitrogen) and against UNC-10 (1:50, RRID:AB_10570332, Developmental Studies Hybridoma Bank). Samples were washed and incubated with secondary antibodies: goat anti-

mouse Alexa 594 (RRID:AB_141372, Invitrogen) or goat anti-rabbit Alexa 488 (RRID:AB_143165, Invitrogen). After washing, samples were mounted with Vectashield mounting media (RRID:AB_2336789, Vector Labs).

MiniSOG photo-oxidation and electron microscopy

To investigate the cellular interactions between the epidermis and the nerve cord, we used miniSOG to label cholinergic neurons using tissue-specific promoters. miniSOG-labeled cellular membranes were visualized by EM after fixation and photo-oxidation. Adult animals were placed in 2% glutaraldehyde, 2% paraformaldehyde solution in 100 mM sodium cacodylate buffer. After the animals stopped moving, they were cut in half and incubated for 1 h on ice. The samples were washed in 100 mM sodium cacodylate buffer and then blocked in 50 mM glycine, 10 mM potassium cyanide, 20 mM aminotriazole in 100 mM sodium cacodylate buffer for 2 h on ice. The samples were placed in a MatTek culture dish containing ice-cold 2.5 mM oxygentated diaminobenzidine and 10 mM HCl in 100 mM sodium cacodylate buffer and illuminated with blue light using a Leica SPEII confocal microscope for 20 min. The samples were then washed in 100 mM sodium cacodylate buffer and post fixed overnight at 4 °C in 2% osmium tetroxide in 100 mM sodium cacodylate buffer. The following day samples were rinsed with ice-cold ddH₂O, dehydrated with ethanol and acetone, embedded in Durcupan, and baked for 3 days at 60 °C.

After embedding, 60-nm-thick serial sections were prepared for analysis. MiniSOG-labeled structures were identified by visual inspection of electron micrographs as vesicular structures appearing darker than background and that persist through at least two consecutive serial sections. For miniSOG::ZIG-10 expressed in cholinergic neurons, the number of miniSOG-labeled structures was counted in the nerve cord and in the epidermis by an individual blinded to the sample genotypes.

Fluorescent microscopy

All images were captured at 63 \times magnification using an LSM710 confocal microscope (Zeiss) using identical settings for each fluorescent protein marker. L4 animals were immobilized in M9 buffer by rolling animals such that their dorsal or ventral surface contacted the coverslip. All image analysis was performed using the Fiji distribution of NIH ImageJ.

For synapse number, dorsal synapses posterior to the vulva were imaged, and a single z-plane of 0.5- μ m thickness was analyzed. Synapse density was determined using the ImageJ Analyze Particles function to count the number of synaptic puncta larger than 0.05 μ m² over a defined length of nerve cord.

For colocalization of phagosomes with dorsal synapses posterior to the vulva, confocal images were manually inspected for colocalization between GFP::FYVE and mCherry::RAB-3, as defined by particles containing pixels from both the red and green channels. The number of

colocalization events per length of nerve cord analyzed was then determined.

For quantifying ZIG-10 near the neuromuscular junctions, synapses were identified by eye as bright spots labeled by the UNC-10 antibody. GFP::ZIG-10 intensity was quantified by drawing a 2- μ m-wide rectangle adjacent to the UNC-10-labeled nerve cord using ImageJ. The integrated density from the green channel was then measured inside of the rectangle. The rectangle was then moved 10 μ m perpendicular to the nerve cord to measure the background integrated density. The background signal was subtracted from the GFP::ZIG-10 signal adjacent to the nerve cord to produce the measured intensity of ZIG-10 at the neuromuscular junction.

Levamisole sensitivity assay

L4 animals were moved to a fresh plate. The following day young adult animals were placed on plates containing 1 mM levamisole (Sigma). Animals were gently touched every 15 min for one hour to assess paralysis; animals that did not move after three touches were considered paralyzed.

Statistical analysis

All quantitative data are displayed as mean with individual data points represented as circles. For comparisons between two groups with normal distributions, Student's *t*-test was used. A Mann-Whitney test was used for comparisons between two groups that did not display Gaussian distribution. For comparisons between multiple groups, an ANOVA was used followed by post-hoc *t*-tests using a Bonferroni correction for multiple comparisons. A *p* values < 0.05 was considered statistically significant. Power analysis was performed to ensure that data was collected from a large enough sample size to provide a beta error ≤ 0.2 .

Acknowledgements

We thank the members of the Jin laboratory for constructive comments. We thank J.S. Dittman for reporter lines. We appreciate Wormbase for genetic and genomic information.

Author contributions

S.J. Cherra and Y. Jin conceived and designed the experiments, interpreted results, and wrote the manuscript. S.J. Cherra also performed experiments and analyzed data. A. Goncharov performed electron microscopy experiments and analyzed data. D. Boassa and M. Ellisman helped develop miniSOG stimulation and detection protocol.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

Some strains were provided by National BioResource Project (Dr. S. Mitani) and the *Caenorhabditis* Genetics Center, which is funded by

the National Institutes of Health Office of Research Infrastructure Programs [P40-OD010440]. The electron micrographs were taken in the UCSD Cellular and Molecular Medicine Electron microscopy core facility, which is supported in part by National Institutes of Health Award [S10-OD023527]. This work was supported by grants from the National Institutes of Health to S.J.C. [K99-NS097638], D.B. [R01-GM086197], M.E. [P41-GM103412], and Y.J. [R01 and R37-NS035546].

ORCID

Salvatore J. Cherra  <http://orcid.org/0000-0003-1581-9150>
Yishi Jin  <http://orcid.org/0000-0002-9371-9860>

References

Albert, M.L., Kim, J.I., & Birge, R.B. (2000). alphavbeta5 integrin recruits the CrkII-Dock180-rac1 complex for phagocytosis of apoptotic cells. *Nature Cell Biology*, 2, 899–905. doi:10.1038/35046549

Allen, N.J. (2013). Role of glia in developmental synapse formation. *Current Opinion in Neurobiology*, 23, 1027–1033. doi:10.1016/j.conb.2013.06.004

Allen, N.J., & Eroglu, C. (2017). Cell biology of astrocyte-synapse interactions. *Neuron*, 96, 697–708. doi:10.1016/j.neuron.2017.09.056

Altun-Gultekin, Z., Andachi, Y., Tsalik, E.L., Pilgrim, D., Kohara, Y., & Hobert, O. (2001). A regulatory cascade of three homeobox genes, ceh-10, ttx-3 and ceh-23, controls cell fate specification of a defined interneuron class in *C. elegans*. *Development (Cambridge, England)*, 128, 1951–1969.

Awasaki, T., Tatsumi, R., Takahashi, K., Arai, K., Nakanishi, Y., Ueda, R., & Ito, K. (2006). Essential role of the apoptotic cell engulfment genes draper and ced-6 in programmed axon pruning during Drosophila metamorphosis. *Neuron*, 50, 855–867. doi:10.1016/j.neuron.2006.04.027

Bangs, P., Franc, N., & White, K. (2000). Molecular mechanisms of cell death and phagocytosis in Drosophila. *Cell Death & Differentiation*, 7, 1027–1034. doi:10.1038/sj.cdd.4400754

Brenner, S. (1974). The genetics of *Caenorhabditis elegans*. *Genetics*, 77, 71–94.

Cao, J., Packer, J.S., Ramani, V., Cusanovich, D.A., Huynh, C., Daza, R., ... Shendure, J. (2017). Comprehensive single-cell transcriptional profiling of a multicellular organism. *Science*, 357, 661–667. doi:10.1126/science.aam8940

Cherra, S.J., 3rd, & Jin, Y. (2015). Advances in synapse formation: Forging connections in the worm. *Wiley Interdisciplinary Reviews: Developmental Biology*, 4, 85–97. doi:10.1002/wdev.165

Cherra, S.J., 3rd, & Jin, Y. (2016). A two-immunoglobulin-domain transmembrane protein mediates an epidermal-neuronal interaction to maintain synapse density. *Neuron*, 89, 325–336. doi:10.1016/j.neuron.2015.12.024

Cho, S.H., Kim, J.Y., Simons, D.L., Song, J.Y., Le, J.H., Swindell, E.C., ... Kim, S. (2012). Genetic ablation of Pals1 in retinal progenitor cells models the retinal pathology of Leber congenital amaurosis. *Human Molecular Genetics*, 21, 2663–2676. doi:10.1093/hmg/dds091

Chung, W.S., Allen, N.J., & Eroglu, C. (2015). Astrocytes control synapse formation, function, and elimination. *Cold Spring Harbor Perspectives in Biology*, 7, a020370. doi:10.1101/cshperspect.a020370

Chung, W.S., Clarke, L.E., Wang, G.X., Stafford, B.K., Sher, A., Chakraborty, C., ... Barres, B.A. (2013). Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways. *Nature*, 504, 394–400. doi:10.1038/nature12776

Clarke, L.E., Liddelow, S.A., Chakraborty, C., Munch, A.E., Heiman, M., & Barres, B.A. (2018). Normal aging induces A1-like astrocyte reactivity. *Proceedings of the National Academy of Sciences of the United States of America*, 115, E1896–E1905. doi:10.1073/pnas.1800165115

Corty, M.M., & Freeman, M.R. (2013). Cell biology in neuroscience: Architects in neural circuit design: glia control neuron numbers and connectivity. *Journal of Cell Biology*, 203, 395–405. doi:10.1083/jcb.201306099

de Wit, J., & Ghosh, A. (2016). Specification of synaptic connectivity by cell surface interactions. *Nature Reviews Neuroscience*, 17, 4–35. doi:10.1038/nrn.2015.3

Duerr, J.S., Frisby, D.L., Gaskin, J., Duke, A., Asermely, K., Huddleston, D., ... Rand, J.B. (1999). The cat-1 gene of *Caenorhabditis elegans* encodes a vesicular monoamine transporter required for specific monoamine-dependent behaviors. *The Journal of Neuroscience*, 19, 72–84. doi:10.1523/JNEUROSCI.19-01-00072.1999

El-Husseini, A.E., Topinka, J.R., Lehrer-Graiwer, J.E., Firestein, B.L., Craven, S.E., Aoki, C., & Bredt, D.S. (2000). Ion channel clustering by membrane-associated guanylate kinases. Differential regulation by N-terminal lipid and metal binding motifs. *Journal of Biological Chemistry*, 275, 23904–23910. doi:10.1074/jbc.M909919199

Frokjaer-Jensen, C., Davis, M.W., Ailion, M., & Jorgensen, E.M. (2012). Improved Mos1-mediated transgenesis in *C. elegans*. *Nature Methods*, 9, 117–118. doi:10.1038/nmeth.1865

Fuentes-Medel, Y., Logan, M.A., Ashley, J., Ataman, B., Budnik, V., & Freeman, M.R. (2009). Glia and muscle sculpt neuromuscular arbors by engulfing destabilized synaptic boutons and shed presynaptic debris. *PLoS Biology*, 7, e1000184. doi:10.1371/journal.pbio.1000184

Hedgecock, E.M., Sulston, J.E., & Thomson, J.N. (1983). Mutations affecting programmed cell deaths in the nematode *Caenorhabditis elegans*. *Science*, 220, 1277–1279. doi:10.1126/science.6857247

Hobert, O. (2013). The neuronal genome of *Caenorhabditis elegans*. *WormBook*, Aug 13:1–106. doi:10.1895/wormbook.1.161.1

Jospin, M., Qi, Y.B., Stawicki, T.M., Boulin, T., Schuske, K.R., Horvitz, H.R., ... Jin, Y. (2009). A neuronal acetylcholine receptor regulates the balance of muscle excitation and inhibition in *Caenorhabditis elegans*. *PLoS Biology*, 7, e1000265. doi:10.1371/journal.pbio.1000265

Lee, R.Y.N., Howe, K.L., Harris, T.W., Arnaboldi, V., Cain, S., Chan, J., ... Sternberg, P.W. (2018). WormBase 2017: molting into a new stage. *Nucleic Acids Research*, 46, D869–D874. doi:10.1093/nar/gkx998

MacDonald, J.M., Beach, M.G., Porpiglia, E., Sheehan, A.E., Watts, R.J., & Freeman, M.R. (2006). The Drosophila cell corpse engulfment receptor Draper mediates glial clearance of severed axons. *Neuron*, 50, 869–881. doi:10.1016/j.neuron.2006.04.028

Mangahas, P.M., & Zhou, Z. (2005). Clearance of apoptotic cells in *Caenorhabditis elegans*. *Seminars in Cell and Developmental Biology*, 16, 295–306. doi:10.1016/j.semcdb.2004.12.005

Mello, C.C., Kramer, J.M., Stinchcomb, D., & Ambros, V. (1991). Efficient gene transfer in *C. elegans*: Extrachromosomal maintenance and integration of transforming sequences. *The EMBO Journal*, 10, 3959–3970. doi:10.1002/j.1460-2075.1991.tb04966.x

Mi, R., Sia, G.M., Rosen, K., Tang, X., Moghekar, A., Black, J.L., McEnery, M., Huganir, R.L., & O'Brien, R.J. (2004). AMPA receptor-dependent clustering of synaptic NMDA receptors is mediated by Stargazin and NR2A/B in spinal neurons and hippocampal interneurons. *Neuron*, 44, 335–349. doi:10.1016/j.neuron.2004.09.029

Olkema, P.G., Harrison, S.W., Plunger, V., Aryana, A., & Fire, A. (1993). Sequence requirements for myosin gene expression and regulation in *Caenorhabditis elegans*. *Genetics*, 135, 385–404.

Ozcelik, M., Cotter, L., Jacob, C., Pereira, J.A., Relvas, J.B., Suter, U., & Tricaud, N. (2010). Pals1 is a major regulator of the epithelial-like polarization and the extension of the myelin sheath in peripheral nerves. *Journal of Neuroscience*, 30, 4120–4131. doi:10.1523/JNEUROSCI.5185-09.2010

Park, B., Alves, C.H., Lundvig, D.M., Tanimoto, N., Beck, S.C., Huber, G., ... Wijnholds, J. (2011). PALS1 is essential for retinal pigment epithelium structure and neural retina stratification. *Journal of Neuroscience*, 31, 17230–17241. doi:10.1523/JNEUROSCI.4430-11.2011

Park, J.Y., Hughes, L.J., Moon, U.Y., Park, R., Kim, S.B., Tran, K., ... Kim, S. (2016). The apical complex protein Pals1 is required to maintain cerebellar progenitor cells in a proliferative state. *Development*, 143, 133–146. doi:10.1242/dev.124180

Rand, J.B. (2007). Acetylcholine. *WormBook*, Jan 30; 1–21.

Rasmussen, J.P., Sack, G.S., Martin, S.M., & Sagasti, A. (2015). Vertebrate epidermal cells are broad-specificity phagocytes that clear sensory axon debris. *The Journal of Neuroscience*, 35, 559–570. doi: [10.1523/JNEUROSCI.3613-14.2015](https://doi.org/10.1523/JNEUROSCI.3613-14.2015)

Shu, X., Lev-Ram, V., Deerinck, T.J., Qi, Y., Ramko, E.B., Davidson, M.W., ... Tsien, R.Y. (2011). A genetically encoded tag for correlated light and electron microscopy of intact cells, tissues, and organisms. *PLoS Biology*, 9, e1001041. doi: [10.1371/journal.pbio.1001041](https://doi.org/10.1371/journal.pbio.1001041)

Stevens, B., Allen, N.J., Vazquez, L.E., Howell, G.R., Christopherson, K.S., Nouri, N., ... Barres, B.A. (2007). The classical complement cascade mediates CNS synapse elimination. *Cell*, 131, 1164–1178. doi: [10.1016/j.cell.2007.10.036](https://doi.org/10.1016/j.cell.2007.10.036)

Sudhof, T.C. (2018). Towards an understanding of synapse formation. *Neuron*, 100, 276–293. doi: [10.1016/j.neuron.2018.09.040](https://doi.org/10.1016/j.neuron.2018.09.040)

Takanayagi-Kiya, S., Zhou, K., & Jin, Y. (2016). Release-dependent feedback inhibition by a presynaptically localized ligand-gated anion channel. *Elife*, 5, pii: e21734. doi: [10.7554/eLife.21734](https://doi.org/10.7554/eLife.21734)

Tepass, U. (2012). The apical polarity protein network in Drosophila epithelial cells: Regulation of polarity, junctions, morphogenesis, cell growth, and survival. *Annual Review of Cell and Developmental Biology*, 28, 655–685. doi: [10.1146/annurev-cellbio-092910-154033](https://doi.org/10.1146/annurev-cellbio-092910-154033)

Thierry-Mieg, D., & Thierry-Mieg, J. (2006). AceView: A comprehensive cDNA-supported gene and transcripts annotation. *Genome Biology*, 7, S12–S14. doi: [10.1186/gb-2006-7-s1-s12](https://doi.org/10.1186/gb-2006-7-s1-s12)

van Rossum, A.G., Aartsen, W.M., Meuleman, J., Klooster, J., Malysheva, A., Versteeg, I., ... Wijnholds, J. (2006). Pals1/Mpp5 is required for correct localization of Crb1 at the subapical region in polarized Muller glia cells. *Human Molecular Genetics*, 15, 2659–2672. doi: [10.1093/hmg/ddl194](https://doi.org/10.1093/hmg/ddl194)

Wang, S.H., Celic, I., Choi, S.Y., Riccomagno, M., Wang, Q., Sun, L.O., ... Kolodkin, A.L. (2014). Dlg5 regulates dendritic spine formation and synaptogenesis by controlling subcellular N-cadherin localization. *The Journal of Neuroscience*, 34, 12745–12761. doi: [10.1523/JNEUROSCI.1280-14.2014](https://doi.org/10.1523/JNEUROSCI.1280-14.2014)

Yu, X., Lu, N., & Zhou, Z. (2008). Phagocytic receptor CED-1 initiates a signaling pathway for degrading engulfed apoptotic cells. *PLoS Biology*, 6, e61. doi: [10.1371/journal.pbio.0060061](https://doi.org/10.1371/journal.pbio.0060061)

Zhang, Y., Chen, K., Sloan, S.A., Bennett, M.L., Scholze, A.R., O'Keeffe, S., ... Wu, J.Q. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *The Journal of Neuroscience*, 34, 11929–11947. doi: [10.1523/JNEUROSCI.1860-14.2014](https://doi.org/10.1523/JNEUROSCI.1860-14.2014)

Zhou, Z., Hartwieg, E., & Horvitz, H.R. (2001). CED-1 is a transmembrane receptor that mediates cell corpse engulfment in *C. elegans*. *Cell*, 104, 43–56. doi: [10.1016/S0092-8674\(01\)00190-8](https://doi.org/10.1016/S0092-8674(01)00190-8)

Zhu, J., Shang, Y., & Zhang, M. (2016). Mechanistic basis of MAGUK-organized complexes in synaptic development and signalling. *Nature Reviews Neuroscience*, 17, 209–223. doi: [10.1038/nrn.2016.18](https://doi.org/10.1038/nrn.2016.18)

Zollinger, D.R., Chang, K.J., Baalman, K., Kim, S., & Rasband, M.N. (2015). The polarity protein Pals1 regulates radial sorting of axons. *The Journal of Neuroscience*, 35, 10474–10484. doi: [10.1523/JNEUROSCI.1593-15.2015](https://doi.org/10.1523/JNEUROSCI.1593-15.2015)